



planetmath.org

Math for the people, by the people.

Lie algebroids

Canonical name	LieAlgebroids
Date of creation	2013-03-22 18:13:47
Last modified on	2013-03-22 18:13:47
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	25
Author	bci1 (20947)
Entry type	Topic
Classification	msc 81P05
Classification	msc 81R15
Classification	msc 81R10
Classification	msc 81R50
Classification	msc 81R05
Synonym	extended quantum symmetry structures
Synonym	generalized double algebras
Related topic	HamiltonianAlgebroids
Related topic	TangentBundle
Related topic	HamiltonianOperatorOfAQuantumSystem
Related topic	Algebroids
Related topic	LieSuperalgebra3
Defines	Lie algebroid
Defines	anchor
Defines	bundle map

## 0.1 Topic on Lie algebroids

This is a topic entry on Lie algebroids that focuses on their quantum applications and extensions of current algebraic theories.

*Lie algebroids* generalize *Lie algebras*, and in certain quantum systems they represent *extended quantum (algebroid) symmetries*. One can think of a *Lie algebroid* as generalizing the idea of a tangent bundle where the tangent space at a point is effectively the equivalence class of curves meeting at that point (thus suggesting a groupoid approach), as well as serving as a site on which to study infinitesimal geometry (see, for example, ref. [?]). The formal definition of a *Lie algebroid* is presented next.

**Definition 0.1** Let  $M$  be a manifold and let  $\mathfrak{X}(M)$  denote the set of vector fields on  $M$ . Then, a *Lie algebroid* over  $M$  consists of a *vector bundle*  $E \rightarrow M$ , equipped with a *Lie bracket*  $[\cdot, \cdot]$  on the space of sections  $\gamma(E)$ , and a *bundle map*  $\Upsilon : E \rightarrow TM$ , usually called the *anchor*. Furthermore, there is an induced map  $\Upsilon : \gamma(E) \rightarrow \mathfrak{X}(M)$ , which is required to be a map of Lie algebras, such that given sections  $\alpha, \beta \in \gamma(E)$  and a differentiable function  $f$ , the following Leibniz rule is satisfied :

$$[\alpha, f\beta] = f[\alpha, \beta] + (\Upsilon(\alpha))\beta . \quad (0.1)$$

**Example 0.1.** A typical example of a Lie algebroid is obtained when  $M$  is a Poisson manifold and  $E = T^*M$ , that is  $E$  is the cotangent bundle of  $M$ .

Now suppose we have a Lie groupoid  $G$ :

$$r, s : G \overset{r}{\underset{s}{\rightrightarrows}} G^{(0)} = M . \quad (0.2)$$

There is an associated Lie algebroid  $\mathcal{A} = \mathcal{A}(G)$ , which in the guise of a vector bundle, it is the restriction to  $M$  of the bundle of tangent vectors along the fibers of  $s$  (ie. the  $s$ -vertical vector fields). Also, the space of sections  $\gamma(\mathcal{A})$  can be identified with the space of  $s$ -vertical, right-invariant vector fields  $\mathfrak{X}_{inv}^s(G)$  which can be seen to be closed under  $[\cdot, \cdot]$ , and the latter induces a bracket operation on  $\gamma(\mathcal{A})$  thus turning  $\mathcal{A}$  into a Lie algebroid. Subsequently, a Lie algebroid  $\mathcal{A}$  is integrable if there exists a Lie groupoid  $G$  inducing  $\mathcal{A}$ .

**Remark 0.1.** Unlike Lie algebras that can be integrated to corresponding Lie groups, not all *Lie algebroids* are ‘smoothly integrable’ to Lie groupoids; the subset of Lie groupoids that have corresponding Lie algebroids are sometimes called ‘*Weinstein groupoids*’.

Note also the relation of the Lie algebroids to Hamiltonian algebroids, also concerning recent developments in (relativistic) quantum gravity theories.

## References

- [1] K. C. H. Mackenzie: *General Theory of Lie Groupoids and Lie Algebroids*, London Math. Soc. Lecture Notes Series, **213**, Cambridge University Press: Cambridge, UK (2005).