

planetmath.org

Math for the people, by the people.

organismic sets

Canonical name OrganismicSets
Date of creation 2013-03-22 18:11:20
Last modified on 2013-03-22 18:11:20

Owner bci1 (20947) Last modified by bci1 (20947)

Numerical id 28

Author bci1 (20947)

Entry type Topic

Classification msc 92C30 Classification msc 92B99 Classification msc 92B20 Classification msc 92B05

Synonym lattices or sets of organismic structures

Related topic OrganismicSet Related topic NicolasRashevsky

Related topic
Re

0.1 Organismic Set Theory and Relational Biology

Organismic sets (OS^n) were defined by Nicolas Rashevsky as simple settheoretical models of organization in living organisms at discrete integer or zero levels by means of sets of several distinct types or order beginning at the zeroth order, and having an upper limit as the fifth or sixth order of roganization. Thus, in the case of organismic sets of zero-th order, S_0 , the elements correspond to genes, and a concrete S_{0c} is defined as the set of all genes $[G_n]$ of a specific organism or organism type (understood as a stable biological species).

Alternatively, S_{0c} can be defined as a set representation of any organismic genome, G_O , consisting of the complete set of active, functional genes of an organism together with controlling genes/operons. The latter are then also considered together with inputs e_i from the environment, as well as their activities a_i and relations R_{ij} among organismic set elements (genes in the case of S_0), where i, j are indices in a countable, index set I. Thus, Rashevsky's organismic set (OS) theory is part of http://planetmath.org/AbstractRelationalBiologyabstrac relational biology, but unlike organismic networks, metabolic-replication systems and organismic supercategories, Rashevsky's OS are only endowed with the discrete topology, and are thus the simplest model whose only connectivities to organization are through the hierarchical lattice structure of the different types of OS, from G_O to (OS^n) , with n>1. At the next level of biological organization, cells are considered as first order organismic sets, S_1 , whereas multi-cellular organisms are modeled by organismic sets of second order, S_2 , whose 'elements' are the first order organismic sets, or cells, S_1 . Attempts were then made by Rashevsky to expand his theory of organismic sets to organizational models of human societies. Results from such studies of relations between sets were considered to be far more important than the numerical or quantitative aspects that play such important roles in physics and chemistry. A number of interesting results were obtained by means of standard (Boolean) logic predicates applied to organismic sets and their relations. Further details can be found in the publications listed below and the references cited therein. Subsequently, autopoietic theories have enlarged upon, and also extended, the application of organismic sets to biological and ecological systems.

References

- [1] Rashevsky, N.: 1965, The Representation of Organisms in Terms of Predicates, *Bulletin of Mathematical Biophysics* 27: 477-491.
- [2] Rashevsky, N.: 1969, Outline of a Unified Approach to Physics, Biology and Sociology., *Bulletin of Mathematical Biophysics* **31**: 159–198.