Foundations of Audio Signal Processing Assignment 1

Giulia Baldini, Luis Fernandes, Agustin Vargas November 2, 2018

Exercise 2.1

a.

$$2e^{\frac{\pi}{2}i}(1+i) = 2(\cos(\frac{\pi}{2}) + i\sin(\frac{\pi}{2}))(1+i)$$

$$= 2i(1+i)$$

$$= -2 + 2i$$

$$a = -2$$

$$b = 2$$

$$r = \sqrt{(-2)^2 + (2)^2}$$

$$= 2\sqrt{2}$$

Knowing that $a = \cos(\phi)$ and $b = \sin(\phi)$:

$$\cos(\phi) = \frac{-2}{2\sqrt{2}} = -\frac{\sqrt{2}}{2}$$
$$\sin(\phi) = \frac{2}{2\sqrt{2}} = \frac{\sqrt{2}}{2}$$
$$\phi = \frac{3}{4}\pi$$
$$z = 2\sqrt{2}e^{\frac{3}{4}\pi i}$$

b. Considering that $z=re^{\phi i},$ $\overline{z}=re^{-\phi i}$ and |z|=r $z\overline{z}=re^{\phi i}re^{-\phi i}$ $=r^2e^0 = |z|^2$

c.

$$\begin{split} \frac{1}{2i}(e^{ia} - e^{-ia}) &= \frac{1}{2i}(\cos(a) + i\sin(a) - (\cos(-a) + i\sin(-a))) \\ &= \frac{1}{2i}(\cos(a) + i\sin(a) - \cos(a) + i\sin(a)) \\ &= \frac{1}{2i}(2i\sin(a)) \end{split}$$

Exercise 2.2

a. For n = 4:

$$\begin{split} &\Omega_4^1 = \cos(\frac{\pi}{2}) + i\sin(\frac{\pi}{2}) = i \\ &\Omega_4^2 = \cos(\pi) + i\sin(\pi) = -1 \\ &\Omega_4^3 = \cos(3\frac{\pi}{2}) + i\sin(3\frac{\pi}{2}) = -i \\ &\Omega_4^4 = 1 \end{split}$$

So the fourth roots of unity are 1, -1, i, -i and, since 4 is the smallest integer for which $\Omega = i, -i$, then these are its primitives.




Figure 1: Illustration of all fourth roots of unity.

For n = 6:

$$\Omega_{6}^{1} = \cos(\frac{\pi}{3}) + i\sin(\frac{\pi}{3}) = \frac{1}{2} + \frac{\sqrt{3}}{2}i$$

$$\Omega_{6}^{2} = \cos(2\frac{\pi}{3}) + i\sin(2\frac{\pi}{3}) = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$$

$$\Omega_{6}^{3} = \cos(\pi) + i\sin(\pi) = -1$$

$$\Omega_{6}^{4} = \cos(4\frac{\pi}{3}) + i\sin(4\frac{\pi}{3}) = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$$

$$\Omega_{6}^{5} = \cos(5\frac{\pi}{3}) + i\sin(5\frac{\pi}{3}) = \frac{1}{2} - \frac{\sqrt{3}}{2}i$$

$$\Omega_{6}^{6} = 1$$

So the sixth roots of unity are $1, -1, \frac{1}{2} + \frac{\sqrt{3}}{2}i, \frac{1}{2} - \frac{\sqrt{3}}{2}i, -\frac{1}{2} - \frac{\sqrt{3}}{2}i, \frac{1}{2} - \frac{\sqrt{3}}{2}i$ and, since 4 is the smallest integer for which $\Omega = -\frac{1}{2} + \frac{\sqrt{3}}{2}i, \frac{1}{2} + \frac{\sqrt{3}}{2}i$, then these are its primitives.




Figure 2: Illustration of all sixth roots of unity.

b.

1 Matlab

a-c. The solutions can be found inside the code folder.

$$\cos(\alpha) = \frac{1}{2} \cdot (e^{ia} + e^{-ia})$$