

Dynamics of contagion and submerged infections during the second wave of the SARS-CoV-2 epidemic in the Italian regions

Giulia Cereda, Cecilia Viscardi, Luca Gherardini, Fabrizia Mealli, Michela Baccini

In this note, we report a description of the SARS-CoV-2 epidemic from August 01, 2020 to February 21, 2021 in the Italian regions. The results are obtained by using a SIRD model calibrated on the notified COVID19-related deaths (<https://github.com/pcm-dpc/COVID-19>). In particular, assuming an infection fatality rate of 0.78% and an average time from infection to recovery or death of 14 days, we explored the behavior of the contagion over time in terms of reproductive number $R_0(t)$ and number of circulating infected individuals. The number of circulating infections, when compared to the number of notified positives, may provide an evaluation of the submerged fraction of the contagion. Details on the methods can be found in [1].

NB. The estimation algorithm used for obtaining the results reported in this note is slightly different from the one described in [1], because it allows a positive constraint on the $R_0(t)$ curve.

In questa nota, riportiamo una descrizione dell'epidemia di SARS-CoV-2 nelle regioni italiane da August 01, 2020 fino al 21/02/2021. I risultati sono ottenuti tramite un modello compartimentale di tipo SIRD calibrato sui decessi COVID-19 (<https://github.com/pcm-dpc/COVID-19>). In particolare, assumendo un tasso di letalità dell'infezione pari a 0.78% ed un tempo medio tra infezione e ricovero di 14 giorni, esploriamo l'andamento del contagio nel tempo attraverso il numero di riproduzione $R_0(t)$ e il numero di infetti circolanti. Quest'ultimo, se confrontato con il numero di infetti notificati, fornisce una indicazione circa il contagio sommerso. Dettagli sul metodo si possono trovare in [1].

NB. L'algoritmo di stima utilizzato per ottenere i risultati riportati in questa nota è leggermente diverso da quello descritto in [1], in quanto permette di vincolare la curva di $R_0(t)$ a valori positivi.




Figure 1: Estimated $R_0(t)$ with pointwise 90% confidence bands from 01/08/2020 to 21/02/2021, by region.
Figura 1: Stima dell'andamento di $R_0(t)$ con bande di confidenza puntuali al 90% dal 01/08/2020 al 21/02/2021 per tutte le regioni italiane




Figure 2: Estimates and 90% confidence intervals of the number of infections (including the undetected ones) present in the Italian regions (from North to South) on 24/01/2021 and 14/02/2021 calculated over 1000 inhabitants. For the analysis, we assumed an infection fatality rate equal to 0.78%.

Figura 2: Stime e intervalli di confidenza al 90% del numero di infezioni ogni 1000 abitanti (incluse quelle non notificate) presenti nelle regioni italiane (da Nord a Sud) al 24/01/2021 and 14/02/2021. Nell'analisi, abbiamo assunto un tasso di letalità dell'infezione pari allo 0.78%.

Table 1: Estimates and confidence intervals of the number of infections (including the undetected ones) present in the Italian regions on 24/01/2021, 07/02/2021, 14/02/2021 calculated over 1000 inhabitants. For the analysis, we assumed an infection fatality rate equal to 0.0078.

Tabella 1: Stime e intervalli di confidenza del numero di contagiati (inclusi i sommersi) presenti nelle varie regioni italiane al 24/01/2021, 07/02/2021, 14/02/2021 calcolati su mille abitanti, e usando un tasso di infezione pari a 0.0078.

	January 24			February 07			February 14		
	prev	90% CI		prev	90% CI		prev	90% CI	
Abruzzo	11.8	9.7	13.4	11.2	9.2	13.5	9.3	7.7	12.5
Basilicata	7.1	5.0	8.9	6.4	4.2	8.7	5.6	3.5	9.0
Calabria	3.7	2.9	4.4	3.6	2.8	4.4	3.0	2.3	4.1
Campania	8.4	7.6	8.9	6.3	5.6	7.0	4.6	4.1	5.5
Emilia-Romagna	21.9	20.5	23.5	17.4	16.0	18.7	13.9	12.4	15.6
Friuli Venezia Giulia	36.1	32.5	38.5	29.4	26.8	32.8	21.9	19.9	26.7
Lazio	12.5	11.5	13.4	11.3	10.3	12.3	10.0	8.8	11.2
Liguria	18.1	15.8	19.5	15.5	13.7	17.5	11.5	10.5	14.3
Lombardia	11.0	10.3	11.4	9.3	8.8	10.0	7.1	6.6	8.0
Marche	14.8	12.8	16.8	12.7	10.9	14.5	11.5	9.2	14.2
Molise	16.2	12.6	20.4	18.0	14.4	23.2	17.6	12.7	25.3
Trentino	16.1	13.4	18.3	12.8	10.4	15.0	11.2	8.3	14.3
Piemonte	11.7	10.7	12.4	10.5	9.5	11.6	8.2	7.5	9.7
Puglia	11.7	10.6	12.8	12.9	11.4	14.0	11.8	10.2	13.3
Sardegna	7.2	5.8	8.4	6.4	4.9	7.6	6.8	4.7	8.4
Sicilia	13.0	11.9	13.8	10.0	9.2	10.9	7.3	6.7	8.6
Toscana	7.5	6.5	8.4	7.8	6.9	8.7	8.1	7.0	9.3
Umbria	12.2	9.9	14.5	18.2	15.1	21.9	21.2	17.0	25.0
Valle d'Aosta	8.1	4.4	13.0	6.6	3.3	10.2	7.8	3.3	13.7
Veneto	24.0	22.5	25.5	14.4	13.2	15.8	10.6	9.4	12.4

References

- [1] Cereda G, Viscardi C, Gherardini L, Mealli F, Baccini M. *Dynamics of contagion and submerged infections during the second wave of the SARS-CoV-2 epidemic in the Italian regions*. doi:10.21203/rs.3.rs-130717/v1