

# DDFV method for Navier-Stokes problem with outflow boundary conditions

Giulia Lissoni

Thesis supervisors: Thierry Goudon, Stella Krell



# Contents

## 1. Motivation

Outflow boundary conditions

## 2. Discrete Duality Finite Volume method

Notations and advantages


## 3. DDFV for Navier-Stokes problem

Theoretical results

Numerical results

# Motivation

# Outflow boundary conditions



- ❑  $\Gamma_1$ : Dirichlet boundary conditions
- ❑  $\Gamma_2$ : Outflow boundary conditions
- ❑  $\Gamma_0 = \partial\Omega \setminus (\Gamma_1 \cup \Gamma_2)$  : no slip boundary conditions

# Outflow boundary conditions

Find  $\mathbf{u} : \Omega \rightarrow \mathbb{R}^2$  and  $p : \Omega \rightarrow \mathbb{R}$  such that:

$$\left\{ \begin{array}{ll} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} - \operatorname{div}(\sigma(\mathbf{u}, p)) = 0; & \text{in } \Omega_T = \Omega \times [0, T] \\ \operatorname{div}(\mathbf{u}) = 0 & \text{in } \Omega_T, \\ \mathbf{u} = \mathbf{g}_1 & \text{on } \Gamma_1 \times (0, T), \\ \mathbf{u} = 0 & \text{on } \Gamma_0 \times (0, T), \\ \text{outflow boundary conditions} & \text{on } \Gamma_2 \times (0, T) \\ \mathbf{u}(0) = \mathbf{u}_{init} & \text{in } \Omega \end{array} \right.$$

with  $T > 0$ ,  $\Omega \subset \mathbb{R}^2$ ,  $\mathbf{u}_{init} \in (L^\infty(\Omega))^2$ ,  $\mathbf{g}_1 \in (H^{\frac{1}{2}}(\Omega))^2$ .

The stress tensor:  $\sigma(\mathbf{u}, p) = \frac{2}{\operatorname{Re}} \mathbf{D}\mathbf{u} - p \mathbf{I}$ , with  $\operatorname{Re} > 0$ .

The strain rate tensor:  $\mathbf{D}\mathbf{u} = \frac{(\nabla \mathbf{u} + {}^t \nabla \mathbf{u})}{2}$ .

# Outflow boundary conditions

We choose a test function  $\Psi$  in the space

$$V = \{\psi \in (H^1(\Omega))^2, \psi|_{\Gamma_1} = 0, \psi|_{\Gamma_0} = 0, \operatorname{div}(\psi) = 0\}.$$

The weak formulation reads:

$$\int_{\Omega} \partial_t \mathbf{u} \cdot \Psi + \int_{\Omega} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \Psi - \int_{\Omega} \operatorname{div}(\sigma(\mathbf{u}, p)) \cdot \Psi = 0$$

# Outflow boundary conditions

We choose a test function  $\Psi$  in the space

$$V = \{\psi \in (H^1(\Omega))^2, \psi|_{\Gamma_1} = 0, \psi|_{\Gamma_0} = 0, \operatorname{div}(\psi) = 0\}.$$

The weak formulation reads:

$$\begin{aligned} \int_{\Omega} \partial_t \mathbf{u} \cdot \Psi + \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \Psi - \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \Psi \cdot \mathbf{u} \\ - \int_{\Omega} \operatorname{div}(\sigma(\mathbf{u}, p)) \cdot \Psi = -\frac{1}{2} \int_{\Gamma_2} (\mathbf{u} \cdot \vec{\mathbf{n}}) \mathbf{u} \cdot \Psi \end{aligned}$$

# Outflow boundary conditions

We choose a test function  $\Psi$  in the space

$$V = \{\psi \in (H^1(\Omega))^2, \psi|_{\Gamma_1} = 0, \psi|_{\Gamma_0} = 0, \operatorname{div}(\psi) = 0\}.$$

The weak formulation reads:

$$\begin{aligned} \int_{\Omega} \partial_t \mathbf{u} \cdot \Psi + \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \Psi - \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \Psi \cdot \mathbf{u} \\ - \int_{\Omega} \operatorname{div} \left( \frac{2}{\operatorname{Re}} D\mathbf{u} - p \mathbf{Id} \right) \cdot \Psi = - \frac{1}{2} \int_{\Gamma_2} (\mathbf{u} \cdot \vec{\mathbf{n}}) \mathbf{u} \cdot \Psi \end{aligned}$$



# Outflow boundary conditions

We choose a test function  $\Psi$  in the space

$$V = \{\psi \in (H^1(\Omega))^2, \psi|_{\Gamma_1} = 0, \psi|_{\Gamma_0} = 0, \operatorname{div}(\psi) = 0\}.$$

The weak formulation reads:

$$\begin{aligned} \int_{\Omega} \partial_t \mathbf{u} \cdot \Psi + \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \Psi - \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \Psi \cdot \mathbf{u} \\ + \frac{2}{\operatorname{Re}} \int_{\Omega} \mathbf{D} \mathbf{u} : \mathbf{D} \Psi - \int_{\Gamma_2} \sigma(\mathbf{u}, p) \cdot \vec{\mathbf{n}} \cdot \Psi = -\frac{1}{2} \int_{\Gamma_2} (\mathbf{u} \cdot \vec{\mathbf{n}}) \mathbf{u} \cdot \Psi \end{aligned}$$

# Outflow boundary conditions

$$\begin{aligned} \int_{\Omega} \partial_t \mathbf{u} \cdot \Psi + \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \Psi - \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \Psi \cdot \mathbf{u} \\ + \frac{2}{\text{Re}} \int_{\Omega} \mathbf{D} \mathbf{u} : \mathbf{D} \Psi - \int_{\Gamma_2} \sigma(\mathbf{u}, p) \cdot \vec{\mathbf{n}} \cdot \Psi = -\frac{1}{2} \int_{\Gamma_2} (\mathbf{u} \cdot \vec{\mathbf{n}}) \mathbf{u} \cdot \Psi \end{aligned}$$

Let  $\mathbf{u}_{ref}, \sigma_{ref}$  be a reference flow. We can impose:

$$\sigma(\mathbf{u}, p) \cdot \vec{\mathbf{n}} = \frac{1}{2} \Theta(\mathbf{u} \cdot \vec{\mathbf{n}}) (\mathbf{u} - \mathbf{u}_{ref}) + \sigma_{ref} \cdot \vec{\mathbf{n}} \quad (1)$$

where

$$\Theta(a) = a, \quad \Theta(a) = -a^-, \quad \Theta(a) = |a|$$

with the notation  $a = a^+ - a^-$ .

(1): [Ch.-H. Bruneau and P. Fabrie, *Effective downstream boundary conditions for incompressible navier stokes equations*, 1994]

# Outflow boundary conditions

We choose:

$$\sigma(\mathbf{u}, p) \cdot \vec{\mathbf{n}} = -\frac{1}{2}(\mathbf{u} \cdot \vec{\mathbf{n}})^-(\mathbf{u} - \mathbf{u}_{ref}) + \sigma_{ref} \cdot \vec{\mathbf{n}}$$

so that the weak formulation becomes:

$$\begin{aligned} \int_{\Omega} \partial_t \mathbf{u} \cdot \Psi + \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \Psi - \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \Psi \cdot \mathbf{u} + \frac{1}{2} \int_{\Gamma_2} (\mathbf{u} \cdot \vec{\mathbf{n}})^+ (\mathbf{u} \cdot \Psi) \\ + \frac{2}{\text{Re}} \int_{\Omega} \mathbf{D}(\mathbf{u}) : \mathbf{D}(\Psi) = \frac{1}{2} \int_{\Gamma_2} (\mathbf{u} \cdot \vec{\mathbf{n}})^- (\mathbf{u}_{ref} \cdot \Psi) + \int_{\Gamma_2} \sigma_{ref} \cdot \vec{\mathbf{n}} \cdot \Psi \end{aligned}$$

# Discrete Duality Finite Volume method

# DDFV meshes

## Primal mesh


$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}} = (\mathbf{u}_K)_{K \in \mathfrak{M}}$$

## Dual mesh

$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}^*} = (\mathbf{u}_{K^*})_{K^* \in \mathfrak{M}^*}$$

## Diamond mesh

$$\rightsquigarrow \nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}}, p^{\mathfrak{D}}$$



Our unknowns are:

$$\mathbf{u}^{\mathfrak{T}} = (\mathbf{u}^{\mathfrak{M}}, \mathbf{u}^{\mathfrak{M}^*}) \text{ and } p^{\mathfrak{D}}$$

# DDFV meshes

## Primal mesh


$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}} = (\mathbf{u}_K)_{K \in \mathfrak{M}}$$

## Dual mesh

$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}^*} = (\mathbf{u}_{K^*})_{K^* \in \mathfrak{M}^*}$$

## Diamond mesh

$$\rightsquigarrow \nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}}, p^{\mathfrak{D}}$$



Our unknowns are:

$$\mathbf{u}^{\mathfrak{T}} = (\mathbf{u}^{\mathfrak{M}}, \mathbf{u}^{\mathfrak{M}^*}) \text{ and } p^{\mathfrak{D}}$$

# DDFV meshes

## Primal mesh


$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}} = (\mathbf{u}_k)_{k \in \mathfrak{M}}$$

## Dual mesh

$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}^*} = (\mathbf{u}_{k^*})_{k^* \in \mathfrak{M}^*}$$

## Diamond mesh

$$\rightsquigarrow \nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}}, p^{\mathfrak{D}}$$



Our unknowns are:

$$\mathbf{u}^{\mathfrak{T}} = (\mathbf{u}^{\mathfrak{M}}, \mathbf{u}^{\mathfrak{M}^*}) \text{ and } p^{\mathfrak{D}}$$

# DDFV meshes

## Primal mesh


$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}} = (\mathbf{u}_k)_{k \in \mathfrak{M}}$$

## Dual mesh

$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}^*} = (\mathbf{u}_{k^*})_{k^* \in \mathfrak{M}^*}$$

## Diamond mesh

$$\rightsquigarrow \nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}}, p^{\mathfrak{D}}$$



Our unknowns are:

$$\mathbf{u}^{\mathfrak{T}} = (\mathbf{u}^{\mathfrak{M}}, \mathbf{u}^{\mathfrak{M}^*}) \text{ and } p^{\mathfrak{D}}$$



# DDFV meshes

## Primal mesh


$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}} = (\mathbf{u}_k)_{k \in \mathfrak{M}}$$

## Dual mesh

$$\rightsquigarrow \mathbf{u}^{\mathfrak{M}^*} = (\mathbf{u}_{k^*})_{k^* \in \mathfrak{M}^*}$$

## Diamond mesh

$$\rightsquigarrow \nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}}, p^{\mathfrak{D}}$$



Our unknowns are:

$$\mathbf{u}^{\mathfrak{T}} = (\mathbf{u}^{\mathfrak{M}}, \mathbf{u}^{\mathfrak{M}^*}) \text{ and } p^{\mathfrak{D}}$$

# DDFV operators

## Discrete gradient

The operator  $\nabla^{\mathfrak{D}} : (\mathbb{R}^2)^{\mathfrak{T}} \mapsto (\mathcal{M}_2(\mathbb{R}))^{\mathfrak{D}}$  where

$$\nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}}(x_L - x_K) = \mathbf{u}_L - \mathbf{u}_K,$$


$$\nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}}(x_{L^*} - x_{K^*}) = \mathbf{u}_{L^*} - \mathbf{u}_{K^*}.$$

$$\nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}} = \frac{1}{2m_D} [m_{\sigma}(\mathbf{u}_L - \mathbf{u}_K) \otimes \vec{\mathbf{n}}_{\sigma K} + m_{\sigma^*}(\mathbf{u}_{L^*} - \mathbf{u}_{K^*}) \otimes \vec{\mathbf{n}}_{\sigma^* K^*}].$$

[Krell '11]

$$\rightsquigarrow \operatorname{div}^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}} = \operatorname{Tr}(\nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}}).$$

$$\rightsquigarrow \mathbf{D}^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}} = \frac{\nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}} + {}^t(\nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}})}{2}.$$



# DDFV operators

## Discrete divergence

$\mathbf{div}^{\mathfrak{T}} : \xi^{\mathfrak{D}} \in (\mathcal{M}_2(\mathbb{R}))^{\mathfrak{D}} \mapsto \mathbf{div}^{\mathfrak{T}} \xi^{\mathfrak{D}} \in (\mathbb{R}^2)^{\mathfrak{T}}$  where:

$$\mathbf{div}^{\mathbf{K}} \xi^{\mathfrak{D}} = \frac{1}{m_{\mathbf{K}}} \sum_{\sigma \subset \partial \mathbf{K}} m_{\sigma} \xi^{\mathfrak{D}} \vec{\mathbf{n}}_{\sigma \mathbf{K}}, \quad \forall \mathbf{K} \in \mathfrak{M}$$

$$\mathbf{div}^{\mathbf{K}^*} \xi^{\mathfrak{D}} = \frac{1}{m_{\mathbf{K}^*}} \sum_{\sigma^* \subset \partial \mathbf{K}^*} m_{\sigma^*} \xi^{\mathfrak{D}} \vec{\mathbf{n}}_{\sigma^* \mathbf{K}^*}, \quad \forall \mathbf{K}^* \in \mathfrak{M}^* \cup \partial \mathfrak{M}^*$$

## Discrete duality property

- ▶ On the continuous level:  $\int_{\Omega} \mathbf{div} \xi \cdot \mathbf{u} = - \int_{\Omega} \xi : \nabla \mathbf{u} + \int_{\partial \Omega} \xi \vec{\mathbf{n}} \cdot \mathbf{u}$
- ▶ On the discrete level:

$$[[\mathbf{div}^{\mathfrak{T}} \xi^{\mathfrak{D}}, \mathbf{u}^{\mathfrak{T}}]]_{\mathfrak{T}} = -(\xi^{\mathfrak{D}} : \nabla^{\mathfrak{D}} \mathbf{u}^{\mathfrak{T}})_{\mathfrak{D}} + (\gamma^{\mathfrak{D}}(\xi^{\mathfrak{D}}) \vec{\mathbf{n}}, \gamma^{\mathfrak{T}}(\mathbf{u}^{\mathfrak{T}}))_{\partial \Omega}$$

[S. Krell, *Stabilized DDFV schemes for the incompressible Navier-Stokes equations*, 2011]

# DDFV for Navier-Stokes problem

# Outflow boundary conditions

Find  $\mathbf{u} : \Omega \rightarrow \mathbb{R}^2$  and  $p : \Omega \rightarrow \mathbb{R}$  such that:

$$\left\{ \begin{array}{ll} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} - \operatorname{div}(\sigma(\mathbf{u}, p)) = 0; & \text{in } \Omega_T = \Omega \times [0, T] \\ \operatorname{div}(\mathbf{u}) = 0 & \text{in } \Omega_T, \\ \mathbf{u} = \mathbf{g}_1 & \text{on } \Gamma_1 \times (0, T), \\ \mathbf{u} = 0 & \text{on } \Gamma_0 \times (0, T), \\ \sigma(\mathbf{u}, p) \cdot \vec{\mathbf{n}} + \frac{1}{2}(\mathbf{u} \cdot \vec{\mathbf{n}})^-(\mathbf{u} - \mathbf{u}_{ref}) = \sigma_{ref} \cdot \vec{\mathbf{n}} & \text{on } \Gamma_2 \times (0, T) \\ \mathbf{u}(0) = \mathbf{u}_{init} & \text{in } \Omega \end{array} \right.$$

with  $T > 0$ ,  $\Omega \subset \mathbb{R}^2$ ,  $\mathbf{u}_{init} \in (L^\infty(\Omega))^2$ ,  $\mathbf{g}_1 \in (H^{\frac{1}{2}}(\Omega))^2$ .

The stress tensor:  $\sigma(\mathbf{u}, p) = \frac{2}{\operatorname{Re}} D\mathbf{u} - p \operatorname{Id}$ , with  $\operatorname{Re} > 0$ .

The strain rate tensor:  $D\mathbf{u} = \frac{(\nabla \mathbf{u} + {}^t \nabla \mathbf{u})}{2}$ .

# Nonlinear convection term $(\mathbf{u} \cdot \nabla)\mathbf{u}$

We construct:

$$\text{+} \quad m_K \mathbf{b}^K(\mathbf{u}^n, \mathbf{u}^{n+1}) \rightsquigarrow \int_K (\mathbf{u}^n \cdot \nabla) \mathbf{u}^{n+1} \text{ when } K \in \mathfrak{M},$$

$$\text{+} \quad m_{K^*} \mathbf{b}^{K^*}(\mathbf{u}^n, \mathbf{u}^{n+1}) \rightsquigarrow \int_{K^*} (\mathbf{u}^n \cdot \nabla) \mathbf{u}^{n+1} \text{ when } K^* \in \mathfrak{M}^* \cup \partial\mathfrak{M}_N^*.$$

We focus on the primal mesh and we observe:

$$\boxed{\int_K (\mathbf{u}^n \cdot \nabla) \mathbf{u}^{n+1} = \sum_{\sigma \subset \partial K} \int_{\sigma} (\mathbf{u}^n \cdot \vec{\mathbf{n}}_{\sigma K}) \mathbf{u}^{n+1}} \quad \forall K \in \mathfrak{M}$$

We define the fluxes:

$$\int_{\sigma} (\mathbf{u}^n \cdot \vec{\mathbf{n}}_{\sigma K}) \rightsquigarrow F_{\sigma, K}(\mathbf{u}^n)$$

# Nonlinear convection term $(\mathbf{u} \cdot \nabla)\mathbf{u}$

We impose:  $F_{\sigma,K}(\mathbf{u}^n) = \begin{cases} - \sum_{\mathfrak{s} \in \mathfrak{S}_K \cap \mathcal{E}_D} G_{\mathfrak{s},D}(\mathbf{u}^n) & \text{if } \sigma \in \mathcal{E}_{int} \\ m_\sigma \gamma^\sigma(\mathbf{u}^n) \cdot \vec{\mathbf{n}}_{\sigma K} & \text{if } \sigma \in \partial\Omega \end{cases},$


where

$$G_{\mathfrak{s},D}(\mathbf{u}^n) = m_{\mathfrak{s}} \frac{\mathbf{u}_K^n + \mathbf{u}_{K^*}^n}{2} \cdot \vec{\mathbf{n}}_{\mathfrak{s}D} \rightsquigarrow \int_{\mathfrak{s}} \mathbf{u}^n \cdot \vec{\mathbf{n}}_{\mathfrak{s}D},$$

$$\gamma_\sigma(\mathbf{u}^n) = \frac{\mathbf{u}_{K^*}^n + 2\mathbf{u}_L^n + \mathbf{u}_{L^*}^n}{4} \quad \forall \sigma = [x_{K^*}, x_{L^*}] \in \partial\mathcal{M}.$$

We have conservativity:

$$F_{\sigma,K} = -F_{\sigma,L}, \quad \forall \sigma = K|L$$



# Nonlinear convection term $(\mathbf{u} \cdot \nabla)\mathbf{u}$

$$m_K \mathbf{b}^K(\mathbf{u}^n, \mathbf{u}^{n+1}) \rightsquigarrow \sum_{\sigma \subset \partial K} \int_{\sigma} (\mathbf{u}^n \cdot \vec{\mathbf{n}}_{\sigma K}) \mathbf{u}^{n+1}$$

So we define  $\forall K \in \mathfrak{M}$ :

$$m_K \mathbf{b}^K(\mathbf{u}^n, \mathbf{u}^{n+1}) = \sum_{D_{\sigma, \sigma^*} \in \mathfrak{D}_K^{int}} F_{\sigma, K}(\mathbf{u}^n) \mathbf{u}_{\sigma^+}^{n+1} + \sum_{D_{\sigma, \sigma^*} \in \mathfrak{D}_K^{ext}} F_{\sigma, K}(\mathbf{u}^n) \gamma^{\sigma}(\mathbf{u}^{n+1})$$

where

$$\mathbf{u}_{\sigma^+}^{n+1} = \begin{cases} \mathbf{u}_K^{n+1} & \text{if } F_{\sigma, K} \geq 0 \\ \mathbf{u}_L^{n+1} & \text{otherwise} \end{cases} \quad \forall \sigma \in \mathcal{E}_{int}$$

and

$$\gamma_{\sigma}(\mathbf{u}^{n+1}) = \frac{\mathbf{u}_{K^*}^{n+1} + 2\mathbf{u}_L^{n+1} + \mathbf{u}_{L^*}^{n+1}}{4} \quad \forall \sigma = [x_{K^*}, x_{L^*}] \in \partial \mathfrak{M}.$$



# Variational formulation

$$\int_{\Omega} \partial_t \mathbf{u} \cdot \Psi + \frac{2}{\text{Re}} \int_{\Omega} \mathbf{D}(\mathbf{u}) : \mathbf{D}(\Psi) + \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \Psi - \frac{1}{2} \int_{\Omega} (\mathbf{u} \cdot \nabla) \Psi \cdot \mathbf{u}$$

$$= -\frac{1}{2} \int_{\Gamma_2} (\mathbf{u} \cdot \vec{\mathbf{n}})^+ (\mathbf{u} \cdot \Psi) + \frac{1}{2} \int_{\Gamma_2} (\mathbf{u} \cdot \vec{\mathbf{n}})^- (\mathbf{u}_{\text{ref}} \cdot \Psi) + \int_{\Gamma_2} \sigma_{\text{ref}} \cdot \vec{\mathbf{n}} \cdot \Psi$$

$\Downarrow$

$$\left[ \left[ \frac{\mathbf{u}^{n+1} - \mathbf{u}^n}{\delta t}, \Psi^{\mathfrak{T}} \right]_{\mathfrak{T}} + \frac{2}{\text{Re}} (\mathbf{D}^{\mathfrak{D}} \mathbf{u}^{n+1}, \mathbf{D}^{\mathfrak{D}} \Psi^{\mathfrak{T}})_{\mathfrak{D}} + \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{u}^n, \mathbf{u}^{n+1}), \Psi^{\mathfrak{T}}]]_{\mathfrak{T}} \right.$$

$$\left. - \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{u}^n, \Psi^{\mathfrak{T}}), \mathbf{u}^{n+1}]]_{\mathfrak{T}} \right] = -\frac{1}{2} \sum_{\mathbf{D} \in \mathbf{D}_{\text{ext}} \cap \Gamma_2} (F_{\sigma, \mathbf{k}}(\mathbf{u}^n))^+ \gamma^{\sigma}(\mathbf{u}^{n+1}) \gamma^{\sigma}(\Psi^{\mathfrak{T}})$$

$$+ \frac{1}{2} \sum_{\mathbf{D} \in \mathbf{D}_{\text{ext}} \cap \Gamma_2} (F_{\sigma, \mathbf{k}}(\mathbf{u}^n))^- \gamma^{\sigma}(\mathbf{u}_{\text{ref}}) \gamma^{\sigma}(\Psi^{\mathfrak{T}}) + \sum_{\mathbf{D} \in \mathbf{D}_{\text{ext}} \cap \Gamma_2} m_{\sigma}(\sigma_{\text{ref}}^{\mathfrak{D}} \cdot \vec{\mathbf{n}}_{\sigma \mathbf{k}}) \cdot \gamma^{\sigma}(\Psi^{\mathfrak{T}}).$$

# The scheme

■ For all  $k \in \mathcal{M}$ :

$$m_k \frac{\mathbf{u}_k^{n+1} - \mathbf{u}_k^n}{\delta t} - m_k \mathbf{div}^k(\sigma^{\mathcal{D}}(\mathbf{u}^{n+1}, \mathbf{p}^{n+1})) + \sum_{\mathcal{D}_{\sigma, \sigma^*} \in \mathcal{D}_k^{int}} F_{\sigma, k}(\mathbf{u}^n) \frac{\mathbf{u}_k^{n+1} + \mathbf{u}_L^{n+1}}{2} + \frac{1}{2} \sum_{\mathcal{D}_{\sigma, \sigma^*} \in \mathcal{D}_k^{ext}} F_{\sigma, k}(\mathbf{u}^n) \left( \mathbf{u}_k^{n+1} + \gamma^\sigma(\mathbf{u}^{n+1}) \right) = 0;$$

■ For all  $\sigma \in \partial \mathcal{O}$ :

$$m_\sigma \sigma^{\mathcal{D}}(\mathbf{u}^{n+1}, \mathbf{p}^{n+1})) \vec{\mathbf{n}}_{\sigma L} - \frac{1}{4} F_{\sigma, L}(\mathbf{u}^n) (\mathbf{u}_k^{n+1} - \mathbf{u}_L^{n+1}) = -\frac{1}{2} (F_{\sigma, L}(\mathbf{u}^n))^- (\gamma^\sigma(\mathbf{u}^{n+1}) - \gamma^\sigma(\mathbf{u}_{ref})) + m_\sigma (\sigma_{ref}^{\mathcal{D}} \cdot \vec{\mathbf{n}}_{\sigma k});$$

■ For all  $\mathcal{D} \in \mathcal{D}$ :

$$\mathbf{div}^{\mathcal{D}}(\mathbf{u}^{n+1}) = 0.$$

# Well-posedness of the scheme

We obtain our scheme by projecting the discrete weak formulation on the DDFV mesh.

## Well-posedness

Let  $\mathfrak{T}$  be a DDFV mesh associated to  $\Omega$  that satisfies inf-sup stability condition. The scheme we obtain has a unique solution  $(\mathbf{u}^{\mathfrak{T},[0,T]}, p^{\mathfrak{D},[0,T]}) \in ((\mathbb{R}^2)^{\mathfrak{T}})^{N+1} \times (\mathbb{R}^{\mathfrak{D}})^{N+1}$ .

# Proof of well-posedness

The scheme is equivalent to a system  $Av = b$  at each time step, with  $A$  square matrix and  $v = (\mathbf{u}^{n+1}, p^{n+1})$ .



We want to show that the square matrix is injective, i.e.

$$Av = 0 \implies v = 0$$

# Proof of well-posedness

If we multiply  $Av = 0$  by a test function  $\Psi^{\mathfrak{T}}$  that satisfies

$$\begin{cases} \Psi^{\mathfrak{T}} \in \mathbb{E}_0^D, \\ \operatorname{div}^{\mathfrak{D}}(\Psi^{\mathfrak{T}}) = 0, \end{cases}$$

this is equivalent to consider the discrete variational formulation in the form:

$$\begin{aligned} & \frac{1}{\delta t} [[\mathbf{u}^{n+1}, \Psi^{\mathfrak{T}}]]_{\mathfrak{T}} + \frac{2}{\operatorname{Re}} (D^{\mathfrak{D}} \mathbf{u}^{n+1}, D^{\mathfrak{D}} \Psi^{\mathfrak{T}})_{\mathfrak{D}} \\ & + \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{u}^n, \mathbf{u}^{n+1}), \Psi^{\mathfrak{T}}]]_{\mathfrak{T}} - \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{u}^n, \Psi^{\mathfrak{T}}), \mathbf{u}^{n+1}]]_{\mathfrak{T}} \\ & = -\frac{1}{2} \sum_{D \in \mathcal{D}_{\text{ext}} \cap \Gamma_2} (F_{\sigma, \mathbf{x}}(\mathbf{u}^n))^+ \gamma^{\sigma}(\mathbf{u}^{n+1}) \gamma^{\sigma}(\Psi^{\mathfrak{T}}). \end{aligned}$$

# Proof of well-posedness

If in this weak formulation:

$$\begin{aligned} \frac{1}{\delta t} [[\mathbf{u}^{n+1}, \Psi^{\mathfrak{T}}]]_{\mathfrak{T}} + \frac{2}{\text{Re}} (D^{\mathfrak{D}} \mathbf{u}^{n+1}, D^{\mathfrak{D}} \Psi^{\mathfrak{T}})_{\mathfrak{D}} \\ + \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{u}^n, \mathbf{u}^{n+1}), \Psi^{\mathfrak{T}}]]_{\mathfrak{T}} - \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{u}^n, \Psi^{\mathfrak{T}}), \mathbf{u}^{n+1}]]_{\mathfrak{T}} \\ = -\frac{1}{2} \sum_{D \in \mathcal{D}_{\text{ext}} \cap \Gamma_2} (F_{\sigma, \kappa}(\mathbf{u}^n))^+ \gamma^{\sigma}(\mathbf{u}^{n+1}) \gamma^{\sigma}(\Psi^{\mathfrak{T}}). \end{aligned}$$

we choose  $\Psi^{\mathfrak{T}} = \mathbf{u}^{n+1}$ , we obtain:

$$\frac{1}{\delta t} \|\mathbf{u}^{n+1}\|_2^2 + \frac{2}{\text{Re}} \|D^{\mathfrak{D}} \mathbf{u}^{n+1}\|_2^2 + \underbrace{\frac{1}{2} \sum_{D \in \mathcal{D}_{\text{ext}}} (F_{\sigma, \kappa}(\mathbf{u}^n))^+ |\gamma^{\sigma}(\mathbf{u}^{n+1})|^2}_{\geq 0} = 0,$$

# Proof of well-posedness

We end up with:

$$\frac{1}{\delta t} \|\mathbf{u}^{n+1}\|_2^2 + \frac{2}{\text{Re}} \|D^{\mathfrak{D}} \mathbf{u}^{n+1}\|_2^2 \leq 0$$

from which we deduce that  $\boxed{\mathbf{u}^{n+1} = 0}$ .

To conclude the proof, we need to show that  $\mathbf{p}^{n+1}$  is equal to zero too. This is true because of the inf-sup stability, that ensures an inequality of the type:

$$\|\mathbf{p}^{n+1} - m(\mathbf{p}^{n+1})\|_2 \leq C \|\nabla^{\mathfrak{D}} \mathbf{u}^{n+1}\|_2,$$

where  $m(\mathbf{p}^{n+1}) = \sum_{D \in \mathfrak{D}} m_D \mathbf{p}^D$ . This inequality implies that  $\mathbf{p}^{n+1}$  is constant.

Thanks to boundary condition on  $\Gamma_2$ , we conclude  $\boxed{\mathbf{p}^{n+1} = 0}$ .

# Discrete energy estimate

Let  $\mathfrak{T}$  be a DDFV mesh associated to  $\Omega$  that satisfies inf-sup stability condition.

Let  $(\mathbf{u}^{\mathfrak{T},[0,T]}, \mathbf{p}^{\mathfrak{D},[0,T]}) \in ((\mathbb{R}^2)^{\mathfrak{T}})^{N+1} \times (\mathbb{R}^{\mathfrak{D}})^{N+1}$  be the solution of the DDFV scheme, where  $\mathbf{u}^{\mathfrak{T},[0,T]} = \mathbf{v}^{\mathfrak{T},[0,T]} + \mathbf{u}_{ref}^{\mathfrak{T}}$ .

For  $N > 1$ , there exists a constant  $C > 0$ , depending on  $\Omega, \mathbf{u}_{ref}^{\mathfrak{T}}, \mathbf{u}_0$ , Re such that:

$$\sum_{j=0}^{N-1} \|\mathbf{v}^{j+1} - \mathbf{v}^j\|_2^2 \leq C, \quad \|\mathbf{v}^N\|_2^2 \leq C,$$

$$\sum_{j=0}^{N-1} \delta t \frac{1}{\text{Re}} \|D^{\mathfrak{D}} \mathbf{v}^{j+1}\|_2^2 \leq C, \quad \delta t \frac{1}{\text{Re}} \|D^{\mathfrak{D}} \mathbf{v}^N\|_2^2 \leq C,$$

$$\sum_{j=0}^{N-1} \delta t \sum_{\mathbf{D} \in \mathfrak{D}_{ext}} (F_{\sigma, \mathbf{k}}(\mathbf{v}^j + \mathbf{u}_{ref}^{\mathfrak{T}}))^+ (\gamma^{\sigma}(\mathbf{v}^{j+1}))^2 \leq C.$$



# Sketch of the proof

1. Rewrite our discrete variational formulation for the unknown  $\mathbf{v}^{n+1} = \mathbf{u}^{n+1} - \mathbf{u}_{ref}^{\mathfrak{T}}$ :

$$\begin{aligned}
 & [[\frac{\mathbf{v}^{n+1} - \mathbf{v}^n}{\delta t}, \Psi^{\mathfrak{T}}]]_{\mathfrak{T}} + \frac{2}{\text{Re}} (D^{\mathfrak{D}} \mathbf{v}^{n+1}, D^{\mathfrak{D}} \Psi^{\mathfrak{T}})_{\mathfrak{D}} \\
 & + \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{v}^n + \mathbf{u}_{ref}^{\mathfrak{T}}, \mathbf{v}^{n+1} + \mathbf{u}_{ref}^{\mathfrak{T}}), \Psi^{\mathfrak{T}}]]_{\mathfrak{T}} - \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{v}^n + \mathbf{u}_{ref}^{\mathfrak{T}}, \Psi^{\mathfrak{T}}), \mathbf{v}^{n+1} + \mathbf{u}_{ref}^{\mathfrak{T}}]]_{\mathfrak{T}} \\
 & + \frac{1}{2} \sum_{\mathfrak{D} \in \mathfrak{D}_{ext}} (F_{\sigma, \mathbf{k}}(\mathbf{v}^n + \mathbf{u}_{ref}^{\mathfrak{T}}))^+ \gamma^{\sigma}(\mathbf{v}^{n+1} + \mathbf{u}_{ref}^{\mathfrak{T}}) \gamma^{\sigma}(\Psi^{\mathfrak{T}}) \\
 & = -\frac{1}{2} \sum_{\mathfrak{D} \in \mathfrak{D}_{ext}} F_{\sigma, \mathbf{k}}(\mathbf{v}^n + \mathbf{u}_{ref}^{\mathfrak{T}})^- \gamma^{\sigma}(\mathbf{u}_{ref}^{\mathfrak{T}}) \gamma^{\sigma}(\Psi^{\mathfrak{T}}).
 \end{aligned}$$

2. The second step consists into selecting  $\Psi^{\mathfrak{T}} = \mathbf{v}^{n+1} = (\mathbf{v}^{n+1} + \mathbf{u}_{ref}^{\mathfrak{T}}) - \mathbf{u}_{ref}^{\mathfrak{T}}$  as a test function. It follows:

$$\begin{aligned}
 E &= [[\frac{\mathbf{v}^{n+1} - \mathbf{v}^n}{\delta t}, \mathbf{v}^{n+1}]]_{\mathfrak{T}} + \frac{2}{\text{Re}} \|D^{\mathfrak{D}} \mathbf{v}^{n+1}\|_2^2 + \frac{1}{2} \sum_{\mathfrak{D} \in \mathfrak{D}_{ext} \cap \Gamma_2} (F_{\sigma, \mathbf{k}}(\mathbf{v}^n + \mathbf{u}_{ref}^{\mathfrak{T}}))^+ (\gamma^{\sigma}(\mathbf{v}^{n+1}))^2 \\
 &\leq \left| \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{v}^n + \mathbf{u}_{ref}^{\mathfrak{T}}, \mathbf{v}^{n+1}), \mathbf{u}_{ref}^{\mathfrak{T}}]]_{\mathfrak{T}} - \frac{1}{2} [[\mathbf{b}^{\mathfrak{T}}(\mathbf{v}^n + \mathbf{u}_{ref}^{\mathfrak{T}}, \mathbf{u}_{ref}^{\mathfrak{T}}), \mathbf{v}^{n+1}]]_{\mathfrak{T}} \right| \\
 &\quad + \left| \frac{1}{2} \sum_{\mathfrak{D} \in \mathfrak{D}_{ext} \cap \Gamma_2} F_{\sigma, \mathbf{k}}(\mathbf{v}^n + \mathbf{u}_{ref}^{\mathfrak{T}})^- \gamma^{\sigma}(\mathbf{u}_{ref}^{\mathfrak{T}}) \gamma^{\sigma}(\mathbf{v}^{n+1}) \right|
 \end{aligned}$$

3. Estimate the RHS.

# Further results: Korn's and trace inequality

## Korn's inequality

Let  $\mathfrak{T}$  be a mesh that satisfies infsup stability condition. Then there exists  $C > 0$  such that :



$$\boxed{\|\nabla^{\mathcal{D}} \mathbf{u}^{\mathfrak{T}}\|_2 \leq C \|D^{\mathcal{D}} \mathbf{u}^{\mathfrak{T}}\|_2} \quad \forall \mathbf{u}^{\mathfrak{T}} \in \mathbb{E}_0^{\mathcal{D}}$$

## Trace theorem

Let  $\mathfrak{T}$  be a DDFV mesh associated to  $\Omega$ . For all  $p > 1$ , there exists a constant  $C > 0$ , depending only on  $p$ ,  $\sin(\alpha_{\mathfrak{T}})$ ,  $\text{reg}(\mathfrak{T})$  and  $\Omega$  such that  $\forall \mathbf{u}^{\mathfrak{T}} \in \mathbb{E}_0^{\mathcal{D}}$  and for all  $s \geq 1$ :

$$\boxed{\|\gamma(\mathbf{u}^{\mathfrak{T}})\|_{s,\partial\Omega}^s \leq C \|\mathbf{u}^{\mathfrak{T}}\|_{1,p} \|\mathbf{u}^{\mathfrak{T}}\|^{\frac{s-1}{p-1}}_{\frac{p(s-1)}{p-1}}}$$

# Numerical results 1





$$\mathbf{g}_1(x, y) = \begin{pmatrix} 6y(1-y) \\ 0 \end{pmatrix} \quad \text{on } \Gamma_1$$

$$\mathbf{u}_{ref}(x, y) = \begin{pmatrix} 6y(1-y) \\ 0 \end{pmatrix} \quad \text{on } \Gamma_2$$

$$\sigma_{ref}(\mathbf{u}, \mathbf{p}) \cdot \vec{\mathbf{n}} = \begin{pmatrix} 0 \\ 6\eta(1-2y) \end{pmatrix} \quad \text{on } \Gamma_2$$

with  $T = 1.5s$ ,  $\text{Re} = 100$ ,  $\eta = 4 \times 10^{-3}$ .

# Numerical results 1



$$\mathbf{g}_1(x, y) = \begin{pmatrix} 6y(1-y) \\ 0 \end{pmatrix} \quad \text{on } \Gamma_1$$

$$\mathbf{u}_{ref}(x, y) = \begin{pmatrix} 6y(1-y) \\ 0 \end{pmatrix} \quad \text{on } \Gamma_2$$


$$\sigma_{ref}(\mathbf{u}, p) \cdot \vec{\mathbf{n}} = \begin{pmatrix} 0 \\ 6\eta(1-2y) \end{pmatrix} \quad \text{on } \Gamma_2$$

with  $T = 3s$ ,  $\text{Re} = 100$ ,  $\eta = 4 \times 10^{-3}$ .

# Numerical results 2

$$C_{d,max} = 2.9754$$


$$C_{l,max} = 0.44902$$



DDFV

$$C_{d,max} = 2.9509$$

$$C_{l,max} = 0.47795$$



Reference values [Volker, '04]

# Conclusions and perspectives

## Conclusions

- ❖ We built a well-posed DDFV scheme
- ❖ We proved a discrete energy estimate
- ❖ We obtained the expected numerical results

## Perspectives

- ❖ investigate the choice of  $\mathbf{u}_{ref}$  for higher Reynold's number
- ❖ **application to Domain Decomposition**

Grazie per l'attenzione!