

Anderson localisation with self-avoiding-walks

Giulia Liconti

Geboren am 21. November 2004 in Bologna

27th August 2026

Bachelor's Thesis Mathematics

Advisor: Prof. Dr. Margerita Disertori

Second advisor: Dr. Davide Macera

INSTITUT OF APPLIED MATHEMATICS

MATHEMATISCH-NATURWISSENSCHAFTLICHE FAKULTÄT DER
RHEINISCHEN FRIEDRICH-WILHELMS-UNIVERSITÄT BONN

Contents

1	Introduction	2
1.1	The Anderson Model	3
1.1.1	The operator	3
1.1.2	The spectrum.	7
1.2	Main results	8
2	Localisation properties	9
2.1	spectral localisation	9
2.1.1	Absence of transport	9
2.2	dynamical localisation	10
2.3	From dynamical to spectral localisation	11
3	Fractional moments	13
3.1	The green function	13
3.2	Localisation at large disorder	13
3.2.1	From fractional moment's exponential bound to dynamical localisation	14
4	SAW representation	16
4.1	SAW Definitions	16
4.2	SAW representation of G	16
4.3	Proof with SAW	16
5	From fractional moments to dynamical localisation	17

1 Introduction

The Anderson localization model was introduced by the physicist P.W. Anderson in 1958 to explain the quantum mechanical effects of disorder in materials. Anderson discusses the behaviour of electrons in dirty crystals, which is the quantum mechanical analogue of a random walk in a random environment. The model considers a d -dimensional grid, in which each site represents an atom. From an heuristical point of view, we also consider electrons that jump from atom to atom and are subject to an external random potential, which influences their motion in the material. When these particles cannot jump, it indicates the absence of quantum transport. This phenomenon is called Anderson localisation. This model deals with conditions for which the electrons stay trapped in a specific region and don't generate electric conduction, defining an insulator. This is in sharp contrast to the behaviour in ideal crystals, which are always conductors.

Anderson's work, and that of N.F.Mott and J.H. van Vleck, won the 1977 physics Nobel prize for "their fundamental theoretical investigations of the electronic structure of magnetic and disordered systems". Since then, the study of random operators has become an important field of mathematical physics, which has led to a tremendous amount of research activity and many mathematical results.

The two important methods to prove Anderson localization are the *multiscale analysis method* (MSA), developed by Froehlich and Spencer [1] in 1983, and the *fractional moment method* (FMM) by Aizenman and Molchanov in 1993. We will deal with the second one, which is more elementary and gives stronger results on *dynamical localization* [Chapter 2.2].

structure of the thesis

Prerequisites

It is useful to have some familiarity with measure theory, basic probabilistic

concepts such as independence and continuous distributions, and foundations of the theory of linear operators in Hilbert spaces, up to the spectral theorem for self adjoint operators and consequences (as spectral types).

1.1 The Anderson Model

Transport phenomena in a disordered material are often described via discrete random Schroedinger operator. On the lattice $\mathbb{Z}^d$, it takes the form of an infinite random matrix called the *Hamiltonian*:

$$H_\omega = -\Delta + \lambda V_\omega.$$

where $-\Delta$ is a deterministic matrix and V_ω is a diagonal matrix with random entries. This operator acts on the Hilbert space

$$l^2(\mathbb{Z}^d) = \{u : \mathbb{Z}^d \rightarrow \mathbb{C} : \sum_{n \in \mathbb{Z}^d} |u(n)|^2 < \infty\}$$

with inner product $\langle u, v \rangle = \sum_n \overline{u(n)}v(n)$.

1.1.1 The operator

The Discrete Laplacian. The kinetic term $-\Delta$ is the discrete analogue of the usual negative Laplacian $(-\sum_j \partial^2/\partial x_j^2)$. For the functions $u \in l^2(\mathbb{Z}^d)$ it is defined as:

$$-\Delta u(x) := \sum_{k \in \mathbb{Z}^d, |x-k|=1} u(x) - u(k) \quad \text{for } x \in \mathbb{Z}^d.$$

where $|\cdot|$ denotes the l^2 norm.

The discrete Laplacian represents the kinetic term and connects neighbour sites on the lattice, such that the electron can transition from one to another.

For each site x , there are $2d$ possible k such that $|x - k| = 1$. We can

now define an infinite matrix such that:

$$(-\Delta)_{x,k} = \begin{cases} 2d & \text{if } x = k \quad (\text{on the diagonal}) \\ -1 & \text{if } |x - k| = 1 \quad (k \sim x, \text{ first neighbors}) \\ 0 & \text{otherwise} \end{cases}$$

For $u \in \ell^2(\mathbb{Z}^d)$, $-\Delta u$ is summable in ℓ^2 , bounded, and self-adjoint.

The first claim (summability) is shown by:

$$\begin{aligned} \sum_{x \in \mathbb{Z}^d} |\Delta u(x)|^2 &= \sum_{x \in \mathbb{Z}^d} \left| \sum_{k \sim x, k \in \mathbb{Z}^d} (u(x) - u(k)) \right|^2 \\ &\leq 2d \sum_{x \in \mathbb{Z}^d} \left(\sum_{k \sim x} |u(x) - u(k)|^2 \right) \\ &\leq 2d \sum_{x \in \mathbb{Z}^d} \left(2 \sum_{k \sim x, k \in \mathbb{Z}^d} |u(x)|^2 + |u(k)|^2 \right) \\ &\leq 2d \cdot 2d \cdot 4 \|u\|_2^2 < \infty. \end{aligned}$$

In each step we used, in order, Cauchy-Schwarz, the bound over the number of neighbors, triangle inequality and the invariance under translation of the l^2 norm.

The boundedness comes from:

$$\frac{\|\Delta u\|_2^2}{\|u\|_2^2} \leq 16d^2 \implies \|\Delta\|_2 \leq 4d.$$

To prove self-adjointness, we only need to show symmetry (given $-\Delta$ is bounded). For $u, v \in \ell^2(\mathbb{Z}^d)$ and $k \in \mathbb{Z}^d$:

$$\begin{aligned} \langle u, -\Delta v \rangle &= \sum_{x \in \mathbb{Z}^d} \overline{u(x)} (-\Delta v(x)) = \sum_{x \in \mathbb{Z}^d} u(x) \left(\sum_{k \sim x} v(x) - v(k) \right) \\ &= \sum_{x \in \mathbb{Z}^d} \left(u(x) 2d v(x) - \sum_{k \sim x} u(x) v(k) \right) = \sum_{x \in \mathbb{Z}^d} 2d u(x) v(x) - \sum_{x \in \mathbb{Z}^d} \sum_{k \sim x} u(k) v(x) \\ &= \sum_{x \in \mathbb{Z}^d} v(x) \left(\sum_{k \sim x} u(x) - u(k) \right) = \sum_{x \in \mathbb{Z}^d} \overline{v(x)} (-\Delta u(x)) = \langle -\Delta u, v \rangle. \end{aligned}$$

The spectrum of the bounded, self-adjoint operator $-\Delta$ is the closed, real-valued interval $[0, 4d]$. In this range, we find the levels of kinetic energy that an electron owns.

In order to understand this results better, let's consider the discrete Fourier transform:

$$\mathcal{F}: \ell^2(\mathbb{Z}^d) \longrightarrow L^2([0, 2\pi)^d)$$

such that:

$$(\mathcal{F}u)(x) := \lim_{N \rightarrow \infty} (2\pi)^{-\frac{d}{2}} \sum_{|n| \leq N} u(n) e^{-ix \cdot n}, \quad x \in [0, 2\pi)^d.$$

For simplicity of the calculations we don't consider the limit and just sum over all $n \in \mathbb{Z}^d$ and u summable.

$$\begin{aligned} (\mathcal{F}\Delta u)(x) &= (2\pi)^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} (\Delta u)(n) e^{-ix \cdot n} \\ &= (2\pi)^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} \left(\sum_{|k-n|=1} (u(n) - u(k)) \right) e^{-ix \cdot n} \\ &= (2\pi)^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} \left(2d u(n) - \sum_{|k-n|=1} u(k) \right) e^{-ix \cdot n} \\ &= 2d (\mathcal{F}u)(x) - (2\pi)^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} \sum_{|k-n|=1} u(k) e^{-ix \cdot n} \\ &= 2d (\mathcal{F}u)(x) - (2\pi)^{-\frac{d}{2}} \sum_{k \in \mathbb{Z}^d} u(k) \sum_{|m|=1} e^{-ix \cdot (k+m)} \quad (\text{for } m = n - k) \\ &= 2d (\mathcal{F}u)(x) - \left((2\pi)^{-\frac{d}{2}} \sum_{k \in \mathbb{Z}^d} u(k) e^{-ix \cdot k} \right) \sum_{|m|=1} e^{-ix \cdot m} \\ &= 2d (\mathcal{F}u)(x) - (\mathcal{F}u)(x) \sum_{j=1}^d (e^{-ix_j} + e^{ix_j}) \\ &= \left(2d - 2 \sum_{j=1}^d \cos(x_j) \right) (\mathcal{F}u)(x) \\ &= \left(2 \sum_{j=1}^d 1 - \cos(x_j) \right) (\mathcal{F}u)(x). \end{aligned}$$

We obtain the multiplication operator:

$$\mathcal{F}\Delta\mathcal{F}^{-1} = 2 \sum_{j=1}^d 1 - \cos(x_j)$$

on $L^2([0, 2\pi)^d)$ in the variable $x = (x_1, \dots, x_d)$. We notice that the Laplacian doesn't have eigenvalues in $l^2(\mathbb{Z}^d)$ Another similarity of Δ with the continuum Laplacian is that it has plane waves as generalized eigenfunctions. To see this, let $x \in [0, 2\pi)^d$ and set

$$\phi_x(n) := e^{-in \cdot x}$$

While $\phi_x \notin l^2(\mathbb{Z}^d)$, Δ acts on it as

$$\begin{aligned} (\Delta\phi_x)(n) &= \sum_{|k|=1} e^{-in \cdot x} - e^{-i(n+k) \cdot x} \\ &= 2d\phi_x(n) - \left(\sum_{j=1}^d 2 \cos(x_j) \right) \phi_x(n) = \left(2 \sum_{j=1}^d 1 - \cos(x_j) \right) \phi_x. \end{aligned}$$

Thus ϕ_x is a bounded generalized eigenfunction of Δ to the spectral value $2 \sum_{j=1}^d 1 - \cos(x_j)$.

...

The Potential Term.

Definition 1.1 (i.i.d. random variables). $(\omega_n)_{n \in \mathbb{Z}^d}$ are identically distributed if there exists a Borel probability measure μ on $\mathbb{R}$ such that $\mathbb{P}(\omega_n \in A) = \mu(A)$ for all $n \in \mathbb{Z}^d$ and Borel sets $A \subset \mathbb{R}$. $(\omega_n)_{n \in \mathbb{Z}^d}$ are independent if

$$\mathbb{P}(\omega_{n_1} \in A_1, \dots, \omega_{n_l} \in A_l) = \prod_{k=1}^l \mathbb{P}(\omega_{n_k} \in A_k) = \prod_{k=1}^l \mu(A_k)$$

for each finite subset $\{n_1, \dots, n_l\}$ of $\mathbb{Z}^d$ and arbitrary Borel sets $A_1, \dots, A_l \subset \mathbb{R}$.

We assume that the distribution μ of ω_v is absolutely continuous with density ρ where ρ is bounded with compact support, i.e.,

$$\mu(B) = \int_B \rho(u) du \quad \text{for } B \subset \mathbb{R} \text{ Borel, } \quad \rho \in L_0^\infty(\mathbb{R}).$$

Physically, V_ω represents the random electric potential created by nuclei at the sites $n \in \mathbb{Z}^d$ **ovviamente il testo da modificare in base al resto delle notazioni** .

$\lambda > 0$ is the disorder parameter

- Definition
- heuristic role generic explanation
- explanation for 0 and very big to give an idea.

1.1.2 The spectrum.

- in $\mathbb{R}$
- bounded $[0, 4d] + \text{supp}\mu$
- spectral division (any use of the spectral theorem should be sent to the Appendix)
- RAGE theorem (or maybe directly in the Appendix or chapter 2)

Dalla teoria spettrale per operatori autoaggiunti su uno spazio di Hilbert $\mathcal{H}$, lo spazio si decompone univocamente nella somma diretta di sottospazi invarianti ortogonali:

$$\mathcal{H} = \mathcal{H}_{pp} \oplus \mathcal{H}_{sc} \oplus \mathcal{H}_{ac} \quad (1.1)$$

a cui corrispondono rispettivamente lo spettro puramente puntuale $\sigma_{pp}(H)$, singolarmente continuo $\sigma_{sc}(H)$ e assolutamente continuo $\sigma_{ac}(H)$, con $\sigma(H) = \sigma_{pp}(H) \cup \sigma_{sc}(H) \cup \sigma_{ac}(H)$.

L'interpretazione fisica di questi sottospazi è chiarita dal fondamentale **Teorema RAGE** (Ruelle-Amrein-Georgescu-Enss):

Theorem 1.2 (RAGE). *Suppose H is a Schrödinger operator acting on $l^2(\mathbb{Z}^d)$. Then:*

- (i) $\phi \in \mathcal{H}_{pp}$ if and only if

$$\lim_{R \rightarrow \infty} \sup_{t \in \mathbb{R}} \|\chi_{(|x| > R)} e^{-itH} \phi\| = 0.$$

- (ii) $\psi \in \mathcal{H}_c = \mathcal{H}_{ac} \oplus \mathcal{H}_{sc}$ if and only if

$$\lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T \|\chi_{(|x| \leq R)} e^{-itH} \psi\| dt = 0$$




Figure 1: Aizenman figure

for all $R > 0$.

Here $\chi_{(|x| \leq R)}$ and $\chi_{(|x| > R)}$ denote the characteristic functions of the ball $\{|x| \leq R\}$ and the complement of the ball, respectively.

1.2 Main results

In this section one could write first of all the main theorem and the spectral results. starting maybe with:

The aim of this thesis is to prove localisation on the whole spectrum of H_ω
...

Poi scrivo il teorema finale con spiegazione che la dimostrazione avverrà nel capitolo 5.

potrebbe essere un'idea anche scrivere una piccola parentesi sulla differenza dei risultati con Frohlich e spencer e con Aizenman e Molchanov.

One of the main results concerns the values of λ

- $\lambda \gg 1$ and $\lambda = 0$
- general case for $d = 1, d \geq 2$

2 Localisation properties

We understood what localisation means from an heuristical point of view. Mathematically, it can be described from spectral and dynamical properties.

2.1 spectral localisation

Definition 2.1 (Spectral Localization). random Schrödinger operator $H = H_\omega$ exhibits **spectral localization** if H_ω almost surely has only pure point spectrum, that is,

$$\sigma(H_\omega) = \sigma_{pp}(H_\omega) \quad \text{and} \quad \sigma_c(H_\omega) = \emptyset \quad \text{almost surely.}$$

Furthermore the eigenfunctions corresponding to all eigenvalues decay exponentially: for almost all ω , H_ω has a complete set of eigenfunctions $(\phi_{\omega,n})_{n \in \mathbb{N}}$ satisfying

$$|\phi_{\omega,n}(x)| \leq C_{\omega,n} e^{-\mu|x-x_{n,\omega}|}$$

where $\mu > 0$ is the localization length parameter, $x_{n,\omega}$ are the centers of localization, and $C_{\omega,n} < \infty$ is a finite constant.

2.1.1 Absence of transport

$$\langle x^2 \rangle_{P_{H_\omega \in [a,b]} \phi_0}(t) = \left\| |x| e^{-itH_\omega} P_{H_\omega \in [a,b]} \phi_0 \right\|^2 \quad (5)$$

If the electrons with energies in $[a, b]$ move ballistically with average velocity v_{av} , then roughly $x(t) \sim v_{av}t$ for large times. Hence $\langle x^2 \rangle_{P_{H_\omega \in [a,b]} \phi_0}(t)$ will be proportional to t^2 for large t .

On the other hand, if the electrons in the energy interval $[a, b]$ are localized, then it should be natural to expect that $\langle x^2 \rangle_{P_{H_\omega \in [a,b]} \phi_0}(t)$ is bounded uniformly in t . It is known (Simon, 1990) that pure point spectrum implies absence of ballistic motion,

$$\lim_{t \rightarrow \infty} \frac{\langle x^2 \rangle_{P_{H_\omega \in [a,b]} \phi_0}(t)}{t^2} = 0$$

for all compactly supported initial conditions ϕ_0 , as soon as the spectrum in $[a, b]$ is pure point. However, del Rio et al. (1996) constructed examples of (non-random) one-dimensional Schrödinger operators with pure point spec-

trum for all energies and exponentially localized eigenfunctions for which

$$\limsup_{t \rightarrow \pm\infty} \frac{\langle x^2 \rangle_{\phi_0}(t)}{t^{2-\delta}} = \infty \quad \text{for all } \delta > 0, \quad (6)$$

for a large class of compactly supported initial conditions ϕ_0 . *Il fallimento risiede nella libertà delle costanti $C_{\omega,n}$: se $C_{\omega,n}$ cresce incontrollatamente con n , le code delle autofunzioni possono sovrapporsi su scale spaziali arbitrariamente grandi.*

2.2 dynamical localisation

Definition 2.2 (Dynamical localisation).

$$\sup_{t \in \mathbb{R}} \| |X|^p e^{-ith_\omega} \psi \| < \infty \quad \text{almost surely}, \quad (2.1)$$

where the position operator $|X|$ is defined by $(|X|\phi)(n) = |n|\phi(n)$ and for all φ with compact support..

Definition 2.3 (Exponential localisation).

$$\mathbb{E} \left(\sup_{t \in \mathbb{R}} |\langle e_j, e^{-itH_\omega} e_k \rangle| \right) \leq C e^{-\mu|j-k|} \quad (2.2)$$

Proposition: *Exponential dynamical localization implies that all moments of the position operator are bounded in time, i.e., dynamical localisation*

Proof. To see how (1) follows from exp. dynamical localization, assume that $\psi(k) = 0$ for $|k| > R$. Then

$$\begin{aligned} \| |X|^p e^{-ith_\omega} \psi \|^2 &= \sum_j |\langle e_j, |X|^p e^{-ith_\omega} \psi \rangle|^2 \\ &= \sum_j |j|^{2p} \left| \sum_{|k| \leq R} \langle e_j, e^{-ith_\omega} e_k \rangle \psi(k) \right|^2 \\ &\leq \sum_j \sum_{|k| \leq R} |j|^{2p} |\langle e_j, e^{-ith_\omega} e_k \rangle|^2 \|\psi\|^2, \end{aligned}$$

where the last step used the Cauchy-Schwarz inequality. We can drop the square from $|\langle e_j, e^{-ith_\omega} e_k \rangle|^2$ (as this number is bounded by 1) and then take

expectations to get

$$\begin{aligned}
\mathbb{E} \left(\sup_t \| |X|^p e^{-ith_\omega} \psi \|^2 \right) &\leq \sum_j \sum_{|k| \leq R} |j|^{2p} \mathbb{E} \left(\sup_t |\langle e_j, e^{-ith_\omega} e_k \rangle| \right) \|\psi\|^2 \\
&\leq C \sum_j \sum_{|k| \leq R} |j|^{2p} e^{-\mu|j-k|} \|\psi\|^2 \\
&< \infty.
\end{aligned}$$

□

2.3 From dynamical to spectral localisation

Proposition 2.4. *Suppose that dynamical localization in the form*

$$\mathbb{E} \left(\sup_{t \in \mathbb{R}} |\langle e_j, e^{-ith_\omega} \chi_I(h_\omega) e_k \rangle| \right) \leq C e^{-\mu|j-k|} \quad (2.3)$$

holds in an open interval I . Then h_ω almost surely has pure point spectrum in I .

Proof. For a discrete Schrödinger operator $h = h_0 + V$ in $l^2(\mathbb{Z}^d)$ let $P_{\text{cont}}(h)$ be the projection onto its continuous spectral subspace. Then the RAGE-Theorem says that for every $\psi \in l^2(\mathbb{Z}^d)$,

$$\|P_{\text{cont}}(h) \chi_I(h) \psi\|^2 = \lim_{R \rightarrow \infty} \lim_{T \rightarrow \infty} \int_0^T \frac{dt}{T} \|\chi_{\{|x| \geq R\}} e^{-ith} \chi_I(h) \psi\|^2. \quad (2.4)$$

If ψ has finite support, say $\text{supp } \psi \subset \{|x| \leq r\}$, then

$$\begin{aligned}
\|\chi_{\{|x| \geq R\}} e^{-ith} \chi_I(h) \psi\|^2 &\leq \|\chi_{\{|x| \geq R\}} e^{-ith} \chi_I(h) \chi_{\{|x| \leq r\}}\|^2 \|\psi\|^2 \\
&\leq \|\chi_{\{|x| \geq R\}} e^{-ith} \chi_I(h) \chi_{\{|x| \leq r\}}\| \|\psi\|^2 \\
&\leq \sum_{|x| \geq R, |y| \leq r} |\langle e_x, e^{-ith} \chi_I(h) e_y \rangle| \|\psi\|^2,
\end{aligned}$$

where dropping a square is allowed as $\|\chi_{\{|x| \geq R\}} e^{-ith} \chi_I(h) \chi_{\{|x| \leq r\}}\| \leq 1$.

Taking expectations in (2) implies, after using Fatou and Fubini,

$$\mathbb{E} (\|P_{\text{cont}}(h_\omega) \chi_I(h_\omega) \psi\|^2) \leq \lim_{R \rightarrow \infty, T \rightarrow \infty} \int_0^T \frac{dt}{T} \sum_{|x| \geq R, |y| \leq r} \mathbb{E} (|\langle e_x, e^{-ith_\omega} \chi_I(h_\omega) e_y \rangle|) \|\psi\|^2. \quad (2.5)$$

By (2.4) we have $\mathbb{E} (|\langle e_x, e^{-ith_\omega} \chi_I(h_\omega) e_y \rangle|) \leq C e^{-\mu|x-y|}$ uniformly in t ,

which bounds the right hand side of (2.6) by

$$\leq \lim_{R \rightarrow \infty} \tilde{C} \sum_{|x| \geq R, |y| \leq r} e^{-\mu|x-y|} = 0.$$

We conclude that $P_{\text{cont}}(h_\omega)\chi_I(h_\omega)\psi = 0$ for almost every ω and every ψ of finite support. The latter are dense in $l^2(\mathbb{Z}^d)$ and thus $P_{\text{cont}}(h_\omega)\chi_I(h_\omega) = 0$ almost surely, meaning that the spectrum in I is pure point. $\square$

3 Fractional moments

In this section we are going to see the important lemmas and theorems that help us proving localisation. In particular we see the power of the fractional moment method.

3.1 The green function

The *resolvent operator* is defined as:

$$G(z) = (H - z)^{-1}$$

3.2 Localisation at large disorder

Theorem 3.1 (Aizenman–Molchanov, 1993). *Let $0 < s < 1$. Then there exists $\lambda_0 > 0$ such that for $\lambda \geq \lambda_0$ there are $C < \infty$ and $\mu > 0$ with*

$$\mathbb{E}(|G_{\omega,\lambda}(x, y; z)|^s) \leq Ce^{-\mu|x-y|} \quad (3.1)$$

uniformly in $x, y \in \mathbb{Z}^d$ and $z \in \mathbb{C} \setminus \mathbb{R}$.

Lemma 3.2 (A priori bound). *There exists a constant $C_1 = C_1(s, \rho) < \infty$ such that*

$$\mathbb{E}_x(|G_{\omega,\lambda}(x, x; z)|^s) \leq C_1 \lambda^{-s} \quad (3.2)$$

for all $x \in \mathbb{Z}^d$, $z \in \mathbb{C} \setminus \mathbb{R}$, and $\lambda > 0$.

Here

$$C_1 = \frac{1}{1-s}$$

for $\omega_x \sim \mathcal{U}[-1, 1]$

Lemma 3.3. RESOLVENT IDENTITY *cómo en el libro*

Proof of a priori bound. This proof demonstrates the simplicity of the fundamental idea underlying the FMM. For fixed $x \in \mathbb{Z}^d$, write $\omega = (\hat{\omega}, \omega_x)$ where $\hat{\omega}$ is short for $(\omega_u)_{u \in \mathbb{Z}^d \setminus \{x\}}$. With $P_{e_x} := \langle e_x, \cdot \rangle e_x$, the orthogonal projection onto the span of e_x , we can separate the ω_x and $\hat{\omega}$ dependence of H_ω as

$$H_\omega = -\Delta + \lambda V_{\hat{\omega}} + \lambda \omega_x P_{e_x} = H_{\hat{\omega}} + \lambda \omega_x P_{e_x}.$$

For the operators $(H_\omega - z)$ and $(H_{\hat{\omega}} - z)$, the resolvent identity yields

$$(H_\omega - z)^{-1} = (H_{\hat{\omega}} - z)^{-1} - \lambda \omega_x (H_{\hat{\omega}} - z)^{-1} P_{e_x} (H_\omega - z)^{-1}. \quad (3.3)$$

Taking matrix-elements we conclude for the corresponding diagonal Green functions that

$$G_{\omega,\lambda}(x, x; z) = G_{\hat{\omega},\lambda}(x, x; z) - \lambda \omega_x G_{\hat{\omega},\lambda}(x, x; z) G_{\omega,\lambda}(x, x; z) \quad (3.4)$$

and

$$G_{\omega,\lambda}(x, x; z) = \frac{1}{a + \lambda \omega_x} \quad \text{with } a = \frac{1}{G_{\hat{\omega},\lambda}(x, x; z)}. \quad (3.5)$$

Note that the latter is well-defined since one can easily check that $\text{Im } G_{\hat{\omega},\lambda}(x, x; z) > 0$ for $\text{Im } z > 0$.

The important fact is that a is a complex number which does not depend on ω_x . Thus, writing $\mathbb{E}_x(\dots) := \int \dots \rho(\omega_x) d\omega_x$ for $\rho(\omega_x) = \frac{1}{2}\chi_{[-1,1]}$, we find that

$$\mathbb{E}_x(|G_{\omega,\lambda}(x, x; z)|^s) = \frac{1}{2\lambda^s} \int_{-1}^1 \frac{d\omega_x}{|\frac{a}{\lambda} + \omega_x|^s} \leq \frac{1}{2\lambda^s} \int_{-1}^1 \frac{d\omega_x}{|\omega_x|^s} = \frac{1}{(1-s)\lambda^s}, \quad (3.6)$$

with $\frac{1}{1-s}$ independent of λ and a , and thus independent of $\hat{\omega}$, z and x . $\square$

3.2.1 From fractional moment's exponential bound to dynamical localisation

Lemma 3.4. *For a bounded and self-adjoint operator A ,*

$$\|(A - z)^{-1}\| \leq \frac{1}{\text{Im}(z)} \quad (3.7)$$

for all $z \in \mathbb{C} \setminus \mathbb{R}$.

Proof. For the definition of the operator norm, we know that

$$\|(A - z)^{-1}\| := \sup_{\mu \in \sigma((A-z)^{-1})} |\mu|$$

Let's consider a $\lambda \in \mathbb{C}$ and write $\mu = \frac{1}{\lambda - z}$. This means:

$$\begin{aligned} \frac{1}{\lambda - z} \in \sigma((A - z)^{-1}) &\iff (A - z)^{-1} - (\lambda - z)^{-1} \text{ not invertible} \\ &\iff (A - z)^{-1}(A - \lambda)(\lambda - z)^{-1} \text{ not invertible} \\ &\iff (A - \lambda) \text{ not invertible} \\ &\iff \lambda \in \sigma(A) \end{aligned}$$

Where the resolvent identity has been used in third Implication.
At this point we obtain

$$\|(A - z)^{-1}\| = \sup_{\lambda \in \sigma(A)} \frac{1}{|\lambda - z|} = \frac{1}{\inf_{\lambda \in \sigma(A)} |\lambda - z|} = \frac{1}{\text{dist}(\sigma(A), z)}$$

For a bounded and self-adjoint operator, we know that the spectrum is real-valued. For $z \in \mathbb{C} \setminus \mathbb{R}$ the minimum distance between z and $\sigma(A)$ is the imaginary component of z :

$$\|(A - z)^{-1}\| \leq \frac{1}{|\text{Im}(z)|}$$

□

4 SAW representation

4.1 SAW Definitions

- 1.

4.2 SAW representation of G

- Theorem of the representation
 1. first prove that $G(x, y) = G(x, x) \sum G(x_1, y)$
 2. then the representation with the R_N
- put figure of the grid and the different paths

4.3 Proof with SAW

5 From fractional moments to dynamical localisation

Theorem 5.1. *Theorem final*

Proposition 5.2.

$$|\Im z| \mathbb{E} (|G(x, y; z)|^2) \leq C_1 \mathbb{E} (|G(x, y; z)|^s)$$

Proof of theorem 5.1.

□