

Patients with $Z > 2$ vs the within-RPre split-half null (out of 10) — cohort screening only, NOT the trace test

$d_S \cdot \text{RPre} \rightarrow \text{TT}$	3/10	3/10	5/10	5/10	4/10	7/10
$d_P \cdot \text{RPre} \rightarrow \text{TT}$	3/10	4/10	6/10	7/10	3/10	8/10
$d_F \cdot \text{RPre} \rightarrow \text{TT}$	5/10	7/10	9/10	10/10	10/10	10/10
$d_S \cdot \text{RPre} \rightarrow \text{RPost}$	3/10	4/10	4/10	5/10	4/10	8/10
$d_P \cdot \text{RPre} \rightarrow \text{RPost}$	5/10	5/10	5/10	7/10	6/10	8/10
$d_F \cdot \text{RPre} \rightarrow \text{RPost}$	6/10	7/10	8/10	7/10	10/10	10/10
$d_S \cdot \text{TT} \rightarrow \text{RPost}$	1/10	2/10	1/10	5/10	4/10	8/10
$d_P \cdot \text{TT} \rightarrow \text{RPost}$	3/10	4/10	2/10	3/10	5/10	9/10
$d_F \cdot \text{TT} \rightarrow \text{RPost}$	3/10	7/10	9/10	10/10	10/10	10/10
	δ	θ	α	β	γ_l	γ_h

■ $n_+ \geq 8/10$ passes cohort screen

■ $5 \leq n_+ \leq 7/10$ marginal

■ $n_+ \leq 4/10$ fails