

Collection of plots

Airas Group

RStudio, PBC

2025/5/9 (updated: 2025-09-21)

Plots for model estimates

Dot-and-whisker plot

Dot-and-whisker plots visually summarize regression results by showing the point estimate (dot) and confidence interval (whiskers) for each predictor. They are useful for quickly comparing effect sizes and significance across multiple variables in a single, clean plot.

- We start from running logistic regression model
- We obtain estimate (with p-values, confidence intervals) for each of the regressor in the model


##	Predictor	OR	CI_low	CI_high	p_value	model
## 1	Age	1.25	1.10	1.40	0.010	Unadjusted
## 2	Age	1.30	1.15	1.50	0.020	Adjusted
## 3	Gender	0.85	0.65	1.05	0.040	Unadjusted
## 4	Gender	0.88	0.70	1.10	0.030	Adjusted
## 5	BMI	1.10	0.90	1.30	0.120	Unadjusted
## 6	BMI	1.20	1.05	1.35	0.100	Adjusted
## 7	HOT_PET	1.40	1.10	1.70	0.001	Unadjusted
## 8	HOT_PET	1.35	1.15	1.55	0.002	Adjusted

Now we need to arrange the data before plotting

From the model output (logistic regression) we need to run this:

```
dw_data <- results_clean %>%  
  
  rename(  
    estimate = OR,  
    conf.low = CI_low,  
    conf.high = CI_high,  
    term = Predictor  
  ) %>%  
  mutate(  
    model = factor(model, levels = c("Unadjusted", "Adjusted")),  
    term = fct_rev(factor(term)),  
    label = sprintf("%.2f", estimate),  
    sig = ifelse(p_value < 0.05, "Significant", "Not significant")  
  )
```

And on the `dw_data` arranged for the plot, we run the `dwplot` function:



Plots for accuracy/ frequency tables

Mosaic plot


Mosaic plots are used to visualize contingency tables. The size of each tile represents the proportion of observations in each group-category combination, allowing for quick comparison of relative frequencies.

We start from a contingency table as the following one:

```
mat <- matrix(c(73, 27, 23, 77),
              nrow = 2,
              byrow = TRUE,
              dimnames = list(
                Group = c("PF%< 2.12", "PF% >= 2.12"),
                Category = c("Low", "High")
              ))
mat
```

##		Category	
##	Group	Low	High
##	PF%< 2.12	73	27
##	PF% >= 2.12	23	77

Next we run the mosaicplot function:



Plots for longitudinal data

Spaghetti plot

A spaghetti plot is a type of line plot used to visualize longitudinal or repeated-measures data for multiple subjects. The key idea is to plot each subject's trajectory over time, so you can see both individual trends and the overall pattern.

- We start from a dataset for example with the following variables:

```
set.seed(42) # the seed allows to generate the same values for rBRLs
df <- data.frame(
  ID = rep(1:20, each = 2),
  Group = rep(c("Treated", "Untreated"), each = 20),
  Time = rep(c("baseline", "follow-up"), times = 20),
  rBRL = c(rnorm(20, 1, 0.2), rnorm(20, 1.5, 0.2))
)
```

- Since some values will coincide, we would like to randomly jitter them (spiking in the 2d-space)
- We will prepare the data before plotting:

```
set.seed(42) # the seed allows to generate the same plot
df_jittered <- df %>%
  mutate(
    Time_num = as.numeric(factor(Time, levels = c("baseline", "follow
    Time_jittered = Time_num + runif(n(), -0.08, 0.08) # jitter for (
  )
```

Before plotting calculate the means:

```
# Group means (without jitter)
group_means <- df %>%
  group_by(Group, Time) %>%
  summarise(mean_rBRL = mean(rBRL), .groups = "drop") %>%
  mutate(Time_num = as.numeric(factor(Time, levels = c("baseline", "1
```

Run entirely the code

