

CERN-EP-2025-193
2025/09/16

CMS-JME-24-001

DEEPMET: Improving missing transverse momentum estimation with a deep neural network

The CMS Collaboration*

Abstract

At hadron colliders, the net transverse momentum of particles that do not interact with the detector (missing transverse momentum, \vec{p}_T^{miss}) is a crucial observable in many analyses. In the standard model, \vec{p}_T^{miss} originates from neutrinos. Many beyond-the-standard-model particles, such as dark matter candidates, are also expected to leave the experimental apparatus undetected. This paper presents a novel \vec{p}_T^{miss} estimator, DEEPMET, which is based on deep neural networks that were developed by the CMS Collaboration at the LHC. The DEEPMET algorithm produces a weight for each reconstructed particle based on its properties. The estimator is based on the negative vector sum of the weighted transverse momenta of all reconstructed particles in an event. Compared with other estimators currently employed by CMS, DEEPMET improves the \vec{p}_T^{miss} resolution by 10–30%, shows improvement for a wide range of final states, is easier to train, and is more resilient against the effects of additional proton-proton interactions accompanying the collision of interest.

Submitted to Physical Review D

1 Introduction

At the CERN LHC, experiments, such as CMS [1] and ATLAS [2], use reconstruction algorithms that estimate the four-momentum, production vertex, and type of all final-state particles to extract the maximum information from the high energy proton-proton (pp) collision data. The reconstructed particles consist of muons, electrons, photons, as well as charged and neutral hadrons, which are then clustered into particle jets to classify quarks, gluons, and tau leptons. The presence of particles that do not interact with the detector, such as standard model (SM) neutrinos or new electrically neutral, weakly interacting particles, e.g., dark matter particles, can only be inferred indirectly, by measuring the so-called missing transverse momentum, denoted as \vec{p}_T^{miss} , with magnitude p_T^{miss} . At a hadron collider, the component transverse to the beam direction of the net momentum of the initial-state partons is close to zero. The negative of the vector sum of all reconstructed particles' transverse momenta therefore estimates \vec{p}_T^{miss} :

$$\vec{p}_T^{\text{miss}} = - \sum_{i \in \text{all reco.}} \vec{p}_{T,i}. \quad (1)$$

The \vec{p}_T^{miss} stemming from weakly interacting particles will be collectively referred to as “genuine \vec{p}_T^{miss} ” in what follows. The accuracy of the \vec{p}_T^{miss} estimate is affected by the particle reconstruction efficiencies and p_T resolutions, detector finite coverage and noise, and the ability of the reconstruction algorithm to include particles from the leading pp interaction, while rejecting the potentially large number of low-momentum particles from additional pp interactions in the same or nearby bunch crossings (pileup). These factors collectively make it challenging to measure \vec{p}_T^{miss} with high accuracy and resolution. Pileup mitigation algorithms have been developed that are effective in reducing this contribution to \vec{p}_T^{miss} resolution [3–5].

The complexity of the \vec{p}_T^{miss} measurement suggests that modern machine-learning (ML) techniques could yield a more accurate estimator. The CMS Collaboration has done previous work showing the utility of this approach. An estimator that is based on multivariate analysis (MVA) technique using boosted decision trees, referred to as MVA- \vec{p}_T^{miss} [6], is employed in CMS Run-1 analyses and demonstrates a relative improvement in precision of around 20% compared with a more traditional calculation [7], particularly for the $H \rightarrow \tau\tau$ process. However, because MVA- \vec{p}_T^{miss} was trained on specific physics processes and their global event properties, it does not, in general, have optimal performance for other physics processes and kinematic regions.

This paper introduces a new algorithm DEEPMET based on a deep neural network (DNN). For each reconstructed particle in the event, the DNN produces a unique weight based on the properties of the particles. The DEEPMET \vec{p}_T^{miss} estimator is based on the negative of the vector sum of the weighted transverse momenta of all particles in an event. In this paper, we describe DEEPMET and compare its performance to that of other \vec{p}_T^{miss} estimators. Our performance metrics are based on the response and resolution for the x - and y -components of a \vec{p}_T^{miss} estimator, the components parallel and perpendicular to the transverse direction of a Z boson (for samples enriched in this process), or the absolute value of p_T^{miss} . The response is defined as the ratio of the measured average to the true average. The response can be smaller than unity due to e.g., particles below reconstruction thresholds or the response of the calorimeters. It can be larger due to resolution effects, noise, and pileup. The resolution can be calculated both with and without correction for the response. Since it can be difficult to know the true \vec{p}_T^{miss} in an event, artificial \vec{p}_T^{miss} can be induced in events with little to no genuine \vec{p}_T^{miss} by excluding leptons from the \vec{p}_T^{miss} calculation (e.g., the muons in a $Z \rightarrow \mu\mu$ event) and comparing with the corresponding precisely measured quantity calculated from the leptons.

Compared with conventional \vec{p}_T^{miss} estimators, DEEPMET shows a relative improvement in the \vec{p}_T^{miss} resolution of 10–30%, and is more resilient to the effects of pileup. Due to the choice of

DNN architecture and its inputs, i.e., the usage of reconstructed particles and the exclusion of event-level observables, the algorithm has limited information about the event topology and is trained only with typical pp events. As a consequence, DEEPMET is trained quickly and generalizes well, showing similarly strong performance in both collision data and a diverse set of simulated physics processes.

The paper is structured as follows. Section 2 gives an overview of the CMS detector. In Section 3, a summary of existing algorithms is presented, along with their strengths and weaknesses. Sections 4 and 5 contain details of the data sets and the object and event selections. After a brief introduction to the \vec{p}_T^{miss} parametrization, Section 6 details the algorithm, its inputs and outputs, and the associated loss functions. Performance metrics and comparisons with alternative algorithms are discussed in Section 7. A method to calibrate simulation performance to align with real collision data is outlined in Section 8. Section 9 explores the source of the enhanced performance of DEEPMET. Benchmarks for the computational needs are given in Section 10. Finally, a summary is presented in Section 11.

Tabulated results are provided in the HEPData record for the performance comparisons in this analysis [8].

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward hadron (HF) calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are detected in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, is reported in Ref. [1].

Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events of interest at a rate of around 100 kHz within a fixed time interval of about 4 μs [9]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to $\mathcal{O}(1 \text{ kHz})$ before data storage [10, 11].

3 Reconstruction, pileup mitigation, and machine learning

3.1 Particle-flow reconstruction

Primary collision vertices (PV) corresponding to pp collisions are found using reconstructed charged particles [12]. The primary pp interaction vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, is referred to as the leading primary vertex (LPV), as described in Section 9.4.1 of Ref. [13]. Other PVs are referred to as pileup vertices.

A particle-flow (PF) algorithm [14] aims to reconstruct and identify each individual particle in an event, using an optimized combination of information from the various elements of the CMS detector. The energy of a photon is obtained from the ECAL measurement. The energy

of an electron is determined from a combination of the electron momentum at the primary interaction vertex as determined by the tracker, the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron track. The p_T of a muon is obtained from the curvature of the corresponding track. The energy of a charged hadron is determined from a combination of the momentum measured in the tracker and the matching ECAL and HCAL energy deposits, corrected for the response function of the calorimeters to hadronic showers. Finally, the energy of a neutral hadron is obtained from the corresponding corrected ECAL and HCAL energies. The reconstructed particles from the PF algorithm are referred to as PF candidates.

Muons are measured in the range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. The single-muon trigger efficiency exceeds 90% over the full η range, and the efficiency to reconstruct and identify muons is greater than 96%. Matching muons to tracks measured in the silicon tracker results in a relative p_T resolution, for muons with p_T up to 100 GeV, of 1% in the barrel and 3% in the endcaps. Measurements made with cosmic ray muons show that, in the central region of the detector, the p_T resolution is better than 7% for muons with p_T up to 1 TeV [15].

For each event, hadronic jets are clustered from these reconstructed particles using the infrared and collinear safe anti- k_T algorithm [16, 17] with a distance parameter of 0.4. Jet momentum is determined as the vector sum of all particle momenta in the jet, and is found from simulation to be, on average, within 5 to 10% of the true momentum over the whole p_T spectrum and detector acceptance. Jet energy corrections are derived from simulation studies so that the average measured energy of jets becomes identical to that of particle level jets. In situ measurements of the momentum balance in dijet, photon+jet, Z+jet, and multijet events are used to determine any residual differences between the jet energy scale (JES) in data and in simulation, and appropriate corrections are made. The jet energy resolution (JER) typically amounts to 15–20% at 30 GeV, 10% at 100 GeV, and 5% at 1 TeV [18]. Additional selection criteria are applied to each jet to remove jets potentially dominated by instrumental effects or reconstruction failures.

The reconstruction also produces a \vec{p}_T^{miss} estimator, PF \vec{p}_T^{miss} , which is calculated using all reconstructed PF candidates in an event [19] and is further corrected for the JES of reconstructed jets. The \vec{p}_T^{miss} relies on the accurate measurement of the reconstructed physics objects, namely muons, electrons, photons, hadronically decaying taus, jets, and unclustered energy (E_U). The E_U is the contribution from the PF candidates not associated with any of the previous physics objects.

3.2 Pileup mitigation

Pileup affects jet and \vec{p}_T^{miss} measurements, since contributions from additional tracks and calorimetric energy depositions increase the apparent jet momentum and change the subsequent \vec{p}_T^{miss} estimation. To mitigate such effects on jets, charged-hadron subtraction (CHS) removes tracks identified as originating from pileup vertices from jets [3]. Corrections are then applied to remove on average the remaining contributions from neutral particles. This correction to the p_T of the jets is then propagated to the \vec{p}_T^{miss} estimation [18, 19].

The pileup-per-particle identification algorithm (PUPPI) [3–5] can additionally be used to mitigate the effect of pileup by making use of local shape information, event pileup properties, and tracking information. Charged particles identified as originating from pileup vertices are discarded. For each neutral particle, a local shape variable, that distinguishes collinear energy deposits originating from the hard scatter from soft/diffuse energy deposits attributable to particles originating from pileup interactions, is defined using the surrounding charged par-

ticles that are compatible with the LPV within the tracker acceptance ($|\eta| < 2.5$) and using all particles in the region outside of the tracker coverage. The momenta of the neutral particles are then rescaled according to their probability to originate from the LPV as deduced from the local shape variable. We denote this probability as the “PUPPI weight” in the following. The rescaling supersedes the need for jet-based offset corrections [3]. The corresponding \vec{p}_T^{miss} is referred to as PUPPI \vec{p}_T^{miss} . In the studies of this paper, “PUPPI v15” tune is used [5].

The CHS and PUPPI algorithms have been shown to improve the precision of \vec{p}_T^{miss} measurements significantly [19]. However, the PUPPI algorithm requires extensive tuning. Its efficacy depends both on the physics process and kinematic region, and it is less effective in low-pileup data. Consequently, its \vec{p}_T^{miss} resolution is not always optimal.

4 Data and simulated samples

This paper uses a data sample collected by the CMS experiment at a center-of-mass energy of 13 TeV during 2016. The data set corresponds to an integrated luminosity of $\mathcal{L} = 16.8 \text{ fb}^{-1}$, with an average pileup of 23 additional pp interactions [20].

Several Monte Carlo (MC) event generators are used to simulate the Z+jets and $t\bar{t}$ samples for the ML training and W+jets samples for the post-training performance studies. The V+jets ($V = W, Z$) samples are simulated at next-to-leading order (NLO) in perturbative quantum chromodynamics (QCD) with the MADGRAPH5_aMC@NLO 2.3.3 event generator [21]. The $t\bar{t}$ samples are simulated with POWHEG 2.0 [22–25]. The PYTHIA 8.230 [26] package is used for parton showering, hadronization, and the underlying event simulation, with the CP5 tune [27, 28]. The NNPDF 3.1 [29] set of parton distribution functions (PDFs) at next-to-NLO (NNLO) in QCD is used as the default.

Additional sets of simulated event samples are used to understand how the performance of the DEEPMET reconstruction depends on the physical process. The processes span a variety of final states, including Higgs bosons, top quarks, heavy-flavor jets, τ leptons, and beyond-the-SM dark-matter candidates. These samples are produced using the same PDFs with the same tunes. Four types of Higgs boson processes are generated with POWHEG at NLO in perturbative QCD: Higgs boson production via vector boson fusion with the Higgs boson decaying to invisible particles ($H \rightarrow \text{invisible}$), Higgs boson pair production via gluon fusion with one Higgs boson decaying to a pair of τ leptons and the other Higgs boson decaying to a pair of b quarks ($HH \rightarrow b\bar{b}\tau\tau$), and Higgs bosons produced in association with a $t\bar{t}$ pair followed by the Higgs boson either decaying to a pair of b quarks or to a pair of muons. Furthermore, two sets corresponding to the production of particles using the Simplified Model Spectra (SMS) framework of supersymmetry are considered [30, 31], both with R parity conservation and a lightest supersymmetric particle, a neutralino $\tilde{\chi}_1^0$ that acts as a dark-matter candidate. One set consists of top squark-antisquark pair production with a top squark mass of 350 GeV and a $\tilde{\chi}_1^0$ mass of 335 GeV. The final state containing two b quarks, four light quarks, and two $\tilde{\chi}_1^0$ particles [32] is selected (denoted SMS T2tt-4bd). The second set is pair production of two of the heavier neutralinos with mass 400 GeV, with each neutralino decaying into a Z boson and a $\tilde{\chi}_1^0$ with a mass of 1 GeV [33] (denoted SMS TChiZZ). Both of these processes are generated with MADGRAPH5_aMC@NLO in version 2.3.3 at LO. An overview of the simulated samples and their respective final states is given in Table 1.

The simulation of the interactions of all final-state particles with the CMS detector is done with GEANT4 [34]. The simulated events are reconstructed using the same algorithms used for the data. The simulated events include pileup, with the number of additional pp interactions

Table 1: Overview of the simulated event samples used for performance studies.

Process identifier	Production mode	Final-state objects
$Z(\ell\ell)+\text{jets}$	—	Charged leptons, jets
$t\bar{t} \rightarrow 2\ell 2\nu 2b$	—	Muons, b quark jets, prompt neutrinos
$W(\ell\nu)+\text{jets}$	—	Charged leptons, jets, prompt neutrinos
$H \rightarrow \text{invisible}$	Vector boson fusion (qqH)	Forward jets, prompt neutrinos
$HH \rightarrow b\bar{b}\tau\tau$	Gluon fusion (ggH)	τ leptons, b quark jets, prompt neutrinos
$t\bar{t}H(H \rightarrow b\bar{b})$	$t\bar{t}H$	Jets, b quark jets, charged leptons, prompt neutrinos
$t\bar{t}H(H \rightarrow \mu\mu)$	$t\bar{t}H$	Muons, jets, b quark jets, prompt neutrinos
SMS T2tt-4bd	$gg \rightarrow t\bar{t} \rightarrow 2b4q2\tilde{\chi}_1^0$	Jets, b quark jets, dark matter candidates
SMS TChiZZ	$gg \rightarrow \tilde{\chi}_2^0\tilde{\chi}_2^0 \rightarrow ZZ\tilde{\chi}_1^0\tilde{\chi}_1^0$	Charged leptons, jets, dark matter candidates

matching that observed in data.

5 Physics object and event selections

To select $W+\text{jets}$, $Z+\text{jets}$, and $t\bar{t}$ processes, candidate events are required to pass the isolated single-muon trigger that requires p_T above 24 GeV.

The selected muon candidates are required to satisfy a set of quality criteria based on the number of spatial measurements in the silicon tracker and the muon system, as well as the fit quality of the combined muon track [15]. Muon candidates are required to have $p_T > 25$ GeV, $|\eta| < 2.4$, and pass the so-called “tight” identification and isolation selections [15].

The $Z+\text{jets}$ and $t\bar{t}$ events used in the algorithm training described in Section 6 are required to have two muons with opposite charge passing the above selections. In the studies quantifying the algorithm performance described in Section 7.1, the dimuon invariant mass is also required to satisfy $80 < m(\mu\mu) < 100$ GeV in order to select $Z \rightarrow \mu\mu$ events and veto those with genuine \vec{p}_T^{miss} (mainly from τ leptons or electroweak decays inside the heavy-flavor jets).

The $W+\text{jets}$ simulated events and data used in Section 7.2 are required to have one muon candidate passing the muon selections.

For the studies presented in Section 7.3 validating the \vec{p}_T^{miss} performance in other simulated physics processes, no object or event selection is applied.

6 The DEEPMET algorithm

The DNN operates on each PF candidate individually. The candidate’s inputs include eight continuous and three categorical features:

- Continuous features:
 - d_{xy} , the impact parameter with respect to the LPV in the transverse plane for charged PF candidates, zero for neutral PF candidates [12];
 - d_z , the impact parameter with respect to the LPV along the beam direction for charged PF candidates, zero for neutral PF candidates;
 - the PUPPI weight;
 - five kinematic features per particle: η , mass, p_x , p_y , and p_T ;
- Categorical features:
 - Particle ID: a unique integer number for the seven different types of PF

candidates, i.e., electrons, muons, photons, charged hadrons, neutral hadrons, and hadronic and electromagnetic PF candidates in the HF;

- charge of the particle (-1 , 0 , or $+1$);
- an LPV association flag, referred to as the `fromLPV` flag. This flag indicates one of four possible associations between the LPV and a charged PF candidate: used in the LPV fit; used in a pileup vertex fit; not used in any of the PV fits, but closest to one of the pileup vertices in the z direction; and not used in any of the PV fits but closest to the LPV in the z direction.

The output of the DNN is a weight (w_i) and two bias terms ($b_{i,x}, b_{i,y}$). The DEEPMET estimator output is the negative of the weighted \vec{p}_T sum of all PF candidates together with their bias contributions. Its x and y components are given by:

$$p_x^{\text{miss}} = - \sum_i (w_i p_{i,x} + b_{i,x}), p_y^{\text{miss}} = - \sum_i (w_i p_{i,y} + b_{i,y}). \quad (2)$$

This calculation is similar to the PUPPI \vec{p}_T^{miss} calculation. The two bias terms in the model were initially intended to enable distinct adjustments in the x and y directions and to correct measurement biases. However, in practice it is found that they have minimal impact on the performance.

6.1 The DNN architecture

Figure 1 shows the DNN architecture. The inputs are the 11 features of each input PF candidate. After the input layer, each of the three categorical features will go through one embedding layer and become an 8-dimensional tensor, so that the feature space can be learned and the differences can be represented by distances in the embedded space.




Figure 1: The DEEPMET DNN architecture. For each event, all PF candidates in the event are considered as input. $N \times n$ represents the number of PF candidates in the event multiplied by the dimensionality of the per-particle feature space, where n can be 1, 2, 3, 8, 16, 32, and 64 in the architecture.

The three 8-dimensional tensors from the embedding layers are then concatenated together with the eight continuous features, and passed to a few dense (or fully-connected) layers with progressively reducing numbers of nodes. These layers act only on each particle individually and can also be considered as one-dimensional convolutions. After each dense layer, batch normalization [35] is applied (not shown in Fig. 1 for simplicity) to reduce the covariate shifts and make the training converge faster. The outputs of these dense layers are the weights w_i , and the biases $b_{i,x}$ and $b_{i,y}$ of all the input PF candidates. In the last layer, the event \vec{p}_T^{miss} is calculated following Eq. 2.

The total number of trainable parameters in the DEEPMET neural network is 4541. Such a small set of trainable parameters allows the training to converge quickly, and also significantly reduces the inference time. Studies have also been conducted using more layers and larger number of nodes per layer, and the effect on the performance was within a few percent.

The DNN architecture is implemented with KERAS [36], which acts as an interface for the TENSORFLOW [37] library. The ADAMW optimizer [38] is used for gradient optimizations.

6.2 Training

The DNN of DEEPMET is trained with the Z+jets and $t\bar{t}$ simulated samples discussed in Section 4. Events are required to pass the dimuon selections described in Section 5. Using two samples allows a wide range of jet multiplicities and particle p_T distributions: Z+jets events tend to have particles with low p_T ; the $t\bar{t}$ events tend to have high p_T and typically contain high- p_T neutrinos.

The \vec{p}_T^{miss} performance in V+jets processes is measured by comparing the momentum of the vector boson \vec{q}_T to that of the hadronic recoil system \vec{u}_T (defined as the vector p_T sum of all PF candidates except for the decay products of the vector boson) [19]. The vector boson momenta \vec{q}_T are measured as the vector p_T sum of the two muons for Z bosons, or taken directly from the generator-level information in simulations of W bosons.

The components of the hadronic recoil parallel and perpendicular to the boson momentum vector are denoted by u_{\parallel} and u_{\perp} , respectively. These are used to study the \vec{p}_T^{miss} response and resolution. The response of \vec{p}_T^{miss} is defined as $-\langle u_{\parallel} \rangle / \langle q_T \rangle$; the resolutions of u_{\parallel} and u_{\perp} are calculated using the root-mean-square of $u_{\parallel} + q_T$ and u_{\perp} distributions, and are denoted $\sigma(u_{\parallel})$ and $\sigma(u_{\perp})$, respectively [19].

Training is performed on an NVIDIA Quadro P6000 graphics card using a few tens of thousands of events from the Z+jets sample and approximately the same number of events from the $t\bar{t}$ sample. The batch size is set to 128, and the training and validation splitting is set to 4:1. The training takes about 30 minutes to converge to stable results.

Further studies are performed by scanning the mixture fractions of the two samples and the hyperparameters (such as the number and size of dense layers, learning rate, training and validation splitting, and the batch size), and by increasing the number of total events in the training and validation samples. DEEPMET performance is found to be only weakly sensitive to modifications of these (hyper)parameters.

6.3 Loss function

The loss function used in the training is the mean squared error loss:

$$L = \frac{1}{2} [(p_x^{\text{miss}} - \hat{p}_x^{\text{miss}})^2 + (p_y^{\text{miss}} - \hat{p}_y^{\text{miss}})^2] \quad (3)$$

where $p_{x,y}^{\text{miss}}$, $\hat{p}_{x,y}^{\text{miss}}$ are the predicted and generator-level components of \vec{p}_T^{miss} in the x, y directions.

7 The DEEPMET performance

The DEEPMET performance in data and MC simulations is discussed in this section. Improvements in resolutions are observed, with the response closer to unity, and more pileup resilience compared with PF \vec{p}_T^{miss} and PUPPI \vec{p}_T^{miss} .

7.1 Performance on Z+jets events

The \vec{p}_T^{miss} performance is studied using the events passing the Z+jets selections in Section 5. Such events should have little or no genuine \vec{p}_T^{miss} . The responses of different \vec{p}_T^{miss} estimators are shown in Fig. 2. Markers correspond to collision data, and dashed lines indicate MC simulations. The DEEPMET algorithm maintains a good, albeit 5-10% worse, response than the PF and PUPPI \vec{p}_T^{miss} estimators.




Figure 2: Recoil responses of different \vec{p}_T^{miss} estimators in data (markers) and MC simulations (dashed) after the $Z \rightarrow \mu\mu$ selections.

To properly account for the response effect on the resolutions, for each \vec{p}_T^{miss} estimator, the average response $-\langle u_{\parallel} \rangle / \langle q_T \rangle$ is calculated using the events in the response plateau region (i.e., $q_T > 150$ GeV), and the resolutions are scaled by the inverse of this average response, referred to as “response-corrected resolutions”. The response-corrected resolutions of u_{\parallel} and u_{\perp} vs. q_T and the number of reconstructed PVs are shown in Figs. 3 and 4, respectively. The DEEPMET algorithm maintains good resolutions and pileup resilience. Despite slightly worse response than the PF and PUPPI \vec{p}_T^{miss} estimators, DEEPMET resolutions are better even after including the response corrections. For example, compared with PUPPI \vec{p}_T^{miss} , the DEEPMET shows a 2-10% worse response but achieves a 20-30% resolution gain around 20 reconstructed primary vertices, significantly outperforming PUPPI in this regime.

The comparisons of response-corrected resolutions of different \vec{p}_T^{miss} estimators in $Z \rightarrow \mu\mu$ events for the low- and high- q_T regions are provided in Figs. 5 and 6, respectively, corresponding to the region where $q_T < (>) 50$ GeV. In both regions, PUPPI \vec{p}_T^{miss} is better than PF \vec{p}_T^{miss} , and DEEPMET outperforms both \vec{p}_T^{miss} estimators.

7.2 The PV-Agnostic DEEPMET and performance on W+jets events

The performance has also been studied in W+jets MC samples, in which the W boson decays into a muon and a(n) (anti)neutrino. Typically, this signal process involves a single isolated muon, genuine \vec{p}_T^{miss} , and several soft tracks as the p_T distribution of W bosons is low with the most probable value around 5 GeV [39]. In this study, the accuracy of LPV identification in approximately 1% of the events in the W+jets simulations is compromised, where the LPV is either reconstructed but incorrectly identified as a pileup vertex, or is not reconstructed at all. This occurs more frequently in events with muons at high $|\eta|$, where the quality of the muon




Figure 3: Response-corrected resolutions of u_{\parallel} (left) and u_{\perp} (right) vs. q_T of different \vec{p}_T^{miss} estimators in data after the $Z \rightarrow \mu\mu$ selections.




Figure 4: Response-corrected resolutions of u_{\parallel} (left) and u_{\perp} (right) vs. number of reconstructed PVs of different \vec{p}_T^{miss} estimators in data (solid) and MC simulations (dashed) after the $Z \rightarrow \mu\mu$ selections. The systematic uncertainties for PUPPI \vec{p}_T^{miss} due to the JES, the JER, and E_U are added in quadrature and displayed with the gray band.




Figure 5: Response-corrected resolutions of u_{\parallel} (left) and u_{\perp} (right) vs. number of reconstructed PVs of different \vec{p}_T^{miss} estimators in data after the $Z \rightarrow \mu\mu$ selections in the region with q_T smaller than 50 GeV.




Figure 6: Response-corrected resolutions of u_{\parallel} (left) and u_{\perp} (right) vs. number of reconstructed PVs of different \vec{p}_T^{miss} estimators in data after the $Z \rightarrow \mu\mu$ selections in the region with q_T larger than 50 GeV.

track is worse. Such mistakes in the LPV identification can result in incorrect PV-track association flags and PUPPI weights, which in turn introduce inaccuracies in subsequent calculations of PUPPI and DEEPMET \vec{p}_T^{miss} .

To handle such scenarios for high-precision $W(\mu\nu)$ analyses, a PV-Agnostic version of DEEPMET is developed, which is robust to the choice of the LPV from the PVs [40]. In this version of the algorithm, the default LPV selection is discarded, and instead the PV closest in z to the leading muon with $dz(\text{PV}, \mu) < 0.2 \text{ cm}$ is chosen. If there is no PV satisfying the requirement (i.e., the LPV is not successfully reconstructed), a “pseudo”-LPV is built, using (x, y) of the beam spot (luminous region of the pp collisions [12]), and the leading muon z position. In the DEEPMET inputs, d_z ’s and the PUPPI weights are re-computed with respect to the new LPV, while fromLPV and d_{xy} are not used, since they cannot be recalculated precisely. The model is retrained using the reduced set of inputs.

Figure 7 illustrates the performance comparisons of various \vec{p}_T^{miss} estimators for W +jets events, with solid (dashed) lines representing events where the LPV is correctly (incorrectly) identified. As anticipated, PF \vec{p}_T^{miss} is the estimator least impacted by LPV identification issues. Conversely, both PUPPI \vec{p}_T^{miss} and the original DEEPMET are significantly influenced. The PV-Agnostic approach helps to mitigate the dependence on LPV identifications, resulting in comparable performance between events with correct and incorrect LPV assignment. The remaining discrepancy is similar to that of PF \vec{p}_T^{miss} , which is primarily due to different kinematic distributions (e.g., larger fraction of LPV-incorrect events for those with muons at higher $|\eta|$).

The \vec{p}_T^{miss} and W boson transverse mass (defined as $m_T = \sqrt{2p_T^\mu p_T^{\text{miss}}[1 - \cos(\Delta\phi)]}$, where $\Delta\phi$ is the azimuthal angle difference between muon p_T and \vec{p}_T^{miss}) distributions are shown in Fig. 8 for data after the $W \rightarrow \mu\nu$ selections. Table 2 gives a \vec{p}_T^{miss} resolution metric: the higher half width at half maximum, defined as the difference between the value of p_T^{miss} (m_T) at the p_T^{miss} (m_T) probability density function (pdf) distribution maximum and the larger of the two values whose pdf is half the maximum value. For both variables, DEEPMET (PV-Agnostic) has a sharper Jacobian peak and thus provides better resolution.

Table 2: Higher half width at half maximum in p_T^{miss} and m_T in data after the $W \rightarrow \mu\nu$ selections.

Higher half width at half maximum / GeV	p_T^{miss}	m_T
PF	25.4	27.0
PUPPI	18.4	16.2
DEEPMET (PV-Agnostic)	15.2	13.0

7.3 Performance for other physics processes

The \vec{p}_T^{miss} resolution of different estimators is also compared for a number of physics processes with genuine \vec{p}_T^{miss} . The samples are listed in Section 4, and are used without any specific event selection applied. Figure 9 shows the p_T^{miss} resolution for the DEEPMET, PUPPI \vec{p}_T^{miss} , and PF \vec{p}_T^{miss} algorithms as a function of the sum of the p_T ’s of generated jets (H_T) for various physics processes for which p_T^{miss} is expected to have a crucial role in the event selection and/or reconstruction, and hence for the final physics sensitivity.

DEEPMET demonstrates superior resolution in p_T^{miss} for all considered physics processes, including Higgs boson production with decays to SM particles, Higgs boson decay to dark matter candidates, and the supersymmetric processes described in Section 4. These processes include a wide range of different jet and lepton multiplicities, jet flavors, and cover a broad kinematic range. The relative improvement in resolution is largest for low values of generator jet H_T , i.e.,




Figure 7: Response (upper), response-corrected resolutions of $u_{||}$ (lower left) and u_{\perp} (lower right) of different \vec{p}_T^{miss} estimators in W+jets MC simulations. Solid (dashed) lines are from events where the LPV is (in)correctly identified.




Figure 8: Distributions of p_T^{miss} (left) and m_T (right) of different \vec{p}_T^{miss} estimators in data after $W \rightarrow \mu\nu$ selections.

for low amounts of hard hadronic activity, indicating that DEEPMET particularly helps with the softer contributions to \vec{p}_T^{miss} .

8 The DEEPMET calibrations

The DEEPMET calibrations can be derived from Z+jets events by matching the performance in MC simulations to data. These corrections can be applied to the DNN outputs.

8.1 The XY correction

Due to asymmetries in the detector responses and resolutions, and also because of the offset of the beam spot position, etc., the reconstructed \vec{p}_T^{miss} response depends on ϕ . The average values of \vec{p}_T^{miss} in the x and y directions are calculated as a function of the number of PVs and then subtracted from the individual \vec{p}_T^{miss} components, centering the distributions at zero. The ϕ distributions of different \vec{p}_T^{miss} estimators before and after the xy correction are shown in Fig. 10. The distribution becomes flat, as expected, for both data and MC simulations after correction.

8.2 Scale and resolution calibration

Since the jet and PF candidate resolutions are typically worse in data than predicted from simulation, a quantile-based calibration is performed. The calibration is done in four main steps.

First the data and MC simulations are binned in q_T of the Z boson. For the data, the contributions from $t\bar{t}$ and diboson production are estimated from MC simulation and subtracted. Each bin's u_{\parallel} and u_{\perp} distribution is fitted with a double-Gaussian function.

In the second step, each of the fitted double-Gaussian functions is integrated to get the corresponding cumulative distribution functions (CDFs).




Figure 9: Comparison of p_T^{miss} resolution for various physics processes in simulated events. The considered processes are HH production via gluon fusion with $HH \rightarrow b\bar{b}\tau\tau$ (upper left), H production via vector boson fusion with $H \rightarrow \text{invisible}$ (upper right), ttH production with either $H \rightarrow b\bar{b}$ (middle left) or $H \rightarrow \mu\mu$ (middle right), the SMS T2b-4bd process (lower left), and the SMS TChiZZ process (lower right).




Figure 10: The ϕ distribution of different \vec{p}_T^{miss} estimators before (left) and after (right) the xy corrections, in data (markers) and MC simulations (dashed) after the $Z \rightarrow \mu\mu$ selections.

In MC simulation, the u_{\parallel} and u_{\perp} CDF values in the i th q_T bin can be found from:

$$\begin{aligned} p_{\parallel} &= \text{CDF}_i^{u_{\parallel}, \text{MC}}(u_{\parallel}^{\text{MC}}), \\ p_{\perp} &= \text{CDF}_i^{u_{\perp}, \text{MC}}(u_{\perp}^{\text{MC}}). \end{aligned} \quad (4)$$

In data, the corresponding u_{\parallel} and u_{\perp} CDF in the same q_T bin is given by:

$$\begin{aligned} u_{\parallel}^{\text{data}} &= (\text{CDF}_i^{u_{\parallel}, \text{data}})^{-1}(p_{\parallel}), \\ u_{\perp}^{\text{data}} &= (\text{CDF}_i^{u_{\perp}, \text{data}})^{-1}(p_{\perp}). \end{aligned} \quad (5)$$

Finally the differences $u_{\parallel}^{\text{data}} - u_{\parallel}^{\text{MC}}$ and $u_{\perp}^{\text{data}} - u_{\perp}^{\text{MC}}$ are applied as the corrections to u_{\parallel} and u_{\perp} in simulation, respectively. The corrected DEEPMET is then calculated from the corrected u_{\parallel} and u_{\perp} .

8.3 Data vs. simulation comparisons after calibration

Figure 11 illustrates the data-simulation comparisons for DEEPMET p_T^{miss} (upper left), recoil p_T (upper right), u_{\parallel} (lower left), and u_{\perp} (lower right) in Z +jets events, after applying the calibrations. A Z boson p_T reweighting is applied to align the boson p_T distribution between data and simulation. Good agreement between data and simulation after the calibrations is observed within the systematic uncertainties, which are detailed in Section 8.4.

Since the calibrations are only applied to MC simulations, all the performance results on data shown in the previous sections are unchanged after calibration.

8.4 Systematic uncertainties from calibration

The systematic uncertainties affecting DEEPMET are estimated during the calibration process. Three main sources are considered:

- *Jet multiplicity binning*: in the default setup, the Z boson p_T is used to derive corrections in different bins. One alternative choice is to have bins in both the p_T of the Z




Figure 11: Data-to-simulation comparisons of DEEPMET p_T^{miss} (upper left), recoil p_T (upper right), u_{\parallel} (lower left), and u_{\perp} (lower right) after the quantile correction. The underflow (overflow) contents are included in the first (last) bin. The gray band represents the systematic uncertainties discussed in Section 8.4.

boson and CHS jet multiplicity. A new set of corrections are derived using the same setup but with two-dimensional binning, and the difference between using both the p_T of the Z boson and the jet multiplicity, and using only the p_T of the Z boson, is chosen as one systematic uncertainty.

- *Choice of the modeling function:* in the default setup, a double-Gaussian function is chosen to model the u_{\parallel} and u_{\perp} distributions. To estimate the possible bias due to this choice, a Gaussian smoothing function is used to smooth the u_{\parallel} and u_{\perp} distributions, and then the CDFs and corrections are derived using the smoothed distributions. The correction difference between the double-Gaussian and Gaussian smoothing functions is chosen as one systematic uncertainty.
- *Background estimation and subtraction.* When fitting the data distributions, the background contributions from $t\bar{t}$ and diboson processes are estimated using MC simulation and then subtracted from data. To estimate the systematic uncertainties from the background subtraction, the contributions estimated from background MC samples are varied by $\pm 10\%$ (typical cross section measurement and theoretical uncertainties), and the difference between the calibrations using the central values and using these are taken as the systematic uncertainty.

The total uncertainties are shown in the gray band of Fig. 11. Around the peak of the Z+jets distributions (low p_T^{miss} and smaller $|u_{\perp}|$), the relative difference is around a few percent. In the tail regions the uncertainty can be as large as 10–15%.

9 Training without PUPPI weights

To understand the role and the importance of the PUPPI weights in DEEPMET’s performance, a version of DEEPMET is trained without the PUPPI weight information. Because of the DNN nature of the architecture, particle-level information is not shared between different PF candidates, and the PUPPI weight is the *only* input that contains local neighborhood information. This comparison sheds light on the relative importance of a particle’s neighborhood in the observed performance improvement.

Figure 12 shows comparisons of the response and response-corrected resolution of DEEPMET trained with (blue) and without (green) PUPPI weights, along with PF \vec{p}_T^{miss} and PUPPI \vec{p}_T^{miss} , after $Z \rightarrow \mu\mu$ selections. As can be seen for the $Z \rightarrow \mu\mu$ physics process, PUPPI weights mostly contribute to response improvements; the resolutions with and without PUPPI weights are similar.

10 Computational performance

The model trained using KERAS is stored as a frozen graph in TENSORFLOW format and integrated into the CMS software framework CMSSW for subsequent inference processes. The CMSSW framework is an open-source software framework utilized in triggering, data reconstruction, simulation, and offline analysis [41, 42]. In this setup, the number of input PF candidates is set to 4500, determined as the maximum count of PF candidates under the current operational conditions (typically ranging from 1500 to 2000). The model inputs consist of the 11 features of each PF candidate, as discussed in Section 6. In cases where there are fewer PF candidates, the input features are padded with zeros. The model processes these inputs to compute \vec{p}_T^{miss} in the x and y directions as the outputs. DEEPMET inference can be performed at any stage of data processing, as long as the list of PF candidates is provided.




Figure 12: Response (upper) and response-corrected resolutions of $u_{||}$ (lower left) and u_{\perp} (lower right) of different \vec{p}_T^{miss} estimators in data after the $Z \rightarrow \mu\mu$ selections.

The computational time is also examined. When running with one thread on a single CPU core, the average processing time for the DEEPMET inference within CMSSW is approximately 10 ms [43], while the total event processing time ranges from 1 to 10 s, depending on the event complexity and the associated data tier. Due to the compact size of the model, the average processing time for DEEPMET is much shorter than the overall CMSSW processing time.

11 Summary

This paper presents a new \vec{p}_T^{miss} estimator, DEEPMET, which is based on a deep neural network. The DEEPMET algorithm utilizes each individual particle reconstructed by the CMS particle-flow algorithm as input and assigns a weight w_i and two bias terms, $b_{i,x}$ and $b_{i,y}$, to each candidate. The estimated \vec{p}_T^{miss} is the negative vector sum of the weighted transverse momenta of all candidates, plus their bias contributions.

With 4541 trainable parameters, the training and deployment of DEEPMET is computationally efficient. DEEPMET is trained using Z+jets and tt samples, but achieves 10–30% better resolution compared with the current PF and PUPPI \vec{p}_T^{miss} estimators across multiple physics processes, such as Z+jets, W+jets, Higgs boson production, and processes with dark matter candidates. Another important feature of the DEEPMET algorithm is its high resilience to pileup, improving the physics reach in LHC Run 2 and Run 3, and future High-Luminosity LHC conditions. Specifically for the measurement of the W boson mass, a PV-Agnostic version of DEEPMET is designed to be more robust in $W \rightarrow \mu\nu$ events where the LPV is not always identified correctly. Events containing $Z \rightarrow \mu\mu$ decays are used to calibrate \vec{p}_T^{miss} , including corrections for asymmetries in detector response, as well as for \vec{p}_T^{miss} scale and resolution.

Good agreement between data and simulation is found. The DEEPMET estimator demonstrates the potential to improve the precision of SM measurements and to achieve higher sensitivity in beyond the standard model physics searches.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, TARISTU24-TK10 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMFTR, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIT and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES, NSC, and NAWA (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Türkiye); NASU (Ukraine); STFC (United Kingdom); DOE

and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, 101001205, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Beijing Municipal Science & Technology Commission, No. Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China under Grant No. 2023YFA1605804, and the Natural Science Foundation of China under Grant No. 12061141002 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ.KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing, FAIR – Future Artificial Intelligence Research, and CUP I53D23001070006 (Mission 4 Component 1), funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369, 2021/43/B/ST2/01552, 2023/49/B/ST2/03273, and the NAWA contract BPN/PPO/2021/1/00011 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, “European Union NextGenerationEU/PRTR”, and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G680009 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [2] ATLAS Collaboration, “The ATLAS experiment at the CERN Large Hadron Collider”, *JINST* **3** (2008) S08003, doi:10.1088/1748-0221/3/08/S08003.
- [3] CMS Collaboration, “Pileup mitigation at CMS in 13 TeV data”, *JINST* **15** (2020) P09018, doi:10.1088/1748-0221/15/09/P09018, arXiv:2003.00503.
- [4] D. Bertolini, P. Harris, M. Low, and N. Tran, “Pileup per particle identification”, *JHEP* **10** (2014) 059, doi:10.1007/JHEP10(2014)059, arXiv:1407.6013.
- [5] CMS Collaboration, “Pileup-per-particle identification: optimisation for Run 2 Legacy and beyond”, CMS Detector Performance Summary CMS-DP-2021-001, 2021.



- [6] CMS Collaboration, “Performance of the CMS missing transverse momentum reconstruction in pp data at $\sqrt{s} = 8$ TeV”, *JINST* **10** (2015) P02006, doi:10.1088/1748-0221/10/02/P02006, arXiv:1411.0511.
- [7] CMS Collaboration, “Evidence for the 125 GeV Higgs boson decaying to a pair of τ leptons”, *JHEP* **05** (2014) 104, doi:10.1007/JHEP05(2014)104, arXiv:1401.5041.
- [8] “HEPData record for this analysis”, 2025. doi:10.17182/hepdata.159179.
- [9] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [10] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [11] CMS Collaboration, “Performance of the CMS high-level trigger during LHC Run 2”, *JINST* **19** (2024) P11021, doi:10.1088/1748-0221/19/11/P11021, arXiv:2410.17038.
- [12] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [13] CMS Collaboration, “Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid”, CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02, 2015.
- [14] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.
- [15] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [16] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- k_T jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:10.1088/1126-6708/2008/04/063, arXiv:0802.1189.
- [17] M. Cacciari, G. P. Salam, and G. Soyez, “FastJet user manual”, *Eur. Phys. J. C* **72** (2012) 1896, doi:10.1140/epjc/s10052-012-1896-2, arXiv:1111.6097.
- [18] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, *JINST* **12** (2017) P02014, doi:10.1088/1748-0221/12/02/P02014, arXiv:1607.03663.
- [19] CMS Collaboration, “Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector”, *JINST* **14** (2019) P07004, doi:10.1088/1748-0221/14/07/P07004, arXiv:1903.06078.
- [20] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [21] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* **07** (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.

-
- [22] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.
 - [23] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with parton shower simulations: the POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
 - [24] S. Alioli, P. Nason, C. Oleari, and E. Re, “NLO vector-boson production matched with shower in POWHEG”, *JHEP* **07** (2008) 060, doi:10.1088/1126-6708/2008/07/060, arXiv:0805.4802.
 - [25] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX”, *JHEP* **06** (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.
 - [26] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
 - [27] P. Skands, S. Carrazza, and J. Rojo, “Tuning PYTHIA 8.1: the Monash 2013 tune”, *Eur. Phys. J. C* **74** (2014) 3024, doi:10.1140/epjc/s10052-014-3024-y, arXiv:1404.5630.
 - [28] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA 8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
 - [29] NNPDF Collaboration, “Parton distributions from high-precision collider data”, *Eur. Phys. J. C* **77** (2017) 663, doi:10.1140/epjc/s10052-017-5199-5, arXiv:1706.00428.
 - [30] J. Alwall, P. Schuster, and N. Toro, “Simplified models for a first characterization of new physics at the LHC”, *Phys. Rev. D* **79** (2009) 075020, doi:10.1103/PhysRevD.79.075020, arXiv:0810.3921.
 - [31] LHC New Physics Working Group, “Simplified models for LHC new physics searches”, *J. Phys. G* **39** (2012) 105005, doi:10.1088/0954-3899/39/10/105005, arXiv:1105.2838.
 - [32] CMS Collaboration, “Search for supersymmetry in proton-proton collisions at 13 TeV in final states with jets and missing transverse momentum”, *JHEP* **10** (2019) 244, doi:10.1007/JHEP10(2019)244, arXiv:1908.04722.
 - [33] CMS Collaboration, “Search for electroweak production of charginos and neutralinos at $\sqrt{s} = 13$ TeV in final states containing hadronic decays of WW, WZ, or WH and missing transverse momentum”, *Phys. Lett. B* **842** (2023) 137460, doi:10.1016/j.physletb.2022.137460, arXiv:2205.09597.
 - [34] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
 - [35] S. Ioffe and C. Szegedy, “Batch normalization: accelerating deep network training by reducing internal covariate shift”, 2015. arXiv:1502.03167.
 - [36] F. Chollet et al., “Keras”. <https://keras.io>, 2015.

















- [37] M. Abadi et al., “TensorFlow: A system for large-scale machine learning”, 2016.
arXiv:1605.08695.
- [38] I. Loshchilov and F. Hutter, “Decoupled weight decay regularization”, 2017.
arXiv:1711.05101.
- [39] CMS Collaboration, “Measurement of the transverse momentum spectra of weak vector bosons produced in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **02** (2017) 096,
doi:10.1007/JHEP02(2017)096, arXiv:1606.05864.
- [40] CMS Collaboration, “High-precision measurement of the W boson mass with the CMS experiment at the LHC”, 2024. arXiv:2412.13872. Submitted to *Nature*.
- [41] CMS Collaboration, “CMS physics: Technical design report volume 1: Detector performance and software”, CMS Technical Design Report CERN-LHCC-2006-001, CMS-TDR-8-1, 2006.
- [42] CMS Collaboration, “CMSSW on Github”. Accessed: 2023-11-08.
<http://cms-sw.github.io/>.
- [43] CMS Collaboration, “Portable acceleration of CMS computing workflows with coprocessors as a service”, *Comput. Softw. Big Sci.* **8** (2024) 17,
doi:10.1007/s41781-024-00124-1, arXiv:2402.15366.

A The CMS Collaboration





Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, V. Makarenko , A. Tumasyan¹ 

















Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , M. Dragicevic , C. Giordano, P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , M. Matthewman, I. Mikulec , J. Schieck² , R. Schöffbeck² , D. Schwarz , M. Shooshtari, M. Sonawane , W. Waltenberger , C.-E. Wulz² 















Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , H. Kwon , D. Ocampo Henao , T. Van Laer , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

J. Bierkens , N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, Y. Hong , P. Kashko , S. Lowette , I. Makarenko , D. Müller , J. Song , S. Tavernier , M. Tytgat³ , G.P. Van Onsem , S. Van Putte , D. Vannerom 














Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , A. Khalilzadeh, F.A. Khan , A. Malara , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer , F. Zhang 











Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , J. Knolle , D. Marckx , K. Skovpen , A.M. Tomaru, N. Van Den Bossche , J. van der Linden , J. Vandenbroeck , L. Wezenbeek 






















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Bein , A. Benecke , A. Bethani , G. Bruno , A. Cappati , J. De Favereau De Jeneret , C. Delaere , A. Giammanco , A.O. Guzel , V. Lemaitre, J. Lidrych , P. Malek , P. Mastrapasqua , S. Turkcapar 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , M. Barroso Ferreira Filho , E. Coelho , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera⁴ , P. Rebello Teles , M. Soeiro , E.J. Tonelli Manganote⁵ , A. Vilela Pereira⁴ 







Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁶ , M. Costa Reis , E.M. Da Costa , G.G. Da Silveira⁷ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza , S. S. Jesus , T. Laux Kuhn⁷ , M. Macedo , K. Mota Amarilo , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel , F. Torres Da Silva De Araujo⁸ 






Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁷ , F. Damas , T.R. Fernandez Perez Tomei , E.M. Gregores , B. Lopes Da Costa , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula , V. Scheurer




Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , P. Danev, R. Hadjiiska , P. Iaydjiev , M. Shopova , G. Sultanov 


University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 



Universidad Tecnica Federico Santa Maria, Valparaiso, Chile

W. Brooks 















Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Wang , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu, X. Wang 

Institute of High Energy Physics, Beijing, China

G.M. Chen⁹ , H.S. Chen⁹ , M. Chen⁹ , Y. Chen , Q. Hou , X. Hou, F. Iemmi , C.H. Jiang, A. Kapoor¹⁰ , H. Liao , G. Liu , Z.-A. Liu¹¹ , J.N. Song¹¹, S. Song, J. Tao , C. Wang⁹, J. Wang , H. Zhang , J. Zhao 


State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, Q. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, C. Quaranta , X. Sun , D. Wang , J. Wang, H. Yang, M. Zhang, Y. Zhao, C. Zhou 

State Key Laboratory of Nuclear Physics and Technology, Institute of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 




Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer^{12,13}, Z. Cui¹³, B. Li¹⁴, H. Wang , K. Yi¹⁵ , J. Zhang 






Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Y. Li





Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao¹⁶ 




Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo , A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega


Universidad de Antioquia, Medellin, Colombia

C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

N. Godinovic , D. Lelas , A. Sculac 









University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic , T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , C. Nicolaou, L. Paizanos , F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 


Charles University, Prague, Czech Republic

M. Finger[†] , M. Finger Jr. 

Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 


Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 








Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

S. Elgammal¹⁷, A. Ellithi Kamel¹⁸ 




Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

A. Hussein, H. Mohammed 


















National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , C. Nielsen , J. Pata , M. Raidal , N. Seeba , L. Tani 

Department of Physics, University of Helsinki, Helsinki, Finland

A. Milieva , K. Osterberg , M. Voutilainen 

Helsinki Institute of Physics, Helsinki, Finland

N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , R. Kumar Verma , T. Lampén , K. Lassila-Perini , B. Lehtela , S. Lehti , T. Lindén , N.R. Mancilla Xinto , M. Myllymäki , M.m. Rantanen , S. Saariokari , N.T. Toikka , J. Tuominiemi 


























Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

H. Kirschenmann , P. Luukka , H. Petrow 







IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France









M. Besancon , F. Couderc , M. Dejardin , D. Denegri, P. Devouge, J.L. Faure , F. Ferri , P. Gagne, S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , S. Ronchi , M.Ö. Sahin , A. Savoy-Navarro¹⁹ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

R. Amella Ranz , F. Beaudette , G. Boldrini , P. Busson , C. Charlot , M. Chiusi , T.D. Cuisset , O. Davignon , A. De Wit , T. Debnath , I.T. Ehle , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , L. Kalipoliti , M. Manoni , M. Nguyen , S. Obraztsov , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , G. Sokmen, L. Urda Gómez , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

















J.-L. Agram²⁰ , J. Andrea , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard 

G. Coulon, S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo , P. Depasse , H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg , G. Grenier , B. Ille , E. Jourdain, I.B. Laktineh, M. Lethuillier , B. Massoteau, L. Mirabito, A. Purohit , M. Vander Donckt , J. Xiao 


























Georgian Technical University, Tbilisi, Georgia

G. Adamov, I. Lomidze , Z. Tsamalaidze²¹ 

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser , P. Nattland , V. Oppenländer, A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 





























































RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

C. Daumann, S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , N. Kumar , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , F. Nowotny, A. Pozdnyakov , W. Redjeb , H. Reithler , U. Sarkar , V. Sarkisovi , A. Schmidt , C. Seth, A. Sharma , J.L. Spah , V. Vaulin, S. Zaleski






























RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany






M.R. Beckers , C. Dziwok , G. Flügge , N. Hoeflich , T. Kress , A. Nowack , O. Pooth , A. Stahl , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany






































H. Aarup Petersen , A. Abel, M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , I. Andreev , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman²² , K. Borras²³ , A. Campbell , S. Chatterjee , L.X. Coll Saravia , G. Eckerlin, D. Eckstein , E. Gallo²² , A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K.-Y. Lin , K. Lipka²⁴ , W. Lohmann²⁵ , J. Malvaso , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , J. Park , E. Ranken , A. Raspereza , D. Rastorguev , L. Rygaard, M. Scham^{26,23} , S. Schnake²³ , P. Schütze , C. Schwanenberger²² , D. Selivanova , K. Sharke , M. Shchedrolosiev , D. Stafford , M. Torkian, F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann, L. Wiens²³ , C. Wissing , Y. Yang , S. Zakharov, A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A.R. Alves Andrade , M. Antonello , S. Bollweg, M. Bonanomi , K. El Morabit , Y. Fischer , M. Frahm, E. Garutti , A. Grohsjean , A.A. Guvenli , J. Haller , D. Hundhausen, G. Kasieczka , P. Keicher , R. Klanner , W. Korcari , T. Kramer , C.c. Kuo, F. Labe , J. Lange , A. Lobanov , L. Moureaux , A. Nigamova , K. Nikolopoulos , A. Paasch , K.J. Pena Rodriguez , N. Prouvost, T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , P. Schleper , M. Schröder , J. Schwandt 

M. Sommerhalder , H. Stadie , G. Steinbrück , R. Ward , B. Wiederspan, M. Wolf 









Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , E. Butz , Y.M. Chen , T. Chwalek , A. Dierlamm , G.G. Dincer , U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann²⁷ , M. Horzela , F. Hummer , U. Husemann , J. Kieseler , M. Klute , R. Kunnilan Muhammed Rafeek, O. Lavoryk , J.M. Lawhorn , A. Lintuluoto , S. Maier , M. Mormile , Th. Müller , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , R. Schmieder, N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa , L. Stockmeier, K. Tauqeer, M. Toms , B. Topko , N. Trevisani , C. Verstege , T. Voigtländer , R.F. Von Cube , J. Von Den Driesch, M. Wassmer , R. Wolf , W.D. Zeuner , X. Zuo 




Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou , G. Daskalakis , A. Kyriakis 







National and Kapodistrian University of Athens, Athens, Greece

G. Melachroinos, Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , E. Tzovara , K. Vellidis , I. Zisopoulos 







National Technical University of Athens, Athens, Greece

T. Chatzistavrou, G. Karapostoli , K. Kousouris , E. Siamarkou, G. Tsipolitis 

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos, I. Evangelou , C. Foudas, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

D. Druzhkin , C. Hajdu , D. Horvath^{28,29} , K. Márton, A.J. Rádl³⁰ , F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³¹ , M.M.A. Gadallah³² , Á. Kadlecik , M. León Coello , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 













HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³¹ , F. Nemes³¹ , T. Novak , I. Szanyi³³ 




Panjab University, Chandigarh, India





S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁴ , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India









































































A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , S. Phor , K. Ranjan , M.K. Saini

















University of Hyderabad, Hyderabad, India

S. Acharya³⁵ , B. Gomber , B. Sahu³⁵ 






Indian Institute of Technology Kanpur, Kanpur, IndiaS. Mukherjee **Saha Institute of Nuclear Physics, HBNI, Kolkata, India**S. Baradia , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar**Indian Institute of Technology Madras, Madras, India**M.M. Ameen , P.K. Behera , S. Chatterjee , G. Dash , A. Dattamunsi, P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri³⁶ , T. Mishra , P.R. Pujahari , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay **IISER Mohali, India, Mohali, India**

B.K. Sirasva
























Tata Institute of Fundamental Research-A, Mumbai, IndiaL. Bhatt, S. Dugad , G.B. Mohanty , M. Shelake , P. Suryadevara**Tata Institute of Fundamental Research-B, Mumbai, India**A. Bala , S. Banerjee , S. Barman³⁷ , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, B.M. Joshi , S. Kumar , M. Maity³⁷, G. Majumder , K. Mazumdar , S. Parolia , R. Saxena , A. Thachayath **National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**S. Bahinipati³⁸ , D. Maity³⁹ , P. Mal , K. Naskar³⁹ , A. Nayak³⁹ , S. Nayak, K. Pal , R. Raturi, P. Sadangi, S.K. Swain , S. Varghese³⁹ , D. Vats³⁹ **Indian Institute of Science Education and Research (IISER), Pune, India**A. Alpana , S. Dube , P. Hazarika , B. Kansal , A. Laha , R. Sharma , S. Sharma , K.Y. Vaish **Indian Institute of Technology Hyderabad, Telangana, India**S. Ghosh **Isfahan University of Technology, Isfahan, Iran**H. Bakhshiansohi⁴⁰ , A. Jafari⁴¹ , V. Sedighzadeh Dalavi , M. Zeinali⁴² **Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**S. Bashiri , S. Chenarani⁴³ , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang⁴⁴ **University College Dublin, Dublin, Ireland**M. Felcini , M. Grunewald **INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy**M. Abbrescia^{a,b} , M. Barbieri^{a,b}, M. Buonsante^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , N. De Filippis^{a,c} , M. De Palma^{a,b} , W. Elmetenawee^{a,b,45} , N. Ferrara^{a,c} , L. Fiore^a , L. Longo^a , M. Louka^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} , S. My^{a,b} , F. Nenna^{a,b} , S. Nuzzo^{a,b} , A. Pellicchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} , Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a , A. Zaza^{a,b} **INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy**G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , P. Capiluppi^{a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} 

P. Giacomelli^a , L. Giommi^{a,b} , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,46} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a , F.L. Navarria^{a,b} , G. Paggi^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,47} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b} , A. Tricomi^{a,b,47} 





INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

J. Altork^{a,b} , P. Assiouras^a , G. Barbagli^a , G. Bardelli^a , M. Bartolini^{a,b} , A. Calandri^{a,b} , B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , L. Damenti^{a,b} , E. Focardi^{a,b} , T. Kello^a , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^a , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b} , G. Sguazzoni^a , L. Viliani^a 




























INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁴⁸ , D. Piccolo 









INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 


























INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , F. Brivio^a , V. Camagni^{a,b} , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , M.R. Kim^a , G. Lavizzari^{a,b} , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} , G. Pizzati^{a,b} , S. Ragazzi^{a,b} , T. Tabarelli de Fatis^{a,b} 










INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a , C. Di Fraia^{a,b} , F. Fabozzi^{a,c} , L. Favilla^{a,d} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,49} , P. Paolucci^{a,27} , B. Rossi^a 















INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università degli Studi di Cagliari^c, Cagliari, Italy

P. Azzi^a , N. Bacchetta^{a,50} , D. Bisello^{a,b} , P. Bortignon^{a,c} , G. Bortolato^{a,b} , A.C.M. Bulla^{a,c} , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^{a,51} , U. Gasparini^{a,b} , S. Giorgetti^a , F. Gonella^a , E. Lusiani^a , M. Margoni^{a,b} , A.T. Meneguzzo^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} , A. Triossi^{a,b} , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b} 

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy






































A. Braghieri^a , S. Calzaferri^a , P. Montagna^{a,b} , M. Pelliccioni^a , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy


























S. Ajmal^{a,b} , M.E. Ascioti^{a,b} , G.M. Bilei^a , C. Carrivale^{a,b} , D. Ciangottini^{a,b} , L. Della Penna^{a,b} , L. Fanò^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,52} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy









































C. Aimè^{a,b} , C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a 

R. Bhattacharya^a , L. Bianchini^{a,b} , T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , L. Calligaris^{a,b} , R. Castaldi^a , F. Cattafesta^{a,c} , M.A. Ciocci^{a,d} , M. Cipriani^{a,b} , R. Dell'Orso^a , S. Donato^{a,b} , R. Forti^{a,b} , A. Giassi^a , F. Ligabue^{a,c} , A.C. Marini^{a,b} , D. Matos Figueiredo^a , A. Messineo^{a,b} , S. Mishra^a , V.K. Muraleedharan Nair Bindhu^{a,b} , S. Nandan^a , F. Palla^a , M. Riggirello^{a,c} , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^{a,53} , T. Sarkar^a , A. Scribano^a , P. Solanki^{a,b} , P. Spagnolo^a , F. Tenchini^{a,b} , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a 









INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Akrap^{a,b} , C. Basile^{a,b} , S.C. Behera^a , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , F. De Ruggi^{a,b} , D. Del Re^{a,b} , E. Di Marco^a , M. Diemoz^a , F. Errico^a , L. Frosina^{a,b} , R. Gargiulo^{a,b} , B. Harikrishnan^{a,b} , F. Lombardi^{a,b} , E. Longo^{a,b} , L. Martikainen^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , N. Palmeri^{a,b} , R. Paramatti^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , V. Vladimirov^{a,b} 

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^{a,c} , R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , L. Lanteri^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , S. Maselli^a , A. Mecca^{a,b} , L. Menzio^{a,b} , P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , A. Staiano^a , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , E. Vlasov^{a,b} , R. White^{a,b} 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar^{a,b} , S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} , R. Delli Gatti^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , J. Kim, T. Kim , D. Lee, H. Lee , J. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , S. Sekmen , B. Tae, Y.C. Yang 






Department of Mathematics and Physics - GWNu, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon , J. Seo 

Hanyang University, Seoul, Korea

E. Asilar , F. Carnevali , J. Choi⁵⁴ , T.J. Kim , Y. Ryou 

Korea University, Seoul, Korea

S. Ha , S. Han, B. Hong , J. Kim , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , J. Shin , S. Yang 

Sejong University, Seoul, Korea








Y. Kang , H. S. Kim , Y. Kim , S. Lee 

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , H. Kim , J. Kim , T. Kim, Y. Kim, 

Y.W. Kim , S. Ko , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , J. Shin , U.K. Yang, I. Yoon 




University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, D. Kim , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

G. Cho, K. Hwang , B. Kim , S. Kim, K. Lee , H.D. Yoo 


Sungkyunkwan University, Suwon, Korea

M. Choi , Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy , Y. Gharbia 


Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 





Riga Technical University, Riga, Latvia

K. Dreimanis , O.M. Eberlins , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs , R. Plese , A. Potrebko , M. Seidel , D. Sidiropoulos Kontos 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , S. Nargelas , A. Rinkevicius , G. Tamulaitis 








National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff⁵⁵ , Z. Zolkapli



Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , A. Cota Rodriguez , L.E. Cuevas Picos, H.A. Encinas Acosta, L.G. Gallegos Maríñez, J.A. Murillo Quijada , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma , R. Lopez-Fernandez , J. Mejia Guisao , R. Reyes-Almanza , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 


Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 





University of Montenegro, Podgorica, Montenegro

I. Bubanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 






National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar , A. Awais , M.I.M. Awan, W.A. Khan 







AGH University of Krakow, Krakow, Poland

V. Avati, L. Forthomme , L. Grzanka , M. Malawski , K. Piotrkowski 

National Centre for Nuclear Research, Swierk, Poland

M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

















Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad 





Warsaw University of Technology, Warsaw, Poland

P. Fokow , K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi , G. Da Molin , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 


Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , L. Markovic , P. Milenovic , V. Milosevic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

D. Devetak , M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic , M. Stojanovic 

















Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Alcalde Martinez , J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , J. Cuchillo Ortega, B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , M. Gonzalez Hernandez , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , R. Paz Herrera , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , J. Vazquez Escobar 




















Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 


Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Ayllon Torresano , A. Cardini , J. Cuevas , J. Del Riego Badas , D. Estrada Acevedo , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , M. Obeso Menendez , E. Palencia Cortezon , J. Prado Pico , A. Soto Rodríguez , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasasa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , C. Quintana San Emeterio , L. Scodellaro , I. Vila , R. Vilar Cortabitarte , J.M. Vizán García 


































University of Colombo, Colombo, Sri Lanka

B. Kailasapathy⁵⁶ , D.D.C. Wickramarathna 













University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna⁵⁷ , K. Liyanage , N. Perera 































CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , R. Ardino , E. Auffray , J. Baechler, D. Barney , J. Bendavid , M. Bianco , A. Bocci , L. Borgonovi , C. Botta , A. Bragagnolo , C.E. Brown , C. Caillol , G. Cerminara , P. Connor , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , P.J. Fernández Manteca , B.A. Fontana Santos Alves , W. Funk , A. Gaddi, S. Giani, D. Gigi, K. Gill , F. Glege , M. Glowacki, A. Gruber , J. Hegeman , J.K. Heikkilä , R. Hofsaess , B. Huber , T. James , P. Janot , O. Kaluzinska , O. Karacheban²⁵ , G. Karathanasis , S. Laurila , P. Lecoq , C. Lourenço , A.-M. Lyon , M. Magherini , L. Malgeri , M. Mannelli , A. Mehta , F. Meijers , J.A. Merlin, S. Mersi , E. Meschi , M. Migliorini , F. Monti , F. Moortgat , M. Mulders , M. Musich , I. Neutelings , S. Orfanelli, F. Pantaleo , M. Pari , G. Petrucciani , A. Pfeiffer , M. Pierini , M. Pitt , H. Qu , D. Rabady , A. Reimers , B. Ribeiro Lopes , F. Riti , P. Rosado , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁵⁸ , A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , E. Vernazza , J. Wanczyk⁵⁹ , J. Wang, S. Wuchterl , M. Zarucki , P. Zehetner , P. Zejdl , G. Zevi Della Porta 


PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua⁶⁰ , L. Caminada⁶⁰ , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , U. Langenegger , L. Noehte⁶⁰ , T. Rohe , A. Samalan 

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'Archiacchiac⁵⁹ , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte , W. Lustermann , M. Malucchi , R.A. Manzoni , M. Marchegiani , L. Marchese , A. Mascellani⁵⁹ , F. Nessi-Tedaldi , F. Pauss , V. Perovic , B. Ristic , R. Seidita , J. Steggemann⁵⁹ , A. Tarabini , D. Valsecchi , R. Wallny 













Universität Zürich, Zurich, Switzerland

C. Amsler⁶¹ , P. Bärttschi , F. Bilandzija , M.F. Canelli , G. Celotto , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , T.H. Kwok , S. Leontsinis , V. Lukashenko , A. Macchiolo , F. Meng , M. Missiroli , J. Motta , P. Robmann, M. Senger , E. Shokr , F. Stäger , R. Tramontano , P. Viscone 

National Central University, Chung-Li, Taiwan

D. Bhowmik, C.M. Kuo, P.K. Rout , S. Taj , P.C. Tiwari³⁶ 


National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tsionou, H.y. Wu , E. Yazgan 
















High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangkuldee , N. Srimanobhas 


Tunis El Manar University, Tunis, Tunisia

Y. Maghrbi 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Dolek , I. Dumanoglu⁶² , Y. Guler⁶³ , E. Gurpinar Guler⁶³ , C. Isik , O. Kara , A. Kayis Topaksu , Y. Komurcu , G. Onengut , K. Ozdemir⁶⁴ , B. Tali⁶⁵ , U.G. Tok , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

M. Yalvac⁶⁶ 



Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi⁶⁷ , E. Gülmez , M. Kaya⁶⁸ , O. Kaya⁶⁹ , M.A. Sarkisla⁷⁰, S. Tekten⁷¹ 







Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak^{62,72} , S. Sen⁷³ 

Istanbul University, Istanbul, Turkey

B. Hacisahinoglu , I. Hos⁷⁴ , B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez 


Yildiz Technical University, Istanbul, Turkey

S. Cerci , C. Dozen⁷⁵ , B. Isildak⁷⁶ , E. Simsek , D. Sunar Cerci , T. Yetkin⁷⁵ 













Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , O. Dadazhanova, B. Grynyov 






















National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 









































University of Bristol, Bristol, United Kingdom

J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , D. Dharmender, H. Flacher , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw, M.S. Sanjrani⁴⁰, J. Segal, V.J. Smith 




Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁷⁷ , C. Brew , R.M. Brown , D.J.A. Cockerill , A. Elliot , K.V. Ellis, J. Gajownik , K. Harder , S. Harper , J. Linacre , K. Manolopoulos, M. Moallemi , D.M. Newbold , E. Olaiya , D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 

Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , O. Buchmuller, C.A. Carrillo Montoya , D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer, G. Fedi , G. Hall , H.R. Hoorani , A. Howard, G. Iles , C.R. Knight , P. Krueper , J. Langford , K.H. Law , J. León Holgado , E. Leutgeb , L. Lyons , A.-M. Magnan , B. Maier , S. Mallios, A. Mastronikolis , M. Mieskolainen , J. Nash⁷⁸ , M. Pesaresi , P.B. Pradeep , B.C. Radburn-Smith , A. Richards, A. Rose , L. Russell , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , T. Virdee²⁷ , M. Vojinovic , N. Wardle , D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA






S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann 

K. Hatakeyama , V. Hegde , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant ,
C. Sutantawibul , J. Wilson 

Bethel University, St. Paul, Minnesota, USA

J.M. Hogan⁷⁹ 


















Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , S. Raj , A.E. Simsek , S.S. Yu 




















The University of Alabama, Tuscaloosa, Alabama, USA

B. Bam , A. Buchot Perraguin , S. Campbell, R. Chudasama , S.I. Cooper , C. Crovella ,
G. Fidalgo , S.V. Gleyzer , A. Khukhunaishvili , K. Matchev , E. Pearson, C.U. Perez ,
P. Rumerio⁸⁰ , E. Usai , R. Yi 

















Boston University, Boston, Massachusetts, USA

S. Cholak , G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , J. Fulcher , F. Golf , S. Jeon , J. O'Cain, I. Reed ,
J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , E. Wurtz,
A.G. Zecchinelli 














Brown University, Providence, Rhode Island, USA

G. Barone , G. Benelli , D. Cutts , S. Ellis , L. Gouskos , M. Hadley , U. Heintz ,
K.W. Ho , T. Kwon , L. Lambrecht , G. Landsberg , K.T. Lau , J. Luo , S. Mondal ,
J. Roloff, T. Russell , S. Sagir⁸¹ , X. Shen , M. Stamenkovic , N. Venkatasubramanian 




University of California, Davis, Davis, California, USA

S. Abbott , B. Barton , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez ,
E. Cannaert, M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , O. Kukral ,
G. Mocellin , S. Ostrom , I. Salazar Segovia, J.S. Tafoya Vargas , W. Wei , S. Yoo 
























University of California, Los Angeles, California, USA

K. Adamidis, M. Bachtis , D. Campos, R. Cousins , A. Datta , G. Flores Avila ,
J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo , E. Manca ,
A. Nunez Del Prado , D. Saltzberg , V. Valuev 




















University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , G. Hanson 











University of California, San Diego, La Jolla, California, USA

A. Aportela , A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , B. D'Anzi ,
D. Diaz , J. Duarte , L. Giannini , Y. Gu, J. Guiang , V. Krutelyov , R. Lee ,
J. Letts , H. Li, M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , D. Primosch,
M. Quinnan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , A. Yagil , Z. Zhao


















University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , S. Carron Montero⁸² , K. Downham ,
C. Grieco , M.M. Hussain, J. Incandela , M.W.K. Lai, A.J. Li , P. Masterson ,
J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart ,
T.Á. Vámi , X. Yan , D. Zhang 

California Institute of Technology, Pasadena, California, USA

A. Albert , S. Bhattacharya , A. Bornheim , O. Cerri, R. Kansal , J. Mao , H.B. Newman ,
G. Reales Gutiérrez, T. Sievert, M. Spiropulu , J.R. Vlimant , R.A. Wynne , S. Xie 


Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , M. Cremonesi, V. Dutta , E.Y. Ertorer , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, M. Kanemura, C. Liu , P. Meiring , T. Mudholkar , S. Murthy , P. Palit , K. Park , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 






























































University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , S. Kwan , J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 




















Cornell University, Ithaca, New York, USA

J. Alexander , X. Chen , J. Dickinson , A. Duquette, J. Fan , X. Fan , J. Grassi , S. Hogan , P. Kotamnives , J. Monroy , G. Niendorf , M. Oshiro , J.R. Patterson , A. Ryd , J. Thom , P. Wittich , R. Zou , L. Zygala 













Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , C. Cosby , G. Cummings , I. Dutta , V.D. Elvira , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , T.C. Herwig , J. Hirschauer , V. Innocente , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijsma , B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , K. Maeshima , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁸³ , C.E. Perez Lara , F. Ravera , A. Reinsvold Hall⁸⁴ , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , C. Wang , I. Zoi 

University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , V. Sulimov , Y. Takahashi , J. Wang 






















Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhimi , A. Askew , S. Bower , R. Goff, R. Hashmi , A. Hassani , R.S. Kim , T. Kolberg , G. Martinez , M. Mazza , H. Prosper , P.R. Prova, R. Yohay 










Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , S. Butalla , S. Das , M. Hohlmann , M. Lavinsky, E. Yanes

University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , N. Barnett, A. Baty , C. Bennett , R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , H. Gupta , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , C. Mills , S. Nanda , G. Nigmatkulov , B. Ozek , T. Phan, D. Pilipovic , R. Pradhan , E. Prifti, P. Roy, T. Roy , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , J. Yoo 























The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend , K. Dilsiz⁸⁵ , O.K. Köseyan , A. Mestvirishvili⁸⁶ , O. Neogi, H. Ogul⁸⁷ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁸⁸ 










Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , S. Sekhar , M.V. Srivastav , M. Swartz 



















The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , S. Popescu⁸⁹ , C. Rogan , C. Royon , S. Rudrabhatla , S. Sanders , C. Smith , G. Wilson 



































Kansas State University, Manhattan, Kansas, USA

B. Allmond , N. Islam , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , G.G. Reddy , D. Roy , G. Sorrentino 


University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim , S. Lascio , P. Major , A.C. Mignerey , C. Palmer , C. Papageorgakis , M.M. Paranjpe , E. Popova⁹⁰ , A. Shevelev , L. Zhang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

C. Baldenegro Barrera , H. Bossi , S. Bright-Thonney , I.A. Cali , Y.c. Chen , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov , G. Grosso , P. Harris , D. Hoang , G.M. Innocenti , D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , T.a. Sheng , G.S.F. Stephans , D. Walter , Z. Wang , B. Wyslouch , T. J. Yang 















University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , W.J. Jackson , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , R. Rusack , O. Sancar , R. Saradhy , N. Strobbe 




















University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , A. Rohilla , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan 






State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava , A. Mandal , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar , A. Williams , P. Young , D. Yu 
































Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma , M. Campana , J. Dervan , Y. Haddad , Y. Han , I. Israr , A. Krishna , M. Lu , N. Manganelli , R. McCarthy , D.M. Morse , T. Orimoto , L. Skinnari , C.S. Thoreson , E. Tsai , D. Wood 











Northwestern University, Evanston, Illinois, USA

S. Dittmer , K.A. Hahn , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliencio , M. Velasco , J. Wang 



















University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci , S. Castells , A. Das , A. Ehnis , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , T. Ivanov , C. Jessop , A. Karneyeu , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano , N. Marinelli , I. Mcalister , T. McCauley , C. Mcgrady , C. Moore , Y. Musienko²¹ , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan , M. Wayne , H. Yockey 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , M. Carrigan , R. De Los Santos , L.S. Durkin , C. Hill , M. Joyce ,
M. Nunez Ornelas , D.A. Wenzl , B.L. Winer , B. R. Yates 




















Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , G. Dezoort , P. Elmer , A. Frankenthal , M. Galli , B. Greenberg ,
N. Haubrich , K. Kennedy , G. Kopp , Y. Lai , D. Lange , A. Loeliger , D. Marlow ,
I. Ojalvo , J. Olsen , F. Simpson , D. Stickland , C. Tully 




University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma 



















Purdue University, West Lafayette, Indiana, USA

S. Chandra , R. Chawla , A. Gu , L. Gutay , M. Jones , A.W. Jung , D. Kondratyev ,
M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , N.R. Saha ,
J.F. Schulte , F. Wang , A. Wildridge , W. Xie , Y. Yao , Y. Zhong 











Purdue University Northwest, Hammond, Indiana, USA

N. Parashar , A. Pathak , E. Shumka 






















Rice University, Houston, Texas, USA

D. Acosta , A. Agrawal , C. Arbour , T. Carnahan , P. Das , K.M. Ecklund , S. Freed,
P. Gardner, F.J.M. Geurts , T. Huang , I. Krommydas , N. Lewis, W. Li , J. Lin ,
O. Miguel Colin , B.P. Padley , R. Redjimi , J. Rotter , M. Wulansatiti , E. Yigitbasi ,
Y. Zhang 

University of Rochester, Rochester, New York, USA

O. Bessidskaia Bylund, A. Bodek , P. de Barbaro[†] , R. Demina , A. Garcia-Bellido ,
H.S. Hare , O. Hindrichs , N. Parmar , P. Parygin⁹⁰ , H. Seo , R. Taus 















Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , S. Donnelly, D. Gadkari , Y. Gershtein ,
E. Halkiadakis , C. Houghton , D. Jaroslawski , A. Kobert , S. Konstantinou ,
I. Laflotte , A. Lath , J. Martins , M. Perez Prada , B. Rand , J. Reichert , P. Saha ,
S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , J. Harris, T. Holmes , A.R. Kanuganti ,
N. Karunarathna , J. Lawless, L. Lee , E. Nibigira , B. Skipworth, S. Spanier 













Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov , A. Bolshov, O. Bouhali⁹¹ ,
A. Cagnotta , V. D'Amante , R. Eusebi , P. Flanagan , J. Gilmore , Y. Guo, T. Kamon ,
S. Luo , R. Mueller , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , Y. Feng , N. Gogate , Y. Kazhykarim, K. Lamichhane ,
S.W. Lee , C. Madrid , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli ,
A. Melo , D. Rathjens , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen ,
J. Zhang

























University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , H. Chung , B. Cox , J. Hakala , G. Hamilton Ilha Machado, R. Hirosky , M. Jose, A. Ledovskoy , C. Mantilla , C. Neu , C. Ramón Álvarez , Z. Wu

Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , S. Lomte , R. Loveless , A. Mallampalli , A. Mohammadi , S. Mondal, T. Nelson, G. Parida , L. Pétré , D. Pinna , A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , Yu. Andreev , T. Aushev , D. Budkouski , R. Chistov , M. Danilov , T. Dimova , A. Ershov , S. Gninenko , I. Gorbunov , A. Gribushin , A. Kamenev , V. Karjavine , M. Kirsanov , V. Klyukhin , O. Kodolova⁹² , V. Koronkov , I. Korsakov, A. Kozyrev , N. Krasnikov , A. Lanev , A. Malakhov , V. Matveev , A. Nikitenko^{93,92} , V. Palichik , V. Perelygin , S. Petrushanko , S. Polikarpov , O. Radchenko , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen , K. Slizhevskiy, V. Smirnov , O. Teryaev , I. Tlisova , A. Toropin , N. Voytishin , A. Zarubin , I. Zhizhin 

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

E. Boos , V. Bunichev , M. Dubinin⁸³ , V. Savrin , A. Snigirev , L. Dudko , K. Ivanov , V. Kim²¹ , V. Murzin , V. Oreshkin , D. Sosnov 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Ghent University, Ghent, Belgium

⁴Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

⁶Also at Universidade Estadual de Campinas, Campinas, Brazil

⁷Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁸Also at The University of the State of Amazonas, Manaus, Brazil

⁹Also at University of Chinese Academy of Sciences, Beijing, China

¹⁰Also at China Center of Advanced Science and Technology, Beijing, China

¹¹Also at University of Chinese Academy of Sciences, Beijing, China

¹²Also at School of Physics, Zhengzhou University, Zhengzhou, China

¹³Now at Henan Normal University, Xinxiang, China

¹⁴Also at University of Shanghai for Science and Technology, Shanghai, China

¹⁵Now at The University of Iowa, Iowa City, Iowa, USA

¹⁶Also at Center for High Energy Physics, Peking University, Beijing, China

¹⁷Now at British University in Egypt, Cairo, Egypt

¹⁸Now at Cairo University, Cairo, Egypt

¹⁹Also at Purdue University, West Lafayette, Indiana, USA

²⁰Also at Université de Haute Alsace, Mulhouse, France

²¹Also at an institute formerly covered by a cooperation agreement with CERN

²²Also at University of Hamburg, Hamburg, Germany

²³Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

- ²⁴Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- ²⁵Also at Brandenburg University of Technology, Cottbus, Germany
- ²⁶Also at Forschungszentrum Jülich, Juelich, Germany
- ²⁷Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ²⁸Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary
- ²⁹Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania
- ³⁰Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
- ³¹Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- ³²Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ³³Also at The University of Kansas, Lawrence, Kansas, USA
- ³⁴Also at Punjab Agricultural University, Ludhiana, India
- ³⁵Also at University of Hyderabad, Hyderabad, India
- ³⁶Also at Indian Institute of Science (IISc), Bangalore, India
- ³⁷Also at University of Visva-Bharati, Santiniketan, India
- ³⁸Also at IIT Bhubaneswar, Bhubaneswar, India
- ³⁹Also at Institute of Physics, Bhubaneswar, India
- ⁴⁰Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴¹Also at Isfahan University of Technology, Isfahan, Iran
- ⁴²Also at Sharif University of Technology, Tehran, Iran
- ⁴³Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁴Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁵Also at Helwan University, Cairo, Egypt
- ⁴⁶Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁷Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁸Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁴⁹Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁵⁰Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵¹Also at Lulea University of Technology, Lulea, Sweden
- ⁵²Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵³Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- ⁵⁴Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵⁵Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁶Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁷Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁵⁸Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁵⁹Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶⁰Also at Universität Zürich, Zurich, Switzerland
- ⁶¹Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶²Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶³Also at Konya Technical University, Konya, Turkey
- ⁶⁴Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁵Also at Adiyaman University, Adiyaman, Turkey
- ⁶⁶Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- ⁶⁷Also at Istanbul Sabahattin Zaim University, Istanbul, Turkey

-
- ⁶⁸Also at Marmara University, Istanbul, Turkey
- ⁶⁹Also at Milli Savunma University, Istanbul, Turkey
- ⁷⁰Also at Informatics and Information Security Research Center, Gebze/Kocaeli, Turkey
- ⁷¹Also at Kafkas University, Kars, Turkey
- ⁷²Now at Istanbul Okan University, Istanbul, Turkey
- ⁷³Also at Hacettepe University, Ankara, Turkey
- ⁷⁴Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁵Also at Istinye University, Istanbul, Turkey
- ⁷⁶Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁷Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁷⁸Also at Monash University, Faculty of Science, Clayton, Australia
- ⁷⁹Also at Bethel University, St. Paul, Minnesota, USA
- ⁸⁰Also at Università di Torino, Torino, Italy
- ⁸¹Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁸²Also at California Lutheran University,, Thousand Oaks, California, USA
- ⁸³Also at California Institute of Technology, Pasadena, California, USA
- ⁸⁴Also at United States Naval Academy, Annapolis, Maryland, USA
- ⁸⁵Also at Bingol University, Bingol, Turkey
- ⁸⁶Also at Georgian Technical University, Tbilisi, Georgia
- ⁸⁷Also at Sinop University, Sinop, Turkey
- ⁸⁸Also at Erciyes University, Kayseri, Turkey
- ⁸⁹Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
- ⁹⁰Now at another institute formerly covered by a cooperation agreement with CERN
- ⁹¹Also at Hamad Bin Khalifa University (HBKU), Doha, Qatar
- ⁹²Also at Yerevan Physics Institute, Yerevan, Armenia
- ⁹³Also at Imperial College, London, United Kingdom