

Probabilistic and Statistical Modeling

Vasilis Gkolemis

ATHENA RC — HUA

June 2025

© Vasilis Gkolemis, 2025. Licensed under CC BY 4.0.

Contents

- 1 Intro & Recap
- 2 Probabilistic vs statistical models
- 3 Bayesian Modeling for Supervised Tasks
- 4 Examples of parametric conditional models

- **Primer on Probabilistic Modeling**

<https://www.inf.ed.ac.uk/teaching/courses/pmr/22-23/assets/notes/probabilistic-modelling-primer.pdf>

Session 1 – Recap

• What we covered:

- ▶ **Probabilistic Modeling:** Model the world using probabilities
- ▶ **Probabilistic Reasoning (Inference):** Use knowns to infer unknowns
- ▶ **Bayesian Analysis:** Modeling and Reasoning with Bayes' rule.
- ▶ **Core Rules of Probability:** The sum, product and Bayes' rule.
- ▶ *Example: Alzheimer's diagnostic test.*

• What's still to explore:

- ▶ Our example was simple
 - ★ X : the test result — a 1D random variable in $\{0, 1\}$
 - ★ Y : the disease status — a 1D random variable in $\{0, 1\}$
- ▶ Real-world problems are more complex
 - ★ Involve high-dimensional random variables
 - ★ Involve complex relationships between variables
- ▶ How can we model these complexities?
 - ★ Session 2 extended our probabilistic toolbox.
 - ★ Session 3 will show how to use it in practice.

Session 2 – Recap

- **What we covered:**

- ▶ **Multivariate Random Variables and Distributions:**

- ★ PDFs, PMFs and CDFs
 - ★ Key properties: expectation and variance.
 - ★ How to sample from these distributions.
 - ★ Key-distributions: Bernoulli, Normal, Poisson.

- ▶ We now have powerful tools to model complexity!

- **What's still to explore:**

- ▶ A glue to connect our probabilistic tools for performing analysis.
 - ▶ A *principled* and *unified* way to:
 - ★ Model complex relationships between variables
 - ★ Infer unknowns from knowns
 - ★ Make predictions about future observations
 - ▶ The **Bayesian framework** is (among others) a powerful glue for this.

Session 3 – Overview

- **What we'll explore:**

- ▶ The Bayesian Framework with each key components:
 - ★ **Prior Distribution:** Our belief before seeing the data.
 - ★ **Likelihood:** How compatible is the observed data is with different parameter values.
 - ★ **Posterior Distribution:** Our updated beliefs after observing the data.
 - ★ **Predictive Distribution:** Make predictions about new, unseen data.
- ▶ How to use the Bayesian framework for predictive tasks.

- **Be confident. You already know important stuff:**

- ▶ Session 1:
 - ★ Intuition about Bayesian modeling → Alzheimer's test case
 - ★ Core probability rules: sum, product, and Bayes' rule
- ▶ Session 2:
 - ★ Multivariate random variables and distributions
 - ★ Key properties: expectation and variance
 - ★ How to sample from these distributions

- The term “model” has multiple meanings, see e.g. <https://en.wikipedia.org/wiki/Model>
- Let's distinguish between three types of models:
 - ▶ probabilistic model
 - ▶ (parametric) statistical model
 - ▶ Bayesian model
- Note: the three types are often confounded, and often just called probabilistic or statistical model, or just “model”.
- [Introduction to Probabilistic Modelling](#) → for further reading.

Probabilistic model

- From first lecture:

A probabilistic model is an abstraction of reality that uses probability theory to quantify the chance of uncertain events.

- Example from the first lecture: cognitive impairment test
 - ▶ Sensitivity of 0.8 and specificity of 0.95 (Scharre, 2010)
 - ▶ Probabilistic model for presence of impairment ($x = 1$) and detection by the test ($y = 1$):
 - ★ $P(x = 1) = 0.11$ (prior)
 - ★ $P(y = 1 \mid x = 1) = 0.8$ (sensitivity)
 - ★ $P(y = 0 \mid x = 0) = 0.95$ (specificity)

Probabilistic model

- More technically:
probabilistic model \equiv *probability distribution (pmf/pdf)*.
- Probabilistic model was written in terms of the probability P .
- In terms of the pmf it is:
 - ▶ $p_x(1) = 0.11$
 - ▶ $p_{y|x}(1 | 1) = 0.8$
 - ▶ $p_{y|x}(0 | 0) = 0.95$
- Commonly written as:
 - ▶ $p(x = 1) = 0.11$
 - ▶ $p(y = 1 | x = 1) = 0.8$
 - ▶ $p(y = 0 | x = 0) = 0.95$
- where the notation for probability measure P and pmf p are confounded.

Statistical model

- If we substitute the numbers with parameters, we obtain a (parametric) statistical model:
 - ▶ $p(x = 1) = \theta_1$
 - ▶ $p(y = 1 \mid x = 1) = \theta_2$
 - ▶ $p(y = 0 \mid x = 0) = \theta_3$
- For each value of the θ_i , we obtain a different pmf.
- Dependency highlighted by writing:
 - ▶ $p(x = 1; \theta_1) = \theta_1$
 - ▶ $p(y = 1 \mid x = 1; \theta_2) = \theta_2$
 - ▶ $p(y = 0 \mid x = 0; \theta_3) = \theta_3$
- $p(x, y; \theta)$ where $\theta = (\theta_1, \theta_2, \theta_3)$ is a vector of parameters.
- or $p(x, y \mid \theta)$, for highlighting that θ is considered a random variable.
- A statistical model corresponds to a set of probabilistic models, here indexed by the parameters θ : $\{p(x; \theta)\}_\theta$

What is Bayesian modeling?

A Bayesian model turns a statistical model into a probabilistic one by treating parameters θ as random variables.

Goal: Learn what we believe about θ after seeing data — and use that to make predictions.

- A Bayesian model is a probabilistic model $p(x, y, \theta)$
- In supervised settings, we consider x as observed, so we care about $p(y, \theta \mid x)$.

Bayesian Modeling in Steps

We don't know the full joint distribution $p(x, y, \theta)$.

- If we did, every analysis would be trivial.

What we do have:

- Observed data $\mathcal{D} = \{(x^{(i)}, y^{(i)})\}_{i=1}^N$ — i.i.d. samples from $p(x, y)$

What we assume:

- $p(y \mid x, \theta)$ — how data is generated given a specific parameter θ
- $p(\theta)$ — our beliefs about the parameters before seeing data

What we want to learn:

- $p(\theta \mid \mathcal{D})$ — what we believe about the parameters after seeing data
- The predictive distribution $p(y \mid x, \mathcal{D})$ — predictions that account for parameter uncertainty
- Possibly others: e.g. marginal likelihood $p(\mathcal{D})$

Bayesian Modeling for Supervised Tasks

- **Supervised learning:** We observe a dataset of input–output pairs $\mathcal{D} = \{(x^{(i)}, y^{(i)})\}_{i=1}^N$ drawn from an unknown joint distribution $p(x, y)$.
- **Bayesian perspective:** Use this data to learn the relationship $x \mapsto y$ while capturing uncertainty in the model parameters.
- **Hypothesis:**
 - ▶ (1): assume a parametric family for $p(y \mid x, \theta)$, such as linear regression, neural networks, etc. (Parametric modeling assumption)
 - ▶ (2): assume a prior belief over the parameters $p(\theta)$ (prior assumption)

Example: Linear Regression

- **Hypothesis:** The outcome depends linearly on some input plus noise.
- **Example:** Predict house price from size.
 - ▶ Y — house price
 - ▶ X — house size
 - ▶ Model: $Y = wX + b + \epsilon$, with $\epsilon \sim \mathcal{N}(0, \sigma^2)$
- $p(Y | X; \theta) = \mathcal{N}(Y | wX + b, \sigma^2)$
- **Parameters:** $\theta = (w, b)$ — 2 variables.

Example: Linear Regression

- **Hypothesis:** The outcome depends linearly on some input plus noise.
- Generalizes to any number of input features
- **Example:** Predict house price from size, number of rooms, and
 - ▶ Y — house price
 - ▶ $X = (X_1, X_2, \dots, X_d)$ — house features (size, number of rooms, etc.)
 - ▶ Model: $Y = wX + b + \epsilon$, with $\epsilon \sim \mathcal{N}(0, \sigma^2)$
- $p(Y | X; \theta) = \mathcal{N}(Y | wX + b, \sigma^2)$
- **Parameters:** $\theta = (w, b)$: $d + 1$ variables.

Example: Non-linear Regression

- **Hypothesis:** The relationship between input and output is complex and nonlinear plus noise.
- **Example:** Predict bike rentals from weather data.
 - ▶ Y — number of bikes rented per hour
 - ▶ X — weather features (temperature, humidity, etc.)
 - ▶ Model $Y = f_{\theta}(X) + \epsilon$, with $\epsilon \sim \mathcal{N}(0, \sigma^2)$
 - ▶ $f_{\theta}(X)$ is a non-linear function parameterized by θ .
- $p(Y | X; \theta) = \mathcal{N}(Y | f_{\theta}(X), \sigma^2)$
- what is $f_{\theta}(X)$?
 - ▶ A neural network with weights θ , normally of thousands of variables.
 - ▶ A random forest with decision trees where the structure and parameters are defined by θ , normally of hundreds of variables.

The Flexibility—and Cost—of the Bayesian Framework

- **Bayesian framework is flexible:** We can assume any model for $p(y \mid x, \theta)$.
 - ▶ Example: $y = f_{\theta}(x) + \epsilon$, where $\epsilon \sim \mathcal{N}(0, \sigma^2)$ and $f_{\theta}(x)$ can be a neural network, random forest, etc.
 - ▶ The output y can follow any distribution — not just Normal: skewed, heavy-tailed, discrete (e.g., Poisson, Bernoulli), etc.
- **But flexibility comes at a cost**
 - ▶ the more complex $p(y \mid x, \theta)$:
 - ★ Complex \rightarrow a high-dimensional θ .
 - ★ Complex \rightarrow a complex $f_{\theta}(x)$
 - ▶ the harder it is to perform inference.