

Regionally Additive Models: Explainable-by-design models minimizing feature interactions

Vasilis Gkolemis^{1,2} Anargyros Tzerefos¹ Theodore Dalamagas¹
Eirini Ntoutsis³ Christos Diou²

¹ATHENA Research and Innovation Center

²Harokopio University of Athens

³Universität der Bundeswehr München

September 2023, Turin, Italy

Generalized Additive Models

Wikipedia says:

In statistics, a generalized additive model (GAM) is a generalized linear model in which the linear response variable depends linearly on unknown smooth functions of some predictor variables, and interest focuses on inference about these smooth functions.

Generalized Additive Models (GAMs)

Wikipedia says:

In statistics, a generalized additive model (GAM) is a generalized linear model in which the linear response variable depends linearly on unknown smooth functions of some predictor variables, and interest focuses on inference about these smooth functions.

Generalized Additive Models (GAMs)

Wikipedia says:

*In statistics, a generalized additive model (GAM) is a generalized linear model in which the linear **response** variable depends linearly on unknown smooth functions of some predictor variables, and interest focuses on inference about these smooth functions.*

y

Generalized Additive Models (GAMs)

Wikipedia says:

*In statistics, a generalized additive model (GAM) is a generalized linear model in which the linear **response** variable depends **linearly** on unknown smooth functions of some predictor variables, and interest focuses on inference about these smooth functions.*

$$y = \cdot + \dots + \cdot$$

Generalized Additive Models (GAMs)

Wikipedia says:

*In statistics, a generalized additive model (GAM) is a generalized linear model in which the linear **response** variable depends **linearly** on unknown **smooth functions of some predictor variables**, and interest focuses on inference about these smooth functions.*

$$y = f_1(x_1) + \dots + f_D(x_D)$$

GAMs are called explainable-by-design, why?

Output/target variable:


- $y_{\text{bike-rentals}}$ the expected number of bike rentals

Input/covariates:

- $X_{\text{temperature}}$
- X_{humidity}
- $X_{\text{is_weekday}}$


GAMs - Interpretability

GAMs are explainable-by-design, why?




GAMs - Interpretability

GAMs are explainable-by-design, why?



GAMs - Interpretability


GAMs are explainable-by-design, why?



Scanned with CamScanner

GAMs - Interpretability

GAMs are explainable-by-design, why?



What if:

- temperature has different effect on week-days vs weekends

GAMs - Limitations

What if:

- temperature has different effect on week-days vs weekends
- Cause: go to work vs go sightseeing

GAMs - Limitations

What if:

- temperature has different effect on week-days vs weekends
- Cause: go to work vs go sightseeing
- Solution 1: $f(x_{\text{temperature}}, x_{\text{is_weekday}})$

What if:

- temperature has different effect on week-days vs weekends
- Cause: go to work vs go sightseeing
- Solution 1: $f(x_{\text{temperature}}, x_{\text{is_weekday}})$
- Solution 2: $f(x_{\text{temperature}} | \text{weekday})$ vs $f(x_{\text{temperature}} | \text{weekend})$

What if:

- temperature has different effect on week-days vs weekends
- Cause: go to work vs go sightseeing
- Solution 1: $f(x_{\text{temperature}}, x_{\text{is_weekday}})$
- Solution 2: $f(x_{\text{temperature}} | \text{weekday})$ vs $f(x_{\text{temperature}} | \text{weekend})$

- Solution 1: GA^2M

What if:

- temperature has different effect on week-days vs weekends
- Cause: go to work vs go sightseeing
- Solution 1: $f(x_{\text{temperature}}, x_{\text{is_weekday}})$
- Solution 2: $f(x_{\text{temperature}} | \text{weekday})$ vs $f(x_{\text{temperature}} | \text{weekend})$

- Solution 1: GA^2M
- Solution 2: RAM

What if:

- Have you ever ridden a bike with cold and humidity?

Solutions:

GAMs - Limitations

What if:

- Have you ever ridden a bike with cold and humidity?
- If it is weekend, cool! Netflix is here!

Solutions:

GAMs - Limitations

What if:

- Have you ever ridden a bike with cold and humidity?
- If it is weekend, cool! Netflix is here!
- But what to do if the only option to go to work?

Solutions:

GAMs - Limitations

What if:

- Have you ever ridden a bike with cold and humidity?
- If it is weekend, cool! Netflix is here!
- But what to do if the only option to go to work?

Solutions:

- $f(x_{\text{temperature}}, x_{\text{humidity}} | x_{\text{is_weekday}})$ *RA²M*

GAMs - Limitations

What if:

- Have you ever ridden a bike with cold and humidity?
- If it is weekend, cool! Netflix is here!
- But what to do if the only option to go to work?


Solutions:

- $f(x_{\text{temperature}}, x_{\text{humidity}} | x_{\text{is_weekday}})$ *RA²M*
- $f(x_{\text{temperature}} | x_{\text{humidity}} = \{high, low\}, x_{\text{is_weekday}})$ *RAM with two conditions*


RAM on toy example

$$f(\mathbf{x}) = 8x_2 \mathbb{1}_{x_1 > 0} \mathbb{1}_{x_3 = 0}$$


$$x_1, x_2 \sim \mathcal{U}(-1, 1), x_3 \sim \text{Bernoulli}(0, 1)$$



(a) $f_2(x_2)$



(b) $f_2(x_2) \mathbb{1}_{x_1 > 0 \text{ and } x_3 = 1}$



(c) $f_2(x_2) \mathbb{1}_{x_1 \leq 0 \text{ or } x_3 \neq 1}$

Figure: (Left) GAM, (Middle and Right) RAM

How RAM works (1)

3-step approach:

- Fit a Black-box model to capture all complex structures
 - ▶ it should be differentiable
 - ▶ neural network
- Use a Regional Effect method to find important interactions
 - ▶ [RHALE - Gkolemis et. al](#)
 - ▶ [Feature Interactions - Herbinger et. al](#)
- Fit a univariate function on each detected subregion

Step 1

- Fit a Black-box model to capture all complex structures
 - ▶ it should be differentiable
 - ▶ A neural network is a good option

Step 2

- Use a Regional Effect method to find important interactions
 - ▶ [RHALE - Gkolemis et. al](#)
 - ▶ [Feature Interactions - Herbinger et. al](#)


Step 3

- Fit a univariate function on each detected subregion




Bike Sharing dataset


Predict bike-rentals (on daily basis) given some features of the day



(a) $f(X_{\text{hour}})$



(b) $f(X_{\text{hour}})\mathbb{1}_{X_{\text{workingday}} \neq 1}$



(c) $f(X_{\text{hour}})\mathbb{1}_{X_{\text{workingday}} = 1}$

Experimental Results

	Black-box	x-by-design			
	all orders	1 st order		2 nd order	
	DNN	GAM	RAM	GA ² M	RA ² M
Bike (MAE)	0.254	0.549	0.430	0.298	0.278
Bike (RMSE)	0.389	0.734	0.563	0.438	0.412
Housing (MAE)	0.373	0.600	0.553	0.554	0.533
Housing (RMSE)	0.533	0.819	0.754	0.774	0.739

What next?

A lot has to be done!

- Results are only preliminary
 - ▶ Experiment with more datasets
 - ▶ Sometimes wide bins are needed
- Can we learn uncertain RAMs?
 - ▶ How do we model uncertainty?

Thank you for the attention

- For more discussion or future ideas on RAM, contact me:
 - ▶ vgkolemis@athenarc.gr
 - ▶ gkolemis@hua.gr
- Questions?