

MATH 32B Problem Set 1

Allan Zhang

January 6, 2025

1 Question 11

The following table gives the approximate height at quarter-meter intervals of a mound of gravel. Estimate the volume of the mound by computing the average of the two Riemann sums $S_{4,3}$ with lower-left and upper-right vertices of the subrectangles as sample points.

| | | | | | |
|------------------|------------------|-------------|------------|-------------|----------|
| 0.75 | 0.1 | 0.2 | 0.2 | 0.15 | 0.1 |
| 0.5 | 0.2 | 0.3 | 0.5 | 0.4 | 0.2 |
| 0.25 | 0.15 | 0.2 | 0.4 | 0.3 | 0.2 |
| $y : \mathbf{0}$ | 0.1 | 0.15 | 0.2 | 0.15 | 0.1 |
| | $x : \mathbf{0}$ | 0.25 | 0.5 | 0.75 | 1 |

Computing area using lower-left vertices as sample points: The vertices we will be using are:

$$\begin{bmatrix} 0.2 & 0.3 & 0.5 & 0.4 \\ 0.15 & 0.2 & 0.4 & 0.3 \\ 0.1 & 0.15 & 0.2 & 0.15 \end{bmatrix}$$

Note that Δx and Δy are both 0.25, meaning the volume of each rectangle will be the height of the rectangles times 0.25^2 , which is 0.0625

$$0.0625 \cdot \begin{bmatrix} 0.2 & 0.3 & 0.5 & 0.4 \\ 0.15 & 0.2 & 0.4 & 0.3 \\ 0.1 & 0.15 & 0.2 & 0.15 \end{bmatrix} = \begin{bmatrix} 0.0125 & 0.01875 & 0.03125 & 0.025 \\ 0.009375 & 0.0125 & 0.025 & 0.01875 \\ 0.00625 & 0.009375 & 0.0125 & 0.009375 \end{bmatrix}$$

Adding up these areas gives us 0.190625

Now, let's repeat, except we will use the upper-right vertices as sample points

$$0.0625 \cdot \begin{bmatrix} 0.2 & 0.2 & 0.15 & 0.1 \\ 0.3 & 0.5 & 0.4 & 0.2 \\ 0.2 & 0.4 & 0.3 & 0.2 \end{bmatrix} = \begin{bmatrix} 0.0125 & 0.0125 & 0.009375 & 0.00625 \\ 0.01875 & 0.03125 & 0.025 & 0.0125 \\ 0.0125 & 0.025 & 0.01875 & 0.03125 \end{bmatrix}$$

Adding up these areas gives us 0.196875. The average of these two sums is **0.19375**, which is our estimate for the volume of the mound.

2 Question 15

Use symmetry to evaluate the double integral

$$\iint_{\mathcal{R}} x^3 dA, \quad \mathcal{R} = [-4, 4] \times [0, 5]$$

Note that x^3 is odd and lacks horizontal or vertical shifts. This means so for some value a , the area between the curve and $[-a, 0]$ will be equal to the area between the curve and $[0, a]$ multiplied by -1 .

Also, note that we can split the domain into two sub-domains. Let us split \mathcal{R} into $\mathcal{R}_1 = [-4, 0] \times [0, 5]$ and $\mathcal{R}_2 = [0, 4] \times [0, 5]$.

$$\iint_{\mathcal{R}} x^3 dA, \quad \mathcal{R} = [-4, 4] \times [0, 5] = \iint_{\mathcal{R}_1} x^3 dA, \quad \mathcal{R}_1 = [-4, 0] \times [0, 5] + \iint_{\mathcal{R}_2} x^3 dA, \quad \mathcal{R}_2 = [0, 4] \times [0, 5]$$

From the observation we stated before, it can be said that

$$\iint_{\mathcal{R}_1} x^3 dA, \quad \mathcal{R}_1 = [-4, 0] \times [0, 5] = - \iint_{\mathcal{R}_2} x^3 dA, \quad \mathcal{R}_2 = [0, 4] \times [0, 5]$$

We can rewrite the original expression as

$$- \iint_{\mathcal{R}_2} x^3 dA + \iint_{\mathcal{R}_2} x^3 dA = 0$$

This double integral evaluates to 0.

3 Question 17

Use symmetry to evaluate the double integral

$$\iint_{\mathcal{R}} \sin x \, dA, \quad \mathcal{R} = [0, 2\pi] \times [0, 2\pi]$$

Note that $\sin x$ oscillates from -1 to 1. Importantly, without any transformations, $\sin x$ completes one whole revolution from 0 to 2π , and the function from 0 to π is identical to the function from π to 2π reflected over the x -axis.

Also remember that we can split the domain into two sub-domains. Let us split \mathcal{R} into $\mathcal{R}_1 = [0, \pi] \times [0, 2\pi]$ and $\mathcal{R}_2 = [\pi, 2\pi] \times [0, 2\pi]$.

$$\iint_{\mathcal{R}} \sin x \, dA, \quad \mathcal{R} = [0, 2\pi] \times [0, 2\pi] = \iint_{\mathcal{R}_1} \sin x \, dA, \quad \mathcal{R}_1 = [0, \pi] \times [0, 2\pi] + \iint_{\mathcal{R}_2} \sin x \, dA, \quad \mathcal{R}_2 = [\pi, 2\pi] \times [0, 2\pi]$$

From what we said before, we can observe that

$$\iint_{\mathcal{R}_1} \sin x \, dA, \quad \mathcal{R}_1 = [0, \pi] \times [0, 2\pi] = - \iint_{\mathcal{R}_2} \sin x \, dA, \quad \mathcal{R}_2 = [\pi, 2\pi] \times [0, 2\pi]$$

Finally, we can state that

$$\iint_{\mathcal{R}} \sin x \, dA, \quad \mathcal{R} = [0, 2\pi] \times [0, 2\pi] = - \iint_{\mathcal{R}_2} \sin x \, dA + \iint_{\mathcal{R}_2} \sin x \, dA = 0$$

The double integral evaluates to 0.

4 Question 19

Evaluate the iterated integral

$$\begin{aligned} & \int_1^3 \int_0^2 x^3 y \, dy \, dx \\ & \int_1^3 \int_0^2 x^3 y \, dy \, dx = \int_1^3 \left(\int_0^2 x^3 y \, dy \right) dx \\ & = \int_1^3 \left(x^3 \cdot \frac{y^2}{2} \Big|_{y=0}^{y=2} \right) dx \\ & = \int_1^3 \left(x^3 \cdot \left(\frac{4}{2} - \frac{0}{2} \right) \right) dx \\ & = \int_1^3 2x^3 \, dx = \frac{x^4}{2} \Big|_{x=1}^{x=3} = \frac{81}{2} - \frac{1}{2} = \frac{80}{2} = 40 \end{aligned}$$

The iterated integral evaluates to 40.

5 Question 21

Evaluate the iterated integral

$$\begin{aligned}\int_4^9 \int_{-3}^8 1 \, dx \, dy \\ \int_4^9 \left(\int_{-3}^8 1 \, dx \right) dy = \\ \int_4^9 x \Big|_{x=-3}^{x=8} dy = \\ \int_4^9 11 \, dy = 11x \Big|_{x=4}^{x=9} = 99 - 44 = 55\end{aligned}$$

The iterated integral evaluates to 55.

6 Question 23

Evaluate the iterated integral

$$\begin{aligned} & \int_{-1}^1 \int_0^{\pi} x^2 \sin y \, dy \, dx \\ & \int_{-1}^1 \left(\int_0^{\pi} x^2 \sin y \, dy \right) dx \\ & \int_{-1}^1 \left(x^2 \cdot -\cos y \Big|_{y=0}^{y=\pi} \right) dx \\ & \int_{-1}^1 x^2 (1 - (-1)) \, dx \\ & \int_{-1}^1 2x^2 \, dx = \frac{2x^3}{3} \Big|_{x=-1}^{x=1} = \frac{2}{3} - \frac{-2}{3} = \frac{4}{3} \end{aligned}$$

The iterated integral evaluates to $\frac{4}{3}$.

7 Question 25

Evaluate the iterated integral

$$\begin{aligned} & \int_2^6 \int_1^4 x^2 \, dx \, dy \\ & \int_2^6 \left(\int_1^4 x^2 \, dx \right) dy \\ & \int_2^6 \left(\frac{x^3}{3} \Big|_{x=1}^{x=4} \right) dy \\ & \int_2^6 \frac{64}{3} - \frac{1}{3} \, dy = \int_2^6 \frac{63}{3} \, dy \\ & \frac{63x}{3} \Big|_{x=2}^{x=6} = \frac{378 - 126}{3} = \frac{252}{3} = 84 \end{aligned}$$

The iterated integral evaluates to 84.

8 Question 37

Evaluate the integral

$$\iint_{\mathcal{R}} \frac{x}{y} dA, \quad \mathcal{R} = [-2, 4] \times [1, 3]$$

$$\iint_{\mathcal{R}} \frac{x}{y} dx dy, \quad \mathcal{R} = [-2, 4] \times [1, 3]$$

$$\int_1^3 \int_{-2}^4 \frac{x}{y} dx dy$$

$$\int_1^3 \left(\frac{x^2}{2y} \Big|_{x=-2}^{x=4} dx \right) dy$$

$$\int_1^3 \frac{1}{2y} (16 - 4) dy$$

$$\int_1^3 \frac{6}{y} dy$$

$$6 \ln y \Big|_{y=1}^{y=3} = 6 \ln 3 - 6 \ln 1 = 6 \ln 3$$

9 Question 40

Evaluate the integral

$$\iint_{\mathcal{R}} \frac{y}{x+1} dA, \quad \mathcal{R} = [0, 2] \times [0, 4]$$

$$\int_0^4 \int_0^2 \frac{y}{x+1} dx dy$$

$$\int_0^4 \int_0^2 \frac{y}{x+1} dx dy$$

$$\int_0^4 \left(y \ln |x+1| \Big|_{x=0}^{x=2} \right) dy$$

$$\int_0^4 y(\ln |3| - \ln |1|) dy$$

$$\int_0^4 y \ln |3| dy$$

$$\frac{\ln |3| y^2}{2} \Big|_{y=0}^{y=4}$$

$$8 \ln 3$$

The solution to this evaluated integral is $8 \ln 3$

10 Question 45

Let $f(x, y) = mxy^2$, where m is a constant. Find a value of m such that $\iint_{\mathcal{R}} f(x, y) \, dA = 1$, where $\mathcal{R} = [0, 1] \times [0, 2]$

11 Question 48

- a. Which is easier, antidifferentiating xe^{xy} with respect to x or with respect to y ? Explain.
- b. Evaluate $\iint_{\mathcal{R}} xe^{xy} dA = 1$, where $\mathcal{R} = [0, 1] \times [0, 1]$

12 Question 49

- a. Which is easier, antidifferentiating $\frac{y}{1+xy}$ with respect to x or with respect to y ? Explain.
- b. Evaluate $\iint_{\mathcal{R}} \frac{y}{1+xy} dA = 1$, where $\mathcal{R} = [0, 1] \times [0, 1]$