

Problem 1

States: $\{s, r, f\}$ where q is our initial state and f is our final state

Our Turing Machine is defined by the following instantaneous descriptions:

$$\delta(q, 0) = (q, 0, R) \qquad \delta(q, 1) = (q, 1, R) \qquad \delta(q, B) = (r, B, L)$$

$$\delta(r, 0) = (f, 1, R) \qquad \delta(r, 1) = (r, 0, L)$$

Problem 2

The method that we will use to perform binary addition is as follows: First of all, we will scan left-to-right copying the symbol from the left-hand operand onto the second tape. Upon reaching a blank, we will switch states. Then, we will move the second tape to its start. Afterwards, we proceed through the second operand, we will print onto the second tape the result of adding the bits at the head of each tape.

Here is an explanation of the states used in this Turing Machine:

- q_0 : This state will be the state that we use to copy the left-hand operand onto the second tape. This is also our start state.
- q_1 : This state will be the state that we use to move the first tape to the last digit on the tape
- q_2 : This is the state that will be used while scanning left-to-right on both the first and second tapes to perform the addition
- q_3 : This is the state that will be used when addition needs to have a carry
- f : This is our final state and will indicate that the addition has been completed.

States: $\{q_0, q_1, q_2, f\}$

We have two tapes. The first tape consists of the two operands, separated by a space. The second tape consists of all blanks, but will be the location where we store the result of the addition.

These are our transitions:

$$\delta(q_0, 0, B) = (q_0, 0, 0, R, R) \qquad \delta(q_0, 1, B) = (q_0, 1, 1, R, R)$$

$$\delta(q_0, B, B) = (q_1, B, B, S, L) \qquad \delta(q_1, 0, B) = (q_1, 0, B, R, S)$$

$$\delta(q_1, 1, B) = (q_1, 1, B, R, S)$$

$$\delta(q_1, B, B) = (q_2, B, B, R, S)$$

$$\delta(q_2, 1, 1) = (q_2, 1, 0, L, L)$$

$$\delta(q_2, 0, 1) = (q_2, 0, 1, L, L)$$

$$\delta(q_2, 0, 0) = (q_2, 0, 0, L, L)$$

$$\delta(q_2, 1, 1) = (q_3, 1, 1, L, L)$$

$$\delta(q_3, 0, 1) = (q_2, 0, 0, L, L)$$

$$\delta(q_3, 1, 0) = (q_2, 1, 0, L, L)$$

$$\delta(q_3, 0, 0) = (q_2, 0, 1, L, L)$$

$$\delta(q_3, 1, 1) = (q_3, 1, 0, L, L)$$

$$\delta(q_2, B, B) = (f, B, B, S, S)$$

Problem 3

Problem 4