

Recollements and Injective Generation of the Derived Category

Charley Cummings

University of Bristol

ICRA 2020/fd-seminar

Injective Generation

Injective Generation

Definition

Let A be a ring. If $\mathcal{D}(A)$ is generated, as a triangulated category with infinite coproducts, by the injective A -modules then we say that **injectives generate for A** .

Injective Generation

Definition

Let A be a ring. If $\mathcal{D}(A)$ is generated, as a triangulated category with infinite coproducts, by the injective A -modules then we say that **injectives generate for A** .

Injectives generate



Keller

Nunke
condition




Generalised
Nakayama
conjecture

Injective Generation

Definition


Let A be a ring. If $\mathcal{D}(A)$ is generated, as a triangulated category with infinite coproducts, by the injective A -modules then we say that **injectives generate for A** .



Injective Generation

Definition

Let A be a ring. If $\mathcal{D}(A)$ is generated, as a triangulated category with infinite coproducts, by the injective A -modules then we say that **injectives generate for A** .



Injective Generation

Definition

Let A be a ring. If $\mathcal{D}(A)$ is generated, as a triangulated category with infinite coproducts, by the injective A -modules then we say that **injectives generate for A** .

Examples

- Rings with finite global dimension.
- Monomial algebras (Rickard 2019) (FDC: Green, Kirkman, Kuzmanovich 1991).
- Commutative noetherian rings (Rickard 2019, Neeman 1992).

Recollements

Recollements

Definition

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j_*} \\ \xleftarrow{j^*} \end{array} \mathcal{D}(C)$$

Let A , B and C be rings. Let $(i^*, i_*, i^!)$ and $(j_!, j^*, j_*)$ be triples of adjoint functors satisfying the properties defined by Beilinson, Bernstein and Deligne (1982).

Recollements

Definition

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j_*} \\ \xleftarrow{j^*} \end{array} \mathcal{D}(C)$$

Let A , B and C be rings. Let $(i^*, i_*, i^!)$ and $(j_!, j^*, j_*)$ be triples of adjoint functors satisfying the properties defined by Beilinson, Bernstein and Deligne (1982).

Idea - $\mathcal{D}(A)$ is 'built from' $\mathcal{D}(B)$ and $\mathcal{D}(C)$.

Recollements

Example (Triangular Matrix Algebra)

Let A be a quiver algebra with quiver Q_A .



Recollements

Example (Triangular Matrix Algebra)

Let A be a quiver algebra with quiver Q_A .



Let B be the quiver algebra associated to Q_B and C be the quiver algebra associated to Q_C .

Recollements

Example (Triangular Matrix Algebra)

Let A be a quiver algebra with quiver Q_A .



Let B be the quiver algebra associated to Q_B and C be the quiver algebra associated to Q_C .

Then there exists a recollement (R) .

$$(R) := \begin{array}{ccccc} & i^* & & j! & \\ & \curvearrowright & & \curvearrowleft & \\ \mathcal{D}(B) & \xrightarrow{i_*} & \mathcal{D}(A) & \xrightarrow{j^*} & \mathcal{D}(C) \\ & \curvearrowleft i^! & & \curvearrowright j_* & \end{array}$$

Recollements

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 1 (C 2020)

Recollements

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 1 (C 2020)

Suppose that i_* preserves **compact objects**.

Recollements

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 1 (C 2020)

Suppose that i_* preserves **compact objects**.

- 1 If injectives generate for both B and C then injectives generate for A .

Recollements

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 1 (C 2020)

Suppose that i_* preserves **compact objects**.

- 1 If injectives generate for both B and C then injectives generate for A .
- 2 If injectives generate for A then injectives generate for B .

Recollements

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 1 (C 2020)

Suppose that i_* preserves **compact objects**.

- 1 If injectives generate for both B and C then injectives generate for A .
- 2 If injectives generate for A then injectives generate for B .

Note that (R) restricts to a recollement of bounded above derived categories (Angeleri H\"ugel, Koenig, Liu, Yang 2017).

Recollements

Example (Triangular Matrix Algebra)

Let A be the quiver algebra with quiver Q_A .



$$(R) := \begin{array}{ccccc} & i^* & & j^! & \\ & \curvearrowright & & \curvearrowright & \\ \mathcal{D}(B) & \xrightarrow{i_*} & \mathcal{D}(A) & \xrightarrow{j^*} & \mathcal{D}(C) \\ & \curvearrowleft i^! & & \curvearrowleft j_* & \end{array}$$

Recollements

Example (Triangular Matrix Algebra)

Let A be the quiver algebra with quiver Q_A .




$$(R) := \begin{array}{ccccc} & i^* & & j^! & \\ & \curvearrowright & & \curvearrowright & \\ \mathcal{D}(B) & \xrightarrow{i_*} & \mathcal{D}(A) & \xrightarrow{j^*} & \mathcal{D}(C) \\ & \curvearrowleft i^! & & \curvearrowleft j_* & \end{array}$$

- 1 If injectives generate for both B and C then injectives generate for A .
- 2 If injectives generate for A then injectives generate for B .


Vertex Removal

(Fuller, Saorín 1992), (Green, Psaroudakis, Solberg 2018).




Vertex Removal


(Fuller, Saorín 1992), (Green, Psaroudakis, Solberg 2018).




Projective A -modules



Let $e = e_2 + e_3$.




Projective eAe -modules




Vertex Removal


(Fuller, Saorín 1992), (Green, Psaroudakis, Solberg 2018).




Projective A -modules



Let $e = e_2 + e_3$.




Projective eAe -modules




Vertex Removal

(Fuller, Saorín 1992), (Green, Psaroudakis, Solberg 2018).




Projective A -modules




$$S(1) = (1 - e)A / \text{rad}((1 - e)A)$$

Let $e = e_2 + e_3$.



Projective eAe -modules



Vertex Removal

Example

Let A be a finite dimensional algebra and $e \in A$ be an idempotent. Let $S := (1 - e)A/\text{rad}((1 - e)A)$.

Vertex Removal

Example

Let A be a finite dimensional algebra and $e \in A$ be an idempotent. Let $S := (1 - e)A/\text{rad}((1 - e)A)$. Suppose that $\text{inj.dim}_A(S) = 1$.

Vertex Removal

Example

Let A be a finite dimensional algebra and $e \in A$ be an idempotent. Let $S := (1 - e)A/\text{rad}((1 - e)A)$. Suppose that $\text{inj.dim}_A(S) = 1$. Then there exists a recollement (R) .
(Green, Psaroudakis, Solberg 2018)

$$(R) := \begin{array}{ccccc} & i^* & & j_! & \\ & \curvearrowright & & \curvearrowleft & \\ \mathcal{D}(A/AeA) & \xrightarrow{i_*} & \mathcal{D}(A) & \xrightarrow{j^*} & \mathcal{D}(eAe) \\ & \curvearrowleft & & \curvearrowright & \\ & i_! & & j_* & \end{array}$$

Vertex Removal

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 2 (C 2020)

Vertex Removal

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 2 (C 2020)

Suppose that i_* preserves complexes that are **quasi-isomorphic to a bounded complex of injectives**.

Vertex Removal

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 2 (C 2020)

Suppose that i_* preserves complexes that are **quasi-isomorphic to a bounded complex of injectives**.

- 1 If injectives generate for both B and C then injectives generate for A .

Vertex Removal

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 2 (C 2020)

Suppose that i_* preserves complexes that are **quasi-isomorphic to a bounded complex of injectives**.

- 1 If injectives generate for both B and C then injectives generate for A .
- 2 If injectives generate for A then injectives generate for C .

Vertex Removal

Let (R) be a recollement of derived categories.

$$(R) := \mathcal{D}(B) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(C)$$

Theorem 2 (C 2020)

Suppose that i_* preserves complexes that are **quasi-isomorphic to a bounded complex of injectives**.

- 1 If injectives generate for both B and C then injectives generate for A .
- 2 If injectives generate for A then injectives generate for C .

Moreover, (R) restricts to a recollement of bounded below derived categories.

Vertex Removal

Example

Let A be a finite dimensional algebra and $e \in A$ be an idempotent. Let $S := (1 - e)A/\text{rad}((1 - e)A)$. Suppose that $\text{inj.dim}_A(S) = 1$.

$$(R) := \mathcal{D}(A/AeA) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}(A) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}(eAe)$$

Vertex Removal

Example

Let A be a finite dimensional algebra and $e \in A$ be an idempotent. Let $S := (1 - e)A/\text{rad}((1 - e)A)$. Suppose that $\text{inj.dim}_A(S) = 1$.

$$(R) := \begin{array}{ccccc} & i^* & & j^! & \\ & \curvearrowright & & \curvearrowright & \\ \mathcal{D}(A/AeA) & \xrightarrow{i_*} & \mathcal{D}(A) & \xrightarrow{j^*} & \mathcal{D}(eAe) \\ & \curvearrowleft & & \curvearrowleft & \\ & i^! & & j_* & \end{array}$$

- 1 If injectives generate for both A/AeA and eAe then injectives generate for A .
- 2 If injectives generate for A then injectives generate for eAe .

Vertex Removal

Example

Let A be a finite dimensional algebra and $e \in A$ be an idempotent. Let $S := (1 - e)A/\text{rad}((1 - e)A)$. Suppose that $\text{inj.dim}_A(S) = 1$.

$$(R) := \begin{array}{ccccc} & i^* & & j^! & \\ & \curvearrowright & & \curvearrowright & \\ \mathcal{D}(A/AeA) & \xrightarrow{i_*} & \mathcal{D}(A) & \xrightarrow{j^*} & \mathcal{D}(eAe) \\ & \curvearrowleft & & \curvearrowleft & \\ & i^! & & j_* & \end{array}$$

Note that A/AeA has finite global dimension.

Injectives generate for A if and only if injectives generate for eAe .

Recollements of dg algebras

Let A be a finite dimensional algebra over a field and $e \in A$ be an idempotent.

Recollements of dg algebras

Let A be a finite dimensional algebra over a field and $e \in A$ be an idempotent.

Then there exists a dg algebra B such that (R) is a recollement.

$$(R) := \begin{array}{ccccc} & i^* & & j_! & \\ & \curvearrowright & & \curvearrowright & \\ \mathcal{D}(B) & \xrightarrow{i_*} & \mathcal{D}(A) & \xrightarrow{j^*} & \mathcal{D}(eAe) \\ & \curvearrowleft i^! & & \curvearrowleft j_* & \end{array}$$

Recollements of dg algebras

Let A a finite dimensional algebra over a field and $e \in A$ be an idempotent.

Theorem 3 (C 2020)

Recollements of dg algebras

Let A a finite dimensional algebra over a field and $e \in A$ be an idempotent.

Theorem 3 (C 2020)

Suppose that $Ae \otimes_{eAe}^L eA$ is bounded in cohomology i.e.
 $\mathrm{Tor}_i^{eAe}(Ae, eA) = 0$ for $i \gg 0$.

Recollements of dg algebras

Let A a finite dimensional algebra over a field and $e \in A$ be an idempotent.

Theorem 3 (C 2020)

Suppose that $Ae \otimes_{eAe}^L eA$ is bounded in cohomology i.e.
 $\mathrm{Tor}_i^{eAe}(Ae, eA) = 0$ for $i \gg 0$. Let $S := (1 - e)A / \mathrm{rad}((1 - e)A)$.
Suppose that $\mathrm{inj.dim}_A(S) < \infty$.


Recollements of dg algebras

Let A a finite dimensional algebra over a field and $e \in A$ be an idempotent.



Theorem 3 (C 2020)

Suppose that $Ae \otimes_{eAe}^L eA$ is bounded in cohomology i.e.
 $\mathrm{Tor}_i^{eAe}(Ae, eA) = 0$ for $i \gg 0$. Let $S := (1 - e)A / \mathrm{rad}((1 - e)A)$.
Suppose that $\mathrm{inj.dim}_A(S) < \infty$. If injectives generate for eAe then
injectives generate for A .


Example



Injective A -modules





Example




	$S(2)$	$S(1)$
inj.dim_A	1	2

Injective A -modules





Example




	$S(2)$	$S(1)$
inj.dim_A	1	2

- $e := e_3 + e_4 + e_5,$

Injective A -modules





Example




	$S(2)$	$S(1)$
inj.dim_A	1	2

- $e := e_3 + e_4 + e_5$,
- $Ae \otimes_{eAe}^L eA$ is bounded in cohomology.

Injective A -modules



Example





	$S(2)$	$S(1)$
inj.dim_A	1	2


- $e := e_3 + e_4 + e_5$,
- $Ae \otimes_{eAe}^L eA$ is bounded in cohomology.

Injectives generate for A if injectives generate for eAe .


Injective A -modules




Example




Injective eAe -modules





Example



Injective eAe -modules




Apply triangular matrix algebra result.




Example

So injectives generate for A .



Injective A -modules



Projective Cogeneration

Projective Cogeneration

Definition

Let A be a ring. If $\mathcal{D}(A)$ is generated, as a triangulated category with infinite **products**, by the **projective** A -modules then we say that **projectives cogenerate for A** .

Projective Cogeneration

Definition

Let A be a ring. If $\mathcal{D}(A)$ is generated, as a triangulated category with infinite **products**, by the **projective** A -modules then we say that **projectives cogenerate for A** .

Let A be a finite dimensional algebra over a field.

Projectives
cogenerate
for A^{op}

\implies
Rickard

Finitistic
dimension
conjecture
holds for A

Projective Cogeneration

Definition

Let A be a ring. If $\mathcal{D}(A)$ is generated, as a triangulated category with infinite **products**, by the **projective** A -modules then we say that **projectives cogenerate for A** .

Let A be a finite dimensional algebra over a field.

Injectives
generate for A $\xRightarrow{\text{Rickard}}$ Projectives
cogenerate
for A^{op} $\xRightarrow{\text{Rickard}}$ Finitistic
dimension
conjecture
holds for A

Thank you

- Lidia Angeleri Hügel, Steffen Koenig, Qunhua Liu, and Dong Yang. Ladders and simplicity of derived module categories. 2017
- Hong Xing Chen and Changchang Xi. Recollements of derived categories III: finitistic dimensions. 2017
- Charley Cummings. Ring constructions and generation of the unbounded derived module category. 2020
- Robert M. Fossum, Phillip A. Griffith, and Idun Reiten. Trivial extensions of abelian categories. 1975
- K. R. Fuller and M. Saorín. On the finitistic dimension conjecture for Artinian rings. 1992
- Edward L. Green, Chrysostomos Psaroudakis, and Øyvind Solberg. Reduction techniques for the finitistic dimension. 2018
- Jeremy Rickard. Unbounded derived categories and the finitistic dimension conjecture. 2019