

Atoms in Singularland

Hankyung Ko

Based on joint works with Ben Elias, Nicolas Libedinsky, Leonardo Patimo

FD Seminar, 8 february 2024

Regularland

Regularland

A *Coxeter group* is a group W , together with a finite generating set S , presented by relations of the following form:

Regularland

A *Coxeter group* is a group W , together with a finite generating set S , presented by relations of the following form:

Regularland

A *Coxeter group* is a group W , together with a finite generating set S , presented by relations of the following form:

- ▶ the *braid relation*

$$\underbrace{st \cdots}_m = \underbrace{ts \cdots}_m, \text{ for each } s, t \in S \text{ with } m = m_{st} = m_{ts} < \infty;$$

Regularland

A *Coxeter group* is a group W , together with a finite generating set S , presented by relations of the following form:

- ▶ the *braid relation*

$$\underbrace{st \cdots}_m = \underbrace{ts \cdots}_m, \text{ for each } s, t \in S \text{ with } m = m_{st} = m_{ts} < \infty;$$

- ▶ the *reflection relation*

$$ss = e$$

for each $s \in S$;

Regularland

A *Coxeter group* is a group W , together with a finite generating set S , presented by relations of the following form:

- ▶ the *braid relation*

$$\underbrace{st \cdots}_m = \underbrace{ts \cdots}_m, \text{ for each } s, t \in S \text{ with } m = m_{st} = m_{ts} < \infty;$$

- ▶ the *reflection relation*

$$ss = e$$

for each $s \in S$;

Example. Symmetric groups S_n with $S = \{s_i := (i, i+1)\}_{1 \leq i \leq n-1}$,

Regularland

A *Coxeter group* is a group W , together with a finite generating set S , presented by relations of the following form:

- ▶ the *braid relation*

$$\underbrace{st \cdots}_m = \underbrace{ts \cdots}_m, \text{ for each } s, t \in S \text{ with } m = m_{st} = m_{ts} < \infty;$$

- ▶ the *reflection relation*

$$ss = e$$

for each $s \in S$;

Example. Symmetric groups S_n with $S = \{s_i := (i, i+1)\}_{1 \leq i \leq n-1}$,
with the braid relations

$$s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1} \quad \text{and} \quad s_i s_j = s_j s_i \quad \text{for } |i - j| > 1.$$

Regular expressions

- ▶ An *expression* of $w \in W$ is a string $[s_1, s_2, \dots, s_r]$ of elements in S such that $w = s_1 s_2 \cdots s_r$.

Regular expressions

- ▶ An *expression* of $w \in W$ is a string $[s_1, s_2, \dots, s_r]$ of elements in S such that $w = s_1 s_2 \cdots s_r$.
- ▶ The *length function*: for $w \in W$, the length $\ell(w)$ is the smallest number of letters in an expression of w .

Regular expressions

- ▶ An *expression* of $w \in W$ is a string $[s_1, s_2, \dots, s_r]$ of elements in S such that $w = s_1 s_2 \cdots s_r$.
- ▶ The *length function*: for $w \in W$, the length $\ell(w)$ is the smallest number of letters in an expression of w .
- ▶ An expression $[s_1, \dots, s_r]$ of $w \in W$ is *reduced* if $r = \ell(w)$.

Regular expressions

- ▶ An *expression* of $w \in W$ is a string $[s_1, s_2, \dots, s_r]$ of elements in S such that $w = s_1 s_2 \cdots s_r$.
- ▶ The *length function*: for $w \in W$, the length $\ell(w)$ is the smallest number of letters in an expression of w .
- ▶ An expression $[s_1, \dots, s_r]$ of $w \in W$ is *reduced* if $r = \ell(w)$.

Matsumoto's Theorem

Any two reduced expressions of $w \in W$ are related by braid relations.

Regular expressions

- ▶ An *expression* of $w \in W$ is a string $[s_1, s_2, \dots, s_r]$ of elements in S such that $w = s_1 s_2 \cdots s_r$.
- ▶ The *length function*: for $w \in W$, the length $\ell(w)$ is the smallest number of letters in an expression of w .
- ▶ An expression $[s_1, \dots, s_r]$ of $w \in W$ is *reduced* if $r = \ell(w)$.

Matsumoto's Theorem

Any two reduced expressions of $w \in W$ are related by braid relations.

Fact: if $|W| < \infty$, there's a unique longest element in W .

Regular expressions

- ▶ An *expression* of $w \in W$ is a string $[s_1, s_2, \dots, s_r]$ of elements in S such that $w = s_1 s_2 \cdots s_r$.
- ▶ The *length function*: for $w \in W$, the length $\ell(w)$ is the smallest number of letters in an expression of w .
- ▶ An expression $[s_1, \dots, s_r]$ of $w \in W$ is *reduced* if $r = \ell(w)$.



Matsumoto's Theorem

Any two reduced expressions of $w \in W$ are related by braid relations.

Fact: if $|W| < \infty$, there's a unique longest element in W .

Fact: the finite Coxeter groups are classified into the types $ABDEFH$ and the dihedral groups.

Example: affine symmetric group \tilde{S}_3



Demazure operators

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

The *Demazure operator* for s_i is the map $D_{s_i} : R \rightarrow R$

$$D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

The *Demazure operator* for s_i is the map $D_{s_i} : R \rightarrow R$

$$D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

Theorem [Demazure, 1973]

The Demazure operators satisfy

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

The *Demazure operator* for s_i is the map $D_{s_i} : R \rightarrow R$

$$D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

Theorem [Demazure, 1973]

The Demazure operators satisfy

$$D_{s_i} D_{s_{i+1}} D_{s_i} = D_{s_{i+1}} D_{s_i} D_{s_{i+1}}, \quad D_{s_i} D_{s_j} = D_{s_j} D_{s_i} \text{ for } |i - j| > 1 \quad (\text{braid relations})$$

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

The *Demazure operator* for s_i is the map $D_{s_i} : R \rightarrow R$

$$D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

Theorem [Demazure, 1973]

The Demazure operators satisfy

$$D_{s_i} D_{s_{i+1}} D_{s_i} = D_{s_{i+1}} D_{s_i} D_{s_{i+1}}, \quad D_{s_i} D_{s_j} = D_{s_j} D_{s_i} \text{ for } |i - j| > 1 \quad (\text{braid relations})$$

$$D_{s_i} D_{s_i} = 0. \quad (\text{nilquadratic relations})$$

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

The *Demazure operator* for s_i is the map $D_{s_i} : R \rightarrow R$

$$D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

Theorem [Demazure, 1973]

The Demazure operators satisfy

$$D_{s_i} D_{s_{i+1}} D_{s_i} = D_{s_{i+1}} D_{s_i} D_{s_{i+1}}, \quad D_{s_i} D_{s_j} = D_{s_j} D_{s_i} \text{ for } |i - j| > 1 \quad (\text{braid relations})$$

$$D_{s_i} D_{s_i} = 0. \quad (\text{nilquadratic relations})$$

Moreover, this gives a presentation of the algebra generated by D_{s_i} ,

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

The *Demazure operator* for s_i is the map $D_{s_i} : R \rightarrow R$

$$D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

Theorem [Demazure, 1973]

The Demazure operators satisfy

$$D_{s_i} D_{s_{i+1}} D_{s_i} = D_{s_{i+1}} D_{s_i} D_{s_{i+1}}, \quad D_{s_i} D_{s_j} = D_{s_j} D_{s_i} \text{ for } |i - j| > 1 \quad (\text{braid relations})$$

$$D_{s_i} D_{s_i} = 0. \quad (\text{nilquadratic relations})$$

Moreover, this gives a presentation of the algebra generated by D_{s_i} , called the *nilCoxeter algebra* (or nilHecke algebra).

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

The *Demazure operator* for s_i is the map $D_{s_i} : R \rightarrow R$

$$D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

Theorem [Demazure, 1973]

The Demazure operators satisfy

$$D_{s_i} D_{s_{i+1}} D_{s_i} = D_{s_{i+1}} D_{s_i} D_{s_{i+1}}, \quad D_{s_i} D_{s_j} = D_{s_j} D_{s_i} \text{ for } |i - j| > 1 \quad (\text{braid relations})$$

$$D_{s_i} D_{s_i} = 0. \quad (\text{nilquadratic relations})$$

Moreover, this gives a presentation of the algebra generated by D_{s_i} , called the *nilCoxeter algebra* (or nilHecke algebra). It has a basis $\{D_w\}_{w \in W}$, where $D_w = D_s D_t \cdots D_u$ if $w = st \cdots u$ is a reduced expression.

Demazure operators

Let $(W, S) = (S_n, \{s_1, \dots, s_{n-1}\})$, for simplicity, and consider its natural action on $R = K[x_1, \dots, x_n]$.

The *Demazure operator* for s_i is the map $D_{s_i} : R \rightarrow R$

$$D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

Theorem [Demazure, 1973]

The Demazure operators satisfy

$$D_{s_i} D_{s_{i+1}} D_{s_i} = D_{s_{i+1}} D_{s_i} D_{s_{i+1}}, \quad D_{s_i} D_{s_j} = D_{s_j} D_{s_i} \text{ for } |i - j| > 1 \quad (\text{braid relations})$$

$$D_{s_i} D_{s_i} = 0. \quad (\text{nilquadratic relations})$$

Moreover, this gives a presentation of the algebra generated by D_{s_i} , called the *nilCoxeter algebra* (or nilHecke algebra). It has a basis $\{D_w\}_{w \in W}$, where $D_w = D_s D_t \cdots D_u$ if $w = st \cdots u$ is a reduced expression.

Demazure operators, general

Let (W, S) be a Coxeter group and consider its natural action on $R = \text{Sym}(V)$, where V is a reasonable (faithful etc) [realization](#).

The *Demazure operator* for $s \in S$ is the map $D_s : R \rightarrow R$

$$D_s(f) = \frac{f - s(f)}{\alpha_s}.$$

Theorem [Demazure, 1973]

The Demazure operators satisfy

$$\underbrace{D_s D_t \cdots}_m = \underbrace{D_t D_s \cdots}_m, \text{ for each } s, t \in S \quad (\text{braid relations})$$

$$D_s D_s = 0. \quad (\text{nilquadratic relations})$$

Moreover, this gives a presentation of the algebra generated by D_s , called the *nilCoxeter algebra* (or nilHecke algebra). It has a basis $\{D_w\}_{w \in W}$, where $D_w = D_s D_t \cdots D_u$ if $w = st \cdots u$ is a reduced expression.

Zoom-in on Demazure operators

Note $D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}$ is s_i -invariant, i.e., belongs to

$$R^{s_i} = \{f \in R \mid s_i(f) = f\}.$$

Zoom-in on Demazure operators

Note $D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}$ is s_i -invariant, i.e., belongs to

$$R^{s_i} = \{f \in R \mid s_i(f) = f\}.$$

Decompose the Demazure operator as

$$D_{s_i} : R \xrightarrow{\delta_{s_i}} R^{s_i} \xrightarrow{\iota_{s_i}} R.$$

Zoom-in on Demazure operators

Note $D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}$ is s_i -invariant, i.e., belongs to

$$R^{s_i} = \{f \in R \mid s_i(f) = f\}.$$

Decompose the Demazure operator as

$$D_{s_i} : R \xrightarrow{\delta_{s_i}} R^{s_i} \xrightarrow{\iota_{s_i}} R.$$

Then for example the nilquadratic relation factors as

$$D_{s_i} D_{s_i} = \iota_{s_i} \circ \delta_{s_i} \circ \iota_{s_i} \circ \delta_{s_i} = 0$$

and follows from the zoomed-in relation

$$\delta_{s_i} \circ \iota_{s_i} = 0.$$

Zoom-in on Demazure operators

Note $D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}$ is s_i -invariant, i.e., belongs to

$$R^{s_i} = \{f \in R \mid s_i(f) = f\}.$$

Decompose the Demazure operator as

$$D_{s_i} : R \xrightarrow{\delta_{s_i}} R^{s_i} \xrightarrow{\iota_{s_i}} R.$$

Then for example the nilquadratic relation factors as

$$D_{s_i} D_{s_i} = \iota_{s_i} \circ \delta_{s_i} \circ \iota_{s_i} \circ \delta_{s_i} = 0$$

and follows from the zoomed-in relation

$$\delta_{s_i} \circ \iota_{s_i} = 0.$$

Geometrically: Considering the algebraic groups $GL_n \supset P_i \supset B(\text{orel})$, the ring R is the B -equivariant cohomology; R^{s_i} is the P_i -equivariant cohomology; δ_{s_i} is the pushforward; ι_{s_i} is the pullback.

Parabolic cosets

Parabolic cosets

Let (W, S) be a Coxeter group. For a subset $I \subseteq S$, denote by W_I the subgroup generated by $I \subseteq W$.

Parabolic cosets

Let (W, S) be a Coxeter group. For a subset $I \subseteq S$, denote by W_I the subgroup generated by $I \subseteq W$. Then (W_I, I) is a Coxeter group.

Parabolic cosets

Let (W, S) be a Coxeter group. For a subset $I \subseteq S$, denote by W_I the subgroup generated by $I \subseteq W$. Then (W_I, I) is a Coxeter group.


Example. Symmetric groups have $W_I = S_a \times S_b \times \cdots \times S_c \subseteq S_n$.

Parabolic cosets

Let (W, S) be a Coxeter group. For a subset $I \subseteq S$, denote by W_I the subgroup generated by $I \subseteq W$. Then (W_I, I) is a Coxeter group.

Example. Symmetric groups have $W_I = S_a \times S_b \times \cdots \times S_c \subseteq S_n$.

The **Coxeter complex** consists of the left cosets for W_I in W for all $I \subseteq S$.



Parabolic cosets

Let (W, S) be a Coxeter group. For a subset $I \subseteq S$, denote by W_I the subgroup generated by $I \subseteq W$. Then (W_I, I) is a Coxeter group.

Example. Symmetric groups have $W_I = S_a \times S_b \times \cdots \times S_c \subseteq S_n$.

The **Coxeter complex** consists of the left cosets for W_I in W for all $I \subsetneq S$.

We assume: $I \subseteq S$ is *finitary* i.e., W_I is finite; so W_I has the longest element w_I .

Parabolic cosets

Let (W, S) be a Coxeter group. For a subset $I \subseteq S$, denote by W_I the subgroup generated by $I \subseteq W$. Then (W_I, I) is a Coxeter group.

Example. Symmetric groups have $W_I = S_a \times S_b \times \cdots \times S_c \subseteq S_n$.

The **Coxeter complex** consists of the left cosets for W_I in W for all $I \subsetneq S$.

We assume: $I \subseteq S$ is *finitary* i.e., W_I is finite; so W_I has the longest element w_I .

For $I, J \subseteq S$ finitary, consider the **finitary (parabolic) double cosets**:

$$p = W_I w W_J = \{xwy \in W \mid x \in W_I, y \in W_J\}$$

Parabolic cosets

Let (W, S) be a Coxeter group. For a subset $I \subseteq S$, denote by W_I the subgroup generated by $I \subseteq W$. Then (W_I, I) is a Coxeter group.

Example. Symmetric groups have $W_I = S_a \times S_b \times \cdots \times S_c \subseteq S_n$.

The **Coxeter complex** consists of the left cosets for W_I in W for all $I \subsetneq S$.

We assume: $I \subseteq S$ is **finitary** i.e., W_I is finite; so W_I has the longest element w_I .

For $I, J \subseteq S$ finitary, consider the **finitary (parabolic) double cosets**:

$$p = W_I w W_J = \{xwy \in W \mid x \in W_I, y \in W_J\}$$

- ▶ Each p contains a unique maximal element \bar{p}

Parabolic cosets

Let (W, S) be a Coxeter group. For a subset $I \subseteq S$, denote by W_I the subgroup generated by $I \subseteq W$. Then (W_I, I) is a Coxeter group.

Example. Symmetric groups have $W_I = S_a \times S_b \times \cdots \times S_c \subseteq S_n$.

The **Coxeter complex** consists of the left cosets for W_I in W for all $I \subsetneq S$.

We assume: $I \subseteq S$ is **finitary** i.e., W_I is finite; so W_I has the longest element w_I .

For $I, J \subseteq S$ finitary, consider the **finitary (parabolic) double cosets**:

$$p = W_I w W_J = \{xwy \in W \mid x \in W_I, y \in W_J\}$$

- ▶ Each p contains a unique maximal element \bar{p}
- ▶ Each p contains a unique minimal element \underline{p}

Zoom-in further on Demazure operators

Zoom-in further on Demazure operators

Take for $I \subset S$ the invariant ring

$$R^I = \{f \in R \mid s_i(f) = f \text{ for } s_i \in I\} = R^{W_I}.$$

Zoom-in further on Demazure operators

Take for $I \subset S$ the invariant ring

$$R^I = \{f \in R \mid s_i(f) = f \text{ for } s_i \in I\} = R^{W_I}.$$

Given $I \subset J \subset S$, the operator $D_{w_J w_I^{-1}} : R \rightarrow R$ is shown to restrict to

$$\partial_{[I,J]} : R^I \rightarrow R^J.$$

Zoom-in further on Demazure operators

Take for $I \subset S$ the invariant ring

$$R^I = \{f \in R \mid s_i(f) = f \text{ for } s_i \in I\} = R^{W_I}.$$

Given $I \subset J \subset S$, the operator $D_{w_J w_I^{-1}} : R \rightarrow R$ is shown to restrict to

$$\partial_{[I,J]} : R^I \rightarrow R^J.$$

Also let $\partial_{[J,I]} : R^J \hookrightarrow R^I$ be the inclusion, where $I \subset J \subset S$.

Zoom-in further on Demazure operators

Take for $I \subset S$ the invariant ring

$$R^I = \{f \in R \mid s_i(f) = f \text{ for } s_i \in I\} = R^{W_I}.$$

Given $I \subset J \subset S$, the operator $D_{w_J w_I^{-1}} : R \rightarrow R$ is shown to restrict to

$$\partial_{[I, J]} : R^I \rightarrow R^J.$$

Also let $\partial_{[J, I]} : R^J \hookrightarrow R^I$ be the inclusion, where $I \subset J \subset S$.

Example: $\partial_{[\emptyset, J]} : R \rightarrow R^J$ is given by

$$\partial_{[\emptyset, J]}(f) = D_{w_J}(f) = \frac{\sum_{w \in W_J} (-1)^{\ell(w)} w(f)}{\prod_{i < j} (x_i - x_j)}.$$

Zoom-in further on Demazure operators

Take for $I \subset S$ the invariant ring

$$R^I = \{f \in R \mid s_i(f) = f \text{ for } s_i \in I\} = R^{W_I}.$$

Given $I \subset J \subset S$, the operator $D_{w_J w_I^{-1}} : R \rightarrow R$ is shown to restrict to

$$\partial_{[I, J]} : R^I \rightarrow R^J.$$

Also let $\partial_{[J, I]} : R^J \hookrightarrow R^I$ be the inclusion, where $I \subset J \subset S$.

Example: $\partial_{[\emptyset, J]} : R \rightarrow R^J$ is given by

$$\partial_{[\emptyset, J]}(f) = D_{w_J}(f) = \frac{\sum_{w \in W_J} (-1)^{\ell(w)} w(f)}{\prod_{i < j} (x_i - x_j)}.$$

Remarks. These too come from geometry.

Zoom-in further on Demazure operators

Take for $I \subset S$ the invariant ring

$$R^I = \{f \in R \mid s_i(f) = f \text{ for } s_i \in I\} = R^{W_I}.$$

Given $I \subset J \subset S$, the operator $D_{w_J w_I^{-1}} : R \rightarrow R$ is shown to restrict to

$$\partial_{[I, J]} : R^I \rightarrow R^J.$$

Also let $\partial_{[J, I]} : R^J \hookrightarrow R^I$ be the inclusion, where $I \subset J \subset S$.

Example: $\partial_{[\emptyset, J]} : R \rightarrow R^J$ is given by

$$\partial_{[\emptyset, J]}(f) = D_{w_J}(f) = \frac{\sum_{w \in W_J} (-1)^{\ell(w)} w(f)}{\prod_{i < j} (x_i - x_j)}.$$

Remarks. These too come from geometry. The inclusion $R^J \subset R^I$ is a **Frobenius extension** with trace $\partial_{[I, J]}$.

$$m = \begin{array}{c} \text{---} I \text{---} \\ \text{---} I s \text{---} \end{array}, \quad \Delta = \begin{array}{c} \text{---} I s \text{---} \\ \text{---} I \text{---} \end{array}, \quad \partial = \begin{array}{c} \text{---} I s \text{---} \\ \text{---} I \text{---} \end{array}, \quad \iota = \begin{array}{c} \text{---} I \text{---} \\ \text{---} I s \text{---} \end{array}.$$

Singular Demazure operators

Definition. Let \mathcal{D} be the K -linear category

- ▶ whose objects are R^I where $I \subset S$ are finitary subsets;
- ▶ whose morphisms are generated by the singular Demazure operators $\partial_{[I,J]}, \partial_{[J,I]}$, for $I \subset J \subset S$ finitary.

Singular Demazure operators

Definition. Let \mathcal{D} be the K -linear category

- ▶ whose objects are R^I where $I \subset S$ are finitary subsets;
- ▶ whose morphisms are generated by the singular Demazure operators $\partial_{[I,J]}, \partial_{[J,I]}$, for $I \subset J \subset S$ finitary.

Notation.

$$\partial_{[I_0, I_1, \dots, I_d]} = \partial_{[I_0, I_1]} \partial_{[I_1, I_2]} \cdots \partial_{[I_{d-1}, I_d]}.$$

Singular Demazure operators

Definition. Let \mathcal{D} be the K -linear category

- ▶ whose objects are R^I where $I \subset S$ are finitary subsets;
- ▶ whose morphisms are generated by the singular Demazure operators $\partial_{[I,J]}, \partial_{[J,I]}$, for $I \subset J \subset S$ finitary.

Notation.

$$\partial_{[I_0, I_1, \dots, I_d]} = \partial_{[I_0, I_1]} \partial_{[I_1, I_2]} \cdots \partial_{[I_{d-1}, I_d]}.$$

We want to present \mathcal{D} by (the above) generators and relations

Singular Demazure operators

Definition. Let \mathcal{D} be the K -linear category

- ▶ whose objects are R^I where $I \subset S$ are finitary subsets;
- ▶ whose morphisms are generated by the singular Demazure operators $\partial_{[I,J]}, \partial_{[J,I]}$, for $I \subset J \subset S$ finitary.

Notation.

$$\partial_{[I_0, I_1, \dots, I_d]} = \partial_{[I_0, I_1]} \partial_{[I_1, I_2]} \cdots \partial_{[I_{d-1}, I_d]}.$$

We want to present \mathcal{D} by (the above) generators and relations

Example. if $I \subset J \subset K$ then $\partial_{[I,J,K]} = \partial_{[I,K]}$ and $\partial_{[K,J,I]} = \partial_{[K,I]}$.

Singular Demazure operators

Definition. Let \mathcal{D} be the K -linear category

- ▶ whose objects are R^I where $I \subset S$ are finitary subsets;
- ▶ whose morphisms are generated by the singular Demazure operators $\partial_{[I,J]}, \partial_{[J,I]}$, for $I \subset J \subset S$ finitary.

Notation.

$$\partial_{[I_0, I_1, \dots, I_d]} = \partial_{[I_0, I_1]} \partial_{[I_1, I_2]} \cdots \partial_{[I_{d-1}, I_d]}.$$

We want to present \mathcal{D} by (the above) generators and relations

Example. if $I \subset J \subset K$ then $\partial_{[I,J,K]} = \partial_{[I,K]}$ and $\partial_{[K,J,I]} = \partial_{[K,I]}$.

Note. It is enough to have $\partial_{[I, I_s]}, \partial_{[I_s, I]}$ as generators, where $I_s := I \sqcup \{s\}$.

Singular Demazure operators

Definition. Let \mathcal{D} be the K -linear category

- ▶ whose objects are R^I where $I \subset S$ are finitary subsets;
- ▶ whose morphisms are generated by the singular Demazure operators $\partial_{[I,J]}, \partial_{[J,I]}$, for $I \subset J \subset S$ finitary.

Notation.

$$\partial_{[I_0, I_1, \dots, I_d]} = \partial_{[I_0, I_1]} \partial_{[I_1, I_2]} \cdots \partial_{[I_{d-1}, I_d]}.$$

We want to present \mathcal{D} by (the above) generators and relations

Example. if $I \subset J \subset K$ then $\partial_{[I,J,K]} = \partial_{[I,K]}$ and $\partial_{[K,J,I]} = \partial_{[K,I]}$.

Note. It is enough to have $\partial_{[I, I_s]}, \partial_{[I_s, I]}$ as generators, where $I_s := I \sqcup \{s\}$.

and give a basis.

Expressions in Singularland

Expressions in Singularland

A (singular) *expression* is a string

$$l_{\bullet} = [l_0, l_1, l_2, \dots, l_r]$$

of finitary subsets of S such that,

Expressions in Singularland

A (singular) *expression* is a string

$$l_{\bullet} = [l_0, l_1, l_2, \dots, l_r]$$

of finitary subsets of S such that, for each i , either

$$l_i = l_{i-1} \setminus s \quad \text{('additive' notation : “ $-s$ ”)}$$

or

$$l_i = l_{i-1} s. \quad \text{('additive' notation : “ $+s$ ”)}$$

Expressions in Singularland

A (singular) *expression* is a string

$$l_{\bullet} = [l_0, l_1, l_2, \dots, l_r]$$

of finitary subsets of S such that, for each i , either

$$l_i = l_{i-1} \setminus s \quad \text{('additive' notation : “ $-s$ ”)}$$

or

$$l_i = l_{i-1} s. \quad \text{('additive' notation : “ $+s$ ”)}$$

Expressions in Singularland

A (singular) *expression* is a string

$$l_{\bullet} = [l_0, l_1, l_2, \dots, l_r]$$

of finitary subsets of S such that, for each i , either

$$l_i = l_{i-1} \setminus s \quad \text{('additive' notation : “ $-s$ ”)}$$

or

$$l_i = l_{i-1} s. \quad \text{('additive' notation : “ $+s$ ”)}$$

An expression l_{\bullet} is *reduced* if $\partial_{l_{\bullet}} \neq 0$. (There's a more natural definition.)

Expressions in Singularland

A (singular) *expression* is a string

$$l_{\bullet} = [l_0, l_1, l_2, \dots, l_r]$$

of finitary subsets of S such that, for each i , either

$$l_i = l_{i-1} \setminus s \quad \text{('additive' notation : “ $-s$ ”)}$$

or

$$l_i = l_{i-1} s. \quad \text{('additive' notation : “ $+s$ ”)}$$

An expression l_{\bullet} is *reduced* if $\partial_{l_{\bullet}} \neq 0$. (There's a more natural definition.)

'Regular' example: $[\emptyset, \{s\}, \emptyset, \{t\}, \emptyset, \{s\}, \emptyset, \{u\}, \emptyset, \{t\}, \emptyset, \{s\}, \emptyset]$

Expressions in Singularland

A (singular) *expression* is a string

$$l_{\bullet} = [l_0, l_1, l_2, \dots, l_r]$$

of finitary subsets of S such that, for each i , either

$$l_i = l_{i-1} \setminus s \quad \text{('additive' notation : “ $-s$ ”)}$$

or

$$l_i = l_{i-1} \cup s. \quad \text{('additive' notation : “ $+s$ ”)}$$

An expression l_{\bullet} is *reduced* if $\partial_{l_{\bullet}} \neq 0$. (There's a more natural definition.)

‘Regular’ example: $[\emptyset, \{s\}, \emptyset, \{t\}, \emptyset, \{s\}, \emptyset, \{u\}, \emptyset, \{t\}, \emptyset, \{s\}, \emptyset]$

in additive notation: $[\emptyset + s - s + t - t + s - s + u - u + t - t + s - s]$

Expressions in Singularland

A (singular) *expression* is a string

$$l_{\bullet} = [l_0, l_1, l_2, \dots, l_r]$$

of finitary subsets of S such that, for each i , either

$$l_i = l_{i-1} \setminus s \quad \text{('additive' notation : “ $-s$ ”)}$$

or

$$l_i = l_{i-1} s. \quad \text{('additive' notation : “ $+s$ ”)}$$

An expression l_{\bullet} is *reduced* if $\partial_{l_{\bullet}} \neq 0$. (There's a more natural definition.)

'Regular' example: $[\emptyset, \{s\}, \emptyset, \{t\}, \emptyset, \{s\}, \emptyset, \{u\}, \emptyset, \{t\}, \emptyset, \{s\}, \emptyset]$

in additive notation: $[\emptyset + s - s + t - t + s - s + u - u + t - t + s - s]$

Other examples: $[\emptyset, \{s\}, \{s, t\}, \{s, t, u\}, \{s, u\}, \{u\}, \emptyset], [\{s, t, u\}, \{s, u\}]$

Expressions in Singularland

A (singular) *expression* is a string

$$l_{\bullet} = [l_0, l_1, l_2, \dots, l_r]$$

of finitary subsets of S such that, for each i , either

$$l_i = l_{i-1} \setminus s \quad \text{('additive' notation : “ $-s$ ”)}$$

or

$$l_i = l_{i-1} s. \quad \text{('additive' notation : “ $+s$ ”)}$$

An expression l_{\bullet} is *reduced* if $\partial_{l_{\bullet}} \neq 0$. (There's a more natural definition.)

'Regular' example: $[\emptyset, \{s\}, \emptyset, \{t\}, \emptyset, \{s\}, \emptyset, \{u\}, \emptyset, \{t\}, \emptyset, \{s\}, \emptyset]$

in additive notation: $[\emptyset + s - s + t - t + s - s + u - u + t - t + s - s]$

Other examples: $[\emptyset, \{s\}, \{s, t\}, \{s, t, u\}, \{s, u\}, \{u\}, \emptyset], [\{s, t, u\}, \{s, u\}]$

in additive notation: $[\emptyset + s + t + u - t - s - u], [\{s, t, u\} - t]$

What does a reduced expression express?

What does a reduced expression express?

Rewrite expressions as

$$[[J_0 \supseteq K_1 \subsetneq J_1 \supsetneq K_2 \subsetneq \cdots \supsetneq K_d \subseteq J_d]]. \quad (\star)$$

What does a reduced expression express?

Rewrite expressions as

$$[[J_0 \supseteq K_1 \subsetneq J_1 \supsetneq K_2 \subsetneq \cdots \supsetneq K_d \subseteq J_d]]. \quad (\star)$$

An alternative formulation of regular reduced expression:

A *reduced expression* of $w \in W$ is a string $[s, t, \dots, u]$ in S such that

$$w = st \cdots u$$

and

$$\ell(st \cdots u) = \ell(s) + \ell(t) + \cdots + \ell(u).$$

What does a reduced expression express?

Rewrite expressions as

$$[[J_0 \supseteq K_1 \subsetneq J_1 \supsetneq K_2 \subsetneq \cdots \supsetneq K_d \subseteq J_d]]. \quad (\star)$$

Definition [Williamson 2008, Elias-K. 2021]

Given $I, J \subset S$ finitary, a *reduced expression of a double coset* $p = W_I \backslash W / W_J$ is a string (\star) , with $J_0 = I$ and $J_d = J$, such that

$$\bar{p} = w_{J_0} w_{K_1}^{-1} w_{J_1} w_{K_2}^{-1} w_{J_2} \cdots w_{K_d}^{-1} w_{J_d}$$

and

$$\ell(\bar{p}) = \ell(w_{J_0}) - \ell(w_{K_1}) + \ell(w_{J_1}) - \ell(w_{K_2}) + \cdots - \ell(w_{K_d}) + \ell(w_{J_d}).$$

Definition [Williamson 2008, Elias-K. 2021]

A *reduced expression* of $p = W_I w W_J$ is a string

$$[[J_0 \supset K_1 \subset J_1 \supset K_2 \subset \cdots \supset K_d \subset J_d]],$$

with $J_0 = I$ and $J_d = J$, such that

$$\bar{p} = w_{J_0} w_{K_1}^{-1} w_{J_1} w_{K_2}^{-1} w_{J_2} \cdots w_{K_d}^{-1} w_{J_d}$$

and

$$\ell(\bar{p}) = \ell(w_{J_0}) - \ell(w_{K_1}) + \ell(w_{J_1}) - \ell(w_{K_2}) + \cdots - \ell(w_{K_d}) + \ell(w_{J_d}).$$

Example. A ‘regular’ singular expression

$$\{w\} \Leftrightarrow [\emptyset + s - s + t - t + \cdots + u - u]$$

Definition [Williamson 2008, Elias-K. 2021]

A *reduced expression* of $p = W_I w W_J$ is a string

$$[[J_0 \supset K_1 \subset J_1 \supset K_2 \subset \cdots \supset K_d \subset J_d]],$$

with $J_0 = I$ and $J_d = J$, such that

$$\bar{p} = w_{J_0} w_{K_1}^{-1} w_{J_1} w_{K_2}^{-1} w_{J_2} \cdots w_{K_d}^{-1} w_{J_d}$$

and

$$\ell(\bar{p}) = \ell(w_{J_0}) - \ell(w_{K_1}) + \ell(w_{J_1}) - \ell(w_{K_2}) + \cdots - \ell(w_{K_d}) + \ell(w_{J_d}).$$

Example. A ‘regular’ singular expression

$$\{w\} \Leftrightarrow [\emptyset + s - s + t - t + \cdots + u - u]$$

is reduced if and only if $w = st \cdots u$ is reduced.

Singular braid relations

As noted above, we have the relations

$$[J + s + t] \Leftrightarrow [J + t + s] \quad (\textit{upup relation})$$

$$[J - s - t] \Leftrightarrow [J - t - s] \quad (\textit{downdown relation})$$

between reduced expressions.

Singular braid relations

As noted above, we have the relations

$$[J + s + t] \Leftrightarrow [J + t + s] \quad (\text{upup relation})$$

$$[J - s - t] \Leftrightarrow [J - t - s] \quad (\text{downdown relation})$$

between reduced expressions.

Some nontrivial relations

The *switchback relations* are of the form

$$[J + u_0 - u_d] \Leftrightarrow [J - u_1 + u_0 - u_2 + u_1 - u_3 + u_2 \cdots - u_{d-1} + u_{d-2} - u_d + u_{d-1}]$$

where u_i is a certain periodic sequence in (finitary) $J' = J \sqcup \{u_0\} \subseteq S$.

Singular braid relations

As noted above, we have the relations

$$[J + s + t] \Leftrightarrow [J + t + s] \quad (\text{upup relation})$$

$$[J - s - t] \Leftrightarrow [J - t - s] \quad (\text{downdown relation})$$

between reduced expressions.

Some nontrivial relations

The *switchback relations* are of the form

$$[J + u_0 - u_d] \Leftrightarrow [J - u_1 + u_0 - u_2 + u_1 - u_3 + u_2 \cdots - u_{d-1} + u_{d-2} - u_d + u_{d-1}]$$

where u_i is a certain periodic sequence in (finitary) $J' = J \sqcup \{u_0\} \subseteq S$.

The u_i sequence and the number d depend on $u_0 \neq u_1$ and the type of J' .

Singular braid relations

As noted above, we have the relations

$$[J + s + t] \Leftrightarrow [J + t + s] \quad (\text{upup relation})$$

$$[J - s - t] \Leftrightarrow [J - t - s] \quad (\text{downdown relation})$$

between reduced expressions.

Some nontrivial relations

The *switchback relations* are of the form

$$[J + u_0 - u_d] \Leftrightarrow [J - u_1 + u_0 - u_2 + u_1 - u_3 + u_2 \cdots - u_{d-1} + u_{d-2} - u_d + u_{d-1}]$$

where u_i is a certain periodic sequence in (finitary) $J' = J \sqcup \{u_0\} \subseteq S$.

The u_i sequence and the number d depend on $u_0 \neq u_1$ and the type of J' .

We call these (singular) *braid relations*.

Singular Coxeter presentation

Singular Matsumoto Theorem [Elias-K.]

For finitary $I, J \subseteq S$ and a double coset $p \in W_I \backslash W / W_J$, any two reduced expressions of p are related by braid relations (upup, downdown, and switchback).

Singular Coxeter presentation

Singular Matsumoto Theorem [Elias-K.]

For finitary $I, J \subseteq S$ and a double coset $p \in W_I \backslash W / W_J$, any two reduced expressions of p are related by braid relations (upup, downdown, and switchback).

It follows that $\partial_p := \partial_{I_\bullet}$, where I_\bullet is a reduced expression for p , is well-defined.

Singular Coxeter presentation

Singular Matsumoto Theorem [Elias-K.]

For finitary $I, J \subseteq S$ and a double coset $p \in W_I \backslash W / W_J$, any two reduced expressions of p are related by braid relations (upup, downdown, and switchback).

It follows that $\partial_p := \partial_{I_\bullet}$, where I_\bullet is a reduced expression for p , is well-defined.

Non-reduced expressions? Recall the following zoomed-in version of $D_s D_s = 0$.

$$\partial_{[Js, J, Js]} = \partial_{[Js-s+s]} = 0. \quad (\text{nilquadratic relation})$$

Singular Coxeter presentation

Singular Matsumoto Theorem [Elias-K.]

For finitary $I, J \subseteq S$ and a double coset $p \in W_I \backslash W / W_J$, any two reduced expressions of p are related by braid relations (upup, downdown, and switchback).

It follows that $\partial_p := \partial_{I_\bullet}$, where I_\bullet is a reduced expression for p , is well-defined.

Non-reduced expressions? Recall the following zoomed-in version of $D_s D_s = 0$.

$$\partial_{[Js, J, Js]} = \partial_{[Js-s+s]} = 0. \quad (\text{nilquadratic relation})$$

Presentation Theorem [Elias-K., Elias-K.-Libedinsky-Patimo]

The braid and nilquadratic relations generate all relations between compositions of Demazure operators. That is, \mathcal{D} has a presentation by the generators $\partial_{[I, Is]}, \partial_{[Is, I]}$ and the above relations.

Singular Coxeter presentation

Singular Matsumoto Theorem [Elias-K.]

For finitary $I, J \subseteq S$ and a double coset $p \in W_I \backslash W / W_J$, any two reduced expressions of p are related by braid relations (upup, downdown, and switchback).

It follows that $\partial_p := \partial_{I_\bullet}$, where I_\bullet is a reduced expression for p , is well-defined.

Non-reduced expressions? Recall the following zoomed-in version of $D_s D_s = 0$.

$$\partial_{[Js, J, Js]} = \partial_{[Js-s+s]} = 0. \quad (\text{nilquadratic relation})$$

Presentation Theorem [Elias-K., Elias-K.-Libedinsky-Patimo]

The braid and nilquadratic relations generate all relations between compositions of Demazure operators. That is, \mathcal{D} has a presentation by the generators $\partial_{[I, Is]}, \partial_{[Is, I]}$ and the above relations. Moreover, the category \mathcal{D} has a basis

$$\{\partial_p \mid p \in W_I \backslash W / W_J, I, J \subseteq S \text{ finitary}\}.$$

Core double cosets

Let $p \in W_I \backslash W / W_J$ be a finitary double coset.

Core double cosets

Let $p \in W_I \backslash W / W_J$ be a finitary double coset.

Definition. The *left (resp. right) redundancy* of p is the subset

$$\text{LR}(p) = I \cap \underline{p} J \underline{p}^{-1} \subset I, \quad \text{resp.}, \quad \text{RR}(p) = \underline{p}^{-1} I \underline{p} \cap J \subset J,$$

Core double cosets

Let $p \in W_I \backslash W / W_J$ be a finitary double coset.

Definition. The *left (resp. right) redundancy* of p is the subset

$$\text{LR}(p) = I \cap \underline{p} J \underline{p}^{-1} \subset I, \quad \text{resp.}, \quad \text{RR}(p) = \underline{p}^{-1} I \underline{p} \cap J \subset J,$$

Definition. We call p a *core coset* if $\text{LR}(p) = I$ and $\text{RR}(p) = J$.

Core double cosets

Let $p \in W_I \backslash W / W_J$ be a finitary double coset.

Definition. The *left (resp. right) redundancy* of p is the subset

$$\text{LR}(p) = I \cap \underline{p} J \underline{p}^{-1} \subset I, \quad \text{resp.}, \quad \text{RR}(p) = \underline{p}^{-1} I \underline{p} \cap J \subset J,$$

Definition. We call p a *core coset* if $\text{LR}(p) = I$ and $\text{RR}(p) = J$.

Remark. A coset $W_I w W_J$ is core if and only if $W_I w W_J = w W_J = W_I w$.

Core double cosets

Let $p \in W_I \backslash W / W_J$ be a finitary double coset.

Definition. The *left (resp. right) redundancy* of p is the subset

$$\text{LR}(p) = I \cap \underline{p} J \underline{p}^{-1} \subset I, \quad \text{resp.}, \quad \text{RR}(p) = \underline{p}^{-1} I \underline{p} \cap J \subset J,$$

Definition. We call p a *core coset* if $\text{LR}(p) = I$ and $\text{RR}(p) = J$.

Remark. A coset $W_I w W_J$ is core if and only if $W_I w W_J = w W_J = W_I w$.

Example. Any $W_\emptyset w W_\emptyset = \{w\}$ is core.

Core double cosets

Let $p \in W_I \backslash W / W_J$ be a finitary double coset.

Definition. The *left (resp. right) redundancy* of p is the subset

$$\text{LR}(p) = I \cap \underline{p} J \underline{p}^{-1} \subset I, \quad \text{resp.}, \quad \text{RR}(p) = \underline{p}^{-1} I \underline{p} \cap J \subset J,$$

Definition. We call p a *core coset* if $\text{LR}(p) = I$ and $\text{RR}(p) = J$.

Remark. A coset $W_I w W_J$ is core if and only if $W_I w W_J = w W_J = W_I w$.

Example. Any $W_\emptyset w W_\emptyset = \{w\}$ is core.


The low road theorem [Elias-K. 2021]

Given $p \in W_I \backslash W / W_J$, the double coset $p^{\text{core}} := W_{\text{LR}(p)} \underline{p} W_{\text{RR}(p)}$ is a core coset. Moreover, if $p^{\text{core}} \Leftrightarrow M_\bullet$ is a reduced expression then


$$p \Leftrightarrow [[I \supset \text{LR}(p)]] \circ M_\bullet \circ [[\text{RR}(p) \subset J]]$$

is a reduced expression.


Left cosets in \tilde{S}_3



Core cosets in \tilde{S}_3




Core cosets in $W_{\{g\}} \backslash W/W_J$, where $J \subset S$ runs



Core cosets in $W_I \backslash W / W_J$, where $J \subset S$ runs

is called *Tits cone intersection* by Iyama-Wemyss whose pictures I paste:



Atomic expressions

Definition. An *atom* is a core coset with a (unique) reduced expression

$$a \Leftrightarrow [l + s - t].$$

Atomic expressions

Definition. An *atom* is a core coset with a (unique) reduced expression

$$a \Leftrightarrow [I + s - t].$$

Example. For each $s \in S$ the coset $\{s\} \Leftrightarrow [\emptyset + s - s]$ is an atom.

Atomic expressions

Definition. An *atom* is a core coset with a (unique) reduced expression

$$a \Leftrightarrow [I + s - t].$$

Example. For each $s \in S$ the coset $\{s\} \Leftrightarrow [\emptyset + s - s]$ is an atom.

Definition. An *atomic (reduced) expression* is a(n reduced) expression of the form

$$p \Leftrightarrow a \circ a' \circ \cdots \circ a'' := [I + s - t + s' - t' + \cdots + s'' - t'']$$

where a, a', \dots, a'' are atoms. (Or think of

$$\partial_p = \partial_a \partial_{a'} \cdots \partial_{a''} = \partial_{[I+s-t+s'-t'+\cdots+s''-t'']}.)$$

Atomic expressions

Definition. An *atom* is a core coset with a (unique) reduced expression

$$a \Leftrightarrow [I + s - t].$$

Example. For each $s \in S$ the coset $\{s\} \Leftrightarrow [\emptyset + s - s]$ is an atom.

Definition. An *atomic (reduced) expression* is a(n reduced) expression of the form


$$p \Leftrightarrow a \circ a' \circ \cdots \circ a'' := [I + s - t + s' - t' + \cdots + s'' - t'']$$

where a, a', \dots, a'' are atoms. (Or think of

$$\partial_p = \partial_a \partial_{a'} \cdots \partial_{a''} = \partial_{[I + s - t + s' - t' + \cdots + s'' - t'']}.)$$

Example. For a regular (reduced) expression $w = st \cdots u$ in W , the expression $\{w\} \Leftrightarrow [\emptyset + s - s + t - t + \cdots + u - u]$ is an atomic (reduced) expression.

Atoms in \tilde{S}_3



An atomic expression



Atomic Coxeter presentation

Atomic Coxeter presentation

Let \mathcal{D}^{at} be the subcategory of \mathcal{D} generated by the atoms, i.e., the morphisms ∂_a where a are atoms.

Atomic Coxeter presentation

Let \mathcal{D}^{at} be the subcategory of \mathcal{D} generated by the atoms, i.e., the morphisms ∂_a where a are atoms.

Theorem [Elias-K.-Libedinsky-Patimo, K.]

The category \mathcal{D}^{at} has a basis

$$\{\partial_p \mid p \text{ is a core double coset}\}$$

Atomic Coxeter presentation

Let \mathcal{D}^{at} be the subcategory of \mathcal{D} generated by the atoms, i.e., the morphisms ∂_a where a are atoms.

Theorem [Elias-K.-Libedinsky-Patimo, K.]

The category \mathcal{D}^{at} has a basis

$$\{\partial_p \mid p \text{ is a core double coset}\}$$

and a presentation by generators (the atoms ∂_a) and the following relations:

- ▶ *atomic braid relations*

$$\underbrace{\partial_a \partial_{b'} \partial_{a''} \cdots}_m = \underbrace{\partial_b \partial_{a'} \partial_{b''} \cdots}_m,$$

where a', a'', \dots and b', b'', \dots are certain twists of the atoms a and b ; and $m = m_{a,b} \geq 2$ is an integer determined by a switchback relation.

- ▶ *atomic nilquadratic relations*

$$\partial_a \partial_{a^{-1}} = 0.$$

Tilting modules for preprojective algebras

Let $Q = (Q_0, Q_1)$ be a quiver and let $\overline{Q} = (Q_0, Q_1 \sqcup Q_1^{op})$ be the double quiver.

Tilting modules for preprojective algebras

Let $Q = (Q_0, Q_1)$ be a quiver and let $\overline{Q} = (Q_0, Q_1 \sqcup Q_1^{op})$ be the double quiver. We will assume that Q is extended Dynkin, i.e., the Dynkin diagram of a (simply-laced) affine Coxeter type.

Tilting modules for preprojective algebras

Let $Q = (Q_0, Q_1)$ be a quiver and let $\overline{Q} = (Q_0, Q_1 \sqcup Q_1^{op})$ be the double quiver. We will assume that Q is extended Dynkin, i.e., the Dynkin diagram of a (simply-laced) affine Coxeter type. Let (W, S) be this Coxeter group.

Tilting modules for preprojective algebras

Let $Q = (Q_0, Q_1)$ be a quiver and let $\overline{Q} = (Q_0, Q_1 \sqcup Q_1^{op})$ be the double quiver. We will assume that Q is extended Dynkin, i.e., the Dynkin diagram of a (simply-laced) affine Coxeter type. Let (W, S) be this Coxeter group.

Definition. The associated *preprojective algebra* Π is the completion of the path algebra for \overline{Q} modulo the ideal $(\sum_{\alpha \in Q_1} \alpha\alpha^* - \alpha^*\alpha)$.

Tilting modules for preprojective algebras

Let $Q = (Q_0, Q_1)$ be a quiver and let $\overline{Q} = (Q_0, Q_1 \sqcup Q_1^{op})$ be the double quiver. We will assume that Q is extended Dynkin, i.e., the Dynkin diagram of a (simply-laced) affine Coxeter type. Let (W, S) be this Coxeter group.

Definition. The associated *preprojective algebra* Π is the completion of the path algebra for \overline{Q} modulo the ideal $(\sum_{\alpha \in Q_1} \alpha \alpha^* - \alpha^* \alpha)$.

Theorem [Iyama-Reiten, Buan-Iyama-Reiten-Scott].

The ideals

$$I_s = (1 - e_s), \quad \text{for } s \in Q_0 = S$$

and their products are *tilting* modules. Moreover, we have a bijection

$$W \rightarrow \{(\text{basic}) \text{ tilting modules for } \Pi\}$$

given by $w \mapsto I_w = I_s I_t \cdots I_u$ where $w = st \cdots u$ is a reduced expression in (W, S) .

Tilting theory for contracted preprojective algebras

In the same setting, for $J \subset S = Q_0$ consider the idempotent $e_J = 1 - \sum_{s \in J} e_s$ and the *contracted preprojective algebra*

$$\Gamma_J = e_J \Pi e_J.$$

Tilting theory for contracted preprojective algebras

In the same setting, for $J \subset S = Q_0$ consider the idempotent $e_J = 1 - \sum_{s \in J} e_s$ and the *contracted preprojective algebra*

$$\Gamma_J = e_J \Pi e_J.$$

Theorem [Iyama-Wemyss, *Tits Cone Intersections and Applications*].

Let Q be extended Dynkin. Then we have a bijection

$$\{\text{core double cosets in } (W, S)\} \rightarrow \bigsqcup_{J \subsetneq S} \{\text{tilting modules for } \Gamma_J\}$$

given by $W_J x W_I \mapsto e_J l_x e_I$.

Tilting theory for contracted preprojective algebras

In the same setting, for $J \subset S = Q_0$ consider the idempotent $e_J = 1 - \sum_{s \in J} e_s$ and the *contracted preprojective algebra*

$$\Gamma_J = e_J \Pi e_J.$$

Theorem [Iyama-Wemyss, *Tits Cone Intersections and Applications*].

Let Q be extended Dynkin. Then we have a bijection

$$\{\text{core double cosets in } (W, S)\} \rightarrow \bigsqcup_{J \subsetneq S} \{\text{tilting modules for } \Gamma_J\}$$

given by $W_J x W_I \mapsto e_J l_x e_I$. Moreover, a reduced composition $- \circ a$ by an atom corresponds to a tilting mutation on the right side.

Theorem [Iyama-Wemyss, *Tits Cone Intersections and Applications*]

Let Q be extended Dynkin and fix $J \subsetneq S$. Then we have a bijection

$$\{\text{core double cosets in } W_J \backslash W / W_I\} \rightarrow \{\text{tilting modules for } \Gamma_J\}.$$

Theorem [Iyama-Wemyss], original version.

Let Q be extended Dynkin and fix $J \subsetneq S$. Then we have a bijection


$$\{\text{chambers in the Tits } J\text{-cone}\} \rightarrow \{\text{tilting modules for } \Gamma_J\}.$$

Theorem [Iyama-Wemyss], original version.

Let Q be extended Dynkin and fix $J \subsetneq S$. Then we have a bijection

$$\{\text{chambers in the Tits } J\text{-cone}\} \rightarrow \{\text{tilting modules for } \Gamma_J\}.$$

If $|S \setminus J| = 3$ then the Tits J -cone is one of the following types:



An atomic Leibniz rule (in progress with ELP)

Recall the regular Demazure operator $D_s : R \rightarrow R$

$$D_s(f) = \frac{f - s(f)}{\alpha_s}, \quad (\text{or, for } s = s_i \in S_n,) \quad D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

An atomic Leibniz rule (in progress with ELP)

Recall the regular Demazure operator $D_s : R \rightarrow R$

$$D_s(f) = \frac{f - s(f)}{\alpha_s}, \quad (\text{or, for } s = s_i \in S_n,) \quad D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

The (*twisted*) *Leibniz rule* is well-known:

$$D_s(f \cdot g) = s(f)D_s(g) + D_s(f)g.$$

An atomic Leibniz rule (in progress with ELP)

Recall the regular Demazure operator $D_s : R \rightarrow R$

$$D_s(f) = \frac{f - s(f)}{\alpha_s}, \quad (\text{or, for } s = s_i \in S_n,) \quad D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

The (*twisted*) *Leibniz rule* is well-known:

$$D_s(f \cdot g) = s(f)D_s(g) + D_s(f)g.$$

Singular analogue.

An atomic Leibniz rule (in progress with ELP)

Recall the regular Demazure operator $D_s : R \rightarrow R$

$$D_s(f) = \frac{f - s(f)}{\alpha_s}, \quad (\text{or, for } s = s_i \in S_n, \quad D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}).$$

The (*twisted*) *Leibniz rule* is well-known:

$$D_s(f \cdot g) = s(f)D_s(g) + D_s(f)g.$$

Singular analogue. An *atomic Leibniz rule* (for an atom a) is

$$\partial_a(f \cdot g) = \underline{p}(f)\partial_a(g) + \sum_{p < a} \partial_p(T_p(f) \cdot g) \quad (\star)$$

where $T_p(f)$ is some element depending on the coset $p < a$ and we use the *Bruhat order* on double cosets.

An atomic Leibniz rule (in progress with ELP)

Recall the regular Demazure operator $D_s : R \rightarrow R$

$$D_s(f) = \frac{f - s(f)}{\alpha_s}, \quad (\text{or, for } s = s_i \in S_n,) \quad D_{s_i}(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$

The (*twisted*) *Leibniz rule* is well-known:

$$D_s(f \cdot g) = s(f)D_s(g) + D_s(f)g.$$

Singular analogue. An *atomic Leibniz rule* (for an atom \mathbf{a}) is

$$\partial_{\mathbf{a}}(f \cdot g) = \underline{p}(f)\partial_{\mathbf{a}}(g) + \sum_{p < \mathbf{a}} \partial_p(T_p(f) \cdot g) \quad (\star)$$

where $T_p(f)$ is some element depending on the coset $p < \mathbf{a}$ and we use the [Bruhat order](#) on double cosets.

Remark. The summands in (\star) have interpretation in terms of [singular light leaves](#), a basis of the [singular Hecke category](#), aka [singular Soergel bimodules](#), and (\star) is equivalent to an essential property of [singular Bott-Samelson bimodules](#) for [singular Soergel calculus](#).

An atomic Leibniz rule (in progress with ELP)

Recall the regular De

$$D_s(f) = \frac{f - s_i(f)}{x_i - x_{i+1}}$$

$$f) = \frac{f - s_i(f)}{x_i - x_{i+1}}.$$


The *(twisted) Leibni*

Singular analogue.

where $T_p(f)$ is some

Bruhat order on dou_s

Remark. The summands in (\star) have interpretation in terms of **singular light leaves**, a basis of the **singular Hecke category**, aka **singular Soergel bimodules**, and (\star) is equivalent to an essential property of **singular Bott-Samelson bimodules** for **singular Soergel calculus**.



\mathfrak{a}) is

$g)$

(\star)

a and we use the

Papers in Singularland (boldface for cited)

Elias-K., **A Singular Coxeter presentation**, arXiv:2105.08563 (2023)

The Singular Land (with Elias, Libedinsky, Patimo), Season 1 (2023-2024)

EKLP, **Demazure operators for double cosets**, arXiv:2307.15021

EKLP, Subexpressions and the Bruhat order for double cosets, arXiv:2307.15726

EKLP, **On reduced expressions for core double cosets** (to be posted)

EKLP, Singular Light Leaves, arXiv:2401.03053

K, **An Atomic Coxeter presentation**, arXiv:2312.16666

The Singular Land Season 2 is coming:

Atomic Leibniz rule, Singular Soergel calculus, ...