

The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

Def (Reid 1983) $R \cong \mathbb{C}[[x, y, z, t]]/(f)$

is a compound Du Val singularity if

$$f = g(x, y, z) + t^k h(x, y, z, t)$$

\uparrow
 $g=0$ Kleinian / Du Val singularity

\hookrightarrow arbitrary

The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

Def (Reid 1983) $R \cong \mathbb{C}[[x, y, z, t]]/(f)$

is a compound Du Val singularity if

$$f = g(x, y, z) + t^k h(x, y, z, t)$$

\uparrow \uparrow arbitrary

$g=0$ Kleinian / Du Val singularity

e.g. $R = \mathbb{C}[[u, v, x, y]]/(uv - xy)$

\uparrow Simplest example

The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

Def (Reid 1983) $R \cong \mathbb{C}[[x, y, z, t]]/(f)$

is a compound Du Val singularity if

$$f = g(x, y, z) + t^k h(x, y, z, t)$$

\uparrow \uparrow arbitrary

$g=0$ Kleinian / Du Val singularity

e.g. $R = \mathbb{C}[[u, v, x, y]]/(uv - xy)$

\uparrow Simplest example

Standing assumptions

- $\text{Spec } R$ has an isolated singularity
- $\exists p: X \rightarrow \text{Spec } R$ crepant resolution
 $p^* w_R \cong w_X$

The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

Def (Reid 1983) $R \cong \mathbb{C}[[x, y, z, t]]/(f)$

is a compound Du Val singularity if

$$f = g(x, y, z) + t^k h(x, y, z, t)$$

\uparrow arbitrary

$g=0$ Kleinian / Du Val singularity

e.g. $R = \mathbb{C}[[u, v, x, y]]/(uv - xy)$

\uparrow Simplest example

Standing assumptions

- $\text{Spec } R$ has an isolated singularity
- $\exists p: X \rightarrow \text{Spec } R$ crepant resolution
 $p^* w_R \cong w_X$

Rmk In general, $\text{Spec } R$ need not admit a crepant resolution and, if it does, it is not unique

The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

Def (Reid 1983) $R \cong \mathbb{C}[[x,y,z,t]]/(f)$

is a compound Du Val singularity if

$$f = g(x,y,z) + t h(x,y,z,t)$$

\uparrow arbitrary

g=0 Kleinian / Du Val singularity


e.g. $R = \mathbb{C}[[u,v,x,y]]/(uv - xy)$

\uparrow Simplest example

Standing assumptions

- $\text{Spec } R$ has an isolated singularity
- $\exists p: X \rightarrow \text{Spec } R$ crepant resolution
 $p^* w_R \cong w_X$

Rmk In general, $\text{Spec } R$ need not admit a crepant resolution and, if it does, it is not unique



The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

Def (Reid 1983) $R \cong \mathbb{C}[[x, y, z, t]]/(f)$

is a compound Du Val singularity if

$$f = g(x, y, z) + t h(x, y, z, t)$$

\uparrow arbitrary

$g=0$ Kleinian / Du Val singularity


$$\text{e.g. } R = \mathbb{C}[[u, v, x, y]]/(uv - xy)$$

\uparrow Simplest example

Standing assumptions

- $\text{Spec } R$ has an isolated singularity
- $\exists p: X \rightarrow \text{Spec } R$ crepant resolution
 $p^* w_R \cong w_X$

Rmk In general, $\text{Spec } R$ need not admit a crepant resolution and, if it does, it is not unique



Thm (Donovan - Wemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

Def (Reid 1983) $R \cong \mathbb{C}[[x,y,z,t]]/(f)$

is a compound Du Val singularity if

$$f = g(x,y,z) + t h(x,y,z,t)$$

\uparrow \uparrow arbitrary

$g=0$ Kleinian / Du Val singularity


e.g. $R = \mathbb{C}[[u,v,x,y]]/(uv - xy)$

\uparrow Simplest example

Standing assumptions

- $\text{Spec } R$ has an isolated singularity
- $\exists p: X \rightarrow \text{Spec } R$ crepant resolution
 $p^* w_R \cong w_X$

Rmk In general, $\text{Spec } R$ need not admit a crepant resolution and, if it does, it is not unique



Thm (Donovan - Wemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

e.g. $R = \mathbb{C}[[u,v,x,y]]/(uv - xy)$

$$\begin{array}{ccc} X & & Y \\ p \searrow & & \swarrow q \\ & \text{Spec } R & \end{array}$$

$\Delta(p) \cong \Delta(q) \cong \mathbb{C}$

misleadingly simple...

The Donovan - Wemyss Conjecture via the Derived Auslander - Iyama Correspondence

(based on joint work with Muro and an insight of Keller)

Def (Reid 1983) $R \cong \mathbb{C}[[x, y, z, t]]/(f)$

is a compound Du Val singularity if

$$f = g(x, y, z) + t h(x, y, z, t)$$

\uparrow arbitrary

$g=0$ Kleinian / Du Val singularity


e.g. $R = \mathbb{C}[[u, v, x, y]]/(uv - xy)$

↑ Simplest example

Standing assumptions

- $\text{Spec } R$ has an isolated singularity
- $\exists p: X \rightarrow \text{Spec } R$ crepant resolution
 $p^* w_R \cong w_X$

Rmk In general, $\text{Spec } R$ need not admit a crepant resolution and, if it does, it is not unique



Thm (Donovan - Wemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

e.g. $R = \mathbb{C}[[u, v, x, y]]/(uv - xy)$

$$\begin{array}{ccc} X & & Y \\ p \searrow & & \swarrow q \\ & \text{Spec } R & \end{array}$$

$$\Delta(p) \cong \Delta(q) \cong \mathbb{C}$$

misleadingly simple...

Rmk $\Delta(p)$ recovers all known numerical invariants associated with $p: X \rightarrow \text{Spec } R$

$\text{Spec } R$: cDW w/ isolated singularity
 $\exists p: X \rightarrow \text{Spec } R$ crepant resolution

$$xxy \xrightarrow{p} \bullet \text{Spec } R \quad p^{-1}(0) = \bigcup_{i=1}^n C_i : \underline{\text{exc. fibre}}$$

$$C_i \cong \mathbb{P}^1$$

Thm (Donovan-Wemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

e.g. $R = \mathbb{C}[u, v, x, y]/(uv - xy)$

$$xxy \xrightarrow{p} \bullet \text{Spec } R \quad q \downarrow \quad X \xrightarrow{Y}$$

$$\Delta(p) \cong \Delta(q) \cong \mathbb{C}$$

misleadingly simple ...

Rank $\Delta(p)$ recovers all known numerical invariants associated with $p: X \rightarrow \text{Spec } R$

$\text{Spec } R$: cDV w/ isolated singularity
 $\exists p: X \rightarrow \text{Spec } R$ crepant resolution

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1,2$)

$$R_1 \cong R_2 \Leftrightarrow D^b(\text{mod } \Delta(p_1)) \simeq D^b(\text{mod } \Delta(p_2))$$

Thm (Donovan-Wemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

e.g. $R = \mathbb{C}[u,v,x,y]/(uv - xy)$

Rank $\Delta(p)$ recovers all known numerical invariants associated with $p: X \rightarrow \text{Spec } R$

$\text{Spec } R$: cDV w/ isolated singularity
 $\exists p: X \rightarrow \text{Spec } R$ crepant resolution

$$p^{-1}(0) = \bigcup_{i=1}^n C_i : \underline{\text{exc. fibre}}$$

$$C_i \cong \mathbb{P}^1$$

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1,2$)

$$R_1 \cong R_2 \Leftrightarrow \mathcal{D}^b(\text{mod } \Delta(p_1)) \simeq \mathcal{D}^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Wemyss (2018) + Dugas (2015)

(\Leftarrow) Reid (1983) in type A

Thm (Donovan-Wemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

e.g. $R = \mathbb{C}[u,v,x,y]/(uv - xy)$

$$\Delta(p) \cong \Delta(q) \cong \mathbb{C}$$

misleadingly simple...

Rank $\Delta(p)$ recovers all known numerical invariants associated with $p: X \rightarrow \text{Spec } R$

$\text{Spec } R$: cDV w/ isolated singularity
 $\exists p: X \rightarrow \text{Spec } R$ crepant resolution

$$p^{-1}(0) = \bigcup_{i=1}^n C_i : \underline{\text{exc. fibre}}$$

$$C_i \cong \mathbb{P}^1$$

Thm (Donovan-Wemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

e.g. $R = \mathbb{C}[u, v, x, y]/(uv - xy)$

$$\Delta(p) \cong \Delta(q) \cong \mathbb{C}$$

misleadingly simple...

Rank $\Delta(p)$ recovers all known numerical invariants associated with $p: X \rightarrow \text{Spec } R$

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1,2$)

$$R_1 \cong R_2 \Leftrightarrow \mathcal{D}^b(\text{mod } \Delta(p_1)) \simeq \mathcal{D}^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Wemyss (2018) + Dugas (2015)

(\Leftarrow) Reid (1983) in type A

Recall $\mathcal{D}_{\text{sg}}(R) := \mathcal{D}^b(\text{mod } R)/\mathcal{K}^b(\text{proj } R)$

is the singularity cat of R

$\text{Spec } R$: cDV w/ isolated singularity
 $\exists p: X \rightarrow \text{Spec } R$ crepant resolution

$$p^{-1}(0) = \bigcup_{i=1}^n C_i : \underline{\text{exc. fibre}}$$

$$C_i \cong \mathbb{P}^1$$

Thm (Donovan-Wemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

e.g. $R = \mathbb{C}[u, v, x, y]/(uv - xy)$

$$\Delta(p) \cong \Delta(q) \cong \mathbb{C}$$

misleadingly simple ...

Rank $\Delta(p)$ recovers all known numerical invariants associated with $p: X \rightarrow \text{Spec } R$

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1, 2$)

$$R_1 \cong R_2 \Leftrightarrow \mathcal{D}^b(\text{mod } \Delta(p_1)) \cong \mathcal{D}^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Wemyss (2018) + Dugas (2015)


(\Leftarrow) Reid (1983) in type A

Recall $\mathcal{D}_{\text{sg}}(R) := \mathcal{D}^b(\text{mod } R)/\mathcal{K}^b(\text{proj } R)$

is the singularity cat of R

- Hom-finite & Krull-Schmidt
- 2-Calabi-Yau $\text{Hom}(Y, X[2]) \cong D\text{Hom}(X, Y)$
- $[2] \cong \mathbb{1}$
- $\forall X \in \mathcal{D}_{\text{sg}}(R)$ $\text{End}(X)$ is symmetric since $\text{Hom}(X, X) \cong \text{Hom}(X, X[2]) \cong D\text{Hom}(X, X)$

$\text{Spec } R$: cDV w/ isolated singularity
 $\exists p: X \rightarrow \text{Spec } R$ crepant resolution



Thm (Donovan - Weemyss 2016)

$\exists \Delta(p)$: basic fin.dim. algebra that "controls" the NC deformations of $\mathcal{O}_{C_i}(-1) \in \text{coh } X$

$\Delta(p)$ is the contraction algebra of p .

e.g. $R = \mathbb{C}[u, v, x, y]/(uv - xy)$

$$\begin{array}{ccc} X & & Y \\ & p \searrow & \swarrow q \\ & \text{Spec } R & \end{array}$$

$\Delta(p) \cong \Delta(q) \cong \mathbb{C}$
misleadingly simple...

Rank $\Delta(p)$ recovers all known numerical invariants associated with $p: X \rightarrow \text{Spec } R$

The Donovan - Weemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1, 2$)

$$R_1 \cong R_2 \Leftrightarrow \mathcal{D}^b(\text{mod } \Delta(p_1)) \cong \mathcal{D}^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Weemyss (2018) + Dugas (2015)

(\Leftarrow) Reid (1983) in type A

Recall $\mathcal{D}_{\text{sg}}(R) := \mathcal{D}^b(\text{mod } R)/\mathcal{K}^b(\text{proj } R)$

is the singularity cat of R

- Hom-finite & Krull-Schmidt
- 2-Calabi-Yau $\text{Hom}(Y, X[2]) \cong D\text{Hom}(X, Y)$
- $[2] \cong \mathbb{1}$
- $\forall X \in \mathcal{D}_{\text{sg}}(R)$ $\text{End}(X)$ is symmetric since $\text{Hom}(X, X) \cong \text{Hom}(X, X[2]) \cong D\text{Hom}(X, X)$

Def (Iyama - Yoshino 2008) $T \in \mathcal{D}_{\text{sg}}(R)$: basic

$T \in \mathcal{D}_{\text{sg}}(R)$ is $2\mathbb{Z}$ -cluster tilting object if
 $\text{add } T = \{X \in \mathcal{D}_{\text{sg}}(R) \mid \text{Hom}(T, X[1]) = 0\}$
& $T \cong T[2]$

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1,2$)

$$R_1 \cong R_2 \Leftrightarrow D^b(\text{mod } \Delta(p_1)) \cong D^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Wemyss (2018) + Dugas (2015)

(\Leftarrow) Reid (1983) in type A

Recall $D_{sg}(R) := D^b(\text{mod } R) / K^b(\text{proj } R)$

is the singularity cat of R

- Hom-linite & Krull-Schmidt
- 2-Calabi-Yau $\text{Hom}(Y, X[2]) \cong D\text{Hom}(X, Y)$
- $[2] \cong \mathbb{1}$
- $\forall X \in D_{sg}(R)$ $\text{End}(X)$ is symmetric since
 $\text{Hom}(X, X) \cong \text{Hom}(X, X[2]) \cong D\text{Hom}(X, X)$

Def (Iyama-Yoshino 2008) $T \in D_{sg}(R)$: basic

$T \in D_{sg}(R)$ is 2~~Z~~-cluster tilting object if

$$\text{add } T = \{ X \in D_{sg}(R) \mid \text{Hom}(T, X[1]) = 0 \}$$

$$\& T \cong T[2]$$

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1,2$)

$$R_1 \cong R_2 \Leftrightarrow D^b(\text{mod } \Delta(p_1)) \cong D^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Wemyss (2018) + Dugas (2015)

(\Leftarrow) Reid (1983) in type A

Recall $D_{sg}(R) := D^b(\text{mod } R) / K^b(\text{proj } R)$

is the singularity cat of R

- Hom-linite & Krull-Schmidt
- 2-Calabi-Yau $\text{Hom}(Y, X[2]) \cong D\text{Hom}(X, Y)$
- $[2] \cong \mathbb{1}$
- $\forall X \in D_{sg}(R)$ $\text{End}(X)$ is symmetric since
 $\text{Hom}(X, X) \cong \text{Hom}(X, X[2]) \cong D\text{Hom}(X, X)$

Def (Iyama-Yoshino 2008) $T \in D_{sg}(R)$: basic

$T \in D_{sg}(R)$ is 2-Calabi-Yau cluster tilting object if

$$\text{add } T = \{ X \in D_{sg}(R) \mid \text{Hom}(T, X[1]) = 0 \}$$

$$\& T \cong T[2]$$

Prop (Iyama-Yoshino 2008)

$T \in D_{sg}(R)$: 2-Calabi-Yau cluster tilting

$$\Rightarrow \forall X \in D_{sg}(R) \exists \underbrace{T^{-1} \rightarrow T^0}_{\text{add } T} \rightarrow X \rightarrow T[1]$$

$$(\Rightarrow \text{thick}(T) = D_{sg}(R))$$

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1,2$)

$$R_1 \cong R_2 \Leftrightarrow \mathcal{D}^b(\text{mod } \Delta(p_1)) \cong \mathcal{D}^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Wemyss (2018) + Dugas (2015)

(\Leftarrow) Reid (1983) in type A

Recall $\mathcal{D}_{\text{sg}}(R) := \mathcal{D}^b(\text{mod } R) / \mathcal{K}^b(\text{proj } R)$

is the singularity cat of R

- Hom-finite & Krull-Schmidt
- 2-Calabi-Yau $\text{Hom}(Y, X[2]) \cong D\text{Hom}(X, Y)$
- $[2] \cong \mathbb{1}$
- $\forall X \in \mathcal{D}_{\text{sg}}(R)$ $\text{End}(X)$ is symmetric since
 $\text{Hom}(X, X) \cong \text{Hom}(X, X[2]) \cong D\text{Hom}(X, X)$

Def (Iyama-Yoshino 2008) $T \in \mathcal{D}_{\text{sg}}(R)$: basic

$T \in \mathcal{D}_{\text{sg}}(R)$ is 2~~Z~~-cluster tilting object if

$$\text{add } T = \{ X \in \mathcal{D}_{\text{sg}}(R) \mid \text{Hom}(T, X[1]) = 0 \}$$

$$\& T \cong T[2]$$

Prop (Iyama-Yoshino 2008)

$T \in \mathcal{D}_{\text{sg}}(R)$: 2~~Z~~-cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{\text{sg}}(R) \exists \underbrace{T^{-1} \rightarrow T^0 \rightarrow X \rightarrow T[1]}_{\text{add } T}$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{\text{sg}}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \xrightarrow{\quad \downarrow \quad} \left\{ T \in \mathcal{D}_{\text{sg}}(R) \mid \text{basic 2~~Z~~-cluster tilting} \right\} / \cong \xrightarrow{\quad T(p) \quad} \mathcal{P}$$

$$\text{Moreover, } \text{End}(T(p)) \cong \Delta(p).$$

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1,2$)

$$R_1 \cong R_2 \Leftrightarrow \mathcal{D}^b(\text{mod } \Delta(p_1)) \cong \mathcal{D}^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Wemyss (2018) + Dugas (2015)

(\Leftarrow) Reid (1983) in type A

Recall $\mathcal{D}_{\text{sg}}(R) := \mathcal{D}^b(\text{mod } R) / \mathcal{K}^b(\text{proj } R)$

is the singularity cat of R

- Hom-finite & Krull-Schmidt
- 2-Calabi-Yau $\text{Hom}(Y, X[2]) \cong D\text{Hom}(X, Y)$
- $[2] \cong \mathbb{1}$
- $\forall X \in \mathcal{D}_{\text{sg}}(R)$ $\text{End}(X)$ is symmetric since $\text{Hom}(X, X) \cong \text{Hom}(X, X[2]) \cong D\text{Hom}(X, X)$

Def (Iyama-Yoshino 2008) $T \in \mathcal{D}_{\text{sg}}(R)$: basic

$T \in \mathcal{D}_{\text{sg}}(R)$ is 2 \mathbb{Z} -cluster tilting object if

$$\text{add } T = \{ X \in \mathcal{D}_{\text{sg}}(R) \mid \text{Hom}(T, X[1]) = 0 \}$$

$$\& T \cong T[2]$$

Prop (Iyama-Yoshino 2008)

$T \in \mathcal{D}_{\text{sg}}(R)$: 2 \mathbb{Z} -cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{\text{sg}}(R) \exists \underbrace{T^{-1} \rightarrow T^0 \rightarrow X \rightarrow T[1]}_{\text{add } T}$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{\text{sg}}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \xrightarrow{\quad p \quad} \left\{ T \in \mathcal{D}_{\text{sg}}(R) \mid \text{basic 2}\mathbb{Z}\text{-cluster tilting} \right\} / \cong T(p)$$

Moreover, $\text{End}(T(p)) \cong \Delta(p)$.

e.g. $R = \mathbb{C}[u, v, x, y]/(uv - xy)$

BMRRRT cluster cat.

$$\begin{array}{ccc} X & Y & \mathcal{D}_{\text{sg}}(R) \cong \mathcal{D}^b(\text{mod } \mathbb{C}) / \mathbb{C}[2] \\ p \downarrow & q \downarrow & T(p) \longmapsto \mathbb{C} \\ \text{Spec } R & & T(q) \longmapsto \mathbb{C}[1] \end{array}$$

The Donovan-Wemyss Conjecture (2016, 2020)

$p_i: X_i \rightarrow \text{Spec } R_i$ crepant res. of cDV's sing ($i=1,2$)

$$R_1 \cong R_2 \Leftrightarrow \mathcal{D}^b(\text{mod } \Delta(p_1)) \cong \mathcal{D}^b(\text{mod } \Delta(p_2))$$

(\Rightarrow) Wemyss (2018) + Dugas (2015)

(\Leftarrow) Reid (1983) in type A

Recall $\mathcal{D}_{sg}(R) := \mathcal{D}^b(\text{mod } R)/\mathcal{K}^b(\text{proj } R)$

is the singularity cat of R

- Hom-linite & Krull-Schmidt
- 2-Calabi-Yau $\text{Hom}(Y, X[2]) \cong D\text{Hom}(X, Y)$
- $[2] \cong \mathbb{1}$
- $\forall X \in \mathcal{D}_{sg}(R)$ $\text{End}(X)$ is symmetric since $\text{Hom}(X, X) \cong \text{Hom}(X, X[2]) \cong D\text{Hom}(X, X)$

Def (Iyama-Yoshino 2008) $T \in \mathcal{D}_{sg}(R)$: basic

$T \in \mathcal{D}_{sg}(R)$ is 2 \mathbb{Z} -cluster tilting object if

$$\text{add } T = \{ X \in \mathcal{D}_{sg}(R) \mid \text{Hom}(T, X[1]) = 0 \}$$

$$\& T \cong T[2]$$

Prop (Iyama-Yoshino 2008)

$T \in \mathcal{D}_{sg}(R)$: 2 \mathbb{Z} -cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{sg}(R) \exists \underbrace{T^{-1} \rightarrow T^0 \rightarrow X \rightarrow T[1]}_{\text{add } T}$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{sg}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \xrightarrow{\quad p \quad} \downarrow$$

$$\left\{ T \in \mathcal{D}_{sg}(R) \mid \text{basic 2 \mathbb{Z} -cluster tilting} \right\} / \cong \xrightarrow{\quad T(p) \quad} T(p)$$

Moreover, $\text{End}(T(p)) \cong \Delta(p)$.

e.g. $R = \mathbb{C}[u, v, x, y]/(uv - xy)$

BMRRRT cluster cat.

$$\begin{array}{ccc} X & Y & \mathcal{D}_{sg}(R) \cong \mathcal{D}^b(\text{mod } \mathbb{C})/\mathbb{C}[2] \xleftarrow{\quad} \\ p \downarrow & \swarrow q & T(p) \longmapsto \mathbb{C} \\ \text{Spec } R & & T(q) \longmapsto \mathbb{C}[1] \end{array}$$

$\mathbb{C}[v^\pm]$, $1+1=-2$: dg algebra with trivial differential

$$\rightsquigarrow \mathcal{D}_{sg}(R) \cong \mathcal{D}^c(\mathbb{C}[v^\pm])$$

\uparrow compact/perfect der. cat.

Prop (Iyama - Yoshino 2008)

$T \in \mathcal{D}_{sg}(R)$: 2 \mathbb{Z} -cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{sg}(R) \quad \exists \underbrace{T^{-1} \rightarrow T^0}_{\text{add } T} \rightarrow X \rightarrow T[1]$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{sg}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \xrightarrow{\quad p \quad} \left\{ T \in \mathcal{D}_{sg}(R) \mid \text{basic 2}\mathbb{Z}\text{-cluster tilting} \right\} / \cong \cong T(p)$$

\downarrow \downarrow

Moreover, $\text{End}(T(p)) \cong \Lambda(p)$.

e.g. $R = \mathbb{C}[u, v, x, y] / (uv - xy)$

BMRRT cluster cat.

$$\begin{array}{ccc} X & Y & \mathcal{D}_{sg}(R) \cong \mathcal{D}^b(\text{mod } \mathbb{C}) / \nu[-2] \\ p \searrow & \swarrow q & T(p) \longmapsto \mathbb{C} \\ \text{Spec } R & & T(q) \longmapsto \mathbb{C}[1] \end{array}$$

$\mathbb{C}[v^\pm]$, $1 \circ 1 = -2$: dg algebra with trivial differential

$$\rightsquigarrow \mathcal{D}_{sg}(R) \cong \mathcal{D}^c(\mathbb{C}[v^\pm])$$

↑ compact / perfect der. cat.

Prop (Iyama - Yoshino 2008)

$T \in \mathcal{D}_{sg}(R)$: 2~~U~~-cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{sg}(R) \exists \underbrace{T^{-1} \rightarrow T^0 \rightarrow X \rightarrow T[1]}_{\text{add } T}$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{sg}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\begin{array}{c} \left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \\ \downarrow \\ \left\{ T \in \mathcal{D}_{sg}(R) \mid \text{basic 2}\cancel{U}\text{-cluster tilting} \right\} / \cong T(p) \end{array}$$

$\overset{p}{\downarrow}$

Moreover, $\text{End}(T(p)) \cong \Lambda(p)$.

e.g. $R = \mathbb{C}[u, v, x, y]/(uv - xy)$

BMRRT cluster cat.

$$\begin{array}{ccc} X & Y & \mathcal{D}_{sg}(R) \cong \mathcal{D}^b(\text{mod } \mathbb{C})/\mathbb{C}[-2] \\ p \searrow & \swarrow q & T(p) \longmapsto \mathbb{C} \\ \text{Spec } R & & T(q) \longmapsto \mathbb{C}[1] \end{array}$$

$\mathbb{C}[v^\pm]$, $1 \circ 1 = -2$: dg algebra with trivial differential

$$\rightsquigarrow \mathcal{D}_{sg}(R) \cong \mathcal{D}^c(\mathbb{C}[v^\pm])$$

↑ compact / perfect der. cat.

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \cong \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Prop (Iyama - Yoshino 2008)

$T \in \mathcal{D}_{sg}(R)$: 2~~U~~-cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{sg}(R) \exists \underbrace{T^{-1} \rightarrow T^0 \rightarrow X \rightarrow T[1]}_{\text{add } T}$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{sg}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \xrightarrow{\quad p \quad} \left\{ T \in \mathcal{D}_{sg}(R) \mid \text{basic 2}\cancel{U}\text{-cluster tilting} \right\} / \cong \xrightarrow{\quad T(p) \quad} \mathcal{T}(p)$$

Moreover, $\text{End}(T(p)) \cong \Lambda(p)$.

e.g. $R = \mathbb{C}[[u, v, x, y]] / (uv - xy)$

BMRRT cluster cat.

$$\begin{array}{ccc} X & Y & \mathcal{D}_{sg}(R) \cong \mathcal{D}^b(\text{mod } \mathbb{C}) / v[-2] \\ p \downarrow & \swarrow q & T(p) \longmapsto \mathbb{C} \\ \text{Spec } R & & T(q) \longmapsto \mathbb{C}[1] \end{array}$$

$\mathbb{C}[v^\pm]$, $1 \circ 1 = -2$: dg algebra with trivial differential

$$\rightsquigarrow \mathcal{D}_{sg}(R) \cong \mathcal{D}^c(\mathbb{C}[v^\pm])$$

↑ compact / perfect der. cat.

Thm (August 2020)

$p: Y \rightarrow \text{Spec } R$ as above & Λ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Lambda) \cong \mathcal{D}^b(\text{mod } \Lambda(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Lambda \cong \Lambda(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i, i=1, 2$ as in DW cong.

$$\mathcal{D}^b(\text{mod } \Lambda(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Lambda(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Lambda(p_2) \cong \Lambda(q)$

Prop (Iyama - Yoshino 2008)

$T \in \mathcal{D}_{sg}(R)$: 2~~U~~-cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{sg}(R) \exists \underbrace{T^{-1} \rightarrow T^0 \rightarrow X \rightarrow T[1]}_{\text{add } T}$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{sg}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \xrightarrow{\quad p \quad} \left\{ T \in \mathcal{D}_{sg}(R) \mid \text{basic 2}\cancel{U}\text{-cluster tilting} \right\} / \cong \cong T(p)$$

Moreover, $\text{End}(T(p)) \cong \Lambda(p)$.

e.g. $R = \mathbb{C}[[u, v, x, y]] / (uv - xy)$

BMRRT cluster cat.

$$\begin{array}{ccc} X & Y & \mathcal{D}_{sg}(R) \cong \mathcal{D}^b(\text{mod } \mathbb{C}) / v[-2] \\ p \downarrow & \swarrow q & T(p) \longmapsto \mathbb{C} \\ \text{Spec } R & & T(q) \longmapsto \mathbb{C}[1] \end{array}$$

$\mathbb{C}[v^\pm]$, $1 \circ 1 = -2$: dg algebra with trivial differential

$$\rightsquigarrow \mathcal{D}_{sg}(R) \cong \mathcal{D}^c(\mathbb{C}[v^\pm])$$

↑ compact/perfect der. cat.

Thm (August 2020)

$p: Y \rightarrow \text{Spec } R$ as above & Λ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Lambda) \cong \mathcal{D}^b(\text{mod } \Lambda(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Lambda \cong \Lambda(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i, i=1, 2$ as in DW cong.

$$\mathcal{D}^b(\text{mod } \Lambda(p_1)) \xleftarrow{\sim} \mathcal{D}^b(\text{mod } \Lambda(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Lambda(p_2) \cong \Lambda(q)$

We may assume that $\Lambda(p_1) \cong \Lambda(p_2)$

Prop (Iyama - Yoshino 2008)

$T \in \mathcal{D}_{sg}(R)$: 2~~U~~-cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{sg}(R) \exists \underbrace{T^{-1} \rightarrow T^0 \rightarrow X \rightarrow T[1]}_{\text{add } T}$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{sg}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \xrightarrow{\quad p \quad} \left\{ T \in \mathcal{D}_{sg}(R) \mid \text{basic 2}\cancel{U}\text{-cluster tilting} \right\} / \cong \xrightarrow{\quad T(p) \quad} \mathcal{D}_{sg}(R)$$

Moreover, $\text{End}(T(p)) \cong \Lambda(p)$.

e.g. $R = \mathbb{C}[[u, v, x, y]] / (uv - xy)$

BMRRT cluster cat.

$$\begin{array}{ccc} X & Y & \mathcal{D}_{sg}(R) \cong \mathcal{D}^b(\text{mod } \mathbb{C}) / v[-2] \\ p \downarrow & \swarrow q & T(p) \longmapsto \mathbb{C} \\ \text{Spec } R & & T(q) \longmapsto \mathbb{C}[1] \end{array}$$

$\mathbb{C}[v^\pm]$, $1 \circ 1 = -2$: dg algebra with trivial differential

$$\rightsquigarrow \mathcal{D}_{sg}(R) \cong \mathcal{D}^c(\mathbb{C}[v^\pm])$$

↑ compact/perfect der. cat.

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Λ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Lambda) \cong \mathcal{D}^b(\text{mod } \Lambda(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Lambda \cong \Lambda(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i, i=1, 2$ as in DW cong.

$$\mathcal{D}^b(\text{mod } \Lambda(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Lambda(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Lambda(p_2) \cong \Lambda(q)$

We may assume that $\Lambda(p_1) \cong \Lambda(p_2)$

Recall $\mathcal{D}_{sg}(R) := \mathcal{D}^b(\text{mod } R)_{dg} / K^b(\text{proj } R)_{dg}$

is the dg singularity cat of R

Prop (Iyama - Yoshino 2008)

$T \in \mathcal{D}_{sg}(R)$: 2~~U~~-cluster tilting

$$\Rightarrow \forall X \in \mathcal{D}_{sg}(R) \exists \underbrace{T^{-1} \rightarrow T^0}_{\text{add } T} \rightarrow X \rightarrow T[1]$$

$$(\Rightarrow \text{thick}(T) = \mathcal{D}_{sg}(R))$$

Thm (Wemyss 2018) There is a bijection

$$\left\{ X \rightarrow \text{Spec } R \mid \text{crepant resolution} \right\} / \cong \xrightarrow{\quad p \quad} \left\{ T \in \mathcal{D}_{sg}(R) \mid \text{basic 2}\cancel{U}\text{-cluster tilting} \right\} / \cong \xrightarrow{\quad T(p) \quad}$$

Moreover, $\text{End}(T(p)) \cong \Lambda(p)$.

e.g. $R = \mathbb{C}[[u, v, x, y]] / (uv - xy)$

BMRRT cluster cat.

$$\begin{array}{ccc} X & Y & \mathcal{D}_{sg}(R) \cong \mathcal{D}^b(\text{mod } \mathbb{C}) / v[-2] \\ p \downarrow & q \downarrow & T(p) \longmapsto \mathbb{C} \\ \text{Spec } R & & T(q) \longmapsto \mathbb{C}[1] \end{array}$$

$\mathbb{C}[v^\pm]$, $1+1=-2$: dg algebra with trivial differential

$$\rightsquigarrow \mathcal{D}_{sg}(R) \cong \mathcal{D}^c(\mathbb{C}[v^\pm])$$

↑ compact/perfect der. cat.

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \cong \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i, i=1, 2$ as in DW cong.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(q)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $\mathcal{D}_{sg}(R) := \mathcal{D}^b(\text{mod } R)_{dg} / K^b(\text{proj } R)_{dg}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$\text{HH}^*(\mathbb{A}) := H^*(\text{IRHom}_{dg}(\mathbb{A}, \mathbb{A}))$ is the Hochschild cohomology of \mathbb{A}

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \simeq \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i$, $i = 1, 2$ as in DW Conj.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(q)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $D_{\text{sg}}(R) := \mathcal{D}^b_{\text{dg}}(\text{mod } R)_{\text{dg}} / K^b(\text{proj } R)_{\text{dg}}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$HH^*(\mathbb{A}) := H^*(\mathbf{R}\text{Hom}_{\mathbb{A}^e}(\mathbb{A}, \mathbb{A}))$ is the

Hochschild cohomology of \mathbb{A}

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \simeq \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i$, $i = 1, 2$ as in DW Conj.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(q)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $D_{sg}(R)_{dg} := \mathcal{D}^b(\text{mod } R)_{dg} / K^b(\text{proj } R)_{dg}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$HH^*(\mathbb{A}) := H^*(\mathbf{R}\text{Hom}_{\mathbb{A}^{op}}(\mathbb{A}, \mathbb{A}))$ is the

Hochschild cohomology of \mathbb{A}

Thm (Hua-Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$$HH^0(D_{sg}(R)_{dg}) \cong T_f : \underline{\text{Tyurina algebra}} \text{ of } f$$

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \simeq \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i$, $i = 1, 2$ as in DW Conj.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists f: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(f)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $D_{sg}(R)_{dg} := \mathcal{D}^b(\text{mod } R)_{dg} / K^b(\text{proj } R)_{dg}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$HH^*(\mathbb{A}) := H^*(\mathbf{R}\text{Hom}_{\mathbb{A}^{op}}(\mathbb{A}, \mathbb{A}))$ is the

Hochschild cohomology of \mathbb{A}

Thm (Hua - Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$$HH^0(D_{sg}(R)_{dg}) \cong T_f : \underline{\text{Tyurina algebra}} \text{ of } f$$

Thm (Mather - You 1982)

$T_f + \dim R$ determine R up to isomorphism

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \simeq \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$ as in DW Conj.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(q)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $D_{sg}(R)_{dg} := \mathcal{D}^b(\text{mod } R)_{dg} / K^b(\text{proj } R)_{dg}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$HH^*(\mathbb{A}) := H^*(\mathbf{R}\text{Hom}_{\mathbb{A}^{op}}(\mathbb{A}, \mathbb{A}))$ is the

Hochschild cohomology of \mathbb{A}

Thm (Hua-Keller 2018) $R \cong \mathbb{C}[x,y,z,t]/(f)$

$$HH^0(D_{sg}(R)_{dg}) \cong T_f : \underline{\text{Tyurina algebra}} \text{ of } f$$

Thm (Mather-Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \simeq \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$ as in DW Conj.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(q)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $\mathcal{D}_{\text{sg}}(R)_{\text{dg}} := \mathcal{D}^b(\text{mod } R)_{\text{dg}} / K^b(\text{proj } R)_{\text{dg}}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$HH^*(\mathbb{A}) := H^*(\mathbf{R}\text{Hom}_{\mathbb{A}^e}(\mathbb{A}, \mathbb{A}))$ is the

Hochschild cohomology of \mathbb{A}

Thm (Hua-Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$$HH^0(\mathcal{D}_{\text{sg}}(R)_{\text{dg}}) \cong T_f : \underline{\text{Tyurina algebra}} \text{ of } f$$

Thm (Mather-Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$$T(p_1) \in \mathcal{D}_{\text{sg}}(R_1)_{\text{dg}} \xrightarrow{\text{ZK-CF}} \Delta(p_1) := R \text{End}(T(p_1))$$

$$T(p_2) \in \mathcal{D}_{\text{sg}}(R_2)_{\text{dg}} \xrightarrow{\text{ZK-CF}} \Delta(p_2) := R \text{End}(T(p_2))$$

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \simeq \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$ as in DW Conj.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(q)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $\mathcal{D}_{\text{sg}}(R) := \mathcal{D}^b(\text{mod } R)_{\text{dg}} / K^b(\text{proj } R)_{\text{dg}}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$HH^*(\mathbb{A}) := H^*(\mathbf{R}\text{Hom}_{\mathbb{A}^e}(\mathbb{A}, \mathbb{A}))$ is the

Hochschild cohomology of \mathbb{A}

Thm (Hua-Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$$HH^0(\mathcal{D}_{\text{sg}}(R)_{\text{dg}}) \cong T_f : \underline{\text{Tyurina algebra}} \text{ of } f$$

Thm (Mather-Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$$T(p_1) \in \mathcal{D}_{\text{sg}}(R_1)_{\text{dg}} \xrightarrow{\text{ZL-CT}} \Delta(p_1) := R \text{End}(T(p_1))$$

$$T(p_2) \in \mathcal{D}_{\text{sg}}(R_2)_{\text{dg}} \xrightarrow{\text{ZL-CT}} \Delta(p_2) := R \text{End}(T(p_2))$$

$$H^*(\Delta(p_1)) \cong H^*(\Delta(p_2)) \cong \Delta[z^\pm], |z| = -2$$

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \simeq \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$ as in DW Conj.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(q)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $\mathcal{D}_{\text{sg}}(R) := \mathcal{D}^b(\text{mod } R)_{\text{dg}} / K^b(\text{proj } R)_{\text{dg}}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$HH^*(\mathbb{A}) := H^*(\text{IHom}_{\mathbb{A}^e}(\mathbb{A}, \mathbb{A}))$ is the

Hochschild cohomology of \mathbb{A}

Thm (Hua-Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$$HH^0(\mathcal{D}_{\text{sg}}(R)_{\text{dg}}) \cong T_f : \underline{\text{Tyurina algebra}} \text{ of } f$$

Thm (Mather-Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$$T(p_1) \in \mathcal{D}_{\text{sg}}(R_1)_{\text{dg}} \xrightarrow{\text{ZL-CT}} \Delta(p_1) := R \text{End}(T(p_1))$$

$$T(p_2) \in \mathcal{D}_{\text{sg}}(R_2)_{\text{dg}} \xrightarrow{\text{ZL-CT}} \Delta(p_2) := R \text{End}(T(p_2))$$

$$H^*(\Delta(p_1)) \cong H^*(\Delta(p_2)) \cong \Delta[z^\pm], |z| = -2$$

$$(\text{Keller 1994}) \quad \mathcal{D}^c(\Delta(p_1))_{\text{dg}} \xrightarrow{\sim} \mathcal{D}_{\text{sg}}(R_1)_{\text{dg}}$$

$$\mathcal{D}^c(\Delta(p_2))_{\text{dg}} \xrightarrow{\sim} \mathcal{D}_{\text{sg}}(R_2)_{\text{dg}}$$

Thm (August 2020)

$p: X \rightarrow \text{Spec } R$ as above & Δ : basic fin. dim. alg.

$$\mathcal{D}^b(\text{mod } \Delta) \simeq \mathcal{D}^b(\text{mod } \Delta(p))$$



$\exists q: Y \rightarrow \text{Spec } R$ crepant resolution s.t. $\Delta \cong \Delta(q)$

Corollary $p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$ as in DW Conj.

$$\mathcal{D}^b(\text{mod } \Delta(p_1)) \xleftrightarrow{\sim} \mathcal{D}^b(\text{mod } \Delta(p_2))$$



$\exists q: Y \rightarrow \text{Spec } R_1$ s.t. $\Delta(p_2) \cong \Delta(q)$

We may assume that $\Delta(p_1) \cong \Delta(p_2)$

Recall $\mathcal{D}_{\text{sg}}(R) := \mathcal{D}^b(\text{mod } R)_{\text{dg}} / K^b(\text{proj } R)_{\text{dg}}$

is the dg singularity cat of R

\mathbb{A} : ess. small dg category

$HH^*(\mathbb{A}) := H^*(\mathbb{I}R\text{Hom}_{\mathbb{A}^{op}}(\mathbb{A}, \mathbb{A}))$ is the Hochschild cohomology of \mathbb{A}

Thm (Hua - Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$$HH^0(\mathcal{D}_{\text{sg}}(R)_{\text{dg}}) \cong T_f : \underline{\text{Tyurina algebra}} \text{ of } f$$

Thm (Mather - You 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1,2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$$T(p_1) \in \mathcal{D}_{\text{sg}}(R_1)_{\text{dg}} \xrightarrow{\text{ZL-CT}} \Delta(p_1) := R\text{End}(T(p_1))$$

$$T(p_2) \in \mathcal{D}_{\text{sg}}(R_2)_{\text{dg}} \xrightarrow{\text{ZL-CT}} \Delta(p_2) := R\text{End}(T(p_2))$$

$$H^*(\Delta(p_1)) \cong H^*(\Delta(p_2)) \cong \Delta[z^\pm], |z| = -2$$

$$(\text{Keller 1994}) \quad \mathcal{D}^c(\Delta(p_1))_{\text{dg}} \xrightarrow{\sim} \mathcal{D}_{\text{sg}}(R_1)_{\text{dg}}$$

$$\mathcal{D}^c(\Delta(p_2))_{\text{dg}} \xrightarrow{\sim} \mathcal{D}_{\text{sg}}(R_2)_{\text{dg}}$$

$$\mathcal{D}^c(\Delta(p_1)_{\text{dg}} \simeq \mathcal{D}^c(\Delta(p_2)_{\text{dg}}) : \text{quasi-eq} \implies \text{DW Conj.}$$

Thm (Hua - Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$HH^0(\mathcal{D}\text{sg}(R)_{dg}) \cong T_f$: Tyurina algebra of f

Thm (Mather - You 1982)

$T_f + \dim R$ determine R up to isomorphism

→ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1, 2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$T(p_1) \in \mathcal{D}\text{sg}(R_1)_{dg}^{Z\text{-CT}} \rightsquigarrow \mathbb{L}\Delta(p_1) := R\text{End}(T(p_1))$

$T(p_2) \in \mathcal{D}\text{sg}(R_2)_{dg}^{Z\text{-CT}} \rightsquigarrow \mathbb{L}\Delta(p_2) := R\text{End}(T(p_2))$

$H^*(\mathbb{L}\Delta(p_1)) \cong H^*(\mathbb{L}\Delta(p_2)) \cong \Delta[z^\pm], |z| = -2$

(Keller 1994) $\mathcal{D}^c(\mathbb{L}\Delta(p_1))_{dg} \xrightarrow{\sim} \mathcal{D}\text{sg}(R_1)_{dg}$

$\mathcal{D}^c(\mathbb{L}\Delta(p_2))_{dg} \xrightarrow{\sim} \mathcal{D}\text{sg}(R_2)_{dg}$

$\mathcal{D}^c(\mathbb{L}\Delta(p_1)_{dg} \cong \mathcal{D}^c(\mathbb{L}\Delta(p_2)_{dg})$: quasi-eq \Rightarrow DW Conj.

Thm (Hua - Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$H^0(\mathcal{D}_{sg}(R)_{dg}) \cong T_f$: Tyurina algebra of f

Thm (Mather - Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1, 2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$T(p_1) \in \mathcal{D}_{sg}(R_1)_{dg}^{\text{2k-CF}} \rightsquigarrow \Delta(p_1) := R\text{End}(T(p_1))$

$T(p_2) \in \mathcal{D}_{sg}(R_2)_{dg}^{\text{2k-CF}} \rightsquigarrow \Delta(p_2) := R\text{End}(T(p_2))$

$H^*(\Delta(p_1)) \cong H^*(\Delta(p_2)) \cong \Delta[z^\pm]$, $|z| = -2$

(Keller 1994) $\mathcal{D}^c(\Delta(p_1))_{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_1)_{dg}$

$\mathcal{D}^c(\Delta(p_2))_{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_2)_{dg}$

Thm (I-Muro 2022)

R : isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ ^{crepant res.}

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique (Karoubian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is 2k-cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm]$, $|z| = -2$

$\mathcal{D}^c(\Delta(p_1))_{dg} \cong \mathcal{D}^c(\Delta(p_2))_{dg}$: quasi-eq \Rightarrow DW Conj.

Thm (Hua - Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$H^0(\mathcal{D}_{sg}(R)_{dg}) \cong T_f$: Tyurina algebra of f

Thm (Mather - Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

→ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1, 2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$T(p_1) \in \mathcal{D}_{sg}(R_1)_{dg}^{\text{2k-CF}} \rightsquigarrow \Delta(p_1) := R\text{End}(T(p_1))$

$T(p_2) \in \mathcal{D}_{sg}(R_2)_{dg}^{\text{2k-CF}} \rightsquigarrow \Delta(p_2) := R\text{End}(T(p_2))$

$H^*(\Delta(p_1)) \cong H^*(\Delta(p_2)) \cong \Lambda[z^\pm]$, $|z| = -2$

(Keller 1994) $\mathcal{D}^c(\Delta(p_1))_{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_1)_{dg}$

$\mathcal{D}^c(\Delta(p_2))_{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_2)_{dg}$

Thm (I-Muro 2022)

R : isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ crepant res.

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique (Karoubian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is 2k-cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm]$, $|z| = -2$

Strategy Prove $R\text{End}(T) = \Delta(p)$

for $T \in \mathcal{D}$ satisfying (1) & (2)

$\mathcal{D}^c(\Delta(p_1))_{dg} \cong \mathcal{D}^c(\Delta(p_2))_{dg}$: quasi-eq \Rightarrow DW Conj.

Thm (Hua - Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$H^0(\mathcal{D}_{sg}(R)_{dg}) \cong T_f$: Tyurina algebra of f

Thm (Mather - Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1, 2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$T(p_1) \in \mathcal{D}_{sg}(R_1)_{dg}^{\text{2k-CF}} \rightsquigarrow \Delta(p_1) := R\text{End}(T(p_1))$

$T(p_2) \in \mathcal{D}_{sg}(R_2)_{dg}^{\text{2k-CF}} \rightsquigarrow \Delta(p_2) := R\text{End}(T(p_2))$

$H^*(\Delta(p_1)) \cong H^*(\Delta(p_2)) \cong \Delta[z^\pm], |z| = -2$

(Keller 1994) $D^c(\Delta(p_1))_{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_1)_{dg}$

$D^c(\Delta(p_2))_{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_2)_{dg}$

$D^c(\Delta(p_1)_{dg} \cong D^c(\Delta(p_2)_{dg} : \text{quasi-eq} \Rightarrow \text{DW Conj.}$

Thm (I-Muro 2022)

R : isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ ^{crepant}
_{res.}

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique
(Karoubian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is 2k-cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm], |z| = -2$

Strategy Prove $R\text{End}(T) = \Delta(p)$

for $T \in \mathcal{A}$ satisfying (1) & (2)

Sketch of proof $\Delta = \Delta(p)$ & $\Delta = \Delta(p)$

$\rightsquigarrow H^*(\Delta) \cong \Delta[z^\pm], |z| = -2$.

Thm (Hua - Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$H^0(\mathcal{D}_{sg}(R)_{dg}) \cong T_f$: Tyurina algebra of f

Thm (Mather - Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1, 2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$T(p_1) \in \mathcal{D}_{sg}(R_1)_{dg}^{\text{2k-CF}} \rightsquigarrow \Delta(p_1) := R\text{End}(T(p_1))$

$T(p_2) \in \mathcal{D}_{sg}(R_2)_{dg}^{\text{2k-CF}} \rightsquigarrow \Delta(p_2) := R\text{End}(T(p_2))$

$H^*(\Delta(p_1)) \cong H^*(\Delta(p_2)) \cong \Delta[z^\pm], |z| = -2$

(Keller 1994) $\mathcal{D}^c(\Delta(p_1))_{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_1)_{dg}$

$\mathcal{D}^c(\Delta(p_2))_{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_2)_{dg}$

$\mathcal{D}^c(\Delta(p_1)_{dg} \cong \mathcal{D}^c(\Delta(p_2)_{dg})$: quasi-eq \Rightarrow DW Conj.

Thm (J-Muro 2022)

R : isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ crepant res.

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique (Karoubian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is 2k-cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm], |z| = -2$

Strategy Prove $R\text{End}(T) = \Delta(p)$

for $T \in \mathcal{A}$ satisfying (1) & (2)

Sketch of proof $\Delta = \Delta(p)$ & $\Delta = \Delta(p)$

$\rightsquigarrow H^*(\Delta) \cong \Delta[z^\pm], |z| = -2$.

(Kadeishvili 1982)

Δ induces a minimal A_∞ -algebra structure on $\Delta[z^\pm]$ s.t. $\Delta \cong \Delta[z^\pm]$ as A_∞ -algebras

Thm (Hua - Keller 2018) $R \cong \mathbb{C}[x, y, z, t]/(f)$

$H^0(\mathcal{D}_{sg}(R)^{dg}) \cong T_f$: Tyurina algebra of f

Thm (Mather - Yau 1982)

$T_f + \dim R$ determine R up to isomorphism

~ Putting everything together ...

$p_i: X_i \rightarrow \text{Spec } R_i$, $i=1, 2$, as in the DW Conjecture

Suppose that $\Delta(p_1) \cong \Delta(p_2) =: \Delta$

$T(p_1) \in \mathcal{D}_{sg}(R_1)^{dg}$ ~~~ $\Delta(p_1) := R\text{End}(T(p_1))$

$T(p_2) \in \mathcal{D}_{sg}(R_2)^{dg}$ ~~~ $\Delta(p_2) := R\text{End}(T(p_2))$

$H^*(\Delta(p_1)) \cong H^*(\Delta(p_2)) \cong \Delta[z^\pm]$, $|z| = -2$

(Keller 1994) $\mathcal{D}^c(\Delta(p_1))^{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_1)^{dg}$

$\mathcal{D}^c(\Delta(p_2))^{dg} \xrightarrow{\sim} \mathcal{D}_{sg}(R_2)^{dg}$

$\mathcal{D}^c(\Delta(p_1))^{dg} \cong \mathcal{D}^c(\Delta(p_2))^{dg}$: quasi-eq \Rightarrow DW Conj.

Thm (J-Muro 2022)

R : isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ crepant res.

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)^{dg}$ is the unique (Karoubian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)^{dg}$ is $2\mathbb{Z}$ -cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm]$, $|z| = -2$

Strategy Prove $R\text{End}(T) = \Delta(p)$

for $T \in \mathcal{A}$ satisfying (1) & (2)

Sketch of proof $\Delta = \Delta(p)$ & $\Delta = \Delta(p)$

~ $H^*(\Delta) \cong \Delta[z^\pm]$, $|z| = -2$.

(Kadeishvili 1982)

Δ induces a minimal A_∞ -algebra structure on $\Delta[z^\pm]$ s.t. $\Delta \cong \Delta[z^\pm]$ as A_∞ -algebras

$m_n: \Delta[z^\pm]^{\otimes n} \rightarrow \Delta[z^\pm]$ of degree $2-n$, $n \geq 2$

~ $\forall n \notin 2\mathbb{Z}$, $m_n = 0$

$\Delta = (\Delta[z^\pm], m_4, m_6, m_8, \dots)$

Thm (]-Muro 2022)

R : isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ ^{crepant}_{res.}

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique
(Kanobian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is $2\mathbb{Z}$ -cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm]$, $|z| = -2$

Strategy Prove $R\text{End}(T) \simeq \Delta(p)$

for $T \in \mathcal{A}$ satisfying (1) & (2)

Sketch of proof $\Delta = \Delta(p)$ & $\Delta = \Delta(p)$

$$\rightsquigarrow H^*(\Delta) \cong \Delta[z^\pm], |z| = -2.$$

(Kadeishvili 1982)

Δ induces a minimal A_∞ -algebra structure

on $\Delta[z^\pm]$ s.t. $\Delta \simeq \Delta[z^\pm]$ as A_∞ -algebras

$$m_n: \Delta[z^\pm]^{\otimes n} \rightarrow \Delta[z^\pm] \text{ of degree } 2-n, n \geq 2$$

$$\rightsquigarrow \forall n \notin 2\mathbb{Z}, m_n = 0$$

$\Delta = (\Delta[z^\pm], m_4, m_6, m_8, \dots)$

Thm (]-Muro 2022)

R: isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ ^{crepant}_{res.}

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique
(Kanobian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is $2\mathbb{Z}$ -cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm]$, $|z| = -2$

Strategy Prove $R\text{End}(T) \simeq \Delta(p)$

for $T \in \mathcal{A}$ satisfying (1) & (2)

Sketch of proof $\Delta = \Delta(p)$ & $\Delta = \Delta(p)$

$$\rightsquigarrow H^*(\Delta) \cong \Delta[z^\pm], |z| = -2.$$

(Kadeishvili 1982)

Δ induces a minimal A_∞ -algebra structure
on $\Delta[z^\pm]$ s.t. $\Delta \simeq \Delta[z^\pm]$ as A_∞ -algebras

$$m_n: \Delta[z^\pm]^{\otimes n} \rightarrow \Delta[z^\pm] \text{ of degree } 2-n, n \geq 2$$

$$\rightsquigarrow \forall n \notin 2\mathbb{Z}, m_n = 0$$

$$\Delta = (\Delta[z^\pm], m_4, m_6, m_8, \dots)$$

$m_4 \in CC^{4,-2}(\Delta[z^\pm], \Delta[z^\pm])$: Hochschild cochain

$$\partial_{Hoch}(m_4) = 0 \rightsquigarrow \{m_4\} \in HH^{4,-2}(\Delta[z^\pm], \Delta[z^\pm])$$

$\{m_4\}$: Universal Maney Product (UMP)

Thm (]-Muro 2022)

R: isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ ^{crepant}_{res.}

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique
(Kanobian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is $2\mathbb{Z}$ -cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm]$, $|z| = -2$

Strategy Prove $R\text{End}(T) = \Delta(p)$

for $T \in \mathcal{A}$ satisfying (1) & (2)

Sketch of proof $\Delta = \Delta(p)$ & $\Delta = \Delta(p)$

$$\rightsquigarrow H^*(\Delta) \cong \Delta[z^\pm], |z| = -2.$$

(Kadeishvili 1982)

Δ induces a minimal A_∞ -algebra structure
on $\Delta[z^\pm]$ s.t. $\Delta \cong \Delta[z^\pm]$ as A_∞ -algebras

$$m_n: \Delta[z^\pm]^{\otimes n} \rightarrow \Delta[z^\pm] \text{ of degree } 2-n, n \geq 2$$

$$\rightsquigarrow \forall n \notin 2\mathbb{Z}, m_n = 0$$

$$\Delta = (\Delta[z^\pm], m_4, m_6, m_8, \dots)$$

$m_4 \in CC^{4,-2}(\Delta[z^\pm], \Delta[z^\pm])$: Hochschild cochain

$\partial_{Hoch}(m_4) = 0 \rightsquigarrow \{m_4\} \in HH^{4,-2}(\Delta[z^\pm], \Delta[z^\pm])$

$\{m_4\}$: Universal Maney Product (UMP)

$j: \Delta \hookrightarrow \Delta[z^\pm]$ inclusion of degree 0 component

$j^*: HH^{\bullet, \bullet}(\Delta[z^\pm], \Delta[z^\pm]) \longrightarrow HH^{\bullet, \bullet}(\Delta, \Delta[z^\pm])$

$j^*\{m_4\}$: restricted Universal Maney Product (rUMP)

Thm (J-Muro 2022)

R : isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ ^{crepant}_{res.}

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique (Kanobian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is $2\mathbb{Z}$ -cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm]$, $|z| = -2$

Strategy Prove $R\text{End}(T) = \Delta(p)$

for $T \in \mathcal{A}$ satisfying (1) & (2)

Sketch of proof $\Delta = \Delta(p)$ & $\Delta = \Delta(p)$

$$\rightsquigarrow H^*(\Delta) \cong \Delta[z^\pm], |z| = -2.$$

(Kadeishvili 1982)

Δ induces a minimal A_∞ -algebra structure on $\Delta[z^\pm]$ s.t. $\Delta \cong \Delta[z^\pm]$ as A_∞ -algebras

$$m_n: \Delta[z^\pm]^{\otimes n} \rightarrow \Delta[z^\pm] \text{ of degree } 2-n, n \geq 2$$

$$\rightsquigarrow \forall n \notin 2\mathbb{Z}, m_n = 0$$

$\Delta = (\Delta[z^\pm], m_4, m_6, m_8, \dots)$

$m_4 \in CC^{4,-2}(\Delta[z^\pm], \Delta[z^\pm])$: Hochschild cochain

$$\partial_{Hoch}(m_4) = 0 \rightsquigarrow \{m_4\} \in HH^{4,-2}(\Delta[z^\pm], \Delta[z^\pm])$$

$\{m_4\}$: Universal Maney Product (UMP)

$j: \Delta \hookrightarrow \Delta[z^\pm]$ inclusion of degree 0 component

$$j^*: HH^{\bullet, \bullet}(\Delta[z^\pm], \Delta[z^\pm]) \longrightarrow HH^{\bullet, \bullet}(\Delta, \Delta[z^\pm])$$

$j^*\{m_4\}$: restricted Universal Maney Product (rUMP)

Thm (J-Muro 2022)

$j^*\{m_4\} \in \underline{HH}^{\bullet, \bullet}(\Delta, \Delta[z^\pm])$ is a unit

↑ Hochschild-Tate cohomology

Thm (J-Muro 2022)

R : isolated cDV singularity & $p: X \rightarrow \text{Spec } R$ ^{crepant}_{res.}

Up to quasi-equivalence, $\mathcal{D}_{sg}(R)_{dg}$ is the unique (Kanobian) pre-triangulated dg category such that

- (1) $\exists T \in \mathcal{D}_{sg}(R)_{dg}$ is $2\mathbb{Z}$ -cluster tilting
- (2) $H^*(R\text{End}(T)) \cong \Delta(p)[z^\pm]$, $|z| = -2$

Strategy Prove $R\text{End}(T) = \Delta(p)$

for $T \in \mathcal{A}$ satisfying (1) & (2)

Sketch of proof $\Delta = \Delta(p)$ & $\Delta = \Delta(p)$

$$\rightsquigarrow H^*(\Delta) \cong \Delta[z^\pm], |z| = -2.$$

(Kadeishvili 1982)

Δ induces a minimal A_∞ -algebra structure on $\Delta[z^\pm]$ s.t. $\Delta \cong \Delta[z^\pm]$ as A_∞ -algebras

$$m_n: \Delta[z^\pm]^{\otimes n} \rightarrow \Delta[z^\pm] \text{ of degree } 2-n, n \geq 2$$

$$\rightsquigarrow \forall n \notin 2\mathbb{Z}, m_n = 0$$

$\Delta = (\Delta[z^\pm], m_4, m_6, m_8, \dots)$

$m_4 \in CC^{4,-2}(\Delta[z^\pm], \Delta[z^\pm])$: Hochschild cochain

$$\partial_{Hoch}(m_4) = 0 \rightsquigarrow \{m_4\} \in HH^{4,-2}(\Delta[z^\pm], \Delta[z^\pm])$$

$\{m_4\}$: Universal Maney Product (UMP)

$j: \Delta \hookrightarrow \Delta[z^\pm]$ inclusion of degree 0 component

$$j^*: HH^{\bullet, \bullet}(\Delta[z^\pm], \Delta[z^\pm]) \longrightarrow HH^{\bullet, \bullet}(\Delta, \Delta[z^\pm])$$

$j^*\{m_4\}$: restricted Universal Maney Product (rUMP)

Thm (J-Muro 2022)

$j^*\{m_4\} \in \underline{HH}^{\bullet, \bullet}(\Delta, \Delta[z^\pm])$ is a unit

↑ Hochschild-Tate cohomology

Moreover, a minimal A_∞ -algebra

$$A = (\Delta[z^\pm], m_4^A, m_6^A, m_8^A, \dots)$$

is A_∞ -isomorphic to Δ if and only if

$$j^*\{m_4^A\} \in \underline{HH}^{\bullet, \bullet}(\Delta, \Delta[z^\pm]) \text{ is a unit}$$