

In [2] we compute the excess for hereditary, radical square zero and monomial triangular algebras. For a bound quiver algebra Λ , a formula for the excess of Λ is obtained. We also give a criterion for Λ to be τ -rigid.

Let $\Lambda = kQ/I$ a bound quiver algebra, and let $Z\Lambda$ be its center. We have

$$\dim_k {}^\tau HH^1(\Lambda) = \dim_k Z\Lambda - \sum_{x \in Q_0} \dim_k x\Lambda x + \sum_{a \in Q_1} \dim_k t(a)\Lambda s(a).$$

Questions arise about Morita invariance, Morita derived invariance or derived invariance of ${}^\tau HH^1(\Lambda)$. Also about a possible Lie structure, and an eventual prolongation towards a cohomological theory.

REFERENCES

- [1] L. Angeleri Hügel *An Introduction to Auslander-Reiten Theory*, Lecture Notes Advanced School on Representation Theory and related Topics, 2006, ICTP Trieste.
<http://docplayer.net/178390377-Advanced-school-and-conference-on-representation-theory-and-related-topics.html>
- [2] C. Cibils; M. Lanzilotta; E.N. Marcos; A. Solotar, *On the first τ -tilting Hochschild cohomology of an algebra*, <https://arxiv.org/abs/2404.06916>
- [3] P. Gabriel *Finite representation type is open*, Represent. Algebr., Proc. int. Conf., Ottawa 1974, Lect. Notes Math. 488, (1975) 132–155.
- [4] B. Marsh *τ -tilting theory and τ -exceptional sequences*, mini-course at the τ -Research School, 5–8 September 2023, University of Cologne.
<https://sites.google.com/view/tau-tilting-school-cologne/schedule-abstracts>
<http://www1.maths.leeds.ac.uk/~marsh/MarshTauTiltingParts1to4.pdf>
- [5] G. Mazzola *The algebraic and geometric classification of associative algebras of dimension five*, Manuscr. Math. 27 (1979), 81–101.
- [6] O. Iyama; I. Reiten *Introduction to τ -tilting theory*, Proc. Natl. Acad. Sci. USA 111, (2014) 9704–9711.

Universal Massey products in representation theory of algebras

GUSTAVO JASSO

(joint work with Fernando Muro)

We work over an arbitrary field. Recall Kadeishvili’s Intrinsic Formality Criterion [Kad88]:

Theorem. Let A be a graded algebra whose Hochschild cohomology vanishes in the following bidegrees:

$$HH^{p+2, -p}(A, A) = 0, \quad p \geq 1.$$

Then, every minimal A_∞ -algebra structure on A is gauge A_∞ -isomorphic to the trivial A_∞ -structure, whose higher operations $m_{p+2} = 0$, $p \geq 1$, vanish.

In our joint work we generalise Kadeishvili’s Criterion as follows.

Definition. Fix an integer $d \geq 1$. A graded algebra is d -sparse if it is concentrated in degrees that are multiples of d (hence this condition is empty if $d = 1$). A d -sparse Massey algebra is a pair (A, m) consisting of a d -sparse graded algebra A and a Hochschild cohomology class

$$m \in \mathrm{HH}^{d+2, -d}(A, A), \quad \mathrm{Sq}(m) = 0,$$

of bidegree $(d+2, -d)$ whose Gerstenhaber square vanishes.

For example, if

$$(A, m_{d+2}, m_{2d+2}, m_{3d+2}, \dots)$$

is a minimal A_∞ -algebra structure on a d -sparse graded algebra A (in which case $m_{i+2} = 0$, $i \notin d\mathbb{Z}$, for degree reasons), then $m_{d+2} \in \mathrm{C}^{d+2, -d}(A, A)$ is a Hochschild cocycle whose associated Hochschild cohomology class

$$\{m_{d+2}\} \in \mathrm{HH}^{d+2, -d},$$

its *universal Massey product* (of length $d+2$), has vanishing Gerstenhaber square

$$\mathrm{Sq}(\{m_{d+2}\}) = 0.$$

Consequently, the pair $(A, \{m_{d+2}\})$ is a d -sparse Massey algebra.

Remark. It is an easy consequence of the d -sparsity assumption that the universal Massey product of a minimal A_∞ -algebra is invariant under A_∞ -isomorphisms.

Remark. Universal Massey products of length 3 have been investigated previously in representation theory, see for example [BKS04].

Definition. The *Hochschild–Massey cohomology* of a d -sparse Massey algebra (A, m) is the cohomology

$$\mathrm{HH}^{\bullet, *}(A, m)$$

of the *Hochschild–Massey (cochain) complex*, that is the bigraded cochain complex with components

$$\mathrm{HH}^{p+2, *}(A, A), \quad p \geq 0,$$

and differential

$$\mathrm{HH}^{\bullet, *}(A, A) \longrightarrow \mathrm{HH}^{\bullet+d+1, *-d}(A, A), \quad x \longmapsto [m, x],$$

in source bidegrees different from $(d+1, -d)$, where the differential is instead given by the formula by

$$\mathrm{HH}^{d+1, -d}(A, A) \longrightarrow \mathrm{HH}^{2(d+1), -2d}(A, A), \quad x \longmapsto [m, x] + x^2.$$

Remark. That the differential of the Hochschild–Massey complex squares to zero is a consequence of the Gerstenhaber relations and the assumption $\mathrm{Sq}(m) = 0$.

Theorem ([JKM22, Theorem B]). Let (A, m) be a d -sparse Massey algebra whose Hochschild–Massey cohomology vanishes in the following bidegrees:

$$\mathrm{HH}^{p+2, -p}(A, m) = 0, \quad p > d.$$

Then, any two minimal A_∞ -algebras

$$(A, m_{d+2}, m_{2d+2}, m_{3d+2}, \dots) \quad \text{and} \quad (A, m'_{d+2}, m'_{2d+2}, m'_{3d+2}, \dots)$$

such that $\{m_{d+2}\} = m = \{m'_{d+2}\}$ are gauge A_∞ -isomorphic.

Remark. Kadeishvili's Intrinsic Formality Criterion is indeed a corollary of the above theorem: Take $d = 1$ and notice that the hypothesis in the criterion implies that every minimal A_∞ -algebra structure on A has vanishing universal Massey product $\{m_3\} = 0$.

The proof of the theorem relies in an essential way on an enhanced A_∞ -obstruction theory developed by F. Muro in [Mur20a]. We also mention that the theorem is one of the key ingredients in the proof of the main theorem in [JKM22] which, as explained by B. Keller in the Appendix to *loc. cit.*, in a special case yields the final step in the proof of the Donovan–Wemyss Conjecture in the context of the Homological Minimal Model Program for threefolds [DW16, Wem23].

The aforementioned applications of the theorem rely on the following observation: The Hochschild–Massey cochain is equipped with a canonical bidegree $(d + 2, -d)$ endomorphism given by

$$\mathrm{HH}^{\bullet,*}(A, A) \longrightarrow \mathrm{HH}^{\bullet+d+2,*-d}(A, A), \quad x \longmapsto m \smile x,$$

in source bidegrees different from $(d + 1, -d)$, where it is given by

$$\mathrm{HH}^{d+1,-d}(A, A) \longrightarrow \mathrm{HH}^{2(d+1)+1,-2d}(A, A), \quad m \smile x + \{\delta_{/d}\} \smile x^2.$$

Here,

$$\delta_{/d} \in C^{1,0}(A, A), \quad x \longmapsto \frac{|x|}{d}x,$$

is the fractional Euler derivation (notice that $\frac{|x|}{d}$ is an integer due to the assumption that the graded algebra A is d -sparse). The above endomorphism is in fact null-homotopic. An explicit bidegree $(1, 0)$ null-homotopy is given by

$$\mathrm{HH}^{\bullet,*}(A, A) \longrightarrow \mathrm{HH}^{\bullet+1,*}(A, A), \quad x \longmapsto \{\delta_{/d}\} \smile x.$$

Thus, a sufficient condition for the assumptions in the theorem to be satisfied is that the components of above endomorphism of the Hochschild–Massey complex of (A, m) are bijective in all non-trivial source bidegrees. The latter condition is satisfied by the d -sparse Massey algebras investigated in [JKM22].

REFERENCES

- [BKS04] David Benson, Henning Krause, and Stefan Schwede. Realizability of modules over Tate cohomology. *Trans. Amer. Math. Soc.*, 356(9):3621–3668, 2004.
- [DW16] Will Donovan and Michael Wemyss. Noncommutative deformations and flops. *Duke Math. J.*, 165(8):1397–1474, 2016.
- [JKM22] Gustavo Jasso and Fernando Muro. The Derived Auslander–Iyama Correspondence, with an appendix by B. Keller, 2022, 2208.14413 [math.RT].
- [Kad88] T. V. Kadeishvili. The structure of the $A(\infty)$ -algebra, and the Hochschild and Harrison cohomologies. *Trudy Tbiliss. Mat. Inst. Razmadze Akad. Nauk Gruzin. SSR*, 91:19–27, 1988.
- [Mur20a] Fernando Muro. Enhanced A_∞ -obstruction theory. *J. Homotopy Relat. Struct.*, 15(1):61–112, 2020.
- [Wem23] Michael Wemyss. A lockdown survey on cDV singularities. In: Yukari Ito, Akira Ishii, and Osamu Iyama, editors, *McKay Correspondence, Mutation and Related Topics*, volume 88 of *Adv. Stud. Pure Math.*, pages 47–93, 2023.