


spectrum [Bal05], the noncommutative spectrum [NVY19] and Matsui's spectrum [Mat19].

However, we note that  $\text{Spent}(K)$  can be empty.

**Example 1.4.** *Let  $k$  be a field. Let  $K = \text{D}^b(\text{mod } kA_2)$  be the bounded derived category of finitely generated right  $kA_2$ -modules. Then  $\text{Thick}(K)$  is the lattice*



and  $\text{CLat}(\text{Thick}(K), \mathbf{2}) = \emptyset$ .

## REFERENCES

- [Bal05] Paul Balmer, *The spectrum of prime ideals in tensor triangulated categories*, J. Reine Angew. Math. **588** (2005), 149–168.
- [GS22] Sira Gratz and Greg Stevenson, *Approximating triangulated categories by spaces*, arXiv, 2022.
- [Mat19] Hiroki Matsui, *Singular equivalences of commutative noetherian rings and reconstruction of singular loci*, Journal of Algebra **522** (2019), 170–194.
- [NVY19] Daniel K Nakano, Kent B Vashaw, and Milen T Yakimov, *Noncommutative tensor triangular geometry*, arXiv preprint arXiv:1909.04304 (2019).

## Derived Endomorphism Algebras in Higher Auslander–Reiten Theory

GUSTAVO JASSO

(joint work with Fernando Muro)

Let  $k$  be a field and  $A$  a finite-dimensional algebra over  $k$ . Suppose that  $A$  is of finite representation type, that is the category  $\text{mod}(A)$  of finite-dimensional (right)  $A$ -modules admits an additive generator  $M$ , say. The algebra  $\Gamma := \text{End}_A(M)$  of endomorphisms of  $M$  is then an *Auslander algebra*, that is  $\Gamma$  has global dimension at most 2 and dominant dimension at least 2 [Aus71]. The basic paradigm of Auslander–Reiten Theory is that the minimal projective resolutions of simple  $\Gamma$ -modules of projective dimension 2 (the largest possible) correspond to almost-split sequences in  $\text{mod}(A)$  [AR75]. More generally, if  $d \geq 1$  and  $M$  is a  $d$ -cluster tilting  $A$ -module, then  $\Gamma$  is a  $(d+1)$ -dimensional Auslander algebra in the sense of Iyama [Iya07], that is  $\Gamma$  has global dimension at most  $d+1$  and dominant dimension at least  $d+1$ . We remind the reader that  $M$  is a  $d$ -cluster tilting  $A$ -module if the following conditions are equivalent for an indecomposable  $A$ -module  $X$ :

- $X$  is a direct summand of  $M$ .
- For all  $0 < i < d$ ,  $\text{Ext}_A^i(X, M) = 0$ .
- For all  $0 < i < d$ ,  $\text{Ext}_A^i(M, X) = 0$ .

Thus, a 1-cluster tilting  $A$ -module is simply an additive generator of  $\text{mod}(A)$  for the latter two conditions are empty in this case. In this more general context, minimal projective resolutions of simple  $\Gamma$ -modules of projective dimension  $d + 1$  correspond to *d-almost-split sequences* in  $\text{add}(M) \subseteq \text{mod}(A)$ , the additive closure of  $M$  in  $\text{mod}(A)$ . Furthermore, up to Morita equivalence, the association  $(A, M) \mapsto \text{End}_A(M)$  induces a bijection between:

- (1) Pairs  $(A, M)$  consisting of a finite-dimensional algebra  $A$  and a  $d$ -cluster tilting  $A$ -module  $M$ .
- (2)  $(d + 1)$ -Auslander algebras  $\Gamma$ .

The above bijective correspondence is known as the *Auslander–Iyama Correspondence* [Aus71, Iya07].

Suppose now that  $\Lambda$  is a finite-dimensional selfinjective algebra; for simplicity, assume  $\Lambda$  to be basic. We wish to interpret the minimal projective resolutions of simple  $\Gamma$ -modules of infinite (!) projective dimension in higher Auslander–Reiten-theoretic terms. For this, it is necessary to enforce a certain periodicity on these resolutions. More precisely, we assume that there exists an exact sequence of  $\Lambda$ -bimodules

$$0 \rightarrow \Lambda_\sigma \rightarrow P_{d+1} \rightarrow P_d \rightarrow \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

with projective middle terms, where  $\sigma$  is an algebra automorphism of  $\Lambda$ ; in this case we say that  $\Lambda$  is *twisted  $(d + 2)$ -periodic with respect to  $\sigma$* . Let  $S$  be a simple  $\Lambda$ -module of infinite projective dimension; applying the tensor product functor  $S \otimes_\Lambda -$  to the above exact sequence yields the first part of a projective resolution of  $S$  that is ‘twisted periodic’ since the  $(d + 2)$ -syzygy of  $S$  is again a simple  $\Lambda$ -module. Thus, the minimal total projective resolution of  $S$  is completely determined by the automorphism  $\sigma$  and the truncation

$$Q_{d+1} \rightarrow Q_d \rightarrow \cdots \rightarrow Q_2 \rightarrow Q_1 \rightarrow Q_0 \rightarrow \nu Q_0,$$

where  $Q_0$  is the projective cover of  $S$  and  $\nu Q_0$  is its injective hull. It is natural to wish to interpret the latter complex as an almost split  $(d + 2)$ -angle [IY08, GKO13]. Indeed, a theorem of Amiot [Ami07] in the case  $d = 1$  and a generalisation by Lin [Lin19] show that the pair  $(\text{proj}(\Lambda), - \otimes_\Lambda \Lambda_{\sigma^{-1}})$  admits a  $(d + 2)$ -angulation, where  $\text{proj}(\Lambda)$  is the category of finite-dimensional projective  $\Lambda$ -modules. Conversely, if  $\text{proj}(\Lambda)$  admits a  $(d + 2)$ -angulated structure, then  $\Lambda$  must be twisted  $(d + 2)$ -periodic with respect to some algebra automorphism [GSS03, GKO13, Han20]. Furthermore, if  $\Lambda$  arises as the endomorphism algebra of a  $d\mathbb{Z}$ -cluster tilting object in a triangulated category<sup>1</sup> with finite-dimensional morphism spaces, then  $\text{proj}(\Lambda)$  admits a  $(d + 2)$ -angulated structure [GKO13]. The main result in [JM22] refines the above to the following more precise statement (the case  $d = 1$  was established in [Mur22]):

<sup>1</sup>That is a (basic)  $d$ -cluster tilting object that is isomorphic to its  $d$ -fold shift.

**Theorem** (Derived Auslander–Iyama Correspondence). *Let  $k$  be a perfect field. There is a bijective correspondence between the following:*

- (1) *Quasi-isomorphism classes of DG algebras  $A$  such that  $H^0(A)$  is a basic finite-dimensional algebra and  $A$  is a  $d\mathbb{Z}$ -cluster tilting object of its perfect derived category  $D^c(A)$ .*
- (2) *Equivalence classes of pairs  $(\Lambda, \sigma)$  consisting of a basic finite-dimensional algebra  $\Lambda$  and  $\sigma$  is an algebra automorphism such that  $\Lambda$  is twisted  $(d+2)$ -periodic with respect to  $\sigma$ .*

*The correspondence is given by  $A \mapsto (H^0(A), \sigma)$ , where  $\sigma$  is a choice of algebra automorphism of  $H^0(A)$  such that  $H^{-d}(A) \cong H^0(A)_\sigma$  as  $H^0(A)$ -bimodules.*

The key ingredient in the proof of the theorem is the *restricted universal Massey product (rUMP)* of length  $d+2$  associated to any minimal  $A_\infty$ -model of  $A$  [Kad82, Kel01, LH]. By definition, the rUMP of  $A$  is the Hochschild cohomology class

$$u_A \in HH^{d+2, -d}(H^0(A), H^*(A))$$

that is the image of the class  $\{m_{d+2}\} \in HH^{d+2, -d}(H^*(A), H^*(A))$  of the higher operation  $m_{d+2}: H^*(A)^{\otimes d+2} \rightarrow H^*(A)[-d]$  under the canonical map

$$HH^{d+2, -d}(H^*(A), H^*(A)) \longrightarrow HH^{d+2, -d}(H^0(A), H^*(A)).$$

Indeed, a further main result in [JM22] is the following variant of the above theorem:

**Theorem.** *Let  $k$  be a perfect field. There is a bijective correspondence between the following:*

- (1) *Quasi-isomorphism classes of DG algebras  $A$  such that  $H^0(A)$  is a basic finite-dimensional algebra and  $A$  is a  $d\mathbb{Z}$ -cluster tilting object of its perfect derived category  $D^c(A)$ .*
- (2)  *$A_\infty$ -isomorphism classes of minimal  $A_\infty$ -algebras  $B$  with the following properties:*
  - *The underlying graded algebra of  $B$  is concentrated in degrees that are multiples of  $d$ , and there exists an invertible element  $\varphi \in B^d$ .*
  - *The rUMP  $u_B \in HH^{d+2, -d}(B^0, B)$  is invertible in the Hochschild–Tate cohomology (bigraded) algebra  $\underline{HH}^{\bullet, *}(B^0, B)$ .*

*The correspondence associates to a DG algebra  $A$  any of its minimal  $A_\infty$ -models.*

It is interesting to investigate in more detail the existence of additional structures on the DG algebras that arise from the Derived Auslander–Iyama Correspondence.

**Conjecture.** Let  $\Lambda$  be a basic finite-dimensional algebra that is twisted  $(d+2)$ -periodic with respect to the Nakayama automorphism  $\nu$  of  $\Lambda$ . Let  $A$  be any DG algebra that corresponds to  $(\Lambda, \nu)$  under the Derived Auslander–Iyama Correspondence. Then,  $A$  admits a right  $d$ -Calabi–Yau structure in the sense of [KS06].

The conjecture is motivated by the existence of a right  $d$ -Calabi–Yau structure on the Amiot–Guo–Keller cluster category [Ami09, Guo11, Kel05a] associated to

the derived  $(d + 1)$ -preprojective algebra [Kel11, IO13] of a  $d$ -representation finite algebra [IO11], see [KL23] for an announcement of the proof of a much more general theorem on Calabi–Yau structures on Drinfeld quotients.

## REFERENCES

- [Ami07] Claire Amiot. On the structure of triangulated categories with finitely many indecomposables. *Bull. Soc. Math. France*, 135(3):435–474, 2007.
- [Ami09] Claire Amiot. Cluster categories for algebras of global dimension 2 and quivers with potential. *Ann. Inst. Fourier (Grenoble)*, 59(6):2525–2590, 2009.
- [AR75] Maurice Auslander and Idun Reiten. Representation theory of artin algebras iii almost split sequences. *Comm. Alg.*, 3:3, 239–294, 1975.
- [Aus71] Maurice Auslander. *Representation dimension of Artin algebras*. Lecture Notes. Queen Mary College, 1971.
- [GKO13] Christof Geiss, Bernhard Keller, and Steffen Oppermann.  $n$ -angulated categories. *J. Reine Angew. Math.*, 675:101–120, 2013.
- [GSS03] Edward L. Green, Nicole Snashall, and Øyvind Solberg. The Hochschild cohomology ring of a selfinjective algebra of finite representation type. *Proc. Amer. Math. Soc.*, 131(11):3387–3393, 2003.
- [Guo11] Lingyan Guo. Cluster tilting objects in generalized higher cluster categories. *J. Pure Appl. Algebra*, 215(9):2055–2071, 2011.
- [Han20] Norihiro Hanihara. Auslander correspondence for triangulated categories. *Algebra Number Theory*, 14(8):2037–2058, 2020.
- [IO11] Osamu Iyama and Steffen Oppermann.  $n$ -representation-finite algebras and  $n$ -APR tilting. *Trans. Amer. Math. Soc.*, 363(12):6575–6614, 2011.
- [IO13] Osamu Iyama and Steffen Oppermann. Stable categories of higher preprojective algebras. *Adv. Math.*, 244:23–68, 2013.
- [IY08] Osamu Iyama and Yuji Yoshino. Mutation in triangulated categories and rigid Cohen-Macaulay modules. *Invent. Math.*, 172(1):117–168, 2008.
- [Iya07] Osamu Iyama. Auslander correspondence. *Adv. Math.*, 210(1):51–82, 2007.
- [JM22] Gustavo Jasso and Fernando Muro. The Derived Auslander–Iyama Correspondence (with an appendix by Bernhard Keller), arXiv:2208.14413 [math.RT].
- [Kad82] T. V. Kadeishvili. The algebraic structure in the homology of an  $A(\infty)$ -algebra. *Soobshch. Akad. Nauk Gruz. SSR*, 108(2):249–252 (1983), 1982.
- [Kel01] Bernhard Keller. Introduction to  $A$ -infinity algebras and modules. *Homology Homotopy Appl.*, 3(1):1–35, 2001.
- [Kel05a] Bernhard Keller. On triangulated orbit categories. *Doc. Math.*, 10:551–581, 2005.
- [Kel11] Bernhard Keller. Deformed Calabi–Yau completions. *J. Reine Angew. Math.*, 654:125–180, 2011. With an appendix by Michel Van den Bergh.
- [KL23] Bernhard Keller and Junyang Liu. Calabi–Yau structures on Drinfeld quotients and Amiot’s Conjecture, arXiv:2208.14413 [math.RT].
- [KS06] M. Kontsevich and Y. Soibelman. Notes on  $A_\infty$ -algebras,  $A_\infty$ -categories and non-commutative geometry. In *Homological mirror symmetry*, volume 757 of *Lecture Notes in Phys.*, pages 153–219. Springer, Berlin, 2006.
- [LH] Kenji Lefèvre-Hasegawa. Sur les  $A$ -infini catégories, math/0310337.
- [Lin19] Zengqiang Lin. A general construction of  $n$ -angulated categories using periodic injective resolutions. *J. Pure Appl. Algebra*, 223(7):3129–3149, 2019.
- [Mur22] Fernando Muro. Enhanced Finite Triangulated Categories. *J. Inst. Math. Jussieu*, 21(3):741–783, 2022.