

In the statement of the Weierstrass Product Theorem, we implicitly assumed that

- there exists a sequence  $\{k_n\} \subset \mathbb{N}$  so that

$$\sum_{n=1}^{\infty} \left( \frac{a}{|z_n|} \right)^{k_n}$$

converges

- within the set of sequences that give convergence, there exists a "minimal" sequence.

We proved the first point in the proof of the theorem— one can see that choosing  $k_n = n$  will guarantee convergence in all cases. As for the question of minimality, we explore in this section what a minimal sequence might be like, and how it reflects the underlying structure of our entire function.

### Definition 0.0.1: Order of Entire Function

Let  $f$  be an entire function. We say that  $f$  is of **order**  $\leq \rho$  for real  $\rho > 0$  if for all  $\varepsilon > 0$ , there exists a constant  $C_\varepsilon$  and a corresponding  $R_{C_\varepsilon} > 0$  such that, for all  $R > R_{C_\varepsilon}$ ,

$$\sup_{z \in D_R} |f(z)| \leq C_\varepsilon^{R^{\rho+\varepsilon}}.$$

We say that  $f$  is of **strict order**  $\leq \rho$  if the inequality holds without the  $\varepsilon$ :

$$\sup_{z \in D_R} |f(z)| \leq C_\varepsilon^{R^\rho}.$$

Finally, the function  $f$  is of **order**  $\rho$  or **strict order**  $\rho$  if  $\rho$  is the greatest lower bound to make the inequality valid.

### Example 0.0.1

The function  $e^z$  is strict order 1 because

$$|e^z| = e^x \leq e^{|z|}.$$

For a more involved example, if we have a canonical Weierstrass product with  $\sup_n \{k_n\} = k$ , then for all  $\rho$  satisfying  $k - 1 < \rho < k$ , the Weierstrass product is of order  $\leq \rho$ . The following lemma allows us to obtain the converse.

### Lemma 0.0.1

Let  $f$  be an entire function of strict order  $\leq \rho$ . Then for  $R$  sufficiently large,

$$\frac{1}{2\pi i} \int_{\partial D_R} \frac{f'}{f} dt \leq R^\rho.$$

Of course because  $f$  is entire, the left-hand side represents the number of zeros within the disc of radius  $R$ .

The following corollary immediately follows:

### Corollary 0.0.1

Let  $f$  have strict order  $\leq \rho$ , and let  $\{z_n\} \subset \mathbb{C} \setminus \{0\}$  be the zeros of  $f$  ordered by increasing modulus. Then for every  $\varepsilon > 0$ , the series

$$\sum_{n=1}^{\infty} \frac{1}{|z_n|^{\rho+\delta}}$$

converges.

This corollary then gives us the converse— every entire function of strict order  $\leq \rho$  has a Weierstrass product form with  $\rho < \sup_n k_n < \rho + 1$ . We summarize these results into the minimum modulus theorem.

**Theorem 0.0.1: Minimum Modulus Theorem**

Let  $f$  be an entire function of order  $\leq \rho$ , and let  $\{z_n\} \subset \mathbb{C}$  be its sequence of zeros ordered by increasing modulus. Then for all  $s > \rho$ ,  $\varepsilon > 0$ , there is a corresponding  $R_0$  so that, for all  $R > R_0$  and  $z \in \mathbb{C} \setminus \overline{D_s}$ :

$$|f(z)| \geq e^{-R^{\rho+\varepsilon}}$$

This ensures the construction we gave in Hadamard's Theorem is valid and unique.