

## MATH 451/551 Midterm 1

Name: \_\_\_\_\_

This exam contains 12 pages (including this cover page) and 11 questions.  
Total of points is 42.

Grade Table (for instructor use only)

Question	Points	Score
1	3	
2	3	
3	4	
4	3	
5	5	
6	3	
7	3	
8	5	
9	5	
10	5	
11	3	
Total:	42	

1. (3 points) Show that  $(A \cup B \cup C)^c = A^c B^c C^c$ . You may use DeMorgan's Law.

$$(A \cup B \cup C)^c = A^c (B \cup C)^c = A^c B^c C^c$$

applying DeMorgan's Law twice.

2. (3 points) Show that

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(AB) - P(AC) - P(BC) + P(ABC).$$

$$\begin{aligned} P(A \cup B \cup C) &= P(A) + P(B \cup C) - P(AB \cup AC) \\ &= P(A) + P(B) + P(C) - P(BC) - P(AB) - P(AC) + P(ABC) \end{aligned}$$

applying our rule for unions twice.

3. (a) (2 points) Let  $P(A) = .5$  and  $P(B^c) = .3$  and  $P(A \cup B) = .8$ . What is  $P(AB)$ ?

$$P(AB) = -P(A \cup B) + P(A) + P(B) = -.8 + .5 + .7 = .4.$$

- (b) (1 point) What is  $P(A^cB)$ ?

$$P(A^c \cup B) = P(B) - P(AB) = .7 - .4 = .3$$

- (c) (1 point) What is  $P(AB^c)$ ?

$$P(AB^c) = P(A) - P(AB) = .5 - .4 = .1$$

4. (3 points) Let  $P$  be a probability measure. Show that if  $A \subset B$  then  $P(A) \leq P(B)$ .

$$0 \leq P(BA^c) = P(B) - P(AB) = P(B) - P(A).$$

hence  $P(A) \leq P(B)$ .

5. (5 points) You are dealt 3 cards from a deck of 52. What is the probability that they all have the same suit?

There are 4 ways to choose the suit and then  $\binom{13}{3}$  ways to choose the cards out of a total of  $\binom{52}{3}$  ways. Hence the probability is

$$\frac{4 * \binom{13}{3}}{\binom{52}{3}}$$

6. (3 points) A box contains eight balls. Four of the balls have an “A” written on them and four of them have a “B” written on them. You draw out the balls one by one (without replacement) and place them in a row. How many distinct strings of “A”s and “B”s are there?

We simply need to choose out of the 8 possible places, the four places for the As to go. So  $\binom{8}{4} = 70$ .

7. (3 points) A group of  $n$  friends contains Larry and Rex. If the friends line up randomly, what is the probability Rex is standing next to Larry?

Consider two possibilities of orderings: (Rex, Larry, ...) and (Larry, Rex, ...). There are  $(n - 2)!$  ways to arrange each of the friends in the remaining spots and so  $2 * (n - 2)!$  ways total. However Rex and Larry don't need to stand at the start of the line. The first of them can be in any of  $n - 1$  positions. Thus there are  $2 * (n - 1)!$  ways for Rex and Larry to stand next to each other. Since there are a total of  $n!$  ways to arrange the friends then this probability is  $2/n$ .



8. (5 points) Of students at W&M, 55% are women and 45% are men. 1 in 100 men have the flu and 2 in 100 women have the flu. If a randomly chosen student has the flu, what is the probability they are a man?

Let  $M$  be the event of being a man and  $F$  be the event of getting the flu.

Then

$$P(M|F) = \frac{P(F|M)P(M)}{P(F)} + = \frac{P(F|M)P(M)}{P(F|M)P(M) + P(F|M^c)P(M^c)} = \frac{(.01)(.45)}{(.01)(.45) + (.02)(.55)}$$

9. (5 points) A car factory has three machines labeled A, B, and C. A makes 20% of the cars, B makes 30% of the cars, and C makes 50% of the cars. 6% of the cars made by A are defective, 7% of the cars made by B are defective, and 8% of the cars made by C are defective. What is the probability that a car is defective if it is made by this factory?

Let  $D$  be the event that the car is defective.

$$P(D) = P(D|A)P(A) + P(D|B)P(B) + P(D|C)P(C) = (.2)(.06) + (.3)(.07) + (.5)(.08) = .073.$$

10. (5 points) Let  $p$  be the probability that I find a pearl when I open an oyster. How many oysters should I open to make sure I have at least a 50% chance of finding at least one pearl? We can assume that finding a pearl in one oyster is independent of finding one in another.

Let  $A_i$  be finding a pearl in the  $i^{th}$  oyster, then

$$\begin{aligned} P(\text{finding at least one pearl}) &= 1 - P(\text{finding no pearls}) = 1 - P(A_1^c A_2^c A_3^c \cdots A_n^c) \\ &= 1 - \prod_{i=1}^n P(A_i^c) \\ &= 1 - (1 - p)^n \end{aligned}$$

and so if  $1/2 \leq 1 - (1 - p)^n$  then  $(1 - p)^n \leq 1/2$  and so  $n \log(1 - p) \leq \log(1/2)$  or  $n \geq -\log(2)/\log(1 - p)$ .

11. (3 points) Let  $A$  and  $B$  be both disjoint and independent. Let  $P(A) = 1/2$ . What is  $P(B)$ ?

$P(AB) = P(\emptyset) = 0$  and so  $0 = P(AB) = P(A)P(B) = 1/2P(B)$  hence  $P(B) = 0$ .