

Almost Sure:

$$P(\{\omega \mid X_n(\omega) \rightarrow X(\omega)\}) = 1$$

In Probability

$$\forall \varepsilon > 0 \quad \lim_{n \rightarrow \infty} P(|X_n - X| \geq \varepsilon) = 0$$

In Distribution:


$$F_n \rightarrow F$$

\uparrow CDF of X_n \uparrow CDF of X

Ex.

$$X_i \stackrel{\text{iid}}{\sim} U(0,1)$$

$$Y_n = \max_{i=1, \dots, n} X_i$$



$$Z_n = n(1 - Y_n)$$

\uparrow distributional limit?

$$F_n(z) = P(Z_n \leq z)$$

$$= P(n(1 - Y_n) \leq z)$$

$$= P(Y_n \geq 1 - z/n)$$

maximum

$$\begin{aligned}
 &= P(Y_n \geq 1 - \delta/n) \\
 &= 1 - P(Y_n \leq 1 - \delta/n)
 \end{aligned}$$

maxim.

$$\begin{aligned}
 &= 1 - P(X_1 \leq 1 - \delta/n, X_2 \leq 1 - \delta/n, \dots, X_n \leq 1 - \delta/n) \\
 &= 1 - \prod_{i=1}^n P(X_i \leq 1 - \delta/n)
 \end{aligned}$$

$X_i \sim U(0,1)$

CDF

$$= 1 - (1 - \delta/n)^n = F_n(z)$$

Recall: $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$

$$\lim_{n \rightarrow \infty} \left(1 + \frac{c}{n}\right)^n = e^c$$

$$\begin{aligned}
 \lim_{n \rightarrow \infty} F_n(z) &= \lim_{n \rightarrow \infty} 1 - (1 - \delta/n)^n \\
 &= 1 - \lim_{n \rightarrow \infty} (1 - \delta/n)^n
 \end{aligned}$$

$$F(z) = 1 - e^{-z}$$

CDF of $\text{Exp}(1)$

So $Z_n \xrightarrow{d} \text{Exp}(1)$


Theorem: a.s. \Rightarrow p \Rightarrow d

Partial Converse:

n is a constant

Partial Converse:

If $X_n \xrightarrow{d} c$ \swarrow c is a constant
i.e. a degenerate RV
then $X_n \xrightarrow{p} c$.



For a seq of numbers:

$$X_n, Y_n \in \mathbb{R}, \text{ and } X_n \rightarrow X, Y_n \rightarrow Y$$

then

$$\rightarrow X_n + Y_n \rightarrow X + Y$$

$$\rightarrow X_n Y_n \rightarrow XY$$

$$\rightarrow X_n / Y_n \rightarrow X / Y$$

$$\rightarrow aX_n + bY_n \rightarrow aX + bY$$

Theorem:

Let $X_n \rightarrow X, Y_n \rightarrow Y$ and $a, b \in \mathbb{R}$

and the convergence is either a.s. or p
(NOT d).

Then

$$(1) aX_n + bY_n \rightarrow aX + bY$$

$$(2) X_n Y_n \rightarrow XY \quad (\text{or } X_n / Y_n \rightarrow X / Y)$$

Note that:

We can treat a seq $C_n \in \mathbb{R}$ as a seq of degenerate RVs

so if $C_n \rightarrow C$ (as numbers)

then $C_n \xrightarrow{\text{a.s.}} C$ (as RVs)

Punchline: $C_n \rightarrow C$ and $X_n \rightarrow X$ (a.s. or i.p.)

then

$$(1) aX_n + bC_n \rightarrow aX + bC$$

$$(2) C_n X_n \rightarrow cX.$$

What about convergence in dist.?

Theorem: Slutsky's Theorem

If $X_n \xrightarrow{d} X$ and $Y_n \xrightarrow{P} c$ ← constant

then

$$(1) X_n + Y_n \xrightarrow{d} X + c$$

$$(2) X_n Y_n \xrightarrow{d} Xc$$

$$(X_n/Y_n \xrightarrow{d} X/c)$$

Theorem: Continuous Mapping Theorem

If $g: \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function

(1) (2) (3) (4) (5) (6) (7) (8) (9) (10) (11) (12) (13) (14) (15) (16) (17) (18) (19) (20) (21) (22) (23) (24) (25) (26) (27) (28) (29) (30) (31) (32) (33) (34) (35) (36) (37) (38) (39) (40) (41) (42) (43) (44) (45) (46) (47) (48) (49) (50) (51) (52) (53) (54) (55) (56) (57) (58) (59) (60) (61) (62) (63) (64) (65) (66) (67) (68) (69) (70) (71) (72) (73) (74) (75) (76) (77) (78) (79) (80) (81) (82) (83) (84) (85) (86) (87) (88) (89) (90) (91) (92) (93) (94) (95) (96) (97) (98) (99) (100)

If $g: \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function
and $X_n \rightarrow X$ (all types of convergence)

then $g(X_n) \rightarrow g(X)$

Pf.


$$\lim_{x \rightarrow a} g(x) = g(\lim_{x \rightarrow a} x)$$

Defn: Consistent Estimator

depends on N
samples

We say an estimator $\hat{\theta}_N$ is consistent for θ

if $\hat{\theta}_N \xrightarrow{P} \theta$
a constant



Another way: consistency \approx asymptotically unbiased \otimes

\otimes For nice enough dist

Ex.

$$S^2 = \frac{1}{N-1} \sum_{n=1}^N (X_n - \bar{X}_N)^2, \quad \mathbb{E}S^2 = \sigma^2$$

$$N-1 \quad n=1$$

$$\hat{\sigma}^2 = \frac{1}{N} \sum_{n=1}^N (X_n - \bar{X}_N)^2, \quad \mathbb{E} \hat{\sigma}^2 = \frac{N-1}{N} \sigma^2$$

notice: $\mathbb{E} \hat{\sigma}^2 \xrightarrow{n} \sigma^2$

Theorem: $\text{MSE} \rightarrow 0$ then $\hat{\theta}$ is consistent

If $\text{MSE}(\hat{\theta}_N) \xrightarrow{N} 0$ then $\hat{\theta}_N \xrightarrow{P} \theta$.

pf.

Want to show:

$$\lim_{n \rightarrow \infty} \underbrace{P(|\hat{\theta}_N - \theta| \geq \epsilon)} = 0$$

Markov's Ineq: $P(X \geq a) \leq \frac{\mathbb{E}X}{a} \quad (X \geq 0)$

$$P(|\hat{\theta}_N - \theta| \geq \epsilon) = P(\overbrace{(\hat{\theta}_N - \theta)^2}^{X \geq 0} \geq \epsilon^2) \quad \text{by Markov's}$$

$$\leq \frac{\mathbb{E}[(\hat{\theta}_N - \theta)^2]}{\epsilon^2}$$


$$= \frac{\text{MSE}(\hat{\theta}_N)}{\epsilon^2}$$

So

$$\text{MSE}(\hat{\theta}_N)$$

$$0 \leq P(|\hat{\theta}_n - \theta| \geq \varepsilon) \leq \frac{\text{MSE}(\hat{\theta}_n)}{\varepsilon^2}$$

If $\text{MSE}(\hat{\theta}_n) \rightarrow 0$ then so does my $P(|\hat{\theta}_n - \theta| \geq \varepsilon) \rightarrow 0$



Intuition:

\bar{X}_N this should be a "good" estimator of μ .

Saw: $E\bar{X}_N = \mu$ and $\text{Var}\bar{X}_N = \frac{\sigma^2}{N}$

Theorem: Weak Law of Large Numbers (WLLN)

If X_n s are uncorrelated and

① $E X_n = \mu$

② $\text{Var}(X_n) = \sigma^2 < \infty$

$$(2) \text{Var}(X_n) = 0 < \infty$$

then $\bar{X}_N \xrightarrow{P} \mu$.

Weak: weak assumptions and converg. in prob.

pf. Chebyshev's $\mu = EY$

$$P(|Y - \mu| \geq \varepsilon) \leq \frac{\text{Var}(Y)}{\varepsilon^2}$$

Want to show:

$$P(|\bar{X}_N - \mu| \geq \varepsilon) \xrightarrow{N} 0$$

Let $Y = \bar{X}_N$ then Chebyshev's says

$$0 \leq P(|\bar{X}_N - \mu| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X}_N)}{\varepsilon^2} = \frac{\sigma^2}{N\varepsilon^2}$$

$\rightarrow 0$

as $n \rightarrow \infty$.

Squeeze theorem

Ex. $X_n \stackrel{\text{iid}}{\sim} \text{Pois}(\lambda)$

$$EX_n = \lambda \text{ and } \text{Var}(X_n) = \lambda < \infty$$

WLLN: $\bar{X}_n \xrightarrow{P} \lambda$.

WLLN: $\bar{X}_N \xrightarrow{P} \lambda$.

Can slightly generalize WLLN

Assume $\text{Var}(X_n) = \sigma_n^2$ but

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \sigma_n^2 < \infty$$

then

Chebyshev's

$$\begin{aligned} P(|\bar{X}_N - \mu| \geq \varepsilon) &\leq \frac{\text{Var}(\bar{X}_N)}{\varepsilon^2} \\ &= \frac{\frac{1}{N^2} \sum_{n=1}^N \sigma_n^2}{\varepsilon^2} \\ &= \frac{1}{N} \left(\frac{\frac{1}{N} \sum_{n=1}^N \sigma_n^2}{\varepsilon^2} \right) \rightarrow 0 \end{aligned}$$

$< \infty$
as $n \rightarrow \infty$

So $\bar{X}_N \xrightarrow{P} \mu$.