

Random Matrix Theory for Machine Learning

Romain Couillet, Jiawei Gu

April 26, 2021

Numerous and large dimensional data is now a default setting in modern machine learning (ML). Standard ML algorithms, starting with kernel methods such as support vector machines and graph-based methods like the PageRank algorithm, were however initially designed out of small dimensional intuitions and tend to misbehave, if not completely collapse, when dealing with real-world large datasets. Random matrix theory has recently developed a broad spectrum of tools to help understand this new curse of dimensionality, to help repair or completely recreate the suboptimal algorithms, and most importantly to provide new intuitions to deal with modern data mining.

This monograph primarily aims to deliver these intuitions, by providing a digest of the recent theoretical and applied breakthroughs of random matrix theory into machine learning. Targeting a broad audience, spanning from undergraduate students interested in statistical learning to AI engineers and researchers alike, the mathematical prerequisites to the book are minimal (basics of probability theory, linear algebra and real and complex analysis are sufficient): as opposed to introductory books in the mathematical literature of random matrix theory and large dimensional statistics, the theoretical focus here is restricted to the essential requirements to machine learning applications. These applications range from detection, statistical inference and estimation, to graph- and kernel-based supervised, semi-supervised and unsupervised classification, as well as neural networks: for these, a precise theoretical prediction of the algorithm performance (often inaccessible when not resorting to a random matrix analysis), large dimensional insights, methods of improvement, along with a fundamental justification of the wide-scope applicability of the methods to real data, are provided.

Most methods, algorithms, and figures proposed in the monograph are coded in MATLAB and Python and made available to the readers. The monograph also contains a series of exercises of two types: short exercises with corrections appended to the end of the book to familiarize the reader with the basic theoretical notions and tools in random matrix analysis, as well as long guided exercises to apply these tools to further concrete machine learning applications.

Contents

1	Introduction	7
1.1	Motivation: the pitfalls of large dimensional statistics	7
1.1.1	The big data era: when n is no longer much larger than p	7
1.1.2	Sample covariance matrices in the large n, p regime	8
1.1.3	Kernel matrices of large dimensional data	11
1.1.4	Summarizing	17
1.2	Random matrix theory as an answer	19
1.2.1	Which theory and why?	19
1.2.2	The double asymptotics: turning the curse of dimensionality into a dimensionality blessing	24
1.2.3	Analyze, understand, and improve large dimensional machine learning methods	27
1.2.4	Exploiting universality: from large dimensional Gaussian vectors to real data	33
1.3	Outline and online toolbox	39
1.3.1	Organization of the manuscript	39
1.3.2	Online codes	42
2	Basics of Random Matrix Theory	43
2.1	Fundamental objects	43
2.1.1	The resolvent	43
2.1.2	Spectral measure and Stieltjes transform	44
2.1.3	Cauchy's integral, linear eigenvalue functionals, and eigenspaces	46
2.1.4	Deterministic and random equivalents	48
2.2	Foundational random matrix results	50
2.2.1	Key lemmas and identities	51
2.2.2	The Marčenko-Pastur and semicircle laws	57
2.2.3	Large dimensional sample covariance matrices and generalized semicircles	75
2.3	Advanced spectrum considerations for sample covariances	87
2.3.1	Limiting spectrum	88
2.3.2	"No eigenvalue outside the support"	93
2.4	Preliminaries on statistical inference	95
2.4.1	Linear eigenvalue statistics	96

2.4.2	Eigenvector projections and subspace methods	103
2.5	Spiked models	110
2.5.1	Isolated eigenvalues	110
2.5.2	Isolated eigenvectors	114
2.5.3	Limiting fluctuations	118
2.5.4	Further discussions and other spiked models	120
2.6	Information-plus-noise, deformed Wigner, and other models	123
2.6.1	Why focus on the sample covariance matrix model?	123
2.6.2	Other models	125
2.6.3	Other statistics	136
2.7	Beyond vectors of independent entries: concentration of measure in RMT	139
2.7.1	Limitations of the i.i.d. assumption	139
2.7.2	Concentrated random vectors as the answer	139
2.7.3	Elements of concentration of measure for random matrices	143
2.7.4	A concentration inequality version of Theorem 2.6	149
2.8	Concluding remarks	155
2.9	Exercises	157
2.9.1	Properties of the Stieltjes transform	157
2.9.2	On limiting laws	158
2.9.3	On eigen-inference	159
2.9.4	Spiked models	160
2.9.5	Deterministic equivalent	160
2.9.6	Concentration of measure	161
2.9.7	Beyond matrices	162
3	Statistical Inference in Linear Models	165
3.1	Detection and estimation in information-plus-noise models	167
3.1.1	GLRT asymptotics	167
3.1.2	Linear and Quadratic Discriminant Analysis	170
3.1.3	Subspace methods: the G-MUSIC algorithm	178
3.2	Covariance matrix distance estimation	183
3.2.1	Distances and divergences between Gaussian laws	183
3.2.2	The random matrix framework	184
3.2.3	Closed-form expressions	188
3.2.4	The Wasserstein and Frobenius distances	192
3.2.5	Application to covariance-based spectral clustering	194
3.3	M-estimators of scatter	195
3.3.1	Reminder on robust statistics	195
3.3.2	The M-estimator of scatter	197
3.3.3	The random matrix framework	198
3.3.4	Extensions	205
3.4	Concluding remarks	210
3.5	Practical course material	212

4 Kernel Methods	219
4.1 Basic setting	220
4.1.1 The non-trivial growth rates	220
4.1.2 Statistical data model	222
4.2 Distance and inner-product random kernel matrices	223
4.2.1 Main intuition	223
4.2.2 Main Results	228
4.3 The α - β random kernel model	232
4.3.1 Motivation	232
4.3.2 Main results	234
4.4 Properly scaling kernel model	238
4.4.1 Motivation	238
4.4.2 Setting	239
4.4.3 Limiting spectrum of the null model	242
4.4.4 Main results	248
4.5 Implications to kernel methods	251
4.5.1 Application to kernel spectral clustering	251
4.5.2 Application to semi-supervised kernel learning	261
4.5.3 Application to kernel ridge regression	275
4.5.4 Summary of Section 4.5	281
4.6 Concluding remarks	282
4.7 Practical course material	284
5 Large Neural Networks	291
5.1 Random neural networks	291
5.1.1 Regression with random neural networks	292
5.1.2 Delving deeper into limiting kernels	302
5.2 Gradient descent dynamics in learning linear neural nets	309
5.3 Recurrent neural nets: echo state networks	314
5.3.1 Preliminaries and echo state networks	314
5.3.2 Results on ESN asymptotics	316
5.4 Concluding remarks	322
5.5 Practical course material	324
6 Optimization-based Methods with Non-explicit Solutions	327
6.1 Generalized linear classifier	328
6.1.1 Basic setting	328
6.1.2 Intuitions and main results	329
6.1.3 Practical consequences and further discussions	334
6.2 Large dimensional support vector machines	339
6.3 Concluding remarks	345
6.4 Practical course material	347

7	Community Detection on Graphs	351
7.1	Community detection in dense graphs	352
7.1.1	The stochastic block model	352
7.1.2	The degree-corrected stochastic block model	362
7.2	From dense to sparse graphs: a different approach	369
7.2.1	The non-backtracking matrix	370
7.2.2	The Bethe Hessian matrix	371
7.2.3	Degree regularization	372
7.2.4	A unifying approach adapted to DC-SBM	372
7.3	Concluding remarks	376
7.4	Practical course material	377
8	Discussions on Universality and Practical Applications	381
8.1	From Gaussian mixtures to concentrated random vectors and GAN images	381
8.1.1	On data models in large dimensions	381
8.1.2	A study of GAN data	383
8.2	Wide-sense universality in large dimensional machine learning .	390
8.3	Discussions and conclusions	393
9	Solutions to exercises in Section 2.9	395

Chapter 1

Introduction

1.1 Motivation: the pitfalls of large dimensional statistics

1.1.1 The big data era: when n is no longer much larger than p

The big data revolution comes along with the challenging need to parse, mine, compress a large amount of large dimensional and possibly heterogeneous data. In many applications, the dimension p of the observations is as large as – if not much larger than – their number n . In array processing and wireless communications, the number of antennas required for fine localization resolution or increased communication throughput may be as large (today in the order of hundreds) as the number of available independent signal observations [Lu et al., 2014, Li and Stoica, 2007]. In genomics, the identification of correlations among hundred of thousands genes based on a limited number of independent (and expensive) samples induces an even larger ratio p/n [Arnold et al., 1994]. In statistical finance, portfolio optimization relies on the need to invest on a large number p of assets to reduce volatility but at the same time to estimate the current (rather than past) asset statistics from a relatively small number n of asset return records [Laloux et al., 2000].

As we shall demonstrate in the next section, the fact that in these problems n is not *much larger* than p annihilates most of the results from standard asymptotic statistics that assume n alone is large [Vaart, 2000]. As a rule of thumb, by “*much larger*” we mean here that n must be at least 100 times larger than p for standard asymptotic statistics to be of practical convenience (see our argument in Section 1.1.2). Many algorithms in statistics, signal processing, and machine learning are precisely derived from this $n \gg p$ assumption that is no longer appropriate today. A major objective of this monograph is to cast some light on the resulting biases and problems incurred and to provide a systematic random matrix framework to improve these algorithms.

Possibly more importantly, we will see along this monograph that (small p) finite-dimensional intuitions which are at the core of many machine learning algorithms (starting with spectral clustering [Ng et al., 2002, Luxburg, 2007]) may strikingly fail when applied in a simultaneously large n, p setting. A compelling example lies in the notion of “distance” between vectors. Most classification methods in machine learning are rooted in the observation that random data vectors arising from a mixture distribution (say Gaussian) gather in “groups” of close-by vectors in Euclidean norm. When dealing with large dimensional data, however, concentration phenomena arise that make Euclidean distances useless, if not counterproductive: vectors from the *same* mixture class may be further away in Euclidean distance than vectors arising from *different* classes: while classification may still be doable, it works in a rather different way from our finite-dimensional intuition. The monograph intends to prepare the reader for the multiple traps caused by this “curse of dimensionality”.

1.1.2 Sample covariance matrices in the large n, p regime

Let us consider the following illustrating example which shows a first elementary, yet counterintuitive, result: for simultaneously large n, p , the sample covariance matrix $\hat{\mathbf{C}} \in \mathbb{R}^{p \times p}$ based on n samples $\mathbf{x}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{C})$ is an *entry-wise* consistent estimator of the population covariance $\mathbf{C} \in \mathbb{R}^{p \times p}$ (i.e., $\|\hat{\mathbf{C}} - \mathbf{C}\|_\infty \rightarrow 0$ as $p, n \rightarrow \infty$ for $\|\mathbf{A}\|_\infty \equiv \max_{ij} |\mathbf{A}_{ij}|$) while overall being an extremely poor estimator in a (more practical) operator norm sense (i.e., $\|\hat{\mathbf{C}} - \mathbf{C}\| \not\rightarrow 0$ with here $\|\cdot\|$ the operator norm). Matrix norms are in particular *not* equivalent in the large n, p scenario.

Let us detail this claim, in the simplest case where $\mathbf{C} = \mathbf{I}_p$. Consider a data set $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ of n independent and identically distributed (i.i.d.) observations from a p -dimensional standard Gaussian distribution, i.e., $\mathbf{x}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ for $i \in \{1, \dots, n\}$. We wish to estimate the population covariance matrix $\mathbf{C} = \mathbf{I}_p$ from the n available samples. The maximum likelihood estimator in this zero-mean Gaussian setting is the sample covariance matrix $\hat{\mathbf{C}}$ defined by

$$\hat{\mathbf{C}} = \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top = \frac{1}{n} \mathbf{X} \mathbf{X}^\top. \quad (1.1)$$

By the strong law of large numbers, for fixed p , $\hat{\mathbf{C}} \rightarrow \mathbf{I}_p$ almost surely as $n \rightarrow \infty$, so that $\|\hat{\mathbf{C}} - \mathbf{I}_p\| \xrightarrow{a.s.} 0$ holds for any standard matrix norm and in particular for the operator norm.

One must be more careful when dealing with the case $n, p \rightarrow \infty$ with ratio $p/n \rightarrow c \in (0, \infty)$ (or, from a practical standpoint, n is *not much larger* than p). First, note that the entry-wise convergence still holds since, invoking the law of large numbers again

$$[\hat{\mathbf{C}}]_{ij} = \frac{1}{n} \sum_{l=1}^n [\mathbf{X}]_{il} [\mathbf{X}]_{jl} \xrightarrow{a.s.} \begin{cases} 1, & i = j \\ 0, & i \neq j. \end{cases}$$

Besides, by a concentration inequality argument, it can even be shown that

$$\max_{1 \leq i, j \leq p} |[\hat{\mathbf{C}} - \mathbf{I}_p]_{ij}| \xrightarrow{a.s.} 0$$

which holds as long as p is no larger than a polynomial function of n , and thus

$$\|\hat{\mathbf{C}} - \mathbf{I}_p\|_\infty \xrightarrow{a.s.} 0.$$

Consider now the case $p > n$. Since $\hat{\mathbf{C}} = \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top$ is the sum of n rank-one matrices, the rank of $\hat{\mathbf{C}}$ is at most equal to n and thus, being a $p \times p$ matrix with $p < n$, the sample covariance matrix $\hat{\mathbf{C}}$ must be a singular matrix having at least $p - n > 0$ null eigenvalues. As a consequence,

$$\|\hat{\mathbf{C}} - \mathbf{I}_p\| \not\rightarrow 0$$

for $\|\cdot\|$ the matrix operator (or spectral) norm. This last result actually extends to the general case where $p/n \rightarrow c \in (0, \infty)$. As such, matrix norms cannot be considered equivalent in the regime where p is not negligible compared to n . This follows from the fact that the (matrix norm) equivalence factors depend on the matrix size p ; here for instance, we have that for symmetric matrices $\mathbf{A} \in \mathbb{R}^{p \times p}$, $\|\mathbf{A}\|_\infty \leq \|\mathbf{A}\| \leq p\|\mathbf{A}\|_\infty$.

Unfortunately, in practice, the (non-converging) operator norm is of more practical interest than the (converging) infinity norm.

Remark 1.1 (On the importance of operator norm). *For practical purposes, this “loss” of norm equivalence for large p raises the question of the relevant matrix norm to consider for a given application. For the purpose of the present monograph, and for most applications in machine learning, the operator (or spectral) norm is the most relevant. First, the operator norm is the matrix norm induced by the Euclidean norm of vectors. Thus, the study of regression vectors or label/score vectors in classification is naturally attached to the spectral study of matrices. Besides, we will often be interested in the asymptotic equivalence of families of large dimensional matrices. If $\|\mathbf{A}_p - \mathbf{B}_p\| \rightarrow 0$ for matrix sequences $\{\mathbf{A}_p\}$ and $\{\mathbf{B}_p\}$, indexed by their dimension p , then according to Weyl’s inequality (see for example Lemma 2.10 in Section 2.2.1),*

$$\max_i |\lambda_i(\mathbf{A}_p) - \lambda_i(\mathbf{B}_p)| \rightarrow 0$$

for $\lambda_1(\mathbf{A}) \geq \lambda_2(\mathbf{A}) \geq \dots$ the eigenvalues of \mathbf{A} in a decreasing order. Besides, for $\mathbf{u}_i(\mathbf{A}_p)$ an eigenvector of \mathbf{A}_p associated with an isolated eigenvalue $\lambda_i(\mathbf{A}_p)$ (i.e., such that $\min\{|\lambda_{i+1}(\mathbf{A}_p) - \lambda_i(\mathbf{A}_p)|, |\lambda_i(\mathbf{A}_p) - \lambda_{i-1}(\mathbf{A}_p)|\} > \varepsilon$ for some $\varepsilon > 0$ uniformly on p),

$$\|\mathbf{u}_i(\mathbf{A}_p) - \mathbf{u}_i(\mathbf{B}_p)\| \rightarrow 0.$$

These results ensure that, as far as spectral properties are concerned, \mathbf{A}_p can be studied equivalently through \mathbf{B}_p . We will often use this argument to investigate intractable random matrices \mathbf{A}_p by means of a more tractable “proxy” \mathbf{B}_p .

The pitfall that consists in assuming that $\hat{\mathbf{C}}$ is a valid estimator of \mathbf{C} since $\|\hat{\mathbf{C}} - \mathbf{C}\|_\infty \xrightarrow{a.s.} 0$ may thus have deleterious practical consequences when n is not significantly larger than p .

Resuming on our norm convergence discussion, it is now natural to ask whether $\hat{\mathbf{C}}$, which badly estimates \mathbf{C} , has a controlled asymptotic behavior. There precisely lay the first theoretical interests of random matrix theory. While $\hat{\mathbf{C}}$ itself does not converge in any useful way, its eigenvalue distribution does exhibit a traceable limiting behavior [Marčenko and Pastur, 1967, Silverstein and Bai, 1995, Bai and Silverstein, 2010]. The seminal result in this direction, due to Marčenko and Pastur, states that, for $\mathbf{C} = \mathbf{I}_p$, as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$, with probability one, the random *discrete eigenvalue/empirical spectral distribution*

$$\mu_p \equiv \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\hat{\mathbf{C}})}$$

converges in law to a non-random *smooth* limit, today referred to as the “Marčenko-Pastur law” [Marčenko and Pastur, 1967]

$$\mu(dx) = (1 - c^{-1})^+ \delta_0(x) + \frac{1}{2\pi cx} \sqrt{(x - E_-)^+(E_+ - x)^+} dx \quad (1.2)$$

where $E_{\pm} = (1 \pm \sqrt{c})^2$ and $(x)^+ \equiv \max(x, 0)$.

Figure 1.1 compares the empirical spectral distribution of $\hat{\mathbf{C}}$ to the limiting Marčenko-Pastur law given in (1.2), for $p = 500$ and $n = 50\,000$.




Figure 1.1: Histogram of the empirical spectral distribution of $\hat{\mathbf{C}}$ versus the Marčenko-Pastur law, for $p = 500$ and $n = 50\,000$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).



The elementary Marčenko-Pastur result is already quite instructive and insightful.

Remark 1.2 (When is one under the random matrix regime?). *Equation (1.2) reveals that the eigenvalues of $\hat{\mathbf{C}}$, instead of concentrating at $x = 1$ as a large- n alone analysis would suggest, are spread from $(1 - \sqrt{c})^2$ to $(1 + \sqrt{c})^2$. As such, the eigenvalues span on a range*

$$(1 + \sqrt{c})^2 - (1 - \sqrt{c})^2 = 4\sqrt{c}.$$

This is a slow decaying behavior with respect to $c = \lim p/n$. In particular, for $n = 100p$, in which case one would expect a sufficiently large number of samples for $\hat{\mathbf{C}}$ to properly estimate $\mathbf{C} = \mathbf{I}_p$, one has $4\sqrt{c} = 0.4$ which is a large spread around the mean (and true) eigenvalue 1. This is visually confirmed by Figure 1.1 for $p = 500$ and $n = 50\,000$, where the histogram of the eigenvalues is nowhere near concentrated at $x = 1$. Therefore, random matrix results will be much more accurate than classical asymptotic statistics even when $n \sim 100p$. As a telling example, estimating the covariance matrix of each digit from the popular MNIST dataset [LeCun et al., 1998], made of no more than 60 000 training samples (and thus about $n = 6\,000$ samples per digit) of size $p = 784$, is likely a hazardous undertaking.

Remark 1.3 (On universality). *Although introduced here in the context of Gaussian distribution for \mathbf{x}_i , the Marčenko-Pastur law applies to much more general cases. Indeed, the result remains valid so long that the \mathbf{x}_i 's have independent normalized entries of zero mean and unit variance (and even beyond this setting, see [El Karoui, 2009, Louart and Couillet, 2018]). Similar to the law of large numbers in standard asymptotic statistics, this universality phenomenon commonly arises in random matrix theory and large dimensional statistics. We will exploit this phenomenon in the monograph to justify the wide applicability of the presented results, even to real datasets. See Chapter 8 for more detail.*

1.1.3 Kernel matrices of large dimensional data

Another less known but equally important example of the curse of dimensionality in machine learning involves the loss of relevance of (the notion of) Euclidean distance between large dimensional data vectors. To be more precise, we will see in the sequel that, *in an asymptotically non-trivial classification setting* (that is, ensuring that asymptotic classification is neither trivially easy nor impossible), large and numerous data vectors $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ extracted from a few-class (say two-class) mixture model tend to be asymptotically at *equal* (Euclidean) distance from one another, irrespective of their corresponding class. Roughly said, in this non-trivial setting and under some reasonable statistical assumptions on the \mathbf{x}_i 's, we have

$$\max_{1 \leq i \neq j \leq n} \left\{ \frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 - \tau \right\} \rightarrow 0 \quad (1.3)$$

for some constant $\tau > 0$ as $n, p \rightarrow \infty$, *independently of the classes (same or different) of \mathbf{x}_i and \mathbf{x}_j* (here the normalization by p is used for compliance with the notations in the remainder of this monograph and has no particular importance).

This asymptotic behavior is extremely counterintuitive and conveys the idea that classification by standard methods ought not to be doable in this large dimensional regime. Indeed, in the conventional finite-dimensional intuition that forged many of the leading machine learning algorithms of everyday use (such as spectral clustering [Ng et al., 2002, Luxburg, 2007]), two data points are assigned to the same class if they are “close” in Euclidean distance. Here we claim that, when p is large, *data pairs are neither close nor far* from each other, regardless of their belonging to the same class or not. Despite this troubling loss of *individual* discriminative power between data pairs, we subsequently show that, thanks to a *collective* behavior of all data belonging to the same (few and thus large) classes, asymptotic data classification or clustering is still achievable. Better, we shall see that, while many conventional methods devised from small dimensional intuitions do fail in this large dimensional regime, some popular approaches (such as the Ng-Jordan-Weiss spectral clustering method [Ng et al., 2002] or the PageRank semi-supervised learning approach [Avrachenkov et al., 2012]) still function. But the core reasons for their functioning are strikingly different from the reasons of their initial designs, and they often operate far from optimally.

1.1.3.1 The non-trivial classification regime

To get a clear picture of the source of Equation (1.3), we first need to clarify what we refer to as the “asymptotically non-trivial” classification setting. Consider the simplest scenario of a binary Gaussian mixture classification: given a training set $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ of n samples independently drawn from the two-class (\mathcal{C}_1 and \mathcal{C}_2) Gaussian mixture

$$\mathcal{C}_1 : \mathbf{x} \sim \mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p), \quad \mathcal{C}_2 : \mathbf{x} \sim \mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p + \mathbf{E}) \quad (1.4)$$

each drawn with probability 1/2, for some deterministic $\boldsymbol{\mu} \in \mathbb{R}^p$ and symmetric $\mathbf{E} \in \mathbb{R}^{p \times p}$, both possibly depending on p . In the ideal case where $\boldsymbol{\mu}$ and \mathbf{E} are perfectly known, one can devise a (decision optimal) Neyman-Pearson test. For an unknown \mathbf{x} , genuinely belonging to \mathcal{C}_1 , the Neyman-Pearson test to decide on the class of \mathbf{x} reads

$$(\mathbf{x} + \boldsymbol{\mu})^\top (\mathbf{I}_p + \mathbf{E})^{-1} (\mathbf{x} + \boldsymbol{\mu}) - (\mathbf{x} - \boldsymbol{\mu})^\top (\mathbf{x} - \boldsymbol{\mu}) \stackrel{\mathcal{C}_1}{\gtrless} -\log \det(\mathbf{I}_p + \mathbf{E}). \quad (1.5)$$

Writing $\mathbf{x} = \boldsymbol{\mu} + \mathbf{z}$ for $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, the above test is equivalent to

$$\begin{aligned} T(\mathbf{x}) \equiv & 4\boldsymbol{\mu}^\top (\mathbf{I}_p + \mathbf{E})^{-1} \boldsymbol{\mu} + 4\boldsymbol{\mu}^\top (\mathbf{I}_p + \mathbf{E})^{-1} \mathbf{z} + \mathbf{z}^\top ((\mathbf{I}_p + \mathbf{E})^{-1} - \mathbf{I}_p) \mathbf{z} \\ & + \log \det(\mathbf{I}_p + \mathbf{E}) \stackrel{\mathcal{C}_2}{\gtrless} 0. \end{aligned} \quad (1.6)$$

Since $\mathbf{U}\mathbf{z}$ for $\mathbf{U} \in \mathbb{R}^{p \times p}$ an eigenvector basis of $(\mathbf{I}_p + \mathbf{E})^{-1}$ (and thus of $(\mathbf{I}_p + \mathbf{E})^{-1} - \mathbf{I}_p$) follows the same distribution as \mathbf{z} , the random variable $T(\mathbf{x})$ can be written as the sum of p independent random variables. Further assuming

that $\|\boldsymbol{\mu}\| = O(1)$ with respect to p , by Lyapunov's central limit theorem (e.g., [Billingsley, 2012, Theorem 27.3]) and the fact that $\text{Var}[\mathbf{z}^\top \mathbf{A} \mathbf{z}] = 2 \text{tr}(\mathbf{A}^2)$ for symmetric $\mathbf{A} \in \mathbb{R}^{p \times p}$ and Gaussian \mathbf{z} , we have, as $p \rightarrow \infty$,

$$V_T^{-1/2}(T(\mathbf{x}) - \bar{T}) \xrightarrow{d} \mathcal{N}(0, 1),$$

where

$$\begin{aligned}\bar{T} &\equiv 4\boldsymbol{\mu}^\top (\mathbf{I}_p + \mathbf{E})^{-1} \boldsymbol{\mu} + \text{tr}(\mathbf{I}_p + \mathbf{E})^{-1} - p + \log \det(\mathbf{I}_p + \mathbf{E}), \\ V_T &\equiv 16\boldsymbol{\mu}^\top (\mathbf{I}_p + \mathbf{E})^{-2} \boldsymbol{\mu} + 2 \text{tr}((\mathbf{I}_p + \mathbf{E})^{-1} - \mathbf{I}_p)^2.\end{aligned}$$

As a consequence, the classification of $\mathbf{x} \in \mathcal{C}_1$ is asymptotically non-trivial (i.e., the classification error neither goes to 0 nor 1 as $p \rightarrow \infty$) if and only if \bar{T} is of the same order as $\sqrt{V_T}$. Considering the (worst case) scenario where $\mathbf{E} = \mathbf{0}$, we must have $\|\boldsymbol{\mu}\| \geq O(1)$ with respect to p (indeed, if instead $\|\boldsymbol{\mu}\| = o(1)$, the classification of \mathbf{x} is asymptotically impossible).

Under the constraint $\|\boldsymbol{\mu}\| = O(1)$, we move on to consider the case $\mathbf{E} \neq \mathbf{0}$ with spectral norm $\|\mathbf{E}\| = o(1)$. By a Taylor expansion of both $(\mathbf{I}_p + \mathbf{E})^{-1}$ and $\log \det(\mathbf{I}_p + \mathbf{E})$ around \mathbf{I}_p we obtain

$$\begin{aligned}\bar{T} &= 4\|\boldsymbol{\mu}\|^2 - \frac{1}{2} \text{tr}(\mathbf{E}^2) + o(1); \\ V_T &= 16\|\boldsymbol{\mu}\|^2 + 2 \text{tr}(\mathbf{E}^2) + o(1),\end{aligned}$$

which demands $\text{tr}(\mathbf{E}^2)$ to be of order $O(1)$ (same as $\|\boldsymbol{\mu}\|$) so as to have discriminative power. Since $\text{tr}(\mathbf{E}^2) \leq p\|\mathbf{E}\|^2$, with equality if and only if \mathbf{E} is proportional to identity, i.e., $\mathbf{E} = \epsilon \mathbf{I}_p$, one must have $\|\mathbf{E}\| \geq O(p^{-1/2})$. Also, by the Cauchy-Schwarz inequality, we have $|\text{tr} \mathbf{E}| \leq \sqrt{\text{tr}(\mathbf{E}^2) \cdot \text{tr} \mathbf{I}_p} = O(\sqrt{p})$, with equality if and only if $\mathbf{E} = \epsilon \mathbf{I}_p$, and we must therefore have $|\text{tr} \mathbf{E}| \geq O(\sqrt{p})$. This allows us to conclude on the following non-trivial classification conditions

$$\|\boldsymbol{\mu}\| \geq O(1), \quad \|\mathbf{E}\| \geq O(p^{-1/2}), \quad |\text{tr}(\mathbf{E})| \geq O(\sqrt{p}), \quad \text{tr}(\mathbf{E}^2) \geq O(1). \quad (1.7)$$

These are the *minimal* conditions for classification in the case of perfectly known means and covariances in the following sense: (i) if none of the inequalities hold (i.e., if the means and covariances from both classes are too close), asymptotic classification must fail, and (ii) if at least one of the inequalities is not tight (say if $\|\boldsymbol{\mu}\| \geq O(\sqrt{p})$), asymptotic classification becomes trivial.¹

We shall subsequently see that (1.7) precisely induces the asymptotic loss of distance discrimination raised in (1.3) but that standard spectral clustering methods based on $n \sim p$ data remain valid.

¹It should be noted here that, unlike in computer science, we will stick in this manuscript with the notation $O(\cdot)$ indifferently from the complexity notations $\Omega(\cdot)$, $O(\cdot)$ and $\Theta(\cdot)$. The exact meaning of $O(\cdot)$ will be clear in context. For instance, under computer science notations, (1.7) would be $\|\boldsymbol{\mu}\| \geq \Theta(1)$, $\|\mathbf{E}\| \geq \Theta(p^{-1/2})$, $|\text{tr}(\mathbf{E})| \geq \Theta(\sqrt{p})$, and $\text{tr}(\mathbf{E}^2) \geq \Theta(1)$.

1.1.3.2 Asymptotic loss of pairwise distance discrimination

Under the equality case for the conditions in (1.7), consider the (normalized) Euclidean distance between two distinct data vectors $\mathbf{x}_i \in \mathcal{C}_a, \mathbf{x}_j \in \mathcal{C}_b, i \neq j$ given by

$$\frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 = \begin{cases} \frac{1}{p} \|\mathbf{z}_i - \mathbf{z}_j\|^2 + Ap^{-1/2}, & \text{for } a = b = 2 \\ \frac{1}{p} \|\mathbf{z}_i - \mathbf{z}_j\|^2 + Bp^{-1/2}, & \text{for } a = 1, b = 2, \end{cases} \quad (1.8)$$

where

$$\begin{aligned} A &= (\mathbf{z}_i^\top \mathbf{E} \mathbf{z}_i + \mathbf{z}_j^\top \mathbf{E} \mathbf{z}_j - 2\mathbf{z}_i^\top \mathbf{E} \mathbf{z}_j) p^{-1/2} \\ B &= (\mathbf{z}_j^\top (\mathbf{E} + \mathbf{E}^2/4) \mathbf{z}_j - \mathbf{z}_i^\top \mathbf{E} \mathbf{z}_j + 4\|\boldsymbol{\mu}\|^2 + 4\boldsymbol{\mu}^\top (\mathbf{z}_i - \mathbf{z}_j) + o(1)) p^{-1/2} \end{aligned}$$

are both of order $O(1)$ (and thus both $Ap^{-1/2}$ and $Bp^{-1/2}$ are of order $O(p^{-1/2})$) while the leading term $\frac{1}{p} \|\mathbf{z}_i - \mathbf{z}_j\|^2$ of (1.8) is of order $O(1)$. As such,

$$\max_{1 \leq i \neq j \leq n} \left\{ \frac{1}{p} \|\mathbf{z}_i - \mathbf{z}_j\|^2 - 2 \right\} \rightarrow 0$$

almost surely as $n, p \rightarrow \infty$ (this follows by exploiting the fact that $\|\mathbf{z}_i - \mathbf{z}_j\|^2$ is a chi-square random variable with p degrees of freedom). As a consequence, as previously claimed in (1.3),

$$\max_{1 \leq i \neq j \leq n} \left\{ \frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 - \tau \right\} \rightarrow 0$$

for $\tau = 2$ here. Besides, let us suppose first that $\text{tr } \mathbf{E} = 0$ (a situation that arises in practice if for example the data all have the same covariance or if they, or their entries $[\mathbf{x}_i]_k$, are normalized). Then, on closer inspection of (1.8), beyond this common value τ of order $O(1)$, the discriminative class information in means $4\|\boldsymbol{\mu}\|^2/p$ and in covariances $\mathbf{z}_j^\top \mathbf{E}^2 \mathbf{z}_j/(4p) \simeq \text{tr}(\mathbf{E}^2)/(4p)$ are both of order $O(p^{-1})$ while, by the central limit theorem, $\|\mathbf{z}_i - \mathbf{z}_j\|^2/p = 2 + O(p^{-1/2})$. The class information is thus largely overtaken by the random fluctuations. As a consequence, asymptotically, the pairwise distance $\|\mathbf{x}_i - \mathbf{x}_j\|^2/p$ contains *no* exploitable information (about $\boldsymbol{\mu}$ or \mathbf{E}) to distinguish if \mathbf{x}_i and \mathbf{x}_j vectors belong to the same or different classes.²

To visually confirm this joint convergence of the data distances, in Figure 1.2 we display the content of the Gaussian (heat) kernel matrix $\mathbf{K} \in \mathbb{R}^{n \times n}$ with $[\mathbf{K}]_{ij} = \exp(-\|\mathbf{x}_i - \mathbf{x}_j\|^2/(2p))$ and the associated second dominant eigenvector \mathbf{v}_2 for a two-class Gaussian mixture $\mathbf{x} \sim \mathcal{N}(\pm \boldsymbol{\mu}, \mathbf{I}_p)$ with $\boldsymbol{\mu} = [2; \mathbf{0}_{p-1}]$. For a constant $n = 500$, we take $p = 5$ in Figure 1.2a and $p = 250$ in Figure 1.2b.

²The case where $\text{tr } \mathbf{E} = O(\sqrt{p})$ leads to slightly different conclusions. There, the term $\mathbf{z}_i^\top \mathbf{E} \mathbf{z}_i/p + \mathbf{z}_j^\top \mathbf{E} \mathbf{z}_j/p \simeq 2 \text{tr } \mathbf{E}/p$ in $Ap^{-1/2}$ dominates over the term $\mathbf{z}_j^\top \mathbf{E} \mathbf{z}_j/p \simeq \text{tr } \mathbf{E}/p$ in $Bp^{-1/2}$ by a factor 2: both are of order $O(p^{-1/2})$, which is the same as the order of the noise fluctuations. It is thus possible to non-trivially extract the class information in this case, merely by comparing the norms $\|\mathbf{x}_1\|, \dots, \|\mathbf{x}_n\|$.

While the “block-structure” in the case of $p = 5$ of Figure 1.2a does agree with the finite-dimensional intuition: data vectors from the same class are “closer” to one another, corresponding to diagonal blocks with larger values (since $\exp(-x/2)$ decreases with x) than in non-diagonal blocks, this intuition collapses when large dimensional data vectors are considered. Indeed, in the large data setting of Figure 1.2b, all entries (except obviously on the diagonal) of \mathbf{K} have approximately the same value, that we now know from (1.3) is $\exp(-1)$.

This is no longer surprising to us. However, what remains surprising in Figure 1.2 at this stage of our analysis is that the eigenvector \mathbf{v}_2 of \mathbf{K} seems not affected by this (asymptotic) loss of class-wise discrimination of individual distances (particularly valid here since $\mathbf{E} = \mathbf{0}$). And spectral clustering seems to work equally well for $p = 5$ and for $p = 250$, despite the radical and intuitively destructive change in the behavior of \mathbf{K} for $p = 250$.




Figure 1.2: Gaussian kernel matrices \mathbf{K} and the second top eigenvectors \mathbf{v}_2 for small and large dimensional data $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ with $\mathbf{x}_1, \dots, \mathbf{x}_{n/2} \in \mathcal{C}_1$ and $\mathbf{x}_{n/2+1}, \dots, \mathbf{x}_n \in \mathcal{C}_2$ for $n = 5\,000$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).



1.1.3.3 Explaining kernel methods with random matrix theory

The fundamental reason behind this surprising behavior lies in the *accumulated* effect of the $n/2$ small “hidden” informative terms $\|\boldsymbol{\mu}\|^2$ and $\text{tr}(\mathbf{E}^2)$ in each class, which collectively “steer” the several top eigenvectors of \mathbf{K} . More explicitly, we shall see in the course of this monograph that the Gaussian kernel matrix \mathbf{K}

can be asymptotically expanded as

$$\mathbf{K} = \exp(-1) \left(\mathbf{1}_n \mathbf{1}_n^\top + \frac{1}{p} \mathbf{Z}^\top \mathbf{Z} \right) + f(\boldsymbol{\mu}, \mathbf{E}) \cdot \frac{1}{p} \mathbf{j} \mathbf{j}^\top + * + o_{\|\cdot\|}(1) \quad (1.9)$$

where $\mathbf{Z} = [\mathbf{z}_1, \dots, \mathbf{z}_n] \in \mathbb{R}^{p \times n}$ is a Gaussian noise matrix, $f(\boldsymbol{\mu}, \mathbf{E}) = O(1)$ and $\mathbf{j} = [\mathbf{1}_{n/2}; -\mathbf{1}_{n/2}]$ is the class-information “label” vector (as in the setting of Figure 1.2). Here ‘*’ symbolizes extra terms of marginal importance to the present discussion and $o_{\|\cdot\|}(1)$ represents terms of asymptotically vanishing operator norm as $n, p \rightarrow \infty$. The important remark to be made here is that

- (i) under this description, $[\mathbf{K}]_{ij} = \exp(-1)(1 + \mathbf{z}_i^\top \mathbf{z}_j/p) \pm f(\boldsymbol{\mu}, \mathbf{E})/p + *$ with $f(\boldsymbol{\mu}, \mathbf{E})/p \ll \mathbf{z}_i^\top \mathbf{z}_j/p = O(p^{-1/2})$; this is consistent with our previous discussion: the statistical information is *entry-wise* dominated by noise;
- (ii) from a *spectral* viewpoint, $\|\mathbf{Z}^\top \mathbf{Z}/p\| = O(1)$, as per the Marčenko-Pastur theorem [Marcenko and Pastur, 1967] discussed in Section 1.1.2 and visually confirmed in Figure 1.1, while $\|f(\boldsymbol{\mu}, \mathbf{E}) \cdot \mathbf{j} \mathbf{j}^\top/p\| = O(1)$: thus, *spectrum-wise*, the information stands on even ground with noise.

The mathematical magic at play here lies in $f(\boldsymbol{\mu}, \mathbf{E}) \cdot \mathbf{j} \mathbf{j}^\top/p$ having entries of order $O(p^{-1})$ while being a low rank (here unit rank) matrix: all its “energy” concentrates in a single non-zero eigenvalue. As for $\mathbf{Z}^\top \mathbf{Z}/p$, with larger $O(p^{-1/2})$ amplitude entries, it is composed of “essentially independent” zero mean random variables and tends to be of full rank and *spread* its energy over its p eigenvalues. Spectrum-wise, both $f(\boldsymbol{\mu}, \mathbf{E}) \cdot \mathbf{j} \mathbf{j}^\top/p$ and $\mathbf{Z}^\top \mathbf{Z}/p$ meet on even ground under the non-trivial classification setting of (1.7).

We shall see in Section 4 that things are actually not as clear-cut and in particular that not all choices of kernel functions can achieve the same non-trivial classification rates. In particular, the popular Gaussian (RBF) kernel will be shown to be largely sub-optimal in this respect.

1.1.3.4 Do real data follow small or large dimensional intuitions?

A first glimpse into this riddle, fundamental for the practical design of machine learning algorithms, is provided here in Figure 1.3. Similar to Figure 1.2 for synthetic Gaussian data, Figure 1.3 depicts the content of kernel matrices built from the MNIST [LeCun et al., 1998] and Fashion-MNIST data [Xiao et al., 2017], with $p = 28 \times 28 = 784$ and $n = 5\,000$ in both cases. In Figure 1.4, instead of raw data, we display the *features* extracted from popular deep neural networks such as VGG-16 [Simonyan and Zisserman, 2014] of the more complex CIFAR-10 images (with $p = 1024$), as well as the so-called “word-embedding” *features* from the popular word2vec method [Mikolov et al., 2013] of the GoogleNews data (with $p = 300$). In all aforementioned cases, we observe a typical large dimensional behavior (that is similar to Figure 1.2b for Gaussian data), not only on raw data but also on efficient features from modern and elaborate machine learning algorithms; even more strikingly, this behavior is consistently observed both for image and natural language data, despite their being of a fundamentally

different nature. Section 1.2.4, at the end of this introductory chapter, provides first clues which justify why this seemingly unexpected observation (recall again that, in the classical motivation behind spectral clustering methods [Ng et al., 2002], we would rather expect a behavior typical of Figure 1.2a) should in fact not be a surprise.




Figure 1.3: Kernel matrices \mathbf{K} and the second top eigenvectors \mathbf{v}_2 for MNIST (class 8 versus 9) and Fashion-MNIST data (class 5 versus 7), with $\mathbf{x}_1, \dots, \mathbf{x}_{n/2} \in \mathcal{C}_1$ and $\mathbf{x}_{n/2+1}, \dots, \mathbf{x}_n \in \mathcal{C}_2$ for $n = 5\,000$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).



1.1.4 Summarizing

In this section we discussed two simple, yet counterintuitive examples of common pitfalls in learning from large dimensional data.

In the sample covariance matrix example of Section 1.1.2, we made the important remark of the *loss of equivalence* between matrix norms in the *random matrix regime* where the data (or feature) dimension p and their number n are both large and comparable, which is at the source of many seemingly striking empirical observations. We in particular insist that, for matrices $\mathbf{A}_n, \mathbf{B}_n \in \mathbb{R}^{n \times n}$ of large sizes

$$\forall i, j, [\mathbf{A}_n - \mathbf{B}_n]_{ij} \rightarrow 0 \not\Rightarrow \|\mathbf{A}_n - \mathbf{B}_n\| \rightarrow 0 \quad (1.10)$$




Figure 1.4: Kernel matrices \mathbf{K} and the second top eigenvectors \mathbf{v}_2 for VGG-16 [Simonyan and Zisserman, 2014] features of CIFAR-10 data (“airplane” versus “bird”) and word2vec [Mikolov et al., 2013] features of GoogleNews-vectors data (“sports” versus “sales”), with $\mathbf{x}_1, \dots, \mathbf{x}_{n/2} \in \mathcal{C}_1$ and $\mathbf{x}_{n/2+1}, \dots, \mathbf{x}_n \in \mathcal{C}_2$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).



in operator norm.

We also realized, from a basic reading of the Marčenko-Pastur theorem, that the random matrix regime arises more often than one may think: while $n/p \sim 100$ may seem large enough a ratio for classical asymptotic statistics to be accurate, random matrix theory is in general a far more appropriate tool (with as much as 20% gain in precision for the estimation of the eigenvalues of sample covariances).

In Section 1.1.3, we gave a concrete machine learning application example of the message (1.10) above. We saw that, in the practically most relevant scenario of non-trivial (not too easy, not too hard) large data classification tasks, the Euclidean distance between any two data vectors “concentrates” around a constant (1.3), regardless of their respective classes. Yet, since again entry-wise convergence $[\mathbf{A}_n]_{ij} \rightarrow \tau$ does not imply operator norm convergence $\|\mathbf{A}_n - \tau \mathbf{1}_n \mathbf{1}_n^\top\| \rightarrow 0$, we understood that, thanks to a collective effect of the small but similarly “oriented” fluctuations in all the entries, spectral clustering remains valid for large dimensional problems.

Possibly most importantly, we discovered that the curse of dimensionality induced by the counterintuitive behavior of large dimensional vectors turns into an asset for mathematical analysis. In the sample covariance matrix example,

we observed that a random-matrix version of the laws of large numbers arises in the convergence of the eigenvalue distributions of large sample covariance matrices to a deterministic limiting measure. As a matter of fact, as we shall see throughout the monograph, the very fact that both p and n are large ensures a generally *fast convergence* of most quantities of practical interest for machine learning: by exploiting $np = O(n^2)$ rather than n degrees of freedom, central limit theorems may converge at $O(1/n)$ rate (instead of the classical $O(1/\sqrt{n})$).

This fast convergence rate further induces another important phenomenon, referred to as the *universality*, which ensures the robustness of the random matrix asymptotics to a vast range of distributions. Essentially, as we shall see in more detail later in this monograph, first- and second-order statistics are often sufficient to describe most asymptotic behaviors, even of complicated data models and methods. This is a first (yet not the most convincing) justification of the repeatedly observed – but quite unexpected – good match between random matrix predictions and experiments on real datasets.

In a nutshell, the fundamentally counter-intuitive yet mathematically addressable changes in behavior of large dimensional data when compared to small dimensional data have two major consequences to statistics and machine learning: (i) most algorithms, originally developed under a finite-dimensional intuition, are likely to fail (as we shall discover in this monograph, many of them do) or at least to perform inefficiently, yet (ii) by benefiting from the extra degrees of freedom offered by large data (in the dimension p), random matrix theory is apt to analyze and improve these methods, but most importantly generates a whole new paradigm for large dimensional learning.

1.2 Random matrix theory as an answer

1.2.1 Which theory and why?

1.2.1.1 A point of history

Random matrix theory originates from the work of John Wishart [Wishart, 1928] on the study of the eigenvalues of the matrix $\mathbf{X}\mathbf{X}^\top$ (now referred to as a Wishart matrix) for $\mathbf{X} \in \mathbb{R}^{p \times n}$ with $[\mathbf{X}]_{ij} \sim \mathcal{N}(0, 1)$. Wishart managed to determine a closed-form expression for the joint eigenvalue distribution of $\mathbf{X}\mathbf{X}^\top$ for every pair of p, n . Few progress however followed, as matrices with non-Gaussian entries are hardly amenable to similar analysis and, even if they were, the actual study of more elaborate functionals of $\mathbf{X}\mathbf{X}^\top$ is at best cumbersome and often simply intractable.

The works of the physicist Eugene Wigner [Wigner, 1955] gave a new impulse to the theory. Interested in the eigenvalues of symmetric matrices $\mathbf{X} \in \mathbb{R}^{n \times n}$ with independent Bernoulli entries (particle spins in his application context), Wigner opted for an *asymptotic* analysis of the eigenvalue distribution, thereby initiating the important and much richer branch of *large dimensional random matrix theory*. Despite this important inspiration, Wigner exploited standard

asymptotic statistics tools (the method of moments) to prove that the *discrete* distribution of the eigenvalues of \mathbf{X} has a *continuous* semicircle looking density in the limit (the now popular semicircular law). This approach was particularly convenient as the limiting law is simple and could be visually anticipated (which is not the case of the next-to-come Marčenko-Pastur limiting distribution of Wishart matrices).

Only until 1967 with the tour-de-force of [Marčenko and Pastur \[1967\]](#) did random matrix theory take a new dimension. Marčenko and Pastur determined the limiting spectral distribution of the sample covariance matrix model \mathbf{XX}^\top of Wishart but under relaxed conditions: $[\mathbf{X}]_{ij}$ are independent entries with zero mean and unit variance, and additional moment assumptions all relaxed in subsequent works. The independence (or weak dependence) property is key to this method. The proof exploits the powerful Stieltjes transform $\frac{1}{p} \text{tr}(\frac{1}{n} \mathbf{XX}^\top - z \mathbf{I}_p)^{-1} = \int (\lambda - z)^{-1} \mu_p(dt)$ of the *empirical spectral distribution* $\mu_p \equiv \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\frac{1}{n} \mathbf{XX}^\top)}$ of $\frac{1}{n} \mathbf{XX}^\top$, a tool borrowed from operator theory in Hilbert spaces [[Akhiezer and Glazman, 2013](#)], rather than the moments $\frac{1}{p} \text{tr}(\frac{1}{n} \mathbf{XX}^\top)^k$ (which may not converge since $\mathbb{E}[\mathbf{X}_{ij}^\ell]$ needs not be finite for $\ell > 2$).

The technical approach devised by Marčenko and Pastur was then largely embraced at the turn of the 21st century by Bai and Silverstein who, in a series of significant breakthroughs (the most noticeable of which are [[Silverstein and Bai, 1995](#), [Bai and Silverstein, 1998](#)]), extended [[Marčenko and Pastur, 1967](#)] to an exhaustive study of sample covariance matrices.

In parallel, another approach to limiting spectral analysis of large random matrices emerged as an application example of the *free probability theory* developed by [Voiculescu et al. \[1992\]](#). Free probability was born as a theory to study random variables in non-commutative algebras, such as the algebra of matrices. Rather than relying on independence assumptions as for the aforementioned Stieltjes transform method, free probability theory relies on a notion of *asymptotic freeness*. In essence, random matrices are asymptotically free if their eigenvector distribution are sufficiently isotropic with respect to each other; for instance, independent Gaussian matrices (matrices with independent Gaussian entries) are free, independent unitary matrices with isotropic eigenvector distributions are free, a deterministic matrix is free with respect to a Gaussian matrix, etc. [[Mingo and Speicher, 2017](#)].

Both free probability and the Stieltjes transform approaches have long lived hand-in-hand, and are essentially capable of proving similar results under various assumptions. A classical example, of great importance to this monograph, is that of *spiked models* (i.e., finite-rank deformation of random matrices, such as the nonzero mean sample covariance $(\mathbf{X} + \boldsymbol{\mu}\mathbf{1}_n^\top)(\mathbf{X} + \boldsymbol{\mu}\mathbf{1}_n^\top)^\top$ or the rank-one perturbed identity covariance $(\mathbf{I}_p + \ell\mathbf{u}\mathbf{u}^\top)^{\frac{1}{2}}\mathbf{XX}^\top(\mathbf{I}_p + \ell\mathbf{u}\mathbf{u}^\top)^{\frac{1}{2}}$ for \mathbf{X} with i.i.d. zero-mean entries) made popular by two key articles [[Baik and Silverstein, 2006](#)] and [[Benaych-Georges and Nadakuditi, 2012](#)], respectively based on a Stieltjes transform and a free probability approach.

These tools are largely sufficient to cover most of the basic statistical prob-

lems in random matrix theory. In particular, the often called *global regime* of random matrices: their limiting eigenvalue spectrum, the behavior of linear statistics of their eigenvalues or eigenvectors, the position of the outlying eigenvalues in spiked models, etc., are all accessible by either method. However, this is often not the case of the *local regime*: the limiting distribution of a specific eigenvalue (notably the largest and smallest, of practical interest) for which more efforts are in general needed. There, researchers have rather resorted to a finite-dimensional analysis of the joint eigenvalue distribution for the Gaussian case (in the spirit of Wishart), and carefully taken the limits of the distribution, exploiting powerful tools such as orthogonal polynomial theory [Johnstone, 2001]. We will not further discuss these approaches in the monograph, which is rather specific and not of direct use to our applications.

1.2.1.2 Resolvents, Gaussian tools, and concentration of measure theory

As we shall see throughout this monograph, realistic data and feature models necessarily contain rich statistical structures and information patterns (to be extracted by the machine learning algorithms). Typical examples include local structures (captured by convolutional filters) in image data, as well as short- and long-term dependences in time series or natural language data. In random matrix terms, this involves dealing with very structured and heterogeneous random matrix models. Although it ebbed and flowed in the past decade, the free probability approach in general requires more effort and more advanced techniques to prove the key asymptotic freeness, if possible at all. For this reason (and also because most research and results are available in the Stieltjes transform-related literature), our focus in this monograph will be on the range of methods surrounding the Stieltjes transform approach.

More exactly, the central object of study in this monograph is the so-called *resolvent* of the (almost always symmetric, or Hermitian in the complex case) random matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ under investigation, that we shall often denote $\mathbf{Q}_{\mathbf{X}}(z)$ or simply $\mathbf{Q}(z)$, and that is defined, for all $z \in \mathbb{C}$ not in the eigenspectrum of \mathbf{X} (i.e., not coinciding with an eigenvalue of \mathbf{X}), by

$$\mathbf{Q}_{\mathbf{X}}(z) \equiv (\mathbf{X} - z\mathbf{I}_n)^{-1}. \quad (1.11)$$

The resolvent is a rich mathematical object that gives access to

- the eigenvalue distribution $\mu_{\mathbf{X}} \equiv \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i(\mathbf{X})}$ of \mathbf{X} through the (inverse) Stieltjes transform relation (for all $a, b \notin \{\lambda_1(\mathbf{X}), \dots, \lambda_n(\mathbf{X})\}$)

$$\int_a^b \mu_{\mathbf{X}}(d\lambda) = \lim_{\epsilon \downarrow 0} \int_a^b \frac{1}{\pi} \Im[m_{\mathbf{X}}(x + i\epsilon)] dx,$$

with

$$m_{\mathbf{X}}(z) \equiv \int \frac{\mu_{\mathbf{X}}(d\lambda)}{\lambda - z} = \frac{1}{n} \sum_{i=1}^n \frac{1}{\lambda_i(\mathbf{X}) - z} = \frac{1}{n} \operatorname{tr} \mathbf{Q}_{\mathbf{X}}(z);$$

- functionals of these eigenvalues $\frac{1}{n} \sum_{i=1}^n f(\lambda_i(\mathbf{X}))$ through Cauchy's integral identity (Theorem 2.2)

$$\frac{1}{n} \sum_{i=1}^n f(\lambda_i(\mathbf{X})) = -\frac{1}{2\pi i n} \oint_{\Gamma} f(z) \operatorname{tr} \mathbf{Q}_{\mathbf{X}}(z) dz,$$

for $\Gamma \subset \mathbb{C}$ a positively oriented contour in the complex plane surrounding all the $\lambda_i(\mathbf{X})$'s and $f(z)$ complex analytic in a neighborhood of the “inside” of Γ ;

- the eigenvectors and subspaces of \mathbf{X} , again through Cauchy's integral relation

$$\mathbf{u}_i(\mathbf{X}) \mathbf{u}_i(\mathbf{X})^T = -\frac{1}{2\pi i} \oint_{\Gamma_{\lambda_i(\mathbf{X})}} \mathbf{Q}_{\mathbf{X}}(z) dz,$$

for $(\lambda_i(\mathbf{X}), \mathbf{u}_i(\mathbf{X}))$ an eigenpair of \mathbf{X} and $\Gamma_{\lambda_i(\mathbf{X})}$ a positively oriented contour surrounding only $\lambda_i(\mathbf{X})$.

As such, the resolvent plays a key role in the analysis of spectral methods such as (kernel) spectral clustering or graph-based community detection, in which case the top eigenvectors of some underlying matrix are exploited.

In addition, the resolvent is a fundamental object that frequently appears in the solutions to linear regression problems (for machine learning applications, in least squares support vector machines, random features and kernel ridge regressions, echo state neural networks, etc.), or to random walk and graph-based semi-supervised learning methods. They will also be shown to appear naturally in not-immediately-related machine learning problems, such as in large dimensional nonlinear regression (logistic or robust M-regression).

The core of the random matrix approach devised in this monograph consists in determining, for various statistical models of random matrices \mathbf{X} a *deterministic equivalent* $\bar{\mathbf{Q}}(z)$ for $\mathbf{Q}(z) = \mathbf{Q}_{\mathbf{X}}(z)$, that is a deterministic matrix $\bar{\mathbf{Q}}(z)$ such that

$$u(\mathbf{Q}(z) - \bar{\mathbf{Q}}(z)) \xrightarrow{a.s.} 0, \quad \text{or} \quad u(\mathbb{E}[\mathbf{Q}(z)] - \bar{\mathbf{Q}}(z)) \rightarrow 0$$

for all 1-Lipschitz linear mapping $u : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$. Of particular interest are the functions $u(\mathbf{X}) = \frac{1}{n} \operatorname{tr}(\mathbf{A}\mathbf{X})$ for $\|\mathbf{A}\| \leq 1$, and $u(\mathbf{X}) = \mathbf{a}^T \mathbf{X} \mathbf{b}$ for $\|\mathbf{a}\|, \|\mathbf{b}\| \leq 1$.³

As an example, in the setting of the Marčenko-Pastur law where the random matrix of interest is $\frac{1}{n} \mathbf{X} \mathbf{X}^T$ with $\mathbf{X} \in \mathbb{R}^{p \times n}$ having i.i.d. zero mean and unit variance entries, the resolvent

$$\mathbf{Q}(z) = \left(\frac{1}{n} \mathbf{X} \mathbf{X}^T - z \mathbf{I}_p \right)^{-1}$$

³Here \mathbf{A} and \mathbf{a}, \mathbf{b} must be understood as “sequences” of deterministic matrices (or vectors) of growing size but with controlled norm; in particular, \mathbf{A} and \mathbf{a}, \mathbf{b} , being deterministic, *cannot* depend on \mathbf{X} (in which case the convergence results may fail: take for instance $\mathbf{a} = \mathbf{b}$ some eigenvector of \mathbf{X} to be convinced).

admits

$$\bar{\mathbf{Q}}(z) = m_\mu(z)\mathbf{I}_p, \quad m_\mu(z) = \int \frac{\mu(d\lambda)}{\lambda - z}, \quad \text{for } \mu \text{ defined in (1.2),}$$

as a deterministic equivalent. Thus, in particular, $\frac{1}{p} \operatorname{tr} \mathbf{Q}(z) - m_\mu(z) \xrightarrow{a.s.} 0$ and $\mathbf{a}^\top \mathbf{Q}(z) \mathbf{b} - m_\mu(z) \mathbf{a}^\top \mathbf{b} \xrightarrow{a.s.} 0$ for deterministic $\mathbf{a}, \mathbf{b} \in \mathbb{R}^p$ of bounded Euclidean norm.

Consequently, the resolvent (Stieltjes transform) approach simultaneously involves notions from three distinct mathematical areas:

- *linear algebra*, and particularly the exploitation of inverse matrix lemmas, the Schur complement, interlacing and low rank perturbation identities [Horn and Johnson, 2012];
- *complex analysis* (the resolvent $\mathbf{Q}(z)$ is a complex analytic matrix-valued function) and particularly the theory of analytic functions, contour integrals and residue calculus [Stein and Shakarchi, 2003];
- *probability theory*, and most specifically notions of convergence, central limit theory, the method of moments, etc. [Billingsley, 2012]. Depending on the underlying random matrix assumptions (independence of entries, Gaussianity, concentration properties), different random matrix-adapted techniques (among others and variations) will be discussed in this monograph: the Gaussian tools developed by Pastur, relying on Stein's lemma and the Nash-Poincaré inequality [Pastur and Shcherbina, 2011], the Bai-Silverstein inductive method [Bai and Silverstein, 2010], the concentration of measure framework developed by Ledoux [2005] and applied to random matrix endeavors successively by El Karoui [2009], Vershynin [2012], and Louart and Couillet [2018], or the double leave-one-out approach devised by El Karoui et al. [2013].

The aforementioned tools are in general used together in a *perturbation approach* in the sense that they exploit the fact that, by eliminating a row or a column (say here both row and column i) of a large random matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ to obtain $\mathbf{X}_{-i} \in \mathbb{R}^{(n-1) \times (n-1)}$, the resolvent $\mathbf{Q}_{-i}(z) = (\mathbf{X}_{-i} - z\mathbf{I}_{n-1})^{-1}$ can be related to the original resolvent $\mathbf{Q}(z)$ through both linear algebraic relations and *asymptotically comparable* statistical behaviors. For instance, in the case of symmetric \mathbf{X} with i.i.d. (properly normalized) entries, it is not difficult to show that $m_{\mathbf{X}}(z) = m_{\mathbf{X}_{-i}}(z) + O(n^{-1})$.

In this regard, Pastur's Gaussian method manages, for models of \mathbf{X} involving Gaussianity (e.g., \mathbf{X} has Gaussian entries or its entries are functions of Gaussian random variables), to obtain asymptotic relations for $\mathbb{E}\mathbf{Q}(z)$. Interpolation methods may then be used to extrapolate the results beyond the Gaussian case. The Bai-Silverstein inductive method, on the opposite, is not restricted to matrices with Gaussian entries but is restricted to the specific analysis of either trace forms $\operatorname{tr} \mathbf{A}\mathbf{Q}(z)$ or bilinear forms $\mathbf{a}^\top \mathbf{Q}(z) \mathbf{b}$ that need be treated

individually (it also suffers to handle exotic forms of dependence within \mathbf{X}). The concentration of measure approach is quite versatile: by merely restricting the matrix under study to be constituted of *concentrated random vectors* (so in particular, Lipschitz maps of standard Gaussian random vectors or of vectors with i.i.d. entries), it allows one to study simultaneously the fluctuations of all Lipschitz (or only all linear) functionals of $\mathbf{Q}(z)$ under light conditions on \mathbf{X} .

1.2.2 The double asymptotics: turning the curse of dimensionality into a dimensionality blessing

1.2.2.1 Why random matrix theory to study the large n, p regime?

Although we have previously made a point that modern data processing and learning involve large dimensions (numerous data, large sample sizes, large number of system parameters) and that large dimensional statistics are a natural class of mathematical tools to turn to, why should one invest in random matrix theory rather than, say, statistical physics,⁴ non-asymptotic random matrix theory,⁵ or compressive sensing?⁶ Large dimensional random matrix theory, as we introduce it in this monograph, has two key distinctive features, making it simultaneously more powerful and more versatile than these alternative tools:

- (i) Unlike non-asymptotic random matrix theory and compressive sensing methods which mostly aim at *bounding* key quantities (from a rather *qualitative* standpoint), large dimensional random matrix theory is able to provide *precise and quantitative* (asymptotically exact) approximations for a host of quantities, defined as functionals of random matrices. As a matter of fact, non-asymptotic random matrix theory is more flexible in its not constraining the system dimensions (p, n) and latent variables (data statistics, model hyperparameters) to increase at a controlled rate. Large dimensional random matrix theory, on the opposite, imposes a controlled

⁴Statistical physics and statistical mechanics are powerful tools to map large dimensional data problems into physics-inspired problems of “interacting particles” [Mézard and Montanari, 2009]. In the early 2000’s, statistical physics has brought inspiring ideas and powerful (but unfortunately often unreliable, since non-rigorous) tools for the analysis of wireless communication and information-theoretic problems, before being caught up by more solid and versatile mathematical techniques. Today, statistical physics has an edge on the study of *sparse* (graph-based) machine learning problems for which random matrix theory still struggles to offer a sound theory.

⁵The recent field of non-asymptotic random matrix theory is based on concentration inequality methods and aims, as such, to provide bounds rather than exact (deterministic) asymptotics on various random matrix quantities [Vershynin, 2018]. This set of concentration inequalities should not to be confused with the *concentration of measure theory* [Ledoux, 2005]: concentration inequalities form a restricted subset of the theory by proving statistical bounds on specific quantities.

⁶Compressive sensing revolves around the assumption that large (p) dimensional data often arise from a manifold in \mathbb{R}^p of much lower intrinsic dimension: under this assumption, the curse of dimensionality (when $p \sim n$ or even $p \gg n$) vanishes if one manages to retrieve the (often unknown) low-dimensional manifold. As an aftermath of the seminal work by Candes and Tao [2005], compressive sensing was possibly the first major breakthrough in the modern field of large dimensional statistical machine learning.

growth on the dimensions *and consequently* on the model statistics to enforce non-trivial limiting behavior. The ensuing drawback of this allowed flexibility is that only qualitative bounds can be obtained on the system behavior, which at best provides “rules of thumbs” and order of magnitudes on the performance of given algorithms. Large dimensional random matrix theory, by providing exact asymptotics, allows one to finely track the system behavior and opens the possibility to improve its (also fully traced) performance.

- (ii) Modern advances in large dimensional random matrix theory, as opposed to statistical physics notably, further provide results for rather generic and complex system models: matrix models involving nonlinearities (kernels, activation functions), structural data dependence (non-identity covariances, heterogeneous mixture models, models of concentrated random vectors with strong nonlinear dependence). These key features bring the random matrix tools much closer to practical settings and algorithms. As such, not only does random matrix theory provide a precise understanding of the behavior of key algorithms in machine learning, but it also predicts their behavior when applied to extremely realistic data models.

These two advantages are decisive to the analysis, improvement and proposition of new machine learning algorithms.

1.2.2.2 The case of machine learning

The major technical difficulty that has long held many machine learning away from *precise quantitative* analysis and theoretical comprehension relates to the *nonlinearity* involved in feature extraction (nonlinear kernels, nonlinear activation functions in neural networks), to the *implicit* nature of some methods (as simple as the logistic regression), and eventually to the difficulty of a proper (statistical) modeling of *complex* realistic data of various natures (starting with images).

An all-encompassing example of these difficulties could be summarized as the following classical problem:

Problem. Determine the *exact* classification performance of logistic regression for n observations of p -dimensional random feature vectors extracted from a set of two-class images (say, images of dogs versus images of cats).

In the conventional wisdom of statistical machine learning, one cannot conceive to solve this problem in an *exact* and *qualitative* manner: the input data (real images) are not easily modeled, the nonlinear features extracted from those data are complex mathematical objects (even in the case where the original data could be modeled as simple Gaussian random vectors), and the logistic regression is an implicit optimization method not easily amenable to mathematical analysis.

We shall demonstrate throughout this monograph that random matrix theory provides a satisfying answer to all these difficulties at once and can actually

solve the **Problem**. This is made possible by the powerful joint *universality* and *determinism* effects brought by large dimensional data models and treatments.

Specifically, in the random matrix regime where n, p grow large at a controlled rate, the following key properties arise:

- *fast asymptotic determinism*: the law of large numbers and the central limit theorem tell us that the average of n i.i.d. random variables converges to a deterministic limit (e.g., the expectation) at an $O(1/\sqrt{n})$ speed. By gathering independence (or degrees of freedom) both in the sample dimension p and size n , functionals of large random matrices (even complex functionals, such as the average of functions of their eigenvalues) also converge to deterministic limits, but at an increased speed of up to $O(1/\sqrt{np})$ which, for $n \sim p$, is $O(1/n)$. In machine learning problems, performance may be expressed in terms of misclassification rates or regression errors (i.e., averaged statistics of sometimes involved random matrix functionals) and can thereby be predicted with high accuracy, even for not too large dimensional datasets;
- *universality with respect to data models*: similarly, again consistently with the law of large numbers and the central limit theorem in the large- n alone setting, the above asymptotic deterministic behavior at large n, p is in general independent of the underlying distribution of the random matrix entries. This phenomenon, referred to in the random matrix literature as *universality*, predicts notably that the asymptotic statistics of even complex machine learning procedures depend on the input data only via the first- and second-order statistics; this is a major distinctive feature when compared to the fixed- p and large- n regime, where the asymptotic performance of algorithms, when accessible, would in general depend on the exact p -dimensional distribution of the data;⁷
- *universality with respect to algorithm nonlinearities*: when nonlinear methods are considered, the nonlinear function f (e.g., the kernel function or the activation function) gets involved in the large dimensional algorithm performance only via a few parameters (e.g., its derivatives $f(\tau), f'(\tau), \dots$ at a precise location τ , its “moments” $\int f^k \mu$ with respect to the Gaussian measure μ , or more elaborate scalars solution to a fixed-point equation involving f). For instance, in the case of kernel random matrices of the type $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$, only the first three derivatives of the kernel function f at the “concentration” point $\tau = \lim_p \|\mathbf{x}_i - \mathbf{x}_j\|^2/p$ matter; the performance of random neural networks depends on the nonlinear activation function $\sigma(\cdot)$ solely through its first Hermite coefficients (i.e., its Gaussian

⁷Compare for instance [Luxburg et al., 2008] on the fixed- p large- n asymptotics of spectral clustering (the main result of which contains nonlinear expressions of the input data distribution) to [Couillet and Benaych-Georges, 2016] on the large p, n asymptotics of the same problem (the main result of which only involving linear and quadratic forms of the statistical mean and covariances of the data, irrespective of the input data distribution, as further confirmed by Seddik et al. [2019]).

moments); in implicit optimization schemes (such as logistic regression), the solution “concentrates” with predictable asymptotics which, despite the nonlinear and implicit nature of the problem, only depend on a few scalar parameters of the logistic loss function. This, together with the asymptotic deterministic behavior of the linear (eigenvalue or eigenvector) statistics discussed above, gives access to the performance of a host of *nonlinear* machine learning algorithms.

- *tractable real data modeling*: possibly the most important aspect of large dimensional random matrix analysis in machine learning practice relates to the counter-intuitive fact that, as p, n grow large, machine learning algorithms tend to treat real data as if they were mere Gaussian mixture models. This statement, to be discussed thoroughly in the subsequent sections, is both supported by empirical observations (with most theoretical findings derived for Gaussian mixtures observed to fit the performances retrieved on real data) and by the theoretical fact that some extremely realistic datasets (in particular artificial images created by the popular generative adversarial networks, or GANs) are by definition *concentrated random vectors* which are (i) amenable to (and in fact extremely well-suited for) random matrix analysis, and (ii) proven to behave as if they were mere Gaussian mixtures.

In a word, in large dimensional problems, data no longer “gather” in groups and do not really “spread” all over their large ambient space neither. But, by accumulation of degrees of freedom, they rather concentrate within a thin lower-dimensional “layer”. Each scalar observation of the data, even through complicated functions (regressors, classifiers for machine learning applications), tends to become deterministic, predictable, and simple functions of first-order statistics of the data distribution. Random matrix theory exploits these effects and is thus able to answer seemingly inaccessible machine learning questions.

1.2.3 Analyze, understand, and improve large dimensional machine learning methods

One of the first elementary objectives of this monograph is to demonstrate that, in a large dimensional and numerous data setting, many standard machine learning (low dimensional) intuitions tend to collapse. As a result, many of the algorithms originally designed for small data sizes fail to perform as expected. Some of these algorithms will be shown to remain valid but for rather unexpected reasons. Some of them will be proven suboptimal, quite largely so sometimes. Finally, some of them will be shown to completely fail to meet their objectives and in need for an adaptation or a complete change of paradigm.

In a second part, the monograph will further show that this “large dimensional” regime, which one may think synonymous to thousands or millions in dimension and sample size, is in reality already visible in much smaller data sizes than the earliest researchers in applied random matrix theory could anticipate.

And, perhaps more importantly, that a large class of “real data” naturally falls under the random matrix theory umbrella.

Our argumentation line and every single treatment of machine learning algorithm analysis and improvement proceed along the following steps: one needs to (i) conceive the limitations of low dimensional intuitions and understand the reach of large dimensional intuitions, (ii) capture the behavior of the main mathematical objects at play in machine learning on large dimensional models so as to (iii) include these objects in a mathematical framework for performance analysis, and (iv) foresee means of improvement based on the newly acquired large dimensional intuitions and mathematical understanding.

In the remainder of this subsection, we will illustrate the above four-step methodology with the examples of kernel methods and the very related random feature maps (which may alternatively be seen as a two-layer neural network model with random weights).

1.2.3.1 From low to large dimensional intuitions

Most of the manuscript focuses on large dimensional data vectors or graph models. By large dimensional, we refer to random vectors $\mathbf{x} \in \mathbb{R}^p$ “built from” numerous (of order $O(p)$) independent degrees of freedom. That is, as opposed to the compressive sensing paradigm [Donoho, 2006], we do not impose the existence of a low dimensional representation of the data.⁸

From this viewpoint, the simplest mixture data model is the Gaussian mixture model $\mathbf{x} \sim \mathcal{N}(\pm\boldsymbol{\mu}, \mathbf{I}_p)$. As we saw previously, for p small (say, $p = 2$ or $p = 3$), classifying n samples of the mixture is easily visualized as grouping two stacks of data: one gathered around $\boldsymbol{\mu} \in \mathbb{R}^p$, the other around $-\boldsymbol{\mu}$. Most of (low dimensional) machine learning algorithms are anchored in this mentally convenient visualization. But the large dimensional image is different. Gaussian vectors $\mathbf{x} \in \mathbb{R}^p$ have a norm of order $\|\mathbf{x}\| \sim O(\sqrt{p})$ but a spread of order $\|\mathbf{x}\| - \mathbb{E}[\|\mathbf{x}\|] \sim O(1)$ and non-trivial classification can be performed as long as $\|\boldsymbol{\mu}\|$ is no smaller than $O(1)$. The mental image is thus one of two spheres in \mathbb{R}^p with an extremely large radius (of order $O(\sqrt{p})$) around which the data of both classes accumulate. Figure 1.5 provides a comparative picture for small versus large dimensions p .

With this image in mind, the Euclidean distance paradigm is shifted: for small p , the information lies in the typical distance from one data point to a “centroid”; for large p , the centroid is far from all data points (it lives in an “empty” region of the space) and the class information is summarized in the accumulated small deterministic deviations of all data points from the class;

⁸The *statistical* information contained in the data such as the mean $\mathbb{E}[\mathbf{x}]$ can be sparse, but the practical large dimensional data vectors must randomly “fluctuate” with sufficiently many degrees of freedom around their possibly low dimensional manifold structure. The large dimensional random fluctuation of the data is essential to produce a statistically “robust” behavior of the algorithms and is key to establishing mathematical convergence in the large n, p regime.




Figure 1.5: Visual representation of classification in (left) small and (right) large dimensions.

this deviation is not (asymptotically) visible for any data vector but can be inferred collectively from the large data matrix.

Consequently, machine learning algorithms based on evaluations of Euclidean distances $\|\mathbf{x}_i - \mathbf{x}_j\|$, inner products $\mathbf{x}_i^\top \mathbf{x}_j$, nonlinear activations $\sigma(\mathbf{w}^\top \mathbf{x}_i)$ and regressions $f(\boldsymbol{\beta}^\top \mathbf{x}_i)$, etc. of data \mathbf{x}_i or data pairs $\mathbf{x}_i, \mathbf{x}_j$ structurally behave differently in large dimensions.

1.2.3.2 Core random matrices in machine learning algorithms

Be it in a supervised, semi-supervised or unsupervised context, machine learning algorithms essentially consist in extracting structural information from some available set of data $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathcal{X}$: this is done in general via one-to-one comparisons of the data. At the heart of most algorithms we notably find affinity matrices of the type

$$\mathbf{K} \equiv \{\kappa(\mathbf{x}_i, \mathbf{x}_j)\}_{i,j=1}^n \in \mathbb{R}^{n \times n} \quad (1.12)$$

where $\kappa : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ evaluates the closeness or affinity between \mathbf{x}_i and \mathbf{x}_j . For graphs, the data \mathbf{x}_i are merely the nodes (or vertices) of the graph, and $\kappa(\mathbf{x}_i, \mathbf{x}_j) = w_{ij}$ is thus the weight of the edge (i, j) which may be real or binary (i.e., $w_{ij} \in \{0, 1\}$ depending on whether node i attaches to node j).

For $\mathcal{X} = \mathbb{R}^p$ and \mathbf{x}_i statistically distributed, this naturally gives rise to a family of *kernel random matrices*, among which are inner-product kernel random matrices with $\kappa(\mathbf{x}_i, \mathbf{x}_j) = f(\mathbf{x}_i^\top \mathbf{x}_j)$, distance-based kernel random matrices with $\kappa(\mathbf{x}_i, \mathbf{x}_j) = f(\|\mathbf{x}_i - \mathbf{x}_j\|^2)$ and correlation random matrices with $\kappa(\mathbf{x}_i, \mathbf{x}_j) = \mathbf{x}_i^\top \mathbf{x}_j / (\|\mathbf{x}_i\| \cdot \|\mathbf{x}_j\|)$. In the first case, f is often taken to be either linear $f(t) = t$ (therefore giving rise to sample covariance matrix models), a polynomial $f(t) = a_k t^k + \dots + a_0$, or of a sigmoid type such as the logistic function $f(t) = (1 + e^{-x})^{-1} - 1/2$ or the hyperbolic tangent $f(t) = \tanh(t)$. In the second case, f can be either linear (and we obtain a Euclidean distance matrix [Dokmanic et al.,

2015]) or more often $f(t) = \exp(-t/(2\sigma^2))$ for some $\sigma > 0$, which is referred to as the *heat kernel*, the *Gaussian kernel*, or even as the *radial basis function kernel*.

When the \mathbf{x}_i 's themselves are not directly separable in their ambient space, they are conventionally mapped into a *feature space* in which they become separable. As feature extraction is possibly the single most important but usually hardest task in machine learning, it comes in a variety of forms. Kernel matrices of the type (1.12) typically plays the role of a feature extraction method which maps the data points into a *reproducing kernel Hilbert space* (RKHS) [Schölkopf and Smola, 2018]. Another closely related, yet equally popular, approach, is random extraction by means of *random feature maps*, which consist in operating $\sigma(\mathbf{W}\mathbf{x})$ for some (usually randomly and independently drawn) matrix $\mathbf{W} \in \mathbb{R}^{N \times p}$ and some nonlinear function $\sigma : \mathbb{R}^p \rightarrow \mathbb{R}^N$ applying entry-wise, i.e., $\sigma(\mathbf{y}) = [\sigma_0(y_1), \dots, \sigma_0(y_N)]^\top$ for some $\sigma_0 : \mathbb{R} \rightarrow \mathbb{R}$ which, with a slight abuse of notation, we simply call σ . Among random feature maps, the most popular is the *random Fourier feature* method proposed in [Rahimi and Recht, 2008] for which $\sigma(t) = \exp(-it)$ (so, formally, $\sigma(\mathbb{R}) \subset \mathbb{C}$ rather than \mathbb{R} in this case).

Neural networks operate likewise. Every size- N layer (that contains N neurons) of a neural network operates $\sigma(\mathbf{W}\mathbf{x})$ for an input \mathbf{x} , a linear mapping $\mathbf{W} \in \mathbb{R}^{N \times p}$ (the neural weights to be learned) and a nonlinear *activation function* $\sigma : \mathbb{R} \rightarrow \mathbb{R}$.⁹ In this setting, σ is usually taken to be a sigmoid function (the logistic function, the tanh or the Gaussian erf function) or, more recently, the rectified linear unit (ReLU) function $\sigma(t) = \max(0, t)$.

Collecting the data in $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$, the sample covariance matrix of the random features of the data then reduces to the Gram matrix

$$\Phi \equiv \sigma(\mathbf{W}\mathbf{X})^\top \sigma(\mathbf{W}\mathbf{X}) \quad (1.13)$$

which is also a central object of interest in this monograph.

The aforementioned kernels and Gram matrices of feature maps are actually much interrelated. For instance, the random Fourier features $\sigma(\mathbf{W}\mathbf{x})$, with $\sigma(t) = \exp(-it)$ and \mathbf{W} with i.i.d. standard Gaussian entries, i.e., $\mathbf{W}_{ij} \sim \mathcal{N}(0, 1)$, are known to have the fundamental property

$$\mathbb{E}_{\mathbf{W}}[\sigma(\mathbf{W}\mathbf{x})^\top \sigma(\mathbf{W}\mathbf{y})] \equiv \exp\left(-\frac{1}{2}\|\mathbf{x} - \mathbf{y}\|^2\right)$$

so that random Fourier features are intricately connected to Gaussian kernel matrices. This property ensures in particular that the Gaussian kernel $\kappa(\mathbf{x}, \mathbf{y}) = \exp(-\|\mathbf{x} - \mathbf{y}\|^2/2)$ is a *nonnegative definite kernel* in the sense that $\mathbf{K} = \{\kappa(\mathbf{x}_i, \mathbf{x}_j)\}_{i,j=1}^n$ is a nonnegative definite matrix (for any n and any set of $\mathbf{x}_1, \dots, \mathbf{x}_n$), a particularly convenient property in both theory and practice. An important subclass of kernel functions, referred to as Mercer kernels [Schölkopf

⁹Sometimes an additional *bias* term is considered and the network operates $\sigma(\mathbf{W}\mathbf{x}) + \mathbf{b}$ for some $\mathbf{b} \in \mathbb{R}^N$ also to be learned.

and Smola, 2018], share this nonnegative definiteness property and have long been privileged in machine learning. We shall see in this monograph that, from a large dimensional perspective, Mercer kernels are in general suboptimal and that simple but less intuitive choices of κ can largely outperform these conventional kernels.

A large body of machine learning algorithms (spectral clustering, linear or logistic regression, support vector machines and neural networks) relates, one way or another, to the aforementioned *global properties* (eigenvalues, content of dominant eigenvectors, linear or nonlinear functionals of the resolvent) of the above matrices \mathbf{K} or Φ . A systematic statistical analysis of these global properties for all finite p, n, N is however often out of reach, even for the simplest standard Gaussian modeling of the data.

In this monograph, we will show that random matrix theory manages to leverage the large dimensional nature of both the data and the learning systems (i.e., large n, p, N), to tackle this statistical analysis. We will see in particular that several conventional models for \mathbf{K} can be “Taylor-expanded” under the form of matrices involving only first- and second-order moments. The Gram matrix Φ cannot be directly Taylor-expanded in this way (it will be “Hermite-polynomially-expanded”) but will also be shown to behave as a kernel random matrix and be decomposed as the sum of more elementary random matrices, the statistical properties of which also become tractable in the large dimensional regime.

In short, the intractable matrices \mathbf{K} and Φ will be approximated by tractable ersatz $\tilde{\mathbf{K}}$ and $\tilde{\Phi}$ which behave asymptotically the same in the sense that

$$\|\mathbf{K} - \tilde{\mathbf{K}}\| \xrightarrow{a.s.} 0, \quad \|\Phi - \tilde{\Phi}\| \xrightarrow{a.s.} 0,$$

in *operator norm* as $n, p, N \rightarrow \infty$ at a similar rate. These matrices $\tilde{\mathbf{K}}$ and $\tilde{\Phi}$ will allow for further and deeper mathematical analysis.

1.2.3.3 Performance analysis: spectral properties and functionals

In a classification context, where conventionally $\mathbf{x}_i \in \mathbb{R}^p$ belongs to one of the k classes $\mathcal{C}_1, \dots, \mathcal{C}_k$ with $k \ll n$ (the number of data samples) and thus $k \ll p$ whenever $p \sim n$, the approximation matrices $\tilde{\mathbf{K}}$ and $\tilde{\Phi}$ will often be shown to take a *spiked random matrix* form. That is, for instance,

$$\tilde{\mathbf{K}} = \mathbf{Z} + \mathbf{P}$$

where $\mathbf{Z} \in \mathbb{R}^{n \times n}$ is a random symmetric matrix in general having entries of zero mean and rather uniform variances, while $\mathbf{P} \in \mathbb{R}^{n \times n}$ is a *low rank* (often related to k) matrix comprising the statistical information about the data-class associations and the statistical properties of the classes.

These spiked random matrix models have been extensively studied and it is possible to extract much information about them. In particular, the dominant eigenvectors of $\tilde{\mathbf{K}}$ are known to relate to the eigenvectors of \mathbf{P} (which carry

the sought-for data-class information) whenever a *phase transition* threshold is exceeded.

In a regression setting where the \mathbf{x}_i 's are assumed identically distributed, the regression vector $\boldsymbol{\beta} \in \mathbb{R}^p$ of interest is a certain functional of \mathbf{K} or Φ . For instance, a random feature regression from the observations $\mathbf{X} \in \mathbb{R}^{p \times n}$ to the desired outputs $\mathbf{y} \in \mathbb{R}^n$ entails the regression vector

$$\boldsymbol{\beta} = \sigma(\mathbf{W}\mathbf{X})(\Phi + \gamma\mathbf{I}_n)^{-1}\mathbf{y}$$

which is thus an (indirect) function of the resolvent $\mathbf{Q}_\Phi(-\gamma) = (\Phi + \gamma\mathbf{I}_n)^{-1}$ of Φ for a certain $\gamma > 0$. Random matrix theory possesses tools to analyze the statistical properties of such vectors $\boldsymbol{\beta}$ as well.

Least squares support vector machines and most conventional algorithms of graph-based semi-supervised learning relate to functionals of the same type. This also holds true (yet less directly) for nonlinear (e.g., logistic) regression where $\boldsymbol{\beta}$ is *implicitly defined* as a function of \mathbf{Q}_Φ . Similarly, in their plain form, support vector machines can be seen as nonlinear regression schemes which also fall within this scope.

Since eigenvalues, eigenvectors and regressor statistics are at the core of machine learning algorithm performance, once these central quantities are accessible, the actual (asymptotic) classification error rates, mean square error of regression, etc. become also accessible. It is important to point out here that not only bounds on performance but *actual accurate estimators* of the performance are provided. Under a random matrix framework, a *precise* characterization of the anticipated performance (as well as its error margins) for the above algorithms becomes available.

Since these performance indicators depend on the various hyperparameters of the problem, themselves being quantifiable from data statistics, in many scenarios, it becomes possible to fine-tune the algorithms without resorting to cross-validation procedures. We shall notably see how some simple instances of neural networks can be fairly well understood: why the rectifier $\max(t, 0)$ is a convenient choice, how the activation function and the data statistics mix up, etc. We will also understand that kernel methods do not function as one may think they should, and that there exists an elegant interplay between data statistics and the successive derivatives of the kernel function at a precise position.

1.2.3.4 Directions of improvement and new ideas

Due to the complete change of paradigm when comparing data from a small versus a large dimensional perspectives, the overall behavior and the ensuing performance of the studied algorithms are often tainted, when large dimensional data are handled.

We shall notably see in the course of the monograph that the conventional heat (or Gaussian) kernel used in various classification contexts is largely sub-optimal. We shall also see that most graph-inspired semi-supervised learning

algorithms of the literature fail to properly accomplish their requested task for n, p large and comparable; yet, we will show that the so-called PageRank approach [Avrachenkov et al., 2012] happens not to fail, although the fundamental reasons behind its non-degrading performance are at odds with the initial inspiration for the method; but most importantly, this popular approach will also be shown to perform quite far from optimal and in particular not to be capable of benefiting from a large addition of unlabeled data. This observation entails the very unpleasant property that purely unsupervised methods tend to outperform semi-supervised ones when the number of unlabeled data is quite large.

For all these applications, the monograph will list a set of recommendations and improved methods which are tailored to large (as well as not so large) dimensional data learning. Among others, optimal but quite counterintuitive kernel functions will be introduced, new regularization procedures for supervised and semi-supervised learning will be discussed that particularly defeat the curse of dimensionality in semi-supervised learning (by fully exploiting the addition of unlabeled data), and some further light on the design of neural networks will be cast.

1.2.4 Exploiting universality: from large dimensional Gaussian vectors to real data

Before delving into the core of the manuscript, we conclude this section by further elaborating on the universality phenomenon briefly discussed above, which is of much greater importance than one may anticipate.

First, let us recall that most random matrix results derived in the literature, even the most recent ones on machine learning applications (discussed in this monograph), are based on the assumption of data either arising from (possibly a mixture of) Gaussian distributions or represented by random vectors with independent entries. These models are generally deemed unsuitable to mimic real data and we will not claim otherwise. It is a fact that real data, such as images, are largely more complex than mere Gaussian vectors.

Yet, what we do claim is that *scalar observations* (regressor or classifier outputs, misclassification rates, etc.) obtained from large dimensional and numerous data *tend to behave as if the data were Gaussian* (mixtures) in the first place. This is a fundamental disruption from small dimensional statistics that random matrix theory structurally exploits: rather than assuming data as fixed entities living in a complex manifold, random matrix theory mostly exploits their numerous degrees of freedom which, by universality, induce deterministic behavior in the large dimensional limit, thus *independently* of the underlying vector data distribution.

We justify this claim below with both empirical and theoretical arguments.

1.2.4.1 Theory versus practice

Our first argument follows after numerous comparative experiments made between theoretical findings on Gaussian versus real data. Indeed, although mostly derived under simple and seemingly unrealistic Gaussian mixture models, many theoretical results mentioned above show an *unexpected close match* when applied to popular real-world (sometimes not so) large dimensional datasets, such as the MNIST handwritten-digit dataset [LeCun et al., 1998], the related Fashion-MNIST [Xiao et al., 2017], Kannada-MNIST [Prabhu, 2019] and Kuzushiji-MNIST [Clanuwat et al., 2018] datasets, the German Traffic Sign dataset [Houben et al., 2013], deep neural network features of the now popular ImageNet dataset [Deng et al., 2009] used for state-of-the-art machine learning and computer vision applications, as well as numerous financial and electroencephalography (EEG) time series datasets. In particular, while most elementary machine learning methods discussed in this monograph cannot be applied directly on raw ImageNet images to yield satisfactory performance, when performed on “deep” features of the data (such as VGG, DenseNet, or ResNet features) obtained from *independent* deep neural networks, these algorithms tend to behave the same as with simple Gaussian mixtures [Seddik et al., 2020]. These seemingly striking empirical observations are indeed theoretically sustained by universality arguments arising from the powerful concentration of measure theory.

To be more precise, the following systematic comparison approach will be pursued in this monograph. An *asymptotically non-trivial* classification or regression problem is studied: that is, we assume that the problem at hand is theoretically neither too easy nor too hard to solve (as the one discussed in Section 1.1.3.1) and practically leads, in general, to, say, (binary) classification error rates of the order of 5% – 30% and of absolute regression errors also of the order 5% – 30%. In particular, we insist that the asymptotic random matrix framework under study is in general incapable to thinly grasp error rates below the 1 – 2% region, which may be the domain of “outliers” and marginal data.

Having posed this non-triviality assumption, we shall generically model the data as drawn from a simple mixture model, for example the Gaussian mixture model that gives access to a large panoply of powerful technical tools. The theoretical results obtained from the proposed analyses (asymptotic performance notably) are thus function of the statistical means and covariances of the mixture distribution. To compare the theoretical results to real data, we then conduct the following procedure:

1. exploiting the numerous (often labeled) samples of the real datasets (such as the $\sim 60\,000$ images of the training MNIST database), we empirically compute statistical means and covariances for each class of the database;
2. we then evaluate the asymptotic performance that a genuine Gaussian mixture model having *these means and covariances* would have;
3. we compare these “theoretical” values to actual simulations.

As the monograph will demonstrate in most scenarios, this procedure systematically leads to the conclusion that *performance of machine learning methods obtained on mere Gaussian mixtures* approximate surprisingly well the performance observed on real data and features. On a side note, we mentioned in Remark 1.2 that it is likely inappropriate to use the sample covariance matrix to estimate the population covariance of the small (i.e., n not much larger than p) databases, such as the MNIST database (for which $n/p \ll 100$). However, it turns out that, as the quantities of interest (e.g., classification or regression errors) are generally *scalar* functionals of the data statistical means and covariances, it is still possible, in the large n, p regime, to derive *consistent* estimators of these quantities without resorting to an exact evaluation of the (large dimensional) moments; see more discussions on this important topic in Section 3.2 and 4.5.

As already mentioned in Remark 1.3, this surprising accordance between theory and practice is possibly due to the *universality* of random matrix theory results, i.e., only the first several order statistics of the problem at hand matter in the large dimensional regime (recall for instance that the limiting eigenvalue distribution of $\frac{1}{n}\mathbf{X}\mathbf{X}^T$ for $\mathbf{X} \in \mathbb{R}^{p \times n}$ having i.i.d. zero mean and unit variance entries is the *same* Marčenko-Pastur law, irrespective of the higher order moments of \mathbf{X}).

Yet, another stronger argument can be made, especially when it comes to machine learning for image processing.

1.2.4.2 Concentrated random vectors and real data modeling

The modeling assumption that the data vectors \mathbf{x}_i are linear or affine maps $\mathbf{x}_i = \mathbf{A}\mathbf{z}_i + \mathbf{b}$ of random vectors \mathbf{z}_i constituted of i.i.d. entries is simultaneously an asset for random matrix analysis (by exploiting the degrees of freedom in the entries of \mathbf{z}_i) but a severe practical limitation as few real datasets are likely of this simplistic form.

El Karoui [2009] provided a first means for random matrix theory to go beyond the “vector of independent entries” assumption.¹⁰ There, relying on elements of the *concentration of measure theory*, extensively developed by Ledoux [2005], El Karoui essentially shows (in a rather technical manner) that some of the early random matrix results from Pastur, Bai, and Silverstein, remain valid under the assumption that the \mathbf{x}_i ’s are *concentrated random vectors*. Roughly speaking, a random vector $\mathbf{x} \in \mathbb{R}^p$ is *concentrated* if, for a certain family of functions $f : \mathbb{R}^p \rightarrow \mathbb{R}$, there exists a deterministic scalar $M_f \in \mathbb{R}$ such that

$$\mathbb{P}(|f(\mathbf{x}) - M_f| > t) \leq \alpha(t) \quad (1.14)$$

for some decreasing function $\alpha : \mathbb{R} \rightarrow \mathbb{R}$; in general, $\alpha(t)$ will be of the form $\alpha(t) = Ce^{-ct^q}$ for some $q > 0$ and $C, c > 0$ constant (which may depend on p

¹⁰See also [Pajor and Pastur, 2009] published the same year under slightly more constrained assumptions.

though). Intuitively, a concentrated random vector is a (random) point in high-dimensional space having “predictable observations” $f(\mathbf{x})$, in the sense that, with (exponentially) high probability, $f(\mathbf{x})$ takes values very close to the deterministic M_f . Thus, in the (one-dimensional) “observable world”, the observation $f(\mathbf{x})$, which may typically be any performance metric of a machine learning algorithm on a test datum \mathbf{x} , appears to be “stable” for any concentrated vector \mathbf{x} .¹¹




Figure 1.6: Multivariate Gaussian distribution $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, a fundamental example of concentrated random vectors. (**Left**) A visual “interpretation” of 500 independent drawings of $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$. (**Right**) Concentration of observations for linear ($f_1(\mathbf{x}) = \mathbf{x}^T \mathbf{1}_p / \sqrt{p}$) and Lipschitz ($f_2(\mathbf{x}) = \|\mathbf{x}\|_\infty$) maps.

Ledoux and El Karoui mostly focus on concentrated random vectors defined on Lipschitz classes of functions f , i.e., \mathbf{x} is *Lipschitz-concentrated* if (1.14) holds for all f such that $|f(\mathbf{x}) - f(\mathbf{y})| \leq \|\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^p$. These stringent constraints however make it very hard to find *any* random vector belonging to this class. As a matter of fact, in this class, the only standard random vectors are the Gaussian random vector $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ and the uniform vector on the sphere $\mathbf{u} = \mathbf{x}/\|\mathbf{x}\| \sim \mathbb{S}^{p-1}$ for $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$. However, quite importantly, *every* $\mathbb{R}^p \rightarrow \mathbb{R}^q$ Lipschitz-mapping $g(\mathbf{x})$ and $g(\mathbf{u})$ of these two random vectors, by definition, also belongs to the class.¹²

¹¹Note that by modeling the input data \mathbf{x} as a concentrated random vector and stating that the output (statistics) of a machine learning algorithm is “stable” implicitly assumes some regularity in the algorithm which, as we shall see, can be shown to hold for many popular methods including deep neural networks (and which often takes the form of a “Lipschitz control”).

¹²Under the more restricted class of *Lipschitz and convex* functions, random vectors with i.i.d. and bounded entries (up to normalization) also create a class of (convexly) concentrated random vectors.

A visual representation of the notion of concentration is presented in Figure 1.6.

Yet, since the widest class of (Lipschitz) concentrated random vectors is restricted to Lipschitz maps of standard Gaussian vectors, at first sight, concentrated random vectors are seemingly no more elaborate models than linear and affine maps of Gaussian vectors. As a consequence, there is a priori no reason to assume that mixtures of concentrated random vectors can model real data any better than Gaussian mixtures.

It turns out that this intuition is again tainted by erroneous finite-dimensional insights. Indeed, there *practically exist extremely data-realistic concentrated random vectors*: the outputs of generative adversarial networks (GANs) [Goodfellow et al., 2014] as shown in Figure 1.7. GANs generate artificial images $g(\mathbf{x})$ from large dimensional standard Gaussian vectors \mathbf{x} where g is a conventional feedforward neural network trained to mimic real data. As such, g is the combination of Lipschitz nonlinear (the neural activations) and linear (the inter-layer connexions) maps, and is thus a Lipschitz mapping.¹³ The output image vectors $g(\mathbf{x})$, see examples in Figure 1.8, are thus concentrated vectors. Modern GANs are so sophisticated that it has become virtually impossible for human beings to tell whether their outputs are genuine or artificial. This, as a result, strongly suggests that concentrated random vectors are accurate models of real-world data.




Figure 1.7: Illustration of a generative adversarial network (GAN).

A strong emphasis has thus lately been given to these models. The monograph will in particular elaborate on the work of Louart and Couillet [2018] which largely generalizes the seminal findings of El Karoui by providing a systematic methodological toolbox of *concentration theory for random matrices*. There, the notion of concentration is generalized by including *linear concentration*, which provides a consistent framework for the important notion of deterministic equivalents in random matrix theory, and by providing a wide range of properties and lemmas of immediate use for random matrix purposes.

¹³In practice, other operations are also performed in neural networks, such as pooling operations, random or deterministic dropouts, various connectivity matrix normalization procedures, so as to achieve better performance. They are all shown to be Lipschitz [Seddik et al., 2020].



Figure 1.8: Images samples generated by BigGAN in [Brock et al., 2019].

An important finding of [Louart and Couillet, 2018] is that *first order statistics* of functionals of random matrices built from concentrated random vectors, so in particular the asymptotic performance of many machine learning methods, are also universal. Specifically, for most conventional machine learning methods (support vector machines, semi-supervised learning, spectral clustering, random feature maps, linear regression, etc.), the asymptotic performance achieved on Gaussian mixtures $\mathcal{N}(\boldsymbol{\mu}_a, \mathbf{C}_a)$, $a \in \{1, \dots, k\}$, coincides with that obtained on concentrated random vectors mixtures $\mathcal{L}_a(\boldsymbol{\mu}_a, \mathbf{C}_a)$, $a \in \{1, \dots, k\}$, having the same means $\boldsymbol{\mu}_a$ and covariances \mathbf{C}_a per class, and are *independent* of the high-order moments of the underlying distribution.

This strongly suggests that Gaussian mixture models, if not appropriate data “models” per se, are largely sufficient statistical assumptions for the theoretical understanding of real data machine learning.

Remark 1.4 (Concentration of measure, concentration inequalities, and non-asymptotic random matrices). *It is important to raise here the fact that the concentration of measure theory is structurally broader than the scope of the popular concentration inequalities regularly used in statistical learning theory [Boucheron et al., 2013, Tropp, 2015, Vershynin, 2018]. Concentration inequalities are merely expressions of (1.14) for specific choices of f and their consequences, and they are in particular not new to random matrix theory. In [Vershynin, 2012, Tao, 2012], Vershynin and Tao exploit the mathematical strength of concentration inequalities (which, thanks to the exponential decay, is stronger and less cumbersome to handle than moment bounds) to prove fundamental results in random matrix theory. Yet, these inequalities are mostly exploited in proofs involving Gaussian or sub-gaussian random vectors (as an instance of concentrated random vector). In particular, Vershynin establishes a non-asymptotic random matrix theory by exploiting concentration inequalities to bound various quantities of theoretical interest (notably bounds on the eigenvalue positions of random matrices). The monograph instead puts forth the interest of concentration of measure theory for data modeling beyond a merely convenient mathematical tool.*

Concentration of measure theory is also all the more suited to machine learning as it structurally relates to linear, Lipschitz, or convex-Lipschitz functionals of random vectors and matrices. These are precisely the core elements of machine learning algorithms (kernels, activation functions, convex optimization

schemes). From this viewpoint, concentration of measure theory is much more adapted to machine learning analysis than seemingly simpler data models. Note for instance that concentrated random vectors are stable (i.e., they remain concentrated) when passed through the layers of a neural network; this is not true of Gaussian random vectors or vectors with independent entries which in general no longer have independent entries when passed through nonlinear layers.

A last but not least convenient aspect of concentration of measure theory is that it flexibly allows one to “decouple” the behavior of the data size p and number n in the large dimensional setting. It is technically much easier to keep track of *independent growth rates* for p and n under a concentration of measure framework than when exploiting more standard random matrix techniques (such as Gaussian tools).

1.3 Outline and online toolbox

1.3.1 Organization of the manuscript

The remainder of the manuscript is divided in two parts.

Chapter 2 introduces the basics of random matrix theory *needed for machine learning applications in this monograph*. In doing so, we shall first revisit the traditional approach found in math-oriented sources, such as [Bai and Silverstein, 2010] based on a Stieltjes transform and truncation machinery, [Pastur and Shcherbina, 2011] based on a Gaussian-method approach, [Tao, 2012, Vershynin, 2012] based on concentration inequalities and a non-asymptotic random matrix approach, and also say a few words on [Mingo and Speicher, 2017] which follows a free probability framework or on [Anderson et al., 2010] which is more oriented towards a determinantal point process and large deviations direction. Unlike most of these references though (at the possible exception of [Pastur and Shcherbina, 2011]), our methodology is primarily centered on the statistical analysis of the *resolvent* (and only secondarily on the Stieltjes transform) of random matrices, which is the chief object of interest to us in most machine learning applications. The particular mathematical toolbox exploited to derive the results is of secondary importance.

In this chapter, we will successively introduce: [** not sure if it is helpful to point to the respect section. **](#)

- the fundamental notion of the *resolvent* $\mathbf{Q}(z) = (\mathbf{X} - z\mathbf{I}_n)^{-1}$ of a (random) matrix \mathbf{X} and its relations to the eigenvalues of \mathbf{X} , the limiting spectrum of \mathbf{X} , the eigenvectors and eigenspaces associated to some specific eigenvalues, as well as its relations to bilinear and quadratic forms often met in applications (linear or kernel regression, linear and quadratic discriminant analysis, support vector machines, and even the performance of some simple neural networks);
- the almost equally important notion of *deterministic equivalents* which extend the notion of the “limiting behavior” of large dimensional random

matrices, when such limits may not exist (which is the case of most structured random matrix models of practical interest); *deterministic equivalents for the resolvent* of random matrix models are at the core of almost all results derived in this monograph;

- the foundational Marčenko-Pastur and Wigner semicircle laws which, as we shall see, serve as a reference “null model” to all random matrix models met in machine learning applications; even quite sophisticated random matrix transformations (through nonlinear kernels, discontinuous activation functions, etc.) will be seen to boil down, in one way or another, to either one (or a mixture of both) of these reference laws;
- a successive presentation of the three main technical tools at our disposal (in this monograph at least) to study random matrix models: the Bai-Silverstein-Stieltjes transform approach, the Pastur-Shcherbina Gaussian tools, and the Louart-Couillet concentration of measure approach;
- the natural extensions of the Marčenko-Pastur- and Wigner-like random matrix models to more structured models: with correlation in either features or samples, with nonzero mean, divided into sub-classes of correlated nonzero mean models, with a variance profile (in the case of heterogeneous graph matrix models), etc.;
- a refined analysis of the large dimensional spectrum of random matrices using tools from complex analysis, based on which statistical inference techniques on covariance matrix models are introduced;
- a thorough treatment of the so-called *spiked models* of random matrices which carry a significant importance in the applications to machine learning: spiked models consist in *low rank deviations* from some elementary or structured random matrix models; this “rank sparsity” property simplifies the analyses and appropriately models the presence of cluster, classes, communities, principal components, etc., in machine learning data models;
- a short exposition of alternative tools and techniques, not of central focus in this monograph, but may have various advantages in specific random matrix structures;
- a short presentation of the very recent concentration of measure theory for random matrices that extends most of the results presented in this chapter to much more realistic (generative) models of data for machine learning applications.

This lengthy chapter provides a vast majority of the necessary tools to conduct the analyses performed in the subsequent chapters of machine learning methods. This second “application” part is organized as follows: [** not sure if it is helpful to point to the respect section. **](#)

- Chapter 3 introduces first applications of the random matrix framework devised in Chapter 2 to detection, estimation and statistical inference; particular emphasis is made on likelihood ratio tests for the detection of information from noise, on linear and quadratic discriminant analysis in a binary hypothesis test, on the estimation of distances between data statistics (particularly here the estimation of distances between unknown covariances and divergences between Gaussian measures of unknown statistics), as well as on the performance of robust estimators of covariance (or scatter) matrices. The estimation of covariance distance is a typical example where the usual large- n alone statistical answer dramatically fails, even when the ratio n/p is quite large, and random matrix analysis provide consistent (and improved) estimators. As for robust M-estimators, it is typical of a scenario where classical statistics fail to perform any satisfying analysis, while random matrix methods exploit concentration of measure phenomena to fully understand and improve their behavior.
- Chapter 4 follows with a detailed exposition of kernel random matrices and their applications to kernel- and graph-based methods in machine learning. This chapter successively exposes the many consequences for these methods of the already several-times discussed *concentration of distances* phenomenon and shows that, as a result, the behavior, performance, and the role of hyperparameters (kernel function, regularization penalty, etc.) become tractable and amenable to improvement. Applications to kernel spectral clustering, graph-based semi-supervised learning (SSL), and kernel ridge-regression (also referred to as least-squares SVM) are investigated, as representative examples of unsupervised, semi-supervised, and supervised learning methods. All these methods will be shown to be theoretically tractable, easy to optimize and thus to improve, with experiments on real data confirming the theoretical findings. The specific example of SSL is quite telling of the limitations of standard finite-dimensional intuitions, and it will be shown that *all known* classical graph-based SSL methods either dramatically fail, or at best, do not exhibit the expected SSL behavior (notably failing to account for the large number of unlabeled data): the proposed random matrix approach is quite simple and is proven to solve this problem.
- Chapter 5 focuses specifically on neural network models. While modern deep neural networks remain difficult to assess, several studies are reported in this chapter that address simpler models of neural networks (with random and few layers, with a possibly recurrent structure) and for which, again, new insights and exact asymptotic performance are provided. An additional discussion of the learning dynamics of gradient descent methods is also exposed in which the step-by-step performance and the importance of early stopping mechanisms are theoretically analyzed.
- Chapter 6 goes a step beyond all previous chapters for which all metrics of interest (algorithm behavior, performance) are *explicit functions* of the

various random matrix models introduced in Chapter 2 (under the form of eigenvalue distribution, eigenvector statistics, bilinear forms on the resolvent, etc.): here we focus on optimization schemes in machine learning having *no explicit solution*. As such, the performance of these algorithms are *implicitly* related to the random data matrix and seem, at first sight, not related to random matrix analysis. The chapter shows instead that most of these methods do exhibit asymptotic (large n, p) performance that can be expressed as an almost explicit function (via a few coupled equations) of random matrix models, thereby opening the door to a wide range of applications (logistic regression, support vector machines, general empirical risk minimization scheme, etc.).

- Chapter 7 discusses spectral methods for community detection on (mostly dense) graphs and networks. As opposed to all previous application chapters for which the elementary random matrix model under study is the Gram matrix $\mathbf{X}^T \mathbf{X}$ for data matrix $\mathbf{X} \in \mathbb{R}^{p \times n}$, the problem of community detection on graphs naturally relates to symmetric graph matrices $\mathbf{X} \in \mathbb{R}^{n \times n}$ with independent Bernoulli entries. The chapter discusses at length the popular stochastic block model (SBM) and degree-corrected SBM which mimic, with a different degree of reality, the behavior of genuine graphs with communities. A short discussion on the (difficult and so far not random matrix-related) modern concern of community detection on the even more realistic case of large dimensional and *sparse* graphs is also made.
- Chapter 8 closes the application chapters with a discussion on the extension of the aforementioned applications to real data modeling. There, using the recent concentration of measure for random matrix framework, simulations of extremely realistic models of data (images mostly) are used to theoretically validate the random matrix results devised in all previous chapters. The chapter notably passes across the fundamental but surprising message that simple data models (as Gaussian mixtures) are often sufficiently rich to account for the large dimensional performance of many existing machine learning algorithms.

1.3.2 Online codes

Matlab as well as Python codes of the algorithms used to obtain most of the visual results (graphs, histograms) provided in the monograph are publicly available at <https://github.com/Zhenyu-LIAO/RMT4ML>. Links to these codes are directly provided in the caption of the corresponding figures.

Chapter 2

Basics of Random Matrix Theory

Random matrix theory, at its inception, primarily dealt with the eigenvalue distribution (also referred to as the spectral measure) of large dimensional random matrices. One of the key technical tools to study these measures is the Stieltjes transform, often presented as the central object of the theory [Bai and Silverstein, 2010, Pastur and Shcherbina, 2011].

But signal processing and machine learning alike are often more interested in subspaces and eigenvectors (which often carry the structural information of the data) than in eigenvalues. Subspace or spectral methods, such as principal component analysis (PCA) [Wold et al., 1987], spectral clustering [Luxburg, 2007] and some semi-supervised learning techniques [Zhu, 2005] are built directly upon the eigenspace spanned by the several top eigenvectors.

Consequently, beyond the Stieltjes transform, a more general mathematical object, the *resolvent* of large random matrices will constitute the cornerstone of the monograph. The resolvent of a matrix gives access to its spectral measure, to the location of its isolated eigenvalues, to the statistical behavior of their associated eigenvectors when random, and consequently provides an entry-door to the performance analysis of numerous machine learning methods.

2.1 Fundamental objects

2.1.1 The resolvent

We first introduce the resolvent of a matrix.

Definition 1 (Resolvent). *For a symmetric matrix $\mathbf{M} \in \mathbb{R}^{n \times n}$, the resolvent $\mathbf{Q}_{\mathbf{M}}(z)$ of \mathbf{M} is defined, for $z \in \mathbb{C}$ not eigenvalue of \mathbf{M} , as*

$$\mathbf{Q}_{\mathbf{M}}(z) \equiv (\mathbf{M} - z\mathbf{I}_n)^{-1}. \quad (2.1)$$

The matrix $\mathbf{Q}_M(z)$ will often simply be denoted $\mathbf{Q}(z)$ when there is no ambiguity.

The resolvent operator is in fact a very classical tool, the use of which goes far beyond random matrix theory. It is for instance exploited in the analysis of linear operators in general Hilbert space [Akhiezer and Glazman, 2013] as well as in monotone operator theory of importance to modern convex optimization theory [Bauschke and Combettes, 2017].

2.1.2 Spectral measure and Stieltjes transform

The first use of the resolvent \mathbf{Q}_M is in its relation to the *empirical spectral measure* μ_M of the matrix M , through the associated *Stieltjes transform* m_{μ_M} , which we all define next.

Definition 2 (Empirical spectral measure). *For a symmetric matrix $M \in \mathbb{R}^{n \times n}$, the spectral measure or empirical spectral measure or empirical spectral distribution (e.s.d.) μ_M of M is defined as the normalized counting measure of the eigenvalues $\lambda_1(M), \dots, \lambda_n(M)$ of M ,*

$$\mu_M \equiv \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i(M)}. \quad (2.2)$$

Since $\int \mu_M(dx) = 1$, the spectral measure μ_M of a matrix $M \in \mathbb{R}^{n \times n}$ (random or not) is a probability measure. For (probability) measures, we can define their associated Stieltjes transforms as follows.

Definition 3 (Stieltjes transform). *For a real probability measure μ with support $\text{supp}(\mu)$, the Stieltjes transform $m_\mu(z)$ is defined, for all $z \in \mathbb{C} \setminus \text{supp}(\mu)$, as*

$$m_\mu(z) \equiv \int \frac{1}{t - z} \mu(dt). \quad (2.3)$$

This definition and the Stieltjes transform framework in effect extend beyond probability measures to σ -finite real measures (i.e., measures μ such that $\mu(\mathbb{R}) < \infty$), which will occasionally be discussed in this monograph.

The Stieltjes transform m_μ has numerous interesting properties: it is complex analytic on its domain of definition $\mathbb{C} \setminus \text{supp}(\mu)$, it is bounded $|m_\mu(z)| \leq 1/\text{dist}(z, \text{supp}(\mu))$, it satisfies $\Im[z] > 0 \Rightarrow \Im[m(z)] > 0$, and it is an increasing function on all connected components of its restriction to $\mathbb{R} \setminus \text{supp}(\mu)$ (since $m'_\mu(x) = \int (t - x)^{-2} \mu(dt) > 0$ with $\lim_{x \rightarrow \pm\infty} m_\mu(x) = 0$ if $\text{supp}(\mu)$ is bounded).

As a transform, m_μ admits an inverse formula to recover μ , as per the following result.

Theorem 2.1 (Inverse Stieltjes transform). *For a, b continuity points of the probability measure μ , we have*

$$\mu([a, b]) = \frac{1}{\pi} \lim_{y \downarrow 0} \int_a^b \Im[m_\mu(x + iy)] dx. \quad (2.4)$$

Besides, if μ admits a density f at x (i.e., $\mu(x)$ is differentiable in a neighborhood of x and $\lim_{\epsilon \rightarrow 0} (2\epsilon)^{-1} \mu([x - \epsilon, x + \epsilon]) = f(x)$),

$$f(x) = \frac{1}{\pi} \lim_{y \downarrow 0} \Im[m_\mu(x + iy)]. \quad (2.5)$$

Also, if μ has an isolated mass at x , then

$$\mu(\{x\}) = \lim_{y \downarrow 0} -iy m_\mu(x + iy). \quad (2.6)$$

Proof. Since $|\frac{y}{(t-x)^2+y^2}| \leq \frac{1}{y}$ for $y > 0$, by Fubini's theorem,

$$\begin{aligned} \frac{1}{\pi} \int_a^b \Im[m_\mu(x + iy)] dx &= \frac{1}{\pi} \int_a^b \left[\int \frac{y}{(t-x)^2+y^2} \mu(dt) \right] dx \\ &= \frac{1}{\pi} \int \left[\int_a^b \frac{y}{(t-x)^2+y^2} dx \right] \mu(dt) \\ &= \frac{1}{\pi} \int \left[\arctan\left(\frac{b-t}{y}\right) - \arctan\left(\frac{a-t}{y}\right) \right] \mu(dt). \end{aligned}$$

As $y \downarrow 0$, the difference in brackets converges either to $\pm\pi$ or 0 depending on the relative position of a, b and t . By the dominated convergence theorem, the limit, as $y \downarrow 0$, is $\int 1_{[a,b]} \mu(dt) = \mu([a, b])$. When μ has an isolated mass at x , say $\mu(dt) = a\delta_x(t)$, we similarly have, again by dominated convergence (using in particular $|y(t-x)| \leq \frac{1}{2}(y^2 + (t-x)^2)$),

$$\lim_{y \downarrow 0} -iy m_\mu(x + iy) = -\lim_{y \downarrow 0} \int \frac{iy(t-x)\mu(dt)}{(t-x)^2+y^2} + \lim_{y \downarrow 0} \int \frac{y^2\mu(dt)}{(t-x)^2+y^2} = a.$$

This concludes the proof of Theorem 2.1. \square

The important relation between the empirical spectral measure $\mu_{\mathbf{M}}$ of $\mathbf{M} \in \mathbb{R}^{n \times n}$, the Stieltjes transform $m_{\mu_{\mathbf{M}}}(z)$ and the resolvent $\mathbf{Q}_{\mathbf{M}}(z)$ lies in the fact that

$$m_{\mu_{\mathbf{M}}}(z) = \frac{1}{n} \sum_{i=1}^n \int \frac{\delta_{\lambda_i(\mathbf{M})}(t)}{t-z} = \frac{1}{n} \sum_{i=1}^n \frac{1}{\lambda_i(\mathbf{M}) - z} = \frac{1}{n} \operatorname{tr} \mathbf{Q}_{\mathbf{M}}(z). \quad (2.7)$$

Combining inverse Stieltjes transform in Theorem 2.1 and the relation above thus provides a link between $\mathbf{Q}_{\mathbf{M}}$ and the eigenvalue distribution of \mathbf{M} . While seemingly contorted at first sight, this link turns out to be a very efficient way to study the spectral measure of *large dimensional random matrices* \mathbf{M} .

In particular, note that Theorem 2.1 raises an interesting fact: the Stieltjes transform $m_\mu(z) = \int (t-z)^{-1} \mu(dt)$ is defined on all $\mathbb{C} \setminus \operatorname{supp}(\mu)$, and as z approaches the support $\operatorname{supp}(\mu)$, the integrand $(t-z)^{-1}$ becomes singular. Yet, this is precisely when $x = \Re[z] \in \operatorname{supp}(\mu)$ while $\Im[z] \downarrow 0$ that one can retrieve the density of μ at x from the Stieltjes transform $m_\mu(z)$. This observation is key to the analysis of the spectrum (both eigenvalues and eigenvectors) of (random) matrices: the singular points of the resolvent of a (random) matrix provide the information about its spectrum.

Remark 2.1 (Resolvent as a matrix-valued Stieltjes transform). *As proposed in [Hachem et al., 2007], it is convenient to extrapolate Definition 3 of Stieltjes transforms to $n \times n$ matrix-valued positive measures $\mathbf{M}(dt)$,¹ in which case Equation (2.7) can be generalized as*

$$\mathbf{Q}_{\mathbf{M}}(z) = \int \frac{\mathbf{M}(dt)}{t - z} = \mathbf{U} \operatorname{diag} \left\{ \frac{1}{\lambda_i(\mathbf{M}) - z} \right\}_{i=1}^n \mathbf{U}^T,$$

where we used the spectral decomposition $\mathbf{M} = \mathbf{U} \operatorname{diag}\{\lambda_i(\mathbf{M})\}_{i=1}^n \mathbf{U}^T$. This definition coincides with the former definition of the resolvent of \mathbf{M} . As such, the resolvent $\mathbf{Q}_{\mathbf{M}}(z)$ is an “improved” Stieltjes transform which enjoys similar properties as Stieltjes transforms on real-valued measures: it is complex analytic on its domain of definition, it is bounded $\|\mathbf{Q}_{\mathbf{M}}(z)\| \leq 1/\operatorname{dist}(z, \operatorname{supp}(\mu_{\mathbf{M}}))$, and $x \mapsto \mathbf{Q}_{\mathbf{M}}(x)$ for $x \in \mathbb{R} \setminus \operatorname{supp}(\mu_{\mathbf{M}})$ is an increasing matrix-valued function with respect to symmetric matrix partial ordering (i.e., $\mathbf{A} \succeq \mathbf{B}$ whenever $\mathbf{z}^T(\mathbf{A} - \mathbf{B})\mathbf{z} \geq 0$ for all \mathbf{z}).

2.1.3 Cauchy’s integral, linear eigenvalue functionals, and eigenspaces

Being complex analytic, the resolvent $\mathbf{Q}_{\mathbf{M}}(z)$ can be manipulated using advanced tools from complex analysis. Of particular interest to this monograph is the relation between the resolvent and Cauchy’s integral theorem.

Theorem 2.2 (Cauchy’s integral formula). *For $\Gamma \subset \mathbb{C}$ a positively (i.e., counterclockwise) oriented simple closed curve and a complex function $f(z)$ analytic in a region containing Γ and its inside, then*

- (i) if $z_0 \in \mathbb{C}$ is enclosed by Γ , $f(z_0) = -\frac{1}{2\pi i} \oint_{\Gamma} \frac{f(z)}{z_0 - z} dz$;
- (ii) if not, $\frac{1}{2\pi i} \oint_{\Gamma} \frac{f(z)}{z_0 - z} dz = 0$.

This result provides an immediate connection between the *linear functionals of the eigenvalues of \mathbf{M}* and the Stieltjes transform $m_{\mu_{\mathbf{M}}}(z)$ through

$$\frac{1}{n} \sum_{i=1}^n f(\lambda_i(\mathbf{M})) = -\frac{1}{2\pi i n} \oint_{\Gamma} f(z) \operatorname{tr}(\mathbf{Q}_{\mathbf{M}}(z)) dz = -\frac{1}{2\pi i} \oint_{\Gamma} f(z) m_{\mu_{\mathbf{M}}}(z) dz,$$

for all f complex analytic in a compact neighborhood of $\operatorname{supp}(\mu_{\mathbf{M}})$, by choosing the contour Γ to enclose $\operatorname{supp}(\mu_{\mathbf{M}})$ (i.e., all the eigenvalues $\lambda_i(\mathbf{M})$). More generally,

$$\frac{1}{n} \sum_{\lambda_i(\mathbf{M}) \in \Gamma^{\circ}} f(\lambda_i(\mathbf{M})) = -\frac{1}{2\pi i} \oint_{\Gamma} f(z) m_{\mu_{\mathbf{M}}}(z) dz,$$

¹Defined by the fact that $\mu(dt; \mathbf{z}) = \mathbf{z}^T \mathbf{M}(dt) \mathbf{z} = \sum_{ij} [\mathbf{z}]_i [\mathbf{z}]_j [\mathbf{M}]_{ij}(dt)$ is a positive real-valued measure for all \mathbf{z} . See [Rozanov, 1967] for an introduction.

for Γ° the inside of the contour Γ . Note that in this case it is sufficient for f to be analytic *in a neighborhood* of $\text{supp}(\mu_{\mathbf{M}}) \cap \Gamma^\circ$; in particular, if one wishes to count the number of eigenvalues in an interval $[a, b]$, one may use the formula for $f(t) = 1_{t \in [a-\varepsilon, b+\varepsilon]}$ for some $\varepsilon > 0$ small, which is of course not analytic on \mathbb{C} but is analytic on an open neighborhood of $[a, b]$.

Another quantity of interest relates to eigenvectors and eigenspaces. Considering the spectral decomposition $\mathbf{M} = \mathbf{U}\Lambda\mathbf{U}^\top$ with $\mathbf{U} = [\mathbf{u}_1, \dots, \mathbf{u}_n] \in \mathbb{R}^{n \times n}$ and $\Lambda = \text{diag}\{\lambda_1(\mathbf{M}), \dots, \lambda_n(\mathbf{M})\}$, we have

$$\mathbf{Q}_{\mathbf{M}}(z) = \sum_{i=1}^n \frac{\mathbf{u}_i \mathbf{u}_i^\top}{\lambda_i(\mathbf{M}) - z}$$

and thus the direct access to the i -th eigenvector \mathbf{u}_i of \mathbf{M} through

$$\mathbf{u}_i \mathbf{u}_i^\top = -\frac{1}{2\pi i} \oint_{\Gamma_{\lambda_i(\mathbf{M})}} \mathbf{Q}_{\mathbf{M}}(z) dz,$$

for $\Gamma_{\lambda_i(\mathbf{M})}$ a contour circling around $\lambda_i(\mathbf{M})$ only. More generally,

$$\mathbf{U}f(\Lambda; \Gamma)\mathbf{U}^\top = -\frac{1}{2\pi i} \oint_{\Gamma} f(z) \mathbf{Q}_{\mathbf{M}}(z) dz,$$

for f analytic in a neighborhood of Γ and its inside Γ° and $f(\Lambda; \Gamma) = \text{diag}\{f(\lambda_i(\mathbf{M})) \cdot 1_{\lambda_i(\mathbf{M}) \in \Gamma^\circ}\}_{i=1}^n$.

Of specific interest in this monograph will be the projection of an individual eigenvectors \mathbf{u}_i of \mathbf{M} onto a deterministic eigenvector \mathbf{v} . In particular, from the above,

$$|\mathbf{v}^\top \mathbf{u}_i|^2 = -\frac{1}{2\pi i} \oint_{\Gamma_{\lambda_i(\mathbf{M})}} \mathbf{v}^\top \mathbf{Q}_{\mathbf{M}}(z) \mathbf{v} dz.$$

In the real case $\mathbf{M} \in \mathbb{R}^{n \times n}$, this gives access to $\mathbf{v}^\top \mathbf{u}_i$, up to a sign (which at any rate is not fixed since both \mathbf{u}_i and $-\mathbf{u}_i$ are valid i -th eigenvectors). The formula extends in the complex by replacing transpose by Hermitian transpose, thus providing access to $\mathbf{v}^\text{H} \mathbf{u}_i$ up to a phase $e^{i\theta}$ for $\theta \in [0, 2\pi)$.

To summarize, the resolvent $\mathbf{Q}_{\mathbf{M}}$ provides access to *scalar observations* of the eigenspectrum of \mathbf{M} through its *linear functionals*, i.e., the scalar observations $\frac{1}{n} \sum_i f(\lambda_i(\mathbf{M}))$ and $|\mathbf{v}^\top \mathbf{u}_i|$ accessible from $\frac{1}{n} \text{tr } \mathbf{Q}_{\mathbf{M}}$ and $\mathbf{v}^\top \mathbf{Q}_{\mathbf{M}} \mathbf{v}$, respectively.

Before proceeding to the application of these results to random matrices, it is worth noticing at this point that the resolvent method automatically enables the powerful tools from *complex analysis*, the Cauchy integral formula being only one instance. Analytic functions, such as the Stieltjes transform and the resolvent, are “extremely smooth” objects, enjoying a host of convenient properties. One such important property is that it suffices to know an analytic function locally to know it globally.

Theorem 2.3 (Vitali's convergence theorem [Titchmarsh, 1939]). *Let f_1, f_2, \dots be a sequence of functions, analytic on a region $D \subset \mathbb{C}$, such that $|f_n(z)| \leq M$ uniformly on n and $z \in D$. Further assume that $f_n(z_j)$ converges for a countable set $z_1, z_2, \dots \in D$ having a limit point inside D . Then $f_n(z)$ converges uniformly in any region bounded by a contour interior to D . This limit is furthermore an analytic function of z .*

Vitali's convergence theorem will be heavily exploited to study the behavior of resolvents $\mathbf{Q}_M(z)$ near the real axis (where it is almost singular but of utmost interest) by instead studying its properties far from the real axis (where it is mathematically more convenient). The theorem is in fact doubly interesting as it states that the knowledge of f_n at a *countable* number of points z_1, z_2, \dots is sufficient to fully characterize the limit f ; in practice we will use this property when proving convergence of functionals $f_n(z) = g(\mathbf{Q}_M(z) - \bar{\mathbf{Q}}(z)) \rightarrow 0$ of *random* resolvents $\mathbf{Q}_M(z)$ to *deterministic* equivalents $\bar{\mathbf{Q}}(z)$ (here n is the growing size of the resolvents): if $f_n(z_j) \rightarrow 0$ almost surely for each z_1, z_2, \dots , then by the countable union of probability one events, $f_n(z_j) \rightarrow 0$ with probability one *uniformly* on the set $\{z_1, z_2, \dots\}$, and by Vitali we obtain that $f_n(z) \rightarrow 0$ with probability one uniformly on a possibly very large subset of \mathbb{C} .

2.1.4 Deterministic and random equivalents

This monograph is concerned with the situation where M is a *large dimensional random matrix*, the eigenvalues and eigenvectors of which need be related to the statistical nature of the model design of M .

In the early days of random matrix theory, the main focus was on the *limiting spectral measure* of $M \in \mathbb{R}^{n \times n}$, that is the characterization of a certain “limit” to the spectral measure μ_M of M as the size of M increases. For this purpose, a natural approach is to study the *random* Stieltjes transform $m_{\mu_M}(z)$ and to show that it admits a limit (in probability or almost surely) $m(z)$ as $n \rightarrow \infty$. However, this method has strong limitations: (i) it supposes that such a limit exists, therefore restricting the study to very regular models for M and (ii) it only quantifies $\frac{1}{n} \text{tr } \mathbf{Q}_M$ (through the Stieltjes transform), thereby discarding all subspace information about M carried in the resolvent \mathbf{Q}_M . As a consequence, a further study of the eigenvectors of M often requires a complete rework.

To avoid these limitations, modern random matrix theory focuses instead on the notion of *deterministic equivalents* which are deterministic matrices – thus finite dimensional objects rather than limits – having (in probability or almost surely) asymptotically the same *scalar observations* as the random ones.² In particular, these scalar observations of deterministic equivalents (e.g., their

²The wide spread of deterministic equivalents in the random matrix literature arose from application needs, primarily in signal processing and wireless communications, involving too structured matrix models for limiting eigenvalue distributions to be meaningful [Hachem et al., 2007, Couillet et al., 2011]. Yet, deterministic equivalents in fact originate from the (much earlier) works of Girko [2001]. They have recently been included as a new feature of free probability theory [Speicher and Vargas, 2012], an alternative approach to the resolvent method, which we will shortly discuss in Section 2.6.2.4.

normalized traces or their bilinear forms) need not themselves admit a limit as the matrix dimension n grows: what only matters is that they deterministically “track” the behavior of their random equivalent with increased accuracy as $n \rightarrow \infty$.

Definition 4 (Deterministic Equivalent). *We say that $\bar{\mathbf{Q}} \in \mathbb{R}^{n \times n}$ is a deterministic equivalent for the symmetric random matrix $\mathbf{Q} \in \mathbb{R}^{n \times n}$ if, for (sequences of) deterministic matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ and vectors $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ of unit norms (operator and Euclidean, respectively), we have, as $n \rightarrow \infty$,*

$$\frac{1}{n} \text{tr } \mathbf{A}(\mathbf{Q} - \bar{\mathbf{Q}}) \rightarrow 0, \quad \mathbf{a}^\top (\mathbf{Q} - \bar{\mathbf{Q}}) \mathbf{b} \rightarrow 0,$$

where the convergence is either in probability or almost sure.

This definition³ has the advantage to bring forth the two key elements giving access to the spectral information about a random matrix \mathbf{M} : traces and bilinear forms (of its resolvent $\mathbf{Q}_\mathbf{M}(z)$ for some z). Deterministic equivalents for the resolvent $\mathbf{Q}_\mathbf{M}$ then encode most of the information necessary to statistically quantify a random matrix \mathbf{M} .

A practical use of deterministic equivalents is to establish that, for a random matrix \mathbf{M} of interest, $\frac{1}{n} \text{tr}(\mathbf{Q}_\mathbf{M}(z) - \bar{\mathbf{Q}}(z)) \rightarrow 0$, say almost surely, for all $z \in \mathcal{C}$ with $\mathcal{C} \subset \mathbb{C}$. Denoting $\bar{m}_n(z) = \frac{1}{n} \text{tr} \bar{\mathbf{Q}}(z)$, this convergence implies that the Stieltjes transform of $\mu_\mathbf{M}$ “converges” in the sense that $m_{\mu_\mathbf{M}}(z) - \bar{m}_n(z) \rightarrow 0$. As we will see, this indicates that $\mu_\mathbf{M}$ gets increasingly well approximated, as n grows large, by a probability measure $\bar{\mu}_n$ having Stieltjes transform $\bar{m}_n(z)$. Identifying $\bar{m}_n(z)$, which uniquely defines $\bar{\mu}_n$, will often be as far as the *Stieltjes transform method* will lead us. But in some rare cases (such as with the Marčenko-Pastur and the semicircle laws), $\bar{\mu}_n$ will be (explicitly) identifiable.

In the remainder of the monograph, we will often characterize the large dimensional behavior of random matrix models \mathbf{M} through the “approximation” offered by the deterministic equivalents $\bar{\mathbf{Q}}(z)$ of their associated resolvents $\mathbf{Q}_\mathbf{M}(z)$, providing simultaneously access to their asymptotic spectral measures as well as to their eigenspaces. We will therefore extrapolate some of the core traditional results in random matrix theory, such as the Marčenko-Pastur law [Marčenko and Pastur, 1967], the sample covariance matrix model [Silverstein and Bai, 1995], etc., under this more general form of deterministic equivalents.

Remark 2.2 ($\bar{\mathbf{Q}}$ versus $\mathbb{E}[\mathbf{Q}]$). *For $\bar{\mathbf{Q}}$ a deterministic equivalent for \mathbf{Q} , the (probabilistic) convergences $\frac{1}{n} \text{tr } \mathbf{A}(\mathbf{Q} - \bar{\mathbf{Q}}) \rightarrow 0$ and $\mathbf{a}^\top (\mathbf{Q} - \bar{\mathbf{Q}}) \mathbf{b} \rightarrow 0$ in general unfold from the fact that*

$$\|\mathbb{E}[\mathbf{Q}] - \bar{\mathbf{Q}}\| \rightarrow 0,$$

³The notion of “deterministic equivalent” has not been formally defined in the literature. The present definition is thus restricted to this monograph and is for the convenience of presentation. Section 2.7 will provide an alternative, possibly more satisfying, definition through the notion of *linear concentration* (Definition 8).

and from a control of the variance of $\frac{1}{n} \text{tr}(\mathbf{AQ})$ and $\mathbf{a}^\top \mathbf{Qb}$; this will often be the strategy followed in our proofs. Note particularly that, if the above relation is met, then $\mathbb{E}[\mathbf{Q}]$ itself is, by Definition 4, a deterministic equivalent for \mathbf{Q} . However, $\mathbb{E}[\mathbf{Q}]$ is often not convenient to work with and a “truly deterministic” matrix $\tilde{\mathbf{Q}}$ involving no integration over probability spaces (and that can be numerically evaluated with ease) will be systematically preferred.

Deterministic equivalents will be used very regularly in the course of this monograph. To avoid heavy notations, particularly in the main theorems and their proofs, we will use the following shortcut notation, valid both for deterministic and random matrix equivalents.

Notation 1 (Matrix Equivalents). *For $\mathbf{X}, \mathbf{Y} \in \mathbb{R}^{n \times n}$ two random or deterministic matrices, we write*

$$\mathbf{X} \leftrightarrow \mathbf{Y},$$

if, for all $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ of unit norms (respectively, operator and Euclidean), we have the simultaneous results

$$\frac{1}{n} \text{tr} \mathbf{A}(\mathbf{X} - \mathbf{Y}) \rightarrow 0, \quad \mathbf{a}^\top (\mathbf{X} - \mathbf{Y})\mathbf{b} \rightarrow 0, \quad \|\mathbb{E}[\mathbf{X} - \mathbf{Y}]\| \rightarrow 0$$

where, for random quantities, the convergence is either in probability or almost sure.

In many situations, deterministic equivalents \mathbf{Y} for a random matrix \mathbf{X} may not be directly accessible with classical random matrix techniques. In these cases, the introduction of an intermediary random matrix $\tilde{\mathbf{X}}$ satisfying $\|\tilde{\mathbf{X}} - \mathbf{X}\| \xrightarrow{a.s.} 0$ in operator norm will help “propagate” the deterministic equivalents. Indeed, if $\tilde{\mathbf{X}} \leftrightarrow \mathbf{Y}$, then necessarily $\mathbf{X} \leftrightarrow \mathbf{Y}$. When the convergence $\|\tilde{\mathbf{X}} - \mathbf{X}\| \xrightarrow{a.s.} 0$ is too demanding, it may of course be sufficient in some cases to prove that $\mathbf{X} \leftrightarrow \tilde{\mathbf{X}}$ (in which case both matrices are random) to ensure that $\mathbf{X} \leftrightarrow \mathbf{Y}$. This justifies the need to apply the notation “ \leftrightarrow ” to arbitrary, random or deterministic, matrices.

2.2 Foundational random matrix results

here!

In this section we introduce the main historical results of random matrix theory (appropriately updated under a deterministic equivalent form), which will serve as supporting models to most applications in machine learning.⁴ For

⁴Although historically and technically, said “Wigner” models of symmetric random matrices with independent entries came first, are simpler to analyze and spurred more mathematical research efforts [Wigner, 1955, Mehta and Gaudin, 1960, Anderson et al., 2010], for the sake of machine learning applications, our focus is primarily set on the slightly more involved sample covariance matrix models [Marcenko and Pastur, 1967, Bai and Silverstein, 2010].

readability and accessibility to the readers new to random matrix theory, we mostly stick to intuitive and short sketches of proofs. Yet, for the readers to have a glimpse on the technical details and modern tools of the field, some of the proof sketches will be appended by a complete exhaustive proof.

Both sketches and detailed proofs rely on a set of elementary lemmas and identities which need be introduced to understand their spirit and cornerstone arguments. This is done below in Section 2.2.1. The main difference between sketches and detailed proofs then relies on additional technical *probability* theory arguments to prove various convergence results. These arguments strongly depend on the underlying random matrix model hypotheses (Gaussian independent, i.i.d., concentrated random vectors, etc.); for readability, we will focus in our proofs on one specific line of arguments (that we claim to be the “historical” one) and will discuss alternative techniques in side remarks. In particular, the specific *concentration of measure* theoretic approach, which is both more “modern” (yet less mature) and more adapted to machine learning applications, will be given a separate treatment in Section 2.7.

2.2.1 Key lemmas and identities

2.2.1.1 Resolvent identities

Most results discussed in this section consist in tools meant to help “approximate” random matrix resolvents $\mathbf{Q}(z)$ via deterministic resolvents $\tilde{\mathbf{Q}}(z)$ in the sense of the deterministic equivalents definition. The following first identity provides a comparison of matrix inverses (often used in practice to compare the aforementioned resolvents).

Lemma 2.1 (Resolvent identity). *For invertible matrices \mathbf{A} and \mathbf{B} , we have*

$$\mathbf{A}^{-1} - \mathbf{B}^{-1} = \mathbf{A}^{-1}(\mathbf{B} - \mathbf{A})\mathbf{B}^{-1}.$$

Proof. This can be easily checked by multiplying both sides on the left by \mathbf{A} and on the right by \mathbf{B} . \square

Another useful lemma that helps directly connect the resolvent of \mathbf{BA} to that of \mathbf{AB} , is given as follows.

Lemma 2.2. *For $\mathbf{A} \in \mathbb{R}^{p \times n}$ and $\mathbf{B} \in \mathbb{R}^{n \times p}$, we have*

$$\mathbf{A}(\mathbf{BA} - z\mathbf{I}_n)^{-1} = (\mathbf{AB} - z\mathbf{I}_p)^{-1}\mathbf{A}$$

for $z \in \mathbb{C}$ distinct from 0 and from the eigenvalues of \mathbf{AB} .

Proof. Left-multiply both ends of the equality by $\mathbf{AB} - z\mathbf{I}_p$ to obtain $\mathbf{A} = \mathbf{A}$. \square

For \mathbf{AB} and \mathbf{BA} symmetric, Lemma 2.2 is a special case of the more general relation $\mathbf{Af(BA)} = f(\mathbf{AB})\mathbf{A}$, with $f(\mathbf{M}) \equiv \mathbf{U}f(\mathbf{\Lambda})\mathbf{U}^\top$ under the spectral decomposition $\mathbf{M} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^\top$ and f complex analytic. Since f is analytic, $f(\mathbf{BA}) = \sum_{i=0}^{\infty} c_i(\mathbf{BA})^i$ for some sequence $\{c_i\}_{i=0}^{\infty}$ and thus $\mathbf{Af(BA)} = \sum_{i=0}^{\infty} c_i(\mathbf{AB})^i\mathbf{A} = f(\mathbf{AB})\mathbf{A}$.

The next lemma, known as *Sylvester's identity* (also known as the Weinstien–Aronszajn identity), similarly relates the resolvents of \mathbf{AB} and \mathbf{BA} through their determinant.

Lemma 2.3 (Sylvester's identity). *For $\mathbf{A} \in \mathbb{R}^{p \times n}$, $\mathbf{B} \in \mathbb{R}^{n \times p}$ and $z \in \mathbb{C} \setminus \{0\}$,*

$$\det(\mathbf{AB} - z\mathbf{I}_p) = \det(\mathbf{BA} - z\mathbf{I}_n) (-z)^{p-n}.$$

Proof. It suffices to develop the block-matrix determinant (recall that $\det \begin{pmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{pmatrix} = \det \mathbf{D} \det(\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C}) = \det \mathbf{A} \det(\mathbf{D} - \mathbf{C}\mathbf{A}^{-1}\mathbf{B})$ when \mathbf{A} and \mathbf{D} are invertible)

$$\det \begin{pmatrix} z\mathbf{I}_p & z\mathbf{A} \\ \mathbf{B} & z\mathbf{I}_n \end{pmatrix} = \det(z\mathbf{I}_p) \det(z\mathbf{I}_n - \mathbf{BA}) = \det(z\mathbf{I}_n) \det(z\mathbf{I}_p - \mathbf{AB}).$$

□

An immediate consequence of Sylvester's identity is that \mathbf{AB} and \mathbf{BA} have the same nonzero eigenvalues (those nonzero z 's for which both left- and right-hand sides vanish). Thus, say $n \geq p$, $\mathbf{AB} \in \mathbb{R}^{p \times p}$ and $\mathbf{BA} \in \mathbb{R}^{n \times n}$ have the same spectrum, except for the additional $n - p$ zero eigenvalues of \mathbf{AB} . This remark implies the next identity.

Lemma 2.4 (Trace of resolvent and co-resolvent). *Let $\mathbf{A} \in \mathbb{R}^{p \times n}$, $\mathbf{B} \in \mathbb{R}^{n \times p}$, and $z \in \mathbb{C}$ not an eigenvalue of \mathbf{AB} nor zero. Then*

$$\text{tr } \mathbf{Q}_{\mathbf{AB}}(z) = \text{tr } \mathbf{Q}_{\mathbf{BA}}(z) + \frac{n-p}{z}.$$

In particular, if \mathbf{AB} and \mathbf{BA} are symmetric,

$$m_{\mu_{\mathbf{AB}}}(z) = \frac{n}{p} m_{\mu_{\mathbf{BA}}}(z) + \frac{n-p}{pz}.$$

It will be customary, if $\mathbf{Q}_{\mathbf{AB}}$ is the resolvent of the matrix model \mathbf{AB} under study to call $\mathbf{Q}_{\mathbf{BA}}$ the *co-resolvent* of \mathbf{AB} . We will see that the resolvent and co-resolvent of random matrix models (in particular the resolvent and co-resolvent of \mathbf{XX}^\top for \mathbf{X} some structured random matrix model) often intervene together, and quite symmetrically, to define their associated deterministic equivalents.

2.2.1.2 Perturbation identities

Quantifying the asymptotic global (e.g., spectral distribution) or local (e.g., isolated eigenvalues or projection on eigenvector) behavior of random matrices \mathbf{M} will systematically involve a *perturbation approach*. The idea often lies in comparing the behavior of the resolvent $\mathbf{Q} = \mathbf{Q}_{\mathbf{M}}$ to the resolvent \mathbf{Q}_{-i} of \mathbf{M}_{-i} , with \mathbf{M}_{-i} defined as \mathbf{M} with either row and column i , or some i -th contribution (e.g., $\mathbf{M}_{-i} = \sum_{j \neq i} \mathbf{x}_j \mathbf{x}_j^\top$ if $\mathbf{M} = \sum_j \mathbf{x}_j \mathbf{x}_j^\top$), discarded. A number of so-called *perturbation* identities are then needed.

The first one involves the segmentation of \mathbf{M} under the form of subblocks, in general consisting of one large block and three small submatrices. The corresponding resolvent $\mathbf{Q}_{\mathbf{M}}$ can correspondingly be segmented in subblocks according to the following block inversion lemma.

Lemma 2.5 (Block matrix inversion). *For $\mathbf{A} \in \mathbb{R}^{p \times p}$, $\mathbf{B} \in \mathbb{R}^{p \times n}$, $\mathbf{C} \in \mathbb{R}^{n \times p}$ and $\mathbf{D} \in \mathbb{R}^{n \times n}$ with \mathbf{D} invertible, we have*

$$\begin{pmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{pmatrix}^{-1} = \begin{pmatrix} \mathbf{S}^{-1} & -\mathbf{S}^{-1}\mathbf{BD}^{-1} \\ -\mathbf{D}^{-1}\mathbf{CS}^{-1} & \mathbf{D}^{-1} + \mathbf{D}^{-1}\mathbf{CS}^{-1}\mathbf{BD}^{-1} \end{pmatrix}$$

where $\mathbf{S} \equiv \mathbf{A} - \mathbf{BD}^{-1}\mathbf{C}$ is the Schur complement (for the block \mathbf{D}) of $(\begin{smallmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{smallmatrix})$.⁵

As a consequence of Lemma 2.5, we get the following explicit form for all diagonal entries of an invertible matrix \mathbf{A} .

Lemma 2.6 (Diagonal entries of matrix inverse). *For invertible $\mathbf{A} \in \mathbb{R}^{p \times p}$ and $\mathbf{A}_{-i} \in \mathbb{R}^{(p-1) \times (p-1)}$, the matrix obtained by removing the i -th row and column from \mathbf{A} , $i = 1, \dots, p$, we have*

$$(\mathbf{A}^{-1})_{ii} = \frac{1}{\mathbf{A}_{ii} - \boldsymbol{\alpha}_i^T (\mathbf{A}_{-i})^{-1} \boldsymbol{\beta}_i}$$

for $\boldsymbol{\alpha}_i^T, \boldsymbol{\beta}_i \in \mathbb{R}^{p-1}$ the i -th row and column of \mathbf{A} with i -th entries removed, respectively.

The result follows from Lemma 2.5 for entry $(1, 1)$ and can then be generalized to an arbitrary diagonal entry (i, i) by pre- and post-multiplication with $[\mathbf{P}]_{ab} = [\mathbf{I}_p]_{\sigma(a), b}$ for the permutation $\sigma(1) = i$, $\sigma(i) = 1$ and $\sigma(j) = j$ otherwise. Alternatively, the result may be obtained from the fact that $\mathbf{A}^{-1} = \frac{\text{adj}(\mathbf{A})}{\det(\mathbf{A})}$, with $\text{adj}(\mathbf{A})$ the adjugate matrix of \mathbf{A} , together with the block determinant formula.

Perturbations by addition or subtraction of low-rank matrices to \mathbf{M} induce modifications in the resolvent \mathbf{Q} that involve Woodbury's identity as follows.

Lemma 2.7 (Woodbury). *For $\mathbf{A} \in \mathbb{R}^{p \times p}$, $\mathbf{U}, \mathbf{V} \in \mathbb{R}^{p \times n}$, such that both \mathbf{A} and $\mathbf{A} + \mathbf{UV}^T$ are invertible, we have*

$$(\mathbf{A} + \mathbf{UV}^T)^{-1} = \mathbf{A}^{-1} - \mathbf{A}^{-1}\mathbf{U}(\mathbf{I}_n + \mathbf{V}^T\mathbf{A}^{-1}\mathbf{U})^{-1}\mathbf{V}^T\mathbf{A}^{-1}.$$

Note importantly that, while $(\mathbf{A} + \mathbf{UV}^T)^{-1}$ is of size $p \times p$, $\mathbf{I}_n + \mathbf{V}^T\mathbf{A}^{-1}\mathbf{U}$ is of size $n \times n$. This will turn out useful, for $n \ll p$, to relate resolvents of large dimensional matrices to resolvents of more elementary fixed small size matrices. In particular, for $n = 1$, i.e., $\mathbf{UV}^T = \mathbf{uv}^T$ for $\mathbf{U} = \mathbf{u} \in \mathbb{R}^p$ and $\mathbf{V} = \mathbf{v} \in \mathbb{R}^p$, Woodbury's identity specializes to the Sherman–Morrison formula.

Lemma 2.8 (Sherman–Morrison). *For $\mathbf{A} \in \mathbb{R}^{p \times p}$ invertible and $\mathbf{u}, \mathbf{v} \in \mathbb{R}^p$, $\mathbf{A} + \mathbf{uv}^T$ is invertible if and only if $1 + \mathbf{v}^T\mathbf{A}^{-1}\mathbf{u} \neq 0$ and*

$$(\mathbf{A} + \mathbf{uv}^T)^{-1} = \mathbf{A}^{-1} - \frac{\mathbf{A}^{-1}\mathbf{uv}^T\mathbf{A}^{-1}}{1 + \mathbf{v}^T\mathbf{A}^{-1}\mathbf{u}}.$$

Besides,

$$(\mathbf{A} + \mathbf{uv}^T)^{-1}\mathbf{u} = \frac{\mathbf{A}^{-1}\mathbf{u}}{1 + \mathbf{v}^T\mathbf{A}^{-1}\mathbf{u}}.$$

⁵The Schur complement $\mathbf{S} = \mathbf{A} - \mathbf{BD}^{-1}\mathbf{C}$ is particularly known for its providing the block determinant formula $\det(\begin{smallmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{smallmatrix}) = \det(\mathbf{D})\det(\mathbf{S})$, already exploited in the proof of Sylvester's identity (Lemma 2.3).

Letting $\mathbf{A} = \mathbf{M} - z\mathbf{I}_p$, $z \in \mathbb{C}$, and $\mathbf{v} = \tau\mathbf{u}$ for $\tau \in \mathbb{R}$ in the previous lemma leads to the following rank-1 perturbation lemma for the resolvent of \mathbf{M} .

Lemma 2.9 ([Silverstein and Bai, 1995, Lemma 2.6]). *For $\mathbf{A}, \mathbf{M} \in \mathbb{R}^{p \times p}$ symmetric, $\mathbf{u} \in \mathbb{R}^p$, $\tau \in \mathbb{R}$ and $z \in \mathbb{C} \setminus \mathbb{R}$,*

$$|\operatorname{tr} \mathbf{A}(\mathbf{M} + \tau\mathbf{u}\mathbf{u}^\top - z\mathbf{I}_p)^{-1} - \operatorname{tr} \mathbf{A}(\mathbf{M} - z\mathbf{I}_p)^{-1}| \leq \frac{\|\mathbf{A}\|}{|\Im(z)|}.$$

Also, for $\mathbf{A}, \mathbf{M} \in \mathbb{R}^{p \times p}$ symmetric and nonnegative definite, $\mathbf{u} \in \mathbb{R}^p$, $\tau > 0$ and $z < 0$,

$$|\operatorname{tr} \mathbf{A}(\mathbf{M} + \tau\mathbf{u}\mathbf{u}^\top - z\mathbf{I}_p)^{-1} - \operatorname{tr} \mathbf{A}(\mathbf{M} - z\mathbf{I}_p)^{-1}| \leq \frac{\|\mathbf{A}\|}{|z|}.$$

Exercise 4 at the end of the chapter proposes a partial proof for the case $z < 0$.

It is interesting (and possibly counterintuitive at first) to note that $\|\mathbf{u}\|$ does not intervene in this inequality. In particular, irrespective of the amplitude of the rank-1 perturbation, under the conditions of the lemma

$$m_{\mu_{\mathbf{M} + \tau\mathbf{u}\mathbf{u}^\top}}(z) = m_{\mu_{\mathbf{M}}}(z) + O(p^{-1})$$

and thus, by the link between spectrum and Stieltjes transform, the spectral measure of \mathbf{M} is asymptotically close to that of $\mathbf{M} + \tau\mathbf{u}\mathbf{u}^\top$ for any \mathbf{u} , in the large p limit. This result can be understood through the following two arguments: (i) for large p , the spectrum of \mathbf{M} (say $\|\mathbf{M}\| = O(1)$ without generality restriction) is only non-trivial if the vast majority of the p eigenvalues of \mathbf{M} are of order $O(1)$: thus, as p eigenvalues use a space of size $O(1)$, they tend to aggregate; (ii) by Weyl's interlacing lemma presented next (Lemma 2.10) for symmetric matrices, the eigenvalues of \mathbf{M} and of $\mathbf{M} + \tau\mathbf{u}\mathbf{u}^\top$ are interlaced. Both arguments thus indicate that, in the large p limit, the spectral measures are indeed asymptotically the same.

Unlike non-symmetric matrices, symmetric matrices indeed enjoy the nice property of having stable spectra with respect to rank-1 perturbations. For $\lambda \in \mathbb{R}$ an eigenvalue of $\mathbf{M} + \tau\mathbf{u}\mathbf{u}^\top$ but not of \mathbf{M} with, say, $\tau > 0$, we indeed have

$$\begin{aligned} 0 &= \det(\mathbf{M} + \tau\mathbf{u}\mathbf{u}^\top - \lambda\mathbf{I}_p) = \det(\mathbf{Q}_{\mathbf{M}}(\lambda)^{-1}) \det(\mathbf{I}_p + \tau\mathbf{Q}_{\mathbf{M}}(\lambda)\mathbf{u}\mathbf{u}^\top) \\ &= \det(\mathbf{Q}_{\mathbf{M}}(\lambda)^{-1}) (1 + \tau\mathbf{u}^\top \mathbf{Q}_{\mathbf{M}}(\lambda)\mathbf{u}) \end{aligned}$$

where the second equality unfolds from factoring out $\mathbf{M} - \lambda\mathbf{I}_p$ (which is not singular as λ is not an eigenvalue of \mathbf{M}) and the third from Sylvester's identity (Lemma 2.3). As a consequence, λ is one of the solutions to

$$-1 = \tau\mathbf{u}^\top \mathbf{Q}_{\mathbf{M}}(\lambda)\mathbf{u} = \tau \sum_{i=1}^p \frac{|\mathbf{v}_i^\top \mathbf{u}|^2}{\lambda_i(\mathbf{M}) - \lambda}, \quad \left(\mathbf{M} = \sum_{i=1}^p \lambda_i(\mathbf{M}) \mathbf{v}_i \mathbf{v}_i^\top \right)$$

which, seen as a function of λ , has asymptotes at each $\lambda_i(\mathbf{M})$ and is increasing (from $-\infty$ to ∞) on the segments $(\lambda_i(\mathbf{M}), \lambda_{i+1}(\mathbf{M}))$ (eigenvalues being sorted in increasing order). The eigenvalues of $\mathbf{M} + \tau \mathbf{u} \mathbf{u}^\top$ are therefore *interlaced* with those of \mathbf{M} (see Figure 2.1 for an illustration). This idea generalizes to finite-rank perturbation as follows.




Figure 2.1: Illustration of the eigenvalues of \mathbf{M} and the rank-one perturbation $\mathbf{M} + \tau \mathbf{u} \mathbf{u}^\top$ of \mathbf{M} , as well as the function $\tau \mathbf{u}^\top \mathbf{Q}_\mathbf{M}(\lambda) \mathbf{u}$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

Lemma 2.10 (Weyl's inequality [Horn and Johnson, 2012, Theorem 4.3.1]). Let $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{p \times p}$ be symmetric matrices and let the respective eigenvalues of \mathbf{A} , \mathbf{B} and $\mathbf{A} + \mathbf{B}$ be arranged in nondecreasing order, i.e., $\lambda_1 \leq \lambda_2 \leq \dots \lambda_{p-1} \leq \lambda_p$. Then, for all $i \in \{1, \dots, p\}$,

$$\begin{aligned}\lambda_i(\mathbf{A} + \mathbf{B}) &\leq \lambda_{i+j}(\mathbf{A}) + \lambda_{p-j}(\mathbf{B}), \quad j = 0, 1, \dots, p - i, \\ \lambda_{i-j+1}(\mathbf{A}) + \lambda_j(\mathbf{B}) &\leq \lambda_i(\mathbf{A} + \mathbf{B}), \quad j = 1, \dots, i,\end{aligned}$$

In particular, taking $i = 1$ in the first equation and $i = p$ in the second equation, together with the fact $\lambda_j(\mathbf{B}) = -\lambda_{p+1-j}(-\mathbf{B})$ for $j = 1, \dots, p$, implies

$$\max_{1 \leq j \leq p} |\lambda_j(\mathbf{A}) - \lambda_j(\mathbf{B})| \leq \|\mathbf{A} - \mathbf{B}\|.$$

This last implication is fundamental as it shows that the difference in *operator norm* $\|\mathbf{A} - \mathbf{B}\|$ controls (uniformly) the pairwise distance in eigenvalues $|\lambda_j(\mathbf{A}) - \lambda_j(\mathbf{B})|$. Since $\|\mathbf{A} - \mathbf{B}\| \leq \|\mathbf{A} - \mathbf{B}\|_F$, the same holds for the numerically simpler Frobenius norm; however, it is in general too demanding to request convergences of matrix differences in Frobenius norm which, as a result, is not used in practice (in particular, for most random matrix models $\mathbf{X} \in \mathbb{R}^{p \times p}$

considered in this monograph and in practice, $\|\mathbf{X}\|_F$ is in general $O(\sqrt{p})$ larger than $\|\mathbf{X}\|$).⁶

2.2.1.3 Probability identities

The results of the previous sections are algebraic identities useful to handle the resolvent \mathbf{Q}_M of the deterministic matrix \mathbf{M} . The second ingredient of random matrix analysis lies in asymptotic probability approximations as the dimensions of \mathbf{M} increase. Quite surprisingly, most results essentially revolve around the convergence of a certain quadratic form, which is often nothing more than a mere extension of the *law of large numbers*.

Those quadratic form convergence results come under multiple forms. The historical form, due to Bai and Silverstein, sometimes referred to as the “trace lemma”, is as follows.

Lemma 2.11 (Quadratic-form-close-to-the-trace, [Bai and Silverstein, 2010, Lemma B.26]). *Let $\mathbf{x} \in \mathbb{R}^p$ have i.i.d. entries of zero mean, unit variance and $\mathbb{E}[|x_i|^K] \leq \nu_K$ for some $K \geq 1$. Then for $\mathbf{A} \in \mathbb{R}^{p \times p}$ and $k \geq 1$,*

$$\mathbb{E} \left[| \mathbf{x}^\top \mathbf{A} \mathbf{x} - \text{tr } \mathbf{A} |^k \right] \leq C_k \left[(\nu_4 \text{tr}(\mathbf{A} \mathbf{A}^\top))^{k/2} + \nu_{2k} \text{tr}(\mathbf{A} \mathbf{A}^\top)^{k/2} \right]$$

for some constant $C_k > 0$ independent of p . In particular, if $\|\mathbf{A}\| \leq 1$ and the entries of \mathbf{x} have bounded eighth-order moment,

$$\mathbb{E} \left[(\mathbf{x}^\top \mathbf{A} \mathbf{x} - \text{tr } \mathbf{A})^4 \right] \leq Cp^2$$

for some $C > 0$ independent of p , and consequently, as $p \rightarrow \infty$,

$$\frac{1}{p} \mathbf{x}^\top \mathbf{A} \mathbf{x} - \frac{1}{p} \text{tr } \mathbf{A} \xrightarrow{\text{a.s.}} 0.$$

This last result is rather intuitive. For $\mathbf{A} = \mathbf{I}_p$, this is simply an instance of the law of large numbers. For generic \mathbf{A} , first note that, by the independence of the entries of \mathbf{x} , $\mathbb{E}[\mathbf{x}^\top \mathbf{A} \mathbf{x}] = \text{tr } \mathbf{A}$. Exploiting the fact that $\text{Var}[\mathbf{x}^\top \mathbf{A} \mathbf{x}/p] = O(p^{-1})$ then ensures that $\mathbf{x}^\top \mathbf{A} \mathbf{x}/p - \text{tr } \mathbf{A}/p \rightarrow 0$, but only in probability; since the variance calculus involves exponentiating the entries x_i of \mathbf{x} up to power 4, they need to be of finite fourth power. The almost sure convergence is achieved by showing the faster moment convergence $\mathbb{E}[(\mathbf{x}^\top \mathbf{A} \mathbf{x}/p - \text{tr } \mathbf{A}/p)^4] = O(p^{-2})$ which is the second statement of the lemma and requires 8-th order exponentiation of the x_i 's. The request for \mathbf{A} to be of bounded norm with respect to p in this case “stabilizes” the quadratic form $\mathbf{x}^\top \mathbf{A} \mathbf{x}$ by maintaining its concentration properties.

⁶This being said, the inequality $\|\mathbf{X}\| \leq (\text{tr}(\mathbf{X} \mathbf{X}^\top))^k)^{1/(2k)}$, which coincides with $\|\mathbf{X}\| \leq \|\mathbf{X}\|_F$ for $k = 1$ and becomes an equality in the $k \rightarrow \infty$ limit, is sometimes used in practice (however with $k \geq 2$) to control the norm $\|\mathbf{X}\|$. The approach is however quite cumbersome as it quickly becomes a heavy combinatorial calculus for not too small k .

Recalling that $\|\mathbf{Q}_M(z)\| \leq (\text{dist}(z, \text{supp}(\mu_M)))^{-1}$, Lemma 2.11 can be exploited for $\mathbf{A} = \mathbf{Q}_M(z)$ for all z away from the support of μ_M and all \mathbf{x} independent of $\mathbf{Q}_M(z)$. The core of the proofs of the main random matrix results is uniquely based on this last remark.

The quadratic-form-close-to-the-trace lemma is fundamental to already obtain *heuristics* on the main random matrix identities, using $\frac{1}{p}\mathbf{x}^\top \mathbf{A}\mathbf{x} \simeq \frac{1}{n}\text{tr } \mathbf{A}$ for \mathbf{x} independent of \mathbf{A} with independent zero-mean unit-variance entries. More rigorously, the lemma allows for a careful control on the fluctuations of $\frac{1}{p}\mathbf{x}^\top \mathbf{A}\mathbf{x}$ for deterministic \mathbf{A} . However, \mathbf{A} may itself be random (as when $\mathbf{A} = \mathbf{Q}_M(z)$ the resolvent of a random matrix M). In this case, the fluctuations of $\frac{1}{n}\text{tr } \mathbf{A}$ will also need to be controlled. The difficulty here, especially when \mathbf{A} takes the form of an inverse matrix $\mathbf{A} = \mathbf{Q}_M(z)$, is to exploit the independence in the columns of M nested inside the matrix inverse (or other elaborate function of M). This is elegantly and universally dealt with using Burkholder's inequality: denoting $\mathbb{E}_i[M]$ the expectation of the random matrix M conditionally to its first (or last) i columns, the sequence $\{(\mathbb{E}_i - \mathbb{E}_{i-1})[M]\}_{i=1}^p$ forms a so-called martingale difference; the fluctuations of such objects (which in a way extend the notion of series of independent random variables) are well controlled by Burkholder's inequality.

Lemma 2.12 (Burkholder inequality, [Bai and Silverstein \[2010, Lemma 2.13\]](#)).
Let $\{X_i\}_{i=1}^\infty$ be a martingale difference for the increasing σ -field $\{\mathcal{F}_i\}$ and denote \mathbb{E}_k the expectation with respect to \mathcal{F}_k . Then, for $k \geq 2$, and some constant C_k only dependent on k ,

$$\mathbb{E} \left[\left| \sum_{i=1}^n X_i \right|^k \right] \leq C_k \left(\mathbb{E} \left[\sum_{i=1}^n \mathbb{E}_{k-1}[|X_i|^2] \right]^{k/2} + \sum_{i=1}^n \mathbb{E}[|X_i|^k] \right).$$

Lemma 2.12 will mostly be used in the context of proof details on the fluctuations of technical random matrix functionals. It may however be substituted by other similar tools such as the Gaussian Nash–Poincaré inequality (Lemma 2.14) which also involves moment bounds but restricted to Gaussian random variables, or more conveniently with concentration inequalities (see Section 2.7 for detail) which no longer involve moments (which can be cumbersome to compute) but exponential bounds.

These identities constitute the main technical ingredients needed to understand the proofs of both historical and recent random matrix results. The next section introduces the most fundamental of those which will repeatedly be called in the remainder of the monograph.

2.2.2 The Marčenko-Pastur and semicircle laws

We start by illustrating how the aforementioned tools are used to prove the two most popular results in random matrix theory: the Marčenko-Pastur law and the Wigner semicircle law.

To simplify the exposition of the results, we will use the notation for deterministic equivalents introduced in Notation 1. That is, for $\mathbf{X}, \mathbf{Y} \in \mathbb{R}^{n \times n}$, we will denote $\mathbf{X} \leftrightarrow \mathbf{Y}$ if, for all unit norm $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$, $\frac{1}{n} \text{tr } \mathbf{A}(\mathbf{X} - \mathbf{Y}) \xrightarrow{\text{a.s.}} 0$, $\mathbf{a}^\top (\mathbf{X} - \mathbf{Y}) \mathbf{b} \xrightarrow{\text{a.s.}} 0$ and $\|\mathbb{E}[\mathbf{X} - \mathbf{Y}]\| \rightarrow 0$.

Most of the results involve Stieltjes transforms $m_\mu(z)$ of a real probability measure with support $\text{supp}(\mu) \subset \mathbb{R}$. Since Stieltjes transforms are such that $m_\mu(z) > 0$ for $z < \inf \text{supp}(\mu)$, $m_\mu(z) < 0$ for $z > \sup \text{supp}(\mu)$ and $\Im[z] \Im[m_\mu(z)] > 0$ if $z \in \mathbb{C} \setminus \mathbb{R}$, it will be convenient in the following to consider the set of scalar pairs

$$\begin{aligned} \mathcal{Z}(\mathcal{A}) = \{(z, m) \in \mathcal{A} \times \mathbb{C}, (\Im[z] \Im[m] > 0 \text{ if } \Im[z] \neq 0) \\ \text{or } (m > 0 \text{ if } z < \inf \mathcal{A}^c \cap \mathbb{R}) \text{ or } (m < 0 \text{ if } z > \sup \mathcal{A}^c \cap \mathbb{R})\}. \end{aligned}$$

In particular, for convenient choices of \mathcal{A} (not always $\mathbb{C} \setminus \text{supp}(\mu)$), many results presented next will involve pairs $(z, m(z))$ defined as the unique solution of an implicit equation *within* $\mathcal{Z}(\mathcal{A})$ (while the implicit equation may in general have more than one solution in $\mathbb{C} \times \mathbb{C}$).

2.2.2.1 The Marčenko-Pastur law

We present the Marčenko-Pastur law under the slightly modified form of a deterministic equivalent for the resolvent $\mathbf{Q}(z)$.

Theorem 2.4 (From [Marcenko and Pastur, 1967]). *Consider the resolvent $\mathbf{Q}(z) = (\frac{1}{n} \mathbf{XX}^\top - z \mathbf{I}_p)^{-1}$, for $\mathbf{X} \in \mathbb{R}^{p \times n}$ having independent entries with zero mean, unit variance and some smooth tail condition.⁷ Then, as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$, we have*

$$\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z), \quad \bar{\mathbf{Q}}(z) = m(z) \mathbf{I}_p \quad (2.8)$$

with $(z, m(z))$ the unique solution in $\mathcal{Z}(\mathbb{C} \setminus [(1 - \sqrt{c})^2, (1 + \sqrt{c})^2])$ of

$$zcm^2(z) - (1 - c - z)m(z) + 1 = 0. \quad (2.9)$$

The function $m(z)$ is the Stieltjes transform of the probability measure μ given explicitly by

$$\mu(dx) = (1 - c^{-1})^+ \delta_0(x) + \frac{1}{2\pi c x} \sqrt{(x - E_-)^+(E_+ - x)^+} dx \quad (2.10)$$

⁷For this result, and those related, various tail conditions may be used: a uniform finite moment of order ℓ for some $\ell > 2$ may be taken (usually $\ell = 4 + \varepsilon$ for any $\varepsilon > 0$ is sufficient). Depending on the proof approach though, stronger conditions may be requested, such as a subgaussian behavior, a concentration of measure condition, etc. Determining the minimalistic conditions for the results to hold has been of long interest to mathematicians, as demonstrated by the huge impact of the complete proof in [Tao and Vu, 2008] of the full-circle law theorem under no other condition than the identical distribution of the zero-mean and unit-variance entries (see also [Bordenave and Chafaï, 2014]). Yet, for machine learning purposes, these are of minor interest: we shall systematically assume “sufficiently smooth” (and technically convenient) conditions to hold, without hampering the practical applicability of the results. This being said, it is already interesting to observe that, here and in the vast majority of the coming results, the matrix entries need *not* be identically distributed, and that only the statistical mean and cross-variance of the entries dictate the limiting behavior.

where $E_{\pm} = (1 \pm \sqrt{c})^2$ and $(x)^+ = \max(0, x)$, and is known as the Marčenko-Pastur distribution. In particular, with probability one, the empirical spectral measure $\mu_{\frac{1}{n}\mathbf{X}\mathbf{X}^\top}$ converges weakly to μ .

Figure 2.2 depicts the density of the Marčenko-Pastur distribution for different values of c . For a fixed dimension p , the ratio c decreases as the number of samples n grows large, so that the eigenvalues of the sample covariance matrix become more “concentrated” (their spread is given by the length of the support $[(1 - \sqrt{c})^2, (1 + \sqrt{c})^2]$) around the unique population covariance matrix eigenvalue (when seeing \mathbf{X} as a collection $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ of p -dimensional data vectors), equal to 1.

Note that the “asymmetric bell” shape of the Marčenko-Pastur law gets increasingly skewed towards large values as c increases and that, as $c = 1$, the left-edge has the very peculiar behavior to diverge. This $c = 1$ setting is referred to as the “hard edge” scenario explained by the fact that the limiting density becomes $\frac{1}{2\pi x} \sqrt{x^+((1 - \sqrt{c})^2 - x)^+} \sim \frac{1}{2\pi\sqrt{x}}$ as $x \downarrow 0$ and thus behaves as $1/\sqrt{x}$ near the left edge (the left edge being at $x = 0$) rather than as $\sqrt{x - (1 + \sqrt{c})^2}$ when $c \neq 0$ (the left edge being at $x = (1 + \sqrt{c})^2$). When $c > 1$, a mass at zero is created (of weight $1 - c^{-1}$) while, possibly unexpectedly, the left edge of the main “bulk” of eigenvalues moves towards the right, leaving the open segment $(0, (1 - \sqrt{c})^2)$ empty.⁸




Figure 2.2: Marčenko-Pastur distribution for different c . Note the peculiar “hard-edge” behavior at $c = 1$, quite unlike other values of c .

⁸This hard-edge phenomenon is in fact not just an amusing artifact of the theory: it indeed has deep consequences in practice and notably explains the so-called *double-descent* phenomenon lately evidenced in large dimensional statistical inference (see e.g., [Nakkiran et al., 2020, Mei and Montanari, 2019, Deng et al., 2019]).

Proof. Before going into the details of the proof, we first give a few intuitive arguments.

Intuitive idea. A first heuristic derivation, essentially due to Bai and Silverstein, consists in iteratively “guessing” the form of $\bar{\mathbf{Q}}(z) = \mathbf{F}(z)^{-1}$ for some matrix $\mathbf{F}(z)$. To this end, from Lemma 2.1, it first appears that, writing $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$,

$$\begin{aligned}\mathbf{Q}(z) - \bar{\mathbf{Q}}(z) &= \mathbf{Q}(z) \left(\mathbf{F}(z) + z\mathbf{I}_p - \frac{1}{n} \mathbf{X}\mathbf{X}^\top \right) \bar{\mathbf{Q}}(z) \\ &= \mathbf{Q}(z) \left(\mathbf{F}(z) + z\mathbf{I}_p - \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top \right) \bar{\mathbf{Q}}(z).\end{aligned}$$

For $\bar{\mathbf{Q}}(z)$ to be a deterministic equivalent for $\mathbf{Q}(z)$, we wish in particular that $\frac{1}{p} \text{tr } \mathbf{A}(\mathbf{Q}(z) - \bar{\mathbf{Q}}(z)) \xrightarrow{a.s.} 0$, for \mathbf{A} arbitrary, deterministic, and such that $\|\mathbf{A}\| = 1$. That is

$$\frac{1}{p} \text{tr}(\mathbf{F}(z) + z\mathbf{I}_p) \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}(z) - \frac{1}{n} \sum_{i=1}^n \frac{1}{p} \mathbf{x}_i^\top \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}(z) \mathbf{x}_i \xrightarrow{a.s.} 0. \quad (2.11)$$

We recognize in $\frac{1}{p} \mathbf{x}_i^\top \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}(z) \mathbf{x}_i$ a quadratic form on which we would like to use Lemma 2.11 to turn it into a trace term independent of \mathbf{x}_i . Yet, Lemma 2.11 cannot be used as $\mathbf{Q}(z)$ depends on \mathbf{x}_i . To counter the difficulty, we then use Lemma 2.8 to write

$$\mathbf{Q}(z) \mathbf{x}_i = \frac{\mathbf{Q}_{-i}(z) \mathbf{x}_i}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i}(z) \mathbf{x}_i}$$

where $\mathbf{Q}_{-i}(z) = (\frac{1}{n} \sum_{j \neq i} \mathbf{x}_j \mathbf{x}_j^\top - z\mathbf{I}_p)^{-1}$ is independent of \mathbf{x}_i . Now legitimately applying Lemma 2.11, we find that

$$\frac{1}{p} \mathbf{x}_i^\top \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}(z) \mathbf{x}_i = \frac{\frac{1}{p} \mathbf{x}_i^\top \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}_{-i}(z) \mathbf{x}_i}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i}(z) \mathbf{x}_i} \simeq \frac{\frac{1}{p} \text{tr } \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}_{-i}(z)}{1 + \frac{1}{n} \text{tr } \mathbf{Q}_{-i}(z)}. \quad (2.12)$$

From Lemma 2.9, normalized traces involving $\mathbf{Q}_{-i}(z)$ or $\mathbf{Q}(z)$ are asymptotically identical (since their inverse only differs by the rank-1 matrix $\frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top$) and thus this further reads

$$\frac{1}{p} \mathbf{x}_i^\top \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}(z) \mathbf{x}_i \simeq \frac{\frac{1}{p} \text{tr } \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}(z)}{1 + \frac{1}{n} \text{tr } \mathbf{Q}(z)}.$$

Getting back to (2.11), we thus end up with the approximation

$$\frac{1}{p} \text{tr}(\mathbf{F}(z) + z\mathbf{I}_p) \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}(z) \simeq \frac{\frac{1}{p} \text{tr } \bar{\mathbf{Q}}(z) \mathbf{A} \mathbf{Q}(z)}{1 + \frac{1}{n} \text{tr } \mathbf{Q}(z)}$$

(the argument of the right-hand side summation over i no longer depends on i , so the sum symbol vanishes). As a consequence, we can now “guess” the form of $\mathbf{F}(z)$. Indeed, if it is to exist, $\mathbf{F}(z)$ must be of the type

$$\mathbf{F}(z) \simeq \left(-z + \frac{1}{1 + \frac{1}{n} \operatorname{tr} \mathbf{Q}(z)} \right) \mathbf{I}_p$$

for the approximation above to hold. To close the loop, taking $\mathbf{A} = \mathbf{I}_p$, $\frac{1}{n} \operatorname{tr} \mathbf{Q}(z)$ appearing in this display must be well approximated by $m(z) \equiv \frac{1}{p} \operatorname{tr} \bar{\mathbf{Q}}(z) = \frac{1}{p} \operatorname{tr} \mathbf{F}(z)^{-1}$ so that

$$\frac{1}{p} \operatorname{tr} \mathbf{Q}(z) \simeq m(z) = \frac{1}{-z + \frac{1}{1 + \frac{p}{n} \operatorname{tr} \mathbf{Q}(z)}} \simeq \frac{1}{-z + \frac{1}{1 + \frac{p}{n} m(z)}} \quad (2.13)$$

and we thus finally have

$$\bar{\mathbf{Q}}(z) = \mathbf{F}(z)^{-1} = m(z) \mathbf{I}_p$$

where, in the large n, p limit, $m(z)$ is solution to

$$m(z) = \frac{1}{-z + \frac{1}{1 + cm(z)}}$$

or equivalently

$$zcm^2(z) - (1 - c - z)m(z) + 1 = 0.$$

This equation has two solutions defined via the two values of the complex square root function (letting $z = \rho e^{i\theta}$ for $\rho \geq 0$ and $\theta \in [0, 2\pi]$, $\sqrt{z} \in \{\pm\sqrt{\rho}e^{i\theta/2}\}$)

$$m(z) = \frac{1 - c - z}{2cz} + \frac{\sqrt{((1 + \sqrt{c})^2 - z)((1 - \sqrt{c})^2 - z)}}{2cz}$$

only one of which is such that $\Im[z]\Im[m(z)] > 0$ as imposed by the definition of Stieltjes transforms. Now, from the inverse Stieltjes transform theorem, Theorem 2.1, we find that $m(z)$ is the Stieltjes transform of the measure μ with

$$\mu([a, b]) = \frac{1}{\pi} \lim_{\epsilon \downarrow 0} \int_a^b \Im[m(x + i\epsilon)] dx$$

for all continuity points $a, b \in \mathbb{R}$ of μ . The term under the square root in $m(z)$ being negative only in the set $[(1 - \sqrt{c})^2, (1 + \sqrt{c})^2]$ (and thus of non-real square root), the latter defines the support of the continuous part of the measure μ with density $\frac{\sqrt{((1 + \sqrt{c})^2 - x)(x - (1 - \sqrt{c})^2)}}{2c\pi x}$ at point x in the set. The case $x = 0$ brings a discontinuity in μ with weight equal to

$$\mu(\{0\}) = -\lim_{y \downarrow 0} iy m(iy) = \frac{c - 1}{2c} \pm \frac{c - 1}{2c}$$

where the sign is established by a second order development of $zm(z)$ in the neighborhood of zero: that is, “+” for $c > 1$ inducing a mass $1 - 1/c$ for $p > n$, or “-” for $c < 1$ in which case $\mu(\{0\}) = 0$ and μ has no mass at zero.

Detailed proof. Having heuristically identified $\bar{\mathbf{Q}}(z)$, we shall now use sound mathematical tools to prove that, indeed, $\bar{\mathbf{Q}}(z)$ is a deterministic equivalent for $\mathbf{Q}(z)$ in the sense of the theorem statement. Let us first show that $\mathbb{E}[\mathbf{Q}(z)] = \bar{\mathbf{Q}}(z) + o_{\|\cdot\|}(1)$, where $o_{\|\cdot\|}(1)$ denotes a matrix term of vanishing operator norm as $n, p \rightarrow \infty$.

Convergence in mean. For mathematical convenience, we will take $z < 0$ in what follows. Since $\mathbf{Q}(z)$ and $\bar{\mathbf{Q}}(z)$ from the theorem statement are complex analytic functions for $z \notin \mathbb{R}^+$ (matrix-valued Stieltjes transforms are analytic), by Vitali's convergence theorem (Theorem 2.3), obtaining the convergence results on \mathbb{R}^- (in fact even on a restricted local subset of \mathbb{R}^-) is equivalent to obtaining the result on all of $\mathbb{C} \setminus \mathbb{R}^+$.

We proceed in two steps by first introducing the intermediate deterministic quantities

$$\alpha(z) \equiv \frac{1}{n} \text{tr } \mathbb{E}[\mathbf{Q}_{-1}(z)], \quad \bar{\mathbf{Q}}(z) \equiv \left(-z + \frac{1}{1 + \alpha(z)} \right)^{-1} \mathbf{I}_p \quad (2.14)$$

where we denote $\mathbf{Q}_{-j}(z) \equiv (\frac{1}{n} \sum_{i \neq j} \mathbf{x}_i \mathbf{x}_i^\top - z \mathbf{I}_p)^{-1}$ the “leave-one-out” version of $\mathbf{Q}(z)$ by removing the \mathbf{x}_j contribution.

From Lemma 2.1, we have (the argument z in $\alpha(z)$, $\mathbf{Q}(z)$ and $\bar{\mathbf{Q}}(z)$ is dropped when confusion is not possible)

$$\begin{aligned} \mathbb{E}[\mathbf{Q} - \bar{\mathbf{Q}}] &= \mathbb{E}\mathbf{Q} \left(\frac{\mathbf{I}_p}{1 + \alpha} - \frac{1}{n} \mathbf{X} \mathbf{X}^\top \right) \bar{\mathbf{Q}} = \frac{\mathbb{E}[\mathbf{Q}]}{1 + \alpha} \bar{\mathbf{Q}} - \frac{1}{n} \mathbb{E}[\mathbf{Q} \mathbf{X} \mathbf{X}^\top] \bar{\mathbf{Q}} \\ &= \frac{\mathbb{E}[\mathbf{Q}]}{1 + \alpha} \bar{\mathbf{Q}} - \sum_{i=1}^n \frac{1}{n} \mathbb{E}[\mathbf{Q} \mathbf{x}_i \mathbf{x}_i^\top] \bar{\mathbf{Q}} = \frac{\mathbb{E}[\mathbf{Q}]}{1 + \alpha} \bar{\mathbf{Q}} - \sum_{i=1}^n \mathbb{E} \left[\frac{\mathbf{Q}_{-i} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i} \right] \bar{\mathbf{Q}} \end{aligned}$$

where we applied Lemma 2.8 to obtain the last equality.

Since we expect $\frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i$ to be close to α (as a consequence of Lemma 2.11), we rewrite

$$\frac{\mathbf{Q}_{-i} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i} = \frac{\mathbf{Q}_{-i} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top}{1 + \alpha} - \frac{\mathbf{Q}_{-i} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top (\frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i - \alpha)}{(1 + \alpha)(1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i)}$$

so that

$$\begin{aligned} \mathbb{E}[\mathbf{Q} - \bar{\mathbf{Q}}] &= \frac{\mathbb{E}[\mathbf{Q}]}{1 + \alpha} \bar{\mathbf{Q}} - \sum_{i=1}^n \frac{\mathbb{E} [\mathbf{Q}_{-i} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top]}{1 + \alpha} \bar{\mathbf{Q}} + \sum_{i=1}^n \frac{\mathbb{E} [\mathbf{Q} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top d_i]}{1 + \alpha} \bar{\mathbf{Q}} \\ &= \frac{\mathbb{E}[\mathbf{Q}]}{1 + \alpha} \bar{\mathbf{Q}} - \sum_{i=1}^n \frac{\mathbb{E} [\mathbf{Q}_{-i} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top]}{1 + \alpha} \bar{\mathbf{Q}} + \frac{\mathbb{E} [\mathbf{Q} \frac{1}{n} \mathbf{D} \mathbf{D}^\top]}{1 + \alpha} \bar{\mathbf{Q}} \end{aligned}$$

where we introduced $\mathbf{D} = \text{diag}\{d_i\}_{i=1}^n$ for $d_i = \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i - \alpha$, and used again Lemma 2.8 to write $\frac{\mathbf{Q}_{-i} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i} = \mathbf{Q} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top$ in the first equality. Since

$\mathbb{E}[\mathbf{Q}_{-i}\mathbf{x}_i\mathbf{x}_i^\top] = \mathbb{E}[\mathbf{Q}_{-i}]$, this further reads

$$\mathbb{E}[\mathbf{Q} - \bar{\mathbf{Q}}] = \frac{1}{n} \sum_{i=1}^n (\mathbb{E}[\mathbf{Q}] - \mathbb{E}[\mathbf{Q}_{-i}]) \frac{\bar{\mathbf{Q}}}{1+\alpha} + \frac{\mathbb{E}[\frac{1}{n}\mathbf{Q}\mathbf{X}\mathbf{D}\mathbf{X}^\top]\bar{\mathbf{Q}}}{1+\alpha}.$$

For the first right-hand side term, again from Lemmas 2.1 and 2.8,

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\mathbf{Q} - \mathbf{Q}_{-i}] &= -\frac{1}{n} \sum_{i=1}^n \mathbb{E}\left[\mathbf{Q} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top \mathbf{Q}_{-i}\right] \\ &= -\frac{1}{n} \sum_{i=1}^n \mathbb{E}\left[\mathbf{Q} \frac{1}{n} \mathbf{x}_i \mathbf{x}_i^\top \mathbf{Q} \left(1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i\right)\right] \\ &= -\frac{1}{n} \mathbb{E}\left[\mathbf{Q} \frac{1}{n} \mathbf{X} \mathbf{D}_2 \mathbf{X}^\top \mathbf{Q}\right] \end{aligned}$$

where $\mathbf{D}_2 = \text{diag}\left\{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i\right\}_{i=1}^n$ and thus

$$\mathbb{E}[\mathbf{Q} - \bar{\mathbf{Q}}] = -\frac{1}{n} \mathbb{E}\left[\mathbf{Q} \frac{1}{n} \mathbf{X} \mathbf{D}_2 \mathbf{X}^\top \mathbf{Q}\right] \frac{\bar{\mathbf{Q}}}{1+\alpha} + \frac{\mathbb{E}[\frac{1}{n}\mathbf{Q}\mathbf{X}\mathbf{D}\mathbf{X}^\top]\bar{\mathbf{Q}}}{1+\alpha}. \quad (2.15)$$

It remains to show that the right-hand side terms vanish in the large p, n limit.

For the first term, note that

$$0 \preceq \mathbf{Q} \frac{1}{n} \mathbf{X} \mathbf{D}_2 \mathbf{X}^\top \mathbf{Q} \preceq \mathbf{Q} \frac{1}{n} \mathbf{X} \mathbf{X}^\top \mathbf{Q} \max_{1 \leq i \leq n} [\mathbf{D}_2]_{ii}$$

in the order of symmetric matrices. Since $\mathbf{Q} \frac{1}{n} \mathbf{X} \mathbf{X}^\top = \mathbf{I}_p + z\mathbf{Q}$ which is of bounded operator norm (by 2) and $\|\mathbf{Q}\| \leq 1/|z|$, controlling $\|\mathbb{E}[\mathbf{Q} \frac{1}{n} \mathbf{X} \mathbf{D}_2 \mathbf{X}^\top \mathbf{Q}]\|$ boils down to controlling $\mathbb{E}[\max_i [\mathbf{D}_2]_{ii}]$. This can be established in various ways. For instance, from the union bound and the i.i.d. nature of the \mathbf{x}_i 's,

$$\mathbb{P}\left(\max_i [\mathbf{D}_2]_{ii} > t\right) \leq n \mathbb{P}([\mathbf{D}_2]_{11} > t).$$

Now, by Markov's inequality $\mathbb{P}(X > t) \leq \mathbb{E}[X^k]/t^k$ for every k (for $X, t > 0$) and the moment inequality in Lemma 2.11 for, say $k = 4$, $\mathbb{P}(\max_i [\mathbf{D}_2]_{ii} > t)$ may be bounded by a function decreasing as t^{-4} , for all t large, and of order n^{-1} . Specifically, for k even,

$$\begin{aligned} \mathbb{P}([\mathbf{D}_2]_{11} > t) &\leq \frac{\mathbb{E}[(1 + \frac{1}{n} \mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1)^k]}{t^k} \\ &= \frac{\mathbb{E}_{\mathbf{Q}_{-1}} \mathbb{E}_{\mathbf{x}_1} [(\frac{1}{n} \mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \frac{1}{n} \text{tr } \mathbf{Q}_{-1} + 1 + \frac{1}{n} \text{tr } \mathbf{Q}_{-1})^k]}{t^k} \\ &\leq 2^{k-1} \frac{\mathbb{E}_{\mathbf{Q}_{-1}} \mathbb{E}_{\mathbf{x}_1} [(\frac{1}{n} \mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \frac{1}{n} \text{tr } \mathbf{Q}_{-1})^k] + \mathbb{E}_{\mathbf{Q}_{-1}} \mathbb{E}_{\mathbf{x}_1} [(1 + \frac{1}{n} \text{tr } \mathbf{Q}_{-1})^k]}{t^k} \end{aligned}$$

where the last inequality uses Minkowski's inequality ($|a+b|^k \leq 2^{k-1}(|a|^k + |b|^k)$) and we isolated the expectation over \mathbf{Q}_{-1} from that over \mathbf{x}_1 to let appear the

difference $\frac{1}{n}\mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \frac{1}{n} \operatorname{tr} \mathbf{Q}_{-1}$ which, conditionally on \mathbf{Q}_{-1} of bounded norm, we know is small and can be controlled via Lemma 2.11:

$$\mathbb{E}_{\mathbf{x}_1} \left[\left| \frac{1}{n} \mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \frac{1}{n} \operatorname{tr} \mathbf{Q}_{-1} \right|^4 \right] \leq \frac{C}{n^4} \operatorname{tr}^2(\mathbf{Q}_{-1}^2)$$

for some constant $C > 0$ which depends on the fourth- and eighth-order moments of the entries of \mathbf{x} , but which is *independent* of n, p , according to Lemma 2.11 with $k = 4$. Since $\frac{1}{n} \operatorname{tr} \mathbf{Q}_{-1} \leq p/(n|z|)$ (note here the key boundedness property of the resolvent which simplifies the analysis, here and in most random matrix proofs), we thus have that

$$\mathbb{P}([\mathbf{D}_2]_{11} > t) \leq \frac{Cp^2}{n^4|z|^4 \cdot t^4}$$

holds for all $t > C'$ for some $C' > 0$ that depends on n, p only via their ratio p/n . Finally, since

$$\begin{aligned} \mathbb{E}[\max_i [\mathbf{D}_2]_{ii}] &= \int_0^{C'} \mathbb{P}(\max_i [\mathbf{D}_2]_{ii} > t) dt + \int_{C'}^\infty \mathbb{P}(\max_i [\mathbf{D}_2]_{ii} > t) dt \\ &\leq C' + n \int_{C'}^\infty \mathbb{P}([\mathbf{D}_2]_{11} > t) dt \leq C' + \frac{Cp^2}{n^3|z|^4} \int_{C'}^\infty t^{-4} dt < \infty \end{aligned}$$

we find that $\mathbb{E}[\max_i [\mathbf{D}_2]_{ii}]$ is bounded. Note in passing that this proves on the way that $\|\mathbb{E}[\mathbf{Q} - \bar{\mathbf{Q}}]\| = O(n^{-1})$. Consequently, due to the leading $1/n$ factor in front of the first right-hand side term of (2.15), this term vanishes as $n, p \rightarrow \infty$.⁹

To now handle the second right-hand side term in (2.15), one needs to control the norm of $\frac{1}{n} \mathbf{Q} \mathbf{X} \mathbf{D} \mathbf{X}^\top \bar{\mathbf{Q}}$. This is not a symmetric matrix, but $\mathbb{E}[\mathbf{Q} - \bar{\mathbf{Q}}]$ is. We may thus rewrite (2.15) as the half-sum of itself and its transpose and we are thus left to controlling the operator norm of $\frac{1}{n} \mathbf{Q} \mathbf{X} \mathbf{D} \mathbf{X}^\top \bar{\mathbf{Q}} + \frac{1}{n} \bar{\mathbf{Q}} \mathbf{X} \mathbf{D} \mathbf{X}^\top \mathbf{Q}$. Using the matrix inequalities $\mathbf{A}\mathbf{B}^\top + \mathbf{B}\mathbf{A}^\top \preceq \mathbf{A}\mathbf{A}^\top + \mathbf{B}\mathbf{B}^\top$ (from $(\mathbf{A} - \mathbf{B})(\mathbf{A} - \mathbf{B})^\top \succeq 0$) and $\mathbf{A}\mathbf{B}^\top + \mathbf{B}\mathbf{A}^\top \succeq -\mathbf{A}\mathbf{A}^\top - \mathbf{B}\mathbf{B}^\top$ (from $(\mathbf{A} + \mathbf{B})(\mathbf{A} + \mathbf{B})^\top \succeq 0$), we are left to bounding the norm of

$$\mathbb{E} \left[\frac{n^\varepsilon}{n} \mathbf{Q} \mathbf{X} \mathbf{D}^2 \mathbf{X}^\top \mathbf{Q} \right] + \mathbb{E} \left[\frac{n^{-\varepsilon}}{n} \bar{\mathbf{Q}} \mathbf{X} \mathbf{X}^\top \bar{\mathbf{Q}} \right]$$

where the division of the n^{-2} constant into $n^{-1+\varepsilon}$ and $n^{-1-\varepsilon}$ for some $\varepsilon \in (0, \frac{1}{2}]$ will appear as essential, since both terms do not have the same orders of magnitude (which depend on the so far unknown magnitude of the entries of

⁹Another proof option could have been to derive a moment inequality for the random variable $|\mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \operatorname{tr} \mathbf{Q}_{-1}|^k$ rather than for $|\mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \operatorname{tr} \mathbf{Q}_{-1}|^k$, which would have involved Burkholder's inequality used a bit later in the proof to control the fluctuations of $\operatorname{tr} \mathbf{Q}_{-1} - \operatorname{tr} \mathbb{E} \mathbf{Q}_{-1}$. But, as we saw, the fundamental boundedness of $\|\mathbf{Q}_{-1}\|$ discards *here* the need to control the fluctuations of \mathbf{Q}_{-1} .

\mathbf{D}). The second term above is easily seen to be of order $O(n^{-\varepsilon})$. As for the first term, we write, similar to the bound on \mathbf{D}_2 ,

$$\begin{aligned} n^\varepsilon \mathbb{E}[\|\mathbf{D}\|^2] &= n^\varepsilon \mathbb{E} \left[\max_i d_i^2 \right] \\ &\leq n^\varepsilon \int_0^{C' n^{-\theta-\varepsilon}} \mathbb{P} \left(\max_i d_i^2 > t \right) dt + n^{1+\varepsilon} \int_{C' n^{-\theta-\varepsilon}}^\infty \mathbb{P} \left(d_1^2 > t \right) dt \\ &\leq C' n^{-\theta} + n^{1+\varepsilon} \int_{C' n^{-\theta-\varepsilon}}^\infty \mathbb{P} \left(\left| \frac{1}{n} \mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \alpha \right|^2 > t \right) dt \end{aligned}$$

for some $C' > 0$ and $\theta \in (0, \frac{1}{2}]$ to be determined, $d_i = \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i - \alpha$ and $\alpha = \frac{1}{n} \text{tr } \mathbb{E}[\mathbf{Q}_{-1}] > 0$. Here, since α involves an expectation over \mathbf{Q}_{-1} (and not \mathbf{Q}_{-1} itself), one needs be more precise in the control of the fluctuations of both \mathbf{x}_1 and \mathbf{Q}_{-1} . Specifically, we write

$$\begin{aligned} &\mathbb{E} \left| \frac{1}{n} \mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \frac{1}{n} \text{tr } \mathbb{E}[\mathbf{Q}_{-1}] \right|^4 \\ &= \mathbb{E} \left| \frac{1}{n} \mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \frac{1}{n} \text{tr } \mathbf{Q}_{-1} + \frac{1}{n} \text{tr}(\mathbf{Q}_{-1} - \mathbb{E}[\mathbf{Q}_{-1}]) \right|^4 \\ &\leq \frac{8}{n^4} \mathbb{E} \left[|\mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \text{tr } \mathbf{Q}_{-1}|^4 \right] + \frac{8}{n^4} \mathbb{E} \left[|\text{tr } \mathbf{Q}_{-1} - \text{tr } \mathbb{E}[\mathbf{Q}_{-1}]|^4 \right] \end{aligned}$$

which we will show to be of order $O(n^{-2})$. For the first right-hand side term, this follows from Lemma 2.11. For the second term, which does not involve a quadratic form but the fluctuations of the columns of \mathbf{X} *inside* the intricate functional $\text{tr } \mathbf{Q}_{-1}$, we will resort to Burkholder's inequality. For the sake of further reuse, we will prove a slightly more general result on $\mathbb{E}[|\text{tr } \mathbf{Q}_{-1} - \text{tr } \mathbb{E}[\mathbf{Q}_{-1}]|^4]$: first note that we may freely replace \mathbf{Q}_{-1} with \mathbf{Q} in the result without altering the desired control, and that we may generalize the control to $\mathbb{E}[|\text{tr } \mathbf{A} \mathbf{Q}_{-1} - \text{tr } \mathbb{E}[\mathbf{A} \mathbf{Q}_{-1}]|^4]$ for arbitrary \mathbf{A} deterministic of bounded norm (again, this will be useful later).

Specifically, under the notation of Lemma 2.12, observe that we may write

$$\begin{aligned} \frac{1}{p} \text{tr } \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) &= \sum_{i=1}^n \mathbb{E}_i \left[\frac{1}{p} \text{tr } \mathbf{A} \mathbf{Q} \right] - \mathbb{E}_{i-1} \left[\frac{1}{p} \text{tr } \mathbf{A} \mathbf{Q} \right] \\ &= \frac{1}{p} \sum_{i=1}^n (\mathbb{E}_i - \mathbb{E}_{i-1}) [\text{tr } \mathbf{A}(\mathbf{Q} - \mathbf{Q}_{-i})] \end{aligned}$$

(since $\mathbb{E}_i[\text{tr } \mathbf{A} \mathbf{Q}_{-i}] = \mathbb{E}_{i-1}[\text{tr } \mathbf{A} \mathbf{Q}_{-i}]$) for \mathcal{F}_i the σ -field generating the columns $\mathbf{x}_{i+1}, \dots, \mathbf{x}_n$ of \mathbf{X} and with the convention $\mathbb{E}_0[f(\mathbf{X})] = f(\mathbf{X})$. This forms a martingale difference sequence so that we fall under the scope of Burkholder's lemma. Now, from the identity $\mathbf{Q} = \mathbf{Q}_{-i} - \frac{1}{n} \frac{\mathbf{Q}_{-i} \mathbf{x}_i \mathbf{x}_i^\top \mathbf{Q}_{-i}}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i}$ (Lemma 2.8),

$$(\mathbb{E}_i - \mathbb{E}_{i-1}) \left[\frac{1}{p} \text{tr } \mathbf{A}(\mathbf{Q} - \mathbf{Q}_{-i}) \right] = -(\mathbb{E}_i - \mathbb{E}_{i-1}) \frac{\frac{1}{pn} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{A} \mathbf{Q}_{-i} \mathbf{x}_i}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i}$$

which is order $O(p^{-1})$. As a consequence, from Lemma 2.12,

$$\mathbb{E} \left[\left| \frac{1}{p} \operatorname{tr} \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) \right|^4 \right] = O(n^{-2}).$$

Of course, this in particular implies that $\mathbb{E}[|\frac{1}{p} \operatorname{tr}(\mathbf{Q}_{-1} - \mathbb{E}\mathbf{Q}_{-1})|^4] = O(n^{-2})$, as desired.

Having obtained this desired control on the moments, it finally follows from Markov's inequality that

$$\mathbb{P} \left(\left| \frac{1}{n} \mathbf{x}_1^\top \mathbf{Q}_{-1} \mathbf{x}_1 - \frac{1}{n} \operatorname{tr} \mathbb{E}[\mathbf{Q}_{-1}] \right|^2 > t \right) \leq Ct^{-2}n^{-2}$$

for all $t > C'$ and for some constant $C', C > 0$. Therefore

$$n^\varepsilon \mathbb{E}[\|\mathbf{D}\|^2] \leq C' n^{-\theta} + CC' n^{2\varepsilon+\theta-1}.$$

By choosing for instance $\varepsilon = \theta = \frac{1}{4}$, we thus conclude that¹⁰

$$\|\mathbb{E}[\mathbf{Q}] - \bar{\mathbf{Q}}\| \leq Cn^{-1/4}, \quad \text{with } \bar{\mathbf{Q}} = \left(-z + \frac{1}{1 + \alpha(z)} \right)^{-1} \mathbf{I}_p. \quad (2.16)$$

The introduction of the intermediate deterministic equivalent $\bar{\mathbf{Q}}$ allowed us to compare \mathbf{Q} to $\bar{\mathbf{Q}}$ by exploiting the more accessible statistical relation between \mathbf{Q} and $\mathbb{E}\mathbf{Q}$. We are now in position to compare the *deterministic* matrices $\bar{\mathbf{Q}}$ and $\bar{\mathbf{Q}}$. To this end, recalling that $\bar{\mathbf{Q}}$ is defined implicitly through $\bar{\mathbf{Q}} = m(z)\mathbf{I}_p$ with $m(z) = (-z + \frac{1}{1+cm(z)})^{-1} = \frac{1}{p} \operatorname{tr} \bar{\mathbf{Q}}(z)$, we write, again with Lemma 2.1,

$$\bar{\mathbf{Q}} - \bar{\mathbf{Q}} = \frac{\alpha(z) - cm(z)}{(1 + cm(z))(1 + \alpha(z))} \bar{\mathbf{Q}} \bar{\mathbf{Q}}$$

so that

$$\begin{aligned} |\alpha(z) - cm(z)| &= \left| \frac{1}{n} \operatorname{tr} (\mathbb{E}[\mathbf{Q}_{-1}(z)] - \bar{\mathbf{Q}}(z)) + \frac{1}{n} \operatorname{tr} (\bar{\mathbf{Q}}(z) - \bar{\mathbf{Q}}(z)) \right| \\ &= |\alpha(z) - cm(z)| \cdot \frac{\frac{1}{n} \operatorname{tr} (\bar{\mathbf{Q}}(z) \bar{\mathbf{Q}}(z))}{(1 + cm(z))(1 + \alpha(z))} + O(n^{-\frac{1}{4}}) \end{aligned}$$

¹⁰The obtained bound has order $O(n^{-\frac{1}{4}})$ which is in fact suboptimal and could at least be improved to $O(n^{-\frac{1}{2}})$. It is interesting to note here that this loss in optimality follows from the very rough bound $\mathbb{P}(\max_i d_i^2 > t) \leq n \mathbb{P}(d_i^2 > t)$, which the moment of order 4 bound in $O(n^{-2})$ applied in Markov's inequality does not optimally compensates. Alternative approaches to avoid this loss are (i) either to evaluate higher moment bounds (in general the moment of order $2k$ is bounded in $O(n^{-k})$) but this comes at the cost of cumbersome calculus, or more conveniently (ii) to obtain exponential decay bounds of $\mathbb{P}(d_i^2 > t)$ of the order $O(e^{-n^\alpha})$ which automatically annihilate the polynomial loss induced by the extra factor n . Item (ii) partially justifies the relevance of a concentration of measure framework for random matrices, which we will detail in Section 2.7.

where we used the fact that $\|\mathbb{E}[\mathbf{Q}_{-1}] - \bar{\mathbf{Q}}\| \leq \|\mathbb{E}[\mathbf{Q}_{-1} - \mathbf{Q}]\| + \|\bar{\mathbf{Q}} - \mathbf{Q}\| = O(n^{-1/4})$ from (2.16). Since $\alpha(z) > 0$ for $z < 0$, we have

$$\mathbf{0} \prec \frac{\bar{\mathbf{Q}}(z)}{1 + \alpha(z)} \prec \frac{\mathbf{I}_p}{1 - z},$$

so that

$$0 < \frac{\frac{1}{n} \text{tr}(\bar{\mathbf{Q}}(z)\bar{\mathbf{Q}}(z))}{(1 + cm(z))(1 + \alpha(z))} < \frac{1}{1 - z} \frac{cm(z)}{1 + cm(z)} < 1,$$

and therefore, since $m(z) > 0$ for $z < 0$,

$$|\alpha(z) - cm(z)| \rightarrow 0,$$

which concludes the proof of (2.8).

Almost sure convergence. To now prove the almost sure convergence $\frac{1}{p} \text{tr} \mathbf{A}(\mathbf{Q} - \bar{\mathbf{Q}}) \xrightarrow{a.s.} 0$ and $\mathbf{a}^\top (\mathbf{Q} - \bar{\mathbf{Q}}) \mathbf{b} \xrightarrow{a.s.} 0$, it suffices to show

$$\frac{1}{p} \text{tr} \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) \xrightarrow{a.s.} 0, \quad \mathbf{a}^\top (\mathbf{Q} - \mathbb{E}\mathbf{Q}) \mathbf{b} \xrightarrow{a.s.} 0.$$

Both results can be proved similarly using Burkholder's inequality (which is the historical approach proposed by [Bai and Silverstein \[2010\]](#)). We have indeed already proved that $\mathbb{E}[|(1/p) \text{tr} \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q})|^4] = O(n^{-2})$ so that, from Markov's inequality (i.e., $\mathbb{P}(|X| > t) \leq \mathbb{E}[|X|^k]/t^k$) and the Borel-Cantelli lemma (i.e., $\mathbb{P}(|X_n| > t) = O(n^{-\ell})$ for some $\ell > 1$ for all $t > 0$ implies $X_n \xrightarrow{a.s.} 0$),

$$\frac{1}{p} \text{tr} \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) \xrightarrow{a.s.} 0$$

as requested. The proof of $\mathbf{a}^\top (\mathbf{Q} - \mathbb{E}\mathbf{Q}) \mathbf{b} \xrightarrow{a.s.} 0$ is obtained similarly. \square

Remark 2.3 (On the convergence rates). *In the course of the proofs above, we saw examples of a general concentration trend for linear statistics and quadratic forms of random matrices. We shall indeed typically have for most of the models of random matrices $\mathbf{X} \in \mathbb{R}^{n \times n}$ under study in this monograph that*

- *linear statistics $\frac{1}{n} \sum_{i=1}^n f(\lambda_i(\mathbf{X}))$ for sufficiently well-behaved f (so for instance $\frac{1}{n} \text{tr} \mathbf{Q}_\mathbf{X}(z) = \frac{1}{n} \sum_i (\lambda_i(\mathbf{X}) - z)^{-1}$) converge at speed $O(1/n)$ (their variance scales like $O(1/n^2)$). From a central-limit theorem viewpoint, this is as fast as it can get. Indeed, \mathbf{X} is maximally composed of $O(n^2)$ “degrees of freedom” and thus, by the central limit theorem, fluctuations are at most at speed $O(1/\sqrt{n^2}) = O(1/n)$.*
- *quadratic forms $\mathbf{a}^\top f(\mathbf{X}) \mathbf{b}$ where $f(\mathbf{X}) = \mathbf{U} \text{diag}(f(\lambda_i(\mathbf{X}))) \mathbf{U}^\top$ (in the spectral decomposition of \mathbf{X}) typically converge at a slower $O(1/\sqrt{n})$ speed.*

This remark is particularly interesting as it indicates from a statistics viewpoint that, for $\mathbf{X} \in \mathbb{R}^{p \times n}$, asymptotic approximations may gain accuracy by doubly exploiting the degrees of freedom in both the sample (n) and feature (p) directions.

Remark 2.4 (On the assumptions on \mathbf{X}). *Let us pursue here on the footnote introduced to clarify the “smooth tail condition” phrase of Theorem 2.4. The Marčenko-Pastur law has been widely generalized and several times proven using different techniques. For instance [Adamczak, 2011, O’Rourke, 2012] assume the \mathbf{X}_{ij} are “weakly” dependent in the sense that their correlation or higher order cross-moments vanish at a certain speed as $n, p \rightarrow \infty$. Alternatively, the works of Bai and Silverstein [2010] tend to assume that the entries of \mathbf{X} are not necessarily identically distributed; in this case, an additional condition on the tails $\mathbb{P}(|\mathbf{X}_{ij}| > t)$ of the probability measures of the entries (for instance a uniform bound on some moment higher than 2) is needed. In [El Karoui, 2009], El Karoui provides a first result which assumes the columns \mathbf{x}_i of $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ are independent concentrated random vectors, an assumption that we will thoroughly discuss in Section 2.7; (very) roughly speaking, concentrated random vectors $\mathbf{x} \in \mathbb{R}^p$ can be written as $\mathbf{x} = \varphi(\tilde{\mathbf{x}})$ where $\tilde{\mathbf{x}}$ has standard i.i.d. entries with either a Gaussian law or a bounded support, and $\varphi : \mathbb{R}^p \rightarrow \mathbb{R}^p$ is any 1-Lipschitz function: this assumption maintains the p degrees of freedom in \mathbf{x} (arising from $\tilde{\mathbf{x}}$) while allowing for strong correlation between the entries of \mathbf{x} . In this case, the Marčenko-Pastur law is indeed still valid if $\varphi(\mathbf{x})$ has zero mean and identity covariance.*

One may wonder how the moment conditions on the entries of \mathbf{X} could be relaxed as this seems to suggest that moment bounds can no longer be used and exponential tail decay no longer hold. The approach historically proposed by Bai and Silverstein (well documented in [Bai and Silverstein, 2010]) relies on a truncation-and-centering approach which consists in replacing \mathbf{X} by a matrix $\tilde{\mathbf{X}}$ defined as $\tilde{\mathbf{X}}_{ij} = \mathbf{X}_{ij} 1_{|\mathbf{X}_{ij}| > t(n)}$ for a certain threshold t , typically (a well chosen) function of n . Being “truncated”, the entries of $\tilde{\mathbf{X}}$ have moments of higher orders (of all orders if $t(n)$ is constant), so that moments bounds can be used on these. It then remains to show that the functional of \mathbf{X} interest (e.g., the empirical measure of the eigenvalues of $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$) is asymptotically the same as that of $\tilde{\mathbf{X}}$ as $n, p \rightarrow \infty$. Other techniques exist which prove a result on \mathbf{X} for Gaussian random variables (for instance using Stein’s identity $\mathbb{E}[wf(w)] = \mathbb{E}[f'(w)]$ for $w \sim \mathcal{N}(0, 1)$; see Lemma 2.13) before using specific controls on the deviations from the Gaussian case (such as generalized Stein’s lemma) to extrapolate between Gaussian and non-Gaussian cases. This is the subject of the next section.

2.2.2.2 The “Gaussian method” alternative

Pastur and Shcherbina [2011] propose an alternative proof scheme for Theorem 2.4, based on a two-step approach: (i) a proof for Gaussian \mathbf{X} and (ii) an interpolation method to non-Gaussian \mathbf{X} ; together known as the “Gaussian

method". Although less intuitive when compared to the Bai and Silverstein's approach presented in the previous section, this method is much more flexible as it can handle more structured random matrix models, in particular when the "guessing" part (of the ultimate deterministic equivalent $\tilde{\mathbf{Q}}$ for \mathbf{Q}) of Bai-Silverstein's method is non trivial.

The Gaussian case itself is handled in two steps (or more precisely is based on two ingredients): (i-a) convergence in means of the resolvent with Stein's lemma, and (i-b) control of the variance with the Nash–Poincaré inequality to establish convergence (in probability or almost surely).

Convergence in mean by Stein's lemma.

Lemma 2.13 (Stein [1981]). *Let $x \sim \mathcal{N}(0, 1)$ and $f : \mathbb{R} \rightarrow \mathbb{R}$ a continuously differentiable function having at most polynomial growth and such that $\mathbb{E}[f'(x)] < \infty$. Then,*

$$\mathbb{E}[xf(x)] = \mathbb{E}[f'(x)]. \quad (2.17)$$

In particular, for $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{C})$ with $\mathbf{C} \in \mathbb{R}^{p \times p}$ and $f : \mathbb{R}^p \rightarrow \mathbb{R}$ a continuously differentiable function with derivatives having at most polynomial growth with respect to p ,

$$\mathbb{E}[[\mathbf{x}]_i f(\mathbf{x})] = \sum_{j=1}^p [\mathbf{C}]_{ij} \mathbb{E}\left[\frac{\partial f(\mathbf{x})}{\partial [\mathbf{x}]_j}\right] \quad (2.18)$$

where $\partial/\partial[\mathbf{x}]_i$ indicates differentiation with respect to the i -th entry of \mathbf{x} .

The lemma, sometimes referred to as the integration-by-parts formula for Gaussian variables, simply follows from

$$\begin{aligned} \mathbb{E}[xf(x)] &= \int xf(x)e^{-\frac{1}{2}x^2} dx \\ &= [-f(x)e^{-\frac{1}{2}x^2}]_{-\infty}^{\infty} + \int f'(x)e^{-\frac{1}{2}x^2} dx = \mathbb{E}[f'(x)] \end{aligned}$$

by integration by parts $\int u'v = [uv] - \int uv'$ for $u(x) = -e^{-\frac{1}{2}x^2}$ and $v(x) = f(x)$.

To exploit Lemma 2.13, let us thus assume \mathbf{X} Gaussian, i.e., $\mathbf{X}_{ij} \sim \mathcal{N}(0, 1)$. Observe that $\mathbf{Q} = \frac{1}{z} \frac{1}{n} \mathbf{X} \mathbf{X}^\top \mathbf{Q} - \frac{1}{z} \mathbf{I}_p$, so that

$$\mathbb{E}[\mathbf{Q}_{ij}] = \frac{1}{zn} \sum_{k=1}^n \mathbb{E}[\mathbf{X}_{ik} [\mathbf{X}^\top \mathbf{Q}]_{kj}] - \frac{1}{z} \delta_{ij}$$

in which $\mathbb{E}[\mathbf{X}_{ik} [\mathbf{X}^\top \mathbf{Q}]_{kj}] = \mathbb{E}[xf(x)]$ for $x = \mathbf{X}_{ik}$ and $f(x) = [\mathbf{X}^\top \mathbf{Q}]_{kj}$. Therefore, from the lemma and the fact that $\partial \mathbf{Q} = -\frac{1}{n} \mathbf{Q} \partial(\mathbf{X} \mathbf{X}^\top) \mathbf{Q}$,¹¹

$$\begin{aligned} \mathbb{E}[\mathbf{X}_{ik} [\mathbf{X}^\top \mathbf{Q}]_{kj}] &= \mathbb{E}\left[\frac{\partial [\mathbf{X}^\top \mathbf{Q}]_{kj}}{\partial \mathbf{X}_{ik}}\right] \\ &= \mathbb{E}[\mathbf{Q}_{ij}] - \mathbb{E}\left[\frac{1}{n} [\mathbf{X}^\top \mathbf{Q} \mathbf{X}]_{kk} \mathbf{Q}_{ij}\right] - \mathbb{E}\left[\frac{1}{n} [\mathbf{X}^\top \mathbf{Q}]_{ki} [\mathbf{X}^\top \mathbf{Q}]_{kj}\right] \end{aligned}$$

¹¹This is the matrix version of $d(1/x) = -1/x^2 dx$.

so that, summing over k ,

$$\begin{aligned} \frac{1}{z} \frac{1}{n} \sum_{k=1}^n \mathbb{E}[\mathbf{X}_{ik} [\mathbf{X}^\top \mathbf{Q}]_{kj}] &= \frac{1}{z} \mathbb{E}[\mathbf{Q}_{ij}] - \frac{1}{z} \frac{1}{n^2} \mathbb{E}[\mathbf{Q}_{ij} \operatorname{tr}(\mathbf{Q} \mathbf{X} \mathbf{X}^\top)] \\ &\quad - \frac{1}{z} \frac{1}{n^2} \mathbb{E}[\mathbf{Q} \mathbf{X} \mathbf{X}^\top \mathbf{Q}]_{ij}. \end{aligned} \quad (2.19)$$

It is not too difficult to see that the rightmost term has vanishing operator norm (of order $O(1/n)$) as $n, p \rightarrow \infty$ (see later Remark 2.5, which shows that for complex-valued Gaussian \mathbf{X} this term does not even appear in the derivation). Also recall that $\operatorname{tr}(\mathbf{Q} \mathbf{X} \mathbf{X}^\top) = np + zn \operatorname{tr} \mathbf{Q}$. As a result, matrix-wise, we obtain

$$\mathbb{E}[\mathbf{Q}] + \frac{1}{z} \mathbf{I}_p = \mathbb{E}[\mathbf{X}_{\cdot k} [\mathbf{X}^\top \mathbf{Q}]_{k \cdot}] = \frac{1}{z} \mathbb{E}[\mathbf{Q}] - \frac{1}{z} \frac{1}{n} \mathbb{E}[\mathbf{Q}(p + z \operatorname{tr} \mathbf{Q})] + o_{\|\cdot\|}(1)$$

where $\mathbf{Y}_{\cdot k}$ and $\mathbf{Y}_{k \cdot}$ are the column vector $[Y_{1k}, \dots, Y_{pk}]^\top$ and row vector $[Y_{k1}, \dots, Y_{kn}]$, respectively. As $\frac{1}{n} \operatorname{tr} \mathbf{Q}$ is expected to converge to some $m(z)$, it can be taken out of the expectation in the limit so that, gathering all terms proportional to $\mathbb{E}[\mathbf{Q}]$ on the left-hand side, we finally have

$$\mathbb{E}[\mathbf{Q}](1 - p/n - z - p/nzm(z)) = \mathbf{I}_p + o_{\|\cdot\|}(1)$$

which, taking the trace to identify $m(z)$, concludes the proof for the Gaussian case.

Almost sure convergence by Nash–Poincaré inequality. To prove the almost sure convergence of traces and bilinear forms of the resolvent in the case of Gaussian \mathbf{X} , one may then use the powerful Nash–Poincaré inequality proposed by Pastur.

Lemma 2.14 (Nash–Poincaré inequality [Pastur, 2005]). *For $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{C})$ with $\mathbf{C} \in \mathbb{R}^{p \times p}$ and $f : \mathbb{R}^p \rightarrow \mathbb{R}$ continuously differentiable with derivatives having at most polynomial growth with respect to p ,*

$$\operatorname{Var}[f(\mathbf{x})] \leq 2 \sum_{i,j=1}^d [\mathbf{C}]_{ij} \mathbb{E} \left[\frac{\partial f(\mathbf{x})}{\partial x_i} \frac{\partial f(\mathbf{x})}{\partial x_j} \right].$$

In the present case, for Gaussian \mathbf{X} with $\mathbf{X}_{ij} \sim \mathcal{N}(0, 1)$,

$$\operatorname{Var} \left[\frac{1}{p} \operatorname{tr} \mathbf{A} \mathbf{Q} \right] \leq \frac{2}{p^2} \sum_{i=1}^p \sum_{j=1}^n \mathbb{E} \left[\left| \frac{\partial \operatorname{tr} \mathbf{A} \mathbf{Q}}{\partial \mathbf{X}_{ij}} \right|^2 \right].$$

Again using $\partial \mathbf{Q} = -\frac{1}{n} \mathbf{Q} \partial(\mathbf{X} \mathbf{X}^\top) \mathbf{Q}$, we find

$$\frac{\partial \operatorname{tr} \mathbf{A} \mathbf{Q}}{\partial \mathbf{X}_{ij}} = -\frac{1}{n} [\mathbf{Q} \mathbf{A} \mathbf{Q} \mathbf{X} + \mathbf{Q} \mathbf{A}^\top \mathbf{Q} \mathbf{X}]_{ij}$$

so that, from $(a + b)^2 \leq 2(a^2 + b^2)$ and $\|\mathbf{A}\| = 1$,

$$\begin{aligned} \frac{2}{p^2} \sum_{i=1}^p \sum_{j=1}^n \mathbb{E} \left[\left| \frac{\partial \text{tr } \mathbf{A}\mathbf{Q}}{\partial \mathbf{X}_{ij}} \right|^2 \right] &\leq \frac{4}{p^2 n^2} (\text{tr}(\mathbf{Q}\mathbf{A}\mathbf{Q}\mathbf{X}\mathbf{X}^\top\mathbf{Q}\mathbf{A}^\top\mathbf{Q}) \\ &\quad + \text{tr}(\mathbf{Q}\mathbf{A}^\top\mathbf{Q}\mathbf{X}\mathbf{X}^\top\mathbf{Q}\mathbf{A}\mathbf{Q})) = O(n^{-2}). \end{aligned}$$

By Markov's inequality and the Borel Cantelli lemma, we thus have that $\frac{1}{p} \text{tr } \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) \xrightarrow{a.s.} 0$.

When it comes to evaluating the fluctuations of $\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}$ with the same approach, it appears that $\text{Var}[\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}] = O(n^{-1})$ which is enough to ensure convergence in probability (by Markov's inequality) but not almost surely (as the Borel Cantelli lemma cannot be applied). Thus one needs to resort to evaluating a higher moment bound, such as $\mathbb{E}[|\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}|^4]$. To this end, we may use the fact that

$$\begin{aligned} \mathbb{E}[|\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}|^4] &= \text{Var}[|\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}|^2] + \mathbb{E}[|\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}|^2]^2 \\ &= \text{Var}[|\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}|^2] + \text{Var}[|\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}|]^2. \end{aligned}$$

Since we know that the rightmost term is of order $O(n^{-2})$, it remains to show, again through Nash–Poincaré inequality, that $\text{Var}[|\mathbf{a}^\top(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b}|^2] = O(n^{-2})$ which is a cumbersome but easily obtained result as well.

Interpolation trick to non-Gaussian \mathbf{X} . To “interpolate” these results from Gaussian \mathbf{X} to non-Gaussian \mathbf{X} , one may then use a generalized version of Stein's lemma to non-Gaussian distributions, for which we have the following lemma.

Lemma 2.15 (Interpolation trick, [Lytova and Pastur, 2009, Corollary 3.1]). *For $x \in \mathbb{R}$ a random variable with zero mean and unit variance, $y \sim \mathcal{N}(0, 1)$, and f a $(k+2)$ -times differentiable function with bounded derivatives,*

$$\mathbb{E}[f(x)] - \mathbb{E}[f(y)] = \sum_{l=2}^k \frac{\kappa_{l+1}}{2l!} \int_0^1 \mathbb{E}[f^{(l+1)}x(t)]t^{(l-1)/2} dt + \epsilon_k$$

where κ_l is the l^{th} cumulant of x , $x(t) = \sqrt{t}x + (1 - \sqrt{t})y$, and $|\epsilon_k| \leq C_k \mathbb{E}[|x|^{k+2}] \sup_t |f^{(k+2)}(t)|$ for some constant C_k only dependent on k .

All Gaussian expectations (means and variance) in the proof above can then be expressed as their non-Gaussian form up to a sum of moment control on the derivatives of f .

Remark 2.5 (Simplification in the complex case). *The Marčenko-Pastur result presented in Theorem 2.4 is also universal with respect to the field (\mathbb{R} or \mathbb{C}) of the entries of \mathbf{X} , where the Gram matrix of interest in the complex case is \mathbf{XX}^**

for \mathbf{A}^* the Hermitian conjugate (transpose conjugate) of \mathbf{A} . The resolvent is in particular now $\mathbf{Q}(z) = (\frac{1}{n}\mathbf{X}\mathbf{X}^* - z\mathbf{I}_p)^{-1}$. More interestingly, Stein's lemma, Lemma 2.13, is simplified in the complex case into

$$\mathbb{E}[\mathbf{X}_{ij}f(\mathbf{X}, \mathbf{X}^*)] = \mathbb{E}\left[\frac{d}{d\bar{\mathbf{X}}_{ij}}f(\mathbf{X}, \mathbf{X}^*)\right]$$

for $f(\mathbf{X}, \mathbf{X}^*)$ a (polynomially bounded) smooth function of both \mathbf{X} and \mathbf{X}^* , and $\bar{\mathbf{X}}_{ij}$ the complex conjugate of \mathbf{X}_{ij} , where the complex derivation rules become $(d/d\bar{x})(x) = 0$ and $(d/d\bar{x})\bar{x} = 1$ (see details in [Pastur and Shcherbina, 2011]). From these identities, we in particular find that

$$\frac{d}{d\mathbf{X}_{ij}}\mathbf{X}\mathbf{X}^* = \mathbf{E}_{ij}\mathbf{X}^*$$

for \mathbf{E}_{ij} the indicator matrix with entry $[\mathbf{E}_{ij}]_{lm} = \delta_{il}\delta_{jm}$. This relation is more convenient to use than in the real case where

$$\frac{d}{d\mathbf{X}_{ij}}\mathbf{X}\mathbf{X}^\top = \mathbf{E}_{ij}\mathbf{X}^\top + \mathbf{X}\mathbf{E}_{ij}^\top$$

and thus two terms instead of one appear; in recollection of the derivation above of the Marčenko-Pastur theorem in the real case with Stein's lemma, this extra term was anticipated to vanish (see Equation (2.19)).

This remark is particularly useful when universality is anticipated (essentially for all such “first order” deterministic equivalents) and when elaborate random matrix models are to be treated. That is, in these settings, it is convenient (at least as a preliminary exploration) to assume \mathbf{X} has complex rather than real Gaussian entries.

2.2.2.3 Wigner's semicircle law

While the Marčenko-Pastur law is at the heart of sample covariance matrix models and thus a starting point in kernel methods for machine learning, Wigner's semicircle law concerns symmetric matrices of independent entries (over and on the diagonal) which is more akin to random graphs and will be used in this monograph almost exclusively to this purpose.¹²

The main result, again presented under the form of a deterministic equivalent for the resolvent, is as follows.

Theorem 2.5 (From [Wigner, 1955]). *Let $\mathbf{X} \in \mathbb{R}^{n \times n}$ be symmetric and such that the $\mathbf{X}_{ij} \in \mathbb{R}$, $j \geq i$, are independent zero mean and unit variance random variables, with smooth tail conditions. Then, for $\mathbf{Q}(z) = (\frac{\mathbf{X}}{\sqrt{n}} - z\mathbf{I}_n)^{-1}$, as $n \rightarrow \infty$,*

$$\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z), \quad \bar{\mathbf{Q}}(z) = m(z)\mathbf{I}_n \tag{2.20}$$

¹²Up to an important exception when dealing with “properly scaling kernels” in Section 4.4.

with $(z, m(z))$ the unique solution in $\mathcal{Z}(\mathbb{C} \setminus [-2, 2])$ of

$$m^2(z) + zm(z) + 1 = 0. \quad (2.21)$$

The function $m(z)$ is the Stieltjes transform of the probability measure

$$\mu(dx) = \frac{1}{2\pi} \sqrt{(4 - x^2)^+} dx \quad (2.22)$$

which is known as the Wigner semicircle law.




Figure 2.3: Histogram of the empirical spectral measure of $\frac{\mathbf{X}}{\sqrt{n}}$ versus the Wigner semicircle law, for $n = 1\,000$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

Figure 2.3 compares the empirical spectral measure of $\frac{\mathbf{X}}{\sqrt{n}}$ defined in Theorem 2.5 with the Wigner semicircle law (which, for a proper scaling of the axes, has a half circular shape as the name suggests), for $n = 1\,000$.

Sketch of proof of Theorem 2.5. Although not the historical method of Wigner,¹³ we propose here to follow exactly the approach detailed in the proof of the Marčenko-Pastur theorem. For pedagogical interest, we provide the main heuristic arguments both for the Bai-Silverstein and for the Gaussian method.

¹³Wigner's proof in [Wigner, 1955] relied on a method of moment approach: having inferred that the limiting measure should be a semicircle, he proved via a combinatorial approach, that the successive "moments" $\frac{1}{n} \text{tr}(n^{-\frac{1}{2}} \mathbf{X})^k$ for $k = 1, 2, \dots$ must converge, as $n \rightarrow \infty$, to the moments of the semicircle measure $\int t^k \mu(dt)$. This method is simple but only useful if indeed the limiting measure μ can be inferred. In the Marčenko-Pastur case of Theorem 2.4 and even worse in more elaborate random matrix settings, the limiting measure μ is less obvious to anticipate.

Bai-Silverstein heuristic. Let $\mathbf{Q} = (\frac{\mathbf{X}}{\sqrt{n}} - z\mathbf{I}_n)^{-1}$ be the resolvent of interest and denote $\alpha = \frac{1}{n} \operatorname{tr} \mathbb{E}\mathbf{Q}$. By Lemma 2.6,

$$\mathbf{Q}_{ii} = \frac{1}{\frac{1}{\sqrt{n}}\mathbf{X}_{ii} - z - \frac{1}{n}\mathbf{x}_i^\top \mathbf{Q}_{-i}\mathbf{x}_i}$$

with $\mathbf{Q}_{-i} = (\frac{\mathbf{X}_{-i}}{\sqrt{n}} - z\mathbf{I}_{n-1})^{-1}$, $\mathbf{X}_{-i} \in \mathbb{R}^{(n-1) \times (n-1)}$ the matrix obtained by deleting the i -th row and column from \mathbf{X} , and $\mathbf{x}_i \in \mathbb{R}^{n-1}$ the i -th column (and thus the i -th row by symmetry) of \mathbf{X} with its i -th entry removed. Taking the sum over i we obtain

$$\frac{1}{n} \operatorname{tr} \mathbf{Q} = \frac{1}{n} \sum_{i=1}^n \frac{1}{\frac{1}{\sqrt{n}}\mathbf{X}_{ii} - z - \frac{1}{n}\mathbf{x}_i^\top \mathbf{Q}_{-i}\mathbf{x}_i} \simeq \frac{1}{n} \sum_{i=1}^n \frac{1}{-z - \frac{1}{n}\mathbf{x}_i^\top \mathbf{Q}_{-i}\mathbf{x}_i}$$

since $\frac{1}{\sqrt{n}}\mathbf{X}_{ii}$ asymptotically vanishes as $n \rightarrow \infty$. By Lemma 2.11 and Lemma 2.9, $\frac{1}{n}\mathbf{x}_i^\top \mathbf{Q}_{-i}\mathbf{x}_i$ should be close to $\alpha = \frac{1}{n} \operatorname{tr} \mathbb{E}\mathbf{Q}$ for large n , so that by taking the expectation we have

$$\alpha \simeq \frac{1}{-z - \alpha} + \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\frac{\frac{1}{n}\mathbf{x}_i^\top \mathbf{Q}_{-i}\mathbf{x}_i - \alpha}{(-z - \frac{1}{n}\mathbf{x}_i^\top \mathbf{Q}_{-i}\mathbf{x}_i)(-z - \alpha)} \right]$$

where we should rigorously evoke moment or concentration arguments to show that the rightmost term vanishes asymptotically as $n \rightarrow \infty$. Together with $m(z) - \alpha(z) \rightarrow 0$, this eventually gives

$$m^2(z) + zm(z) + 1 = 0$$

the solution of which is explicitly given by

$$m(z) = \frac{1}{2}(-z + \sqrt{z^2 - 4})$$

with $\sqrt{\cdot}$ chosen as the branch of the square root for which $m(z)$ is a Stieltjes transform. Taking the imaginary part and the limit when $z \rightarrow x \in \mathbb{R}$ (which is only non-zero if $x^2 - 4 < 0$) gives the form of the density $\mu(dx)$ in the theorem statement.

Gaussian method heuristic. Similar to the proof of the Marčenko-Pastur law with Gaussian methods in Section 2.2.2.2, observe that, for $\mathbf{Q} = (\frac{\mathbf{X}}{\sqrt{n}} - z\mathbf{I}_n)^{-1}$, we have

$$\frac{1}{\sqrt{n}} \mathbb{E}[\mathbf{X}\mathbf{Q}] = \mathbf{I}_n + z\mathbb{E}[\mathbf{Q}] \tag{2.23}$$

so that by Lemma 2.13 and the fact that $\partial \mathbf{Q} = -\frac{1}{\sqrt{n}} \mathbf{Q}(\partial \mathbf{X})\mathbf{Q}$,

$$\begin{aligned}\mathbb{E}[\mathbf{Q}_{ij}] &= \frac{1}{z} \frac{1}{\sqrt{n}} \sum_{k=1}^n \mathbb{E}[\mathbf{X}_{ik} \mathbf{Q}_{kj}] - \frac{1}{z} \delta_{ij} \\ &= \frac{1}{z} \frac{1}{\sqrt{n}} \sum_{k=1}^n \mathbb{E} \left[\frac{\partial \mathbf{Q}_{kj}}{\partial \mathbf{X}_{ik}} \right] - \frac{1}{z} \delta_{ij} \\ &= -\frac{1}{z} \frac{1}{n} \sum_{k=1}^n \mathbb{E}[\mathbf{Q}_{ki} \mathbf{Q}_{kj} + \mathbf{Q}_{kk} \mathbf{Q}_{ij}] - \frac{1}{z} \delta_{ij} \\ &= -\frac{1}{z} \frac{1}{n} \mathbb{E}[\mathbf{Q}_{ij}^2 + \mathbf{Q}_{ij} \operatorname{tr} \mathbf{Q}] - \frac{1}{z} \delta_{ij}\end{aligned}$$

which can be summarized in matrix form as

$$\mathbb{E}[\mathbf{Q}] \simeq -\frac{1}{z} \frac{1}{n} \mathbb{E}[\mathbf{Q}^2] - \frac{1}{z} \mathbb{E}[\mathbf{Q}] m(z) - \frac{1}{z} \mathbf{I}_n \quad (2.24)$$

where we used the fact that the random quantity $\frac{1}{n} \operatorname{tr} \mathbf{Q} \simeq \frac{1}{n} \operatorname{tr} \mathbb{E} \mathbf{Q}$ is expected to converge to some deterministic $m(z)$ and can therefore be taken out of the expectation.

Since the first term on the right-hand side asymptotically vanishes (of operator norm order $O(1/n)$) as $n, p \rightarrow \infty$,¹⁴ we reach

$$\mathbb{E}[\mathbf{Q}] \simeq -\frac{1}{z} \left(1 + \frac{1}{z} \frac{1}{n} \operatorname{tr} \mathbb{E} \mathbf{Q} \right)^{-1} \mathbf{I}_n$$

which, after taking the trace and using $m(z) \simeq \frac{1}{n} \operatorname{tr} \mathbb{E} \mathbf{Q}(z)$, gives the limiting formula

$$m^2(z) + zm(z) + 1 = 0.$$

The rest of the development is then identical to the Bai-Silverstein approach above. \square

2.2.3 Large dimensional sample covariance matrices and generalized semicircles

The Marčenko-Pastur and semicircle theorems have long been the gold-standard in both theoretical and applied random matrix theory, in the sense that most mathematical studies and practical results concerned the Wishart and Wigner random matrix models.¹⁵ But the assumption of data \mathbf{X} with i.i.d., let alone standard Gaussian, entries is often limiting. In statistics where one is interested

¹⁴Again, we could even more simply have exploited Remark 2.5 to not even produce the term $\mathbb{E}[\mathbf{Q}_{ki} \mathbf{Q}_{kj}]$ in the early development of the calculus.

¹⁵Among those studies are generalizations of the data model assumptions to matrices \mathbf{X} with non independent entries [Pajor and Pastur, 2009], studies and characterization of the limiting spectra [Silverstein and Choi, 1995], deeper considerations on the local behavior of eigenvalues [Johnstone, 2001, 2008], etc.

in the sample correlation matrix $\mathbf{X}\mathbf{X}^\top$, it is expected that the columns $\mathbf{x}_i \in \mathbb{R}^p$ of \mathbf{X} exhibit a correlation structure and be non-necessarily independent (in particular when they are samples from a time series). In graph theory, where the affinity matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ is the central object of study, one may wish to model graph patterns, degree heterogeneity, community structures, etc., which go against the i.i.d. Bernoulli assumption of so-called Erdős-Rényi graphs.

This section introduces generalizations of these results, to a level that is convenient to machine learning applications. In particular, in order to model the existence of classes within the data, \mathbf{X} will often be subdivided into subblocks that can be identified with each class.¹⁶

2.2.3.1 Large sample covariance matrix model and its generalizations

Our first result generalizes the Marčenko-Pastur law to sample covariance matrices and is originally due to a long line of works by Bai and Silverstein [Silverstein and Bai, 1995].

Theorem 2.6 (Sample covariance matrix, from [Silverstein and Bai, 1995]). *Let $\mathbf{X} = \mathbf{C}^{\frac{1}{2}}\mathbf{Z} \in \mathbb{R}^{p \times n}$ with symmetric nonnegative definite $\mathbf{C} \in \mathbb{R}^{p \times p}$ of bounded operator norm (i.e., $\limsup_p \|\mathbf{C}\| < \infty$),¹⁷ $\mathbf{Z} \in \mathbb{R}^{p \times n}$ having i.i.d. zero mean and unit variance entries with smooth tail conditions. Then, as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$, letting $\mathbf{Q}(z) = (\frac{1}{n}\mathbf{X}\mathbf{X}^\top - z\mathbf{I}_p)^{-1}$ and $\tilde{\mathbf{Q}}(z) = (\frac{1}{n}\mathbf{X}^\top\mathbf{X} - z\mathbf{I}_n)^{-1}$, we have*

$$\begin{aligned}\mathbf{Q}(z) &\leftrightarrow \bar{\mathbf{Q}}(z) = -\frac{1}{z}(\mathbf{I}_p + \tilde{m}_p(z)\mathbf{C})^{-1} \\ \tilde{\mathbf{Q}}(z) &\leftrightarrow \tilde{\bar{\mathbf{Q}}}(z) = \tilde{m}_p(z)\mathbf{I}_n\end{aligned}$$

where $(z, \tilde{m}_p(z))$ is the unique solution in $\mathcal{Z}(\mathbb{C} \setminus \mathbb{R}^+)$ of

$$\tilde{m}_p(z) = \left(-z + \frac{1}{n} \operatorname{tr} \mathbf{C} (\mathbf{I}_p + \tilde{m}_p(z)\mathbf{C})^{-1} \right)^{-1}.$$

In particular, if $\mu_{\mathbf{C}} \rightarrow \nu$ as $p \rightarrow \infty$, then $\mu_{\frac{1}{n}\mathbf{X}\mathbf{X}^\top} \xrightarrow{a.s.} \mu$, $\mu_{\frac{1}{n}\mathbf{X}^\top\mathbf{X}} \xrightarrow{a.s.} \tilde{\mu}$ as $p, n \rightarrow \infty$ where $\mu, \tilde{\mu}$ are the unique measures having Stieltjes transforms $m(z)$ and $\tilde{m}(z)$, respectively, with

$$m(z) = \frac{1}{c} \tilde{m}(z) + \frac{1-c}{cz}, \quad \tilde{m}(z) = \left(-z + c \int \frac{t\nu(dt)}{1+\tilde{m}(z)t} \right)^{-1}.$$

¹⁶A host of other results for more elaborate models exist in the literature. Many are gathered in the books [Tulino and Verdú, 2004, Couillet and Debbah, 2011]: these monographs are applied to the field of wireless communications, which has known an immense burst of interest into random matrix models in the 2000-2010 period. Some of these results have effectively been reused to form the base ground of the current wave of machine learning-applied random matrix models.

¹⁷In the original article [Silverstein and Bai, 1995], the constraint on the bounded norm of $\|\mathbf{C}\|$ is relaxed and unnecessary. Yet, this complicates the proof and is never of actual use for the purpose of this monograph.

A few remarks are in order to better understand the statement of the theorem.

Remark 2.6 (On the implicit statement). *As opposed to Theorem 2.4, the statement of the theorem is here implicit in the sense that μ is only defined through $m_\mu(z)$, itself implicitly defined as the solution of a fixed-point equation. The main reason for the explicit nature of Theorem 2.4 is that Equation (2.13), which provides the connection between $m(z)$ and a function of itself, boils down to a quadratic equation in $m(z)$ which can be solved and from which Theorem 2.1 can be applied. Due to the presence of \mathbf{C} , in the present situation, the form equivalent to (2.13) here remains implicit. This will in fact be the case of almost all generalizations of the Marčenko-Pastur and Wigner theorems to be introduced in this monograph.*

Note importantly that the uniqueness of $(z, \tilde{m}_p(z))$ is stated within the set $\mathcal{Z}(\mathbb{C} \setminus \mathbb{R}^+)$. In particular, for $z \in \mathbb{C}^+$, there exists a unique $\tilde{m}_p(z) \in \mathbb{C}^+$ solution to the implicit equation; however, nothing prevents the existence of another solution (say in $\mathbb{C}^- = \{z \mid \Im[z] < 0\}$) to exist: this solution would not correspond to the sought-for $\tilde{m}_p(z)$. Possibly most importantly, we will see in Section 2.3 that, for $(z, \tilde{m}_p(z)) \in \mathbb{R}^+ \setminus \text{supp}(\mu) \times \mathbb{R}$ (a set excluded from $\mathcal{Z}(\mathbb{C} \setminus \mathbb{R}^+)$ but where $(z, \tilde{m}_p(z))$ can be formally defined by continuity), there may exist multiple solutions to the implicit equation! Fortunately, we will see that, here again, the correct solution can be identified.

Another fortunate realization is that the sought-for $\tilde{m}_p(z)$ solution also often happens to be the only stable one, in the sense that it will often be the only one discovered by numerical methods. See Remark 2.7 below for detail.

In applications to machine learning, the value of $\tilde{m}_p(z)$ will rarely be a priority. Renaming $\tilde{\delta}_p(z) = \tilde{m}_p(z)$, we will instead more often meet the quantity $\delta_p(z) \equiv \frac{1}{n} \text{tr } \mathbf{C}(-z[\mathbf{I}_p + \tilde{\delta}_p(z)\mathbf{C}])^{-1}$ (in the vast majority of cases, for $z = -\gamma$, $\gamma \in \mathbb{R}^+$ some regularization parameter) which, from Lemma 2.11, corresponds to a deterministic equivalent to $\frac{1}{n} \mathbf{x}_0^\top \mathbf{Q}(z) \mathbf{x}_0$ for all $\mathbf{x}_0 = \mathbf{C}^{\frac{1}{2}} \mathbf{z}_0$ with $\mathbf{z}_0 \in \mathbb{R}^p$ independent of \mathbf{Z} having i.i.d. zero mean and unit variance entries: this last quantity appears in the analysis of most regularized (not necessarily linear) regression problems. Interestingly, from the theorem statement, $\delta_p(z)$ satisfies the following very elegant symmetrical coupled equation

$$\begin{aligned}\delta_p(z) &= \frac{1}{n} \text{tr } \mathbf{C}(-z[\mathbf{I}_p + \tilde{\delta}_p(z)\mathbf{C}])^{-1} \\ \tilde{\delta}_p(z) &= \frac{1}{n} \text{tr } \mathbf{I}_n(-z[\mathbf{I}_n + \delta_p(z)\mathbf{I}_n])^{-1} = -\frac{1}{z} \frac{1}{1 + \delta_p(z)}.\end{aligned}$$

Theorem 2.7 below will generalize this expression to the bi-correlated $\mathbf{C}^{\frac{1}{2}} \mathbf{Z} \tilde{\mathbf{C}}^{\frac{1}{2}}$ with \mathbf{I}_n formally replaced by $\tilde{\mathbf{C}}$ in the coupled equation.

Remark 2.7 (Numerical evaluation of $m(z)$). *Due to its implicit nature, determining $m(z)$ for $z \in \mathbb{C} \setminus \mathbb{R}^+$ requires to solve an implicit equation. Using*

contraction and analyticity arguments, it can be shown that the standard fixed-point algorithm converges, i.e.,

$$m(z) = \lim_{\ell \rightarrow \infty} m^{(\ell)}(z)$$

for $\tilde{m}^{(0)}(z) = 0$ (say) and, for $\ell \geq 0$, $m^{(\ell)}(z) = \frac{1}{c} \tilde{m}^{(\ell)}(z) + \frac{1-c}{cz}$, $\tilde{m}^{(\ell+1)}(z) = (-z + c \int \frac{t\nu(dt)}{1+\tilde{m}^{(\ell)}(z)t})^{-1}$ (or the equivalent finite dimensional version with \mathbf{C}).

One must be careful here that, since $m(z)$ is not formally defined for $z \in \text{supp}(\mu)$, the above argument does not hold in this set. Yet, the argument extends to $(\sup \text{supp}(\mu), \infty)$ where the numerical fixed-point algorithm is also stable, but trying to solve for $m(z)$ with $z \in \text{supp}(\mu)$ numerically leads to a non-converging $m^{(\ell)}(z)$ sequence. This last remark can be effectively used in practice to actually determine the right edge $\sup \text{supp}(\mu)$ of the support as being the smallest $z > 0$, starting from $+\infty$, for which the algorithm fails to converge (this can be done fast by dichotomy, starting from a left value $z_- > 0$ known to belong to the support and a large enough right value z_+).

Numerically, when evaluating $m(z)$ for $z \in \mathbb{C}^+$ close to the real axis (say for $z = x + i\epsilon$, $|\epsilon| \ll 1$), the convergence can appear to be quite slow for $x \in \text{supp}(\mu)$. A convenient workaround is to sequentially evaluate $m(z)$ for all z 's of the form $x + i\epsilon$, starting from some $z_0 = x_0 + i\epsilon$ with $x_0 \notin \text{supp}(\mu)$, then moving on to $z_1 = (x_0 + \epsilon') + i\epsilon$, then $z_2 = (x_0 + 2\epsilon') + i\epsilon$, etc., (for some $\epsilon' > 0$ small) and systematically initializing the fixed-point iterations at position z_i with the value $m(z_{i-1})$. Proceeding this way, the z_i 's for which $\Re[z_i] \in \text{supp}(\mu)$ are initialized close to the (non real) solution (however note that the procedure may fail close to a mass of the spectrum of μ , typically at $z = 0$, and may keep accumulating errors if it happens to fail to converge at any given position of the spectrum).

Remark 2.8 (Drawing μ). As shall be seen in Section 2.3, the limiting measure μ in Theorem 2.6 admits a density, which, from the inverse Stieltjes transform formula in Theorem 2.1 and Remark 2.7 above, can be approximated by solving for $m(z)$ with $z \in \mathbb{R} + i\epsilon$ for some $\epsilon > 0$ small (say $\epsilon = 10^{-5}$) and retrieving the density at x as $\frac{1}{\pi} \Im[m(x + i\epsilon)]$.

This procedure however only allows for a numerical approximation (rather than a theoretical evaluation) of μ and of its support (in particular, the support consists approximately in all values of x 's such that $|\frac{1}{\pi} \Im[m(x + i\epsilon)]| \sim \epsilon \ll 1$). Section 2.3 will go beyond this naive approach and provide an exact determination of both $\lim_{z \in \mathbb{C}^+ \rightarrow x \in \mathbb{R}^*} m(z)$ for all $x \in \mathbb{R}^*$ and the support of μ .¹⁸

Figure 2.4 depicts the empirical versus limiting measure of $\mu_{\frac{1}{n}\mathbf{X}\mathbf{X}^\top}$ for \mathbf{C} having three distinct and evenly numerous eigenvalues. In this particular setting, the limiting spectrum is composed of several connected components, with

¹⁸One may be surprised at the implicit statement that $\lim_{z \in \mathbb{C}^+ \rightarrow x \in \mathbb{R}^*} m(z)$ exists for all $x \in \mathbb{R}^*$, so in particular for $x \in \text{supp}(\mu)$ while we also stated that $m(x) = \int (t - x)^{-1} \mu(dt)$ is not formally defined for $x \in \text{supp}(\mu)$. This is not a contradiction and is, we recall, at the core of the inverse Stieltjes transform formula: the spectrum μ is precisely determined by looking at $\Im[m(z)]/\pi$ for z arbitrarily close to the real axis. We will see in Section 2.3 that, at least for the sample covariance matrix model, $\lim_{z \in \mathbb{C}^+ \rightarrow x \in \mathbb{R}^*} m(z)$ (as well as $\lim_{z \in \mathbb{C}^- \rightarrow x \in \mathbb{R}^*} m(z)$ but whose value may be different!) indeed exists while $m(x)$ itself needs not be defined.

shapes akin to the Marčenko-Pastur law. For sufficiently distinct eigenvalues of \mathbf{C} , these components are disjoint (top) while for close eigenvalues they tend to merge (middle), and for $n < p$ a Dirac mass at zero is observed and the eigenvalues spread out even further into a single large component (bottom).




Figure 2.4: Histogram of the eigenvalues of $\frac{1}{n}\mathbf{XX}^\top$, $\mathbf{X} = \mathbf{C}^{\frac{1}{2}}\mathbf{Z} \in \mathbb{R}^{p \times n}$, $Z_{ij} \sim \mathcal{N}(0, 1)$, $n = 3000$; for $p = 300$ and \mathbf{C} having spectral measure $\mu_{\mathbf{C}} = \frac{1}{3}(\delta_1 + \delta_3 + \delta_7)$ (**top**), $\mu_{\mathbf{C}} = \frac{1}{3}(\delta_1 + \delta_3 + \delta_5)$ (**middle**) and $p = 4500$ with $\mu_{\mathbf{C}} = \frac{1}{3}(\delta_1 + \delta_3 + \delta_5)$ (**bottom**). Link to code: [\[Matlab\]](#) and [\[Python\]](#).

Remark 2.9 (Deterministic equivalent for $\mu_{\frac{1}{n}\mathbf{XX}^\top}$). *The convergence result $\mu_{\frac{1}{n}\mathbf{XX}^\top} \xrightarrow{a.s.} \mu$ in Theorem 2.6 imposes that there exists a limit ν to which $\mu_{\mathbf{C}}$ converges as $p \rightarrow \infty$: this may not be practically meaningful. In generalized versions of Theorem 2.6 (see e.g., Theorem 2.8 below), even if the spectral measure of the covariance matrix does converge, $\mu_{\frac{1}{n}\mathbf{XX}^\top}$ may not have a limit.*

One may instead consider the deterministic equivalent μ_p for $\mu_{\frac{1}{n}\mathbf{XX}^\top}$ which is a sequence of probability measures for which $\text{dist}(\mu_{\frac{1}{n}\mathbf{XX}^\top}, \mu_p) \xrightarrow{a.s.} 0$ for some distance between distributions (for instance, such that $\mu_{\frac{1}{n}\mathbf{XX}^\top} - \mu_p \xrightarrow{a.s.} 0$ vaguely) as $n, p \rightarrow \infty$.

Practically speaking, since the data dimension p is in general a fixed quan-

tity and \mathbf{C} a given covariance matrix (rather than specific values in a growing sequence of p 's and \mathbf{C} 's), one will always consider that the “effective” limiting measure ν actually coincides with (or is “frozen” to) $\mu_{\mathbf{C}} = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C})}$.

Sketch of proof of Theorem 2.6. The proof of Theorem 2.6 generally follows the same line of arguments as that of Theorem 2.4. The main difference is that (2.16) here becomes

$$\frac{1}{n} \mathbf{x}_i^\top \bar{\mathbf{Q}} \mathbf{A} \mathbf{Q} \mathbf{x}_i = \frac{\frac{1}{n} \mathbf{x}_i^\top \bar{\mathbf{Q}} \mathbf{A} \mathbf{Q}_{-i} \mathbf{x}_i}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i} \simeq \frac{\frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \mathbf{A} \mathbf{Q}_{-i} \mathbf{C}}{1 + \frac{1}{n} \operatorname{tr} \mathbf{Q}_{-i} \mathbf{C}}$$

where we used the fact that, denoting $\mathbf{x}_i = \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$ for \mathbf{z}_i the i -th column of \mathbf{Z} having i.i.d. zero mean and unit variance entries, by Lemma 2.11,

$$\frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i = \frac{1}{n} \mathbf{z}_i^\top \mathbf{C}^{\frac{1}{2}} \mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i \simeq \frac{1}{n} \operatorname{tr} \mathbf{Q}_{-i} \mathbf{C}.$$

Again with Lemma 2.9 and the fact that $\frac{1}{n} \operatorname{tr} \mathbf{Q}_{-i} \mathbf{C} \leq \|\mathbf{C}\| \frac{1}{n} \operatorname{tr} \mathbf{Q}_{-i}$, we obtain the approximation

$$\frac{1}{n} \operatorname{tr} (\mathbf{F} + z \mathbf{I}_p) \bar{\mathbf{Q}} \mathbf{A} \mathbf{Q} \simeq \frac{\frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}} \mathbf{A} \mathbf{Q}}{1 + \frac{1}{n} \operatorname{tr} \mathbf{Q} \mathbf{C}}$$

with $\mathbf{F}(z) = \bar{\mathbf{Q}}^{-1}(z)$ the sought-for deterministic equivalent, which then must admit the form

$$\mathbf{F}(z) \simeq \frac{\mathbf{C}}{1 + \frac{1}{n} \operatorname{tr} \mathbf{Q} \mathbf{C}} - z \mathbf{I}_p$$

for the previous approximation to hold. Unlike in the proof of the Marčenko-Pastur theorem, we see here the new term $\frac{1}{n} \operatorname{tr} \mathbf{Q} \mathbf{C}$ appear, which thus needs be studied. Fortunately, taking $\mathbf{A} = \mathbf{C}$ in $\frac{1}{n} \operatorname{tr} \mathbf{A}(\mathbf{Q} - \bar{\mathbf{Q}}) \xrightarrow{a.s.} 0$ induces a closed form equation: specifically,

$$\frac{1}{n} \operatorname{tr} \mathbf{C} \mathbf{Q} \simeq \frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}} \simeq \frac{1}{n} \operatorname{tr} \mathbf{C} \left(-z \mathbf{I}_p + \frac{\mathbf{C}}{1 + \frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}}} \right)^{-1} \quad (2.25)$$

from which we obtain

$$\tilde{m}_p(z) = \left(-z + \frac{1}{n} \operatorname{tr} \mathbf{C} (\mathbf{I}_p + \tilde{m}_p(z) \mathbf{C})^{-1} \right)^{-1}$$

if we denote $\tilde{m}_p(z) = -\frac{1}{z} (1 + \frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}}(z))^{-1}$, as requested. Note that we implicitly used here the fact that $\|\mathbf{C}\|$ is bounded.

With a deterministic equivalent $\bar{\mathbf{Q}} = \mathbf{F}^{-1}$ for \mathbf{Q} at hand, a corresponding deterministic equivalent for $\tilde{\mathbf{Q}}$ follows from the direct observation that $\tilde{\mathbf{Q}} = \frac{1}{z} \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q} \mathbf{x}_j - \frac{1}{z} \mathbf{I}_n$, so that

$$\begin{aligned} \tilde{\mathbf{Q}}_{ij} &= \frac{1}{z} \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q} \mathbf{x}_j - \frac{1}{z} \delta_{ij} = \frac{1}{z} \frac{\frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_j}{1 + \frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i} \mathbf{x}_i} - \frac{1}{z} \delta_{ij} \\ &\simeq \frac{1}{z} \frac{\frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}}}{1 + \frac{1}{n} \operatorname{tr} \mathbf{C} \mathbf{Q}} \delta_{ij} - \frac{1}{z} \delta_{ij} = -\frac{1}{z} \left(1 + \frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}} \right)^{-1} \delta_{ij} = \tilde{m}_p(z) \delta_{ij} \end{aligned}$$

and thus, for $\mathbf{A} \in \mathbb{R}^{n \times n}$ deterministic of bounded norm, applying the operator $\frac{1}{n} \sum_{i,j=1}^n A_{ij} \cdot$ on both sides (one must be careful to ensure that the pointwise-‘ \simeq ’ symbol indeed still holds under this operator), we confirm that $\tilde{\mathbf{Q}}(z) \equiv \tilde{m}_p(z)\mathbf{I}_n$ is indeed a deterministic equivalent for $\tilde{\mathbf{Q}}$. \square

Remark 2.10 (On singular population covariances). *It is interesting to note from Theorem 2.6 that, if the population covariance \mathbf{C} contains some zero eigenvalues, for example if $\mu_{\mathbf{C}} \rightarrow \nu$ as $p \rightarrow \infty$ with*

$$\nu(dx) = (1 - c_{\nu})\delta_0(x) + c_{\nu}\tilde{\nu}(dx)$$

for $c_{\nu} \in (0, 1)$ and $\tilde{\nu}$ another probability measure, then (as properly shown in [Silverstein and Choi, 1995]) $\tilde{\mu}(\{0\}) = \max(0, 1 - cc_{\nu})$. This further implies

$$\mu(\{0\}) = \begin{cases} 1 - c_{\nu} & \text{for } cc_{\nu} \leq 1 \\ 1 - c^{-1} & \text{otherwise.} \end{cases}$$

This result differs from the systematic $\mu(\{0\}) = \max(0, 1 - c^{-1})$ in the Marčenko-Pastur scenario, and takes into consideration the intrinsic dimension $c_{\nu}p$ of the random vector $\mathbf{C}^{\frac{1}{2}}\mathbf{z}_i \in \mathbb{R}^p$.

In machine learning applications, the data covariance structure \mathbf{C} may contain a wide range of very small eigenvalues, a behavior suggesting that the data representation is of much smaller effective dimension. It is interesting to observe that Theorem 2.6, in its expression function of \mathbf{C} , in fact does not depend on p/n itself but on $\frac{1}{n} \operatorname{tr} \mathbf{C}(\mathbf{I}_p + \tilde{m}_p(z)\mathbf{C})^{-1}$: the effective data dimension is thus encapsulated within \mathbf{C} in (a non-trivial manner in) the fixed-point expression.

When the data $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ arise from a time series, or when each data sample is weighted by an independent coefficient (as shall be seen in Section 3.3 on robust statistical methods), the sample covariance matrix model is not sufficiently expressive but can be generalized to the *bi-correlated* model

$$\frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \tilde{\mathbf{C}} \mathbf{Z}^T \mathbf{C}^{\frac{1}{2}}.$$

for $\mathbf{C} \in \mathbb{R}^{p \times p}$ and $\tilde{\mathbf{C}} \in \mathbb{R}^{n \times n}$ two nonnegative definite matrices and \mathbf{Z}_{ij} i.i.d. random variables with zero mean and unit variance. In particular, for \mathbf{Z} Gaussian and $\tilde{\mathbf{C}}^{\frac{1}{2}}$ Toeplitz (i.e., such that $[\tilde{\mathbf{C}}^{\frac{1}{2}}]_{ij} = \alpha_{|i-j|}$ for some sequence $\alpha_0, \dots, \alpha_{n-1}$), the columns of $\mathbf{Z} \tilde{\mathbf{C}}^{\frac{1}{2}}$ model a first order auto-regressive process [Hamilton, 1994].¹⁹

For this model, we have the following theorem.

¹⁹In passing, Toeplitz matrices involved in time series analyses also exhibit interesting large dimensional behavior. In [Gray, 2006], Gray shows that, under some decay condition on the sequence $\{\alpha_i\}_{i=1}^n$, their spectral behavior is the same as that of equivalent *circulant* matrices, the latter having the nice property to be diagonalizable in the Fourier basis: the asymptotic eigenvalues of the Toeplitz matrix are in particular the coefficients of the discrete Fourier transform of the series $\alpha_0, \alpha_1, \dots$

Theorem 2.7 (Bi-correlated model, from [Paul and Silverstein, 2009]). *Let $\mathbf{Z} \in \mathbb{R}^{p \times n}$ be a random matrix with i.i.d. zero mean, unit variance and smooth tail entries, and $\mathbf{C} \in \mathbb{R}^{p \times p}$, $\tilde{\mathbf{C}} \in \mathbb{R}^{n \times n}$ be symmetric nonnegative definite matrices with bounded operator norm. Then, as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$, letting $\mathbf{Q}(z) = (\frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \tilde{\mathbf{C}} \mathbf{Z}^T \mathbf{C}^{\frac{1}{2}} - z \mathbf{I}_p)^{-1}$ and $\tilde{\mathbf{Q}}(z) = (\frac{1}{n} \tilde{\mathbf{C}}^{\frac{1}{2}} \mathbf{Z}^T \mathbf{C} \mathbf{Z} \tilde{\mathbf{C}}^{\frac{1}{2}} - z \mathbf{I}_n)^{-1}$, we have*

$$\begin{aligned}\mathbf{Q}(z) &\leftrightarrow \bar{\mathbf{Q}}(z) = -\frac{1}{z} \left(\mathbf{I}_p + \tilde{\delta}_p(z) \mathbf{C} \right)^{-1} \\ \tilde{\mathbf{Q}}(z) &\leftrightarrow \bar{\tilde{\mathbf{Q}}}(z) = -\frac{1}{z} \left(\mathbf{I}_n + \delta_p(z) \tilde{\mathbf{C}} \right)^{-1}\end{aligned}$$

with $(z, \delta_p(z)), (z, \tilde{\delta}_p(z)) \in \mathcal{Z}(\mathbb{C} \setminus \mathbb{R}^+)$ unique solutions to

$$\delta_p(z) = \frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}}(z), \quad \tilde{\delta}_p(z) = \frac{1}{n} \operatorname{tr} \tilde{\mathbf{C}} \bar{\tilde{\mathbf{Q}}}(z).$$

In particular, if $\mu_{\mathbf{C}} \rightarrow \nu$ and $\mu_{\tilde{\mathbf{C}}} \rightarrow \tilde{\nu}$, then $\mu_{\frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \tilde{\mathbf{C}} \mathbf{Z}^T \mathbf{C}^{\frac{1}{2}}} \rightarrow \mu$ and $\mu_{\frac{1}{n} \tilde{\mathbf{C}}^{\frac{1}{2}} \mathbf{Z}^T \mathbf{C} \mathbf{Z} \tilde{\mathbf{C}}^{\frac{1}{2}}} \rightarrow \tilde{\mu}$ where $\mu, \tilde{\mu}$ are defined by their Stieltjes transforms $m(z)$ and $\tilde{m}(z)$ given by

$$m(z) = -\frac{1}{z} \int \frac{\nu(dt)}{1 + \tilde{\delta}(z)t}, \quad \tilde{m}(z) = -\frac{1}{z} \int \frac{\tilde{\nu}(dt)}{1 + \delta(z)t}$$

where $(z, \delta(z)), (z, \tilde{\delta}(z))$ are the unique solutions in $\mathcal{Z}(\mathbb{C} \setminus \mathbb{R}^+)$ to

$$\delta(z) = -\frac{c}{z} \int \frac{t\nu(dt)}{1 + \tilde{\delta}(z)t}, \quad \tilde{\delta}(z) = -\frac{1}{z} \int \frac{t\tilde{\nu}(dt)}{1 + \delta(z)t}.$$

Sketch of proof of Theorem 2.7. For simplicity and readability, only the case where both \mathbf{C} and $\tilde{\mathbf{C}}$ are diagonal is presented here.²⁰ In this case, similar to the decomposition performed in Theorem 2.6, one has the following symmetric re-expression of $\mathbf{Q}(z)$ and $\tilde{\mathbf{Q}}(z)$

$$\begin{aligned}\mathbf{Q}(z) &= \left(\frac{1}{n} \sum_{i=1}^n \mathbf{C}^{\frac{1}{2}} \tilde{\mathbf{y}}_i (\mathbf{C}^{\frac{1}{2}} \tilde{\mathbf{y}}_i)^T - z \mathbf{I}_p \right)^{-1} \\ \tilde{\mathbf{Q}}(z) &= \left(\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{C}}^{\frac{1}{2}} \mathbf{y}_i (\tilde{\mathbf{C}}^{\frac{1}{2}} \mathbf{y}_i)^T - z \mathbf{I}_n \right)^{-1}\end{aligned}$$

where we denote $\tilde{\mathbf{y}}_i \in \mathbb{R}^p$ the i -th column of $\mathbf{Z} \tilde{\mathbf{C}}^{\frac{1}{2}}$ and $\mathbf{y}_i \in \mathbb{R}^n$ the i -th column of $\mathbf{Z}^T \mathbf{C}^{\frac{1}{2}}$ so that, for \mathbf{C} and $\tilde{\mathbf{C}}$ both diagonal, one has $\tilde{\mathbf{y}}_i = \tilde{\mathbf{C}}_{ii}^{\frac{1}{2}} \mathbf{z}_i$ and $\mathbf{y}_i = \mathbf{C}_{ii}^{\frac{1}{2}} \tilde{\mathbf{z}}_i$ with $\mathbf{z}_i \in \mathbb{R}^p$ the i -th column and $\tilde{\mathbf{z}}_i \in \mathbb{R}^n$ the i -th row of $\mathbf{Z} \in \mathbb{R}^{p \times n}$.

²⁰Note that, if \mathbf{Z} is standard Gaussian, then $\mathbf{Z} \tilde{\mathbf{C}} \mathbf{Z}^T$ has the same distribution as $\mathbf{Z} \mathbf{U} \tilde{\mathbf{C}} \mathbf{U}^T \mathbf{Z}^T$ for any unitary matrix \mathbf{U} (because $\mathbf{Z} \sim \mathbf{Z} \mathbf{U}$ in law). We may then allow $\tilde{\mathbf{C}}$ to be diagonal by specifically choosing \mathbf{U} to be a matrix of eigenvectors of $\tilde{\mathbf{C}}$. By the universality of random matrix results with respect to the law of the independent entries of \mathbf{Z} , this should be sufficient to retrieve the result for any \mathbf{Z} . The same remark symmetrically holds for \mathbf{C} .

As a consequence, with $\bar{\mathbf{Q}}(z) = \mathbf{F}(z)^{-1}$ and $\tilde{\bar{\mathbf{Q}}}(z) = \tilde{\mathbf{F}}(z)^{-1}$, one obtains again with Lemma 2.8 that

$$\begin{aligned}\mathbf{Q}(z) - \bar{\mathbf{Q}}(z) &= \mathbf{Q}(z) \left(\mathbf{F}(z) + z\mathbf{I}_p - \frac{1}{n} \sum_{i=1}^n \mathbf{C}^{\frac{1}{2}} \tilde{\mathbf{y}}_i (\mathbf{C}^{\frac{1}{2}} \tilde{\mathbf{y}}_i)^T \right) \bar{\mathbf{Q}}(z) \\ &= \mathbf{Q}(\mathbf{F} + z\mathbf{I}_p) \bar{\mathbf{Q}} - \frac{1}{n} \sum_{i=1}^n \frac{\mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} \tilde{\mathbf{C}}_{ii} \mathbf{z}_i \mathbf{z}_i^T \mathbf{C}^{\frac{1}{2}} \bar{\mathbf{Q}}}{1 + \frac{1}{n} \mathbf{C}_{ii} \mathbf{z}_i^T \mathbf{C}^{\frac{1}{2}} \mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i} \\ \tilde{\mathbf{Q}}(z) - \tilde{\bar{\mathbf{Q}}}(z) &= \tilde{\mathbf{Q}}(z) \left(\tilde{\mathbf{F}}(z) + z\mathbf{I}_n - \frac{1}{n} \sum_{i=1}^p \tilde{\mathbf{C}}^{\frac{1}{2}} \mathbf{y}_i (\tilde{\mathbf{C}}^{\frac{1}{2}} \mathbf{y}_i)^T \right) \tilde{\bar{\mathbf{Q}}}(z) \\ &= \tilde{\mathbf{Q}}(\tilde{\mathbf{F}} + z\mathbf{I}_n) \tilde{\bar{\mathbf{Q}}} - \frac{1}{n} \sum_{i=1}^p \frac{\tilde{\mathbf{Q}}_{-i} \tilde{\mathbf{C}}^{\frac{1}{2}} \mathbf{C}_{ii} \tilde{\mathbf{z}}_i \tilde{\mathbf{z}}_i^T \tilde{\mathbf{C}}^{\frac{1}{2}} \tilde{\bar{\mathbf{Q}}}}{1 + \frac{1}{n} \tilde{\mathbf{C}}_{ii} \tilde{\mathbf{z}}_i^T \tilde{\mathbf{C}}^{\frac{1}{2}} \tilde{\mathbf{Q}}_{-i} \tilde{\mathbf{C}}^{\frac{1}{2}} \tilde{\mathbf{z}}_i}\end{aligned}$$

where we denote $\mathbf{Q}_{-i}(z) \equiv \left(\frac{1}{n} \sum_{j \neq i}^n \mathbf{C}^{\frac{1}{2}} \tilde{\mathbf{C}}_{jj} \mathbf{z}_j \mathbf{z}_j^T \mathbf{C}^{\frac{1}{2}} - z\mathbf{I}_p \right)^{-1}$ and symmetrically $\tilde{\mathbf{Q}}_{-i}(z) \equiv \left(\frac{1}{n} \sum_{j \neq i}^p \tilde{\mathbf{C}}^{\frac{1}{2}} \mathbf{C}_{jj} \tilde{\mathbf{z}}_j \tilde{\mathbf{z}}_j^T \tilde{\mathbf{C}}^{\frac{1}{2}} - z\mathbf{I}_n \right)^{-1}$, which are independent of \mathbf{z}_i and $\tilde{\mathbf{z}}_i$, respectively.

With this independence of \mathbf{Q}_{-i} on \mathbf{z}_i and $\tilde{\mathbf{Q}}_{-i}$ on $\tilde{\mathbf{z}}_i$, one deduces again with Lemma 2.11 that

$$\begin{aligned}\frac{1}{n} \tilde{\mathbf{C}}_{ii} \mathbf{z}_i^T \mathbf{C}^{\frac{1}{2}} \mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i - \frac{1}{n} \tilde{\mathbf{C}}_{ii} \text{tr}(\mathbf{Q}_{-i} \mathbf{C}) &\xrightarrow{a.s.} 0 \\ \frac{1}{n} \mathbf{C}_{ii} \tilde{\mathbf{z}}_i^T \tilde{\mathbf{C}}^{\frac{1}{2}} \tilde{\mathbf{Q}}_{-i} \tilde{\mathbf{C}}^{\frac{1}{2}} \tilde{\mathbf{z}}_i - \frac{1}{n} \mathbf{C}_{ii} \text{tr}(\tilde{\mathbf{Q}}_{-i} \tilde{\mathbf{C}}) &\xrightarrow{a.s.} 0\end{aligned}$$

so that $\mathbf{F}(z)$ and $\tilde{\mathbf{F}}(z)$ must take the forms

$$\begin{aligned}\mathbf{F}(z) &= \frac{1}{n} \sum_{i=1}^n \frac{\tilde{\mathbf{C}}_{ii}}{1 + \tilde{\mathbf{C}}_{ii} \frac{1}{n} \text{tr}(\mathbf{Q}_{-i} \mathbf{C})} \mathbf{C} - z\mathbf{I}_p \simeq \frac{1}{n} \sum_{i=1}^n \frac{\tilde{\mathbf{C}}_{ii}}{1 + \tilde{\mathbf{C}}_{ii} \frac{1}{n} \text{tr} \mathbf{C} \bar{\mathbf{Q}}} \mathbf{C} - z\mathbf{I}_p \\ \tilde{\mathbf{F}}(z) &= \frac{1}{n} \sum_{i=1}^p \frac{\mathbf{C}_{ii}}{1 + \mathbf{C}_{ii} \frac{1}{n} \text{tr}(\tilde{\mathbf{Q}}_{-i} \tilde{\mathbf{C}})} \tilde{\mathbf{C}} - z\mathbf{I}_n \simeq \frac{1}{n} \sum_{i=1}^p \frac{\mathbf{C}_{ii}}{1 + \mathbf{C}_{ii} \frac{1}{n} \text{tr} \tilde{\mathbf{C}} \tilde{\bar{\mathbf{Q}}}} \tilde{\mathbf{C}} - z\mathbf{I}_n.\end{aligned}$$

Denoting $\delta_p(z) = \frac{1}{n} \text{tr} \mathbf{C} \bar{\mathbf{Q}}(z)$ and $\tilde{\delta}_p(z) = \frac{1}{n} \text{tr} \tilde{\mathbf{C}} \tilde{\bar{\mathbf{Q}}}(z)$, this can be further reduced to

$$\begin{aligned}\bar{\mathbf{Q}}(z) &= \mathbf{F}(z)^{-1} \simeq -\frac{1}{z} \left(\mathbf{I}_p - \frac{1}{z} \frac{1}{n} \sum_{i=1}^n \frac{\tilde{\mathbf{C}}_{ii}}{1 + \tilde{\mathbf{C}}_{ii} \delta_p(z)} \mathbf{C} \right)^{-1} \\ \tilde{\bar{\mathbf{Q}}}(z) &= \tilde{\mathbf{F}}(z)^{-1} \simeq -\frac{1}{z} \left(\mathbf{I}_n - \frac{1}{z} \frac{1}{n} \sum_{i=1}^p \frac{\mathbf{C}_{ii}}{1 + \mathbf{C}_{ii} \tilde{\delta}_p(z)} \tilde{\mathbf{C}} \right)^{-1}.\end{aligned}$$

To eventually close the loop and obtain the sought-for relation on $(\delta_p, \tilde{\delta}_p)$, one may plug the above approximation into the definition of δ_p and $\tilde{\delta}_p$ to obtain the

following symmetric equation

$$\delta_p(z) \simeq -\frac{1}{z} \frac{1}{n} \sum_{i=1}^p \frac{\mathbf{C}_{ii}}{1 - \frac{1}{z} \frac{1}{n} \sum_{j=1}^n \frac{\tilde{\mathbf{C}}_{jj}}{1 + \tilde{\mathbf{C}}_{jj} \delta_p(z)}}$$

$$\tilde{\delta}_p(z) \simeq -\frac{1}{z} \frac{1}{n} \sum_{i=1}^n \frac{\tilde{\mathbf{C}}_{ii}}{1 - \frac{1}{z} \frac{1}{n} \sum_{j=1}^p \frac{\mathbf{C}_{jj}}{1 + \mathbf{C}_{jj} \delta_p(z)}}$$

which retrieves the expressions of Theorem 2.7. \square

As already hinted at when commenting on Theorem 2.6, it is quite interesting to note the almost perfect symmetry in the equations for the resolvent and co-resolvent in the bi-correlated model. From a machine learning perspective, wherein $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \tilde{\mathbf{C}}^{\frac{1}{2}}$ are the observed data, this symmetry between “space” and “time” correlations, or between the sample covariance matrix \mathbf{XX}^T and the kernel matrix $\mathbf{X}^T \mathbf{X}$, will often allow for a natural connection between results in the spatial (e.g., PCA, subspace methods) and in the temporal (classification, regression) domains.

From a technical angle, by the trace lemma (Lemma 2.11), we immediately find that the functions $\delta_p(z)$ and $\tilde{\delta}_p(z)$ (which also happen to be Stieltjes transforms of finite measures on \mathbb{R}^+) are respectively deterministic equivalents for $\frac{1}{n} \mathbf{x}_0^T \mathbf{Q}(z) \mathbf{x}_0$ and $\frac{1}{n} \tilde{\mathbf{x}}_0^T \tilde{\mathbf{Q}}(z) \tilde{\mathbf{x}}_0$ for $\mathbf{x}_0 = \mathbf{C}^{\frac{1}{2}} \mathbf{z}_0$, $\tilde{\mathbf{x}}_0 = \tilde{\mathbf{C}}^{\frac{1}{2}} \tilde{\mathbf{z}}_0$ and $\mathbf{z}_0 \in \mathbb{R}^p$, $\tilde{\mathbf{z}}_0 \in \mathbb{R}^n$ vectors of independent zero mean and unit variance entries, both independent of \mathbf{Z} . These quadratic forms will naturally arise in various applications of statistical inference and regression: particularly for $z = -\gamma$ with $\gamma > 0$ a regularization parameter, $\mathbf{C}^{\frac{1}{2}} \mathbf{z}_i \in \mathbb{R}^p$ the data and $\tilde{\mathbf{C}}_{ii}$ ($\tilde{\mathbf{C}}$ will usually be diagonal) an effective weight parameter induced by the algorithm under study on data $\mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$.

As pointed out above, the Gram matrix $\mathbf{X}^T \mathbf{X}$ is directly connected to kernel matrices of the type $\mathbf{K} = \{\frac{1}{p} \mathbf{x}_i^T \mathbf{x}_j\}_{i,j=1}^n = \frac{1}{p} \mathbf{X}^T \mathbf{X}$ (inner-product kernels) and even to $\mathbf{K} = \{\frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2\}_{i,j=1}^n$ (distance kernels) since $\frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 = \frac{1}{p} \|\mathbf{x}_i\|^2 + \frac{1}{p} \|\mathbf{x}_j\|^2 - 2 \frac{1}{p} \mathbf{x}_i^T \mathbf{x}_j$ which also involves the matrix $\frac{1}{p} \mathbf{X}^T \mathbf{X}$.²¹ Assuming, as is the basic setting in a multi-class machine learning classification context, that the vectors \mathbf{x}_i arise from a mixture model, the following generalization of Theorem 2.6 will thus be of practical relevance.

Theorem 2.8 (Sample covariance of k -class mixture models, from [Benaych-Georges and Couillet, 2016]). *Let $\mathbf{X} = [\mathbf{X}^{(1)}, \dots, \mathbf{X}^{(k)}] \in \mathbb{R}^{p \times n}$ with $\mathbf{X}^{(a)} = [\mathbf{x}_1^{(a)}, \dots, \mathbf{x}_{n_a}^{(a)}] \in \mathbb{R}^{p \times n_a}$ and $\mathbf{x}_i^{(a)} = \mathbf{C}_a^{\frac{1}{2}} \mathbf{z}_i^{(a)}$ for $\mathbf{z}_i^{(a)}$ a vector with i.i.d. zero mean, unit variance and smooth tail entries. Then, as $n_a, p \rightarrow \infty$ in such a way that k is fixed, $p/n \rightarrow c \in (0, \infty)$, and $n_a/n \rightarrow c_a \in (0, 1)$ for $a \in \{1, \dots, k\}$,*

²¹The prefactor $1/p$ is necessary under our notation framework to ensure that the spectrum of \mathbf{K} remains of order $O(1)$ as p, n increase.

letting $\mathbf{Q}(z) = (\frac{1}{n}\mathbf{XX}^\top - z\mathbf{I}_p)^{-1}$ and $\tilde{\mathbf{Q}}(z) = (\frac{1}{n}\mathbf{X}^\top\mathbf{X} - z\mathbf{I}_n)^{-1}$, we have²²

$$\begin{aligned}\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z) &= -\frac{1}{z} \left(\mathbf{I}_p + \sum_{a=1}^k c_a \tilde{g}_a(z) \mathbf{C}_a \right)^{-1} \\ \tilde{\mathbf{Q}}(z) \leftrightarrow \bar{\tilde{\mathbf{Q}}}(z) &= \text{diag}\{\tilde{g}_a(z)\mathbf{1}_{n_a}\}_{a=1}^k\end{aligned}$$

with $(z, \tilde{g}_a(z))$, $a \in \{1, \dots, k\}$, the unique solutions in $\mathcal{Z}(\mathbb{C} \setminus \mathbb{R}^+)$ of

$$\tilde{g}_a(z) = -\frac{1}{z}(1 + g_a(z))^{-1}, \quad g_a(z) = -\frac{1}{z} \frac{1}{n} \text{tr } \mathbf{C}_a \left(\mathbf{I}_p + \sum_{b=1}^k c_b \tilde{g}_b(z) \mathbf{C}_b \right)^{-1}.$$

Sketch of proof of Theorem 2.8. Similar to the proof of Theorem 2.6, we obtain, with the initial guess $\bar{\mathbf{Q}}(z) = \mathbf{F}^{-1}(z)$, that

$$\mathbf{Q} - \bar{\mathbf{Q}} = \mathbf{Q} \left(\mathbf{F} + z\mathbf{I}_p - \frac{1}{n} \sum_{a=1}^k \sum_{i=1}^{n_a} \mathbf{x}_i^{(a)} (\mathbf{x}_i^{(a)})^\top \right) \bar{\mathbf{Q}}$$

which, unlike in the proof of Theorem 2.6, contains a sum over a due to the different class covariances \mathbf{C}_a . To establish $\frac{1}{n} \text{tr } \mathbf{A}(\mathbf{Q} - \bar{\mathbf{Q}}) \xrightarrow{a.s.} 0$, one must have

$$\frac{1}{n} \text{tr}(\mathbf{F} + z\mathbf{I}_p) \bar{\mathbf{Q}} \mathbf{A} \mathbf{Q} - \frac{1}{n} \sum_{a=1}^k \sum_{i=1}^{n_a} \frac{1}{n} (\mathbf{x}_i^{(a)})^\top \bar{\mathbf{Q}} \mathbf{A} \mathbf{Q} \mathbf{x}_i^{(a)} \xrightarrow{a.s.} 0.$$

Applying Lemma 2.8 to remove the dependence in \mathbf{Q} of $\mathbf{x}_i^{(a)}$, together with Lemma 2.9, we deduce

$$\frac{1}{n} \sum_{a=1}^k \sum_{i=1}^{n_a} \frac{1}{n} (\mathbf{x}_i^{(a)})^\top \bar{\mathbf{Q}} \mathbf{A} \mathbf{Q} \mathbf{x}_i^{(a)} \simeq \sum_{a=1}^k \frac{n_a}{n} \frac{\frac{1}{n} \text{tr } \mathbf{C}_a \bar{\mathbf{Q}} \mathbf{A} \bar{\mathbf{Q}}}{1 + \frac{1}{n} \text{tr } \bar{\mathbf{Q}} \mathbf{C}_a}$$

so that \mathbf{F} must be written as the sum over a

$$\mathbf{F} \simeq \sum_{a=1}^k c_a \frac{\mathbf{C}_a}{1 + \frac{1}{n} \text{tr } \bar{\mathbf{Q}} \mathbf{C}_a} - z\mathbf{I}_p$$

which produces the k terms $\frac{1}{n} \text{tr } \bar{\mathbf{Q}} \mathbf{C}_a$. To identify these terms and close the loop, we take $\mathbf{A} = \mathbf{C}_b$ for each $b \in \{1, \dots, k\}$ to establish

$$\begin{aligned}\frac{1}{n} \text{tr } \mathbf{C}_b \mathbf{Q} &\simeq \frac{1}{n} \text{tr } \mathbf{C}_b \bar{\mathbf{Q}} \equiv g_b(z) \simeq \frac{1}{n} \text{tr } \mathbf{C}_b \left(-z\mathbf{I}_p + \sum_{a=1}^k c_a \frac{\mathbf{C}_a}{1 + \frac{1}{n} \text{tr } \bar{\mathbf{Q}} \mathbf{C}_a} \right)^{-1} \\ &\equiv -\frac{1}{z} \frac{1}{n} \text{tr } \mathbf{C}_b \left(\mathbf{I}_p + \sum_{a=1}^k c_a \tilde{g}_a(z) \mathbf{C}_a \right)\end{aligned}$$

²²Here $\text{diag}(\mathbf{v}_a)_{a=1}^k$ is a diagonal matrix with, on the diagonal the concatenated vector $\mathbf{v} = [\mathbf{v}_1^\top, \dots, \mathbf{v}_k^\top]^\top$; and $\mathbf{1}_{n_a} \in \mathbb{R}^{n_a}$ is the n_a -dimensional vector of all ones.

where we denoted $\tilde{g}_a(z) \equiv -\frac{1}{z} \left(1 + \frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \mathbf{C}_a\right)^{-1} = -\frac{1}{z} (1 + g_a(z))^{-1}$, as desired. This thus produces a k -dimensional *vector* equation linking the $g_a(z)$'s rather than a scalar one.

To finally derive a deterministic equivalent of $\tilde{\mathbf{Q}}$ from that of \mathbf{Q} , we use the fact that $\tilde{\mathbf{Q}} = \frac{1}{z} \frac{1}{n} \mathbf{X}^\top \mathbf{Q} \mathbf{X} - \frac{1}{z} \mathbf{I}_n$ and therefore, indexing the set $\{1, \dots, n\}$ as $\{(1)1, \dots, (1)n_1, \dots, (k)1, \dots, (k)n_k\}$,

$$\begin{aligned} \mathbf{Q}_{(a)i,(b)j} &= \frac{1}{z} \frac{1}{n} (\mathbf{x}_i^{(a)})^\top \mathbf{Q} \mathbf{x}_j^{(b)} - \frac{1}{z} \delta_{(a)i,(b)j} \\ &\simeq -\frac{1}{z} \left(1 + \frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \mathbf{C}_a\right)^{-1} \delta_{(a)i,(b)j} = \tilde{g}_a(z) \delta_{(a)i,(b)j} \end{aligned}$$

which, after applying $\frac{1}{n} \operatorname{tr} \mathbf{A} \cdot$ on both sides for \mathbf{A} deterministic bounded, concludes the proof of Theorem 2.8. \square

With some further control, Theorem 2.8 may in fact be extended to $k = n$, i.e., each data vector \mathbf{x}_i has its own, possibly distinct, covariance matrix, as shown in [Wagner et al., 2012]. When the covariance matrices are diagonal, this is then equivalent to letting \mathbf{X} have a *variance profile*, i.e., the \mathbf{X}_{ij} 's are then all independent with zero mean and variance $\sigma_{ij}^2 \equiv [\mathbf{C}_i]_{jj}$ (with $\mathbf{C}_i = \mathbb{E}[\mathbf{x}_i \mathbf{x}_i^\top]$), a setting studied in depth in [Hachem et al., 2007] but originally found in [Girko, 2001].

The application of a variance profile to random matrices with independent entries finds an even more relevant application to Wigner matrices, as detailed next.

2.2.3.2 Generalized semicircle law with a variance profile

Similar to the large sample covariance matrix model, generalizations also exist for the Wigner semicircle law in Theorem 2.5. In the following theorem, a variance profile for the entries of the symmetric random matrix is taken into account.

Theorem 2.9 (From [Pastur and Shcherbina, 2011]). *Let $\mathbf{X} \in \mathbb{R}^{n \times n}$ be symmetric and such that \mathbf{X}_{ij} , $j \geq i$, is of zero mean, bounded variance $\operatorname{Var}[\mathbf{X}_{ij}] = \sigma_{ij}^2$, and smooth tail conditions. Then, for $\mathbf{Q}(z) = (\frac{\mathbf{X}}{\sqrt{n}} - z\mathbf{I}_n)^{-1}$, we have*

$$\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z), \quad \bar{\mathbf{Q}}(z) = \operatorname{diag} \left\{ \frac{1}{-z - g_i(z)} \right\}_{i=1}^n \quad (2.26)$$

with $(z, g_i(z)) \in \mathcal{Z}(\mathbb{C} \setminus \mathbb{R}^+)$, $i \in \{1, \dots, n\}$, uniquely determined by

$$g_i(z) = \frac{1}{n} \sum_{j=1}^n \frac{\sigma_{ij}^2}{-z - g_j(z)}.$$

Sketch of proof of Theorem 2.9. Basing ourselves on the Gaussian approach, the proof of Theorem 2.9 differs from that of Theorem 2.5 in the application

of Lemma 2.13. Taking into consideration the variance $\mathbb{E}[\mathbf{X}_{ik}^2] = \sigma_{ik}^2$, Equation (2.23) gives

$$\begin{aligned}\mathbb{E}[\mathbf{Q}_{ij}] &= \frac{1}{z} \frac{1}{\sqrt{n}} \sum_{k=1}^n \mathbb{E}[\mathbf{X}_{ik}^2] \mathbb{E}\left[\frac{\partial \mathbf{Q}_{kj}}{\partial \mathbf{X}_{ik}}\right] - \frac{1}{z} \delta_{ij} \\ &= -\frac{1}{z} \frac{1}{n} \sum_{k=1}^n \sigma_{ik}^2 \mathbb{E}[\mathbf{Q}_{ki} \mathbf{Q}_{kj} + \mathbf{Q}_{kk} \mathbf{Q}_{ij}] - \frac{1}{z} \delta_{ij} \\ &= -\frac{1}{z} \frac{1}{n} \mathbb{E}[\mathbf{Q} \Sigma_i \mathbf{Q}]_{ij} - \frac{1}{z} \frac{1}{n} \mathbb{E}[\text{tr}(\Sigma_i \mathbf{Q}) \mathbf{Q}_{ij}] - \frac{1}{z} \delta_{ij}\end{aligned}$$

with $\Sigma_i \equiv \text{diag}(\sigma_{ik}^2)_{k=1}^n$, so that $\|\Sigma_i\| = O(1)$ uniformly over all i .

Note that the semicircle law in Theorem 2.5 is indeed a special case with $\sigma_{ij}^2 = \delta_{ij}$ and $\Sigma_i = \mathbf{I}_n$. As a consequence, similar to the term $\frac{1}{n} \mathbb{E}[\mathbf{Q}^2]$ in (2.24), the first term on the right-hand side vanishes as $n, p \rightarrow \infty$. Following the same reasoning, the random variable $\frac{1}{n} \text{tr} \Sigma_i \mathbf{Q}(z)$ essentially plays the role of $\frac{1}{n} \text{tr} \mathbf{Q}(z)$ in (2.24) and is expected to converge to some deterministic $g_i(z) \equiv \frac{1}{n} \text{tr} \Sigma_i \bar{\mathbf{Q}}(z)$ which can be taken out of the expectation. This gives, in matrix form

$$\mathbb{E}[\mathbf{Q}(z)] \simeq -\frac{1}{z} \text{diag}\{g_i(z)\}_{i=1}^n \mathbb{E}[\mathbf{Q}(z)] - \frac{1}{z} \mathbf{I}_n.$$

Solving this equation for $\mathbb{E}[\mathbf{Q}(z)] \simeq \bar{\mathbf{Q}}(z)$ we conclude the proof of Theorem 2.9. \square

Theorem 2.9 plays a significant role in the study of random graphs, with applications to community detection in large graphs or networks. We shall come back to this model in more detail later in Section 7.1.

Summarizing, this lengthy first technical section provided the necessary technical ingredients, along with several key results, to study the (large n, p) spectrum of “data sample matrices” from the data population statistics. In Section 2.4, we will seek to go backwards, trying to infer the population statistics from the observed *empirical* spectrum of the samples. To this end though, subtle supplementary results on the limiting spectra must be introduced. This is the objective of the next section.

The subsequent section, possibly the most technical of this part of the monograph, may be skipped at first read, the main ideas of Section 2.4 being understandable if some results are admitted. Yet, for a clear and rigorous treatment of the limitations of statistical inference, the reader will need to grasp the notions of Section 2.3 below.

2.3 Advanced spectrum considerations for sample covariances

As opposed to the Marčenko-Pastur result, Theorem 2.4, the generalized sample covariance matrix model of Theorem 2.6 (and beyond) only provides a characterization of the limiting spectral measure μ of $\mu_{\frac{1}{n} \mathbf{XX}^\top}$ (or a deterministic

equivalent μ_p for it) through its Stieltjes transform $m(z)$ for $z \in \mathbb{C} \setminus \mathbb{R}^+$ (respectively, through a sequence $m_p(z)$ of Stieltjes transforms) which itself assumes an implicit form. Since the Stieltjes transform inversion formula (Theorem 2.1) involves the limit of $m(z)$ for $z \rightarrow x \in \mathbb{R}$, the sole information about $m(z)$ for all $z \in \mathbb{C} \setminus \mathbb{R}^+$ does not immediately quantifies the measure μ .

From a theoretical standpoint, one may wonder whether μ admits a density as in the Marčenko-Pastur case and, if so, whether one can determine this density and its exact support. As recalled in Remarks 2.7 and 2.8, the density of μ (provided it exists) can be “numerically depicted” by solving for $m(z)$ with z close to, but formally *away* from, the real axis. We aim here at a more theoretical and precise characterization of μ .

From a practical standpoint, a fundamental byproduct of this characterization is the introduction of the function $z \mapsto -\frac{1}{m(z)}$ which plays a key role in statistical inference. Indeed, we shall see in Section 2.4 and the many applications of Chapter 3 that statistical information related to the population covariance \mathbf{C} (such as functionals of its eigenvalues, projections on its eigenvectors) can be accessed from the measurement matrix \mathbf{X} by means of a complex integral method involving the change of variable $z \mapsto -\frac{1}{m(z)}$.

2.3.1 Limiting spectrum

In [Silverstein and Choi, 1995] (generalized later in [Couillet and Hachem, 2014] with a more systematic approach), the authors prove that, for any measure ν (the limiting spectral distribution of \mathbf{C}), the limiting measures μ and $\tilde{\mu}$ introduced in Theorem 2.6 indeed have a density with a well-defined support.²³

2.3.1.1 Density and support of μ (and $\tilde{\mu}$)

Precisely, recall that $\mu = \frac{1}{c}\tilde{\mu} + (1 - \frac{1}{c})\delta_0$ (with δ_x the Dirac mass at x) with $\tilde{\mu}$ defined by its Stieltjes transform $\tilde{m}(z)$ solution to

$$\tilde{m}(z) = \left(-z + c \int \frac{t\nu(dt)}{1+t\tilde{m}(z)} \right)^{-1}.$$

This functional expression has the interesting key property of being invertible, in the sense that it is formally equivalent to

$$z = -\frac{1}{\tilde{m}(z)} + c \int \frac{t\nu(dt)}{1+t\tilde{m}(z)}.$$

²³It may come as very surprising but very few works in the random matrix literature have actually studied the actual spectrum of the limiting measure μ of advanced random matrix models. The few exceptions are the works [Silverstein and Choi, 1995, Couillet and Hachem, 2014] which study the defining equation of the Stieltjes transform m_μ of μ associated to the sample covariance matrix models $\mathbf{C}^{\frac{1}{2}}\mathbf{XX}^\top\mathbf{C}^{\frac{1}{2}}$ and $\mathbf{C}^{\frac{1}{2}}\mathbf{X}\tilde{\mathbf{C}}\mathbf{X}^\top\mathbf{C}^{\frac{1}{2}}$, respectively, as well as the very extensive work [Ajanki et al., 2019] on the defining equation of m_μ attached to generalized Wigner models (for instance the deformed semicircle law for Wigner models with a variance profile, Theorem 2.9). The small number of these studies testifies of the greater importance of the Stieltjes transform relation defining m_μ over the measure μ itself which, both in theory and in practice, is quite often of lesser interest.

As a consequence, the function $\tilde{m}(\cdot) : \mathbb{C} \setminus \text{supp}(\tilde{\mu}) \rightarrow \mathbb{C}$, $z \mapsto \tilde{m}(z)$ admits the functional inverse

$$\begin{aligned} z(\cdot) : \tilde{m}(\mathbb{C} \setminus \text{supp}(\tilde{\mu})) &\rightarrow \mathbb{C} \\ \tilde{m} &\mapsto -\frac{1}{\tilde{m}} + c \int \frac{t\nu(dt)}{1+t\tilde{m}}. \end{aligned}$$

The important point to notice here is that $z(\cdot)$, seen as the functional inverse of $\tilde{m}(\cdot)$, is only defined on the domain $\tilde{m}(\mathbb{C} \setminus \text{supp}(\tilde{\mu}))$. Yet, formally, this function could be *extended* to all values $\tilde{m} \in \mathbb{C}$ such that $0 \notin 1 + \tilde{m} \cdot \text{supp}(\nu)$ (i.e., all values that do not cancel the denominator $1 + t\tilde{m}$ for some $t \in \text{supp}(\nu)$).

The idea of Silverstein and Choi, originally expressed in the seminal work of Marčenko and Pastur [Marcenko and Pastur, 1967], is twofold:

- **Outside the support.** (i) the Stieltjes transform $m_\mu(x) = \int (t-x)^{-1} \mu(dt)$ of a measure μ is well defined and an increasing function on its restriction to $x \in \mathbb{R} \setminus \text{supp}(\mu)$ (it has positive derivative there), hence (ii) so must be its functional inverse $x(\cdot)$ on its restriction to $m_\mu(\mathbb{R} \setminus \text{supp}(\mu))$, (iii) consequently, if $x(\cdot)$ admits an extension to some domain \mathcal{S} with $m_\mu(\mathbb{R} \setminus \text{supp}(\mu)) \subset \mathcal{S} \subset \mathbb{R}$, $x(\cdot)$ should only be increasing on $m_\mu(\mathbb{R} \setminus \text{supp}(\mu))$;²⁴ (iv) therefore, the complementary $\mathbb{R} \setminus \text{supp}(\mu)$ to the support of μ can be determined as the union of the image of all increasing sections of $x(\cdot)$. See Figure 2.5, commented below, for a simplified visual understanding.

In our setting, this thus defines the support of the limiting measure μ of $\mu_{\frac{1}{n}\mathbf{XX}^\top}$.

- **In the support.** Inside this support, one then needs to determine the density of μ . To this end, one may first prove the existence of $\tilde{m}^\circ(x) = \lim_{\epsilon \rightarrow 0} \tilde{m}(x + i\epsilon)$. Upon existence, since $\Im[\tilde{m}^\circ(x)] > 0$ for $x \in \text{supp}(\mu)$, dominated convergence can be applied on the defining equation for $\tilde{m}(z)$ to find that $\tilde{m}^\circ(x)$ is a solution with positive imaginary part of

$$\tilde{m}^\circ(x) = \left(-x + c \int \frac{t\nu(dt)}{1+\tilde{m}^\circ(x)t} \right)^{-1}$$

which is then shown to be unique.

These arguments are formally stated in the following theorem.

Theorem 2.10 (From [Silverstein and Choi, 1995]). *Under the setting of Theorem 2.6 with $\mu_C \rightarrow \nu$, define*

$$\begin{aligned} x(\cdot) : \mathbb{R} \setminus \{\tilde{m} \mid (-1/\tilde{m}) \in \text{supp}(\nu)\} &\rightarrow \mathbb{R} \\ \tilde{m} &\mapsto -\frac{1}{\tilde{m}} + c \int \frac{t\nu(dt)}{1+\tilde{m}t}. \end{aligned}$$

²⁴Formally, it is clear that all decreasing sections of (the extended version of) $x(\cdot)$ cannot correspond to the functional inverse of a Stieltjes transform. It is less evident though that all increasing sections do correspond to the inverse of a Stieltjes transform; this was settled in [Silverstein and Choi, 1995].

Then, $\tilde{\mu}$ has a density \tilde{f} on $\mathbb{R} \setminus \{0\}$ and

- for $y \in \text{supp}(\tilde{\mu})$, $\tilde{f}(y) = \frac{1}{\pi} \Im[\tilde{m}^\circ(y)]$ with $\tilde{m}^\circ(y)$ the unique solution with positive imaginary part of $x(\tilde{m}^\circ(y)) = y$;
- the support $\text{supp}(\tilde{\mu}) \setminus \{0\}$, which coincides with $\text{supp}(\mu) \setminus \{0\}$, is defined by

$$\begin{aligned} & \text{supp}(\mu) \setminus \{0\} \\ &= \mathbb{R} \setminus \{x(\tilde{m}) \mid (-1/\tilde{m}) \in \mathbb{R} \setminus \{\text{supp}(\nu) \cup \{0\}\} \text{ and } x'(\tilde{m}) > 0\}. \end{aligned}$$

Figure 2.5 depicts the function $x(\tilde{m})$ under a similar setting as Figure 2.4 with ν composed of three Dirac masses. The top display shows four increasing regions of $x(\cdot)$, thus corresponding (on the y -axis) to four connected components of $\mathbb{R} \setminus \text{supp}(\mu)$. The complementary, depicted in blue on the y -axis, corresponds to the (three) connected components of $\text{supp}(\mu)$. The middle display only shows three growing regions for $x(\cdot)$, thus restricting the support of μ to two connected components. Analogously, in the bottom display there is only one growing region for $x(\cdot)$ (close to the y -axis from above), which now corresponds to a single connected component for $\text{supp}(\mu) \setminus \{0\}$. This is in accordance with the observations made in Figure 2.4, when altering either ν or c .

A careful analysis of the function $x(\cdot)$ actually reveals additional interesting properties:

1. the restriction of $x(\cdot)$ to its growing sections is a growing function. This follows from the fact that, there, $x(\cdot)$ is the functional inverse of $\tilde{m}(\cdot)$ restricted to $\mathbb{R} \setminus \text{supp}(\mu)$ which is a growing function.
2. in the case of Figure 2.5, since ν is discrete, $x(\cdot)$ presents asymptotes at each $-1/t$, $t \in \text{supp}(\nu)$. Thus, from the previous item, $\text{supp}(\mu)$ is here determined by the union $\cup_k [\tilde{m}_k^-, \tilde{m}_k^+]$ for $\tilde{m}_1^- < \tilde{m}_1^+ < \tilde{m}_2^- < \dots$ the successive values of \tilde{m} such that $x'(\tilde{m}) = 0$. This remark may however not hold for ν with continuous support. Detailed conditions for this characterization to hold are discussed in [Coullet and Hachem, 2014].
3. the derivative of $x(\cdot)$ is given by

$$x'(\tilde{m}) = \frac{1}{\tilde{m}^2} - c \int \frac{t^2 \nu(dt)}{(1 + t\tilde{m})^2}$$

and thus $\tilde{m}^2 x'(\tilde{m})$ converges to $1 - c$ as $|\tilde{m}| \rightarrow \infty$, while $x(\tilde{m}) \rightarrow 0$. Thus $x(\cdot)$ is either decreasing or increasing at $\pm\infty$ depending on whether $c < 1$ or $c > 1$. In particular, the pre-image by $x(\cdot)$ of 0^+ is $-\infty$ if $c < 1$ (top and middle displays of Figure 2.5) and some positive value if $c > 1$ (bottom display of Figure 2.5): this remark is fundamental for the next section.




Figure 2.5: $x(\tilde{m})$ for $-1/\tilde{m} \in \mathbb{R} \setminus \text{supp}(\nu)$, with $\nu = \frac{1}{3}(\delta_1 + \delta_3 + \delta_7)$ (**top**) and $\nu = \frac{1}{3}(\delta_1 + \delta_3 + \delta_5)$ (**middle**), $c = 1/10$ in both cases, and $\nu = \frac{1}{3}(\delta_1 + \delta_3 + \delta_5)$ with $c = 2$ (**bottom**). Local extrema are marked by circles, inflection points by squares. The support of μ can be read on the vertical axes. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

2.3.1.2 Variable change: relating $\text{supp}(\nu)$ and $\text{supp}(\mu)$.

An important side consequence of the study above of $z(\cdot)$ (and its restriction $x(\cdot)$ to the real axis) is that the function

$$\begin{aligned} \gamma : \mathbb{C} \setminus \{\text{supp}(\mu) \cup \{0\}\} &\rightarrow \mathbb{C} \\ z = z(\tilde{m}) &\mapsto -\frac{1}{\tilde{m}} \end{aligned} \tag{2.27}$$

provides an *injective* mapping between points outside the support of μ and points outside the support of ν with the property that

$$\gamma(\mathbb{C} \setminus \mathbb{R}) \subset \mathbb{C} \setminus \mathbb{R} \quad \text{and} \quad \gamma(\mathbb{R} \setminus \text{supp}(\mu)) \subset \mathbb{R} \setminus \text{supp}(\nu)$$

but where the *inclusion is strict* in general.

To understand this statement, first consider $z \in \mathbb{C} \setminus \mathbb{R}^+$. Then, by Theorem 2.6, there exists a unique pair $(z, \tilde{m}(z)) \in \mathcal{Z}$ and we may thus write $z = z(\tilde{m})$ for the value $\tilde{m} \in \mathbb{C} \setminus \mathbb{R}^-$ given by $\tilde{m} = \tilde{m}(z)$. For $z = x \in \mathbb{R}^+ \setminus \text{supp}(\mu)$, we have just seen in our discussion of Theorem 2.10 and in Figure 2.5 that there also exists $\tilde{m} \in \mathbb{R}^-$ (it must be real because $\Im[\tilde{m}(x)] = 0$ outside the support) such that $x = x(\tilde{m})$. As a consequence, for $z \in \mathbb{C} \setminus \mathbb{R}$, $\tilde{m} = \tilde{m}(z) \in \mathbb{C} \setminus \mathbb{R}$ and thus $-1/\tilde{m} \in \mathbb{C} \setminus \mathbb{R}$. Similarly, for $x \in \mathbb{R} \setminus \text{supp}(\mu)$, from Figure 2.5, $-1/\tilde{m} \in \mathbb{R} \setminus \text{supp}(\nu)$. The map is however only injective (in general not surjective) as not all values of $\mathbb{C} \setminus \text{supp}(\nu)$ can be reached. For instance, in Figure 2.5, the sets $(-1/\tilde{m}_1^-, 1)$ and $(1, -1/\tilde{m}_1^+)$ cannot be reached by γ . This remark will constitute a fundamental limitation to statistical inference methods.

More visually, Figure 2.6 depicts in blue the complementary to the image $\gamma(\mathbb{C} \setminus \text{supp}(\mu))$. This blue region is inaccessible in the sense that no point in $\mathbb{C} \setminus \text{supp}(\mu)$ can have an image by $\gamma(\cdot)$ in it. In red are depicted typical images by $\gamma(\cdot)$ of rectangular contours surrounding $\text{supp}(\mu)$. Intuitively, we observe that, as c increases (compare left to right displays), the exclusion region increases in size and one thus cannot get “too close” to the support of ν (which is here the discrete union of three point masses): this “pushes” the image of the red contour further away from the real axis.

In particular, for $c > 1$, the exclusion region includes $\{0\}$. This is a consequence of Item 3 in the remarks of the previous paragraph: while the right real crossing of a contour $\Gamma_\mu \subset \{z \in \mathbb{C}, \Re[z] > 0\}$ surrounding the support of μ will have an image by $\gamma(\cdot)$ somewhere on the right side of $\text{supp}(\nu)$, (i) for $c < 1$, the left real crossing will have 0^+ for image, and (ii) for $c > 1$, the left real crossing will have a negative value for image.

This, we shall see next in Section 2.4, is an important problem when it comes to estimating certain functionals $\int f d\nu$ of ν based on the sample measure $\mu_n^\perp \mathbf{X}\mathbf{X}^\top$.




Figure 2.6: Domain of validity of variable changes, for $\nu = \frac{1}{3}(\delta_1 + \delta_3 + \delta_5)$, with $c = \frac{1}{10}$ (**left**) and $c = 2$ (**right**). The filled region in the bottom display is the (inaccessible) complementary to the image of $-1/\tilde{m}(\cdot)$. The **red** contour Γ_ν is the image by $-1/\tilde{m}(\cdot)$ of a rectangular contour Γ_μ surrounding $\text{supp}(\mu)$. [Link to code]

2.3.2 “No eigenvalue outside the support”

Before exploiting the aforementioned change of variable (the mapping $z \mapsto -1/\tilde{m}(z)$) for statistical inference, an important extension of Theorem 2.6 is needed.

It must be stressed that the limiting results of Theorem 2.6 are *weak convergences* for the *normalized counting measure* $\frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\frac{1}{n}\mathbf{XX}^\top)}$ (i.e., the spectral measure) of the eigenvalues of $\frac{1}{n}\mathbf{XX}^\top$. This, by definition, means that, for every continuous bounded f ,

$$\frac{1}{p} \sum_{i=1}^p f\left(\lambda_i\left(\frac{1}{n}\mathbf{XX}^\top\right)\right) - \int f(t)\mu(dt) \xrightarrow{\text{a.s.}} 0.$$

Letting for instance f be a smoothed version of the indicator $1_{[a,b]}$ for $a, b \notin$

$\text{supp}(\mu)$, this thus only says that the *averaged* number of eigenvalues of $\frac{1}{n}\mathbf{XX}^\top$ within $[a, b]$ converges to $\mu([a, b])$.

In the example of Figure 2.4 (top or middle), if p_1 is the number of eigenvalues falling in the neighborhood of the leftmost connected component of μ , we thus only know that $p_1/p = 1/3 + o(1)$ (almost surely), that is $p_1 = p/3 + o(p)$. This thus does *not* guarantee that $p_1 - p/3 \xrightarrow{a.s.} 0$ exactly as $n, p \rightarrow \infty$.

Worse, Theorem 2.6 only guarantees that, for $[a, b]$ a connected component of $\mathbb{R} \setminus \text{supp}(\mu)$, the number of eigenvalues of $\frac{1}{n}\mathbf{XX}^\top$ inside $[a, b]$ is asymptotically of order $o(p)$. As such, $[a, b]$ may never be empty, even for arbitrarily large n, p (it can contain a fixed finite number of eigenvalues or even a growing number of eigenvalues, so long that this number is much less than $O(p)$).

The following result, again originally due to Bai and Silverstein, settles this non-trivial issue.

Theorem 2.11 (“No eigenvalue outside the support” and “exact separation”: from [Bai and Silverstein, 1998, 1999, Bai et al., 1988]). *Under the setting of Theorem 2.6,²⁵ let $\|\mathbf{C}\|$ be bounded with $\mu_{\mathbf{C}} \rightarrow \nu$ and $\max_{1 \leq i \leq p} \text{dist}(\lambda_i(\mathbf{C}), \text{supp}(\nu)) \rightarrow 0$ as $p \rightarrow \infty$. Consider also $-\infty \leq a < b \leq \infty$ such that $a, b \in \mathbb{R}^+ \setminus \text{supp}(\mu)$. Then the following results hold*

- if $\mathbb{E}[|\mathbf{Z}_{ij}|^4] < \infty$, then

$$\left| \left\{ \lambda_i \left(\frac{1}{n} \mathbf{XX}^\top \right) \in [a, b] \right\} - |\{\lambda_i(\mathbf{C}) \in [\gamma(a), \gamma(b)]\}| \right| \xrightarrow{a.s.} 0$$

with $\gamma(\cdot)$ defined by (2.27). In particular, if $[a, b]$ is a connected component of $\mathbb{R}^+ \setminus \text{supp}(\mu)$, then

$$\left| \left\{ \lambda_i \left(\frac{1}{n} \mathbf{XX}^\top \right) \in [a, b] \right\} \right| \xrightarrow{a.s.} 0.$$

- if $\mathbb{E}[\mathbf{Z}_{ij}^4] = \infty$,

$$\max_{1 \leq i \leq p} \lambda_i \left(\frac{1}{n} \mathbf{XX}^\top \right) \xrightarrow{a.s.} \infty.$$

In plain words, the theorem precisely states that:

- under the condition that $\mathbb{E}[\mathbf{Z}_{ij}^4] < \infty$ and that no eigenvalue of \mathbf{C} isolates from its associated limiting spectrum ν , (i) there asymptotically exists no eigenvalue outside the support of μ and (ii) the eigenvalues assembled in asymptotically contiguous “bulks” are found in asymptotically expected numbers. For instance, in the setting of Figure 2.4, it can be verified that not a single eigenvalue is found away from the support of μ and,

²⁵Here formally, the theorem statement must be understood with the “smooth tail condition” discarded, i.e., the only condition on \mathbf{Z} is that it is composed of i.i.d. entries with zero mean and unit variance.

in addition, that the exact number of eigenvalues in the neighborhood of each connected component of μ is in exact proportion (for the top figure, exactly $p/3$ eigenvalues in each component, and for the middle figure, exactly $2p/3$ eigenvalues in the largest component).

- if $\mathbb{E}[\mathbf{Z}_{ij}^4] = \infty$ (for instance for a Student t-distribution with low degree of freedom), this “exact separation” collapses: while in correct asymptotic proportion, up to $o(p)$ eigenvalues may be found away from the support of μ , with in particular the largest eigenvalue going to infinity.

For future reference, we insist on the condition

$$\max_{1 \leq i \leq p} \text{dist}(\lambda_i(\mathbf{C}), \text{supp}(\nu)) \rightarrow 0$$

which is also fundamental for the theorem to hold. Not surprisingly, if a single eigenvalue of \mathbf{C} were to diverge as $p \rightarrow \infty$, it is expected that an eigenvalue of $\frac{1}{n}\mathbf{XX}^\top$ would also diverge. For instance, say $\lambda_1(\mathbf{C}) = p$ and $\lambda_2(\mathbf{C}) = \dots = \lambda_p(\mathbf{C}) = 1$; then $\mu_{\mathbf{C}} \rightarrow \delta_1$ so that Theorems 2.6 and 2.4 ensure that $\mu_{\frac{1}{n}\mathbf{XX}^\top}$ converges weakly to the Marčenko-Pastur law, while the largest eigenvalue of $\frac{1}{n}\mathbf{XX}^\top$ is strongly expected to diverge to infinity (which it indeed does in this case). Section 2.5 on *spiked models* is strongly inspired by this remark.

2.4 Preliminaries on statistical inference

Section 2.3 provides the necessary ingredients for basic statistical inference considerations of large dimensional sample covariance matrix models.

In this section, we will successively consider the estimation (i) of linear eigenvalue statistics²⁶ of the type $\frac{1}{p} \sum_{i=1}^p f(\lambda_i(\mathbf{C}))$ and (ii) of eigenvector projections $\mathbf{a}^\top \mathbf{u}_i$ (\mathbf{u}_i an eigenvector of \mathbf{C}) for deterministic vectors \mathbf{a} , from the sample observation $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$, $\mathbf{x}_i = \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$ and \mathbf{z}_i with standard i.i.d. entries, as defined in Theorem 2.6.

Before entering the topic, it must be mentioned that *large dimensional statistical inference*, from a random matrix approach, has stood for long as a complex problem. In particular, retrieving information about a population covariance \mathbf{C} from the samples $\mathbf{C}^{\frac{1}{2}} \mathbf{X}$ may be seen as inverting Theorem 2.6, a problem tentatively tackled in [El Karoui, 2008] and later in [Bun et al., 2017] using convex optimization (thus non exact) schemes, but with limited success. Some specific objects, such as traces of powers of \mathbf{C} , traces of its resolvent, quadratic forms, etc., may be estimated by detoured means and formed the extensive database of the more-than-fifty *G-estimators* due to Girko [Girko, 2001] (the phrase ‘G-estimator’ should be understood to stand for ‘generalized estimators’, according to Girko). In this section, we instead concentrate on a contour integral approach to *systematically* estimate a broad class of functionals of \mathbf{C} : the idea, found scattered in the literature, was revived by Mestre in [Mestre, 2008]. The content of

²⁶These are called *linear statistics* although f will of course in general not be linear. What is linear here is the mapping $(f(\lambda_1), \dots, f(\lambda_p)) \mapsto \frac{1}{n} \sum_{i=1}^p f(\lambda_i)$.

this section is not easily found in the literature but is strongly inspired by (a simplified treatment of) [Mestre, 2008].

2.4.1 Linear eigenvalue statistics

2.4.1.1 Relating population and sample Stieltjes transforms

A first observation is that the defining equation for $\tilde{m}(z)$ in Theorem 2.6, i.e.,

$$\tilde{m}(z) = \left(-z + c \int \frac{t\nu(dt)}{1+t\tilde{m}(z)} \right)^{-1}$$

can be equivalently rewritten under the form

$$m_\nu \left(-\frac{1}{\tilde{m}(z)} \right) = -z m(z) \tilde{m}(z) \quad (2.28)$$

where we recall that $m(z) = \frac{1}{c}\tilde{m}(z) + \frac{1-c}{c}\frac{1}{z}$. This simply follows from noticing that

$$\begin{aligned} \int \frac{t\nu(dt)}{1+t\tilde{m}(z)} &= \frac{1}{\tilde{m}(z)} \int \frac{t\tilde{m}(z)\nu(dt)}{1+t\tilde{m}(z)} \\ &= \frac{1}{\tilde{m}(z)} \left(1 - \int \frac{\nu(dt)}{1+t\tilde{m}(z)} \right) \\ &= \frac{1}{\tilde{m}(z)} \left(1 - \frac{1}{\tilde{m}(z)} \int \frac{\nu(dt)}{t - (-1/\tilde{m}(z))} \right) \end{aligned}$$

where, from Definition 3, we recognize $\int \frac{\nu(dt)}{t - (-1/\tilde{m}(z))}$ to be the Stieltjes transform m_ν of ν evaluated at $-1/\tilde{m}(z)$.

Theorem 2.6 thus indirectly establishes a relation between the population statistics \mathbf{C} and the sample covariance matrix $\frac{1}{n}\mathbf{XX}^\top$, through the Stieltjes transforms of their *limiting* measures, and we can already anticipate that $-1/\tilde{m}(z)$ will indeed play the role of a variable change to move from z in $m(z), \tilde{m}(z)$ to z' in $m_\nu(z')$ if $z' = -1/\tilde{m}(z)$.

2.4.1.2 Eigen-inference

Now, observe that, for $f : \mathbb{C} \rightarrow \mathbb{C}$ a function analytic in a neighborhood of the eigenvalues of \mathbf{C} , by Cauchy's integral theorem, the linear statistics

$\frac{1}{p} \sum_{i=1}^p f(\lambda_i(\mathbf{C}))$ of the eigenvalues of \mathbf{C} can be expressed as²⁷

$$\begin{aligned} \frac{1}{p} \sum_{i=1}^p f(\lambda_i(\mathbf{C})) &\simeq \int f(t) \nu(dt) \\ &= \int \left[\frac{1}{2\pi i} \oint_{\Gamma_\nu} \frac{f(z)}{z-t} dz \right] \nu(dt) \\ &= -\frac{1}{2\pi i} \oint_{\Gamma_\nu} f(z) \left[\int \frac{\nu(dt)}{t-z} \right] dz \\ &= -\frac{1}{2\pi i} \oint_{\Gamma_\nu} f(z) m_\nu(z) dz \end{aligned} \quad (2.29)$$

where $\Gamma_\nu \subset \mathbb{C}$ is a (positively oriented) contour encircling the support of ν but no singularity of f . Here the integral exchange comes at no difficulty because Γ_ν is a closed compact contour carefully avoiding the support of ν (so that $t-z$ in the denominator is uniformly far from zero) and $\text{supp}(\nu)$ is bounded. Thus, one can express (smooth) linear statistics of the eigenvalues of \mathbf{C} by means of a complex integral involving the Stieltjes transform $m_\nu(z)$.

As a consequence of (2.28), it is now possible to relate the non-observable $m_\nu(z)$ to $\tilde{m}(z)$, which is the large n, p limit of the observable Stieltjes transform $m_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(z)$. To be able to plug (2.28) in (2.29), one needs to perform the change of variable $z \mapsto -1/\tilde{m}(z)$. This is however only possible if there indeed exists a $\Gamma_\nu \subset \mathbb{C}$ (the contour in (2.29)) such that $\Gamma_\nu = -1/\tilde{m}(\Gamma_\mu)$ for some well defined complex path Γ_μ . The discussions in Section 2.3.1 have clarified the conditions under which such a Γ_ν exists.

But let us assume that Γ_ν is indeed well defined as $\Gamma_\nu = -1/\tilde{m}(\Gamma_\mu)$ for a valid Γ_μ . Then, Equation (2.29) along with (2.28) imply

$$\begin{aligned} \int f(t) \nu(dt) &= -\frac{1}{2\pi i} \oint_{\Gamma_\mu} f\left(-\frac{1}{\tilde{m}(\omega)}\right) m_\nu\left(-\frac{1}{\tilde{m}(\omega)}\right) \frac{\tilde{m}'(\omega)}{\tilde{m}(\omega)^2} d\omega \\ &= \frac{1}{2\pi i} \oint_{\Gamma_\mu} f\left(-\frac{1}{\tilde{m}(\omega)}\right) \omega \frac{m(\omega)\tilde{m}'(\omega)}{\tilde{m}(\omega)} d\omega \end{aligned}$$

where we wrote $z = -1/\tilde{m}(\omega)$. Recalling that $m(\omega) = \frac{1}{c}\tilde{m}(\omega) + (1-c)/(c\omega)$, this further reads

$$\begin{aligned} \int f(t) \nu(dt) &= \frac{1}{2c\pi i} \oint_{\Gamma_\mu} f\left(-\frac{1}{\tilde{m}(\omega)}\right) \frac{(\omega\tilde{m}(\omega) + (1-c))\tilde{m}'(\omega)}{\tilde{m}(\omega)} d\omega \\ &= \frac{1}{2c\pi i} \oint_{\Gamma_\mu} f\left(-\frac{1}{\tilde{m}(\omega)}\right) \omega \tilde{m}'(\omega) d\omega - \frac{1-c}{c} f(0) \mathbf{1}_{\{0 \in \Gamma_\nu^\circ\}} \end{aligned}$$

where Γ_ν° is the inside of Γ_ν , and where for the last equality we used

$$\oint_{\Gamma_\mu} f\left(-\frac{1}{\tilde{m}(\omega)}\right) \frac{\tilde{m}'(\omega)}{\tilde{m}(\omega)} d\omega = - \oint_{\Gamma_\nu} z^{-1} f(z) dz = -f(0) \mathbf{1}_{\{0 \in \Gamma_\nu^\circ\}}$$

²⁷Here again the “ \simeq ” sign can be turned into an equality if one assumes $\nu = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C})}$.

by residue calculus, assuming again that f is analytic on a sufficiently large region (in particular here in zero).

To complete the statistical inference framework, one finally needs to relate the above expression to the observation \mathbf{X} . The idea is to use the fact that $m_{\frac{1}{n}\mathbf{X}^T\mathbf{X}}(\omega) \xrightarrow{a.s.} \tilde{m}(\omega)$. To ensure that $\tilde{m}(\omega)$ can be replaced by $m_{\frac{1}{n}\mathbf{X}^T\mathbf{X}}(\omega)$ in the above expression, one however needs to ensure that dominated convergence on the compact set Γ_μ holds. For this, two ingredients are needed: (i) first guarantee that the convergence $m_{\frac{1}{n}\mathbf{X}^T\mathbf{X}}(\omega) \xrightarrow{a.s.} \tilde{m}(\omega)$ is uniform on Γ_μ , which easily follows from the analytic nature of Stieltjes transforms, and most importantly (ii) prove that $f(-1/m_{\frac{1}{n}\mathbf{X}^T\mathbf{X}}(\omega))\omega m'_{\frac{1}{n}\mathbf{X}^T\mathbf{X}}(\omega)$ is uniformly bounded on Γ_μ . This second item follows from Theorem 2.11 which guarantees that all eigenvalues remain in the vicinity of $\text{supp}(\mu)$ under the additional conditions (i) $\mathbb{E}[|\mathbf{X}_{ij}|^4] < \infty$ and (ii) $\max_i \text{dist}(\lambda_i(\mathbf{C}), \text{supp}(\nu)) \rightarrow 0$.

As a consequence, accounting now for the conditions of validity of the variable change discussed in the previous section, we have the following fundamental statistical inference result, the original ideas of which are due to Mestre.

Theorem 2.12 (Inspired from [Mestre, 2008]). *Under the setting of Theorem 2.6 with $\mathbb{E}[|\mathbf{X}_{ij}|^4] < \infty$ and $\max_{1 \leq i \leq p} \text{dist}(\lambda_i(\mathbf{C}), \text{supp}(\nu)) \rightarrow 0$, let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a complex function analytic on the complement of $\gamma(\mathbb{C} \setminus \text{supp}(\mu))$ in \mathbb{C} with γ defined in (2.27). Then,*

$$\frac{1}{p} \sum_{i=1}^p f(\lambda_i(\mathbf{C})) - \frac{1}{2c\pi i} \oint_{\Gamma_\mu} f\left(\frac{-1}{m_{\frac{1}{n}\mathbf{X}^T\mathbf{X}}(\omega)}\right) \omega m'_{\frac{1}{n}\mathbf{X}^T\mathbf{X}}(\omega) d\omega \xrightarrow{a.s.} 0$$

for some complex positively oriented contour $\Gamma_\mu \subset \mathbb{C}$ surrounding $\text{supp}(\mu) \setminus \{0\}$. In particular, if $c < 1$, the result holds for any f analytic on $\{z \in \mathbb{C}, \Re[z] > 0\}$ with Γ_μ chosen as any such contour within $\{z \in \mathbb{C}, \Re[z] > 0\}$.

From a numerical standpoint, for $c < 1$, Theorem 2.12 is rather simple: it indicates that any complex contour Γ_μ in $\{z \in \mathbb{C}, \Re[z] > 0\}$ guarantees the result. For $c > 1$, the choice of Γ_μ is less trivial. For safety, it is advised to take Γ_μ a contour closely fitting the support of $\mu_{\frac{1}{n}\mathbf{X}^T\mathbf{X}}$, excluding zero (such as a small rectangle). Figure 2.5 and Figure 2.6 visually explain this issue and the technical request regarding the analytic nature of f . From Figure 2.6, since the “tightest-to-the-real-line” (red) contours Γ_ν in the bottom displays must avoid the blue areas (to be well defined images of valid contours Γ_μ from the top displays), the minimal request is for f to be analytic on those blue areas enclosed in the red contour; if not analytic there, the complex integral would have additional residues, thereby altering the result of the theorem.

In practice, the most problematic case occurs when 0 falls within a blue area and one has to deal with functions $f(z)$ involving $\log(z)$, \sqrt{z} , $1/z$, all of which are singular at $z = 0$. A typical way out of this situation would be to add an extra term in the result to compensate for the extra residue; this compensating term would then have to be estimated. This however appears not always to be possible as discussed next.

Remark 2.11 (On the $c > 1$ case). *For $c > 1$ and for f not analytic at zero (for instance $f(z) = \log(z)$, $f(z) = z^{-1}$, or $f(z) = \sqrt{z}$), Theorem 2.12 cannot be applied. That is, for these functions,*

$$\frac{1}{p} \sum_{i=1}^p f(\lambda_i(\mathbf{C}))$$

cannot be consistently estimated directly from the theorem statement. Using the above compensation by the residue at zero workaround, however, it appears that the compensating term is at least as hard to estimate as $\frac{1}{p} \sum_{i=1}^p f(\lambda_i(\mathbf{C}))$ itself. This somehow suggests that, when $p > n$ and thus the sample covariance matrix $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ is of rank $n < p$, one lacks information to estimate some functionals of the p eigenvalues of \mathbf{C} . A similar problem will be discussed in Remark 3.3 on the application to between-covariance matrix distance estimation.

2.4.1.3 Application example: estimating population eigenvalues of large multiplicity

Figure 2.4 present three scenarios where the population measure $\mu_{\mathbf{C}}$ (or equivalently its limit ν) is a discrete sum of three distinct eigenvalues. A natural concern in the large n, p dimension is whether it is possible to estimate these eigenvalues consistently from the sample data \mathbf{X} of size n .

In the top display of Figure 2.4, it a priori appears that averaging the sample eigenvalues of each connected component of μ_p may provide such a consistent estimator. This is however not the case: this estimator is indeed biased. The framework devised in the previous section will provide a consistent estimator: the idea is now to design a contour Γ_ν which would encircle *only one* of the three masses in the spectrum (rather than encircling the whole support of ν); one must then find a corresponding valid contour Γ_μ such that $\Gamma_\nu = -1/\tilde{m}(\Gamma_\mu)$; not surprisingly, this contour Γ_μ will encircle the “hump” in the spectrum of μ associated to the corresponding sought-for eigenvalue of \mathbf{C} (this being a consequence of Section 2.3 and particularly of Figure 2.5). In the middle display, a problem arises for the two eigenvalues (3 and 5) of \mathbf{C} associated to the same connected component of μ : for these, no complex contour Γ_ν exists that would be a proper image $\Gamma_\nu = -1/\tilde{m}(\Gamma_\mu)$ and that would circle around either 3 or 5 alone. We will see that a more involved procedure can nonetheless consistently estimate them both. In the bottom display of Figure 2.4, the difficulty further increases: here again, it remains possible to estimate 1, 3 and 5 but at the cost of a more involved method.

Consider then the following generalized setting of Figure 2.4, where

$$\nu_{\mathbf{C}} = \frac{1}{p} \sum_{i=1}^k p_i \delta_{\ell_i} \rightarrow \sum_{i=1}^k c_i \delta_{\ell_i}$$

for $\ell_1 > \dots > \ell_k > 0$, k fixed with respect to n, p , and $p_i/p \rightarrow c_i > 0$ as $p \rightarrow \infty$ (i.e., each eigenvalue has a large multiplicity).

Fully separable case. We additionally assume for the moment that the sample size $n > p$ of $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ (where $\mathbf{x}_i = \mathbf{C}^{\frac{1}{2}}\mathbf{z}_i$, \mathbf{z}_i having standard i.i.d. entries and bounded fourth order moment) is sufficiently large for the number of connected components in μ to be exactly k , i.e., each eigenvalue of \mathbf{C} is “mapped” to a single connected component of $\text{supp}(\mu)$.

Then, to estimate a given ℓ_a , Theorem 2.12 may be applied to the mere function $f(z) = z$, however for Γ_μ changed into $\Gamma_\mu^{(a)}$ a contour circling around the a -th connected component of $\text{supp}(\mu)$ only (sorted descendingly from ∞ to 0). Adapting Theorem 2.12 according to Theorem 2.11 and our previous line of reasoning, we then have

$$\ell_a - \hat{\ell}_a \xrightarrow{a.s.} 0, \quad \hat{\ell}_a = -\frac{n}{p_a} \frac{1}{2\pi i} \oint_{\Gamma_\mu^{(a)}} \omega \frac{m'_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(\omega)}{m_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(\omega)} d\omega \xrightarrow{a.s.} 0.$$

The estimator $\hat{\ell}_a$ can be numerically evaluated. However, recalling that $m_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(\omega)$ (and its derivative) are rational functions, this integral is prone to estimation by a simple residue calculus. Indeed, first observe that the integrand in the expression of $\hat{\ell}_a$ has two types of poles: (i) the $\lambda_i = \lambda_i(\frac{1}{n}\mathbf{X}^\top \mathbf{X})$ falling inside the surface described by $\Gamma_\mu^{(a)}$, since in the neighborhood of λ_i ,

$$\begin{aligned} -\frac{n}{p_a} \omega \frac{m'_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(\omega)}{m_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(\omega)} &= -\frac{n}{p_a} \omega \frac{\frac{1}{n} \sum_{i=1}^n \frac{1}{(\lambda_i - \omega)^2}}{\frac{1}{n} \sum_{i=1}^n \frac{1}{\lambda_i - \omega}} \\ &\sim_{\omega \sim \lambda_i} -\frac{n}{p_a} \frac{\omega}{\lambda_i - \omega} \end{aligned}$$

and (ii) the zeros of $m_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}$ falling within $\Gamma_\mu^{(a)}$.

For readability in what follows, we sort the eigenvalues of $\frac{1}{n}\mathbf{X}^\top \mathbf{X}$ as $\lambda_1 \geq \dots \geq \lambda_n$ (these are almost surely distinct but for the possible zero eigenvalues). Dealing with the first poles is easy: the λ_i falling within $\Gamma_\mu^{(1)}$ are precisely the p_1 largest, within $\Gamma_\mu^{(2)}$ the next p_2 largest, etc., as per Theorem 2.11. The residue associated to λ_i is then

$$\lim_{\omega \rightarrow \lambda_i} (\omega - \lambda_i) \frac{n}{p_a} \frac{-\omega}{\lambda_i - \omega} = \frac{n}{p_a} \lambda_i.$$

The second set of poles is less immediate to retrieve. An important remark is that the zeros, call them η_j (sorted also as $\eta_1 \geq \eta_2 \geq \dots$), of $m_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(\omega)$ are necessarily real (since the Stieltjes transform has nonzero imaginary part for $\Re[\omega] \neq 0$) and satisfy

$$\frac{1}{n} \sum_{i=1}^n \frac{1}{\lambda_i - \eta_j} = 0.$$

Since the function

$$x \mapsto \frac{1}{n} \sum_{i=1}^n \frac{1}{\lambda_i - x}$$

is increasing and has ∞ and $-\infty$ asymptotes at $x = \lambda_i - 0$ and $x = \lambda_i + 0$, respectively, each η_j falls exactly in one of the intervals $[\lambda_i, \lambda_{i+1}]$ and thus each λ_i pole is accompanied by its η_i pole (if sorted similarly; see Figure 2.7 for an illustration). The residue calculus then gives, by Taylor expanding the denominator,

$$\lim_{\omega \rightarrow \eta_j} (\omega - \eta_j) \frac{n}{p_a} \frac{-\omega m'_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(\omega)}{0 + m'_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(\eta_j)(\omega - \eta_j)} = -\frac{n}{p_a} \eta_j.$$

As a result, we finally have the estimator

$$\hat{\ell}_a = \frac{n}{p_a} \sum_{i=p_1+\dots+p_{a-1}+1}^{p_1+\dots+p_a} \lambda_i - \eta_i.$$

Surprisingly at first, it appears that the estimator is the sum of $p_a = O(p)$ terms, which may seem to conduct to an estimate of order $O(p)$. However, recall that $\lambda_1, \dots, \lambda_p$ are “compacted” in a support of size $O(1)$ and that $\lambda_i < \eta_i < \lambda_{i+1}$ so that $\lambda_i - \eta_i = O(1/p)$, which settles the problem.




Figure 2.7: Illustration of the zeros (η_i) and poles (λ_i) of the (restriction to the real axis of the) Stieltjes transform $m_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}(x)$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

This formulation is nonetheless still not fully closed in the sense that the η_i are so far only provided in terms of the zeros of $m_{\frac{1}{n}\mathbf{X}^\top \mathbf{X}}$. The following remark provides an explicit form.

Remark 2.12 (Explicit expression for the zeros of $m_{\mathbf{X}}(z)$). *For $\mathbf{X} \in \mathbb{R}^{n \times n}$ symmetric with eigenvalues $\lambda_1 > \dots > \lambda_n$, the zeros $\eta_1 > \eta_2 > \dots$ of $m_{\mathbf{X}}(z)$*

satisfy the following equivalence relations

$$\begin{aligned}
\frac{1}{n} \sum_{i=1}^n \frac{1}{\lambda_i - \eta_j} = 0 &\Leftrightarrow \frac{1}{n} \sum_{i=1}^n \frac{-\eta_j}{\lambda_i - \eta_j} = 0 \\
&\Leftrightarrow \frac{1}{n} \sum_{i=1}^n \frac{\lambda_i}{\lambda_i - \eta_j} - 1 = 0 \\
&\Leftrightarrow \frac{1}{n} \sqrt{\boldsymbol{\lambda}}^\top (\boldsymbol{\Lambda} - \eta_j \mathbf{I}_n)^{-1} \sqrt{\boldsymbol{\lambda}} - 1 = 0 \\
&\Leftrightarrow \det \left(\frac{1}{n} \sqrt{\boldsymbol{\lambda}} \sqrt{\boldsymbol{\lambda}}^\top (\boldsymbol{\Lambda} - \eta_j \mathbf{I}_n)^{-1} - \mathbf{I}_n \right) = 0 \\
&\Leftrightarrow \det \left(\frac{1}{n} \sqrt{\boldsymbol{\lambda}} \sqrt{\boldsymbol{\lambda}}^\top - \boldsymbol{\Lambda} + \eta_j \mathbf{I}_n \right) = 0
\end{aligned}$$

where we denoted $\sqrt{\boldsymbol{\lambda}} \in \mathbb{R}^p$ the (column) vector of the $\sqrt{\lambda_i}$'s and $\boldsymbol{\Lambda} \in \mathbb{R}^{p \times p}$ the diagonal matrix $\text{diag}(\lambda_1, \dots, \lambda_p)$, sorted in the same way, and used Lemma 2.3 as well as the fact that $\det(\boldsymbol{\Lambda} - \eta_j \mathbf{I}_n) \neq 0$ according to our discussion above.

Consequently, the zeros of $m_{\mathbf{X}}$ are exactly the eigenvalues of

$$\boldsymbol{\Lambda} - \frac{1}{n} \sqrt{\boldsymbol{\lambda}} \sqrt{\boldsymbol{\lambda}}^\top.$$

Figure 2.8 depicts the estimation error in the setting of two population eigenvalues ℓ_1 and ℓ_2 (with $\ell_1 = 1$ and $p/n = 1/4$), as a function of the difference $\Delta\lambda = \ell_2 - \ell_1$. The error grows rapidly once $\Delta\lambda < 1$: this is a typical “avalanche effect” which appears below the phase transition threshold when the two connected components of the support of the empirical measure μ are no longer separable.

Non-separable case. The estimator introduced above is only valid if the contour $\Gamma_\mu^{(a)}$ is licit, in the sense that its image by the variable change $z \mapsto -1/\tilde{m}(z)$ leads to a valid contour surrounding ℓ_a only. However, we have seen (in Figure 2.6 notably) that there may not exist any such licit $\Gamma_\mu^{(a)}$. In our present setting, Figures 2.5 (both middle and bottom) and Figure 2.6 (both left and right) reveal that, if say ℓ_1 and ℓ_2 are associated to a single connected component of $\text{supp}(\mu)$, then all contours $\Gamma_\mu^{(1)}$ surrounding the p_1 largest λ_i are illicit.

In order to estimate both ℓ_1 and ℓ_2 individually, one must then resort to using at least *two* estimates of linear functionals of the couple (ℓ_1, ℓ_2) . One approach is to estimate simultaneously both $\frac{p_1}{p} \ell_1 + \frac{p_2}{p} \ell_2$ and $\frac{p_1}{p} \ell_1^2 + \frac{p_2}{p} \ell_2^2$, which are accessible from our present adaptation of Theorem 2.12 for $f(z) = z$ and $f(z) = z^2$, with a contour $\Gamma_\mu^{(1,2)}$ surrounding the connected component of μ encompassing the $p_1 + p_2$ largest λ_i .

Assuming p_1 and p_2 are known, this thus boils down to solving a second order polynomial in $\hat{\ell}_1$ and $\hat{\ell}_2$. This procedure however has several limitations:




Figure 2.8: Eigenvalue estimation error as a function of $\Delta\lambda$, for $\ell_1 = 1$, $\ell_2 = 1 + \Delta\lambda$ and $p/n = 1/4$. Results averaged over 10 runs. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

(i) the polynomial equations may lead to nonreal solutions (recall that, while asymptotically this will not occur, the procedure is based on the *finite dimensional* random realization \mathbf{X} , so that nonreal solutions may arise with non-zero probability), and (ii) assuming that p_1 and p_2 are known is quite demanding as they cannot be easily estimated from the λ_i themselves (an additional third equation is then needed here), (iii) a further classical issue in statistics is that estimates of high order moments are increasingly prone to large variances as the moment order increases: as such, the need for additional equations to estimate the individual ℓ_a and their multiplicity must pass through generalized (non polynomial) moments, which are possibly cumbersome to estimate.

2.4.2 Eigenvector projections and subspace methods

In the previous section on the inference methods for the linear statistics (of eigenvalues) of the population covariance \mathbf{C} , we exploited the relation

$$m_\nu(-1/\tilde{m}(z)) = -zm(z)\tilde{m}(z)$$

between the Stieltjes transform m_ν of the population measure ν and the Stieltjes transform m (and \tilde{m}) of the sample measure μ (and $\tilde{\mu} = c\mu + (1-c)\delta_0$), which are immediate consequences of Theorem 2.6.

The deterministic equivalent statements $\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z)$ (as well as $\tilde{\mathbf{Q}}(z) \leftrightarrow \bar{\tilde{\mathbf{Q}}}(z)$) in Theorem 2.6 go beyond Stieltjes transform relations as they connect the whole resolvent $\mathbf{Q}(z) = (\frac{1}{n}\mathbf{XX}^\top - z\mathbf{I}_p)^{-1}$ of the sample covariance (almost directly) to the resolvent $(\mathbf{C} - z\mathbf{I}_p)^{-1}$ of the population covariance.

These relations allow for two main estimates: (i) when \mathbf{C} is known, they provide asymptotic characterizations of some functionals of \mathbf{X} involving its singular vectors (i.e., the eigenvectors $\mathbf{u}_i(\mathbf{X}^\top \mathbf{X})$ of $\mathbf{X}^\top \mathbf{X}$ or $\mathbf{u}_i(\mathbf{X}\mathbf{X}^\top)$ of $\mathbf{X}\mathbf{X}^\top$), in particular projections $\mathbf{u}_i(\mathbf{X}\mathbf{X}^\top)^\top \mathbf{u}(\mathbf{C})$ onto the corresponding eigenvectors $\mathbf{u}(\mathbf{C})$ of \mathbf{C} ; (ii) when \mathbf{C} is unknown, they provide estimates for some functionals of the eigenvectors of \mathbf{C} , notably projections $\mathbf{a}^\top \mathbf{u}(\mathbf{C})$ onto deterministic vectors \mathbf{a} . The latter case is particularly suited to the so-called subspace methods, based on the fact that $\mathbf{u}(\mathbf{C})$ is known to be aligned (or be equal) to some vector \mathbf{a}_θ parametrized by θ and one aims to solve for θ maximizing this alignment. Another scenario is significance is spectral clustering where the dominant eigenvectors of the kernel matrix $\mathbf{X}^\top \mathbf{X}$ are used to estimate the dominant population eigenvectors, themselves precisely providing the data classes: knowing the asymptotic alignment between both in turn provides an estimate on the performance of spectral clustering.

2.4.2.1 Estimates of functionals of \mathbf{X}

In some applications, the observed data \mathbf{X} will be processed in a nonlinear fashion that may nonetheless preserve its eigenvector structure. The spectral behavior of the resulting matrix may here be typically evaluated by means of its projection onto specific vector structures. This is for instance the case of some simple gradient descent mechanisms for supervised learning discussed in Section 5.2, where the learning performance can be measured from the alignment between the gradient descent iterates and the classification vectors (such as the vector $[-\mathbf{1}_{n_1}, \mathbf{1}_{n_2}]$ in a binary classification setting).

For $\mathbf{M} \in \mathbb{R}^{p \times p}$ a symmetric matrix with spectral decomposition $\mathbf{M} = \mathbf{U}\Lambda\mathbf{U}^\top$ and $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_p)$, and $f : \mathbb{R} \rightarrow \mathbb{R}$, we shall here denote

$$f(\mathbf{M}) = \mathbf{U} \text{diag}(f(\lambda_i))_{i=1}^p \mathbf{U}^\top.$$

Assume f is extensible to a complex function $f : \mathbb{C} \rightarrow \mathbb{C}$, analytic on a neighborhood of $\lambda_1, \dots, \lambda_p$. Then, we have that

$$f(\mathbf{M}) = -\frac{1}{2\pi i} \oint_{\Gamma} f(z) \mathbf{Q}_{\mathbf{M}}(z) dz$$

for $\Gamma \subset \mathbb{C}$ a contour closely encompassing $\lambda_1, \dots, \lambda_p$ but no singularity of f . This result arises from a simple residue calculus. Indeed, writing

$$\mathbf{Q}_{\mathbf{M}} = \mathbf{U}(\Lambda - z\mathbf{I}_p)^{-1} \mathbf{U}^\top = \sum_{i=1}^p \frac{\mathbf{u}_i \mathbf{u}_i^\top}{\lambda_i - z}$$

with $\mathbf{U} = [\mathbf{u}_1, \dots, \mathbf{u}_p]$, each λ_j is a pole of the integrand and the associated residue is

$$\lim_{z \rightarrow \lambda_j} (z - \lambda_j) f(z) \sum_{i=1}^p \frac{\mathbf{u}_i \mathbf{u}_i^\top}{\lambda_i - z} = -f(\lambda_j) \mathbf{u}_j \mathbf{u}_j^\top.$$

Summing this result over j gives the result.

Now, assuming $\mathbf{Q}_M(z)$ has a deterministic equivalent $\bar{\mathbf{Q}}(z)$, we have in particular, for $\mathbf{A} \in \mathbb{R}^{p \times p}$ and $\mathbf{a}, \mathbf{b} \in \mathbb{R}^p$, deterministic and of bounded norms,

$$\begin{aligned}\frac{1}{p} \operatorname{tr} \mathbf{A} f(\mathbf{M}) &= -\frac{1}{2\pi i} \oint_{\Gamma} f(z) \frac{1}{p} \operatorname{tr} \mathbf{A} \mathbf{Q}_M(z) dz \\ &= -\frac{1}{2\pi i} \oint_{\Gamma} f(z) \frac{1}{p} \operatorname{tr} \mathbf{A} \bar{\mathbf{Q}}(z) dz + o(1) \\ \mathbf{a}^\top f(\mathbf{M}) \mathbf{b} &= -\frac{1}{2\pi i} \oint_{\Gamma} f(z) \mathbf{a}^\top \mathbf{Q}_M(z) \mathbf{b} dz \\ &= -\frac{1}{2\pi i} \oint_{\Gamma} f(z) \mathbf{a}^\top \bar{\mathbf{Q}}(z) \mathbf{b} dz + o(1)\end{aligned}$$

thereby giving access to the asymptotics of these eigenvector functionals.

Under the notations of Theorem 2.6, for $\mathbf{M} = \frac{1}{n} \mathbf{X} \mathbf{X}^\top$ the sample covariance matrix under study, we have in particular

$$\begin{aligned}\frac{1}{p} \operatorname{tr} \mathbf{A} f\left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top\right) &= \frac{1}{2\pi i} \oint_{\Gamma_\mu} \frac{f(z)}{z} \frac{1}{p} \operatorname{tr} \mathbf{A} (\mathbf{I}_p + \tilde{m}(z) \mathbf{C})^{-1} dz + o(1) \quad (2.30) \\ \mathbf{a}^\top f\left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top\right) \mathbf{b} &= \frac{1}{2\pi i} \oint_{\Gamma_\mu} \frac{f(z)}{z} \mathbf{a}^\top (\mathbf{I}_p + \tilde{m}(z) \mathbf{C})^{-1} \mathbf{b} dz + o(1)\end{aligned}$$

for Γ_μ a contour circling around $\operatorname{supp}(\mu)$.

Example: Eigenspace correlation. Returning to Figure 2.4, we have seen that, when ν is a discrete measure $\nu = \sum_{a=1}^k \frac{p_a}{p} \delta_{\ell_a}$ and c is small enough, μ has a density that spreads in k connected components $\operatorname{supp}(\mu) = \mathcal{S}_1 \cup \dots \cup \mathcal{S}_k$, with \mathcal{S}_a mapped to the atom ℓ_a of ν ; these connected components spread more when c increases. A natural subsequent question would be to know whether the eigenvectors $\hat{\mathbf{u}}_i$ associated to the p_a eigenvalues of $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ of a given connected component \mathcal{S}_a share the same eigenspace as that spanned by the eigenvectors \mathbf{u}_i of \mathbf{C} corresponding to population eigenvalue ℓ_a (with multiplicity p_a) of ν .

This question is readily answered by evaluating

$$\frac{1}{p_a} \operatorname{tr} \Pi_a \hat{\Pi}_a, \quad \Pi_a = \sum_{\lambda_i(\mathbf{C})=\ell_a} \mathbf{u}_i \mathbf{u}_i^\top, \quad \hat{\Pi}_a = \sum_{i \sim \mathcal{S}_a} \hat{\mathbf{u}}_i \hat{\mathbf{u}}_i^\top$$

and where the relation $i \sim \mathcal{S}_a$ stands for $\operatorname{dist}(\lambda_i, \mathcal{S}_a) \rightarrow 0$ (i.e., those eigenvalues of $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ converging to the limiting connected component \mathcal{S}_a).

This quantity can be evaluated by letting $\mathbf{A} = \Pi_a$, $f(z) = 1$ and changing Γ_μ into $\Gamma_{\mathcal{S}_a}$ a contour surrounding only the component \mathcal{S}_a of $\operatorname{supp}(\mu)$ in (2.30).

We precisely get

$$\begin{aligned} \frac{1}{p_a} \operatorname{tr} \boldsymbol{\Pi}_a \hat{\boldsymbol{\Pi}}_a &= \frac{1}{2\pi i} \oint_{\Gamma_{S_a}} \frac{1}{z} \frac{1}{p_a} \operatorname{tr} \boldsymbol{\Pi}_a (\mathbf{I}_p + \tilde{m}(z) \mathbf{C})^{-1} dz + o(1) \\ &= \frac{1}{2\pi i} \oint_{\Gamma_{S_a}} \frac{1}{z} \frac{1}{1 + \tilde{m}(z) \ell_a} dz + o(1) \end{aligned} \quad (2.31)$$

which, for a given population eigenvalue ℓ_a , can be evaluated numerically with the following two-step procedure:

1. with Theorem 2.10, determine the support of μ , which is assumed to have exactly k disjoint components, i.e., $\operatorname{supp}(\mu) = \bigcup_{i=1}^k \mathcal{S}_i$ with say $\mathcal{S}_i = [s_i^-, s_i^+]$ and $s_i^+ < s_{i+1}^-$;
2. choose *any* licit contour Γ_{S_a} that carefully circles around *only* the component \mathcal{S}_a , for instance the rectangular Γ_{S_a} as depicted in Figure 2.6 (which was adopted e.g., in [Bai and Silverstein, 2004]); then evaluated the integral numerically over this contour by solving the fixed-point defining equation of $\tilde{m}(z)$.




Figure 2.9: Typical contour Γ_{S_a} , for $\nu = \frac{1}{3}(\delta_1 + \delta_3 + \delta_7)$ with $c = 1/10$.

But we may go beyond numerical evaluation and obtain an explicit expression of the integral. To this end, for the chosen rectangular contour Γ_{S_a} (Figure 2.9), this consists in evaluating the sum of four line integrals (two “horizontal” and two “vertical”). We provide here the full derivation as it is instrumental of many such calculus arising in similar inference problems and, to the best of our knowledge, this specific calculus was not derived elsewhere in the random matrix literature. Let us first focus on the sum of the two horizontal integrals

$$\int_{s_a^+ + \epsilon_x}^{s_a^- - \epsilon_x} g(x + i\epsilon_y) dx + \int_{s_a^- - \epsilon_x}^{s_a^+ + \epsilon_x} g(x - i\epsilon_y) dx$$

for $g(z) \equiv \frac{1}{z} \frac{1}{1+\tilde{m}(z)\ell_a}$ our object of interest here. Note from the definition of Stieltjes transform (Definition 3) that

$$\Re[m(x+iy)] = \Re[m(x-iy)], \quad \Im[m(x+iy)] = -\Im[m(x-iy)]$$

for any Stieltjes transform $m(z)$ and, consequently,

$$\Re[g(x+iy)] = \Re[g(x-iy)], \quad \Im[g(x+iy)] = -\Im[g(x-iy)].$$

A direct consequence of this observation is that

$$\int_{s_a^+ + \epsilon_x}^{s_a^- - \epsilon_x} g(x + i\epsilon_y) dx + \int_{s_a^- - \epsilon_x}^{s_a^+ + \epsilon_x} g(x - i\epsilon_y) dx = -2i \int_{s_a^- - \epsilon_x}^{s_a^+ + \epsilon_x} \Im[g(x + i\epsilon_y)] dx$$

and thus only the imaginary part of $g(z) = \frac{1}{z} \frac{1}{1+\tilde{m}(z)\ell_a}$ remains, which is explicitly given by

$$\Im[g(x + i\epsilon_y)] = -\frac{\epsilon_y + \ell_a (x\Im[\tilde{m}(x + i\epsilon_y)] + \epsilon_y\Re[\tilde{m}(x + i\epsilon_y)])}{(x^2 + \epsilon_y^2)(1 + 2\ell_a\Re[\tilde{m}(x + i\epsilon_y)] + \ell_a^2|\tilde{m}(x + i\epsilon_y)|^2)}.$$

As for the two vertical integrals (from $-\epsilon_y$ to ϵ_y), we expect that, in the limit $\epsilon_y \rightarrow 0$, they can be neglected. This is indeed the case as we know from Theorem 2.10 that the limit

$$\tilde{m}^\circ(x) = \lim_{\epsilon_y \downarrow 0} \tilde{m}(x + i\epsilon_y) = \lim_{\epsilon_y \downarrow 0} (\tilde{m}(x - i\epsilon_y))^*$$

(with $(\cdot)^*$ the complex conjugate) exists and is real for $x \notin \text{supp}(\mu)$, so that $g(z)$ is continuous on the vertical lines and the vertical integrals thus vanish as $\epsilon_y \rightarrow 0$. The resulting complex integral thus corresponds to the limit of the horizontal integrals for $\epsilon_y \rightarrow 0$. As opposed to the vertical integrals though, for every $x \in \text{supp}(\mu)$, $\tilde{m}^\circ(x)$ is of positive imaginary part, so that the limits of $m(z)$ and thus of $g(z)$, for $z = x \pm i\epsilon_y$, come in conjugate pairs as $\epsilon_y \downarrow 0$. This finally leads to

$$\frac{1}{p_a} \text{tr } \boldsymbol{\Pi}_a \hat{\boldsymbol{\Pi}}_a = \frac{1}{\pi} \int_{s_a^-}^{s_a^+} \frac{\ell_a \Im[\tilde{m}^\circ(x)]}{1 + 2\ell_a \Re[\tilde{m}^\circ(x)] + \ell_a^2 |\tilde{m}^\circ(x)|^2} \frac{dx}{x} + o(1) \quad (2.32)$$

where we recall that, for x inside the support, $\tilde{m}^\circ(x)$ is the unique solution *with positive imaginary part* of

$$\tilde{m}^\circ(x) = \left(-x + c \int \frac{t\nu(dt)}{1 + \tilde{m}^\circ(x)t} \right)^{-1}.$$

We will show in Section 2.5 on ‘‘spiked models’’ that, when the multiplicity p_a of atom ℓ_a is small – technically, if one assumes that $p_a = O(1)$ with respect to p –, the alignment $\text{tr } \boldsymbol{\Pi}_a \hat{\boldsymbol{\Pi}}_a$ just derived assumes a much simpler and fully explicit form (see Theorem 2.14). Yet, the present estimate, which we set under

the opposite scenario where $p_a = O(p)$, turns out (as numerical observations suggest) to be more precise even when p_a is small.

To make this claim more visual, consider the setting where the population covariance $\mathbf{C} \in \mathbb{R}^{p \times p}$ has its $p - m$ eigenvalues equal to 1 and the remaining m eigenvalues equal to $\ell > 1$, so that the population spectral measure ν is a discrete measure having two components: $\nu = \frac{p-m}{p}\delta_1 + \frac{m}{p}\delta_\ell$. In the case where $m, n, p \rightarrow \infty$ with $\lim m/p, p/n \in (0, \infty)$, the correlation of eigenspaces that correspond to the leading eigenvalues of \mathbf{C} (equal to ℓ with multiplicity m) and of $\frac{1}{n}\mathbf{X}\mathbf{X}^\top$, can be fully characterized by (2.32). Figure 2.10 compares the empirical eigenspace correlation with different limiting behavior predicted by the “separate bulk” model in (2.32) versus the spiked model introduced later in Theorem 2.14. For small values of m , both limiting predictions are close, although (2.32) already shows a surprisingly marked advantage over the spiked model, even though $m \ll p$ (which goes against our assumptions). But as m increases, the spiked model-based Theorem 2.14 tends to overestimate the correlation where the prediction (2.32) is a close match to the empirical output.

This observation, which appears to be quite systematic in random matrix theory, is interesting for applicative considerations: in practice, \mathbf{C} is fixed (instead of growing large) and so are m, p and n . Yet, the random matrix predictions based on simultaneously large m, p, n are always extremely accurate and, most importantly, systematically more accurate than when one assumes one of the dimensions (be it m, p , or n) is fixed.




Figure 2.10: Empirical versus limiting eigenspace correlation as a function of $\frac{p}{n} = \frac{p}{m} \frac{m}{n}$ for $\nu = (1 - \frac{m}{p})\delta_1 + \frac{m}{p}\delta_2$, $\frac{m}{p} = \frac{1}{16}$ and $n = 1\,024$. [Link to code]

As a side remark, if we only have access to the empirical covariance $\frac{1}{n}\mathbf{X}^\top\mathbf{X}$ and its Stieltjes transform (i.e., if \mathbf{C} is unknown), then the contour integration

in (2.31) asymptotically and practically reduces to residue calculus as

$$\begin{aligned} \frac{1}{p_a} \operatorname{tr} \boldsymbol{\Pi}_a \hat{\boldsymbol{\Pi}}_a &= \frac{1}{2\pi i} \oint_{\Gamma_{\mathcal{S}_a}} \frac{1}{z} \frac{1}{1 + m_{\frac{1}{n}} \mathbf{X}^\top \mathbf{X}(z) \ell_a} dz + o(1) \\ &= \sum_{i \sim \mathcal{S}_a} \frac{-m_{\frac{1}{n}} \mathbf{X}^\top \mathbf{X}(\zeta_i)}{\zeta_i m'_{\frac{1}{n}} \mathbf{X}^\top \mathbf{X}(\zeta_i)} + o(1) \end{aligned}$$

with ζ_i the roots of $m_{\frac{1}{n}} \mathbf{X}^\top \mathbf{X}(\zeta_i) = -1/\ell_a$, which, by an argument similar to Remark 2.12, are the (sorted) eigenvalues of $\Lambda + \frac{\ell_a}{n} \mathbf{1}_n \mathbf{1}_n^\top$, for Λ the diagonal matrix containing the eigenvalue of $\frac{1}{n} \mathbf{X}^\top \mathbf{X}$. The residue calculus technique performed in the last equation was described in the previous section.

2.4.2.2 Eigenvector inference and subspace methods

The second interest of the deterministic equivalent $\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z)$ of Theorem 2.6, already underlined in the previous example, now concerns the statistical inference of the eigenvectors and eigenspaces of \mathbf{C} . Unless a strong a priori structure is imposed, the eigenvectors themselves cannot be consistently estimated from \mathbf{X} (especially large eigenspaces involving $O(p^2)$ parameters, which cannot be estimated from the $O(pn)$ data observations). But their *scalar* projections onto deterministic vectors are accessible. Precisely, for $\mathbf{a}, \mathbf{b} \in \mathbb{R}^p$ of bounded Euclidean norm, denoting $\boldsymbol{\Pi}_i$ a projector on the eigenspace associated to the eigenvalue $\lambda_i(\mathbf{C})$,

$$\mathbf{a}^\top \boldsymbol{\Pi}_i \mathbf{b} = -\frac{1}{2\pi i} \oint_{\Gamma_\nu^i} \mathbf{a}^\top (\mathbf{C} - z \mathbf{I}_p)^{-1} \mathbf{b} dz$$

for Γ_ν^i a contour circling around $\lambda_i(\mathbf{C})$ only. From Theorem 2.6 and our subsequent discussions of Section 2.3, it is strongly desirable to use again the variable change $z = -1/\tilde{m}(\omega)$ in order to estimate $\mathbf{a}^\top \boldsymbol{\Pi}_i \mathbf{b}$ from an integral over $\mathbf{a}^\top \mathbf{Q}(z) \mathbf{b}$. However, this is again only possible if there exists a pair of contours (Γ_ν^i, Γ) such that $-1/\tilde{m}(\Gamma) = \Gamma_\nu^i$. This is in general not possible unless $\lambda_i(\mathbf{C})$ “induces” its own associated connected component in $\operatorname{supp}(\mu)$. This is precisely the case again of the example of Figure 2.4. Assuming the validity of the variable change, we thus have

$$\begin{aligned} \mathbf{a}^\top \boldsymbol{\Pi}_i \mathbf{b} &= -\frac{1}{2\pi i} \oint_{\Gamma} \mathbf{a}^\top \left(\mathbf{C} + \frac{1}{\tilde{m}(\omega)} \mathbf{I}_p \right)^{-1} \mathbf{b} \frac{\tilde{m}'(\omega)}{\tilde{m}(\omega)^2} d\omega \\ &= \frac{1}{2\pi i} \oint_{\Gamma} \mathbf{a}^\top \mathbf{Q}(\omega) \mathbf{b} \frac{\omega \tilde{m}'(\omega)}{\tilde{m}(\omega)} d\omega + o(1). \end{aligned}$$

This formula reveals handy when testing whether an expected “structure” vector $\mathbf{a} \in \mathbb{R}^p$ is present in the dominant subspace associated to the largest eigenvalue (possibly with multiplicity) $\lambda_1(\mathbf{C})$ of the data covariance structure \mathbf{C} . The value $\mathbf{a}^\top \boldsymbol{\Pi}_1 \mathbf{a} / \|\mathbf{a}\|^2 \in [0, 1]$ precisely evaluates a score for the structure vector \mathbf{a} to be in the span of the dominant eigenvectors of \mathbf{C} .

This analysis finds several applications in detection and estimation, notably in the field of array processing. A concrete example, the G-MUSIC algorithm, is discussed in Section 3.1.3 but more results are available in the dedicated array processing literature [Mestre and Lagunas, 2008, Kammoun et al., 2017].

2.5 Spiked models

The statistical methods discussed in the previous sections for the sample covariance matrix model offer a flexible estimation and inference framework, which can be extended to a large spectrum of random matrix models. However, they have a certain number of practical limitations: (i) they rely on the implicit nature of Theorem 2.6 and thus their behavior is not easily understood, (ii) the complex integration framework, while theoretically satisfying, may be difficult to handle in practice (conditions of existence of valid contours need be ensured, the complex integrals do not necessarily lend themselves to simple analytical evaluation, etc.).

In this section, we will consider a very special, yet practically far reaching, case of sample covariance matrix models for which the limiting spectral measure coincides with the Marčenko-Pastur law, while the population covariance matrix has a non-trivial informative structure. Since the Marčenko-Pastur law assumes an explicit well-understood expression (recall Theorem 2.4), the various estimates of interest will be explicit, and thus intuitions on their behavior are easily derived. Besides, the various change of variable difficulties for contour integral methods met in the previous sections are greatly simplified in this setting.

These special models fundamentally rely on letting the covariance matrix \mathbf{C} be a *low rank* perturbation of the identity matrix \mathbf{I}_p , i.e., $\mathbf{C} = \mathbf{I}_p + \mathbf{P}$ for $\mathbf{P} \in \mathbb{R}^{p \times p}$ with $\text{rank}(\mathbf{P}) = k$ fixed with respect to n, p .

Such statistical models corresponding to a *low rank update* of a classical random matrix model with well-known behavior are generically called *spiked models*.

2.5.1 Isolated eigenvalues

Let us then consider again the model $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ with $\mathbf{x}_i = \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$, $\mathbf{z}_i \in \mathbb{R}^p$ with standard i.i.d. entries and where

$$\mathbf{C} = \mathbf{I}_p + \mathbf{P}, \quad \mathbf{P} = \sum_{i=1}^k \ell_i \mathbf{u}_i \mathbf{u}_i^\top$$

with k and $\ell_1 \geq \dots \geq \ell_k > 0$ fixed with respect to n, p .

According to Theorem 2.6, $\mu_{\frac{1}{n} \mathbf{XX}^\top}$ has a limiting measure μ defined through

the limiting measure ν of $\mu_{\mathbf{C}}$. But clearly $\nu = \delta_1$ here since

$$\mu_{\mathbf{C}} = \frac{p-k}{p} \delta_1 + \frac{1}{p} \sum_{i=1}^k \delta_{1+\ell_i} \rightarrow \delta_1.$$

As a consequence, while \mathbf{C} is not the identity matrix, μ is the Marčenko-Pastur law introduced in Theorem 2.4. But note importantly that the assumptions of the “no eigenvalue outside the support theorem”, Theorem 2.11, do not hold here since $\text{dist}(1 + \ell_1, \text{supp}(\nu)) \not\rightarrow 0$. Therefore, one cannot claim that *all* the eigenvalues of $\frac{1}{n}\mathbf{X}\mathbf{X}^T$ will converge to the support $\text{supp}(\mu)$.

We will precisely show here that, depending on the values of ℓ_i and the ratio $c = \lim p/n$, the i -th largest eigenvalue of $\frac{1}{n}\mathbf{X}\mathbf{X}^T$ may indeed *isolate* from $\text{supp}(\mu)$. As such, since most of the eigenvalues of $\frac{1}{n}\mathbf{X}\mathbf{X}^T$ aggregate, except possibly for a few ones (up to k of them), the latter isolated eigenvalues are seen as isolated “spikes” in the histogram of eigenvalues. See Figure 2.11, commented next, for a visual representation of these “spikes”.

The specific result, here due to Baik (not Bai) and Silverstein, is given in the following theorem.²⁸

Theorem 2.13 (Spiked models, from [Baik and Silverstein, 2006]). *Under the setting of Theorem 2.6 with $\mathbb{E}[\mathbf{Z}_{ij}^4] < \infty$, let $\mathbf{C} = \mathbf{I}_p + \mathbf{P}$ with $\mathbf{P} = \sum_{i=1}^k \ell_i \mathbf{u}_i \mathbf{u}_i^T$ its spectral decomposition, where k and $\ell_1 \geq \dots \geq \ell_k > 0$ are fixed with respect to n, p . Then, denoting $\lambda_1 \geq \dots \geq \lambda_p$ the eigenvalues of $\frac{1}{n}\mathbf{X}\mathbf{X}^T$, as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$,*

$$\lambda_i \xrightarrow{\text{a.s.}} \begin{cases} \rho_i = 1 + \ell_i + c \frac{1+\ell_i}{\ell_i} > (1 + \sqrt{c})^2 & , \ell_i > \sqrt{c} \\ (1 + \sqrt{c})^2 & , \ell_i \leq \sqrt{c}. \end{cases}$$

The theorem thus identifies an abrupt change in the behavior of the i -th dominant eigenvalue λ_i of $\frac{1}{n}\mathbf{X}\mathbf{X}^T$: if $\ell_i \leq \sqrt{c}$, λ_i converges to the right-edge $(1 + \sqrt{c})^2$ of the support of the Marčenko-Pastur law μ and thus does *not* isolate. However, as soon as $\ell_i > \sqrt{c}$, λ_i converges to a limit *beyond* the right-edge of μ and thus *does isolate*. Note in passing that the transition is smooth as the limit of ρ_i as $\ell_i \rightarrow \sqrt{c}$ indeed coincides with $(1 + \sqrt{c})^2$ (so there is no “sudden jump” of the limiting eigenvalue location at the \sqrt{c} transition).

With a physics inspiration, this phenomenon is often referred to as the *phase transition* of the spiked models.

From a statistical viewpoint, the fact that the i -th eigenvalue λ_i of the sample covariance matrix “macroscopically” exceeds or not the other eigenvalues depending on whether $\ell_i > \sqrt{c}$ or $\ell_i \leq \sqrt{c}$ can be interpreted as a test of whether

²⁸These same results were later retrieved using a free probability approach (so formally under slightly different assumptions on \mathbf{Z}) in the work [Benaych-Georges and Nadakuditi, 2011] and were generalized to a larger class of random matrix models. This last article, a richer set of dedicated and better digested techniques (such as those exposed presently), as well as the growing evidence of fundamental applications of the results [Bianchi et al., 2011, Donoho et al., 2018, Candès et al., 2015, Couillet, 2015, Couillet and Hachem, 2013], triggered a renewed wave of interest for spiked models [Loubaton and Vallet, 2011, Capitaine, 2014].

the “signal strength” ℓ_i of the structured data exceeds the minimal *detectability* threshold \sqrt{c} : this can be achieved if the signal strength ℓ_i is strong enough, or alternatively if the number of observed independent data n is large enough (so that $c = \lim p/n$ is small), as common sense would suggest. Indeed, if $\ell_1 < \sqrt{c}$, the eigenvalues of $\frac{1}{n}\mathbf{XX}^\top$ are all asymptotically compacted in the support $[(1 - \sqrt{c})^2, (1 + \sqrt{c})^2]$ and thus it is theoretically (asymptotically) impossible to tell whether $\mathbf{C} = \mathbf{I}_p$ or \mathbf{C} is more structured from the mere observation of the spectral measure of $\frac{1}{n}\mathbf{XX}^\top$. This phase transition effect, for all successive spikes, is well illustrated in Figure 2.11.

Remark 2.13 (The case of negative ℓ_i 's). *In [Baik and Silverstein, 2006], the authors in fact generalize the result of Theorem 2.13 to account for possibly negative ℓ_i 's, i.e., $\ell_i \in (-1, 0)$. In this situation, the following interesting phenomenon occurs: (i) if $c < 1$ and $\ell_i < -\sqrt{c}$, there exists an associated eigenvalue of $\frac{1}{n}\mathbf{XX}^\top$ which converges to $1 + \ell_i + c(1 + \ell_i)/\ell_i \in (0, (1 - \sqrt{c})^2)$ so on the left edge of the limiting support; (ii) if $c < 1$ and $\ell_i \geq -\sqrt{c}$, the associated eigenvalue converges to the left edge $(1 - \sqrt{c})^2$; (iii) if $c > 1$ (so in particular, since $\ell_i > -1$, we cannot have $\ell_i < -\sqrt{c}$), the corresponding eigenvalue tends to 0: so it is never possible to find isolated eigenvalues in the empty space $(0, (1 - \sqrt{c})^2)$ when $c > 1$.*

This negative- ℓ_i setting in effect finds no practical applications that we are aware of, and would additionally be cumbersome to integrate (due to heavier indexing) into a more general statement of Theorem 2.13.

Proof of Theorem 2.13. When it comes to assessing the eigenvalues of a given matrix \mathbf{M} , the first thing that comes to mind is to solve the determinant equation $\det(\mathbf{M} - \lambda\mathbf{I}) = 0$. This approach is not convenient for \mathbf{M} of increasing dimensions and we have seen that the Stieltjes transform method is an appropriate substitute in that case. Here, since the low rank matrix \mathbf{P} only induces a low rank perturbation of $\frac{1}{n}\mathbf{ZZ}^\top$, the use of Sylvester's identity (Lemma 2.3) will turn the large dimensional determinant equation into a small (fixed) dimensional one, and the determinant equation method is now valid. This is the approach we pursue here.

Specifically, let us seek for the presence of an eigenvalue λ of $\frac{1}{n}\mathbf{XX}^\top$ that is asymptotically greater than $(1 + \sqrt{c})^2$. Our approach is to “isolate” the low rank contribution due to \mathbf{P} from the “whitened” sample covariance matrix model $\frac{1}{n}\mathbf{ZZ}^\top$ with identity covariance. To this end, we use the following sequence of equivalences

$$\begin{aligned} 0 &= \det\left(\frac{1}{n}\mathbf{XX}^\top - \lambda\mathbf{I}_p\right) \\ &= \det\left(\frac{1}{n}(\mathbf{I}_p + \mathbf{P})^{\frac{1}{2}}\mathbf{ZZ}^\top(\mathbf{I}_p + \mathbf{P})^{\frac{1}{2}} - \lambda\mathbf{I}_p\right) \\ &= \det(\mathbf{I}_p + \mathbf{P}) \det\left(\frac{1}{n}\mathbf{ZZ}^\top - \lambda(\mathbf{I}_p + \mathbf{P})^{-1}\right) \end{aligned}$$

where we denoted $\mathbf{X} = \mathbf{C}^{\frac{1}{2}}\mathbf{Z}$. Obviously, $\det(\mathbf{I}_p + \mathbf{P}) \neq 0$ so that the first determinant can be discarded. For the second determinant, first recall from the resolvent identity (Lemma 2.1) that

$$(\mathbf{I}_p + \mathbf{P})^{-1} = \mathbf{I}_p - (\mathbf{I}_p + \mathbf{P})^{-1}\mathbf{P}$$

so that we can isolate the (now well-understood) resolvent of the “whitened” model. That is, letting $\mathbf{Q}(\lambda) = (\frac{1}{n}\mathbf{Z}\mathbf{Z}^T - \lambda\mathbf{I}_p)^{-1}$, we write

$$\begin{aligned} 0 &= \det\left(\frac{1}{n}\mathbf{Z}\mathbf{Z}^T - \lambda\mathbf{I}_p + \lambda(\mathbf{I}_p + \mathbf{P})^{-1}\mathbf{P}\right) \\ &= \det\mathbf{Q}^{-1}(\lambda)\det(\mathbf{I}_p + \lambda\mathbf{Q}(\lambda)(\mathbf{I}_p + \mathbf{P})^{-1}\mathbf{P}). \end{aligned}$$

Thanks to Theorem 2.11, inverting the matrix $\frac{1}{n}\mathbf{Z}\mathbf{Z}^T - \lambda\mathbf{I}_p$ is (almost surely) licit for all large n, p as we demanded $\lambda > (1 + \sqrt{c})^2$. Now, denoting $\mathbf{P} = \mathbf{U}\mathbf{L}\mathbf{U}^T$ with $\mathbf{L} = \text{diag}(\ell_1, \dots, \ell_k)$ and $\mathbf{U} = [\mathbf{u}_1, \dots, \mathbf{u}_k] \in \mathbb{R}^{p \times k}$, we further have

$$(\mathbf{I}_p + \mathbf{P})^{-1}\mathbf{P} = (\mathbf{I}_p + \mathbf{U}\mathbf{L}\mathbf{U}^T)^{-1}\mathbf{U}\mathbf{L}\mathbf{U}^T = \mathbf{U}(\mathbf{I}_k + \mathbf{L})^{-1}\mathbf{L}\mathbf{U}^T.$$

Plugging into the above determinant equation, this is

$$\begin{aligned} 0 &= \det\mathbf{Q}^{-1}(\lambda)\det(\mathbf{I}_p + \lambda\mathbf{Q}(\lambda)\mathbf{U}(\mathbf{I}_k + \mathbf{L})^{-1}\mathbf{L}\mathbf{U}^T) \\ &= \det\mathbf{Q}^{-1}(\lambda)\det(\mathbf{I}_k + \lambda\mathbf{U}^T\mathbf{Q}(\lambda)\mathbf{U}(\mathbf{I}_k + \mathbf{L})^{-1}\mathbf{L}) \end{aligned}$$

where in the last equality we applied Sylvester’s identity (Lemma 2.3). Since $\det\mathbf{Q}^{-1}(\lambda) = \det(\frac{1}{n}\mathbf{Z}\mathbf{Z}^T - \lambda\mathbf{I}_p)$ does not vanish for all large n, p at $\lambda > (1 + \sqrt{c})^2$, we finally have for all large n, p ,

$$0 = \det(\mathbf{I}_k + \lambda\mathbf{U}^T\mathbf{Q}(\lambda)\mathbf{U}(\mathbf{I}_k + \mathbf{L})^{-1}\mathbf{L}).$$

From Theorem 2.4 applied entry-wise to the $k \times k$ entries of $\mathbf{U}^T\mathbf{Q}(\lambda)\mathbf{U}$ (this is where it is fundamental that k remain finite), we now know that

$$\mathbf{U}^T\mathbf{Q}(\lambda)\mathbf{U} = m(\lambda)\mathbf{I}_k + o_{\|\cdot\|}(1)$$

almost surely, for $m(z)$ the Stieltjes transform of the Marčenko-Pastur law μ (the term \mathbf{I}_k arises from the fact that $\mathbf{U}^T\mathbf{U} = \mathbf{I}_k$). Consequently, by continuity of the determinant (this is a polynomial of its entries), we have

$$0 = \det(\mathbf{I}_k + \lambda m(\lambda)(\mathbf{I}_k + \mathbf{L})^{-1}\mathbf{L}) + o(1)$$

and thus, if such a λ exists, it must satisfy

$$\lambda m(\lambda) = -\frac{1 + \ell_i}{\ell_i} + o(1)$$

for some $i \in \{1, \dots, k\}$. We thus need to understand when this equation has a solution. To this end, observe that the function $\mathbb{R} \setminus \text{supp}(\mu) \rightarrow \mathbb{R}$, $x \mapsto xm(x) =$

$\int \frac{x}{t-x} \mu(dt)$ is increasing on its domain of definition and that $xm(x) \rightarrow -1$ as $x \rightarrow \infty$. Solving for $m(z)$ in Theorem 2.4, i.e.,

$$\begin{aligned} zcm^2(z) - (1 - c - z)m(z) + 1 &= 0 \\ \Leftrightarrow zm(z) &= \frac{z + czm(z)}{1 - z - czm(z)} \end{aligned} \quad (2.33)$$

we can isolate $zm(z)$ and express it as a function of z alone (alternatively, we could use the explicit solution for $m(z)$ in the proof of the Marčenko-Pastur law, but this is slightly more cumbersome), so to obtain

$$\lim_{x \in \mathbb{R} \downarrow (1+\sqrt{c})^2} xm(x) = -\frac{1 + \sqrt{c}}{\sqrt{c}}.$$

Thus, $xm(x)$ increases from $-\frac{1+\sqrt{c}}{\sqrt{c}}$ to -1 on the set $((1 + \sqrt{c})^2, \infty)$. The equation $\lambda m(\lambda) = -\frac{1+\ell_i}{\ell_i}$ thus only has a solution if and only if

$$-\frac{1 + \ell_i}{\ell_i} > -\frac{1 + \sqrt{c}}{\sqrt{c}}$$

that is, whenever $\ell_i > \sqrt{c}$. Assuming this holds, we may then use again (2.33) (replacing $zm(z)$ by $-(1 + \ell_i)/\ell_i$) to obtain

$$\lambda \rightarrow 1 + \ell_i + c \frac{1 + \ell_i}{\ell_i}$$

which concludes the proof of Theorem 2.13. \square

Figure 2.11 depicts the eigenvalues of $\frac{1}{n}\mathbf{XX}^\top$ versus the Marčenko-Pastur law, in the scenario where $\mathbf{C} = \mathbf{I}_p + \mathbf{P}$ with \mathbf{P} of rank four, for various ratios p/n . As predicted by Theorem 2.13, the number of visible ‘spikes’ outside the limiting support of $\frac{1}{n}\mathbf{XX}^\top$ varies with p/n : as the ratio decreases, less spikes are visible.

2.5.2 Isolated eigenvectors

From a practical standpoint, we have seen that the presence of isolated eigenvalues in the spectrum of the sample covariance $\frac{1}{n}\mathbf{XX}^\top$ reveals the presence of some ‘structure’ in the population covariance \mathbf{C} in the sense that $\mathbf{C} \neq \mathbf{I}_p$. We have however also seen that the converse is not true: assuming a spiked model for \mathbf{C} , the absence of isolated eigenvalue does not imply $\mathbf{C} = \mathbf{I}_p$.

More interestingly, whether this ‘structure’ is detected or not, one may wonder whether it can be estimated at all. More specifically, for $\mathbf{C} = \mathbf{I}_p + \mathbf{P}$ with $\mathbf{P} = \sum_{i=1}^k \ell_i \mathbf{u}_i \mathbf{u}_i^\top$, are the eigenvectors $\hat{\mathbf{u}}_1, \dots, \hat{\mathbf{u}}_k$ of $\frac{1}{n}\mathbf{XX}^\top$ associated to the k largest eigenvalues $\lambda_1 \geq \dots \geq \lambda_k$ good estimators of $\mathbf{u}_1, \dots, \mathbf{u}_k$?

Not surprisingly, the answer is here again twofold: (i) if $\ell_i \leq \sqrt{c}$ then $\hat{\mathbf{u}}_i$ tends to be totally uncorrelated from and thus asymptotically orthogonal to \mathbf{u}_i ,




Figure 2.11: Eigenvalues of $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ (blue crosses) and the Marčenko-Pastur law (red line) for $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{Z}$, $\mathbf{C} = \mathbf{I}_p + \mathbf{P}$ with $\mu_{\mathbf{P}} = \frac{p-4}{p} \delta_0 + \frac{1}{p} (\delta_1 + \delta_2 + \delta_3 + \delta_4)$, for $p = 512$ and different values of n . Link to code: [\[Matlab\]](#) and [\[Python\]](#).

while (ii) if $\ell_i > \sqrt{c}$, $\hat{\mathbf{u}}_i$ is to some extent aligned to \mathbf{u}_i . The following theorem, due to Paul, quantifies this ‘‘to some extent’’.²⁹

Theorem 2.14 (Eigenvector alignment, from [Paul, 2007]). *Under the setting of Theorem 2.13, let $\hat{\mathbf{u}}_1, \dots, \hat{\mathbf{u}}_k$ be the eigenvectors associated with the largest k eigenvalues $\lambda_1 > \dots > \lambda_k$ of $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$. Further assume that $\ell_1 > \dots > \ell_k > 0$ are all distinct. Then, for $\mathbf{a}, \mathbf{b} \in \mathbb{R}^k$ unit norm deterministic vectors*

$$\mathbf{a}^\top \hat{\mathbf{u}}_i \hat{\mathbf{u}}_i^\top \mathbf{b} - \mathbf{a}^\top \mathbf{u}_i \mathbf{u}_i^\top \mathbf{b} \frac{1 - c\ell_i^{-2}}{1 + c\ell_i^{-1}} \mathbf{1}_{\ell_i > \sqrt{c}} \xrightarrow{a.s.} 0. \quad (2.34)$$

²⁹Here again, a large body of literature and modernized tools were set in place to study asymptotic eigenvector behaviors. Some of them are only valid (or only convenient) for rank-one spike models [Paul, 2007, Benaych-Georges and Nadakuditi, 2011, 2012], but the techniques now widely used (such as the contour-integral method presented in this monograph) generally apply to an arbitrary number of spikes [Couillet and Hachem, 2013, Baik et al., 2005].

In particular, with $\mathbf{a} = \mathbf{b} = \mathbf{u}_i$ we obtain

$$|\mathbf{u}_i^\top \hat{\mathbf{u}}_i|^2 \xrightarrow{a.s.} \zeta_i \equiv \frac{1 - c\ell_i^{-2}}{1 + c\ell_i^{-1}} 1_{\ell_i > \sqrt{c}}. \quad (2.35)$$

Proof of Theorem 2.14. We may first write that, for all large n, p almost surely,

$$\mathbf{a}^\top \hat{\mathbf{u}}_i \hat{\mathbf{u}}_i^\top \mathbf{b} = -\frac{1}{2\pi i} \oint_{\Gamma_{\rho_i}} \mathbf{a}^\top \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top - z \mathbf{I}_p \right)^{-1} \mathbf{b} dz$$

for Γ_{ρ_i} a small contour enclosing the almost sure limit $\rho_i = 1 + \ell_i + c\frac{1+\ell_a}{\ell_i}$ of the eigenvalue λ_i of $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ given in Theorem 2.13 only. Isolating $\frac{1}{n} \mathbf{Z} \mathbf{Z}^\top$ from $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ as in the proof of Theorem 2.13, we have from Woodbury's identity (Lemma 2.7) that

$$\begin{aligned} & \mathbf{a}^\top \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top - z \mathbf{I}_p \right)^{-1} \mathbf{b} \\ &= \mathbf{a}^\top \left(\frac{1}{n} (\mathbf{I}_p + \mathbf{P})^{\frac{1}{2}} \mathbf{Z} \mathbf{Z}^\top (\mathbf{I}_p + \mathbf{P})^{\frac{1}{2}} - z \mathbf{I}_p \right)^{-1} \mathbf{b} \\ &= \mathbf{a}^\top (\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} \left(\frac{1}{n} \mathbf{Z} \mathbf{Z}^\top - z \mathbf{I}_p - z \mathbf{P} \right)^{-1} (\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} \mathbf{b} \\ &= \mathbf{a}^\top (\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} \mathbf{Q}(z) (\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} \mathbf{b} \\ &+ z \mathbf{a}^\top (\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} \mathbf{Q}(z) \mathbf{U} \mathbf{L} (\mathbf{I}_k - z \mathbf{U}^\top \mathbf{Q}(z) \mathbf{U} \mathbf{L})^{-1} \mathbf{U}^\top \mathbf{Q}(z) (\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} \mathbf{b} \\ &= \mathbf{a}^\top (\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} \mathbf{Q}(z) (\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} \mathbf{b} \\ &+ zm^2(z) \mathbf{a}^\top \mathbf{U} (\mathbf{I}_k + \mathbf{L})^{-\frac{1}{2}} \mathbf{L} (\mathbf{I}_k - zm(z) \mathbf{L})^{-1} (\mathbf{I}_k + \mathbf{L})^{-\frac{1}{2}} \mathbf{U}^\top \mathbf{b} + o(1) \end{aligned}$$

where $\mathbf{Q}(z) = (\frac{1}{n} \mathbf{Z} \mathbf{Z}^\top - z \mathbf{I}_p)^{-1}$ and for the last equality we used $(\mathbf{I}_p + \mathbf{P})^{-\frac{1}{2}} = \mathbf{U}(\mathbf{I}_k + \mathbf{L})^{-\frac{1}{2}} \mathbf{U}^\top$ and $\mathbf{U}^\top \mathbf{Q}(z) \mathbf{U} = m(z) \mathbf{I}_k + o_{\parallel \cdot \parallel}(1)$, as per Theorem 2.4. The complex integration of $\mathbf{Q}(z)$ on the contour Γ_{ρ_i} only brings a non-trivial residue for the second right-hand side term owing to the inverse $(\mathbf{I}_k - zm(z) \mathbf{L})^{-1}$ which is singular at $z = \rho_i$. We thus finally have

$$\begin{aligned} \mathbf{a}^\top \hat{\mathbf{u}}_i \hat{\mathbf{u}}_i^\top \mathbf{b} &= -\frac{1}{2\pi i} \oint_{\Gamma_{\rho_i}} zm^2(z) \mathbf{a}^\top \mathbf{U} (\mathbf{I}_k + \mathbf{L})^{-\frac{1}{2}} \mathbf{L} (\mathbf{I}_k - zm(z) \mathbf{L})^{-1} \\ &\quad \times (\mathbf{I}_k + \mathbf{L})^{-\frac{1}{2}} \mathbf{U}^\top \mathbf{b} dz + o(1). \end{aligned}$$

This expression is then evaluated by residue calculus at $z = \rho_i$ (the only singularity of the integrand). The corresponding residue is simple and thus equals

$$\lim_{z \rightarrow \rho_i} (z - \rho_i) (\mathbf{I}_k - zm(z) \mathbf{L})^{-1} = -(\rho_i m'(\rho_i) + m(\rho_i))^{-1} \ell_i^{-1} \mathbf{e}_i \mathbf{e}_i^\top$$

with $\mathbf{e}_i \in \mathbb{R}^k$ the canonical basis vector defined as $[\mathbf{e}_i]_j = \delta_{ij}$. Using the (here most convenient) form

$$m(z) = \frac{1}{-z + \frac{1}{1 + cm(z)}}$$

of the Stieltjes transform of the Marčenko-Pastur law gives³⁰

$$m'(z) = \frac{m(z)^2}{1 - \frac{cm(z)^2}{(1+cm(z))^2}}$$

from which we obtain in particular that $m(\rho_i) = -1/(\ell_i + c)$ and $m'(\rho_i) = \ell_i^2(\ell_i + c)^{-2}(\ell_i^2 - c)^{-1}$. It unfolds that the residue associated to $(\mathbf{I}_k - zm(z)\mathbf{L})^{-1}$ reads

$$(-\rho_i m'(\rho_i) - m(\rho_i))^{-1} \ell_i^{-1} \mathbf{e}_i \mathbf{e}_i^\top = -(\ell_i^2 - c) \ell_i^{-1} \mathbf{e}_i \mathbf{e}_i^\top$$

and we finally get

$$\mathbf{a}^\top \hat{\mathbf{u}}_i \hat{\mathbf{u}}_i^\top \mathbf{b} = \mathbf{a}^\top \mathbf{u}_i \mathbf{u}_i^\top \mathbf{b} \frac{\rho_i m^2(\rho_i) \ell_i}{1 + \ell_i} (\ell_i^2 - c) \ell_i^{-1} + o(1) = \mathbf{a}^\top \mathbf{u}_i \mathbf{u}_i^\top \mathbf{b} \frac{1 - c \ell_i^{-2}}{1 + c \ell_i^{-1}} + o(1)$$

which concludes the proof of Theorem 2.14. \square




Figure 2.12: Empirical versus limiting $|\hat{\mathbf{u}}_1^\top \mathbf{u}_1|^2$ for $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{Z}$, $\mathbf{C} = \mathbf{I}_p + \ell_1 \mathbf{u}_1 \mathbf{u}_1^\top$, $p/n = 1/3$, for varying ℓ_1 . Results obtained by averaging over 100 runs. In black dashed line the local behavior around \sqrt{c} . Link to code: [\[Matlab\]](#) and [\[Python\]](#).

Figure 2.12 compares, in a single-spike scenario, the theoretical limit of $|\hat{\mathbf{u}}_1^\top \mathbf{u}_1|^2$ (in red) versus its empirical value for different ℓ_1 and different p, n with

³⁰In passing, note that $m'(z)$ assumes an *explicit form* as a function of z and $m(z)$. While not surprising in the Marčenko-Pastur case, this turns out to be also true of more elaborate models where $m(z)$ does not have an explicit expression.

constant ratio p/n . It is important to note that the theoretical *asymptotic* phase transition phenomenon at $\ell_1 = \sqrt{c}$ corresponds to a sharp non-differentiable change in the function $\ell_1 \mapsto \zeta_1 = (1 - c\ell_1^{-2})1 + c\ell_1^{-1}1_{\ell_1 \geq \sqrt{c}}$; a local analysis in the limit of $\ell_1 = \sqrt{c} + \epsilon$ reveals that ζ_1 (and thus in the limit $|\hat{\mathbf{u}}_1^\top \mathbf{u}_1|^2$) is locally equal to $\zeta_1 \simeq \frac{2\epsilon}{\sqrt{c}(1+\sqrt{c})}$ and thus, for sufficiently large n, p ,

$$|\hat{\mathbf{u}}_1^\top \mathbf{u}_1| =_{\ell_1=\sqrt{c}+\epsilon} \sqrt{\frac{2}{\sqrt{c}(1+\sqrt{c})}}\sqrt{\epsilon} + O(\epsilon)$$

thus having an infinite derivative in the limit $\ell_1 \downarrow \sqrt{c}$. On real data of finite size, this sharp transition is only observed for extremely large values of n, p . This in particular means that, in practice, residual information is present below the phase transition.

2.5.3 Limiting fluctuations

Theorem 2.13 on the *limiting* presence and position of isolated eigenvalues in the spectrum of $\frac{1}{n}\mathbf{XX}^\top$ establishes that it suffices to evaluate whether the largest eigenvalue λ_1 of $\frac{1}{n}\mathbf{XX}^\top$ “isolates from the other eigenvalues $\lambda_2 > \dots > \lambda_p$ ” to determine the presence of a structure in the population covariance \mathbf{C} .

However, in practice, from the empirical observation of $\lambda_1, \dots, \lambda_p$, how can one decide whether λ_1 is isolated? On a random realization of \mathbf{X} , λ_1 may haphazardly be found “rather far” from λ_2 by a mere finite-dimensional probability effect. The natural question is then to determine whether the rate of occurrence of such “haphazard” events can be evaluated.

A whole line of works, based on significantly different tools from the Stieltjes transform approach adopted in this monograph,³¹ settles this question by evaluating, for $\mathbf{C} = \mathbf{I}_p$ or $\mathbf{C} = \mathbf{I}_p + \mathbf{P}$ with the eigenvalues of \mathbf{P} below the phase transition, the asymptotic probability for λ_1 to escape its limiting value $(1 + \sqrt{c})^2$.

The main result of importance is the following.

Theorem 2.15 (Fluctuation of the largest eigenvalue, from [Baik et al., 2005]).
Under the setting of Theorem 2.13, assume $0 \leq \ell_k < \dots < \ell_1 < \sqrt{c}$. Then,

$$n^{\frac{2}{3}} \frac{\lambda_1 - (1 + \sqrt{c})^2}{(1 + \sqrt{c})^{\frac{4}{3}} c^{-\frac{1}{6}}} \rightarrow \text{TW}_1$$

³¹Unlike the Stieltjes transform method, these tools start from the explicit (finite-dimensional) formula of the joint eigenvalue distribution of matrix models \mathbf{X} or \mathbf{XX}^\top , which is known in the Gaussian case (and only in this case) and given by (2.39). Exploiting the theory of orthogonal polynomials and determinantal processes, (2.39) can be marginalized so to retrieve the exact (finite-dimensional) law of one or several specific eigenvalues (inside the bulk or on the edge). Taking the large dimensional limit relates the law of the eigenvalues to the determinant of a specific kernel [Soshnikov, 2000] (Airy kernel for the edge eigenvalues [Johnstone, 2001, Soshnikov, 1999], sine kernel in the bulk [Arous and Péché, 2005, Erdős et al., 2010]). Details on these techniques can be found in [Anderson et al., 2010].

in law, where TW_1 is the (real) Tracy-Widom distribution, historically defined in [Tracy and Widom, 1996].³²

Specifically, the theorem is here placed under the setting where the “population spikes” are *below* the phase transition and thus asymptotically *not isolated* (as per Theorem 2.13). In this setting, the largest eigenvalue λ_1 thus converges to the right edge $(1 + \sqrt{c})^2$ of the Marčenko-Pastur law and more precisely behaves, according to the theorem, as $\lambda_1 = (1 + \sqrt{c})^2 + n^{-\frac{2}{3}}T$ where T is a (scaled) Tracy-Widom random variable.

This result is of practical interest as it allows one to estimate, for sufficiently large n, p , the probability for λ_1 to be found away from its theoretical limit $(1 + \sqrt{c})^2$ below the phase transition.

Precisely, the result shows that the limiting fluctuations of λ_1 are not Gaussian but follow the Tracy-Widom distribution and that, possibly surprisingly, the rate of this fluctuation is of order $O(n^{-\frac{2}{3}})$ (instead of $O(n^{-\frac{1}{2}})$ or $O(n^{-1})$ as one would usually expect). This rate is strongly related to the following result, initially observed by Silverstein and Choi in [Silverstein and Choi, 1995]: close to the right-edge of its support, the Marčenko-Pastur law behaves proportionally to $\sqrt{(1 + \sqrt{c})^2 - x}$. As such, the typical number of eigenvalues in a space of size ϵ in the neighborhood of the edge is

$$\int_{(1 + \sqrt{c})^2 - \epsilon}^{(1 + \sqrt{c})^2} \sqrt{(1 + \sqrt{c})^2 - x} dx \propto \epsilon^{\frac{3}{2}}.$$

This explains the typical $O(n^{-\frac{2}{3}})$ fluctuation of the eigenvalues in this neighborhood.

The original result from [Baik et al., 2005] also provides the limiting fluctuations of $\lambda_1, \dots, \lambda_k$ *beyond* the phase transition (i.e., when $\ell_i > \sqrt{c}$). Interestingly, after the transition, the fluctuation of λ_1 is now a classical central limit with speed $O(n^{-\frac{1}{2}})$. The surprising “transition” from $O(n^{-\frac{2}{3}})$ to $O(n^{-\frac{1}{2}})$ of the fluctuations of λ_1 (which has little meaning or interpretability for finite n, p) is often called the *BBP phase transition* after the names of Baik et al. [2005]. Couillet and Hachem [2013] go beyond by providing the joint fluctuations of the eigenvalues and eigenvector projections as follows.

Theorem 2.16 (Joint fluctuations beyond the phase transition (from [Couillet and Hachem, 2013])). *Under the setting and notations of Theorem 2.13 and Theorem 2.14, assume $\ell_1 > \dots > \ell_k > \sqrt{c}$ and define $L = \sqrt{p}[\lambda_1 - \rho_1, \dots, \lambda_k - \rho_k]^T$ and $V = \sqrt{p}[|\mathbf{u}_1^T \hat{\mathbf{u}}_1|^2 - \zeta_1, \dots, |\mathbf{u}_k^T \hat{\mathbf{u}}_k|^2 - \zeta_k]^T$. Then, as $p, n \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$,*

$$\begin{pmatrix} L \\ V \end{pmatrix} \rightarrow \mathcal{N} \left(\mathbf{0}_{2k}, \text{BlockDiag} \left(\begin{bmatrix} \frac{c^2(1+\ell_i)^2 \left(1 + c \frac{(1+\ell_i)^2}{(\ell_i+c)^2} \right)}{(c+\ell_i)^2(\ell_i^2-c)} & \frac{(1+\ell_i)^3 c^2}{(\ell_i+c)^2 \ell_i} \\ \frac{(1+\ell_i)^3 c^2}{(\ell_i+c)^2 \ell_i} & \frac{c(1+\ell_i)^2 (\ell_i^2-c)}{\ell_i^2} \end{bmatrix} \right)_{i=1}^k \right)$$

³²The Tracy-Widom distribution does not have an explicit form. Several works have provided approximated forms as well as tables of TW_1 [Chiani, 2014, Ma, 2012].

where $\text{BlockDiag}(\cdot)$ is the “block-diagonal” operator.

The theorem notably states that, in the large n, p limit, while each eigenvalue-eigenvector projector pair fluctuates together, the k pairs fluctuate independently (which would no longer be the case if some ℓ_i ’s had multiplicity greater than one; the request $\ell_1 > \dots > \ell_k > \sqrt{c}$ in the theorem statement avoids this difficulty, which is treated in [Bai and Yao, 2008, Couillet and Hachem, 2013] but gives rise to a more complex result).

Remark 2.14 (Tracy-Widom law: beyond the real field and universality). *The Tracy-Widom law was first introduced in the context of Wigner random matrices (Theorem 2.5). More precisely, Tracy and Widom [1996] showed that the fluctuation of the largest eigenvalue of a real Gaussian Wigner random matrix ($\frac{1}{\sqrt{n}}\mathbf{X}$ with $\mathbf{X} \in \mathbb{R}^{n \times n}$ of i.i.d. zero mean and unit variance Gaussian entries) asymptotically follows a Tracy-Widom distribution*

$$n^{\frac{2}{3}}(\lambda_1 - 2) \rightarrow \text{TW}_1.$$

The Tracy-Widom law also extends beyond the largest eigenvalue: it holds true for the finitely many largest as well as smallest eigenvalues of the Wigner and the Wishart matrix (in the latter case only if $p/n \rightarrow c < 1$). It also goes beyond real-valued symmetric Gaussian matrices (often referred to as the Gaussian orthogonal ensemble (GOE)) and the real-valued Wishart random matrices, to complex (Gaussian Unitary Ensemble (GUE)) and quaternionic (Gaussian Symplectic Ensemble (GSE)) Gaussian matrices: in these scenarios, the limiting laws are respectively the TW_2 and TW_4 Tracy-Widom distributions [Tracy and Widom, 2000]. See Figure 2.13 for an illustration.

The Tracy-Widom law has also been proven, to some extent, to be universal with respect to the Gaussian distribution. In [Soshnikov, 1999] and [Erdős, 2011], the authors prove that, for fast decaying distributions, it is sufficient to match the first two moments of the entries of \mathbf{X} to obtain asymptotic Tracy-Widom fluctuations.

Finally, while the fluctuations of the (finitely many) largest or smallest eigenvalues of $\frac{1}{n}\mathbf{XX}^\top$ are not independent (they give rise, both for the k largest or for the k smallest, to joint fluctuations), Bianchi et al. [2010] showed that the joint fluctuations of the one largest and one smallest eigenvalues of $\frac{1}{n}\mathbf{XX}^\top$ are independent. This last result has the interesting consequence that the fluctuations of the condition number of $\frac{1}{n}\mathbf{XX}^\top$ (defined as the ratio between largest and smallest eigenvalues) around $(1 + \sqrt{c})^2 / (1 - \sqrt{c})^2$ are easily obtained (using for instance the so-called delta method [Vaart, 2000] on the ratio $\lambda_1/\lambda_p = f((\lambda_1 - (1 + \sqrt{c})^2, \lambda_p - (1 - \sqrt{c})^2))$ for some rather elementary function f).

2.5.4 Further discussions and other spiked models

The “spiked model” terminology goes beyond sample covariance matrix models with $\mathbf{C} = \mathbf{I}_p + \mathbf{P}$, for \mathbf{P} a low rank matrix. In the literature, spiked models loosely refer to as “low rank perturbation” models in the following sense: there exists an




Figure 2.13: **(Top)** Empirical histogram of $n^{\frac{2}{3}} \frac{\lambda_1 - (1 + \sqrt{c})^2}{(1 + \sqrt{c})^{\frac{4}{3}} c^{-\frac{1}{6}}}$ for $p = 256$, $n = 512$ versus the real Tracy-Widom law TW_1 . Histogram obtained over 5000 independent runs. **(Bottom)** Tracy-Widom distribution TW_β for $\beta = 1$ (real \mathbf{X}_{ij}), 2 (complex \mathbf{X}_{ij}), and 4 (symplectic \mathbf{X}_{ij}). Link to code: [\[Matlab\]](#) and [\[Python\]](#).

underlying random matrix model \mathbf{X} , the spectral measure of which converges to a well-defined measure with compact support *and* having eigenvalues converging to the support (i.e., no single eigenvalue isolates) which is then modified in some way by a low rank perturbation matrix \mathbf{P} ; the resulting matrix has the same limiting spectral measure as that of \mathbf{X} but with possibly some spurious (isolated)

eigenvalues.

Baik and Silverstein [2006] were the first to study spiked models, but their approach relied on applying the results on sample covariance matrix models (i.e., Theorem 2.6) to the specific case where $\mathbf{C} = \mathbf{I}_p + \mathbf{P}$. This approach requires to have a full understanding of a “more complex” statistical model before particularizing it to a low rank perturbation. Pursuing on the footnote introducing Theorem 2.13, more modern tools launched a second wave of advances in spiked models, mostly triggered by the ideas found in [Benaych-Georges and Nadakuditi, 2012] (with a free probability approach), which is based on relating the perturbation matrix model to the underlying simple (non perturbed) matrix; this is mathematically simpler and opened the path to a broader scope of generalizations to more advanced random matrix models.

Among the popular spiked models, we have the following cases:

- the *information-plus-noise* model of the type

$$\frac{1}{n}(\mathbf{X} + \mathbf{P})(\mathbf{X} + \mathbf{P})^\top$$

with $\mathbf{X} \in \mathbb{R}^{p \times n}$ having i.i.d. standard entries (zero mean, unit variance and finite fourth moment) and $\mathbf{P} \in \mathbb{R}^{p \times n}$ deterministic (or at least independent of \mathbf{X}) of fixed rank k ;

- the *additive* model of the type

$$\mathbf{M} + \mathbf{P}$$

where $\mathbf{M} \in \mathbb{R}^{p \times p}$ is either of the type $\mathbf{M} = \frac{1}{n}\mathbf{XX}^\top$, $\mathbf{X} \in \mathbb{R}^{p \times n}$ with standard i.i.d. entries, or of $\mathbf{M} = \frac{1}{\sqrt{n}}\mathbf{X}$ with \mathbf{X} symmetric with standard i.i.d. entries above and on the diagonal and $\mathbf{P} \in \mathbb{R}^{n \times n}$ deterministic of low rank.

Each of these models has its own phase transition threshold (i.e., the value that eigenvalues of \mathbf{P} must exceed for a spike to be observed), dominant eigenvalue limits, and eigenvector projections. These can all be determined with the aforementioned proof approaches (see examples in the exercise list, Section 2.9).

However, we will see in several applications in Chapter 4 that, in practice, we will be confronted with more general forms of low rank perturbation models that do not fit this conventional “random matrix \mathbf{X} and deterministic perturbation \mathbf{P} ” assumption.

In particular, \mathbf{P} will often be a (possibly elaborate) function of \mathbf{X} . Also, \mathbf{X} itself, which will often stand for the “noisy” part of the data model (while \mathbf{P} will in general comprise both the relevant information and possibly some extra noise), may induce its own isolated eigenvalues. For instance, we shall see that, depending on the ratios p/n and $\text{tr } \mathbf{C}^4 / (\text{tr } \mathbf{C}^2)^2$, the random matrix $\{[\mathbf{XX}^\top]_{ij}^2 1_{i \neq j}\}_{i,j=1}^p$ where $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{Z}$ and \mathbf{Z} with i.i.d. standard entries, may have two isolated eigenvalues although all the eigenvalues of \mathbf{C} remain in their limiting

support. Also, in the context of robust estimation of covariance matrices, it will not be natural for the statistical model to impose that all its population eigenvalues converge to their limiting support (in particular to mimic the action of a few outliers).

Yet, despite these technical differences, the proof approaches of Theorem 2.13 and Theorem 2.14 remain essentially valid. We thus propose here to generalize the notion of “spiked models” to models of the type $\mathbf{X} + \mathbf{P}$ where \mathbf{X} is some reference, well understood, random matrix model (possibly inducing its own spikes) and \mathbf{P} is a low rank matrix, possibly depending on \mathbf{X} .

With this definition, the aforementioned *sample covariance, information-plus-noise* and *additive* models are in fact all equivalent to an additive model. Precisely, we may write

$$\frac{1}{n}(\mathbf{X} + \mathbf{P})(\mathbf{X} + \mathbf{P})^\top = \mathbf{M} + \mathbf{P}'$$

with $\mathbf{M} = \frac{1}{n}\mathbf{XX}^\top$, $\mathbf{P}' = \frac{1}{n}(\mathbf{XP}^\top + \mathbf{PX}^\top + \mathbf{PP}^\top)$

and

$$\frac{1}{n}(\mathbf{I}_p + \mathbf{P})^{\frac{1}{2}}\mathbf{XX}^\top(\mathbf{I}_p + \mathbf{P})^{\frac{1}{2}} = \mathbf{M} + \mathbf{P}'$$

with $\mathbf{M} = \frac{1}{n}\mathbf{XX}^\top$, $\mathbf{P}' = \frac{1}{n}(\mathbf{XP}''^\top + \mathbf{P}''\mathbf{X} + \mathbf{P}''\mathbf{P}''^\top)$

where we introduced $\mathbf{P}'' = \mathbf{U}((\mathbf{I}_k + \mathbf{L})^{\frac{1}{2}} - \mathbf{I}_k)\mathbf{U}^\top$ with $\mathbf{P} = \mathbf{ULU}^\top$. In the remainder of the monograph, we shall systematically exploit this generic modeling approach to treat all spiked models.

2.6 Information-plus-noise, deformed Wigner, and other models

2.6.1 Why focus on the sample covariance matrix model?

The previous sections have mostly been concerned with the sample covariance matrix (as well as more marginally with Wigner matrices), as an instrumental statistical model for the introduction of the main technical tools of interest to the monograph: the Stieltjes transform method, the spiked model approach, and statistical inference based on contour integrals.

Several other classical random matrix models, of interest in statistics, will be listed in this section. The technical methods required to study these models are however not very different and thus not worth detailing. Only pointers to relevant references will be provided here for the interested reader.

It is in particular important to stress that many statistical models arising in machine learning applications are so specific that they may not (strictly) fall in any of the conventional models discussed above. Yet, up to some additional fine-tuning and tricks, the analytical tools required to study these models are

in general not much different from those presented in this chapter. Among examples met in the next chapters of this monograph, we may list:

- Graph Laplacian matrices

$$\mathbf{D} - \mathbf{A}, \quad \mathbf{D}^{-\frac{1}{2}} \mathbf{A} \mathbf{D}^{-\frac{1}{2}}, \quad \mathbf{D}^{-1} \mathbf{A}$$

for $\mathbf{A} \in \mathbb{R}^{n \times n}$ a symmetric matrix with independent entries (up to symmetry) and $\mathbf{D} = \text{diag}(\mathbf{A}\mathbf{1}_n)$. The dependence between \mathbf{A} and \mathbf{D} makes these random matrices slightly different from *deformed Wigner matrices* (see below) of the type $\mathbf{A} + \mathbf{D}$ where \mathbf{A} has independent entries and \mathbf{D} is *deterministic*.

- Kernel random matrices of the inner-product or distance type

$$\mathbf{K} = \{f(\mathbf{x}_i^\top \mathbf{x}_j)\}_{i,j=1}^n, \quad \mathbf{K} = \{f(\|\mathbf{x}_i - \mathbf{x}_j\|^2)\}_{i,j=1}^n.$$

There, the non-trivial dependence between the entries of \mathbf{K} differs significantly from sample covariance models (except for the linear kernel function $f(t) = t$ in the inner-product case).

- Robust estimators of scatter $\hat{\mathbf{C}}$ defined by the solutions to

$$\hat{\mathbf{C}} = \frac{1}{n} \sum_{i=1}^n u\left(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i\right) \mathbf{x}_i \mathbf{x}_i^\top$$

for some non-increasing function $u(t)$. There, due to the implicit nature of $\hat{\mathbf{C}}$, sample covariance matrix results cannot be strictly applied.

- F-matrix models $\hat{\mathbf{C}}_1^{-1} \hat{\mathbf{C}}_2$ and product models $\hat{\mathbf{C}}_1 \hat{\mathbf{C}}_2$ for $\hat{\mathbf{C}}_a = \frac{1}{n} \mathbf{X}_a \mathbf{X}_a^\top$, $a \in \{1, 2\}$, with $\mathbf{X}_1, \mathbf{X}_2$ independent (notably Gaussian) random matrices, used in whitening methods, or in covariance matrix distance evaluation (e.g., Fisher distance, KL divergence, Wasserstein distance, etc.). By successive conditioning, these models are more directly related to the sample covariance matrix models, although not strictly equivalent.

The models and applications listed above appear to be strongly related, one way or another, to sample covariance matrices. Of the examples above, kernel matrices, robust estimators, F-matrices and sample covariance products all relate to sample covariance matrices. The graph Laplacian (as well, to some extent, as the kernel random matrix models) is rather connected to Wigner matrices. This justifies the particularly focused vision of this chapter.³³

³³In effect and to the best of our knowledge, most, if not all, random matrix results directly related to machine learning applications boil down, in simple data model settings at least, to combinations (usually sums, sometimes products) of asymptotically independent matrices of the Wishart (Marčenko-Pastur related) and Wigner (semicircle related) types.

2.6.2 Other models

2.6.2.1 Advanced sample covariance matrices

From a historical standpoint, the model studied by [Silverstein and Bai \[1995\]](#) is slightly more general than that presented in Theorem 2.6. This model indeed assumes the presence of an additional deterministic matrix \mathbf{A} :

$$\mathbf{A} + \frac{1}{n} \mathbf{X}^\top \mathbf{C} \mathbf{X}$$

for $\mathbf{X} \in \mathbb{R}^{p \times n}$ with independent entries and \mathbf{A}, \mathbf{C} deterministic matrices (in fact, \mathbf{C} was imposed to be diagonal in [\[Silverstein and Bai, 1995\]](#) but this condition was later relaxed).

The bi-correlated model of the type $\frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{X} \tilde{\mathbf{C}} \mathbf{X}^\top \mathbf{C}^{\frac{1}{2}}$ introduced in Theorem 2.7 was later studied in [\[Paul and Silverstein, 2009\]](#), where not only the limiting spectrum but also the exact separation of the eigenvalues is studied. The extension of the spectral analysis of [\[Silverstein and Choi, 1995\]](#) for this model was then provided in [\[Couillet and Hachem, 2014\]](#); a convenient explicit Stieltjes transform inverse $z(\tilde{m})$ no longer exists for this model (due to the presence of a coupled system of equations), but inverse mapping theorems guarantee its existence and enable similar results.

For wireless communication purposes, this model was further extended in [\[Couillet et al., 2011\]](#) to

$$\sum_{i=1}^k \frac{1}{n_i} \mathbf{R}_i^{\frac{1}{2}} \mathbf{X}_i \mathbf{T}_i \mathbf{X}_i^\top \mathbf{R}_i^{\frac{1}{2}}$$

where $\mathbf{T}_i \in \mathbb{R}^{n_i \times n_i}$ and $\mathbf{R}_i \in \mathbb{C}^{p \times p}$ are symmetric nonnegative definite matrices standing respectively for the transmit (T) and receive (R) correlation matrices at each end of a communication channel between k devices equipped with n_1, \dots, n_k antennas and a single receiver equipped with p antennas. Establishing the limiting spectral measure of this model allows one to estimate the maximally achievable communication rates between k simultaneously transmitting mobiles terminals (phones, laptops, IoT devices) and a local base station. Further extensions of this model were then proposed to account for more involved wireless communication models, but they mostly all consist in summing independent versions of Gram matrices $\mathbf{Z}_i \mathbf{Z}_i^\top$ where \mathbf{Z}_i is a Gaussian (or beyond Gaussian) matrix with possibly nonzero mean, side correlations, a variance profile, etc.; see e.g., [\[Wen et al., 2013, Hachem et al., 2007, Wagner et al., 2012, Papazafeiropoulos and Ratnarajah, 2015\]](#) out of a much larger list of articles on the topic.

Of interest to statistics is also the *information-plus-noise* model of the type

$$\frac{1}{n} (\mathbf{X} + \mathbf{A})(\mathbf{X} + \mathbf{A})^\top$$

which is the sample “correlation” matrix between non-centered independent data $\mathbf{X} + \mathbf{A}$. This model also finds some interest in wireless communications where $[\mathbf{X} + \mathbf{A}]_{ij}$ models the statistical link between transmit antenna j and receive antenna i which may not be at the same mean-distance (controlled by \mathbf{A}_{ij}) than another antenna pair. This model was first studied by [Dozier and Silverstein \[2007\]](#) who established the (unique) canonical equation ruling the limiting spectral distribution of the model, as a function of the limiting Stieltjes transform of $\mu_{\mathbf{A}}$. Surprisingly enough, this model induces specific technical difficulties that left open for long the question of the exact location of the eigenvalues. Only much later in [\[Loubaton and Vallet, 2011\]](#) for the Gaussian case and then in [\[Capitaine, 2014\]](#) for the generic i.i.d. setting was the result fully obtained: that is, as for the sample covariance matrix result (Theorem 2.11), under compactness assumption on the eigenvalues of \mathbf{A} , none of the eigenvalues of $\frac{1}{n}(\mathbf{X} + \mathbf{A})(\mathbf{X} + \mathbf{A})^T$ asymptotically escapes the limiting support with high probability.

Yet, for practical applications, if the vectors of means $\mathbf{A}_{\cdot 1}, \dots, \mathbf{A}_{\cdot n}$ in the model $\mathbf{X} + \mathbf{A}$ are equal (to say vector $\boldsymbol{\mu}$), then $\mathbf{A} = \boldsymbol{\mu}\mathbf{1}_n^T$ reduces to a rank-one matrix, and $\frac{1}{n}(\mathbf{X} + \mathbf{A})(\mathbf{X} + \mathbf{A})^T$ is a mere spiked model, which does not necessitate the technical intricacies of the aforementioned articles. If instead the entries \mathbf{A}_{ij} are distinct with no specific structure, then it is in general not natural to assume that the \mathbf{X}_{ij} have equal variance (as the variance should scale with the mean). To handle this setting, [Hachem et al. \[2007\]](#), [Dumont et al. \[2010\]](#) study the generic *non-centered variance profile* model

$$\frac{1}{n}(\mathbf{B} \odot \mathbf{X} + \mathbf{A})(\mathbf{B} \odot \mathbf{X} + \mathbf{A})^T$$

where \mathbf{B} is a symmetric matrix and \odot is the entry-wise Hadamard product. There is no natural limiting spectral measure for this model (even when requiring the spectra of \mathbf{A}, \mathbf{B} to converge) but deterministic equivalents can be established. Those rely on a set of pn fixed-point equations. To our knowledge, no result on the conditions for the exact spectrum separation has been obtained in this setting. In the “separable case” where $\mathbf{B} = \mathbf{b}_1\mathbf{b}_2^T$ for some vectors $\mathbf{b}_1, \mathbf{b}_2$ (in which case $\mathbf{B} \odot \mathbf{X} = \text{diag}(\mathbf{b}_1)\mathbf{X} \text{diag}(\mathbf{b}_2)$), the solution reduces to two fixed-point equations and the exact asymptotic location of the eigenvalues is almost a direct application of the “no-eigenvalue outside the support” theorem for the bi-correlated and the information-plus-noise models (i.e., the extension of Theorem 2.11 to these models).

2.6.2.2 Advanced Wigner matrices

The generalizations of the Wigner random matrix model ($\frac{1}{\sqrt{n}}\mathbf{X}$ with \mathbf{X} having i.i.d. zero mean and unit variance entries) have been studied quite in parallel to the generalizations of the sample covariance matrix model $\frac{1}{n}\mathbf{XX}^T$, as the technical tools and proofs are quite alike (if not simpler).

The first extended model of historical interest was that of the deformed (i.e.,

nonzero mean) Wigner model of the type

$$\frac{1}{\sqrt{n}}(\mathbf{X} + \mathbf{A})$$

for \mathbf{A} symmetric and deterministic [Khorunzhy and Pastur, 1994]. Yet again, of utmost interest in practice is the case where the independent entries of \mathbf{X} have differing variances, which brings forth the model

$$\frac{1}{\sqrt{n}}(\mathbf{B} \odot \mathbf{X} + \mathbf{A}).$$

The set of the n^2 canonical implicit (Stieltjes transform related) equations induced for this model, or for its separable version ($\mathbf{B} = \mathbf{b}_1 \mathbf{b}_2^\top$), have been thoroughly investigated in [Ajanki et al., 2019].

In practice, these models are directly applicable to the adjacency matrices of random graphs ($[\mathbf{X} + \mathbf{A}]_{ij}$ is the connectivity between node i and node j) with independent link probabilities. The elementary case of such random graph models is the so-called Erdős–Rényi graph for which $\mathbf{X} + \mathbf{A}$ has i.i.d. Bernoulli $\{0, 1\}$ entries with parameter p . In this case $\mathbf{A} = p\mathbf{1}_n \mathbf{1}_n^\top$ is a rank-one matrix and \mathbf{X} has i.i.d. $\{-p, 1 - p\}$ entries such that $\mathbb{P}(\mathbf{X}_{ij} = 1 - p) = p$ and $\mathbb{P}(\mathbf{X}_{ij} = -p) = 1 - p$. This here boils down to a spiked model. Assuming the graph has *heterogeneous degrees*, in the sense that every particular node has its own probability q_i to connect to any other arbitrary node in the graph, we end up with the model $\text{diag}(\mathbf{q})\mathbf{X}\text{diag}(\mathbf{q}) + \mathbf{A}$ with $\mathbf{q} = [q_1, \dots, q_n]$, $\mathbf{X}_{ij} \in \{-q_i q_j, 1 - q_i q_j\}$ and $\mathbf{A}_{ij} = q_i q_j$. Here again $\mathbf{A} = \mathbf{q}\mathbf{q}^\top$ is a rank-one matrix.

2.6.2.3 (Real) Haar random matrices

Many algorithms and techniques in machine learning and data (or signal) processing involve random projections, in general on lower dimensional subspaces. This naturally calls for the study of matrix models involving random isometric matrices $\mathbf{U} \in \mathbb{R}^{p \times n}$, $n \leq p$, such that $\mathbf{U}^\top \mathbf{U} = \mathbf{I}_n$ (because then $\mathbf{U}\mathbf{U}^\top \in \mathbb{R}^{p \times p}$ is a projector on the n -dimensional subspace spanned by the columns of \mathbf{U}). These can be alternatively seen as the concatenated n columns of an underlying orthogonal matrix $\tilde{\mathbf{U}} \in \mathbb{R}^{p \times p}$.

Assuming $\tilde{\mathbf{U}}$ to be drawn uniformly in the space of unitary $p \times p$ matrices (this is called the *Haar* measure), \mathbf{U} is an *orthogonally invariant* random matrix, i.e., $\mathbf{V}\mathbf{U}\mathbf{W}$ has the same law as \mathbf{U} for any pair of deterministic orthogonal matrices $\mathbf{V} \in \mathbb{R}^{p \times p}$, $\mathbf{W} \in \mathbb{R}^{n \times n}$. However, unlike Gaussian random matrices $\mathbf{X} \in \mathbb{R}^{p \times n}$, which are also orthogonally invariant, the entries of \mathbf{U} are *not* independent as they must satisfy $\mathbf{U}^\top \mathbf{U} = \mathbf{I}_n$. This makes the study of the family of *Haar random matrices* more involved than the classical Gaussian case.

Yet, strong analogies exist between the Gaussian and the Haar random matrices. To start with, note that \mathbf{U} can be constructed from Gaussian random matrices by letting $\mathbf{U} = \mathbf{Z}(\mathbf{Z}^\top \mathbf{Z})^{-\frac{1}{2}}$ where $\mathbf{Z} \in \mathbb{R}^{p \times n}$, $n \leq p$, is a random matrix

with i.i.d. standard Gaussian entries (it suffices to verify that $\mathbf{U}^\top \mathbf{U} = \mathbf{I}_n$). Using this property, the fundamental trace lemma, Lemma 2.11, can be extended to a Haar-matrix equivalent [Debbah et al., 2003, Couillet et al., 2012].

Lemma 2.16 (Trace lemma for isometric matrices, [Couillet et al., 2012, Lemma 5]). *Let $\mathbf{U} \in \mathbb{R}^{p \times n}$ be $n < p$ columns of a $p \times p$ Haar random matrix and let $\mathbf{u} \in \mathbb{R}^p$ be a column of \mathbf{U} . Then, for $\mathbf{X} \in \mathbb{R}^{p \times p}$ a matrix function of the columns of \mathbf{U} , except \mathbf{u} , and of bounded operator norm,*

$$\mathbb{E} \left[\left| \mathbf{u}^\top \mathbf{X} \mathbf{u} - \frac{1}{p-n} \operatorname{tr} \mathbf{\Pi} \mathbf{X} \right|^4 \right] \leq \frac{C}{p^2}$$

where $\mathbf{\Pi} = \mathbf{I}_p - \mathbf{U} \mathbf{U}^\top + \mathbf{u} \mathbf{u}^\top$ (i.e., a projector on the complementary to the subspace spanned by the columns of \mathbf{U} , except \mathbf{u}) and C a constant depending only on the operator norm $\|\mathbf{X}\|$ and the ratio n/p .

Of course, since $\mathbf{U} \mathbf{U}^\top$ is a projection matrix, all its eigenvalues are 1 and 0 and there is thus no interest in the spectrum of $\mathbf{U} \mathbf{U}^\top$ itself. The above trace lemma however becomes handy when dealing with more structured models, such as $\mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{U}^\top \mathbf{C}^{\frac{1}{2}}$ for some deterministic \mathbf{C} matrix; the latter may be seen as a generalization of the sample covariance matrix model of Theorem 2.6. Specifically, we have the following result which provides a deterministic equivalent for this model.

Theorem 2.17 (Haar sample covariance [Couillet et al., 2012, Theorem 1]). *Let $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{U} \in \mathbb{R}^{p \times n}$, where $\mathbf{U} \in \mathbb{R}^{p \times n}$ are the $n < p$ columns of a $p \times p$ Haar random matrix and $\mathbf{C} \in \mathbb{R}^{p \times p}$ be symmetric nonnegative definite with bounded operator norm. Then, for $z < 0$, as $p/n \rightarrow c \in (1, \infty)$, letting $\mathbf{Q}(z) = (\frac{p}{n} \mathbf{X} \mathbf{X}^\top - z \mathbf{I}_p)^{-1}$, we have the deterministic equivalent*

$$\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z) = -\frac{1}{z} (\mathbf{I}_p + \tilde{m}_p(z) \mathbf{C})^{-1}$$

where $\tilde{m}_p(z)$ is the unique positive solution to

$$\tilde{m}_p(z) = \left(-z + (1 + zc^{-1} \tilde{m}_p(z)) \frac{1}{n} \operatorname{tr} \mathbf{C} (\mathbf{I}_p + \tilde{m}_p(z) \mathbf{C})^{-1} \right)^{-1}. \quad (2.36)$$

In the statement of the theorem, we used a correction $\frac{p}{n}$ in front of $\mathbf{X} \mathbf{X}^\top$ to ensure the correspondence between $\mathbb{E}[\mathbf{U} \mathbf{U}^\top] = \frac{n}{p} \mathbf{I}_p$ and the setting of Theorem 2.6 where $\mathbb{E}[\frac{1}{n} \mathbf{Z} \mathbf{Z}^\top] = \mathbf{I}_p$. Indeed, it is quite interesting to observe the tight relation between Theorem 2.6 and Theorem 2.17 which, despite the major difference imposed by the strongly dependent structure of \mathbf{U} versus the independent structure of \mathbf{Z} , leads almost to the same deterministic equivalent. The only difference lies in the extra term $zc^{-1} \tilde{m}_p(z)$ in the defining equation for $\tilde{m}_p(z)$.

Similar to the case of random matrices with i.i.d. entries versus Gaussian entries, it is nonetheless more convenient to work with Gaussian-specific identities

rather than the “independence”-related trace lemma. Specifically, an equivalent for Stein’s lemma, Lemma 2.13, also exists for Haar matrices.

Lemma 2.17 (Stein’s lemma for Haar matrices, from [Pastur and Shcherbina, 2011, Chapter 8]). *Let $\tilde{\mathbf{U}} \in \mathbb{R}^{p \times p}$ be a Haar matrix and $f : \mathbb{R}^{p \times p} \rightarrow \mathbb{R}$ a function admitting an analytic extension in the neighborhood of the set of unitary matrices in $\mathbb{R}^{p \times p}$. Then we have, for all $j, j' \in \{1, \dots, p\}$,*

$$\mathbb{E} \left[\sum_{i=1}^p f'_{ij}(\tilde{\mathbf{U}}) \tilde{\mathbf{U}}_{ij'} - f'_{ij'}(\tilde{\mathbf{U}}) \tilde{\mathbf{U}}_{ij} \right] = 0$$

where f'_{ij} is the classical derivative over $\tilde{\mathbf{U}}_{ij}$ (not accounting for the dependence of the other entries in $\tilde{\mathbf{U}}$). In the complex case ($\tilde{\mathbf{U}} \in \mathbb{C}^{p \times p}$ and $f(\tilde{\mathbf{U}}) \in \mathbb{C}$), this reduces to³⁴

$$\mathbb{E} \left[\sum_{i=1}^p f'_{ij}(\tilde{\mathbf{U}}) \tilde{\mathbf{U}}_{ij'} \right] = 0.$$

Similarly a Nash-Poincaré inequality for Haar matrix models is defined.

Lemma 2.18 (Nash-Poincaré for Haar matrices). *Under the setting of Lemma 2.17, we have*

$$\text{Var}(f(\tilde{\mathbf{U}})) \leq \frac{1}{p} \sum_{i,j=1}^p \mathbb{E} \left[|f'_{ij}(\tilde{\mathbf{U}})|^2 \right].$$

Although seemingly less exploitable, Stein’s lemma for Haar matrices is in fact quite convenient and easily leads to results such as the aforementioned Theorem 2.17 (for instance, by considering matrix functions of the form $f(\tilde{\mathbf{U}}\mathbf{D})$ for $\mathbf{D} \in \mathbb{R}^{p \times p}$ diagonal with $\mathbf{D}_{ii} = \delta_{i \leq n}$ — so that $f(\tilde{\mathbf{U}}\mathbf{D})$ only selects $n < p$ columns of $\tilde{\mathbf{U}} \in \mathbb{R}^{p \times p}$). Exercise 12 proposes to retrieve the result of Theorem 2.17 using both Gaussian (so applying Lemmas 2.17 and 2.18) and i.i.d. (so applying Lemma 2.16) approaches.

As a major difference between the i.i.d. and the Haar settings, note that Theorem 2.17 is stated under the constraint that z be real negative. In effect, Couillet et al. [2012] shows that it is far from trivial to extend the result to z beyond the negative real axis: in particular, unlike in the classical sample covariance setting of Theorem 2.6, in the “Haar sample covariance”, the fixed-point iteration in (2.36) fails to converge for $z = x + i\epsilon$ with $x > 0$ and $\epsilon \ll 1$. This particularly makes it difficult to exploit the result to retrieve (both theoretically³⁵ and numerically) the limiting spectral measure of $\mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{U}^T \mathbf{C}^{\frac{1}{2}}$.

³⁴Similar to what we saw in Remark 2.5, it is in practice convenient to work under a complex (unitary) \mathbf{U} setting, even in the real (orthogonal) case, as deterministic equivalents are universal with respect to the underlying field (real or complex) of the entries of \mathbf{U} .

³⁵One may claim that, since convergence holds for all $z < 0$, as per Vitali’s convergence theorem (Theorem 2.3), it can then be extended to all of $\mathbb{C} \setminus \mathbb{R}^+$. This is however not so simple as it is difficult to ensure that $\tilde{m}_p(z)$ in Theorem 2.17, as defined through its fixed-point equation, is indeed analytic in a certain cone $\{z = e^{i\theta} \mid \theta \in (-\theta^\circ, \theta^\circ) \setminus \{0\}\}$ ($\theta^\circ \in (0, \pi)$).

There in fact exists a whole other branch of tools in the random matrix literature, called *free probability theory*, which much more easily recovers Theorem 2.17 (under a different formulation though) and also obtains the limiting spectral measure of $\frac{p}{n}\mathbf{C}^{\frac{1}{2}}\mathbf{U}\mathbf{U}^T\mathbf{C}^{\frac{1}{2}}$: specifically, using an interesting extension to random matrices of classical probability theory on scalar random variables, free probability theory demonstrates that the limiting measure is the so-called *free multiplicative convolution* (see Section 2.6.2.4 below for a proper definition) of the limiting spectral measure ν of \mathbf{C} and of the limiting measure of $\mathbf{U}\mathbf{U}^T$ (i.e., the discrete measure $\delta_1 + \frac{p-n}{n}\delta_0$). The next section provides a short introduction to the main ingredients of free probability theory.

Turning to machine learning applications of results on (derivatives of) Haar random matrix models, to the best of our knowledge, very few works have so far fully exploited the strength of these identities. For this reason, we will not elaborate much more on these aspects and will only briefly introduce free probability theory, which has many advantages (especially when dealing with Haar or permutation-invariant random matrices) but also strong limitations when compared to the Stieltjes transform method. We thus point the interested reader to the literature on random Haar matrices for more detail on the topic. Those tools may nonetheless reveal fundamental in the future for the analysis of specific random projection or random permutation-based methods with isometric constraints in machine learning and AI.

2.6.2.4 The free probability approach

Free probability theory is a drastically different approach to study random matrices. It is particularly efficient in some scenarios, such as when the sum or product of random matrices are involved. The theory was developed in parallel to the Stieltjes transform method discussed in this monograph and originates from the works of Voiculescu et al. [1992], who originally aimed to describe a theory of probabilities on non-commutative algebras. A detailed introduction of the theory is beyond the scope of this monograph and we refer the interested readers to [Hiai and Petz, 2006, Biane, 1998] and [Couillet and Debbah, 2011, Chapter 4 and Chapter 5]. Although free probability theory is rooted in a combinatorial approach (see e.g., [Nica and Speicher, 2006]), it also contains some elegant analytic results, which can be related to the Stieltjes transform: in the sequel, we emphasize those useful results.

For μ and ν two compactly supported probability measures on $[0, \infty)$, Hiai and Petz [2006] proved that there always exist two *free random variables* a and b in some non-commutative probability space having distributions μ and ν , respectively. The distribution of $a + b$ and ab depend solely on μ, ν and can be associated with probability measures called *free additive convolution* and *free multiplicative convolution* of the distributions μ and ν , denoted $\mu \boxplus \nu$ and $\mu \boxtimes \nu$, respectively. These measures are both compactly supported on $[0, \infty)$ [Voiculescu et al., 1992].

These free additive and multiplicative convolutions satisfy convenient ana-

lytic expressions, through the so-called R - and S -transforms introduced below.

Definition 5 (R - and S -transform). *Let μ be a real probability measure with support $\text{supp}(\mu)$ and Stieltjes transform $m_\mu(z)$, for $z \in \mathbb{C}^+$. The R -transform of μ , denoted R_μ , is defined as the solution to*

$$m_\mu(R_\mu(z) + z^{-1}) = -z$$

or equivalently

$$m_\mu(z) = \frac{1}{R_\mu(-m_\mu(z)) - z}.$$

Next, let $\psi_\mu(z)$ be defined as

$$\psi_\mu(z) = \int \frac{zt}{1-zt} \mu(dt) = -1 - z^{-1} m_\mu(z^{-1})$$

and let χ_μ be its unique functional inverse, analytic in the neighborhood of zero, i.e., $\chi_\mu(\psi_\mu(z)) = z$ for $|z|$ small enough. Then, the S -transform of μ , denoted S_μ , is given by

$$S_\mu(z) = \chi_\mu(z) \frac{1+z}{z}.$$

In particular, $S_\mu(z)$ satisfies

$$m_\mu\left(\frac{z+1}{zS_\mu(z)}\right) = -zS_\mu(z).$$

The main property of R - and S -transforms is summarized below, and requires the notion of *freeness* between non-commutative random variables. Freeness is not an easy notion, and is defined through a series of moment conditions and combinatorial calculus, which we will not go into here (see again [Hiai and Petz, 2006, Biane, 1998] for details). One needs just remember at this point that freeness extends the notion of independence to non-commutative random variables.

Lemma 2.19 (R - and S -transforms of sums and products). *For a and b two free random variables with compactly supported distributions μ and ν , respectively, the law $\mu \boxplus \nu$ of $a + b$ satisfies*

$$R_{\mu \boxplus \nu}(z) = R_\mu(z) + R_\nu(z).$$

Similarly, the law $\mu \boxtimes \nu$ of ab satisfies

$$S_{\mu \boxtimes \nu}(z) = S_\mu(z)S_\nu(z).$$

Of interest to the present monograph is that “asymptotically large random matrices” are typical examples of non-commutative random variables for which freeness can be ensured. To avoid dealing with infinite-size linear operators, it is more appropriate to define a notion of *asymptotic freeness* for finite-dimensional

random matrices, which translates the freeness of their respective limiting operators.³⁶

As such, the main result of interest to us is the following: for $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\mathbf{B} \in \mathbb{R}^{n \times n}$ two asymptotically free random matrices with respective *limiting spectral measures* μ_A and μ_B , the limiting spectral measure μ_{A+B} ($A+B$ is here merely a notation with no formal meaning) of $\mathbf{A}+\mathbf{B}$ exists and satisfies

$$\mu_{A+B} = \mu_A \boxplus \mu_B, \quad R_{A+B}(z) = R_A(z) + R_B(z)$$

for $R_A(z)$, $R_B(z)$ and $R_{A+B}(z)$ the R -transforms of μ_A , μ_B and μ_{A+B} , respectively. Similarly, μ_{AB} , the limiting spectral measure of the matrix product \mathbf{AB} , exists and satisfies

$$\mu_{AB} = \mu_A \boxtimes \mu_B, \quad S_{AB}(z) = S_A(z)S_B(z)$$

for $S_A(z)$, $S_B(z)$ and $S_{A+B}(z)$ the S -transforms of μ_A , μ_B and μ_{A+B} . The above equalities should be understood to hold in the almost sure sense.

Clearly, the asymptotically freeness assumption plays a key role in relating the limiting spectrum of $\mathbf{A}+\mathbf{B}$ or \mathbf{AB} to that of \mathbf{A} and \mathbf{B} , which unfortunately in practice only applies to a limited range of random matrices. Essentially, \mathbf{A} and \mathbf{B} are asymptotically free if they are both independent and if the distribution of their respective eigenvectors are sufficiently “isotropic” with respect to one another: so essentially, when one of the two matrices is invariant by left and right product by arbitrary unitary matrices. Essentially, the two major cases of matrix pairs known to be asymptotically free are: (i) a Gaussian standard random matrix and any other independent random matrix (for instance a deterministic matrix or another Gaussian standard random matrix, independent of the first), (ii) a Haar random matrix and any other independent random matrix. One may for instance easily determine the limiting spectral measure of models of the type $\mathbf{X} + \mathbf{A}$ for \mathbf{X} a Wigner matrix or a Wishart matrix and \mathbf{A} deterministic, or even $\mathbf{X}\mathbf{A}\mathbf{X}^\top$ with \mathbf{X} Gaussian or Haar distributed. These objects are few because it is technically difficult to establish asymptotic freeness, the formal definition of which is a matter of heavy combinatorial calculus (see e.g., [Biane, 1998, Section 3]). As a result, free probability theory is more complex to use when summing or multiplying two random matrices with structured eigenvectors, such as simple models like $\mathbf{X} \odot \mathbf{B} + \mathbf{A}$ for \mathbf{X} a Wigner matrix and \mathbf{B} a deterministic variance profile: these matrices are not free with respect

³⁶One must be careful in the whole monograph that we never define large dimensional random matrices as being of “infinite” dimensions (which would turn them, when correctly defined, into operators in an infinite-dimensional Hilbert space): all the objects treated throughout the book are finite dimensional objects, some functionals of which are studied when the size of the matrices increase. For actual works on operators, seen as limit of random matrices in infinite-dimensional spaces, see [Pastur and Figotin, 1992] on almost-periodic random operators. Aside from a few exceptions though [Hachem et al., 2015], these elegant works find little practical applications in systems and software engineering. This being said, it is theoretically interesting to observe that the Stieltjes transform approach thoroughly developed in the present monograph shares many common grounds with the more general theory of linear operators in Hilbert spaces [Akhiezer and Glazman, 2013].

to deterministic matrices, so that the R - and S -transform formulas cannot be exploited; this very fact has strongly limited the reach and importance of free probability theory in the past decade.

A fundamental result to efficiently use the additive and product rules of Lemma 2.19 are the basic forms of the R - and S -transforms of elementary random matrix models. Specifically, the R - and S -transforms of the Marčenko-Pastur and semicircle distributions are known in closed-form.

Lemma 2.20 (R and S -transforms of Marčenko-Pastur and semicircle law). *The R -transform $R_{\text{MP},c}(z)$ and S -transform $S_{\text{MP},c}(z)$ of the Marčenko-Pastur law $\mu_{\text{MP},c}$ of parameter c , i.e., the limiting spectral measure of $\frac{1}{n}\mathbf{Z}\mathbf{Z}^\top$, $\mathbf{Z} \in \mathbb{R}^{p \times n}$ with i.i.d. zero mean, unit variance entries, as $p/n \rightarrow c \in (0, \infty)$, given explicitly by (2.10), read*

$$R_{\text{MP},c}(z) = \frac{1}{1 - cz}, \quad S_{\text{MP},c}(z) = \frac{1}{1 + cz}. \quad (2.37)$$

As for the R -transform $R_{\text{SC}}(z)$ of the semicircle law μ_{SC} , given by (2.22), it is simply

$$R_{\text{SC}}(z) = z. \quad (2.38)$$

With Lemma 2.20, one is able to derive, with a free probability approach, the limiting spectral measure of the information-plus-noise-type random matrix model $\mathbf{M} = \mathbf{A} + \frac{1}{n}\mathbf{X}\mathbf{X}^\top$ for $\mathbf{X} \in \mathbb{R}^{p \times n}$ having i.i.d. standard Gaussian entries and $\mathbf{A} \in \mathbb{R}^{p \times p}$ a deterministic matrix. Specifically, calling μ_A and μ_M the limiting spectral measure of A and M as $n, p \rightarrow \infty$ with $p/n \rightarrow c$, we have

$$\mu_M = \mu_A \boxplus \mu_{\text{MP},c}, \quad R_M(z) = R_A(z) + R_{\text{MP},c}(z)$$

so that, by Definition 5 and Lemma 2.20,

$$m_M(z) = \frac{1}{R_M(-m_M(z)) - z} = \frac{1}{R_A(-m_M(z)) + \frac{1}{1 + cm_M(z)} - z}$$

or equivalently

$$R_A(-m_M(z)) + \frac{1}{-m_M(z)} = z - \frac{1}{1 + cm_M(z)}$$

which, by taking the Stieltjes transform $m_A(\cdot)$ of the limiting law of \mathbf{A} on both sides, together with Definition 5, gives

$$m_M(z) = m_A \left(z - \frac{1}{1 + cm_M(z)} \right).$$

The same result would have been more painstaking to derive using a purely Stieltjes transform approach. However, very few matrix models being asymptotically free, the free probability framework quickly fails to operate in more structured random matrix models.

Recent works try to cope with these limitations as well as open the range of applicability of free probability theory to handle sums and products of matrices under weaker forms of asymptotic freeness conditions (to characterize for instance the limiting spectrum of the sum of random matrices with row and columns permutation invariance [Au et al., 2018], or to extend the notion of deterministic equivalents to a free probability setting [Speicher and Vargas, 2012]). Despite these efforts, the major trend and most flexible tools in random matrix theory remain the approaches linked to the resolvent and Stieltjes transform approaches, which are the focus of the monograph.

2.6.2.5 Full circle law, β -ensembles, sparse random matrices, etc.

Mathematicians have long been intrigued by the “simplest” random matrix model in appearance that is $\frac{1}{\sqrt{n}}\mathbf{X} \in \mathbb{R}^{n \times n}$ (non-symmetric) with i.i.d. zero mean and unit variance entries. Being non-symmetric (at least with probability one), the eigenvalues of $\frac{1}{\sqrt{n}}\mathbf{X}$ are complex. They have long been known to spread uniformly on the unit complex disc $\{z \in \mathbb{C}, |z| < 1\}$. Surprisingly though, despite its simple statement, this result, known as the *full circle law* or the *circular law*, has only been proven in full generality very recently by Tao and Vu [2008]. To explain the difficulties: (i) the Stieltjes transform method cannot be applied directly as the spectrum is complex (and thus taking a limit $z \rightarrow z_0$ for z_0 in the support does not allow to “enter” the complex support as in the real eigenvalue case); there the solution was provided earlier by Girko who introduced the alternative V-transform; (ii) the V-transform involves the limit of an integral form on the logarithm of the singular values of \mathbf{X} which, being square, tends to have a lot of singular values tending to zero (the singular values of \mathbf{X} are the square roots of the eigenvalues of $\mathbf{X}\mathbf{X}^\top$ with \mathbf{X} of size $p \times n$ where $p = n$: i.e., this is the technically most difficult *hard edge* scenario of the Marčenko-Pastur law depicted in Figure 2.2 with $c = 1$); this technical difficulty, previously worked around by invoking the existence of high order moments for \mathbf{X}_{ij} was solved by Tao and Vu by means of the *ϵ -net technique*, popular today in compressive sensing.

From the perspective of the present monograph, the eigenvalues of non-symmetric models are of marginal interest. These could be used for the analysis of directed random graphs although, to our knowledge, not much works exist in this direction.

Another purely mathematical interest relates to the fact that Gaussian random matrices are much better known than random matrices with i.i.d. entries and, consequently, come along with a host of other technical tools. In particular, not only the limiting spectral measure, but actually the exact *finite-dimensional joint distribution* $\mathbb{P}(\lambda_1, \dots, \lambda_n)$ of (real, complex, or quaternionic) Gaussian symmetric random matrix \mathbf{X} and $\frac{1}{n}\mathbf{X}\mathbf{X}^\top$ (with \mathbf{X} having i.i.d. entries) are known. The expressions of $\mathbb{P}(\lambda_1, \dots, \lambda_n)$ for all these cases are quite related.

In particular, the joint eigenvalue distribution for the Gaussian Wigner ma-

trix $\mathbf{X} \in \mathbb{R}^{n \times n}$ is explicitly given by

$$\mathbb{P}(\lambda_1, \dots, \lambda_n) \propto \prod_{i=1}^n e^{-\frac{1}{4}\beta n \lambda_i} \prod_{1 \leq i < j \leq n} |\lambda_i - \lambda_j|^\beta \quad (2.39)$$

for real Gaussian \mathbf{X} when $\beta = 1$ (the random ensemble is called here the Gaussian orthogonal ensemble, GOE), complex Gaussian \mathbf{X} when $\beta = 2$ (referred to as the Gaussian unitary ensemble, GUE), and quaternionic Gaussian \mathbf{X} when $\beta = 4$ (the Gaussian symplectic ensemble, GSE). Much work has been devoted to the study of the asymptotics of the joint law of this now called β -ensemble of random matrices. In particular, the Tracy-Widom limit for the largest eigenvalue introduced in Theorem 2.15 is obtained by marginalizing the joint measure to obtain the probability $\mathbb{P}(\lambda_1 > x)$. See [Anderson et al., 2010] for an introduction to these quite different methods.

Most of the aforementioned random matrix models however share as a common denominator their relying on $O(n^2)$ “degrees of freedom”, in the sense that they are designed out of $O(n^2)$ independent random variables. For the sample covariance matrix model $\frac{1}{n}\mathbf{XX}^\top$, we have $\mathbf{X} = \mathbf{C}^{\frac{1}{2}}\mathbf{Z} \in \mathbb{R}^{p \times n}$ with \mathbf{Z} made of np independent entries. For the Wigner model, $\mathbf{X} \in \mathbb{R}^{n \times n}$ has $n(n+1)/2$ independent entries on and above the diagonal. This large number of degrees of freedom is the *major asset of random matrix theory*, as presented in this monograph: they trigger (i) concentration properties that do not appear if p is fixed and only $n \rightarrow \infty$ (such as quadratic form concentration $\frac{1}{p}\|\mathbf{z}_i\|^2 \xrightarrow{a.s.} 1$), (ii) fast convergence rates with typical central limit speeds of order up to $O(1/n)$ and, possibly most importantly, (iii) *universality* with respect to the underlying distribution of the independent entries (i.e., asymptotic statistics loosely depend on the actual law of the entries) which simplifies the analysis and provides robustness of the studied objects to deviations from the statistical model.

Yet, a host of practical random matrix models demand less degrees of freedom. Realistic networks for instance (social nets, brain connectivity, molecular networks, etc.) are naturally modeled by *sparse* random (say symmetric) adjacency matrices $\mathbf{A} \in \mathbb{R}^{n \times n}$ with typical number of nonzero elements scaling as $O(n)$ rather than $O(n^2)$. Every row/column \mathbf{a}_i of \mathbf{A} typically has $O(1)$ nonzero elements (corresponding to the neighbors or contacts of node i in the underlying graph), and thus $\|\mathbf{a}_i\|$ does not concentrate. Kernel random matrices $\mathbf{K} = \{f(\|\mathbf{x}_i - \mathbf{x}_j\|^2)\}_{i,j=1}^n$ of finite-dimensional vectors $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ with p small (such as used in the context of classification or clustering of 2D or 3D points) are also more challenging to study than their large- p counterpart, as every entry of \mathbf{K} remains a random variable which does not concentrate in the large- n alone limit. The consequences are numerous: (i) the analysis of these objects is more difficult, if doable at all, and only few large- n statistics can be studied, (ii) universality and robustness to model assumptions are lost: large- n asymptotics remain a function of the law of the data vectors \mathbf{x}_i . The hard-to-obtain results are thus additionally less applicable in practice.

Nevertheless, a branch of random matrix theory is interested in these important models. Stieltjes transform methods are here mostly ineffective so that,

for lack of better alternatives, one has to rely essentially on moment approaches and combinatorics. A particularly interesting approach when it comes to sparse random graphs of size n is that, as $n \rightarrow \infty$, the graph has a “tree-like” structure; indeed, with a probability $O(1/n)$ for each node to reach out any another node, the probability of presence of cycles in the graph is vanishingly small. This has motivated the independent development of a graph-based random matrix framework, strongly pushed by Bordenave and Lelarge in e.g., [Bordenave et al., 2011, Bordenave and Lelarge, 2010]. The results are however generally “weak” from a practical standpoint. For instance, while it has long been known that the spectral measure of a *dense* Erdős–Rényi random graph \mathbf{A} with Bernoulli i.i.d. entries (i.e., with $O(n^2)$ degrees of freedom) converges to the semicircle law, it is still unknown to which measure a *sparse* random graph \mathbf{A} converges: the limiting law is known to exist, to be decomposed as the sum of a (known) discrete measure and a (unknown) continuous measure, and to have an unbounded support (as opposed to the semi-circular distribution) [Salez, 2011].

The specific kernel random matrix $\mathbf{K} = \{\|\mathbf{x}_i - \mathbf{x}_j\|^2\}_{i,j=1}^n$ with \mathbf{x}_i of fixed dimension, known as a Euclidean random matrix, has also been studied in [Bordenave, 2008], but again with results of limited practical reach.

Aside from side comments, the monograph will not dig into these fundamentally different problems, tools, and results. We exclusively concentrate on dense random matrix models.

2.6.3 Other statistics

Most of the statistics of practical interest in the application chapters are directly related to deterministic equivalents of the resolvent of random matrices and to their linear statistics. For instance, we shall see that the performance of classification methods (correct classification rates) of n data vectors $\mathbf{x}_1, \dots, \mathbf{x}_n$ in a k -class ($\mathcal{C}_1, \dots, \mathcal{C}_k$) problem can in general be estimated from the k -dimensional matrix of quadratic forms

$$\frac{1}{n} \mathbf{J}^\top \mathbf{Q}(z) \mathbf{J}$$

(or some closely related statistics) where $\mathbf{Q}(z)$ is the resolvent of the underlying affinity matrix of the data (kernel, graph Laplacian, etc.) and $\mathbf{J} = [\mathbf{j}_1, \dots, \mathbf{j}_k] \in \mathbb{R}^{n \times k}$ with $[\mathbf{j}_a]_i = \delta_{\mathbf{x}_i \in \mathcal{C}_a}$ the canonical vector of the classes.

Yet, some specific results (such as the classification rate of some random neural networks, the exact proof of the asymptotic Gaussian behavior of the dominant eigenvector entries in graph adjacency and kernel matrices, etc.) demand more than just *first order* limiting statistics. A further common statistics of interest lies in the *second order fluctuations*, i.e., in central limit theorems, of the objects under study.

These statistics have long been studied in the random matrix literature, starting with the works of Bai and Silverstein [2004] who, under the sample

covariance setting of Theorem 2.6, established a central limit of the type

$$n \int f(t) (\mu_{\frac{1}{n}} \mathbf{X} \mathbf{X}^\top - \mu)(dt) \rightarrow \mathcal{N}(M(f), \sigma^2(f))$$

for all analytic functions f . This result (and all similar results in related models) has the following noteworthy properties:

- the convergence speed is of order $O(n^{-1})$. This however only stands for linear statistics of the eigenvalues; fluctuations of bilinear forms $\mathbf{a}^\top (\mathbf{Q}(z) - \bar{\mathbf{Q}}(z)) \mathbf{b}$ fluctuate at a slower $O(n^{-\frac{1}{2}})$ speed;
- the mean (or bias term) $M(f)$ and variance $\sigma^2(f)$ depend on $\mathbb{E}[|\mathbf{Z}_{ij}|^4]$ (which, thus, must be finite). Both can be decomposed as a sum $A + \kappa B$ with κ the kurtosis of \mathbf{Z}_{ij} . The mean in particular vanishes in the complex Gaussian case and the variance is twice as large in the real Gaussian as in the complex Gaussian case.

Many results on central limit theorems for a vast spectrum of linear statistics of random models have been established, for instance in [Hachem et al., 2008] for sample covariance matrices $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ with \mathbf{X} having a variance profile, in [Lytova and Pastur, 2009] to Wigner matrix model for less smooth functions f (five-times differentiable), or in [Zheng et al., 2017] for F-matrix models of the type $(\frac{1}{n_1} \mathbf{X}_1 \mathbf{X}_1^\top)^{-1} \frac{1}{n_2} \mathbf{X}_2 \mathbf{X}_2^\top$. A generalization to three-times differentiable f is proposed in [Najim and Yao, 2016]. Central limit theorems for bilinear forms are found for instance in [Kammoun et al., 2009]. Fluctuations of the isolated eigenvalues and eigenvector projections in a spiked random matrix model can also be found in [Baik et al., 2005, Bai and Yao, 2008, Couillet and Hachem, 2013]. These fluctuations are at the slowest $O(n^{-\frac{1}{2}})$ rate.

A central limit result for the linear statistical inference method of Theorem 2.12 has also been established in [Yao et al., 2013]. There again it is shown that the convergence speed is of order $O(n^{-1})$ with a bias and a variance of the form $A + \kappa B$ with κ the kurtosis of the underlying distribution (and, again, the bias is only zero in the complex Gaussian case). An estimation method is also proposed for the means and variances, which is of practical interest to empirically assess the confidence interval of the estimator.

Due to a strong motivation from the field of wireless communications, some specific linear statistics have been particularly widely studied in the random matrix literature. This is notably the case of the logarithm. Statistics of the type

$$\int \log(1 + st) \mu_{\frac{1}{n}} \mathbf{X} \mathbf{X}^\top(dt)$$

for $s > 0$ are particularly important in wireless communications as they give access to the achievable communication rate over a linear wireless communication channel \mathbf{X} . This $\log(1 + st)$ term arises from the entropy of Gaussian random variables and is also found in many other applications, such as with the

estimation of the Kullback-Liebler divergence between two multivariate Gaussian vectors discussed in Chapter 3. A particularly convenient feature of the integral form $\int \log(1+st)\mu(dt)$ is that its derivative with respect to s (that is $\int t/(1+st)\mu(dt)$) is immediately related to the Stieltjes transform of μ .³⁷ It is thus not required to use a complex contour integral method to assess these quantities (a real integration is sufficient). See [Tulino and Verdú, 2004, Couillet and Debbah, 2011] for a detailed account of these findings.

In technical terms, there are essentially two major methods to obtain central limit theorems of random matrix quantities. Recalling that linear functionals $u(\mathbf{Q})$ of the resolvent $\mathbf{Q} = (\mathbf{X} - z\mathbf{I}_n)^{-1}$ of the random matrix \mathbf{X} under study (e.g., bilinear forms $\mathbf{a}^T \mathbf{Q} \mathbf{b}$ or traces $\text{tr } \mathbf{A} \mathbf{Q}$) is usually the central object of interest, it cannot in general be expressed as a sum of independent random variables. Instead, Silverstein and Bai propose to use the martingale difference approach, which we previously exploited in the detailed proof of the Marčenko-Pastur theorem (Theorem 2.4). Recall indeed that, if \mathbf{X} has independent columns, it is convenient to write

$$u(\mathbf{Q}) - \mathbb{E}u(\mathbf{Q}) = \sum_{i=1}^n \mathbb{E}_{i-1}[u(\mathbf{Q})] - \mathbb{E}_i[u(\mathbf{Q})]$$

where \mathbb{E}_i is the expectation conditioned on the columns $\mathbf{x}_1, \dots, \mathbf{x}_i$ of \mathbf{X} , with the convention $\mathbb{E}_0[u(\mathbf{Q})] = u(\mathbf{Q})$. This is a sum of martingale differences, for which [Billingsley, 2012, Theorem 35.12] provides a central limit theorem (see also [Bai and Silverstein, 2010, Chapter 9]).

Alternatively, Pastur proposes to use Gaussian techniques, which we also explored in the alternative proof of the Marčenko-Pastur theorem, along with a *characteristic function approach* (see examples in [Pastur and Shcherbina, 2011]) to show that

$$\mathbb{E} \left[e^{-\imath t u(\mathbf{Q})} \right] \rightarrow e^{-\imath t \mu - \frac{1}{2} t^2 \sigma^2}$$

which is the Gaussian characteristic function. To reach this convergence, the approach consists in exploiting the Gaussian integration-by-parts formula (Stein's lemma) on the differentiated (along t) left-hand expectation, i.e.,

$$\mathbb{E} \left[-\imath u(\mathbf{Q}) e^{-\imath t u(\mathbf{Q})} \right].$$

Exploiting the fact that u is linear and that $\mathbf{Q} = -\frac{1}{z}\mathbf{I}_n - \frac{1}{z}\mathbf{Q}\mathbf{X}$, this expectation can be reduced as a function of the type $\mathbb{E}[\mathbf{X}f(\mathbf{X})]$ on which the Gaussian integration-by-parts method can be applied. The objective is then to show that this differentiated characteristic function converges to the derivative of the limiting Gaussian characteristic function, i.e., $(-\imath\mu - t\sigma^2)e^{-\imath t \mu - \frac{1}{2} t^2 \sigma^2}$. This is in particular performed by controlling the small terms using the Nash–Poincaré inequality.

³⁷Specifically, $\int t/(1+st)\mu(dt) = s^{-1}(1 - s^{-1}m_\mu(-s^{-1}))$.

2.7 Beyond vectors of independent entries: concentration of measure in RMT

2.7.1 Limitations of the i.i.d. assumption

In the previous sections, we have shown that the Stieltjes transform and resolvent approaches are quite versatile tools which, in a way, form a surrounding “complex analysis and linear algebra core” for random matrix theory analysis. This core must, however, must be *independently supplemented* by appropriate probabilistic tools (which ensure the necessary convergences for linear algebra and complex analysis methods to be exploited).

When it comes to these probabilistic methods, we have seen that a major driver for most of the results lies in exploiting the *independence both in sample and features* of the entries of the underlying random matrix \mathbf{X} . It is thus no wonder that a natural and long-standing assumption in the early works in random matrix theory was to request for \mathbf{X} to have all independent (or linearly dependent as in models such as $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \tilde{\mathbf{C}}^{\frac{1}{2}} + \mathbf{A}$) entries. Most generalizations of these results usually assume mere deviations from this setting (by allowing weak, or asymptotically vanishing, correlation between the entries for instance).

However, while for random graphs it is largely conceivable to request independent “noise” associated to each link and for random vector observations it is natural to ask for these observations to be independent, requesting that every single observation is made of independent entries is very constraining. Note in particular that what we referred to as the *sample covariance matrix model* in Theorem 2.6 is in fact a very restricted model where every observation \mathbf{x}_i needs be of the form $\mathbf{x}_i = \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$ for some \mathbf{z}_i with independent entries. This model is mostly convenient only for the Gaussian case where $\mathbf{z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ and as a result $\mathbf{x}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{C})$. Most multivariate random vectors \mathbf{x}_i with zero mean and covariance \mathbf{C} (elliptical distributions, correlated vectors of Bernoulli entries, etc.) cannot be factorized under this form.

Most importantly, the “real data” \mathbf{x}_i (images, sounds, videos, DNA sequences, population features) met in machine learning applications tend to live in very contorted manifolds that cannot be linearly “whitened” into a vector of independent entries by merely operating $\mathbf{C}^{-\frac{1}{2}} \mathbf{x}_i$.

2.7.2 Concentrated random vectors as the answer

In [El Karoui, 2009] and [Pajor and Pastur, 2009], El Karoui and Pajor-Pastur were the first to realize (or at least to fully exploit the fact) that, from a probability standpoint, the proof of the sample covariance matrix result in Theorem 2.6 from [Silverstein and Bai, 1995], only relies on (i) the independence between the (*column*) vectors \mathbf{x}_i composing $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{Z}$ (and not of all the entries), and (ii) the convergence

$$\frac{1}{n} \mathbf{x}_i^\top \mathbf{Q}_{-i}(z) \mathbf{x}_i - \frac{1}{n} \operatorname{tr} \mathbf{Q}_{-i} \mathbf{C} \rightarrow 0 \quad (2.40)$$

in some probabilistic sense, where $\mathbf{Q}_{-i}(z) = (\frac{1}{n}\mathbf{XX}^\top - \frac{1}{n}\mathbf{x}_i\mathbf{x}_i^\top - z\mathbf{I}_p)^{-1}$. For the latter, it is sufficient but *not necessary* for $\mathbf{z}_i = \mathbf{C}^{-\frac{1}{2}}\mathbf{x}_i$ to have standard i.i.d. entries. In particular, El Karoui showed that this convergence also holds if \mathbf{x}_i is a *concentrated random vector*: a fundamental property at the core of our present concern and which, we will show, has far reaching consequences to the application of random matrix results to real data.

In a nutshell, the concentration of measure theory, as extensively developed by Ledoux [2005], considers random vectors $\mathbf{x} \in \mathbb{R}^p$ having the property that every 1-Lipschitz functional $\phi : \mathbb{R}^p \rightarrow \mathbb{R}$ of \mathbf{x} is “predictable” in the sense that there exists a deterministic value $M_\phi \in \mathbb{R}$ such that the random variable $\phi(\mathbf{x})$ remains in the neighborhood of M_ϕ and that the diameter of this neighborhood vanishes as $p \rightarrow \infty$. This notion must *not* be confused with the fact that \mathbf{x} itself converges, which is in general largely wrong: only the *scalar projections* $\phi(\mathbf{x})$ do converge (we will say in this case that \mathbf{x} “concentrates”).³⁸ More formally, assuming $M_\phi = O(1)$ with respect to p (otherwise, it needs to be appropriately scaled), there exists a function $\alpha(t, p)$ decreasing to zero in both t and p such that

$$\mathbb{P}(|\phi(\mathbf{x}) - M_\phi| > t) \leq \alpha(t, p). \quad (2.41)$$

Of particular interest is the family of functions $\alpha(t, p) = e^{-t^\beta p^\gamma}$ for some $\beta, \gamma > 0$ which, since the exponential grows faster than any polynomial, provides a more powerful and much more flexible inequality than the moment bounds introduced in the proof of the Marčenko-Pastur law.³⁹ The mapping $\mathbf{x}_i \rightarrow \frac{1}{n}\mathbf{x}_i^\top \mathbf{Q}_{-i}(z)\mathbf{x}_i$ in (2.40) is however not Lipschitz, and thus more profound technical considerations are requested to show that Theorem 2.6 extends to the case where the vectors \mathbf{x}_i are independent concentrated random vectors. This is performed in an intricate manner in [El Karoui, 2009]. A more systematic approach has been recently developed in [Louart and Couillet, 2018], the basics of which are discussed in the next section.

Paradoxically, very few “classical” multivariate distributions are known to produce concentrated random vectors, and yet, this is enough to bring an outstanding practical competitive advantage against vectors with independent entries, when it comes to modeling real data for machine learning.

Of these popular distributions, only the Gaussian random vector $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, the uniformly distributed vector \mathbf{x} on the unit sphere of \mathbb{R}^p , and the vector \mathbf{x} with i.i.d. entries with bounded support (i.e., $|x_i| < K$ for some $K > 0$) are

³⁸As a matter of fact, as we will see, the concept of concentration is even more general in that it allows one to control the rate of fluctuations of $\phi(\mathbf{x})$, for arbitrary $\phi : \mathbb{R}^p \rightarrow \mathbb{R}^{p'}$ (so with $p' \geq 1$ generic), even when the latter does not converge. Lipschitz functions being stable operators through composition, iterated controls of Lipschitz functions with various Lipschitz constants enables a thin tracking of the behavior of sometimes intricate non-linear functionals of \mathbf{x} (such as through the layers of a neural network).

³⁹Of course, as a compensation for this simplification, this imposes more constraints on the entries of the random vector \mathbf{x} , such as the existence of moments of all orders. But, as far as practical statistical machine learning considerations are concerned, this is far from a heavy request.

2.7. BEYOND VECTORS OF INDEPENDENT ENTRIES: CONCENTRATION OF MEASURE IN RMT141

known to be concentrated random vectors. Worse, for the latter, the definition (2.41) only holds for all 1-Lipschitz *and convex* maps, which is practically inconvenient (since, as opposed to Lipschitz maps, Lipschitz-convex functions are not stable through composition).

Let us thus stick for the moment to the example of $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$. The major advantage of being a concentrated random vector is that this concentration property is stable under any 1-Lipschitz map $f : \mathbb{R}^p \rightarrow \mathbb{R}^q$. So, if \mathbf{x} is concentrated, so is $\mathbf{x}' = f(\mathbf{x})$, which can be a vector with *intricate nonlinear dependence* between its entries (as we shall see right after, this intricate dependence may be such that photo-realistic images, able to deceive the human eye, can be generated from Lipschitz maps of white Gaussian vectors).

Now, the key reasons why the class of random vectors $\{f(\mathbf{x})\}$ spanned by 1-Lipschitz maps f is so fundamental to machine learning are that:

- (i) there exist machine learning techniques that *learn* to produce artificial but highly realistic data, exclusively based on Lipschitz maps. The most popular of these methods are the generative adversarial networks (GANs) initially proposed by Goodfellow et al. [2014]. Those are feedforward neural networks which, after training, generate highly realistic data $f(\mathbf{x})$ from a standard Gaussian vector input $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ (so realistic that even human beings cannot tell synthetic data from real ones). Since a feedforward neural network is a sequence of linear operators (inter-layer connections and convolution operators) and Lipschitz nonlinear activation functions (sigmoid, rectified linear, etc.), f is indeed Lipschitz (as the composition of Lipschitz operators remains Lipschitz);
- (ii) feature extraction procedures in machine learning are also mostly Lipschitz maps. The most popular of these today are convolutional neural networks, which are again feedforward neural nets and thus, by definition, Lipschitz maps of the input data. But this is also mostly true of all conventional methods, such as support vector machines, semi-supervised graph learning, spectral clustering, etc.

As a consequence of (i) and (ii), since real data can trust-worthily be approximated by outputs \mathbf{x}' of some Lipschitz function $\mathbf{x}' = f(\mathbf{x})$ of random Gaussian vectors \mathbf{x} , the class of concentrated random vectors encompasses a broad “set of realistic data”. Furthermore, in practice, the actual data features exploited in most machine learning algorithms can be seen as yet another Lipschitz mapping $g(\mathbf{x}')$ of the data \mathbf{x}' . Since \mathbf{x}' is well modeled under the form $\mathbf{x}' = f(\mathbf{x})$ for standard Gaussian \mathbf{x} , $\mathbf{x}'' = (g \circ f)(\mathbf{x})$ is again a Lipschitz map of a standard Gaussian vector.

It then becomes natural to model a wide range of realistic data as concentrated random vectors, e.g., as Lipschitz functions of standard Gaussian vectors.

Remark 2.15 (Concentration inequalities versus concentration of measure theory). *The concentration of measure theory developed by Ledoux provides as corollaries a list of popular concentration inequalities such as Gaussian concentration*

inequalities, Bernstein’s and Talagrand’s inequalities for random variables with bounded entries,⁴⁰ McDiarmid’s inequalities for functionals of bounded deviations of independent random variables, etc. These few results, quite popular in statistics, can however only marginally be used as a full-fledged concentration of measure-oriented random matrix framework. As already pointed out, quadratic forms are not naturally handled by these concentration inequalities. More importantly, while quadratic form concentration is essentially sufficient to prove the convergence of Stieltjes transforms, proving the resolvent convergence $\mathbb{E}[\mathbf{Q}] - \bar{\mathbf{Q}} \rightarrow 0$ under a concentration inequality setting actually demands to further expand the works of Ledoux, as will be shown next.

It must also be stressed that, in [Tao, 2012, Vershynin, 2012], Tao and Vershynin provide an introduction to what Vershynin refers to as non-asymptotic random matrix theory based on popular concentration inequalities. The approach followed by the authors however significantly differs from the present $n, p \rightarrow \infty$ with $p/n \rightarrow c$ large dimensional random matrix considerations. In non-asymptotic random matrix theory, the variables n, p are left “free” to grow at any relative speed to infinity and the use of concentration inequalities aims at retrieving bounds on largest or smallest eigenvalues or singular values of the underlying random matrices, without resorting to the Stieltjes transform approach. For Tao, this control step is the crux of the proof of the circular law (based on the ϵ -net theory developed by the author) for non-symmetric matrices \mathbf{X} with i.i.d. entries. For Vershynin, these non-asymptotic spectrum controls are exploited in applications to compressive sensing, where random matrix theory also plays a key role – for instance in providing “typical” matrices fulfilling the popular restricted isometry property [Candès, 2008].

The approach proposed in this monograph also provides a set of inequalities where n, p have an untied growth to infinity, but the application of these convergence results are mostly of interest in a joint growth rate for n, p . Besides, additional tools to Ledoux’s original framework, such as the notion of linear concentration, will be needed.

Remark 2.16 (Limitations of the concentration of measure framework). It is important to raise here (somewhat ironically) that the concentration of measure framework, which finds important corollaries to the field of compressive sensing [Donoho, 2006, Baraniuk, 2007], is, as presented here, at odds with the compressive sensing framework. Indeed, compressive sensing is a major field of research in large dimensional statistics and machine learning which assumes that large dimensional data are intrinsically of low dimension. That is, in the simplest linear setting, data vectors $\mathbf{x} \in \mathbb{R}^p$ can be written as $\mathbf{x} = \mathbf{A}\mathbf{y}$ for some matrix $\mathbf{A} \in \mathbb{R}^{p \times q}$ (generally unknown) and $\mathbf{y} \in \mathbb{R}^q$ for q very small. From there, the idea of compressive sensing is that meaningful statistical inference on \mathbf{y} can be performed based on few independent realizations $n \ll p$ (which is convenient if p is extremely large). There, concentration inequalities are mostly used to deal with the (usually random) observation matrix \mathbf{A} , rather than with the data.

⁴⁰To be more exact, Talagrand’s work was developed in parallel to Ledoux’s theory and are rather complementary than a consequence of one another.

In the present random matrix framework, concentration of measure is used to model the data, not the data operating matrices. These data however must not be of intrinsic low dimension $q \ll n$. Or, at least, if they were, we would impose in our framework that $n \sim q$ and $n, q, p \rightarrow \infty$ with a small but $O(1)$ ratio q/p . If instead $q = O(1) \ll n$, then we would fall back under the (difficult) sparse setting discussed at the end of Section 2.6.2.5 where our present framework is mostly ineffective.

*As shall be seen in the concrete applications presented in this monograph, large dimensional images are very appropriately modeled by concentrated random vectors of intrinsically large dimensions. However, feature vectors such as bag-of-words (also known as tf*idf features) for text classification, which are very large but extremely sparse vectors, cannot be handled by random matrix theory so far.*

This however does not mean that compressive sensing is complementary to random matrix theory. Compressive sensing indeed tackles the “difficult” problem analyzing sparse recovery algorithms by somehow “loose” inequalities and bounds: that is, it cannot accurately predict the exact performance of a given algorithm (however it can ensure its convergence and its efficiency as $n \rightarrow \infty$ at a certain rate with respect to p , while q is in general fixed). Random matrix theory instead requests that the intrinsic dimension $q \rightarrow \infty$, even slowly so, but manages in exchange (by exploiting the q degrees of freedom in the feature space) to provide accurate performance estimates of machine learning algorithms for all finite (but at least moderately large) n, q .

2.7.3 Elements of concentration of measure for random matrices

We recall here basic elements of the concentration of measure theory of immediate interest to random matrix applications. More advanced considerations can be found in [Ledoux, 2005] from a mathematical standpoint, and in [Louart and Couillet, 2018] with a more random matrix-oriented flavor.

2.7.3.1 Concentration of random variables

Before getting into generic multivariate concentration of measure, we need to start with the concept of concentration of a (univariate) random variable. Concentration of measure can be defined in two parallel ways.

Definition 6 (Concentration of a random variable). *Let $\alpha : \mathbb{R}^+ \rightarrow [0, 1]$ be a non-increasing function with $\alpha(\infty) = 0$. A random variable x is α -concentrated and we write $x \propto \alpha$ if, for an independent copy x' of x , and all $t > 0$,*

$$\mathbb{P}(|x - x'| > t) \leq \alpha(t).$$

The definition suggests that any two independent realizations of x cannot live far from one another. Alternatively, we may define x as concentrated if there exists a deterministic pivot a close to which x remains.

Definition 7 (Concentration around a pivot). *Let $\alpha : \mathbb{R}^+ \rightarrow [0, 1]$ be a non-increasing function and $a \in \mathbb{R}$. Then x is α -concentrated around the pivot a , denoted $x \in a \pm \alpha$, if for all $t > 0$,*

$$\mathbb{P}(|x - a| > t) \leq \alpha(t).$$

These two definitions are not formally equivalent. However, we have the implication

$$x \propto \alpha \Rightarrow x \in M_x \pm 2\alpha \Rightarrow x \propto 4\alpha(\cdot/2)$$

where $M_x \in \mathbb{R}$ is a *median* of x , i.e., is such that $\mathbb{P}(x \geq M_x) \geq \frac{1}{2}$ and $\mathbb{P}(x \leq M_x) \geq \frac{1}{2}$. The loss of a factor $1/2$ arises here from the bound $\mathbb{P}(|x - x'| > t) \leq \mathbb{P}(|x - a| > t/2) + \mathbb{P}(|x' - a| > t/2)$. Up to constants, it is then possible to use either definition interchangeably (the proofs of subsequent results are usually more accessible to one or the other definition).

A particularly appealing result is that 1-Lipschitz maps $f : \mathbb{R} \rightarrow \mathbb{R}$ of a concentrated random variable x maintain the concentration, i.e.,

$$x \propto \alpha \Rightarrow f(x) \propto \alpha. \quad (2.42)$$

This is a particularly fundamental result which suggests that every smooth map of sub-linear growth of x satisfies the same concentration property. This result naturally arises from the fact that $|f(x) - f(x')| \leq |x - x'|$ and thus $\mathbb{P}(|f(x) - f(x')| > t) \leq \mathbb{P}(|x - x'| > t)$.

Evidently, sums of concentrated random variables are concentrated:

$$\begin{aligned} x_1 \propto \alpha, x_2 \propto \beta &\Rightarrow x_1 + x_2 \propto \alpha(\cdot/2) + \beta(\cdot/2) \\ x_1 \in a \pm \alpha, x_2 \in b \pm \beta &\Rightarrow x_1 + x_2 \in a + b \pm [\alpha(\cdot/2) + \beta(\cdot/2)] \end{aligned}$$

where in the first line the factor $1/2$ unfolds from the bound $\mathbb{P}(|x_1 + x_2 - x'_1 - x'_2| > t) \leq \mathbb{P}(|x_1 - x'_1| > t/2) + \mathbb{P}(|x_2 - x'_2| > t/2)$, and similarly for the second line.

However, products, particularly of *dependent* random variables, are less obvious to tackle, as one needs to avoid conditioning. The problem can be worked around using the following two relations

$$\begin{aligned} x_1 x_2 - ab &= (x_1 - a)(x_2 - b) + a(x_2 - b) + b(x_1 - a) \\ |x_1 - a||x_2 - b| > t &\Rightarrow (|x_1 - a| > \sqrt{t}) \text{ or } (|x_2 - b| > \sqrt{t}) \end{aligned}$$

so to obtain

$$\begin{aligned} x_1 \in a \pm \alpha, x_2 \in b \pm \beta &\Rightarrow x_1 x_2 \in ab \pm \\ \begin{cases} \alpha(\sqrt{\cdot/3}) + \alpha(\cdot/3|b|) + \beta(\sqrt{\cdot/3}) + \beta(\cdot/3|a|), & a, b \neq 0 \\ \alpha(\sqrt{\cdot/2}) + \alpha(\cdot/2|b|) + \beta(\sqrt{\cdot/3}), & a = 0, b \neq 0 \\ \alpha(\sqrt{\cdot}) + \beta(\sqrt{\cdot}), & a = b = 0. \end{cases} \end{aligned}$$

For large t , the probability $\mathbb{P}(|x_1 x_2| > t)$ is here dominated by the terms $\alpha(\sqrt{\cdot})$ and $\beta(\sqrt{\cdot})$ which is not surprising. In the particular case where $x_1 = x_2 = x$, or more generally for powers x^k of concentrated random variables x , we have

$$x \in a \pm \alpha \Rightarrow x^k \in a^k \pm \left[\alpha(\cdot/2^k |a|^{k-1}) + \alpha((\cdot/2)^{\frac{1}{k}}) \right] \quad (2.43)$$

with $\alpha(\cdot/0) = \alpha(\infty)$ by convention, which is based on noticing that

$$|x^k - a^k| \leq (2|a|)^k \left(\frac{x-a}{a} + \frac{|x-a|^k}{|a|^k} \right).$$

This result will be particularly useful for random matrix applications to quadratic forms.

Remark 2.17 (Exponential concentration). *Of utmost interest is the case where $\alpha(t) = Ce^{-(t/\sigma)^q}$ for some $C, \sigma, q > 0$. In particular, it is known that standard random Gaussian variables x satisfy*

$$x \sim \mathcal{N}(0, 1) \Rightarrow x \in 0 \pm 2e^{-(\cdot)^2/2}.$$

Exponential concentrations are fast and induce a lot of convenient properties. In particular, using the formula $\mathbb{E}[|x|^k] = \int_0^\infty \mathbb{P}(|x|^k > t) dt$, it appears that all (absolute) moments of exponentially concentrated random variables exist. In particular,

$$x \propto Ce^{-(\cdot/\sigma)^q} \Rightarrow x \in \mathbb{E}[x] \pm e^{\frac{Cq}{q}} e^{-(\cdot/2\sigma)^q} \quad (2.44)$$

so that an exponentially concentrated random variable concentrates around its mean. But most importantly, we have the implications

$$\begin{aligned} x \in a \pm Ce^{-(\cdot/\sigma)^q} &\Rightarrow \forall r \geq q, \mathbb{E}[|x-a|^r] \leq C\Gamma(r/q+1)\sigma^r \\ &\Rightarrow x \in a \pm Ce^{-(\cdot/\sigma)^q/e} \end{aligned}$$

with Γ the gamma-distribution. Thus, exponential concentration is “equivalent” to controlled growth by σ^r of all moments $r \geq q$. This is particularly appealing when moments occasionally turn out more convenient to deal with than bounds on tail probabilities.

2.7.3.2 Concentration of random vectors

The concept of concentration of random variables, stating that x does not deviate much from a given pivot a , cannot be straightforwardly extended to that of random vectors. Indeed random vectors (in particular large dimensional ones) rather tend to “avoid” their statistical means or medians: e.g., Gaussian random vectors $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ are of zero mean but they “concentrate” on a $O(1)$ -thick layer around the sphere of \mathbb{R}^p of diameter \sqrt{p} .

Instead, for a normed vector space $(E, \|\cdot\|)$, we will consider that a random vector $\mathbf{x} \in E$ is concentrated *for some class of functions* $\mathcal{F} : \mathbb{R}^p \rightarrow \mathbb{R}$

if, for all $f \in \mathcal{F}$, $f(\mathbf{x})$ is a concentrated random *variable*. Depending on the “broadness” of the class, being a concentrated random vector can be more demanding. [Ledoux \[2005\]](#) originally defined two such classes \mathcal{F} : the class of 1-Lipschitz maps (appropriate for Gaussian or random unitary vectors) and the class of convex (or weakly convex) 1-Lipschitz maps (adapted to vectors of independent bounded entries). There, the Lipschitz definition (i.e., the fact that $|f(\mathbf{x}) - f(\mathbf{y})| \leq \|\mathbf{x} - \mathbf{y}\|$) is with respect to the norm $\|\cdot\|$ in E , and thus the concentration rates may depend on $\|\cdot\|$. In order to better encompass random matrices in the concentration inequality framework, a looser additional class \mathcal{F} will be introduced here: that of unit-norm linear functionals.

Linear concentration Linear concentration is an important concept in random matrix theory as it provides a quite general and flexible definition for the notion of *deterministic equivalents*.

Definition 8 (Linear concentration). *A random vector $\mathbf{x} \in E$ is linearly α -concentrated around the deterministic equivalent $\bar{\mathbf{x}}$, with respect to the norm $\|\cdot\|$ in E , if, for all unit norm linear functional $u : E \rightarrow \mathbb{R}$ (i.e., $|u(\mathbf{x})| \leq \|\mathbf{x}\|$),*

$$u(\mathbf{x}) \in u(\bar{\mathbf{x}}) \pm \alpha.$$

The expectation being a linear operator (from E to E), an advantage of linear concentration is that, upon existence, $\mathbb{E}[\mathbf{x}]$ is a deterministic equivalent for the concentrated random vector \mathbf{x} . In particular, if \mathbf{Q} is a random matrix (e.g., the resolvent of some other underlying random matrix) in the “vector space” $(\mathbb{R}^{p \times p}, \|\cdot\|)$, with $\|\cdot\|$ the operator norm, and that \mathbf{Q} is linearly concentrated with respect to $\|\cdot\|$, then $\mathbb{E}\mathbf{Q}$ is a deterministic equivalent for \mathbf{Q} and we have in particular, for all $\mathbf{A} \in \mathbb{R}^{p \times p}$ and $\mathbf{a}, \mathbf{b} \in \mathbb{R}^p$ of bounded norms,

$$\frac{1}{p} \operatorname{tr} \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) \rightarrow 0, \quad \mathbf{a}^\top (\mathbf{Q} - \mathbb{E}\mathbf{Q}) \mathbf{b} \rightarrow 0, \quad \|\mathbf{Q} - \mathbb{E}\mathbf{Q}\| \rightarrow 0$$

where the convergence is in probability and, if $\alpha(t) = Ce^{-t^q}$ for some $q > 0$, the convergence is also almost sure.⁴¹ This result implies that the newly defined notion of *deterministic equivalents* from a linear concentration standpoint automatically induces the former Definition 4.

Lipschitz concentration Lipschitz concentration is the most popular type of concentrations (due to its compatibility with (2.42)). This notion is even in general merely called “concentration” (rather than Lipschitz concentration); it is defined as follows.

Definition 9 (Lipschitz concentration). *A random vector $\mathbf{x} \in E$ is Lipschitz α -concentrated with respect to the norm $\|\cdot\|$ if, for every 1-Lipschitz function*

⁴¹Here we exploit the fact that, for $u(\mathbf{Z}) = \frac{1}{p} \operatorname{tr} \mathbf{A}\mathbf{Z}$, $|u(\mathbf{Z})| \leq \|\mathbf{Z}\|$ when $\|\mathbf{A}\| \leq 1$ and that, for $u(\mathbf{Z}) = \mathbf{a}^\top \mathbf{Z} \mathbf{b} = \operatorname{tr}(\mathbf{b}\mathbf{a}^\top \mathbf{Z})$, $|u(\mathbf{Z})| \leq \|\mathbf{Z}\|$ for $\|\mathbf{a}\|, \|\mathbf{b}\| \leq 1$.

$f : E \rightarrow \mathbb{R}$, we have either of the conditions

$$\begin{aligned} f(\mathbf{x}) &\propto \alpha, \text{ denoted } \mathbf{x} \propto \alpha \\ f(\mathbf{x}) &\in M_f \pm \alpha, \text{ denoted } \mathbf{x} \stackrel{M}{\propto} \alpha \\ f(\mathbf{x}) &\in \mathbb{E}[f(\mathbf{x})] \pm \alpha, \text{ denoted } \mathbf{x} \stackrel{\mathbb{E}}{\propto} \alpha \end{aligned}$$

hold, where M_f is a median of $f(\mathbf{x})$.

Similar to the concentration of random variables, the three notions are not fully equivalent. For generic α -concentration,

$$\mathbf{x} \propto \alpha \Rightarrow \mathbf{x} \stackrel{\mathbb{E}}{\propto} \alpha \Rightarrow \mathbf{x} \propto 4\alpha(\cdot/2)$$

and, in the case of exponential concentrations, the expectation is well defined and we further have

$$\mathbf{x} \stackrel{\mathbb{E}}{\propto} C e^{-(\cdot/\sigma)^q} \Rightarrow \mathbf{x} \stackrel{M}{\propto} e^{C^q/q} e^{-(\cdot/2\sigma)^q} \Rightarrow \mathbf{x} \stackrel{\mathbb{E}}{\propto} e^{C^q/q} e^{-(\cdot/4\sigma)^q}.$$

The most fundamental result at the very heart of the concentration of measure theory is that Gaussian random vectors $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ are Lipschitz concentrated in $(\mathbb{R}^p, \|\cdot\|)$ for $\|\cdot\|$ the Euclidean norm, that is

$$\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p) \Rightarrow \mathbf{x} \stackrel{M}{\propto} 2e^{-(\cdot/2)^2} \text{ and } \mathbf{x} \stackrel{\mathbb{E}}{\propto} 2e^{-(\cdot/2)^2}.$$

A fundamental fact about the above concentration is that it does *not* depend on the size p of the ambient space (neither in the tail nor in the head parameters). As such, arbitrarily large standard Gaussian vectors (and thus concatenation of n such vectors, as well as matrices $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ built from independent standard Gaussian vectors \mathbf{x}_i endowed with the Frobenius norm) also concentrate with no dependence on p, n .

This is in fact far from natural as, even for independent vectors $\mathbf{x}_1, \dots, \mathbf{x}_n$, all of which being concentrated, the joint concentration of $(\mathbf{x}_1, \dots, \mathbf{x}_n)$ with respect to the Euclidean norm in the product space generally comes along with a loss of concentration rate proportional to n . Besides, if the vectors \mathbf{x}_1 and \mathbf{x}_2 are both concentrated but not independent, the vector $(\mathbf{x}_1, \mathbf{x}_2)$ may not even be concentrated.

Remark 2.18 (On the location of Gaussian vectors). *To clearly understand the relation between standard Gaussian $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ and its dimension, note that, $\|\mathbf{x}\|$ having a chi-distribution with median $\sqrt{p} + O(1/\sqrt{p})$, its exponential concentration precisely implies*

$$\mathbb{P}(|\|\mathbf{x}\| - \sqrt{p}| > t) \leq 2e^{-(t+O(1/p))^2/4}.$$

Thus, \mathbf{x} is a random variable that essentially lives close to a sphere of radius $O(\sqrt{p})$ and of thickness $O(1)$ or, equivalently, \mathbf{x}/\sqrt{p} is a random vector distributed close to the unit sphere in \mathbb{R}^p with actual distance to the sphere vanishing as $O(1/\sqrt{p})$. The vector \mathbf{x} is thus nowhere near its expected value $\mathbf{0}$.

This remark is fundamental as it disrupts with the finite-dimensional case where \mathbf{x} lives close to its mean. In 1D to 3D, one indeed visualizes (independent) Gaussian random vectors as densely “concentrated” around their mean (close to the center of the bell-shaped distribution). The intuitive extension of this visualization to larger dimensions would be erroneous.

As for concentrated random variables, Lipschitz concentrated random vectors are stable through Lipschitz mapping in the sense that, for all 1-Lipschitz $\phi : E \rightarrow E'$ with respect to norms $\|\cdot\|_E$ and $\|\cdot\|_{E'}$,

$$\mathbf{x} \stackrel{M}{\propto} \alpha \Rightarrow \phi(\mathbf{x}) \stackrel{M}{\propto} \alpha. \quad (2.45)$$

Convex (Lipschitz) concentration. To define convex concentration, we need to recall the notion of quasi-convex functions: $f : E \rightarrow \mathbb{R}$ is quasi-convex if, for all $t \in \mathbb{R}$, the sets $\{\mathbf{x} \in E \mid f(\mathbf{x}) \leq t\}$ are convex sets, i.e., for all $t \in [0, 1]$ and $\mathbf{x}, \mathbf{y} \in E$, $f(t\mathbf{x} + (1-t)\mathbf{y}) \leq \max\{f(\mathbf{x}), f(\mathbf{y})\}$. In particular, convex functions are quasi-convex (thus the notion generalizes convexity) and, for $E = \mathbb{R}$, all monotonous functions (even concave ones) are quasi-convex.

Then we have the following definition of convex concentration.

Definition 10 (Convex concentration). *A vector $\mathbf{x} \in E$ is (Lipschitz) convexly concentrated for the norm $\|\cdot\|$ if, for any 1-Lipschitz and quasi-convex function $f : \mathbb{E} \rightarrow \mathbb{R}$, we have either of the conditions*

$$\begin{aligned} f(\mathbf{x}) &\propto \alpha, \text{ denoted } \mathbf{x} \propto_c \alpha \\ f(\mathbf{x}) &\in M_f \pm \alpha, \text{ denoted } \mathbf{x} \stackrel{M}{\propto}_c \alpha \\ f(\mathbf{x}) &\in \mathbb{E}[f(\mathbf{x})] \pm \alpha, \text{ denoted } \mathbf{x} \stackrel{\mathbb{E}}{\propto}_c \alpha \end{aligned}$$

where M_f is a median of $f(\mathbf{x})$.

Obviously, all Lipschitz convex functions being Lipschitz, Lipschitz concentration implies convex concentration (which itself implies the even less demanding linear concentration); for instance, in the case of exponential concentration,

$$\mathbf{x} \stackrel{\mathbb{E}}{\propto} C e^{-(\cdot/\sigma)^q} \Rightarrow \mathbf{x} \stackrel{\mathbb{E}}{\propto}_c C e^{-(\cdot/\sigma)^q} \Rightarrow \mathbf{x} \in \mathbb{E}[\mathbf{x}] \pm e^{-(\cdot/\sigma)^q}.$$

The interest for convex concentration is related to the following result due to Talagrand [Talagrand, 1995, Theorem 4.1.1]: let $\mathbf{x} \in [0, 1]^p$ be a vector of independent entries, then

$$\mathbf{x} \stackrel{M}{\propto}_c 4e^{-\cdot^2/4}.$$

However, convex concentration has the major limitation that quasi-convex functions are *not* stable by composition. This prevents the simple adaptation of numerous results obtained for Lipschitz (or linear) concentration. Yet, for f quasi-convex and g affine, $f \circ g$ is still quasi-convex.

Nonetheless, the results necessary to our present random matrix analysis of sample covariance matrix models can fortunately be extended.

Convex concentration transversally to a group action. A last convenient notion of concentration, dedicated to random matrix theory, consists in transferring concentration from \mathbf{X} to the vector of its singular values. This will help transfer concentration from the data to linear statistics of the eigenvalues of the sample covariance matrix. To this end though, convex concentration is too demanding and we need to further restrict the space of functions as follows.

Definition 11 (Convex concentration transversally to group action). *Let $\mathbf{x} \in E$ and G a group acting on E . Then, \mathbf{x} is convexly α -concentrated transversally to the action of G if, for all quasi-convex 1-Lipschitz and G -invariant function f (i.e., $f(g \cdot \mathbf{x}) = f(\mathbf{x})$ for $g \in G$), $f(\mathbf{x}) \propto \alpha$. This is denoted $\mathbf{x} \propto_G^T \alpha$.*

In particular, denote $\sigma(\mathbf{X}) = (\sigma_1(\mathbf{X}), \dots, \sigma_{\min\{p,n\}}(\mathbf{X}))$ the vector of the singular values of $\mathbf{X} \in \mathbb{R}^{p \times n}$ (i.e., $\sigma_i(\mathbf{X}) = \sqrt{\lambda_i(\mathbf{X}\mathbf{X}^\top)}$ for $i \leq \min\{p, n\}$), and define the group $\mathcal{O}_{p,n} = \{(\mathbf{U}, \mathbf{V}) \in \mathbb{R}^{p \times p} \times \mathbb{R}^{n \times n} \text{ orthonormal}\}$ acting on $\mathbb{R}^{p \times n}$ by $(\mathbf{U}, \mathbf{V}) \cdot \mathbf{M} = \mathbf{U}\mathbf{M}\mathbf{V}^\top$ and the group \mathcal{S}_p of permutations of size p acting on \mathbb{R}^p by $\tau \cdot \mathbf{y} = (\mathbf{y}_{\tau(1)}, \dots, \mathbf{y}_{\tau(p)})$. Then, we have the following result, inspired by [Davis, 1957],

$$\mathbf{X} \propto_{\mathcal{O}_{p,n}}^T \alpha \Leftrightarrow \sigma(\mathbf{X}) \propto_{\mathcal{S}_{\min\{p,n\}}}^T \alpha. \quad (2.46)$$

2.7.4 A concentration inequality version of Theorem 2.6

Equipped with these elementary results, we can now provide an extension of Theorem 2.6 to the case of concentrated data vectors.

Before getting to the main result, we introduce some preliminary lemmas, which generalize classical random matrix results to the concentration of measure framework. Most of these results and their corresponding proofs can be found in [Louart and Couillet, 2018].

2.7.4.1 Trace lemma

A first result of importance concerns the extension of the “quadratic-form-close-to-the-trace” lemma, Lemma 2.11, from a moment-based version to a concentration of measure setting. The result consists in a generalization of a popular result in concentration of measure theory known as Hanson-Wright’s theorem (see e.g., [Vershynin, 2018, Theorem 6.2.1]).

Lemma 2.21 (Trace lemma for concentrated vectors). *Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ and $\mathbf{x} \in \mathbb{R}^p$ such that $\mathbf{x} \stackrel{\mathbb{E}}{\propto_c} C e^{-(\cdot/\sigma)^q}$. Then,*

$$\mathbf{x}^\top \mathbf{A} \mathbf{x} \in \text{tr}(\mathbb{E}[\mathbf{x} \mathbf{x}^\top] \mathbf{A}) \pm 2C \left(e^{-(\cdot/4\sigma\|\mathbf{A}\|)\mathbb{E}[\|\mathbf{x}\|]^q} + e^{-(\cdot/2\|\mathbf{A}\|\sigma^2)^{\frac{q}{2}}} \right).$$

This lemma follows almost automatically from two elementary ingredients of the concentration of measure theory: (i) assuming first that \mathbf{A} is nonnegative definite, $\mathbf{x}^\top \mathbf{A} \mathbf{x} = \|\mathbf{A}^{\frac{1}{2}} \mathbf{x}\|^2$ with $\|\mathbf{A}^{\frac{1}{2}} \mathbf{x}\|$ a concentrated random variable (it is a

Lipschitz and convex function of \mathbf{x}) which, (ii) from the concentration of powers of concentrated random variables (2.43) for $k = 2$, gives the result. For generic \mathbf{A} , it suffices to write \mathbf{A} as the sum of its symmetric nonnegative and symmetric negative parts (which explains the leading factor 2).

This lemma particularly stresses the technical convenience of the concentration of measure framework. The key random matrix results, such as Lemma 2.11 for vectors of i.i.d. entries, often rely on dedicated tools and possibly heavy (combinatorial) proof techniques. Here, the concentration of measure alternative to Lemma 2.11 follows from a mere two-line argument (once the elementary tools of the theory are in place). Besides the exponential rate of convergence is very versatile and particularly ensures the uniform convergence of $\{\mathbf{x}_i^\top \mathbf{A} \mathbf{x}_i, i = 1, \dots, n\}$, for n any polynomial in p ; using the moment method would demand to systematically compute high order moments of $\mathbf{x}_i^\top \mathbf{A} \mathbf{x}_i$ to obtain uniform convergence over large n (using Markov's inequality).

2.7.4.2 Concentration of the Stieltjes transform

Next, we generalize the convergence of Stieltjes transforms in a generic concentration of measure form.

Lemma 2.22 (Trace of Resolvent). *For $\mathbf{X} \in \mathbb{R}^{p \times n}$ equipped with the Frobenius norm, and $\mathbf{Q}(z) = (\frac{1}{n} \mathbf{X} \mathbf{X}^\top - z \mathbf{I}_p)^{-1}$ for $z < 0$,*

$$\mathbf{X} \propto_c \alpha \text{ in } (\mathbb{R}^{p \times n}, \|\cdot\|_F) \Rightarrow \text{tr } \mathbf{Q}(z) \propto 2\alpha \left(\frac{\sqrt{n|z|^3}}{8 \min\{p, n\}} \right).$$

To prove this lemma, first recall that $\mathbf{X} \propto_c \alpha \Rightarrow \sigma(\mathbf{X}) \propto_{\mathcal{S}_d}^T \alpha$ with $d = \min\{p, n\}$. Also, $\text{tr } \mathbf{Q}(z) = \sum_{i=1}^d f(\sigma_i(\mathbf{X}))$ for $f : \mathbb{R}_+ \rightarrow \mathbb{R}$, $s \mapsto 1/(s^2 - z)$. This function f is $(2|z|^{-\frac{2}{3}})$ -Lipschitz (checked by bounding its derivative) and the mapping $(s_1, \dots, s_d) \mapsto \sum_{i=1}^d s_i$ is evidently \mathcal{S}_d -invariant. However, f is not quasi-convex but can be written as the sum $f = g - h$ of two quasi-convex $4|z|^{-\frac{2}{3}}$ -Lipschitz functions $(h(s) = (s/|z| - 1/\sqrt{|z|})^2 \mathbf{1}_{\{s \in [0, \sqrt{|z|}]\}})$ and $g = f + h$. Consequently, since $\mathbf{X} \propto_c \alpha \Rightarrow \mathbf{X} \propto_{\mathcal{O}_{p,n}}^T \alpha$, we have from (2.46) both the concentration of $\sum_i g(\sigma_i(\mathbf{X}))$ and of $\sum_i h(\sigma_i(\mathbf{X}))$, and it then remains to apply the result on the concentration of the sum of two concentrated random variables to obtain the result.

Again here, the proof is elegant and immediate although the mapping $\mathbf{X} \mapsto \text{tr } \mathbf{Q}(z)$ is highly non-trivial from a statistical standpoint. Note in particular that the technical difficulty raised by the non-convexity of f would not have been a problem if we had rather assumed Lipschitz concentration $\mathbf{X} \propto \alpha$ for \mathbf{X} (which we recall is more demanding for \mathbf{X} and would exclude the case of \mathbf{X} with bounded i.i.d. entries).

2.7.4.3 Concentration of the resolvent \mathbf{Q} and deterministic equivalents

The approach followed in the previous lemma uses the convenient expression of $f : \mathbb{R}^{p \times n} \rightarrow \mathbb{R}$ as $f = g - h$ for two convex and Lipschitz functions g and h . It does not seem that the mapping $f(\mathbf{X}) = \mathbf{Q}_{\frac{1}{n}\mathbf{XX}^T}(z)$ from $\mathbb{R}^{n \times p}$ to $\mathbb{R}^{p \times p}$ can be treated similarly, as no such Lipschitz function division can be exploited. One must therefore resort to the additional strength of exponential concentration to divide the space $\mathbb{R}^{p \times n}$ into a compact space for the operator norm $\{\mathbf{X} \mid \|\mathbf{X}\| \leq K\sqrt{n}\}$ where f will be shown to be automatically Lipschitz (as its image is bounded) and the complement space $\{\mathbf{X} \mid \|\mathbf{X}\| > K\sqrt{n}\}$ which is of vanishing probability for all large $K > 0$.

Regrouping these two results, we have the following concentration for the resolvent.

Lemma 2.23 (Concentration of $\mathbf{Q}_{\frac{1}{n}\mathbf{XX}^T}$). *For $\mathbf{X} \in \mathbb{R}^{p \times n}$ and $z < 0$, let $\mathbf{Q}(z) = (\frac{1}{n}\mathbf{XX}^T - z\mathbf{I}_p)^{-1}$. Then we have the following two results*

$$\begin{aligned} \mathbf{X} \propto \alpha &\Rightarrow \mathbf{Q}(z) \propto \alpha \left(\sqrt{n|z|^3} \cdot /2 \right) \\ \mathbf{X} \stackrel{\mathbb{E}}{\propto} C e^{-\cdot)^q} &\Rightarrow \mathbf{Q}(z) \in \mathbb{E}\mathbf{Q}(z) \pm 2C e^{-\left(\sqrt{|z|^3 n} \cdot /4\sigma \right)^q} \end{aligned}$$

where the left-hand side concentrations are understood in $(\mathbb{R}^{p \times n}, \|\cdot\|_F)$ and the right-hand side in $(\mathbb{R}^{p \times p}, \|\cdot\|_F)$.

This result is in fact quite powerful and automatically induces (and vastly generalizes) the notion of deterministic equivalent of Definition 4, i.e., it implies that $\frac{1}{n} \text{tr } \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) \xrightarrow{a.s.} 0$ and $\mathbf{a}^T(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b} \xrightarrow{a.s.} 0$ for all $\mathbf{A}, \mathbf{a}, \mathbf{b}$ of unit norm. Indeed, first recall that the first statement (of Lipschitz concentration) implies that

$$\mathbf{X} \stackrel{\mathbb{E}}{\propto} \alpha \Rightarrow \mathbf{Q}(z) \in \mathbb{E}\mathbf{Q}(z) \pm \alpha \left(\sqrt{n|z|^3} \cdot /2 \right)$$

(since Lipschitz concentration around the mean implies linear concentration around the mean). Next, note that the linear concentrations of \mathbf{Q} (under either Lipschitz or convex-Lipschitz-exponential concentration for \mathbf{X}) hold here with respect to the Frobenius norm of $\mathbf{X} \in \mathbb{R}^{p \times n}$. That is, for $\mathbf{A} \in \mathbb{R}^{p \times p}$ of bounded Frobenius (rather than only spectral) norm,⁴²

$$\text{tr } \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) = O(n^{-\frac{1}{2}}).$$

In particular, letting $p/n \rightarrow c > 0$, from $\|\mathbf{A}\|_F \leq \sqrt{\text{rank}(\mathbf{A})}\|\mathbf{A}\|$ (with $\|\cdot\|$ the operator norm) and $\|\mathbf{A}\| \leq \|\mathbf{A}\|_F$, we have (i) if $\mathbf{A} = \mathbf{ab}^T$ is of unit rank, $\text{tr } \mathbf{ab}^T(\mathbf{Q} - \mathbb{E}\mathbf{Q}) = \mathbf{a}^T(\mathbf{Q} - \mathbb{E}\mathbf{Q})\mathbf{b} = O(n^{-1/2})$, while (ii) if \mathbf{A} is of arbitrary rank

⁴²One must be careful not to confuse the steps of the proof which, as shown next, use a smart division of $\mathbb{R}^{p \times n}$ into bounded and unbounded operator norm $\|\mathbf{X}\|$, and the fact that the ultimate concentration results hold with respect to the Frobenius norm.

(say $\text{rank}(\mathbf{A}) = p$), $p^{-\frac{1}{2}} \text{tr } \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) = O(n^{-1/2})$ so that $\frac{1}{p} \text{tr } \mathbf{A}(\mathbf{Q} - \mathbb{E}\mathbf{Q}) = O(n^{-1})$.

Of course, since $\|\cdot\| \leq \|\cdot\|_F$ in $\mathbb{R}^{p \times p}$, Lemma 2.23 applies to \mathbf{Q} in $(\mathbb{R}^{p \times p}, \|\cdot\|)$ as well.

The proof of the first part of the lemma is again extremely simple, once the basic concentration of measure arguments are in place. Here we simply use the fact that the mapping $f : \mathbb{R}^{p \times n} \rightarrow \mathbb{R}^{p \times p}$, $\mathbf{X} \mapsto \mathbf{Q}(z)$ is $(2/\sqrt{|z|^3 n})$ -Lipschitz. Indeed, by the resolvent identity, Lemma 2.1,

$$f(\mathbf{X} + \mathbf{H}) - f(\mathbf{X}) = \frac{1}{n} f(\mathbf{X} + \mathbf{H})((\mathbf{X} + \mathbf{H})\mathbf{H}^\top + \mathbf{H}\mathbf{X}^\top)f(\mathbf{X})$$

so that, from $\|f(\mathbf{X})\mathbf{X}\| \leq \sqrt{n/|z|}$, $\|f(\mathbf{X})\| \leq 1/|z|$ and $\|\mathbf{AB}\|_F \leq \|\mathbf{A}\|\|\mathbf{B}\|_F$ (where $\|\cdot\|$ is the operator norm), we have $\|f(\mathbf{X} + \mathbf{H}) - f(\mathbf{X})\|_F \leq 2\|\mathbf{H}\|_F/\sqrt{|z|^3 n}$ and thus the result.

The proof of the second part is less immediate. Since the result is a linear concentration of the resolvent, one needs to control the concentration of the random variable $\text{tr } \mathbf{AQ}$ obtained for arbitrary $\mathbf{A} \in \mathbb{R}^{p \times p}$ with $\|\mathbf{A}\|_F \leq 1$. This is obtained by considering the mapping $f : \mathbf{X} \mapsto \text{tr } \mathbf{AQ}$, with the major difference from Lemma 2.22 that $f(\mathbf{Q})$ is now not a mere combination of the singular values of \mathbf{Q} . The function f is not convex (as already discussed in Lemma 2.22) but can be divided as $f = h - g$ with $g : \mathbf{X} \mapsto \frac{1}{n|z|^2} \text{tr } \mathbf{XX}^\top$ and $h = f + g$ both convex, with h Lipschitz and g Lipschitz on the bounded region $\{\mathbf{X} \mid \|\mathbf{X}\| \leq K\sqrt{n}\}$. Using a truncation method by considering $(\mathbf{X}^K)_{ij} = \mathbf{X}_{ij} 1_{\{\mathbf{X}_{ij} < K\sqrt{n}\}}$ for growing K , one obtains that the sequence of concentrated random variables $\text{tr } \mathbf{AQ}^K = \text{tr } \mathbf{A}(\frac{1}{n} \mathbf{X}^K (\mathbf{X}^K)^\top - z \mathbf{I}_p)^{-1}$ converges in law to $\text{tr } \mathbf{AQ}$, which can then be shown to imply that $\text{tr } \mathbf{AQ}$ is also a concentrated random variable.

2.7.4.4 Main result

Let us rephrase the setting of Theorem 2.6 by letting $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ be n i.i.d. random vectors with law \mathcal{L} such that

$$\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \propto C e^{-(\cdot)^q/c}$$

for some $C, c, q > 0$ with respect to the Frobenius norm (which implies in particular, by the action of the 1-Lipschitz mapping $f : (\mathbf{x}_1, \dots, \mathbf{x}_n) \mapsto \mathbf{x}_i$, that each \mathbf{x}_i is concentrated). This request of joint rather than individual vector concentration may be considered demanding, but is at least satisfied by (i) $\mathbf{x}_i = \phi(\mathbf{y}_i)$ with 1-Lipschitz maps $\phi : \mathbb{R}^{p'} \rightarrow \mathbb{R}^p$ for (i-a) $\mathbf{y}_i \sim \mathcal{N}(0, \mathbf{I}_{p'})$ or (i-b) \mathbf{y}_i uniformly distributed on the $\sqrt{p'}$ -radius sphere of $\mathbb{R}^{p'}$, or (ii) for \mathbf{x}_i composed of (an affine mapping of) i.i.d. entries with support in $[-1, 1]$.

With the above results, and some specific technical arguments, we have the following *concentration of measure version* of Theorem 2.6.

Theorem 2.18 (Sample covariance of concentrated random vectors). *Let $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \propto C e^{-(\cdot)^q/c}$ with $\mathbf{x}_i \in \mathbb{R}^p$, and $z < 0$. Further assume that $\frac{1}{\sqrt{p}} \mathbb{E} \|\mathbf{x}_i\|$ (or, if $q \geq 2$, simply $\frac{1}{\sqrt{p}} \|\mathbb{E}[\mathbf{x}_i]\|$), $\frac{1}{p} \text{tr } \Phi$ with $\Phi = \mathbb{E}[\frac{1}{n} \mathbf{X} \mathbf{X}^\top]$, as well as p/n are all bounded. Then, for all large n ,*

$$\mathbf{Q}(z) \in \bar{\mathbf{Q}}(z) \pm C' e^{-(\sqrt{n} \cdot)^q/c'} \text{ in } (\mathbb{R}^{p \times n}, \|\cdot\|)$$

for some $C', c' > 0$, where

$$\bar{\mathbf{Q}}(z) = \left(\frac{\Phi}{1 + \delta(z)} - z \mathbf{I}_p \right)^{-1}$$

and $\delta(z)$ is the unique positive solution to $\delta(z) = \frac{1}{n} \text{tr } \Phi \bar{\mathbf{Q}}(z)$.

Remark 2.19 (On real $z < 0$). *It must be noted here that the concentration framework devised in this section is only valid for real-valued matrices and thus Theorem 2.18 holds here for $z < 0$ only. Using additional arguments (of complex analytic extension of $\mathbf{Q}(z)$ and $\bar{\mathbf{Q}}(z)$), Theorem 2.18 naturally extends to all $z \in \mathbb{C} \setminus \mathbb{R}^+$.*

Denoting $\delta(z) = -1 - \frac{1}{z \tilde{m}_p(z)}$ and $\Phi = \mathbf{C}$, it comes immediately that the deterministic equivalent $\bar{\mathbf{Q}}$ in Theorem 2.18 above has the same ‘‘formal statement’’ as in Theorem 2.6; we shall see that using $\delta(z)$ rather than $\tilde{m}_p(z)$ is more convenient under the concentration of measure framework. Yet, there are a few key differences to raise between both theorems. First, $\Phi = \mathbb{E}[\frac{1}{n} \mathbf{X} \mathbf{X}^\top]$ is not a covariance matrix as the present concentration of measure on \mathbf{X} does not impose that $\mathbb{E}[\mathbf{X}] = \mathbf{0}$. Also, the deterministic equivalent $\bar{\mathbf{Q}}(z)$ comes along with a convergence speed and an exponential tail, which are both more practical than a mere almost sure convergence of specific statistics.

Theorem 2.18 unfolds from the same idea introduced in the proof of the Marčenko-Pastur law (Theorem 2.4), by successively introducing two deterministic equivalents. We provide here the basic arguments of the proof. We already know from Lemma 2.23 that $\mathbf{Q}(z) \in \mathbb{E}\mathbf{Q}(z) \pm C e^{-c(\sqrt{n} \cdot)^q}$ for some $C, c > 0$ and it only remains to show that $\|\mathbb{E}\mathbf{Q}(z) - \bar{\mathbf{Q}}(z)\|$ is small.

To this end, we introduce the first deterministic equivalent

$$\bar{\mathbf{Q}}(z) = \left(\frac{\Phi}{1 + \delta'(z)} - z \mathbf{I}_p \right)^{-1}$$

where $\delta'(z) = \frac{1}{n} \mathbb{E}[\mathbf{x}^\top \mathbf{Q}_-(z) \mathbf{x}] = \frac{1}{n} \text{tr } \Phi \mathbb{E}\mathbf{Q}_-$ for $\mathbf{Q}_- \in \mathbb{R}^{p \times p}$ the resolvent of $\frac{1}{n} \mathbf{X} \mathbf{X}^\top - \frac{1}{n} \mathbf{x} \mathbf{x}^\top$ and \mathbf{x} any column of \mathbf{X} . Applying the same ideas as in the proof of Theorem 2.4, we obtain (we discard the z ’s for readability)

$$\begin{aligned} \mathbb{E}\mathbf{Q} - \bar{\mathbf{Q}} &= \mathbb{E} \left[\mathbf{Q} \left(\frac{\Phi}{1 + \delta'} - \frac{1}{n} \mathbf{X} \mathbf{X}^\top \right) \right] \bar{\mathbf{Q}} \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\mathbf{Q} \left(\frac{\Phi}{1 + \delta'} - \mathbf{x}_i \mathbf{x}_i^\top \right) \right] \bar{\mathbf{Q}} = \mathbb{E} \left[\mathbf{Q} \left(\frac{\Phi}{1 + \delta'} - \mathbf{x} \mathbf{x}^\top \right) \right] \bar{\mathbf{Q}} \end{aligned}$$

which, along with $\mathbf{Q} = \mathbf{Q}_- - \frac{1}{n} \frac{\mathbf{Q}_{-\mathbf{x}\mathbf{x}^\top} \mathbf{Q}_-}{1 + \frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}}}$ and $\mathbf{Q}\mathbf{x} = \frac{\mathbf{Q}_{-\mathbf{x}}}{1 + \frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}}}$, gives

$$\begin{aligned}\mathbb{E}\mathbf{Q} - \bar{\mathbf{Q}} &= \mathbb{E}[\mathbf{E}_1] - \mathbb{E}[\mathbf{E}_2] \\ \mathbf{E}_1 &= \mathbf{Q}_- \left(\frac{\Phi}{1 + \delta'} - \frac{\mathbf{x}\mathbf{x}^\top}{1 + \frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}}} \right) \bar{\mathbf{Q}} \\ \mathbf{E}_2 &= \frac{1}{n(1 + \delta')} \mathbf{Q}_{-\mathbf{x}\mathbf{x}^\top} \mathbf{Q} \Phi \bar{\mathbf{Q}}.\end{aligned}$$

To bound $\|\mathbb{E}\mathbf{Q} - \bar{\mathbf{Q}}\|$ it suffices to bound $|\mathbf{a}^\top (\mathbb{E}\mathbf{Q} - \bar{\mathbf{Q}})\mathbf{a}|$ for any unit norm \mathbf{a} . From Hölder's inequality

$$\begin{aligned}|\mathbf{a}^\top \mathbb{E}[\mathbf{E}_1]\mathbf{a}| &= \left| \mathbb{E} \left[\mathbf{a}^\top \mathbf{Q}_{-\mathbf{x}\mathbf{x}^\top} \bar{\mathbf{Q}} \mathbf{a} \frac{\frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}} - \delta'}{(1 + \delta')(1 + \frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}})} \right] \right| \\ &\leq \mathbb{E} \left[|\mathbf{a}^\top \mathbf{Q}_{-\mathbf{x}}| |\mathbf{x}^\top \bar{\mathbf{Q}} \mathbf{a}| \left| \frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}} - \delta' \right| \right] \\ &\leq \mathbb{E}[|\mathbf{a}^\top \mathbf{Q}_{-\mathbf{x}}|] \mathbb{E}[|\mathbf{x}^\top \bar{\mathbf{Q}} \mathbf{a}|] \mathbb{E} \left[\left| \frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}} - \delta' \right| \right] \\ &= O(n^{-\frac{1}{2}})\end{aligned}$$

where we used here: (i) $\mathbf{a}^\top \bar{\mathbf{Q}} \mathbf{x} \propto C e^{-(\cdot)^q}$ and $\mathbf{a}^\top \mathbf{Q}_{-\mathbf{x}} \propto C e^{-c(\cdot)^q}$ (from which $\mathbb{E}[|\mathbf{a}^\top \bar{\mathbf{Q}} \mathbf{x}|^k] = O(1)$ and $\mathbb{E}[|\mathbf{a}^\top \mathbf{Q}_{-\mathbf{x}}|^k] = O(1)$) and (ii) $\frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}} \in \delta' \pm C e^{-c(n \cdot)^{q/2}} + C e^{-c(\sqrt{n \cdot})^q}$ (from which $\mathbb{E}[|\frac{1}{n} \mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}} - \delta'|^k] = O(n^{-\frac{k}{2}})$). The concentration results (i) and (ii) themselves unfold from the previous generic results on concentration of vectors and bilinear forms. Similarly,

$$|\mathbf{a}^\top \mathbb{E}[\mathbf{E}_2]\mathbf{a}| \leq \frac{1}{n} \mathbb{E}[|\mathbf{a}^\top \mathbf{Q}_{-\mathbf{x}}|] \mathbb{E}[|\mathbf{x}^\top \mathbf{Q}_{-\mathbf{x}} \Phi \bar{\mathbf{Q}} \mathbf{a}|] = O(n^{-1}).$$

Letting \mathbf{a} be the dominant singular vector of $\mathbb{E}\mathbf{Q} - \bar{\mathbf{Q}}$, we thus find that $\|\mathbb{E}\mathbf{Q} - \bar{\mathbf{Q}}\| = O(n^{-\frac{1}{2}})$. Integrated into $\mathbf{Q}(z) \in \mathbb{E}\mathbf{Q}(z) \pm C e^{-c(\sqrt{n \cdot})^q}$, this gives $\mathbf{Q}(z) \in \bar{\mathbf{Q}} \pm C e^{-c(\sqrt{n \cdot})^q}$.

It thus remains to show similarly that $\|\bar{\mathbf{Q}} - \bar{\bar{\mathbf{Q}}}\|$ is small. We have

$$\|\bar{\mathbf{Q}} - \bar{\bar{\mathbf{Q}}}\| = \frac{|\delta' - \delta|}{(1 + \delta)(1 + \delta')} \|\bar{\mathbf{Q}} \Phi \bar{\bar{\mathbf{Q}}}\| \leq \frac{|\delta - \delta'|}{|z|}$$

and it thus suffices to control $\delta - \delta'$, which, by the implicit form of δ , satisfies

$$\begin{aligned}|\delta - \delta'| &= \frac{1}{n} |\operatorname{tr} \Phi(\bar{\mathbf{Q}} - \bar{\bar{\mathbf{Q}}} + \bar{\bar{\mathbf{Q}}} - \mathbb{E}\mathbf{Q} + \mathbb{E}[\mathbf{Q} - \mathbf{Q}_-])| \\ &\leq \frac{1}{n} |\operatorname{tr} \Phi(\bar{\mathbf{Q}} - \bar{\bar{\mathbf{Q}}})| + \frac{1}{n} \operatorname{tr} \Phi \|\bar{\bar{\mathbf{Q}}} - \mathbb{E}\mathbf{Q}\| + \frac{1}{n} \operatorname{tr} \Phi \|\mathbb{E}[\mathbf{Q} - \mathbf{Q}_-]\| \\ &\leq \sqrt{\frac{1}{n(1 + \delta)^2} \operatorname{tr} \Phi^2 \bar{\mathbf{Q}}^2} \sqrt{\frac{1}{n(1 + \delta')^2} \operatorname{tr} \Phi^2 \bar{\bar{\mathbf{Q}}}^2} |\delta - \delta'| + O(n^{-\frac{1}{2}})\end{aligned}$$

where we used $\text{tr } \mathbf{AB} \leq \|\mathbf{B}\| \text{tr } \mathbf{A}$ for symmetric and nonnegative definite $\mathbf{A} \in \mathbb{R}^{p \times p}$, and $\|\mathbb{E}[\mathbf{Q} - \mathbf{Q}_-]\| = O(n^{-1/2})$, which unfolds from

$$\|\mathbb{E}[\mathbf{Q} - \mathbf{Q}_-]\| = \frac{1}{n} \left\| \mathbb{E} \frac{\mathbf{Q}_- \mathbf{x} \mathbf{x}^\top \mathbf{Q}_-}{1 + \frac{1}{n} \mathbf{x}^\top \mathbf{Q}_- \mathbf{x}} \right\| = \frac{1}{n} \left\| \frac{\mathbb{E}[\mathbf{Q}_- \Phi \mathbf{Q}_-]}{1 + \delta'} \right\| + O(n^{-\frac{1}{2}}).$$

The prefactor of $|\delta - \delta'|$ is strictly less than 1 for all large n , and thus $|\delta - \delta'| = O(n^{-\frac{1}{2}})$, which concludes the proof.

2.8 Concluding remarks

This section explored basic to advanced spectral properties of a family of random matrix models, with a strong emphasis on the sample covariance matrix model, in the regime of large and commensurable data number n and dimension p . Despite the simplicity of its definition, we saw that the limiting eigenvalue distribution of the sample covariance matrix is far from trivial, that quite advanced techniques from complex analysis are needed to perform statistical inference, and that, unlike in the classical $n \rightarrow \infty$ and p fixed regime, phase transition phenomena arise, below which some inference problems are asymptotically insoluble.

Fortunately, even if the statistical models used in concrete machine learning applications are often more involved, we will see that the main techniques and tools used to understand and improve machine learning algorithms and methods are essentially the same as those presented so far. In particular, we will see in the following sections that

- in (not necessarily linear) regression problems, the resolvent (of sample covariance matrices, of kernel matrices, of the Gram matrix of random feature maps, etc.) will systematically appear as the central object of interest (which is reminiscent of the fact that regression is an inverse problem);
- in classification problems, the spectrum of kernel random matrices and Laplacian random matrices (for spectral clustering or spectral community detection), or alternatively functionals of these kernel and Laplacian random matrices (for supervised or semi-supervised graph-based learning) will play an important role; the performance achieved by these methods, given in terms of misclassification rates, probability of false alarms, etc., will in particular demand the evaluation of the limiting means and variances of these functionals;
- in the specific case of spectral or subspace methods, such as PCA, manifold-based clustering, spectral clustering, and community detection, the aforementioned phase transition phenomena will arise and show that there exist strict limitations for these methods: in particular, a minimal samples-over-dimension ratio exists below which no detection or classification is possible;

- even for optimization-based machine learning problems, such as generalized linear models [Nelder and Wedderburn, 1972], that rarely offer a solution explicitly defined from (the resolvent of) a particular random matrix, their large dimensional (limiting) performance will be shown ultimately to depend in an almost explicit way on some slightly more involved random matrices; there, the twist will be to realize that some random quantities (not always easy to identify) converge and can asymptotically be replaced by deterministic equivalents obtained from a perturbation analysis (e.g., some sort of a local “linearization”).

Before delving into these applications, it is important to recall that we shall purposely place ourselves under the “realistic” situation where the number of samples n cannot be chosen arbitrarily large (samples do not always come for free in practice) and particularly not overwhelmingly larger than the typical dimension p of the data. More importantly, we also impose that the problem being addressed is not “asymptotically trivial”, i.e., for p, n realistically large, the misclassification probability or the cost to be minimized will not vanish. This way, the asymptotic analysis ($n, p \rightarrow \infty$) will be a realistic representative of the finite (but not too small) dimensional and (moderately) difficult machine learning problem. This is quite different from many parallel theoretical machine learning works which often aim at concluding (usually through the evaluation of error bounds, rather than exact results) that the algorithm under analysis provides an asymptotic perfect performance (vanishing misclassification rate or cost) in a certain growth regime of n with respect to p . Our vision instead is that, for a realist and meaningful analysis, n and p must be considered as both fixed (only not to too small values).

As such, to best appreciate the many results to come in the next chapters, these must be seen under this “finite-dimensional and realistic” lens.

2.9 Exercises

In this section, we provide short exercises to familiarize the reader with various useful notions and properties of random matrix calculus discussed this far, with detailed solutions deferred to the end of the monograph in Chapter 9.

[**copy the version in Chapter 9 here!**](#)

2.9.1 Properties of the Stieltjes transform

Exercise 1 (Stieltjes transform and moments). *Show that the Stieltjes transform $m_\mu(z)$, of a probability measure μ with bounded support (and thus finite moments), is a moment generating function in the sense that, for all $z \in \mathbb{C}$ such that $|z| > \max\{|\inf(\text{supp}(\mu))|, |\sup(\text{supp}(\mu))|\}$,*

$$m_\mu(z) = -\frac{1}{z} \sum_{n=0}^{\infty} M_n z^{-n}$$

where $M_n = \int t^n \mu(dt)$.

From this formulation, propose a method to evaluate the successive moments of μ .

Exercise 2 (Non-immediate Stieltjes transforms). *Let $\mathbf{X} \in \mathbb{R}^{n \times n}$ be a Hermitian matrix and $\mathbf{Q}(z) = (\mathbf{X} - z\mathbf{I}_n)^{-1}$ its resolvent. Show that, for any $\mathbf{u} \in \mathbb{R}^n$ unitary ($\|\mathbf{u}\| = 1$) and any \mathbf{A} nonnegative definite and with $\text{tr } \mathbf{A} = 1$, the quantities $\mathbf{u}^\top \mathbf{Q}(z) \mathbf{u}$ and $\text{tr } \mathbf{A} \mathbf{Q}(z)$ are the Stieltjes transform of probability measures.*

What are these measures and what are their supports?

Exercise 3 (Stieltjes transform and singular values). *Let μ be a probability measure on \mathbb{R}^+ and ν, ν' be the measures defined by*

$$\begin{aligned} \int f(t) \nu(dt) &= \int f(\sqrt{t}) \mu(dt) \\ \int f(t) \nu'(dt) &= \frac{1}{2} \left(\int f(t) \nu(dt) + \int f(-t) \nu(dt) \right) \end{aligned}$$

for all bounded continuous f .

What are ν, ν' when $\mu = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$ for $x_1, \dots, x_n \geq 0$?

Show that the Stieltjes transform $m_{\nu'}$ of ν' satisfies

$$m_{\nu'}(z) = zm_\mu(z^2).$$

Letting $\mathbf{X} \in \mathbb{R}^{n \times n}$ and μ be the empirical spectral distribution of $\mathbf{X} \mathbf{X}^\top$, relate the Stieltjes transform of the matrix

$$\Gamma = \begin{bmatrix} 0 & \mathbf{X} \\ \mathbf{X}^\top & 0 \end{bmatrix}$$

to that of the measure μ , and conclude on the nature of this Stieltjes transform.

Exercise 4 (Partial proof of Lemma 2.9). *For \mathbf{A}, \mathbf{M} symmetric nonnegative definite matrices, $\tau > 0$ and $z < 0$, using inversion lemmas and norm inequalities, prove that*

$$\left| \operatorname{tr} \mathbf{A} (\mathbf{M} + \tau \mathbf{u} \mathbf{u}^\top - z \mathbf{I}_p)^{-1} - \operatorname{tr} \mathbf{A} (\mathbf{M} - z \mathbf{I}_p)^{-1} \right| \leq \frac{\|\mathbf{A}\|}{|z|}.$$

2.9.2 On limiting laws

Exercise 5 (The $\sqrt{|x-b|}$ behavior of the edges). *Show that both the semicircle law and the Marčenko-Pastur law (for $c \neq 1$) have a local $\sqrt{|x-b|}$ behavior at each of the edges b of their support.*

Conclude on the typical number of eigenvalues of the Wishart matrix $\frac{1}{n} \mathbf{X} \mathbf{X}^\top \in \mathbb{R}^{p \times p}$ with $\mathbf{X}_{ij} \sim \mathcal{N}(0, 1)$ independent, and the Wigner $\mathbf{X} \in \mathbb{R}^{n \times n}$ with $\mathbf{X}_{ij} = \mathbf{X}_{ji} \sim \mathcal{N}(0, 1)$ independent up to symmetry, found near the edges of their respective supports.

Relate this finding to the fluctuations of the Tracy-Widom distribution of the largest and smallest eigenvalues.

What happens for the left-edge of the support of the Marčenko-Pastur law and to the associated smallest eigenvalues of Wishart matrices when $\lim p/n = c = 1$? How many eigenvalues are then found close to the left edge in this so-called “hard-edge” setting? Conclude on the typical fluctuations of these eigenvalues and confirm numerically.

Exercise 6 (The $\sqrt{|x-b|}$ behavior in elaborate models). *We here seek to extend the results of Exercise 5 to the sample covariance matrix model $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ where $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{Z}$ with \mathbf{Z} having independent standard Gaussian entries and \mathbf{C} having a bounded limiting spectral measure ν with fast decaying tails. We denote $\tilde{m}(z)$ the Stieltjes transform of the limiting spectral measure $\tilde{\mu}$ of $\frac{1}{n} \mathbf{X}^\top \mathbf{X}$.*

Using Figure 2.5 as a reference and recalling the formulation for the inverse Stieltjes transform

$$x(\tilde{m}) = -\frac{1}{\tilde{m}} + c \int \frac{t \nu(dt)}{1 + t \tilde{m}}$$

visually justify that $x''(\tilde{m})$ can be (complex) analytically extended in the neighborhood of each point \tilde{m} where $x'(\tilde{m}) = 0$ into a function $z(\tilde{m})$ which must locally coincide with the inverse Stieltjes transform of $\tilde{m}(z)$.

Deduce that $\tilde{m}(z)$ must be of the form $\sqrt{z-b}$ near an edge and conclude.

Exercise 7 (Further results on $x(\tilde{m})$). *We aim in this exercise to justify some of the visual observations made on Figure 2.5.*

Show that, for $\tilde{m}_1 \neq \tilde{m}_2$ such that $x'(\tilde{m}_1), x'(\tilde{m}_2) > 0$, we cannot have $x(\tilde{m}_1) = x(\tilde{m}_2)$: that is, the increasing segments of $x(\tilde{m})$ never “overlap”.

Besides, show that, if $\tilde{m}_1 < \tilde{m}_2$ are both of the same sign, and $x'(\tilde{m}_1), x'(\tilde{m}_2) > 0$, then $x(\tilde{m}_1) < x(\tilde{m}_2)$: that is, the increasing segments of $x(\tilde{m})$ never “swap”.

To this end, we may prove the intermediary result

$$(\tilde{m}_1 - \tilde{m}_2) \left(1 - \int \frac{c\tilde{m}_1\tilde{m}_2 t^2 \nu(dt)}{(1+t\tilde{m}_1)(1+t\tilde{m}_2)} \right) = \tilde{m}_1\tilde{m}_2(x(\tilde{m}_1) - x(\tilde{m}_2))$$

and use the Cauchy-Schwarz inequality to control the left-hand side parenthesis.

Finally show that, if ν has bounded support, then $x(\tilde{m}) \rightarrow 0$ as $\tilde{m} \rightarrow \pm\infty$.

As a final remark, note that the only important observation about Figure 2.5 which we have not shown here is the fact that the points \tilde{m} where $x'(\tilde{m}) = 0$ must exist. In fact, this is not always the case and heavily depends on the nature of the tails of the underlying measure ν . Justify in particular that, for some ν , there may be no asymptote on the edges of the domain of definition of $x(\cdot)$ (as opposed to what is seen in Figure 2.5).

2.9.3 On eigen-inference

Exercise 8 (Alternative estimates of $\frac{1}{p} \text{tr}(\frac{1}{n} \mathbf{X}\mathbf{X}^\top)^2$). Let $\mathbf{X} = \mathbf{C}^{\frac{1}{2}} \mathbf{Z}$ for $\mathbf{Z} \in \mathbb{R}^{p \times n}$ with independent standard Gaussian entries, and \mathbf{C} deterministic symmetric nonnegative definite, of bounded spectral norm, and limiting eigenvalue distribution ν .

By a direct calculus, determine the limit, as $n, p \rightarrow \infty$ and $p/n \rightarrow c > 0$ of the second order moment

$$M_2 = \frac{1}{p} \text{tr} \left(\frac{1}{n} \mathbf{X}\mathbf{X}^\top \right)^2$$

as a function of the moments of ν .

Retrieve the same result using the results of Exercise 1 along with the expression of the Stieltjes transform $m(z)$ of the limiting spectrum μ of $\frac{1}{n} \mathbf{X}\mathbf{X}^\top$. Hint: It may be useful here to first show that $m(z)$ is solution to

$$m(z) = \int \frac{\nu(dt)}{-z(1 + ctm(z)) + (1 - c)t}$$

with ν the limiting spectral measure of \mathbf{C} .

Exercise 9 (Location of the zeros of $\tilde{m}(z)$). Figure 2.7 and Remark 2.12 both show that the zeros η_1, \dots, η_n of $m_{\mathbf{X}}(z)$, the Stieltjes transform of a symmetric matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$, are interlaced with the eigenvalues $\lambda_1, \dots, \lambda_n$ of \mathbf{X} .

In the sample covariance matrix case $\frac{1}{n} \mathbf{Z}^\top \mathbf{C} \mathbf{Z}$ with $\mathbf{Z} \in \mathbb{R}^{p \times n}$ having independent standard Gaussian entries and \mathbf{C} with limited spectral measure ν of bounded and connected support, this means that (up to zero eigenvalues) the roots η_i of $m_{\frac{1}{n} \mathbf{Z}^\top \mathbf{C} \mathbf{Z}}(z)$ are all found in the limiting support $\tilde{\mu}$ of the empirical spectral distribution, at the possible exception of the leftmost η_1 .

Using a variable change involving $\tilde{m}(z)$ on the formula

$$0 = \frac{1}{2\pi i} \oint_{\Gamma} \frac{1}{w} dw$$

for all Γ not enclosing zero, then the approximation $\tilde{m}(z) = m_{\frac{1}{n}\mathbf{Z}^\top \mathbf{C} \mathbf{Z}}(z) + o(1)$ and finally a residue calculus, show that no root of $m_{\frac{1}{n}\mathbf{Z}^\top \mathbf{C} \mathbf{Z}}(z)$ can be found at macroscopic distance from the limiting support of μ . Conclude.

2.9.4 Spiked models

Exercise 10 (Additive spiked model). Similar to Theorem 2.13, show the phase transition threshold for the additive model $\mathbf{Y} \equiv \frac{1}{n}\mathbf{X}\mathbf{X}^\top + \mathbf{P}$ for \mathbf{X} having i.i.d. entries of zero mean, unit variance and low rank $\mathbf{P} = \sum_{i=1}^k \ell_i \mathbf{u}_i \mathbf{u}_i^\top$, with $\ell_1 > \dots > \ell_k > 0$, is determined by the condition

$$\ell_i > \sqrt{c}(1 + \sqrt{c})$$

with $c = \lim p/n$ as $p, n \rightarrow \infty$. Under this condition, show that the (almost sure) limiting value of the corresponding isolated eigenvalue $\hat{\gamma}_i$ of \mathbf{Y} is given by

$$\hat{\gamma}_i \xrightarrow{a.s.} \gamma_i = 1 + \ell_i + \frac{c}{\ell_i - c}.$$

Further show that, letting $\hat{\mathbf{u}}_i$ be the eigenvector of \mathbf{Y} associated with $\hat{\gamma}_i$, we have the convergence

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 \xrightarrow{a.s.} 1 - \frac{c}{(\ell_i - c)^2}. \quad (2.47)$$

Exercise 11 (Additive spiked model: the Wigner case). Let \mathbf{X} be symmetric with X_{ij} , $i \geq j$, i.i.d. with zero mean and unit variance. As in Exercise 10, show that the “spiked” phase transition threshold for the model $\mathbf{Y} \equiv \frac{1}{\sqrt{n}}\mathbf{X} + \mathbf{P}$, where $\mathbf{P} = \sum_{i=1}^k \ell_i \mathbf{u}_i \mathbf{u}_i^\top$, with $\ell_1 > \dots > \ell_k > 0$, is determined by the condition

$$\ell_i > 1$$

and that, under this condition, the isolated eigenvalue $\hat{\gamma}_i$ of \mathbf{Y} associated with ℓ_i is given by

$$\hat{\gamma}_i \xrightarrow{a.s.} \gamma_i = \ell_i + \frac{1}{\ell_i}.$$

Show finally that, for $\hat{\mathbf{u}}_i$ the eigenvector of \mathbf{Y} associated with $\hat{\gamma}_i$, we have

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 \xrightarrow{a.s.} 1 - \frac{1}{\ell_i^2}.$$

2.9.5 Deterministic equivalent

Exercise 12 (Proof of Theorem 2.17). Inspired by the proofs of Theorem 2.6, prove Theorem 2.17 using

1. the trace method adapted to Haar random matrices, Lemma 2.16; and
2. the Stein’s lemma adapted to Haar random matrices, Lemma 2.17.

Exercise 13 (Higher-order deterministic equivalent). *Theorem 2.4 provides an deterministic equivalent for the resolvent $\mathbf{Q} = \left(\frac{1}{n}\mathbf{XX}^\top - z\mathbf{I}_p\right)^{-1}$ for $\mathbf{X} \in \mathbb{R}^{p \times n}$ having i.i.d. zero mean and unit variance entries, which, according to Notation 1, provides access to the asymptotic behavior of $\mathbf{a}^\top \mathbf{Q} \mathbf{b}$. In many machine learning applications, however, the object of natural interest (e.g., mean squared error in regression analysis and variance analysis in a classification context) often involves the asymptotic behavior of $\mathbf{a}^\top \mathbf{Q} \mathbf{A} \mathbf{Q} \mathbf{b}$ which requires a deterministic equivalent for random matrices of the type $\mathbf{Q} \mathbf{A} \mathbf{Q}$, for arbitrary \mathbf{A} independent of \mathbf{Q} . In particular, for $\mathbf{Q} \leftrightarrow \bar{\mathbf{Q}}$ (such that $\|\mathbb{E}[\mathbf{Q}] - \bar{\mathbf{Q}}\| \rightarrow 0$), $\bar{\mathbf{Q}} \mathbf{A} \bar{\mathbf{Q}}$ is in general not a deterministic equivalent for $\mathbf{Q} \mathbf{A} \mathbf{Q}$, due to the fact that*

$$\mathbb{E}[\mathbf{Q} \mathbf{A} \mathbf{Q}] \not\simeq \mathbb{E}[\mathbf{Q}] \mathbf{A} \mathbb{E}[\mathbf{Q}].$$

Instead, prove that, in the setting of Theorem 2.4, one has

$$\mathbf{Q}(z) \mathbf{A} \mathbf{Q}(z) \leftrightarrow m(z)^2 \mathbf{A} + \frac{1}{n} \operatorname{tr} \mathbf{A} \cdot \frac{m'(z)m(z)^2}{(1+cm(z))^2} \mathbf{I}_p. \quad (2.48)$$

For sanity check, using the fact that $\partial \mathbf{Q}(z)/\partial z = \mathbf{Q}^2(z)$ and taking $\mathbf{A} = \mathbf{I}_p$ in the equation above, confirm that

$$\mathbf{Q}^2(z) \leftrightarrow m'(z) \mathbf{I}_p$$

for $m'(z) = \frac{m(z)^2}{1 - \frac{cm(z)^2}{(1+cm(z))^2}}$ obtained by differentiating the Marčenko–Pastur equation (2.9).

2.9.6 Concentration of measure

Exercise 14 (Concentration of matrix quadratic forms). *Recalling the definitions and notations of Section 2.7, let $\mathbf{X} \in \mathbb{R}^{p \times n}$ be a random matrix satisfying*

$$\mathbf{X} \propto C e^{c \cdot^2}, \text{ and } \|\mathbb{E}[\mathbf{X}]\| \leq K$$

for some $K, C, c > 0$. Given $\mathbf{A} \in \mathbb{R}^{p \times p}$ deterministic, we aim to prove the linear concentration of $\mathbf{X}^\top \mathbf{A} \mathbf{X}$ in $(\mathbb{R}^{n \times n}, \|\cdot\|_F)$. To this end, we consider a deterministic matrix $\mathbf{B} \in \mathbb{R}^{n \times n}$ such that $\|\mathbf{B}\|_F \leq 1$ and study the behavior of $\operatorname{tr}(\mathbf{B} \mathbf{X}^\top \mathbf{A} \mathbf{X})$. Consider first the singular value decomposition

$$\mathbf{A} = \mathbf{U}_\mathbf{A} \Lambda_\mathbf{A} \mathbf{V}_\mathbf{A}^\top, \quad \mathbf{B} = \mathbf{U}_\mathbf{B} \Lambda_\mathbf{B} \mathbf{V}_\mathbf{B}^\top,$$

with $\mathbf{U}_\mathbf{A}, \mathbf{V}_\mathbf{A} \in \mathbb{R}^{p \times p}$ and $\mathbf{U}_\mathbf{B}, \mathbf{V}_\mathbf{B} \in \mathbb{R}^{n \times n}$ orthogonal matrices, $\Lambda_\mathbf{A} \in \mathbb{R}^{p \times p}$, $\Lambda_\mathbf{B} \in \mathbb{R}^{n \times n}$ diagonal matrices, and define $\tilde{\mathbf{X}}_1 = \mathbf{U}_\mathbf{A}^\top \mathbf{X} \mathbf{V}_\mathbf{B}$, $\tilde{\mathbf{X}}_2 = \mathbf{V}_\mathbf{A}^\top \mathbf{X} \mathbf{U}_\mathbf{B} \in \mathbb{R}^{p \times n}$. In the sequel, the constants $K', C', c' > 0$ are understood only depending on K, C, c and might change from line to line.

First show that there exist $K', C', c' > 0$ such that, for $t > K' \sqrt{\log(np)}$ and $\tilde{\mathbf{X}} \in \{\tilde{\mathbf{X}}_1, \tilde{\mathbf{X}}_2\}$

$$\mathbb{P}\left(\|\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}]\|_\infty \geq t\right) \leq C' e^{-c' t^2 / \log(np)}.$$

Deduce from the bound $\|\mathbb{E}[\mathbf{X}]\| \leq K$ that there exists a constant $K' > 0$ depending only on K, C, c such that

$$\mathbb{E}[\|\tilde{\mathbf{X}}\|_\infty] \leq K' \sqrt{\log(np)}.$$

This established, introduce the set $\mathcal{A}_\theta = \{\mathbf{X} \in \mathbb{R}^{p \times n}, \max\{\|\tilde{\mathbf{X}}_1\|_\infty, \|\tilde{\mathbf{X}}_2\|_\infty\} \leq \theta\} \subset \mathbb{R}^{p \times n}$ and show that for all $\theta \geq K' \sqrt{\log(np)}$ with $K' > 1$, we have

$$\mathbb{P}(\mathbf{X} \in \mathcal{A}_\theta^c) \leq C' e^{-c'\theta^2}$$

and that the mapping $\mathbf{X} \mapsto \text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})$ is $\theta\|\mathbf{A}\|_F$ -Lipschitz on \mathcal{A}_θ .

Introduce M , a median of $\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})$, and note that

$$\mathbb{P}(|\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X}) - M| \geq t, \mathbf{X} \in \mathcal{A}_\theta) \leq C' e^{-c't^2/(\theta\|\mathbf{A}\|_F)^2}.$$

Conclude by carefully choosing the parameter $\theta \geq K' \sqrt{\log(np)}$ and showing that

$$\mathbf{X}^\top \mathbf{A}\mathbf{X} \in \mathbb{E}[\mathbf{X}^\top \mathbf{A}\mathbf{X}] \pm C' e^{-c'^2/(\log(np)\|\mathbf{A}\|_F^2)} + C' e^{-c'/\|\mathbf{A}\|_F}.$$

2.9.7 Beyond matrices

Exercise 15 (Towards Spiked Models in Random Tensors). Let $\mathcal{Y} \in \mathbb{R}^{n \times n \times n}$ be a three-way symmetric tensor, i.e., such that \mathcal{Y}_{ijk} is constant to exchanges of its indexes, defined by

$$\mathcal{Y} = \lambda \mathbf{x} \otimes \mathbf{x} \otimes \mathbf{x} + \frac{1}{\sqrt{n}} \mathcal{W}$$

where $\mathcal{W} \in \mathbb{R}^{n \times n \times n}$ has independent $\mathcal{N}(0, 1)$ entries up to symmetry, $\mathbf{x} \in \mathbb{R}^n$ is of unit norm, and $[\mathbf{a} \otimes \mathbf{b} \otimes \mathbf{c}]_{ijk} = a_i b_j c_k$.

A possible definition of the “eigenvalue-eigenvector” pair (ℓ, \mathbf{u}) with $\|\mathbf{u}\| = 1$ of the symmetric tensor \mathcal{Y} is given by the solutions to

$$\mathcal{Y} \cdot \mathbf{u} \cdot \mathbf{u} = \ell \mathbf{u}$$

where $\mathcal{A} \cdot \mathbf{a} \cdot \mathbf{b} = \sum_{ijk} \mathcal{A}_{ij} a_i b_j \in \mathbb{R}^n$ is the contraction of tensor \mathcal{A} on the vectors \mathbf{a}, \mathbf{b} . The objective is to characterize the largest eigenvalue ℓ_{\max} as well as the associated alignment $|\mathbf{u}_{\max}^\top \mathbf{x}|$ between the dominant eigenvector and the spike \mathbf{x} .

Show first that the matrix $\mathbf{Y}_x = \mathcal{Y} \cdot \mathbf{x} = \sum_{i=1}^n \mathcal{Y}_{i,\cdot,\cdot} x_i$ is given by

$$\mathbf{Y}_x = \lambda \mathbf{x} \mathbf{x}^\top + \frac{1}{\sqrt{n}} \sum_{i=1}^n x_i \mathbf{W}_{i,\cdot,\cdot}$$

where $\mathbf{W}_{i,\cdot,\cdot} \in \mathbb{R}^{n \times n}$ is the i -th “layer” matrix of the tensor \mathcal{W} with $[\mathbf{W}_{i,\cdot,\cdot}]_{ab} = \mathcal{W}_{iab}$.

Using Pastur’s Stein approach, show that the limiting empirical spectral measure of \mathbf{Y}_x is the semicircle distribution supported on $[-2, 2]$ (we may discard the rank-one matrix $\lambda \mathbf{x} \mathbf{x}^\top$ to retrieve this result). Then, using a spiked analysis, show that

- for all $\lambda > 0$, there must exist an isolated eigenvalue $\hat{\gamma}$ of \mathbf{Y}_x (thus no phase transition) asymptotically equal to (with high probability)

$$\hat{\gamma} \rightarrow \gamma = \sqrt{\lambda^2 + 4};$$

- the eigenvector $\hat{\mathbf{u}}$ associated with $\hat{\gamma}$ satisfies (with high probability)

$$|\hat{\mathbf{u}}^\top \mathbf{x}|^2 \rightarrow \frac{\lambda}{\sqrt{\lambda^2 + 4}}.$$

Conclude on a (asymptotic) bound for the quantity $\ell_{\max} |\mathbf{u}_{\max}^\top \mathbf{x}|$.

Chapter 3

Statistical Inference in Linear Models

Sections 2.2 through 2.5 provided the basic material to perform fundamental signal and data processing tasks such as detection (hypothesis testing) and estimation (statistical inference) for sample covariance matrix models.

These sections can be summarized as follows: if no a priori information is known about the population covariance matrix $\mathbf{C} \in \mathbb{R}^{p \times p}$ (i.e., it is not known to be sparse, Toeplitz, etc.), the observation of i.i.d. (say zero mean) samples $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ with covariance $\text{Cov}(\mathbf{x}_i) = \mathbf{C}$ is not sufficient to estimate \mathbf{C} itself if p and n are of the same order of magnitude (essentially because the np degrees of freedom in \mathbf{X} are not enough to evaluate the $O(p^2)$ distinct elements of \mathbf{C}). As such, the standard methods for detection and estimation involving \mathbf{C} which conventionally substitute $\hat{\mathbf{C}} = \frac{1}{n}\mathbf{X}\mathbf{X}^\top$ for \mathbf{C} are bound to fail when p is not too small compared to n .

Yet, \mathbf{C} itself may not be the object of central interest. One is often rather interested in a scalar functional of \mathbf{C} : the binary answer to a signal detection procedure, the probability of an hypothesis test, the class-label in a classification method, or more generally the estimation of a certain more-or-less involved function of \mathbf{C} (its dominant eigenvalue, the content of its dominant eigenvector, etc.).

Section 2.3 showed that, while $\hat{\mathbf{C}} \not\rightarrow \mathbf{C}$ in the random matrix regime, there exist complex analytic relations between the resolvents $\mathbf{Q}_{\hat{\mathbf{C}}}(z)$ of $\hat{\mathbf{C}}$ and $\mathbf{Q}_{\mathbf{C}}(z)$ of \mathbf{C} . These relations enable the connection of a large class of functionals of \mathbf{C} (linear functionals of its eigenvalues, subspace projections of some of its eigenvectors) to those of $\hat{\mathbf{C}}$, allowing for random-matrix improved estimates of these functionals. However, these estimates can be quite involved and limited in practice by complex integration boundaries. In many cases of practical interest where information and noise can be decoupled in a *low rank* information and a *high rank* noise, spiked models previously discussed in Section 2.5 give access to much simplified versions of these inference methods.

In this chapter, examples of concrete problems involving covariance matrix-based estimators (and beyond) are discussed:

1. **Hypothesis testing in signal-plus-noise model:** assuming $\mathbf{x}_i = \mathbf{z}_i$ is pure noise or $\mathbf{x}_i = \boldsymbol{\mu} + \mathbf{z}_i$ contains an unknown signal and noise, we discuss in Section 3.1.1 the generalized likelihood ratio test (GLRT) aiming to detect the presence of a signal, using asymptotic results on the eigenvalues of the sample covariance model.
2. **Linear and quadratic discriminant analysis:** the objective of LDA and QDA discussed in Section 3.1.2 is to classify a test datum \mathbf{x} into one of the two Gaussian hypotheses $\mathcal{N}(\boldsymbol{\mu}_0, \mathbf{C}_0)$ or $\mathcal{N}(\boldsymbol{\mu}_1, \mathbf{C}_1)$, based on some (linear) discrimination rule $T(\mathbf{x})$ obtained from the available training set. Random matrix results will be used here to analyze the (surprisingly far from obvious) performance of LDA/QDA.
3. **Estimation with subspace methods:** for signals parameterized by some specific “direction of arrival” angles $\theta_1, \dots, \theta_k$, the so-called subspace methods, in particular the MUSIC algorithm, can be used to recover this “angular” information from the signal (containing noise) sample covariance. In Section 3.1.3, we introduce the random matrix improved G-MUSIC algorithm and discuss its significant performance gain in the large dimensional signal scenario.
4. **Distance estimation:** as a concrete example of statistical inference procedures commonly used in machine learning, in Section 3.2 we will estimate the distance between two centered (population) data distributions (or covariance matrices) based on few samples from each distribution. Here random matrix theory provides original estimators which are consistent in the regime of simultaneously large sample size and data dimension.
5. **Robust covariance estimation:** in presence of outliers, sample covariance matrices are known to be non-robust estimators of population covariance matrices, already in the $n \gg p$ case: robust estimators of scatter are an efficient alternative in these situations. These objects are hard to theoretically grasp in the classical $n \gg p$ setting; we present here a much more tractable random matrix analysis of these estimates in Section 3.3.

These topics not being *core* machine learning algorithms but rather statistical tools broadly surrounding machine learning (and for most of them already well documented in other textbooks [Tulino and Verdú, 2004, Couillet and Debbah, 2011]), each subject will be mostly “brushed over” rather than exhaustively explored. Pointers to external articles and research works are provided for the interested readers.

3.1 Detection and estimation in information-plus-noise models

3.1.1 GLRT asymptotics

Possibly one of the most immediate and telling applications of random matrix theory, and particularly of spiked models, in statistics deals with the decision of the presence of some “information” buried in white noise.

Denoting $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ a matrix with i.i.d. columns \mathbf{x}_i , the decision problem is formulated as the binary hypothesis test:

$$\mathbf{X} = \begin{cases} \sigma \mathbf{Z}, & \mathcal{H}_0 \\ \mathbf{a} \mathbf{s}^\top + \sigma \mathbf{Z}, & \mathcal{H}_1 \end{cases}$$

where $\mathbf{Z} = [\mathbf{z}_1, \dots, \mathbf{z}_n]$ with $\mathbf{z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, $\mathbf{a} \in \mathbb{R}^p$ deterministic with unit norm $\|\mathbf{a}\| = 1$, $\mathbf{s} = [s_1, \dots, s_n]^\top \in \mathbb{R}^n$ with s_i i.i.d. random scalars, and $\sigma > 0$. We also denote $c = p/n$ (and demand as usual that $0 < \liminf c \leq \limsup c < \infty$).

This model describes the observation of either pure noise data $\sigma \mathbf{z}_i$ with zero mean and covariance $\sigma^2 \mathbf{I}_p$ or of a deterministic information \mathbf{a} possibly modulated by a scalar (random) signal s_i (which could simply be ± 1) added to the noise. Obviously, if the parameters \mathbf{a} , σ as well as the statistics of s_i are known, a mere Neyman-Pearson test allows one to discriminate between \mathcal{H}_0 and \mathcal{H}_1 with optimal detection probability, for all finite n, p ; precisely, one will decide on the genuine hypothesis according to the ratio of posterior probabilities

$$\frac{\mathbb{P}(\mathbf{X} | \mathcal{H}_1)}{\mathbb{P}(\mathbf{X} | \mathcal{H}_0)} \underset{\mathcal{H}_0}{\gtrless} \alpha$$

for some $\alpha > 0$ controlling the desired Type I and Type II error rates (that is, the probability of false positives and of false negatives).

However, in practice, unless the existence of a set of previous pure-noise acquisitions is assumed, it is quite unlikely that σ be already known. Similarly, if the ultimate objective (post-decision) is to estimate the data structure \mathbf{a} under \mathcal{H}_1 , \mathbf{a} is naturally assumed partially or completely unknown (it may be known to belong to a subset of \mathbb{R}^p in which case more elaborate procedures than proposed here can be carried on). In the most generic scenario where \mathbf{a} is fully unknown, assuming additionally the data of zero mean, we may thus impose without generality restriction that $\mathbb{E}[s_i] = 0$ and $\mathbb{E}[s_i^2] = 1$. Under this very restricted prior knowledge, instead of the maximum likelihood test, one may resort to a *generalized likelihood ratio test* (GLRT) defined as

$$\frac{\sup_{\sigma, \mathbf{a}} \mathbb{P}(\mathbf{X} | \sigma, \mathbf{a}, \mathcal{H}_1)}{\sup_{\sigma, \mathbf{a}} \mathbb{P}(\mathbf{X} | \sigma, \mathcal{H}_0)} \underset{\mathcal{H}_0}{\gtrless} \alpha.$$

Under both a Gaussian noise and signal s_i assumption, the GLRT has an explicit expression which appears to be a monotonous increasing function of

$\|\mathbf{X}\mathbf{X}^T\|/\text{tr}(\mathbf{X}\mathbf{X}^T)$. That is, the test is equivalent to

$$T_p \equiv \frac{\left\| \frac{1}{n} \mathbf{X} \mathbf{X}^T \right\|}{\frac{1}{p} \text{tr} \left(\frac{1}{n} \mathbf{X} \mathbf{X}^T \right)} \stackrel{\mathcal{H}_1}{\gtrless} f(\alpha) \quad \stackrel{\mathcal{H}_0}{\leq}$$

(the result is immediate but may be found in detail in [Wax and Kailath, 1985, Anderson et al., 1963]) for some known monotonously increasing function f ,¹ where we introduced the scales $1/p$ and $1/n$ to maintain both numerator and denominator of order $O(1)$ as $n, p \rightarrow \infty$.

Obviously, since the ratio has limit $(1 + \sqrt{c})^2$ under the \mathcal{H}_0 asymptotics, $f(\alpha)$ must be of the form $f(\alpha) = (1 + \sqrt{c})^2 + g(\alpha)$ for some $g(\alpha) > 0$. Also, as we know that $\frac{1}{p} \text{tr} \left(\frac{1}{n} \mathbf{X} \mathbf{X}^T \right)$ fluctuates at the (fastest) speed $O(n^{-1})$, while $\left\| \frac{1}{n} \mathbf{X} \mathbf{X}^T \right\|$ fluctuates at the slower speed $O(n^{-2/3})$ (as per Theorem 2.15), the global fluctuation is dominated by the numerator at a rate of order $O(n^{-2/3})$, i.e., under \mathcal{H}_0 we have

$$T_p \stackrel{\mathcal{H}_0}{=} (1 + \sqrt{c})^2 + O(n^{-2/3}).$$

Since the denominator essentially converges while the numerator still fluctuates, despite the dependence between both, only the fluctuations of $\left\| \frac{1}{n} \mathbf{X} \mathbf{X}^T \right\|$ influence the behavior of the ratio, and thus

$$T_p \stackrel{\mathcal{H}_0}{\sim} (1 + \sqrt{c})^2 + (1 + \sqrt{c})^{\frac{4}{3}} c^{-\frac{1}{6}} n^{-\frac{2}{3}} \text{TW}_1 + o_p(n^{-2/3})$$

where $\stackrel{\mathcal{H}_0}{\sim}$ denotes equality in law under \mathcal{H}_0 .

As a consequence, in order to set a maximum false alarm rate (or false positive, or Type I error) of $r > 0$ in the limit of large n, p , one must choose a threshold $f(\alpha)$ for T_p such that

$$\mathbb{P}(T_p \leq f(\alpha)) = r$$

that is, such that

$$\mu_{\text{TW}_1}((-\infty, A_p]) = r, \quad A_p = (f(\alpha) - (1 + \sqrt{c})^2)(1 + \sqrt{c})^{-\frac{4}{3}} c^{\frac{1}{6}} n^{\frac{2}{3}} \quad (3.1)$$

with $\mu_{\text{TW}_1}(dt)$ the Tracy-Widom measure.

Figure 3.1 compares the empirical false alarm rate obtained from different choices of thresholds $f(\alpha) = (1 + \sqrt{c})^2 + O(n^{-2/3})$ to the asymptotic estimate $\text{TW}_1(A_p)$. For a given maximum false alarm rate r , one can thus numerically determine the threshold $f(\alpha)$ for which $\text{TW}_1(A_p(f(\alpha))) = r$.

Assume now that, under the \mathcal{H}_1 hypothesis, $\mathbf{s} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$. We may then write

$$\mathbf{X} = \mathbf{a} \mathbf{s}^T + \sigma \mathbf{Z} = [\mathbf{a} \quad \sigma \mathbf{I}_p] \tilde{\mathbf{Z}}, \quad \tilde{\mathbf{Z}} = \begin{bmatrix} \mathbf{s}^T \\ \mathbf{Z} \end{bmatrix}$$

¹Specifically, f is the functional inverse of $g : x \mapsto (1 - 1/p)^{(1-p)n} x^{-n} (1 - x/p)^{n(1-p)}$. See also [Bianchi et al., 2011] for detail.




Figure 3.1: Comparison between empirical false alarm rates and $\text{TW}_1(A_p)$ for A_p defined in (3.1), as a function of the threshold $f(\alpha) \in [(1+\sqrt{c})^2 - 5n^{-2/3}, (1+\sqrt{c})^2 + 5n^{-2/3}]$, for $p = 256$, $n = 1024$ and $\sigma = 1$. Results obtained from 100 runs. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

with $\tilde{\mathbf{Z}} \in \mathbb{R}^{(p+1) \times n}$ having i.i.d. $\mathcal{N}(0, 1)$ entries. Hence $\sigma^{-1}\mathbf{X}$ has independent columns with zero mean and covariance

$$\mathbf{C} \equiv \mathbb{E} \left[\sigma^{-2} \frac{1}{n} \mathbf{X} \mathbf{X}^\top \right] = \mathbf{I}_p + \sigma^{-2} \mathbf{a} \mathbf{a}^\top.$$

This is a spiked model with population eigenvalues (i) $\sigma^{-2} + 1$ with unit multiplicity, and (ii) 1 with multiplicity $p - 1$. We thus know from Theorem 2.13 that T_p converges to a quantity strictly greater than $(1 + \sqrt{c})^2$ if and only if the “signal-to-noise ratio” σ^{-2} satisfies $\sigma^{-2} > \sqrt{c}$.

Assuming $\sigma = \sigma_p$ depends on p, n , we thus have that, for the signal detection to be asymptotically non-trivial, σ_p^{-2} must be of the form $\sqrt{c} + O(n^{-2/3})$, in which case $T_p = (1 + \sqrt{c})^2 + O(n^{-2/3})$. From there, the probability of correct detection under \mathcal{H}_1 (or *power of the test*) can be derived, and in turn the receiver operator curve (ROC) of the detection test. We do not pursue these considerations further – which, for the interested reader, are given a detailed account in [Bianchi et al., 2011] –, as (i) they require more elaborate technical material (specifically, large deviation theory) not discussed in Chapter 2, being of marginal use in practice, and (ii) retrieving the power of the test has little practical algorithmic relevance as it can only be obtained when σ and $\|\mathbf{a}\|$ are known.

We instead move on in the next section to discriminant analysis, a familiar tool in statistical machine learning for which obtaining a full account of the

asymptotic performance is a crucial step to properly parameterize (and thus optimize) the method.

3.1.2 Linear and Quadratic Discriminant Analysis

The application of the random matrix framework to linear (LDA) and quadratic (QDA) discriminant analyses is a very telling example of the counterintuitive behavior of large dimensional statistics.

Specifically, LDA and QDA aim at selecting one out of two Gaussian model hypotheses $\mathcal{N}(\boldsymbol{\mu}_0, \mathbf{C}_0)$ (hypothesis \mathcal{H}_0) versus $\mathcal{N}(\boldsymbol{\mu}_1, \mathbf{C}_1)$ (hypothesis \mathcal{H}_1) for an observed vector $\mathbf{x} \in \mathbb{R}^p$.

The means and covariances under \mathcal{H}_0 and \mathcal{H}_1 are however unknown, and are instead directly estimated from two sets of training data $\mathbf{x}_1^{(\ell)}, \dots, \mathbf{x}_{n_\ell}^{(\ell)} \sim \mathcal{N}(\boldsymbol{\mu}_\ell, \mathbf{C}_\ell)$ with $\ell \in \{0, 1\}$, as per the standard empirical estimators

$$\begin{aligned}\hat{\boldsymbol{\mu}}_\ell &\equiv \frac{1}{n_\ell} \sum_{i=1}^{n_\ell} \mathbf{x}_i^{(\ell)} \\ \hat{\mathbf{C}}_\ell &\equiv \frac{1}{n_\ell - 1} \sum_{i=1}^{n_\ell} (\mathbf{x}_i^{(\ell)} - \hat{\boldsymbol{\mu}}_\ell)(\mathbf{x}_i^{(\ell)} - \hat{\boldsymbol{\mu}}_\ell)^\top.\end{aligned}$$

The test decision for an unknown new observation \mathbf{x} is then carried out using a standard Neyman–Pearson likelihood-ratio procedure under the assumption that $\boldsymbol{\mu}_\ell$ and \mathbf{C}_ℓ are correctly estimated by $\hat{\boldsymbol{\mu}}_\ell$ and $\hat{\mathbf{C}}_\ell$. In the context of LDA, it is assumed that $\mathbf{C}_0 = \mathbf{C}_1$ (which may be an invalid assumption), in which case the discrimination is based on the test:²

$$T_{\text{LDA}}(\mathbf{x}) \equiv (\mathbf{x} - \hat{\boldsymbol{\mu}})^\top \hat{\mathbf{C}}^{-1} (\hat{\boldsymbol{\mu}}_0 - \hat{\boldsymbol{\mu}}_1) \stackrel{\mathcal{H}_0}{\gtrless} \stackrel{\mathcal{H}_1}{\gtrless} 0$$

where $\hat{\boldsymbol{\mu}} = \frac{1}{2}(\hat{\boldsymbol{\mu}}_0 + \hat{\boldsymbol{\mu}}_1)$, $\hat{\mathbf{C}} = \frac{n_0-1}{n-2} \hat{\mathbf{C}}_0 + \frac{n_1-1}{n-2} \hat{\mathbf{C}}_1$, and we implicitly assumed that \mathcal{H}_0 and \mathcal{H}_1 are equally probable. As for QDA, it instead fully accounts for the possible difference between \mathbf{C}_0 and \mathbf{C}_1 , and the corresponding test is³

$$\begin{aligned}T_{\text{QDA}}(\mathbf{x}) &\equiv -\frac{1}{2}(\mathbf{x} - \hat{\boldsymbol{\mu}}_0)^\top \hat{\mathbf{C}}_0^{-1} (\mathbf{x} - \hat{\boldsymbol{\mu}}_0) + \frac{1}{2}(\mathbf{x} - \hat{\boldsymbol{\mu}}_1)^\top \hat{\mathbf{C}}_1^{-1} (\mathbf{x} - \hat{\boldsymbol{\mu}}_1) \\ &+ \frac{1}{2} \log \frac{|\hat{\mathbf{C}}_0|}{|\hat{\mathbf{C}}_1|} \stackrel{\mathcal{H}_0}{\gtrless} \stackrel{\mathcal{H}_1}{\gtrless} 0.\end{aligned}$$

Of course, due to the presence of the matrix inverses $\hat{\mathbf{C}}_0^{-1}$ and $\hat{\mathbf{C}}_1^{-1}$, these estimators are only defined (almost surely) for $n_0, n_1 \geq p$. If this condition is

²To retrieve this result, it suffices to verify that $\mathbb{P}_{\mathbf{x} \sim \mathcal{N}(\hat{\boldsymbol{\mu}}_0, \hat{\mathbf{C}})}(\mathbf{x}) \gtrless \mathbb{P}_{\mathbf{x} \sim \mathcal{N}(\hat{\boldsymbol{\mu}}_1, \hat{\mathbf{C}})}(\mathbf{x})$ is equivalent to $(\mathbf{x} - \hat{\boldsymbol{\mu}})^\top \hat{\mathbf{C}}^{-1} (\hat{\boldsymbol{\mu}}_0 - \hat{\boldsymbol{\mu}}_1) \gtrless 0$.

³Here, as opposed to the LDA setting where $\hat{\mathbf{C}}$ is common to both hypotheses, the comparison $\mathbb{P}_{\mathbf{x} \sim \mathcal{N}(\hat{\boldsymbol{\mu}}_0, \hat{\mathbf{C}}_0)}(\mathbf{x}) \gtrless \mathbb{P}_{\mathbf{x} \sim \mathcal{N}(\hat{\boldsymbol{\mu}}_1, \hat{\mathbf{C}}_1)}(\mathbf{x})$ maintains the determinants $|\hat{\mathbf{C}}_0|$ and $|\hat{\mathbf{C}}_1|$ from the Gaussian density formula.

not met, the estimates of $\hat{\mathbf{C}}_\ell$ are generally regularized as $\hat{\mathbf{C}}_\ell^{(\gamma)} \equiv \hat{\mathbf{C}}_\ell + \gamma \mathbf{I}_p$ for some $\gamma > 0$. As a matter of fact, even when $n_0, n_1 \geq p$, the large conditioning of the empirical inverses (i.e., the ratio between their largest to smallest eigenvalues) significantly degrades the performance and imposes this regularization in practice.

The objective of this section is to analyze the impact of a large dimensional assumption on n_0, n_1, p on the performance of (regularized) LDA and QDA.

In a nutshell, and quite surprisingly at first sight, we will observe that LDA almost systematically outperforms QDA, even when $\mathbf{C}_0 \neq \mathbf{C}_1$: specifically, the minimal regime for $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\|$, when seen as a function of p , under which hypotheses \mathcal{H}_0 and \mathcal{H}_1 can be discriminated is $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| \geq O(1)$ for LDA but only $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| \geq O(\sqrt{p})$ for QDA. In other words, quite paradoxically, in possibly wrongly assuming that $\mathbf{C}_0 = \mathbf{C}_1$, LDA is in general capable to discriminate \mathcal{H}_0 from \mathcal{H}_1 when $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| = O(1)$, where QDA must fail.

This remark is all the more counterintuitive that we will see later in Section 4.5.3 that a mere least-squares regression method, which does not even need to know that the data are Gaussian distributed, will in the present setting outperform QDA. This fundamental statement must be understood as follows: the performance gain induced by a perfect modeling of the data statistics (Neyman-Pearson test over two Gaussian hypotheses when the data are genuinely Gaussian) is insufficient to outweigh the huge loss incurred by the inappropriate estimation of \mathbf{C}_ℓ by $\hat{\mathbf{C}}_\ell$; this entails that ill-matched procedures (not fitting the genuine Gaussian nature of the data, but most importantly not trying to estimate and invert the data population covariance) may perform better.

In the following two sections, we first provide a full account of the large dimensional behavior and performance of regularized LDA. We will then only justify the main reasons behind the comparatively poor performance of QDA, or at least its inability to perform at the same optimal $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| = O(1)$ rate. A detailed analysis of regularized QDA is provided in [Elkhalil et al., 2020].

3.1.2.1 Linear discriminant analysis

The performance of LDA is provided by the statistics $\mathbb{P}(T_{\text{LDA}}(\mathbf{x}) > 0 \mid \mathbf{x} \sim \mathcal{H}_\ell)$ for $\ell \in \{0, 1\}$.

In the large n_0, n_1, p regime where $n_\ell/n \rightarrow c_\ell \in (0, 1)$ and $p/n \rightarrow c \in (0, \infty)$, in order for this quantity to remain non-trivial (i.e., neither converging to 0, 1 or 1/2), some growth rate constraints need be set on the distance $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\|$. Assuming $\|\mathbf{C}_\ell\| = O(1)$ (without generality restriction), it appears in the calculus that this non-trivial regime corresponds to $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| = O(1)$. Recalling Equation 1.7 in Section 1.1.3, this regime happens to be the minimal possible growth rate for detection in the oracle case where $\boldsymbol{\mu}_0$ and $\boldsymbol{\mu}_1$ would be perfectly known; as such, in terms of optimally allowed growth rates for $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\|$, LDA does not loose in performance.

Under this assumption, one then needs to evaluate the statistics of $T_{\text{LDA}}(\mathbf{x})$. This study is performed in [Elkhalil et al., 2020] with a (slightly different form

of) regularization γ . It is precisely shown that

$$T_{\text{LDA}}^{(\gamma)}(\mathbf{x}) = (\mathbf{x} - \hat{\boldsymbol{\mu}})^T [\hat{\mathbf{C}}^{(\gamma)}]^{-1} (\hat{\boldsymbol{\mu}}_0 - \hat{\boldsymbol{\mu}}_1)$$

satisfies a central limit theorem in the large n_0, n_1, p limit. To obtain this limiting behavior of $T_{\text{LDA}}^{(\gamma)}$, let us assume that $\mathbf{x} = \boldsymbol{\mu}_\ell + \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{z} \sim \mathcal{H}_\ell$ and observe that we may write the complete training data set as $\mathbf{X} = [\mathbf{C}_0^{\frac{1}{2}} \mathbf{Z}_0 + \boldsymbol{\mu}_0 \mathbf{1}_{n_0}^\top, \mathbf{C}_1^{\frac{1}{2}} \mathbf{Z}_1 + \boldsymbol{\mu}_1 \mathbf{1}_{n_1}^\top]$ and its ‘‘hypothesis-wise’’ empirically centered version as $\mathbf{X}^\circ = [\mathbf{C}_0^{\frac{1}{2}} \mathbf{Z}_0, \mathbf{C}_1^{\frac{1}{2}} \mathbf{Z}_1] - [\frac{1}{n_0} \mathbf{C}_0^{\frac{1}{2}} \mathbf{Z}_0 \mathbf{1}_{n_0} \mathbf{1}_{n_0}^\top, \frac{1}{n_1} \mathbf{C}_1^{\frac{1}{2}} \mathbf{Z}_1 \mathbf{1}_{n_1} \mathbf{1}_{n_1}^\top]$ (needed to evaluate $\hat{\mathbf{C}}^{(\gamma)}$), with $\mathbf{Z}_0, \mathbf{Z}_1$ having i.i.d. $\mathcal{N}(0, 1)$ entries. Developing $T_{\text{LDA}}^{(\gamma)}(\mathbf{x})$ using these notations, we find

$$T_{\text{LDA}}^{(\gamma)}(\mathbf{x}) = \left(\mathbf{C}_\ell^{\frac{1}{2}} \mathbf{z} + \frac{1}{2} \mathbf{U} \begin{bmatrix} (-1)^\ell \\ (-1)^{\ell+1} \\ -1 \\ -1 \end{bmatrix} \right)^T \mathbf{Q} \mathbf{U} \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix}$$

where

$$\begin{aligned} \mathbf{Q}^{-1} &= \frac{1}{n-2} \mathbf{C}_0^{\frac{1}{2}} \mathbf{Z}_0 \mathbf{Z}_0^\top \mathbf{C}_0^{\frac{1}{2}} + \frac{1}{n-2} \mathbf{C}_1^{\frac{1}{2}} \mathbf{Z}_1 \mathbf{Z}_1^\top \mathbf{C}_1^{\frac{1}{2}} \\ &\quad - \mathbf{U} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \otimes \begin{bmatrix} \frac{n_0}{n-2} & 0 \\ 0 & \frac{n_1}{n-2} \end{bmatrix} \mathbf{U}^\top + \gamma \mathbf{I}_p \\ \mathbf{U} &= \begin{bmatrix} \boldsymbol{\mu}_0 & \boldsymbol{\mu}_1 & \frac{1}{n_0} \mathbf{C}_0^{\frac{1}{2}} \mathbf{Z}_0 \mathbf{1}_{n_0} & \frac{1}{n_1} \mathbf{C}_1^{\frac{1}{2}} \mathbf{Z}_1 \mathbf{1}_{n_1} \end{bmatrix} \end{aligned}$$

and where ‘ \otimes ’ is the Kronecker product.

The matrix \mathbf{Q}^{-1} takes the form of a ‘‘spiked’’ random matrix model, and we may therefore use Woodbury’s identity to isolate the low rank from the large rank parts in \mathbf{Q} as

$$\begin{aligned} \mathbf{Q} &= \mathbf{Q}^\circ + \mathbf{Q}^\circ \mathbf{U} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \otimes \begin{bmatrix} \frac{n_0}{n-2} & 0 \\ 0 & \frac{n_1}{n-2} \end{bmatrix} \\ &\quad \times \left(\mathbf{I}_4 - \mathbf{U}^\top \mathbf{Q}^\circ \mathbf{U} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \otimes \begin{bmatrix} \frac{n_0}{n-2} & 0 \\ 0 & \frac{n_1}{n-2} \end{bmatrix} \right)^{-1} \mathbf{U}^\top \mathbf{Q}^\circ \end{aligned}$$

in which $\mathbf{Q}^\circ = \mathbf{Q}^\circ(-\gamma) = (\frac{1}{n-2} \mathbf{C}_0^{\frac{1}{2}} \mathbf{Z}_0 \mathbf{Z}_0^\top \mathbf{C}_0^{\frac{1}{2}} + \frac{1}{n-2} \mathbf{C}_1^{\frac{1}{2}} \mathbf{Z}_1 \mathbf{Z}_1^\top \mathbf{C}_1^{\frac{1}{2}} + \gamma \mathbf{I}_p)^{-1}$. We may then invoke Theorem 2.8 for which we have in particular that

$$\begin{aligned} \mathbf{Q}^\circ(z) &\leftrightarrow \bar{\mathbf{Q}}^\circ(z) \equiv -\frac{1}{z} \left(\mathbf{I}_p + \sum_{\ell=0}^1 c_\ell \tilde{g}_\ell(z) \mathbf{C}_\ell \right)^{-1} \\ \tilde{\mathbf{Q}}^\circ(z) &\leftrightarrow \bar{\tilde{\mathbf{Q}}}^\circ(z) \equiv \text{diag}(\{\tilde{g}_\ell(z) \mathbf{1}_{n_\ell}\}_{\ell=0}^1) \end{aligned}$$

where $(g_\ell(z), \tilde{g}_\ell(z))_{\ell=0}^1$ are solutions to

$$g_\ell(z) = \frac{1}{n} \text{tr} \mathbf{C}_\ell \bar{\mathbf{Q}}^\circ(z), \quad \tilde{g}_\ell(z) = -\frac{1}{z} \frac{1}{1 + g_\ell(z)}.$$

Developing $T_{\text{LDA}}^{(\gamma)}$, multiple instances of the form $\mathbf{U}^\top \mathbf{Q}^\circ \mathbf{U}$ appear, which can be expressed in the large n_0, n_1, p limit as

$$\mathbf{U}^\top \mathbf{Q}^\circ \mathbf{U} = \begin{bmatrix} \mu_0^\top \mathbf{Q}^\circ \mu_0 & \mu_0^\top \mathbf{Q}^\circ \mu_1 & 0 & 0 \\ \mu_1^\top \mathbf{Q}^\circ \mu_0 & \mu_1^\top \mathbf{Q}^\circ \mu_1 & 0 & 0 \\ 0 & 0 & \frac{1-\gamma\tilde{g}_0(-\gamma)}{c_0} & 0 \\ 0 & 0 & 0 & \frac{1-\gamma\tilde{g}_1(-\gamma)}{c_1} \end{bmatrix} + o(1).$$

Plugging this result into the expression of $T_{\text{LDA}}^{(\gamma)}(\mathbf{x})$, we find that in the large n_0, n_1, p limit,

$$\begin{aligned} T_{\text{LDA}}^{(\gamma)}(\mathbf{x}) &= \frac{(-1)^\ell}{2} (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1)^\top \bar{\mathbf{Q}}^\circ (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1) - \frac{1}{2} g_0(-\gamma) + \frac{1}{2} g_1(-\gamma) \\ &\quad + \mathbf{z}^\top \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Q}^\circ \mathbf{U} \begin{bmatrix} 1 \\ -1 \\ \frac{1}{\gamma\tilde{g}_0(-\gamma)} \\ -\frac{1}{\gamma\tilde{g}_1(-\gamma)} \end{bmatrix} + o(1) \end{aligned}$$

where we used in particular the fact that $\frac{1-\gamma\tilde{g}_0(-\gamma)}{\gamma\tilde{g}_0(-\gamma)} = g_0(-\gamma)$.

Remark 3.1 (Optimal decision threshold). *Since $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, it is clear that $\mathbb{E}[T_{\text{LDA}}^{(\gamma)}(\mathbf{x})]$ is dominated by the quantity $\pm(\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1)^\top \bar{\mathbf{Q}}^\circ (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1)$ which is positive when $\ell = 0$ and negative when $\ell = 1$, as expected. Yet, the term $\frac{1}{2}(g_1(-\gamma) - g_0(-\gamma))$ intervenes as a bias. If $\mathbf{C}_0 = \mathbf{C}_1$, then $g_0 = g_1$ and this bias disappears; however, for $\mathbf{C}_0, \mathbf{C}_1$ distinct, this bias remains and must be accounted for in the decision threshold which, therefore, should not be zero.*

In passing, note that the first three terms in the expansion of $T_{\text{LDA}}^{(\gamma)}(\mathbf{x})$ are of order $O(1)$ with respect to p , while the fourth term is Gaussian conditionally to \mathbf{X} and of variance also of order $O(1)$ (see below for detail). This thus justifies the need for $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\|$ to be of order $O(1)$ (if instead $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| = O(p^t)$ for $t > 0$, the first term dominates and $T_{\text{LDA}}^{(\gamma)}(\mathbf{x})$ becomes deterministic: the decision is trivial; while if $t < 0$ the first term vanishes when compared to the other three and the decision is asymptotically equivalent to a random guess).

To now estimate the variance of $T_{\text{LDA}}^{(\gamma)}(\mathbf{x})$ (which will provide the LDA performance), one now needs to evaluate the second order moment of the term involving z , that is

$$\text{Var}(T_{\text{LDA}}^{(\gamma)}(\mathbf{x})) = \begin{bmatrix} 1 \\ -1 \\ \frac{1}{\gamma\tilde{g}_0(-\gamma)} \\ \frac{-1}{\gamma\tilde{g}_1(-\gamma)} \end{bmatrix}^\top \mathbf{U}^\top \mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ \mathbf{U} \begin{bmatrix} 1 \\ -1 \\ \frac{1}{\gamma\tilde{g}_0(-\gamma)} \\ \frac{-1}{\gamma\tilde{g}_1(-\gamma)} \end{bmatrix} + o(1).$$

To proceed, the deterministic equivalent for \mathbf{Q}° is not sufficient and we must resort to a deterministic equivalent for $\mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ$ (similar to Exercise 13).

This result was derived in [Benaych-Georges and Couillet, 2016] and states

$$\mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ \leftrightarrow \overline{\mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ} \equiv \bar{\mathbf{Q}}^\circ \mathbf{C}_\ell \bar{\mathbf{Q}}^\circ + \bar{\mathbf{Q}}^\circ (R_{0\ell} \mathbf{C}_0 + R_{1\ell} \mathbf{C}_1) \bar{\mathbf{Q}}^\circ$$

$$\text{with } R_{ij} = \frac{c_i}{c_j} [(\mathbf{I}_2 - \mathbf{S})^{-1} \mathbf{S}]_{i+1,j+1}, [\mathbf{S}]_{i+1,j+1} = c_j \gamma^2 \tilde{g}_i(-\gamma)^2 \frac{1}{n} \operatorname{tr} \mathbf{C}_i \bar{\mathbf{Q}}^\circ \mathbf{C}_j \bar{\mathbf{Q}}^\circ.$$

This deterministic equivalent immediately provides a deterministic approximation for the upper-left 2×2 submatrix of $\mathbf{U}^\top \mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ \mathbf{U}$. As for the off-diagonal 2×2 blocks, they are easily seen to vanish. Finally, the bottom-right 2×2 matrix involves the inner products

$$\frac{1}{n_\ell^2} \mathbf{1}_{n_\ell}^\top \mathbf{Z}_\ell^\top \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ \mathbf{C}_{\ell'}^{\frac{1}{2}} \mathbf{Z}_{\ell'} \mathbf{1}_{n_{\ell'}}.$$

By asymmetry, this is non-vanishing only for $\ell = \ell'$. And when $\ell = \ell'$, one must be careful in evaluating the quadratic form as \mathbf{Z}_ℓ and \mathbf{Q}° are not independent. To deal with this case, we may write,

$$\mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Z}_\ell \mathbf{Z}_\ell^\top \mathbf{C}_\ell^{\frac{1}{2}} = \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Z}_\ell \left(\mathbf{I}_{n_\ell} - \frac{1}{n_\ell} \mathbf{1}_{n_\ell} \mathbf{1}_{n_\ell}^\top \right) \mathbf{Z}_\ell \mathbf{C}_\ell^{\frac{1}{2}} + \frac{1}{n_\ell^2} \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Z}_\ell \mathbf{1}_{n_\ell} \mathbf{1}_{n_\ell}^\top \mathbf{Z}_\ell^\top \mathbf{C}_\ell^{\frac{1}{2}}$$

in which the columns of $\mathbf{Z}_\ell (\mathbf{I}_{n_\ell} - \frac{1}{n_\ell} \mathbf{1}_{n_\ell} \mathbf{1}_{n_\ell}^\top)$ and $\mathbf{Z}_\ell \frac{1}{n_\ell} \mathbf{1}_{n_\ell}$ are orthogonal and thus independent Gaussian vectors. As such, with a rank-one perturbation argument,

$$\mathbf{1}_{n_\ell}^\top \mathbf{Z}_\ell^\top \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Q}^\circ = \frac{\mathbf{1}_{n_\ell}^\top \mathbf{Z}_\ell^\top \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Q}_{-\ell}^\circ}{1 + \frac{1}{n_\ell(n-2)} \mathbf{1}_{n_\ell}^\top \mathbf{Z}_\ell^\top \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Q}_{-\ell}^\circ \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Z}_\ell \mathbf{1}_{n_\ell}}$$

with $\mathbf{Q}_{-\ell}^\circ$ the matrix \mathbf{Q}° with contribution from $\frac{1}{n_\ell^2} \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Z}_\ell \mathbf{1}_{n_\ell} \mathbf{1}_{n_\ell}^\top \mathbf{Z}_\ell^\top \mathbf{C}_\ell^{\frac{1}{2}}$ discarded. By the detoured manner to induce independence, we may now apply the trace lemma, Lemma 2.11, to obtain

$$\begin{aligned} \frac{1}{n_\ell^2} \mathbf{1}_{n_\ell}^\top \mathbf{Z}_\ell^\top \mathbf{C}_\ell^{\frac{1}{2}} \mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ \mathbf{C}_{\ell'}^{\frac{1}{2}} \mathbf{Z}_{\ell'} \mathbf{1}_{n_{\ell'}} &= \delta_{\ell\ell'} \frac{\frac{1}{n_\ell} \operatorname{tr} (\mathbf{C}_\ell \mathbf{Q}^\circ)^2}{(1 + \frac{1}{n} \operatorname{tr} \mathbf{C}_\ell \mathbf{Q}^\circ)^2} + o(1) \\ &= \delta_{\ell\ell'} \gamma^2 \tilde{g}_\ell(-\gamma)^2 \frac{1}{n_\ell} \operatorname{tr} \mathbf{C}_\ell \overline{\mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ} + o(1). \end{aligned}$$

Plugging these results in the expression of the variance, we thus conclude that, for $\mathbf{x} \sim \mathcal{H}_\ell$,

$$\begin{aligned} \operatorname{Var}(T_{\text{LDA}}^{(\gamma)}(\mathbf{x})) &= (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1)^\top \overline{\mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ} (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1) \\ &\quad + \frac{1}{n_0} \operatorname{tr} \mathbf{C}_0 \overline{\mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ} + \frac{1}{n_1} \operatorname{tr} \mathbf{C}_1 \overline{\mathbf{Q}^\circ \mathbf{C}_\ell \mathbf{Q}^\circ}. \end{aligned}$$

We finally conclude that, for a decision threshold $\xi \in \mathbb{R}$,

$$\begin{aligned} & \mathbb{P}\left(T_{\text{LDA}}^{(\gamma)}(\mathbf{x}) > \xi \mid \mathbf{x} \sim \mathcal{H}_\ell\right) \\ &= Q\left(\frac{\xi - \frac{1}{2} [(-1)^\ell (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1)^\top \bar{\mathbf{Q}}^\circ (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1) - g_0(-\gamma) + g_1(-\gamma)]}{\sqrt{(\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1)^\top \bar{\mathbf{Q}}^\circ \mathbf{C}_\ell \bar{\mathbf{Q}}^\circ (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1) + \text{tr}\left(\frac{\mathbf{C}_0}{n_0} + \frac{\mathbf{C}_1}{n_1}\right) \bar{\mathbf{Q}}^\circ \mathbf{C}_\ell \bar{\mathbf{Q}}^\circ}}\right) + o(1) \end{aligned}$$

where $Q(t) = \frac{1}{\sqrt{2\pi}} \int_t^\infty e^{-u^2/2} du$ is the Gaussian Q -function. This expression emphasizes again the optimality of the decision threshold $\xi = \frac{1}{2}g_1(-\gamma) - \frac{1}{2}g_0(-\gamma)$ pointed out in Remark 3.1.

Figure 3.2 depicts the histograms of the output of $T_{\text{LDA}}^{(\gamma)}(\mathbf{x})$ versus theory in the practical case of Fashion-MNIST and Kannada-MNIST data (thus not Gaussian vectors but realistic data!), here for $\gamma = 1$ and $n_0 = n_1 = 1024$ samples per class. The optimal decision threshold, which assumes instead that the data are genuinely Gaussian, appears to be quite close to $\xi = 0$ in this balanced training sample setting, thereby suggesting that the quantities $g_0(-\gamma) = \frac{1}{n} \text{tr} \mathbf{C}_0 \bar{\mathbf{Q}}^\circ(-\gamma)$ and $g_1(-\gamma) = \frac{1}{n} \text{tr} \mathbf{C}_1 \bar{\mathbf{Q}}^\circ(-\gamma)$ are very similar. This does not necessarily implies that \mathbf{C}_0 and \mathbf{C}_1 are similar, but rather that their “structures” are similar so that $g_0(-\gamma) \approx g_1(-\gamma)$.

It is quite remarkable to observe that the empirical performance obtained on the real MNIST is strongly coincident with the Gaussian theory. This observation (that the algorithm performance on real data is an astounding match to Gaussian predictions) will follow us far in our progression to the successive applications of random matrix theory to practical machine learning algorithm, before being theoretically justified in Chapter 8.

We complete this study of LDA by importantly mentioning that, in practice, the quantities $g_\ell(-\gamma)$ are simple to estimate. Indeed, it suffices to notice that, from the trace lemma (Lemma 2.11), for every sample $\mathbf{x}_i \sim \mathcal{H}_\ell$ of the training set,

$$\frac{1}{p} (\mathbf{x}_i - \hat{\boldsymbol{\mu}}_\ell)^\top [\hat{\mathbf{C}}_{-i}^{(\gamma)}]^{-1} (\mathbf{x}_i - \hat{\boldsymbol{\mu}}_\ell) = g_\ell(-\gamma) + O(p^{-\frac{1}{2}})$$

where $\hat{\mathbf{C}}_{-i}^{(\gamma)} = \hat{\mathbf{C}} - \frac{1}{n-2} \mathbf{x}_i \mathbf{x}_i^\top + \gamma \mathbf{I}_p$. Or equivalently, using the rank-one perturbation lemma,

$$\frac{\frac{1}{p} (\mathbf{x}_i - \hat{\boldsymbol{\mu}}_\ell)^\top [\hat{\mathbf{C}}^{(\gamma)}]^{-1} (\mathbf{x}_i - \hat{\boldsymbol{\mu}}_\ell)}{1 - \frac{1}{n-2} (\mathbf{x}_i - \hat{\boldsymbol{\mu}}_\ell)^\top [\hat{\mathbf{C}}^{(\gamma)}]^{-1} (\mathbf{x}_i - \hat{\boldsymbol{\mu}}_\ell)} = g_\ell(-\gamma) + O(p^{-\frac{1}{2}})$$

which, averaging over $i = 1, \dots, n_\ell$, gives the even more accurate estimate

$$\frac{1}{n_\ell} \sum_{i=1}^{n_\ell} \frac{\frac{1}{p} (\mathbf{x}_i^{(\ell)} - \hat{\boldsymbol{\mu}}_\ell)^\top [\hat{\mathbf{C}}^{(\gamma)}]^{-1} (\mathbf{x}_i^{(\ell)} - \hat{\boldsymbol{\mu}}_\ell)}{1 - \frac{1}{n-2} (\mathbf{x}_i^{(\ell)} - \hat{\boldsymbol{\mu}}_\ell)^\top [\hat{\mathbf{C}}^{(\gamma)}]^{-1} (\mathbf{x}_i^{(\ell)} - \hat{\boldsymbol{\mu}}_\ell)} = g_\ell(-\gamma) + O(p^{-1}).$$

This thus means that one can practically estimate the detection threshold ξ necessary to achieve a desired level of prediction accuracy. This also means that the hyperparameter γ can be a priori optimized so to maximize a target performance (e.g., maximize over γ the quantity $\mathbb{P}(T_{\text{LDA}}^{(\gamma)}(\mathbf{x}) > \xi(\gamma) \mid \mathbf{x} \sim \mathcal{H}_1)$ given that $\xi = \xi(\gamma)$ is set such that $\mathbb{P}(T_{\text{LDA}}^{(\gamma)}(\mathbf{x}) > \xi(\gamma) \mid \mathbf{x} \sim \mathcal{H}_0) \leq \alpha$). In addition to the strong fit between theory and practice on real dataset, this full control and estimation of the performance (even before running the algorithm!) has a strong incidence on the optimal use of LDA for real data classification.




Figure 3.2: Empirical histogram of $T_{\text{LDA}}^{(\gamma)}(\mathbf{x})$ versus the Gaussian limiting prediction (dashed), \mathcal{H}_0 in **blue** and \mathcal{H}_1 in **red**, $n_0 = n_1 = 1024$, $p = 784$, $\gamma = 0.1$, for Fashion-MINST (**left**) and Kannada-MNIST (**right**) data, class 3 versus 4. Empirical results averaged over 30 runs on test sets of size 128. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

3.1.2.2 Quadratic discriminant analysis

To best understand the large dimensional behavior of (regularized) QDA and to fine-tune the non-trivial assumptions on $\boldsymbol{\mu}_0, \boldsymbol{\mu}_1$ and $\mathbf{C}_0, \mathbf{C}_1$, an important preliminary step of order of magnitude estimation is needed.

Under the regularized setting, let us define

$$\begin{aligned} T_{\text{QDA}}^{(\gamma)}(\mathbf{x}) \equiv & \frac{1}{2\sqrt{p}} \log \frac{|\hat{\mathbf{C}}_0^{(\gamma)}|}{|\hat{\mathbf{C}}_1^{(\gamma)}|} - \frac{1}{2\sqrt{p}} (\mathbf{x} - \hat{\boldsymbol{\mu}}_0)^T [\hat{\mathbf{C}}_0^{(\gamma)}]^{-1} (\mathbf{x} - \hat{\boldsymbol{\mu}}_0) \\ & + \frac{1}{2\sqrt{p}} (\mathbf{x} - \hat{\boldsymbol{\mu}}_1)^T [\hat{\mathbf{C}}_1^{(\gamma)}]^{-1} (\mathbf{x} - \hat{\boldsymbol{\mu}}_1) \end{aligned}$$

where $\hat{\mathbf{C}}_\ell^{(\gamma)} = \hat{\mathbf{C}}_\ell + \gamma \mathbf{I}_p$, and the division by $1/\sqrt{p}$ is chosen here so that $T_{\text{QDA}}^{(\gamma)}(\mathbf{x})$ be of order $O(1)$ in the non-trivial regime, as we shall see.

First observe that $T_{\text{QDA}}^{(\gamma)}(\mathbf{x})$ is the sum of two quadratic forms $(\frac{1}{2\sqrt{p}}(\mathbf{x} - \hat{\boldsymbol{\mu}}_\ell)^\top [\hat{\mathbf{C}}_\ell^{(\gamma)}]^{-1}(\mathbf{x} - \hat{\boldsymbol{\mu}}_\ell))$ and of two linear statistics $(\frac{1}{\sqrt{p}} \log |\hat{\mathbf{C}}_\ell^{(\gamma)}|)$ of the resolvent of large dimensional random matrices. Under the present $1/\sqrt{p}$ normalization, it is not difficult to see that the leading order of the quadratic forms is $O(\sqrt{p})$ while their fluctuations of order $O(1)$; as for the linear statistics, their means are of order $O(\sqrt{p})$ are their fluctuations of order $O(1/\sqrt{p})$ (recall from the discussion in Section 2.6.3 that linear statistics have a fast convergence rate with central limit theorems of speed $O(1/\sqrt{pn}) = O(1/p)$).

As such, if \mathbf{C}_0 and \mathbf{C}_1 are sufficiently ‘distinct’ in the large n, p regime, that is $\|\mathbf{C}_0 - \mathbf{C}_1\| \geq O(1)$, the sum of these means remains of order $O(\sqrt{p})$ and the fluctuations at most of order $O(1)$: the (random) $T_{\text{QDA}}^{(\gamma)}(\mathbf{x})$ is asymptotically deterministic and the problem becomes trivially easy. One thus needs a stronger assumption on the difference $\|\mathbf{C}_0 - \mathbf{C}_1\|$ for the decision not to be trivial. Not surprisingly, since the objective is to lower the dominant order by a factor $1/\sqrt{p}$, we must thus demand here that

$$\|\mathbf{C}_0 - \mathbf{C}_1\| = O(1/\sqrt{p}).$$

Yet, due to the independence of \mathbf{Z}_0 and \mathbf{Z}_1 , and the fact that p/n remains away from zero, this condition still implies that

$$\|\hat{\mathbf{C}}_0 - \hat{\mathbf{C}}_1\| = O(1)$$

which holds even when $\mathbf{C}_0 = \mathbf{C}_1$. To see why $\|\hat{\mathbf{C}}_0 - \hat{\mathbf{C}}_1\|$ must be $O(1)$, it suffices to note that, for say $\mathbf{C}_0 = \mathbf{C}_1 = \mathbf{I}_p$, $\hat{\mathbf{C}}_0 - \hat{\mathbf{C}}_1 = [\mathbf{Z}_0, \mathbf{Z}_1] \text{diag}(\frac{1}{n_0} \mathbf{1}_{n_0}^\top, \frac{1}{n_1} \mathbf{1}_{n_1}^\top) [\mathbf{Z}_0, \mathbf{Z}_1]^\top$, the eigenvalues of which are known from Theorem 2.6 to be of order $O(1)$.

As a consequence of this critical remark, observe that in the case $\mathbf{C}_0 = \mathbf{C}_1 = \mathbf{I}_p$, for say $\mathbf{x} = \boldsymbol{\mu}_0 + \mathbf{z}$ with $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$,

$$\begin{aligned} & (\mathbf{x} - \hat{\boldsymbol{\mu}}_1)^\top [\hat{\mathbf{C}}_1^{(\gamma)}]^{-1} (\mathbf{x} - \hat{\boldsymbol{\mu}}_1) - (\mathbf{x} - \hat{\boldsymbol{\mu}}_0)^\top [\hat{\mathbf{C}}_0^{(\gamma)}]^{-1} (\mathbf{x} - \hat{\boldsymbol{\mu}}_0) \\ &= \left(\mathbf{z} + \boldsymbol{\mu}_0 - \boldsymbol{\mu}_1 - \frac{1}{n_0} \mathbf{Z}_0 \mathbf{1}_{n_0} \right)^\top [\hat{\mathbf{C}}_1^{(\gamma)}]^{-1} \left(\mathbf{z} + \boldsymbol{\mu}_0 - \boldsymbol{\mu}_1 - \frac{1}{n_1} \mathbf{Z}_1 \mathbf{1}_{n_1} \right) \\ &\quad - \left(\mathbf{z} - \frac{1}{n_0} \mathbf{Z}_0 \mathbf{1}_{n_0} \right)^\top [\hat{\mathbf{C}}_0^{(\gamma)}]^{-1} \left(\mathbf{z} - \frac{1}{n_1} \mathbf{Z}_1 \mathbf{1}_{n_1} \right). \end{aligned}$$

Note that $\|\frac{1}{n_\ell} \mathbf{Z}_\ell \mathbf{1}_{n_\ell}\| = O(1)$ which is thus negligible when compared to $\|\mathbf{z}\| = O(\sqrt{p})$. Further developing, we find that, if $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| = O(1)$, the dominant term is

$$\mathbf{z}^\top [\hat{\mathbf{C}}_1^{(\gamma)}]^{-1} (\hat{\mathbf{C}}_0 - \hat{\mathbf{C}}_1) [\hat{\mathbf{C}}_0^{(\gamma)}]^{-1} \mathbf{z} \tag{3.2}$$

which is of order $O(p)$, while the informative means-discriminating term

$$(\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1)^\top [\hat{\mathbf{C}}_1^{(\gamma)}]^{-1} (\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1) \tag{3.3}$$

is of order $O(1)$ and thus negligible. When in particular $\mathbf{C}_0 = \mathbf{C}_1$ and $n_0 = n_1$, while $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| = O(1)$, the dominant term of Equation (3.2) is a random noise term of arbitrary sign, and thus leads the asymptotic detection performance to be no better than random guess.

To avoid this trivial scenario it is thus required to let

$$\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| = O(\sqrt{p})$$

thereby turning the informative term of Equation (3.3) to be comparable with the term of Equation (3.2). Note however that, if \mathbf{C}_0 and \mathbf{C}_1 had been perfectly known, letting $\hat{\mathbf{C}}_\ell = \mathbf{C}_\ell$, the dominant term in Equation (3.2) would vanish and detection would be achievable at the optimal $\|\boldsymbol{\mu}_0 - \boldsymbol{\mu}_1\| = O(1)$ rate. So the key limiting factor of QDA is effectively due to the inappropriate estimation of population covariance matrices by their sample counterparts.

Being largely suboptimal (when compared to simpler methods developed in the course of the monograph, such as the LS-SVM approach to be discussed in Section 4.5.3) and bringing little additional intuition, we do not further expose the technical development of the regularized QDA performance. An exhaustive account is proposed in [Elkhalil et al., 2020]. It is worth mentioning that, while not solving this deleterious problem, in a further work [Bejaoui et al., 2020], the authors manage to obtain significant improvement to QDA by applying different regularizations $\gamma_0 \neq \gamma_1$ to the sample covariance matrices $\hat{\mathbf{C}}_0^{(\gamma_0)}$ and $\hat{\mathbf{C}}_1^{(\gamma_1)}$ (particularly in the case of unbalanced classification).

In the next section, we move away binary hypothesis testing to consider spectral-based estimation problems in statistics, starting with the popular subspace estimation methods.

3.1.3 Subspace methods: the G-MUSIC algorithm

In several applied contexts, such as in array processing, or brain signal processing, the statistical covariance of a sequence of multivariate observations in \mathbb{R}^p testifies of specific “directions of arrival” of a sought signal (arising from radar bounces in array processing, or brain regions in brain signal processing). In these scenarios, the covariance matrix is quite structured and, if few (say $k \ll p$) distinct signals, or directions of arrival, are to be retrieved (compared to the dimension p of the data collecting array), this population covariance matrix is both structured and of a “spiked model type”.

The so-called subspace methods are used to retrieve this structural information in the covariance, and to infer the directions of arrival, from the dominant eigenvectors of the sample covariance matrix. These methods are known to perform well only when the typical angular distance between the angles $\theta_1, \dots, \theta_k$ to be estimated is sufficiently large, but fail to be discriminative otherwise.

We shall see in this section that these algorithms, and particularly the most popular of them – the MUSIC algorithm [Schmidt, 1986] –, are again based on the assumption that the population covariance can be consistently estimated by the sample covariance matrix. Paradoxically, this approximation, which we

now know is quite rough and hazardous, will not alter the consistency of the classical MUSIC algorithm in the individual estimation of the angles $\theta_1, \dots, \theta_k$. However, we will see that a random matrix analysis allows for an improvement of MUSIC providing a much more accurate discrimination and estimation of close angles.

3.1.3.1 The MUSIC algorithm

We consider the multivariate data (or signal) model of the form

$$\mathbf{x}_i = \sum_{\ell=1}^k \mathbf{a}(\theta_\ell) s_{\ell,i} + \sigma \mathbf{w}_i$$

for $i \in \{1, \dots, n\}$, where $\mathbf{a}(\theta_\ell) \in \mathbb{R}^p$ is a deterministic normalized ($\|\mathbf{a}(\theta)\| = 1$) ‘steering vector’ parameterized by the scalar angle $\theta_\ell \in (-\pi, \pi]$, $s_{\ell,i} \in \mathbb{R}$ is a deterministic or random signal carried in the direction θ_ℓ at time instant i , $\sigma > 0$ and $\mathbf{w}_i \in \mathbb{R}^p$ is an independent random thermal noise at time i .⁴

Various hypotheses can be formulated on prior knowledge on the signal $s_{\ell,i}$ and its dependence across time instants and across array elements. We consider here the simple setting where $\mathbf{s}_i = [s_{1,i}, \dots, s_{k,i}]^\top \sim \mathcal{N}(\mathbf{0}, \mathbf{P})$ for diagonal $\mathbf{P} \in \mathbb{R}^{k \times k}$ with $\mathbf{P}_{\ell\ell} > 0$ collecting the energy of the sources, and $\mathbf{s}_1, \dots, \mathbf{s}_n$ drawn independently. We also assume that $\mathbf{w}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ are independent and independent of the \mathbf{s}_j ’s. In particular, source ℓ has an associated signal-to-noise ratio (SNR) $\mathbf{P}_{\ell\ell}/\sigma^2$.

As a consequence, we obtain for each i

$$\mathbb{E}[\mathbf{x}_i \mathbf{x}_i^\top] = \sum_{\ell=1}^k \mathbf{P}_{\ell\ell} \mathbf{a}(\theta_\ell) \mathbf{a}(\theta_\ell)^\top + \sigma^2 \mathbf{I}_p = \mathbf{A}_\theta \mathbf{P} \mathbf{A}_\theta^\top + \sigma^2 \mathbf{I}_p$$

where $\mathbf{A}_\theta = [\mathbf{a}(\theta_1), \dots, \mathbf{a}(\theta_k)] \in \mathbb{R}^{p \times k}$.

Let $\mathbb{E}[\mathbf{x}_i \mathbf{x}_i^\top] = \mathbf{U} \Lambda \mathbf{U}^\top$ be the spectral decomposition of $\mathbb{E}[\mathbf{x}_i \mathbf{x}_i^\top]$ with $\mathbf{U} = [\mathbf{u}_1, \dots, \mathbf{u}_p] \in \mathbb{R}^{p \times p}$, and $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_p)$ where $\lambda_1 \geq \dots \geq \lambda_p$. The MUSIC algorithm is fundamentally based on the observation that the sought steering vectors $\mathbf{a}(\theta_1), \dots, \mathbf{a}(\theta_k)$ live in the k -dimensional subspace spanned by the k dominant eigenvectors $\mathbf{U}_S = [\mathbf{u}_1, \dots, \mathbf{u}_k]$ of $\mathbb{E}[\mathbf{x}_i \mathbf{x}_i^\top]$. The $\mathbf{a}(\theta_\ell)$ ’s are therefore all orthogonal to the complementary subspace $\mathbf{I}_p - \mathbf{U}_S \mathbf{U}_S^\top$ and

$$\mathbf{a}(\theta_\ell)^\top (\mathbf{I}_p - \mathbf{U}_S \mathbf{U}_S^\top) \mathbf{a}(\theta_\ell) = 0$$

for $\ell \in \{1, \dots, k\}$. This equality is then turned into a detection criterion for $\theta_1, \dots, \theta_\ell$ since, if the steering vectors $\mathbf{a}(\theta)$ are linearly independent across

⁴In wireless applications, such as radar array processing, \mathbf{x}_i is rather a vector in \mathbb{C}^p than of \mathbb{R}^p , and $\mathbf{a}(\theta)$ is often supposed assumed to be the complex-valued steering vector with entries $\mathbf{a}(\theta)_j = \frac{1}{\sqrt{p}} \exp(i d j \sin(\theta))$ for some real d (this is precisely the case for *uniform linear arrays*). For consistency with the rest of the monograph, we stick here with real notations, although all results hold identically in the complex case.

$(-\pi, \pi]$, we have the equivalence

$$\eta(\theta) \equiv \mathbf{a}(\theta)^T \mathbf{U}_S \mathbf{U}_S^T \mathbf{a}(\theta) = \|\mathbf{a}(\theta)\|^2 = 1 \Leftrightarrow \theta \in \{\theta_1, \dots, \theta_\ell\}.$$

Would $\mathbb{E}[\mathbf{x}_i \mathbf{x}_i^T]$ be perfectly known, the identification criterion for the θ_ℓ 's would then consist in scanning $\eta(\theta)$ over $(-\pi, \pi]$ and extract the k zeros of the function. Being a quadratic form, all other values of $\eta(\theta)$ are positive. In practice, however, the population covariance $\mathbb{E}[\mathbf{x}_i \mathbf{x}_i^T]$ is substituted by the sample estimation $\frac{1}{n} \mathbf{X} \mathbf{X}^T$, where $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ and the criterion consists in retrieving the local minima of

$$\mathbf{a}(\theta)^T (\mathbf{I}_p - \hat{\mathbf{U}}_S \hat{\mathbf{U}}_S^T) \mathbf{a}(\theta) = 1 - \mathbf{a}(\theta)^T \hat{\mathbf{U}}_S \hat{\mathbf{U}}_S^T \mathbf{a}(\theta)$$

or alternatively the local maxima of the spike “MUSIC” estimator

$$\hat{\eta}_{\text{MUSIC}}(\theta) \equiv \mathbf{a}(\theta)^T \hat{\mathbf{U}}_S \hat{\mathbf{U}}_S^T \mathbf{a}(\theta)$$

for $\hat{\mathbf{U}}_S \in \mathbb{R}^{p \times k}$ the collection of eigenvectors associated with the k largest eigenvalues of the sample covariance $\frac{1}{n} \mathbf{X} \mathbf{X}^T$.

In the presence of few samples, i.e., if n is not much larger than p , $\hat{\eta}(\theta)$ is *not* a consistent estimator for $\eta(\theta)$. Surprisingly though, it has been shown (using random matrix arguments) that, despite this inconsistency in estimating $\eta(\theta)$, for non-trivial ratios p/n , the corresponding estimates $\hat{\theta}_1, \dots, \hat{\theta}_k$ of the angles $\theta_1, \dots, \theta_k$ are consistent estimates [Vallet et al. \[2015\]](#). This means that, while $\hat{\eta}(\theta) - \eta(\theta) \not\rightarrow 0$ as $n, p \rightarrow \infty$, we still have that $\arg \max_{\theta \in \partial\theta_\ell} \hat{\eta}(\theta) - \arg \max_{\theta \in \partial\theta_\ell} \eta(\theta) \rightarrow 0$ (where $\partial\theta_\ell$ is a sufficiently small neighborhood of the genuine angle θ_ℓ), i.e., the local maxima of $\hat{\eta}(\theta)$ do asymptotically coincide with the local maxima of $\eta(\theta)$.

This remark possibly explains the widespread usage of the MUSIC algorithm, despite its inherently using an erroneous estimate of $\eta(\theta)$. This ill-estimate however has the major defect of exhibiting only one local minimum when two steering angles appear to be too close to one another (see an illustration in the right display of Figure 3.3), and is thus not able to *resolve* close angles. The next section revisits the large n, p estimation of $\eta(\theta)$ using a spiked covariance matrix approach, thus exploiting the results established in Section 2.5. The section shows that random matrix can effectively improve the MUSIC algorithm (into a so-called G-MUSIC alternative) by providing a consistent estimate for $\eta(\theta)$, with the main advantage to largely increase the resolution power of MUSIC.

The following results are immediate consequences of Section 2.5 but were primarily developed, under a broader scope of assumptions, in a long series of works on the topic (see in particular [\[Mestre, 2008, Loubaton and Vallet, 2011, Vallet et al., 2015\]](#)).

3.1.3.2 Spiked G-MUSIC

Assuming that the ratio p/n between the number of sensors p (antenna array elements, electrodes, etc.) and the number of independent snapshots n of \mathbf{x}_i 's

is not small, estimating \mathbf{U}_S by $\hat{\mathbf{U}}_S$ is quite inappropriate, as $\|\mathbf{U}_S - \hat{\mathbf{U}}_S\| \not\rightarrow 0$ as $p, n \rightarrow \infty$ with a non-trivial p/n ratio. This unfolds directly from $\frac{1}{n}\mathbf{XX}^\top$ not being a consistent estimator for $\mathbb{E}[\mathbf{x}_i \mathbf{x}_i^\top]$.

However, the objective of interest is $\eta(\theta) = \mathbf{a}(\theta)^\top \mathbf{U}_S \mathbf{U}_S^\top \mathbf{a}(\theta)$, which is a quadratic form involving the deterministic vector $\mathbf{a}(\theta)$ and the rank- k matrix $\mathbf{U}_S \mathbf{U}_S^\top \equiv \sum_{i=1}^k \mathbf{u}_{S,i} \mathbf{u}_{S,i}^\top$, with k in general small compared to n, p . This quantity can thus be estimated consistently using the results of Section 2.5 on spiked random matrices.

More precisely, note that $\frac{1}{\sigma^2 n} \mathbf{XX}^\top$ is a Wishart random matrix with $\mathbf{x}_i/\sigma \sim \mathcal{N}(\mathbf{0}, \sigma^{-2} \mathbf{A}_\theta \mathbf{P} \mathbf{A}_\theta^\top + \mathbf{I}_p)$ and $\text{rank}(\sigma^{-2} \mathbf{A}_\theta \mathbf{P} \mathbf{A}_\theta^\top) = k$, which thus falls under the setting of Theorems 2.13 and 2.14. Writing

$$\sigma^{-2} \mathbf{A}_\theta \mathbf{P} \mathbf{A}_\theta^\top = \mathbf{U}_S \mathbf{L}_S \mathbf{U}_S^\top = \sum_{i=1}^k \ell_i \cdot \mathbf{u}_{S,i} \mathbf{u}_{S,i}^\top$$

for some diagonal matrix $\mathbf{L}_S = \text{diag}(\ell_1, \dots, \ell_k) \in \mathbb{R}^{k \times k}$ and assuming that $\ell_i > \sqrt{c}$ for each i (high SNR scenario), by Theorem 2.14, for all deterministic unit-norm vector $\mathbf{a} \in \mathbb{R}^p$, as $p, n \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$,

$$\mathbf{a}^\top \mathbf{U}_S \mathbf{U}_S^\top \mathbf{a} = \sum_{i=1}^k |\mathbf{a}^\top \mathbf{u}_{S,i}|^2 = \sum_{i=1}^k |\mathbf{a}^\top \hat{\mathbf{u}}_{S,i}|^2 \frac{1 + c\ell_i^{-1}}{1 - c\ell_i^{-2}} + o(1).$$

In the specific case of array processing, it is most convenient to assume that $\|\mathbf{a}(\theta)\| = 1$ and $\mathbf{a}(\theta)^\top \mathbf{a}(\theta') \rightarrow 0$ for all fixed $\theta \neq \theta'$ as $p \rightarrow \infty$. This is particularly valid for the canonical example of a “uniform linear array” (that is, an array of sensor evenly spaced on a line). Under this assumption, one can identify

$$\ell_i = \frac{\mathbf{P}_{ii}}{\sigma^2} + o(1)$$

which corresponds to the SNR of the i -th signal source. As a result, we have a first estimator:

$$\sum_{i=1}^k |\mathbf{a}(\theta)^\top \hat{\mathbf{u}}_{S,i}|^2 \frac{1 + \frac{p}{n} \left(\frac{\mathbf{P}_{ii}}{\sigma^2} \right)^{-1}}{1 - \frac{p}{n} \left(\frac{\mathbf{P}_{ii}}{\sigma^2} \right)^{-2}} = \eta(\theta) + o(1).$$

However, \mathbf{P}_{ii}/σ^2 is in general not known and also needs to be estimated. To this end, one may use Theorem 2.13 by noticing that, still under the condition that $\ell_i = \mathbf{P}_{ii}/\sigma^2 + o(1) > \sqrt{c}$, the i -th largest eigenvalue λ_i of $\frac{1}{\sigma^2 n} \mathbf{XX}^\top$ satisfies

$$\lambda_i \xrightarrow{a.s.} 1 + \ell_i + c \frac{1 + \ell_i}{\ell_i} = 1 + c + \frac{\mathbf{P}_{ii}}{\sigma^2} + c \frac{\sigma^2}{\mathbf{P}_{ii}} + o(1).$$

By inverting the expression,

$$\hat{\ell}_i \equiv \frac{\lambda_i - (1 + p/n)}{2} + \frac{1}{2} \sqrt{\left(\lambda_i - \left(1 + \frac{p}{n} \right) \right)^2 - \frac{4p}{n}} = \frac{\mathbf{P}_{ii}}{\sigma^2} + o(1) \quad (3.4)$$

which entails the final spike “G-MUSIC” estimate

$$\hat{\eta}_{\text{GMUSIC}}(\theta) \equiv \sum_{i=1}^k |\mathbf{a}(\theta)^T \hat{\mathbf{u}}_{S,i}|^2 \frac{1 + \frac{p}{n} \hat{\ell}_i^{-1}}{1 - \frac{p}{n} \hat{\ell}_i^{-2}} = \eta(\theta) + o(1)$$

with $\hat{\ell}_i$ given in (3.4) (which is only defined under the condition that $\lambda_i > (1 + \sqrt{p/n})^2$). This approach demands to know σ^2 since λ_i is the i -th largest (and isolated) eigenvalue of $\frac{1}{\sigma^2 n} \mathbf{X} \mathbf{X}^T$. Yet, it is also known that the limiting spectrum of $\frac{1}{\sigma^2 n} \mathbf{X} \mathbf{X}^T$ is the Marčenko-Pastur distribution with right edge equal to (and thus, here, with $(k+1)$ -th largest eigenvalue converging to) $(1 + \sqrt{c})^2 = (1 + \sqrt{p/n})^2 + o(1)$. As such, σ^2 can be estimated from the fact that $\lambda_{k+1}(\frac{1}{n} \mathbf{X} \mathbf{X}^T) = \sigma^2(1 + \sqrt{p/n})^2 + o(1)$. This estimator is however possibly less accurate as λ_{k+1} fluctuates at a rate $O(n^{-2/3})$ by the Tracy-Widom theorem, Theorem 2.15. A better estimate consists in noticing that $\frac{1}{\sigma^2} \frac{1}{p} \text{tr}(\frac{1}{n} \mathbf{X} \mathbf{X}^T) = 1 + O(n^{-1})$, or even more accurately $\frac{1}{\sigma^2} \frac{1}{p-k} \text{tr}(\frac{1}{n} \mathbf{X} \mathbf{X}^T) - \frac{1}{p-k} \sum_{i=1}^k \lambda_i = 1 + O(n^{-1})$ (this second choice discards the isolated signal eigenvalues which in the limit do not contribute but may induce a bias in finite dimension).

The resulting G-MUSIC estimator is nothing more than a “weighted” version of the standard MUSIC algorithm where, instead of projecting $\mathbf{a}(\theta)$ against each of the k dominant eigenvectors of $\frac{1}{n} \mathbf{X} \mathbf{X}^T$, $\mathbf{a}(\theta)$ is now projected against an appropriate weighted sum of these eigenvectors. However, these weighted projections no longer form a projector onto a subspace, and one must be even particularly careful that $1 - \hat{\eta}_{\text{GMUSIC}}(\theta)$ might, with low but non zero probability, be negative (since, unlike MUSIC which enjoys the projector property $\hat{\eta}_{\text{MUSIC}}(\theta) \leq \|\mathbf{a}(\theta)\| = 1$, nothing prevents $\hat{\eta}_{\text{GMUSIC}}(\theta)$ to be greater than 1). This is a common unfortunate artifact of various random matrix approximations of quadratic forms which may result in negative estimates of positive quantities. This has no deleterious consequences here though, as the resulting GMUSIC algorithm looks for the deepest minima (that could then be negative) of $1 - \hat{\eta}(\theta)$.

Figure 3.3 depicts, for $\|\mathbf{a}(\theta)\| = 1$, the performance of the estimator of $1 - \hat{\eta}(\theta)$ (more conventionally used than $\hat{\eta}(\theta)$) for both the classical MUSIC and the improved G-MUSIC algorithms. The ground truth $\eta(\theta)$ has zeros precisely at the location of the genuine angles (here -10° , 35° and 37°). It is observed that deep local minima are precisely exhibited by both MUSIC and G-MUSIC around these positions (on a log scale). However, (i) the minima reached by G-MUSIC are deeper and, most importantly, (ii) the precision of the estimates is largely improved with G-MUSIC, leading in particular to an increased resolution capability of close angles estimation, as observed in the vicinity of the two angles 35° and 37° in the right display of Figure 3.3.

The improved spike G-MUSIC algorithm is a typical example of the possibility to (quite elementarily) improve over a classical and largely used algorithm (MUSIC), long known to suffer from large p/n ratios. This being said, as we illustrated in Figure 2.12 when comparing the accuracy of *spiked* (where k is assumed fixed as p, n increase) versus *non-spiked* (where k is assumed to grow




Figure 3.3: **Left:** MUSIC versus G-MUSIC for $k = 3$ sources, $p = 30$, $n = 150$ samples, $\sigma^2 = 0.1$ (-10 dB). Angles of arrival of 10° , 35° , and 37° . **Right:** zoom on the region of interest of close angles. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

with p, n) estimates, the spiked GMUSIC algorithm would be in fact further improved by considering a non-spiked version of the sample covariance matrix model $\frac{1}{n}\mathbf{XX}^T$ (that is, by considering $\mathbb{E}[\mathbf{x}_i\mathbf{x}_i^T]$ as a generic covariance matrix rather than a low-rank perturbation of the identity). This is the approach carried out originally in [Mestre, 2008], however leading to a more complex (and thus less intuitive) formulation of the GMUSIC estimate of $\eta(\theta)$. Interestingly though, while the spiked-model approach fully “decorrelates” the individual k projection $|\mathbf{a}(\theta)^T \mathbf{u}_{S,i}|^2$ estimates and only exploits the dominant k eigenvectors of the sample covariance, the non-spike approach exploits *all* eigenvalues and eigenvectors in a rather intricate (but in the end more powerful) manner.⁵

3.2 Covariance matrix distance estimation

3.2.1 Distances and divergences between Gaussian laws

Most statistical detection, estimation, and classification methods rely exclusively on the first order statistics of the data model. The various notions of “distances” between classes of data are then strongly related to these first moments.

Since means and covariances are often sufficiently discriminating, especially in large dimensions (see Chapter 4), distances and divergences revolving around Gaussian distributions are of common interest. Among these, the Kullback-Liebler divergence d_{KL} or the Rényi divergence $d_{\alpha R}$ (parameterized by a factor $\alpha \in \mathbb{R}$) between two Gaussian $\mathcal{N}(\mu_1, \mathbf{C}_1)$ and $\mathcal{N}(\mu_2, \mathbf{C}_2)$, as well as Bhattacharyya distance d_B and the Fisher distance d_F (the length of the geodesic in the

⁵To this end, one now needs to use the techniques detailed in Section 2.4.2, and thus deal with questions of bulk separation in the limiting spectrum of $\frac{1}{n}\mathbf{XX}^T$ in place of the phase transitions of spiked models.

“natural” Riemannian space of positive definite matrices) between two covariance matrices \mathbf{C}_1 and \mathbf{C}_2 are among the most popular.

Assuming the observed data vectors have zero mean (or equal mean) and positive definite covariance, these distances and divergences, that we shall generically denote $d(\mathbf{C}_1, \mathbf{C}_2; f)$, share the property that

$$d(\mathbf{C}_1, \mathbf{C}_2; f) = \int f(t) \nu_p(dt)$$

for $\nu_p = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C}_1^{-1}\mathbf{C}_2)}$ and some specific function f . Table 3.1 lists the mappings between distances⁶ and functions f and shows in particular that, in order to evaluate all of these distances, it suffices to assess $\int f(t) \nu_p(dt)$ for $f(t)$ one of the functions $t \mapsto t$, $t \mapsto \log(1+st)$ ($s \in (0, \infty)$), with $\log(t) = \lim_{s \rightarrow \infty} \log(1+st) - \log(s)$) and $t \mapsto \log^2(t)$.

Divergences	$f(z)$
d_F	$\log^2(z)$
d_B	$-\frac{1}{4} \log(z) + \frac{1}{2} \log(1+z) - \frac{1}{2} \log(2)$
d_{KL}	$\frac{1}{2}z - \frac{1}{2} \log(z) - \frac{1}{2}$
$d_{\alpha,R}$	$\frac{-1}{2(\alpha-1)} \log(\alpha + (1-\alpha)z) + \frac{1}{2} \log(z)$

Table 3.1: Distances and divergences, and their corresponding $f(z)$.

Remark 3.2 (The Wasserstein distance). *Of increasing interest in machine learning lately is the Wasserstein distance which, for the laws $\mathcal{N}(\mathbf{0}, \mathbf{C}_1)$ and $\mathcal{N}(\mathbf{0}, \mathbf{C}_2)$, reduces (up to normalization by $1/p$) to*

$$\begin{aligned} d_W(\mathbf{C}_1, \mathbf{C}_2) &= \frac{1}{p} \text{tr}(\mathbf{C}_1) + \frac{1}{p} \text{tr}(\mathbf{C}_2) - \frac{2}{p} \text{tr} \left[(\mathbf{C}_1^{\frac{1}{2}} \mathbf{C}_2 \mathbf{C}_1^{\frac{1}{2}})^{\frac{1}{2}} \right] \\ &= \frac{1}{p} \text{tr}(\mathbf{C}_1) + \frac{1}{p} \text{tr}(\mathbf{C}_2) - 2 \int \sqrt{t} \nu'_p(dt), \quad \nu'_p = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C}_1 \mathbf{C}_2)}. \end{aligned}$$

The Wasserstein distance therefore does not enter the present scheme as it involves the eigenvalues of the product $\mathbf{C}_1 \mathbf{C}_2$ rather than $\mathbf{C}_1^{-1} \mathbf{C}_2$. The same remark holds for the Frobenius distance $d_{Fro}(\mathbf{C}_1, \mathbf{C}_2)^2 = \frac{1}{p} \text{tr}(\mathbf{C}_1 - \mathbf{C}_2)^2$. Yet, the derivations in this section easily extend to this setup. Section 3.2.4 below reports the corresponding results.

3.2.2 The random matrix framework

In practice, to evaluate the “distance” between data classes $\mathcal{C}_1, \dots, \mathcal{C}_k$, one needs access to a certain number n_a of training vectors $\mathbf{X}^{(a)} = [\mathbf{x}_1^{(a)}, \dots, \mathbf{x}_{n_a}^{(a)}] \in \mathbb{R}^{p \times n_a}$ from each class \mathcal{C}_a from which the pairwise distances $d(\mathbf{C}_a, \mathbf{C}_b; f)$ may

⁶We slightly abuse the definitions here as the Fisher and Bhattacharya distances and in fact the square roots of $\int f(t) \nu_p(dt)$ and not the integrals themselves.

be estimated. Assuming $n_a > p$ and the $\mathbf{x}_i^{(a)}$ independent with zero mean (and covariance \mathbf{C}_a), $d(\mathbf{C}_a, \mathbf{C}_b; f)$ is conventionally estimated through the empirical estimate $d(\hat{\mathbf{C}}_a, \hat{\mathbf{C}}_b; f)$ where $\hat{\mathbf{C}}_a = \frac{1}{n_a} \mathbf{X}^{(a)} (\mathbf{X}^{(a)})^\top$.

However, for n_a not much larger than p , $\hat{\mathbf{C}}_a$ is known to be a poor estimator for \mathbf{C}_a and $d(\hat{\mathbf{C}}_a, \hat{\mathbf{C}}_b; f)$ is likely a poor estimator for $d(\mathbf{C}_a, \mathbf{C}_b; f)$. To convince oneself, for say $f(t) = \log(t)$ and $\mathbf{C}_1 = \mathbf{C}_2$, $\mathbf{C}_1^{-1}\mathbf{C}_2 = \mathbf{I}_p$ so that $d(\mathbf{C}_1, \mathbf{C}_2; f) = 0$, while $\hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2$ is distributed as a F-matrix with eigenvalues asymptotically supported on a compact interval around 1 and with left edge converging to zero as $n_1/p > 1$ or $n_2/p > 1$ is close to 1 [Silverstein, 1985], so that $d(\hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2; f)$ may be arbitrarily large as either $n_1/p, n_2/p \downarrow 1$.

The idea of the random matrix framework is to evaluate $d(\mathbf{C}_a, \mathbf{C}_b; f)$ consistently from $\mathbf{X}_a, \mathbf{X}_b$ in the spirit of Section 2.4. Since only pairwise distances are of interest, in the following we focus on evaluating the metric $d(\mathbf{C}_1, \mathbf{C}_2; f)$ for some arbitrary analytic function f .

To this end, similar to (2.29), we introduce the Stieltjes transform $m_{\nu_p}(z)$ of the spectral measure $\nu_p = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C}_1^{-1}\mathbf{C}_2)}$ and write

$$d(\mathbf{C}_1, \mathbf{C}_2; f) = \int f(t) \nu_p(dt) = -\frac{1}{2\pi i} \oint_{\Gamma_\nu} f(z) m_{\nu_p}(z) dz$$

for Γ_ν a contour surrounding the support of ν_p , i.e., surrounding the eigenvalues of $\mathbf{C}_1^{-1}\mathbf{C}_2$, but surrounding none of the singularities of f . Using the finite-dimensional trick that consists in letting $\nu = \nu_p$ be a fictitious asymptotic limit for ν_p as $p \rightarrow \infty$, this becomes

$$d(\mathbf{C}_1, \mathbf{C}_2; f) = -\frac{1}{2\pi i} \oint_{\Gamma_\nu} f(z) m_\nu(z) dz.$$

Similarly, we will in the following denote $c_1 = n_1/p = \lim n_1/p$, $c_2 = n_2/p = \lim n_2/p$ the fictitious limiting data sample/size ratios.

To connect the unknown ν to the observed $\mu_p = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2)}$, we first need to establish a link between m_ν and m_μ , with μ the (almost sure) weak limit of μ_p as $n, p \rightarrow \infty$. For this, it suffices to proceed as follows:

- by Sylvester's identity (Lemma 2.3), $\hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2$ has the same eigenvalues as the symmetric matrix $\hat{\mathbf{C}}_2^{\frac{1}{2}}\hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2^{\frac{1}{2}}$; which are the inverse eigenvalues of $\hat{\mathbf{C}}_2^{-\frac{1}{2}}\hat{\mathbf{C}}_1\hat{\mathbf{C}}_2^{-\frac{1}{2}}$;
- conditioned on \mathbf{X}_2 , $\hat{\mathbf{C}}_2^{-\frac{1}{2}}\hat{\mathbf{C}}_1\hat{\mathbf{C}}_2^{-\frac{1}{2}}$ is a sample covariance matrix with population covariance $\mathbb{E}_{\mathbf{X}_1}[\hat{\mathbf{C}}_2^{-\frac{1}{2}}\hat{\mathbf{C}}_1\hat{\mathbf{C}}_2^{-\frac{1}{2}}] = \hat{\mathbf{C}}_2^{-\frac{1}{2}}\mathbf{C}_1\hat{\mathbf{C}}_2^{-\frac{1}{2}}$, for which Theorem 2.6 provides a deterministic equivalent, as a function of the “deterministic” limiting spectral measure of $\hat{\mathbf{C}}_2^{-\frac{1}{2}}\mathbf{C}_1\hat{\mathbf{C}}_2^{-\frac{1}{2}}$;

- similarly, the matrix $\hat{\mathbf{C}}_2^{-\frac{1}{2}} \mathbf{C}_1 \hat{\mathbf{C}}_2^{-\frac{1}{2}}$ has the same eigenvalues as $\mathbf{C}_1^{\frac{1}{2}} \hat{\mathbf{C}}_2^{-1} \mathbf{C}_1^{\frac{1}{2}}$, which are the inverse eigenvalues of $\mathbf{C}_1^{-\frac{1}{2}} \hat{\mathbf{C}}_2 \mathbf{C}_1^{-\frac{1}{2}}$, the latter being a sample covariance matrix with population covariance $\mathbf{C}_1^{-\frac{1}{2}} \mathbf{C}_2 \mathbf{C}_1^{-\frac{1}{2}}$ for which Theorem 2.6 also establishes the limiting spectrum.

Thus, iterating Theorem 2.6 twice can be shown to result in the following fixed-point system

$$\begin{aligned} m_\nu(-1/m_{\tilde{\zeta}}(z)) &= -zm_{\zeta}(z)m_{\tilde{\zeta}}(z) \\ zm_\mu(z) &= \varphi(z)m_\zeta(\varphi(z)) \end{aligned} \quad (3.5)$$

where $\varphi(z) = z(1 + c_1 zm_\mu(z))$ and $m_{\tilde{\zeta}}(z) = c_2 m_\zeta(z) - (1 - c_2)/z$ for ζ the intermediary limiting spectral measure of $\frac{1}{n_2} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \mathbf{Z}^\top \mathbf{C}^{\frac{1}{2}}$ with $\mathbf{Z} \in \mathbb{R}^{p \times n_2}$ having i.i.d. standard Gaussian entries and $\mathbf{C} = \mathbf{C}_1^{-\frac{1}{2}} \mathbf{C}_2 \mathbf{C}_1^{-\frac{1}{2}}$. While the first of the two equations follows immediately from (2.28), here for a sample covariance matrix model with $\mathbf{C} = \mathbf{C}_1^{-\frac{1}{2}} \mathbf{C}_2 \mathbf{C}_1^{-\frac{1}{2}}$ (of limiting spectrum ν) and $\hat{\mathbf{C}} = \frac{1}{n_2} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \mathbf{Z}^\top \mathbf{C}^{\frac{1}{2}}$ (having the same limiting spectrum ζ as $\mathbf{C}_1^{-1} \hat{\mathbf{C}}_2$), obtaining (3.5) is not so immediate. To this end, first write

$$m_{\zeta^{-1}}(-1/m_{\tilde{\mu}^{-1}}(z)) = -zm_{\mu^{-1}}(z)m_{\tilde{\mu}^{-1}}(z)$$

which relates the Stieltjes transform of the limiting spectral measure of $\hat{\mathbf{C}}_2^{-1} \hat{\mathbf{C}}_1$ (denoted μ^{-1}) to that of $\hat{\mathbf{C}}_2^{-1} \mathbf{C}_1$ (i.e., ζ^{-1} , considered to be the population measure when conditioned on \mathbf{X}_2), with the convention that, for a probability measure θ , θ^{-1} is defined through $\theta^{-1}([a, b]) = \theta([b^{-1}, a^{-1}])$ for $0 < a < b$, and where we used the now standard notation $m_{\tilde{\mu}^{-1}}(z) = c_1 m_{\mu^{-1}}(z) - (1 - c_1)/z$. The result is finally obtained by linking the Stieltjes transform of a measure to that of its inverse, through

$$m_{\zeta^{-1}}(z) = -\frac{1}{z} - \frac{1}{z^2} m_\zeta\left(\frac{1}{z}\right)$$

which is a direct consequence of $m_{\zeta^{-1}}(z) = \int \frac{\zeta(dt)}{t^{-1}-z} = -\frac{1}{z} - \frac{1}{z^2} \int \frac{\zeta(dt)}{t-z^{-1}}$. Stitching these results together, we finally reach (3.5), as desired.

For further need, note that the derivative along z of (3.5) gives

$$m'_\zeta(\varphi(z)) = \frac{1}{\varphi(z)} \left(-\frac{\psi'(z)}{c_2 \varphi'(z)} - m_\zeta(\varphi(z)) \right)$$

where we introduced the function $\psi(z) = 1 - c_2 - c_2 zm_\mu(z)$.

Two successive changes of variable ($\omega \mapsto z = -1/m_{\tilde{\zeta}}(\omega)$ and $u \mapsto \omega = \varphi(u)$) are then needed to relate m_ν first to m_ζ and then m_ζ to m_μ . Assuming the existence of a contour Γ_μ with valid pre-image Γ_ν by these changes of variable,

we find

$$\begin{aligned} d(\mathbf{C}_1, \mathbf{C}_2; f) &= -\frac{1}{2\pi i} \oint_{\Gamma_\nu} f(z) m_\nu(z) dz \\ &= \frac{1}{2\pi i} \oint_{\Gamma_\mu} f\left(\frac{\varphi(u)}{\psi(u)}\right) \frac{\psi(u)}{c_2} \left[\frac{\varphi'(u)}{\varphi(u)} - \frac{\psi'(u)}{\psi(u)}\right] du \\ &\quad - \frac{1-c_2}{c_2} \frac{1}{2\pi i} \oint_{\Gamma_\mu} f\left(\frac{\varphi(u)}{\psi(u)}\right) \left[\frac{\varphi'(u)}{\varphi(u)} - \frac{\psi'(u)}{\psi(u)}\right] du \end{aligned}$$

where we recall that $\varphi(u) = u(1 + c_1 u m_\mu(u))$ and $\psi(u) = 1 - c_2 - c_2 u m_\mu(u)$. Performing the variable changes backwards, the term in the last line writes

$$\begin{aligned} &- \frac{1-c_2}{c_2} \frac{1}{2\pi i} \oint_{\Gamma_\mu} f\left(\frac{\varphi(u)}{\psi(u)}\right) \left[\frac{\varphi'(u)}{\varphi(u)} - \frac{\psi'(u)}{\psi(u)}\right] du \\ &= -\frac{1-c_2}{c_2} \frac{1}{2\pi i} \oint_{\Gamma_\mu} f\left(\frac{\varphi(u)}{\psi(u)}\right) \left(\frac{\varphi(u)}{\psi(u)}\right)^{-1} \left(\frac{\varphi(u)}{\psi(u)}\right)' du \\ &= -\frac{1-c_2}{c_2} \frac{1}{2\pi i} \oint_{\Gamma_\nu} \frac{f(z)}{z} dz. \end{aligned}$$

The contour change analyses performed in Section 2.3.1 are fundamental at this point. Since we here operate twice sample-covariance matrix variable changes, it can be shown that, if $c_1, c_2 < 1$ (i.e., $n_1, n_2 > p$), then any contour $\Gamma_\mu \subset \{z \in \mathbb{C}, \Re[z] > 0\}$ enclosing the support of μ has $\Gamma_\nu \subset \{z \in \mathbb{C}, \Re[z] > 0\}$ as pre-image by the variable changes. Importantly, being subsets of $\{z \in \mathbb{C}, \Re[z] > 0\}$, both Γ_μ and Γ_ν both *exclude* $z = 0$.

Thus, if f is analytic on $\{z \in \mathbb{C}, \Re[z] > 0\}$, we have

$$-\frac{1-c_2}{c_2} \frac{1}{2\pi i} \oint_{\Gamma_\nu} \frac{f(z)}{z} dz = 0.$$

It then suffices to replace the limiting measure μ by its empirical version $\mu_p = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\hat{\mathbf{C}}_1^{-1} \hat{\mathbf{C}}_2)}$ to obtain the final estimate.

Theorem 3.1 (Covariance distance estimate, from [Couillet et al., 2019]). *Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a real function with a complex analytic extension on $\{z \in \mathbb{C}, \Re[z] > 0\}$ and $c_1, c_2 < 1$. Then, with the above notations,⁷*

$$d(\mathbf{C}_1, \mathbf{C}_2; f) - \hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) \xrightarrow{a.s.} 0$$

where

$$\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) = \frac{1}{2\pi i} \oint_{\Gamma_\mu} f\left(\frac{\varphi_p(u)}{\psi_p(u)}\right) \frac{\psi_p(u)}{c_2} \left[\frac{\varphi'_p(u)}{\varphi_p(u)} - \frac{\psi'_p(u)}{\psi_p(u)}\right] du$$

where $\varphi_p(z) = z(1 + c_1 z m_{\mu_p}(z))$ and $\psi_p(z) = 1 - c_2 - c_2 z m_{\mu_p}(z)$.

⁷To avoid too heavy notations, we maintain c_1, c_2 in the empirical estimates (and in the subsequent discussions) but one should in reality replace them systematically with n_1/n and n_2/n .

3.2.3 Closed-form expressions

Theorem 3.1 is quite generic, as valid for any f analytic on $\{z \in \mathbb{C}, \Re[z] > 0\}$. Yet, it practically demands a numerical complex integration procedure and thus conveys little insights on the actual estimate being computed. Yet, most distances of practical interest (recall Table 3.1) involve linear combinations of the functions $f(t) = t$, $f(t) = \log(t)$, $f(t) = \log(1 + st)$ and $f(t) = \log^2(t)$, and it thus suffices to compute these integrals for the complex analytic extensions of these few functions.

Since $m_{\mu_p}(z) = \frac{1}{p} \sum_{i=1}^p (\lambda_i(\hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2) - z)^{-1}$ is a rational function, for f also a rational function, the integrand in Theorem 3.1 is itself a rational function for which residue calculus can be performed. Among the functions above, this is the case only for $f(t) = t$. The other functions (involving logarithms) are “multi-valued” complex functions for which the integral must be computed using more advanced complex analytic calculus.

In all cases, a first requirement is to precisely understand the function $\varphi_p(z)/\psi_p(z)$ on $\{z \in \mathbb{C}, \Re[z] > 0\}$ where f is evaluated. This function can be shown to only have null imaginary part on the real axis and one thus needs to investigate $\varphi_p(z)/\psi_p(z)$ for z real. Also, similar to the proof of Remark 2.12, it can be shown that $\varphi_p(z)$ vanishes exactly at $0 < \eta_1 < \dots < \eta_p$ the eigenvalues of $\mathbf{\Lambda} + \frac{\sqrt{\lambda}\sqrt{\lambda}^\top}{n_1-p}$ while $\psi_p(z)$ vanishes exactly at $0 < \zeta_1 < \dots < \zeta_p$ the eigenvalues of $\mathbf{\Lambda} - \frac{\sqrt{\lambda}\sqrt{\lambda}^\top}{n_2}$, for diagonal $\mathbf{\Lambda} \in \mathbb{R}^{p \times p}$ and $\lambda = (\lambda_1, \dots, \lambda_p)^\top \in \mathbb{R}^p$ the increasingly sorted eigenvalues of $\hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2$ (this follows from Remark 2.12 adapted to the present scenario). Figure 3.4 depicts the function $x \mapsto xm_{\mu_p}(x)$ at the core of the definition of both $\varphi_p(x)$ and $\psi_p(x)$. The ordering of the triplets $\zeta_i < \lambda_i < \eta_i$ is easily established and it appears that $\varphi_p(z)/\psi_p(z)$ is everywhere positive on \mathbb{R}_+ but on the p intervals $[\zeta_i, \eta_i]$. These segments are important as they correspond to *branch cuts* for the multivalued functions $z \mapsto \log^a(\varphi_p(z)/\psi_p(z))$ ($a \in \{1, 2\}$), i.e., they are discontinuity intervals for the function, of central importance to evaluate the sought complex integral.

For $f(t) = t$, a mere residue calculus accounting for the singularities at ζ_i , λ_i and η_i allows one to establish the following simple corollary.

Corollary 3.1 (Case $f(t) = t$). *Under the setting of Theorem 3.1, for $f(t) = t$,*

$$\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) = (1 - c_1) \int t \mu_p(dt).$$

That is, this “simplest” metric ($\int t \nu_p(dt)$) is consistently estimated by a scaled version (with a $1 - c_1$ prefactor) of the standard estimator $\int t \mu_p(dt) = \frac{1}{p} \text{tr } \hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2$. We notably recover the standard large- n estimator when $c_1, c_2 \rightarrow 0$. It may be surprising at first not to see c_2 appearing in this expression; this is explained by the fact that $\frac{1}{p} \text{tr } \mathbf{A}\hat{\mathbf{C}}_2$ is a consistent estimator of $\frac{1}{p} \text{tr } \mathbf{AC}_2$ for all \mathbf{A} of bounded norm, as long as $\liminf n_2/p > 0$ (but the same is not true for $\frac{1}{p} \text{tr } \mathbf{A}\hat{\mathbf{C}}_1^{-1}$ which is not a consistent estimate of $\frac{1}{p} \text{tr } \mathbf{AC}_1^{-1}$).




Figure 3.4: Typical behavior of the function $x \mapsto xm_{\mu_p}(x)$. The Blue bar stresses the range of negative values for $\varphi_p(x)/\psi_p(x)$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

To handle the case $f(t) = \log(t)$, one needs to “deform” the contour Γ_μ to avoid the aforementioned branch cuts. A natural (although admittedly contorted) approach is to proceed as depicted in Figure 3.5 by appending Γ_μ into a circuit surrounding the whole segment $[\zeta_1, \eta_p]$ slightly from above and from below in the complex plane, with small half-circles of vanishing radius around all possible singularities (ζ_i ’s, λ_i ’s and η_i ’s). The whole circuit, call it Γ , has zero integral (as it encompasses no singularity) and is the sum of the desired original integral over Γ_μ , of several line (asymptotically real) integrals, and of half-circle integrals around the poles (evaluated by a variable change $z = \epsilon e^{i\theta}$ and then taking $\epsilon \rightarrow 0$).

The result for the case $f(t) = \log(t)$ again assumes a simple form and is as follows.⁸

Corollary 3.2 (Case $f(t) = \log(t)$). *Under the setting of Theorem 3.1, for $f(t) = \log(t)$,*

$$\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) = \int \log(t) \mu_p(dt) - \frac{1 - c_1}{c_1} \log(1 - c_1) + \frac{1 - c_2}{c_2} \log(1 - c_2).$$

Interestingly, while the case $f(t) = t$ led to a ultimate estimator corresponding to a mere scaling of the large- n estimator, here the estimate is a biased version of the large- n estimator by a constant. Besides, for $c_1 = c_2$, the con-

⁸In effect, the result could have been recovered independently using the much simpler fact that $\int \log(t) \mu_p(dt) = \frac{1}{p} \log |\hat{\mathbf{C}}_1^{-1} \hat{\mathbf{C}}_2| = \frac{1}{p} \log |\hat{\mathbf{C}}_2| - \frac{1}{p} \log |\hat{\mathbf{C}}_2|$; this simple separation trick can however no longer be applied to the functions $\log(1 + st)$ or $\log^2(t)$.




Figure 3.5: Deformed contour Γ to evaluate Theorem 3.1 for $f(z) = \log^a(z)$, $a \in \{1, 2\}$. Branch cuts are displayed in blue bars.

stant vanishes and thus, quite surprisingly, the standard large- n estimator is consistent.

The case $f(t) = \log^2(t)$ is technically more involved to evaluate than $f(t) = \log(t)$. The core of both results lies in the evaluation of the real integrals right above and under the branch cuts (as illustrated in Figure 3.5). The sum of every pair of integrals is of the form $\int_{\zeta_i+\imath 0}^{\eta_i+\imath 0} - \int_{\zeta_i-\imath 0}^{\eta_i-\imath 0} \log^a([\varphi_p/\psi_p](z))g(z)dz$ for some rational function $g(z)$ and $a \in \{1, 2\}$. Using the fact that $\log(\omega) = \log|\omega| + \imath \arg(\omega)$ (i.e., for $a = 1$), $\log([\varphi_p/\psi_p](x + \imath 0)) - \log([\varphi_p/\psi_p](x - \imath 0)) = 2\imath\pi$ and the resulting real integral is thus still a rational function. For $a = 2$ though, $\log^2([\varphi_p/\psi_p](x + \imath 0)) - \log^2([\varphi_p/\psi_p](x - \imath 0)) = 2\imath\pi \log|[\varphi_p/\psi_p](x)|$ and thus the resulting integral involves products of (real) logarithm and rational functions.

After careful calculus, the following corollary is obtained.

Corollary 3.3 (Case $f(t) = \log^2(t)$). *Under the setting of Theorem 3.1, for $f(t) = \log^2(t)$,*

$$\begin{aligned} \hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) &= \frac{c_1 + c_2 - c_1 c_2}{c_1 c_2} \left[\sum_{i=1}^p \log^2((1 - c_1)\eta_i) - \log^2((1 - c_1)\lambda_i) \right. \\ &\quad \left. + 2 \sum_{i,j=1}^p \left(\text{Li}_2\left(1 - \frac{\zeta_i}{\lambda_j}\right) - \text{Li}_2\left(1 - \frac{\eta_i}{\lambda_j}\right) + \text{Li}_2\left(1 - \frac{\eta_i}{\eta_j}\right) - \text{Li}_2\left(1 - \frac{\zeta_i}{\eta_j}\right) \right) \right] \\ &\quad - \frac{1 - c_2}{c_2} \left[\log^2(1 - c_2) - \log^2(1 - c_1) + \sum_{i=1}^p (\log^2(\eta_i) - \log^2(\zeta_i)) \right] \\ &\quad - \frac{1}{p} \left[2 \sum_{i,j=1}^p \left(\text{Li}_2\left(1 - \frac{\zeta_i}{\lambda_j}\right) - \text{Li}_2\left(1 - \frac{\eta_i}{\lambda_j}\right) \right) - \sum_{i=1}^p \log^2((1 - c_1)\lambda_i) \right] \end{aligned}$$

where $\text{Li}_2(x) = - \int_0^x \frac{\log(1-u)}{u} du$ is the dilogarithm function and the η_i 's and ζ_i 's have been previously defined.

The case $f(t) = \log(1+st)$, with $s \in (0, \infty)$, is treated similarly as the case $f(t) = \log(t)$ with the main difference being a modification in the branch cuts that now occur when $\varphi_p(x)/\psi_p(x) < -1/s$ (rather than < 0). A new set of singularities $\kappa_0 < 0 < \kappa_1 < \dots < \kappa_p$, the zeros of $\varphi_p(x)/\psi_p(x) + 1/s$, are then naturally introduced. A simplification nonetheless allows one to express the resulting expression of the integral only as a function of κ_0 , as follows.

Corollary 3.4 (Case $f(t) = \log(1+st)$). *Under the setting of Theorem 3.1, let $s > 0$ and $f(t) = \log(1+st)$. Then,*

$$\begin{aligned}\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) &= \frac{c_1 + c_2 - c_1 c_2}{c_1 c_2} \log \left(\frac{c_1 + c_2 - c_1 c_2}{(1 - c_1)(c_2 - sc_1 \kappa_0)} \right) \\ &\quad + \frac{1}{c_2} \log(-s\kappa_0(1 - c_1)) + \int \log(1 - t/\kappa_0) \mu_p(dt)\end{aligned}$$

where $\kappa_0 < 0$ is the unique negative solution to $\varphi_p(\kappa_0)/\psi_p(\kappa_0) = -1/s$.

Details on the derivation of these results are available in [Couillet et al., 2019].

Table 3.2 illustrates, for the Fisher distance, the comparative performance gain of the large n, p -consistent estimator $\hat{d}_F(\mathbf{X}_1, \mathbf{X}_2) = \hat{d}(\mathbf{X}_1, \mathbf{X}_2; \log^2)$ proposed in Theorem 3.1 with respect to the traditional sample covariance plug-in estimator $d_F(\hat{\mathbf{C}}_1, \hat{\mathbf{C}}_2)$. It clearly appears that, as p/n_1 and p/n_2 become large (bottom part of the table), the standard large- n estimator dramatically fails, while the large- n, p consistent estimator remains quite accurate. More surprisingly is the fact that, even for small p , the large- n, p estimator still overtakes the large- n estimator on average. An evaluation of the respective estimate variances also reveals that both standard and RMT-improved approaches have similar fluctuations around their mean estimate.

Remark 3.3 (On the cases $c_1, c_2 > 1$). *While the large- n standard estimator requires $n \gg p$ and thus $n > p$, most random matrix analyses only demand that n, p be simultaneously large. Yet, Theorem 3.1 explicitly demands that $c_1 = \lim p/n_1 < 1$ and $c_2 = \lim p/n_2 < 1$. A careful control of the two successive changes of variable indeed reveals that, for say $c_2 > 1$, the pre-image Γ_ν of a contour Γ_μ around $\text{supp}(\mu)$ necessarily encloses zero. For $f(z)$ analytic in a neighborhood of $z = 0$, this has no consequence. But for $f(z) = \log^a(z)$, this annihilates the derivation and there seems to exist no simple workaround in this case.⁹ The case $f(z) = \log(1+sz)$ may still be valid, however only for sufficiently small values of s (that depend on c_1, c_2).*

⁹At the exception of the tentative alternative by polynomial approximation performed in [Tiomoko and Couillet, 2019b].

p	$d_F(\mathbf{C}_1, \mathbf{C}_2)$	$\hat{d}_F(\mathbf{X}_1, \mathbf{X}_2)$	$d_F(\hat{\mathbf{C}}_1, \hat{\mathbf{C}}_2)$
2	0.0533	0.0524	0.0568
4	0.0796	0.0840	0.0913
8	0.0927	0.0917	0.1048
16	0.0992	0.1007	0.1253
32	0.1025	0.1029	0.1509
64	0.1042	0.1049	0.2009
128	0.1050	0.1044	0.3023
256	0.1054	0.1057	0.5341
512	0.1056	0.1086	1.1556

Table 3.2: Estimation of the Fisher distance $d_F(\mathbf{C}_1, \mathbf{C}_2)$. Simulation example for $\mathbf{x}_i^{(a)} \sim \mathcal{N}(\mathbf{0}, \mathbf{C}_a)$ with $[\mathbf{C}_1]_{ij} = .2^{|i-j|}$, $[\mathbf{C}_2]_{ij} = .4^{|i-j|}$, $n_1 = 1\,024$, $n_2 = 2\,048$, as a function of p , results averaged over 30 runs. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

Remark 3.4 (Fluctuations). *Being a linear statistics (although a rather involved one) of the eigenvalues of $\hat{\mathbf{C}}_1^{-1}\hat{\mathbf{C}}_2$ with n_1, n_2, p of similar order, the estimate $\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f)$ can be shown to satisfy a central limit theorem with optimal speed $O(1/p)$, i.e.,*

$$\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) = d(\mathbf{C}_1, \mathbf{C}_2; f) + \frac{1}{p}\mathcal{N}(M, \sigma^2) + o(p^{-1})$$

for some $M, \sigma^2 = O(1)$. Besides, in the complex Gaussian case (i.e., $\mathbf{x}_i^{(a)} \sim \mathcal{CN}(\mathbf{0}, \mathbf{C}_a)$), $M = 0$.

Remark 3.5 (On nonnegativity). *It is important to stress that, similar to the GMUSIC estimator discussed in Section 3.1.3, although $\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) - d(\mathbf{C}_1, \mathbf{C}_2; f) \xrightarrow{a.s.} 0$, the nonnegativity of the distance $d(\mathbf{C}_1, \mathbf{C}_2; f)$ does not imply that of $\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f)$. In particular, for f such that $d(\cdot, \cdot; f)$ is an actual distance, if $\mathbf{C}_1 = \mathbf{C}_2 = \mathbf{C}$, $d(\mathbf{C}, \mathbf{C}; f) = 0$ while $\hat{d}(\mathbf{X}_1, \mathbf{X}_2; f) = 0 + \frac{1}{p}\mathcal{N}(M, \sigma^2) + o(p^{-1})$ which is thus often negative (with nonzero probability). This is another instance of the typical price to be paid for asymptotic consistency of random matrix estimators.*

3.2.4 The Wasserstein and Frobenius distances

As recalled in Remark 3.2, the Wasserstein distance between two centered Gaussian measures $\mathcal{N}(\mathbf{0}, \mathbf{C}_1)$ and $\mathcal{N}(\mathbf{0}, \mathbf{C}_2)$ is defined as

$$d_W(\mathbf{C}_1, \mathbf{C}_2) = \frac{1}{p} \text{tr}(\mathbf{C}_1) + \frac{1}{p} \text{tr}(\mathbf{C}_2) - 2 \int \sqrt{t} \nu_p^+(dt), \quad \nu_p^+ = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C}_1 \mathbf{C}_2)}$$

where the + sign in the exponent is here to recall that we take the product $\mathbf{C}_1^{+1} \mathbf{C}_2^{+1}$ rather than $\mathbf{C}_1^{-1} \mathbf{C}_2^{+1}$ as in the previous section.

In a similar manner, the Frobenius distance between \mathbf{C}_1 and \mathbf{C}_2 can be written as

$$d_{\text{Fro}}(\mathbf{C}_1, \mathbf{C}_2) = \frac{1}{p} \text{tr} \mathbf{C}_1^2 + \frac{1}{p} \text{tr} \mathbf{C}_2^2 - 2 \int t \nu_p^+(dt), \quad \nu_p^+ = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C}_1 \mathbf{C}_2)}.$$

Here, $\frac{1}{p} \text{tr} \hat{\mathbf{C}}_1 \hat{\mathbf{C}}_2$ is known to be a consistent estimate for $\frac{1}{p} \text{tr} \mathbf{C}_1 \mathbf{C}_2 = \int t \nu_p^+(dt)$ (which follows by first conditioning on, say, $\hat{\mathbf{C}}_1$ to obtain $\frac{1}{p} \text{tr} \hat{\mathbf{C}}_1 \hat{\mathbf{C}}_2 - \frac{1}{p} \text{tr} \hat{\mathbf{C}}_1 \mathbf{C}_2 \xrightarrow{a.s.} 0$ and operate similarly on $\hat{\mathbf{C}}_1$ to obtain the trivial estimator), so that the present framework is of marginal interest for the Frobenius distance between covariance matrices.

For the square-root function in the case of the Wasserstein distance and for generic functions f , the same framework as above may be used to estimate integral forms of the type

$$\int f(t) \nu_p^+(dt), \quad \nu_p^+ = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C}_1 \mathbf{C}_2)}.$$

This is performed in [Tiomoko and Couillet, 2019a] with the following result.

Theorem 3.2 (Covariance distance estimate for product matrices, from [Tiomoko and Couillet, 2019a]). *Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a real function with a complex analytic extension on $\{z \in \mathbb{C}, \Re[z] > 0\}$ and $c_1, c_2 < 1$. Then, with the same notations as in Theorem 3.1,*

$$d_+(\mathbf{C}_1, \mathbf{C}_2; f) - \hat{d}_+(\mathbf{X}_1, \mathbf{X}_2; f) \xrightarrow{a.s.} 0$$

where

$$d_+(\mathbf{C}_1, \mathbf{C}_2; f) = \int f(t) \nu_p^+(dt), \quad \nu_p^+ = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C}_1 \mathbf{C}_2)}$$

and

$$\hat{d}_+(\mathbf{X}_1, \mathbf{X}_2; f) = \frac{1}{2\pi i} \oint_{\Gamma_{\mu_p^+}} f\left(\frac{\varphi_p^+(u)}{\psi_p^+(u)}\right) \frac{\psi_p^+(u)}{c_2} \left[\frac{\varphi_p^{+'}(u)}{\varphi_p^+(u)} - \frac{\psi_p^{+'}(u)}{\psi_p^+(u)} \right] du$$

where $\varphi_p^+(z) = z/(1 - c_1 - c_1 z m_{\mu_p^+}(z))$, $\psi_p^+(z) = 1 - c_2 - c_2 z m_{\mu_p^+}(z)$ and μ_p^+ is the empirical spectral measure of $\hat{\mathbf{C}}_1 \hat{\mathbf{C}}_2$.

It is interesting to note that, “formally”, Theorem 3.2 only differs from Theorem 3.1 from the expression of $\psi_p^+(z)$ (called $\psi_p(z)$ in Theorem 3.1), but of course μ_p is also now changed into μ_p^+ which is a whole different function. Applied to the Wasserstein metric, a suitable complex integration calculus for the function $f(t) = \sqrt{t}$ leads to the following corollary.

Corollary 3.5 (Wasserstein distance estimate). *Under the setting of Theorem 3.2, let $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_p)^\top$ and $\boldsymbol{\Lambda} = \text{diag}(\boldsymbol{\lambda})$, for $\lambda_1 \leq \dots \leq \lambda_p$ the eigenvalues of $\hat{\mathbf{C}}_1 \hat{\mathbf{C}}_2$. Then,*

$$d_+(\mathbf{C}_1, \mathbf{C}_2; \sqrt{\cdot}) - \hat{d}_+(\mathbf{X}_1, \mathbf{X}_2; \sqrt{\cdot}) \xrightarrow{a.s.} 0$$

where, for $n_1 \neq n_2$,

$$\hat{d}_+(\mathbf{X}_1, \mathbf{X}_2; \sqrt{\cdot}) = \frac{2}{\sqrt{c_1 c_2}} \sum_{j=1}^p \sqrt{\lambda_j} + \frac{2}{\pi c_2} \sum_{j=1}^p \int_{\xi_j}^{\eta_j} \sqrt{-\frac{\varphi_p^+(x)}{\psi_p^+(x)}} \psi_p^{+'}(x) dx$$

while, for $n_1 = n_2$,

$$\hat{d}_+(\mathbf{X}_1, \mathbf{X}_2; \sqrt{\cdot}) = \frac{2}{c_2} \sum_{j=1}^p (\sqrt{\lambda_j} - \sqrt{\xi_j})$$

with $\{\xi_j\}_{j=1}^p$ and $\{\eta_j\}_{j=1}^p$ the increasing eigenvalues of $\boldsymbol{\Lambda} - \frac{\sqrt{\boldsymbol{\lambda}} \sqrt{\boldsymbol{\lambda}}^\top}{n_1}$ and $\boldsymbol{\Lambda} - \frac{\sqrt{\boldsymbol{\lambda}} \sqrt{\boldsymbol{\lambda}}^\top}{n_2}$, respectively.

The formula is particularly attractive in the case where $n_1 = n_2$, although its formal interpretation is not obvious. In Section 3.5, the detailed derivation and empirical evaluation of Corollary 3.5 above are provided in the form of a practical lecture material.

3.2.5 Application to covariance-based spectral clustering

In machine learning, the distance $d(\cdot, \cdot)$ between statistical covariance matrices \mathbf{C}_i is a popular *feature* to compare and classify data sets of, say, m data matrices $\mathbf{X}_1, \dots, \mathbf{X}_m$, where each datum \mathbf{X}_i is a collection of n_i vectors $\mathbf{X}_i = [\mathbf{x}_1^{(i)}, \dots, \mathbf{x}_{n_i}^{(i)}]$ with $\mathbb{E}[\mathbf{x}_j^{(i)}] = 0$ and $\mathbb{E}[\mathbf{x}_j^{(i)} \mathbf{x}_j^{(i)\top}] = \mathbf{C}_i$ (for instance, \mathbf{X}_i is n_i consecutive samples from a multivariate time series). These data can then be discriminated based on their differing covariance structures. Here, we will not be concerned with the size of m (which may be small or growing with n_i, p) but will impose that n_1, \dots, n_m, p are all large and comparable (and also that $n_1, \dots, n_m > p$ in accordance with Remark 3.3).

With these m observations $\mathbf{X}_1, \dots, \mathbf{X}_m$, classification based on a standard Gaussian kernel method would typically consist in assessing the kernel matrix

$$\mathbf{K} = \left\{ \exp \left(-\frac{1}{2} d(\mathbf{C}_i, \mathbf{C}_j) \right) \right\}_{i,j=1}^m$$

and then typically proceed to either (kernel) support vector classification (when in a supervised setting) or spectral clustering (when unsupervised). As we now know that estimating $d(\mathbf{C}_i, \mathbf{C}_j)$ by $d(\hat{\mathbf{C}}_i, \hat{\mathbf{C}}_j)$ may lead to dramatically erroneous results, \mathbf{K}_{ij} may be more appropriately estimated via

$$\hat{\mathbf{K}}_{ij} = \left\{ \exp \left(-\frac{1}{2} \hat{d}(\mathbf{X}_i, \mathbf{X}_j) \right) \right\}_{i,j=1}^m.$$

Figure 3.6 illustrates this idea in the context of spectral clustering of two classes, where $\mathbf{C}_1 = \dots = \mathbf{C}_{m/2} = \mathbf{C}^{(1)}$ (class 1) and $\mathbf{C}_{m/2+1} = \dots = \mathbf{C}_m = \mathbf{C}^{(2)}$ (class 2). There are depicted the dominant two eigenvectors of $\hat{\mathbf{K}}$, compared with those obtained by the classical estimates $d(\hat{\mathbf{C}}_i, \hat{\mathbf{C}}_j)$. The difference between these two methods is particularly remarkable as the number of samples n_i for each data differs:¹⁰ in this case, the estimation bias induced by $d(\hat{\mathbf{C}}_i, \hat{\mathbf{C}}_j)$ strongly depends on n_i, n_j and thus differently affects each single estimate. This is seen in the left display by the important spread of the pairs of eigenvector entries when the n_i 's are all different, and in the right display by the singular behavior of the only pair with different n_i value (which has the deleterious effect to affect the classification of all other data points!). As a result, spectral clustering performs significantly better with the random matrix-improved kernel.

The results introduced thus far on improved estimates for detection and estimation rely on an improved use of the sample covariance matrix. This is theoretically natural as, in the Gaussian case, sample covariance matrices are maximum-likelihood estimators for the population covariance: there is, as such, very little one can do better to retrieve information about the population covariance (unless prior information is available). The sample covariance estimator may however be far from optimal *away* from the Gaussian setting: when the observation vectors either emerge from “heavy-tailed” distributions or contain outliers. In this setting, refined versions of the sample covariance exist, the (small or large dimensional) behavior of which is technically more difficult to grasp. The next section introduces basic notions on the field known as *robust statistics*, originally designed to handle these non-Gaussian scenarios, and provides (again quite counter-intuitive) results retrieved by random matrix theory in the large n, p regime.

3.3 M-estimators of scatter

3.3.1 Reminder on robust statistics

Most probabilistic data models revolve around a Gaussian assumption: in signal processing additive noise models are mostly Gaussian, and in machine learning the Gaussian mixture model is one of the basic data models in classification. As already mentioned in Section 2.7, Gaussian models may adequately be replaced by a larger class of concentrated random vector models in large dimensions with little impact on the behavior of many machine learning algorithms. Gaussian and concentrated random vectors share the property that they “behave in a smooth and controllable way”. Instead, data polluted by outliers, missing entries, duplicates, etc., can typically not be accounted for by Gaussian or even

¹⁰In fact, for non-trivial tasks, where $\mathbf{C}^{(1)}$ and $\mathbf{C}^{(2)}$ are rather “close” to each other, *and for n_i all equal*, here again it can be surprisingly shown that the spectral clustering performance based on the inappropriate estimator $d(\hat{\mathbf{C}}_i, \hat{\mathbf{C}}_j)$ of $d(\mathbf{C}_i, \mathbf{C}_j)$ is as good as spectral clustering with the improved $\hat{d}(\mathbf{X}_i, \mathbf{X}_j)$; this a priori strange phenomenon vanishes as soon as the n_i 's are different.




Figure 3.6: First and second eigenvectors of $\hat{\mathbf{K}}$ with $\hat{\mathbf{K}}_{ij} = \exp(-\hat{d}(\mathbf{X}_i, \mathbf{X}_j)/2)$ (blue crosses) versus $\hat{\mathbf{K}}_{ij} = \exp(-d(\hat{\mathbf{C}}_i, \hat{\mathbf{C}}_j)/2)$ (red circles); with random number of snapshots n_i (left) and $n_1 = \dots = n_{m-1} = 512$ and $n_m = 256$ (right). [\[Link to code\]](#)

concentrated vector models.

To consider these outlying data in the models, Huber and his successors developed in the sixties the field of *robust statistics* [Huber, 2011, Maronna et al., 2018]. The basic observation of Huber lies in the lack of “robustness” of sample estimators (sample mean, sample variance and covariance matrix) to the presence of a single arbitrarily deviant outlying sample. Typically, for i.i.d. scalar samples $x_1, \dots, x_n \in \mathbb{R}$ with mean $M = \mathbb{E}[x_i]$, $\frac{1}{n} \sum_{i=1}^n x_i \xrightarrow{a.s.} M$ by the law of large numbers. Yet, the addition of x_0 with arbitrarily large amplitude to the sample average can drive $\frac{1}{n+1} \sum_{i=0}^n x_i$ arbitrarily far from M .

Huber proposed a min-max statistical mean and covariance estimation procedure in [Huber, 2011] that reduces the negative impact of outliers. The underlying assumption of Huber’s work is that the data x_i arise from a mixture of laws $(1 - \epsilon)\mu + \epsilon\mu'$ with μ a known “well-behaved” measure, μ' an unknown arbitrary measure and $\epsilon > 0$ small. The work of Maronna in [Maronna, 1976] generalizes that of Huber by letting x_i belong to a class of (multivariate) generalized Gaussian distributions, and notably of elliptical measures. A vector $\mathbf{x}_i \in \mathbb{R}^p$ is elliptically distributed if it can be written under the form¹¹

$$\mathbf{x}_i = \boldsymbol{\mu} + \sqrt{\tau_i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i \quad (3.6)$$

where \mathbf{z}_i is drawn uniformly at random on the sphere centered at zero and of radius \sqrt{p} , $\tau_i > 0$ is also drawn at random independently of \mathbf{z}_i , and the

¹¹In the literature, elliptical vectors are rather defined through their probability density which is of the type $\exp(-g((\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{C}^{-1}(\mathbf{x} - \boldsymbol{\mu})))$ for some function g , median vector $\boldsymbol{\mu}$ and scatter matrix \mathbf{C} ($\boldsymbol{\mu}$ coincides with the mean when it exists, and \mathbf{C} is proportional to the covariance matrix when it exists). The identity (3.6) is equivalent to the original definition, for g related to the law of the τ_i ’s.

nonnegative definite $\mathbf{C} \in \mathbb{R}^{p \times p}$ is the so-called *scatter matrix*. The law of the parameters τ_i (and notably its moments) controls the degree of “impulsiveness” of the data. When $\boldsymbol{\mu} = \mathbf{0}$ and $\mathbb{E}[\tau_i]$ is finite, then \mathbf{C} is proportional to the population covariance of \mathbf{x}_i ; if $\mathbb{E}[\tau_i] = \infty$, the covariance is not defined. Multivariate Gaussian (for which $\tau_i \xrightarrow{a.s.} 1$ as $p \rightarrow \infty$) and Student-t distributions belong to the class of elliptical distributions. Maronna derived the maximum likelihood estimators for $\boldsymbol{\mu}$ and \mathbf{C} for generic measures \mathcal{T} of the τ_i 's. These generalize the sample covariance matrix, include Huber's estimator as a special case, and are particularly resilient to “outlying” samples from the dataset.

3.3.2 The M-estimator of scatter

Of particular interest to the sample covariance matrix model thoroughly explored in this monograph is its relation to the M-estimators of scatter matrices. Under both Huber and Maronna's framework, for $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ data samples with $n > p$, Maronna's estimator of scatter $\hat{\mathbf{C}} \in \mathbb{R}^{p \times p}$ is defined as a solution to the fixed-point equation

$$\hat{\mathbf{C}} = \frac{1}{n} \sum_{i=1}^n u\left(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i\right) \mathbf{x}_i \mathbf{x}_i^\top \quad (3.7)$$

for $u : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a non-increasing function such that $\varphi(x) = xu(x)$ is non-decreasing and bounded. Typical examples of such functions are $u(x) = (1 + \alpha)/(x + \alpha)$ for some $\alpha > 0$ (this is the prototype of functions met in the maximum likelihood estimator of scatter for Student-t distributions) and $u(x) = \min\{\alpha/x, \beta\}$ for some $\alpha, \beta > 0$ (this is the prototype of Huber's robust estimators).

Under these assumptions on u , if $n > p$ and the \mathbf{x}_i 's are linearly independent, the solution $\hat{\mathbf{C}}$ to (3.7) can be shown to exist and be unique [Maronna, 1976]. Besides, the iterative fixed-point algorithm consisting in letting $\hat{\mathbf{C}}_0 = \mathbf{I}_p$ and, for $t \geq 0$,

$$\hat{\mathbf{C}}_{t+1} = \frac{1}{n} \sum_{i=1}^n u\left(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_t^{-1} \mathbf{x}_i\right) \mathbf{x}_i \mathbf{x}_i^\top$$

converges to $\hat{\mathbf{C}}$ as $t \rightarrow \infty$.

Due to their implicit definition, the behavior of these estimators is particularly difficult to apprehend. Still, it is interesting to observe the mode of action of $\hat{\mathbf{C}}$: u being decreasing, $\hat{\mathbf{C}}$ essentially reduces the impact of those \mathbf{x}_i 's such that $\mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i$ is large; that is, (i) those \mathbf{x}_i 's having too large amplitude (i.e., having large values of τ_i in the case of elliptical distributions) or (ii) those \mathbf{x}_i 's having a weak correlation to the dominant eigenvectors of $\hat{\mathbf{C}}$.

Under a finite n, p regime though, no much more can be said about $\hat{\mathbf{C}}$. However, under a large sample setting where $n \rightarrow \infty$ alone, it was shown that, if the \mathbf{x}_i 's are i.i.d. and elliptically distributed, $\hat{\mathbf{C}}$ converges almost surely to a matrix equal, up to a multiplicative constant, to \mathbf{C} .

A particular difficulty in handling the large- n alone scenario is that the quadratic form $\frac{1}{p}\mathbf{x}_i^\top \hat{\mathbf{C}}^{-1}\mathbf{x}_i$ does not concentrate and thus remains a rather involved random variable. This problem is completely alleviated in the large n, p regime. However, since $\hat{\mathbf{C}}$ depends on \mathbf{x}_i in a non-trivial manner, the large n, p limit of $\frac{1}{p}\mathbf{x}_i^\top \hat{\mathbf{C}}^{-1}\mathbf{x}_i$ is not as simple as for the sample covariance matrix model.

3.3.3 The random matrix framework

It is not easy to directly obtain a deterministic equivalent, or any asymptotic spectral behavior, for the robust estimator of scatter $\hat{\mathbf{C}}$ from (3.7), due to its implicit definition.

In such scenarios, which will also be the case for kernel random matrices met in Chapter 4, the objective is to substitute the intractable random matrix under study by a *random equivalent*, more amenable to random matrix analysis by conventional tools and techniques. In the present section, we show that, as $n, p \rightarrow \infty$, $\hat{\mathbf{C}}$ asymptotically behaves similarly to another random matrix $\hat{\mathbf{S}}$ in the sense that $\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| \xrightarrow{a.s.} 0$. The *random equivalent* $\hat{\mathbf{S}}$ is not observable but tractable to random matrix analysis. Since the convergence $\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| \xrightarrow{a.s.} 0$ transfers many spectral properties from $\hat{\mathbf{S}}$ to $\hat{\mathbf{C}}$, most of the (large dimensional) behavior of $\hat{\mathbf{C}}$ becomes accessible: in particular, deterministic equivalents for $\hat{\mathbf{C}}$ will be obtained by retrieving deterministic equivalents for $\hat{\mathbf{S}}$.

The random equivalent $\hat{\mathbf{S}}$ will however depend on the statistical distribution of the data \mathbf{x}_i and must in general be redesigned for each data model. We consider here the setting where $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ arise from a zero-mean elliptical distribution, i.e., $\mathbf{x}_i = \sqrt{\tau_i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$ as above with $\mathbf{z}_i \in \mathbb{R}^p$ uniformly drawn on the sphere centered at zero and of radius \sqrt{p} , $\boldsymbol{\mu} = \mathbf{0}$, \mathbf{C} positive definite, and $n > p$. Besides, the τ_i 's are positive i.i.d. random variables with measure \mathcal{T} having finite first order moment.

A first key observation is that one may already assume $\mathbf{C} = \mathbf{I}_p$ in the study of $\hat{\mathbf{C}}$. Indeed, by definition

$$\mathbf{C}^{-\frac{1}{2}} \hat{\mathbf{C}} \mathbf{C}^{-\frac{1}{2}} = \frac{1}{n} \sum_{i=1}^n u \left(\frac{1}{p} \mathbf{x}_i^\top \mathbf{C}^{-\frac{1}{2}} (\mathbf{C}^{-\frac{1}{2}} \hat{\mathbf{C}} \mathbf{C}^{-\frac{1}{2}})^{-1} \mathbf{C}^{-\frac{1}{2}} \mathbf{x}_i \right) \mathbf{C}^{-\frac{1}{2}} \mathbf{x}_i \mathbf{x}_i^\top \mathbf{C}^{-\frac{1}{2}}$$

where, in the inner parentheses, we simply wrote $\hat{\mathbf{C}} = \mathbf{C}^{-\frac{1}{2}} (\mathbf{C}^{-\frac{1}{2}} \hat{\mathbf{C}} \mathbf{C}^{-\frac{1}{2}})^{-1} \mathbf{C}^{-\frac{1}{2}}$. It is thus equivalent for $\hat{\mathbf{C}}$ to be the solution to the original problem for the data \mathbf{x}_i and for $\mathbf{C}^{-\frac{1}{2}} \hat{\mathbf{C}} \mathbf{C}^{-\frac{1}{2}}$ to be the solution to the same problem but with data $\mathbf{C}^{-\frac{1}{2}} \mathbf{x}_i$ in place of \mathbf{x}_i . We may then simply assume from now on that

$$\mathbf{x}_i = \sqrt{\tau_i} \mathbf{z}_i.$$

We mostly provide here the intuitive derivation of the main result, with a few words on the actual rigorous proof approach at the end. The idea starts

with the following intuition: letting

$$\hat{\mathbf{C}}_{-i} = \hat{\mathbf{C}} - \frac{1}{n} u \left(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i \right) \mathbf{x}_i \mathbf{x}_i^\top = \frac{1}{n} \sum_{j \neq i} u \left(\frac{1}{p} \mathbf{x}_j^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_j \right) \mathbf{x}_j \mathbf{x}_j^\top$$

it is clear that \mathbf{x}_i depends on $\hat{\mathbf{C}}_{-i}$ (because \mathbf{x}_i is part of $\hat{\mathbf{C}}$ appearing in each quadratic form $\frac{1}{p} \mathbf{x}_j^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_j$); yet, this dependence is seemingly “asymptotically weak” as \mathbf{x}_i only accounts for one out of n constitutive elements in $\hat{\mathbf{C}}$ and thus the quadratic forms $\frac{1}{p} \mathbf{x}_j^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_j$, $j \neq i$, barely depend on \mathbf{x}_i .

If this intuition is correct, we may expect the trace-lemma, Lemma 2.11, to hold for $\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i$; that is, we expect

$$\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i = \tau_i \frac{1}{p} \mathbf{z}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{z}_i \simeq \tau_i \frac{1}{p} \operatorname{tr} \hat{\mathbf{C}}_{-i}^{-1} \simeq \tau_i \frac{1}{p} \operatorname{tr} \hat{\mathbf{C}}^{-1} \equiv \tau_i \gamma_p$$

where in the last line we applied the rank-one perturbation Lemma 2.9 and introduced the notation $\gamma_p \equiv \frac{1}{p} \operatorname{tr} \hat{\mathbf{C}}^{-1}$.

In order to exploit the concentration $\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i \simeq \tau_i \gamma_p$, we now need to express $\hat{\mathbf{C}}$ as a function of such quadratic forms. To this end, by Lemma 2.8, first observe that

$$\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i = \frac{\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i}{1 + \frac{1}{n} u(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i) \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i}$$

which we may equivalently rewrite as

$$\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i = \frac{\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i}{1 - \frac{p}{n} \varphi(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i)} \quad (3.8)$$

assuming $1 - \frac{p}{n} \varphi(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i) \neq 0$, where we recall that $\varphi(x) = xu(x)$.

Consequently, if the mapping

$$\begin{aligned} g : \mathbb{R}^+ &\rightarrow \mathbb{R}^+ \\ x &\mapsto \frac{x}{1 - c\varphi(x)} \end{aligned}$$

with $c = p/n$, is bijective, one can express $\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i$ as

$$\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i = g^{-1} \left(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i \right).$$

This is indeed the case ($g'(x) > 0$ and $g(0) = 0$, $g(\infty) = \infty$) so long that $\|\varphi\|_\infty < c^{-1}$.

We will pose this assumption from now on and we may thus now rewrite $\hat{\mathbf{C}}$ under the form

$$\hat{\mathbf{C}} = \frac{1}{n} \sum_{i=1}^n v \left(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i \right) \mathbf{x}_i \mathbf{x}_i^\top$$

where we introduced the notation $v = u \circ g^{-1}$, a non-increasing function (graphically very similar to u). As $\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i \simeq \tau_i \gamma_p$, we further have

$$\begin{aligned}\hat{\mathbf{C}} &= \frac{1}{n} \sum_{i=1}^n v(\tau_i \gamma_p) \mathbf{x}_i \mathbf{x}_i^\top + o_{\|\cdot\|}(1) \\ &= \frac{1}{n} \sum_{i=1}^n \tau_i v(\tau_i \gamma_p) \mathbf{z}_i \mathbf{z}_i^\top + o_{\|\cdot\|}(1) \\ &= \frac{1}{\gamma_p} \frac{1}{n} \sum_{i=1}^n \psi(\tau_i \gamma_p) \mathbf{z}_i \mathbf{z}_i^\top + o_{\|\cdot\|}(1)\end{aligned}$$

where we defined $\psi(x) = xv(x)$ which, similar to φ , is increasing and bounded. It finally remains to evaluate γ_p . By definition

$$\gamma_p = \frac{1}{p} \text{tr } \hat{\mathbf{C}}^{-1} \simeq \frac{1}{p} \text{tr} \left(\frac{1}{\gamma_p} \frac{1}{n} \sum_{i=1}^n \psi(\tau_i \gamma_p) \mathbf{z}_i \mathbf{z}_i^\top \right)^{-1}$$

or equivalently

$$1 \simeq \frac{1}{p} \text{tr} \left(\frac{1}{n} \sum_{i=1}^n \psi(\tau_i \gamma_p) \mathbf{z}_i \mathbf{z}_i^\top \right)^{-1}. \quad (3.9)$$

As we may additionally expect that $\gamma_p \rightarrow \gamma$ for some deterministic γ , as $n, p \rightarrow \infty$, the trace above is merely the Stieltjes transform evaluated at zero of the matrix $\frac{1}{n} \mathbf{Z} \mathbf{D} \mathbf{Z}^\top$ where $\mathbf{Z} = [\mathbf{z}_1, \dots, \mathbf{z}_n]$ and \mathbf{D} is diagonal with $\mathbf{D}_{ii} = \psi(\tau_i \gamma)$. Since ψ is bounded, $c = p/n < 1$ and \mathbf{z}_i is a concentrated random vector (that can be seen as a normalized random Gaussian vector with norm tending to one), the Stieltjes transform for this model is well defined at zero and, from Theorem 2.6 or Theorem 2.18, has limit

$$m_{\frac{1}{n} \mathbf{Z} \mathbf{D} \mathbf{Z}^\top}(0) \xrightarrow{a.s.} m(0), \quad \text{with} \quad \frac{1}{m(0)} = \int \frac{\psi(t\gamma) \mathcal{T}(dt)}{1 + c\psi(t\gamma)m(0)}$$

where we recall that \mathcal{T} is the law of the τ_i 's. Moreover, since $m(0) = 1$ by (3.9), we conclude that γ is solution to:

$$1 = \int \frac{\psi(t\gamma) \mathcal{T}(dt)}{1 + c\psi(t\gamma)}.$$

This heuristic derivation allows us to conclude on the following asymptotic behavior for $\hat{\mathbf{C}}$.

Theorem 3.3 (Asymptotic equivalent for $\hat{\mathbf{C}}$, from [Couillet et al., 2015]). *Let $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$, $c = p/n < 1$, with $\mathbf{x}_i = \sqrt{\tau_i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$, τ_i i.i.d. with law \mathcal{T} of bounded moment of order $1 + \epsilon$ (for some $\epsilon > 0$), $\mathbf{C} \in \mathbb{R}^{p \times p}$ positive definite*

and $\mathbf{z}_i \in \mathbb{R}^p$ i.i.d. uniformly drawn at random on the sphere of mean zero and radius \sqrt{p} . Further let $u : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a non-increasing function such that $\varphi(x) = xu(x)$ is increasing and bounded by c^{-1} . Then

$$\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| \xrightarrow{a.s.} 0$$

where

$$\hat{\mathbf{C}} = \frac{1}{n} \sum_{i=1}^n u \left(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i \right) \mathbf{x}_i \mathbf{x}_i^\top, \quad \hat{\mathbf{S}} = \frac{1}{n} \sum_{i=1}^n v(\tau_i \gamma) \mathbf{x}_i \mathbf{x}_i^\top$$

with $v = u \circ g^{-1}$, $g(x) = x/(1 - c\phi(x))$ and, for $\psi(x) = xv(x)$, γ the unique positive solution to

$$1 = \int \frac{\psi(t\gamma)}{1 + c\psi(t\gamma)} \mathcal{T}(dt).$$

The fundamental result behind Theorem 3.3 is that, under an elliptical data model (the result would of course vary under other statistical assumptions), $\hat{\mathbf{C}}$ has the same asymptotic spectral behavior as the random matrix $\hat{\mathbf{S}}$. Now, unlike $\hat{\mathbf{C}}, \hat{\mathbf{S}}$ follows a quite elementary statistical model:

$$\hat{\mathbf{S}} = \frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \mathbf{D} \mathbf{Z}^\top \mathbf{C}^{\frac{1}{2}}, \quad \mathbf{Z} = [\mathbf{z}_1, \dots, \mathbf{z}_n], \quad \mathbf{D} = \text{diag}\{\tau_i v(\tau_i \gamma)\}_{i=1}^n.$$

Since the τ_i 's are independent of the \mathbf{z}_i 's, $\hat{\mathbf{S}}$ is merely a sample covariance matrix model with weights \mathbf{D}_{ii} on the samples. This model is thus completely characterized by Theorem 2.7 which in particular provides the limiting spectral distribution for $\hat{\mathbf{S}}$ which is identical to that of $\hat{\mathbf{C}}$.¹² These are depicted in Figure 3.8 and can be compared to the limiting spectral measure of the sample covariance matrix $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ in Figure 3.7, here for τ_i i.i.d. following a Gamma distribution. Of utmost interest from these figures is to remark that, while the limiting support of $\mu_{\frac{1}{n} \mathbf{X} \mathbf{X}^\top}$ is (provably) unbounded, since the Gamma distribution itself has unbounded support, the limiting support of $\mu_{\hat{\mathbf{C}}}$ (and of $\mu_{\hat{\mathbf{S}}}$) is bounded.

Besides, it can be easily checked that $\|\hat{\mathbf{C}}\|$ is also almost surely bounded since $\|\hat{\mathbf{S}}\| = \|\frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \mathbf{D} \mathbf{Z}^\top \mathbf{C}^{\frac{1}{2}}\|$, where $\|\mathbf{C}\|$ is bounded and $\mathbf{D} = \text{diag}\{\tau_i v(\tau_i \gamma)\}_{i=1}^n$ with entries bounded as $\tau_i v(\tau_i \gamma) = \psi(\tau_i \gamma)/\gamma < \|\psi\|_\infty/\gamma$ (which is finite).

The spectrum boundedness has one key consequence to spiked-model extensions of the model.

Remark 3.6 (Robust Spiked Model). *The model $\mathbf{x}_i = \sqrt{\tau_i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$ with $\mathbf{C} = \mathbf{I}_p$ is appropriate to model impulsive noise in signal processing applications, particularly so in array processing where radar signals are likely impulsive. In*

¹²Again, the entries of the \mathbf{z}_i 's are not i.i.d. but can be assumed as such by writing $\mathbf{z}_i = \sqrt{p} \tilde{\mathbf{z}}_i / \|\tilde{\mathbf{z}}_i\|$ with $\tilde{\mathbf{z}}_i \sim \mathcal{N}(0, \mathbf{I}_p)$, for which we have $\|\tilde{\mathbf{z}}_i\|/\sqrt{p} \xrightarrow{a.s.} 1$ uniformly on $i \in \{1, \dots, n\}$.

this array processing context, the natural extension to an information-plus-noise model reads

$$\mathbf{x}_i = \mathbf{a}s_i + \sqrt{\tau_i}\mathbf{z}_i$$

for a certain information vector $\mathbf{a} \in \mathbb{R}^p$ (to be detected and estimated) and possibly some scalar random signal $s_i \in \mathbb{R}$. Writing $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ leads to

$$\mathbf{X} = \mathbf{a}\mathbf{s}^\top + \mathbf{Z}\mathbf{T}^{\frac{1}{2}}, \quad \mathbf{T} = \text{diag}\{\tau_i\}_{i=1}^n, \quad \mathbf{s} = [s_1, \dots, s_n]^\top$$

which follows a spiked model.

However, due to the presence of the unbounded norm \mathbf{T} matrix, the sample covariance $\frac{1}{n}\mathbf{X}\mathbf{X}^\top$ has unbounded limiting support and thus no visible spike for all large n, p . As a main deleterious consequence, the signal \mathbf{a} can be neither detected nor estimated with spectral methods from the sample covariance.

Letting instead $\hat{\mathbf{C}} = \frac{1}{n} \sum_{i=1}^n u(\frac{1}{p}\mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i) \mathbf{x}_i \mathbf{x}_i^\top$ be the robust estimator of scatter, it is easily shown that $\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| \xrightarrow{a.s.} 0$ with $\hat{\mathbf{S}} = \frac{1}{n}\mathbf{Z}\mathbf{D}\mathbf{Z}^\top$ for $\mathbf{D} = \text{diag}\{\tau_i v(\tau_i \gamma)\}_{i=1}^n$ for the same γ defined in the noise-alone model; one must be careful though that $\hat{\mathbf{S}}$ differs from its expression in Theorem 3.3, as the present definition of \mathbf{Z} has changed. Since \mathbf{D} is bounded, this model for $\hat{\mathbf{S}}$ now is a proper spiked-model, allowing for the detection and estimation of \mathbf{a} . Details are provided in [Couillet, 2015].

Unlike all spiked models discussed in Section 2.5, a fundamental particularity of this spiked model is that, since the τ_i 's tend to spread due to their impulsive nature, even for large n , the support of \mathbf{D} may be quite scattered (all the more so when $u(x)$ is close to 1). This is a problem in practice as \mathbf{D} may induce its own “spikes” (isolated eigenvalues) into the spectrum of $\hat{\mathbf{C}}$, and these spikes may be confused with the genuine informative spikes. Here, a very peculiar phenomenon arises: since $\|\mathbf{D}\| \leq \|\psi\|_\infty / \gamma$, the noise-only model satisfies $\limsup \|\frac{1}{n}\mathbf{Z}\mathbf{D}\mathbf{Z}^\top\| \leq \|\psi\|_\infty (1 + \sqrt{c})^2 / \gamma \equiv S^+$ almost surely. Therefore, the noise-induced spikes can (asymptotically) not be found beyond S^+ , while the genuine spikes may. We thus have the typical picture of Figure 3.9 where (i) between the right-edge S_μ^+ of the support of the limiting spectrum μ of $\hat{\mathbf{C}}$ and S^+ , one can find both genuine and noise-driven spikes, while (ii) beyond S^+ only genuine informative spikes can be found.

3.3.3.1 A few words on the rigorous proof

The derivation leading up to Theorem 3.3 strongly relies on the claim that the trace lemma concentration

$$\frac{1}{p} \mathbf{z}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{z}_i \simeq \frac{1}{p} \text{tr } \hat{\mathbf{C}}_{-i}^{-1} \simeq \gamma$$

effectively holds true, uniformly so on $1 \leq i \leq n$, despite the dependence between \mathbf{z}_i and $\hat{\mathbf{C}}_{-i}$. The strategy proposed in [Couillet et al., 2015] to prove




Figure 3.7: Histogram of the eigenvalues of $\frac{1}{n} \mathbf{X} \mathbf{X}^T$, $\mathbf{x}_i = \sqrt{\tau_i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$, \mathbf{z}_i uniform on the \sqrt{p} -sphere, for $n = 2500$, $p = 500$, $\mathbf{C} = \text{diag}\{\mathbf{I}_{p/4}, 3\mathbf{I}_{p/4}, 10\mathbf{I}_{p/2}\}$, τ_i with $\Gamma(0.5, 2)$ -distribution. Link to code: [\[Matlab\]](#) and [\[Python\]](#).




Figure 3.8: Histogram of the eigenvalues of $\hat{\mathbf{C}}$ (left) and random equivalent $\hat{\mathbf{S}}$ (right) in the same setting of Figure 3.7, for $u(x) = (1 + \alpha)/(\alpha + x)$ with $\alpha = 0.2$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

this result is to “sandwich” the quantities $\frac{1}{p} \mathbf{z}_i^T \hat{\mathbf{C}}_{-i}^{-1} \mathbf{z}_i$ for $1 \leq i \leq n$ between two quantities with no dependence problem and which are easily shown to converge to γ .

However, as $\frac{1}{p} \mathbf{z}_i^T \hat{\mathbf{C}}_{-i}^{-1} \mathbf{z}_i$ appears as arguments of the function v , this sandwiching idea must be extended to a more convenient quantity. Precisely, recall-




Figure 3.9: Histogram of the eigenvalues of $\hat{\mathbf{C}}$ in a single-spike model, $\mathbf{x}_i = \sqrt{\tau_i} \mathbf{z}_i$ with \mathbf{z}_i uniform on the \sqrt{p} -sphere, for $u(x) = (1 + \alpha)/(\alpha + x)$ with $\alpha = 0.2$, $p = 256$, $n = 1024$, Student-t τ_i 's. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

ing that we may assume $\mathbf{C} = \mathbf{I}_p$ in the proof, we let

$$e_i = \frac{v\left(\frac{1}{p}\mathbf{x}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{x}_i\right)}{v(\tau_i \gamma)}$$

and relabel the e_i 's indices in such a way that $e_1 \leq \dots \leq e_n$. The main idea is then to observe that, letting $d_i = \frac{1}{p}\mathbf{z}_i^\top \hat{\mathbf{C}}_{-i}^{-1} \mathbf{z}_i$, we have $v(\tau_i d_i) = e_i v(\tau_i \gamma)$, and thus

$$\begin{aligned} e_i &= \frac{v\left(\tau_i \frac{1}{p} \mathbf{z}_i^\top \left(\frac{1}{n} \sum_{j \neq i} \tau_j v(\tau_j \gamma) \mathbf{z}_j \mathbf{z}_j^\top\right)^{-1} \mathbf{z}_i\right)}{v(\tau_i \gamma)} \\ &= \frac{v\left(\tau_i \frac{1}{p} \mathbf{z}_i^\top \left(\frac{1}{n} \sum_{j \neq i} \tau_j v(\tau_j \gamma) e_j \mathbf{z}_j \mathbf{z}_j^\top\right)^{-1} \mathbf{z}_i\right)}{v(\tau_i \gamma)}. \end{aligned}$$

Using $e_1 \leq e_i \leq e_n$ and the nondecreasing nature of v , we have both

$$\begin{aligned} e_i &\leq \frac{v\left(\frac{1}{e_n} \tau_i \frac{1}{p} \mathbf{z}_i^\top \left(\frac{1}{n} \sum_{j \neq i} \tau_j v(\tau_j \gamma) \mathbf{z}_j \mathbf{z}_j^\top\right)^{-1} \mathbf{z}_i\right)}{v(\tau_i \gamma)}, \\ e_i &\geq \frac{v\left(\frac{1}{e_1} \tau_i \frac{1}{p} \mathbf{z}_i^\top \left(\frac{1}{n} \sum_{j \neq i} \tau_j v(\tau_j \gamma) \mathbf{z}_j \mathbf{z}_j^\top\right)^{-1} \mathbf{z}_i\right)}{v(\tau_i \gamma)}. \end{aligned}$$

Focusing on the first inequality, being valid for each i , it is particularly valid for $i = n$, and thus

$$e_n \leq \frac{v\left(\frac{1}{e_n}\tau_n \frac{1}{p}\mathbf{z}_n^\top \left(\frac{1}{n} \sum_{j \neq n} \tau_j v(\tau_j \gamma) \mathbf{z}_j \mathbf{z}_j^\top\right)^{-1} \mathbf{z}_n\right)}{v(\tau_n \gamma)}.$$

The quadratic form in the numerator above has been cleared of its dependence problems and it is thus only a matter of standard random matrix theory to show that

$$\frac{1}{p}\mathbf{z}_n^\top \left(\frac{1}{n} \sum_{j \neq n} \tau_j v(\tau_j \gamma) \mathbf{z}_j \mathbf{z}_j^\top\right)^{-1} \mathbf{z}_n \xrightarrow{a.s.} \gamma$$

(based on the same Stieltjes transform argument as previously). As a side but important comment, note that, due to the relabeling of e_1, \dots, e_n , \mathbf{z}_n is effectively no longer independent of $\mathbf{z}_1, \dots, \mathbf{z}_{n-1}$ and thus the convergence above in fact follows from a *uniform* convergence to γ of the quadratic forms for all $i = 1, \dots, n$. We thus conclude that, for $\epsilon > 0$ arbitrarily small, for all large n, p ,

$$e_n \leq \frac{v\left(\frac{1}{e_n}\tau_n(\gamma - \epsilon)\right)}{v(\tau_n \gamma)}$$

almost surely which we can equivalently write, by dividing left and right sides by $\tau_n \gamma$,

$$\psi(\tau_n \gamma) \leq v\left(\frac{1}{e_n}\tau_n(\gamma - \epsilon)\right) \frac{1}{e_n} \tau_n \gamma.$$

Let us now assume that $\limsup_n e_n > 1$ (which we want to disprove) and let us restrict ourselves to a subsequence over which e_n is away from one. For simplicity, we consider also that the τ_i 's are all bounded and that ψ is strictly increasing (with the general case treated in [Coullet et al., 2015]). We may thus extract a further subsequence over which $\tau_n \rightarrow \tau_\infty$, $e_n \rightarrow e_\infty \in (1, \infty]$, $\epsilon \rightarrow 0$, and then in the limit

$$\psi(\tau_\infty \gamma) \leq \psi(e_\infty^{-1} \tau_\infty \gamma).$$

But since ψ is strictly increasing, this implies that $e_\infty \leq 1$ which contradicts the assumption that $\limsup_n e_n > 1$. Thus, $\limsup_n e_n \leq 1$ (almost surely).

With the same arguments, we show that $\liminf_n e_1 \geq 1$ almost surely, so that finally we have that $e_i \xrightarrow{a.s.} 1$ for all $i = 1, \dots, n$ which completes the proof.

3.3.4 Extensions

Tyler's rotational invariant estimator. The class of robust estimators of scatter of the type of $\hat{\mathbf{C}}$ in (3.7) developed by Huber and Maronna imposes that

the robustness function u be such that $u(0)$ be defined and that $x \mapsto xu(x)$ be increasing and bounded.

In [Tyler, 1983], Tyler proposed another version of $\hat{\mathbf{C}}$, which can be thought of as a limiting version of (3.7) for $u(x) = 1/x$, i.e.,

$$\hat{\mathbf{C}} = \frac{1}{n} \sum_{i=1}^n \frac{\mathbf{x}_i \mathbf{x}_i^\top}{\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i}. \quad (3.10)$$

The apparently slight modification of Maronna's conditions (here $u(0)$ is not defined and $xu(x)$ is not increasing) dramatically disrupts the behavior of $\hat{\mathbf{C}}$. First, note that $\hat{\mathbf{C}}$ is *no longer unique* as a solution to (3.10): indeed, if $\hat{\mathbf{C}}$ is solution, it is easy to see that so is $\alpha\hat{\mathbf{C}}$ for all $\alpha > 0$. It can be shown that these are all the solutions. That is, for $n > p$ and \mathbf{x}_i linearly independent, there exists $\hat{\mathbf{C}}_0$ solution to (3.10) and the set of solutions $\hat{\mathbf{C}}$ is exactly $\{\alpha\hat{\mathbf{C}}_0, \alpha > 0\}$.

The main advantage of this robust formulation lies in its invariance with respect to the amplitude of the outliers. Under this formulation, all the \mathbf{x}_i 's are “normalized” since (3.10) features the ratio $\mathbf{x}_i \mathbf{x}_i^\top / \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i$. This advantage however turns into a drawback if one needs to maintain and compare the norms of the data.

The asymptotic analysis of Tyler's estimator in the large n, p regime does not unfold from the proof of Theorem 3.3 which strongly exploits the fact that $x \mapsto xu(x)$ is increasing (while here $xu(x) = 1$). In [Zhang et al., 2014], the authors exploit a different approach to prove that, for elliptical data with scatter matrix $\mathbf{C} = \mathbf{I}_p$, Tyler's estimator asymptotically behaves as a sample covariance matrix composed of i.i.d. random vectors with zero mean and identity covariance. Consequently, the limiting spectral measure of $\hat{\mathbf{C}}$ is simply the Marčenko-Pastur law.

In a sense, Tyler's estimator is “as robust as robust estimators can get” since its null-hypothesis spectrum (when $\mathbf{C} = \mathbf{I}_p$) leads to the “most compact” spectral distribution. For all other u functions, the limiting spectrum is more spread. This at first seems more advantageous in a spiked extension of the model, as isolated eigenvalues are likely more visible under this setting. Yet, the harsh normalization of all data simultaneously “breaks” the low rank eigenspaces *maximally* for Tyler's estimator. A compromise for a suitable u function is demanded in this case. These aspects are discussed in [Couillet and Benaych-Georges, 2016].

The $p > n$ scenario. The robust estimator of scatter $\hat{\mathbf{C}}$ as defined in (3.7) has the major inconvenience of not being well defined for $p > n$ as $\hat{\mathbf{C}}$ is expressed through its inverse while being the sum of the n rank-one matrices $\frac{1}{n} u(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i) \mathbf{x}_i \mathbf{x}_i^\top$ and is thus of maximum rank $n < p$.

To cover the scenario $p > n$, one usually resorts to a *linear-shrinkage* version

of the original $\hat{\mathbf{C}}$ by instead defining $\hat{\mathbf{C}}$ as the solution to

$$\hat{\mathbf{C}} = (1 - \rho) \frac{1}{n} \sum_{i=1}^n u \left(\frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i \right) \mathbf{x}_i \mathbf{x}_i^\top + \rho \mathbf{I}_p$$

for some $\rho \in (0, 1]$. Thanks to the $\rho \mathbf{I}_p$ addition, the right-hand side term is positive definite (i.e., $\hat{\mathbf{C}} \succeq \rho \mathbf{I}_p$ in the sense of symmetric matrices) and it can be shown that, under similar conditions on u as in the previous paragraphs, $\hat{\mathbf{C}}$ is well defined as the unique solution to the equation.

Under this setting, the asymptotic analysis of $\hat{\mathbf{C}}$ essentially boils down to the control of the minimal eigenvalue of $\hat{\mathbf{C}}$ (which could be close to zero, thus leading to an explosion of $\mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i$). The works [Coullet and McKay, 2014, Auguin et al., 2018] extend the results in Theorem 3.3 to this setting.

Robustness to arbitrary outliers. It is important to insist that the asymptotic equivalence $\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| \xrightarrow{a.s.} 0$ in Theorem 3.3 is only valid for the specific elliptic model of the data \mathbf{x}_i , a scenario mostly motivated by Maronna's original works [Maronna, 1976] on the maximum likelihood estimator for elliptical datasets and by the adequate modeling of impulsive noise environments beyond the Gaussian noise model in practice.

In the original works of Huber on robust statistics though, the initial objective of $\hat{\mathbf{C}}$ was to cope with the presence of outlying data in the samples. To this end, from a large dimensional random matrix viewpoint, it is more convenient to assume that the data observations $\mathbf{X}_{\mathbf{A}} \in \mathbb{R}^{p \times n}$ are composed in part of *clean data* and in part of *outliers*. We may write

$$\mathbf{X}_{\mathbf{A}} = [\mathbf{X}, \mathbf{A}] = [\mathbf{x}_1, \dots, \mathbf{x}_{(1-\epsilon_n)n}, \mathbf{a}_1, \dots, \mathbf{a}_{\epsilon_n n}]$$

for a proportion ϵ_n (a small or large multiple of $1/n$) of deterministic unknown outliers $\mathbf{A} = [\mathbf{a}_1, \dots, \mathbf{a}_{\epsilon_n n}]$ and $(1 - \epsilon_n)$ of genuine data $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_{(1-\epsilon_n)n}]$.

In [Morales-Jimenez et al., 2015], letting \mathbf{x}_i be independent $\mathcal{N}(\mathbf{0}, \mathbf{C})$ and the \mathbf{a}_i 's be such that $\frac{1}{n} \mathbf{C}^{-\frac{1}{2}} \mathbf{A} \mathbf{A}^\top \mathbf{C}^{-\frac{1}{2}}$ has bounded norm, Theorem 3.3 is turned into the following result.

Theorem 3.4 (Robust estimator with outliers, from [Morales-Jimenez et al., 2015]). *Let $\mathbf{X}_{\mathbf{A}} = [\mathbf{X}, \mathbf{A}]$ be as above. Then, under the assumptions and notations of Theorem 3.3 and with $\epsilon_n \rightarrow \epsilon \in (0, 1 - c)$,*

$$\|\hat{\mathbf{C}} - \hat{\mathbf{S}}_{\mathbf{A}}\| \xrightarrow{a.s.} 0, \quad \hat{\mathbf{S}}_{\mathbf{A}} = v(\gamma_n) \frac{1}{n} \mathbf{X} \mathbf{X}^\top + \frac{1}{n} \sum_{i=1}^{\epsilon_n n} v(\alpha_{i,n}) \mathbf{a}_i \mathbf{a}_i^\top$$

where $(\gamma_n, \alpha_{1,n}, \dots, \alpha_{\epsilon n, n})$ are the unique solution to

$$\begin{aligned}\gamma_n &= \frac{1}{p} \operatorname{tr} \mathbf{C} \left(\frac{(1-\epsilon)v(\gamma_n)}{1+cv(\gamma_n)\gamma_n} \mathbf{C} + \frac{1}{n} \sum_{i=1}^{\epsilon n n} v(\alpha_{i,n}) \mathbf{a}_i \mathbf{a}_i^\top \right)^{-1} \\ \alpha_{i,n} &= \mathbf{a}_i^\top \left(\frac{(1-\epsilon)v(\gamma_n)}{1+cv(\gamma_n)\gamma_n} \mathbf{C} + \frac{1}{n} \sum_{j \neq i} v(\alpha_{j,n}) \mathbf{a}_j \mathbf{a}_j^\top \right)^{-1} \mathbf{a}_i.\end{aligned}$$

It is already interesting to see that the random equivalent $\hat{\mathbf{S}}_{\mathbf{A}}$ (and thus the robust estimator $\hat{\mathbf{C}}$) properly weighs all genuine data by a constant $v(\gamma_n)$ and then weighs all outliers with a parameter $v(\alpha_{i,n})$ proportional to its “outlying” character.

Of particular interest is the case of a vanishing proportion of outliers with $\epsilon_n \rightarrow 0$ (for instance $\epsilon_n = k/n$, corresponding to exactly k outliers), which slightly extends the theorem statement. Then, it is easily seen that $\gamma_n \rightarrow \gamma$ given in the statement of Theorem 3.3 for $\tau_i = 1$, and thus

$$\gamma_n \rightarrow \gamma = \frac{\varphi^{-1}(1)}{1-c}.$$

In this case, we have

$$\hat{\mathbf{S}}_{\mathbf{A}} = \frac{1}{\varphi^{-1}(1)} \frac{1}{n} \mathbf{X} \mathbf{X}^\top + \frac{1}{n} \sum_{i=1}^n v(\alpha_{i,n}) \mathbf{a}_i \mathbf{a}_i^\top.$$

If, in addition, $\epsilon_n n = k$ and $\mathbf{a}_1 = \dots = \mathbf{a}_k \equiv \mathbf{a}$, then $\alpha_{1,n} = \dots = \alpha_{\epsilon n, n} \equiv \alpha_n$ is given by the unique positive solution of

$$\alpha_n = \frac{\gamma \frac{1}{p} \mathbf{a}^\top \mathbf{C}^{-1} \mathbf{a}}{1 + c\gamma(k-1)v(\alpha_n) \frac{1}{p} \mathbf{a}^\top \mathbf{C}^{-1} \mathbf{a}}.$$

Several conclusions can be drawn here: first note that the outlying data are weighed by a factor proportional to $\mathbf{a}^\top \mathbf{C}^{-1} \mathbf{a}$. Thus, the robust estimator behaves *as if knowledgeable of the quantity* $\mathbf{a}^\top \mathbf{C}^{-1} \mathbf{a}$ although \mathbf{C} itself is not accessible; it thus performs much as expected by only discarding from the samples those data vectors \mathbf{a} not aligned with \mathbf{C} . But also note that, if $\mathbf{C} = \mathbf{I}_p$, then $\mathbf{a}^\top \mathbf{C}^{-1} \mathbf{a} = \|\mathbf{a}\|^2$, in which case the robust estimator only evaluates the amplitude of the outlier, rather than its characteristic covariance structure, to decide on the allocated weight. Consequently, robust estimators are mostly resilient to outliers if the genuine data is quite structured (and outliers are misaligned to this structure, so for instance when they have i.i.d. entries), but the converse is not true.

But a much more troubling and unexpected effect is that, if k (so finitely many) outliers are *identical*, then α_n scales as $1/k$ and thus quickly vanishes. Ultimately, only few outliers suffice to obtain $\alpha_n < \gamma$ and thus $v(\alpha_n) > v(\gamma)$:

the outliers are then given *more weight* than the genuine data, going in a stark opposite direction as originally intended. It is interesting that random matrix theory so easily reveals such behaviors which conventional $n \gg 1$ statistics is generally unable to see (or only through approximations).

Second order statistics. The convergence $\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| \rightarrow 0$ (Theorems 3.3 and 3.4) is convenient to transfer the *first order* spectral properties of $\hat{\mathbf{S}}$ to $\hat{\mathbf{C}}$. That is, this convergence implies (i) $\max_{1 \leq i \leq p} |\lambda_i(\hat{\mathbf{C}}) - \lambda_i(\hat{\mathbf{S}})| \rightarrow 0$ and (ii) $\|\mathbf{u}_i(\hat{\mathbf{C}}) - \hat{\mathbf{u}}_i(\hat{\mathbf{S}})\| \rightarrow 0$ for all *isolated* eigenpair $(\lambda_i(\hat{\mathbf{C}}), \mathbf{u}_i(\hat{\mathbf{C}}))$ (i.e., such that $|\lambda_i(\hat{\mathbf{C}}) - \lambda_{i \pm 1}(\hat{\mathbf{C}})|$ does not vanish).

As such, $\hat{\mathbf{C}}$ and $\hat{\mathbf{S}}$ have the same limiting spectral distribution, their eigenvalues are point-wise asymptotically equal and they share the same *isolated* eigenvectors in the limit. From a practical standpoint, this in particular means that the asymptotic threshold for signal detection (based on isolated eigenvalues) can be transferred from the statistics of $\hat{\mathbf{S}}$ to $\hat{\mathbf{C}}$ and that the informative content in the eigenvectors of $\hat{\mathbf{C}}$ can be understood from those of $\hat{\mathbf{S}}$.

However, this is as far as the $\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| \rightarrow 0$ convergence goes. The question of the asymptotic local fluctuations of the individual eigenvalues and of the eigenvector projection statistics cannot be transferred straightforwardly. In particular, it is believed that $\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| = O(1/\sqrt{n})$. Since the dominant eigenvalues $\lambda_i(\hat{\mathbf{S}})$ and eigenvector projections $\mathbf{u}_i(\hat{\mathbf{S}})^T \mathbf{a}$ for deterministic \mathbf{a} satisfy central limit theorems at this $O(1/\sqrt{n})$ rate precisely, we can at least tell that $\lambda_i(\hat{\mathbf{C}})$ and $\mathbf{u}_i(\hat{\mathbf{C}})^T \mathbf{a}$ also fluctuate at a $O(1/\sqrt{n})$ rate but it is impossible to infer whether a central limit theorem holds and even to estimate the limiting mean and variance of the fluctuation. The problem is even exacerbated when it comes to faster statistics, such as linear functionals $\frac{1}{n} \sum_i f(\lambda_i(\hat{\mathbf{C}}))$: while $\frac{1}{n} \sum_i f(\lambda_i(\hat{\mathbf{S}}))$ is known to satisfy a central limit theorem at rate $O(1/n)$, the convergence $\|\hat{\mathbf{C}} - \hat{\mathbf{S}}\| = O(1/\sqrt{n})$ only allows one to say that $\frac{1}{n} \sum_i f(\lambda_i(\hat{\mathbf{C}}))$ fluctuates at speed $O(1/\sqrt{n})$.

In [Couillet et al., 2016a], it is shown that more can be said for precise statistics. Assuming the case where $u(x) = 1/x$, $\mathbf{C} = \mathbf{I}_p$, and $\hat{\mathbf{C}}$ is regularized by $\rho \mathbf{I}_p$ for any $\rho > 0$, it is proved that, for all deterministic vectors $\mathbf{a}, \mathbf{b} \in \mathbb{R}^p$ and for all $k \in \mathbb{Z}$,

$$\mathbf{a}^T \hat{\mathbf{C}}^k \mathbf{b} - \mathbf{a}^T \hat{\mathbf{S}}^k \mathbf{b} = O(n^{-1+\epsilon}) \quad (3.11)$$

for all $\epsilon > 0$. This result can be straightforwardly used to show that $\mathbf{u}_i(\hat{\mathbf{C}})^T \mathbf{a}$ satisfies the same asymptotic fluctuations as $\mathbf{u}_i(\hat{\mathbf{S}})^T \mathbf{a}$.

As a concrete application example, the robust estimator $\hat{\mathbf{C}}$ may be used for the following hypothesis testing problem

$$\mathbf{x} = \begin{cases} \sqrt{\tau} \mathbf{z} & , \mathcal{H}_0 \\ s \mathbf{a} + \sqrt{\tau} \mathbf{z} & , \mathcal{H}_1 \end{cases}, \quad \text{given } \mathbf{x}_i = \sqrt{\tau_i} \mathbf{z}_i, \quad 1 \leq i \leq n$$

where \mathbf{z}, \mathbf{z}_i are, say, independent and uniformly distributed on the \sqrt{p} -sphere, $\mathbf{a} \in \mathbb{R}^p$ is some known vector and $s \in \mathbb{R}$ is unknown. Here, one assumes to have access to a single observation \mathbf{x} but also to some prior “pure noise” data $\mathbf{x}_1, \dots, \mathbf{x}_n$, and it is to be tested whether the vector \mathbf{a} is present in \mathbf{x} .

This setting corresponds to that of an impulsive background noise environment within which an informative data \mathbf{a} is expected to be eventually detected (\mathbf{a} could be a signal signature such as a steering vector in array processing).

Since \mathbf{a} is known, the generalized likelihood ratio test (GLRT) in this setting (recall Section 3.1.1 for a definition of the GLRT) reads

$$T_p \equiv \frac{|\mathbf{x}^\top \hat{\mathbf{C}}^{-1} \mathbf{a}|^2}{(\mathbf{x}^\top \hat{\mathbf{C}}^{-1} \mathbf{x})(\mathbf{a}^\top \hat{\mathbf{C}}^{-1} \mathbf{a})} \stackrel{\mathcal{H}_1}{\gtrless} \stackrel{\mathcal{H}_0}{\lessgtr} \alpha.$$

In the absence of \mathbf{a} within \mathbf{x} , $\mathbf{x}^\top \hat{\mathbf{C}}^{-1} \mathbf{a} = O(1/\sqrt{n})$ while in the presence of \mathbf{a} , $\mathbf{x}^\top \hat{\mathbf{C}}^{-1} \mathbf{a}$ is of the order of α . The test is therefore asymptotically non-trivial only if α is of order $O(1/\sqrt{n})$. Under these conditions, the performance of the test (its asymptotic errors of Type I and II) are given by the asymptotic behavior of T_p under both \mathcal{H}_0 and \mathcal{H}_1 hypotheses.

This behavior is accessible by showing a central limit theorem for T_p , which itself follows (by the delta-method [Vaart, 2000]) from a central limit theorem on the vector $(\mathbf{x}^\top \hat{\mathbf{C}}^{-1} \mathbf{a}, \mathbf{x}^\top \hat{\mathbf{C}}^{-1} \mathbf{x}, \mathbf{a}^\top \hat{\mathbf{C}}^{-1} \mathbf{a})$. From (3.11), this is asymptotically equivalent to establishing a central limit theorem for $(\mathbf{x}^\top \hat{\mathbf{S}}^{-1} \mathbf{a}, \mathbf{x}^\top \hat{\mathbf{S}}^{-1} \mathbf{x}, \mathbf{a}^\top \hat{\mathbf{S}}^{-1} \mathbf{a})$ which, given the elementary modeling of $\hat{\mathbf{S}}$, is within reach of random matrix theory.

A thorough investigation of this GLRT asymptotics is performed in [Couillet et al., 2016a, Kammoun et al., 2017].

3.4 Concluding remarks

All the methods presented in this chapter, from discriminant analysis down to robust covariance estimation, all consist, one way or another, in improving the mismatched estimation of a covariance matrix \mathbf{C} by its sample covariance $\hat{\mathbf{C}}$.

However, as opposed to the conventional idea that one must before everything improve this mismatched estimate $\hat{\mathbf{C}}$ into a “better” plug-in estimate of \mathbf{C} , the random matrix approach developed in this chapter rather consists in the first place in identifying the ultimate scalar (or small dimensional) parameter to be optimized, and only then, adapt the estimate of \mathbf{C} appropriately. Specifically, we saw that:

- in the discriminant analysis scenario, we estimate \mathbf{C} through a “linear shrinkage” version $\hat{\mathbf{C}} + \gamma \mathbf{I}_p$ of $\hat{\mathbf{C}}$, and then aim at optimizing γ in such a way to maximize the ultimate detection probability of the underlying hypothesis test;
- in the spike G-MUSIC improvement of the MUSIC algorithm, one aims primarily at estimating quadratic forms of the type $\mathbf{a}^\top \mathbf{u} \mathbf{u}^\top \mathbf{a}$ where \mathbf{u} is

an eigenvector (associated to the largest eigenvalues) of \mathbf{C} . There, the covariance \mathbf{C} is never estimated, and only the quadratic form $\mathbf{a}^\top \mathbf{u} \mathbf{u}^\top \mathbf{a}$ is retrieved as a function of $\mathbf{a}^\top \hat{\mathbf{u}} \hat{\mathbf{u}}^\top \mathbf{a}$ with $\hat{\mathbf{u}}$ the corresponding eigenvector in $\hat{\mathbf{C}}$;

- in the distance estimation framework between two covariance matrices, again, the ultimate target is the distance $d(\mathbf{C}_1, \mathbf{C}_2)$; instead of correcting the quite erroneous but natural estimate $d(\hat{\mathbf{C}}_1, \hat{\mathbf{C}}_2)$, the strategy consists in characterizing $d(\mathbf{C}_1, \mathbf{C}_2)$ as a function of the resolvents of \mathbf{C}_1 and \mathbf{C}_2 , before connecting them to $\hat{\mathbf{C}}_1$ and $\hat{\mathbf{C}}_2$; in the end, the estimators depend in a non-trivial manner on some functional of the eigenvalues of $\hat{\mathbf{C}}_1 \hat{\mathbf{C}}_2$ (or $\hat{\mathbf{C}}_1^{-1} \hat{\mathbf{C}}_2$);
- finally, for the robust covariance estimator method, the asymptotics of the robust estimator are not so trivially related to the underlying covariance matrices being estimated, but a deep investigation of the quadratic forms $\mathbf{x}_i^\top \hat{\mathbf{C}}^{-1} \mathbf{x}_i$ at the core of the estimator fully reveals the robust estimator behavior.

While estimating the $p(p - 1)/2$ elements of \mathbf{C} from the np samples $[\mathbf{X}]_{ij}$ of the data matrix $\mathbf{X} \in \mathbb{R}^{p \times n}$ cannot be performed consistently in the regime where $n \sim p$ (at least when no strong a priori structure is supposed on \mathbf{C}), this does not necessarily mean that there is no means to improve over the sample covariance matrix.

Specifically, a recent direction is being followed which consists in generalizing the notion of “linear shrinkage”, i.e., estimating \mathbf{C} through the matrix $\hat{\mathbf{C}} + \gamma \mathbf{I}_p$ for some $\gamma > 0$, to “nonlinear shrinkage”. The idea behind nonlinear shrinkage is to design an estimator of the type $\hat{\mathbf{U}} f(\hat{\Lambda}) \hat{\mathbf{U}}^\top$, where $\hat{\mathbf{U}} \hat{\Lambda} \hat{\mathbf{U}}^\top = \hat{\mathbf{C}}$ is the spectral decomposition of $\hat{\mathbf{C}}$ and $f(\cdot)$ is a function applied entry-wise on the diagonal elements. The function f is then selected in such a way that a distance criterion, such as the Frobenius norm error $\mathbb{E}[\|\mathbf{C} - \hat{\mathbf{U}} f(\hat{\Lambda}) \hat{\mathbf{U}}^\top\|_F^2]$, is minimized or alternatively such that $f(\hat{\lambda}_i)$ estimates the corresponding i -th eigenvalue of \mathbf{C} . In a series of works [Ledoit and Péché, 2011, Ledoit and Wolf, 2012, Bun et al., 2017], the authors proposed several such functions f .

Overall, it is interesting to take a step back and realize the number and diversity of findings which were obtained since the seminal article of Marčenko and Pastur in 1967 surrounding the sample covariance matrix model. Clearly ubiquitous in machine learning, second order statistics have for long never been the subject of so deep investigations (as the sample covariance was considered a good estimator for the population covariance) until this fundamental first random matrix work. It is now fully admitted by many research communities (statistics, statistical physics, electrical engineering), and increasingly by the machine learning experts, that all methods and algorithms derived from a mere replacement of the population covariance matrix by the sample covariance are at best hazardous, and often counterproductive.

This is another instance of the curse of dimensionality in large and numerous data processing problems, which is being more and more efficiently tackled. The next chapter goes a step further, beyond the linear and quadratic setting, into kernel-based algorithms (and thus nonlinear functions of the data, more akin to most machine learning algorithms).

3.5 Practical course material

Practical Lecture Material 1 (The Wasserstein distance estimation). *This exercise aims to formally derive the proof of Theorem 3.2 in the specific case of the Wasserstein distance. That is, we aim at estimating, for two sets of independent centered Gaussian samples with covariance matrices \mathbf{C}_1 and \mathbf{C}_2 , the quantity*

$$d_W(\mathbf{C}_1, \mathbf{C}_2) = \frac{1}{p} \left(\text{tr}(\mathbf{C}_1) + \text{tr}(\mathbf{C}_2) - 2 \text{tr} \left[(\mathbf{C}_1^{\frac{1}{2}} \mathbf{C}_2 \mathbf{C}_1^{\frac{1}{2}})^{\frac{1}{2}} \right] \right).$$

Prove that only the rightmost term is the challenging one to estimate, i.e., from n_i independent samples $\mathbf{X}_i = [\mathbf{x}_{i1}, \dots, \mathbf{x}_{in_i}] \in \mathbb{R}^{p \times n_i}$ and with $p \sim n_i$, show that $\frac{1}{p} \text{tr} \hat{\mathbf{C}}_i$ is a consistent estimate for $\frac{1}{p} \text{tr} \mathbf{C}_i$ with $\hat{\mathbf{C}}_i = \frac{1}{n_i} \mathbf{X}_i \mathbf{X}_i^\top$ the sample covariance estimate of \mathbf{C}_i , so that only the quantity $\frac{1}{p} \text{tr}[(\mathbf{C}_1^{\frac{1}{2}} \mathbf{C}_2 \mathbf{C}_1^{\frac{1}{2}})^{\frac{1}{2}}]$ remains to be estimated.

In the following we thus aim to estimate $d = \frac{1}{p} \text{tr}[(\mathbf{C}_1^{\frac{1}{2}} \mathbf{C}_2 \mathbf{C}_1^{\frac{1}{2}})^{\frac{1}{2}}]$. First show that $d = \int \sqrt{t} \nu_p(dt)$ with $\nu_p = \frac{1}{p} \sum_{i=1}^p \delta_{\lambda_i(\mathbf{C}_1 \mathbf{C}_2)}$ and deduce, by Cauchy's integration formula (Theorem 2.2), that $d = \frac{-1}{2\pi i} \oint_{\Gamma_\nu} \sqrt{z} m_{\nu_p}(z) dz$, with $m_{\nu_p}(z)$ the Stieltjes transform of ν_p , Γ_ν an appropriate complex contour, and \sqrt{z} some complex analytic extension of the real square root (letting $z = re^{i\theta}$ for $r \geq 0$ and $\theta \in (-\pi, \pi]$, we will consider here the principal root of z , defined as $\sqrt{z} = \sqrt{r} e^{i\theta/2}$).

With the help of Bai-Silverstein's theorem (Theorem 2.6), show that the Stieltjes transform m_{μ_p} of the spectral measure μ_p of $\hat{\mathbf{C}}_1 \hat{\mathbf{C}}_2$ relates to m_{ν_p} through the set of equations

$$zm_{\mu_p}(z) = \varphi_p(z)m_{\zeta_p}(\varphi_p(z)) + o(1) \quad (3.12)$$

$$m_{\nu_p} \left(\frac{z}{\Psi_p(z)} \right) = m_{\zeta_p}(z)\Psi_p(z) + o(1). \quad (3.13)$$

where ζ_p is the spectral measure of $\mathbf{C}_2^{\frac{1}{2}} \hat{\mathbf{C}}_1 \mathbf{C}_2^{\frac{1}{2}}$, $\Psi_p(z) \equiv 1 - \frac{p}{n_2} - \frac{p}{n_2} zm_{\zeta_p}(z)$ and $\varphi_p(z) = z/(1 - \frac{p}{n_1} - \frac{p}{n_1} zm_{\mu_p}(z))$.

By means of two successive changes of variables, prove that d can be consistently estimated, as $n, p \rightarrow \infty$, by

$$\hat{d} \equiv \frac{n_2}{2\pi i p} \oint_{\Gamma} \sqrt{\frac{\varphi_p(z)}{\psi_p(z)}} \left[\frac{\varphi'_p(z)}{\varphi_p(z)} - \frac{\psi'_p(z)}{\psi_p(z)} \right] \psi_p(z) dz$$

where $\psi_p(z) \equiv 1 - \frac{p}{n_2} - \frac{p}{n_2} zm_{\mu_p}(z)$, and Γ is some complex contour to be carefully positioned. You may refer to the steps of the derivation performed in Section 3.2 for help.

The functions $\varphi_p(z)$ and $\psi_p(z)$ are rational functions (as rational functions of m_{μ_p} , itself a relational function). By identifying zeros, poles and limiting behavior as $|z| \rightarrow \infty$, show that they can be expressed under the rational forms

$$\varphi_p(z) = z \frac{\prod_{i=1}^p z - \lambda_i}{\prod_{i=1}^p z - \eta_i}, \quad \psi_p(z) = \frac{\prod_{i=1}^p z - \xi_i}{\prod_{i=1}^p z - \lambda_i}$$

where $\lambda_1 \leq \dots \leq \lambda_p$ are the increasingly sorted eigenvalues of $\hat{\mathbf{C}}_1 \hat{\mathbf{C}}_2$, and $\{\xi_i\}_{i=1}^p$, $\{\eta_i\}_{i=1}^p$ the increasingly sorted eigenvalues of $\mathbf{\Lambda} - \frac{1}{n_1} \sqrt{\mathbf{\lambda}} \sqrt{\mathbf{\lambda}}^\top$ and $\mathbf{\Lambda} - \frac{1}{n_2} \sqrt{\mathbf{\lambda}} \sqrt{\mathbf{\lambda}}^\top$, respectively, where $\mathbf{\lambda} = (\lambda_1, \dots, \lambda_p)^\top$, $\mathbf{\Lambda} = \text{diag}(\mathbf{\lambda})$ and $\sqrt{\cdot}$ is understood entry wise.

Find the singularities, the poles and the branch cuts of the complex integrand in the expression of \hat{d} and represent them on the complex plane. You may refer to Figure 3.5, but must be extremely careful on the relative ordering of the $(\xi_i, \eta_i, \lambda_i)$ triplets. Based on this representation, and as again following Figure 3.5, deform the contour Γ into a more convenient contour for integration. Prove then that the resulting integrals over ϵ -radius circles around ξ_i are null in the small ϵ limit (using for instance the change of variable $z = \xi_i + \epsilon e^{i\theta}$). Next prove that the resulting integrals over the real axis (again in the $\epsilon \rightarrow 0$ limit) between $\xi_j + \epsilon$ and $\eta_j + \epsilon$ sum up to

$$A_1 = \frac{2n_2}{\pi p} \sum_{j=1}^p \int_{\xi_j}^{\eta_j} \sqrt{-\frac{\varphi_p(x)}{\psi_p(x)}} \psi'_p(x) dx - \frac{2n_2}{\pi p} \sum_{j=1}^p \frac{1}{\sqrt{\epsilon \frac{d}{dx} \left(\frac{1}{\varphi_p(x)\psi_p(x)} \right) (\eta_j)}}.$$

Continue the calculus by proving, using the change of variable $z = \eta_j + \epsilon e^{i\theta}$, that integrals over the ϵ -radius circles around η_j do not vanish but convey a second contribution summing up in the $\epsilon \rightarrow 0$ limit to

$$A_2 = \frac{2n_2}{\pi p} \sum_{j=1}^p \frac{1}{\sqrt{\epsilon \frac{d}{dx} \left(\frac{1}{\varphi_p(x)\psi_p(x)} \right) (\eta_j)}}.$$

Finally prove that the residues associated to the ‘‘poles’’ λ_i (that is the integral over small circles surrounding the λ_i ’s) sum up to

$$A_3 = \frac{2n_2}{p} \sqrt{\frac{n_1}{n_2}} \sum_{j=1}^p \sqrt{\lambda_j}.$$

Conclude from these three contributions that d can be estimated by the real-line integral form

$$\hat{d}(\mathbf{X}_1, \mathbf{X}_2) \equiv 2\sqrt{n_1 n_2} \frac{1}{p} \sum_{j=1}^p \sqrt{\lambda_j} + \frac{2n_2}{\pi p} \sum_{j=1}^p \int_{\xi_j}^{\eta_j} \sqrt{-\frac{\varphi_p(x)}{\psi_p(x)}} \psi'_p(x) dx.$$

Show in particular that, when $n_1 = n_2 = n/2$, this estimate further simplifies as

$$\hat{d}(\mathbf{X}_1, \mathbf{X}_2) = \frac{n}{p} \sum_{j=1}^p \left(\sqrt{\lambda_j} - \sqrt{\xi_j} \right)$$

using the fact that $\xi_j \rightarrow \eta_j$ in the limit where $n_1/n - n_2/n \rightarrow 0$, and that, by deforming the “real line” part of the contour,

$$\frac{1}{\pi} \lim_{t \rightarrow \xi_j} \int_{\xi_j}^t \sqrt{-\frac{\varphi_p(x)}{\psi_p(x)}} \psi'_p(x) dx = \frac{1}{2\pi i} \lim_{\epsilon \rightarrow 0} \oint_{\Gamma_{\xi_j}^\epsilon} \sqrt{-\varphi_p(z)\psi_p(z)} \frac{\psi'_p(z)}{\psi_p(z)} dz$$

where $\Gamma_{\xi_j}^\epsilon$ is an ϵ -radius circular contour around ξ_j .

Deduce the final expression of an n, p -consistent estimate of the Wasserstein distance for Gaussian samples, and confirm by reproducing Table 3.3.

p	$d(\mathbf{C}_1, \mathbf{C}_2)$	$\hat{d}(\mathbf{X}_1, \mathbf{X}_2)$	$d(\hat{\mathbf{C}}_1, \hat{\mathbf{C}}_2)$
2	0.0110	0.0109	0.0115
4	0.0175	0.0189	0.0203
8	0.0208	0.0213	0.0240
16	0.0225	0.0232	0.0286
32	0.0233	0.0237	0.0343
64	0.0237	0.0243	0.0454
128	0.0239	0.0240	0.0663
256	0.0240	0.0243	0.1092
512	0.0241	0.0245	0.1954

Table 3.3: Estimation of the Wasserstein distance for $\mathbf{x}_i^{(a)} \sim \mathcal{N}(\mathbf{0}, \mathbf{C}_a)$ with $[\mathbf{C}_1]_{ij} = 0.2^{|i-j|}$, $[\mathbf{C}_2]_{ij} = 0.4^{|i-j|}$, $n_1 = 1024$, $n_2 = 2048$, as a function of p , results averaged over 30 runs. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

Practical Lecture Material 2 (Robust portfolio optimization via Tyler estimator). In computational finance, one of the problems of the popular Markowitz’s mean-variance optimization framework consists in determining a portfolio vector $\mathbf{w} \in \mathbb{R}^p$, to allocate to p assets (stock market indices), in such a way to maximize the expected return and/or minimize the risk of the investment. From a statistical perspective, \mathbf{w} is thus set to minimize a certain cost function based on past observations $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ of the market evolution (the returns) of the p assets. Those are often assumed independent for simplicity, but cannot be considered Gaussian due to the possibly erratic nature of the market.

We consider here for simplicity the problem of minimizing the risk, without constraining the expected return. We assume independent (centered) elliptically distributed $\mathbf{x}_i = \sqrt{\tau_i} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$ return vectors, where $\mathbf{z}_i \in \mathbb{R}^p$ is uniform on the sphere of radius \sqrt{p} and $\tau_i > 0$ are random i.i.d. impulses independent of \mathbf{z}_i , as in (3.6). We wish to determine

$$\mathbf{w}^* = \arg \min_{\mathbf{w} \in \mathbb{R}^p, \mathbf{w}^\top \mathbf{1}_p = 1} \mathbb{E}[|\mathbf{w}^\top \mathbf{x}|^2] \quad (3.14)$$

where the constraint $\mathbf{w}^\top \mathbf{1}_p = 1$ ensures that the total investment remains constant.

Assuming that $\mathbb{E}[\tau] = 1$ (which we can set for convenience and without generality restriction), show via the Lagrangian multipliers method that the solution to (3.14) is explicitly given by

$$\mathbf{w}^* = \frac{\mathbf{C}^{-1} \mathbf{1}_p}{\mathbf{1}_p^\top \mathbf{C}^{-1} \mathbf{1}_p}$$

thus with associated minimal (expected) risk $\mathbb{E}[|(\mathbf{w}^*)^\top \mathbf{x}|^2] = \frac{1}{\mathbf{1}_p^\top \mathbf{C}^{-1} \mathbf{1}_p}$.

The covariance \mathbf{C} being unknown, and the historical returns \mathbf{x}_i being impulsive in nature, we wish to estimate \mathbf{w}^* via a robust estimator of scatter approach as in Section 3.3. That is, we replace \mathbf{C} in the formulas above by the robust (shrinkage) Tyler estimator $\hat{\mathbf{C}}(\gamma)$ defined, for $\gamma \in (\max\{0, 1 - n/p\}, 1]$, as the unique solution to

$$\hat{\mathbf{C}}(\gamma) = (1 - \gamma) \frac{1}{n} \mathbf{X} \mathbf{D}^{-1} \mathbf{X}^\top + \gamma \mathbf{I}_p, \quad \mathbf{D} = \text{diag} \left\{ \frac{1}{p} \mathbf{x}_i^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{x}_i \right\}_{i=1}^n.$$

The regularization term $\gamma \mathbf{I}_p$ allows for $p > n$ and offers an additional degree of freedom, and the choice of a Tyler estimator (i.e., $u(t) = 1/t$ in our previous notations) is made here for computational convenience.

First show that, replacing the unknown \mathbf{C} by $\hat{\mathbf{C}}(\gamma)$, letting $\hat{\mathbf{w}} = \frac{\hat{\mathbf{C}}(\gamma)^{-1} \mathbf{1}_p}{\mathbf{1}_p^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{1}_p}$, the resulting portfolio risk is given by

$$\mathbb{E}_{\mathbf{x}}[|\hat{\mathbf{w}}^\top \mathbf{x}|^2] = \frac{\mathbf{1}_p^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{C} \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{1}_p}{(\mathbf{1}_p^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{1}_p)^2} \quad (3.15)$$

and confirm that we retrieve the correct result as $\hat{\mathbf{C}}(\gamma)$ coincides with \mathbf{C} . Our objective here is to estimate this quantity to retrieve the performance of this robust portfolio design as a function of γ .

Similar to the intuitive approach developed in Section 3.3.3 for generic $u(t)$ functions (but without regularization), show that the following random equivalent asymptotics hold:

$$\begin{aligned} \|\hat{\mathbf{C}}(\gamma) - \hat{\mathbf{S}}(\gamma)\| &\xrightarrow{a.s.} 0 \\ \hat{\mathbf{S}}(\gamma) &\equiv \frac{1}{\delta(\gamma)} \frac{1 - \gamma}{1 - (1 - \gamma)c} \frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \mathbf{Z}^\top \mathbf{C}^{\frac{1}{2}} + \gamma \mathbf{I}_p \end{aligned}$$

where $c = \lim p/n$, $\mathbf{Z} = [\mathbf{z}_1, \dots, \mathbf{z}_n]$ and $\delta(\gamma)$ is the unique solution to

$$1 = \frac{1}{p} \text{tr } \mathbf{C} ((1 - \gamma) \mathbf{C} + \delta(\gamma) \gamma \mathbf{I}_p)^{-1}$$

and that we have the following deterministic equivalent for the inverse of $\hat{\mathbf{C}}(\gamma)$:

$$\hat{\mathbf{C}}(\gamma)^{-1} \leftrightarrow \left(\frac{1 - \gamma}{\delta(\gamma)} \mathbf{C} + \gamma \mathbf{I}_p \right)^{-1}.$$

To this end, one may first evaluate $\hat{\mathbf{C}}(\gamma)^{-1} - \hat{\mathbf{C}}_{-i}(\gamma)^{-1}$, where $\hat{\mathbf{C}}_{-i}$ is defined as $\hat{\mathbf{C}}$ but with the summation over $1 \leq j \neq i \leq n$, i.e.,

$$\hat{\mathbf{C}}_{-i}(\gamma) = (1 - \gamma) \frac{1}{n} \sum_{j \neq i} \frac{\mathbf{x}_j \mathbf{x}_j^\top}{\frac{1}{p} \mathbf{x}_j^\top \hat{\mathbf{C}}(\gamma) \mathbf{x}_j} + \gamma \mathbf{I}_p$$

then show that

$$\frac{1}{p} \mathbf{z}_i^\top \mathbf{C}^{\frac{1}{2}} \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i = \left(1 - (1 - \gamma) \frac{p}{n}\right) \frac{1}{p} \mathbf{z}_i^\top \mathbf{C}^{\frac{1}{2}} \hat{\mathbf{C}}_{-i}(\gamma)^{-1} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i \quad (3.16)$$

and use the intuition that

$$\frac{1}{p} \mathbf{z}_i^\top \mathbf{C}^{\frac{1}{2}} \hat{\mathbf{C}}_{-i}(\gamma)^{-1} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i \simeq \frac{1}{p} \operatorname{tr} \mathbf{C} \hat{\mathbf{C}}(\gamma)^{-1}$$

the right-end side term corresponding in the limit to $\delta(\gamma)$. Complete the proof by using appropriately Theorem 2.6.

With this result and (3.15) at hand, show that

$$\mathbb{E}[|(\hat{\mathbf{w}})^\top \mathbf{x}|^2] = \sigma^2(\gamma) + o(p^{-1}) \quad (3.17)$$

where

$$\begin{aligned} \sigma^2(\gamma) &\equiv \frac{\delta(\gamma)^2}{\delta(\gamma)^2 - c\beta(\gamma)(1 - \gamma)^2} \frac{\mathbf{1}_p^\top \left(\frac{1-\gamma}{\delta(\gamma)} \mathbf{C} + \gamma \mathbf{I}_p \right)^{-1} \mathbf{C} \left(\frac{1-\gamma}{\delta(\gamma)} \mathbf{C} + \gamma \mathbf{I}_p \right)^{-1} \mathbf{1}_p}{\left(\mathbf{1}_p^\top \left(\frac{1-\gamma}{\delta(\gamma)} \mathbf{C} + \gamma \mathbf{I}_p \right)^{-1} \mathbf{1}_p \right)^2} \\ \beta(\gamma) &\equiv \frac{1}{p} \operatorname{tr} \mathbf{C}^2 \left(\frac{1-\gamma}{\delta(\gamma)} \mathbf{C} + \gamma \mathbf{I}_p \right)^{-2}. \end{aligned}$$

To this end, one may first demonstrate (for the more technical numerator) that

$$\hat{\mathbf{C}}(\gamma)^{-1} \mathbf{C} \hat{\mathbf{C}}(\gamma)^{-1} = -\frac{d}{d\omega} \left\{ \left(\hat{\mathbf{C}}(\gamma) + \omega \mathbf{C} \right)^{-1} \right\}_{\omega=0}$$

and construct (for instance based on the proof of Theorem 2.6) a deterministic equivalent for $(\hat{\mathbf{C}}(\gamma) + \omega \mathbf{C})^{-1}$, or equivalently for $(\hat{\mathbf{S}}(\gamma) + \omega \mathbf{C})^{-1}$, which we may then differentiate and evaluate at $\omega = 0$ to retrieve the result.

Hint: In detail, obtaining this deterministic equivalent may be performed with the following steps: (i) show, using the Bai-Silverstein approach of the proof of Theorem 2.6, that

$$(\hat{\mathbf{S}} + \omega \mathbf{C})^{-1} \leftrightarrow (\alpha_\omega \mathbf{C} + \gamma \mathbf{I}_p + \omega \mathbf{C})^{-1}$$

where

$$\alpha_\omega = \frac{1 - \gamma}{\delta(\gamma)(1 - (1 - \gamma)c) + (1 - \gamma)c\Delta_\omega},$$

and Δ_ω is solution to

$$\Delta_\omega = \frac{1}{p} \operatorname{tr} \mathbf{C} (\alpha_\omega \mathbf{C} + \gamma \mathbf{I}_p + \omega \mathbf{C})^{-1}.$$

and in particular confirm that $\Delta_0 = \delta(\gamma)$ and $\alpha_0 = (1 - \gamma)/\delta(\gamma)$. Then proceed (ii) by showing that

$$\begin{aligned} & -\frac{d}{d\omega} (\hat{\mathbf{S}} + \omega \mathbf{C})^{-1} \\ & \leftrightarrow -\frac{d}{d\omega} (\alpha_\omega \mathbf{C} + \gamma \mathbf{I}_p + \omega \mathbf{C})^{-1} \\ & = \left(1 - c\alpha_\omega^2 \frac{d}{d\omega} \Delta_\omega\right) (\alpha_\omega \mathbf{C} + \gamma \mathbf{I}_p + \omega \mathbf{C})^{-1} \mathbf{C} (\alpha_\omega \mathbf{C} + \gamma \mathbf{I}_p + \omega \mathbf{C})^{-1} \end{aligned}$$

and prove that

$$\frac{d}{d\omega} \Delta_\omega = -\frac{\frac{1}{p} \operatorname{tr} \mathbf{C}^2 (\alpha_\omega \mathbf{C} + \gamma \mathbf{I}_p + \omega \mathbf{C})^{-2}}{1 - c\alpha_\omega^2 \frac{1}{p} \operatorname{tr} \mathbf{C}^2 (\alpha_\omega \mathbf{C} + \gamma \mathbf{I}_p + \omega \mathbf{C})^{-2}}.$$

Put all things together (iii) and set ω to zero to conclude.

For the expression of $\sigma^2(\gamma)$ to be of practical interest, one needs a consistent estimate for $\sigma^2(\gamma)$ for all $\gamma > 0$, upon which an estimate of the optimal γ (i.e., the one achieving the minimum estimated risk) can be obtained. We will proceed here in two steps. From (3.16), first establish that

$$\hat{\delta}(\gamma) - \frac{\delta(\gamma)}{\frac{1}{p} \operatorname{tr} \mathbf{C}} \xrightarrow{a.s.} 0$$

where

$$\begin{aligned} \hat{\delta}(\gamma) & \equiv \frac{1}{1 - (1 - \gamma) \frac{p}{n}} \frac{1}{n} \operatorname{tr} \mathbf{X}^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{X} \operatorname{diag}\{\|\mathbf{x}_i\|^{-2}\}_{i=1}^n \\ & = \frac{1}{1 - (1 - \gamma) \frac{p}{n}} \frac{1}{n} \sum_{i=1}^n \frac{\mathbf{x}_i^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{x}_i}{\|\mathbf{x}_i\|^2} = \frac{1}{1 - (1 - \gamma) \frac{p}{n}} \frac{1}{n} \sum_{i=1}^n \frac{\mathbf{x}_i^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{x}_i}{\|\mathbf{x}_i\|^2}. \end{aligned}$$

Then, establish that

$$\begin{aligned} \hat{\sigma}^2(\gamma) - \frac{\sigma^2(\gamma)}{\frac{1}{p} \operatorname{tr} \mathbf{C}} & \xrightarrow{a.s.} 0 \\ \hat{\sigma}^2(\gamma) & \equiv \frac{\hat{\delta}(\gamma)}{1 - \gamma - (1 - \gamma)^2 \frac{p}{n}} \frac{\mathbf{1}_p^\top \hat{\mathbf{C}}(\gamma)^{-1} (\hat{\mathbf{C}}(\gamma) - \gamma \mathbf{I}_p) \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{1}_p}{\left(\mathbf{1}_p^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{1}_p\right)^2} \end{aligned} \tag{3.18}$$

by developing $\hat{\mathbf{C}}(\gamma)^{-1} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i \mathbf{z}_i^\top \mathbf{C}^{\frac{1}{2}} \hat{\mathbf{C}}(\gamma)^{-1}$ (or equivalently $\hat{\mathbf{S}}(\gamma)^{-1} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i \mathbf{z}_i^\top \mathbf{C}^{\frac{1}{2}} \hat{\mathbf{S}}(\gamma)^{-1}$) as a function of the matrix form $\hat{\mathbf{S}}_{-i}(\gamma)^{-1} \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i \mathbf{z}_i^\top \mathbf{C}^{\frac{1}{2}} \hat{\mathbf{S}}_{-i}(\gamma)^{-1}$ and taking the expectation over \mathbf{z}_i , together with the asymptotic approximation

$$\mathbf{1}_n^\top \hat{\mathbf{S}}_{-i}(\gamma)^{-1} \mathbf{C} \hat{\mathbf{S}}_{-i}(\gamma)^{-1} \mathbf{1}_n \simeq \mathbf{1}_n^\top \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{C} \hat{\mathbf{C}}(\gamma)^{-1} \mathbf{1}_n.$$

Confirm the results in (3.17) and (3.18) by reproducing Figure 3.10 in the setting where $\sqrt{\tau_i} = \sqrt{\chi_d^2/d}$ for χ_d^2 a Chi-square random variable with $d = 3$ degree of freedom (so that $\mathbb{E}[\tau_i] = 1$) and $\mathbf{C} = 5 \cdot \mathbf{u}\mathbf{u}^\top + \mathbf{I}_p$ for \mathbf{u} with uniformly distributed and normalized $[\mathbf{u}]_i \sim \text{Unif}(0.5, 1.5)/\sqrt{p}$ entries.

These results may also be simulated on real financial time series from leading international markets (e.g., based on daily historical returns from the NYSE, HSI, CAC-40, etc., over a time window of typically a few years). An exhaustive analysis is provided in [Yang et al., 2015].




Figure 3.10: Portfolio risk, the asymptotic approximation $\sigma^2(\gamma)$, and the estimate $\frac{1}{p} \text{tr } \mathbf{C} \cdot \hat{\sigma}^2(\gamma)$ versus regularization penalty γ , for chi-square τ , $p = 256$ and $n = 512$. Results averaged over 50 runs. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

Chapter 4

Kernel Methods

In a broad sense, kernel methods are at the core of many, if not most, machine learning algorithms. Given a set of data $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$, most learning mechanisms rely on extracting the structural data information from direct pairwise comparisons $\kappa(\mathbf{x}_i, \mathbf{x}_j)$ for some *affinity metric* $\kappa(\cdot, \cdot)$. Gathered in an $n \times n$ matrix

$$\mathbf{K} = \{\kappa(\mathbf{x}_i, \mathbf{x}_j)\}_{i,j=1}^n$$

the “cumulative” effect of these comparisons for numerous ($n \gg 1$) data is at the heart of various supervised, semi-supervised, or unsupervised methods such as support vector machines, graph Laplacian-based learning, spectral clustering, and has deep connections to neural networks.

These applications will be thoroughly discussed in Section 4.5. For the moment though, our main interest lies in the mathematical characterization of the kernel matrix \mathbf{K} for various choices of affinity functions κ and for various statistical models of the data \mathbf{x}_i .

Clearly, from a purely machine learning perspective, the choice of the affinity function $\kappa(\cdot, \cdot)$ is central to a good performance of the learning method under study. Since real data in general have highly complex structures, a typical viewpoint is to assume that any pair of data \mathbf{x}_i and \mathbf{x}_j are not directly comparable in their ambient space but that there exists a convenient *feature extraction* function $\phi : \mathbb{R}^p \rightarrow \mathbb{R}^q$ ($q \in \mathbb{N} \cup \{+\infty\}$) such that $\phi(\mathbf{x}_i)$ and $\phi(\mathbf{x}_j)$ are more amenable to comparison. Otherwise stated, in the image of $\phi(\cdot)$, the data are more “linear” (or more “linearly separable” if one seeks to group the data in affinity classes). The simplest affinity function between \mathbf{x}_i and \mathbf{x}_j would then read $\phi(\mathbf{x}_i)^\top \phi(\mathbf{x}_j)$.

Since q is often much larger than p , the mere cost of evaluating $\phi(\mathbf{x}_i)^\top \phi(\mathbf{x}_j)$ can be deleterious to practical implementation. The so-called kernel trick is anchored in the remark that, for a wide class of such functions ϕ , $\phi(\mathbf{x}_i)^\top \phi(\mathbf{x}_j) = f(\|\mathbf{x}_i - \mathbf{x}_j\|^2)$ or $= f(\mathbf{x}_i^\top \mathbf{x}_j)$ for some function $f : \mathbb{R} \rightarrow \mathbb{R}$ and it suffices to simply evaluate $\|\mathbf{x}_i - \mathbf{x}_j\|^2$ or $\mathbf{x}_i^\top \mathbf{x}_j$ in the ambient space and then apply some f in an entry-wise manner, leading to more practically convenient methods.

Although the class of such functions f is inherently restricted by the need for a mapping ϕ to exist such that, say, $\phi(\mathbf{x})^\top \phi(\mathbf{y}) = f(\|\mathbf{x} - \mathbf{y}\|^2)$ (these are sometimes called Mercer kernel functions),¹ with time, practitioners have started to use arbitrary functions f and worked with generic kernel matrices of the form

$$\mathbf{K} = \{f(\|\mathbf{x}_i - \mathbf{x}_j\|^2)\}_{i,j=1}^n, \quad \text{or} \quad \mathbf{K} = \{f(\mathbf{x}_i^\top \mathbf{x}_j)\}_{i,j=1}^n$$

for diverse functions f . There are in particular empirical evidences showing that well-chosen “indefinite” (i.e., non-Mercer type) kernels can sometimes outperform conventional positive definite kernels that satisfy the Mercer’s condition [Haasdonk, 2005, Luss and D’aspremont, 2008].

Remark 4.1 (Typical family of functions f and finite-dimensional setting). *It is important to raise here a direct consequence of the “finite-dimensional intuitions” inherent to kernel methods in machine learning. For an affinity of the type $\kappa(\mathbf{x}_i, \mathbf{x}_j) = f(\|\mathbf{x}_i - \mathbf{x}_j\|^2)$, it is natural to assume that f be a non-increasing function, as close-by data $\mathbf{x}_i, \mathbf{x}_j$ should have a stronger affinity than distant $\mathbf{x}_i, \mathbf{x}_j$. The popular choice $f(t) = \exp(-t/2)$ (known as the Gaussian or heat kernel, related to an infinite-dimensional map ϕ) is particularly appealing as it brings arbitrarily close data to a unit affinity ($\kappa(\mathbf{x}_i, \mathbf{x}_j) \rightarrow 1$ as $\|\mathbf{x}_i - \mathbf{x}_j\| \rightarrow 0$) and far data to a null affinity ($\kappa(\mathbf{x}_i, \mathbf{x}_j) \rightarrow 0$ as $\|\mathbf{x}_i - \mathbf{x}_j\| \rightarrow \infty$).*

We will subsequently show in this section that this natural reasoning often collapses when dealing with realistic large dimensional data, leading to erroneous intuitions and disrupting many conventional ideas behind kernel-based machine learning.

4.1 Basic setting

As pointed out in Remark 4.1 and shall become evident from the coming analysis, the finite-dimensional intuition according to which f should be a “valid” Mercer function becomes rather meaningless when dealing with large dimensional data.

To fully capture this aspect, a first important consideration is to deal with “non-trivial” relative growth rates between p, n and the statistical data parameters. By non-trivial, we mean that the underlying classification or regression problem for which the kernel method is designed should neither be impossible nor too easy to solve as $p, n \rightarrow \infty$. In this section we will mostly focus on the use of kernel methods for classification, and thus the non-trivial settings are given in terms of the growth rate of the “distance” between data classes.

4.1.1 The non-trivial growth rates

In classical large- n only asymptotic statistics, laws of large numbers demand a scaling by $1/n$ of the summed observations. When centered, central limit

¹In particular, since the matrix $\{\phi(\mathbf{x}_i)^\top \phi(\mathbf{x}_j)\}_{i,j=1}^n$ is nonnegative definite, f must be such that $\{f(\|\mathbf{x}_i - \mathbf{x}_j\|^2)\}_{i,j=1}^n$ is also nonnegative definite irrespective of n and $\mathbf{x}_1, \dots, \mathbf{x}_n$.

theorems then occur after multiplication of the average by \sqrt{n} . A similar requirement is needed when we now assume that the dimension p of the data is also large. In particular, we will demand that the norm of each observation remains bounded. Assuming $\mathbf{x} \in \mathbb{R}^p$ is a vector of (bounded) entries, i.e., each of order $O(1)$ with respect to p , the correct normalization is typically \mathbf{x}/\sqrt{p} .

In the context of kernel methods, one wishes that the argument of $f(\cdot)$ in the inner-product kernel $f(\mathbf{x}_i^\top \mathbf{x}_j)$ or the distance kernel $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2)$ be of order $O(1)$, when f is assumed fixed with respect to p .

The “correct” scaling appears not to be fully immediate. Letting \mathbf{x}_i have entries of order $O(1)$, one naturally has that $\|\mathbf{x}_i - \mathbf{x}_j\|^2 = O(p)$ and thus it seems natural to scale $\|\mathbf{x}_i - \mathbf{x}_j\|^2$ by $1/p$. Similarly, if the norm of the mean $\|\mathbb{E}[\mathbf{x}_i]\|$ of \mathbf{x}_i has the same order of magnitude as $\|\mathbf{x}_i\|$ itself (as it should in general), then for $\mathbf{x}_i, \mathbf{x}_j$ independent, $\mathbb{E}[\mathbf{x}_i^\top \mathbf{x}_j] = O(p)$. So again, one should scale the inner-product also by $1/p$. Hence the kernels

$$\mathbf{K} = \left\{ f\left(\frac{1}{p}\|\mathbf{x}_i - \mathbf{x}_j\|^2\right) \right\}_{i,j=1}^n, \quad \text{and} \quad \left\{ f\left(\frac{1}{p}\mathbf{x}_i^\top \mathbf{x}_j\right) \right\}_{i,j=1}^n.$$

Section 4.2 (and most applications thereafter) will be placed under these kernel forms. The most commonly used Gaussian (or heat, or radial-basis-function) kernel, defined as $\mathbf{K} = \{\exp(-\|\mathbf{x}_i - \mathbf{x}_j\|^2/2\sigma^2)\}_{i,j=1}^n$, falls into this family as one usually demands that $\sigma^2 \sim \mathbb{E}[\|\mathbf{x}_i - \mathbf{x}_j\|^2]$ (to avoid evaluating the exponential close to zero or infinity).

However, as already demonstrated in Section 1.1.3.1, if n scales like p , then, for the resulting classification problem to be asymptotically non-trivial, the difference $\|\mathbb{E}[\mathbf{x}_i] - \mathbb{E}[\mathbf{x}_j]\|^2$ need to scale as $O(1)$ rather than $O(p)$ (otherwise data would be too easy to cluster), resulting in $\|\mathbf{x}_i - \mathbf{x}_j\|^2/p$ possibly converging to a constant value irrespective of the data classes (of \mathbf{x}_i and \mathbf{x}_j), with a typical “spread” of order $O(1/\sqrt{p})$. Similarly, up to re-centering,² $\mathbf{x}_i^\top \mathbf{x}_j/p$ scales like $O(1/\sqrt{p})$ rather than $O(1)$. As such, it seems more appropriate to normalize the kernels as

$$[\mathbf{K}]_{ij} = f\left(\frac{\|\mathbf{x}_i - \mathbf{x}_j\|^2}{\sqrt{p}} - \frac{1}{n^2} \sum_{i',j'} \frac{\|\mathbf{x}_{i'} - \mathbf{x}_{j'}\|^2}{\sqrt{p}}\right), \quad \text{or} \quad [\mathbf{K}]_{ij} = f\left(\frac{1}{\sqrt{p}} \mathbf{x}_i^\top \mathbf{x}_j\right)$$

in order here to avoid evaluating f essentially at a single value (equal to zero for the inner-product kernel and the average data distance for the distance kernel).

This “properly scaling” setting is in fact much richer than the $1/p$ normalization when n, p are of the same order of magnitude. Sections 4.3 and 4.4 elaborate on this scenario.

²More precisely, $[\mathbf{P}\mathbf{X}^\top \mathbf{X}\mathbf{P}]_{ij}$ with $\mathbf{P} = \mathbf{I}_n - \frac{1}{n}\mathbf{1}_n\mathbf{1}_n^\top$ the data centering projector.

4.1.2 Statistical data model

In the remainder of this section, we assume the observation of n independent data vectors from in total k classes gathered as $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ where

$$\begin{aligned}\mathbf{x}_1, \dots, \mathbf{x}_{n_1} &\sim \mathcal{N}(\boldsymbol{\mu}_1, \mathbf{C}_1) \\ &\vdots \quad \vdots \\ \mathbf{x}_{n-n_k+1}, \dots, \mathbf{x}_n &\sim \mathcal{N}(\boldsymbol{\mu}_k, \mathbf{C}_k)\end{aligned}$$

which is a k -class *Gaussian mixture model* (GMM) with a fixed cardinality n_1, \dots, n_k in each class. The fact that the data are labeled according to classes simplifies the notation but has no practical consequence in the analysis.

We will denote

$$\mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{C}_a) \Leftrightarrow \mathbf{x}_i \in \mathcal{C}_a$$

for $a \in \{1, \dots, k\}$ and use the matrix

$$\mathbf{J} = [\mathbf{j}_1, \dots, \mathbf{j}_k] \in \mathbb{R}^{n \times k}, \quad \mathbf{j}_a = [\underbrace{0, \dots, 0}_{n_1+\dots+n_{a-1}}, \underbrace{1, \dots, 1}_{n_a}, \underbrace{0, \dots, 0}_{n_{a+1}+\dots+n_k}]^\top$$

to summarize the class label information (which is known in a supervised learning setting and is to be recovered in unsupervised learning).

We shall systematically make the following simplifying growth rate assumption for p, n, n_1, \dots, n_k .

Assumption 1 (Growth rate of data size and number). *As $n \rightarrow \infty$, we have $p/n \rightarrow c \in (0, \infty)$ and $n_a/n \rightarrow c_a \in (0, 1)$.*

This assumption in particular implies that each class is “large” in the sense that their cardinality increases with n .³

Accordingly with the discussions in Chapter 2, from a random matrix “universal” perspective, the Gaussian mixture assumption will often (yet not always) turn out equivalent to demanding that

$$\mathbf{x}_i = \boldsymbol{\mu}_a + \mathbf{C}_a^{\frac{1}{2}} \mathbf{z}_i$$

for $\mathbf{x}_i \in \mathcal{C}_a$, with $\mathbf{z}_i \in \mathbb{R}^p$ a random vector with i.i.d. entries of zero mean, finite variance and bounded (say fourth) order moment.

This hypothesis is thus seemingly quite restrictive as it imposes that the data, up to centering and linear scaling, are composed of i.i.d. entries. Equivalently, this suggests that only data that are the resultant of affine transformations of

³If we were to relax the assumption by letting, say \mathcal{C}_a to be of much smaller cardinality than $O(n)$, it would then be necessary to counterbalance this lack of redundancy by increasing the rate of the “distances” between the statistics of \mathcal{C}_a (mean and covariance in particular) and the other classes.

vectors with i.i.d. entries can be studied. This is obviously quite restrictive in practice as “real data” are deemed much more complex.

Exploring the notion of concentrated random vectors introduced in Section 2.7, Chapter 8 will open up this discussion by showing that a much larger class of (statistical) data models embrace the same asymptotic statistics.

4.2 Distance and inner-product random kernel matrices

The most widely used kernel model in machine learning applications is the heat kernel $\mathbf{K} = \{\exp(-\|\mathbf{x}_i - \mathbf{x}_j\|^2/2\sigma^2)\}_{i,j=1}^n$, for some $\sigma > 0$. It is thus natural to start our large dimensional analysis of kernel random matrices by focusing on this model.

As mentioned in the previous sections, for the Gaussian mixture model above, as the dimension p increases, σ^2 needs to scale as $O(p)$, so say $\sigma^2 = \tilde{\sigma}^2 p$, to avoid evaluating the exponential at increasingly large values. As such, the prototypical kernel of present interest is

$$\mathbf{K} = \left\{ f \left(\frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 \right) \right\}_{i,j=1}^n$$

for f a sufficiently smooth function (e.g., $f(t) = \exp(-t/2\tilde{\sigma}^2)$ for the heat kernel).

4.2.1 Main intuition

Euclidean random matrix with equal covariances. In order to get a first picture of the large dimensional behavior of \mathbf{K} , let us develop the distance $\|\mathbf{x}_i - \mathbf{x}_j\|^2/p$ for $\mathbf{x}_i \in \mathcal{C}_a$ and $\mathbf{x}_j \in \mathcal{C}_b$, with $i \neq j$.

For simplicity, let us assume for the moment $\mathbf{C}_1 = \dots = \mathbf{C}_k = \mathbf{I}_p$ and recall the notation $\mathbf{x}_i = \boldsymbol{\mu}_a + \mathbf{z}_i$. We have, for $i \neq j$,

$$\begin{aligned} \frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 &= \frac{1}{p} \|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 + \frac{2}{p} (\boldsymbol{\mu}_a - \boldsymbol{\mu}_b)^\top (\mathbf{z}_i - \mathbf{z}_j) \\ &\quad + \frac{1}{p} \|\mathbf{z}_i\|^2 + \frac{1}{p} \|\mathbf{z}_j\|^2 - \frac{2}{p} \mathbf{z}_i^\top \mathbf{z}_j. \end{aligned}$$

For $\|\mathbf{x}_i\|$ of order $O(\sqrt{p})$, if $\|\boldsymbol{\mu}_a\| = O(\sqrt{p})$ for all $a \in \{1, \dots, k\}$ (which would be natural), then $\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2/p$ is a priori of order $O(1)$ while, by the central limit theorem, $\|\mathbf{z}_i\|^2/p = 1 + O(p^{-1/2})$. Also, again by the central limit theorem, $\mathbf{z}_i^\top \mathbf{z}_j/p = O(p^{-1/2})$ and $2(\boldsymbol{\mu}_a - \boldsymbol{\mu}_b)^\top (\mathbf{z}_i - \mathbf{z}_j)/p = O(p^{-1/2})$.

As a consequence, as $p \rightarrow \infty$, $\|\mathbf{x}_i - \mathbf{x}_j\|^2/p$ is dominated by $\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2/p + 2$ and easily discriminates classes from the pairwise observations of $\mathbf{x}_i, \mathbf{x}_j$, making the classification problem asymptotically trivial (without having to resort to any kernel method). It is thus of interest to understand how kernels come into play in the more practical scenario where the class distances are less significant.

To this end, we now demand that $\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\| = O(1)$, which is the minimal distance rate that can be discriminated from a mere Bayesian inference analysis as discussed in Section 1.1.3.1. Since the kernel operates only on the distances $\|\mathbf{x}_i - \mathbf{x}_j\|$, we may even request (up to centering all data by, say, the constant vector $\frac{1}{n} \sum_{i=1}^n n_a \boldsymbol{\mu}_a$) that $\|\boldsymbol{\mu}_a\| = O(1)$ for each a .

In this case though, $\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2/p = O(p^{-1})$ which is dominated by the noise terms $2\mathbf{z}_i^\top \mathbf{z}_j/p$ and $\|\mathbf{z}_i\|^2/p + \|\mathbf{z}_j\|^2/p - 2$, both of order $O(p^{-1/2})$. It thus seems at first that $\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\| = O(1)$ is too demanding a constraint.

However, “matrix-wise”, the situation appears to be quite different. Indeed, we have

$$\begin{aligned} \left\{ \frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 \right\}_{i,j=1}^n &= 2 \cdot \mathbf{1}_n \mathbf{1}_n^\top + \frac{1}{p} \mathbf{J} \left\{ \|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 \right\}_{i,j=1}^k \mathbf{J}^\top \\ &\quad + \boldsymbol{\psi} \mathbf{1}_n^\top + \mathbf{1}_n \boldsymbol{\psi}^\top - \frac{2}{p} \mathbf{Z}^\top \mathbf{Z} \\ &\quad + \frac{2}{p} (\mathbf{d} \mathbf{1}_n^\top + \mathbf{1}_n \mathbf{d}^\top) - \frac{2}{p} (\mathbf{J} \mathbf{M}^\top \mathbf{Z} + \mathbf{Z}^\top \mathbf{M} \mathbf{J}) - \text{diag}(\cdot) \end{aligned} \quad (4.1)$$

where $\mathbf{M} = [\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_k] \in \mathbb{R}^{p \times k}$, $\boldsymbol{\psi} \in \mathbb{R}^n$ is the vector with independent (asymptotically Gaussian) entries $\psi_i = 1 - \|\mathbf{z}_i\|^2/p = O(p^{-1/2})$, $\mathbf{d} = \text{diag}(\mathbf{J} \mathbf{M}^\top \mathbf{Z}) \in \mathbb{R}^n$ having independent entries of zero mean and variance $\|\boldsymbol{\mu}_a\|^2 = O(1)$ if the i -th entry has mean $\mathbb{E}[\mathbf{x}_i] = \boldsymbol{\mu}_a$, and the operator $\mathbf{X} - \text{diag}(\cdot)$ returns matrix \mathbf{X} with diagonal entries replaced by zeros.

From a spectral viewpoint, observe that this matrix is largely dominated by the matrix $2 \cdot \mathbf{1}_n \mathbf{1}_n^\top$ which has norm $2n$. Next in norm comes the rank-2 matrix $\boldsymbol{\psi} \mathbf{1}_n^\top + \mathbf{1}_n \boldsymbol{\psi}^\top$. The sum of these being of rank 2 (since $\mathbf{1}_n$ is common), these matrices marginally affect the spectrum of the whole matrix. What is particularly interesting now is to observe that the resulting matrix is an (in fact not quite standard) “properly normalized” spiked model, as studied in Section 2.5. Indeed, $2\mathbf{Z}^\top \mathbf{Z}/p$ is a Wishart matrix having limiting spectral measure the Marčenko-Pastur distribution with support of order $O(1)$ and all eigenvalues remaining asymptotically close to the support (Theorem 2.11); then $\mathbf{J} \left\{ \|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 \right\}_{i,j=1}^k \mathbf{J}^\top/p + 2(\mathbf{d} \mathbf{1}_n^\top + \mathbf{1}_n \mathbf{d}^\top)/p - 2(\mathbf{J} \mathbf{M}^\top \mathbf{Z} + \mathbf{Z}^\top \mathbf{M} \mathbf{J})/p$ is a rank at most $2k + 2$ matrix with the fundamental property also to be of norm $O(1)$ (which is easily verified from our assumptions).

This may seem surprising at first. Indeed, while $\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2/p = O(p^{-1})$ is largely dominated by $2\mathbf{z}_i^\top \mathbf{z}_j/p = O(p^{-1/2})$ (and thus the class information is asymptotically *not* accessible from *any* entry $\|\mathbf{x}_i - \mathbf{x}_j\|^2/p$ alone), matrix-wise, the operator norm of $\mathbf{J} \left\{ \|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 \right\}_{i,j=1}^k \mathbf{J}^\top/p$ is of the same order as that of $2\mathbf{Z}^\top \mathbf{Z}/p$. This can be understood by the “redundant” effect of the multiple copies (of order $O(n)$) of \mathbf{x}_i sharing the same mean $\boldsymbol{\mu}_a$, for each a , which together “coherently align” into a matrix with all “energy” concentrated into few nonzero eigenvalues (up to $k \ll n$); this is to be opposed to $2\mathbf{Z}^\top \mathbf{Z}/p$ the entries of which are all centered (except on the diagonal) with essentially independent fluctuations, that result in a more or less even spread of the matrix energy in its

n eigenvalues. Altogether, this redundancy effect compensates the individual weak information carried by each \mathbf{x}_i as opposed to the noise.

From the results of Section 2.5 on spiked models, it is thus expected that, as $p, n \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$, if the dominant eigenvalues of the matrix $\{\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2\}_{a,b=1}^k$ exceed a certain threshold (that depends on c), the Euclidean distance matrix of (4.1) asymptotically has isolated eigenvalues.

More importantly, the eigenvectors associated with these eigenvalues are, as a consequence, to some extent aligned to the eigenvectors of the rank- k matrix $\mathbf{J} \{\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2\}_{i,j=1}^k \mathbf{J}^\top / p$. The latter precisely lie in the span of the columns $\mathbf{j}_1, \dots, \mathbf{j}_k$ of \mathbf{J} . Thus, the (possible isolated) eigenvectors shall correlate to the linear combinations of the indicator vectors \mathbf{j}_a , which is exactly what is observed in practice: the class information can be recovered (in a fully unsupervised manner) from the dominant eigenvectors of \mathbf{K} . This suggests that, while the class of any \mathbf{x}_i cannot be retrieved from a mere pairwise comparison of the distances $\|\mathbf{x}_i - \mathbf{x}_j\|$, matrix-wise, the eigenvectors of \mathbf{K} provide this information. This remark is at the heart of the large dimensional analysis of the spectral clustering algorithms discussed in Section 4.5.1.

Including covariance structures. The kernel matrix \mathbf{K} in (4.1) has so far only been described under the setting where (i) $f(t) = t$ and (ii) $\mathbf{C}_a = \mathbf{I}_p$. A first observation is that, for $\mathbf{C}_1, \dots, \mathbf{C}_k$ distinct (from \mathbf{I}_p), we naturally have $\text{tr}(\mathbf{C}_a - \mathbf{C}_b)/p = O(1)$ (with $\text{tr} \mathbf{C}/p \leq \|\mathbf{C}\| = O(1)$) and thus, defining $\mathbf{C}^\circ = \sum_{a=1}^k \frac{n_a}{n} \mathbf{C}_a$ and $\mathbf{C}_a^\circ = \mathbf{C}_a - \mathbf{C}^\circ$ the centered covariance, we necessarily find $\text{tr} \mathbf{C}_a^\circ/p = O(1)$ for $a = 1, \dots, k$.

Accounting for covariance matrices, for $\mathbf{x}_i = \boldsymbol{\mu}_a + \mathbf{C}_a^{1/2} \mathbf{z}_i$, in the development of $\|\mathbf{x}_i - \mathbf{x}_j\|^2/p$, we thus now have $\mathbf{z}_i^\top \mathbf{C}_a \mathbf{z}_i/p + \mathbf{z}_j^\top \mathbf{C}_b \mathbf{z}_j/p = \text{tr} \mathbf{C}_a/p + \text{tr} \mathbf{C}_b/p + O(p^{-\frac{1}{2}}) = 2 \text{tr} \mathbf{C}^\circ/p + \text{tr} \mathbf{C}_a^\circ/p + \text{tr} \mathbf{C}_b^\circ/p + O(p^{-1/2})$. As a consequence, here again, $\text{tr} \mathbf{C}_a^\circ/p + \text{tr} \mathbf{C}_b^\circ/p = O(1)$ predominates the noise terms $2\mathbf{z}_i^\top \mathbf{C}_a^{1/2} \mathbf{C}_b^{1/2} \mathbf{z}_j/p = O(p^{-1/2})$ and $\mathbf{z}_i^\top \mathbf{C}_a \mathbf{z}_i/p - \text{tr} \mathbf{C}_a/p = O(p^{-1/2})$, and classification becomes trivial. For this not to arise, we further demand that $\text{tr} \mathbf{C}_a^\circ/p = O(p^{-1/2})$, i.e., that “trace-wise” the covariances $\mathbf{C}_1, \dots, \mathbf{C}_k$ are at most distinct by $O(\sqrt{p})$ rather than $O(p)$. This appears to be the correct (and again, from the analysis of Section 1.1.3, this Bayesian minimal) regime of non-trivial classification when n, p scale proportionally.

Remark that this constraint is quite interesting as, in an effort not to trivialize classification, it in turn has the effect to “entry-wise trivialize the Euclidean

distance matrix". Indeed, under this assumption, for $i \neq j$,

$$\begin{aligned} \frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 &= \frac{2}{p} \operatorname{tr} \mathbf{C}^\circ + \frac{1}{p} \|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 + \frac{1}{p} \operatorname{tr}(\mathbf{C}_a^\circ + \mathbf{C}_b^\circ) - \frac{2}{p} \mathbf{z}_i^\top \mathbf{C}_a^{\frac{1}{2}} \mathbf{C}_b^{\frac{1}{2}} \mathbf{z}_j \\ &\quad + \left(\frac{1}{p} \mathbf{z}_i^\top \mathbf{C}_a \mathbf{z}_i - \frac{1}{p} \operatorname{tr} \mathbf{C}_a \right) + \left(\frac{1}{p} \mathbf{z}_i^\top \mathbf{C}_b \mathbf{z}_i - \frac{1}{p} \operatorname{tr} \mathbf{C}_b \right) \\ &\quad + \frac{2}{p} (\boldsymbol{\mu}_a - \boldsymbol{\mu}_b)^\top (\mathbf{C}_a^{\frac{1}{2}} \mathbf{z}_i - \mathbf{C}_b^{\frac{1}{2}} \mathbf{z}_j) \\ &= \frac{2}{p} \operatorname{tr} \mathbf{C}^\circ + O(p^{-\frac{1}{2}}). \end{aligned}$$

As such, all of the entries are dominated by the constant $2 \operatorname{tr} \mathbf{C}^\circ / p$.

On the other hand, note that similar to (4.1), matrix-wise, it appears that

$$\begin{aligned} \left\{ \frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 \right\}_{i,j=1}^n &= \frac{2}{p} \operatorname{tr} \mathbf{C}^\circ \cdot \mathbf{1}_n \mathbf{1}_n^\top + \frac{1}{p} \mathbf{J} \{ \|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 \}_{i,j=1}^k \mathbf{J}^\top \\ &\quad + (\boldsymbol{\psi} + \mathbf{J}\mathbf{t}) \mathbf{1}_n^\top + \mathbf{1}_n (\boldsymbol{\psi} + \mathbf{J}\mathbf{t})^\top - \frac{2}{p} \mathbf{W}^\top \mathbf{W} \\ &\quad + \frac{2}{p} (\mathbf{d} \mathbf{1}_n^\top + \mathbf{1}_n \mathbf{d}^\top) - \frac{2}{p} (\mathbf{J} \mathbf{M}^\top \mathbf{W} + \mathbf{W}^\top \mathbf{M} \mathbf{J}) \\ &\quad - \operatorname{diag}(\cdot) \end{aligned} \tag{4.2}$$

where we denoted $\mathbf{W} = [\mathbf{C}_1^{\frac{1}{2}} \mathbf{Z}_1, \dots, \mathbf{C}_k^{\frac{1}{2}} \mathbf{Z}_k] \in \mathbb{R}^{p \times n}$, $\mathbf{t} = \{\operatorname{tr} \mathbf{C}_a^\circ / p\}_{a=1}^k \in \mathbb{R}^k$, $\boldsymbol{\psi}_i = \mathbf{z}_i^\top \mathbf{C}_a \mathbf{z}_i / p - \operatorname{tr} \mathbf{C}_a / p = O(p^{-1/2})$ for $\mathbf{x}_i \in \mathcal{C}_a$, and similar to previously used $\mathbf{d} = \operatorname{diag}(\mathbf{J} \mathbf{M}^\top \mathbf{W})$. Again, the matrix is dominated by the $O(n)$ -norm matrix $2 \operatorname{tr} \mathbf{C}^\circ / p \cdot \mathbf{1}_n \mathbf{1}_n^\top$, but the second dominant term, the $O(\sqrt{n})$ -norm rank-2 matrix $(\boldsymbol{\psi} + \mathbf{J}\mathbf{t}) \mathbf{1}_n^\top + \mathbf{1}_n (\boldsymbol{\psi} + \mathbf{J}\mathbf{t})^\top$ is now informative as $\mathbf{J}\mathbf{t}$ is block-wise composed of elements from each class. This matrix being norm-dominant (once the irrelevant rank-1 matrix $2 \operatorname{tr} \mathbf{C}^\circ / p \cdot \mathbf{1}_n \mathbf{1}_n^\top$ discarded), the vector $\boldsymbol{\psi} + \mathbf{J}\mathbf{t}$ is directly accessible (for all large p, n). Thus, a mere principle component analysis gives access to its entries (for $\mathbf{x}_i \in \mathcal{C}_a$):

$$t_a + [\boldsymbol{\psi}]_i \stackrel{\mathcal{L}}{=} \frac{1}{p} \operatorname{tr} \mathbf{C}_a^\circ + \sqrt{\frac{3}{p^2} \operatorname{tr} \mathbf{C}_a^2} \cdot \mathcal{N}(0, 1) \tag{4.3}$$

in law, where the factor 3 arises from the fourth order moment of the standard Gaussian, which follows from the central limit theorem on $\|\mathbf{z}_i\|^2$. Since both $\operatorname{tr} \mathbf{C}_a^\circ / p$ and $\sqrt{3 \operatorname{tr} \mathbf{C}_a^2 / p^2}$ are of order $O(p^{-1/2})$, non-trivial clustering can be performed based on the covariance traces directly using the second dominant eigenvector of the Euclidean matrix.

Setting $(\boldsymbol{\psi} + \mathbf{J}\mathbf{t}) \mathbf{1}_n^\top + \mathbf{1}_n (\boldsymbol{\psi} + \mathbf{J}\mathbf{t})^\top$ aside, we are then left with the smaller order terms

$$\begin{aligned} &- \frac{2}{p} \mathbf{W} \mathbf{W}^\top + \frac{1}{p} \mathbf{J} \{ \|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 \}_{i,j=1}^k \mathbf{J}^\top + \frac{2}{p} (\mathbf{d} \mathbf{1}_n^\top + \mathbf{1}_n \mathbf{d}^\top) \\ &- \frac{2}{p} (\mathbf{J} \mathbf{M}^\top \mathbf{W} + \mathbf{W}^\top \mathbf{M} \mathbf{J}) \end{aligned}$$

from which discrimination based on the means μ_1, \dots, μ_k can be performed.

This is as far as the Euclidean distance random matrix can go. In particular, consider the case where $\mu_1 = \dots = \mu_k$ and $\text{tr } \mathbf{C}_1 = \dots = \text{tr } \mathbf{C}_k$ while the $\mathbf{C}_1, \dots, \mathbf{C}_k$ are different. Then, asymptotically, no spectral information can be retrieved from the Euclidean distance matrix which can be used to discriminate the data classes (the covariance matrices appear in \mathbf{W} but the singular vectors of \mathbf{W} do not provide straightforward access to this information).

This is where the limitations of the linear kernel $f(t) = t$ first appear. To go further and be capable to classify data with different covariance structures, f must be taken nonlinear.

Non-linear kernel model. To analyze the general nonlinear kernel, we start from (4.2) of the Euclidean matrix $\{\|\mathbf{x}_i - \mathbf{x}_j\|^2/p\}_{i,j=1}^n$. We recall that, entrywise, the matrix is dominated by $\tau_p \equiv 2 \text{tr } \mathbf{C}^0/p$. As such, if f is smooth around τ_p , a Taylor-expansion can be performed to obtain (for $i \neq j$)

$$\begin{aligned} f\left(\frac{1}{p}\|\mathbf{x}_i - \mathbf{x}_j\|^2\right) &= f(\tau_p) + f'(\tau_p)\left(\frac{1}{p}\|\mathbf{x}_i - \mathbf{x}_j\|^2 - \tau_p\right) \\ &\quad + \frac{f''(\tau_p)}{2}\left(\frac{1}{p}\|\mathbf{x}_i - \mathbf{x}_j\|^2 - \tau_p\right)^2 \\ &\quad + O\left(\left(\frac{1}{p}\|\mathbf{x}_i - \mathbf{x}_j\|^2 - \tau_p\right)^3\right). \end{aligned}$$

Let us assume for the moment that $f'(\tau_p) \neq 0$, that is f does not have a minimum at (or in a vanishing vicinity of) τ_p . Since τ_p is of order $O(1)$ and f is smooth around τ_p , all derivatives $f^{(\ell)}(\tau_p)$ are of order $O(1)$. As for $(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p - \tau_p)^\ell$, from our previous calculus, it is of order $O(p^{-\ell/2})$.

As a consequence, $(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p - \tau_p)^3 = O(p^{-3/2})$ so that, in the best case the matrix $\{(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p - \tau_p)^3\}_{i,j=1}^n$ has operator norm of order $O(p^{-1/2})$. Given the presence of the full-rank “noise” term $-2f'(\tau_p)\mathbf{W}\mathbf{W}^\top/p$ in the Taylor expansion, this third-order term vanishes in operator norm. This is not the case for the second-order term that may be of operator norm of order $O(1)$ (which, as we shall see, it indeed does). This is valid provided that $f'(\tau_p) \neq 0$: the case where $f'(\tau_p) \simeq 0$ has a fundamentally different behavior and will be treated subsequently in Section 4.3.

As such, for $f'(\tau_p) \neq 0$, a second order Taylor expansion of the kernel matrix \mathbf{K} is sufficient to fully characterize the spectral behavior of \mathbf{K} in the large n, p regime.

Remark 4.2 (On data/feature centering). *Replacing all \mathbf{x}_i by $\mathbf{x}_i - \mathbf{u}$ for some fixed vector \mathbf{u} should not impede classification. As such, one may freely work with $\mathbf{x}_i^\circ = \mathbf{x}_i - \frac{1}{n} \sum_{j=1}^n \mathbf{x}_j$. This is particularly relevant when dealing with inner-product kernels of the type $f(\mathbf{x}_i^\top \mathbf{x}_j)$ since then f is applied in other positions, but is irrelevant for distance kernels since $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2) = f(\|\mathbf{x}_i^\circ - \mathbf{x}_j^\circ\|^2)$.*

Now, recall that for Mercer kernels, $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2) = \phi(\mathbf{x}_i)^\top \phi(\mathbf{x}_j)$ for some function $\phi : \mathbb{R}^p \rightarrow \mathbb{R}^q$ (say). Then it may seem natural to also center data in the feature space \mathbb{R}^q . That is, instead of working with \mathbf{K} one may equivalently work with \mathbf{PKP} where $\mathbf{P} = \mathbf{I}_n - \frac{1}{n}\mathbf{1}_n\mathbf{1}_n^\top$ is the centering operator.

This has a major technical advantage: since $\mathbf{P}\mathbf{1}_n = \mathbf{0}$, from the calculus above, many terms that are irrelevant to classification vanish and the study of \mathbf{PKP} is made simpler. It is however not so clear what the impact of this centering operation to non-Mercer kernels really is.

4.2.2 Main Results

4.2.2.1 Kernel random equivalent.

A careful derivation of all terms in the second-order Taylor expansion of \mathbf{K} above is rigorously performed in [Couillet and Benaych-Georges, 2016]. The result is summarized as follows.

Theorem 4.1 (Couillet and Benaych-Georges [2016]). *Let f be three-times continuously differentiable in the vicinity of $\tau_p = \frac{2}{p} \text{tr } \mathbf{C}^\circ$ and such that $0 < \liminf_p |f'(\tau_p)| \leq \limsup_p |f'(\tau_p)| < \infty$. Further assume the growth rates*

$$\begin{aligned} \mathbf{M} &= [\boldsymbol{\mu}_1^\circ, \dots, \boldsymbol{\mu}_k^\circ] = O_{\|\cdot\|}(1), \quad \boldsymbol{\mu}_\ell^\circ = \boldsymbol{\mu}_\ell - \sum_{a=1}^k \frac{n_a}{n} \boldsymbol{\mu}_a \\ \mathbf{t} &= [t_1, \dots, t_k]^\top = O_{\|\cdot\|}(1), \quad t_a = \frac{1}{\sqrt{p}} \text{tr}(\mathbf{C}_a - \mathbf{C}^\circ) \\ \mathbf{S} &= \{S_{ab}\}_{a,b=1}^k = O_{\|\cdot\|}(1), \quad S_{ab} = \frac{1}{p} \text{tr } \mathbf{C}_a \mathbf{C}_b. \end{aligned}$$

Then, with previous notations, as $p, n \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$ and $n_a/n \rightarrow c_a \in (0, 1)$,

$$\|\mathbf{K} - \tilde{\mathbf{K}}\| \xrightarrow{a.s.} 0$$

where $\mathbf{K} = \left\{ f\left(\frac{1}{p}\|\mathbf{x}_i - \mathbf{x}_j\|^2\right) \right\}_{i,j=1}^n$ and

$$\tilde{\mathbf{K}} = -2f'(\tau_p) \left(\frac{1}{p} \mathbf{W}^\top \mathbf{W} + \mathbf{V} \mathbf{A} \mathbf{V}^\top \right) + (f(0) - f(\tau_p) + \tau_p f'(\tau_p)) \mathbf{I}_n$$

with $\mathbf{W} = [\mathbf{W}_1, \dots, \mathbf{W}_k]$, $\mathbf{W}_a \in \mathbb{R}^{p \times n_a}$,

$$\begin{aligned} \mathbf{V} &= \left[\frac{\mathbf{J}}{\sqrt{p}}, \frac{\mathbf{W}^\top \mathbf{M}}{\sqrt{p}}, \tilde{\mathbf{v}}, \boldsymbol{\psi}, \sqrt{p}\boldsymbol{\psi}^2, \tilde{\boldsymbol{\psi}} \right], \quad \tilde{\mathbf{v}} = \left\{ \frac{1}{\sqrt{p}} \mathbf{W}_a^\top \boldsymbol{\mu}_a \right\}_{a=1}^k, \\ \tilde{\boldsymbol{\psi}} &= \text{diag}(t_a \mathbf{1}_{n_a}) \boldsymbol{\psi}, \quad \boldsymbol{\psi} = \frac{1}{p} \{ \|\mathbf{w}_i\|^2 - \mathbb{E}[\|\mathbf{w}_i\|^2] \}_{i=1}^n \end{aligned}$$

with $\psi^2 \equiv [\psi_1^2, \dots, \psi_n^2]$ and

$$\mathbf{A} = \mathbf{A}_n + \mathbf{A}_{\sqrt{n}} + \mathbf{A}_1$$

with $\mathbf{A}_{n^\alpha} \in \mathbb{R}^{(2k+4) \times (2k+4)}$ the symmetric matrices of operator norm of order $O(n^\alpha)$ described as

$$\begin{aligned} \mathbf{A}_n &= -\frac{f(\tau_p)}{2f'(\tau_p)} p \begin{bmatrix} \mathbf{1}_k \mathbf{1}_k^\top & \mathbf{0} \\ * & \mathbf{0} \end{bmatrix} \\ \mathbf{A}_{\sqrt{n}} &= -\frac{1}{2}\sqrt{p} \begin{bmatrix} \{t_a + t_b\}_{a,b=1}^k & \mathbf{0} & \mathbf{0} & \mathbf{1}_k & \mathbf{0} & \mathbf{0} \\ * & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ * & * & 0 & 0 & 0 & 0 \\ * & * & * & 0 & 0 & 0 \\ * & * & * & 0 & 0 & 0 \\ * & * & * & * & 0 & 0 \\ * & * & * & * & * & 0 \end{bmatrix} \\ \mathbf{A}_1 &= \begin{bmatrix} \mathbf{A}_{1,11} & \mathbf{I}_k & -\mathbf{1}_k & -\frac{f''(\tau_p)}{2f'(\tau_p)} \mathbf{t} & -\frac{f''(\tau_p)}{4f'(\tau_p)} \mathbf{1}_k & -\frac{f''(\tau_p)}{2f'(\tau_p)} \mathbf{1}_k \\ * & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ * & * & 0 & 0 & 0 & 0 \\ * & * & * & -\frac{f''(\tau_p)}{2f'(\tau_p)} & 0 & 0 \\ * & * & * & * & 0 & 0 \\ * & * & * & * & * & 0 \end{bmatrix} \\ \mathbf{A}_{1,11} &= \left\{ -\frac{1}{2} \|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 - \frac{f''(\tau_p)}{4f'(\tau_p)} (t_a + t_b)^2 - \frac{f''(\tau_p)}{f'(\tau_p)} S_{ab} \right\}_{a,b=1}^k. \end{aligned}$$

The first two leading-order terms in $\tilde{\mathbf{K}}$ are $\mathbf{V}\mathbf{A}_n\mathbf{V}^\top = O_{\|\cdot\|}(n)$ and $\mathbf{V}\mathbf{A}_{\sqrt{n}}\mathbf{V}^\top = O_{\|\cdot\|}(\sqrt{n})$. Note that they are (i) low rank matrices and in particular (ii) orthogonal to the projection matrix \mathbf{P} . As a consequence of Remark 4.2, the following corollary provides an asymptotic equivalent for the centered distance random kernel matrix \mathbf{PKP} that takes a much simpler form.

Corollary 4.1 (Centered distance random kernel matrix). *With the same notations and assumptions as in Theorem 4.1, as $p, n \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$ and $n_a/n \rightarrow c_a \in (0, 1)$,*

$$\|\mathbf{PKP} - \mathbf{P}\tilde{\mathbf{K}}\mathbf{P}\| \xrightarrow{a.s.} 0$$

where

$$\tilde{\mathbf{K}} = -2f'(\tau_p) \left(\frac{1}{p} \mathbf{W}^\top \mathbf{W} + \mathbf{V} \mathbf{A} \mathbf{V}^\top \right) + (f(0) - f(\tau_p) + \tau_p f'(\tau_p)) \mathbf{I}_n$$

with $\mathbf{W} = [\mathbf{W}_1, \dots, \mathbf{W}_k]$, $\mathbf{W}_a \in \mathbb{R}^{p \times n_a}$,

$$\mathbf{V} = \left[\frac{\mathbf{J}}{\sqrt{p}}, \frac{\mathbf{W}^\top \mathbf{M}}{\sqrt{p}}, \tilde{\mathbf{v}}, \psi \right], \quad \tilde{\mathbf{v}} = \left\{ \frac{1}{\sqrt{p}} \mathbf{W}_a^\top \boldsymbol{\mu}_a \right\}_{a=1}^k, \quad \psi = \frac{1}{p} \{ \|\mathbf{w}_i\|^2 - \mathbb{E}[\|\mathbf{w}_i\|^2] \}_{i=1}^n$$

and

$$\mathbf{A} = \mathbf{A}_1 = \begin{bmatrix} \mathbf{A}_{1,11} & \mathbf{I}_k & -\mathbf{1}_k & -\frac{f''(\tau_p)}{2f'(\tau_p)}\mathbf{t} \\ * & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ * & * & 0 & 0 \\ * & * & * & -\frac{f''(\tau_p)}{2f'(\tau_p)} \end{bmatrix}$$

$$\mathbf{A}_{1,11} = \left\{ -\frac{1}{2}\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|^2 - \frac{f''(\tau_p)}{2f'(\tau_p)}t_a t_b - \frac{f''(\tau_p)}{f'(\tau_p)}S_{ab} \right\}_{a,b=1}^k.$$

Remark 4.3 (Loss of first informative eigenvector due to centering). *Note that, compared to Theorem 4.1 without centering, in the asymptotic approximation of PKP in Corollary 4.1 the information on the covariance traces \mathbf{t} (in $\mathbf{A}_{\sqrt{n}}$ in Theorem 4.1), which takes the form $\boldsymbol{\psi} + \mathbf{J}\mathbf{t}$ as in (4.3), is now lost. This is, however, not really an issue: this information is readily accessible through a simple evaluation of the norms $\|\mathbf{x}_i^\circ\|^2$ with the same accuracy.*

To fully understand this result, let us make a list of successive observations:

- discarding the term proportional to \mathbf{I}_n (which follows from the treatment of the diagonal elements of \mathbf{K}), after centering, $\tilde{\mathbf{K}}$ is the sum of the full-rank $O(1)$ -norm $\frac{1}{p}\mathbf{W}^\top \mathbf{W}$ matrix and of the low rank (up to $2k+2$) matrix $\mathbf{V}\mathbf{A}\mathbf{V}^\top$. This matrix is of the family of spiked random matrix models and can be studied as per Section 2.5: its asymptotic spectrum, eigenvalue positions, phase transitions, angles between its eigenvectors and those of $\mathbf{V}\mathbf{A}\mathbf{V}^\top$, etc., can all be studied.
- the matrix \mathbf{V} is built such in a way that its vector components are $O(1)$ -norm and asymptotically “essentially” orthogonal (in the sense that $\mathbf{V}_{\cdot a}^\top \mathbf{V}_{\cdot b} \xrightarrow{a.s.} 0$ for $a \neq b$ as $p, n \rightarrow \infty$). Besides, \mathbf{V} mainly contains two types of submatrices: the class (informative) matrix \mathbf{J} and the “noise” (uninformative) remaining random vectors of zero mean. The latter are claimed uninformative in a spectral sense, as their class-wise means are all zero (only the variances of their entries depend on the classes, but a spectral method cannot directly detect this information).
- being a spiked model, the relevance of the dominant eigenvectors of $\tilde{\mathbf{K}}$ for classification depends on the ratio between the largest “informative” eigenvalues of $\mathbf{V}\mathbf{A}\mathbf{V}^\top$ and the typical spread of the eigenvalues of the noise matrix $\frac{1}{p}\mathbf{W}^\top \mathbf{W}$. The useful vector here is \mathbf{J} which contains the class information. As such, by examining the block entry $(1, 1)$ of the matrices \mathbf{A}_{n^α} above, and in particular the block entry $\mathbf{A}_{1,11}$ in the centered case, it appears that asymptotically, the eigenspectrum of the distance random kernel matrix \mathbf{K} *only* depends on:
 - the first three derivatives $f(\tau_p)$, $f'(\tau_p)$ and $f''(\tau_p)$ of the function f ;
 - the vector \mathbf{t} and the matrices \mathbf{M} and \mathbf{S} .

As a consequence of these observations, it appears that when performed on large dimensional data, spectral classification methods based on the kernel matrix \mathbf{K} do not exploit any other information than those contained in \mathbf{M} , \mathbf{t} and \mathbf{S} . Besides, since $\tilde{\mathbf{K}}$ only depends on $f(\tau_p)$, $f'(\tau_p)$ and $f''(\tau_p)$, *most* (if not all) classification methods based on \mathbf{K} asymptotically perform equivalently for f taken as a polynomial of order two having the same first derivatives at τ_p .

Now, further note that the coefficients $f(\tau_p)$, $f'(\tau_p)$ and $f''(\tau_p)$ are prefactors of \mathbf{M} , \mathbf{t} , \mathbf{S} , as well as of the (noisy) random matrix $\mathbf{W}^\top \mathbf{W}$. This leads to several fundamental remarks:

- letting $f''(\tau_p) = 0$, the term \mathbf{S} vanishes from $\tilde{\mathbf{K}}$. As such, spectral methods based on \mathbf{K} cannot distinguish different classes from their covariance “shapes” (but they can still discriminate clusters having different covariance traces, if no centering is applied). This in particular explains why spectral clustering with the linear kernel $f(t) = t$ does not allow to distinguish Gaussian mixtures of equal means and covariance traces;
- more fundamentally, the analysis reveals a non-trivial fact: letting $f'(\tau_p) \rightarrow 0$, the noisy term $-2f'(\tau_p)\mathbf{W}^\top \mathbf{W}/p$ vanishes, while the informative term $-2f'(\tau_p)\mathbf{V}\mathbf{A}\mathbf{V}^\top$ remains. Indeed, since $\mathbf{A}_{1,11}$ contains terms having $f'(\tau_p)$ in the denominator, in this $f'(\tau_p) \rightarrow 0$ limit with $\mathbf{t} = o_{\|\cdot\|}(1)$ (e.g., for normalized data vectors such that $\|\mathbf{x}_i\| = \sqrt{p}$ for all i , of particular interest in a subspace clustering context [Couillet and Kamoun, 2016]), $\tilde{\mathbf{K}}$ becomes essentially deterministic with the covariance shape information \mathbf{S} remaining (\mathbf{M} being discarded), indicating that (asymptotic) perfect classification can be achieved in this case. See also for an application to the LS-SVM classifier in Figure 4.22 later in Section 4.5.3.

The latter remark is quite surprising: it indicates that, for f a kernel function having a local extremum (minimum or maximum) at τ_p , up to data normalization, for $\mathbf{S} = O_{\|\cdot\|}(1)$ as requested in our previous “non-trivial classification” analysis, the classification does become trivially easy. This is not met for any commonly used monotonously decreasing function in the machine learning literature, such as the Gaussian kernel function $f(t) = \exp(-t/2)$. This somewhat changes the perspective of kernel methods that originally request f to define a positive definite kernel, i.e., that f be such that there exists a function $\phi : \mathbb{R}^p \rightarrow \mathbb{R}^q$ (for some $q \in \mathbb{N} \cup \{\infty\}$) for which $f(\|\mathbf{x} - \mathbf{y}\|^2) = \phi(\mathbf{x})^\top \phi(\mathbf{y})$. For f satisfying $f'(\tau_p) = 0$, it is unlikely that f defines a positive definite kernel.

Going further, the possibility to turn the non-trivial classification problem into a trivial one means that, instead of requiring $\mathbf{M}, \mathbf{t}, \mathbf{S}$ to be of order $O(1)$, it might be possible to perform classification for more stringent discriminative rates. Since \mathbf{M} and \mathbf{t} are already rate-optimal in the Neyman-Pearson test analysis, only the growth rate of \mathbf{S} can be improved. More precisely, denote $\mathbf{C}_a^\circ \equiv \mathbf{C}_a - \mathbf{C}^\circ$ and assume $\|\mathbf{C}_a^\circ\| = O_{\|\cdot\|}(p^{-1/2})$ (such that $t_a = \text{tr } \mathbf{C}_a^\circ / \sqrt{p} = O(1)$ that achieves the Bayes optimal rate), we have

$$[\mathbf{S}]_{ab} \equiv \frac{1}{p} \text{tr } \mathbf{C}_a \mathbf{C}_b = \frac{1}{p} \text{tr}(\mathbf{C}^\circ)^2 + \frac{1}{p} \text{tr } \mathbf{C}^\circ (\mathbf{C}_a^\circ + \mathbf{C}_b^\circ) + \frac{1}{p} \text{tr } \mathbf{C}_a^\circ \mathbf{C}_b^\circ \quad (4.4)$$

where we recall $\mathbf{C}^\circ \equiv \sum_{a=1}^k \frac{n_a}{n} \mathbf{C}_a = O_{\|\cdot\|}(1)$ so that $\frac{1}{p} \text{tr}(\mathbf{C}^\circ)^2 = O(1)$ and $\frac{1}{p} \text{tr} \mathbf{C}^\circ (\mathbf{C}_a^\circ + \mathbf{C}_b^\circ) \leq \|\mathbf{C}^\circ\| (t_a + t_b) / \sqrt{p} = O(p^{-1/2})$.

Based on this expansion, we shall show in the subsequent sections that, for a careful choice of f , it is indeed possible, in the $n \sim p$ regime, to perform non-trivial classification down to $\mathbf{S} = \frac{1}{p} \text{tr}(\mathbf{C}^\circ)^2 \mathbf{1}_k \mathbf{1}_k^\top + O_{\|\cdot\|}(p^{-1/2})$ (which is still not Neyman-Pearson optimal, as recall from (1.7), but likely the optimal rate of unsupervised classification methods).

Remark 4.4 (Estimation of τ_p). *The above results suggest that, depending on the discriminating information practitioners wish to emphasize, it suffices to tune the kernel function f by properly selecting its successive derivatives at τ_p . This thus demands an estimate of τ_p in the large n, p setting.*

It can be shown that

$$\frac{1}{n(n-1)} \sum_{i,j=1}^n \frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2 - \tau_p = O(p^{-1})$$

in probability. Thus τ_p is fast and easy to estimate. For future use, we raise here the importance of the small fluctuations (of order $1/p$) of the estimate which, in the above derivation in the successive orders $1, 1/\sqrt{p}, 1/p$, etc., is a small (second) order fluctuation.

Remark 4.5 (The inner-product kernel case). *The case where $\mathbf{K} = \{f((\mathbf{x}_i^\circ)^\top \mathbf{x}_j^\circ / p)\}_{i,j=1}^n$ (with \mathbf{x}_i centered as per Remark 4.2) gives an inner-product kernel that is simpler to deal with. Note that $\|\mathbf{x}_i - \mathbf{x}_j\|^2 / p = \|\mathbf{x}_i^\circ\|^2 / p + \|\mathbf{x}_j^\circ\|^2 / p - 2(\mathbf{x}_i^\circ)^\top \mathbf{x}_j^\circ / p$, the Taylor expansion of the inner-product case is now performed around the limit (equal to 0) of $(\mathbf{x}_i^\circ)^\top \mathbf{x}_j^\circ / p$, with much less terms involved than for the distance kernel model. For instance, the covariance trace information $\mathbf{t} = \{\text{tr } \mathbf{C}_a^\circ / \sqrt{p}\}_{a=1}^k$ disappears in the (asymptotic) expansion of inner-product kernel matrices and, as a consequence, the inner-product kernels are not able to separate two “nested balls” $\mathcal{N}(\mathbf{0}, (1 \pm \epsilon / \sqrt{p}) \mathbf{I}_p)$ while the distance kernel can.⁴ Figure 4.1 illustrates this remark by displaying the eigenvalues and top eigenvectors of distance and inner-product kernel matrices, respectively. While an informative spike and the associated eigenvector are observed for the distance kernel, this is not the case for the inner-product kernel.*

4.3 The α - β random kernel model

4.3.1 Motivation

We have seen above that, with the particular choice $f'(\tau_p) = 0$, the classification becomes (asymptotically) trivial if the eigenvalues of \mathbf{S} are of order $O(1)$. In a two-class setting, this is equivalent to $\text{tr}(\mathbf{C}_1 - \mathbf{C}_2)^2 / p = O(1)$ (which is also the normalized Frobenius distance $\|\mathbf{C}_1 - \mathbf{C}_2\|_F^2 / p$ between covariances \mathbf{C}_1 and

⁴Note here that, the covariance “shape” $\mathbf{S} = \mathbf{1}_2 \mathbf{1}_2^\top + O_{\|\cdot\|}(p^{-1/2})$ is not discriminative.




Figure 4.1: Eigenvalues and top eigenvector of recentered distance $f(\|\mathbf{x}_i^\circ - \mathbf{x}_j^\circ\|^2/p)$ and inner product $f((\mathbf{x}_i^\circ)^\top \mathbf{x}_j^\circ/p)$ kernel matrices \mathbf{PKP} , with $f(t) = \exp(-t/2)$, $p = 1024$, $n = 512$, $\mu = \mathbf{0}$ and $\mathbf{C}_a = (1 + 5 \cdot (-1)^a/\sqrt{p})\mathbf{I}_p$ for $a \in \{1, 2\}$, $\mathbf{x}_1, \dots, \mathbf{x}_{n/2} \in \mathcal{C}_1$ and $\mathbf{x}_{n/2+1}, \dots, \mathbf{x}_n \in \mathcal{C}_2$. [Link to code]

\mathbf{C}_2). It is thus theoretically possible to discriminate closer covariance matrices. As we shall see next, in the $n \sim p$ context, one can lower the constraint on \mathbf{S} to $\mathbf{S} = \frac{1}{p} \text{tr}(\mathbf{C}^\circ)^2 \mathbf{1}_k \mathbf{1}_k^\top + O_{\|\cdot\|}(p^{-1/2})$, that is, to be able to discriminate data classes under the weaker condition

$$\text{tr}(\mathbf{C}_a - \mathbf{C}_b)^2 = O(\sqrt{p}).$$

However, in the random kernel models discussed in Section 4.2 (with f independent of n, p), choosing $f'(\tau_p) = 0$ simultaneously discards the information in \mathbf{M} about the class means.

A careful analysis of our previous derivations reveals the fact that the information about $\mathbf{M} = O_{\|\cdot\|}(1)$ and $\mathbf{S} = \frac{1}{p} \text{tr}(\mathbf{C}^\circ)^2 \mathbf{1}_k \mathbf{1}_k^\top + O_{\|\cdot\|}(p^{-1/2})$ can be set on “even grounds” by letting f depend on p in such a way that

$$f(\tau_p) = O(1), \quad f'(\tau_p) = O(p^{-\frac{1}{2}}), \quad f''(\tau_p) = O(1).$$

That is, instead of requesting $f'(\tau_p) = 0$, we merely demand $f'(\tau_p) = \alpha/\sqrt{p}$ and $f''(\tau_p) = 2\beta$ (where the factor 2 is here for future convenience) for some $\alpha, \beta \in (0, \infty)$. Examples of such kernel functions include

$$f(t) = \beta \left(t - \tau_p + \frac{\alpha}{2\beta\sqrt{p}} \right)^2, \quad f(t) = \exp \left(- \left(t - \tau_p + \frac{\alpha}{2\beta\sqrt{p}} \right)^2 \right).$$

The first function may be seen as a generalized second-order polynomial kernel, and the second as a generalized (properly normalized) Gaussian heat-kernel.

Note importantly that, from Remark 4.4, τ_p can be estimated within $O(p^{-1})$. Therefore, writing $f'(\tau_p) \simeq f'(\hat{\tau}_p) + (\tau_p - \hat{\tau}_p)f''(\hat{\tau}_p)$ with $f'(\tau_p) = O(p^{-1/2})$ and $f''(\hat{\tau}_p) = O(1)$, the relative estimation error $(f'(\tau_p) - f'(\hat{\tau}_p))/f'(\tau_p)$ vanishes as $O(p^{-1/2})$. This means one can still accurately select f to fulfill the above conditions on derivatives, with the estimate $\hat{\tau}_p$ proposed in Remark 4.4.

For simplicity of analysis (see Remarks 4.2 and 4.5), we here consider the inner-product random kernel matrix with feature centering

$$\mathbf{PKP} = \mathbf{P} \left\{ f \left(\frac{1}{p} (\mathbf{x}_i^\circ)^\top \mathbf{x}_j^\circ \right) \right\}_{i,j=1}^n \mathbf{P} \quad (4.5)$$

where we recall that $\mathbf{x}_i^\circ = \mathbf{x}_i - \frac{1}{n} \sum_{j=1}^n \mathbf{x}_j$ and $\mathbf{P} = \mathbf{I}_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^\top$. This double-centering has the advantage of both ensuring that the inner products concentrate at 0 and also eliminating many terms in the Taylor expansion around 0 thanks to the projection matrix \mathbf{P} (as in Corollary 4.1). We now demand that f depend on p and satisfy the conditions

$$f(0) = O(1), \quad f'(0) = \frac{\alpha}{\sqrt{p}}, \quad f''(0) = 2\beta$$

with $\alpha, \beta \in \mathbb{R}$ fixed with respect to p . Here, typical kernel functions are

$$f(t) = \beta \left(t + \frac{\alpha}{2\beta\sqrt{p}} \right)^2, \quad f(t) = \exp \left(- \left(t + \frac{\alpha}{2\beta\sqrt{p}} \right)^2 \right).$$

In terms of data statistics, we consider here the same setting for means $\mathbf{M} = O_{\|\cdot\|}(1)$ as in the previous section. For covariances, recalling the expansion of $S_{ab} = \frac{1}{p} \text{tr } \mathbf{C}_a \mathbf{C}_b$ in (4.4), we introduce the *centered and rescaled* (by \sqrt{p}) covariance “shape” information

$$\mathbf{S}^\circ = \left\{ \frac{1}{\sqrt{p}} \text{tr } \mathbf{C}_a^\circ \mathbf{C}_b^\circ \right\}_{a,b=1}^k$$

for which we demand $\mathbf{S}^\circ = O_{\|\cdot\|}(1)$, rather than $O(\sqrt{p})$ as in the previous setting.

4.3.2 Main results

For the inner-product random kernel matrix \mathbf{K} above, we have the following asymptotics.

Theorem 4.2 (α - β kernel [Tiomoko Ali et al., 2018]). *Let $\mathbf{K} \in \mathbb{R}^{n \times n}$ be defined as in (4.5) with $\mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{C}_a)$ for $\mathbf{x}_i \in \mathcal{C}_a$ satisfying the following growth rate conditions*

$$\mathbf{M} = O_{\|\cdot\|}(1), \quad \mathbf{S}^\circ = \frac{1}{\sqrt{p}} \{ \text{tr } \mathbf{C}_a^\circ \mathbf{C}_b^\circ \}_{a,b=1}^k = O_{\|\cdot\|}(1)$$

Then, as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$,

$$\left\| \sqrt{p}(\mathbf{PKP} + (f(0) + (\text{tr } \mathbf{C}^\circ/p) \cdot f'(0))\mathbf{P}) - \mathbf{PK}\tilde{\mathbf{K}}\mathbf{P} \right\| \xrightarrow{a.s.} 0$$

where

$$\begin{aligned} \tilde{\mathbf{K}} &= \alpha \cdot \frac{1}{p} \mathbf{W}^\top \mathbf{W} + \beta \cdot \Phi + \mathbf{V} \mathbf{A} \mathbf{V}^\top \\ \mathbf{A} &= \begin{bmatrix} \alpha \cdot \mathbf{M}^\top \mathbf{M} + \beta \cdot \mathbf{S}^\circ & \alpha \mathbf{I}_k \\ \alpha \mathbf{I}_k & \mathbf{0} \end{bmatrix} \\ \mathbf{V} &= \begin{bmatrix} \mathbf{J} \\ \frac{\mathbf{W}^\top \mathbf{M}}{\sqrt{p}} \end{bmatrix} \\ \frac{\Phi}{\sqrt{p}} &= \left\{ \left(\frac{1}{p} \mathbf{w}_i^\top \mathbf{w}_j \right)^2 \right\}_{i,j=1}^n - \left\{ \frac{1}{p^2} \text{tr}(\mathbf{C}_a \mathbf{C}_b) \mathbf{1}_{n_a} \mathbf{1}_{n_b}^\top \right\}_{a,b=1}^k - \text{diag}(\cdot) \end{aligned}$$

where we recall that $\mathbf{Z} - \text{diag}(\cdot)$ sets the diagonal entries of the matrix \mathbf{Z} to zero.

It is worth mentioning that, similar to Remark 4.5 for “classical” inner-product kernel with f independent of p , the covariance trace information \mathbf{t} also disappear in the α - β inner-product kernel model and the covariance “shape” information \mathbf{S} in Theorem 4.1 appears here in the form of \mathbf{S}° .

It is fundamental to see here that, by reducing the amplitude of $f'(\tau_p)$ by a factor \sqrt{p} , the formerly leading noise term $\mathbf{W}^\top \mathbf{W}/p$ in Theorem 4.1 is now on even grounds to the second-order noise term Φ . Thus, \mathbf{K} can be here seen to asymptotically behave like a very special spiked model, for which the full rank (or noise) matrix is the sum $\alpha \mathbf{W}^\top \mathbf{W}/p + \beta \Phi$ constituted of the *dependent* \mathbf{W} and Φ .

Individually, $\mathbf{W}^\top \mathbf{W}/p$ has a limiting spectrum akin to the Bai-Silverstein law (Theorem 2.6), with \mathbf{C}° as population covariance (or to the Marčenko-Pastur law for $\mathbf{C}^\circ = \mathbf{I}_p$). Indeed, taking $\mathbf{S}^\circ = O_{\|\cdot\|}(1)$ ensures that the covariance matrices $\mathbf{C}_1, \dots, \mathbf{C}_k$ cannot be too different from \mathbf{C}° (e.g., all eigenvalues of $\mathbf{C}_a^\circ = \mathbf{C}_a - \mathbf{C}^\circ$ are of the order $O(p^{-1/4})$) and it can then be shown that the limiting spectrum of $\mathbf{W}^\top \mathbf{W}/p$ is the same as if all columns of \mathbf{W} have the same covariance matrix \mathbf{C}° . This largely simplifies the theoretical analysis.

As for Φ , note that it has identically distributed entries (but on the diagonal) of zero mean that are however not independent. Yet, it can be shown [Kammoun and Couillet, 2017] that the limiting spectrum of Φ is indeed a semicircle distribution, similar to the Wigner case (Theorem 2.5) where the entries are independent; there is nonetheless a major difference in the spectrum of Φ to that of a matrix with i.i.d. entries: Φ may present *up to two* isolated eigenvalues outside the semicircle bulk. This unfolds from the fact that the deterministic equivalent for the resolvent $(\Phi - z\mathbf{I}_n)^{-1}$ is of the form

$$(\Phi - z\mathbf{I}_n)^{-1} \leftrightarrow m(z)\mathbf{I}_n + \frac{\Omega^2 m^3(z)}{c^2 - \Omega^2 m^2(z)} \cdot \frac{\mathbf{1}_n \mathbf{1}_n^\top}{n} \quad (4.6)$$

where

$$\omega = \frac{\sqrt{2}}{p} \text{tr}(\mathbf{C}^\circ)^2, \quad \Omega = \sqrt{\frac{2}{p} \text{tr}(\mathbf{C}^\circ)^4}$$

and $m(z)$ is the Stieltjes transform of the (rescaled) semicircle law with support $[-2\omega/\sqrt{c}, 2\omega/\sqrt{c}]$ which is the solution to

$$m(z) = -\frac{1}{z + c^{-1}\omega^2 m(z)}$$

and admits the (limiting) spectral measure having density

$$\mu(dx) = \frac{c}{2\omega^2\pi} \sqrt{(4\omega^2/c - x^2)^+} dx.$$

In particular, if $\Omega \leq \sqrt{c}\omega$, the contribution $\mathbf{1}_n \mathbf{1}_n^\top / n$ in the deterministic equivalent does not induce a (symmetric) pair of isolated spikes in the eigenspectrum of Φ . Otherwise, spikes can be found at the position $\pm(c^{-1}\Omega + \omega^2/\Omega)$ (see for an illustration in Figure 4.2).⁵ Yet, this is of little relevance here since $\mathbf{P}\Phi\mathbf{P}$ naturally discards the contribution from $\mathbf{1}_n \mathbf{1}_n^\top$ and thus no spurious spike appears in $\mathbf{P}\Phi\mathbf{P}$.




Figure 4.2: Eigenvalues of Φ versus the (rescaled) semicircle law, for $p = 800$ and $n = 2400$. Two spikes appear here which carry *no information*, with red crosses indicating the location of $\pm(\Omega/c + \omega^2/\Omega)$. [Link to code]

Now, the main issue in determining the limiting spectral measure of $\alpha \mathbf{W}^\top \mathbf{W}/p + \beta \Phi$ (and its centered version) is to deal with the dependence between \mathbf{W} and Φ . Intuitively, note that the main “driving randomness” in $\mathbf{W}^\top \mathbf{W}/p$ are the

⁵This follows from solving the determinant equation $\det(\Phi - z\mathbf{I}_n) = 0$.

first-order fluctuations of the inner products $\mathbf{w}_i^\top \mathbf{w}_j / p$ ($i \neq j$) around 0, while for Φ this driving randomness are the *second-order fluctuations* of the type $(\mathbf{w}_i^\top \mathbf{w}_j / p)^2 - \mathbb{E}[(\mathbf{w}_i^\top \mathbf{w}_j / p)^2]$. These two fluctuations (of different order) essentially “behave” independently in the large n, p limit. Rigorously, it can be proven that the limiting spectral measure of $\alpha \mathbf{W}^\top \mathbf{W} / p + \beta \Phi$ is a so-called *free additive convolution* (recall from Definition 5) of the limiting measures of each component, i.e., the same limiting distribution as that of the *independent* sum $\alpha \mathbf{Z}_1^\top \mathbf{Z}_1 / p + \beta \mathbf{Z}_2 / \sqrt{p}$ for \mathbf{Z}_1 with i.i.d. zero mean columns of covariance \mathbf{C}° and \mathbf{Z}_2 a symmetric matrix of i.i.d. zero mean unit variance entries (up to symmetry and with zeros on the diagonal).

Precisely, we have the following result.

Theorem 4.3 (Limiting spectrum of α - β kernel). *Under the conditions of Theorem 4.2, the empirical spectral measure $\mu_{\tilde{\mathbf{K}}}$ of $\tilde{\mathbf{K}} \equiv \sqrt{p}(\mathbf{P}\mathbf{K}\mathbf{P} + (f(0) + (\text{tr } \mathbf{C}^\circ / p) \cdot f'(0))\mathbf{P})$ (after centering and rescaling as per Theorem 4.2) satisfies $\mu_{\tilde{\mathbf{K}}} - \mu \xrightarrow{a.s.} 0$ weakly, for a probability measure μ defined by its Stieltjes transform $m(z)$ as the unique solution to*

$$\frac{1}{m(z)} = -z + \frac{\alpha}{p} \text{tr } \mathbf{C}^\circ (\mathbf{I}_p + c^{-1} \alpha m(z) \mathbf{C}^\circ)^{-1} - \beta^2 c^{-1} \omega^2 m(z)$$

where we recall $\omega = \frac{\sqrt{2}}{p} \text{tr}(\mathbf{C}^\circ)^2$.

We recognize a semicircle equation for $\alpha = 0$ and a Bai-Silverstein equation for $\beta = 0$. Figure 4.3 illustrates the transition from the Marčenko-Pastur to the semicircle law limit for the eigenvalues of \mathbf{K} with different values of α, β .

Back to the statement of Theorem 4.2, note now that α and β also weigh the relative impact of the statistical means (through $\mathbf{M}^\top \mathbf{M}$) and covariances (through \mathbf{S}°) of the data. The spectrum of \mathbf{K} thus has two extreme scenarios: for $\beta = 0$, the main bulk of \mathbf{K} forms a Marčenko-Pastur distribution and isolated eigenvalues can be found only if $\mathbf{M}^\top \mathbf{M}$ is large, with the information in \mathbf{S}° unused; for $\alpha = 0$, the main bulk is a semicircle law, with isolated eigenvalues only induced by \mathbf{S}° and the information in \mathbf{M} being discarded.

This kernel is deemed “optimal” in the sense that it allows for a separation of different classes at the minimal discriminating rate for $\mathbf{M}^\top \mathbf{M}$ and quasi-optimal rate for \mathbf{S}° (as a significant improvement from the kernel models in the previous sections). This is of particular interest in scenarios where the covariance information is critical to classification, while not discarding the (usually equally important) mean statistics.

While (quasi-)optimal in its discriminating power, the “ α - β ” kernel still has its limitations: (i) it acts solely in the neighborhood of zero so that all functions f having the same derivatives at zero produce *asymptotic* equivalent kernels, regardless of their behavior on the rest of \mathbb{R} ; (ii) this differentiability request automatically discards a large class of kernel functions of more practical interest, such as the sign function, the rectified linear unit (ReLU) in neural networks, etc.




Figure 4.3: Eigenvalues of $\check{\mathbf{K}}$ as defined in Theorem 4.3 versus the limiting laws, for $p = 512$, $n = 1024$, with zero mean and identity covariance, $f(t) = \beta(t + \frac{\alpha}{2\beta}p^{-1/2})^2$. [Link to code]

The next section generalizes the idea of the “ α - β ” kernel, however with a more “proper” scaling of the kernel function. That is, we shall now demand that f operates on $\mathbf{x}_i^\top \mathbf{x}_j / \sqrt{p}$ which does not converge, rather than on $\mathbf{x}_i^\top \mathbf{x}_j / p$ (that “concentrates” around zero for large p). Quite astonishingly, we will see in Section 4.4 that the α - β kernel with f restricted to a second-order polynomial (i) corresponds to a “properly scaling” polynomial kernel and that (ii) in some respect, is in fact optimal in the class of nonlinear kernels.

4.4 Properly scaling kernel model

4.4.1 Motivation

The concentration of distances $\|\mathbf{x}_i - \mathbf{x}_j\|^2/p - \tau_p \xrightarrow{a.s.} 0$ or of inner products $\mathbf{x}_i^\top \mathbf{x}_j/p \xrightarrow{a.s.} 0$, as $n, p \rightarrow \infty$ discussed in Section 4.2 and 4.3, is advantageous from a computational perspective as it allowed for (entry-wise) Taylor expansions of nonlinear kernel functions. On the downside though, these concentration phenomena strongly restrict the effective impact of the kernel function f :

as shown previously, only the first two successive derivatives of f at the point τ_p or zero really affect the kernel (and, as a result, the kernel-based classification or regression performance).

This might be interpreted as an incorrect “scaling” of $\|\mathbf{x}_i - \mathbf{x}_j\|^2$ or $\mathbf{x}_i^\top \mathbf{x}_j$. For the latter, it is in fact rather immediate, from central limit arguments, that, assuming $\mathbb{E}[\mathbf{x}_i] = 0$ for each i , $\mathbf{x}_i^\top \mathbf{x}_j = O(\sqrt{p})$ for $i \neq j$; it is therefore more natural to evaluate f at $(\mathbf{x}_i^\circ)^\top (\mathbf{x}_j^\circ)/\sqrt{p}$ for $\mathbf{x}_i^\circ = \mathbf{x}_i - \frac{1}{n} \sum_{j=1}^n \mathbf{x}_j$. In this case, the whole support of f can be exploited. Similarly, for $\|\mathbf{x}_i - \mathbf{x}_j\|^2$, although not naturally considered in the literature, it turns out to be more appropriate to center and scale it as $(\|\mathbf{x}_i - \mathbf{x}_j\|^2 - p\tau_p)/\sqrt{p}$.

This section precisely studies this “properly scaling” kernels which, as opposed to previous models, cannot be dealt with by means of entry-wise Taylor expansions. A more refined approach, based on orthogonal polynomials for Gaussian measure (here arising as a consequence of the central limit $(\mathbf{x}_i^\circ)^\top (\mathbf{x}_j^\circ)/\sqrt{p} \xrightarrow{d} \mathcal{N}(0, \sigma^2)$), needs to be devised.

Surprisingly enough, despite the initial motivation to avoid concentrating the effect of f at a single point (τ_p or zero, so that the kernel matrix depends on f only via its successive derivative at that point), we will see in the sequel that, under the same non-trivial classification condition, the asymptotics of these “properly scaling” kernels still only depend on two or three key parameters of f and are thus essentially no more powerful than the previously studied kernels. Yet, a few advantages are worth discussing; as we shall see:

- properly scaling kernels can detect differences in covariances down to the rates of the α - β kernels discussed in Section 4.3; in this respect, they are more powerful than the conventional Gaussian heat kernels;
- we will no longer need to demand f to be smooth and in particular it will not necessarily be differentiable at τ_p or zero. This has the substantial advantage that binary kernels, i.e., kernels such that $f \in \{0, 1\}$, can be shown to meet the same asymptotic performance as the “optimal” α - β kernels when properly defined.

4.4.2 Setting

For simplicity and readability, we will exclusively focus on the inner-product kernel \mathbf{K} , the (i, j) entry of which is given by

$$\mathbf{K}_{ij} = \frac{1}{\sqrt{p}} f \left(\frac{1}{\sqrt{p}} (\mathbf{x}_i^\circ)^\top (\mathbf{x}_j^\circ) \right) \delta_{i \neq j}. \quad (4.7)$$

Note here the importance of discarding the diagonal elements, since $\|\mathbf{x}_i\|^2/\sqrt{p} = O(\sqrt{p})$, the diagonal elements essentially evaluate f at ∞ . The leading $1/\sqrt{p}$ term is here to ensure that the main support of \mathbf{K} is of order $O(1)$, for most functions f .

We will show that, under the same (optimal) growth rate assumptions on the data statistics as in the previous sections, and under mild regularity conditions

for f , the spectrum of \mathbf{K} defined in (4.7) (asymptotically) still only depends on *three* parameters, that are, however, no longer the derivatives of f at zero.

The loss of “concentration” of $(\mathbf{x}_i^\circ)^\top(\mathbf{x}_j^\circ)/\sqrt{p}$ clearly breaks down the Taylor expansion approach used so far to assess the kernel spectral behavior in the large n, p regime.

The cornerstone idea in this regime is to exploit the fact that $(\mathbf{x}_i^\circ)^\top(\mathbf{x}_j^\circ)\sqrt{p}$ converges *in law* to a Gaussian random variable so that \mathbf{K} may be viewed in the limit as a matrix with *dependent* Gaussian entries to which f is entry-wise applied. The main problem posed by this non-trivial dependence is that many elementary tools to determine the limiting eigenvalue distribution or a deterministic equivalent for the resolvent now collapse. Most of all, it is not possible to extract a row/column from \mathbf{K} (or from any simple random matrix asymptotically equivalent to \mathbf{K}) while ensuring its independence with respect to the other columns.

In the seminal works [Cheng and Singer, 2013, Do and Vu, 2013], the authors manage to work the problem around by smartly expanding f in its series of Hermite polynomials; i.e., by approximating f by a sum of *orthogonal polynomials* with respect to the Gaussian measure. In essence, the orthogonal polynomials restore the aforementioned lost independence between the rows/columns of \mathbf{K} . The need to use orthogonal polynomials with respect to the Gaussian measure arises from $(\mathbf{x}_i^\circ)^\top(\mathbf{x}_j^\circ)/\sqrt{p}$ being essentially Gaussian in the $p \rightarrow \infty$ limit.

For ease of presentation, we let here $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ be drawn independently from one of k classes $\mathcal{C}_1, \dots, \mathcal{C}_k$ (of cardinality n_1, \dots, n_k) with now

$$\mathbf{x}_i \in \mathcal{C}_a \Leftrightarrow \mathbf{x}_i = \boldsymbol{\mu}_a + (\mathbf{I}_p + \mathbf{E}_a)^{\frac{1}{2}} \mathbf{z}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{I}_p + \mathbf{E}_a) \quad (4.8)$$

so that $\mathbf{C}^\circ = \mathbf{I}_p$. The non-trivial classification assumptions under this setting are as follows (in essence, the same as for the α - β kernel in Section 4.3).

$$\begin{aligned} \mathbf{M} &= [\boldsymbol{\mu}_1^\circ, \dots, \boldsymbol{\mu}_k^\circ] = O_{\|\cdot\|}(1), \quad \boldsymbol{\mu}_\ell^\circ = \boldsymbol{\mu}_\ell - \sum_{a=1}^k \frac{n_a}{n} \boldsymbol{\mu}_a \\ \mathbf{t} &= [t_1, \dots, t_k]^\top = O_{\|\cdot\|}(1), \quad t_a = \frac{1}{\sqrt{p}} \operatorname{tr} \mathbf{C}_a^\circ = \frac{1}{\sqrt{p}} \operatorname{tr} \mathbf{E}_a \\ \mathbf{S}^\circ &= \{S_{ab}^\circ\}_{a,b=1}^k = O_{\|\cdot\|}(1), \quad S_{ab}^\circ = \frac{1}{\sqrt{p}} \operatorname{tr} \mathbf{C}_a^\circ \mathbf{C}_b^\circ = \frac{1}{\sqrt{p}} \operatorname{tr} \mathbf{E}_a \mathbf{E}_b. \end{aligned}$$

It will appear convenient in the following to first consider the “null model”, that is, for each a , $\boldsymbol{\mu}_a = \mathbf{0}$ and $\mathbf{E}_a = \mathbf{0}$, before discussing the general case.

Under the null model, we write $\mathbf{K} = \mathbf{K}_N$ which is defined as

$$[\mathbf{K}_N]_{ij} = \delta_{i \neq j} f(\mathbf{z}_i^\top \mathbf{z}_j / \sqrt{p}) / \sqrt{p} \quad (4.9)$$

for $\mathbf{z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ such that $\mathbf{z}_i^\top \mathbf{z}_j / \sqrt{p} \xrightarrow{d} \mathcal{N}(0, 1)$.

As announced, in order to analyze the spectral behavior of \mathbf{K}_N , we will resort to the theory of orthogonal polynomials and particularly of the class of Hermite polynomials [Lozier, 2003], as discussed below.

For a probability measure μ , we denote the set of orthogonal polynomials with respect to the scalar product $\langle f, g \rangle = \int f \cdot g d\mu$ as $\{P_l(x), l = 0, 1, \dots\}$, obtained from the Gram-Schmidt procedure on the monomials $\{1, x, x^2, \dots\}$ such that $P_0(x) = 1$, P_l is of degree l and $\langle P_{l_1}, P_{l_2} \rangle = \delta_{l_1-l_2}$. By the Riesz-Fisher theorem [Rudin, 1964, Theorem 11.43], for any function $f \in L^2(\mu)$, with $L^2(\mu)$ the set of square-integrable functions with respect to $\langle \cdot, \cdot \rangle$, one can formally expand f as

$$f(x) \sim \sum_{l=0}^{\infty} a_l P_l(x), \quad a_l = \int f(x) P_l(x) d\mu(x) \quad (4.10)$$

where " $f \sim \sum_{l=0}^{\infty} P_l$ " indicates that $\|f - \sum_{l=0}^N P_l\| \rightarrow 0$ as $N \rightarrow \infty$ (and $\|f\|^2 = \langle f, f \rangle$).

For our kernel matrix purpose, we demand that f is sufficiently smooth in that it can be well approximated by a sequence of orthogonal polynomials with respect to close-to-Gaussian measures.

Assumption 2. For each p , let $\xi_p = \mathbf{z}_i^T \mathbf{z}_j / \sqrt{p}$ and let $\{P_{l,p}(x), l \geq 0\}$ be the set of orthogonal polynomials with respect to the probability measure μ_p of ξ_p . For $f \in L^2(\mu_p)$ for each p , we denote

$$f(x) \sim \sum_{l=0}^{\infty} a_{l,p} P_{l,p}(x), \quad a_{l,p} = \int f(x) P_{l,p}(x) d\mu(x)$$

and we demand that

(i) $\sum_{l=0}^{\infty} a_{l,p} P_{l,p}(x) \mu_p(dx)$ converges in $L^2(\mu_p)$ to $f(x)$ uniformly over large p , i.e., for any $\epsilon > 0$ there exists L such that for all p large,

$$\left\| f - \sum_{l=0}^L a_{l,p} P_{l,p} \right\|_{L^2(\mu_p)}^2 = \sum_{l=L+1}^{\infty} |a_{l,p}|^2 \leq \epsilon,$$

(ii) as $p \rightarrow \infty$, $\sum_{l=1}^{\infty} |a_{l,p}|^2 \rightarrow \nu \in [0, \infty)$. Moreover, for $l = 0, 1, 2$, $a_{l,p}$ converges and we denote a_0 , a_1 and a_2 their limits, respectively.

(iii) $a_0 = 0$.

Since $\xi_p \rightarrow \mathcal{N}(0, 1)$, the limiting parameters a_0, a_1, a_2 and ν are indeed (generalized) moments of the standard Gaussian measure involving f . Precisely,

$$a_0 = \mathbb{E}[f(\xi)], \quad a_1 = \mathbb{E}[\xi f(\xi)], \quad \sqrt{2}a_2 = \mathbb{E}[(\xi^2 - 1)f(\xi)], \quad \nu = \text{Var}[f(\xi)] \geq a_1^2 + a_2^2 \quad (4.11)$$

for $\xi \sim \mathcal{N}(0, 1)$. These parameters will be of central significance to determine the eigenspectrum behavior of \mathbf{K} .

Item (iii) above is a simplifying assumption to avoid the existence of a constant component in all (non-diagonal) entries of \mathbf{K} . It suffices to subtract the constant $\mathbb{E}[f(\xi)]$ from f , without affecting the classification or regression performance of kernel methods.

4.4.3 Limiting spectrum of the null model

As mentioned previously, we start by investigating the null model inner-product kernel matrix $\mathbf{K} = \mathbf{K}_N$ with

$$[\mathbf{K}]_{ij} = \begin{cases} f(\mathbf{z}_i^\top \mathbf{z}_j / \sqrt{p}) / \sqrt{p} & \text{for } i \neq j \\ 0 & \text{for } i = j \end{cases}$$

for i.i.d. $\mathbf{z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$. We are, as usual, interested in the associated resolvent

$$\mathbf{Q}(z) \equiv (\mathbf{K} - z\mathbf{I}_n)^{-1}$$

Following the Marčenko-Pastur and Bai-Silverstein approaches (in Theorem 2.4 and 2.6), we first remove the i -th row and the i -th column of the symmetric matrix \mathbf{K} to decompose it as

$$\begin{aligned} \mathbf{K} &= \begin{bmatrix} \mathbf{K}_{-i} & f(\mathbf{Z}_{-i}^\top \mathbf{z}_i / \sqrt{p}) / \sqrt{p} \\ f(\mathbf{z}_i^\top \mathbf{Z}_{-i} / \sqrt{p}) / \sqrt{p} & 0 \end{bmatrix} \\ \mathbf{K}_{-i} &\equiv f(\mathbf{Z}_{-i}^\top \mathbf{Z}_{-i} / \sqrt{p}) / \sqrt{p} - \text{diag}(\cdot) \in \mathbb{R}^{(n-1) \times (n-1)} \end{aligned}$$

(that is, with zeros on the diagonal of \mathbf{K}_{-i}) where $\mathbf{Z}_{-i} \in \mathbb{R}^{p \times (n-1)}$ is the Gaussian matrix \mathbf{Z} without the i -th column \mathbf{z}_i . As such, \mathbf{K}_{-i} is independent of \mathbf{z}_i , and is asymptotically close to \mathbf{K} for n large. We similarly define the resolvent of \mathbf{K}_{-i} as

$$\mathbf{Q}_{-i} \equiv (\mathbf{K}_{-i} - z\mathbf{I}_{n-1})^{-1}.$$

With Lemma 2.6, the (i, i) -th diagonal entry of \mathbf{Q} is given by

$$\mathbf{Q}_{ii} = \frac{1}{-z - \frac{1}{p} f(\mathbf{z}_i^\top \mathbf{Z}_{-i} / \sqrt{p}) \mathbf{Q}_{-i} f(\mathbf{Z}_{-i}^\top \mathbf{z}_i / \sqrt{p})} \equiv \frac{1}{-z - \delta_i} \quad (4.12)$$

where we recall that the diagonals of both \mathbf{K} and \mathbf{K}_{-i} are all zero. To evaluate the Stieltjes transform $\frac{1}{n} \sum_{i=1}^n \mathbf{Q}_{ii}(z)$, the key object is the (nonlinear) quadratic form

$$\delta_i \equiv \frac{1}{p} f(\mathbf{z}_i^\top \mathbf{Z}_{-i} / \sqrt{p}) \mathbf{Q}_{-i} f(\mathbf{Z}_{-i}^\top \mathbf{z}_i / \sqrt{p}). \quad (4.13)$$

We then wish to relate δ_i to the (normalized) trace of \mathbf{Q}_{-i} (and then to $m(z)$), for instance in the spirit of the trace lemma, Lemma 2.11. However, here Lemma 2.11 does not apply since, the ‘leave-one-out’ kernel matrix \mathbf{K}_{-i} , and thus its resolvent \mathbf{Q}_{-i} , is *not independent* of the random vector $f(\mathbf{z}_i^\top \mathbf{Z}_{-i} / \sqrt{p}) / \sqrt{p}$.

To handle the *nonlinear* random vector $f(\mathbf{Z}_{-i}^\top \mathbf{z}_i / \sqrt{p})$, Cheng and Singer [2013] propose to perform the following orthogonal decomposition of \mathbf{z}_j :

$$\mathbf{z}_j = \alpha_j \frac{\mathbf{z}_i}{\|\mathbf{z}_i\|} + \mathbf{z}_j^\perp, \quad \alpha_j = \frac{\mathbf{z}_i^\top \mathbf{z}_j}{\|\mathbf{z}_i\|} \quad (4.14)$$

for $j \neq i$, where $\mathbf{z}_i / \|\mathbf{z}_i\|$ is the unit vector in the direction of \mathbf{z}_i and \mathbf{z}_j^\perp lies in the $(p-1)$ -dimensional subspace orthogonal to \mathbf{z}_i . Since $\mathbf{z}_i, \mathbf{z}_j$ are independent

standard Gaussian vectors, we have, in the large p limit that, $\alpha_j \sim \mathcal{N}(0, 1)$, $\mathbf{z}_j^\perp \sim \mathcal{N}(0, \mathbf{I}_{p-1})$ and they are *independent*.

The fact that α_j and \mathbf{z}_j^\perp are independent is of crucial significance in the analysis of \mathbf{K} and can be checked by showing that, conditioned on \mathbf{z}_i , they are uncorrelated Gaussian variables.

With this decomposition of \mathbf{z}_i , taking $k \neq j$ and $k \neq i$, the term $\mathbf{z}_j^\top \mathbf{z}_k$ can be expanded as

$$\mathbf{z}_j^\top \mathbf{z}_k = \alpha_j \alpha_k + (\mathbf{z}_j^\perp)^\top \mathbf{z}_k^\perp \equiv \alpha_j \alpha_k + \Phi_{jk}^\perp \quad (4.15)$$

where the cross terms in the product expansion disappear by orthogonality. Note from (4.14) that, both \mathbf{z}_j and \mathbf{z}_j^\perp are of (Euclidean) norm $O(\sqrt{p})$, while $\alpha_j \cdot \|\mathbf{z}_i\| / \|\mathbf{z}_i\| = O(1)$. Similarly, in (4.15), both $\mathbf{z}_j^\top \mathbf{z}_k$ and Φ_{jk}^\perp are of order $O(\sqrt{p})$, while $\alpha_j \alpha_k = O(1)$. In this sense, Φ_{jk}^\perp is asymptotically close to the inner product $\mathbf{z}_j^\top \mathbf{z}_k$, with only the contribution from \mathbf{z}_i excluded and explicitly given by $\alpha_j \alpha_k$.

We further denote $\boldsymbol{\alpha}_{-i} = [\alpha_1, \dots, \alpha_{i-1}, \alpha_{i+1}, \dots, \alpha_n]^\top \in \mathbb{R}^{n-1}$ and $\mathbf{K}_{-i}^\perp \in \mathbb{R}^{(n-1) \times (n-1)}$ with its (j, k) entry given by

$$[\mathbf{K}_{-i}^\perp]_{jk} \equiv \delta_{j \neq k} \cdot f((\mathbf{z}_j^\perp)^\top \mathbf{z}_k^\perp / \sqrt{p}) / \sqrt{p} = \delta_{j \neq k} \cdot f(\Phi_{jk}^\perp / \sqrt{p}) / \sqrt{p} \quad (4.16)$$

so that the nonlinear random vector $f(\mathbf{Z}_{-i}^\top \mathbf{z}_i / \sqrt{p}) = f(\boldsymbol{\alpha}_{-i} \|\mathbf{z}_i\| / \sqrt{p}) \simeq f(\boldsymbol{\alpha}_{-i})$ for p large. It is also worth remarking here that, the random vector $\boldsymbol{\alpha}_{-i}$ is merely a standard Gaussian random vector $\boldsymbol{\alpha}_{-i} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_{n-1})$ in the large n, p limit in the sense that each entry is asymptotically Gaussian and uncorrelated.

The advantage of introducing Φ^\perp and \mathbf{K}_{-i}^\perp is that, $\boldsymbol{\alpha}_{-i}$ is “essentially” asymptotically independent of Φ^\perp in the sense that the expectations $\mathbb{E}[\Phi^\perp \boldsymbol{\alpha}_{-i}]$ and $\mathbb{E}[\mathbf{K}_{-i}^\perp \boldsymbol{\alpha}_{-i}]$ vanish in the large n, p limit. This is, however, not the case for the (original) “leave-one-out” kernel matrix \mathbf{K}_{-i} as previously discussed in (4.13), i.e., $\|\mathbb{E}[\mathbf{K}_{-i} \boldsymbol{\alpha}_{-i}]\| \neq 0$.

The study of \mathbf{K}_{-i} thus boils down to that of \mathbf{K}_{-i}^\perp , we will need in the remainder its resolvent

$$\mathbf{Q}_{-i}^\perp \equiv (\mathbf{K}_{-i}^\perp - z \mathbf{I}_{n-1})^{-1}$$

that is therefore also “asymptotically” independent of $\boldsymbol{\alpha}_{-i}$.

With these preliminary derivations, we now focus on the “leave-one-out” kernel matrix $\mathbf{K}_{-i} \equiv f(\mathbf{Z}_{-i}^\top \mathbf{Z}_{-i} / \sqrt{p}) / \sqrt{p} - \text{diag}(\cdot)$. With (4.15), its (k, l) -entry is given by

$$[\mathbf{K}_{-i}]_{jk} = \frac{1}{\sqrt{p}} f \left(\frac{1}{\sqrt{p}} \alpha_j \alpha_k + \frac{1}{\sqrt{p}} \Phi_{jk}^\perp \right)$$

where we recall that $\Phi_{jk}^\perp = O(\sqrt{p})$, $\alpha_j \alpha_k = O(1)$ and they are (asymptotically) independent. As a consequence, with a Taylor expansion of $f(\alpha_j \alpha_k / \sqrt{p} +$

Φ_{jk}^\perp/\sqrt{p}) around the leading term Φ_{jk}^\perp/\sqrt{p} , we obtain⁶

$$f\left(\frac{1}{\sqrt{p}}\alpha_j\alpha_k + \frac{1}{\sqrt{p}}\Phi_{jk}^\perp\right) = f\left(\frac{1}{\sqrt{p}}\Phi_{jk}^\perp\right) + f'\left(\frac{1}{\sqrt{p}}\Phi_{jk}^\perp\right)\frac{1}{\sqrt{p}}\alpha_j\alpha_k + O(p^{-1})$$

so that, for $j \neq k$,

$$\begin{aligned} [\mathbf{K}_{-i}]_{jk} &= \frac{1}{\sqrt{p}}f\left(\frac{1}{\sqrt{p}}\Phi_{jk}^\perp\right) + \frac{a_1}{p}\alpha_j\alpha_k + \frac{1}{p}g\left(\frac{1}{\sqrt{p}}\Phi_{jk}^\perp\right)\alpha_j\alpha_k + O(p^{-3/2}) \\ &= \left[\mathbf{K}_{-i}^\perp + \frac{a_1}{p}\boldsymbol{\alpha}_{-i}\boldsymbol{\alpha}_{-i}^\top + \frac{1}{\sqrt{p}}\text{diag}(\boldsymbol{\alpha}_{-i})\mathbf{G}\text{diag}(\boldsymbol{\alpha}_{-i})\right]_{jk} + O(p^{-3/2}) \end{aligned}$$

with the shortcut $g(x) = f'(x) - a_1$, for a_1 the first Hermite coefficient in (4.11), \mathbf{K}_{-i}^\perp defined in (4.16), and another ‘kernel matrix’ \mathbf{G} with its (j, k) -entry given by

$$\mathbf{G}_{jk} \equiv g(\Phi_{jk}^\perp/\sqrt{p})/\sqrt{p}$$

with kernel function $g(x) = f'(x) - a_1$ so that in particular, $\mathbb{E}_{\xi \sim \mathcal{N}(0,1)}[g(\xi)] = 0$.

Note that we intentionally extract the constant a_1 from the nonlinear function $f'(x)$ since

- for $\boldsymbol{\alpha}_{-i} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_{n-1})$ in the large n, p limit, we have $\frac{1}{p}\boldsymbol{\alpha}_{-i}\boldsymbol{\alpha}_{-i}^\top = O_{\|\cdot\|}(1)$;
- on the other hand, for $\mathbf{G}-\text{diag}(\cdot) = O_{\|\cdot\|}(1)$, we have $\frac{1}{\sqrt{p}}\text{diag}(\boldsymbol{\alpha}_{-i})\mathbf{G}\text{diag}(\boldsymbol{\alpha}_{-i}) = O_{\|\cdot\|}(p^{-1/2})$ that is of vanishing operator norm as $n, p \rightarrow \infty$.

The fact that $\mathbf{G}-\text{diag}(\cdot) = O_{\|\cdot\|}(1)$ is closely related to Item (iii) of Assumption 2: for $a_0 = \mathbb{E}[f(\xi)] = 0$, it can indeed be shown that the kernel matrix \mathbf{K} has bound operator norm (see, e.g., [Fan and Montanari, 2019, Theorem 1.7]) for n, p large. And similarly for the (slightly different) kernel matrix $\mathbf{G}-\text{diag}(\cdot)$ with $\mathbb{E}[g(\xi)] = 0$. It allows us to conclude that, in matrix form,

$$\mathbf{K}_{-i} = \mathbf{K}_{-i}^\perp + \frac{a_1}{p}\boldsymbol{\alpha}_{-i}\boldsymbol{\alpha}_{-i}^\top + o_{\|\cdot\|}(1) \quad (4.17)$$

with $\boldsymbol{\alpha}_{-i}$ ‘essentially’ asymptotically independent of \mathbf{K}_{-i}^\perp .

As a consequence of the above approximation, we have, for $\mathbf{Q}_{-i} \equiv (\mathbf{K}_{-i} - z\mathbf{I}_{n-1})^{-1}$,

$$\begin{aligned} \mathbf{Q}_{-i} &= \left(\mathbf{K}_{-i}^\perp + \frac{a_1}{p}\boldsymbol{\alpha}_{-i}\boldsymbol{\alpha}_{-i}^\top - z\mathbf{I}_{n-1}\right)^{-1} + o_{\|\cdot\|}(1) \\ &= \mathbf{Q}_{-i}^\perp - \frac{a_1\mathbf{Q}_{-i}^\perp \frac{1}{p}\boldsymbol{\alpha}_{-i}\boldsymbol{\alpha}_{-i}^\top \mathbf{Q}_{-i}^\perp}{1 + \frac{a_1}{p}\boldsymbol{\alpha}_{-i}^\top \mathbf{Q}_{-i}^\perp \boldsymbol{\alpha}_{-i}} + o_{\|\cdot\|}(1) \\ &= \mathbf{Q}_{-i}^\perp - \frac{a_1\mathbf{Q}_{-i}^\perp \frac{1}{p}\boldsymbol{\alpha}_{-i}\boldsymbol{\alpha}_{-i}^\top \mathbf{Q}_{-i}^\perp}{1 + \frac{a_1}{p}\text{tr } \mathbf{Q}_{-i}^\perp} + o_{\|\cdot\|}(1) \end{aligned}$$

⁶We consider for the moment f to be a Hermite polynomial (and thus differentiable), and extend to square-summable f (with respect to Gaussian measure) under Assumption 2.

where we recall the definition $\mathbf{Q}_{-i}^\perp \equiv (\mathbf{K}_{-i}^\perp - z\mathbf{I}_{n-1})^{-1}$ that is asymptotically independent of $\boldsymbol{\alpha}_{-i}$. Here we used Lemma 2.8 for the second line and Lemma 2.11 for the approximation in the third line. Also, with (4.17) and Lemma 2.9 we deduce

$$\frac{1}{p} \operatorname{tr} \mathbf{Q}_{-i}^\perp = \frac{1}{p} \operatorname{tr} \mathbf{Q}_{-i} + o(1) = \frac{1}{p} \operatorname{tr} \mathbf{Q} + o(1) \simeq \frac{n}{p} m(z) \equiv \frac{1}{p} \operatorname{tr} \bar{\mathbf{Q}}(z)$$

for $\bar{\mathbf{Q}}(z)$ a deterministic equivalent for $\mathbf{Q}(z)$ (the form of which to be specified) and $m(z) = \frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}}(z)$.

To form an equation for $m(z)$, we would like to use (4.12) and it remains to investigate the nonlinear quadratic form $\delta_i = \frac{1}{p} f(\mathbf{z}_i^\top \mathbf{Z}_{-i}/\sqrt{p}) \mathbf{Q}_{-i} f(\mathbf{Z}_{-1}^\top \mathbf{z}_i/\sqrt{p})$ in (4.13). Note from (4.14) that, for $j \neq i$,

$$\begin{aligned} f(\mathbf{z}_j^\top \mathbf{z}_i/\sqrt{p}) &= f(\alpha_j \|\mathbf{z}_i\|/\sqrt{p}) = f(\alpha_j) + f'(\alpha_j) \cdot \alpha_j (\|\mathbf{z}_i\|/\sqrt{p} - 1) + O(p^{-1}) \\ &= f(\alpha_j) + O(p^{-1/2}) \end{aligned}$$

and therefore

$$\begin{aligned} \delta_i &= \frac{1}{p} f(\boldsymbol{\alpha}_{-i})^\top \mathbf{Q}_{-i}^\perp f(\boldsymbol{\alpha}_{-i}) - a_1 \frac{\left(\frac{1}{p} \boldsymbol{\alpha}_{-i}^\top \mathbf{Q}_{-i}^\perp f(\boldsymbol{\alpha}_{-i})\right)^2}{1 + a_1 \cdot \frac{n}{p} m(z)} + o(1) \\ &= \frac{a_1^2}{p} \boldsymbol{\alpha}_{-i}^\top \mathbf{Q}_{-i}^\perp \boldsymbol{\alpha}_{-i} + \frac{1}{p} f_{>1}(\boldsymbol{\alpha}_{-i}) \mathbf{Q}_{-i}^\perp f_{>1}(\boldsymbol{\alpha}_{-i}) - a_1 \frac{\left(\frac{a_1}{p} \boldsymbol{\alpha}_{-i}^\top \mathbf{Q}_{-i}^\perp \boldsymbol{\alpha}_{-i}\right)^2}{1 + a_1 \cdot \frac{n}{p} m(z)} + o(1) \end{aligned}$$

where we decompose again $f(x)$ as the sum of its linear part $a_1 x$ and its ‘purely’ nonlinear part $f_{>1}(x) = f(x) - a_1 x$ that is *orthogonal* to $a_1 x$ in the sense that $\mathbb{E}[\xi f_{>1}(\xi)] = \langle x, f_{>1}(x) \rangle = 0$. As a consequence, we have

$$\begin{aligned} \frac{1}{p} \boldsymbol{\alpha}_{-i}^\top \mathbf{A} \boldsymbol{\alpha}_{-i} &= \frac{1}{p} \operatorname{tr} \mathbf{A} + o(1) \\ \frac{1}{p} f_{>1}(\boldsymbol{\alpha}_{-i})^\top \mathbf{A} f_{>1}(\boldsymbol{\alpha}_{-i}) &= \operatorname{Var}[f_{>1}(\alpha_j)] \cdot \frac{1}{p} \operatorname{tr} \mathbf{A} + o(1) = (\nu - a_1^2) \cdot \frac{1}{p} \operatorname{tr} \mathbf{A} + o(1) \\ \frac{1}{p} \boldsymbol{\alpha}_{-i}^\top \mathbf{A} f_{>1}(\boldsymbol{\alpha}_{-i}) &= o(1) \end{aligned}$$

for \mathbf{A} independent of $\boldsymbol{\alpha}_{-i}$ of bounded operator norm, where we recall the definition $\nu = \operatorname{Var}[f(\xi)]$ from (4.11). This leads to the following approximation for the quadratic form

$$\delta_i = \frac{1}{p} f(\mathbf{z}_i^\top \mathbf{Z}_{-i}/\sqrt{p}) \mathbf{Q}_{-i} f(\mathbf{Z}_{-1}^\top \mathbf{z}_i/\sqrt{p}) = \frac{a_1^2 \cdot \frac{n}{p} m(z)}{1 + a_1 \cdot \frac{n}{p} m(z)} + (\nu - a_1^2) \cdot \frac{n}{p} m(z) + o(1).$$

Ultimately, recalling (4.12), we deduce

$$m(z) = \frac{1}{n} \sum_{i=1}^n \frac{1}{-z + \delta_i} + o(1) = \frac{1}{-z - \frac{(a_1^2/c)m(z)}{1+(a_1/c)m(z)} - \frac{\nu-a_1^2}{c}m(z)} + o(1)$$

which entails the following result.

Theorem 4.4 (Limiting spectrum of properly scaling kernel, [Cheng and Singer, 2013, Do and Vu, 2013]). Let $p/n \rightarrow c \in (0, \infty)$ and Assumption 2 hold. Then, the empirical spectral measure of the null model \mathbf{K}_N defined in (4.9) converges weakly and almost surely to a probability measure μ defined by its Stieltjes transform $m(z)$ as the unique solution to

$$-\frac{1}{m(z)} = z + \frac{a_1^2 m(z)}{c + a_1 m(z)} + \frac{\nu - a_1^2}{c} m(z). \quad (4.18)$$

Despite derived here only in the Gaussian case, Theorem 4.4 holds more generally for the large family of sub-exponential distribution (having finite higher order moments) [Do and Vu, 2013]. While universality is classical in random matrix results, with mostly first and second order statistics involved, the present universality result is much less obvious, due to the nonlinearity and the strong dependence (between entries) in the model under study.

Remark 4.6 (Connection to α - β kernel). Note that, taking $f(t) = \beta t^2 + \alpha t + \alpha^2/(4\beta)$ in (4.9), one obtains, up to centering, the second-order polynomial α - β kernel discussed in Section 4.3. In a sense, the properly scaling model is a natural extension of the α - β kernel model, allowing for a much larger family of nonlinear functions f (including even non-smooth function).

Similar to the α - β kernel model in Theorem 4.3, here the limiting spectral measure μ of the null model \mathbf{K}_N is the *free additive convolution* (denoted as ‘ \boxplus ’, see Definition 5) between the Marčenko–Pastur law (denote $\mu_{MP,c}$ of shape parameter $c = \lim p/n$) and the semicircle law (μ_{SC}) as

$$\mu = a_1(\mu_{MP,c^{-1}} - 1) \boxplus \sqrt{(\nu - a_1^2)c^{-1}}\mu_{SC}$$

where we denote $a_1(\mu_{MP,c^{-1}} - 1)$ the law of $a_1(x - 1)$ for $x \sim \mu_{MP,c^{-1}}$ and $\sqrt{(\nu - a_1^2)c^{-1}}\mu_{SC}$ the law of $\sqrt{(\nu - a_1^2)c^{-1}} \cdot x$ for $x \sim \mu_{SC}$. Intuitively speaking, the Marčenko–Pastur law characterizes the linear part (a_1x) of the kernel function $f(x)$, while the higher-order “purely” nonlinear part $f(x) - a_1x$ contributes to the semicircle law. These two contributions are asymptotically “independent” so that the resulting limiting spectrum is the free additive convolution of each component. As an illustration, Figure 4.4 compares, for $f(t) = \tanh(t)$, the empirical spectral measure of the null model \mathbf{K}_N to the limiting law μ in Theorem 4.4 that appears to be the “sum” of a Marčenko–Pastur and a semicircle distribution.

It is worth mentioning that Theorem 4.4 only characterizes the limiting eigenspectrum and does not provide a description of the possible spikes. Again, akin to the α - β kernel (recall Figure 4.2), the null model \mathbf{K}_N may present *up to two* isolated eigenvalues outside the limiting support in Theorem 4.4. This happens only when the second Hermite coefficient $a_2 \neq 0$ and is precisely described in the following theorem.




Figure 4.4: Eigenvalues of \mathbf{K}_N versus the limiting law from Theorem 4.4, for $p = 800$, $n = 3200$ and $f(t) = \tanh(t)$. [Link to code]

Theorem 4.5 (Deterministic equivalent for properly scaling kernel, from [Fan and Montanari, 2019]). *With the notations and assumptions in Theorem 4.4, we have*

$$(\mathbf{K}_N - z\mathbf{I}_n)^{-1} \leftrightarrow m(z)\mathbf{I}_n + \frac{a_2^2(m_4 - 1)m^3(z)}{2c^2 - a_2^2(m_4 - 1)m^2(z)} \cdot \frac{1}{n}\mathbf{1}_n\mathbf{1}_n^\top$$

with $m(z)$ the solution to (4.18) and m_4 the fourth moment of the entries of \mathbf{z} .

As a result of Theorem 4.5, if $a_2 = 0$ or $m_4 = 1$ (for instance for Bernoulli ± 1 entries), there is asymptotically no spike outside the limiting support of the null model \mathbf{K}_N . With $a_2 \neq 0$ and $m_4 > 1$, however, one may have up to two spikes at

$$\lambda = \mp \frac{a_2}{c\delta} \mp \frac{a_2\nu \pm a_1(\nu - a_1^2)\delta}{a_2(a_2 \pm a_1\delta)}\delta, \quad \delta = \sqrt{\frac{2}{m_4 - 1}}$$

which results from solving the determinant equation $\det(\mathbf{K}_N - z\mathbf{I}_n) = 0$.

Note that (i) for $a_1 = 0$, the two spikes are at the position $\pm \left(\frac{a_2}{c\delta} + \frac{\nu\delta}{a_2} \right)$ and are symmetric as in the case of the α - β kernel; we recall that in this case the limiting spectrum is a rescaled semicircle law

$$\mu(dx) = \frac{c}{2\nu\pi} \sqrt{(4\nu/c - x^2)^+} dx$$

and (ii) while the limiting spectrum is universal with respect to the distribution (of the entries), here the spike *does* depend on the fourth moment of the distribution. In Figure 4.5 we observe a farther (left) spike for the Student-t distribution (left) than for the Gaussian distribution (right), with the same limiting law.

In particular, in the case of the Gaussian distribution one has $m_4 = 3$ and by taking $a_2 = \sqrt{2}$ one obtains the same rank-one structure as in (4.6), while the limiting spectrum can be very different.

** we may want to update the reference and add the phase transition etc later**




Figure 4.5: Eigenvalues of \mathbf{K}_N versus the limiting laws and spikes in Theorem 4.5, for Student-t (with 7 degrees of freedom, **left**) and Gaussian distribution (**right**) with $p = 512$, $n = 2048$, and $f(t) = \max(t, 0)$. The position of the isolated (non-informative) eigenvalue *depends* on the kurtosis of the random matrix entries. [Link to code]

4.4.4 Main results

Having covered the analysis of the (pure-noise) null mode kernel matrix \mathbf{K}_N , we will establish, in this section, an “information-plus-noise” asymptotic equivalent for the kernel matrix \mathbf{K} , again under the non-trivial classification assumptions for the mixture $\mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{I}_p + \mathbf{E}_a) \in \mathcal{C}_a$, $a \in \{1, \dots, k\}$.

The main idea for this “information-plus-noise” decomposition comes in two steps: (i) first, by an expansion of $\mathbf{x}_i^\top \mathbf{x}_j$ as a function of $\mathbf{z}_i, \mathbf{z}_j$ and the statistical mixture model parameters $\boldsymbol{\mu}, \mathbf{E}$, one can decompose $\mathbf{x}_i^\top \mathbf{x}_j$ into successive orders of magnitudes with respect to p ; this, as we will show, further allows for a Taylor expansion of $f(\mathbf{x}_i^\top \mathbf{x}_j / \sqrt{p})$ for at least twice differentiable functions f around its dominant term $f(\mathbf{z}_i^\top \mathbf{z}_j / \sqrt{p})$. Then, (ii) we rely on the orthogonal polynomial approach of the previous section to “linearize” the resulting matrix terms $\{f(\mathbf{x}_i^\top \mathbf{x}_j / \sqrt{p})\}$, $\{f'(\mathbf{x}_i^\top \mathbf{x}_j / \sqrt{p})\}$ and $\{f''(\mathbf{x}_i^\top \mathbf{x}_j / \sqrt{p})\}$ (all terms corresponding to higher order derivatives asymptotically vanish) and use Assumption 2 to extend the result to all square-summable f .

The main conclusion is that the kernel matrix \mathbf{K} asymptotically behaves like the sum $\tilde{\mathbf{K}} = \mathbf{K}_N + \tilde{\mathbf{K}}_I$ of the full-rank “noise” matrix \mathbf{K}_N (characterized in Theorem 4.4 and 4.5) and a low-rank “information” matrix $\tilde{\mathbf{K}}_I$. This is stated in the following theorem.

Theorem 4.6 (Properly scaling kernel, [Liao and Couillet, 2019a]). *Let Assumption 2 hold and let $\mathbf{K} \in \mathbb{R}^{n \times n}$ be the properly scaling kernel defined in (4.7) with $\mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{I}_p + \mathbf{E}_a)$, for $\mathbf{x}_i \in \mathcal{C}_a$ satisfying the following growth rate conditions*

$$\mathbf{M} = O_{\|\cdot\|}(1), \quad \mathbf{t} = O_{\|\cdot\|}(1), \quad \mathbf{S}^\circ = O_{\|\cdot\|}(1).$$

Then, as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$,

$$\|\mathbf{K} - \tilde{\mathbf{K}}\| \xrightarrow{a.s.} 0, \quad \tilde{\mathbf{K}} = \mathbf{K}_N + \mathbf{V}\mathbf{A}\mathbf{V}^\top$$

with \mathbf{K}_N defined in (4.9) and

$$\begin{aligned} \mathbf{A} &= \begin{bmatrix} a_1 \cdot \mathbf{M}^\top \mathbf{M} + \frac{a_2}{\sqrt{2}} \cdot (\mathbf{t}\mathbf{1}_k^\top + \mathbf{1}_k \mathbf{t}^\top + \mathbf{S}^\circ) & a_1 \mathbf{I}_k \\ a_1 \mathbf{I}_k & \mathbf{0} \end{bmatrix} \\ \mathbf{V} &= \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{p}}, & \frac{\mathbf{Z}^\top \mathbf{M}}{\sqrt{p}} \end{bmatrix} \end{aligned}$$

where we recall the first two Hermite coefficients $a_1 = \mathbb{E}[\xi f(\xi)]$ and $a_2 = \mathbb{E}[(\xi^2 - 1)f(\xi)]/\sqrt{2}$ for $\xi \sim \mathcal{N}(0, 1)$.

Interestingly, different from the noisy spikes in Theorem 4.5 and Figure 4.5, which depends on the fourth moment of the distribution, here the informative spikes in Theorem 4.6 are shown to be universal [Liao and Couillet, 2019a], as in the case of limiting spectrum in Theorem 4.4.

Figure 4.6 compares the spectra, and in particular the isolated eigenvalues of \mathbf{K} and $\tilde{\mathbf{K}}$ for $f(t) = \text{sign}(t), |t|$ and $\max(t, 0)$, for random vectors having Gaussian or Student-t entries. These numerical evidences validate Theorem 4.6. Note that we have $a_2 = 0$ for $f(t) = \text{sign}(t)$ and $a_1 = 0$ for $f(t) = |t|$. As a result, different types of spikes (due to means and covariances) are present. More generally, for f odd ($f(-t) = -f(t)$), $a_2 = 0$ and the statistical information on covariances (through \mathbf{E}) asymptotically vanishes in \mathbf{K} ; for f even ($f(-t) = f(t)$), $a_1 = 0$ and information about the means $\boldsymbol{\mu}$ vanishes. Thus, only f neither odd nor even can preserve both first and second order discriminating statistics (e.g., the popular ReLU function $f(t) = \max(t, 0)$). Besides, these isolated eigenvalues are informative and, the corresponding eigenvectors are, as expected, noisy versions of linear combinations of the columns of \mathbf{J} .

Practical consequences: universality of binary kernels. As a direct consequence of Theorem 4.6, similar to the α - β kernel, the performance of spectral methods with properly scaling kernel only depends on the three parameters of the nonlinear function f : $a_1 = \mathbb{E}[\xi f(\xi)]$, $a_2 = \mathbb{E}[\xi^2 f(\xi)]/\sqrt{2}$ and $\nu = \mathbb{E}[f^2(\xi)]$. More precisely, the parameters a_1, ν determine the limiting spectral measure while a_1, a_2 determine the low rank informative structure.

Therefore, arbitrary (square-summable) kernel functions f (with $a_0 = 0$) are asymptotically equivalent to the simple cubic function $c_3 x^3 + c_2 x^2 + c_1 x - c_2$ having the same Hermite polynomial coefficients a_1, a_2, ν , with the coefficients being related through

$$a_1 = 3c_3 + c_1, \quad a_2 = \sqrt{2}c_2, \quad \nu = (3c_3 + c_1)^2 + 6c_3^2 + 2c_2^2.$$




Figure 4.6: Eigenvalues of \mathbf{K} (blue) versus the limiting laws and spikes of $\tilde{\mathbf{K}}$ in Theorem 4.6 (red), for $p = 1024$, $n = 2048$, with two classes $n_1 = n_2 = n/2$ and $\boldsymbol{\mu}_1 = -[2; \mathbf{0}_{p-1}] = -\boldsymbol{\mu}_2$ and $\mathbf{E}_1 = -5 \cdot \mathbf{I}_p / \sqrt{p} = -\mathbf{E}_2$. [Link to code]

It is thus possible to design a prototypical family \mathcal{F} of functions f having (i) universal properties with respect to (a_1, a_2, ν) , i.e., for each (a_1, a_2, ν) there exists $f \in \mathcal{F}$ with these Hermite coefficients and (ii) having numerically advantageous properties. Thus, any arbitrary kernel function f can be mapped, through (a_1, a_2, ν) , to a function in \mathcal{F} with good numerical properties. One such prototypical family \mathcal{F} is the set of f , parametrized by (t, s_-, s_+) , and defined as

$$f(x) = \begin{cases} -rt & x \leq \sqrt{2}s_- \\ 0 & \sqrt{2}s_- < x \leq \sqrt{2}s_+ \\ t & x > \sqrt{2}s_+ \end{cases}, \quad \begin{cases} a_1 = \frac{t}{\sqrt{2\pi}}(e^{-s_+^2} + re^{-s_-^2}) \\ a_2 = \frac{t}{\sqrt{2\pi}}(s_+e^{-s_+^2} + rs_-e^{-s_-^2}) \\ \nu = \frac{t^2}{2}(1 - \text{erf}(s_+))(1 + r) \end{cases} \quad (4.19)$$

where $r \equiv \frac{1 - \text{erf}(s_+)}{1 + \text{erf}(s_-)}$. Figure 4.7 displays such f in (4.19) together with the cubic function $c_3x^3 + c_2(x^2 - 1) + c_1x$ sharing the same coefficients (a_1, a_2, ν) , with the gains in storage size shown in Table 4.1.

The class of equivalence of kernel functions induced by this mapping (i.e.,

Table 4.1: Storage size (in Mb) of \mathbf{K} for piecewise constant and cubic f , in the setting of Figure 4.6.

f	Size
\mathcal{F} -family	4.15
Cubic	16.75




Figure 4.7: Piecewise constant \mathcal{F} -family (green) versus cubic (blue) function with equal (a_1, a_2, ν) .

those having asymptotically equivalent spectral properties) is quite unlike the equivalence class of the previous section for the “improper” scaling $f(\mathbf{x}_i^\top \mathbf{x}_j/p)$ regime. In the latter, functions $f(x)$ of the same class of equivalence are those having common $f'(0)$ and $f''(0)$ values while here these functions may have no similar local behavior (as shown in the example of Figure 4.7).

** performance and complexity trade-off TBD here! **

4.5 Implications to kernel methods

By simply “plugging” the random matrix equivalents of the kernel matrices studied in the previous sections into kernel-based learning algorithms, it is now possible to analyze the asymptotic performance of these algorithms. The present section is dedicated to this analysis, successively for unsupervised (kernel spectral clustering in Section 4.5.1), semi-supervised (with kernel graph Laplacian in Section 4.5.2), and fully supervised (kernel ridge regression in Section 4.5.3) learning.

We shall discover that, as a result of the curse of dimensionality $\|\mathbf{x}_i - \mathbf{x}_j\|^2/p \xrightarrow{a.s.} \tau_p$ and of the resulting inappropriate (low dimensional) intuitions in the large dimensional setting, all these algorithms (i) behave differently from what is expected, (ii) sometimes fail to perform as intended and, (iii) are often far from optimal. Luckily, the random matrix analyses in previous section provide new intuitions and allow for a proper adaptation (such as improved hyperparameter setting) and improvement of the algorithms.

4.5.1 Application to kernel spectral clustering

From a machine learning perspective, spectral clustering is often seen as a *discrete-to-continuous relaxation* of a graph min-cut problem [Luxburg, 2007]. More precisely, assuming \mathbf{K} to be the adjacency matrix of a graph with nodes $\mathbf{x}_1, \dots, \mathbf{x}_n$ and edges $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$, the min-cut problem consists in determining a k -class partition $\mathcal{S}_1 \cup \dots \cup \mathcal{S}_k$ of $\{1, \dots, n\}$ that minimizes the affinity

across classes, i.e.,

$$(\mathcal{S}_1, \dots, \mathcal{S}_k) \in \arg \min_{\mathcal{S}_1 \cup \dots \cup \mathcal{S}_k = \{1, \dots, n\}} \sum_{a=1}^k \sum_{\substack{i \in \mathcal{S}_a \\ j \notin \mathcal{S}_a}} \frac{f(\frac{1}{p} \|\mathbf{x}_i - \mathbf{x}_j\|^2)}{|\mathcal{S}_a|} \quad (4.20)$$

where the division by the cardinality $|\mathcal{S}_a|$ ensures that classes have approximately balanced weights (this is formally known as the *ratio-cut* adaptation of the original min-cut problem for which the denominator is simply 1). This optimization problem has been shown to be equivalent to finding the orthonormal matrix $\mathbf{S} = [\mathbf{s}_1, \dots, \mathbf{s}_k] \in \mathbb{R}^{n \times k}$ (i.e., $\mathbf{S}^\top \mathbf{S} = \mathbf{I}_k$) with columns defined as $[\mathbf{s}_a]_i = |\mathcal{S}_a|^{-\frac{1}{2}} \cdot \delta_{i \in \mathcal{S}_a}$ which minimize

$$\text{tr } \mathbf{S}^\top (\mathbf{D} - \mathbf{K}) \mathbf{S}$$

where $\mathbf{D} = \text{diag}(\mathbf{K} \mathbf{1}_n)$. Solving this discrete problem is NP-hard [Luxburg, 2007], but relaxing \mathbf{S} to be merely an orthonormal matrix with no structure constraint gives the straightforward solution that \mathbf{S} is the collection of the k eigenvectors associated to the smallest eigenvalues of $\mathbf{D} - \mathbf{K}$. This precisely leads to the spectral clustering algorithm.

Yet, performing spectral clustering directly on $\mathbf{D} - \mathbf{K}$ is observed to lead to poor performance in practice. It has instead been proposed to replace $|\mathcal{S}_a|$ in the min-cut cost function by $\text{vol}(\mathcal{S}_a) = \sum_{i \in \mathcal{S}_a} \mathbf{D}_{ii}$ which is the total weight of the edges connecting the nodes of class \mathcal{S}_a (rather than the number of nodes). With this normalization, the problem now becomes equivalent to minimizing

$$\text{tr } \mathbf{S}^\top (\mathbf{I}_n - \mathbf{D}^{-\frac{1}{2}} \mathbf{K} \mathbf{D}^{-\frac{1}{2}}) \mathbf{S}$$

which led to the most popular Ng-Weiss-Jordan spectral clustering algorithm [Ng et al., 2002].

From a random matrix standpoint, the aforementioned heuristic considerations to choose whether 1, $|\mathcal{S}_a|$ or $\text{vol}(\mathcal{S}_a)$ as normalization is somewhat irrelevant, as we now know that the large dimensional behavior of \mathbf{K} is prone to many erroneous intuitive interpretations. Indeed, the above reasoning is fundamentally based on the fact that f is a decreasing “affinity” function and that $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$ ought to be large for $\mathbf{x}_i, \mathbf{x}_j$ in the same class, and small otherwise; however, at least as far as non-trivial large dimensional Gaussian mixtures are concerned, this intuition is inappropriate.

Instead, we now need to resort to our novel “large dimensional spectral intuition”, for the moment in the case of standard “improperly scaling” kernels (such as the popular Gaussian kernel).

4.5.1.1 The case of standard distance-kernels

We have seen in Theorem 4.1 that the dominant eigenvectors of \mathbf{K} contain class label information \mathbf{J} and can thus be used for spectral clustering. Yet, \mathbf{K}

has the inconvenience that its first two dominant eigenvalues scale like $O(n)$ and $O(\sqrt{n})$. As for the matrix $\mathbf{D} - \mathbf{K}$, it can be readily seen as quite inappropriate for clustering. Indeed, note that, while the informative spectrum of \mathbf{K} is essentially of order $O(1)$ (if we exclude the little informative two dominant eigenvectors), the matrix \mathbf{D} has diagonal elements

$$[\mathbf{D}]_{ii} = nf(\tau_p) + \zeta_i + O(1), \quad \zeta_i = nf'(\tau_p)[\psi]_i = O(\sqrt{n})$$

where the random ζ_i terms are “essentially” of zero mean and asymptotically independent across i . Consequently, the spectrum of $\mathbf{D} - \mathbf{K}$ is largely dominated by the non-informative diagonal elements of \mathbf{D} . As such, $\mathbf{D} - \mathbf{K}$ is not appropriate for spectral clustering, as confirmed by empirical results.

The matrix $\mathbf{D}^{-\frac{1}{2}}\mathbf{K}\mathbf{D}^{-\frac{1}{2}}$ advocated by Ng-Weiss-Jordan is more interesting. First, since $[\mathbf{D}]_{ii} = O(n)$, it is convenient to consider the normalized matrix

$$\mathbf{L} = n\mathbf{D}^{-\frac{1}{2}}\mathbf{K}\mathbf{D}^{-\frac{1}{2}}. \quad (4.21)$$

Note that, $\mathbf{D}^{\frac{1}{2}}\mathbf{1}_n$ is an eigenvector for \mathbf{L} with corresponding eigenvalue n , since

$$n\mathbf{D}^{-\frac{1}{2}}\mathbf{K}\mathbf{D}^{-\frac{1}{2}}(\mathbf{D}^{\frac{1}{2}}\mathbf{1}_n) = \mathbf{D}^{-\frac{1}{2}}\mathbf{K}\mathbf{1}_n = n\mathbf{D}^{-\frac{1}{2}}\mathbf{D}\mathbf{1}_n = n\mathbf{D}^{\frac{1}{2}}\mathbf{1}_n.$$

This is also the largest eigenvalue of \mathbf{L} . Moreover, and quite surprisingly, a thorough Taylor expansion of $\mathbf{D}^{-\frac{1}{2}}$ pre- and post-multiplying the random equivalent $\tilde{\mathbf{K}}$ of \mathbf{K} , reveals that

$$\mathbf{L}' \equiv n\mathbf{D}^{-\frac{1}{2}}\mathbf{K}\mathbf{D}^{-\frac{1}{2}} - n\frac{\mathbf{D}^{\frac{1}{2}}\mathbf{1}_n\mathbf{1}_n^\top\mathbf{D}^{\frac{1}{2}}}{\mathbf{1}_n^\top\mathbf{D}\mathbf{1}_n} = n\mathbf{D}^{-\frac{1}{2}}\left(\mathbf{K} - \frac{\mathbf{d}\mathbf{d}^\top}{\mathbf{1}_n^\top\mathbf{D}\mathbf{1}_n}\right)\mathbf{D}^{-\frac{1}{2}} \quad (4.22)$$

with $\mathbf{d} = \mathbf{D}\mathbf{1}_n$, is asymptotically (with high probability) of bounded operator norm. That is, both matrices \mathbf{A}_n and $\mathbf{A}_{\sqrt{n}}$ (of operator norm of order $O(n)$ and $O(\sqrt{n})$, respectively) from Theorem 4.1 asymptotically disappear after normalization by $\mathbf{D}^{-\frac{1}{2}}$ and projection against the dominant eigen-direction $\mathbf{D}^{\frac{1}{2}}\mathbf{1}_n$. This makes \mathbf{L}' easier to handle mathematically (as no spurious eigenvalue evades from the spectrum at a fast rate) and more “stable” from a statistical viewpoint.

Precisely, we have the following result.

Theorem 4.7 (Random equivalent of normalized Laplacian [Coullet and Benaych-Georges, 2016]). *Under the notations and assumptions of Theorem 4.1, for \mathbf{L}' defined in (4.22), we have*

$$\begin{aligned} & \|\mathbf{L}' - \tilde{\mathbf{L}}'\| \xrightarrow{a.s.} 0 \\ & \frac{\mathbf{D}^{\frac{1}{2}}\mathbf{1}_n}{\sqrt{\mathbf{1}_n^\top\mathbf{D}\mathbf{1}_n}} = \frac{\mathbf{1}_n}{\sqrt{n}} + \frac{1}{n\sqrt{c}} \frac{f'(\tau_p)}{2f(\tau_p)} [\{t_a\mathbf{1}_{n_a}\}_{a=1}^k + \boldsymbol{\psi}] + o(n^{-1}) \end{aligned}$$

where

$$\tilde{\mathbf{L}}' = -2 \frac{f'(\tau_p)}{f(\tau_p)} \left(\frac{1}{p} \mathbf{W}^\top \mathbf{W} + \mathbf{U} \mathbf{B} \mathbf{U}^\top \right) + \frac{f(0) - f(\tau_p) + \tau_p f'(\tau_p)}{f(\tau_p)} \mathbf{I}_n$$

and

$$\begin{aligned}\mathbf{U} &= \left[\frac{\mathbf{J}}{\sqrt{p}}, \frac{\mathbf{W}^\top \mathbf{M}}{\sqrt{p}}, \boldsymbol{\psi} \right] \\ \mathbf{B} &= \begin{bmatrix} \mathbf{B}_{11} & \mathbf{I}_k - \mathbf{1}_k \mathbf{c}^\top & \left(\frac{5f'(\tau_p)}{8f(\tau_p)} - \frac{f''(\tau_p)}{2f'(\tau_p)} \right) \mathbf{t} \\ * & \mathbf{0} & \mathbf{0} \\ * & * & \frac{5f'(\tau_p)}{8f(\tau_p)} - \frac{f''(\tau_p)}{2f'(\tau_p)} \end{bmatrix} \\ \mathbf{B}_{11} &= \mathbf{M}^\top \mathbf{M} + \left(\frac{5f'(\tau_p)}{8f(\tau_p)} - \frac{f''(\tau_p)}{2f'(\tau_p)} \right) \mathbf{t} \mathbf{t}^\top - \frac{f''(\tau_p)}{f'(\tau_p)} \frac{\mathbf{S}^\circ}{\sqrt{p}} \\ &\quad + c \cdot \frac{f(0) - f(\tau_p) + \tau_p f'(\tau_p)}{2f'(\tau_p)} \mathbf{1}_k \mathbf{1}_k^\top \\ \frac{\mathbf{S}^\circ}{\sqrt{p}} &= \left\{ \frac{1}{p} \operatorname{tr} \mathbf{C}_a^\circ \mathbf{C}_b^\circ \right\}_{a,b=1}^k, \quad \mathbf{C}_a^\circ = \mathbf{C}_a - \mathbf{C}^\circ\end{aligned}$$

with $\mathbf{c} = [c_1, \dots, c_k]^\top$.

It first provides an explicit characterization of the (exactly known) dominant eigenvector of \mathbf{L}' associated with the eigenvalue n : up to a dominant constant $1/\sqrt{n}$, the eigenvector entries contain small deviations (of order $1/n$) that are discriminative class information when the t_a 's are of order $O(1)$. These information can be exploited for clustering: indeed, although the deviations t_a/n are small compared to the dominant term $1/\sqrt{n}$, the latter is strictly constant and is thus merely a constant shift and is inconsequential to classification. However, if the t_a 's are equal or only differ by $o(1)$, these information are “buried” in the noisy zero-mean and asymptotically Gaussian vector $\boldsymbol{\psi}$ and cannot be used for clustering.

The projection \mathbf{L}' of the normalized Laplacian matrix \mathbf{L} to the space orthogonal to $\mathbf{D}^{\frac{1}{2}} \mathbf{1}_n$, is then well approximated by an up-to-($2k+1$) rank spiked model (of order $O_{\parallel \cdot \parallel}(1)$). The information appears here as a combination of the statistical information $\mathbf{M}^\top \mathbf{M}$, $\mathbf{t} \mathbf{t}^\top$ and \mathbf{S}° , again modulated by the successive derivatives of f at τ_p .

A complete analysis of the asymptotic spectrum of \mathbf{L} is then tractable, leading to the following remarks [Couillet and Benaych-Georges, 2016]:

- due to the presence of the non-informative vector $\boldsymbol{\psi}$ in \mathbf{U} , one *isolated non-informative* eigenvalue may be found in the spectrum of \mathbf{L}' . This has two main consequences: (i) even in the absence of classes, \mathbf{L}' , and thus \mathbf{L} , may contain an isolated eigenvalue which may be confused as information; (ii) in the presence of classes, not all eigenvectors are informative and there is no deciding which one of the isolated eigenvalues is possibly not informative. Figure 4.8 depicts the typical behavior of the spectrum for a Gaussian mixture with equal identity covariance, with emphasis on the non-informative eigenvalue-eigenvector pair;
- unlike $\mathbf{M} \mathbf{M}^\top$ and \mathbf{S}° which are matrices of rank at most $k-1$, $\mathbf{t} \mathbf{t}^\top$ is a rank-one matrix; as such, if data are mostly discriminable by the trace

of their covariances (i.e., the information in \mathbf{t}), then only one informative eigenvector of \mathbf{L}' (in addition to the eigenvector $\mathbf{D}^{\frac{1}{2}}\mathbf{1}_n$ of \mathbf{L}) can be exploited. This is again counterintuitive since, irrespective of the number of classes, the information is “concentrated” into a single eigenvector. The rule of thumb according to which the number of relevant eigenvectors matches the number of classes thus fails in this case. Figure 4.9 shows the difference between the two informative eigenvectors of \mathbf{L} (the second and third) under a Gaussian mixture with different means and equal covariance, versus the two informative eigenvectors of \mathbf{L} (the first and second) under a common-mean and different-covariance trace scenario. In particular, the bottom display confirms that the discriminative covariance trace information is carried along a one-dimensional axis;

- similar to \mathbf{K} , selecting f such that $f'(\tau_p) \simeq 0$ simultaneously discards the discriminative information of the statistical means across classes as well as the noise terms. The matrix $\mathbf{S}^\circ/\sqrt{p}$ emerges alone and classification becomes asymptotically trivial if $\mathbf{S}^\circ = O_{\|\cdot\|}(\sqrt{p})$.




Figure 4.8: Eigenvalues of \mathbf{L} and top four eigenvectors for $\mathbf{C}_a = \mathbf{I}_p$, $f(t) = 4(t - \tau_p)^2 - (t - \tau_p) + 4$ with $\tau_p = 2$, $f(0) = 22$, $f(\tau_p) = 4$, $f'(\tau_p) = -1$, $f''(\tau_p) = 8$, $p = 2048$, $n = 512$, three classes with $n_1 = n_2 = 128$, $n_3 = 256$ and $[\boldsymbol{\mu}_a]_i = 5\delta_{ai}$. Emphasis made on the (third) “non-informative” eigenvalue-eigenvector pair in red. [Link to code]

Implementation on real data. The above results are quite telling of the many misconceptions of, as simple and widely spread an algorithm in machine learning as kernel spectral clustering.




Figure 4.9: Two dimensional representation of (top): eigenvectors two versus three of \mathbf{L} , $[\mu_a]_i = 5\delta_{ai}$, $\mathbf{C}_a = \mathbf{I}_p$, and of (bottom): eigenvectors one versus two of \mathbf{L} , $\mu_a = \mathbf{0}$, $\mathbf{C}_a = (1+4(a-1)/\sqrt{p})\mathbf{I}_p$. In both cases, $p = 3072$, $n_1 = n_2 = 192$, $n_3 = 384$, $f(t) = 1.5(t - \tau_p)^2 - (t - \tau_p) + 5$. [\[Link to code\]](#)

These novel insights can be summarized as follows: under a *large dimensional Gaussian mixture model* setting, (i) the Euclidean distance between data vectors tend to be the same while spectral clustering still performs in a non-trivial fashion, (ii) the number of isolated eigenvalues needs not necessarily match the number of classes, (iii) some dominant eigenvectors may not be informative at all, and possibly most-importantly, (iv) the kernel function f needs not be decreasing and only operates through its first derivatives in the vicinity of τ_p .

Yet, the Gaussian mixture assumption is somewhat fundamental to our anal-

ysis as it brings forth the necessary degrees of independence that induce the data concentration. One may wonder whether the results still hold when applied to real-world datasets instead of Gaussian vectors.

Figure 4.11 depicts the four dominant eigenvectors of \mathbf{L} for $\mathbf{K}_{ij} = \exp(-\|\mathbf{x}_i - \mathbf{x}_j\|^2/(2p))$ for $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ extracted from three classes (zeros, ones, and twos) of the popular MNIST dataset, each class containing 64 vectorized images of size 28×28 pixels (so that $n = 192$ and $p = 784$). Each data vector \mathbf{x}_i is preprocessed by centering and scaling by the empirical mean and variance computed from the whole MNIST database (and then by \sqrt{p} to adhere to the normalization of the theorem statement). An image example from each class are depicted in Figure 4.10.

Figure 4.11 precisely shows in red lines the genuine four dominant eigenvectors of \mathbf{L} and in black the eigenvectors of $\tilde{\mathbf{L}}$ from Theorem 4.7. To obtain $\tilde{\mathbf{L}}$, the statistical means and covariances are empirical computed from statistical averaging over the whole set of images of zeros, ones and twos of the MNIST database; as for \mathbf{W} (and thus ψ), it is computed by discarding from \mathbf{X} the evaluated average. Finally, in blue are shown the theoretical class-wise eigenvector means and ± 1 standard deviations obtained from a spiked-model analysis of $\tilde{\mathbf{L}}$ (see for details in [Coulillet and Benaych-Georges, 2016]).

It is surprising to see that, despite the obvious non-Gaussianity of the MNIST dataset, the theoretical predictions are in almost perfect accordance with empirical observations.



Figure 4.10: Samples from the MNIST database.

4.5.1.2 The case of “ α - β ” inner-product kernels

The previous section demonstrated that, despite the phenomenon of distance concentration, spectral clustering on the normalized Laplacian \mathbf{L} remains valid under large dimensional data assumption, at the expense of a few unexpected outcomes (presence of non-informative isolated eigenvectors, incoherence between number of classes and number of informative eigenvectors, etc.). These are immediate consequences of the theoretical study performed in Section 4.2 and were shown to adequately match the actual performance of spectral clustering on, not only Gaussian, but also real-world data.

However, it was also showed in Section 4.2 that kernels of the type $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$, despite their wide popularity, are suboptimal when it comes to classifying data down to their minimal statistical discrimination. We then proven in Section 4.3 that the α - β kernels, that satisfy $f(\tau_p) = O(1)$, $f'(\tau_p) = \alpha p^{-1/2}$ and $f''(\tau_p) = 2\beta$ for some $\alpha, \beta = O(1)$, are more powerful in discriminating data having close (even equal) means and slightly differing covariances.




Figure 4.11: Leading four eigenvectors of \mathbf{L} (red) versus $\tilde{\mathbf{L}}$ (black) and theoretical class-wise means and standard deviations (blue) for MNIST data. [Link to code]

We consider this setting here.

Specifically, Figure 4.12 displays the comparative performance of Gaussian versus $\alpha\beta$ inner-product kernels in the setting of a two-class Gaussian mixture with equal means but slightly differing covariances (thus here with $\alpha = 0$). We observe that the Gaussian kernel is incapable of resolving the two classes while the $\alpha\beta$ kernel is fully adapted. Figure 4.13 then extends the same analysis for an EEG dataset (epileptic versus sane patients)⁷ specifically chosen since, being a more or less stationary zero mean time series, the critical class discriminating features lie more in the second order statistics (i.e., covariance matrices) rather than in the first (means). The dataset vectors were appropriately centered and normalized to be fully compliant with the kernel models. In this case, the Gaussian kernel is observed to have less discriminating power compared to the $\alpha\beta$ kernel (chosen here again with $\alpha = 0$, i.e., $f'(\tau_p) = 0$).

We next move to a more general analysis of the $\alpha\beta$ kernel for all values of α/β , rather than just for $\alpha = 0$. For a simplified comparison, for varying val-

⁷<http://www.meb.uni-bonn.de/epileptologie/science/physik/eegdata.html>.




Figure 4.12: Comparison of 2D representation of eigenvectors one and two of (**top**) the Gaussian kernel $\mathbf{K}_{ij} = \exp(-\|\mathbf{x}_i - \mathbf{x}_j\|^2/(2p))$ and (**bottom**) the $\alpha\beta$ kernel $\mathbf{K}_{ij} = (\|\mathbf{x}_i - \mathbf{x}_j\|^2/p - 2)^2$ with $\alpha = 0$. Here $k = 2$ classes of even size, $p = 400$, $n = 1000$, $\mu_a = \mathbf{0}$, $\mathbf{C}_a = 0.1\mathbf{I}_p + 2\mathbf{Z}_a\mathbf{Z}_a^T/p$ where $\mathbf{Z}_a \in \mathbb{R}^{p \times p/2}$ have independent standard Gaussian entries. [Link to code]

ues of α/β , k-means clustering is performed on the two dominant eigenvectors of the kernels under study. Figure 4.3 and Figure 4.15 provide a comparison of the spectral clustering performance for the “ $\alpha\beta$ ” kernel, versus the standard Gaussian kernel on two real datasets: the MNIST image database and the previous EEG database. Figure 4.15 clearly evidences this fact with a strong improvement of the “ $\alpha\beta$ ” kernel performance for α/β close to zero (thus when voluntarily disregarding the information about the means and focusing on the




Figure 4.13: Comparison of 2D representation of eigenvectors one and two of (**top**) the Gaussian kernel $\mathbf{K}_{ij} = \exp(-\|\mathbf{x}_i - \mathbf{x}_j\|^2/(2p))$ and (**bottom**) the α - β kernel $\mathbf{K}_{ij} = (\|\mathbf{x}_i - \mathbf{x}_j\|^2/p - 2)^2$ with $\alpha = 0$. Here $k = 2$ classes of even size from the EEG dataset (class ? versus class ?), $p = 100$ and $n = 1\,000$. [Link to code]

covariances instead); the Gaussian kernel in this case has a mediocre performance. But more strikingly, the classification performance on MNIST data in Figure 4.3, where statistical means are (supposedly) more informative than covariances, in general favors a large value for the ratio α/β , but not always: in the case of the pair of digits (3, 8) (which are understandably harder to discriminate than digits (3, 6)), the second order statistics are more valued by the kernel, with a boost of up to 10% of classification rate over the Gaussian kernel.

Setting up the proper value for α, β beforehand is however not immediate as the performance depends on the statistics of each class. Being unknown under a fully unsupervised setting, only iterative procedures (whereby a first iteration provides a crude classification and thus the possibility to estimate the sufficient statistics) can seemingly be exploited to selectively adapt the algorithm and improve its performance. Alternatively, an informed guess of the relative importance of means versus covariances may be used to adapt the algorithm.




Figure 4.14: Spectral clustering of the MNIST database with the “ α - β ” kernel $\mathbf{K}_{ij} = (\frac{1}{p}\|\mathbf{x}_i - \mathbf{x}_j\|^2 - 2)^2$ for varying $\frac{\alpha}{\beta}$ versus Gaussian kernel ($\mathbf{K}_{ij} = \exp(-\frac{1}{2p}\|\mathbf{x}_i - \mathbf{x}_j\|^2)$). k-means clustering on dominant two eigenvectors. [Link to code]

figures to be done! and y-axis always be misclassification rate.

4.5.2 Application to semi-supervised kernel learning

Semi-supervised learning is possibly the most natural, but paradoxically the least studied, framework in machine learning in that it assumes the existence of a large set of data, say to be classified, with only some of the data already manually labeled. This both encompasses unsupervised learning in the extreme case of no labeled data and supervised learning at the other extreme.

We will see in this section that the reason behind its not being profoundly studied may precisely lie in a misunderstanding of the (not so) large dimensional behavior of the devised methods. These lead to often quite erroneous outcomes, which have been worked around in the literature by various intuitions, rarely sustained by theory.

The random matrix approach clarifies the main problems, demonstrates that some of the popular methods are indeed inconsistent and *must fail*, and most importantly, allows for a novel improved (again in a very counterintuitive fashion) scheme that is provably consistent.




Figure 4.15: Spectral clustering of the EEG database with the “ α - β ” kernel \mathbf{K} for varying $\frac{\alpha}{\beta}$ versus Gaussian kernel ($\mathbf{K}_{ij} = \exp(-\frac{1}{2p}\|\mathbf{x}_i - \mathbf{x}_j\|^2)$). [Link to code]

4.5.2.1 Semi-supervised graph Laplacian and random walk approaches

The likely most common approach to semi-supervised learning are graph-based methods, which have a dual interpretation. In these methods, one considers data $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$ as vertices of a weighted graph with edge weights $\mathbf{K}_{ij} \equiv \kappa(\mathbf{x}_i, \mathbf{x}_j)$ encoding the similarity between \mathbf{x}_i and \mathbf{x}_j . As usual, we consider $\mathbf{K} = \{\mathbf{K}_{ij}\}_{i,j=1}^n$ to be of kernel form, e.g.,

$$\mathbf{K}_{ij} = f\left(\frac{1}{p}\|\mathbf{x}_i - \mathbf{x}_j\|^2\right).$$

From a finite-dimensional intuition, vectors \mathbf{x}_i that are alike should aggregate in clusters, with some nodes in these clusters already labeled. As such, a natural idea to recover the classes of the unknown nodes consists in “spreading” the labels throughout the graph by means of random walks on the graph: either starting from an unlabeled node and having it walk randomly according to edge weights, until it reaches a labeled node, or conversely, walking from labeled nodes to unlabeled nodes; this procedure is iterated randomly on the graph until convergence. In this spirit, assuming k data classes $\mathcal{C}_1, \dots, \mathcal{C}_k$, the random walk method [Szummer and Jaakkola, 2002] or the label propagation approach [Zhu and Ghahramani, 2002] allocates “soft scores” $(\mathbf{S}_{i1}, \dots, \mathbf{S}_{ik}) \in \mathbb{R}^k$ for an unlabeled node \mathbf{x}_i to belong to each class. These k scores are then compared and the argument of the highest score is assigned to \mathbf{x}_i . Of utmost interest for us in the sequel is the particularly popular PageRank approach [Avrachenkov et al., 2012], because of its robustness and high performance.

The alternative viewpoint is more related to the optimization schemes (such as the graph-cut minimization (4.20)) in unsupervised spectral clustering. There

again, a matrix scores $\mathbf{S} = \{\mathbf{S}_{ia}\}_{1 \leq i \leq n, 1 \leq a \leq k}$ for the n data vectors in each of the k classes needs to be filled, by solving the optimization problem of the type

$$\begin{aligned} \mathbf{S} &= \arg \min_{\mathbf{S} \in \mathbb{R}^{n \times k}} \sum_{a=1}^k \sum_{i,j=1}^n [\mathbf{K}]_{ij} ([\mathbf{S}]_{ia} d_i^\alpha - [\mathbf{S}]_{ja} d_j^\alpha)^2 \\ \text{s.t. } \mathbf{S}_{ia} &= \delta_{\mathbf{x}_i \in \mathcal{C}_a} \text{ for labeled nodes.} \end{aligned} \quad (4.23)$$

Here $d_i = \sum_{j=1}^n \mathbf{K}_{ij}$ is the degree of the node i in the graph representation and $\alpha \in \mathbb{R}$ is some parameter to be specified. Note that the optimization scheme imposes $\mathbf{K}_{ij} \geq 0$ to be meaningful (otherwise, an arbitrarily small negative solution could be found). With this constraint, the optimization scheme naturally induces the scores \mathbf{S}_{ia} and \mathbf{S}_{ja} to be close for \mathbf{K}_{ij} large, and allows them to be distinct if \mathbf{K}_{ij} is close to zero.

Yet, it was empirically observed that, for $\alpha = 0$, the algorithm tends to fail. From a large dimensional perspective, we now understand the fundamental reason behind this observation is the erroneous assumption that \mathbf{K}_{ij} is either “small” or “large”, while past attempts rather blamed this on the unbalances in the graph. To avoid some (too strongly connected) nodes to create biases, the natural first solution has been to weigh the scores \mathbf{S}_{ij} by a negative power of the degree d^α for some $\alpha < 0$. This is the approach essentially followed, for different choices of α , in [Zhu et al., 2003, Belkin et al., 2004, Joachims, 2003, Zhou et al., 2004].

Quite interestingly, the explicit solutions of these (quadratic under linear constraint) optimization problems can essentially be mapped to the stationary points of the aforementioned label propagation or random walk on graphs, as shown in [Avrachenkov et al., 2012] for $\alpha = 0, -1/2$ and -1 . The case $\alpha = -1$, which we shall discuss in depth in the following, precisely corresponds to the PageRank algorithm.

Remark 4.7 (Laplacian versus manifold methods). *Aside from graph-Laplacian approaches, another popular semi-supervised learning scheme is the manifold-based method [Belkin and Niyogi, 2004, Goldberg et al., 2009, Moscovitch et al., 2016]. These algorithms however rely on a first step of “manifold learning” which imposes the unsupervised projection on a dominant subspace that we know, from the previous section, to possibly lead (below a certain phase transition) to a complete loss of information. The Laplacian-based approaches, as shall be seen in this section, do not suffer from this phase transition issue and are thus more robust.*

The solution to (4.23) is explicitly given by

$$\mathbf{S}_{[u]} = \left(\mathbf{I}_{n_u} - \mathbf{D}_{[u]}^{-1-\alpha} \mathbf{K}_{[uu]} \mathbf{D}_{[u]}^\alpha \right)^{-1} \mathbf{D}_{[u]}^{-1-\alpha} \mathbf{K}_{[ul]} \mathbf{D}_{[l]}^\alpha \mathbf{S}_{[l]}. \quad (4.24)$$

where $\mathbf{D} = \text{diag}(d_1, \dots, d_n) \in \mathbb{R}^{n \times n}$ and we divide \mathbf{S} , \mathbf{K} and \mathbf{D} in sub-blocks of labeled (l) versus unlabeled (u) data indices

$$\mathbf{S} = \begin{bmatrix} \mathbf{S}_{[l]} \\ \mathbf{S}_{[u]} \end{bmatrix}, \quad \mathbf{K} = \begin{bmatrix} \mathbf{K}_{[ll]} & \mathbf{K}_{[lu]} \\ \mathbf{K}_{[ul]} & \mathbf{K}_{[uu]} \end{bmatrix}, \quad \text{and } \mathbf{D} = \begin{bmatrix} \mathbf{D}_{[l]} & 0 \\ 0 & \mathbf{D}_{[u]} \end{bmatrix}. \quad (4.25)$$

The final decision, i.e., the allocated class index $\hat{\mathcal{C}}_{\mathbf{x}_i}$ for data \mathbf{x}_i , is then given by

$$\hat{\mathcal{C}}_{\mathbf{x}_i} = \mathcal{C}_{\hat{a}} \text{ for } \hat{a} = \arg \max_{1 \leq a \leq k} \mathbf{S}_{ia}.$$

4.5.2.2 Large dimensional performance analysis

As in the previous section for unsupervised classification, we place ourselves under a “non-trivial” Gaussian mixture model, that is

$$\mathbf{x}_i \in \mathcal{C}_a \Leftrightarrow \mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{C}_a).$$

for $\boldsymbol{\mu}_a \in \mathbb{R}^p$ and $\mathbf{C}_a \in \mathbb{R}^{p \times p}$. We assume that there exists k classes $\mathcal{C}_1, \dots, \mathcal{C}_k$ of sizes n_1, \dots, n_k , with in total $n_{[l]}$ labeled and $n_{[u]}$ unlabeled nodes. We denote $n_{[l]a}, n_{[u]a}$ the number of labeled and unlabeled nodes of class \mathcal{C}_a that are all assumed to be of order $O(n)$.

Since our focus lies in the understanding of the behavior of the semi-supervised learning algorithm, so in particular (but not only) in the statistics of the score matrix \mathbf{S} and the impact of the hyperparameter α , here we merely consider the case $\mathbf{K}_{ij} = f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$ for $f'(\tau_p)$ away from zero, where we recall

$$\tau_p = \frac{2}{p} \operatorname{tr} \mathbf{C}^\circ, \quad \mathbf{C}^\circ = \sum_{a=1}^k \frac{n_a}{n} \mathbf{C}_a$$

and demand for every $a, b \in \{1, \dots, k\}$ that

$$\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\| = O(1), \quad \operatorname{tr}(\mathbf{C}_a - \mathbf{C}_b) = O(\sqrt{p}), \quad \operatorname{tr}(\mathbf{C}_a - \mathbf{C}_b)^2 = O(p).$$

The fact that these distances are possibly suboptimal for certain classification tasks will not be our primary focus here.

First intuitions. A first key observation is that, under the non-trivial growth rate, since $\max_{ij} |\mathbf{K}_{ij} - \tau_p| \xrightarrow{a.s.} 0$, there are strong reasons to believe that the optimization scheme (4.23), through its solution in (4.24), is bound to fail. However, while the algorithm always behaves differently from our low dimensional intuition, it may not fail in some cases. To see this, let us set $\alpha = -1$ and perform semi-supervised graph learning with a Gaussian kernel on two even-sized classes $\mathcal{N}(\pm \boldsymbol{\mu}, \mathbf{I}_p)$. Figure 4.16 illustrates the scores \mathbf{S} for $p = 1$ (small dimensional case) versus $p = 20$ (moderately large dimensional case).

Recalling that the finite-dimensional intuition behind the optimization framework in (4.23) (or its equivalent walk on graph and label propagation interpretation) is that, the unlabeled data scores should be “drawn” to the scores of neighbors effectively from the same class. This expected behavior of the score vector $\mathbf{S}_{\cdot a} \in \mathbb{R}^n$ of class \mathcal{C}_a is displayed at the top of Figure 4.16 for data of dimension $p = 1$. Yet, as soon as p is slightly larger, this behavior is largely disrupted, as observed in the bottom display for $p = 20$. Note in particular that:

- the unlabeled data scores do not seem affected by the labeled data scores (0 or 1); indeed, a pairwise comparison of \mathbf{S}_{i1} and \mathbf{S}_{i2} reveals that the scores are extremely close to one another but their average is not a fixed value (one would expect 0.5);
- despite this completely different behavior between $p = 1$ and $p = 20$ scenario, the algorithm seems to work properly in that $\mathbf{S}_{i1} > \mathbf{S}_{i2}$ for most $i \leq n/2$ (so for data genuinely in \mathcal{C}_1) and conversely.

As a consequence, although the finite-dimensional intuition is largely disrupted here, the semi-supervised learning scheme does not completely fail, at least in this very elementary Gaussian mixture toy model.




Figure 4.16: Scores \mathbf{S} for Laplacian-based semi-supervised learning with two classes $\mathbf{x}_i \sim \mathcal{N}(\pm\boldsymbol{\mu}, \mathbf{I}_p)$ with $\boldsymbol{\mu} = [2, \mathbf{1}_{p-1}]$, Gaussian kernel, $\alpha = -1$, $n = 200$, $n_{[l]} = 60$, $n_{[u]} = 140$ and (top) $p = 1$, (bottom) $p = 20$. [\[Link to code\]](#)




Figure 4.17: Scores \mathbf{S} of MNIST data (ones versus twos) with $p = 784$, in the same setting as Figure 4.16. [\[Link to code\]](#)

What about real data scenarios? Figure 4.17 proposes the same setting as Figure 4.16 for the MNIST dataset. The situation appears much closer to the bottom than the top display of Figure 4.16, thereby suggesting a closer fit with the large dimensional viewpoint. The situation is even worse since the unlabeled data scores are less close to 1/2. Yet, the algorithm again works decently as the comparison between \mathbf{S}_{i1} and \mathbf{S}_{i2} shows a clear advantage of class \mathcal{C}_1 on the first half of the unlabeled data and class \mathcal{C}_2 on the second half.

But all this is valid here for $\alpha = -1$. The same simulation with for example $\alpha = 0$ or $\alpha = -1/2$ reveals a total failure of the algorithm with all data mapped to the same class.

Derivations and main results. To understand the behavior of Laplacian-based semi-supervised learning observed above, we shall first focus on the large n, p asymptotics of the score vectors $\mathbf{S}_{\cdot a} \in \mathbb{R}^n$ for $1 \leq a \leq k$. Characterizing the ultimate performance of the algorithm will consist instead in studying, for each \mathbf{x}_i unlabeled, the “joint” vector of scores $\mathbf{S}_i \in \mathbb{R}^k$.

Recall from Theorem 4.1 that, the kernel matrix \mathbf{K} under study here admits the random matrix equivalent $\tilde{\mathbf{K}}$ in the sense that $\|\mathbf{K} - \tilde{\mathbf{K}}\| \xrightarrow{a.s.} 0$ in the large n, p limit. In particular, $\|\mathbf{K}_{[ul]} - \tilde{\mathbf{K}}_{[ul]}\| \xrightarrow{a.s.} 0$ and similarly for all sub-blocks of \mathbf{K} under the decomposition in (4.25). From the explicit form (4.24) of the unlabeled data scores $\mathbf{S}_{[u]}$, it is thus tempting to replace all sub-blocks of \mathbf{K} by those of $\tilde{\mathbf{K}}$. This is indeed justified since, almost surely, for all large n , the resolvent $(\mathbf{I}_{n_u} - \mathbf{D}_{[u]}^{-1-\alpha} \mathbf{K}_{[uu]} \mathbf{D}_{[u]}^\alpha)^{-1}$ has bounded spectrum. After calculus, it indeed comes that

$$\mathbf{D}_{[u]}^{-1-\alpha} \mathbf{K}_{[uu]} \mathbf{D}_{[u]}^\alpha = \frac{1}{n} \mathbf{1}_{n_{[u]}} \mathbf{1}_{n_{[u]}}^\top + O_{\|\cdot\|}(n^{-\frac{1}{2}})$$

so that the resolvent can be further expanded as

$$\left(\mathbf{I}_{n_{[u]}} - \mathbf{D}_{[u]}^{-1-\alpha} \mathbf{K}_{[uu]} \mathbf{D}_{[u]}^\alpha \right)^{-1} = \mathbf{I}_{n_{[u]}} + \frac{1}{n_{[l]}} \mathbf{1}_{n_{[u]}} \mathbf{1}_{n_{[u]}}^\top + O_{\|\cdot\|}(n^{-\frac{1}{2}}). \quad (4.26)$$

We will see later, although the $O_{\|\cdot\|}(n^{-\frac{1}{2}})$ term contains statistical information about the classes, this seemingly “trivial” linearization of the resolvent is at the source of various counterintuitive phenomena observed in large dimensional semi-supervised learning.

Once the linearization of the resolvent performed, characterizing the asymptotic behavior of $\mathbf{S}_{[u]}$ becomes a matter of algebraic calculus from the results in Theorem 4.1. This calculus leads to the following first central result (see proof details in [Mai and Couillet, 2018])

$$(\mathbf{S}_{[u]})_{\cdot a} = \frac{n_{[l]a}}{n} \left(\mathbf{1}_{n_{[u]}} + \mathbf{v} + (\alpha + 1) \frac{f'(\tau_p)}{f(\tau_p)\sqrt{c}} \frac{t_a \mathbf{1}_{n_{[u]}}}{\sqrt{n}} \right) + O(n^{-1}) \quad (4.27)$$

where we recall $t_a = \text{tr } \mathbf{C}_a^\circ / \sqrt{p} = O(1)$, $O(n^{-1})$ is understood entry-wise and $\mathbf{v} \in \mathbb{R}^{n_{[u]}}$ is a zero-mean *random vector* with entries of order $O(n^{-1/2})$ which is independent on the class index a .

This first result states that:

1. the score $(\mathbf{S}_{[u]})_{ia}$ of an unlabeled datum \mathbf{x}_i is largely dominated by the constant $n_{[l]a}/n$; as such, if $n_{[l]1}, \dots, n_{[l]k}$ are distinct, all the unlabeled data will be allocated to the class \mathcal{C}_a corresponding to the largest (or smallest) value of $n_{[l]a}$;
2. the next dominant term in $(\mathbf{S}_{[u]})_{ia}$ is the sum of two terms of order $O(n^{-1/2})$: the zero-mean random noise $[\mathbf{v}]_i$ and a term proportional to $(\alpha+1)t_a$ which does not depend on i : as such, provided the $n_{[l].}$ are equal, all unlabeled data \mathbf{x}_i will likely be classified in the class \mathcal{C}_a for which t_a is maximal (or minimal), unless the t_a 's are all equal (or distinct by at most $O(n^{-1/2})$) or $\alpha = -1$ (or at least equal to $-1 + O(n^{-1/2})$); since the former cannot be guaranteed in practice, one must set the parameter α close to -1 ; this here explains the long observed advantage of the PageRank algorithm over Laplacian methods;
3. once all these constraints are set, what remains in $(\mathbf{S}_{[u]})_{ia}$ is a term of order $O(n^{-1})$, leading to the vector $(\mathbf{S}_{[u]})._a$ being of norm $O(n^{-1/2})$: this “weak” $O(n^{-1})$ term, as we shall see, is the one containing the *relevant classification information* which *must* be protected from the dominant higher $O(n^{-1/2})$ order noise!

From Item 1, we thus conclude that $\mathbf{S}_{[u]}$ is *not* the appropriate metric, and one must instead consider the normalized score matrix

$$\hat{\mathbf{S}}_{[u]} = \mathbf{S}_{[u]} \operatorname{diag} \left(\frac{n}{n_{[l]1}}, \dots, \frac{n}{n_{[l]k}} \right). \quad (4.28)$$

From Item 3, α must be set to $-1 + \beta/\sqrt{p}$ for some $\beta = O(1)$. This preprocessing step discards all *dominant noise* in $\mathbf{S}_{[u]}$ and therefore now reveals the “hidden” (otherwise buried in noise) class information structure from the residual $O(n^{-1})$ term in the expansion of $(\mathbf{S}_{[u]})._a$ in (4.27). A careful random matrix analysis leads to the following result.

Theorem 4.8 ([Mai and Couillet, 2018, Theorem 2]). *For $\mathbf{x}_i \in \mathcal{C}_b$ an unlabeled data point, let $\hat{\mathbf{S}}$ be defined in (4.28) and $\alpha = -1 + \beta/\sqrt{p}$ for some $\beta \in \mathbb{R}$. Then*

$$p\hat{\mathbf{S}}_{ii} = p(1 + [\mathbf{v}]_i)\mathbf{1}_k + \mathbf{g}_i + o(1), \quad \mathbf{g}_i \sim \mathcal{N}(\mathbf{E}_b, \mathbf{V}_b)$$

where $[\mathbf{v}]_i = O(n^{-1/2})$ only depends on i and

$$\begin{aligned} [\mathbf{E}_b]_a &= -\frac{2f'(\tau_p)}{f(\tau_p)} \tilde{\boldsymbol{\mu}}_a^\top \tilde{\boldsymbol{\mu}}_b + \left(\frac{f''(\tau_p)}{f(\tau_p)} - \frac{f'(\tau_p)^2}{f(\tau_p)^2} \right) \tilde{t}_a \tilde{t}_b \\ &\quad + \frac{2f''(\tau_p)}{f(\tau_p)} \frac{1}{p} \operatorname{tr} \tilde{\mathbf{C}}_a \tilde{\mathbf{C}}_b + \frac{n\beta}{n_{[l]}} \frac{f'(\tau_p)}{f(\tau_p)} t_a \end{aligned} \quad (4.29)$$

$$\begin{aligned} [\mathbf{V}_b]_{a_1 a_2} &= 2 \left(\frac{f''(\tau_p)}{f(\tau_p)} - \frac{f'(\tau_p)^2}{f(\tau_p)^2} \right)^2 t_{a_1} t_{a_2} \cdot \frac{1}{p} \operatorname{tr} \mathbf{C}_b \mathbf{C}_b \\ &\quad + 4 \frac{f'(\tau_p)^2}{f(\tau_p)^2} \left[(\boldsymbol{\mu}_{a_1}^\circ)^\top \mathbf{C}_b \boldsymbol{\mu}_{a_2}^\circ + \frac{n\delta_{a_1 a_2}}{n_{[l]} n_{[l]}} \operatorname{tr} \mathbf{C}_{a_1} \mathbf{C}_b \right]. \end{aligned} \quad (4.30)$$

where we introduce the “labeled-data centered” statistics

$$\tilde{\boldsymbol{\mu}}_a \equiv \boldsymbol{\mu}_a - \sum_{b=1}^k \frac{n_{[l]} b}{n_{[l]}} \boldsymbol{\mu}_b, \quad \tilde{t}_a \equiv \frac{1}{\sqrt{p}} \operatorname{tr} \tilde{\mathbf{C}}_a, \quad \tilde{\mathbf{C}}_a \equiv \mathbf{C}_a - \sum_{b=1}^k \frac{n_{[l]} b}{n_{[l]}} \mathbf{C}_b.$$

The main message of Theorem 4.8 is that, up to the irrelevant dominant term $p(1 + z_i \mathbf{v}_i) \mathbf{1}_k$, the normalized score vector $\hat{\mathbf{S}}_i \in \mathbb{R}^k$ of \mathbf{x}_i has a limiting Gaussian behavior with precisely characterized mean and covariance. Those not surprisingly depend on the statistical means $\boldsymbol{\mu}_a$ and covariance matrices \mathbf{C}_a of the data classes and on the first derivatives of f at τ_p . The parameter β also plays a non-trivial “biasing” role that may be helpful in correcting some inherent unbalance between classes.

Yet, generally speaking, most conclusions drawn in the previous section on spectral clustering remain valid (in particular that taking $f'(\tau_p) = 0$ leads to trivial classification of covariance-based classes), at the noticeable exception of the following surprising remark.

Remark 4.8 (Sub-optimality of the Gaussian kernel). *It is interesting to observe that the term $f''(\tau_p)/f(\tau_p) - f'(\tau_p)^2/f(\tau_p)^2$ plays a dominant role in discriminating classes having various “amplitudes” (i.e., distinct values of t_a). For the Gaussian kernel $f(t) = \exp(-t/2\sigma^2)$, this term is exactly zero for all choices of τ_p and thus the Gaussian kernel, in this semi-supervised context, fails for instance to separate two “nested balls” $\mathcal{N}(\mathbf{0}, (1 \pm \epsilon/\sqrt{p}) \mathbf{I}_p)$, as in the case of Remark 4.5 for “regular” inner-product kernels, while most polynomial kernels succeed. This is illustrated in Figure 4.18.*

Looking now more specifically into the “semi-supervised” aspect of the algorithm, a major problem arises immediately: up to renaming β into $\beta n / n_{[l]}$ which is a free parameter, the mean \mathbf{E}_b in Theorem 4.8 depends neither on $n_{[l]}$ nor on $n_{[u]}$. As for the variance \mathbf{V}_b , note that its diagonal elements decrease as $n_{[l]}$ increases (for fixed $p, n_{[u]}$) but *does not decrease* as $n_{[u]}$ increases (for fixed $p, n_{[l]}$). This suggests that the semi-supervised Laplacian approach *does not* learn from unlabeled data. This surprising outcome is in fact well documented in the semi-supervised learning literature: see in particular [Olivier et al., 2006, Chapter 4] which we quote here




Figure 4.18: Empirical misclassification rate of semi-supervised Laplacian approach on two-class Gaussian mixtures with $\mu_1 = \mu_2$, $\mathbf{C}_1 = \mathbf{I}_p$ and $\mathbf{C}_2 = (1 + 3/\sqrt{p})\mathbf{I}_p$, for (left) Gaussian kernel $f(t) = \exp(-t/(2\sigma^2))$ and (right) polynomial kernel of degree two with $f(\tau_p) = f''(\tau_p) = 1$, $n = 1024$, $p = 512$, $n_{[l]}/n = 1/16$, $n_{[l]1} = n_{[l]2}$ and $\alpha = -1$. [Link to code]

“Our concern is this: it is frequently the case that we would be better off just discarding the unlabeled data and employing a supervised method, rather than taking a semi-supervised route. Thus we worry about the embarrassing situation where the addition of unlabeled data degrades the performance of a classifier.”

The situation is depicted in Figure 4.20 where the performance of the Laplacian approach (blue triangles) is indeed confirmed not to increase with $n_{[u]}$: the popular Laplacian method is thus merely reduced, at least in the large n, p regime, to supervised classification. A particularly problematic consequence is that the fully unsupervised spectral clustering method studied in Section 4.5.1 tends to overtake the Laplacian method.

The next section is dedicated to a more profound analysis of the problem, leading to a random matrix-inspired solution that benefits from more unlabeled data (red circles in Figure 4.20).

4.5.2.3 Improving semi-supervised learning

As pointed out previously, the linearization (4.26) of the resolvent in the expression of $\mathbf{S}_{[u]}$ holds the key to the inefficiency of the use of the unlabeled data in the graph-based semi-supervised learning algorithm.

Main intuition. The situation may be loosely summarized as follows: $\mathbf{S}_{[u]} = \mathbf{A}_{[uu]}\mathbf{A}_{[ul]}$ with $\mathbf{A}_{[uu]}$ the “unsupervised part” of the algorithm and $\mathbf{A}_{[ul]}$ the supervised part. The (operator-norm) dominant-order term of $\mathbf{A}_{[uu]}$ contains the identity matrix $\mathbf{I}_{n_{[u]}}$, while the dominant term of $\mathbf{A}_{[ul]}$ only contains $\frac{1}{n}\mathbf{1}_{n_{[u]}}\mathbf{1}_{n_{[l]}}^\top$; as for the informative terms of least order, call them $\mathbf{B}_{[uu]}$ and $\mathbf{B}_{[ul]}$, they are both such that their (i, a) -entry depends on the class of \mathbf{x}_i and on the class

index a . In order for $(\mathbf{S}_{[u]})_{ia}$ to be informative, it must also depend on both the class of \mathbf{x}_i and on a . However, taking the product $\mathbf{A}_{[uu]} \mathbf{A}_{[ul]}$, the only non-vanishing informative terms are the cross-products between the dominant- and least-order terms, so in particular $\mathbf{I}_{n_{[u]}} \mathbf{B}_{[ul]}$ and $\mathbf{B}_{[uu]} \frac{1}{n} \mathbf{1}_{n_{[u]}} \mathbf{1}_{n_{[l]}}^\top$. While the former has entries (i, a) depending on both the class of \mathbf{x}_i and a , this is not true for the latter which does not depend on a . As a consequence, the (unlabeled) informative $\mathbf{B}_{[uu]}$ vanishes from the output and the unsupervised information is not used.

In order to remedy this situation, one must discard the dominant matrices of the type $\mathbf{1}_{n_{[u]}} \mathbf{1}_{n_{[l]}}^\top$ from the derivation. This term is the first order approximation of $\mathbf{K}_{[ul]} \mathbf{D}_{[l]}^{-1}$, and mainly unfolds from the *non-negativity constraint* of (the entries of) \mathbf{K} , which creates this large “bias”. From a purely mathematical standpoint, it stands to reason to remove this bias, for example by changing \mathbf{K} into $\hat{\mathbf{K}}$ with

$$\hat{\mathbf{K}} = \mathbf{P} \mathbf{K} \mathbf{P}, \quad \mathbf{P} = \mathbf{I}_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^\top \quad (4.31)$$

which is then “orthogonal” to the bias vector $\mathbf{1}_n$. This is not so simple though, as this induces that $\mathbf{D} = \text{diag}(\hat{\mathbf{K}} \mathbf{1}_n) = \mathbf{0}$. Also, replacing \mathbf{K} by $\hat{\mathbf{K}}$ in the original optimization problem (4.23), now that $\hat{\mathbf{K}}$ has negative entries, and leads to an unbounded negative solution.

Adapted optimization framework. The proposed workaround in [Mai and Couillet, 2020] consists in starting from the optimization problem (4.23), replacing \mathbf{K} with $\hat{\mathbf{K}}$ (with thus $\alpha = 0$ since $d_i = 0$ for all i) and imposing an additional constraint on the Frobenius norm $\|\mathbf{S}\|_F$ to avoid the trivial unbounded negative solution.

The optimization framework reads

$$\hat{\mathbf{S}} = \arg \min_{\hat{\mathbf{S}} \in \mathbb{R}^{n \times k}} \sum_{a=1}^k \sum_{i,j=1}^n [\hat{\mathbf{K}}]_{ij} \left([\hat{\mathbf{S}}]_{ia} - [\hat{\mathbf{S}}]_{ja} \right)^2 \quad (4.32)$$

$$\text{s.t. } \begin{cases} \hat{\mathbf{S}}_{[l]} = \left(\mathbf{I}_{n_{[l]}} - \frac{1}{n_{[l]}} \mathbf{1}_{n_{[l]}} \mathbf{1}_{n_{[l]}}^\top \right) \mathbf{S}_{[l]} \\ [\mathbf{S}]_{ia} = \delta_{\mathbf{x}_i \in \mathcal{C}_a} \text{ for labeled nodes,} \\ \|\hat{\mathbf{S}}_{[u]}\|_F^2 = n_{[u]} \gamma, \text{ for some } e > 0 \end{cases} \quad (4.33)$$

the solution of which is explicitly given by

$$\hat{\mathbf{S}}_{[u]} = \left(\alpha \mathbf{I}_{n_{[u]}} - \hat{\mathbf{K}}_{[uu]} \right)^{-1} \hat{\mathbf{K}}_{[ul]} \hat{\mathbf{S}}_{[l]} \quad (4.34)$$

where α is the Lagrangian multiplier associated with the constraint $\|\hat{\mathbf{S}}_{[u]}\|_F^2 = n_{[u]} \gamma$ and satisfies $\alpha > \|\hat{\mathbf{K}}_{[uu]}\|$.

Performance analysis. The performance of the random matrix-improved semi-supervised learning approach in (4.34) is studied in [Mai and Couillet, 2020], under the simplified setting of $k = 2$ classes with $\mathbf{C}_1 = \mathbf{C}_2 \equiv \mathbf{C}$ and with the “one-hot” $\hat{\mathbf{S}} \in \mathbb{R}^{n \times k}$ replaced by the “sign” vector $\hat{\mathbf{s}} \in \mathbb{R}^n$ such that

$$\hat{\mathbf{s}}_{[l]} = \left(\mathbf{I}_{n_{[l]}} - \frac{1}{n_{[l]}} \mathbf{1}_{n_{[l]}} \mathbf{1}_{n_{[l]}}^\top \right) \mathbf{s}_{[l]}, \quad [\hat{\mathbf{s}}]_i = (-1)^a \text{ for labeled node } \mathbf{x}_i \in \mathcal{C}_a. \quad (4.35)$$

The derivation of the asymptotic performance is more technically challenging as the resolvent $(\alpha \mathbf{I}_{n_{[u]}} - \hat{\mathbf{K}}_{[uu]})^{-1}$ no longer trivially expands around a leading (non-informative) matrix. For $\hat{\mathbf{K}} = \mathbf{P}\hat{\mathbf{K}}\mathbf{P}$, in the notations of Corollary 4.1, it is easily seen that, the random equivalent $\mathbf{P}\tilde{\mathbf{K}}\mathbf{P}$ is of order $O_{\parallel \cdot \parallel}(1)$ which is the order of the informative terms. Therefore, $\mathbf{P}\tilde{\mathbf{K}}\mathbf{P}$ may be seen as a low rank perturbation of the (full rank) random matrix $-2f'(\tau_p)\frac{1}{p}\mathbf{P}\mathbf{W}^\top\mathbf{W}\mathbf{P}$, with known deterministic equivalents for its resolvent (e.g., Theorem 2.8).

This leads to the following performance asymptotics.

Theorem 4.9 (From [Mai and Couillet, 2020]). *Let $\hat{\mathbf{s}}_{[u]} = (\alpha \mathbf{I}_{n_{[u]}} - \hat{\mathbf{K}}_{[uu]})^{-1} \hat{\mathbf{K}}_{[ul]} \hat{\mathbf{s}}_{[l]}$ with $\hat{\mathbf{s}}_{[l]}$ defined in (4.35) such that $\|\hat{\mathbf{s}}_{[u]}\|^2 = n_{[u]}\gamma$. Then, for an unlabeled data point $\mathbf{x}_i \in \mathcal{C}_b$ with (a priori) probability $\mathbb{P}(\mathbf{x}_i \in \mathcal{C}_b) = c_b$, $b \in \{1, 2\}$,*

$$[\hat{\mathbf{s}}]_i = g_i + o(1), \quad g_i \sim \mathcal{N}((-1)^b(1 - \rho_b)E, V)$$

with $E \equiv E(\xi)$ and $V \equiv V(\xi)$ for ξ the unique solution to $c_1c_2E(\xi)^2 + V(\xi) = e^2 = \gamma$ with positive functions $E(t)$ and $V(t)$ defined as

$$E(t) = \frac{2n_{[l]}\theta(t)}{n_{[u]}(1 - \theta(t))}, \quad V(t) = c_1c_2 \frac{(2n_{[l]} + E(t)n_{[u]})^2\zeta(t) + (4n_{[l]} + E(t)^2n_{[u]})p\eta(t)}{n_{[u]}(n_{[u]} - p\eta(t))}$$

where $\Delta\boldsymbol{\mu} \equiv \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2$ and

$$\begin{aligned} \theta(t) &= c_1c_2t\Delta\boldsymbol{\mu}^\top(\mathbf{I}_p - t\mathbf{C})^{-1}\Delta\boldsymbol{\mu}, \\ \eta(t) &= t^2 \operatorname{tr}(\mathbf{C}^2(\mathbf{I}_p - t\mathbf{C})^{-2})/p, \\ \zeta(t) &= c_1c_2t^2\Delta\boldsymbol{\mu}^\top(\mathbf{I}_p - t\mathbf{C})^{-1}\mathbf{C}(\mathbf{I}_p - t\mathbf{C})^{-1}\Delta\boldsymbol{\mu}. \end{aligned}$$

The formulations of Theorem 4.9, not being explicit, are not immediate to interpret. In the proof of Theorem 4.9, it is shown that θ is of the order of $\|\mathbf{s}_{[u]}\|/\|\mathbf{s}_{[l]}\|$. As such, θ increases with the constraint $e > 0$, itself inversely proportional to the Lagrangian multiplier α . Consequently, raising $\alpha \rightarrow \infty$ brings $\theta \rightarrow 0$ and m/σ no longer depends on $n_{[u]}$: semi-supervised learning is merely a supervised learning. On the opposite, as $\alpha \downarrow \|\hat{\mathbf{K}}_{[uu]}\|$, $\theta \rightarrow \infty$ and now m/σ only depends on $n_{[u]}$: only unlabeled data are used, making the algorithm fully unsupervised. In fact, [Mai and Couillet, 2020] precisely shows that the limit $\alpha \downarrow \|\hat{\mathbf{K}}_{[uu]}\|$ perfectly recovers spectral clustering; this is not difficult to intuit: the resolvent $(\alpha \mathbf{I}_{n_{[u]}} - \hat{\mathbf{K}}_{[uu]})^{-1}$ is strongly dominated by the inverse of the projector $\mathbf{v}\mathbf{v}^\top$ with \mathbf{v} the eigenvector associated with the largest eigenvalue

of $\hat{\mathbf{K}}_{[uu]}$, and thus $\hat{\mathbf{s}}_{[u]} \propto \mathbf{v}(\mathbf{v}^T \hat{\mathbf{K}}_{[ul]} \hat{\mathbf{s}}_{[l]}) \propto \mathbf{v}$, i.e., this is a mere spectral clustering algorithm.

Theorem 4.9, despite taking a rather implicit form, suggests that:

- The classification accuracy is propositional to the ratio E^2/V with

$$\frac{E^2}{V} = \frac{1}{c_1 c_2} \frac{n_{[u]}(n_{[u]} - p\eta(t))E^2}{(2n_{[l]} + E(t)n_{[u]})^2 \zeta(t) + (4n_{[l]} + E(t)^2 n_{[u]})p\eta(t)} \quad (4.36)$$

which, for a well-chosen constraint $\gamma > 0$, is a monotonically increasing function of both $n_{[l]}$ and $n_{[u]}$. As such, the random matrix-improved semi-supervised learning approach is guaranteed to benefit from additional labeled and more importantly, unlabeled data, at least in the simple $\mathcal{N}(\boldsymbol{\mu}_1, \mathbf{C})$ versus $\mathcal{N}(\boldsymbol{\mu}_2, \mathbf{C})$ scenario.

- It can be shown that the quantity $\theta \equiv \theta(\xi) \in (0, 1)$ reflects how much the semi-supervised learning procedure relies on unlabeled data. Precisely, taking $\gamma \rightarrow 0$ leads to $\theta \rightarrow 0$ as well as $\alpha \rightarrow \infty$ and one retrieves (the performance of) the classical normalized Laplacian approach discussed in Theorem 4.8 that is totally *supervised* in high dimensions; on the other hand, taking $\gamma \rightarrow \infty$ results in $\theta \rightarrow 1$ with $\alpha \downarrow \|\hat{\mathbf{K}}_{[uu]}\|$, and one obtains the same the performance as the *unsupervised* spectral clustering approach.⁸ As a result, tuning the constraint parameter γ allows one to find the best trade-off between the supervised and unsupervised learning schemes. This is confirmed in Figure 4.19 for fixed and varying values of $n_{[u]}$ and $n_{[l]}$.

Figure 4.20, as already discussed in the previous section, shows that the random matrix-improved semi-supervised learning method significantly improves over the standard Laplacian method, overtaking both the semi-supervised Laplacian and the non-supervised spectral clustering.

Application to real-world datasets. One may wonder why the after-all quite simple solution proposed in [Mai and Couillet, 2020] has not appeared earlier in the literature. A first reason was mentioned previously: the fact that $\hat{\mathbf{K}}$ has negative entries, when placed in the optimization framework of (4.32) is counterintuitive.

A second reason might follow from the actual output of simulations on real data. The bottom left display of Figure 4.21 compares the performance of the Laplacian versus RMT-improved Laplacian method for a increasing number of unlabeled data $n_{[u]}$: while the RMT-improved method *consistently* outperforms the standard Laplacian, the anticipated incapacity of the latter to use unlabeled data is not observed in practice. This is explained, in the top left display, by the slight but already too large average distance between intra- and inter-class data. Adding Gaussian white noise to the data (middle and right displays), the gap

⁸Since the resolvent $(\alpha \mathbf{I}_{n_{[u]}} - \hat{\mathbf{K}}_{[uu]})^{-1} \simeq (\alpha - \lambda_{\max}(\hat{\mathbf{K}}_{[uu]}))^{-1} \mathbf{v} \mathbf{v}^T$ as $\alpha \downarrow \|\hat{\mathbf{K}}_{[uu]}\|$ with \mathbf{v} the eigenvector associated with the largest eigenvalue of $\hat{\mathbf{K}}_{[uu]}$, we have $\hat{\mathbf{s}}_{[u]} \propto \mathbf{v}(\mathbf{v}^T \hat{\mathbf{K}}_{[ul]} \hat{\mathbf{s}}_{[l]}) \propto \mathbf{v}$ and this is a mere spectral clustering algorithm.




Figure 4.19: Asymptotic misclassification rate as a function of $\theta \equiv \theta(\xi)$ with $c_1 = c_2 = 1/2$, $p = 100$, $\Delta\mu = [2; \mathbf{0}_{p-1}]$, $[\mathbf{C}]_{i,j} = .1^{|i-j|}$. (**Left**): different $n_{[u]}$ with $n_{[l]} = 100$. (**Right**): different $n_{[l]}$ with $n_{[u]} = 800$. Minimal errors are marked in circles. [\[Link to code\]](#)




Figure 4.20: Empirical misclassification rate as a function of $n_{[u]}$ with $n_{[l]} = 200$, $p = 100$, $c_1 = c_2 = 1/2$, $\mathbf{x}_i \sim \mathcal{N}(\pm\mu, \mathbf{I}_p)$ for $\mu = [1; \mathbf{0}_{p-1}]$. $\alpha = \|\hat{\mathbf{K}}_{[uu]}\| + 1$ for RMT-improved and $\alpha = -1$ for classical Laplacian method. Gaussian kernel with $f(t) = \exp(-t/2)$. Results averaged over 50 runs. [\[Link to code\]](#)

between intra- and inter-class distances vanishes and the performance gain of the RMT-improved method increases significantly, with the standard Laplacian now saturating for larger $n_{[u]}$.

Attempts to reduce the observed limitations of the Laplacian method had in fact been reported in the earlier literature, such as in [Zhou and Belkin, 2011], where an ‘‘iterated Laplacian’’ approach was devised. The basic idea is to replace the kernel matrix \mathbf{K} (or variant of its Laplacian, e.g., $\mathbf{D}^{-1}\mathbf{K}$) by powers of the type \mathbf{K}^m : from a label propagation or graph random walk viewpoint, this consists in ‘‘iterating’’ m propagation steps at once, thereby partially avoiding the problem of uninformative direct neighbors. The performance of the iterated Laplacian approach, for a well-chosen m , in general outperform those of

the standard Laplacian. Yet, from a purely theoretical standpoint, a random matrix analysis would also reveal that the problem of (asymptotic) uselessness of additional unlabeled data remains (although, for finite p, n , it might be “pushed” and appear only at larger values of $n_{[u]}$). In Table 4.2 and Table 4.3, comprehensive simulations are performed on two-class semi-supervised learning for the popular MNIST (from raw data) and then more advanced German Traffic Signs (from HOG features)⁹ datasets to compare the standard Laplacian, and its RMT-improved version, as well as the iterated Laplacian and its correspondingly RMT-improved equivalent. Performance of manifold-based semi-supervised learning are also reported. For fair comparison, the hyperparameters taken for all these simulations (the α parameter and the power m) are optimized in an oracle manner. For both MNIST and German Traffic Signs, the iterated RMT-improved method (which likely benefits from the ideas of [Zhou and Belkin, 2011]) outperforms all other methods.

These numerical evidences on real-world datasets again confirm the strong resilience of the large dimensional statistics approach to real data (or, at least, to some appropriate representation of these data).




Figure 4.21: (**Top**) Histogram of normalized pairwise distances $\|\mathbf{x}_i - \mathbf{x}_j\|^2, i \neq j$ with additive white Gaussian noise of intra- and inter-class MNIST digits (8 versus 9). (**Bottom**) Average accuracy as a function of $n_{[u]}$ with $n_{[l]} = 10$, computed over 1 000 random realizations with 99% confidence intervals represented by shaded regions. [\[Link to code\]](#)

⁹<http://benchmark.ini.rub.de/?section=gtsrb&subsection=dataset>.

Digits	(0, 8)	(2, 7)	(6, 9)
$n_{[u]} = 100$			
RMT-improved	89.5±3.6	89.5±3.4	85.3±5.9
Iterated RMT-improved	89.5±3.6	89.5±3.4	85.3±5.9
Laplacian	75.5±5.6	74.2±5.8	70.0±5.5
Iterated Laplacian	87.2±4.7	86.0±5.2	81.4±6.8
Manifold	88.0±4.7	88.4±3.9	82.8±6.5
$n_{[u]} = 1000$			
RMT-improved	92.2±0.9	92.5±0.8	92.6±1.6
Iterated RMT-improved	92.3±0.9	92.5±0.8	92.9±1.4
Laplacian	65.6±4.1	74.4±4.0	69.5±3.7
Iterated Laplacian	92.2±0.9	92.4±0.9	92.0±1.6
Manifold	91.1±1.7	91.4±1.9	91.4±2.0

Table 4.2: Classification accuracy on MNIST datasets with $n_{[l]} = 10$. Results obtained from 1 000 runs for $n_{[u]} = 100$ and $n_{[u]} = 1000$.

Class ID	(2, 7)	(9, 10)	(11, 18)
$n_{[u]} = 100$			
RMT-improved	79.0±10.4	77.5±9.2	78.5±7.1
Iterated RMT-improved	85.3±5.9	89.2±5.6	90.1±6.7
Laplacian	73.8±9.8	77.3±9.5	78.6±7.2
Iterated Laplacian	83.7±7.2	88.0±6.8	87.1±8.8
Manifold	77.6±8.9	81.4±10.4	82.3±10.8
$n_{[u]} = 1000$			
RMT-improved	83.6±2.4	84.6±2.4	88.7±9.4
Iterated RMT-improved	84.8±3.8	88.0±5.5	96.4±3.0
Laplacian	72.7±4.2	88.9±5.7	95.8±3.2
Iterated Laplacian	83.0±5.5	88.2±6.0	92.7±6.1
Manifold	77.7±5.8	85.0±9.0	90.6±8.1

Table 4.3: Classification accuracy on HOG features of German Traffic Sign datasets with $n_{[l]} = 10$. Results obtained from 1 000 random runs for $n_{[u]} = 100$ and $n_{[u]} = 1000$.

4.5.3 Application to kernel ridge regression

We have discussed applications of kernel methods to unsupervised learning (kernel spectral clustering in Section 4.5.1) and semi-supervised learning (the graph-based approaches of Section 4.5.2). This section closes the investigation of kernel methods by considering now the most popular supervised learning scenario. The first natural method to supervised learning with kernels is kernel ridge regression, which can be used for both regression and classification purposes. In classification applications, it is also referred to as the least-squares support vector machine, or LS-SVM [Suykens and Vandewalle, 1999], and is considered

as a computationally efficient alternative of the classical SVM method (to be discussed later in Chapter 6).

In a binary classification scenario, consider a training set $\{(\mathbf{x}_i, y_i)\}_{i=1}^n$ of size n with data $\mathbf{x}_i \in \mathbb{R}^p$ and labels $y_i = \pm 1$. We denote $\mathbf{x}_i \in \mathcal{C}_1$ if $y_i = -1$ and $\mathbf{x}_i \in \mathcal{C}_2$ if $y_i = +1$. The objective of LS-SVM is to devise a decision function

$$g(\mathbf{x}) = \mathbf{w}^\top \phi(\mathbf{x}_i) + b \quad (4.37)$$

that ideally maps all the (features $\phi(\mathbf{x}_i)$ of the) training data \mathbf{x}_i to the corresponding y_i , and subsequently an unknown test datum \mathbf{x} to its corresponding y value, by solving the following optimization problem

$$\begin{aligned} \arg \min_{\mathbf{w}, b} \quad & \|\mathbf{w}\|^2 + \frac{\gamma}{n} \sum_{i=1}^n e_i^2 \\ \text{s.t.} \quad & y_i = \mathbf{w}^\top \phi(\mathbf{x}_i) + b + e_i, \quad i = 1, \dots, n \end{aligned} \quad (4.38)$$

for some penalty factor $\gamma > 0$ that weights the structural risk $\|\mathbf{w}\|^2$ against the empirical risk $\frac{1}{n} \sum_{i=1}^n e_i^2$.

By introducing the Lagrange multipliers $\{\alpha_i\}_{i=1}^n$, the solution to (4.38) can be expressed as $\mathbf{w} = \sum_{i=1}^n \alpha_i \phi(\mathbf{x}_i)$, where

$$\begin{cases} \boldsymbol{\alpha} &= \mathbf{Q} \left(\mathbf{I}_n - \frac{\mathbf{1}_n \mathbf{1}_n^\top \mathbf{Q}}{\mathbf{1}_n^\top \mathbf{Q} \mathbf{1}_n} \right) \mathbf{y} \\ b &= \frac{\mathbf{1}_n^\top \mathbf{Q} \mathbf{y}}{\mathbf{1}_n^\top \mathbf{Q} \mathbf{1}_n} \end{cases} \quad (4.39)$$

for $\mathbf{y} = [y_1, \dots, y_n]^\top$, $\mathbf{Q} \equiv (\mathbf{K} + \frac{n}{\gamma} \mathbf{I}_n)^{-1}$ and $\mathbf{K} \equiv \{\phi(\mathbf{x}_i)^\top \phi(\mathbf{x}_j)\}_{i,j=1}^n$ the kernel matrix from training data, which is assumed to take the following form

$$\mathbf{K} = \{f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)\}_{i,j=1}^n$$

for some smooth function f .

Given $\boldsymbol{\alpha}$ and b , a new datum \mathbf{x} is then classified into class \mathcal{C}_1 or \mathcal{C}_2 depending on the value of the following decision function

$$g(\mathbf{x}) = \boldsymbol{\alpha}^\top \mathbf{k}(\mathbf{x}) + b, \quad \mathbf{k}(\mathbf{x}) \equiv \{f(\|\mathbf{x} - \mathbf{x}_i\|^2/p)\}_{i=1}^n. \quad (4.40)$$

One of the most popular choices is to use the sign of $g(\mathbf{x})$, and assign \mathbf{x} to class \mathcal{C}_1 if $g(\mathbf{x}) < 0$ and to class \mathcal{C}_2 otherwise. As we shall see, this decision criterion can be highly biased in some cases, when large dimensional data are considered.

Thanks to the ‘‘kernel trick’’, the decision function $g(\mathbf{x})$ no longer needs the evaluation of $\phi(\cdot)$, and can be fully expressed through the kernel function $(\mathbf{x}_i, \mathbf{x}_j) \mapsto f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$, that is indirectly related to $\phi(\cdot)$ through $\phi(\mathbf{x}_i)^\top \phi(\mathbf{x}_j) = f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$. Yet, to allow for a much larger range of functions $f(\cdot)$ (in particular functions f which do not necessarily arise from a feature mapping $\phi(\cdot)$), in the remainder of this section, we shall allow for arbitrary f with a minimalist set of constraints. We shall in particular observe that some functions f , not positive definite and thus not necessarily deriving from a feature map ϕ , prove extremely powerful in some specific scenarios.

4.5.3.1 Large dimensional performance analysis and its implications

As in the previous sections on unsupervised or semi-supervised classification methods, we place ourselves under the following “non-trivial” Gaussian mixture model

$$\mathbf{x}_i \in \mathcal{C}_a \Leftrightarrow \mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{C}_a), \quad a \in \{1, 2\}$$

for $\boldsymbol{\mu}_a \in \mathbb{R}^p$ and $\mathbf{C}_a \in \mathbb{R}^{p \times p}$ such that $\|\mathbf{C}_a\| = O(1)$ and

$$\|\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2\| = O(1), \quad \text{tr}(\mathbf{C}_1 - \mathbf{C}_2) = O(\sqrt{p}), \quad \text{tr}(\mathbf{C}_1 - \mathbf{C}_2)^2 = O(p). \quad (4.41)$$

We assume a training set of n_1 samples in class \mathcal{C}_1 and n_2 samples in class \mathcal{C}_2 so that $n_1 + n_2 = n$, and that n_1, n_2 and p grow at the same rate (i.e., p/n_a remains away from 0 and ∞ in the large n, p limit). We recall from previous sections on kernel methods that, letting $\tau_p = \frac{2}{p} \text{tr } \mathbf{C}^\circ$ with $\mathbf{C}^\circ = \frac{n_1}{n} \mathbf{C}_1 + \frac{n_2}{n} \mathbf{C}_2$, these non-trivial growth rate conditions ensure that

$$\max_{i,j} \{[\mathbf{K}]_{ij} - \tau_p\} \xrightarrow{a.s.} 0. \quad (4.42)$$

As such, the kernel matrix is dominated by the rank-one matrix $f(\tau_p) \mathbf{1}_n \mathbf{1}_n^\top$, which is of operator norm order $O(n)$, and thus of the same of order as $\frac{n}{\gamma} \mathbf{I}_n$ for $\gamma = O(1)$.

As a result, with the (asymptotic) Taylor expansion of the kernel matrix \mathbf{K} derived in Theorem 4.1, it is possible to similarly “linearize” the resolvent $\mathbf{Q} \equiv \left(\mathbf{K} + \frac{n}{\gamma} \mathbf{I}_n\right)^{-1}$ with a Taylor expansion around the leading $f(\tau_p) \mathbf{1}_n \mathbf{1}_n^\top + \frac{n}{\gamma} \mathbf{I}_n$ term. Since the decision function $g(\mathbf{x})$ depends on $\boldsymbol{\alpha}$ and b , that both depend explicitly on \mathbf{Q} , we can then work out an asymptotic linearization of $g(\mathbf{x})$. An asymptotic expression of the misclassification rate of LS-SVM then unfolds, which is a function of the local behavior of kernel function f around τ_p , as well as the data statistics $\boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \mathbf{C}_1, \mathbf{C}_2$. This result is detailed in the following theorem.

Theorem 4.10 ([Liao and Couillet, 2019b, Theorem 2]). *Under the non-trivial Gaussian mixture model in (4.41), we have, for $g(\mathbf{x})$ defined in (4.40),*

$$V_a^{-\frac{1}{2}}(g(\mathbf{x}) - E_a) \xrightarrow{d} \mathcal{N}(0, 1), \quad a \in \{1, 2\}$$

for

$$E_a = c_2 - c_1 + \frac{2}{p}(-1)^a(1 - c_a)\gamma c_1 c_2 \mathcal{D}, \quad V_a = \frac{8}{p^2} \gamma^2 c_1^2 c_2^2 \mathcal{V}_a$$

where $c_a = n_a/n$ and

$$\begin{aligned} \mathcal{D} &= -2f'(\tau_p)\|\Delta\boldsymbol{\mu}\|^2 + \frac{f''(\tau_p)}{p} ((\text{tr } \Delta\mathbf{C})^2 + 2\text{tr}(\Delta\mathbf{C}^2)) \\ \mathcal{V}_a &= \frac{(f''(\tau_p))^2}{p^2} (\text{tr } \Delta\mathbf{C})^2 \cdot \text{tr}(\mathbf{C}_a^2) + 2(f'(\tau_p))^2 \left(\Delta\boldsymbol{\mu}^\top \mathbf{C}_a \Delta\boldsymbol{\mu} + \frac{1}{n} \text{tr } \mathbf{C}_a \left(\frac{\mathbf{C}_1}{c_1} + \frac{\mathbf{C}_2}{c_2} \right) \right) \end{aligned}$$

in which we denoted $\Delta\boldsymbol{\mu} \equiv \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2$ and $\Delta\mathbf{C} \equiv \mathbf{C}_1 - \mathbf{C}_2$.

An immediate remark from Theorem 4.10 is that, since under the setting of (4.41) both \mathcal{D} and \mathcal{V}_a are of order $O(1)$, the decision function is of order $g(\mathbf{x}) = c_2 - c_1 + O(n^{-1})$. This result contradicts the classical ‘‘sign-based’’ decision criterion, by which the decision threshold ξ equals zero, i.e., the new datum \mathbf{x} is assigned to \mathcal{C}_1 if $g(\mathbf{x}) < \xi = 0$ and to \mathcal{C}_2 otherwise. When $c_1 - c_2 \neq 0$ (in unbalanced classification scenarios), this would lead to an asymptotic classification of all new data into one of the two classes. Two options to alleviate this issue are:

1. taking the decision threshold ξ , instead of $\xi = 0$ in the sign-based criterion, to be $\xi = \xi_n = c_2 - c_1 + O(n^{-1})$;
2. normalizing the labels $y_i \in \{-1, +1\}$ as $y_i^* \in \{-n/n_1, +n/n_2\}$, while maintaining the decision threshold to $\xi = 0$ (or technically speaking $O(n^{-1})$). This is also referred to as the Fisher’s targets in the context of kernel Fisher discriminant analysis [Mika et al., 1999]. It can be shown that, when trained with y_i^* , the associated decision function satisfies $g^*(\mathbf{x}) = 0 + O(n^{-1})$.

As a corollary of Theorem 4.10, the asymptotic misclassification rate is a function of the decision threshold ξ_n , the local behavior of kernel function f , the dimensions p, n_1, n_2 as well as the data statistics $\mu_1, \mu_2, \mathbf{C}_1, \mathbf{C}_2$.

Corollary 4.2 (Asymptotic misclassification error rate). *Under the setting of Theorem 4.10, for a decision threshold ξ_n that may depend on n , as $n \rightarrow \infty$,*

$$\begin{aligned} \mathbb{P}(g(\mathbf{x}) > \xi_n \mid \mathbf{x} \in \mathcal{C}_1) - Q\left(\frac{\xi_n - E_1}{\sqrt{V_1}}\right) &\xrightarrow{a.s.} 0 \\ \mathbb{P}(g(\mathbf{x}) < \xi_n \mid \mathbf{x} \in \mathcal{C}_2) - Q\left(\frac{E_2 - \xi_n}{\sqrt{V_2}}\right) &\xrightarrow{a.s.} 0 \end{aligned}$$

for E_a and V_a given in Theorem 4.10 and $Q(x) = \frac{1}{\sqrt{2\pi}} \int_x^\infty \exp(-t^2/2) dt$.

Interestingly, Corollary 4.2 implies that, if one takes $\xi_n = c_2 - c_1 = \frac{n_2}{n} - \frac{n_1}{n}$, the asymptotic classification error is independent of the regularization parameter γ . It is however worth noting that this remark is only valid for $\gamma = O(1)$, i.e., γ is considered to remain a constant as $n, p \rightarrow \infty$, and the threshold is taken to be exactly $c_2 - c_1$.

Remark 4.9 (On non-trivial γ choices). *Since \mathbf{K} is dominated by $f(\tau_p)\mathbf{1}_n\mathbf{1}_n^\top$, taking $\gamma = O(1)$ is a mandatory choice to avoid the asymptotic singularity of the resolvent $\mathbf{Q} = (\mathbf{K} + \frac{n}{\gamma}\mathbf{I}_n)^{-1}$. An alternative approach may consist in working with \mathbf{PKP} for $\mathbf{P} = \mathbf{I}_n - \frac{1}{n}\mathbf{1}_n\mathbf{1}_n^\top$ instead of \mathbf{K} (as in [Elkhalil et al., 2019]), thereby discarding the dominant (and non-informative) matrix $f(\tau_p)\mathbf{1}_n\mathbf{1}_n^\top$ and allowing for γ to be chosen of order $\gamma = O(n)$. In this case, its specific choice would have a nontrivial impact on the classification performance, as in the case of semi-supervised learning in Section 4.5.2. We do not further elaborate on this setting as this moves us rather far from the conventional LS-SVM formulation.*

Due to the concentration of Euclidean distances in large dimensions, the performance of LS-SVM depends on the kernel function f solely via its successive derivatives at τ_p (which, as recalled from Remark 4.4, can be consistently estimated from the data). More discussions on the choice of f are in order.

1. Note that with $f'(\tau_p) = 0$, the difference in statistical means $\Delta\mu$ vanishes from the expressions of \mathcal{D} and \mathcal{V}_a in Theorem 4.10 and the classification can only be performed based on the covariance structures. In this situation, rather surprisingly, if one further assumes $\text{tr}(\mathbf{C}_1 - \mathbf{C}_2) = \text{tr} \Delta \mathbf{C} = o(\sqrt{p})$ (which is below the minimum “distance” $\text{tr} \Delta \mathbf{C} = O(\sqrt{p})$ in (4.41)), then $\mathcal{D} = 2f''(\tau_p) \text{tr}(\Delta \mathbf{C}^2)/p + o(1)$ and $\mathcal{V}_a = o(1)$ so that with, say $\text{tr}(\Delta \mathbf{C}^2) = O(p)$, perfect classification can be achieved. This remark is of particular interest for example when data in different classes are of zero mean, unit Euclidean norm and thus have indistinguishable $\mathbb{E}[\|\mathbf{x}\|^2] = \text{tr} \mathbf{C}_a$. In this case, the covariance “shape” information can be better exploited with the family of kernels such that $f'(\tau_p) = 0$. Figure 4.22 compares the empirical classification error rate for $p = 512$ and $p = 1024$ to the theoretical asymptotic error predicted in Corollary 4.2, and confirms, when $\text{tr} \Delta \mathbf{C} = 0$, the rapid drop of classification error (which ultimately vanishes) as $f'(\tau_p)$ gets close to zero.
2. Since $|E_1 - E_2|$ is proportional to \mathcal{D} and should, for fixed V_a (which does not depend on the signs of $f'(\tau_p)$ and $f''(\tau_p)$), be made as large as possible to achieve optimal classification performance, the kernel function must satisfy $f'(\tau_p) < 0$ and $f''(\tau_p) > 0$. Incidentally, this condition is naturally satisfied by the popular Gaussian kernel $f(x) = \exp(-x/\sigma^2)$ for any σ , but this is not always the case of polynomial kernels.
3. When the difference in statistical means $\|\Delta\mu\|$ is largely dominant over the difference in covariances ($\text{tr} \Delta \mathbf{C}^2/p$ and $\text{tr}(\Delta \mathbf{C}^2)/p$, from Theorem 4.10, both $E_a - (c_2 - c_1)$ and $\sqrt{V_a}$ are approximately proportionally to $f'(\tau_p)$, making the choice of the kernel function eventually irrelevant in this case, so long that $f'(\tau_p) \neq 0$.

4.5.3.2 Application to real-world data

Although derived from a simple Gaussian mixture model, the previous theoretical results, when applied to popular large dimensional real-world datasets, again show a (at first unexpected) similar behavior. Figure 4.23 considers the classification of (two from the ten classes of) MNIST and Fashion-MNIST data. Despite the obvious non-Gaussianity as well as the clearly different natures of the data (from the two datasets), the empirical histogram of the decision function $g(\mathbf{x})$ always behaves surprisingly close to its limiting behavior predicted by Theorem 4.10.

In Figure 4.24, the classification error rate is displayed as a function of the decision threshold ξ , again for both MNIST and Fashion-MINIST data. The




Figure 4.22: Classification error of LS-SVM, $p = 512$, $n = 2048$, $c_1 = c_2 = 1/2$, $\gamma = 1$, second order polynomial kernel with $f(\tau_p) = 4$ and $f''(\tau_p) = 2$. For Gaussian data $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{C}_a)$ with $\mathbf{C}_1 = \mathbf{I}_p$ and $[\mathbf{C}_2]_{ij} = .4^{|i-j|}$. Empirical results averaged over 30 runs. [Link to code]




Figure 4.23: Empirical histogram of $g(\mathbf{x})$ versus the Gaussian limiting behavior predicted in Theorem 4.10, $n = 2048$, $p = 784$, $\gamma = 1$ with Gaussian kernel, for MINST (left, 7 versus 9) and Fashion-MNIST (right, 8 versus 9) data. Results averaged over 20 runs. [Link to code]

conclusion that the optimal decision threshold should approximately be $c_2 - c_1$ rather than 0 is conclusively observed to hold true in both cases.




Figure 4.24: Classification error rate as a function of decision threshold ξ , with $n = 512, p = 784, c_2 - c_1 = 1/8$ in **purple**, $\gamma = 1$, Gaussian kernel for MNIST (**left**) and Fashion-MNIST data (**right**). With empirical optimal decision thresholds $\xi_* = 0.12$ (**left**) and 0.11 (**right**) in **red**. [Link to code]

4.5.4 Summary of Section 4.5

In this section, we discussed the implications of random matrix analyses to unsupervised, semi-supervised, and supervised learning methods. By assuming the data vectors independently drawn from a k -class Gaussian mixture $\mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{C}_a)$, $a \in \{1, \dots, k\}$, the precise asymptotic performance of various algorithms (spectral clustering in Theorem 4.7, graph-based semi-supervised learning in Theorem 4.8, and the LS-SVM classification approach in Theorem 4.10) are derived, as a function of the dimensionality ratio p/n , the data (discriminative) statistics, generally under the form of

$$\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|, \quad \text{tr}(\mathbf{C}_a - \mathbf{C}_b)/\sqrt{p}, \quad \text{tr}(\mathbf{C}_a - \mathbf{C}_b)^2/p \quad (4.43)$$

as well as the hyperparameters of the algorithm such as the choice of kernel function, the graph regularization parameter α in (4.23), or the (ridge) regularization penalty γ in (4.38). As a result, with the access to (estimates of) these key statistics, one can then “optimally” tune these hyperparameters by optimizing over the theoretical performance formulas, which avoids cross-validation procedures that may “consume” a certain amount of training data. In the semi-supervised and supervised cases, these key statistics can be estimated from the labeled data, as described in the following remark.

Remark 4.10 (Estimation of GMM discriminative statistics). Denote $\mathbf{X}_a \in \mathbb{R}^{p \times n_a}, \mathbf{X}_b \in \mathbb{R}^{p \times n_b}$ the data (sub)matrix of class C_a and C_b , respectively, then, $\|\boldsymbol{\mu}_a - \boldsymbol{\mu}_b\|$ and $\text{tr}(\mathbf{C}_a - \mathbf{C}_b)/\sqrt{p}$ can be consistently estimated via their associated empirical estimators, i.e., for $\hat{\boldsymbol{\mu}}_a \equiv \frac{1}{n_a} \mathbf{X}_a \mathbf{1}_{n_a}$ and $\hat{\mathbf{C}}_a \equiv \frac{1}{n_a-1} (\mathbf{X}_a -$

$$\hat{\mu}_a \mathbf{1}_{n_a}^T) (\mathbf{X}_a - \hat{\mu}_a \mathbf{1}_{n_a}^T)^T, \\ \|\hat{\mu}_a - \hat{\mu}_b\| - \|\mu_a - \mu_b\| = o(1), \quad \text{tr}(\hat{\mathbf{C}}_a - \hat{\mathbf{C}}_b)/\sqrt{p} - \text{tr}(\mathbf{C}_a - \mathbf{C}_b)/\sqrt{p} = o(1). \quad (4.44)$$

For the covariance (Frobenius) distance $\text{tr}(\mathbf{C}_a - \mathbf{C}_b)^2/p$, it follows from Exercise 8 that

$$\left(\frac{1}{p} \text{tr}(\hat{\mathbf{C}}_a^2 + \hat{\mathbf{C}}_b^2) - \frac{(\text{tr} \hat{\mathbf{C}}_a)^2}{pn_a} - \frac{(\text{tr} \hat{\mathbf{C}}_b)^2}{pn_b} - \frac{2}{p} \text{tr}(\hat{\mathbf{C}}_a \hat{\mathbf{C}}_b) \right) - \frac{1}{p} \text{tr}(\mathbf{C}_a - \mathbf{C}_b)^2 = o(1). \quad (4.45)$$

In particular, for kernels of the type $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$ and $f(\mathbf{x}_i^T \mathbf{x}_j/p)$, the performance of kernel-based methods depends on the (smooth) function f only via the successive derivatives $f(\tau_p)$, $f'(\tau_p)$ and $f''(\tau_p)$ (with τ_p obtained from Remark 4.4). As a consequence, all such f yield asymptotically the same performance as the simple quadratic function $f(t) = at^2 + bt + c$ and it thus suffices to perform a three-dimensional optimization procedure (of the coefficients) of the “universal” quadratic function with the above estimated statistics. This remark also holds for α - β and properly scaling kernels, discussed in Section 4.3 and 4.4.

In the unsupervised case, however, while it is always possible to estimate the key parameter τ in a consistent manner (as in Remark 4.4 which does not need labeled data), it is unlikely to make a good estimate of the class discriminative statistics (as in Remark 4.10 that depends on the data classes) without performing any form of clustering beforehand.

To conclude, let us insist again on the capability of random matrix analyses to unveil critical but systematic behavior in large dimensional learning problems, such as the crucial choice of hyperparameter $\alpha = -1 + O(p^{-1/2})$ in graph-based semi-supervised learning (Theorem 4.8), and the decision bias in unbalanced kernel LS-SVM classification (Theorem 4.10): such behavior is *independent* of the precise values of the data statistics, empirically observed in real-world datasets that are not necessarily Gaussian, and, as discussed later in Chapter 8, theoretically supported by universality arguments of large dimensional data.

4.6 Concluding remarks

Before the present chapter, the first part of the monograph was mostly concerned with the sample covariance matrix model \mathbf{XX}^T/n (as well more marginally as on the Wigner model \mathbf{X}/\sqrt{n} for symmetrical \mathbf{X}), where the columns of \mathbf{X} are independent and the entries of each column are independent or linearly dependent. Historically, this model and its numerous variants (with a variance profile, with right-side correlation, summed up to other independent matrices of the same form, etc.) have covered most of the mathematical and applied interest of the first two decades (since the early nineties) of intense random matrix advances. The main drivers for these early developments were statistics, signal processing, and wireless communications. The present chapter leaped much further in

considering now random matrix models with possibly highly correlated entries, with a specific focus on kernel matrices: for the first time, the machine learning applications clearly steered mathematical research towards these models. The main reason lies in the poor understanding and tractability of kernel matrices in finite-dimensional settings: where only (as we saw, sometimes wrong) intuitions and heuristics were available, random matrix theory now provides accurate (and asymptotically exact) performance assessment along with the possibility to largely improve the performance of kernel-based machine learning methods. This, in effect, creates a small revolution (not to the extent of the deep learning revolution though) in our understanding of machine learning on realistic large datasets.

A first important finding of the analysis of large dimensional kernel statistics reported here is the ubiquitous character of the Marčenko-Pastur and the semi-circular laws. As a matter of fact, all random matrix models studied in this chapter, and in particular the kernel regimes $f(\mathbf{x}_i^\top \mathbf{x}_j/p)$ (which concentrate around $f(0)$) and $f(\mathbf{x}_i^\top \mathbf{x}_j/\sqrt{p})$ (which tends to $f(\mathcal{N}(0, 1))$), have a limiting eigenvalue distribution akin to a combination of the two laws. This combination may vary from case to case (compare for instance the results of Exercise 3 to Theorem 4.4), but is often parametrized in a such way that the Marčenko-Pastur and semicircle laws appear as limiting cases (in the context of Exercise 3, they correspond to the limiting cases of dense versus sparse kernels, and in Theorem 4.4 to the limiting cases of linear versus “purely” nonlinear kernels).

A second point of importance, worth recalling, is the range of valid kernels analyzed: for the “natural” scaling $1/p$ (so kernels of the type $f(\mathbf{x}_i^\top \mathbf{x}_j/p)$ or $f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)$), all kernel functions $f(t)$ are equivalent so long that they are differentiable and have the same first three derivatives at the concentration point $t = 0$ (for inner-product) or τ_p (for distance kernels); for the “proper” scaling $1/\sqrt{p}$ (so kernels of the form $f(\mathbf{x}_i^\top \mathbf{x}_j/\sqrt{p})$ or $f(\sqrt{p}[\|\mathbf{x}_i - \mathbf{x}_j\|^2/p - \tau_p])$), kernel functions f are equivalent if they share in common three specific Gaussian moments (i.e., $\int P(t)f(t)\mu(dt)$ for μ the Gaussian measure). As such, while it is not necessarily surprising to see the naturally scaling kernels being equivalent if they have similar behavior near the concentration point, it is quite surprising at first that properly scaling kernel functions f , which “see” values in the whole real axis, do not offer more flexibility or diversity. For both types of kernels, only the first elementary moments of the data are accounted for. The main difference between naturally and properly scaling kernels though, is the possibility of the latter to study very discontinuous kernel functions, such as the sign (or more generally quantization) and thresholding functions: this finds a broad range of applications in low-cost kernel techniques. On this aspect, Chapter 5 recalls the intimate connection between neural networks, random projections, and their limiting kernels: a direct connection can be established between sparsification techniques in neural networks (such as the popular random or deterministic dropout procedure) and sparsification techniques in kernel methods. The present chapter may then provide some keys to understanding and improving computationally constrained methods in the performance-complexity tradeoff of

methods beyond kernels.

This being said, the analysis of kernel methods is far from over. Many kernel matrices remain out of analytical reach by the current random matrix machinery. This is in particular the case of “rank correlation” (also referred to as U-statistics) matrices, such as Spearman- ρ [Spearman, 1987] or Kendall- τ [Kendall, 1938]. For data vectors $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$, the entry \mathbf{K}_{ij} of a rank correlation matrix \mathbf{K} evaluates a function on the *ranks* of the entries $[\mathbf{x}_i]_1, \dots, [\mathbf{x}_i]_p$ and $[\mathbf{x}_j]_1, \dots, [\mathbf{x}_j]_p$ of \mathbf{x}_i and \mathbf{x}_j . For Kendall- τ , \mathbf{K}_{ij} evaluates the number of pairs $[\mathbf{x}_i]_a, [\mathbf{x}_i]_b$ such that $\text{sign}([\mathbf{x}_i]_a - [\mathbf{x}_i]_b) = \text{sign}([\mathbf{x}_j]_a - [\mathbf{x}_j]_b)$. As for Spearman- ρ , \mathbf{K}_{ij} evaluates the empirical correlation between the ranks $\text{rg}([\mathbf{x}_i]_a)$ and $\text{rg}([\mathbf{x}_j]_a)$, for $a = 1, \dots, p$. For these kernels, the rank variables create correlations between the entries of the data matrix \mathbf{X} which prevent classical random matrix tools to immediately apply. The problem was worked around in [Bandeira et al., 2017] for \mathbf{X} having fully i.i.d. entries, exploiting there the very fact that the i.i.d. nature of variables induces an i.i.d. distribution of the ranks. The generalization to non-i.i.d. random variables would however fail in general.

Another important family of difficult-to-grasp kernels is that of k-nearest neighbor (k-NN) kernels [Fix and Hodges, 1989]. These kernel matrices \mathbf{K} are such that $[\mathbf{K}]_{ij}$ is only nonzero if \mathbf{x}_j is one of the k-nearest neighbors of \mathbf{x}_i , among $\mathbf{x}_1, \dots, \mathbf{x}_n$ in the sense of a given metric (typically of the Euclidean distance). The induced dependence between the entries of such matrices is more subtle to grasp, leaving for the moment the study of these kernels, widely used in machine learning applications largely open.

4.7 Practical course material

Practical Lecture Material 3 (Complexity-performance trade-off in spectral clustering with sparse kernel, from [Zarrouk et al., 2020]). *In this exercise, we study the spectrum of a “punctured” version $\mathbf{K} = \mathbf{B} \odot (\mathbf{X}^\top \mathbf{X}/p)$ (with the Hadamard product $[\mathbf{A} \odot \mathbf{B}]_{ij} = [\mathbf{A}]_{ij} [\mathbf{B}]_{ij}$) of the linear kernel $\mathbf{X}^\top \mathbf{X}/p$, with $\mathbf{X} \in \mathbb{R}^{p \times n}$ and a symmetric random mask-matrix $\mathbf{B} \in \{0, 1\}^{n \times n}$ having independent $[\mathbf{B}]_{ij} \sim \text{Bern}(\epsilon)$ entries for $i \neq j$ (up to symmetry) and $[\mathbf{B}]_{ii} = b \in \{0, 1\}$ fixed, in the limit $p, n \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$, thereby mimicking the behavior of computing only a proportion $\epsilon \in (0, 1)$ of the entries of $\mathbf{X}^\top \mathbf{X}/n$. Letting $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ with \mathbf{x}_i independently and uniformly drawn from the following symmetric two-class Gaussian mixture*

$$\mathcal{C}_1 : \mathbf{x}_i \sim \mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p), \quad \mathcal{C}_2 : \mathbf{x}_i \sim \mathcal{N}(+\boldsymbol{\mu}, \mathbf{I}_p) \quad (4.46)$$

for $\boldsymbol{\mu} \in \mathbb{R}^p$ such that $\|\boldsymbol{\mu}\| = O(1)$ with respect to n, p . The idea is to study the effect of uniformly “zeroing-out” with the Bernoulli \mathbf{B} on the presence of an isolated spike in the spectrum of \mathbf{K} , and thus on the spectral clustering performance of data (class \mathcal{C}_1 versus \mathcal{C}_2).

We will study the spectrum of \mathbf{K} using Stein’s lemma and the Gaussian method discussed in Section 2.2.2.2. Let $\mathbf{Z} = [\mathbf{z}_1, \dots, \mathbf{z}_n] \in \mathbb{R}^{p \times n}$ for $\mathbf{z}_i = \mathbf{x}_i -$

$(-1)^a \boldsymbol{\mu} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ with $\mathbf{x}_i \in \mathcal{C}_a$ and $\mathbf{M} = \boldsymbol{\mu} \mathbf{j}^\top$ with $\mathbf{j} = [-\mathbf{1}_{n/2}, \mathbf{1}_{n/2}]^\top \in \mathbb{R}^n$ so that $\mathbf{X} = \mathbf{M} + \mathbf{Z}$. First show that, for $\mathbf{Q} \equiv \mathbf{Q}(z) = (\mathbf{K} - z\mathbf{I}_n)^{-1}$,

$$\begin{aligned}\mathbf{Q} &= -\frac{1}{z}\mathbf{I}_n + \frac{1}{z}\left(\frac{\mathbf{Z}^\top \mathbf{Z}}{p} \odot \mathbf{B}\right)\mathbf{Q} + \frac{1}{z}\left(\frac{\mathbf{Z}^\top \mathbf{M}}{p} \odot \mathbf{B}\right)\mathbf{Q} \\ &\quad + \frac{1}{z}\left(\frac{\mathbf{M}^\top \mathbf{Z}}{p} \odot \mathbf{B}\right)\mathbf{Q} + \frac{1}{z}\left(\frac{\mathbf{M}^\top \mathbf{M}}{p} \odot \mathbf{B}\right)\mathbf{Q}.\end{aligned}$$

To proceed, we need to go slightly beyond the study of these four terms. Specifically, using Stein's lemma, Lemma 2.11, show that, for arbitrary matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ of bounded norm,

$$\begin{aligned}\mathbb{E}\left[\left(\frac{\mathbf{Z}^\top \mathbf{Z}}{p} \odot \mathbf{A}\right)\mathbf{Q}\right]_{ij} &= \mathbf{A}_{ii}\mathbb{E}[\mathbf{Q}_{ij}] - \mathbb{E}\left[\frac{1}{n}\text{tr}\left(\mathbf{Q}\mathbf{D}_{\mathbf{b}_i}\frac{\mathbf{Z}^\top \mathbf{Z}}{p}\mathbf{D}_{\mathbf{a}_i}\right)\mathbf{Q}_{ij}\right] \\ &\quad - \mathbb{E}\left[\frac{1}{n}\text{tr}\left(\mathbf{Q}\mathbf{D}_{\mathbf{b}_i}\frac{\mathbf{M}^\top \mathbf{Z}}{p}\mathbf{D}_{\mathbf{a}_i}\right)\mathbf{Q}_{ij}\right] \\ \mathbb{E}\left[\left(\frac{\mathbf{Z}^\top \mathbf{M}}{p} \odot \mathbf{A}\right)\mathbf{Q}\right]_{ij} &= -\mathbb{E}\left[\frac{1}{n}\text{tr}\left(\mathbf{Q}\mathbf{D}_{\mathbf{b}_i}\frac{\mathbf{Z}^\top \mathbf{M}}{p}\mathbf{D}_{\mathbf{a}_i}\right)\mathbf{Q}_{ij}\right] \\ &\quad - \mathbb{E}\left[\frac{1}{n}\text{tr}\left(\mathbf{Q}\mathbf{D}_{\mathbf{b}_i}\frac{\mathbf{M}^\top \mathbf{M}}{p}\mathbf{D}_{\mathbf{a}_i}\right)\mathbf{Q}_{ij}\right] \\ \mathbb{E}\left[\left(\frac{\mathbf{M}^\top \mathbf{Z}}{p} \odot \mathbf{A}\right)\mathbf{Q}\right]_{ij} &= -\mathbb{E}\left[\frac{\mathbf{M}^\top \mathbf{Z}}{p}\mathbf{D}_{\mathbf{a}_i, \mathbf{B}}\mathbf{Q}\right]_{ij} - \mathbb{E}\left[\frac{\mathbf{M}^\top \mathbf{M}}{p}\mathbf{D}_{\mathbf{a}_i, \mathbf{B}}\mathbf{Q}\right]_{ij}\end{aligned}$$

where $\mathbf{a}_i \in \mathbb{R}^n$ is the i -th (transposed) row of \mathbf{A} , $\mathbf{b}_i \in \mathbb{R}^n$ the i -th column of \mathbf{B} , $\mathbf{D}_{\mathbf{a}_i} \equiv \text{diag}(\mathbf{a}_i)$, $\mathbf{D}_{\mathbf{b}_i} \equiv \text{diag}(\mathbf{b}_i)$ and

$$\mathbf{D}_{\mathbf{a}_i, \mathbf{B}} = \text{diag}\left\{\frac{1}{n}\text{tr}(\mathbf{Q}\mathbf{D}_{\mathbf{a}_i}\mathbf{D}_{\mathbf{b}_k})\right\}_{k=1}^n = \text{diag}\left\{\frac{1}{n}\text{tr}(\mathbf{Q} \odot \mathbf{a}_i \mathbf{b}_k^\top)\right\}_{k=1}^n \quad (4.47)$$

and conclude that

$$\begin{aligned}\mathbb{E}[\mathbf{Q}_{ij}] &\simeq -\frac{1}{z}\delta_{ij} + \frac{1}{z}\left[\mathbf{B}_{ii} - \frac{1}{n}\text{tr}\left(\mathbf{Q}\mathbf{D}_{\mathbf{B}_{\cdot i}}\frac{1}{p}(\mathbf{Z} + \mathbf{M})^\top(\mathbf{Z} + \mathbf{M})\mathbf{D}_{\mathbf{B}_{i \cdot}}\right)\right]\mathbb{E}[\mathbf{Q}_{ij}] \\ &\quad + \frac{1}{z}\mathbb{E}\left[\left(\frac{\mathbf{M}^\top \mathbf{M}}{p} \odot \mathbf{B}\right)\mathbf{Q}\right]_{ij} - \frac{1}{z}\mathbb{E}\left[\frac{\mathbf{M}^\top(\mathbf{Z} + \mathbf{M})}{p}\mathbf{D}_{i, \mathbf{B}}\mathbf{Q}\right]_{ij} \quad (4.48)\end{aligned}$$

for $\mathbf{B}_{\cdot i} = \mathbf{b}_i$ the i -th column of \mathbf{B} , $\mathbf{B}_{i \cdot}$ the i -th row of \mathbf{B} and

$$\mathbf{D}_{i, \mathbf{B}} = \mathbf{D}_{\mathbf{B}_{\cdot i}, \mathbf{B}} = \text{diag}\left\{\frac{1}{n}\text{tr}(\mathbf{Q} \odot \mathbf{B}_{\cdot i} \mathbf{B}_{k \cdot})\right\}_{k=1}^n \quad (4.49)$$

The main difficulty here lies in the last term $\frac{1}{n}\mathbb{E}[\mathbf{M}^\top(\mathbf{Z} + \mathbf{M})\mathbf{D}_{i, \mathbf{B}}\mathbf{Q}]_{ij}$, for which we will admit the following result

$$[\mathbf{D}_{i, \mathbf{B}}]_{kk} = \frac{1}{n}\text{tr}(\mathbf{Q} \odot \mathbf{b}_i \mathbf{b}_k^\top) = \begin{cases} \frac{\epsilon}{n_2} \text{tr } \mathbf{Q} + o(1), & \text{for } i = k \\ \frac{\epsilon^2}{n} \text{tr } \mathbf{Q} + o(1), & \text{otherwise,} \end{cases} \quad (4.50)$$

which follows from the symmetry of \mathbf{B} and a concentration argument. From this result, along with the remark that $\mathbf{A} = \mathbf{A} \odot \mathbf{1}_n \mathbf{1}_n^\top$ and $\|\boldsymbol{\mu}\| = O(1)$, show that

$$\mathbb{E} \left[\frac{\mathbf{M}^\top \mathbf{M} \odot \mathbf{B}}{p} \mathbf{Q} \right]_{ij} = -\mathbb{E} \left[\frac{\mathbf{M}^\top \mathbf{M} \mathbf{Q}}{p} \right]_{ij} \frac{1}{1 + \frac{\epsilon}{n} \text{tr } \mathbf{Q}} \frac{\epsilon^2}{n} \text{tr } \mathbf{Q} + o(1).$$

To obtain a self-consistent equation, we need to find a recursive relation for the quantities

$$L_{ij} \equiv \frac{1}{n} \text{tr} \left(\mathbf{Q} \left(\frac{1}{p} (\mathbf{Z} + \mathbf{M})^\top (\mathbf{Z} + \mathbf{M}) \odot \mathbf{b}_i \mathbf{b}_j^\top \right) \right)$$

which appeared in the development of (4.48). By interchangeability, observe that $L_{ij} = L_{\neq} + o(1)$ all $i \neq j$ and $L_{ii} = L_{=} + o(1)$ for all i , for some L_{\neq} and $L_{=}$. Show further that

$$L_{\neq} = \frac{\frac{\epsilon^2}{n} \text{tr } \mathbf{Q}}{1 + \frac{\epsilon}{n} \text{tr } \mathbf{Q}}, \quad L_{=} = \frac{\epsilon}{n} \text{tr } \mathbf{Q} - \frac{\epsilon^3 (\frac{1}{n} \text{tr } \mathbf{Q})^2}{1 + \frac{\epsilon}{n} \text{tr } \mathbf{Q}}. \quad (4.51)$$

Conclude from these developments that the following deterministic equivalent relation holds

$$\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z) = m(z) \left(\mathbf{I}_n + \|\boldsymbol{\mu}\|^2 \frac{\epsilon m(z)}{c + \epsilon m(z)} \frac{\mathbf{j} \mathbf{j}^\top}{n} \right)^{-1} \quad (4.52)$$

where $m(z)$ is the Stieltjes transform (the limit of $\frac{1}{n} \text{tr } \mathbf{Q}(z)$) solution to

$$z = b - \frac{1}{m(z)} - \frac{\epsilon}{c} m(z) + \frac{\epsilon^3 m^2(z)}{c(c + \epsilon m(z))} \quad (4.53)$$

in which we recall that $c = \lim p/n$ and $b = [\mathbf{B}]_{ii}$ for all i . Show in particular that, up to shift and scaling, we retrieve the Marčenko-Pastur in the limit $\epsilon = 1$ and the semicircle law in the limit $\epsilon \rightarrow 0$. Confirm numerically the ‘‘transition’’ from semicircle to Marčenko-Pastur behavior by tuning ϵ as in Figure 4.25.

Using a spiked model approach (Section 2.5), define the functions

$$\begin{aligned} F(x) &= x^4 + 2x^3 + \left(1 - \frac{c}{\epsilon}\right)x^2 - 2cx - c, \\ G(x) &= b + \frac{\epsilon}{c}(1+x) + \frac{1}{1+x} + \frac{\epsilon}{x(1+x)}, \end{aligned}$$

and let Γ be the largest real solution to $F(\Gamma) = 0$. Show that the largest eigenvalue-eigenvector pair $(\hat{\lambda}, \hat{\mathbf{u}})$ of \mathbf{K} satisfy

$$\hat{\lambda} \rightarrow \lambda = \begin{cases} G(\rho), & \rho > \Gamma \\ G(\Gamma), & \rho \leq \Gamma \end{cases} \quad \frac{1}{n} |\mathbf{j}^\top \hat{\mathbf{u}}|^2 \rightarrow \zeta = \begin{cases} \frac{F(\rho)}{\rho(1+\rho)^3}, & \rho > \Gamma \\ 0, & \rho \leq \Gamma \end{cases}$$

almost surely, where we denote $\rho = \lim \|\boldsymbol{\mu}\|^2$.




Figure 4.25: Eigenvalues of sparse \mathbf{K} versus the limiting laws, for $\boldsymbol{\mu} = [1, \mathbf{0}_{p-1}]$, $b = 0$, $p = 512$ and $n = 2048$. [\[Link to code\]](#)

notations to clean, and hopefully, add some simulations!

Practical Lecture Material 4 (Towards transfer learning with kernel regression, [Tiomoko et al., 2021]). In this exercise, we consider the extension of the supervised kernel regression (or LS-SVM) approach to a transfer learning scenario. Specifically, consider two datasets $\mathbf{X}_1 \in \mathbb{R}^{p \times n_1}$ and $\mathbf{X}_2 \in \mathbb{R}^{p \times n_2}$ associated to two learning tasks T_1 and T_2 . The exercise consists in “transferring” the binary classification of task T_1 with training examples $\mathbf{X}_1 = [\mathbf{X}_1^{(1)}, \mathbf{X}_1^{(2)}]$, $\mathbf{X}_i^{(j)} \in \mathbb{R}^{p \times n_{ij}}$, and labels $\mathbf{y}_1 \in \mathbb{R}^{n_1}$, towards the learning of the binary classification of task T_2 with training examples $\mathbf{X}_2 = [\mathbf{X}_2^{(1)}, \mathbf{X}_2^{(2)}]$ and labels $\mathbf{y}_2 \in \mathbb{R}^{n_2}$. Furthermore, $\mathbf{X}_i^{(j)} = [\mathbf{x}_{i1}, \dots, \mathbf{x}_{in_{ij}}] \in \mathbb{R}^{p \times n_{ij}}$ and we assume, as usual, that $\mathbf{x}_{ij} \sim \mathcal{N}(\boldsymbol{\mu}_{ij}, \mathbf{I}_p)$.

Extending the kernel regression (or LS-SVM) approach of Section 4.5.3, we here seek two hyperplanes $\boldsymbol{\omega}_1 = \boldsymbol{\omega}_0 + \mathbf{v}_1$ and $\boldsymbol{\omega}_2 = \boldsymbol{\omega}_0 + \mathbf{v}_2$ and biases b_1 and b_2 which solves the joint optimization problem:

$$\min_{(\boldsymbol{\omega}_0, \mathbf{V}, \mathbf{b}) \in \mathbb{R}^p \times \mathbb{R}^{p \times 2} \times \mathbb{R}^2} \frac{1}{2\lambda} \|\boldsymbol{\omega}_0\|^2 + \frac{1}{2} \sum_{i=1}^2 \frac{\|\mathbf{v}_i\|^2}{\gamma_i} + \frac{1}{2} \sum_{i=1}^2 \|\boldsymbol{\xi}_i\|^2$$

$$\boldsymbol{\xi}_i = \mathbf{y}_i - (\mathbf{X}_i^\top \boldsymbol{\omega}_i + b_i \mathbf{1}_{n_i}), \quad i \in \{1, 2\}.$$

First show that the problem solves two parallel learning tasks as $\lambda \rightarrow 0$ and a single learning task for $\lambda \rightarrow \infty$, and thus conclude on the role of the hyperparameters γ_i .

Using the Lagrangian multipliers method, show that

$$\mathbf{y}_i = (\lambda + \gamma_i) \mathbf{X}_i^\top \boldsymbol{\alpha}_i + \lambda \sum_{j \neq i} \mathbf{X}_i^\top \mathbf{X}_j \boldsymbol{\alpha}_j + b_i \mathbf{1}_{n_i} + \boldsymbol{\alpha}_i$$

$$0 = \mathbf{1}_{n_i}^\top \boldsymbol{\alpha}_i$$

for $\boldsymbol{\alpha} = [\boldsymbol{\alpha}_1^\top, \boldsymbol{\alpha}_2^\top]^\top \in \mathbb{R}^n$ some appropriately chosen Lagrange multipliers. Then, gathering $\mathbf{y} = [\mathbf{y}_1^\top, \mathbf{y}_2^\top]^\top \in \mathbb{R}^n$, $\boldsymbol{\alpha} = [\boldsymbol{\alpha}_1^\top, \boldsymbol{\alpha}_2^\top] \in \mathbb{R}^n$, and using the shortcut

notations

$$\mathbf{Z} = \begin{pmatrix} \mathbf{X}_1 & 0 \\ 0 & \mathbf{X}_2 \end{pmatrix} \in \mathbb{R}^{2p \times n}, \quad \text{and} \quad \mathbf{P} = \begin{pmatrix} \mathbf{1}_{n_1} & 0 \\ 0 & \mathbf{1}_{n_2} \end{pmatrix} \in \mathbb{R}^{n \times 2},$$

confirm that we then obtain the compact form for $\boldsymbol{\alpha}$ and \mathbf{b}

$$\mathbf{P}\mathbf{b} + \mathbf{Q}^{-1}\boldsymbol{\alpha} = \mathbf{y} \quad \text{and} \quad \mathbf{P}^T\boldsymbol{\alpha} = \mathbf{0}_2$$

where

$$\begin{aligned} \mathbf{Q} &\equiv \left(\frac{1}{2p} \mathbf{Z}^T \mathbf{A} \mathbf{Z} + \mathbf{I}_n \right)^{-1} \\ \mathbf{A} &\equiv (\mathbf{D}_\gamma + \lambda \mathbf{1}_2 \mathbf{1}_2^T) \otimes \mathbf{I}_p \in \mathbb{R}^{2p \times 2p} \end{aligned}$$

in which ‘ \otimes ’ is the Kronecker product of matrices and \mathbf{D}_x denotes a generic diagonal matrix with diagonal entries x_1, x_2, \dots

Conclude on the final expressions for $\boldsymbol{\alpha}$ and \mathbf{b} and show finally that

$$\boldsymbol{\omega}_i = \left(\mathbf{e}_i^{[2]T} \otimes \mathbf{I}_p \right) \mathbf{A} \mathbf{Z} \boldsymbol{\alpha}$$

where $\mathbf{e}_i^{[2]} \in \mathbb{R}^2$ is the canonical basis vector $(0, 1)$ with entry 1 in position i . Further show that the classification score for task T_i of any test point $\hat{\mathbf{x}}$ is given by

$$g_i(\hat{\mathbf{x}}) = \frac{1}{2p} \left(\mathbf{e}_i^{[2]} \otimes \hat{\mathbf{x}}_i \right)^T \mathbf{A} \mathbf{Z} \boldsymbol{\alpha} + b_i$$

in which $\hat{\mathbf{x}}_i = \hat{\mathbf{x}} - \frac{1}{n_i} \mathbf{X}_i \mathbf{1}_{n_i}$. From these results and the matrix identity $(\mathbf{I} + \mathbf{DB})^{-1} \mathbf{D} = \mathbf{D}(\mathbf{I} + \mathbf{BD})^{-1}$, show that $g_i(\hat{\mathbf{x}})$ can be also expressed as

$$g_i(\hat{\mathbf{x}}) = \frac{1}{2p} \left(\mathbf{e}_i^{[2]} \otimes \hat{\mathbf{x}} \right)^T \mathbf{A}^{\frac{1}{2}} \tilde{\mathbf{Q}} \mathbf{A}^{\frac{1}{2}} \mathbf{Z} (\mathbf{y} - \mathbf{P}\mathbf{b}) + b_i$$

where $\tilde{\mathbf{Q}} = (\frac{1}{2p} \mathbf{A}^{\frac{1}{2}} \mathbf{Z} \mathbf{Z}^T \mathbf{A}^{\frac{1}{2}} + \mathbf{I}_{2p})^{-1}$.

Using either of the results on deterministic equivalents in the monograph, show then that

$$\mathbf{A}^{\frac{1}{2}} \tilde{\mathbf{Q}} \mathbf{A}^{\frac{1}{2}} \mathbf{Z} \leftrightarrow \mathbf{A}^{\frac{1}{2}} \bar{\tilde{\mathbf{Q}}} \mathbf{A}^{\frac{1}{2}} \mathbf{M}_\Delta \mathbf{J}^T$$

where $\bar{\tilde{\mathbf{Q}}}$ is a deterministic equivalent for $\tilde{\mathbf{Q}}$ given by

$$\begin{aligned} \tilde{\mathbf{Q}} \leftrightarrow \bar{\tilde{\mathbf{Q}}} &\equiv \left(\sum_{i=1}^2 \sum_{j=1}^2 \tilde{\Delta}_{ij} \mathbf{C}_{ij} + \mathbf{I}_{2p} \right)^{-1} \\ \mathbf{C}_{ij} &= \mathbf{A}^{\frac{1}{2}} \left(\mathbf{e}_i^{[2]} \mathbf{e}_i^{[2]T} \otimes (\mathbf{I}_p + \boldsymbol{\mu}_{ij} \boldsymbol{\mu}_{ij}^T) \right) \mathbf{A}^{\frac{1}{2}} \end{aligned}$$

and

$$\begin{aligned}\mathbf{M}_\Delta &\equiv \mathbf{M} \sum_{ij} \frac{1}{1 + \Delta_i} \mathbf{e}_{ij}^{[4]} \mathbf{e}_{ij}^{[4]\top} \\ \mathbf{M} &= \left(e_1^{[2]} \otimes [\boldsymbol{\mu}_{11}, \boldsymbol{\mu}_{12}], e_2^{[2]} \otimes [\boldsymbol{\mu}_{21}, \boldsymbol{\mu}_{22}] \right) \\ \Delta_i &= \frac{1}{2p} \text{tr} \left(\mathbf{A}^{\frac{1}{2}} \left(\mathbf{e}_i^{[2]} \mathbf{e}_i^{[2]\top} \otimes \mathbf{I}_p \right) \mathbf{A}^{\frac{1}{2}} \bar{\mathbf{Q}} \right) \\ \tilde{\Delta}_{ij} &= \frac{n_{ij}}{2p(1 + \Delta_i)} \\ \mathbf{J} &= \begin{bmatrix} \mathbf{1}_{n_{11}} & 0 & 0 & 0 \\ 0 & \mathbf{1}_{n_{12}} & 0 & 0 \\ 0 & 0 & \mathbf{1}_{n_{21}} & 0 \\ 0 & 0 & 0 & \mathbf{1}_{n_{22}} \end{bmatrix}.\end{aligned}$$

To go further, using Woodbury's matrix identity, show that

$$\bar{\mathbf{Q}} = \bar{\mathbf{Q}}_0 - \bar{\mathbf{Q}}_0 \bar{\mathbf{M}} \left(\mathbf{I}_{2p} + \bar{\mathbf{M}}^\top \bar{\mathbf{Q}} \bar{\mathbf{M}} \right)^{-1} \bar{\mathbf{M}}^\top \bar{\mathbf{Q}}$$

where $\bar{\mathbf{Q}}_0 = \left[(\mathbf{D}_\gamma + \lambda \mathbf{1}_2 \mathbf{1}_2^\top)^{\frac{1}{2}} \mathbf{D}_{\tilde{\Delta}} (\mathbf{D}_\gamma + \lambda \mathbf{1}_2 \mathbf{1}_2^\top)^{\frac{1}{2}} + \mathbf{I}_2 \right]^{-1} \otimes \mathbf{I}_p$ and $\bar{\mathbf{M}} = A^{\frac{1}{2}} \mathbf{M} \mathbf{D}_{\tilde{\Delta}}^{\frac{1}{2}}$
with $\tilde{\Delta} = [\tilde{\Delta}_{11}, \dots, \tilde{\Delta}_{22}]$ and $\tilde{\Delta} = [\tilde{\Delta}_{11} + \tilde{\Delta}_{12}, \tilde{\Delta}_{21} + \tilde{\Delta}_{22}]$.

Finally conclude by deducing the expression of the statistical mean m_{ij} of $g_i(\hat{\mathbf{x}})$ and simulate to verify. To this end, show that

$$m_{ij} = \mathbf{e}_{ij}^\top \mathbf{D}_{\tilde{\Delta}}^{-\frac{1}{2}} \left[\mathbf{I}_4 - \left(\mathbf{I}_4 + \bar{\mathbf{M}}^\top \bar{\mathbf{Q}}_0 \bar{\mathbf{M}} \right)^{-1} \right] \mathbf{D}_{\tilde{\Delta}}^{\frac{1}{2}} \tilde{\mathbf{y}} + b_i$$

where $\tilde{\mathbf{y}} = [\tilde{y}_{11} - b_1, \dots, \tilde{y}_{22} - b_2]^\top$ in the compacted expression $\mathbf{y} = [\tilde{y}_{11} \mathbf{1}_{n_{11}}^\top, \dots, \tilde{y}_{22} \mathbf{1}_{n_{22}}^\top]^\top$
and $\mathbf{e}_{ij}^{[4]} \in \mathbb{R}^4$ is the canonical vector with entry 1 in position $2(i-1) + j$.

Chapter 5

Large Neural Networks

5.1 Random neural networks

Neural networks, and particularly today's popular deep neural networks, are extremely challenging to study. Even in a large dimensional regime, several technical boundaries are to this day seemingly unbreakable. The most important among these is the highly non-convex nature of their underlying optimization framework. While Chapter 6 later shows that the asymptotic performance analyses accessible to the random matrix framework is not limited to algorithms assuming explicit output functionals (such as linear regressions) and that some implicit (but convex) optimization schemes can be studied in the limit, neural network learning, which involves highly non-convex optimization, is still out of reach.

Although much less popular than modern deep neural networks, neural networks with random fixed weights are simpler to analyze. These have frequently arisen in the past decades as an appropriate solution to handle the possibly restricted number of training data, to reduce the computational and memory complexity and, from another viewpoint, can be seen as efficient *random feature extractors*. These neural networks in fact find their roots in Rosenblatt's perceptron [Rosenblatt, 1958] and have then been many times revisited, rediscovered, and analyzed in a number of works, both in their feedforward [Schmidt et al., 1992] and recurrent [Gelenbe, 1993] versions. The simplest modern versions of these random networks are the so-called extreme learning machine [Huang et al., 2012] for the feedforward case, which one may see as a mere linear regression method on random nonlinear features, and the echo state network [Jaeger, 2001] for the recurrent case, see [Scardapane and Wang, 2017] for a more exhaustive overview of randomness in neural networks.

It is also to be noted that deep neural networks are initialized at random and that random operations (such as random node deletions or voluntarily not-learning a large proportion of randomly-initialized neural network weights) are common and efficient in neural network learning [Srivastava et al., 2014, Fran-

kle and Carbin, 2019]. We may also point the recent endeavor towards neural network “learning without backpropagation” which, inspired by biological neural networks (that do not operate backpropagation learning), proposes learning mechanisms with fixed random *backward* weights and asymmetric *forward* learning procedure [Lillicrap et al., 2016, Nøkland, 2016, Baldi et al., 2018, Frenkel et al., 2019, Han et al., 2019]. As such, the study of random neural network structures may be instrumental to future improved understanding of advanced neural network procedures.

As shall be seen subsequently, the simple models of random neural networks are to a large extent connected to *kernel random matrices*. More specifically, the classification or regression performance at the output of these random neural networks are functionals of random matrices that fall into the wide class of kernel random matrices, yet of a slightly different form from those studied in Section 4. Perhaps more surprisingly, this connection still exists for *deep neural networks* that are (i) randomly initialized and (ii) then trained with gradient descent, via the so-called *neural tangent kernel* [Jacot et al., 2018], by considering the “infinite-neurons” limit, that is, the limit where the network widths of all layers go to infinity simultaneously. This close connection between neural networks and kernels has triggered a renewed interest for the theoretical investigation of deep neural networks from various perspectives including optimization [Du et al., 2019, Chizat et al., 2019], generalization [Allen-Zhu et al., 2019, Arora et al., 2019a, Bietti and Mairal, 2019], and learning dynamics [Lee et al., 2020, Advani et al., 2020]. These works shed new light on our theoretical understanding of deep neural network models and specifically demonstrate the significance of studying simple networks with random weights and their associated kernels to assess the intrinsic mechanisms of more elaborate and practical deep networks.

5.1.1 Regression with random neural networks

Throughout this section, we consider a feedforward single-hidden-layer neural network, as illustrated in Figure 5.1 (for convenience from right to left). Another important class of neural network models, the (random) recurrent network, will be discussed later in Section 5.3.




Figure 5.1: Illustration of a single-hidden-layer neural network (from right to left).

For input data $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$, we denote $\Sigma \equiv \sigma(\mathbf{W}\mathbf{X}) \in \mathbb{R}^{N \times n}$ the output of the first layer comprising N neurons. This output arises from the pre-multiplication of \mathbf{X} by some random weight matrix $\mathbf{W} \in \mathbb{R}^{N \times p}$ with i.i.d. (say standard Gaussian) entries and the *entry-wise* application of the nonlinear activation function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$. As such, the columns $\sigma(\mathbf{W}\mathbf{x}_i)$ of Σ can be seen as random nonlinear *features* of \mathbf{x}_i . The second layer weight $\beta \in \mathbb{R}^{N \times d}$ is then learned to adapt the feature matrix Σ to the associated target $\mathbf{Y} = [\mathbf{y}_1, \dots, \mathbf{y}_n] \in \mathbb{R}^{d \times n}$, for instance by minimizing the Frobenius norm $\|\mathbf{Y} - \beta^\top \Sigma\|_F^2$.

Remark 5.1 (Random neural networks, random feature maps and random kernels). *The columns of Σ may be seen as the output of the $\mathbb{R}^p \rightarrow \mathbb{R}^N$ random feature map $\phi : \mathbf{x}_i \mapsto \sigma(\mathbf{W}\mathbf{x}_i)$ for some given matrix \mathbf{W} . In [Rahimi and Recht, 2008], it is shown that, for every nonnegative definite “shift-invariant” kernel of the form $(\mathbf{x}, \mathbf{y}) \mapsto f(\|\mathbf{x} - \mathbf{y}\|^2)$, there exist appropriate choices for σ and the law of the entries of \mathbf{W} so that, as $N \rightarrow \infty$,*

$$\sigma(\mathbf{W}\mathbf{x}_i)^\top \sigma(\mathbf{W}\mathbf{x}_j) \xrightarrow{a.s.} f(\|\mathbf{x}_i - \mathbf{x}_j\|^2).$$

As such, for large enough N (that in general must scale with n, p), the bivariate function $(\mathbf{x}, \mathbf{y}) \mapsto \sigma(\mathbf{W}\mathbf{x})^\top \sigma(\mathbf{W}\mathbf{y})$ approximates a kernel function of the type $f(\|\mathbf{x} - \mathbf{y}\|^2)$ studied in the previous chapter. This result is then generalized, in subsequent works, to a larger family of kernels including inner-product kernels [Kar and Karnick, 2012], additive homogeneous kernels [Vedaldi and Zisserman, 2012], just to name a few. Another, possibly more marginal, connection with the previous sections is that $\sigma(\mathbf{w}^\top \mathbf{x})$ can be interpreted as a “properly scaling” inner-product kernel function applied to the “data” pair $\mathbf{w}, \mathbf{x} \in \mathbb{R}^p$. This technically induces another strong relation between the study of kernels and that of neural networks.

If the network weight β is designed to minimize the regularized mean squared error (MSE) $L(\beta) = \frac{1}{n} \sum_{i=1}^n \|\mathbf{y}_i - \beta^\top \sigma(\mathbf{W}\mathbf{x}_i)\|^2 + \gamma \|\beta\|_F^2$, for some regularization factor $\gamma > 0$, we obtain the explicit *ridge-regressor*¹

$$\beta \equiv \frac{1}{n} \Sigma \left(\frac{1}{n} \Sigma^\top \Sigma + \gamma \mathbf{I}_n \right)^{-1} \mathbf{Y}^\top \quad (5.1)$$

which follows from differentiating $L(\beta)$ with respect to β to obtain $0 = \gamma\beta + \frac{1}{n} \Sigma (\Sigma^\top \beta - \mathbf{Y}^\top)$ so that $(\frac{1}{n} \Sigma \Sigma^\top + \gamma \mathbf{I}_N) \beta = \frac{1}{n} \Sigma \mathbf{Y}^\top$ which, along with $(\frac{1}{n} \Sigma \Sigma^\top + \gamma \mathbf{I}_N)^{-1} \Sigma = \Sigma (\frac{1}{n} \Sigma^\top \Sigma + \gamma \mathbf{I}_n)^{-1}$ for $\gamma > 0$, gives the result.

The single-hidden-layer random neural net model presented above, with fixed random first layer and second layer performing a ridge regression, is sometimes referred to as an “extreme learning machine” in the literature [Huang et al., 2012], but from Remark 5.1 this is merely a random feature-based ridge regression.

¹Here we use a lowercase β to illustrate the fact that $\beta \in \mathbb{R}^{N \times d}$ is a “tall” matrix with $N \gg d$ (often $d = 1$), and acts like a vector from a random matrix point of view.

Note that, for β defined in (5.1), the training MSE (on the given training set (\mathbf{X}, \mathbf{Y})) reads

$$E_{\text{train}} = \frac{1}{n} \|\mathbf{Y}^\top - \Sigma^\top \beta\|_F^2 = \frac{\gamma^2}{n} \text{tr } \mathbf{Y} \mathbf{Q}^2(\gamma) \mathbf{Y}^\top, \quad \mathbf{Q}(\gamma) \equiv \left(\frac{1}{n} \Sigma^\top \Sigma + \gamma \mathbf{I}_n \right)^{-1} \quad (5.2)$$

for $\mathbf{Q}(\gamma)$ the resolvent of $\frac{1}{n} \Sigma^\top \Sigma$. Similarly, the test MSE on the test set $(\hat{\mathbf{X}}, \hat{\mathbf{Y}}) \in \mathbb{R}^{p \times \hat{n}} \times \mathbb{R}^{d \times \hat{n}}$ of size \hat{n} is given by

$$E_{\text{test}} = \frac{1}{\hat{n}} \|\hat{\mathbf{Y}}^\top - \hat{\Sigma}^\top \beta\|_F^2, \quad \hat{\Sigma} = \sigma(\mathbf{W} \hat{\mathbf{X}}) \quad (5.3)$$

with $\beta \in \mathbb{R}^{N \times d}$ the same as used in (5.2) which only depends on \mathbf{W} , the training set (\mathbf{X}, \mathbf{Y}) and γ .

The objective of this section is to understand the asymptotic behavior of the training and test MSE, in the large dimensional limit where $n, p, N \rightarrow \infty$ at the same rate, and how they depend on the law of (the entries of) \mathbf{W} , the activation function $\sigma(\cdot)$, the regularization penalty γ , as well as the training and test data (\mathbf{X}, \mathbf{Y}) and $(\hat{\mathbf{X}}, \hat{\mathbf{Y}})$.

5.1.1.1 Intuition and main results

Consider first the training error E_{train} defined in (5.2). Since

$$\text{tr } \mathbf{Y} \mathbf{Q}^2(\gamma) \mathbf{Y}^\top = -\frac{\partial}{\partial \gamma} \text{tr } \mathbf{Y} \mathbf{Q}(\gamma) \mathbf{Y}^\top, \quad (5.4)$$

a deterministic equivalent for the resolvent $\mathbf{Q}(\gamma)$ is sufficient to access the asymptotic behavior of E_{train} .

With a linear activation $\sigma(t) = t$, the resolvent of interest

$$\mathbf{Q}(\gamma) = \left(\frac{1}{n} \sigma(\mathbf{W} \mathbf{X})^\top \sigma(\mathbf{W} \mathbf{X}) + \gamma \mathbf{I}_n \right)^{-1} \quad (5.5)$$

is the same as in Theorem 2.6. In a sense, the evaluation of E_{train} calls for an extension of Theorem 2.6 to handle the case of nonlinear activations. Recall now that the main ingredients to derive a deterministic equivalent for (the linear case) $\mathbf{Q} = (\mathbf{X}^\top \mathbf{W}^\top \mathbf{W} \mathbf{X}/n + \gamma \mathbf{I}_n)^{-1}$ are (i) $\mathbf{X}^\top \mathbf{W}^\top$ has i.i.d. columns, and (ii) its i -th column $[\mathbf{W}^\top]_i$ has i.i.d. entries so that the key Lemma 2.11 applies. These hold due to the i.i.d. property of the entries of \mathbf{W} .

However, while for (i) the nonlinear $\Sigma^\top = \sigma(\mathbf{W} \mathbf{X})^\top$ still has i.i.d. columns, for (ii) its i -th column $\sigma([\mathbf{X}^\top \mathbf{W}^\top]_i)$ no longer has i.i.d. entries. Therefore, the main technical difficulty here is to obtain a nonlinear version of the trace lemma, Lemma 2.11. That is, we expect that the concentration of quadratic forms around their expectation remains valid despite the application of the entry-wise nonlinear σ . This naturally falls into the concentration of measure theory discussed in Section 2.7 and is given by the following lemma.

Lemma 5.1 (Concentration of nonlinear quadratic form, [Louart et al., 2018, Lemma 1]). For $\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, 1-Lipschitz $\sigma(\cdot)$, and $\mathbf{A} \in \mathbb{R}^{n \times n}, \mathbf{X} \in \mathbb{R}^{p \times n}$ such that $\|\mathbf{A}\| \leq 1$ and $\|\mathbf{X}\|$ bounded with respect to p, n , then

$$\mathbb{P} \left(\left| \frac{1}{n} \sigma(\mathbf{w}^\top \mathbf{X}) \mathbf{A} \sigma(\mathbf{X}^\top \mathbf{w}) - \frac{1}{n} \operatorname{tr} \mathbf{A} \mathbf{K} \right| > t \right) \leq C e^{-cn \min(t, t^2)}$$

for some $C, c > 0$ with

$$\mathbf{K} \equiv \mathbf{K}_{\mathbf{XX}} \equiv \mathbb{E}_{\mathbf{w}}[\sigma(\mathbf{X}^\top \mathbf{w}) \sigma(\mathbf{w}^\top \mathbf{X})] \in \mathbb{R}^{n \times n}. \quad (5.6)$$

In particular, for p, n large together, $\frac{1}{n} \sigma(\mathbf{w}^\top \mathbf{X}) \mathbf{A} \sigma(\mathbf{X}^\top \mathbf{w}) = \frac{1}{n} \operatorname{tr} \mathbf{A} \mathbf{K} + O(n^{-1/2})$ as in the linear case: the convergence rate of the linear case is thus not affected by 1-Lipschitz σ functions.

Equation (5.1) is the core ingredient to generalize Theorem 2.6 to the nonlinear setting, leading to the following result.

Theorem 5.1 (Nonlinear Gram matrix, [Louart et al., 2018]). Let $\mathbf{W} \in \mathbb{R}^{N \times p}$ be a random matrix with i.i.d. standard Gaussian entries, $\sigma(\cdot)$ be 1-Lipschitz continuous, and $\mathbf{X} \in \mathbb{R}^{p \times n}$ be of bounded operator norm (i.e., $\limsup_{n,p} \|\mathbf{X}\| < \infty$). Then, as $n, p, N \rightarrow \infty$ with p/n and N/n bounded away from zero and infinity, for $\mathbf{Q} = (\sigma(\mathbf{X}^\top \mathbf{W}^\top) \sigma(\mathbf{W} \mathbf{X})/n + \gamma \mathbf{I}_n)^{-1}$ with $\gamma > 0$,

$$\mathbf{Q} \leftrightarrow \bar{\mathbf{Q}} = \left(\frac{N}{n} \frac{\mathbf{K}}{1 + \delta} + \gamma \mathbf{I}_n \right)^{-1}$$

for δ the unique positive solution to $\delta = \frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \mathbf{K}$ and \mathbf{K} defined in (5.6).

As a direct consequence of Theorem 5.1, we have the following results on the asymptotic training and test mean square errors of a single-hidden-layer random neural network model in Figure 5.1. We refer the interested readers to [Louart et al., 2018] for the detailed proof and more discussions therein.

Corollary 5.1 (Asymptotic training and test MSEs, [Louart et al., 2018]). Under the setting and notations of Theorem 5.1, for training and test mean square errors defined in (5.2) and (5.3), respectively, as $n, p, N \rightarrow \infty$,

$$E_{\text{train}} - \bar{E}_{\text{train}} \xrightarrow{a.s.} 0, \quad E_{\text{test}} - \bar{E}_{\text{test}} \xrightarrow{a.s.} 0$$

with

$$\begin{aligned} \bar{E}_{\text{train}} &= \frac{\gamma^2}{n} \operatorname{tr} \mathbf{Y} \bar{\mathbf{Q}} \left(\frac{\frac{1}{N} \operatorname{tr} \bar{\mathbf{Q}} \mathbf{K} \bar{\mathbf{Q}}}{1 - \frac{1}{N} \operatorname{tr} \bar{\mathbf{Q}} \mathbf{K} \bar{\mathbf{Q}}} \bar{\mathbf{K}} + \mathbf{I}_n \right) \bar{\mathbf{Q}} \mathbf{Y}^\top \\ \bar{E}_{\text{test}} &= \frac{1}{\hat{n}} \|\hat{\mathbf{Y}}^\top - \bar{\mathbf{K}}_{\mathbf{XX}}^\top \bar{\mathbf{Q}} \mathbf{Y}^\top\|_F^2 \\ &\quad + \frac{\frac{1}{N} \operatorname{tr} \mathbf{Y} \bar{\mathbf{Q}} \bar{\mathbf{K}} \bar{\mathbf{Q}} \mathbf{Y}^\top}{1 - \frac{1}{N} \operatorname{tr} \bar{\mathbf{Q}} \mathbf{K} \bar{\mathbf{Q}}} \left(\frac{1}{\hat{n}} \operatorname{tr} \bar{\mathbf{K}}_{\mathbf{XX}} - \frac{1}{\hat{n}} \operatorname{tr} (\mathbf{I}_n + \gamma \bar{\mathbf{Q}}) (\bar{\mathbf{K}}_{\mathbf{XX}} \bar{\mathbf{K}}_{\mathbf{XX}}^\top \bar{\mathbf{Q}}) \right) \end{aligned}$$

where, similarly to $\mathbf{K} = \mathbb{E}[\sigma(\mathbf{X}^\top \mathbf{w})\sigma(\mathbf{w}^\top \mathbf{X})]$ in (5.6), we denoted

$$\mathbf{K}_{\mathbf{X}\hat{\mathbf{X}}} \equiv \mathbb{E}[\sigma(\mathbf{X}^\top \mathbf{w})\sigma(\mathbf{w}^\top \hat{\mathbf{X}})], \quad \mathbf{K}_{\hat{\mathbf{X}}\hat{\mathbf{X}}} \equiv \mathbb{E}[\sigma(\hat{\mathbf{X}}^\top \mathbf{w})\sigma(\mathbf{w}^\top \hat{\mathbf{X}})] \quad (5.7)$$

and

$$\bar{\mathbf{K}} \equiv \frac{N}{n} \frac{\mathbf{K}}{1+\delta}, \quad \bar{\mathbf{K}}_{\mathbf{X}\hat{\mathbf{X}}} \equiv \frac{N}{n} \frac{\mathbf{K}_{\mathbf{X}\hat{\mathbf{X}}}}{1+\delta}, \quad \bar{\mathbf{K}}_{\hat{\mathbf{X}}\hat{\mathbf{X}}} \equiv \frac{N}{n} \frac{\mathbf{K}_{\hat{\mathbf{X}}\hat{\mathbf{X}}}}{1+\delta}.$$

The proof of Corollary 5.1 is based on a higher-order deterministic equivalent of the type $\mathbf{Q}\mathbf{A}\mathbf{Q}$ for some deterministic or structured random matrices \mathbf{A} , as in Exercise 13. More precisely, for the training MSE E_{train} , we have from (5.4) that it suffices to derive a deterministic equivalent for $\mathbf{Q}^2(\gamma)$ (or, alternatively, by considering the derivative with respect to γ). For the test MSE E_{test} , we deduce from (5.3) that

$$E_{\text{test}} = \frac{1}{\hat{n}} \text{tr } \hat{\mathbf{Y}} \hat{\mathbf{Y}}^\top - \frac{2}{n\hat{n}} \text{tr } \mathbf{Y} \mathbf{Q} \Sigma^\top \hat{\Sigma} \hat{\mathbf{Y}}^\top + \frac{1}{n^2 \hat{n}} \text{tr } \mathbf{Y} \mathbf{Q} \Sigma^\top \hat{\Sigma} \hat{\Sigma}^\top \Sigma \mathbf{Q} \mathbf{Y}^\top \quad (5.8)$$

which involves the assessment of, for instance, the more involved expectation $\mathbb{E}_\mathbf{W}[\mathbf{Q} \Sigma^\top \hat{\Sigma} \hat{\Sigma}^\top \Sigma \mathbf{Q}]$ with $\hat{\Sigma} = \sigma(\mathbf{W} \hat{\mathbf{X}})$.

Clearly, to evaluate the asymptotic training and test errors in Corollary 5.1, the computation of the expectation of the form \mathbf{K} in (5.6) is needed. To go further, \mathbf{W} being standard Gaussian, the (i, j) entry of matrix \mathbf{K} writes

$$\begin{aligned} \mathbf{K}_{ij} &\equiv \kappa(\mathbf{x}_i, \mathbf{x}_j) \equiv \mathbb{E}[\sigma(\mathbf{w}^\top \mathbf{x}_i)\sigma(\mathbf{w}^\top \mathbf{x}_j)] \\ &= (2\pi)^{-\frac{p}{2}} \int \sigma(\mathbf{w}^\top \mathbf{x}_i)\sigma(\mathbf{w}^\top \mathbf{x}_j)e^{-\frac{1}{2}\|\mathbf{w}\|^2} d\mathbf{w} \end{aligned}$$

and indeed defines the limiting kernel of the random feature map $\mathbf{x}_i \mapsto \sigma(\mathbf{W}\mathbf{x}_i)$ as discussed in Remark 5.1. For a set of commonly used activation functions σ (that are not necessarily Lipschitz), the corresponding kernel matrix \mathbf{K} can be computed explicitly via an integration projection trick (see e.g., [Williams, 1997] for similar calculus). These results are provided in Table 5.1.

Given some data \mathbf{X} , Table 5.1 allows one to compute the “limiting” kernel \mathbf{K} for the listed activation functions $\sigma(\cdot)$. Then, by iterating the fixed-point equation in Theorem 5.1, one obtains the *effective kernel* $\bar{\mathbf{K}} \equiv \frac{N}{n} \frac{\mathbf{K}}{1+\delta}$ in the more practical large n, p, N setting (compared to the $N \rightarrow \infty$ alone limiting kernel \mathbf{K}). In this sense, Theorem 5.1 characterizes the impact of the effective kernel $\bar{\mathbf{K}}$ on the regression performance as detailed in Corollary 5.1.

5.1.1.2 Implications for learning with large neural networks

To validate the asymptotic analysis in Theorem 5.1 and Corollary 5.1 on real-world finite-dimensional data, Figure 5.2 and 5.3 compare the empirical MSEs with their limiting behaviors predicted in Corollary 5.1, for a random network having $N = 512$ neurons, with various types of Lipschitz and non-Lipschitz activations $\sigma(\cdot)$, respectively. The regressor $\beta \in \mathbb{R}^p$ maps the vectorized images from the Fashion-MNIST dataset (class 1 and 2) to their corresponding

unidimensional ($d = 1$) output labels $\mathbf{Y}_{1i}, \hat{\mathbf{Y}}_{1j} \in \{\pm 1\}$. For n, p, N only of order a few hundreds, a close match between theory and practice is observed for Lipschitz activations in Figure 5.2. The precision is less accurate but still not poor for the case of non-Lipschitz activations (Figure 5.3) which are not supported by the theorem statement – here for $\sigma(t) = 1 - t^2/2$, $\sigma(t) = 1_{t>0}$ and $\sigma(t) = \text{sign}(t)$. For all activations, the deviation from theory is more acute for small values of γ .

$\sigma(t)$	$\kappa(\mathbf{x}, \mathbf{y})$
t	$\mathbf{x}^\top \mathbf{y}$
$ t $	$\frac{2}{\pi} \ \mathbf{x}\ \ \mathbf{y}\ (\angle \cdot \arcsin(\angle) + \sqrt{1 - \angle^2})$
$\text{ReLU}(t) \equiv \max(t, 0)$	$\frac{1}{2\pi} \ \mathbf{x}\ \ \mathbf{y}\ (\angle \arccos(-\angle) + \sqrt{1 - \angle^2})$
$a_+ \max(t, 0) + a_- \max(-t, 0)$	$\frac{1}{2} (a_+^2 + a_-^2) \mathbf{x}^\top \mathbf{y} + \frac{1}{2\pi} \ \mathbf{x}\ \ \mathbf{y}\ (a_+ + a_-)^2 (-\angle \arccos(\angle) + \sqrt{1 - \angle^2})$
$a_2 t^2 + a_1 t + a_0$	$a_2^2 (2(\mathbf{x}^\top \mathbf{y})^2 + \ \mathbf{x}\ ^2 \ \mathbf{y}\ ^2) + a_1^2 \mathbf{x}^\top \mathbf{y} + a_2 a_0 (\ \mathbf{x}\ ^2 + \ \mathbf{y}\ ^2) + a_0^2$
$\text{erf}(t)$	$\frac{2}{\pi} \arcsin \left(\frac{2\mathbf{x}^\top \mathbf{y}}{\sqrt{(1+2\ \mathbf{x}\ ^2)(1+2\ \mathbf{y}\ ^2)}} \right)$
$1_{t>0}$	$\frac{1}{2} - \frac{1}{2\pi} \arccos(\angle)$
$\text{sign}(t)$	$\frac{2}{\pi} \arcsin(\angle)$
$\cos(t)$	$\exp \left(-\frac{1}{2} (\ \mathbf{x}\ ^2 + \ \mathbf{y}\ ^2) \right) \cosh(\mathbf{x}^\top \mathbf{y})$
$\sin(t)$	$\exp \left(-\frac{1}{2} (\ \mathbf{x}\ ^2 + \ \mathbf{y}\ ^2) \right) \sinh(\mathbf{x}^\top \mathbf{y})$
$\exp(-t^2/2)$	$\frac{1}{\sqrt{(1+\ \mathbf{x}\ ^2)(1+\ \mathbf{y}\ ^2)-(x^\top y)^2}}$

Table 5.1: Limiting kernel $\kappa(\mathbf{x}, \mathbf{y})$ for standard Gaussian \mathbf{W} , with $\angle \equiv \frac{\mathbf{x}^\top \mathbf{y}}{\|\mathbf{x}\| \|\mathbf{y}\|}$.




Figure 5.2: Neural network regression errors for Lipschitz $\sigma(\cdot)$ as a function of γ , $\sigma(t) = t$ in red, $\sigma(t) = \text{erf}(t)$ in green, and $\sigma(t) = \text{ReLU}(t)$ in blue, for 2-class Fashion-MNIST data (class 1 versus 2), $N = 512$, $n = 1024$, $\hat{n} = 512$, $p = 784$. Results averaged over 30 runs. [Link to code]




Figure 5.3: Neural network regression errors for non-Lipschitz $\sigma(\cdot)$ as a function of γ , $\sigma(t) = 1_{t>0}$ in red, quadratic $\sigma(t)$ in green, and $\sigma(t) = \text{sign}(t)$ in blue, for 2-class Fashion-MNIST data (class 1 versus 2), $N = 512$, $n = 1024$, $\hat{n} = 512$, $p = 784$. Results averaged over 30 runs. [Link to code]

Figures 5.2 and 5.3 confirm that, while the training error is a monotonically increasing function of the regularization parameter γ , there always exists an optimal value for γ which minimizes the test error. In particular, the theoretical formulas derived in Corollary 5.1 allow for a (data-dependent) fast offline tuning of the hyperparameter γ of the network, in the setting when n, p, N are not too small and comparable. In terms of activation functions (those listed here), we observe that, on the Fashion-MNIST dataset, the ReLU nonlinearity $\sigma(t) =$

$\max(t, 0)$ is optimal and achieves the minimum test error, while the quadratic activation $\sigma(t) = 1 - t^2/2$ is the worst and produces much higher training and test errors compared to others. This observation can be theoretically explained with a deeper analysis of the underlying kernel matrix \mathbf{K} , as performed in Section 5.1.2. Lastly, although not immediate at first sight, the training and test error curves of $\sigma(t) = 1_{t>0}$ and $\sigma(t) = \text{sign}(t)$ are indeed the same, up to a shift in γ , as a consequence of the relation $\text{sign}(t) = 2 \cdot 1_{t>0} - 1$.

Model complexity and the double descent phenomenon. The deterministic equivalent of the limiting performance provided in Corollary 5.1 explicitly depends on the ratio feature-to-sample N/n (as well as, more implicitly through the data, on the size p of the data). The quantity N/n is of crucial significance from a machine learning perspective, as it characterizes the (relative) *model complexity* of the neural network model under investigation. For a training set of size n , the increase of the number of neurons N represents the growth of model complexity and, as a consequence, the increase of its capacity to fit the given training set.

According to the golden “bias-variance tradeoff” rule [Friedman et al., 2001], it is necessary to control the model complexity (N here) to achieve optimal *test* performance. Essentially, as the model size increases, the model tends to better fit the given training set, resulting in a smaller “bias”, while on the other hand, gradually overfits the training set and may perform poorly on an independent test set due to the large “variance”. To prevent overfitting, explicit regularization schemes such as Tikhonov-type regularization or early stopping are proposed to control the model capacity by avoiding too-small training errors.

It has thus been long believed that the optimal choice of model complexity should produce a small but nonzero training error, but the success of deep learning seems to contradict this conventional wisdom. Modern deep neural networks often have a huge number of parameters and are routinely trained to fit the training data almost perfectly, while still yielding remarkably good test performance in many cases [Zhang et al., 2016]. This particularly means that, in some scenarios, it is possible to have good or even optimal models that are much larger than intuitively needed (typically with $N \gg n$).

This counterintuitive phenomenon is empirically observed for various large-scale machine learning models, and more recently, extensively investigated from a theoretical standpoint [Belkin et al., 2019, Hastie et al., 2019, Mei and Montanari, 2019]. More precisely, it is observed that, as the model becomes larger, the test error decreases and then increases, following the traditional “bias versus variance” U-shaped curve, until the interpolation threshold where the model fits perfectly the training set and achieves zero training error, typically at $N = n$. Then, very surprisingly, in the over-parameterization $N > n$ regime, the test error starts to decrease again as N further grows, reaching an error that can (but not always) be even smaller than the optimal error in the under-parametrization $N < n$ regime.

This so-called “double-descent” phenomenon (due to its W-shaped curve)

is depicted in Figure 5.4 for the random neural network model in Figure 5.1, with $\gamma = 10^{-7}$ to mimic the unregularized case. Observe that, when $N = n$, while the training error vanishes, the test error blows up. But then, in the over-parameterization $N > n$ regime, the test error monotonically decreases as N further increases and reaches an even smaller error than the optimal error in the $N < n$ regime, at least in this particular setting.




Figure 5.4: Training and test MSEs of random neural network model as a function of the ratio N/n on Fashion-MNIST data (class 1 versus 2), with $p = 784$, $n = 1000$, $\sigma(t) = \text{ReLU}(t)$ and $\lambda = 10^{-7}$. Results averaged over 30 runs. [\[Link to code\]](#)

This unexpected double-descent behavior with a test error singularity at $N = n$ can readily be anticipated from Theorem 5.1 in the unregularized $\gamma = 0$ case. Depending on whether $N > n$ or $N < n$, we indeed have the following *phase transition* behavior:

1. in the over-parameterization $N > n$ regime, by taking $\gamma \rightarrow 0$ in Theorem 5.1, we obtain

$$\delta = \frac{1}{n} \text{tr } \mathbf{K} \left(\frac{N}{n} \frac{\mathbf{K}}{1 + \delta} \right)^{-1} = \frac{n}{N - n} \quad (5.9)$$

where we assume \mathbf{K} to be invertible, so that $\bar{\mathbf{Q}} \equiv (\frac{N}{n} \frac{\mathbf{K}}{1 + \delta} + \gamma \mathbf{I}_n)^{-1} = \frac{n}{N - n} \mathbf{K}^{-1}$ is well defined at $\gamma = 0$;

2. on the other hand, in the under-parameterization $N < n$ regime, δ diverges when $\gamma \rightarrow 0$, but we remark that

$$\gamma \delta = \frac{1}{n} \text{tr } \mathbf{K} \left(\frac{N}{n} \frac{\mathbf{K}}{\gamma + \gamma \delta} + \mathbf{I}_n \right)^{-1} \quad (5.10)$$

converges, as $\gamma \rightarrow 0$, to $\gamma \delta \rightarrow \theta = \frac{1}{n} \text{tr } \mathbf{K} (\frac{N}{n} \frac{\mathbf{K}}{\theta} + \mathbf{I}_n)^{-1}$; that is, $\delta, \bar{\mathbf{Q}}$ both scale like γ^{-1} . We have in particular $\mathbb{E}[\gamma \mathbf{Q}] \sim \gamma \bar{\mathbf{Q}} \sim (\frac{N}{n} \frac{\mathbf{K}}{\theta} + \mathbf{I}_n)^{-1}$. This

is in accordance with the fact that the Gram matrix $\Sigma^\top \Sigma \in \mathbb{R}^{n \times n}$ is of rank at most $\min(N, n)$ and is thus not invertible for $N < n$ (such that $\mathbf{Q} \equiv (\Sigma^\top \Sigma/n + \gamma \mathbf{I}_n)^{-1}$ scales as γ^{-1} as $\gamma \rightarrow 0$).

With this remark, the behavior of the asymptotic test error in Corollary 5.1 as N approaches n , from both sides of $N < n$ and $N > n$ is better understood. In particular, the denominator of \bar{E}_{test} reads

$$\begin{aligned} 1 - \frac{1}{N} \text{tr} \bar{\mathbf{K}} \bar{\mathbf{Q}} \bar{\mathbf{K}} \bar{\mathbf{Q}} \\ = \begin{cases} 1 - \frac{n}{N} + \frac{2\gamma}{N} \text{tr} \left(\frac{N}{n} \frac{\mathbf{K}}{1+\delta} + \gamma \mathbf{I}_n \right)^{-1} - \frac{\gamma^2}{N} \text{tr} \left(\frac{N}{n} \frac{\mathbf{K}}{1+\delta} + \gamma \mathbf{I}_n \right)^{-2}, & N > n \\ 1 - \frac{n}{N} + \frac{2}{N} \text{tr} \left(\frac{N}{n} \frac{\mathbf{K}}{\gamma+\theta} + \mathbf{I}_n \right)^{-1} - \frac{1}{N} \text{tr} \left(\frac{N}{n} \frac{\mathbf{K}}{\gamma+\theta} + \mathbf{I}_n \right)^{-2}, & N < n \end{cases} \end{aligned}$$

which, as N approaches n from either side, becomes $1 - \frac{1}{N} \text{tr} \bar{\mathbf{K}} \bar{\mathbf{Q}} \bar{\mathbf{K}} \bar{\mathbf{Q}} \sim \gamma$ (in the small γ regime).

The fact that this denominator scales like γ as $\gamma \rightarrow 0$ explains the major difference between the training and test error behavior in Figure 5.4. Due to the γ^2 prefactor in \bar{E}_{train} , the training error is guaranteed to be finite (in fact to be zero). But for the test error, if the numerator term $\frac{1}{n} \text{tr} \bar{\mathbf{K}}_{\hat{\mathbf{X}}\hat{\mathbf{X}}} - \frac{1}{n} \text{tr} (\mathbf{I}_n + \gamma \bar{\mathbf{Q}}) (\bar{\mathbf{K}}_{\mathbf{X}\hat{\mathbf{X}}} \bar{\mathbf{K}}_{\hat{\mathbf{X}}\hat{\mathbf{X}}}^\top \bar{\mathbf{Q}})$ does not scale like γ as $\gamma \rightarrow 0$, then \bar{E}_{test} diverges to infinity at $N = n$. A first counterexample is of course when $\hat{\mathbf{X}} = \mathbf{X}$, for which the numerator term of \bar{E}_{test} is now

$$\frac{1}{n} - \frac{1}{n} \text{tr} (\mathbf{I}_n + \gamma \bar{\mathbf{Q}}) (\bar{\mathbf{K}}_{\mathbf{X}\hat{\mathbf{X}}} \bar{\mathbf{K}}_{\hat{\mathbf{X}}\hat{\mathbf{X}}}^\top \bar{\mathbf{Q}}) = \frac{\gamma^2}{n} \text{tr} \bar{\mathbf{Q}} \bar{\mathbf{K}} \bar{\mathbf{Q}}$$

where we exploited the fact that $\bar{\mathbf{K}} \bar{\mathbf{Q}} = \mathbf{I}_n - \gamma \bar{\mathbf{Q}}$; of course, the test error \bar{E}_{test} coincides with the training error \bar{E}_{train} in this case. In general though, when $\hat{\mathbf{X}} \neq \mathbf{X}$, the numerator does not vanish with γ and the test error diverges at $N = n$ and $\gamma \rightarrow 0$.

Therefore, this double-descent singularity at $N = n$ only occurs: (i) in the unregularized case as $\gamma \rightarrow 0$ where some sort of invertibility issue arises, and (ii) when the test data $\hat{\mathbf{X}}$ is sufficiently “distinct” from the training data \mathbf{X} , in a kernel matrix sense, and this is fully *independent* of the targets \mathbf{Y} and $\hat{\mathbf{Y}}$.

5.1.2 Delving deeper into limiting kernels

To understand the behavior of different activation functions in Figure 5.2 and 5.3, we would like to build up a *data-dependent* theory that can provide the performance of different nonlinearities in a more explicit manner. According to our previous discussion, the performance depends on the nonlinear activation $\sigma(\cdot)$ only via the kernel matrix of the form

$$\mathbf{K} \equiv \mathbb{E}_{\mathbf{w}} [\sigma(\mathbf{X}^\top \mathbf{w}) \sigma(\mathbf{w}^\top \mathbf{X})] \quad (5.11)$$

given explicitly in Table 5.1. The entries of \mathbf{K} are nonlinear functions of $\|\mathbf{x}_i\|$, $\|\mathbf{x}_j\|$ or $\mathbf{x}_i^\top \mathbf{x}_j$, reminiscent of the distance and inner-product random kernel matrices studied in Section 4.2. Following the same idea, under the k -class

Gaussian mixture model for the data \mathbf{x}_i :

$$\mathbf{x}_i \in \mathcal{C}_a \Leftrightarrow \sqrt{p}\mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}_a, \mathbf{C}_a)$$

for $a \in \{1, \dots, k\}$ and the non-trivial classification condition

$$\begin{aligned} \mathbf{M} &= [\boldsymbol{\mu}_1^\circ, \dots, \boldsymbol{\mu}_k^\circ] = O_{\|\cdot\|}(1), \quad \boldsymbol{\mu}_\ell^\circ = \boldsymbol{\mu}_\ell - \sum_{a=1}^k \frac{n_a}{n} \boldsymbol{\mu}_a \\ \mathbf{t} &= [t_1, \dots, t_k]^\top = O_{\|\cdot\|}(1), \quad t_a = \frac{1}{\sqrt{p}} \text{tr}(\mathbf{C}_a - \mathbf{C}^\circ) \\ \mathbf{S} &= \{S_{ab}\}_{a,b=1}^k = O_{\|\cdot\|}(1), \quad S_{ab} = \frac{1}{p} \text{tr} \mathbf{C}_a \mathbf{C}_b. \end{aligned}$$

for $\mathbf{C}^\circ = \sum_{a=1}^k \frac{n_a}{n} \mathbf{C}_a$, we have, for $\mathbf{x}_i \in \mathcal{C}_a$ and $\mathbf{x}_j \in \mathcal{C}_b$, $i \neq j$ that

$$\mathbf{x}_i^\top \mathbf{x}_j = 0 + \frac{1}{p} \mathbf{z}_i^\top \mathbf{C}_a^{\frac{1}{2}} \mathbf{C}_b^{\frac{1}{2}} \mathbf{z}_j + \frac{1}{p} (\boldsymbol{\mu}_a^\top \mathbf{C}_b^{\frac{1}{2}} \mathbf{z}_j + \boldsymbol{\mu}_b^\top \mathbf{C}_a^{\frac{1}{2}} \mathbf{z}_i) + \frac{1}{p} \boldsymbol{\mu}_a^\top \boldsymbol{\mu}_b$$

and

$$\|\mathbf{x}_i\|^2 = \mathbf{x}_i^\top \mathbf{x}_i = \frac{1}{p} \text{tr} \mathbf{C}^\circ + \frac{1}{p} \text{tr}(\mathbf{C}_a - \mathbf{C}^\circ) + [\boldsymbol{\psi}]_i + \frac{1}{p} \|\boldsymbol{\mu}_a\|^2 + \frac{2}{p} \boldsymbol{\mu}_a^\top \mathbf{C}_a^{\frac{1}{2}} \mathbf{z}_i$$

with $\text{tr} \mathbf{C}^\circ / p = O(1)$ and $\boldsymbol{\psi} \in \mathbb{R}^n$ such that $[\boldsymbol{\psi}]_i = \mathbf{z}_i^\top \mathbf{C}_a \mathbf{z}_i / p - \text{tr} \mathbf{C}_a / p = O(p^{-1/2})$ as in Theorem 4.1. As a consequence, a Taylor expansion (around 0 or $-\text{tr} \mathbf{C}^\circ / p$) can be performed, as in Section 4.2, to asymptotically ‘‘linearize’’ these kernel matrices arising from different nonlinear activations in Table 5.1, leading to the following result, which extends Theorem 4.1.

Theorem 5.2 ([Liao and Couillet, 2018b]). *For all $\sigma(\cdot)$ in Table 5.1 and $\mathbf{P} = \mathbf{I}_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^\top$, we have, as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$ and $n_a/n \rightarrow c_a \in (0, 1)$ that*

$$\|\mathbf{PKP} - \mathbf{PK}\tilde{\mathbf{K}}\mathbf{P}\| \xrightarrow{a.s.} 0$$

where

$$\tilde{\mathbf{K}} = d_1 \cdot \frac{1}{p} (\mathbf{W} + \mathbf{MJ}^\top)^\top (\mathbf{W} + \mathbf{MJ}^\top) + d_2 \cdot \mathbf{UBU}^\top + d_0 \cdot \mathbf{I}_n$$

for $\mathbf{W} \equiv [\mathbf{W}_1, \dots, \mathbf{W}_k] \in \mathbb{R}^{p \times n}$, $\mathbf{W}_a \equiv \mathbf{C}_a^{\frac{1}{2}} \mathbf{Z}_a \in \mathbb{R}^{p \times n_a}$ and

$$\mathbf{U} = \begin{bmatrix} \mathbf{J} \\ \sqrt{p} \end{bmatrix} \in \mathbb{R}^{n \times (k+1)}, \quad \mathbf{B} = \begin{bmatrix} \mathbf{tt}^\top + 2\mathbf{S} & \mathbf{t} \\ \mathbf{t}^\top & 1 \end{bmatrix}, \quad \boldsymbol{\psi} \equiv \frac{1}{p} \{ \|\mathbf{w}_i\|^2 - \mathbb{E}[\|\mathbf{w}_i\|^2] \}_{i=1}^n$$

with the corresponding coefficients d_0 and d_1, d_2 given in Table 5.2.

Following the discussions in Remark 4.2 and 4.5, for simplicity of analysis, the result is presented here under the form of *centered* kernel matrix \mathbf{PKP} , that essentially performs a feature centering in the kernel space.

Theorem 5.2 tells us that, for the (non-trivial) classification of a mixture of k Gaussians, the (centered) kernel matrix \mathbf{K} depends on the nonlinear function $\sigma(\cdot)$ solely via *three* scalars d_0 , d_1 and d_2 , for all nonlinearities listed in Table 5.1. Moreover, note that the coefficient d_0 only introduces a constant shift of *all* eigenvalues of the kernel matrix \mathbf{K} , and has thus no impact on the performance of kernel-based algorithms such as kernel spectral clustering or kernel ridge regression. These two “universal” parameters (d_1, d_2) are reminiscent of the parameters α and β of the α - β kernel in Theorem 4.2 and the properly scaling kernel parameterized by (a_1, a_2) in Theorem 4.6. This is another evidence of large dimensional universality that will be discussed at length in Chapter 8.

Letting the term $d_0\mathbf{I}_n$ aside, the kernel matrix $\tilde{\mathbf{K}}$ has two “information-plus-noise” type components, $\mathbf{W} + \mathbf{M}\mathbf{J}^\top$ and $\mathbf{U} = [\mathbf{J}/\sqrt{p}, \boldsymbol{\psi}]$, weighted by d_1 and d_2 , respectively, where we recall the class structure information matrix $\mathbf{J} = [\mathbf{j}_1, \dots, \mathbf{j}_k] \in \mathbb{R}^{n \times k}$. It is therefore impossible to get rid of the noisy terms (\mathbf{W} and $\boldsymbol{\psi}$) by wisely choosing the function $\sigma(\cdot)$ without affecting \mathbf{J} . This is in sharp contrast to more general kernels (i.e., not arising from random neural networks) as in Corollary 4.1, which allow for a more flexible treatment of information versus noise.

Moreover, since the matrix of statistical means \mathbf{M} is multiplied by d_1 and covariance information \mathbf{t} and \mathbf{S} by d_2 , Theorem 5.2 provides practical instructions in a “data-dependent” choice of the nonlinearity. More precisely, the functions $\sigma(\cdot)$ in Table 5.2 can be divided into the following three groups

1. *mean-oriented*, where $d_1 \neq 0$ while $d_2 = 0$: this is the case of the functions $\sigma(t) = t$, $\text{sign}(t)$ and $\sin(t)$ and $\text{erf}(t)$, which asymptotically capture only the difference in means, with the information in covariance discarded;
2. *covariance-oriented*, where $d_1 = 0$ while $d_2 \neq 0$: this concerns the functions $\sigma(t) = |t|$, $\cos(t)$ and $\exp(-t^2/2)$ which only exploit the information in covariances (\mathbf{t} and \mathbf{S});
3. *balanced*, where both $d_1, d_2 \neq 0$: here for the ReLU $\sigma(t) = \max(t, 0)$, Leaky ReLU (L-ReLU) $a_+ \max(t, 0) + a_- \max(-t, 0)$ [Maas et al., 2013] and the quadratic function $a_2 t^2 + a_1 t + a_0$.

Note that, all mean-oriented functions are odd and all covariance-oriented functions are even. This remark is also reminiscent of a similar conclusion in the α - β kernel discussed in Section 4.3 as well as of the properly scaling inner-product kernel in Section 4.4 on a closely related, yet different, model.

$\sigma(t)$	d_1	d_2	d_0
t	1	0	0
$ t $	0	$\frac{1}{\pi\tau_p}$	$(\frac{1}{2} - \frac{1}{\pi})\tau_p$
$\text{ReLU}(t) \equiv \max(t, 0)$	$\frac{1}{4}$	$\frac{1}{4\pi\tau_p}$	$(\frac{1}{8} - \frac{1}{4\pi})\tau_p$
$\text{LReLU}(t) \equiv a_+ \max(t, 0) + a_- \max(-t, 0)$	$\frac{1}{4}(a_+ - a_-)^2$	$\frac{1}{4\pi\tau_p}(a_+ + a_-)^2$	$\frac{\pi-2}{8\pi}(a_+ + a_-)^2\tau_p$
$a_2 t^2 + a_1 t + a_0$	a_1^2	a_2^2	$\frac{1}{2}\tau_p^2 a_2^2$
$\text{erf}(t)$	$\frac{4}{\pi} \frac{1}{\tau_p + 1}$	0	$\frac{2}{\pi} \left(\arccos \frac{\tau_p}{\tau_p + 1} - \frac{\tau_p}{\tau_p + 1} \right)$
$1_{t>0}$	$\frac{1}{\pi\tau_p}$	0	$\frac{1}{4} - \frac{1}{2\pi}$
$\text{sign}(t)$	$\frac{4}{\pi\tau_p}$	0	$1 - \frac{2}{\pi}$
$\cos(t)$	0	$\frac{1}{4} \cdot e^{-\tau_p/2}$	$\frac{1}{2}(1 + e^{-\tau_p}) - e^{-\tau_p/2}$
$\sin(t)$	$e^{-\tau_p/2}$	0	$\frac{1}{2}(1 - e^{-\tau_p}) - \frac{\tau_p}{2}e^{-\tau_p/2}$
$\exp(-t^2/2)$	0	$\frac{2}{(\tau_p + 2)^3}$	$\frac{1}{\sqrt{\tau_p + 1}} - \frac{2}{\tau_p + 2}$

Table 5.2: Coefficients d_1, d_2 and d_0 for different $\sigma(\cdot)$, with $\tau_p = \frac{2}{p} \text{tr } \mathbf{C}^\circ$.

Also, similarly to the random kernel matrices discussed in Section 4.5.1, $\tilde{\mathbf{K}}$ in Theorem 5.2 follows a spiked random matrix model and contains the class structural information matrix \mathbf{J} . As a result, the top eigenvectors of the kernel matrix \mathbf{K} are expected to align to (the subspace spanned by the columns of) \mathbf{J} and can be used for spectral clustering.

To corroborate the above classification of different nonlinearities, we perform kernel spectral clustering with **PKP**, in Figure 5.5, on four classes of Gaussian data: $\mathcal{N}(\mu_1, \mathbf{C}_1)$, $\mathcal{N}(\mu_1, \mathbf{C}_2)$, $\mathcal{N}(\mu_2, \mathbf{C}_1)$ and $\mathcal{N}(\mu_2, \mathbf{C}_2)$ with the LReLU function $\text{LReLU}(t) \equiv a_+ \max(t, 0) + a_- \max(-t, 0)$ that takes different values for a_+ and a_- . For $b = 1, 2$, $\mu_b = [\mathbf{0}_{b-1}; 5; \mathbf{0}_{p-b}]$ and $\mathbf{C}_b = (1 + 15(b-1)/\sqrt{p})\mathbf{I}_p$. By choosing $a_+ = -a_- = 1$ (equivalent to $\sigma(t) = |t|$) and $a_+ = a_- = 1$ (equivalent to the linear function $\sigma(t) = t$), with the leading two eigenvectors we always recover two classes instead of four, as each setting of parameters only allows for a part of the statistical information (only mean or only covariance) of the data for clustering. However, by taking $a_+ = 1, a_- = 0$ (i.e., the ReLU function) we distinguish all four classes with the leading two eigenvectors, to which the k-means method can then be applied for a final classification, as shown in Figure 5.6.




Figure 5.5: Leading two eigenvectors of **PKP** for the LReLU function with $a_+ = -a_- = 1$ (top) and $a_+ = a_- = 1$ (bottom), performed on four classes Gaussian mixture data with $p = 512$, $n = 256$, $c_b = 1/4$ and $\mathbf{j}_b = [\mathbf{0}_{n_{b-1}}; \mathbf{1}_{n_b}; \mathbf{0}_{n-n_b}]$, for $b \in \{1, 2, 3, 4\}$. [Link to code]

Again, although derived from a simple k -class Gaussian mixture model, Theorem 5.2 establishes an unexpected close match in practice when applied to real-world datasets. To illustrate this claim, we consider two different types of classification tasks: one on the MNIST [LeCun et al., 1998] database (number 6 and 8), and the other on epileptic EEG time series data [Andrzejak et al., 2001]




Figure 5.6: Leading two eigenvectors of \mathbf{PKP} (top) for the LReLU function with $a_+ = 1$, $a_- = 0$ (equivalent to $\text{ReLU}(t)$) and two dimensional representation of these eigenvectors (bottom), in the same setting as in Figure 5.5. [Link to code]

(set B and E). These two datasets are typical examples of means-dominant (handwritten digits recognition) and covariances-dominant (EEG times series classification) tasks. This is numerically confirmed in Table 5.3.

Table 5.3: Empirical estimation of (normalized) differences in means and covariances of the MNIST and epileptic EEG datasets.

	$\ \mathbf{M}^\top \mathbf{M}\ $	$\ \mathbf{t}\mathbf{t}^\top + 2\mathbf{S}\ $
MNIST data	172.4	86.0
EEG data	1.2	182.7

Recall from (5.11) that the random feature Gram matrix $\frac{1}{N}\sigma(\mathbf{X}^\top \mathbf{W}^\top)\sigma(\mathbf{W}\mathbf{X}) = \frac{1}{N}\Sigma^\top \Sigma$ is an (empirical) estimate of the (expected) kernel matrix \mathbf{K} , and is therefore expected to behave similarly to \mathbf{K} for not-too-small N .

Here we perform random feature-based spectral clustering on data matrices \mathbf{X} that consist of $n = 32, 64$ and 128 randomly selected vectorized images of size $p = 784$ from the MNIST dataset. The k-means method is then applied to the leading two eigenvectors of the Gram matrix $\frac{1}{n}\Sigma^\top \Sigma$ that consists of $N = 32$ random features to perform unsupervised classification, with resulting accuracies (averaged over 50 runs) reported in Table 5.4. As seen from Table 5.3, the mean-oriented $\sigma(t)$ functions are expected to outperform the covariance-oriented functions in this task, which is consistent with the results in Table 5.4.

Table 5.4: Classification accuracies for random feature-based spectral clustering with different $\sigma(t)$ on the MNIST dataset.

	$\sigma(t)$	$n = 32$	$n = 64$	$n = 128$
mean-oriented	t	85.31%	88.94%	87.30%
	$1_{t>0}$	86.00%	82.94%	85.56%
	$\text{sign}(t)$	81.94%	83.34%	85.22%
	$\sin(t)$	85.31%	87.81%	87.50%
	$\text{erf}(t)$	86.50%	87.28%	86.59%
cov-oriented	$ t $	62.81%	60.41%	57.81%
	$\cos(t)$	62.50%	59.56%	57.72%
	$\exp(-t^2/2)$	64.00%	60.44%	58.67%
balanced	$\text{ReLU}(t)$	82.87%	85.72%	82.27%

For the epileptic EEG dataset [Andrzejak et al., 2001] that is more “covariance-dominant” per Table 5.4, random feature-based spectral clustering on $n = 32, 64$ and 128 randomly picked EEG segments of length $p = 100$ from the dataset. After k-means classification on the leading two eigenvectors of the (centered) Gram matrix composed of $N = 32$ random features, the accuracies (averaged over 50 runs) are reported in Table 5.5 ,where we observe that covariance-oriented functions (i.e., $\sigma(t) = |t|, \cos(t)$ and $\exp(-t^2/2)$) far outperform any other with almost perfect classification accuracies. It is particularly interesting to note that the popular ReLU function is suboptimal in both tasks, but never performs very badly, thereby offering a good risk-performance tradeoff.

Table 5.5: Classification accuracies for random feature-based spectral clustering with different $\sigma(t)$ on the epileptic EEG dataset.

	$\sigma(t)$	$n = 32$	$n = 64$	$n = 128$
mean-oriented	t	71.81%	70.31%	69.58%
	$1_{t>0}$	65.19%	65.87%	63.47%
	$\text{sign}(t)$	67.13%	64.63%	63.03%
	$\sin(t)$	71.94%	70.34%	68.22%
	$\text{erf}(t)$	69.44%	70.59%	67.70%
cov-oriented	$ t $	99.69%	99.69%	99.50%
	$\cos(t)$	99.00%	99.38%	99.36%
	$\exp(-t^2/2)$	99.81%	99.81%	99.77%
balanced	$\text{ReLU}(t)$	84.50%	87.91%	90.97%

The classification of “mean-oriented”, “covariance-oriented” and “balanced” nonlinearities for nonlinear Gram matrix $\frac{1}{n}\Sigma^\top\Sigma$ also helps explain the performance of different activation functions in neural network models in Figure 5.2 and 5.3. Since the MNIST data have more information in the statistical means, it is not surprising to observe in Figure 5.3 that both training

and test errors of the (covariance-oriented) quadratic activation $\sigma(t) = 1 - t^2/2$ are much higher than the others and the flexible ReLU function achieves the minimal training and test errors in Figure 5.2.

5.2 Gradient descent dynamics in learning linear neural nets

In Section 5.1 we considered the single-hidden-layer neural network model with random first layer, which, as pointed out in Remark 5.1, is closely connected to random feature maps and kernel methods. More precisely, the analyses in Theorem 5.1 and Corollary 5.1 provide access to the model performance (training and test MSEs) as a function of the nonlinear activation function $\sigma(\cdot)$, through the underlying kernel matrix \mathbf{K} .

Besides these *nonlinear* transformations discussed in Section 5.1, another salient feature of neural network models is that they are routinely trained with the (stochastic) gradient descent method. As a consequence, the network, and thus its performance, are functions of the training time (or the number of descent steps).

In this section, the gradient descent dynamics (GDDs) of learning a ridge regression model (i.e., a single-layer linear network) are considered. For a given training data matrix $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ with associated labels $\mathbf{y} = [y_1, \dots, y_n] \in \mathbb{R}^n$, a weight vector $\mathbf{w} \in \mathbb{R}^p$ is learned by minimizing the following (ridge-regularized) square loss

$$L(\mathbf{w}) = \frac{1}{2n} \|\mathbf{y} - \mathbf{X}^\top \mathbf{w}\|^2 + \frac{\gamma}{2} \|\mathbf{w}\|^2 \quad (5.12)$$

with some regularization penalty $\gamma \geq 0$. The gradient of L with respect to \mathbf{w} is given by $\nabla_{\mathbf{w}} L(\mathbf{w}) = -\frac{1}{n} \mathbf{X}(\mathbf{y} - \mathbf{X}^\top \mathbf{w}) + \gamma \mathbf{w}$ so that with small gradient descent step (or, learning rate) α , we obtain, by performing a continuous-time approximation, the following differential equation

$$\frac{d\mathbf{w}(t)}{dt} = -\alpha \nabla_{\mathbf{w}} L(\mathbf{w}) = \frac{\alpha}{n} \mathbf{X} \mathbf{y} - \alpha \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p \right) \mathbf{w}$$

the solution of which is explicitly given by

$$\mathbf{w}(t) = e^{-\alpha t \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p \right)} \mathbf{w}_0 + \left(\mathbf{I}_p - e^{-\alpha t \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p \right)} \right) \mathbf{w}_\infty \quad (5.13)$$

where we denote $\mathbf{w}_0 = \mathbf{w}(t=0)$ the initialization of gradient descent and

$$\mathbf{w}_\infty = \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p \right)^{-1} \frac{1}{n} \mathbf{X} \mathbf{y} \quad (5.14)$$

the ridge regression solution with regularization parameter γ . We recall that the exponential of a symmetric matrix \mathbf{A} is given by $e^{\mathbf{A}} = \sum_{k=0}^{\infty} \frac{1}{k!} \mathbf{A}^k = \mathbf{V} e^{\Lambda} \mathbf{V}^\top$, with $\mathbf{A} = \mathbf{V} \Lambda \mathbf{V}^\top$ the spectral decomposition of \mathbf{A} .

We consider the following symmetric binary Gaussian mixture model for the input data

$$\mathcal{C}_1 : \mathbf{x}_i \sim \mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p) \quad \mathcal{C}_2 : \mathbf{x}_i \sim \mathcal{N}(\boldsymbol{\mu}, \mathbf{I}_p)$$

with associated labels $y_i = -1$ and $y_i = 1$, respectively, and we assume as in previous sections the non-trivial setting where $\|\boldsymbol{\mu}\| = O(1)$.

For the linear classifier with $\mathbf{w}(t)$ given by (5.13), the GDDs can be studied by considering the training and test misclassification error rates as

$$\mathbb{P}(\mathbf{x}_i^\top \mathbf{w}(t) > 0 \mid y_i = -1), \quad \mathbb{P}(\hat{\mathbf{x}}^\top \mathbf{w}(t) > 0 \mid \hat{y} = -1)$$

for $\hat{\mathbf{x}} \sim \mathcal{N}(-\boldsymbol{\mu}, \mathbf{I}_p)$ a new test datum (that is independent of the training set (\mathbf{X}, \mathbf{y})), genuinely of label $\hat{y} = -1$.

To evaluate the test performance, since the test datum $\hat{\mathbf{x}}$ is independent of (\mathbf{X}, \mathbf{y}) (and thus of $\mathbf{w}(t)$), conditioned on $\mathbf{w}(t)$, $\mathbf{w}(t)^\top \hat{\mathbf{x}}$ is a Gaussian random variable of mean $\mathbf{w}(t)^\top \boldsymbol{\mu}$ and variance $\|\mathbf{w}(t)\|^2$. The misclassification probability can be retrieved with the Gaussian Q -function and we thus resort to the computation of $\mathbf{w}(t)^\top \boldsymbol{\mu}$ as well as $\mathbf{w}(t)^\top \mathbf{w}(t)$ to assess the classification performance.

For $\boldsymbol{\mu}^\top \mathbf{w}(t)$, with Cauchy's integration formula we have

$$\begin{aligned} \boldsymbol{\mu}^\top \mathbf{w}(t) &= \boldsymbol{\mu}^\top e^{-\alpha t(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p)} \mathbf{w}_0 + \boldsymbol{\mu}^\top \left(\mathbf{I}_p - e^{-\alpha t(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p)} \right) \mathbf{w}_\infty \\ &= -\frac{1}{2\pi i} \oint_{\Gamma} f_t(z) \boldsymbol{\mu}^\top \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p - z \mathbf{I}_p \right)^{-1} \mathbf{w}_0 dz \\ &\quad - \frac{1}{2\pi i} \oint_{\Gamma} \frac{1 - f_t(z)}{z} \boldsymbol{\mu}^\top \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p - z \mathbf{I}_p \right)^{-1} \frac{1}{n} \mathbf{X} \mathbf{y} dz \end{aligned}$$

with $f_t(z) \equiv \exp(-\alpha t z)$ and Γ a positive closed path circling around all the eigenvalues of the (shifted) sample covariance $\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p$.

Noting that $\mathbf{X} = \boldsymbol{\mu} \mathbf{y}^\top + \mathbf{Z}$ for \mathbf{Z} with i.i.d. standard Gaussian entries, we have, with Lemma 2.7 that

$$\left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \gamma \mathbf{I}_p - z \mathbf{I}_p \right)^{-1} = \mathbf{Q}(z) - \mathbf{Q}(z) \mathbf{U} \begin{bmatrix} \|\boldsymbol{\mu}\|^2 m(z) & 1 \\ 1 & -\frac{1}{1 + c m(z)} \end{bmatrix}^{-1} \mathbf{U}^\top \mathbf{Q}(z) + o(1)$$

for $\mathbf{U} = [\boldsymbol{\mu} \quad \frac{1}{n} \mathbf{Z} \mathbf{y}] \in \mathbb{R}^{p \times 2}$ and $\mathbf{Q}(z) \equiv (\frac{1}{n} \mathbf{Z} \mathbf{Z}^\top + \gamma \mathbf{I}_p - z \mathbf{I}_p)^{-1}$ admitting the deterministic equivalent relation (from Theorem 2.4) $\mathbf{Q}(z) \leftrightarrow m(z) \mathbf{I}_p$, with $m(z)$ the unique solution to

$$c(z - \gamma) m^2(z) - (1 - c - z + \gamma) m(z) + 1 = 0. \quad (5.15)$$

5.2.0.1 Main results

Based on the above deterministic equivalent, the following results on the asymptotic training and test performance can be derived.

Theorem 5.3 (Training and test performance of GDD, [Liao and Couillet, 2018a]). *For random initialization $\mathbf{w}_0 \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I}_p/p)$ independent of \mathbf{X} , as $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$, we have*

$$\begin{aligned}\mathbb{P}(\hat{\mathbf{x}}^\top \mathbf{w}(t) > 0 \mid \hat{y} = -1) - Q\left(\frac{E_{\text{test}}}{\sqrt{V_{\text{test}}}}\right) &\rightarrow 0, \\ \mathbb{P}(\mathbf{x}^\top \mathbf{w}(t) > 0 \mid y = -1) - Q\left(\frac{E_{\text{train}}}{\sqrt{V_{\text{train}}}}\right) &\rightarrow 0,\end{aligned}$$

almost surely, where

$$\begin{aligned}E_{\text{test}} &\equiv -\frac{1}{2\pi i} \oint_{\Gamma} \frac{1-f_t(z)}{z} \frac{\|\boldsymbol{\mu}\|^2 m(z) dz}{(\|\boldsymbol{\mu}\|^2 + c) m(z) + 1} \\ V_{\text{test}} &\equiv \frac{1}{2\pi i} \oint_{\Gamma} \left[\frac{\frac{1}{z^2} (1-f_t(z))^2}{(\|\boldsymbol{\mu}\|^2 + c) m(z) + 1} - \sigma^2 f_t^2(z) m(z) \right] dz \\ E_{\text{train}} &\equiv -\frac{1}{2\pi i} \oint_{\Gamma} \frac{1-f_t(z)}{z} \frac{dz}{(\|\boldsymbol{\mu}\|^2 + c) m(z) + 1} \\ V_{\text{train}} &\equiv \frac{1}{2\pi i} \oint_{\Gamma} \left[\frac{\frac{1}{z} (1-f_t(z))^2}{(\|\boldsymbol{\mu}\|^2 + c) m(z) + 1} - \sigma^2 f_t^2(z) z m(z) \right] dz - E_{\text{train}}^2 - \gamma V_{\text{test}}\end{aligned}$$

with Γ a positive contour surrounding the support of the Marčenko–Pastur law (shifted by $\gamma \geq 0$) and the points $(\gamma, 0)$ and $(\gamma + \lambda_s, 0)$ with $\lambda_s = c + 1 + \|\boldsymbol{\mu}\|^2 + c\|\boldsymbol{\mu}\|^{-2}$, $f_t(z) \equiv \exp(-atz)$ and $m(z)$ given by (5.15).

Note that the point $(\gamma + \lambda_s, 0)$ is the (possible) spike due to the low rank structure $\boldsymbol{\mu}\mathbf{y}^\top$ in \mathbf{X} . We have in particular that

$$\lambda_s = c + 1 + \|\boldsymbol{\mu}\|^2 + \frac{c}{\|\boldsymbol{\mu}\|^2} \geq (\sqrt{c} + 1)^2 \quad (5.16)$$

with equality if and only if $\|\boldsymbol{\mu}\|^2 = \sqrt{c}$. As in Theorem 2.13, here we see that the spike isolates from the main bulk of the Marčenko–Pastur support (with right edge $(\sqrt{c} + 1)^2$) only when $\|\boldsymbol{\mu}\|^2 \geq \sqrt{c}$.

However, the expressions in Theorem 5.3 of contour integrations are not easily analyzable nor interpretable. To obtain a more explicit expression, we choose the rectangular contour as in Figure 2.9 that circles around both the main bulk and the isolated eigenvalue (if any). For a shifted version (by γ) of the main bulk of the Marčenko–Pastur law (between $\lambda_- \equiv (1 - \sqrt{c})^2$ and $\lambda_+ \equiv (1 + \sqrt{c})^2$) given by

$$\nu(dx) = \frac{\sqrt{(x - \lambda_-)^+(\lambda_+ - x)^+}}{2\pi c x} dx + (1 - c^{-1})^+ \delta(x) \quad (5.17)$$

we know from Theorem 2.10 that the limit $\lim_{\epsilon_y \rightarrow 0} m(x + i\epsilon_y)$ exists for x within the support, while the second part which encloses the isolated eigenvalue

at $(\gamma_s + \gamma, 0)$ can be computed by residue calculus. This together leads to the expressions of $(E_{\text{test}}, V_{\text{test}})$ and $(E_{\text{train}}, V_{\text{train}})$ as follows.

$$E_{\text{test}} = \int \frac{1 - f_t(x + \gamma)}{x + \gamma} \omega(dx) \quad (5.18)$$

$$V_{\text{test}} = \frac{\|\mu\|^2 + c}{\|\mu\|^2} \int \frac{(1 - f_t(x + \gamma))^2 \omega(dx)}{(x + \gamma)^2} + \sigma^2 \int f_t^2(x + \gamma) \nu(dx) \quad (5.19)$$

$$E_{\text{train}} = \frac{\|\mu\|^2 + c}{\|\mu\|^2} \int \frac{1 - f_t(x + \gamma)}{x + \gamma} \omega(dx) \quad (5.20)$$

$$V_{\text{train}} = \frac{\|\mu\|^2 + c}{\|\mu\|^2} \int \frac{x(1 - f_t(x + \gamma))^2 \omega(dx)}{(x + \gamma)^2} + \sigma^2 \int x f_t^2(x + \gamma) \nu(dx) - E_{\text{train}}^2 \quad (5.21)$$

where we recall $f_t(x) = \exp(-\alpha t x)$, $\nu(x)$ is given by (5.17) and where we introduced the measure

$$\omega(dx) \equiv \frac{\sqrt{(x - \lambda_-)^+(\lambda_+ - x)^+}}{2\pi(\lambda_s - x)} dx + \frac{(\|\mu\|^4 - c)^+}{\|\mu\|^2} \delta_{\lambda_s}(x) \quad (5.22)$$

for λ_s defined in (5.16). The expressions in (5.18)-(5.21) are supported by the numerical results in Figure 5.7 with $p = 256$ and $n = 512$.




Figure 5.7: Training and test misclassification rate of a linear network as a function of the gradient descent training time t , for $p = 256$, $n = 512$, $\gamma = 0$, $\alpha = 10^{-2}$, $\sigma^2 = 0.1$ and $\mu = [-\mathbf{1}_{p/2}, \mathbf{1}_{p/2}] / \sqrt{p}$. Empirical results averaged over 50 runs. [Link to code]

5.2.0.2 Practical implications

With the simplified expressions in (5.18)-(5.21), several remarks are in order.

Remark 5.2 (Optimal performance). Note that, by the Cauchy-Schwarz inequality and the fact that $\int \omega(dx) = \|\boldsymbol{\mu}\|^2$, we have

$$E_{\text{test}}^2 \leq \int \frac{(1 - f_t(x + \gamma))^2}{(x + \gamma)^2} \omega(dx) \cdot \int \omega(dx) \leq \frac{\|\boldsymbol{\mu}\|^4}{\|\boldsymbol{\mu}\|^2 + c} V_{\text{test}}$$

with equality in the rightmost inequality only when the variance of the random initialization is $\sigma^2 = 0$. We thus have $E_{\text{test}}/\sqrt{V_{\text{test}}} \leq \|\boldsymbol{\mu}\|^2/\sqrt{\|\boldsymbol{\mu}\|^2 + c}$ so that the optimal test performance (lowest misclassification rate) is given by $Q(\|\boldsymbol{\mu}\|^2/\sqrt{\|\boldsymbol{\mu}\|^2 + c})$. This performance bound holds for any classifier obtained by minimizing the ridge-regularization squared loss in (5.12) with gradient descent (for any $t \geq 0$ and any regularization $\gamma \geq 0$). As seen later in Section 6.1, the bound in fact holds for an even larger class of generalized linear classifiers.

Remark 5.3 (Ridge regression and early-stopping). Note that the measures ν and ω defined in (5.17) and (5.22) do not depend on the regularization penalty γ . Since $(1 - f_t(x))/x$ and $f_t(x)$ are both decreasing functions of x , adding a positive ridge regularization $\gamma > 0$ is equivalent to early-stopping the gradient descent algorithm at some $t_\gamma < t$ without regularization (i.e., with $\gamma = 0$). The close connection between ridge regression and early-stopping in gradient descent has been explored in various models and settings [Yao et al., 2007, Suggala et al., 2018, Ali et al., 2019].

Remark 5.4 (Double descent in linear network). As $t \rightarrow \infty$, one obtains the ridge regression solution $\mathbf{w}_\infty = (\mathbf{X}\mathbf{X}^\top + n\gamma\mathbf{I}_p)^{-1}\mathbf{X}\mathbf{y}$, $\lambda \geq 0$ with

$$\frac{\boldsymbol{\mu}^\top \mathbf{w}_\infty}{\|\mathbf{w}_\infty\|} \rightarrow \frac{\|\boldsymbol{\mu}\|^2}{\sqrt{\|\boldsymbol{\mu}\|^2 + c}} \cdot \sqrt{\frac{c\gamma m^2(-\gamma) + 1}{cm(-\gamma) + 1}}$$

for $m(-\gamma)$ the unique positive solution to $-c\gamma m^2(-\gamma) - (1 - c + \gamma)m(-\gamma) + 1 = 0$.

In particular, taking $\lambda \rightarrow 0$ gives the “ridgeless” least squares solution $\mathbf{w}_{LS} = (\mathbf{X}^+)^{\top} \mathbf{y}$ with \mathbf{X}^+ the Moore-Penrose pseudoinverse of \mathbf{X} , that is equivalent to $\mathbf{w}_{LS} = (\mathbf{X}\mathbf{X}^\top)^{-1}\mathbf{X}\mathbf{y}$ for invertible $\mathbf{X}\mathbf{X}^\top$ and $\mathbf{w}_{LS} = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1}\mathbf{y}$ for invertible $\mathbf{X}^\top \mathbf{X}$ so that

$$\frac{\boldsymbol{\mu}^\top \mathbf{w}_{LS}}{\|\mathbf{w}_{LS}\|} \rightarrow \frac{\|\boldsymbol{\mu}\|^2}{\sqrt{\|\boldsymbol{\mu}\|^2 + c}} \cdot \sqrt{1 - \min(c, c^{-1})} \quad (5.23)$$

when one of $\mathbf{X}\mathbf{X}^\top$ and $\mathbf{X}^\top \mathbf{X}$ is invertible (which is almost surely the case for $c = \lim p/n \neq 1$). This means that the least squares solution \mathbf{w}_{LS} overfits the training set and the performance drops by a factor $\sqrt{1 - \min(c, c^{-1})}$ compared to the theoretical optimal value in Remark 5.2. This becomes even worse when the ratio p/n gets close to 1 and can be viewed as another manifestation of the “double descent” behavior discussed at the end of Section 5.1.1.2 (here the parameter dimension of $\mathbf{w} \in \mathbb{R}^p$ coincides with the input data dimension p).

Since positive ridge regularization $\gamma > 0$ is known to help alleviate this sharp performance drop at $c = 1$ [Hastie et al., 2019, Mei and Montanari, 2019], we compare, in Figure 5.8, the misclassification rate as a function of the

training time, for $\gamma = 0$ and $\gamma = 0.1$. As t grows large, the regularized classifier with $\gamma = 0.1$ has a test error that saturates at a much lower level than the unregularized classifier with $\gamma = 0$, the test error of which continues to grow (to 0.5 that is equivalent to random guess) as predicted in (5.23).




Figure 5.8: Test performance as a function of the training time t , for $n = p = 512$, $\gamma \in \{0, 0.1\}$, $\alpha = 0.1$, $\sigma^2 = 0.1$ and $\mu = [\sqrt{2}, \mathbf{0}_{p-1}]$. Empirical results averaged over 50 runs. [Link to code]

Following a similar direction as presented here, the authors of [Advani et al., 2020] considered the problem of learning, with a gradient descent approach, a similar linear regressor (or a linear single-layer neural network) from n training samples of dimension p , which are however generated by a noisy teacher network; that is, the training labels/targets \mathbf{y} are generated from an independent teacher network taking the same training data \mathbf{X} as input, and then corrupted with additive Gaussian noise. Under the setting of $n, p \rightarrow \infty$ with $p/n \rightarrow c \in (0, \infty)$, they found, by studying the training and test MSEs as a function of the training time (i.e., the number of descent steps), that the overfitting problem becomes most critical when c is close to 1, namely, when the number of parameters of the linear regressor (i.e., the model complexity) p gets close to the number of training data n . Although their theoretical results are limited to the linear regression model, they additionally carried experiments which provide strong evidence that these conclusions extend to deep linear and nonlinear neural networks, as has been discussed in Section 5.1.1.

5.3 Recurrent neural nets: echo state networks

5.3.1 Preliminaries and echo state networks

Echo-state neural networks (ESNs), popularized by Jaeger [2001], are elementary and simply parametrized, yet already quite efficient, recurrent neural networks. Also referred to as *reservoir computing* networks, they consist of a




Figure 5.9: Illustration of an echo state network.

single-hidden layer of size n with state $\mathbf{s}_t \in \mathbb{R}^n$ at time t , which evolves according to:

$$\mathbf{s}_{t+1} = \sigma(\mathbf{W}\mathbf{s}_t + \mathbf{W}_{\text{in}}\mathbf{x}_{t+1} + \eta\boldsymbol{\varepsilon}_{t+1})$$

for $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ an entry-wise activation function, $\mathbf{W} \in \mathbb{R}^{n \times n}$ the neuron connectivity matrix (which induces the recursion), $\mathbf{W}_{\text{in}} \in \mathbb{R}^{n \times p}$ the input layer connection and $\mathbf{x}_t \in \mathbb{R}^p$ the input data at time t . Added to the state is sometimes an independent noise term $\eta\boldsymbol{\varepsilon}_{t+1}$ for $\eta \in \mathbb{R}$, $\boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ mimicking thermal noise inside the network (of relevance in biological modeling of short-term memory neural networks).

In particular, for $\mathbf{W} = \mathbf{0}$ and $\eta = 0$, the network reduces to a nonlinear single-layer projection map, which boils down to a random feature map if \mathbf{W}_{in} is randomly designed.

Assuming the existence of a training dataset $\{(\mathbf{x}_t, \mathbf{y}_t)\}_{t=0}^{T-1}$ over a “time” window T , where $\mathbf{y}_t \in \mathbb{R}^d$ is the expected output at time t , the echo state network learning consists in a mere linear regression from the state \mathbf{s}_t into the output \mathbf{y}_t by minimizing

$$L(\boldsymbol{\beta}) = \frac{1}{T} \|\mathbf{Y} - \boldsymbol{\beta}^T \mathbf{S}\|_F^2$$

over the regression matrix $\boldsymbol{\beta} \in \mathbb{R}^{n \times d}$, where $\mathbf{X} = [\mathbf{x}_0, \dots, \mathbf{x}_{T-1}] \in \mathbb{R}^{p \times T}$, $\mathbf{Y} = [\mathbf{y}_0, \dots, \mathbf{y}_{T-1}] \in \mathbb{R}^{d \times T}$ and $\mathbf{S} = [\mathbf{s}_0, \dots, \mathbf{s}_{T-1}] \in \mathbb{R}^{n \times T}$. The explicit form of $\boldsymbol{\beta}$ is given by

$$\boldsymbol{\beta} = \begin{cases} \mathbf{S}(\mathbf{S}^T \mathbf{S})^{-1} \mathbf{Y}^T & , n > T; \\ (\mathbf{S} \mathbf{S}^T)^{-1} \mathbf{S} \mathbf{Y}^T & , n < T. \end{cases} \quad (5.24)$$

assuming $\mathbf{S}^T \mathbf{S}$ and $\mathbf{S} \mathbf{S}^T$ is invertible, respectively. Once $\boldsymbol{\beta}$ is set, the corresponding test error on a test set $\hat{\mathbf{X}} \in \mathbb{R}^{p \times \hat{T}}$ with underlying associated output $\hat{\mathbf{Y}} \in \mathbb{R}^{d \times \hat{T}}$ is then given by

$$E_{\text{test}} = \frac{1}{\hat{T}} \|\hat{\mathbf{Y}} - \boldsymbol{\beta}^T \hat{\mathbf{S}}\|_F^2$$

where $\hat{\mathbf{S}} = [\hat{\mathbf{s}}_0, \dots, \hat{\mathbf{s}}_{\hat{T}-1}] \in \mathbb{R}^{n \times \hat{T}}$ and $\hat{\mathbf{s}}_{t+1} = \sigma(\mathbf{W}\hat{\mathbf{s}}_t + \mathbf{W}_{\text{in}}\hat{\mathbf{x}}_{t+1} + \eta\hat{\boldsymbol{\varepsilon}}_{t+1})$, with $\hat{\boldsymbol{\varepsilon}}_t$ independent of $\boldsymbol{\varepsilon}_t$.

Being recursive networks, typical tasks of ESNs are time series regression when \mathbf{y}_t is a function of $\mathbf{x}_t, \mathbf{x}_{t-1}, \dots$, or time series prediction when $\mathbf{y}_t = \mathbf{x}_{t+\tau}$ for a certain $\tau > 0$.

As opposed to more involved recursive networks, such as the popular long short term memory (LSTM) nets [Hochreiter and Schmidhuber, 1997], echo state networks are extremely easy to train. Yet, they are hyper-parametrized by several key variables: the reservoir connectivity matrix $\mathbf{W} \in \mathbb{R}^{n \times n}$, the input layer $\mathbf{W}_{\text{in}} \in \mathbb{R}^{n \times p}$ and the activation function $\sigma(\cdot)$. Due to the recursive structure though, even in the simplest settings, establishing theoretically the training and test performance remains complex.

Possibly the most important hyperparameter is the spectral radius $\rho(\mathbf{W}) = \max_{1 \leq i \leq n} |\lambda_i(\mathbf{W})|$ (note that the square matrix \mathbf{W} is not imposed to be symmetric so the spectral radius is not necessarily the spectral norm). Indeed, in spirit, the recursive nature of ESNs involves the successive powers \mathbf{W}^t which, for t large, either decays exponentially with t for $\rho(\mathbf{W}) < 1$, thereby only maintaining short term memory in this reservoir, or diverges exponentially for $\rho(\mathbf{W}) > 1$, leading to quickly unstable behavior. The key property of echo state networks though is that, with a carefully chosen nonlinear activation $\sigma(\cdot)$, values of $\rho(\mathbf{W})$ slightly greater than 1 are still allowed and preserve the stability of the network, placing it in a so-called *near-chaotic* mode.

Yet, the successive iterations involving the nonlinearity $\sigma(\cdot)$ are much less tractable and we shall discuss here the simplified, yet already quite theoretically elaborate, setting where $\sigma(t) = t$ (hence the linear network case), and where \mathbf{W} and \mathbf{W}_{in} are successively fixed deterministic and then set randomly. We also assume the scalar setting where $p = d = 1$ for both input and output variables, letting in particular $\mathbf{W}_{\text{in}} = \mathbf{w}_{\text{in}} \in \mathbb{R}^n$, $\mathbf{X}^\top = \mathbf{x} = [x_0, \dots, x_{T-1}]^\top \in \mathbb{R}^T$, $\mathbf{Y}^\top = \mathbf{y} = [y_0, \dots, y_{T-1}]^\top \in \mathbb{R}^T$, and work under the simultaneously large n, T regime.

5.3.2 Results on ESN asymptotics

Gathering the simplifications above, we consider here the model

$$\mathbf{s}_{t+1} = \mathbf{W}\mathbf{s}_t + \mathbf{w}_{\text{in}}x_{t+1} + \eta\boldsymbol{\varepsilon}_{t+1}$$

with the associated training and test errors given by

$$E_{\text{train}} = \frac{1}{T}\|\mathbf{y} - \mathbf{S}^\top\boldsymbol{\beta}\|^2, \quad E_{\text{test}} = \frac{1}{\hat{T}}\|\hat{\mathbf{y}} - \hat{\mathbf{S}}^\top\boldsymbol{\beta}\|^2 \quad (5.25)$$

and $\boldsymbol{\beta} \in \mathbb{R}^n$ such that

$$\boldsymbol{\beta} = \begin{cases} \mathbf{S}(\mathbf{S}^\top\mathbf{S})^{-1}\mathbf{y} & , n > T; \\ (\mathbf{S}\mathbf{S}^\top)^{-1}\mathbf{S}\mathbf{y} & , n < T. \end{cases} \quad (5.26)$$

For further simplicity of exposition, we particularly focus here on the training performance, which already conveys quite inspiring results. The complete analyses of both train and test performance are available in [Coullet et al., 2016b].

To investigate the large n, T asymptotics of the training error E_{train} defined in (5.25), first remark that, letting

$$\mathbf{Q}_\gamma \equiv \left(\frac{1}{T} \mathbf{S} \mathbf{S}^\top + \gamma \mathbf{I}_n \right)^{-1} \quad \text{and} \quad \tilde{\mathbf{Q}}_\gamma \equiv \left(\frac{1}{T} \mathbf{S}^\top \mathbf{S} + \gamma \mathbf{I}_T \right)^{-1} \quad (5.27)$$

we have, irrespective of the sign of $n - T$,

$$E_{\text{train}} = \lim_{\gamma \downarrow 0} \frac{\gamma}{T} \mathbf{y}^\top \tilde{\mathbf{Q}}_\gamma \mathbf{y}$$

thereby formulating the estimation problem as the characterization of a quadratic form over the resolvent of $\frac{1}{T} \mathbf{S}^\top \mathbf{S}$.

The specific difficulty induced by \mathbf{S} lies in the intricate dependence between its columns, as successive observations of a multivariate time series. In particular, in order to simplify the analysis and to avoid edge problems at time $t = 0$, we assume (as is conventionally done in practice) that a sufficiently long “washout period” is performed preliminary to observing x_0 , i.e., the considered time series x_0, \dots, x_{T-1} is a finite time extraction of an infinite series $\dots, x_{-1}, x_0, x_1, \dots$; this ensures in particular the stationarity of the random network states $\mathbf{s}_0, \dots, \mathbf{s}_{T-1}$.

With these in mind, we may first describe the state evolution $\mathbf{S} = [\mathbf{s}_0, \dots, \mathbf{s}_{T-1}]$ through the following convenient relation

$$\begin{aligned} \mathbf{S} &= \sqrt{T}(\mathbf{A} + \mathbf{Z}), \quad \mathbf{A} = \mathbf{M}\mathbf{X}, \quad \mathbf{M} = \{\mathbf{W}^j \mathbf{w}_{\text{in}}\}_{j=0}^{T-1} \in \mathbb{R}^{n \times T}, \\ \mathbf{X} &= \frac{\{\mathbf{x}_{j-i}\}_{i,j=0}^{T-1}}{\sqrt{T}} \in \mathbb{R}^{T \times T}, \quad \mathbf{Z} = \frac{\eta}{\sqrt{T}} \left\{ \sum_{k \geq 0} \mathbf{W}^k \boldsymbol{\varepsilon}_{j-k} \right\}_{j=0}^{T-1} \in \mathbb{R}^{n \times T}. \end{aligned}$$

This relation isolates the random part of \mathbf{S} into \mathbf{Z} and the deterministic time series \mathbf{x} into the Toeplitz matrix \mathbf{X} . This expression clearly evidences the important stability condition of the network dynamics, i.e., $\rho(\mathbf{W}) < 1$. We impose this condition from now on.

Using the Gaussian method and Stein’s lemma, Lemma 2.13, we can then evaluate the behavior of the resolvents $\mathbf{Q} \equiv \mathbf{Q}_\gamma$ and $\tilde{\mathbf{Q}} \equiv \tilde{\mathbf{Q}}_\gamma$ by remarking that, for \mathbf{Q} (similar relations being obtained for $\tilde{\mathbf{Q}}$),

$$\mathbb{E}[\mathbf{Q}]_{ij} = \frac{1}{\gamma} \delta_{ij} - \frac{1}{\gamma} \left(\underbrace{\mathbb{E}[\mathbf{Z} \mathbf{Z}^\top \mathbf{Q}]_{ij}}_{(I)} + \underbrace{\mathbb{E}[\mathbf{Z} \mathbf{A}^\top \mathbf{Q}]_{ij}}_{(II)} + \underbrace{\mathbb{E}[\mathbf{A} \mathbf{Z}^\top \mathbf{Q}]_{ij}}_{(III)} + \underbrace{\mathbb{E}[\mathbf{A} \mathbf{A}^\top \mathbf{Q}]_{ij}}_{(IV)} \right)$$

and handle the terms $(I), \dots, (IV)$ subsequently. From the expansion

$$[\mathbf{Z}]_{ab} = \frac{\eta}{\sqrt{T}} \sum_{k \geq 0} \sum_{m=1}^n [\mathbf{W}^k]_{am} \cdot \boldsymbol{\varepsilon}_{m,b-k}$$

we have in particular that

$$\frac{\partial [\mathbf{Z}]_{ab}}{\partial \boldsymbol{\varepsilon}_{il}} = \frac{\eta}{\sqrt{T}} \sum_{k \geq 0} \sum_{q=1}^n \delta_{qi} \delta_{l,b-k} [\mathbf{W}^k]_{aq}$$

and

$$\begin{aligned} \frac{\partial[\mathbf{Q}]_{mj}}{\partial \varepsilon_{il}} = & -\frac{\eta}{\sqrt{T}} \sum_{q=1}^n \delta_{l \leq q} \left([\mathbf{Q}(\mathbf{Z} + \mathbf{A})]_{mq} [(\mathbf{W}^{q-l})^\top \mathbf{Q}]_{ij} \right. \\ & \left. + [(\mathbf{Z} + \mathbf{A})^\top \mathbf{Q}]_{qj} [\mathbf{Q} \mathbf{W}^{q-l}]_{mi} \right). \end{aligned}$$

These two relations are then exploited to develop the terms $(I), \dots, (IV)$, however with a specific difficulty: quite unlike the random matrix models studied so far in the monograph, this formula involves a large sum (over the index q) of successive powers of \mathbf{W} . Fortunately, the exponentially fast decrease of \mathbf{W}^k , due to $\rho(\mathbf{W}) < 1$, makes most of these powers negligible and only roughly $O(\log T)$ of them effectively remain. This, as such, does not impede the technical development and the control of small terms via the Nash-Poincaré inequality, Lemma 2.14, the development is only more cumbersome.

To facilitate the computations and present the results in a simpler form, it is convenient to define the shift matrix $\mathbf{J} \in \mathbb{R}^{T \times T}$ with $[\mathbf{J}^q]_{ij} \equiv \delta_{i+q,j}$, for which $[\mathbf{J}^q \mathbf{B}]_{ij} = [\mathbf{B}]_{q+i,j}$. A careful control of the development ultimately leads to the following deterministic equivalents for \mathbf{Q}_γ and $\tilde{\mathbf{Q}}_\gamma$, established in the limit of simultaneously large n, T ,

$$\begin{aligned} \mathbf{Q}_\gamma \leftrightarrow \bar{\mathbf{Q}}_\gamma &\equiv \frac{1}{\gamma} \left(\mathbf{I}_n + \eta^2 \tilde{\mathbf{R}}_\gamma + \frac{1}{\gamma} \mathbf{A} (\mathbf{I}_T + \eta^2 \mathbf{R}_\gamma)^{-1} \mathbf{A}^\top \right)^{-1} \\ \tilde{\mathbf{Q}}_\gamma \leftrightarrow \tilde{\bar{\mathbf{Q}}}_\gamma &\equiv \frac{1}{\gamma} \left(\mathbf{I}_T + \eta^2 \mathbf{R}_\gamma + \frac{1}{\gamma} \mathbf{A}^\top (\mathbf{I}_n + \eta^2 \tilde{\mathbf{R}}_\gamma)^{-1} \mathbf{A} \right)^{-1} \end{aligned}$$

where $\mathbf{R}_\gamma \in \mathbb{R}^{T \times T}$ and $\tilde{\mathbf{R}}_\gamma \in \mathbb{R}^{n \times n}$ are solutions to

$$\mathbf{R}_\gamma = \left\{ \frac{1}{T} \text{tr} (\mathcal{W}_{i-j} \bar{\mathbf{Q}}_\gamma) \right\}_{i,j=1}^T, \quad \tilde{\mathbf{R}}_\gamma = \sum_{q=-\infty}^{\infty} \frac{1}{T} \text{tr} (\mathbf{J}^q \tilde{\bar{\mathbf{Q}}}_\gamma) \cdot \mathcal{W}_q$$

where $\mathcal{W}_q \equiv \sum_{k \geq 0} \mathbf{W}^{k+(-q)^+} (\mathbf{W}^{k+(q)^+})^\top$ with $(a)^+ = \max(a, 0)$. A detailed development is provided in [Coullet et al., 2016b].

Taking the limit $\lim_{\gamma \downarrow 0} \gamma \tilde{\mathbf{Q}}_\gamma$, we thus find as an immediate corollary that, as $n, T \rightarrow \infty$ with $n/T \rightarrow c \in (0, \infty) \setminus \{1\}$, $E_{\text{train}} - \bar{E}_{\text{train}} \rightarrow 0$ almost surely, with

$$\bar{E}_{\text{train}} = \frac{1}{T} \mathbf{y}^\top \tilde{\mathbf{Q}} \mathbf{y} \cdot 1_{c < 1} + 0 \cdot 1_{c > 1} \tag{5.28}$$

where

$$\begin{aligned} \tilde{\mathbf{Q}} &\equiv \left(\mathbf{I}_T \cdot \delta_{c < 1} + \mathcal{R} + \frac{1}{\eta^2} \mathbf{A}^\top \left(\mathbf{I}_n \cdot \delta_{c > 1} + \tilde{\mathcal{R}} \right)^{-1} \mathbf{A} \right)^{-1} \\ \mathbf{Q} &\equiv \left(\mathbf{I}_n \cdot \delta_{c > 1} + \tilde{\mathcal{R}} + \frac{1}{\eta^2} \mathbf{A} (\mathbf{I}_T \cdot \delta_{c < 1} + \mathcal{R})^{-1} \mathbf{A}^\top \right)^{-1} \end{aligned}$$

and

$$\begin{aligned}\mathcal{R} &= \left\{ \frac{1}{T} \text{tr} \left(\mathcal{W}_{i-j} \left(\mathbf{I}_n \cdot \delta_{c>1} + \tilde{\mathcal{R}} \right)^{-1} \right) \right\}_{i,j=1}^T \\ \tilde{\mathcal{R}} &= \sum_{q=-\infty}^{\infty} \frac{1}{T} \text{tr} (\mathbf{J}^q (\mathbf{I}_T \cdot \delta_{c<1} + \mathcal{R})^{-1}) \cdot \mathcal{W}_q.\end{aligned}$$

Due to the ill-defined nature of some of these limits when $\gamma \downarrow 0$, the relations between \mathcal{R} , $\tilde{\mathcal{R}}$, \mathcal{Q} , $\tilde{\mathcal{Q}}$, and their associated \mathbf{R}_γ , $\tilde{\mathbf{R}}_\gamma$, \mathbf{Q}_γ , $\tilde{\mathbf{Q}}_\gamma$ are not mere corresponding limits as $\gamma \downarrow 0$. Specifically, when $c < 1$, we have, as $\gamma \downarrow 0$ that: $\eta^2 \mathbf{R}_\gamma \rightarrow \mathcal{R}$, $\gamma \tilde{\mathbf{R}}_\gamma \rightarrow \tilde{\mathcal{R}}$, $\eta^2 \mathbf{Q}_\gamma \rightarrow \mathcal{Q}$, and $\gamma \tilde{\mathbf{Q}}_\gamma \rightarrow \tilde{\mathcal{Q}}$. When instead $c > 1$, we have correspondingly: $\gamma \mathbf{R}_\gamma \rightarrow \mathcal{R}$, $\eta^2 \tilde{\mathbf{R}}_\gamma \rightarrow \tilde{\mathcal{R}}$, $\gamma \tilde{\mathbf{Q}}_\gamma \rightarrow \tilde{\mathcal{Q}}$, and $\eta^2 \tilde{\mathbf{Q}}_\gamma \rightarrow \tilde{\mathcal{Q}}$.

It is particularly useful to expand the resolvent $\tilde{\mathcal{Q}}$ in the expression of \bar{E}_{train} in (5.28) to retrieve some intuition on this result. Indeed, for $c < 1$, we precisely have

$$\bar{E}_{\text{train}} = \frac{1}{T} \mathbf{y}^\top \left(\mathbf{I}_T + \mathcal{R} + \frac{1}{\eta^2} \mathbf{X}^\top \left\{ \mathbf{w}_{\text{in}}^\top (\mathbf{W}^i)^\top \tilde{\mathcal{R}}^{-1} \mathbf{W}^j \mathbf{w}_{\text{in}} \right\}_{i,j=0}^{T-1} \mathbf{X} \right)^{-1} \mathbf{y}.$$

As such, the memory (training) performance of the network particularly depends on the typical speed of decay of the entries of $\left\{ \mathbf{w}_{\text{in}}^\top (\mathbf{W}^i)^\top \tilde{\mathcal{R}}^{-1} \mathbf{W}^j \mathbf{w}_{\text{in}} \right\}$ as one moves away from its $(1, 1)$ entry. A particularly telling example is the pure-memory task by which $\mathbf{x} = [\sqrt{T}, \mathbf{0}_{T-1}]$ (all the vector energy is concentrated at time 0) and $\mathbf{y} = [\mathbf{0}_{\tau-1}, \sqrt{T}, \mathbf{0}_{T-\tau}]$, that is we wish to recover at time τ the value of x_0 . Then, we find that

$$\bar{E}_{\text{train}} = \left[\left(\mathbf{I}_T + \mathcal{R} + \frac{1}{\eta^2} \mathbf{X}^\top \left\{ \mathbf{w}_{\text{in}}^\top (\mathbf{W}^i)^\top \tilde{\mathcal{R}}^{-1} \mathbf{W}^j \mathbf{w}_{\text{in}} \right\}_{i,j=0}^{T-1} \mathbf{X} \right)^{-1} \right]_{\tau+1, \tau+1}.$$

As shown subsequently, for \mathbf{W} (non-symmetric) having random independent zero mean entries, all off-diagonal entries of \mathcal{R} and $\left\{ \mathbf{w}_{\text{in}}^\top (\mathbf{W}^i)^\top \tilde{\mathcal{R}}^{-1} \mathbf{W}^j \mathbf{w}_{\text{in}} \right\}$ asymptotically vanish, and we have in this case

$$\bar{E}_{\text{train}} = \frac{\eta^2}{\eta^2 (1 + [\mathcal{R}]_{11}) + \mathbf{w}_{\text{in}}^\top (\mathbf{W}^\tau)^\top \tilde{\mathcal{R}}^{-1} \mathbf{W}^\tau \mathbf{w}_{\text{in}}}.$$

For arbitrary \mathbf{W} and \mathbf{w}_{in} , these performance asymptotics may not be quite expressive though. Simpler forms of the performance emerge when considering randomly drawn connectivity matrices. In particular, for $c < 1$, letting \mathbf{w}_{in} be independent of \mathbf{W} and of unit norm and $\mathbf{W} = \alpha \mathbf{W}^\circ$ where $\mathbf{W}^\circ \in \mathbb{R}^{n \times n}$ is a Haar matrix (see Section 2.6.2.3) and $\alpha < 1$, we find that

$$\bar{E}_{\text{train}} = (1 - c) \frac{1}{T} \mathbf{y}^\top \left(\mathbf{I}_T + \frac{1}{\eta^2} \mathbf{X}^\top \mathbf{D} \mathbf{X} \right)^{-1} \mathbf{y} \quad (5.29)$$

where $\mathbf{D} = \text{diag}\{(1 - \alpha^2)\alpha^{2(i-1)}\}_{i=1}^T$ and $\alpha < 1$ plays the role of a (short-term) memory depth parameter.

This result may be further expanded into a “multi-modal memory” structure for $\mathbf{W} \in \mathbb{R}^{n \times n}$ by letting $\mathbf{W} = \text{diag}(\mathbf{W}_1, \dots, \mathbf{W}_k)$ where $\mathbf{W}_i = \alpha_i \mathbf{W}_i^\circ$ and $\mathbf{W}_i^\circ \in \mathbb{R}^{n_i \times n_i}$ is a Haar matrix independent of all other \mathbf{W}_j° 's as well as $\sum_{i=1}^k n_i = n$. Writing $c_i = \lim_n n_i/n \in (0, 1)$, \bar{E}_{train} has the same form as in (5.29) above, however with $\mathbf{D} = \text{diag}\{[\mathbf{D}]_{ii}\}_{i=1}^T$ now given by

$$[\mathbf{D}]_{ii} = \frac{\sum_{j=1}^k c_j \alpha_j^{2(i-1)}}{\sum_{j=1}^k c_j (1 - \alpha_j^2)^{-1}}.$$

By letting $\alpha_1 < \dots < \alpha_k < 1$, this form of \mathbf{D} is interesting as it exhibits a “controlled” decay of the memory capacity for a range of memory depths α_j .

In particular, defining formally the memory capacity $\text{MC}(\tau; \mathbf{W})$ as the inverse training error for a pure data recovery shifted by τ in the noiseless limit, i.e.,

$$\text{MC}(\tau; \mathbf{W}) = \lim_{\eta \downarrow 0} \eta^2 E_{\text{train}}^{-1}, \quad \text{for } \mathbf{x} = [\sqrt{T}, \mathbf{0}_{T-1}], \mathbf{y} = [\mathbf{0}_{\tau-1}, \sqrt{T}, \mathbf{0}_{T-\tau}] \quad (5.30)$$

we obtain the typical memory curves depicted in Figure 5.10. This expression of the “memory capacity” of the network is tightly connected with the so-called Fisher memory curve proposed in [Ganguli et al., 2008] as an alternative measure of the network memory.




Figure 5.10: Memory curves for $\mathbf{W} = \text{diag}(\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_3)$, $\mathbf{W}_j = \alpha_j \mathbf{W}_j^\circ$, $\mathbf{W}_j^\circ \in \mathbb{R}^{n_j \times n_j}$ Haar distributed, $\alpha_1 = 0.99$, $n_1/n = 0.01$, $\alpha_2 = 0.9$, $n_2/n = 0.1$, and $\alpha_3 = 0.5$, $n_3/n = 0.89$, compared to single-mode memory with \mathbf{W}_i^+ such that $\mathbf{W}_i^+ = \alpha_i \mathbf{W}_i^{+\circ}$ for Haar $\mathbf{W}_i^{+\circ} \in \mathbb{R}^{n \times n}$, $n/T = 0.75$. [Link to code]




Figure 5.11: Upper 9×9 part of \mathcal{R} for $c = 1/2$ for $\mathbf{W} = \alpha \mathbf{W}^\circ$, $\alpha = 0.9$, and \mathbf{W}° with i.i.d. zero mean and variance $1/n$ non-symmetric Gaussian entries (**left**) and symmetric-Gaussian (**right**). Linear grayscale representation with black being 1 and white being 0. [\[Link to code\]](#)




Figure 5.12: Symmetric versus non-symmetric \mathbf{W} performance ($\mathbf{W} = \alpha \mathbf{W}^\circ$, \mathbf{W}° with i.i.d. $\mathcal{N}(0, 1/n)$ entries possibly up to symmetry; $\alpha = 0.9$): (**Left**) Memory curve $MC(\tau; \mathbf{W})$; (**Right**) Training performance of a τ -delay task for $\tau \in \{1, \dots, 4\}$ on the popular Mackey-Glass near-chaotic model [[Glass and Mackey, 1979](#)].

In these recurrent network structures, symmetry constraints in \mathbf{W} were shown in the literature not to be generally profitable, although few arguments are proposed to support this claim [Canaday, 2019, Section 5.5]. The deterministic equivalents obtained above allow for a better understanding, as exemplified by Figure 5.11 which compares the matrices \mathcal{R} for \mathbf{W} Gaussian symmetric or non-symmetric: the symmetric \mathbf{W} case exhibits an erratic behavior with a quite specific and structured correlation of the time-delayed source data; the associated loss in performance in both mean squared error and memory capacity of symmetric reservoirs is corroborated by the theoretical results displayed in Figure 5.12.

These asymptotic results established in this section, along with the further studies carried out in [Couillet et al., 2016b], allow for a thorough understanding and further improvement of the design of random connectivity matrices aiming for enhanced (possibly selective) memory performance of the networks.

The above study, and the theoretical literature on echo state network performance as a whole, is nonetheless limited to the case of linear nets, while we claimed above the very interest of these networks to lie in the combination of a nonlinear function $\sigma(\cdot)$ and of a carefully chosen spectral radius $\rho(\mathbf{W}) > 1$. In the same way as the performance of deep feedforward neural networks (even trained ones with fully random weight matrices) is difficult to study, notably due to their accumulating nonlinearities under the form $\sigma(\mathbf{W}_L\sigma(\mathbf{W}_{L-1}\cdots\sigma(\mathbf{W}_1\mathbf{x}_i)))$, the performance of nonlinear echo state networks remains an open riddle.

5.4 Concluding remarks

In this chapter, by leveraging tools from the concentration of measure theory (see Section 2.7), we went beyond the (mere) Taylor expansion-based approach to the understanding of nonlinear transformations for kernel methods in Chapter 4, and evaluated the large dimensional asymptotics, of a single-hidden-layer *nonlinear* neural network model in Section 5.1 for real-world data. As we shall see later in Chapter 8, this “universal large dimensional concentration” argument has an even more significant impact in practical applications and will be exploited to extend the current analyses and insights (on the choice of kernel function and activation function) to a much broader and more realistic setting.

There has been a growing interest in the eigenspectra, or more generally the large dimensional asymptotics of (random) neural network models. These investigations include the (limiting) eigenspectral measure of the nonlinear Gram matrices [Pennington and Worah, 2017, Benigni and Péché, 2019] (similar to Section 5.1), as well as that of the input-output Jacobian matrices [Pennington et al., 2017, Pastur, 2020, Pastur and Slavin, 2020] (closely connected to the behavior of back propagation gradients) of a multilayer neural network with random Gaussian or orthogonal weights. These analyses are not limited to classical feedforward fully connected networks but have been performed on networks with convolutional [Novak et al., 2019, Xiao et al., 2018], recurrent [Chen et al.,

2018, Gilboa et al., 2019] and skip-connection structures [Ling and Qiu, 2019] (as in the case of the popular residual network architecture [He et al., 2016]), to name a few. Since random (Gaussian) initializations are widely used in training such deep networks [Glorot and Bengio, 2010], these works shed important light on the initial “landscape” of deep neural networks as well as on the impact of nonlinear activations.

The investigations on randomly weighted neural networks are of even greater interest through the *neural tangent kernel* recently introduced by Jacot et al. [2018]: while the weight matrices *after* training are no longer random, the underlying neural tangent kernel is determined only by the random initialization and remains unchanged during the whole training procedure in the “infinite-neurons” limit. As such, the eigenspectral assessments of the neural tangent kernel of randomly weighted deep networks go beyond the initial stage of training and leads to much richer results on, for example, the learning dynamics of networks [Fan and Wang, 2020, Adlam and Pennington, 2020], at least in this so-called neural tangent limit where the networks widths are *much larger* than both the number of training data n and their dimension p . Nonetheless, most of these works are concerned with random noise-like input data (e.g., i.i.d. Gaussian data or almost orthonormal data [Fan and Wang, 2020, Adlam et al., 2019]) with no information structure, and thus fail to provide sharp qualitative predictions on real-world datasets such as the MNIST dataset. In this respect, considering more involved structural data models, together with the “universal large dimensional concentration” to be discussed later in Chapter 8, is expected to bring greater insights into these elaborate learning systems in a more precise and qualitative manner that better match real-world observations.

Focusing on the optimization methods used in neural network training, we discussed in Section 5.2 the gradient descent dynamics in learning a simple ridge regression model arising from the minimization of the (ridge-regularized) square loss, which enjoys the advantage of having an explicit solution. In the more general context of modern machine learning, however, this is rarely the case: (i) the optimization problem often has no closed-form solution and, perhaps even worse, (ii) it may not be guaranteed to have a *unique* solution, due to the non-convex nature of the problem, for instance in the case of deep neural network models. The non-convexity has long been one of the main technical difficulties for a more solid theoretical understanding of the deep learning method.

In pursuit of the theoretical understanding of the non-convex “loss landscape” of deep networks, Bray and Dean [2007] studied the nature of critical points of *random Gaussian fields* in a large dimensional setting from a statistical physics approach, by evaluating for instance the fraction of negative eigenvalues of the Hessian at critical points. This type of analyses provides precise descriptions on the “chance” of continuing to decrease the loss function as well as on the number of saddle points (with indefinite Hessian) or “bad” local minima (that have significantly higher loss values) [Dauphin et al., 2014, Choromanska et al., 2015, Pennington and Bahri, 2017]. Nonetheless, these works are formally based on statistical physics models quite different from actual neural networks (such

as the spin glass model) and require quite restrictive assumptions. A clear and more precise picture of the large dimensional statistical loss landscape of deep networks that accounts for the influence of the network depth, width, choice of nonlinear activation, etc., is still in need.

While optimization problems with non-explicit solutions are ubiquitous in machine learning and pose additional technical difficulty in evaluating their performance, many of the convex problems (that admit a unique solution) still remain accessible in the large dimensional regime. As a telling example, when the more commonly used *cross-entropy* loss (rather than the square loss)

$$L(\boldsymbol{\beta}) = -\frac{1}{n} \sum_{i=1}^n [y_i \log(\sigma(\boldsymbol{\beta}^\top \mathbf{x}_i)) + (1 - y_i) \log(1 - \sigma(\boldsymbol{\beta}^\top \mathbf{x}_i))] \quad (5.31)$$

is adopted to learn a (generalized) linear classifier $\boldsymbol{\beta}$ from a training set $\{(\mathbf{x}_i, y_i)\}_{i=1}^n$, with $\sigma(t) = (1 + e^{-t})^{-1}$ the *logistic sigmoid* function and label $y_i \in \{0, 1\}$, no closed-form solution exists for the loss-minimizing solution $\boldsymbol{\beta}$. In fact, due to the intricate dependence of the learned classifier $\boldsymbol{\beta}$ on $\{\mathbf{x}_i, y_i\}_{i=1}^n$, the statistical behavior of $\boldsymbol{\beta}$ is highly non-trivial to tackle and this gets even worse with the application of the logistic sigmoid nonlinearity $\sigma(\cdot)$. Despite all these technical difficulties, it is nonetheless possible to pursue a large dimensional stochastic description of $\boldsymbol{\beta}$ and consequently to evaluate the performance of, not only the logistic regression classifier, but also any smooth convex loss L that falls into the general *empirical risk minimization* framework discussed in the next chapter.

5.5 Practical course material

Practical Lecture Material 5 (Effective kernel of large dimensional random Fourier features). *As discussed in Remark 5.1, instead of the single-type nonlinearity setting in Figure 5.1 thoroughly investigated in Section 5.1.1, from a random feature map and kernel approximation perspective, a mixture of nonlinearities such as ‘cos + sin’ in the case of random Fourier features [Rahimi and Recht, 2008] turns out to be a more natural choice. Specifically, for $\mathbf{W} \in \mathbb{R}^{N \times p}$ with independent standard Gaussian entries, the random Fourier features refer to the cascade of the random features from both ‘cos’ and ‘sin’ activations as*

$$\boldsymbol{\Sigma}^\top = [\cos(\mathbf{W}\mathbf{X})^\top \quad \sin(\mathbf{W}\mathbf{X})^\top] \in \mathbb{R}^{n \times 2N}. \quad (5.32)$$

Using the fact that $\mathbb{E}_{\mathbf{w}}[\cos(\mathbf{w}^\top \mathbf{x}_i) \sin(\mathbf{w}^\top \mathbf{x}_j)] = 0$ for $\mathbf{w} \sim \mathcal{N}(0, \mathbf{I}_p)$ and that both $\cos(\cdot)$ and $\sin(\cdot)$ are 1-Lipschitz, show, with the help of Lemma 5.1 and similar to Theorem 5.1, that the random Fourier resolvent $(\frac{1}{n} \boldsymbol{\Sigma}^\top \boldsymbol{\Sigma} + \gamma \mathbf{I}_n)^{-1}$ admits the following deterministic equivalent

$$\mathbf{Q} \leftrightarrow \bar{\mathbf{Q}}, \quad \bar{\mathbf{Q}} \equiv \left(\frac{N}{n} \left(\frac{\mathbf{K}_{\text{cos}}}{1 + \delta_{\text{cos}}} + \frac{\mathbf{K}_{\text{sin}}}{1 + \delta_{\text{sin}}} \right) + \gamma \mathbf{I}_n \right)^{-1}$$

for $(\delta_{\cos}, \delta_{\sin})$ the unique positive solution to

$$\delta_{\cos} = \frac{1}{n} \operatorname{tr} \mathbf{K}_{\cos} \bar{\mathbf{Q}}, \quad \delta_{\sin} = \frac{1}{n} \operatorname{tr} \mathbf{K}_{\sin} \bar{\mathbf{Q}}$$

with \mathbf{K}_{\cos} and \mathbf{K}_{\sin} the limiting kernels of ‘cos’ and ‘sin’ nonlinearities enlisted in Table 5.1, respectively.

Then, similar to Corollary 5.1, show that the asymptotic training and test MSEs takes the following forms

$$\begin{aligned}\bar{E}_{\text{train}} &= \frac{\gamma^2}{n} \operatorname{tr} \mathbf{Y} \bar{\mathbf{Q}}^2 \mathbf{Y}^\top \\ &\quad + \frac{N}{n} \frac{\gamma^2}{n} \left[\frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\cos} \bar{\mathbf{Q}} - \frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\sin} \bar{\mathbf{Q}} \right] \Omega \begin{bmatrix} \operatorname{tr} \mathbf{Y} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\cos} \bar{\mathbf{Q}} \mathbf{Y}^\top \\ \operatorname{tr} \mathbf{Y} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\sin} \bar{\mathbf{Q}} \mathbf{Y}^\top \end{bmatrix} \\ \bar{E}_{\text{test}} &= \frac{1}{\hat{n}} \|\hat{\mathbf{Y}}^\top - \Phi_{\mathbf{X}\hat{\mathbf{X}}}^\top \bar{\mathbf{Q}} \mathbf{Y}^\top\|_F^2 + \left(\frac{N}{n} \right)^2 \frac{1}{\hat{n}} \begin{bmatrix} \Theta_{\cos} & \Theta_{\sin} \end{bmatrix} \Omega \begin{bmatrix} \operatorname{tr} \mathbf{Y} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\cos} \bar{\mathbf{Q}} \mathbf{Y}^\top \\ \operatorname{tr} \mathbf{Y} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\sin} \bar{\mathbf{Q}} \mathbf{Y}^\top \end{bmatrix}\end{aligned}$$

with $\bar{\mathbf{K}}_{\cos} \equiv \frac{\mathbf{K}_{\cos}}{1+\delta_{\cos}}$, $\bar{\mathbf{K}}_{\sin} \equiv \frac{\mathbf{K}_{\sin}}{1+\delta_{\sin}}$,

$$\Omega^{-1} = \mathbf{I}_2 - \frac{N}{n} \begin{bmatrix} \frac{\frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\cos} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\cos}}{(1+\delta_{\cos})^2} & \frac{\frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\cos} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\sin}}{(1+\delta_{\sin})^2} \\ \frac{\frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\sin} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\cos}}{(1+\delta_{\cos})^2} & \frac{\frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\sin} \bar{\mathbf{Q}} \bar{\mathbf{K}}_{\sin}}{(1+\delta_{\sin})^2} \end{bmatrix}$$

and, for $\sigma \in \{\cos, \sin\}$,

$$\begin{aligned}\Theta_\sigma &\equiv \frac{1}{1+\delta_\sigma} \left(\frac{1}{N} \operatorname{tr} \bar{\mathbf{K}}_\sigma \hat{\mathbf{X}} \hat{\mathbf{X}}^\top + \frac{N}{n} \frac{1}{n} \operatorname{tr} \bar{\mathbf{Q}} \Phi_{\mathbf{X}\hat{\mathbf{X}}} \Phi_{\mathbf{X}\hat{\mathbf{X}}}^\top \bar{\mathbf{Q}} \mathbf{K}_\sigma - \frac{2}{N} \operatorname{tr} \bar{\mathbf{Q}} \Phi_{\mathbf{X}\hat{\mathbf{X}}} (\mathbf{K}_\sigma^{\mathbf{X}\hat{\mathbf{X}}})^\top \right) \\ \Phi &= \frac{N}{n} (\bar{\mathbf{K}}_{\cos} + \bar{\mathbf{K}}_{\sin}), \quad \Phi_{\mathbf{X}\hat{\mathbf{X}}} = \frac{N}{n} (\bar{\mathbf{K}}_{\cos}^{\mathbf{X}\hat{\mathbf{X}}} + \bar{\mathbf{K}}_{\sin}^{\mathbf{X}\hat{\mathbf{X}}}).\end{aligned}$$

Confirm numerically that for not-too-large ratio N/n as in Figure 5.13, a significant gap exists between empirical training MSE of random Fourier ridge regression and the classical Gaussian kernel prediction on MNIST data, while the large N, p, n asymptotics derived above consistently fits empirical observations almost perfectly.

Check also that a double-descent singularity behavior occurs when $\gamma \rightarrow 0$, however at $2N = n$ (rather than $N = n$ as in the case of single nonlinearity discussed at the end of Section 5.1.1.2), due the singular behavior of the two by two matrix Ω^{-1} .




Figure 5.13: Training MSEs of RFF ridge regression on MNIST data (class 3 versus 7), as a function of regression penalty λ , for $p = 784$, $n = 1\,024$, $N = 256$ and 512. Empirical results displayed in **blue** circles; Gaussian kernel predictions (assuming $N \rightarrow \infty$ alone) in **black** dashed lines; and RMT predictions in **red** solid lines. Results obtained by averaging over 30 runs. [\[Link to code\]](#)

Chapter 6

Optimization-based Methods with Non-explicit Solutions

Unlike kernel eigenspectra-based methods in Section 4 or simple neural network models in Section 5, where the objects of study (e.g., kernel matrices and random feature ridge regressors) are given in a rather *explicit* manner, many other machine learning algorithms take the form of (the solutions of) optimization problems, having in general *no closed-form* expressions. A first example is the popular logistic regression algorithm, where one aims to find, in a binary classification setting, an optimal linear classifier $\beta \in \mathbb{R}^p$ by minimizing the logistic loss $\frac{1}{n} \sum_{i=1}^n \log(1 + e^{-y_i \beta^\top \mathbf{x}_i})$ over a training set $\{(\mathbf{x}_i, y_i)\}_{i=1}^n$ with labels $y_i \in \{-1, +1\}$. More generally, by choosing other loss functions rather than the logistic loss, logistic regression can be viewed as a special case of the *empirical risk minimization* [Vapnik, 1992] problem

$$\arg \min_{\beta \in \mathbb{R}^p} \frac{1}{n} \sum_{i=1}^n L(y_i \beta^\top \mathbf{x}_i) + \frac{\gamma}{2} \|\beta\|^2 \quad (6.1)$$

for some convex loss $L : \mathbb{R} \rightarrow \mathbb{R}^+$ and regularization factor $\gamma \geq 0$. With the logistic loss $L(t) = \log(1 + e^{-t})$ one gets the logistic regression, while the least-squares classifier (or ridge regressor) can be obtained with the square loss $L(t) = (t - 1)^2$. Other popular choices of $L(\cdot)$ include the exponential loss $L(t) = e^{-t}$ widely used in boosting algorithms [Schapire, 1999] and the hinge loss $L(t) = \max(0, 1 - t)$ in the case of support vector machines (SVMs) [Rosasco et al., 2004]. Figure 6.1 illustrates these different losses.

Except for the least-squares solution where $L(t) = (t - 1)^2$, the minimization of a generic loss L generally leads to a classifier β which only takes an *implicit* form. It is thus less clear how the resulting β depends on the data \mathbf{X} , and makes it more challenging to investigate its (large dimensional) statistical behavior.

The technical challenge from implicit optimization problems appears not only in the analysis of logistic regression, but also in most other machine learning algorithms of daily use, starting with the popular deep learning schemes.




Figure 6.1: Different loss functions for classification: 0–1 loss (**red**), logistic loss (**green**), exponential loss (**purple**), square loss (**blue**) and hinge loss (**black**).

It is therefore of crucial importance to adapt the random matrix-based analysis framework discussed in the previous chapters to assess the performance of optimization-based learning methods. In this chapter, we focus on the empirical risk minimization example of (6.1) and evaluate the large dimensional behavior of the classifier β . Technically, a major emphasis will be cast on the “leave-one-out” approach, which aims to “decouple” the intricate statistical dependencies within the learning system.

6.1 Generalized linear classifier

6.1.1 Basic setting

For simplicity of exposition, we consider the problem of classifying a binary symmetric Gaussian mixture of the form

$$\mathcal{C}_1 : \mathbf{x}_i \sim \mathcal{N}(-\boldsymbol{\mu}, \mathbf{C}), \quad y_i = -1 \quad \mathcal{C}_2 : \mathbf{x}_i \sim \mathcal{N}(+\boldsymbol{\mu}, \mathbf{C}), \quad y_i = +1 \quad (6.2)$$

each with a class prior probability of 1/2, for some $\boldsymbol{\mu} \in \mathbb{R}^p$ and positive definite $\mathbf{C} \in \mathbb{R}^{p \times p}$. As in the previous chapters, we ensure the classification problem is asymptotically non-trivial by specifying the following growth rate assumptions.

$$\|\boldsymbol{\mu}\| = O(1), \quad \max\{\|\mathbf{C}\|, \|\mathbf{C}^{-1}\|\} = O(1). \quad (6.3)$$

Note that the mixture model in (6.2) satisfies the logistic model in the sense that the conditional class probability is

$$\begin{aligned} P(y | \mathbf{x}) &= \frac{P(y)P(\mathbf{x} | y)}{P(\mathbf{x})} = \frac{e^{-\frac{1}{2}(\mathbf{x}-y\boldsymbol{\mu})\mathbf{C}^{-1}(\mathbf{x}-y\boldsymbol{\mu})}}{e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})\mathbf{C}^{-1}(\mathbf{x}-\boldsymbol{\mu})} + e^{-\frac{1}{2}(\mathbf{x}+\boldsymbol{\mu})\mathbf{C}^{-1}(\mathbf{x}+\boldsymbol{\mu})}} \\ &= \frac{1}{1 + e^{-2y\boldsymbol{\mu}^\top \mathbf{C}^{-1}\mathbf{x}}} \equiv \sigma(\boldsymbol{\beta}_*^\top y\mathbf{x}), \quad \boldsymbol{\beta}_* \equiv 2\mathbf{C}^{-1}\boldsymbol{\mu} \end{aligned} \quad (6.4)$$

with $\sigma(t) = (1 + e^{-t})^{-1}$ the *logistic sigmoid* function and the optimal Bayes solution $\beta_* = 2\mathbf{C}^{-1}\mu$. By the symmetry of (6.2), it is convenient to use the shortcut notation $\tilde{\mathbf{x}}_i \equiv y_i \mathbf{x}_i$ so that

$$\tilde{\mathbf{x}}_i \sim \mathcal{N}(\mu, \mathbf{C})$$

regardless of the class of \mathbf{x}_i .

To investigate the large dimensional asymptotics of the implicit classifier that minimizes the empirical risk in (6.1), the main technical difficulty lies in the fact that β , as the solution of a convex optimization problem, depends on all the random $\tilde{\mathbf{x}}_i$'s in a rather involved manner. Nonetheless, by canceling the loss function derivative with respect to β in (6.1), we can still obtain the following seemingly simple equation satisfied by β ,

$$\gamma\beta = \frac{1}{n} \sum_{i=1}^n -L'(\beta^\top \tilde{\mathbf{x}}_i) \tilde{\mathbf{x}}_i \quad (6.5)$$

where we assume the loss function L is convex and at least three-times continuously differentiable (making β unique for $\gamma > 0$). We provide, in the following section, the main “leave-one-out” intuition of crucial significance in building up the system of equations which (asymptotically) characterizes the statistical behavior of β .

6.1.2 Intuitions and main results

The proof of the main results on the asymptotic characterization of β is based on the “leave-one-out” intuition similar to the proof of Theorem 2.4, to solve a “leave-one-version” of (6.5) (without the i -th datum $\tilde{\mathbf{x}}_i$), the solution of which should be asymptotically close to the original solution β , so as to form a self-consistent equation for β . Here we focus on providing a few intuitive arguments to convey the main idea, with the proof details given in [Mai and Liao, 2019].

From (6.5), β can be viewed as a linear combination of all $\tilde{\mathbf{x}}_i$'s, weighted by the coefficient $-L'(\beta^\top \tilde{\mathbf{x}}_i)$. The idea is to understand how $\tilde{\mathbf{x}}_i$ affects the corresponding coefficient $-L'(\beta^\top \tilde{\mathbf{x}}_i)$. To handle the complex dependence of β on all $\tilde{\mathbf{x}}_i$'s, we establish a “leave-one-out” version of β , denoted β_{-i} , that is (i) asymptotically close to β by removing the contribution of a single datum and (ii) that is independent of $\tilde{\mathbf{x}}_i$, by solving (6.1) for all data $\tilde{\mathbf{x}}_j$ different from $\tilde{\mathbf{x}}_i$, that is

$$\gamma\beta_{-i} = \frac{1}{n} \sum_{j \neq i} -L'(\beta_{-i}^\top \tilde{\mathbf{x}}_j) \tilde{\mathbf{x}}_j.$$

As a consequence, the difference $\lambda(\beta - \beta_{-i})$ satisfies

$$\gamma(\beta - \beta_{-i}) = \frac{1}{n} \sum_{j \neq i} (L'(\beta_{-i}^\top \tilde{\mathbf{x}}_j) - L'(\beta^\top \tilde{\mathbf{x}}_j)) \tilde{\mathbf{x}}_j - \frac{1}{n} L'(\beta^\top \tilde{\mathbf{x}}_i) \tilde{\mathbf{x}}_i. \quad (6.6)$$

Intuitively, the difference $\|\beta - \beta_{-i}\|$ must vanish as $n, p \rightarrow \infty$ so that, by Taylor expanding L' around $\beta_{-i}^\top \tilde{\mathbf{x}}_j$, $j \neq i$, one obtains

$$L'(\beta^\top \tilde{\mathbf{x}}_j) = L'(\beta_{-i}^\top \tilde{\mathbf{x}}_j) + L''(\beta_{-i}^\top \tilde{\mathbf{x}}_j)(\beta - \beta_{-i})^\top \tilde{\mathbf{x}}_j + O(\|\beta - \beta_{-i}\|^2).$$

Ignoring higher order terms, together with (6.6), this leads to the following equation for $\beta - \beta_{-i}$

$$\gamma(\beta - \beta_{-i}) \simeq -\frac{1}{n} \sum_{j \neq i} L''(\beta_{-i}^\top \tilde{\mathbf{x}}_j) \tilde{\mathbf{x}}_j \tilde{\mathbf{x}}_j^\top (\beta - \beta_{-i}) - \frac{1}{n} L'(\beta^\top \tilde{\mathbf{x}}_i) \tilde{\mathbf{x}}_i$$

or equivalently

$$\left(\frac{1}{n} \tilde{\mathbf{X}}_{-i} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top + \gamma \mathbf{I}_p \right) (\beta - \beta_{-i}) \simeq -\frac{1}{n} L'(\beta^\top \tilde{\mathbf{x}}_i) \tilde{\mathbf{x}}_i$$

with $\tilde{\mathbf{X}}_{-i} = [\tilde{\mathbf{x}}_1, \dots, \tilde{\mathbf{x}}_{i-1}, \tilde{\mathbf{x}}_{i+1}, \dots, \tilde{\mathbf{x}}_n] \in \mathbb{R}^{p \times (n-1)}$ and diagonal $\mathbf{D}_{-i} \in \mathbb{R}^{(n-1) \times (n-1)}$ with $[\mathbf{D}_{-i}]_{jj} = L''(\beta_{-i}^\top \tilde{\mathbf{x}}_j)$ that are both independent of \mathbf{x}_i .

Note further by the convexity of L that $L''(t) \geq 0$ for all t , so that for any $\gamma > 0$, the matrix $\frac{1}{n} \tilde{\mathbf{X}}_{-i} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top + \gamma \mathbf{I}_p$ is positive definite (and invertible). Thus we may write

$$\beta - \beta_{-i} \simeq -\frac{1}{n} L'(\beta^\top \tilde{\mathbf{x}}_i) \left(\frac{1}{n} \tilde{\mathbf{X}}_{-i} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top + \gamma \mathbf{I}_p \right)^{-1} \tilde{\mathbf{x}}_i$$

which, while not solving a linear regression problem, once again involves the *resolvent* of (a “weights” version of) the sample covariance matrix of the training data.

In particular, projecting against $\tilde{\mathbf{x}}_i$ gives

$$(\beta - \beta_{-i})^\top \tilde{\mathbf{x}}_i \simeq -\frac{1}{n} L'(\beta^\top \tilde{\mathbf{x}}_i) \tilde{\mathbf{x}}_i^\top \left(\frac{1}{n} \tilde{\mathbf{X}}_{-i} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top + \gamma \mathbf{I}_p \right)^{-1} \tilde{\mathbf{x}}_i. \quad (6.7)$$

At this point, note that $(\frac{1}{n} \tilde{\mathbf{X}}_{-i} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top + \gamma \mathbf{I}_p)^{-1}$ is of bounded spectral norm (by $1/\gamma$) and independent of $\tilde{\mathbf{x}}_i = \mu + \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i$ for $\mathbf{z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$. By the trace lemma (Lemma 2.11) and Theorem 2.7, one must therefore have under the growth rate (6.3) that, as $n \rightarrow \infty$

$$\frac{1}{n} \tilde{\mathbf{x}}_i^\top \left(\frac{1}{n} \tilde{\mathbf{X}}_{-i} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top + \gamma \mathbf{I}_p \right)^{-1} \tilde{\mathbf{x}}_i - \delta \xrightarrow{a.s.} 0 \quad (6.8)$$

with δ the unique positive solution to¹

$$\delta = \frac{1}{n} \text{tr } \mathbf{C} \left(\mathbb{E} \left[\frac{L''(\beta^\top \tilde{\mathbf{x}})}{1 + \delta L''(\beta^\top \tilde{\mathbf{x}})} \right] \mathbf{C} + \gamma \mathbf{I}_p \right)^{-1}. \quad (6.9)$$

¹In the present setting, \mathbf{D}_{-i} plays the role of matrix $\tilde{\mathbf{C}}$ in Theorem 2.7 and the equality $-z\delta_p(z) = \frac{1}{n} \text{tr } \tilde{\mathbf{C}}(\mathbf{I}_n + \delta_p(z)\tilde{\mathbf{C}})^{-1}$ thus becomes, here for $z = -\gamma$, $\gamma\tilde{\delta}_p = \frac{1}{n} \sum_{j=1}^n [\mathbf{D}_{-i}]_{jj}/(1 + \delta[\mathbf{D}_{-i}]_{jj}) \xrightarrow{a.s.} \mathbb{E}[L''(\beta^\top \tilde{\mathbf{x}})/(1 + \delta L''(\beta^\top \tilde{\mathbf{x}}))]$. Formally though, \mathbf{D}_{-i} is *not independent* of \mathbf{X}_{-i} , so that Lemma 2.11 and Theorem 2.7 do not straightforwardly lead to the form in (6.9); yet, with additional arguments detailed in [Mai and Liao, 2019], this dependence can be shown to asymptotically vanish.

The positiveness of δ follows from the quadratic form in (6.8). As for uniqueness, it is guaranteed by rewriting the trace relation under the form

$$1 = \frac{1}{n} \sum_{i=1}^p \frac{\lambda_i}{\left(1 - \mathbb{E}\left[\frac{1}{1+\delta L''(\cdot)}\right]\right) \lambda_i + \gamma \delta} \equiv g(\delta)$$

with $\{\lambda_i\}_{i=1}^p$ the eigenvalues of \mathbf{C} and noticing that, for $L''(\cdot) \geq 0$, $g(\delta)$ is a decreasing function of δ with $\lim_{\delta \rightarrow 0} g(\delta) \rightarrow \infty$ and $\lim_{\delta \rightarrow \infty} g(\delta) \rightarrow 0$. This ensures the uniqueness of δ as g is thus one-to-one from $(0, \infty)$ to $(0, \infty)$.

Back to the expression of (6.8), it now unfolds from (6.7) that

$$\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i \simeq \boldsymbol{\beta}_{-i}^\top \tilde{\mathbf{x}}_i - \delta L'(\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i).$$

Solving for $\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i$, we may remark that $\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i$ is (approximately) explicitly given by the proximal operator of $\boldsymbol{\beta}_{-i}^\top \tilde{\mathbf{x}}_i$ for the function δL , i.e.,

$$\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i \simeq \text{prox}_{\delta L}(\boldsymbol{\beta}_{-i}^\top \tilde{\mathbf{x}}_i)$$

with $\text{prox}_{\delta L}(t)$ the *proximal mapping* [Bauschke and Combettes, 2017], defined as the *unique* solution of the minimization problem

$$\text{prox}_{\delta L}(t) \equiv \arg \min_{x \in \mathbb{R}} \left\{ \delta L(x) + \frac{1}{2}(x - t)^2 \right\}. \quad (6.10)$$

As a consequence, replacing in (6.5) gives the approximation

$$\gamma \boldsymbol{\beta} \simeq \frac{1}{n} \sum_{i=1}^n -L'(\text{prox}_{\delta L}(\boldsymbol{\beta}_{-i}^\top \tilde{\mathbf{x}}_i)) \tilde{\mathbf{x}}_i \equiv \frac{1}{n} \sum_{i=1}^n f(\boldsymbol{\beta}_{-i}^\top \tilde{\mathbf{x}}_i) \tilde{\mathbf{x}}_i, \quad (6.11)$$

where we denoted $f(x) \equiv -L'(\text{prox}_{\delta L}(x)) = (\text{prox}_{\delta L}(x) - x)/\delta$, which follows from differentiating the (convex) right-hand side of (6.10).

Equation (6.11) establishes an asymptotic connection between $\boldsymbol{\beta}$ and (the average of) the “leave-one-out” $\boldsymbol{\beta}_{-i}$. One may want for instance to take the expectation (6.11) to retrieve $\mathbb{E}[\boldsymbol{\beta}]$ as a function of the expectation of $\tilde{\mathbf{x}}_i$. However, the expectation of the type $\mathbb{E}_{\tilde{\mathbf{x}}_i}[f(\boldsymbol{\beta}_{-i}^\top \tilde{\mathbf{x}}_i) \tilde{\mathbf{x}}_i]$ is delicate to handle due to the $\tilde{\mathbf{x}}_i$ inside the nonlinear function $f(\cdot)$. This can be resolved by using the Gaussianity of $\tilde{\mathbf{x}}_i$ with the following decomposition: recall now that $\tilde{\mathbf{x}}_i = \boldsymbol{\mu} + \mathbf{C}^{1/2} \mathbf{z}_i$ for $\mathbf{z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, so that by conditioning on $\boldsymbol{\beta}_{-i}$ (which is independent of \mathbf{z}_i), one can decompose \mathbf{z}_i under the form

$$\mathbf{z}_i = \eta_i \frac{\mathbf{C}^{1/2} \boldsymbol{\beta}_{-i}}{\sqrt{\boldsymbol{\beta}_{-i}^\top \mathbf{C} \boldsymbol{\beta}_{-i}}} + \mathbf{z}_i^\perp, \quad \eta_i = \frac{\boldsymbol{\beta}_{-i}^\top \mathbf{C}^{1/2} \mathbf{z}_i}{\sqrt{\boldsymbol{\beta}_{-i}^\top \mathbf{C} \boldsymbol{\beta}_{-i}}} \quad (6.12)$$

with $\mathbf{C}^{1/2} \boldsymbol{\beta}_{-i} / \sqrt{\boldsymbol{\beta}_{-i}^\top \mathbf{C} \boldsymbol{\beta}_{-i}}$ the unit vector oriented in the direction of $\mathbf{C}^{1/2} \boldsymbol{\beta}_{-i}$ and \mathbf{z}_i^\perp lying on the $(p-1)$ -dimensional subspace orthogonal to $\mathbf{C}^{1/2} \boldsymbol{\beta}_{-i}$. By the

orthogonal invariance of the standard multivariate Gaussian distribution, both η_i and \mathbf{z}_i^\perp are Gaussian and orthogonal, thus independent. This is a natural decomposition of \mathbf{z}_i to reduce the fluctuation $\beta_{-i}^\top \mathbf{C}^{1/2} \mathbf{z}_i$ in the expansion of $\beta_{-i}^\top \tilde{\mathbf{x}}_i$ to the scalar fluctuation of $\eta_i \sim \mathcal{N}(0, 1)$ (since $\beta_{-i}^\top \mathbf{C}^{1/2} \mathbf{z}_i^\perp = 0$).

Exploiting this decomposition, (6.11) becomes

$$\begin{aligned}\gamma\beta &\simeq \frac{1}{n} \sum_{i=1}^n f(\beta_{-i}^\top \tilde{\mathbf{x}}_i)(\mu + \mathbf{C}^{1/2} \mathbf{z}_i) = \frac{1}{n} \sum_{i=1}^n f(\beta_{-i}^\top \mu + \beta_{-i}^\top \mathbf{C}^{1/2} \mathbf{z}_i)(\mu + \mathbf{C}^{1/2} \mathbf{z}_i) \\ &= \frac{1}{n} \sum_{i=1}^n f\left(\beta_{-i}^\top \mu + \sqrt{\beta_{-i}^\top \mathbf{C} \beta_{-i}} \eta_i\right) \left(\mu + \frac{\eta_i \mathbf{C} \beta_{-i}}{\sqrt{\beta_{-i}^\top \mathbf{C} \beta_{-i}}} + \mathbf{C}^{1/2} \mathbf{z}_i^\perp\right)\end{aligned}$$

for $\eta_i \sim \mathcal{N}(0, 1)$ (conditioned on β_{-i}) and *independent* of \mathbf{z}_i^\perp , which is more convenient to work with.

By construction, β_{-i} is independent of $\tilde{\mathbf{x}}_i$, so that, conditioning on β_{-i} the norm of which should converge to a limit (the same as that of $\|\beta\|$), one expects to have $\beta_{-i}^\top \tilde{\mathbf{x}}_i \sim \mathcal{N}(M, \sigma^2)$ in the large p, n limit. The deterministic pair (M, σ^2) is however so far unknown, but it must satisfy

$$M \simeq \mathbb{E}[\beta_{-i}^\top \mu] \simeq \beta_{-i}^\top \mu, \quad \sigma^2 \simeq \mathbb{E}[\beta_{-i}^\top \mathbf{C} \beta_{-i}] \simeq \beta_{-i}^\top \mathbf{C} \beta_{-i} \quad (6.13)$$

with a concentration argument. Intuitively, the variable variable $\beta^\top \mu$ characterizes the (expected) projection of β on a new datum (of mean μ) and determines the classification performance of β that should be asymptotically deterministic. A similar argument holds for $\beta_{-i}^\top \mathbf{C} \beta_{-i}$.

Assuming this indeed holds, we have, as $n, p \rightarrow \infty$,

$$\begin{aligned}\frac{1}{n} \sum_{i=1}^n f(\beta_{-i}^\top \tilde{\mathbf{x}}_i) \mu &\simeq \mathbb{E}[r] \mu, \\ \frac{1}{n} \sum_{i=1}^n f(\beta_{-i}^\top \tilde{\mathbf{x}}_i) \frac{\eta_i \mathbf{C} \beta_{-i}}{\sqrt{\beta_{-i}^\top \mathbf{C} \beta_{-i}}} &= \frac{1}{n} \sum_{i=1}^n f\left(\beta_{-i}^\top \mu + \sqrt{\beta_{-i}^\top \mathbf{C} \beta_{-i}} \cdot \eta_i\right) \frac{\eta_i \mathbf{C} \beta_{-i}}{\sqrt{\beta_{-i}^\top \mathbf{C} \beta_{-i}}} \\ &\simeq \left(\frac{1}{n} \sum_{i=1}^n f(M + \sigma \eta_i) \eta_i\right) \cdot \frac{\mathbf{C} \beta}{\sigma} \\ &\simeq \frac{\mathbb{E}[f(r)(r - M)]}{\sigma^2} \mathbf{C} \beta,\end{aligned}$$

for $r \sim \mathcal{N}(M, \sigma^2)$ with the law of large numbers, where for the second term we use that $\beta_{-i} \simeq \beta$. As such, (6.11) further reads

$$\gamma\beta \simeq \mathbb{E}[f(r)] \mu + \frac{\mathbb{E}[f(r)(r - M)]}{\sigma^2} \mathbf{C} \beta + \frac{1}{n} \sum_{i=1}^n f(M + \sigma \cdot \eta_i) \mathbf{C}^{1/2} \mathbf{z}_i^\perp \quad (6.14)$$

for η_i behaves asymptotically like a standard Gaussian variable, where we retain the last term under the form of (the sum of) random vectors \mathbf{z}_i^\perp since, unlike

the second term where all β_{-i} are asymptotically “in the same direction”, the \mathbf{z}_i^\perp ’s asymptotically behave like \mathbf{z}_i and are totally random (in fact, uniformly distributed on the unit sphere of radius \sqrt{p}), so we do not see “the effect of averaging” for the third term as for the second term. Further remark that with the decomposition in (6.12), \mathbf{z}_i^\perp is independent of η_i , so that by denoting $\mathbf{u} = \frac{1}{n} \sum_{i=1}^n f(M + \sigma \cdot \eta_i) \mathbf{C}^{\frac{1}{2}} \mathbf{z}_i^\perp$, we should have

$$\mathbb{E}[\mathbf{u}] = \mathbf{0}, \quad \mathbb{E}[\mathbf{u}\mathbf{u}^\top] \simeq \frac{\mathbb{E}[f^2(r)]}{n} \mathbf{C} \quad (6.15)$$

which, together with an asymptotically Gaussian fluctuation argument, leads to

$$\mathbf{u} \sim \mathcal{N} \left(\mathbf{0}, \frac{\mathbb{E}[f^2(r)]}{n} \mathbf{C} \right)$$

which is of the same amplitude as $\boldsymbol{\mu}$, and thus not negligible.

Solving (6.14) for $\boldsymbol{\beta}$ gives

$$(\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C}) \boldsymbol{\beta} \simeq \mathbb{E}[f(r)] \boldsymbol{\mu} + \mathbf{u} \quad (6.16)$$

which unfolds from an integration by parts to write $\mathbb{E}[f(r)(r - M)]/\sigma^2 = \mathbb{E}[f'(r)]$. This provides a *stochastic* characterization of $\boldsymbol{\beta}$ as the sum of a deterministic ($\mathbb{E}[f(r)]\boldsymbol{\mu}$) and a random (\mathbf{u}) part.

To close the loop and connect (M, σ^2) to the data statistics $\boldsymbol{\mu}, \mathbf{C}$ and particularly to those of $\boldsymbol{\beta}$, recall from (6.13) and $\boldsymbol{\beta}_{-i} \simeq \boldsymbol{\beta}$ that

$$M \simeq \mathbb{E}[\boldsymbol{\beta}]^\top \boldsymbol{\mu}, \quad \sigma^2 \simeq \text{tr}(\mathbf{C} \mathbb{E}[\boldsymbol{\beta} \boldsymbol{\beta}^\top]).$$

Therefore, taking the expectation of both sides of (6.16) and solving for $\mathbb{E}[\boldsymbol{\beta}]$, one reaches

$$\mathbb{E}[\boldsymbol{\beta}] \simeq \mathbb{E}[f(r)] (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C})^{-1} \boldsymbol{\mu}.$$

Recalling the definition $f : x \mapsto -(x - \text{prox}_{\delta L}(x))/\delta$ where $\delta > 0$, by the firmly non-expansive nature of the proximal mapping (see detail in [Bauschke and Combettes, 2017, Chapter 12]), it unfolds that $f'(\cdot) \leq 0$ and that the inverse in (6.16) is well defined for any $\gamma > 0$. [check the reasoning here!](#)

Lastly, using again (6.16),

$$\begin{aligned} \mathbb{E}[\boldsymbol{\beta} \boldsymbol{\beta}^\top] &\simeq (\mathbb{E}[f(r)])^2 (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C})^{-1} \boldsymbol{\mu} \boldsymbol{\mu}^\top (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C})^{-1} \\ &\quad + (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C})^{-1} \mathbb{E}[\mathbf{u} \mathbf{u}^\top] (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C})^{-1} \end{aligned}$$

so that

$$\begin{aligned} \sigma^2 &\simeq (\mathbb{E}[f(r)])^2 \cdot \boldsymbol{\mu}^\top (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C})^{-1} \mathbf{C} (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C})^{-1} \boldsymbol{\mu} \\ &\quad + \mathbb{E}[f^2(r)] \cdot \frac{1}{n} \text{tr} [(\gamma \mathbf{I}_p - \mathbb{E}[f'(r)] \mathbf{C})^{-2} \mathbf{C}^2]. \end{aligned}$$

The above heuristic derivation is summarized in the following theorem.

Theorem 6.1 (Asymptotic behavior of β , [Mai and Liao, 2019, Theorem 1]). For $\max\{\|\mu\|, \|\mathbf{C}\|, \|\mathbf{C}^{-1}\|\} = O(1)$, convex and three-times continuously differentiable L and β the unique solution to (6.1), we have, as $n, p \rightarrow \infty$,

$$\|\beta - \tilde{\beta}\| \rightarrow 0, \quad (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)]\mathbf{C})\tilde{\beta} \sim \mathcal{N}(\mathbb{E}[f(r)]\mu, \mathbb{E}[f^2(r)]\mathbf{C}/n)$$

where $f(r) = -L'(\text{prox}_{\delta L}(r))$, $r \sim \mathcal{N}(M, \sigma^2)$, and (M, σ^2) solution to

$$\begin{aligned} M &= \mathbb{E}[f(r)] \cdot \mu^\top (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)]\mathbf{C})^{-1} \mu, \\ \sigma^2 &= (\mathbb{E}[f(r)])^2 \cdot \mu^\top (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)]\mathbf{C})^{-1} \mathbf{C} (\gamma \mathbf{I}_p - \mathbb{E}[f'(r)]\mathbf{C})^{-1} \mu \\ &\quad + \mathbb{E}[f^2(r)] \cdot \frac{1}{n} \text{tr}[(\gamma \mathbf{I}_p - \mathbb{E}[f'(r)]\mathbf{C})^{-2} \mathbf{C}^2] \end{aligned}$$

with δ the unique positive solution to

$$\delta = \frac{1}{n} \text{tr} \mathbf{C} \left(\mathbb{E} \left[\frac{L''(\text{prox}_{\delta L}(r))}{1 + \delta L''(\text{prox}_{\delta L}(r))} \right] \mathbf{C} + \gamma \mathbf{I}_p \right)^{-1}.$$

Note that here $\tilde{\beta}$ (and thus β) is of norm order $O(1)$ (like μ), so if μ (and thus the mean $\mathbb{E}[f(t)](\gamma \mathbf{I}_p - \mathbb{E}[f'(r)]\mathbf{C})^{-1} \mu$ of $\tilde{\beta}$) is *delocalized* in the sense that its entries are of order $O(1/\sqrt{p})$, the entries of $\tilde{\beta}$ then fluctuate as $O(1/\sqrt{p})$, thus at the same order as their means: the vector is therefore not asymptotically deterministic. We will come back to this point in more detail later.

From the above ‘leave-one-out’ derivation, since β_{-i} is by construction independent of $\tilde{\mathbf{x}}_i$, the random variable $r_i = \beta_{-i}^\top \tilde{\mathbf{x}}_i$ is asymptotically close to $\beta^\top \tilde{\mathbf{x}}$, that is, the projection of the classifier β on a new datum $\tilde{\mathbf{x}} \sim \mathcal{N}(\mu, \mathbf{C})$. This gives access to the asymptotic test classification error rate

$$\mathbb{P}(\beta^\top \tilde{\mathbf{x}} < 0) - Q(M/\sigma) \rightarrow 0 \tag{6.17}$$

with $Q(\cdot)$ the Gaussian Q -function. Similarly, since $\beta^\top \tilde{\mathbf{x}}_i \simeq \text{prox}_{\delta L}(\beta_{-i}^\top \tilde{\mathbf{x}}_i)$, the training classification error is given by

$$\mathbb{P}(\beta^\top \tilde{\mathbf{x}}_i < 0) - \mathbb{P}(\text{prox}_{\delta L}(r) < 0) \rightarrow 0 \tag{6.18}$$

for $r \sim \mathcal{N}(M, \sigma^2)$ given in Theorem 6.1.

6.1.3 Practical consequences and further discussions

To validate the asymptotic results given in Theorem 6.1 for n, p of reasonable sizes, Figure 6.2 compares the empirical distribution of $\{\beta_{-i}^\top \tilde{\mathbf{x}}_i\}_{i=1}^n$ to the limiting Gaussian distribution $\mathcal{N}(M, \sigma^2)$ from the system of fixed-point equations in Theorem 6.1. The theoretical results are seen to fit the simulations almost perfectly, already for $p = 256$ and $n = 1024$.




Figure 6.2: Histogram of $\beta_{-i}^\top \tilde{\mathbf{x}}_i$ and the Gaussian distribution $\mathcal{N}(M, \sigma^2)$ defined in Theorem 6.1 with $\boldsymbol{\mu} = \mathbf{1}_p / \sqrt{p}$, $\mathbf{C} = \text{diag}[\mathbf{1}_{p/4}; 3 \cdot \mathbf{1}_{p/4}; 5 \cdot \mathbf{1}_{p/2}]$, for logistic loss $L(t) = \log(1 + e^{-t})$, $\lambda = 0.1$, $p = 256$ and $n = 1024$. [Link to code]

6.1.3.1 The existence and uniqueness of empirical risk minimizer

We now interpret the results in Theorem 6.1. First, let us for simplicity restrict ourselves to the unregularized case where $\gamma = 0$ and assume the existence and uniqueness of the solution to the (unregularized) optimization problem

$$\arg \min_{\boldsymbol{\beta} \in \mathbb{R}^p} \frac{1}{n} \sum_{i=1}^n L(y_i \boldsymbol{\beta}^\top \mathbf{x}_i) \quad (6.19)$$

and the asymptotic boundedness of the solution (i.e., $\limsup_p \|\boldsymbol{\beta}\| < \infty$). This assumption is necessary since, in the unregularized $\gamma = 0$ case, such a minimizer may not exist, and if it does, may not be unique or may have a diverging behavior. A well-known counterexample in the logistic regression $L(t) = \log(1 + e^{-t})$ setting is as follows: if the training data $\{(\mathbf{x}_i, y_i)\}_{i=1}^n$ are *linearly separable* in the sense that there exists a linear decision boundary $\mathbf{b} \in \mathbb{R}^p$ so that

$$y_i \mathbf{x}_i^\top \mathbf{b} \geq 0, \quad i \in \{1, \dots, n\}$$

then, since $L(t)$ is strictly decreasing, one can always decrease the objective function and have $\frac{1}{n} \sum_{i=1}^n L(y_i \alpha_1 \mathbf{b}^\top \mathbf{x}_i) < \frac{1}{n} \sum_{i=1}^n L(y_i \alpha_2 \mathbf{b}^\top \mathbf{x}_i)$ as long as $\alpha_1 > \alpha_2 > 0$, and the (global) minimizer of (6.19) does not exist.

Remark 6.1 (Existence and uniqueness of a well-defined empirical risk minimizer). *In the large dimensional setting under consideration, the existence of a unique and well-behaved minimizer can be characterized, as a function of the problem setting ($\boldsymbol{\mu}, \mathbf{C}$ and the loss L) and of the ratio p/n . For instance, it was shown in [Candès and Sur, 2020] for the logistic regression that a sharp phase transition exists for the existence of the minimizer of (6.19) in the sense that, for $g(\cdot)$ some decreasing function, if $p/n > g(\boldsymbol{\mu}^\top \mathbf{C}^{-1} \boldsymbol{\mu})$, the minimizer exists with probability approaching zero; but if $p/n < g(\boldsymbol{\mu}^\top \mathbf{C}^{-1} \boldsymbol{\mu})$, the probability*

approaches one. The function g additionally has the property that $g(\cdot) \leq 1/2$, meaning that the minimizer (asymptotically) does not exist if $n < 2p$.² This result was then extended in [Taheri et al., 2019] to more general convex losses.

According to the above discussion, assume the existence of a bounded solution to the unregularized optimization problem (6.19), from which one may simplify the expression in Theorem 6.1 as

$$\tilde{\beta} \sim \mathcal{N}\left(\frac{\mathbb{E}[f(r)]}{\mathbb{E}[f'(r)]}\mathbf{C}^{-1}\mu, \frac{\mathbb{E}[f^2(r)]}{(\mathbb{E}[f'(r)])^2}\frac{\mathbf{C}^{-1}}{n}\right) \quad (6.20)$$

for $r \sim \mathcal{N}(M, \sigma^2)$, $f(r) = -L'(\text{prox}_{\delta L}(r))$ with

$$M = \frac{\mathbb{E}[f(r)]}{\mathbb{E}[f'(r)]}\mu^\top \mathbf{C}^{-1}\mu, \quad \sigma^2 = \left(\frac{\mathbb{E}[f(r)]}{\mathbb{E}[f'(r)]}\right)^2 \mu^\top \mathbf{C}^{-1}\mu + \frac{\mathbb{E}[f^2(r)]}{(\mathbb{E}[f'(r)])^2}\frac{p}{n}$$

and δ the unique positive solution of

$$\mathbb{E}\left[\frac{1}{1 + \delta L''(\text{prox}_{\delta L}(r))}\right] = 1 - \frac{p}{n}. \quad (6.21)$$

Note already from the convexity of L that one must have (at least) $n > p$ so as to have $\delta > 0$, in accordance with the existence and uniqueness condition in [Candès and Sur, 2020, Taheri et al., 2019]. Also, taking $L''(t) = 0$, which excludes the existence of $\delta > 0$ for $p/n > 0$, is not allowed.

6.1.3.2 Implications to large dimensional empirical risk minimization

Debiasing the estimator in large dimensions. Recall from (6.4) that the underlying model follows a logistic model, with the optimal Bayes solution given by $\beta_* = 2\mathbf{C}^{-1}\mu$. However, from the asymptotic characterization in (6.20) the expectation of the minimizer β of the logistic loss $L(t) = \log(1 + e^{-t})$, despite being the maximum likelihood estimator, instead of being equal to β_* as the large- n asymptotic suggests [Rosasco et al., 2004], is only a scaled version of β_* , due to the non-vanishing ratio p/n .

Although in a classification context one has $\text{sign}(\beta^\top \mathbf{x}) = \text{sign}(\alpha\beta^\top \mathbf{x})$ for $\alpha > 0$, so that a positive constant rescaling of the classifier β does not affect the classification performance, it is often still desirable to have a large- (n, p) consistent estimator of β_* , for inference or risk management purposes. For instance, such an estimator is necessary to predict the variability of the obtained solution by means of standard errors, confidence intervals or p-values [Sur and Candes, 2019].

To this end, we see from (6.20) that it suffices to estimate $\mathbb{E}[f(r)]$ and $\mathbb{E}[f'(r)]$ in a consistent manner, which, according to the derivation in the previous section, can be evaluated as follows.

²This remark is reminiscent of the similar, yet formally different, phenomenon where the minimizer to (6.19) for square loss $L(t) = (t - 1)^2/2$ and $\gamma = 0$ exists but is *not unique* in the under-determined regime $n < p$.

Lemma 6.1. *Under the notations of the conditions of Theorem 6.1, we have*

$$\begin{aligned} -\frac{1}{n} \sum_{i=1}^n L'(\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i) - \mathbb{E}[f(t)] &\xrightarrow{a.s.} 0 \\ \frac{1}{n} \sum_{i=1}^n [L'(\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i)]^2 - \mathbb{E}[f^2(t)] &\xrightarrow{a.s.} 0 \\ \frac{1}{n} \sum_{i=1}^n \frac{L'(\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i)(r_i - \frac{1}{n} \sum_{i=1}^n r_i)}{\frac{1}{n} \sum_{i=1}^n (r_i - \frac{1}{n} \sum_{i=1}^n r_i)^2} - \mathbb{E}[f'(t)] &\xrightarrow{a.s.} 0 \end{aligned}$$

for $r_i = \boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i + \hat{\delta} L'(\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i)$ with $\hat{\delta}$ the (unique) positive solution to

$$\hat{\delta} = \frac{p}{n} \left(\frac{1}{n} \sum_{i=1}^n \frac{L''(\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i)}{1 + \hat{\delta} L''(\boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i)} \right)^{-1}.$$

We have in particular $|\hat{\delta} - \delta| \xrightarrow{a.s.} 0$.

Figure 6.3 examines the empirical mean (as an estimate of the expectation) of $\boldsymbol{\beta}$, the proposed rescaling version and the optimal Bayes solution $\boldsymbol{\beta}_*$ and confirms that, by rescaling $\boldsymbol{\beta}$ with the plug-in estimators in Lemma 6.1, one retrieves on average the optimal $\boldsymbol{\beta}_*$.




Figure 6.3: Comparison of the averaged $\boldsymbol{\beta}$ (over 500 realizations, to estimate the expectation), the optimal Bayes solution $\boldsymbol{\beta}_*$ and the (averaged) rescaled solution $\boldsymbol{\beta}$ proposed in Lemma 6.1, for logistic loss with $\boldsymbol{\mu} = [\mathbf{1}_{p/4}, -\mathbf{1}_{p/4}, \frac{3}{4}\mathbf{1}_{p/2}]/\sqrt{p}$, $\mathbf{C} = 2\mathbf{I}_p$, for $p = 64$ and $n = 512$. [\[Link to code\]](#)

Let us now focus on the random fluctuations of $\boldsymbol{\beta}$, which are (asymptotically) Gaussian with covariance \mathbf{C}^{-1}/n (so that the entry-wise noise is typically of the order $O(n^{-1/2})$) under (6.3). Therefore, since $\|\boldsymbol{\beta}_*\| = O(1)$, depending on the nature of $\boldsymbol{\beta}_*$, either of the following two situations may arise:

1. $\boldsymbol{\beta}_* \in \mathbb{R}^p$ is *sparse* in the sense that the number of its nonzero entries (i.e., its ℓ_0 norm) is of order $O(1)$ and each nonzero element is of order $O(1)$. In

this case, one may wish to perform some sort of soft-thresholding and to set noise-like small valued entries to zero, with a wisely chosen threshold;

2. $\beta_* \in \mathbb{R}^p$ is more “delocalized” with $\|\beta_*\|_0 = O(n)$ and $\|\beta_*\|_\infty = O(n^{-1/2})$, which is of the same order as the random noise. Intuitively, the energy of β_* is spread over all $O(n)$ entries and it is unlikely to recover the desired β_* from a single realization in this case. An alternative, yet computational more burdensome, approach is to solve n times the “leave-one-out” version of (6.19) by removing different datum each time, and then average the obtained solutions.

Optimal loss for classification. From a classification standpoint, it follows from (6.17) that the optimal design of the loss L is the function that maximizes the ratio M^2/σ^2 or, equivalently, as per (6.20), the function solution to

$$\max_f \frac{M^2}{\sigma^2} = \max_f \frac{(\mathbb{E}[f(r)])^2 (\mu^\top C^{-1} \mu)^2}{(\mathbb{E}[f(r)])^2 (\mu^\top C^{-1} \mu)^2 + \mathbb{E}[f^2(r)] \frac{p}{n}}.$$

An immediate remark is that, by the Cauchy-Schwarz inequality,

$$(\mathbb{E}[f(r)])^2 \leq \mathbb{E}[f^2(r)]$$

with equality if and only if $f(r) = -L'(\text{prox}_{\delta L}(r))$ is constant (which is however unattainable with the current framework as discussed previously below (6.21)). Yet, a theoretical misclassification error rate “lower bound” is given by

$$Q\left(\frac{M}{\sigma}\right) = Q\left(\frac{\mu^\top C^{-1} \mu}{\sqrt{\mu^\top C^{-1} \mu + p/n}}\right).$$

This extends the optimal performance bound for linear classifier from gradient descent minimizing squared loss in Remark 5.2 to more general loss functions.

For a given problem with fixed $\mu^\top C^{-1} \mu$ and p/n , it thus suffices to maximize the ratio $(\mathbb{E}[f(r)])^2 / \mathbb{E}[f^2(r)]$ or, alternatively, according to Lemma 6.1, to maximize the following empirical version for n, p large

$$\max_L \frac{|L'(\beta^\top \tilde{X}) \mathbf{1}_n|}{\sqrt{L'(\beta^\top \tilde{X}) L'(\tilde{X}^\top \beta)}} \quad (6.22)$$

where the function $L'(\cdot)$ is applied entry-wise to the vector $\beta^\top \tilde{X} \in \mathbb{R}^n$. At this point, note from (6.5) that in the unregularized case ($\gamma = 0$) one must have

$$\tilde{X} L'(\tilde{X}^\top \beta) = \mathbf{0}$$

so that, by considering the singular value decomposition of \tilde{X}

$$\tilde{X} = \mathbf{U} \Sigma \mathbf{V}^\top = \mathbf{U} [\mathbf{S} \quad \mathbf{0}] \begin{bmatrix} \mathbf{V}_1^\top \\ \mathbf{V}_2^\top \end{bmatrix}$$

for $\tilde{\mathbf{X}} \in \mathbb{R}^{p \times n}$ with $\Sigma \in \mathbb{R}^{p \times n}$, orthogonal $\mathbf{U} \in \mathbb{R}^{p \times p}$, $\mathbf{V} \in \mathbb{R}^{n \times n}$, diagonal $\mathbf{S} \in \mathbb{R}^{p \times p}$ and $\mathbf{V}_1 \in \mathbb{R}^{n \times p}$, $\mathbf{V}_2 \in \mathbb{R}^{n \times (n-p)}$ (recall the necessary $n > p$ condition when $\gamma = 0$ case), one must have

$$\mathbf{V}_1^T L'(\tilde{\mathbf{X}}^T \boldsymbol{\beta}) = \mathbf{0}.$$

That is, $L'(\tilde{\mathbf{X}}^T \boldsymbol{\beta})$ lies on the subspace spanned by the column vectors of \mathbf{V}_2 , i.e., there exists $\mathbf{a} \in \mathbb{R}^{n-p}$ for which

$$L'(\tilde{\mathbf{X}}^T \boldsymbol{\beta}) = \mathbf{V}_2 \mathbf{a}$$

and thus (6.22) further simplifies to

$$\max_L \frac{\mathbf{a}^T \mathbf{V}_2^T \mathbf{1}_n}{\|\mathbf{a}\|}$$

which attains the maximum if and only if \mathbf{a} is aligned to $\mathbf{V}_2^T \mathbf{1}_n$, i.e., $\mathbf{a} = a \mathbf{V}_2^T \mathbf{1}_n$ for some $a > 0$. This optimality condition is met with the square loss $L(t) = (t - 1)^2/2$.

Consequently, the surprising conclusion of this section is that, in the unregularized case, among all convex and smooth functions, the simplest square loss turns out to be optimal. In particular, it *uniformly* outperforms the maximum likelihood solution induced by the logistic loss, as it systematically reaches lower classification error. Perhaps even more surprisingly, a similar conclusion can be reached in the *regularized case*, where is more subtle to establish as one needs to consider the possible different levels of regularization for different losses; we refer the interested readers to [Mai and Liao, 2019] for more details.³

The analysis framework presented in this section is based on a “local” Taylor expansion of the loss L . This analysis however excludes the non differentiable hinge loss of the popular and important method of support vector machines (SVMs). The following section develops the intuitive ideas to handle this non-smooth implicit optimization case.

6.2 Large dimensional support vector machines

As one of most popular classification methods in the machine learning literature, the support vector machine (SVM), and in particular the hard-margin SVM, is based on the intuition of finding two separating hyperplanes with a maximum “margin” between them, so as to make robust future predictions.

Assuming the *linear separability* of the training set $\{(\mathbf{x}_i, y_i)\}_{i=1}^n$ with $y_i \in \{-1, +1\}$, the max margin classifier given by the solution to the hard-margin

³The (approximate) optimality of the square loss among all *convex* losses was recently established by Taheri et al. [2020c,a,b] on similar, yet formally different, models, showing the (somewhat unexpected) advantageous performance of the simple square loss in large dimensional problems.

SVM, is formulated as the following optimization problem

$$\begin{aligned} \arg \min_{\beta \in \mathbb{R}^p} \quad & \|\beta\|^2, \\ \text{s.t.} \quad & y_i \beta^\top \mathbf{x}_i \geq 1, \quad i \in \{1, \dots, n\}. \end{aligned} \quad (6.23)$$

It is interesting to note that, unlike for logistic regression where the training set *must not be* linearly separable as mentioned in Remark 6.1, the hard-margin SVM, on the contrary, assumes explicitly the *linear separability* of data.

Remark 6.2 (Connection between linear, logistic regression and SVM). *It has been shown in [Soudry et al., 2018] that, on a linearly separable dataset, the gradient descent method, when applied to minimize the unregularized logistic loss, converges in direction to the max margin classifier, that is, the solution to the hard-margin SVM.*

This can be understood by drawing an analogy to the linear (ridgeless) regression on the pair (\mathbf{X}, \mathbf{y}) for $\mathbf{X} \in \mathbb{R}^{p \times n}$, $\mathbf{y} \in \mathbb{R}^n$: in the under-parameterized $p < n$ regime there exists a unique solution $\beta = (\mathbf{X}\mathbf{X}^\top)^{-1}\mathbf{X}\mathbf{y}$ that minimizes the unregularized square loss $L(\beta) = \|\mathbf{y}^\top - \beta^\top \mathbf{X}\|^2$, when $\mathbf{X}\mathbf{X}^\top \in \mathbb{R}^{p \times p}$ is invertible; while for $p > n$ there are infinitely many solutions, and by running gradient descent (for almost all initializations) we obtain the minimal norm solution given by the Moore–Penrose pseudo-inverse $(\mathbf{X}\mathbf{X}^\top)^+ \mathbf{X}\mathbf{y} = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1}\mathbf{y}$ when $\mathbf{X}^\top \mathbf{X} \in \mathbb{R}^{n \times n}$ is invertible. In the linear regression case, having $p > n$ implies that there always exists $\beta_0 \in \mathbb{R}^p$ that can interpolate all the n training samples so that $\mathbf{X}^\top \beta_0 = \mathbf{y}$. Analogously, when the training set is linearly separable, there also exists β_0 such that $\text{sign}(\beta_0^\top \mathbf{x}_i) = y_i$ for $i \in \{1, \dots, n\}$ that perfectly classifies all training samples. In this respect, the hard-margin SVM solution is nothing more than the minimum norm solution (as per (6.23)) among all “interpolation” classifiers.

When the training set is *not linearly separable*, the “hard” constraints in (6.23) are not attainable. Nonetheless, by allowing residual small errors in the linear “separation”, the “soft-margin” alternative can be introduced

$$\begin{aligned} \arg \min_{\beta \in \mathbb{R}^p} \quad & \frac{1}{2} \|\beta\|^2 + \frac{\gamma}{n} \sum_{i=1}^n \varepsilon_i \\ \text{s.t.} \quad & y_i \beta^\top \mathbf{x}_i \geq 1 - \varepsilon_i, \quad \varepsilon_i \geq 0, \quad i \in \{1, \dots, n\} \end{aligned} \quad (6.24)$$

with some regularization parameter $\gamma > 0$, which is, as one may note, equivalent to solving the general empirical risk minimization problem in (6.1) with the hinge loss $L(t) = \max(0, 1 - t)$, by introducing the variable $\varepsilon_i = \max(0, 1 - y_i \beta^\top \mathbf{x}_i)$ which is the smallest nonnegative number satisfying $y_i \beta^\top \mathbf{x}_i \geq 1 - \varepsilon_i$.

Solving the associated Lagrangian dual, (6.24) can be further reduced to the

following simplified optimization problem

$$\begin{aligned} \max_{c_i, i \in \{1, \dots, n\}} \quad & \sum_{i=1}^n c_i - \frac{1}{2n} \sum_{i,j=1}^n c_i c_j \tilde{\mathbf{x}}_i^\top \tilde{\mathbf{x}}_j \\ \text{s.t.} \quad & \sum_{i=1}^n c_i y_i = 0, \quad 0 \leq c_i \leq \gamma, \quad i \in \{1, \dots, n\} \end{aligned} \quad (6.25)$$

where we use again the shortcut $\tilde{\mathbf{x}}_i \equiv y_i \mathbf{x}_i$ and assume for simplicity the symmetric Gaussian mixture

$$\mathcal{C}_1 : \mathbf{x}_i \sim \mathcal{N}(-\boldsymbol{\mu}, \mathbf{C}), \quad y_i = -1 \quad \mathcal{C}_2 : \mathbf{x}_i \sim \mathcal{N}(+\boldsymbol{\mu}, \mathbf{C}), \quad y_i = +1 \quad (6.26)$$

with class prior probability of 1/2 as in the previous section. The hard-margin solution, if it exists, can be retrieved by taking $\gamma \rightarrow \infty$. With the dual variables c_i , the SVM classifier $\boldsymbol{\beta}$ is given by

$$\boldsymbol{\beta} = \frac{1}{n} \sum_{i=1}^n c_i \tilde{\mathbf{x}}_i \quad (6.27)$$

which is very reminiscent of (6.5) in the previous section, with the dependence of c_i on the $\boldsymbol{\beta}$ and all $\tilde{\mathbf{x}}_i$'s seemingly more involved. Yet, the core idea remains the same: we need to “unwrap” this complex statistical dependence by introducing a series of self-consistent *nonlinear* equations as in (6.11) of the previous section, which allows for the evaluation of the statistics of $\boldsymbol{\beta}$.

Precisely, with the Karush–Kuhn–Tucker (KKT) conditions, the variable c_i satisfies the following constraints

$$\begin{cases} c_i = 0, & \text{for } \boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i > 1 \\ 0 < c_i < \gamma, & \text{for } \boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i = 1 \\ c_i = \gamma, & \text{for } \boldsymbol{\beta}^\top \tilde{\mathbf{x}}_i < 1 \end{cases} \quad (6.28)$$

which, by (6.27), can be further compactly rewritten as

$$c_i = f \left(\frac{1 - \frac{1}{n} \sum_{j \neq i} c_j \tilde{\mathbf{x}}_j^\top \tilde{\mathbf{x}}_i}{\|\tilde{\mathbf{x}}_i\|^2/n} \right) \equiv f \left(\frac{1 - \boldsymbol{\alpha}^\top \tilde{\mathbf{x}}_i}{\|\tilde{\mathbf{x}}_i\|^2/n} \right), \quad \boldsymbol{\alpha} = \frac{1}{n} \sum_{j \neq i} c_j \tilde{\mathbf{x}}_j \quad (6.29)$$

for $f(t) = \max(0, \min(t, \gamma))$. Note importantly here that $\boldsymbol{\alpha}$ is already a (first) “leave-one-out” version of the solution $\boldsymbol{\beta}$ in the sense that

$$\boldsymbol{\beta} = \boldsymbol{\alpha} + \frac{c_i}{n} \tilde{\mathbf{x}}_i \quad (6.30)$$

with the contribution from a *single* datum $c_i \tilde{\mathbf{x}}_i$ of negligible impact to $\boldsymbol{\beta}$.

Similar to the derivation in Section 6.1, we introduce a second (standard) “leave-one-out” solution $\boldsymbol{\beta}_{-i}$ by solving the optimization problem (6.24) for all training data except \mathbf{x}_i . As a consequence, we shall have

$$\boldsymbol{\beta}_{-i} = \frac{1}{n} \sum_{j \neq i}^n c_j^{-i} \tilde{\mathbf{x}}_j, \quad c_j^{-i} = f \left(\frac{1 - \frac{1}{n} \sum_{l \neq i, j} c_l^{-i} \tilde{\mathbf{x}}_l^\top \tilde{\mathbf{x}}_j}{\|\tilde{\mathbf{x}}_j\|^2/n} \right) \quad (6.31)$$

with c_j^{-i} the associated dual coefficients. Despite taking similar forms, α and β_{-i} crucially differ from each other in the fact that α depends on the \mathbf{x}_i (through the coefficients c_j), while β_{-i} not. Our objective is, again similar to Section 6.1, to derive the (asymptotic) relation between the dual coefficient c_j^{-i} and c_j , as well as α and β_{-i} (that should both be asymptotically “close to” β).

We start by writing, for all $j \neq i$,

$$c_j - c_j^{-i} = f\left(\frac{1 - \frac{1}{n} \sum_{l \neq i,j} c_l \tilde{\mathbf{x}}_l^\top \tilde{\mathbf{x}}_j - \frac{1}{n} c_i \tilde{\mathbf{x}}_i^\top \tilde{\mathbf{x}}_j}{\|\tilde{\mathbf{x}}_j\|^2/n}\right) - f\left(\frac{1 - \frac{1}{n} \sum_{l \neq i,j} c_l^{-i} \tilde{\mathbf{x}}_l^\top \tilde{\mathbf{x}}_j}{\|\tilde{\mathbf{x}}_j\|^2/n}\right).$$

Since $f(t) = \max(0, \min(t, \gamma))$, there exists $d \in [0, 1]$ such that for $t_1, t_2 \in \mathbb{R}$ we have $f(t_1) - f(t_2) = d(t_1 - t_2)$. Denote $d_j \in [0, 1]$ the constant that satisfies⁴

$$c_j - c_j^{-i} = d_j \left(\frac{-\frac{1}{n} \sum_{l \neq i,j} (c_l - c_l^{-i}) \tilde{\mathbf{x}}_l^\top \tilde{\mathbf{x}}_j - \frac{1}{n} c_i \tilde{\mathbf{x}}_i^\top \tilde{\mathbf{x}}_j}{\|\tilde{\mathbf{x}}_j\|^2/n} \right) \equiv [\Delta c]_j \quad (6.32)$$

so that, for $\Delta c \in \mathbb{R}^{n-1}$ the column vector composed of the differences $c_j - c_j^{-i}$ and $\mathbf{D}_{-i} \in \mathbb{R}^{(n-1) \times (n-1)}$ the diagonal matrix of all nonzero d_j 's, we have the following (matrix form) relation

$$\left(\frac{1}{n} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top \tilde{\mathbf{X}}_{-i} + \text{diag} \left(\frac{1}{n} \tilde{\mathbf{X}}_{-i}^\top \tilde{\mathbf{X}}_{-i} - \frac{1}{n} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top \tilde{\mathbf{X}}_{-i} \right) \right) \Delta c = -c_i \frac{1}{n} \mathbf{D}_{-i} \tilde{\mathbf{X}}_{-i}^\top \tilde{\mathbf{x}}_i$$

so that

$$\Delta c = -\frac{c_i}{n} \mathbf{M}_{-i}^{-1} \tilde{\mathbf{X}}_{-i}^\top \tilde{\mathbf{x}}_i, \quad \mathbf{M}_{-i} \equiv \frac{1}{n} \tilde{\mathbf{X}}_{-i}^\top \tilde{\mathbf{X}}_{-i} + \text{diag} \left((\mathbf{D}_{-i}^{-1} - \mathbf{I}_{n-1}) \frac{1}{n} \tilde{\mathbf{X}}_{-i}^\top \tilde{\mathbf{X}}_{-i} \right). \quad (6.33)$$

As a consequence, we deduce

$$\alpha - \beta_{-i} = \frac{1}{n} \sum_{j \neq i} (c_j - c_j^{-i}) \tilde{\mathbf{x}}_j = \frac{1}{n} \tilde{\mathbf{X}}_{-i} \Delta c = -\frac{c_i}{n^2} \tilde{\mathbf{X}}_{-i} \mathbf{M}_{-i}^{-1} \tilde{\mathbf{X}}_{-i}^\top \tilde{\mathbf{x}}_i \quad (6.34)$$

which again takes a similar form as (6.7) in the previous section. As such

$$\beta^\top \tilde{\mathbf{x}}_i = \alpha^\top \tilde{\mathbf{x}}_i + \frac{c_i}{n} \|\mathbf{x}_i\|^2 = \beta_{-i}^\top \tilde{\mathbf{x}}_i + c_i \tilde{\delta}_i, \quad c_i = f \left(\frac{1 - \beta_{-i}^\top \tilde{\mathbf{x}}_i}{\tilde{\delta}_i} \right) \quad (6.35)$$

where we recall $f(t) = \max(0, \min(t, \gamma))$ and denote the random variable

$$\tilde{\delta} \equiv \frac{1}{n} \tilde{\mathbf{x}}_i^\top \left(\mathbf{I}_p - \frac{1}{n} \tilde{\mathbf{X}}_{-i} \mathbf{M}_{-i}^{-1} \tilde{\mathbf{X}}_{-i}^\top \right) \tilde{\mathbf{x}}_i \quad (6.36)$$

⁴Here we observe a similar form as (6.6) in the previous section, the main difference being that, due to the non-differentiability of the hinge loss, the auxiliary variables d_j 's are introduced, the statistics of which remain to be determined.

which, as in (6.8) in the previous section, is expected to “converge” to a deterministic limit δ as $n, p \rightarrow \infty$, by the trace lemma, Lemma 2.11. As such, by (6.35) the random variable c_i is expected to behave as a nonlinear transformation of a Gaussian random variable (depending on the data statistics μ, \mathbf{C} and δ) in the large $n, p \rightarrow \infty$ limit (since $\beta_{-i}^\top \tilde{\mathbf{x}}_i$ is Gaussian conditioned on β_{-i} , the norm of which should converge to a limit). Nonetheless, the self-consistent equation of δ must involve the statistics of the d_j ’s that remain unknown for the moment.

To investigate the statistics of the d_j ’s as well as that of $\tilde{\delta}$, note that with a Taylor expansion-type argument, we have for $t \in \mathbb{R} \setminus \{0, \gamma\}$ that

$$f(t + \varepsilon) - f(t) = f'(t) \cdot \varepsilon + O(\varepsilon^2)$$

for $f'(t) = 1$ for $t \in (0, \gamma)$ and 0 otherwise. Therefore, by (6.32) we have that d_j is nonzero only when the associated c_j^{-i} lies in $(0, \gamma)$.

With (6.36) we thus conclude that

$$\tilde{\delta}_i = \frac{1}{n} \tilde{\mathbf{x}}_i^\top \left(\mathbf{I}_p - \tilde{\mathbf{X}}_{-i} (\tilde{\mathbf{X}}_{-i,S}^\top \tilde{\mathbf{X}}_{-i,S})^{-1} \tilde{\mathbf{X}}_{-i}^\top \right) \tilde{\mathbf{x}}_i \quad (6.37)$$

where the j -th column of $\tilde{\mathbf{X}}_{-i,S}$ is $\tilde{\mathbf{x}}_j$ such that the associated coefficient c_j^{-i} lies in $(0, \gamma)$ and $\tilde{\delta} \xrightarrow{a.s.} \delta$ for δ the unique solution to

$$\delta = \frac{1}{n} \text{tr } \mathbf{C} \left(\mathbf{I}_p + \frac{\mathbb{E}[c \cdot 1_{(0,\gamma)}]}{\delta} \mathbf{C} \right)^{-1}, \quad c \sim f \left(\frac{1-r}{\delta} \right) \quad (6.38)$$

for $r \sim \mathcal{N}(M, \sigma^2)$ with

$$M = \mathbb{E}[\beta]^\top \mu, \quad \sigma^2 = \text{tr}(\mathbf{C} \mathbb{E}[\beta \beta^\top]) \quad (6.39)$$

which presents a close analogy to (6.9) and (6.13) in the previous section, respectively.

To close the loop, it remains, as in the previous section, to express the statistics of β via (6.27) as a function of those of c_i and μ, \mathbf{C} . This leads to the following result on the (asymptotically) statistical behavior of SVM, as an extension of Theorem 6.1 in the previous section.

[ref below to be updated](#)

Theorem 6.2 (Asymptotic behavior of β for SVM, [Mai, 2019, Theorem 5.3.1]). *For $\max\{\|\mu\|, \|\mathbf{C}\|, \|\mathbf{C}^{-1}\|\} = O(1)$ and β the unique solution to (6.24), we have, as $n, p \rightarrow \infty$,*

$$\|\beta - \tilde{\beta}\| \rightarrow 0, \quad \left(\mathbf{I}_p + \frac{\mathbb{E}[c \cdot 1_{(0,\gamma)}]}{\delta} \mathbf{C} \right) \tilde{\beta} \sim \mathcal{N}(\mathbb{E}[c] \mu, \mathbb{E}[c^2] \mathbf{C} / n)$$

where $c \sim f((1-r)/\delta)$ for $f(t) = \max(0, \min(t, \gamma))$ and $r \sim \mathcal{N}(M, \sigma^2)$ for (M, σ^2) solution to

$$M = \mathbb{E}[\tilde{\beta}]^\top \mu, \quad \sigma^2 = \text{tr}(\mathbf{C} \mathbb{E}[\tilde{\beta} \tilde{\beta}^\top]) \quad (6.40)$$

with δ the unique positive solution to

$$\delta = \frac{1}{n} \operatorname{tr} \mathbf{C} \left(\mathbf{I}_p + \frac{\mathbb{E}[c \cdot 1_{(0,\gamma)}]}{\delta} \mathbf{C} \right)^{-1}.$$

One may wish to take the limit of $\gamma \rightarrow \infty$ in the above theorem to retrieve the solution of hard-margin SVM. However, attention must be paid here since, as in Remark 6.1 for logistic regression, a *sharp* phase transition on the ratio p/n also exists in the case of hard-margin SVM, see for instance [Kammoun and Alouini, 2021] for a more detailed discussion on this point.

Analogy to (6.17) and (6.18), the asymptotic test classification error rates of SVM can also be given, for a new datum $\tilde{\mathbf{x}}$, by

$$\mathbb{P}(\boldsymbol{\beta}^T \tilde{\mathbf{x}} < 0) - Q(M/\sigma) \rightarrow 0 \quad (6.41)$$

with $Q(\cdot)$ the Gaussian Q -function and similarly the asymptotic training classification error

$$\mathbb{P}(\boldsymbol{\beta}^T \tilde{\mathbf{x}}_i < 0) - \mathbb{P}(r + c\delta < 0) \rightarrow 0 \quad (6.42)$$

for $r \sim \mathcal{N}(M, \sigma^2)$ and $c \sim f((1-r)/\delta)$ defined in Theorem 6.2.

*** it would be good if we could find a simple argument to show that least-squares outperforms SVM***

In this section, we provided the heuristic derivation of the asymptotic performance of the soft-margin SVM classifier defined in (6.24) arising from the minimization of the non smooth hinge loss $L(t) = \max(0, 1 - t)$, with a focus on the similarities and differences from the large dimensional analysis of, say, the logistic regression, in Section 6.1 where the loss is at least three-times differentiable.

Recall from the discussions at the end of Section 6.1.3 that the simple square loss is proven to be optimal in the classification of the Gaussian mixture $\mathcal{N}(\pm \boldsymbol{\mu}, \mathbf{C})$ and beyond [Taheri et al., 2020c,a,b]. The theoretical analyses in these two sections unveil the surprising and counterintuitive facts that, for large dimensions problems, classical empirical risk minimization solutions may fail to exist (e.g., the maximum likelihood solution discussed in Remark 6.1 or the hard-margin SVM), and if they exist, may fail to meet the performance of the simple square loss. These results challenge the conventional wisdom from classical statistical learning theory where the data dimension p is considered negligibly small with respect to the sample size n , in which case the maximum likelihood solution is believed to produce the minimum regression error and the hinge loss is proven to be optimal in a classification context [Rosasco et al., 2004]. The theoretical analysis proposed in this section, although performed on relatively simple statistical models, opens the door to a renewed understanding of machine learning basics, starting with the long-proclaimed SVM method, when considering large (and even moderately large) dimensional data.

6.3 Concluding remarks

After covering the analysis of machine learning algorithms involving optimization methods with *explicit* solutions (such as least squares regression or spectral methods), this chapter pushed into the large dimensional analysis of algorithms with solution expressed as the (*unique*) *minimum of a convex optimization problem*. Unlike in the explicit case where the structure of dependence within the solution (such as between the entries of a least squares regression vector) more-or-less easily relates to a functional of the underlying random matrix model, the main difficulty involved in the implicit case lies in a possibly very intricate structure of dependence. The chapter proposed to “break” this dependence by exploiting a “leave-one-out” approach, by which an arbitrary (large dimensional) data vector is isolated from the dataset to perturb the solution in a controlled manner.

Other options in the large dimensional statistics literature exist which may tackle the same problem in a different manner. Among these, we find

- the “double leave-one-out” approach adopted early on in [El Karoui et al., 2013], in order to study the statistical behavior of a family of (robust) M-estimators. This approach hinges on the intuition that, as both dimensions n, p grow, the “leave-one-sample-out” approach, popular in asymptotic (large- n alone) statistics and upon which we based our derivations in the present chapter, could be extended into a double “leave-one-sample-out” and “leave-one-feature-out” approach. Specifically, having reached step (6.11) in Section 6.1, in order to evaluate the statistics of β , the double leave-one-out method would not decompose $\tilde{\mathbf{x}}_i$ as in (6.12) but rather extract an arbitrary entry j from each vector $\tilde{\mathbf{x}}_i$ (and accordingly from β) to understand the statistics of the entry $[\beta]_j$. However, this approach is only valid (or easily adaptable) when the $[\beta]_j$ ’s are statistically exchangeable or linearly related, so essentially for “*unstructured*” feature vectors with *no informative* pattern. This is in particular not directly applicable to data mixture models as simple as $\mathcal{N}(\pm\mu, \mathbf{C})$ discussed at length in this chapter;
- the convex Gaussian min-max theorem (CGMT), first introduced by Gordon [1985] and then largely popularized by Thrampoulidis et al. [2018]. The idea of the CGMT lies in an opportunistic result relating two “Gaussian” min-max optimization problems:

$$\Phi(\mathbf{G}) = \min_{\mathbf{v}} \max_{\mathbf{u}} \{ \mathbf{u}^T \mathbf{G} \mathbf{v} + \psi(\mathbf{v}, \mathbf{u}) \} \quad (6.43)$$

$$\phi(\mathbf{g}, \mathbf{h}) = \min_{\mathbf{v}} \max_{\mathbf{u}} \{ \|\mathbf{v}\| \cdot \mathbf{g}^T \mathbf{u} + \|\mathbf{u}\| \cdot \mathbf{h}^T \mathbf{v} + \psi(\mathbf{v}, \mathbf{u}) \} \quad (6.44)$$

where $\mathbf{G} \in \mathbb{R}^{n \times p}$, $\mathbf{g} \in \mathbb{R}^n$ and $\mathbf{h} \in \mathbb{R}^p$ are matrix and vectors composed of i.i.d. $\mathcal{N}(0, 1)$ entries, and $\psi : \mathbb{R}^p \times \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous and convex-concave, and for which it can be shown that, for any $x, t \in \mathbb{R}$,

$$\mathbb{P}(|\Phi(\mathbf{G}) - x| > t) \leq 2\mathbb{P}(|\phi(\mathbf{g}, \mathbf{h}) - x| > t).$$

Evidently, the second optimization being simpler than the first, if $\phi(\mathbf{g}, \mathbf{h})$ concentrates around x as $n, p \rightarrow \infty$, then so does $\Phi(\mathbf{G})$. The idea to apply the above CGMT inequality is then to express the studied optimization problem under the form $\Phi(\mathbf{G})$. This is usually performed by introducing an additional slack variable \mathbf{u} to enforce the constraints. For instance, in the hard-margin SVM setting, one may rewrite (6.23) under the form

$$\begin{aligned} & \min_{\boldsymbol{\beta}} \max_{\mathbf{u} \leq \mathbf{0}} \left\{ \|\boldsymbol{\beta}\|^2 + \frac{1}{n} \sum_{i=1}^n u_i (y_i \boldsymbol{\beta}^\top \mathbf{x}_i - 1) \right\} \\ &= \min_{\boldsymbol{\beta}} \max_{\mathbf{u} \leq \mathbf{0}} \left\{ \frac{1}{n} \mathbf{u}^\top \tilde{\mathbf{X}}^\top \boldsymbol{\beta} + \|\boldsymbol{\beta}\|^2 - \frac{1}{n} \mathbf{u}^\top \mathbf{1}_n \right\} \end{aligned}$$

for $\tilde{\mathbf{X}} = [y_1 \mathbf{x}_1, \dots, y_n \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ and $\mathbf{u} \leq \mathbf{0}$ understood in an entry-wise sense, in which we recognize the form $\mathbf{u}^\top \mathbf{G} \mathbf{v}$ of $\Phi(\mathbf{G})$ in the term $\frac{1}{n} \mathbf{u}^\top \tilde{\mathbf{X}}^\top \boldsymbol{\beta}$. A thorough analysis of SVM and logistic regression under the CGMT framework is performed in [Deng et al., 2019].

As such, the CGMT approach does not solve the “zero-gradient” equation implied by the optimization problem, but rather rewrites the optimization problem itself to make it simpler to solve. Specifically, the major interest here is to “break” the bilinear form $\mathbf{u}^\top \mathbf{G} \mathbf{v}$ in the primal optimization (6.43) into the sum of two linear optimizations $\|\mathbf{v}\| \mathbf{g}^\top \mathbf{u} + \|\mathbf{u}\| \mathbf{h}^\top \mathbf{v}$ (conditionally on the norms of \mathbf{v} and then \mathbf{u}) in (6.44). This also interestingly breaks the very random matrix structure of the original problem, turning the random Gaussian matrix \mathbf{G} into the random Gaussian vectors \mathbf{g} and \mathbf{h} . The method however strongly relies on the CGMT inequality and thus requires \mathbf{G} to have strictly i.i.d. $\mathcal{N}(0, 1)$ entries. Any deviation from this setting (for instance by adding correlation structures within \mathbf{G}) may pose new challenges to the CGMT framework;

- the approximate message passing (AMP) and the state evolution techniques [Donoho and Montanari, 2016]. Unlike the previous methods, AMP proposes to directly solve the optimization problem by following the dynamics of an iterative fixed-point algorithm ultimately converging to the sought-for solution. The method is reminiscent and indeed strongly related with the message passing (also known as belief propagation [Pearl, 1986]) approach from statistical physics, yet enjoys a mathematical sound framework (unlike belief propagation). The AMP approach however requires the introduction of many additional tools which would drive us too far from our present topic.

Having in hand numerous tools to handle convex optimization problems in machine learning, a natural next step would be to deal with *non-convex optimization problems*. These problems are deemed hard to solve (and consequently to study) due to the possible existence of multiple, even perhaps infinity many, critical points which can all be candidate solutions of the problem. In practice,

initialization may play a crucial role in gradient-based learning and one may wish to start the algorithm at a “good guess” starting point, sufficient close to the sought-for (global) optimal solution. In this respect, RMT analyses provide efficient *spectral-based initialization* schemes widely used in many non-convex problems, such as phase retrieval [Fienup, 1982], matrix completion [Keshavan et al., 2010], low-rank matrix recovery [Jain et al., 2013], blind deconvolution [Lee et al., 2017], sparse coding [Arora et al., 2015], to name a few; a detailed example on the problem of phase retrieval is developed as a practical course exercise in the next section. Yet, many non-convex optimization problems need to be confronted with directly, without resorting to first approximations. This is the case notably of neural network training, and particularly in the large dimensional setting of deep learning, as well as of other popular, yet still ill-understood, machine learning methods such as stochastic neighborhood embedding (SNE [Hinton and Roweis, 2003] or its popular t-SNE variant [Maaten and Hinton, 2008]). For these challenging problems, the strong hope is that, while as $n, p \rightarrow \infty$, the number of critical points in these problems typically grows at a fast (possibly exponential) rate, these critical points share numerous symmetries and their associated performance (the “depth” of a given local minimum for instance) often reduce to a few states. This is indeed to be expected as one of the strong arguments in favor of the observed very stable behavior of deep neural networks [Choromanska et al., 2015]: irrespective of the learning initialization point, gradient descent learning in these highly non-convex optimization problems is expected to converge to a local minimum with essentially the same performance level as the vast majority of the local minima. The article [Choromanska et al., 2015] provides tentative theoretical insights in this direction, yet for an overly simplified version of a deep learning network (modeled as a mere Gaussian random field), using the Kac-Rice formula (which assesses the number of critical points of a problem) in conjunction to random matrix arguments.

6.4 Practical course material

Practical Lecture Material 6 (Phase retrieval). *The object of phase retrieval is to recover an unknown (deterministic) signal $\mathbf{a} \in \mathbb{R}^p$ (say with $\|\mathbf{a}\| = 1$) from magnitude measurements of the type $y_i = (\mathbf{a}^\top \mathbf{x}_i)^2$, for i.i.d. Gaussian sensing vectors $\mathbf{x}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, $i \in \{1, \dots, n\}$. One popular algorithm to solve this problem is the so-called “Wirtinger Flow algorithm” [Candès et al., 2015] which comes in two steps: (i) a careful initialization obtained by means of a spectral method, and (ii) gradient descent updates to “fine-tune” this initial estimate on a target cost function. Due to the non-convex nature of the underlying problem, the initialization step (i) is of crucial significance for good performance. Candès et al. [2015] proposed to use the dominant eigenvector of the sample covariance-type matrix $\mathbf{H} = \frac{1}{n} \sum_{i=1}^n y_i \mathbf{x}_i \mathbf{x}_i^\top$ as a first estimate. The objective of this exercise is to understand the relevance of this idea through the analysis of the spectral property of \mathbf{H} .*

To this end, first decompose (the columns of) $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ into the

sum of a component aligned to \mathbf{a} and a component orthogonal to \mathbf{a} , and show that \mathbf{X} can be expressed under the form $\mathbf{X} = \mathbf{aa}^\top \mathbf{X} + \mathbf{X}^\perp$ with $\mathbf{X}^\perp = (\mathbf{I}_p - \mathbf{aa}^\top) \mathbf{X} \in \mathbb{R}^{p \times n}$; confirm in particular that $\mathbf{a}^\top \mathbf{X}^\perp = \mathbf{0}$ and, more importantly, that $\mathbf{X}^\top \mathbf{a} = \mathbf{v} \in \mathbb{R}^n$ and \mathbf{X}^\perp are jointly Gaussian and independent of each other.

Based on the above decomposition, show, using Lemma 2.9, that the limiting spectral measure of $\frac{1}{n} \mathbf{X}^\perp \mathbf{D}(\mathbf{X}^\perp)^\top$ for diagonal $\mathbf{D} = \text{diag}\{y_i\}_{i=1}^n$, if it exists, coincides with that of the original $\mathbf{H} = \frac{1}{n} \mathbf{X} \mathbf{D} \mathbf{X}^\top = \frac{1}{n} \sum_{i=1}^n y_i \mathbf{x}_i \mathbf{x}_i^\top$.

With this decomposition in mind, and using similar steps as in the proof of Theorem 2.6, determine the limiting spectrum of $\frac{1}{n} \mathbf{X} \mathbf{D} \mathbf{X}^\top$ via its Stieltjes transform. Show that it is asymptotically close to that of $\frac{1}{n} \sum_{i=1}^n \tau_i \mathbf{x}_i \mathbf{x}_i^\top$ for i.i.d. τ_i following a chi-square distribution with one degree of freedom which is independent of \mathbf{x}_i . In other words, the dependence between $y_i = (\mathbf{x}_i^\top \mathbf{a})^2$ and \mathbf{x}_i does not “contribute” to the limiting spectral measure of $\frac{1}{n} \mathbf{X} \mathbf{D} \mathbf{X}^\top$.

Recalling that a chi-square distribution with one degree of freedom admits the density

$$\nu(dt) = \frac{1}{\sqrt{2\Gamma(1/2)}} \frac{e^{-\frac{t}{2}}}{\sqrt{t}}, \quad t > 0 \quad (6.45)$$

and has unbounded support. Conclude, using the argument in Section 2.3.1, that the support of limiting spectrum of $\mathbf{H} = \frac{1}{n} \mathbf{X} \mathbf{D} \mathbf{X}^\top$ is also unbounded and that it may not be possible to have (an almost sure) isolated eigenvalue “jumping out” in the large n, p limit.

Further consider the “truncated” model $\frac{1}{n} \mathbf{X} f(\mathbf{D}) \mathbf{X}^\top = \frac{1}{n} \sum_{i=1}^n f(y_i) \mathbf{x}_i \mathbf{x}_i^\top$ for some (entry-wise) processing function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ that is of bounded support, for instance $f(t) = 1_{t \leq \tau}$ for some (predefined) threshold $\tau > 0$. This is indeed the trimming strategy proposed in [Chen and Candès, 2017] and was shown there to play a crucial role in the success of the algorithm in the large n, p regime. Again with the decomposition used above and Theorem 2.6, show that the resolvent $\mathbf{Q} = (\frac{1}{n} \mathbf{X}^\perp f(\mathbf{D})(\mathbf{X}^\perp)^\top - z \mathbf{I}_p)^{-1}$ admits the following deterministic equivalents

$$\mathbf{Q} \leftrightarrow m(z)(\mathbf{I}_p - \mathbf{aa}^\top) - \frac{1}{z} \mathbf{aa}^\top,$$

for $(z, m(z))$ the unique solution in $\mathcal{Z}(\mathbb{C} \setminus \mathbb{R}^+)$ to

$$m(z) = \left(-z + \frac{1}{n} \text{tr } f(\mathbf{D})(\mathbf{I}_n + cm(z)f(\mathbf{D}))^{-1} \right)^{-1}$$

or equivalently

$$m(z) = \left(-z + \int \frac{f(t)\nu(dt)}{1 + cm(z)f(t)} \right)^{-1}$$

for ν the chi-square distribution with one degree of freedom (recall that $\|\mathbf{a}\| = 1$ and $\mathbb{E}[\mathbf{x}^\perp(\mathbf{x}^\perp)^\top] = \mathbf{I}_p - \mathbf{aa}^\top$). With the deterministic equivalents derived above, following the proof of Theorem 2.13, solve $\det(\frac{1}{n} \mathbf{X} f(\mathbf{D}) \mathbf{X}^\top - \lambda \mathbf{I}_p) = 0$ to find an

(hypothetically) isolated spike λ and check that the associated $m(\lambda)$ then satisfies

$$-\frac{1}{m(\lambda)} + \int \frac{f(t)\nu(dt)}{1+cf(t)m(\lambda)} = \int \frac{tf(t)\nu(dt)}{1+cf(t)m(\lambda)} \quad (6.46)$$

with $\nu(dt)$ defined in (6.45). Determine then μ , the limiting spectral support of $\frac{1}{n}\mathbf{X}^\perp f(\mathbf{D})(\mathbf{X}^\perp)^\top$ (and thus of $\frac{1}{n}\mathbf{X}f(\mathbf{D})\mathbf{X}^\top$), with the help of the functional inverse

$$x(m) = -\frac{1}{m} + \int \frac{f(t)\nu(dt)}{1+cmf(t)}$$

introduced in Section 2.3.1. Conclude on the (phase transition) condition for the spike λ to exist, using the sign of the derivative $x'(m_*)$, with m_* the solution to (6.46).

Denote the shortcut $\alpha \equiv -\frac{1}{cm}$, $\alpha_* \equiv -\frac{1}{cm_*}$ and

$$x(m) = \psi_c \left(\alpha \equiv -\frac{1}{cm} \right) = \alpha \left(c + \int \frac{f(t)\nu(dt)}{\alpha - f(t)} \right)$$

as well as

$$\phi(\alpha) = \alpha \int \frac{tf(t)\nu(dt)}{\alpha - f(t)}$$

so that (6.46) can be compactly rewritten as $\phi(\alpha_*) = \psi_c(\alpha_*)$.

Check that, on the interval $\alpha \in (\tau, \infty)$ (recall τ the upper bound of the truncation function f , i.e., $f(\cdot) \leq \tau$), $\phi(\alpha)$ is a non-increasing function and $\psi_c(\alpha)$ is a convex function, attaining its unique minimum at $\bar{\alpha}$ that satisfies

$$\psi'_c(\bar{\alpha}) = 0 \Leftrightarrow \int \frac{f^2(t)\nu(dt)}{(\bar{\alpha} - f(t))^2} = c.$$

Check that the phase transition condition on the sign of $x'(m_*)$ derived above is equivalent to

$$\begin{cases} \phi(\bar{\alpha}) > \psi_c(\bar{\alpha}) \Leftrightarrow \bar{\alpha} < \alpha_* \text{ and } \psi'_c(\alpha_*) > 0; \\ \phi(\bar{\alpha}) \leq \psi_c(\bar{\alpha}) \Leftrightarrow \bar{\alpha} \geq \alpha_* \text{ and } \psi'_c(\alpha_*) \leq 0. \end{cases}$$

As a consequence, in pursuit of an optimal design for the truncation function $f(\cdot)$ with maximum phase transition threshold c_{th} , it suffices to find $f(\cdot)$ such that

$$\int \frac{f^2(t)\nu(dt)}{(\bar{\alpha} - f(t))^2} = c, \quad \int \frac{f(t)(t-1)\nu(dt)}{\bar{\alpha} - f(t)} > c \quad (6.47)$$

holds for a maximal value of c . Using the Cauchy-Schwarz inequality, show that we must then have

$$c^2 \leq 2c$$

with equality if and only if $\int \frac{f^2(t)\nu(dt)}{(\bar{\alpha} - f(t))^2} = \int (t-1)^2\nu(dt)$. Deduce that the optimal phase transition threshold is $c_{th} = 2$, in the sense that there is (almost surely)

no spike in the spectrum of $\frac{1}{n}\mathbf{X}f(\mathbf{D})\mathbf{X}^\top$ so long that $c < c_{th} = 2$. Conclude then that the associated optimal truncation function is therefore given by

$$f(t) = \frac{\max(t, 0) - 1}{\max(t, 0) + \sqrt{2/c} - 1} \quad (6.48)$$

such that $\frac{f(t)}{1-f(t)} - (t-1) \rightarrow 0$ as $c \rightarrow c_{th} = 2$. Confirm the theoretical findings above numerically as in Figure 6.4.

We refer the interested readers to [Lu and Li, 2019] for the asymptotic behavior of the associated isolated eigenvector and [Mondelli and Montanari, 2019] for the complex sensing matrix case.




Figure 6.4: Empirical alignment $(\mathbf{a}^\top \hat{\mathbf{u}})^2$ for $\hat{\mathbf{u}}$ the dominant eigenvector of $\frac{1}{n}\mathbf{X}f(\mathbf{D})\mathbf{X}^\top$ as a function of the ratio p/n , for different processing functions f with $p = 512$ and $\mathbf{a} = [-\mathbf{1}_{p/2}, \mathbf{1}_{p/2}]/\sqrt{p}$. Results averaged over 50 runs. [\[Link to code\]](#)

Chapter 7

Community Detection on Graphs

In the previous chapters, our attention has been long cast on numerous applications involving the sample covariance matrix of the type $\mathbf{XX}^\top/n \in \mathbb{R}^{p \times p}$ for some random matrix $\mathbf{X} \in \mathbb{R}^{p \times n}$ following a certain statistical model, as well as involving the quite related Gram matrix $\mathbf{X}^\top\mathbf{X}/n \in \mathbb{R}^{n \times n}$ and kernel matrices $f(\mathbf{X}^\top\mathbf{X}/n)$ (with f applied here entry-wise). The starting point of the asymptotic analysis of machine learning algorithms for most of these models is the Marčenko-Pastur law (Theorem 2.4) and its various generalizations (e.g., Theorem 2.6).

When it comes to studying the statistical behavior of randomly generated graphs and networks, starting with the so-called Erdős-Rényi graph, which randomly and independently draws links between each pair of nodes in the graph according to a Bernoulli law, the related random matrix model will be instead a Wigner matrix (for undirected graphs) and theoretical analyses will rely instead on Wigner's semicircle law (Theorem 2.5) and its variations (e.g., Theorem 2.9).

In this chapter, we will be particularly interested in the question of *community detection* on large dimensional and *dense* undirected and unweighted n -node graphs. By unweighted, we mean that an edge between node i and node j carries a unit weight 1, and zero otherwise. By undirected, we mean that, if node i connects to node j , then node j connects to node i , which in particular implies that the *adjacency matrix* $\mathbf{A} \in \{0, 1\}^{n \times n}$ of the graph is symmetric (i.e., $\mathbf{A}_{ij} = \mathbf{A}_{ji}$). By dense graphs, we mean graphs for which the typical number of neighbors of each node scales proportionally to the graph size n as $n \rightarrow \infty$. This will cover the main Section 7.1 of this chapter. A few notes and remarks on the more involved case of *sparse* graphs, for which each node instead has $O(1)$ neighbors, will be made in Section 7.2.

7.1 Community detection in dense graphs

7.1.1 The stochastic block model

7.1.1.1 Erdős-Rényi random graphs and the modularity matrix

The adjacency matrix. The most natural random undirected graph is the Erdős-Rényi graph, defined by the fact that its adjacency matrix $\mathbf{A} \in \{0, 1\}^{n \times n}$ is composed, up to symmetry ($\mathbf{A}_{ij} = \mathbf{A}_{ji}$) and to a null diagonal ($\mathbf{A}_{ii} = 0$), of *i.i.d.* Bernoulli entries: for all $i \neq j$,

$$\mathbf{A}_{ij} = \mathbf{A}_{ji} \sim \text{Bern}(p)$$

where $p \in (0, 1)$. To ensure that the graph is dense, we demand that, for each i , $\sum_j \mathbf{A}_{ij} = O(n)$, which implies that $p = O(1)$ with respect to n . Note that, unlike in previous sections, p denotes here the Bernoulli parameter rather than a vector dimension: no confusion will arise in this section where the only large dimension is the size n of the graph.

This setting implies that, for all $i < j$, the \mathbf{A}_{ij} are independent with $\mathbb{E}[\mathbf{A}_{ij}] = p$ and $\text{Var}[\mathbf{A}_{ij}] = p(1-p)$. In particular, by the central limit theorem, the degree d_i of node i , satisfies

$$d_i \equiv \sum_{j=1}^n \mathbf{A}_{ij} = np + \sqrt{p(1-p)n} \cdot (\mathcal{N}(0, 1) + o(1)).$$

and the average degree d_i/n converges almost surely to p .

Writing in matrix form

$$\mathbf{A} = \mathbb{E}[\mathbf{A}] + \sqrt{p(1-p)}\mathbf{X} - \text{diag}(\cdot) = p(\mathbf{1}_n \mathbf{1}_n^\top - \mathbf{I}_n) + \sqrt{p(1-p)}(\mathbf{X} - \text{diag}(\mathbf{X})) \quad (7.1)$$

where $\mathbf{X} \in \mathbb{R}^{n \times n}$ has independent entries of zero mean and unit variance and $\text{diag}(\mathbf{X})$ is the diagonal matrix containing only the diagonal entries of \mathbf{X} , we find that

$$\frac{\mathbf{A}}{\sqrt{n}} = \frac{p}{\sqrt{n}} \mathbf{1}_n \mathbf{1}_n^\top + \frac{\sqrt{p(1-p)}}{\sqrt{n}} \mathbf{X} + O_{\|\cdot\|}(n^{-\frac{1}{2}})$$

is a rank-one perturbation of a rescaled Wigner matrix (where $O_{\|\cdot\|}(\cdot)$ is in probability). Theorem 2.5 therefore applies and we have

- in the first order, $\frac{1}{n}\mathbf{A} = \frac{p}{n}\mathbf{1}_n \mathbf{1}_n^\top + O_{\|\cdot\|}(n^{-1/2})$ is essentially a unit-rank matrix with eigen-direction $\mathbf{1}_n$ and eigenvalue p .
- the limiting spectral measure of $\frac{1}{\sqrt{n}}\mathbf{A}$ is a semicircle law scaled by $\sqrt{p(1-p)}$, so in particular supported on $[-2\sqrt{p(1-p)}, 2\sqrt{p(1-p)}]$.

The modularity matrix. To avoid the technically problematic, and practically irrelevant, largely dominant $\frac{p}{\sqrt{n}} \mathbf{1}_n \mathbf{1}_n^\top$ matrix in \mathbf{A} , it is customary to rather work with the so-called *modularity* matrix

$$\mathbf{B} = \frac{1}{\sqrt{n}} \left(\mathbf{A} - \frac{\mathbf{d}\mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n} \right)$$

introduced in [Newman, 2006], where $\mathbf{d} = [d_1, \dots, d_n]^\top = \mathbf{A}\mathbf{1}_n$ is the degree vector. The advantage of working with \mathbf{B} versus \mathbf{A} is that $\mathbf{B}\mathbf{1}_n = \mathbf{0}$. This eliminates the dominant contribution of the vector $\mathbf{1}_n$. However, as a negative side effect, the matrix $\mathbf{A} - \frac{\mathbf{d}\mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n}$ is the summation of two strongly dependent random matrices. Let us try to decipher this dependence and study the spectrum of \mathbf{B} for n large.

First, the fact that $\mathbf{d} = \mathbf{A}\mathbf{1}_n$ gives

$$\mathbf{d} = pn\mathbf{1}_n + \sqrt{p(1-p)}\mathbf{X}\mathbf{1}_n + O(1)$$

where $O(1)$ is understood in an entry-wise sense, from which it follows that

$$\begin{aligned} \mathbf{d}^\top \mathbf{1}_n &= pn^2 + O(n) \\ \mathbf{d}\mathbf{d}^\top &= p^2 n^2 \mathbf{1}_n \mathbf{1}_n^\top + pn\sqrt{p(1-p)}(\mathbf{X}\mathbf{1}_n \mathbf{1}_n^\top + \mathbf{1}_n \mathbf{1}_n^\top \mathbf{X}) + O_{\|\cdot\|}(n^2) \end{aligned}$$

so that

$$\frac{1}{\sqrt{n}} \frac{\mathbf{d}\mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n} = p \frac{\mathbf{1}_n \mathbf{1}_n^\top}{\sqrt{n}} + \sqrt{p(1-p)} \frac{\mathbf{1}_n \mathbf{1}_n^\top \mathbf{X} + \mathbf{X} \mathbf{1}_n \mathbf{1}_n^\top}{n\sqrt{n}} + O_{\|\cdot\|}(n^{-\frac{1}{2}}).$$

Thus, we obtain

$$\frac{\mathbf{B}}{\sqrt{p(1-p)}} = \frac{\mathbf{X}}{\sqrt{n}} - \frac{\mathbf{1}_n \mathbf{1}_n^\top \mathbf{X} + \mathbf{X} \mathbf{1}_n \mathbf{1}_n^\top}{n\sqrt{n}} + O_{\|\cdot\|}(n^{-\frac{1}{2}}). \quad (7.2)$$

As a consequence, the modularity matrix \mathbf{B} may be asymptotically seen as a rank-two perturbation of a random Wigner matrix \mathbf{X}/\sqrt{n} . Solving $\det(\mathbf{B} - \lambda \mathbf{I}_n) = 0$ for $\lambda \notin [-2\sqrt{p(1-p)}, 2\sqrt{p(1-p)}]$ (the support of the limiting spectral measure of \mathbf{B}) reveals that there asymptotically exists *no* solution. As such, as one would have expected, all the eigenvalues of \mathbf{B} are compactly supported within the limiting semicircle support. A detailed derivation will be provided in the next section for the more interesting stochastic block model.

7.1.1.2 From Erdős-Rényi to the SBM

In order to account for the presence of communities of nodes in the graphs, we introduce now the stochastic block model (SBM) by assuming the possibility for the Bernoulli parameter of \mathbf{A}_{ij} to depend on the pair of nodes (i, j) . Specifically, letting $\mathcal{C}_1, \dots, \mathcal{C}_k$ be k communities of cardinalities $n_a \equiv |\mathcal{C}_a|$, we define $\mathbf{C} \in$

$\mathbb{R}^{k \times k}$ the matrix of Bernoulli parameters such that, if node i belongs to class \mathcal{C}_a and node $j \neq i$ belongs to class \mathcal{C}_b with $a, b \in \{1, \dots, k\}$,

$$\mathbf{A}_{ij} = \mathbf{A}_{ji} \sim \text{Bern}(\mathbf{C}_{ab}).$$

We further consider that all classes are of “large size” in the sense that $n_a/n \rightarrow c_a \in (0, 1)$ as $n \rightarrow \infty$.

As in the case of spectral clustering discussed in Chapter 4, in order to avoid trivial scenarios in the large- n asymptotics, a careful control of the differences between the elements of the matrix \mathbf{C} is needed. As we shall see next, the proper normalization is such that

$$[\mathbf{C}]_{ab} = p \left(1 + \frac{[\mathbf{M}]_{ab}}{\sqrt{n}} \right) \quad (7.3)$$

for $\mathbf{M} \in \mathbb{R}^{k \times k}$ a deterministic matrix, independent of n , and $p \in (0, 1)$ as above, also independent of n . That is, as n increases, communities with Bernoulli parameter differences scaling as $1/\sqrt{n}$ can still be distinguished in SBM.

Following the same analysis as above, it follows that

$$\mathbb{E}[\mathbf{A}] = p \left(\mathbf{1}_n \mathbf{1}_n^\top + \frac{1}{\sqrt{n}} \mathbf{J} \mathbf{M} \mathbf{J}^\top \right) \quad (7.4)$$

where, as in the previous chapters, we defined $\mathbf{J} = [\mathbf{j}_1, \dots, \mathbf{j}_k] \in \mathbb{R}^{n \times k}$ for $[\mathbf{j}_a]_i = \delta_{[\text{node } i] \in \mathcal{C}_a}$ the canonical vector of community \mathcal{C}_a . Also,

$$\text{Var}[\mathbf{A}_{ij}] = p \left(1 + \frac{\mathbf{M}_{ab}}{\sqrt{n}} \right) \left[1 - p \left(1 + \frac{\mathbf{M}_{ab}}{\sqrt{n}} \right) \right] = p(1-p) + O(n^{-\frac{1}{2}}). \quad (7.5)$$

As such, since $\mathbf{J} \mathbf{1}_k = \mathbf{1}_n$ and thus $\mathbf{1}_n \mathbf{1}_n^\top = \mathbf{J} \mathbf{1}_k \mathbf{1}_k^\top \mathbf{J}^\top$, we can anticipate from the previous section that $\frac{1}{\sqrt{np(1-p)}} \mathbf{A}$ is well approximated by a rank (at most) k perturbation of a random Wigner matrix $\frac{1}{\sqrt{n}} \mathbf{X}$ having i.i.d. entries of zero mean and unit variance. The perturbation matrix has a largely dominant eigenvalue of order $O(\sqrt{n})$ and up to $k-1$ isolated eigenvalues outside the bulk of the limiting spectrum of $\frac{1}{\sqrt{n}} \mathbf{X}$; here “up to” translates the fact that, $\frac{1}{n} \mathbf{J} \mathbf{M} \mathbf{J}^\top$ being of operator norm $O(1)$ (same as $\frac{1}{\sqrt{n}} \mathbf{X}$), phase transition phenomena are bound to occur.

As for $\mathbf{B} = \frac{1}{\sqrt{n}} (\mathbf{A} - \frac{\mathbf{d}\mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n})$, it will likely have the same effect of discarding the dominant $O(\sqrt{n})$ eigenvalue-eigenvector pair, thereby ensuring that $\|\mathbf{B}\| = O(1)$ and that the possibly isolated and bulk eigenvalues of \mathbf{B} are comparable.

Performing the same analysis as in the Erdős-Rényi setting brings additional terms which, after carefully discarding the terms of vanishing norms (as usual, special care is needed when taking the product of matrices or vectors with

different norms and different levels of dependence), gives the SBM version of (7.2):

$$\frac{\mathbf{B}}{\sqrt{p(1-p)}} = \frac{\mathbf{X}}{\sqrt{n}} + \sqrt{\frac{p}{1-p}} \frac{\mathbf{JM}^\circ \mathbf{J}^\top}{n} - \frac{\mathbf{1}_n \mathbf{1}_n^\top \mathbf{X} + \mathbf{X} \mathbf{1}_n \mathbf{1}_n^\top}{n\sqrt{n}} + O_{\|\cdot\|}(n^{-\frac{1}{2}}) \quad (7.6)$$

where we defined

$$\mathbf{M}^\circ = (\mathbf{I}_k - \mathbf{1}_k \mathbf{c}^\top) \mathbf{M} (\mathbf{I}_k - \mathbf{c} \mathbf{1}_k^\top) \quad (7.7)$$

with $\mathbf{c} = [\frac{n_1}{n}, \dots, \frac{n_k}{n}]^\top \in \mathbb{R}^k$ the community size ratios. The matrix \mathbf{M}° is a “centered” version of \mathbf{M} accounting for the community sizes in the sense that $\mathbf{M}^\circ \mathbf{c} = \mathbf{0}$, i.e., for all a , $\sum_{b=1}^k n_b [\mathbf{M}^\circ]_{ab} = 0$.

Consequently, \mathbf{B} is again (up to scaling) the sum of a symmetric random matrix \mathbf{X} with zero mean unit variance entries and a perturbation matrix of rank up to k (since $\mathbf{1}_n$ is in the span of the columns of \mathbf{J}). Clearly, in expectation,

$$\mathbb{E}[\mathbf{B}] = p \frac{\mathbf{JM}^\circ \mathbf{J}^\top}{n} + O_{\|\cdot\|}(n^{-\frac{1}{2}})$$

so that the dominant eigenvectors of \mathbf{B} are, as one would expect, linear combinations of the class structure vectors $\mathbf{j}_1, \dots, \mathbf{j}_k$, weighted here by the coefficients of \mathbf{M}° . The additional random fluctuations being due to the matrices \mathbf{X} and $\mathbf{1}_n \mathbf{1}_n^\top \mathbf{X} + \mathbf{X} \mathbf{1}_n \mathbf{1}_n^\top$, which are “isotropic” with respect to the class structures, they should not affect the quality of a k-means or expectation-maximization (EM) clustering of the informative eigenvectors of \mathbf{B} . We will see in the next section that this important remark no longer holds for stochastic block models with heterogeneous degrees.

Pursuing our derivation, to stress the presence of a rank- k perturbation, note that \mathbf{B} can be rewritten compactly as

$$\frac{\mathbf{B}}{\sqrt{p(1-p)}} = \frac{\mathbf{X}}{\sqrt{n}} + \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{X} \mathbf{1}_n}{n} \end{bmatrix} \begin{bmatrix} \sqrt{\frac{p}{1-p}} \mathbf{M}^\circ & -\mathbf{1}_k \\ -\mathbf{1}_k^\top & 0 \end{bmatrix} \begin{bmatrix} \frac{\mathbf{J}^\top}{\sqrt{n}} \\ \frac{\mathbf{1}_n^\top \mathbf{X}^\top}{n} \end{bmatrix} + O_{\|\cdot\|}(n^{-\frac{1}{2}}).$$

It is interesting to note that \mathbf{B} is here a rank- k perturbation of \mathbf{X} that is *not* independent of \mathbf{X} , from the presence of the vector $\mathbf{X} \mathbf{1}_n$ (although, as we will see subsequently, the term $\mathbf{X} \mathbf{1}_n$ will have asymptotically no effect on the limiting spectral properties of \mathbf{B}).

Studying the (limiting) location of the (possibly) isolated eigenvalues of \mathbf{B} thus consists in solving, for $\lambda > 2$ (i.e., the right edge of the semicircle law, as the limiting support of the eigenvalues of \mathbf{X}/\sqrt{n}),

$$\begin{aligned} & \det \left(\frac{\mathbf{X}}{\sqrt{n}} - \lambda \mathbf{I}_n + \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{X} \mathbf{1}_n}{n} \end{bmatrix} \begin{bmatrix} \sqrt{\frac{p}{1-p}} \mathbf{M}^\circ & -\mathbf{1}_k \\ -\mathbf{1}_k^\top & 0 \end{bmatrix} \begin{bmatrix} \frac{\mathbf{J}^\top}{\sqrt{n}} \\ \frac{\mathbf{1}_n^\top \mathbf{X}^\top}{n} \end{bmatrix} \right) = 0 \\ & \Leftrightarrow \det \left(\mathbf{I}_n + \mathbf{Q} \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{X} \mathbf{1}_n}{n} \end{bmatrix} \begin{bmatrix} \sqrt{\frac{p}{1-p}} \mathbf{M}^\circ & -\mathbf{1}_k \\ -\mathbf{1}_k^\top & 0 \end{bmatrix} \begin{bmatrix} \frac{\mathbf{J}^\top}{\sqrt{n}} \\ \frac{\mathbf{1}_n^\top \mathbf{X}^\top}{n} \end{bmatrix} \right) = 0 \\ & \Leftrightarrow \det \left(\mathbf{I}_{k+1} + \begin{bmatrix} \frac{\mathbf{J}^\top}{\sqrt{n}} \\ \frac{\mathbf{1}_n^\top \mathbf{X}^\top}{n} \end{bmatrix} \mathbf{Q} \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{X} \mathbf{1}_n}{n} \end{bmatrix} \begin{bmatrix} \sqrt{\frac{p}{1-p}} \mathbf{M}^\circ & -\mathbf{1}_k \\ -\mathbf{1}_k^\top & 0 \end{bmatrix} \right) = 0 \end{aligned}$$

where we introduced the resolvent $\mathbf{Q} = \mathbf{Q}(\lambda) = (\frac{\mathbf{X}}{\sqrt{n}} - \lambda \mathbf{I}_n)^{-1}$.

From the deterministic equivalent $\mathbf{Q} \leftrightarrow m(\lambda) \mathbf{I}_n$ in Theorem 2.5, where $m(\lambda)$ is here the unique *negative* solution to $m^2(\lambda) + \lambda m(\lambda) + 1 = 0$, we find that

$$\begin{bmatrix} \frac{\mathbf{J}^\top}{\sqrt{n}} \\ \frac{\mathbf{1}_n^\top \mathbf{X}^\top}{n} \end{bmatrix} \mathbf{Q} \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{X} \mathbf{1}_n}{n} \end{bmatrix} = \begin{bmatrix} m(\lambda) \mathbf{D}_c & (1 + \lambda m(\lambda)) \mathbf{c} \\ (1 + \lambda m(\lambda)) \mathbf{c}^\top & \lambda(1 + \lambda m(\lambda)) \end{bmatrix} + o(1)$$

(almost surely) where $\mathbf{D}_c = \text{diag}(\mathbf{c})$ and the bottom right entry follows from a repeated use of the relation $\mathbf{Q}\mathbf{X}/\sqrt{n} = \mathbf{I}_n + \lambda\mathbf{Q}$ and of the above deterministic equivalent. This gives the (asymptotically) equivalent determinantal equation

$$\det \left(\begin{bmatrix} \mathbf{I}_k + m(\lambda) \sqrt{\frac{p}{1-p}} \mathbf{D}_c \mathbf{M}^\circ & (1 + \lambda m(\lambda)) \mathbf{c} \mathbf{1}_k^\top & -m(\lambda) \mathbf{c} \\ -\lambda(1 + \lambda m(\lambda)) \mathbf{1}_k^\top & -\lambda m(\lambda) \end{bmatrix} \right) = 0 \quad (7.8)$$

which, using the block-determinant relation $\det(\frac{\mathbf{A}^\top}{\mathbf{v}^\top} \frac{\mathbf{u}}{w}) = \det(\mathbf{A} - \frac{1}{w} \mathbf{u} \mathbf{v}^\top)$, is simply

$$\det \left(\mathbf{I}_k + m(\lambda) \sqrt{\frac{p}{1-p}} \mathbf{D}_c \mathbf{M}^\circ \right) = \det \left(\mathbf{I}_k + m(\lambda) \sqrt{\frac{p}{1-p}} \mathbf{D}_{\sqrt{c}} \mathbf{M}^\circ \mathbf{D}_{\sqrt{c}} \right) = 0$$

where we denote $\sqrt{\mathbf{c}} = (\sqrt{\frac{n_1}{n}}, \dots, \sqrt{\frac{n_k}{n}})^\top$. The cancellation of the term proportional to $\mathbf{c} \mathbf{1}_k^\top$ stresses the asymptotic “inaction” of the vector $\mathbf{X} \mathbf{1}_n$ in (7.6) on the *informative* (for community detection purpose) eigenvalues of \mathbf{B} .

This leads to the following result on isolated eigenvalues.

Theorem 7.1 (Isolated eigenvalues in the SBM, from [Ali and Couillet, 2018]).
Under the setting of (7.3) and $n_a/n \rightarrow c_a \in (0, 1)$, for each eigenvalue ℓ of $\mathbf{D}_c \mathbf{M}^\circ$ satisfying

$$|\ell| > \sqrt{\frac{1-p}{p}}$$

there exists an associated eigenvalue λ_ℓ of $\frac{1}{\sqrt{p(1-p)}} \mathbf{B}$ such that, for all large n almost surely, $|\lambda_\ell| > 2$ and

$$\lambda_\ell = m^{-1} \left(-\sqrt{\frac{1-p}{p}} \frac{1}{\ell} \right) + o(1) = \frac{p\ell + \frac{1-p}{\ell}}{\sqrt{p(1-p)}} + o(1)$$

for $m^{-1}(\cdot) : (-1, 0) \rightarrow (2, \infty)$, $t \mapsto (-1 - t^2)/t$ the local inverse of $m(\cdot)$.

Obviously, as in the case of spectral clustering in Section 4.5.1, isolated eigenvalues in the spectrum of \mathbf{B} are associated with eigenvectors aligned to the informative linear combination of the class-vectors $\mathbf{j}_1, \dots, \mathbf{j}_k$.

To assess the (limiting) projection of these eigenvectors $\hat{\mathbf{u}}_\ell$ of $\mathbf{B}/\sqrt{p(1-p)}$ on each (normalized) direction $\mathbf{j}_1, \dots, \mathbf{j}_k$, we may next evaluate

$$\frac{1}{n} \mathbf{D}_c^{-\frac{1}{2}} \mathbf{J}^\top \hat{\mathbf{u}}_\ell \hat{\mathbf{u}}_\ell^\top \mathbf{J} \mathbf{D}_c^{-\frac{1}{2}} = -\frac{1}{2\pi i} \oint_{\Gamma_\ell} \frac{1}{n} \mathbf{D}_c^{-\frac{1}{2}} \mathbf{J}^\top \left(\frac{\mathbf{B}}{\sqrt{p(1-p)}} - z \mathbf{I}_n \right)^{-1} \mathbf{J} \mathbf{D}_c^{-\frac{1}{2}} dz \quad (7.9)$$

where $\frac{1}{n} \mathbf{J} \mathbf{D}_{\mathbf{c}}^{-1} \mathbf{J}^T$ is a projector on span of $\mathbf{j}_1, \dots, \mathbf{j}_k$ and Γ_ℓ a fixed complex contour circling around λ_ℓ only (for all large n). Similar to the determination of the limiting location of the isolated eigenvalues, by isolating $\mathbf{X}/\sqrt{n} - z\mathbf{I}_n$ from $\mathbf{B}/\sqrt{p(1-p)} - z\mathbf{I}_n$ with the matrix inversion lemmas, Lemma 2.7 and 2.5, we obtain

$$\frac{1}{n} \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}} \mathbf{J}^T \hat{\mathbf{u}}_\ell \hat{\mathbf{u}}_\ell^T \mathbf{J} \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}} = -\frac{1}{2\pi i} \oint_{\Gamma_\ell} m(z) \left(\mathbf{I}_k + \frac{\sqrt{pm}(z) \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}} \mathbf{M}^\circ \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}}}{\sqrt{1-p}} \right)^{-1} dz + o(1).$$

In detail, to obtain this result, note that the desired matrix $\frac{1}{n} \mathbf{J}^T \left(\frac{\mathbf{B}}{\sqrt{p(1-p)}} - z\mathbf{I}_n \right)^{-1} \mathbf{J}$ is the $(1, 1)$ block of

$$\begin{aligned} & \begin{bmatrix} \frac{\mathbf{J}^T}{\sqrt{n}} \\ \frac{\mathbf{1}_n^T \mathbf{X}^T}{n} \end{bmatrix} \left(\frac{\mathbf{B}}{\sqrt{p(1-p)}} - z\mathbf{I}_n \right)^{-1} \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{X}\mathbf{1}_n}{n} \end{bmatrix} \\ &= \left(\mathbf{I}_{k+1} + \begin{bmatrix} \frac{\mathbf{J}^T}{\sqrt{n}} \\ \frac{\mathbf{1}_n^T \mathbf{X}^T}{n} \end{bmatrix} \mathbf{Q} \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{X}\mathbf{1}_n}{n} \end{bmatrix} \begin{bmatrix} \sqrt{\frac{p}{1-p}} \mathbf{M}^\circ & -\mathbf{1}_k \\ -\mathbf{1}_k^T & 0 \end{bmatrix} \right)^{-1} \\ &\quad \times \begin{bmatrix} \frac{\mathbf{J}^T}{\sqrt{n}} \\ \frac{\mathbf{1}_n^T \mathbf{X}^T}{n} \end{bmatrix} \mathbf{Q} \begin{bmatrix} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{X}\mathbf{1}_n}{n} \end{bmatrix} + o(1) \end{aligned}$$

where we recall $\mathbf{Q} = \mathbf{Q}(\lambda) = (\frac{\mathbf{X}}{\sqrt{n}} - \lambda\mathbf{I}_n)^{-1}$. Again with the deterministic equivalent result $\mathbf{Q} \leftrightarrow m(\lambda)\mathbf{I}_n$ in Theorem 2.5, we deduce

$$\begin{aligned} & \frac{1}{n} \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}} \mathbf{J}^T \hat{\mathbf{u}}_\ell \hat{\mathbf{u}}_\ell^T \mathbf{J} \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}} \\ &= -\frac{1}{2\pi i} \oint_{\Gamma_\ell} m(z) \left(\mathbf{I}_k + \frac{\sqrt{pm}(z) \mathbf{D}_{\sqrt{\mathbf{c}}} \mathbf{M}^\circ \mathbf{D}_{\sqrt{\mathbf{c}}}}{\sqrt{1-p}} \right)^{-1} \left(\mathbf{I}_k - \sqrt{\mathbf{c}} \sqrt{\mathbf{c}}^T \right) dz + o(1) \end{aligned}$$

where we neglected all terms leading to a vanishing residue in the large n limit. Since $\mathbf{M}^\circ \mathbf{D}_{\sqrt{\mathbf{c}}} \sqrt{\mathbf{c}} = \mathbf{M}^\circ \mathbf{c} = \mathbf{0}$, it follows after development that the term $\left(\mathbf{I}_k + \frac{\sqrt{pm}(z) \mathbf{D}_{\sqrt{\mathbf{c}}} \mathbf{M}^\circ \mathbf{D}_{\sqrt{\mathbf{c}}}}{\sqrt{1-p}} \right)^{-1} \sqrt{\mathbf{c}}$ will not have a residue in Γ_ℓ , so that $(\mathbf{I}_k - \sqrt{\mathbf{c}} \sqrt{\mathbf{c}}^T)$ above can be replaced by \mathbf{I}_k .

Further residue calculus leads to the existence of a unique pole of order 1 with associated residue given by

$$\begin{aligned} \frac{1}{n} \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}} \mathbf{J}^T \hat{\mathbf{u}}_\ell \hat{\mathbf{u}}_\ell^T \mathbf{J} \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}} &= \lim_{z \rightarrow \lambda_\ell} (\lambda_\ell - z) m(z) \left(\mathbf{I}_k + \frac{\sqrt{pm}(z) \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}} \mathbf{M}^\circ \mathbf{D}_{\mathbf{c}}^{-\frac{1}{2}}}{\sqrt{1-p}} \right)^{-1} + o(1) \\ &= \frac{-m(\lambda_\ell)}{\sqrt{\frac{p}{1-p}} \ell m'(\lambda_\ell)} \mathbf{u}_\ell \mathbf{u}_\ell^T + o(1) \end{aligned}$$

where \mathbf{u}_ℓ is the eigenvector of $\mathbf{D}_{\sqrt{\mathbf{c}}} \mathbf{M}^\circ \mathbf{D}_{\sqrt{\mathbf{c}}}$ associated with eigenvalue ℓ .

Recalling that λ_ℓ is solution to $1 + \sqrt{p/(1-p)}\ell m(\lambda_\ell) = 0$ and that $m(z)^2 + zm(z) = -1$ (which we can differentiate along z), this can be further simplified as

$$\frac{1}{n} \mathbf{D}_c^{-\frac{1}{2}} \mathbf{J}^\top \hat{\mathbf{u}}_\ell \hat{\mathbf{u}}_\ell^\top \mathbf{J} \mathbf{D}_c^{-\frac{1}{2}} = (1 - m(\lambda_\ell)^2) \mathbf{u}_\ell \mathbf{u}_\ell^\top + o(1) = \left(1 - \frac{1-p}{p\ell^2}\right) \mathbf{u}_\ell \mathbf{u}_\ell^\top + o(1).$$

These two alternative formulas have nice interpretations: outside the support of the semicircle law, $\lambda \mapsto 1 - m(\lambda)^2$ is positive, increasing and maps $(2, \infty)$ onto $(0, 1)$. In particular, the alignment of $\hat{\mathbf{u}}_\ell$ onto the subspace spanned by $\mathbf{j}_1, \dots, \mathbf{j}_k$ is given by $\frac{1}{n} \|\mathbf{D}_c^{-\frac{1}{2}} \mathbf{J}^\top \hat{\mathbf{u}}_\ell\|^2 = 1 - m(\lambda_\ell)^2 + o(1)$. Equivalently, recalling that one needs $|\ell| > \sqrt{(1-p)/p}$ to have isolated eigenvalues, as $|\ell|$ increases in $(\sqrt{(1-p)/p}, \infty)$, the alignment increases to 1 at rate $1/\ell^2$.

This is summarized in the following result.

Theorem 7.2 (Eigenvector alignment in the SBM). *Under the setting and notations of Theorem 7.1, if $|\ell| > \sqrt{(1-p)/p}$ for (ℓ, \mathbf{u}_ℓ) an eigenvalue-eigenvector pair of $\mathbf{D}_{\sqrt{c}} \mathbf{M}^\circ \mathbf{D}_{\sqrt{c}}$, then the eigenvalue-eigenvector pair $(\lambda_\ell, \hat{\mathbf{u}}_\ell)$ of \mathbf{B} satisfies*

$$\frac{1}{n} \mathbf{D}_c^{-\frac{1}{2}} \mathbf{J}^\top \hat{\mathbf{u}}_\ell \hat{\mathbf{u}}_\ell^\top \mathbf{J} \mathbf{D}_c^{-\frac{1}{2}} = (1 - m(\lambda_\ell)^2) \mathbf{u}_\ell \mathbf{u}_\ell^\top + o(1) = \left(1 - \frac{1-p}{p\ell^2}\right) \mathbf{u}_\ell \mathbf{u}_\ell^\top + o(1).$$

as $n \rightarrow \infty$, almost surely.

7.1.1.3 Case study: 2-class symmetric SBM

Let us observe the consequences of the results in Theorem 7.1 and 7.2 on the case of the two-class symmetric SBM. In this setting, we define the connection probability matrix \mathbf{C} in (7.3) as

$$\mathbf{C} = \begin{bmatrix} p_{\text{in}} & p_{\text{out}} \\ p_{\text{out}} & p_{\text{in}} \end{bmatrix}$$

for some $p_{\text{in}}, p_{\text{out}} \in (0, 1)$ the inner-class and outer-class connection probabilities. We also set the class cardinalities as $\mathbf{c} = [1/2, 1/2]^\top$. By an exchangeability argument, the statistics of the eigenvectors of \mathbf{B} are in this case symmetric and thus more expressive.

In the context of the previous section, this choice implies that

$$p = p_{\text{out}}, \quad \mathbf{M} = \sqrt{n} \cdot \frac{p_{\text{in}} - p_{\text{out}}}{p_{\text{out}}} \mathbf{I}_2 \tag{7.10}$$

which indicates, of course, that p_{in} depends on n and scales as $p_{\text{out}} + O(n^{-\frac{1}{2}})$ for \mathbf{M} to remain of bounded norm as $n \rightarrow \infty$. As a consequence,

$$\mathbf{D}_{\sqrt{c}} \mathbf{M}^\circ \mathbf{D}_{\sqrt{c}} = \frac{\sqrt{n}(p_{\text{in}} - p_{\text{out}})}{p_{\text{out}}} \frac{1}{2} \left(\mathbf{I}_2 - \frac{1}{2} \mathbf{1}_2 \mathbf{1}_2^\top \right) \tag{7.11}$$

which has a unique nonzero eigenvalue, equal to

$$\ell = \frac{\sqrt{n}(p_{\text{in}} - p_{\text{out}})}{2p_{\text{out}}}$$

and with associated eigenvector (up to a sign)

$$\mathbf{u}_\ell = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

It follows from Theorem 7.1 that the community detectability phase transition occurs under the condition

$$|p_{\text{in}} - p_{\text{out}}| > \frac{2\sqrt{p_{\text{out}}(1-p_{\text{out}})}}{\sqrt{n}}. \quad (7.12)$$

The isolated eigenvalue λ_ℓ of $\frac{1}{\sqrt{p_{\text{out}}(1-p_{\text{out}})}}\mathbf{B}$ is thus defined via its Stieltjes transform as

$$m(\lambda_\ell) = -\frac{2\sqrt{p_{\text{out}}(1-p_{\text{out}})}}{\sqrt{n}(p_{\text{in}} - p_{\text{out}})} + o(1)$$

or equivalently, using $m(\cdot)^{-1}(t) = (-1 - t^2)/t$,

$$\lambda_\ell = \frac{\sqrt{n}(p_{\text{in}} - p_{\text{out}})}{2\sqrt{p_{\text{out}}(1-p_{\text{out}})}} + \frac{2\sqrt{p_{\text{out}}(1-p_{\text{out}})}}{\sqrt{n}(p_{\text{in}} - p_{\text{out}})} + o(1).$$

Consequently, the asymptotic projection of the associated eigenvector $\hat{\mathbf{u}}_\ell$ on $\mathbf{j}_1, \mathbf{j}_2$ is given by

$$\frac{2}{n} [\mathbf{j}_1 \quad \mathbf{j}_2]^\top \hat{\mathbf{u}}_\ell \hat{\mathbf{u}}_\ell^\top [\mathbf{j}_1 \quad \mathbf{j}_2] = \frac{1}{2} \left(1 - \frac{4p_{\text{out}}(1-p_{\text{out}})}{n(p_{\text{in}} - p_{\text{out}})^2} \right) \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} + o(1).$$

By exchangeability and symmetry, these results also give access to the mean and variance of every entry $[\hat{\mathbf{u}}_\ell]_i$ of the vector $\hat{\mathbf{u}}_\ell$. Specifically,

$$\frac{2}{n} (\mathbf{j}_1^\top \hat{\mathbf{u}}_\ell)^2 = \frac{n}{2} \left(\frac{1}{n/2} \sum_{i=1}^{n/2} [\hat{\mathbf{u}}_\ell]_i \right)^2$$

gives access to the empirical average value of the entries $[\hat{\mathbf{u}}_\ell]_i$ that are identically distributed on $1 \leq i \leq n/2$ and on $n/2 + 1 \leq i \leq n$. Also

$$\frac{2}{n} (\mathbf{j}_1^\top \hat{\mathbf{u}}_\ell)(\mathbf{j}_2^\top \hat{\mathbf{u}}_\ell) = \frac{n}{2} \left(\frac{1}{n/2} \sum_{i=1}^{n/2} [\hat{\mathbf{u}}_\ell]_i \right) \left(\frac{1}{n/2} \sum_{i=n/2+1}^n [\hat{\mathbf{u}}_\ell]_i \right)$$

gives access to the (empirical) correlation between the first $n/2$ and last $n/2$ elements of $\hat{\mathbf{u}}_\ell$. We thus find that

$$\begin{aligned}\mathbb{E}[\hat{\mathbf{u}}_\ell]_i &= \begin{cases} \sqrt{\frac{1}{n} \left(1 - \frac{4p_{\text{out}}(1-p_{\text{out}})}{n(p_{\text{in}}-p_{\text{out}})^2} \right)} + o(1), & 1 \leq i \leq \frac{n}{2} \\ -\sqrt{\frac{1}{n} \left(1 - \frac{4p_{\text{out}}(1-p_{\text{out}})}{n(p_{\text{in}}-p_{\text{out}})^2} \right)} + o(1), & \frac{n}{2} + 1 \leq i \leq n \end{cases} \\ \text{Var}[\hat{\mathbf{u}}_\ell]_i &= \frac{4p_{\text{out}}(1-p_{\text{out}})}{n^2(p_{\text{in}}-p_{\text{out}})^2} + o(1)\end{aligned}$$

where the result on the expectation is valid up to sign (since eigenvectors are defined up to a sign) and the result on the variance exploits the constraint $\sum_{i=1}^n [\hat{\mathbf{u}}_\ell]_i^2 = 1$.

It can be further shown that the fluctuations of $[\hat{\mathbf{u}}_\ell]_i$ are asymptotically Gaussian and asymptotically independent across i , see Remark 7.1 below, (only asymptotically since the constraint $\|\hat{\mathbf{u}}_\ell\| = 1$ creates a finite-dimensional dependence), the above results on the expectation and variance thus immediately lead to the evaluation of the classification error based on $\hat{\mathbf{u}}_\ell$. Letting $\hat{\mathcal{C}}_i = \text{sign}([\hat{\mathbf{u}}_\ell]_i)$ be the estimate of the underlying community \mathcal{C}_i of the node i , with the sign convention $[\hat{\mathbf{u}}_\ell]_1 \geq 0$, the classification error rate satisfies

$$\frac{1}{n} \sum_{i=1}^n \delta_{\hat{\mathcal{C}}_i \neq \mathcal{C}_i} - Q \left(\sqrt{\frac{n(p_{\text{in}}-p_{\text{out}})^2}{4p_{\text{out}}(1-p_{\text{out}})}} - 1 \right) \xrightarrow{\text{a.s.}} 0 \quad (7.13)$$

for $Q(t) = \frac{1}{\sqrt{2\pi}} \int_t^\infty e^{-\frac{u^2}{2}} du$. Note in particular that this classification error does not depend on n as $n(p_{\text{in}}-p_{\text{out}})^2 = O(1)$ under (7.10).

Remark 7.1 (On the asymptotic Gaussianity of the error rate). *It is interesting to realize that the asymptotic Gaussianity of the misclassification probability, despite depending on all the entries of $\hat{\mathbf{u}}_\ell$, only requires to prove the asymptotic two-dimensional Gaussianity of any pair $([\hat{\mathbf{u}}_\ell]_i, [\hat{\mathbf{u}}_\ell]_j)$ of the entries of $\hat{\mathbf{u}}_\ell$. It suffices indeed to proceed as follows [Kadavankandy and Couillet, 2019]:*

1. **Pairwise Gaussianity using the resolvent.** As usual, we first seek to express the quantities of interest (here $[\hat{\mathbf{u}}_\ell]_i$) as a function of the resolvent $\mathbf{Q}(z) = (\mathbf{X}/\sqrt{n} - z\mathbf{I}_n)^{-1}$. We start by writing

$$\frac{1}{\sqrt{p_{\text{out}}(1-p_{\text{out}})}} \mathbf{B} \hat{\mathbf{u}}_\ell = \lambda_\ell \hat{\mathbf{u}}_\ell$$

which, with (7.6) and basic algebraic manipulations, leads to

$$\sqrt{n}[\hat{\mathbf{u}}_\ell]_i = -\sqrt{\frac{p_{\text{out}}}{1-p_{\text{out}}}} \left(\sqrt{n} \mathbf{e}_i^\top \mathbf{Q}(\lambda) \frac{\mathbf{J}}{\sqrt{n}} \right) \left(\mathbf{M}^\circ \frac{\mathbf{J}^\top}{\sqrt{n}} \hat{\mathbf{u}}_\ell \right) + o(1)$$

with $[\mathbf{e}_i]_j = \delta_{ij}$ the canonical basis vector. The right parenthesis term converges to known limits (from standard eigenvector alignment results,

e.g., Theorem 2.14), while the first parenthesis term “fluctuates”. Using central limit arguments for random matrices (either based on a martingale difference approach as in [Bai and Silverstein, 2010] or a Gaussian integration-by-part technique as in [Pastur and Sheferbina, 2011]; see Section 2.6.3 for more discussions), it can be further shown that the vector $[\sqrt{n}[\hat{\mathbf{u}}_\ell]_i, \sqrt{n}[\hat{\mathbf{u}}_\ell]_j]^\top$ has a two-dimensional Gaussian limit.

2. From there, the misclassification rate in the left-hand side of (7.13) corresponds to

$$S \equiv \frac{1}{n} \sum_{i=1}^n \delta_{\hat{c}_i \neq c_i} = \frac{1}{n} \sum_{i \leq n/2} \delta_{\sqrt{n}[\hat{\mathbf{u}}_\ell]_i < 0} + \frac{1}{n} \sum_{i > n/2} \delta_{\sqrt{n}[\hat{\mathbf{u}}_\ell]_i > 0}.$$

Writing $S = S - \mathbb{E}[S] + \mathbb{E}[S]$, by exchangeability we have from the previous item that $\mathbb{E}[S] = \mathbb{P}(\sqrt{n}[\hat{\mathbf{u}}_\ell]_1 < 0)$ for $\sqrt{n}[\hat{\mathbf{u}}_\ell]_1$ having a known Gaussian limit, while the fluctuation $\mathbb{P}(|S - \mathbb{E}[S]| > t) \leq \text{Var}[S]/t^2$ which exclusively depends on $\mathbb{E}[n[\hat{\mathbf{u}}_\ell]_i[\hat{\mathbf{u}}_\ell]_j]$ for any pair (i, j) , is bound to vanish. This completes the proof of (7.13) without having to resort to any further joint statistics of the entries of $\hat{\mathbf{u}}_\ell$.

Figure 7.1 depicts the probability of correct classification for a 2-class SBM under the present symmetric setting. The asymptotic predictions in (7.13) closely match the empirical performance, with a slight mismatch for small n around the phase transition (around 1 in the x-axis). This is not surprising as the limiting discontinuity can hardly be observed in finite (especially small) dimensions. This is a typical example where the convergence to random matrix asymptotics tends to be slow, as in the case of Figure 2.12 in Section 2.5.

The limiting results derived in this section are quite simple and have the advantage of being in closed form. The SBM setting is however quite unrealistic in the sense that the average degree of each node is constant (converging to p), which does not translate the heterogeneity of node connectivity in real graphs and, as a result, cannot provide a typical “power-law” scaling of the degrees, that is of more practical interest for real-world graph problems [Adamic and Glance, 2005, Borgs et al., 2019].

The next section brings the present analysis into more realistic graph models by considering the so-called degree-corrected SBM (DC-SBM, Coja-Oghlan and Lanka [2010], Karrer and Newman [2011]) which takes into account the degree heterogeneity. This has several non-trivial consequences on: (i) the “shape” of the limiting eigenvalue distribution of \mathbf{B} (no longer a scaled semicircle in general), (ii) the resulting phase transition condition, and (iii) the “content” of the dominant eigenvectors that do not lead straightforwardly to the classes as in the SBM case. The expressions of these limiting behaviors are less simple but provide a sufficiently clear account of the (mis)behavior of spectral clustering to envision several directions of improvements.




Figure 7.1: Classification performance for a 2-class SBM with $n_1 = n_2$, as a function of $p_{in} - p_{out}$ with $p_{out} = 0.4$. Simulations averaged over 100 realizations.
[\[Link to code\]](#)

7.1.2 The degree-corrected stochastic block model

In this section, we generalize the stochastic block model by allowing, in addition to the existence of communities, different ‘‘intrinsic’’ degrees for the nodes in the graph. This better translates the nature of real-world graphs in which nodes have possibly very heterogeneous degrees.

Precisely, we here demand that

$$\mathbf{A}_{ij} = \mathbf{A}_{ji} \sim \text{Bern}(q_i q_j \mathbf{C}_{ab})$$

for $q_i > 0$ some weight factor accounting for the connectivity of node i and $\mathcal{C}_a, \mathcal{C}_b \in \{1, \dots, k\}$ the communities of node i and j , respectively. Similar to before, we assume the cardinality $n_a = |\mathcal{C}_a|$ of class \mathcal{C}_a to be of the same order as n , i.e., $n_a/n \rightarrow c_a \in (0, 1)$. For the moment, we consider the q_i ’s to be deterministic, but we will soon take them random i.i.d., yet independent of the Bernoulli realization.

As in the SBM case in (7.3), we also request the non-trivial clustering setting corresponding to

$$[\mathbf{C}]_{ab} = 1 + \frac{[\mathbf{M}]_{ab}}{\sqrt{n}} \quad (7.14)$$

where, as opposed to the SBM setting, the parameter p is no longer necessary.

A first important remark is that, similar to (7.4) for SBM, we have in this configuration,

$$\mathbb{E}[\mathbf{A}] = \mathbf{D}_{\mathbf{q}} \left(\mathbf{1}_n \mathbf{1}_n^\top + \frac{1}{\sqrt{n}} \mathbf{J} \mathbf{M} \mathbf{J}^\top \right) \mathbf{D}_{\mathbf{q}} \quad (7.15)$$

where $\mathbf{q} = [q_1, \dots, q_n]^\top \in \mathbb{R}^n$ and $\mathbf{D}_\mathbf{q} = \text{diag}(\mathbf{q})$. In particular, observe that the eigenvectors of $\mathbb{E}[\mathbf{A}]$ are no longer linear combinations of $\mathbf{j}_1, \dots, \mathbf{j}_k$ but are “deformed” by the (usually unknown) weights q_1, \dots, q_n . Compensating for this eigenvector deformation is not completely obvious and will be the major technical point of interest in this section.

As for the variance of the elements of \mathbf{A} , similar to the SBM setting in (7.5), we find that

$$\text{Var}[\mathbf{A}_{ij}] = q_i q_j (1 - q_i q_j) + O(n^{-\frac{1}{2}})$$

which does not depend on the communities of nodes i and j .

As such, up to low-rank perturbation, \mathbf{A} is a matrix with independent entries of zero mean and variance $q_i q_j (1 - q_i q_j)$. Consequently, the limiting spectral measure of $\frac{1}{\sqrt{n}}\mathbf{A}$ (and of its rank-one perturbation $\mathbf{B} = \frac{1}{\sqrt{n}}(\mathbf{A} - \frac{\mathbf{d}\mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n})$), is that of a *deformed* Wigner matrix with variance profile.

It is instructive to first analyze this spectrum. From Theorem 2.9 in which we set $\sigma_{ij}^2 = q_i q_j (1 - q_i q_j)$, we find that the Stieltjes transform of the eigenvalue distribution of \mathbf{B} satisfies

$$\frac{1}{n} \text{tr}(\mathbf{B} - z\mathbf{I}_n)^{-1} - \frac{1}{n} \text{tr}(\mathbf{D}_\mathbf{g} - z\mathbf{I}_n)^{-1} \xrightarrow{a.s.} 0$$

where $\mathbf{g} = [g_1, \dots, g_n]^\top$ and

$$\begin{aligned} g_i &= -\frac{1}{n} \sum_{j=1}^n \frac{q_i q_j - q_i^2 q_j^2}{-z + g_j} = -q_i \frac{1}{n} \sum_{j=1}^n \frac{q_j}{-z + g_j} + q_i^2 \frac{1}{n} \sum_{j=1}^n \frac{q_j^2}{-z + g_j} \\ &\equiv -q_i g_{10} + q_i^2 g_{20} \end{aligned}$$

where we introduced g_{10} and g_{20} the solutions to

$$g_{10} = \frac{1}{n} \sum_{i=1}^n \frac{q_i}{-z - q_i g_{10} + q_i^2 g_{20}}, \quad g_{20} = \frac{1}{n} \sum_{i=1}^n \frac{q_i^2}{-z - q_i g_{10} + q_i^2 g_{20}}. \quad (7.16)$$

Thus, finally,

$$\frac{1}{n} \text{tr}(\mathbf{B} - z\mathbf{I}_n)^{-1} - \frac{1}{n} \text{tr}(-g_{10}\mathbf{D}_\mathbf{q} + g_{20}\mathbf{D}_{\mathbf{q}^2} - z\mathbf{I}_n)^{-1} \xrightarrow{a.s.} 0$$

where $\mathbf{q}^2 = [q_1^2, \dots, q_n^2]^\top$ and g_{10}, g_{20} are defined as the solution to (7.16).

As shown in Figure 7.2, unlike in the SBM setting, the spectrum of \mathbf{B} can be more spread out than a semicircle, when the q_i 's are independently drawn from a bimodal law. As a consequence, it is expected that phase transitions for the appearance of isolated eigenvalues due to the presence of communities will occur more or less easily depending on this eigenvalue spreading. Here in Figure 7.2, depending on \mathbf{M} , either two isolated eigenvalues or only one are found in the spectrum (with corresponding eigenvectors displaying more or less informative structure).




Figure 7.2: Two graphs generated from DCSBM with $k = 3$ communities, $n = 3000$, $c_1 = 0.1$, $c_2 = 0.3$, $c_3 = 0.6$, q_i 's drawn i.i.d. from the measure $\frac{1}{2}\delta_{q_{(1)}} + \frac{1}{2}\delta_{q_{(2)}}$ with affinity matrix \mathbf{M} . (**Left**): $q_{(1)} = 0.8$, $q_{(2)} = 0.9$ and $\mathbf{M} = 10 \cdot \mathbf{I}_3$; (**Right**): $q_{(1)} = 0.1$, $q_{(2)} = 0.9$ and $\mathbf{M} = 5 \cdot \mathbf{I}_3$. Eigenvalue distribution (**top**) and first two eigenvectors of \mathbf{B} (**bottom**). [Link to code]

An intuitive way to reduce this spread is to pre-process the matrix \mathbf{B} in such a way that its spectrum is “as close as possible” to a semicircle. For not too large q_i , $q_i q_j (1 - q_i q_j) \simeq q_i q_j$, and at the same time $d_i / \sqrt{\mathbf{d}^\top \mathbf{1}_n} \simeq q_i$ (see Lemma 7.1 below), so an idea is to pre- and post-multiply \mathbf{B} by \mathbf{D}^{-1} for $\mathbf{D} = \text{diag}\{d_i\}_{i=1}^n$, as proposed in [Coja-Oghlan and Lanka, 2010, Gulikers et al., 2017].

However, while affecting positively the spread of the spectrum of \mathbf{B} , its effect on isolated eigenvalues is not trivial and, as we will see next, may be deleterious. An improved strategy consists in pre- and post-multiplying \mathbf{B} by $\mathbf{D}^{-\alpha}$ for some wisely chosen α and then multiplying the retrieved eigenvectors by $\mathbf{D}^{\alpha-1}$ (see below why). We will hereafter denote

$$\mathbf{L}_\alpha \equiv \frac{(\mathbf{d}^\top \mathbf{1}_n)^\alpha}{\sqrt{n}} \mathbf{D}^{-\alpha} \left(\mathbf{A} - \frac{\mathbf{d} \mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n} \right) \mathbf{D}^{-\alpha} \quad (7.17)$$

for which the normalization by $(\mathbf{d}^\top \mathbf{1}_n)^\alpha$ will appear later to be the natural one. This strategy in particular allows one to retrieve, for $\alpha = 0$ the modularity matrix \mathbf{B} , for $\alpha = 1/2$ the normalized Laplacian matrix [Qin and Rohe, 2013, Chung, 1996]

$$\mathbf{L}_{\frac{1}{2}} = \sqrt{\frac{\mathbf{d}^\top \mathbf{1}_n}{n}} \mathbf{D}^{-\frac{1}{2}} \left(\mathbf{A} - \frac{\mathbf{d} \mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n} \right) \mathbf{D}^{-\frac{1}{2}}$$

and for $\alpha = 1$ the bi-lateral random walk Laplacian matrix [Coja-Oghlan and Lanka, 2010, Gulikers et al., 2017].

$$\mathbf{L}_1 = \frac{\mathbf{d}^\top \mathbf{1}_n}{\sqrt{n}} \left(\mathbf{D}^{-1} \mathbf{A} \mathbf{D}^{-1} - \frac{\mathbf{1}_n \mathbf{1}_n^\top}{\mathbf{d}^\top \mathbf{1}_n} \right).$$

In a similar manner as in the previous decomposition of \mathbf{A} and \mathbf{B} for the Erdős-Rényi and SBM cases, it can be shown (see details in [Ali and Couillet, 2018]) that, in the large n regime,

$$\mathbf{L}_\alpha = \frac{1}{\sqrt{n}} \mathbf{D}_{\mathbf{q}}^{-\alpha} \mathbf{X} \mathbf{D}_{\mathbf{q}}^{-\alpha} + \begin{bmatrix} \mathbf{D}_{\mathbf{q}}^{1-\alpha} \frac{\mathbf{J}}{\sqrt{n}} & \frac{\mathbf{D}_{\mathbf{q}}^{-\alpha} \mathbf{X} \mathbf{1}_n}{\mathbf{q}^\top \mathbf{1}_n} \\ -\mathbf{1}_k^\top & 0 \end{bmatrix} \begin{bmatrix} \mathbf{M}^\circ & -\mathbf{1}_k \\ -\mathbf{1}_k^\top & 0 \end{bmatrix} \begin{bmatrix} \frac{\mathbf{J}^\top \mathbf{D}_{\mathbf{q}}^{1-\alpha}}{\sqrt{n}} \\ \frac{\mathbf{1}_n^\top \mathbf{X} \mathbf{D}_{\mathbf{q}}^{-\alpha}}{\mathbf{q}^\top \mathbf{1}_n} \end{bmatrix} + o_{\|\cdot\|}(1)$$

where we recall $\mathbf{M}^\circ = (\mathbf{I}_k - \mathbf{1}_k \mathbf{c}^\top) \mathbf{M} (\mathbf{I}_k - \mathbf{c} \mathbf{1}_k^\top)$.

We immediately see from this expression that, for high SNRs (i.e., when the nonzero eigenvalues of \mathbf{M}° dominate those of \mathbf{X}), the dominant eigenvectors of \mathbf{L}_α are aligned to the linear combination of the vectors $\mathbf{D}_{\mathbf{q}}^{1-\alpha} \mathbf{j}_a$ for $a = 1, \dots, k$. To retrieve the sought-for \mathbf{j}_a , it is thus necessary to post-process the obtained eigenvectors by $\mathbf{D}_{\mathbf{q}}^{\alpha-1}$ which, in the absence of a perfect knowledge of the vector \mathbf{q} , can be performed empirically by post-processing the eigenvectors by $\mathbf{D}^{\alpha-1}$ instead (see again Lemma 7.1).

The resulting algorithm for spectral clustering under realistic heterogeneous degree graphs is thus summarized as follows:

1. select a scalar $\alpha \in \mathbb{R}$;
2. identify isolated eigenvalues in the spectrum of \mathbf{L}_α defined in (7.17) and extract the corresponding eigenvectors, say $\mathbf{V} = [\mathbf{v}_1, \dots, \mathbf{v}_m] \in \mathbb{R}^{n \times m}$, where $m < k$;
3. perform a k -class k-means (or expectation-maximization) clustering based on the m -dimensional row vectors of the matrix $\mathbf{D}^{\alpha-1} \mathbf{V} \in \mathbb{R}^{n \times m}$.

By an asymptotic analysis similar to the SBM case (see [Ali and Couillet, 2018] for details) as in the previous section, this method is granted to outperform standard spectral clustering approaches. Yet, it remains to properly identify an appropriate value for α . An idea would be to select the value α which maximizes the asymptotic classification performance as $n \rightarrow \infty$: however, this choice strongly depends on \mathbf{M} which is naturally unknown (and cannot be estimated without performing some sort of clustering in the first place).

Instead, we may choose α to be the value for which the “worse case detectability” is achieved. That is, for each α , there exists a smallest value of $\|\mathbf{M}^\circ\|$ for which community detection performs asymptotically better than random guess. We thus decide to choose this value of α such that under the constraint that community detection remains doable, $\|\mathbf{M}^\circ\|$ is the smallest possible. This does not require any information on the actual value of \mathbf{M}° .

To identify this value of α , it suffices to evaluate the limiting spectrum of \mathbf{L}_α and the condition under which (informative) isolated eigenvalue-eigenvectors appear (by solving $\det(\mathbf{L}_\alpha - \lambda \mathbf{I}_n) = 0$ as in the SBM case). The result is summarized as follows.

Theorem 7.3 (Limiting spectrum and isolated eigenvalues for \mathbf{L}_α , [Ali and Couillet, 2018]). *For $\alpha \in \mathbb{R}$, as $n \rightarrow \infty$, the empirical spectrum measure $\mu_{\mathbf{L}}$ of \mathbf{L}_α satisfies $\mu_{\mathbf{L}} - \mu_\alpha \xrightarrow{a.s.} 0$ weakly, where μ_α is defined by its Stieltjes transform $m_{\mu_\alpha}(z)$ as*

$$m_{\mu_\alpha}(z) = \frac{1}{n} \sum_{i=1}^n \frac{1}{-z - g_\alpha(z)q_i^{1-2\alpha} + \tilde{g}_\alpha(z)q_i^{2-2\alpha}}$$

with $(g_\alpha(z), \tilde{g}_\alpha(z))$ the unique solution to

$$\begin{aligned} g_\alpha(z) &= \frac{1}{n} \sum_{i=1}^n \frac{q_i^{1-2\alpha}}{-z - g_\alpha(z)q_i^{1-2\alpha} + \tilde{g}_\alpha(z)q_i^{2-2\alpha}}, \\ \tilde{g}_\alpha(z) &= \frac{1}{n} \sum_{i=1}^n \frac{q_i^{2-2\alpha}}{-z - g_\alpha(z)q_i^{1-2\alpha} + \tilde{g}_\alpha(z)q_i^{2-2\alpha}}. \end{aligned}$$

The limiting spectrum μ_α is continuous and of compact symmetric support $[-S_\alpha, S_\alpha]$. Moreover, if there exists an eigenvalue ℓ of $\mathbf{D}_{\mathbf{c}} \mathbf{M}^\circ$ such that

$$|\ell| > \tau_\alpha \equiv - \lim_{x \downarrow S_\alpha} \frac{1}{\tilde{g}_\alpha(x)}$$

then there exists a corresponding isolated eigenvalue λ_ℓ of \mathbf{L}_α that satisfies

$$\lambda_\ell = \tilde{g}_\alpha^{-1} \left(-\frac{1}{\ell} \right) + o(1).$$

In particular, taking $\alpha = 0$ in the fixed-equation above we obtain (7.16) as a special case.

When compared to the SBM case in Theorem 7.1, the main difference is that the Stieltjes transform m_{μ_α} and its inverse do not assume closed-form formulas.

The optimal value for α discussed above is thus defined as

$$\alpha_* \in \arg \min_{\alpha \in \mathbb{R}} \tau_\alpha$$

where we used an inclusion (rather than equality) sign in case the minimum is not unique (which is for instance the case in the SBM setting where all q_i 's are equal). Under this definition, α_* is indeed the smallest possible phase transition value which ensures in the worst case the existence of isolated eigenvalues, as desired.

From a practical standpoint, of course, since the q_i 's are unknown, it is not possible to identify α_* precisely. Yet, as mentioned several times above, it can be shown that $d_i / \sqrt{\mathbf{d}^\top \mathbf{1}_n} \xrightarrow{a.s.} q_i$ uniformly over i . More specifically, we have

Lemma 7.1. *Under the setting of (7.14), assume that $0 < \liminf_n \min_{1 \leq i \leq n} \{q_i\} \leq \limsup_n \max_{1 \leq i \leq n} \{q_i\} < 1$, then*

$$\max_{1 \leq i \leq n} \left| \frac{d_i}{\sqrt{\mathbf{d}^\top \mathbf{1}_n}} - q_i \right| \xrightarrow{a.s.} 0.$$

It is important to understand here that the condition $[\mathbf{C}]_{ab} = 1 + [\mathbf{M}]_{ab}/\sqrt{n}$ in (7.14) and thus $[\mathbf{C}]_{ab} - [\mathbf{C}]_{a'b'} = O(1/\sqrt{n})$, plays a fundamental role in the above estimate: d_i is, up to scaling, a consistent estimate of q_i , *irrespective* of the class affinities of node i because the difference between the affinities is asymptotically negligible. Note also that the condition for q_i to be bounded away from zero, which ensures that the graph is nowhere sparse, is somewhat limited when applied to realistic graph models (typically having power laws for their degrees [Adamic and Glance, 2005, Borgs et al., 2019]), but is theoretically necessary here.

With Lemma 7.1, one is able to estimate the desired τ_α (in pursuit of the optimal α_*), by substituting the q_i 's in Theorem 7.3 with the estimate $\hat{q}_i = d_i/\sqrt{\mathbf{d}^\top \mathbf{1}_n}$. The last difficulty consists in estimating the right edge S_α of the support as one essentially needs $\lim_{x \downarrow S_\alpha} 1/\tilde{g}_\alpha(x)$. This is unfortunately not easily performed, and to our knowledge there exists no standard estimate of S_α (or of any limiting spectrum edge based on the defining fixed-point equations in general). Numerically, the idea implemented in [Ali and Couillet, 2018] consists in solving the fixed-point equation in $(g_\alpha(x), \tilde{g}_\alpha(x))$ for decreasing values of x until convergence fails (indeed, the fixed-point equation must not have a solution inside the support of μ_α). More practically, a dichotomy approach can be pursued to identify the pivotal value of x for which solving for $(g_\alpha(x), \tilde{g}_\alpha(x))$ becomes possible. This value (the first one for which the fixed-point algorithm does converge) is used as an estimate for S_α .

To evaluate the performance gains incurred by the improved choice of α , Figure 7.3 and Figure 7.4 depict the “overlap” metric (adapted to $k > 2$ classes) Krzakala et al. [2013] defined as¹

$$\text{Overlap} = \frac{\frac{1}{n} \sum_{i=1}^n \delta_{\hat{\mathcal{C}}_i = \mathcal{C}_i} - \frac{1}{k}}{1 - \frac{1}{k}}$$

where $\hat{\mathcal{C}}_i$ is the community allocated by the algorithm to node i and \mathcal{C}_i the associated ground truth, compared for various algorithms (notably against the recent Bethe Hessian approach [Saade et al., 2014] discussed in the next section). Figure 7.3 considers a DCSBM with fixed \mathbf{M} , while 3/4 of the nodes connect with a fixed weight $q_{(1)} = 0.1$ and 1/4 with a higher varying weight $q_{(2)}$. In Figure 7.4, a more realistic synthetic graph setting is considered with the q_i 's following a power law truncated to support on $[0.05, 0.3]$ (to avoid nodes with no neighbors) and with varying \mathbf{M} proportional to the identity matrix. In both

¹Note that this overlap metric is, up to normalization, a generalization of the classification error rate defined in (7.13) under the 2-class symmetric SBM setting.

cases, choosing α optimally (at least in such a way that phase transitions are observed at the lowest values of $\|\mathbf{M}\|$) largely overtakes the performance of standard methods, even beyond the phase transition point.




Figure 7.3: Classification overlap for $n = 3000$, $k = 3$ with $c_1 = c_2 = c_3 = 1/3$, q_i 's i.i.d. with law $\frac{3}{4}\delta_{q(1)} + \frac{1}{4}\delta_{q(2)}$ for $q(1) = 0.1$ and different $q(2)$, \mathbf{M} defined by $[\mathbf{M}]_{ii} = 10$ and $[\mathbf{M}]_{ij} = -10$ for $i \neq j$. Results averaged over 50 runs. [Link to code]

The DC-SBM setting is another telling example of a scenario where the conventional algorithms (here spectral clustering on the adjacency or modularity matrix) may severely fail. Spectral clustering on the matrix $\mathbf{D}^{-\alpha} \mathbf{A} \mathbf{D}^{-\alpha}$ provides a work around, but does not come along with a proof of optimality (more elaborate algorithms may perform better, and even improve the phase transition point).

More generally, another important limitation of the aforementioned analysis of spectral clustering for graphs and data (Section 4.5.1) is that they fundamentally rely on “dense” graphs and affinity matrices, which are possibly unrealistic in practice: e.g., real graphs tend to be rather sparse with each node having a number of neighbors irrespective of the size of the graph [Decelle et al., 2011]. It is indeed central to the random matrix framework that the rows and columns of the adjacency matrices have $O(n)$ degrees of freedom (i.e., are constituted from $O(n)$ independent random variables), so that the matrix itself has $O(n^2)$ degrees of freedom. If instead the number of degrees of freedom per row or column scales as $O(1)$, most random matrix results collapse. Remark for instance that the trace lemma, Lemma 2.11, according to which $\frac{1}{n} \mathbf{x}^\top \mathbf{A} \mathbf{x} \simeq \frac{1}{n} \text{tr } \mathbf{A}$, that is at the core of most of the derivations in this monograph, would no longer be valid if $\mathbf{x} \in \mathbb{R}^n$ had independent Bernoulli entries with parameter $1/n$: in this case, $\mathbb{E}[\mathbf{x}^\top \mathbf{A} \mathbf{x}] = \frac{1}{n} \text{tr } \mathbf{A}$ remains valid but $\mathbf{x}^\top \mathbf{A} \mathbf{x}$ no longer converges; for




Figure 7.4: Overlap for $n = 3\,000$, $k = 3$, $c_1 = c_2 = c_3 = 1/3$, q_i 's following a power law with exponent 3 and support $[0.05, 0.3]$, $\mathbf{M} = \Delta \cdot \mathbf{I}_3$. Here $\hat{\alpha}_* = 0.28$. Circles indicate the theoretical phase transition positions. Results averaged over 50 runs. [Link to code]

instance, we have for $\mathbf{A} = \mathbf{I}_n$ that $\text{Var}[\mathbf{x}^\top \mathbf{x}] = 1 - \frac{1}{n}$ which is of the same order as the mean $\mathbb{E}[\mathbf{x}^\top \mathbf{x}] = 1$ and thus $\mathbf{x}^\top \mathbf{A} \mathbf{x}$ cannot converge.

Handling sparse random matrices requires fundamentally different approaches and the mathematical tools under this setting are not well established yet. These will not be presented in detail in this monograph, as they would demand an altogether very different set of prerequisites. Instead, the subsequent section discusses the few findings arising either from these alternative mathematical tools or, more often, from strikingly different intuitions from statistical physicists (however in general non rigorous).

7.2 From dense to sparse graphs: a different approach

In sparse graph settings, spectral clustering on the adjacency matrix is largely suboptimal, even under a stochastic block model for the graph. This follows from the fact that, for an Erdős-Rényi graph with $O(1)$ node degrees (that is, $\mathbf{A}_{ij} \sim \text{Bern}(p/n)$ where $p = O(1)$), the limiting spectrum of \mathbf{A} is no longer a semicircle law. Surprisingly enough, while a limiting spectrum does exist, very little is known about it. The main (striking) result obtained so far is that, as opposed to the semicircle law, the limiting spectrum has an *unbounded support* and has localized point masses [Salez, 2019].

The unboundedness of the support in the sparse regime is problematic for

spectral clustering in presence of communities and explains why spectral clustering on \mathbf{A} (or the modularity matrix \mathbf{B}) is bound to fail. One must then devise other methods and possibly find alternative matrices to the adjacency.

7.2.1 The non-backtracking matrix

The first convincing idea arose from a statistical physics interpretation [Krzakala et al., 2013]: the nodes of a graph may be seen as interacting particles with interaction strength given by the entries of the adjacency matrix (in the binary case, particles i and j interact if $\mathbf{A}_{ij} = 1$). If let free of external “force fields”, the system tends to minimize its energy, which corresponds to falling into a state of (local) maximal probability. By establishing expressions for the probability of each state and performing linear approximations around the so-called “ground state” solution, it appears that the dominant eigenvectors of the so-called *non-backtracking matrix* must be correlated to the communities of the graph. The non-backtracking matrix \mathbf{N} is defined on the set \mathcal{E} of edges of the graph as

$$\mathbf{N}_{(ij)(kl)} = \delta_{jk}(1 - \delta_{il}), \quad \forall (ij), (kl) \in \mathcal{E}. \quad (7.18)$$

It is thus a non-symmetric matrix. Its limiting spectrum is mostly unknown but, in the SBM case, it has been importantly known that all eigenvalues are asymptotically found inside a disc of controlled radius, with a possible exception for finitely many real eigenvalues of larger amplitude: the associated eigenvectors are those correlated to the communities. Precisely, letting \mathbf{v} be such an eigenvector (of size the number of edges in the graph), the vector $\tilde{\mathbf{v}} \in \mathbb{R}^n$ defined by

$$\tilde{\mathbf{v}}_i = \sum_{j \in \partial i} \mathbf{v}_{(ij)}, \quad \partial i \equiv \{j \mid (i, j) \in \mathcal{E}\} \quad (7.19)$$

provides a clustering vector of the graph communities. As in the dense setting, the presence of isolated eigenvalues is ruled by a phase transition phenomenon. In the symmetric SBM where $\mathbf{A}_{ij} \sim \text{Bern}(p_{\text{in}}/n)$ if i, j are in the same community and $\mathbf{A}_{ij} \sim \text{Bern}(p_{\text{out}}/n)$ otherwise, this phase transition has been rigorously proven in [Mossel et al., 2015, Massoulié, 2014] to occur when

$$\frac{|p_{\text{in}} - p_{\text{out}}|}{\sqrt{\frac{1}{2}(p_{\text{in}} + p_{\text{out}})}} > 2.$$

Note in particular that, as opposed to the dense regime in (7.12) where $(p_{\text{in}} - p_{\text{out}})/(p_{\text{in}} + p_{\text{out}})$ would be requested to scale as $O(n^{-1/2})$, here it is necessary to have $(p_{\text{in}} - p_{\text{out}})/(p_{\text{in}} + p_{\text{out}}) = O(1)$ for communities to arise in spectral clustering: in the absence of strong redundancy, the classification task is thus, not surprisingly, much harder.

The non-backtracking approach is however quite expensive to implement as the matrix is non-symmetric and possibly of large dimensions (scaling as the number of edges rather than the number of nodes in the graph). Also, the vector




Figure 7.5: Complex spectrum of the non-backtracking matrix \mathbf{N} ; $n = 1\,000$, $p_{\text{in}} = 12$, $p_{\text{out}} = 1$. Emphasized in red circles are the two informative eigenvalues. [Link to code]

$\tilde{\mathbf{v}}$ defined in (7.19), while correlated to the node classes, is seen in simulations to be largely affected by heterogeneity in the node degrees: that is, it becomes largely inconsistent beyond the SBM setting.

It in fact turns out though that the spectrum of the non-backtracking matrix is intimately related to that of another matrix, called the Bethe Hessian matrix, also familiar of statistical physicists and which, as shown later in Section 7.2.4, can be exploited to naturally fight against degree heterogeneity.

7.2.2 The Bethe Hessian matrix

It can be shown that, for $\mathbf{N}\mathbf{v} = \gamma\mathbf{v}$, letting $\tilde{\mathbf{v}}_i = \sum_{j \in \partial_i} \mathbf{v}_{(ij)}$ as in (7.19),

$$((\gamma^2 - 1)\mathbf{I}_n + \mathbf{D} - \gamma\mathbf{A})\tilde{\mathbf{v}} = \mathbf{0}.$$

Thus, $\tilde{\mathbf{v}}$ is also an eigenvector (associated to a zero eigenvalue) of the so-called *Bethe Hessian* matrix

$$\mathbf{H}_\gamma \equiv (\gamma^2 - 1)\mathbf{I}_n + \mathbf{D} - \gamma\mathbf{A}. \quad (7.20)$$

The parameter γ defining \mathbf{H}_γ is however unknown, since it requires to solve an eigenvector equation for \mathbf{N} , which we precisely would like to avoid.

But the Bethe Hessian \mathbf{H}_γ also finds a parallel origin, again from a statistical physics interpretation: the isolated eigenvectors of \mathbf{H}_γ (associated to the *smallest* eigenvalues) correspond to particle states of minimal *Bethe free energy*, where $1/\gamma$ is the *temperature* of the system of interacting particles. Under this interpretation, [Saade et al., 2014] heuristically propose to chose $\gamma = \sqrt{\rho(\mathbf{N})}$ with $\rho(\mathbf{N})$ the spectral radius (largest eigenvalue in amplitude) of \mathbf{N} and to perform clustering on the eigenvector associated with the *second smallest* eigenvalue

of \mathbf{H}_γ . Figure 7.6 reports the histogram of the eigenvalues and the informative eigenvector of \mathbf{H}_γ for two choices of γ .

In the specific case of SBM, the choice $\gamma = \sqrt{\rho(\mathbf{N})}$ corresponds in the limit to $\gamma = \sqrt{(p_{\text{in}} + p_{\text{out}})/2}$. This choice of γ , inspired by a SBM analysis, seems indeed rather optimal in this setting. Yet, the same remark on degree heterogeneity reported for the non-backtracking matrix still holds here: for $\gamma = \sqrt{(p_{\text{in}} + p_{\text{out}})/2}$, spectral clustering on \mathbf{H}_γ is tainted by the heterogeneity of node degrees and therefore appears to be suboptimal for DC-SBM as in the left plot of Figure 7.6.

7.2.3 Degree regularization

An alternative approach to improve the adjacency matrix \mathbf{A} or the various normalized Laplacian matrices $\mathbf{D}^{-1}\mathbf{A}$ or $\mathbf{D}^{-\frac{1}{2}}\mathbf{A}\mathbf{D}^{-\frac{1}{2}}$ consists in observing that their main defect in dealing with sparse graphs is due to: (i) the instability in \mathbf{D}^{-1} caused by nodes with low connectivity and (ii) the existence of spurious “hubs”, that is, nodes i with exceptionally high degrees (very rare in dense graphs but not so uncommon in sparse graphs), which “pull” their own eigenvectors ($\mathbf{v}_j \simeq \delta_{ij}$).

The non-backtracking matrix \mathbf{N} precisely handles item (ii) by reducing the number of rows with large “degrees” (through the non-backtracking walks that escape hubs without returning to them).

Alternatively, several authors proposed (heuristically) to correct the adjacency or normalized Laplacian matrices by adding a regularization term: e.g., $\mathbf{A}_\tau = \mathbf{A} + \tau \mathbf{1}_n \mathbf{1}_n^\top$ in [Amini et al., 2013] or $\mathbf{L}_\tau = (\mathbf{D} + \tau \mathbf{I}_n)^{-\frac{1}{2}} \mathbf{A} (\mathbf{D} + \tau \mathbf{I}_n)^{-\frac{1}{2}}$ in [Qin and Rohe, 2013]. In the latter, the authors, still heuristically, propose to take $\tau = (p_{\text{in}} + p_{\text{out}})/2$ (despite their few theoretical results suggesting to take much larger values for τ).

Interestingly, as opposed to the Bethe Hessian and non-backtracking methods described above which sometimes fail on realistic (especially heterogeneous) graphs, spectral clustering on \mathbf{L}_τ with this particular choice of τ is seen in simulations to be extremely efficient and resilient to realistic graph clustering.

7.2.4 A unifying approach adapted to DC-SBM

In [Dall’Amico et al., 2019], a unified approach is proposed to explain how the Bethe Hessian and regularized Laplacian relate to each other, and most importantly, to provide control of the hyperparameters γ and τ (for Bethe Hessian and regularized Laplacian, respectively) that make spectral clustering insensitive to degree heterogeneity.

The authors simply observe that, in a two-class symmetric DC-SBM setting, letting $\mathbf{j} = [\mathbf{1}_{n/2}, -\mathbf{1}_{n/2}]^\top$, one has

$$[(\mathbf{D} - \gamma \mathbf{A})\mathbf{j}]_i = d_i [\mathbf{j}]_i \left[1 - \gamma \left(\frac{|\partial_i^{(\text{in})}|}{d_i} - \frac{|\partial_i^{\text{out}}|}{d_i} \right) \right]$$

with $\partial_i^{(\text{in})}$ the nodes connected to i within the same community and $\partial_i^{(\text{out})}$ the nodes connected to i within the other community. Assuming the average degree not too small, this gives

$$[(\mathbf{D} - \gamma \mathbf{A})\mathbf{j}]_i \simeq d_i[\mathbf{j}]_i \left(1 - \gamma \cdot \frac{p_{\text{in}} - p_{\text{out}}}{p_{\text{in}} + p_{\text{out}}}\right).$$

Thus, \mathbf{j} is an approximate eigenvector of $\mathbf{D} - \gamma \mathbf{A}$ if one chooses

$$\gamma = \frac{p_{\text{in}} + p_{\text{out}}}{p_{\text{in}} - p_{\text{out}}}.$$

As opposed to the regularization values ($\gamma = \sqrt{(p_{\text{in}} + p_{\text{out}})/2}$ and τ) in the previous sections, it is interesting to note that the newly proposed γ depends on the clustering task difficulty (via $p_{\text{in}} - p_{\text{out}}$).




Figure 7.6: Histogram of the empirical spectral distribution of \mathbf{H}_γ and informative eigenvector under a DC-SBM model for (left) $\gamma = \sqrt{(p_{\text{in}} + p_{\text{out}})/2}$ and (right) $\gamma = (p_{\text{in}} + p_{\text{out}})/(p_{\text{in}} - p_{\text{out}})$. Here $n = 1000$ nodes, $p_{\text{in}} = 35$, $p_{\text{out}} = 5$ and $\mathbf{q} = [\text{linspace}(0.2, 0.9), \text{linspace}(0.2, 0.9)]$. [Link to code]

Since $\mathbf{D} - \gamma \mathbf{A}$ has the same eigenvectors as \mathbf{H}_γ per (7.20), this choice of γ offers a new value for the Bethe Hessian parameter which is now insensitive to degree heterogeneity. Yet, as opposed to $\gamma = \sqrt{(p_{\text{in}} + p_{\text{out}})/2}$ which can be estimated consistently by evaluating the average node degree in the graph, $\gamma = (p_{\text{in}} + p_{\text{out}})/(p_{\text{in}} - p_{\text{out}})$ cannot be directly estimated from the graph (since p_{in} and p_{out} are unknown). Nonetheless, Dall'Amico et al. [2019] showed that $\gamma = (p_{\text{in}} +$

$p_{\text{out}})/(p_{\text{in}} - p_{\text{out}})$ corresponds (asymptotically) to the smallest value for which $\lambda_2(\mathbf{H}_\gamma) = 0$ (with $\lambda_2(\cdot)$ the second smallest eigenvalue). The eigenvector \mathbf{v} carrying the class information is then the one associated with the zero eigenvalue of \mathbf{H}_γ (i.e., such that $\mathbf{H}_\gamma \mathbf{v} = \mathbf{0}$). The right-hand side display of Figure 7.6 demonstrates the better resilience of this choice to graph heterogeneity.

In a k -class setting, it is similarly shown that spectral clustering can be performed, no longer on a single matrix \mathbf{H}_γ , but on the matrices $\mathbf{H}_{\gamma_2}, \dots, \mathbf{H}_{\gamma_k}$ where γ_p is the value of γ such that $\lambda_p(\mathbf{H}_\gamma) = 0$ (with $\lambda_p(\cdot)$ the p -th smallest eigenvalue) and the corresponding informative eigenvector \mathbf{v}_p is the one for which $\mathbf{H}_\gamma \mathbf{v}_p = \mathbf{0}$.

Besides, it is observed that the following two equations are equivalent:

$$[(\gamma_p^2 - 1)\mathbf{I}_n + \mathbf{D} - \gamma_p \mathbf{A}] \mathbf{v}_p = \mathbf{0} \Leftrightarrow (\mathbf{D} + (\gamma_p^2 - 1)\mathbf{I}_n)^{-1} \mathbf{A} \mathbf{v}_p = \frac{\mathbf{v}_p}{\gamma_p}, \quad (7.21)$$

meaning that \mathbf{v}_p is also an eigenvector of the *regularized* random-walk Laplacian $(\mathbf{D} + (\gamma_p^2 - 1)\mathbf{I}_n)^{-1} \mathbf{A}$. Since the eigenvalues of the latter are the same as the eigenvalues of $(\mathbf{D} + (\gamma_p^2 - 1)\mathbf{I}_n)^{-\frac{1}{2}} \mathbf{A} (\mathbf{D} + (\gamma_p^2 - 1)\mathbf{I}_n)^{-\frac{1}{2}}$ and that the associated eigenvectors are just scaled by the normalized degrees, we also find a natural connection to the regularized Laplacian matrix of [Qin and Rohe, 2013] discussed in the previous section.

Using an efficient procedure to estimate the number of classes \hat{k} and the values $\gamma_2, \dots, \gamma_{\hat{k}}$ (without resorting to expensive line searches), Dall'Amico et al. [2019] provide a comparative performance table of all aforementioned spectral clustering procedures on real benchmark graphs. This is reported in Table 7.1, in which \bar{d} denotes the average node degree.

Dataset	n	d	k	$\{\mathbf{H}_{\gamma_p}\}$	\mathbf{A}	\mathbf{H}_γ	\mathbf{N}	$\mathbf{D}^{-1}\mathbf{A}$	\mathbf{L}_τ
Dolphins	62	5	<u>2</u>	0.38	0.21	0.34	0.22	0.38	0.38
Polbook	105	8.4	<u>3</u>	0.5	0.44	0.5	0.45	0.5	0.5
Mail	1133	9.6	21	0.5	0.32	0.4	0.37	0.5	0.5
Polblogs	1222	27.4	<u>2</u>	0.43	0.23	0.27	0.24	0	0.43
Tv	3892	8.9	41	0.8	0.51	0.58	0.55	0.55	0.8
Facebook	4039	43.7	55	0.78	0.43	0.49	0.49	0.78	0.57
Power grid	4941	2.7	25	0.93	0.18	0.37	0.31	0.93	0.85
GrQc	5242	5.5	29	0.53	0.45	0.49	0.49	0.42	0.79
Politicians	5908	14.1	62	0.85	0.48	0.54	0.5	0.83	0.74
GNutella P2P	6301	6.6	4	0.26	0.16	0.14	0.19	0	0.35
Wikipedia	7115	28.3	22	0.23	0.15	0.17	0.17	0.23	0.27
Vip	11565	11.6	53	0.62	0.27	0.33	0.3	0.55	0.54
HepPh	12008	19.7	60	0.37	0.42	0.42	0.42	0.11	0.52
Croatia	57573	18.3	<u>84</u>	0.65	0.33	0.39	0.34	0.69	0.62

Table 7.1: Modularity comparison on real networks [Leskovec and Krevl, 2014]. k (unless underlined) and ground truth labels are unknown. Special emphasis is made on $\{\mathbf{H}_{\gamma_p}\}_{p=1}^{\hat{k}}$ proposed in [Dall’Amico et al., 2019] (to be distinguished from the “classical” \mathbf{H}_γ with $\gamma = \sqrt{(p_{in} + p_{out})/2}$), which outperforms most competing approaches. (Here for \mathbf{L}_τ we choose $\tau = \frac{1}{n}\mathbf{1}_n^\top \mathbf{A} \mathbf{1}_n$, the average degree.)

These studies remain largely at a heuristic level. To the noticeable exception of [Massoulié, 2014] which theoretically proves that the phase transition proposed by the physicists for the non-backtracking operator is indeed optimal, very few random matrix analyses exist that are able to tackle the spectrum of sparse graphs. Here, Stieltjes transform approaches collapse, and are mostly replaced by more burdensome combinatorics and random graph techniques.

Yet, the analysis of sparse graphs is fundamental for at least two reasons: (i) as said, the reality of real networks tends more towards the sparse than the dense side, and (ii) sparsification techniques may also be used in practice to reduce computational costs: in clustering data using kernel methods, one may use k-NN (k-nearest neighbors) with a relatively small value of k, or alternatively only compute few entries of the kernel matrix. The spectral and, more generally, algorithmic implications of sparse data and sparsification procedures will surely be a subject of active future interest in large dimensional statistics and random matrices for machine learning.

7.3 Concluding remarks

Spectral methods for community detection are the “Wigner semicircle” counterpart of spectral clustering for large dimensional data (which, in its simplest setting, is the “Wishart Marčenko-Pastur” equivalent) discussed in Chapter 4. The random matrix tools and proof methods being equally applicable to each setting, their ultimate study is quite similar.

A second difference relates to the matrix entries: the entries of the adjacency matrices are typically Bernoulli distributed (at least in unweighted graphs) where instead kernel matrices tend to be filled with continuous variables (aside from k-NN kernels). Nonetheless, in dense (or moderately dense) graphs, from the universality of random matrix results, this difference also vanishes. In particular, the case of weighted dense graphs (including the DC-SBM), although not quite studied in the literature, would be easily handled.

Major differences start to appear when considering sparse graph settings. It is likely that the limiting spectral measure of the adjacency matrix \mathbf{A} of such a graph, as well as its associated Laplacian, depends on the law of the entries beyond its first and second moments. This may be understood from the fact that the columns of \mathbf{A} no longer “concentrate” in the sparse regime (e.g., their norm does not converge but remains a random variable fully dependent on the law of the entries). This behavior, although possibly averaged over the columns to some extent, breaks the convenient universality phenomena arising in dense random matrices.

An alternative approach to “partially” account for the sparse regime while remaining tractable to classical tools in random matrix theory is to assume that the average degree of the nodes in the graph grows slowly (e.g., as $O(\log n)$) with the size n of the graph. In doing so, slow convergence behavior appears, with Wigner’s semicircle law being valid again. The major problem though is that, under the classical SBM, for which $p_{\text{in}} - p_{\text{out}} = O(1)$, classification

becomes asymptotically trivial: that is, the dominant eigenvalue of \mathbf{A} grows unbounded, yet at a very slow rate, as the graph size n grows large. Studying this setting remains interesting, as one is able to precisely “control” the evolution, for all finite but large n , of the spectrum of \mathbf{A} and of the Laplacian, Bethe Hessian, non-backtracking matrices, etc. Under the not completely unsatisfying $O(\log n) \approx O(1)$ approximation, for large but finite size n , these studies may provide a sufficiently accurate picture of the behavior of real sparse graphs. This path is currently at the central focus of many random matrix researches for graphs, see e.g., [Coste and Zhu, 2020] in which results on the position of the real eigenvalues of the non-backtracking matrix are “tracked”.

7.4 Practical course material

Practical Lecture Material 7 (Asymptotic Gaussian fluctuations of the SBM dominant eigenvector). *This exercise aims to complete Remark 7.1 on the asymptotic joint Gaussian fluctuations of (the entries of) the dominant eigenvector for the modularity matrix $\mathbf{B} = \frac{1}{\sqrt{n}}(\mathbf{A} - \frac{\mathbf{d}\mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n})$ in a stochastic block model setting under (7.3), thereby leading to the asymptotic classification error rate in the form of Gaussian Q -function.*

We consider for simplicity the case of two classes $\mathcal{C}_1, \mathcal{C}_2$ with $|\mathcal{C}_1| = |\mathcal{C}_2|$ where the affinity matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ has i.i.d. Bernoulli entries $[\mathbf{A}]_{ij} \sim \text{Bern}(\mathbf{C}_{ab})$ with [node i] $\in \mathcal{C}_a$, [node j] $\in \mathcal{C}_b$ and $\mathbf{C} = [\begin{smallmatrix} p_{\text{in}} & p_{\text{out}} \\ p_{\text{out}} & p_{\text{in}} \end{smallmatrix}]$, where $p_{\text{in}} - p_{\text{out}} = O(n^{-1/2})$. We denote $\mathbf{d} = [d_1, \dots, d_n]^\top$ where $d_i = \sum_j [\mathbf{A}]_{ij}$.

For $\mathbf{B} = \frac{1}{\sqrt{n}}(\mathbf{A} - \frac{\mathbf{d}\mathbf{d}^\top}{\mathbf{d}^\top \mathbf{1}_n})$, first establish, from the results of this chapter (in particular (7.6) and (7.11)) that

$$\begin{aligned} \frac{1}{\sqrt{p_{\text{out}}(1-p_{\text{out}})}} \mathbf{B} &= \frac{\mathbf{X}}{\sqrt{n}} + \frac{p_{\text{in}} - p_{\text{out}}}{2\sqrt{n}\sqrt{p_{\text{out}}(1-p_{\text{out}})}} \mathbf{j}\mathbf{j}^\top \\ &\quad - \left(\frac{\mathbf{1}_n \mathbf{1}_n^\top}{n} \frac{\mathbf{X}}{\sqrt{n}} + \frac{\mathbf{X}}{\sqrt{n}} \frac{\mathbf{1}_n \mathbf{1}_n^\top}{n} \right) + O_{\|\cdot\|}(n^{-\frac{1}{2}}) \end{aligned}$$

where $\mathbf{j} = [\mathbf{1}_{n/2}, -\mathbf{1}_{n/2}]^\top \in \mathbb{R}^n$ and \mathbf{X} is a symmetric matrix with i.i.d. zero mean and unit variance entries. Since $p_{\text{in}} - p_{\text{out}} = O(n^{-1/2})$, we focus on the matrix

$$\mathbf{Y} \equiv \frac{\mathbf{X}}{\sqrt{n}} + \frac{\gamma}{n} \mathbf{j}\mathbf{j}^\top - \left(\frac{\mathbf{1}_n \mathbf{1}_n^\top}{n} \frac{\mathbf{X}}{\sqrt{n}} + \frac{\mathbf{X}}{\sqrt{n}} \frac{\mathbf{1}_n \mathbf{1}_n^\top}{n} \right) \quad (7.22)$$

for some $\gamma > 0$ and \mathbf{X} an arbitrary random matrix with i.i.d. zero mean and unit variance entries. We further assume γ large enough (in fact $\gamma > 1$) so that an isolated eigenvalue-eigenvector pair $(\hat{\lambda}, \hat{\mathbf{u}})$ emerges (almost surely) in the spectrum of \mathbf{Y} .

Using $\mathbf{j}^\top \mathbf{1}_n = 0$, establish that $\mathbf{1}_n$ is an eigenvector of \mathbf{Y} associated with an eigenvalue tending to zero (as $n \rightarrow \infty$) and conclude that $\hat{\mathbf{u}}^\top \mathbf{1}_n = 0$.

Based on this observation and $\mathbf{Y}\hat{\mathbf{u}} = \hat{\lambda}\hat{\mathbf{u}}$, show that

$$\hat{\mathbf{u}} = \frac{\mathbf{1}_n^\top \mathbf{X} \hat{\mathbf{u}}}{n\sqrt{n}} \mathbf{Q}(\hat{\lambda}) \mathbf{1}_n - \frac{\gamma \mathbf{j}^\top \hat{\mathbf{u}}}{n} \mathbf{Q}(\hat{\lambda}) \mathbf{j}$$

where $\mathbf{Q}(z) \equiv (\frac{\mathbf{X}}{\sqrt{n}} - z\mathbf{I}_n)^{-1}$ and that, in particular, for $\mathbf{e}_i \in \mathbb{R}^n$ the canonical basis vector with $[\mathbf{e}_i]_j = \delta_{ij}$,

$$\sqrt{n} \cdot [\hat{\mathbf{u}}]_i = \frac{\mathbf{1}_n^\top \mathbf{X} \hat{\mathbf{u}}}{n} \mathbf{e}_i^\top \mathbf{Q}(\hat{\lambda}) \mathbf{1}_n - \frac{\gamma \mathbf{j}^\top \hat{\mathbf{u}}}{\sqrt{n}} \mathbf{e}_i^\top \mathbf{Q}(\hat{\lambda}) \mathbf{j}. \quad (7.23)$$

Following a spiked model approach as in (7.12), establish that

$$\hat{\lambda} = \lambda + o(1) \equiv \gamma + \frac{1}{\gamma} + o(1), \quad \left| \frac{1}{\sqrt{n}} \mathbf{j}^\top \hat{\mathbf{u}} \right| = \sqrt{1 - \frac{1}{\gamma^2}} + o(1), \quad (7.24)$$

almost surely, for $\lambda = \gamma + \gamma^{-1}$, as well as

$$\frac{1}{\sqrt{n}} \mathbf{1}_n^\top \mathbf{X} \hat{\mathbf{u}} \xrightarrow{a.s.} 0. \quad (7.25)$$

Note in particular that the behavior of $\hat{\lambda}$ and $|\mathbf{j}^\top \hat{\mathbf{u}}/\sqrt{n}|$ is the same asymptotically as the isolated eigenvalue-eigenvector pair of the model $\frac{\mathbf{X}}{\sqrt{n}} + \frac{\gamma}{n} \mathbf{j} \mathbf{j}^\top$: that is, the additional term $\mathbf{1}_n \mathbf{1}_n^\top \mathbf{X} + \mathbf{X} \mathbf{1}_n \mathbf{1}_n^\top$ in (7.22) has asymptotically no impact on the spectrum of \mathbf{Y} .

Next, using the resolvent identity, Lemma 2.1, establish for any deterministic vectors $\mathbf{v}_1, \mathbf{v}_2$ and scalars ℓ_1, ℓ_2 that

$$\mathbf{v}_1^\top \mathbf{Q}(\ell_1) \mathbf{v}_2 = \mathbf{v}_1^\top \mathbf{Q}(\ell_2) \mathbf{v}_2 - (\ell_1 - \ell_2) \mathbf{v}_1^\top \mathbf{Q}(\ell_1) \mathbf{Q}(\ell_2) \mathbf{v}_2.$$

Next, we would like to show that the bilinear form $\mathbf{e}_i^\top \mathbf{Q}(\hat{\lambda}) \mathbf{1}_n / \sqrt{n}$ is of order $O(1)$, which, together with (7.25), allows us to (asymptotically) discard the first term in (7.23). To this end, note that the term $\mathbf{e}_i^\top \mathbf{Q}(\hat{\lambda}) \mathbf{1}_n / \sqrt{n}$, despite being a “classical” bilinear form, contains a “vanishing” dependence between \mathbf{X} and $\hat{\lambda}$ (as an eigenvalue of \mathbf{Y} that depends on \mathbf{X}). To get rid of this dependence, establish, with the resolvent identity, Lemma 2.1, that for any deterministic vectors $\mathbf{v}_1, \mathbf{v}_2$ and scalars ℓ_1, ℓ_2

$$\mathbf{v}_1^\top \mathbf{Q}(\ell_1) \mathbf{v}_2 = \mathbf{v}_1^\top \mathbf{Q}(\ell_2) \mathbf{v}_2 - (\ell_1 - \ell_2) \mathbf{v}_1^\top \mathbf{Q}(\ell_1) \mathbf{Q}(\ell_2) \mathbf{v}_2$$

so that $|\mathbf{e}_i^\top \mathbf{Q}(\hat{\lambda}) \mathbf{1}_n - \mathbf{e}_i^\top \mathbf{Q}(\lambda) \mathbf{1}_n| / \sqrt{n} \leq |\hat{\lambda} - \lambda| \cdot \|\mathbf{Q}(\hat{\lambda}) \mathbf{Q}(\lambda)\|$ for $\lambda = \gamma + \gamma^{-1}$. Apply similarly this result to $\mathbf{e}_i^\top \mathbf{Q}(\hat{\lambda}) \mathbf{j}$.

Assume for the moment that \mathbf{X} has independent $\mathcal{N}(0, 1)$ entries and show, by a diagonalization of \mathbf{X} (recalling that eigenvalues and eigenvectors are independent in the Gaussian case) and a standard central limit theorem argument, that

$$\frac{1}{\sqrt{n}} \mathbf{e}_i^\top \mathbf{Q}(z) \mathbf{1}_n = \mathcal{N}(m(z), m'(z) - m(z)^2) + O(n^{-1})$$

in probability, for $m(z)$ the Stieltjes transform of the semicircle law (Theorem 2.5) and $m'(z)$ its derivative. Confirm by a universality argument that this result holds true beyond the Gaussian distribution, so long that \mathbf{X} has entries of zero mean and unit variance.

Since $m^2(z) + zm(z) + 1 = 0$ and that $m(\lambda) = -1/\gamma$, conclude that

$$\sqrt{n} \cdot [\hat{\mathbf{u}}]_i = \pm \sqrt{1 - \gamma^{-2}} + \gamma^{-1} w_i + o(1)$$

where $w_i \sim \mathcal{N}(0, 1)$.

Generalize now this result to a d -dimensional setting by showing that, for any k entries $[\hat{\mathbf{u}}]_{i_1}, \dots, [\hat{\mathbf{u}}]_{i_d}$ of $\hat{\mathbf{u}}$, with the same line of arguments

$$\sqrt{n} \begin{bmatrix} [\hat{\mathbf{u}}]_{i_1} \\ \vdots \\ [\hat{\mathbf{u}}]_{i_d} \end{bmatrix} = \pm \sqrt{1 - \gamma^{-2}} \cdot \mathbf{1}_d + \gamma^{-1} \mathbf{w} + o(1)$$

where $\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_k)$. In particular, the fluctuations of the entries of $\hat{\mathbf{u}}$ are asymptotically decorrelated under the SBM setting.

Conclude, from this result an Item 2 of Remark 7.1, that the probability of misclassification is asymptotically given by $Q(\sqrt{\gamma^2 - 1})$, with $Q(\cdot)$ the Gaussian Q -function, and translate this result in terms of the parameters p_{out} and $\sqrt{n}(p_{\text{in}} - p_{\text{out}})$ to recover, as expected, the asymptotic error rate

$$Q \left(\sqrt{\frac{n(p_{\text{in}} - p_{\text{out}})^2}{4p_{\text{out}}(1 - p_{\text{out}})}} - 1 \right)$$

established in Equation (7.13).

Chapter 8

Discussions on Universality and Practical Applications

8.1 From Gaussian mixtures to concentrated random vectors and GAN images

8.1.1 On data models in large dimensions

In the previous chapters, we have repeatedly worked under the assumption that data arise from a Gaussian mixture model to elaborate asymptotic performance analyses of a wide range of machine learning algorithms. This assumption primarily arises for mathematical convenience: the Gaussian model has many mathematical virtues: it is parameterized only through its first two moments, specific mathematical tools are available, Gaussian vectors are (up to centering and scaling) vectors with independent entries, etc.

From a small dimensional viewpoint (p small and fixed), it is clear that Gaussian vectors $\mathbf{x} \in \mathbb{R}^p$ are extremely limited models for most realistic datasets: Gaussian vectors of small dimensions are restricted to ellipsoid-shaped distributions and cannot account for the possibly complex dependence relations between the entries of \mathbf{x} . By extrapolation, the many possible interactions between the entries of a large dimensional vector \mathbf{x} are even less prone to modeling by means of a Gaussian vector.

Yet, we have seen in the previous chapters a systematic, sometimes seemingly perfect, match between the performance machine learning algorithms achieve on *real datasets* and those predicted on Gaussian (mixture) models sharing the same statistical means and covariances as the real data (these means and covariances being estimated empirically from the whole dataset).

The objective of this chapter is to demonstrate that this is far from a coincidence. It is indeed possible to prove mathematically that many of the results in this monograph *do extend* to a wide range of “almost” real data. More specifi-

cally, we will successively show in this chapter that

- as already hinted at in Theorem 2.18 which proves that Theorem 2.6 not only holds for vectors with independent entries (up to centering and scaling), but also for the much larger class of *concentrated random vectors*, many core results from the previous chapters *hold* almost identically under a data model of (mixture of) concentrated random vectors. In particular, it appears that the salient information, that dictates the behavior of most machine learning algorithms, lies in the first two statistical moments: those are sufficient to capture the essence of most learning mechanisms;
- the class of concentrated random vectors naturally contains all random vectors arising from a Lipschitz transformation of large standard Gaussian vectors, which in particular comprises all random vectors produced by generative adversarial networks (better known as GANs, i.e., deep neural networks that are designed to generate fake, but extremely close to real, data) [Goodfellow et al., 2014]: as a consequence of the previous item, *the performance of many machine learning algorithms on data produced by GANs is theoretically predictable*;
- extensive simulations have been run on state-of-the-art classification frameworks for real versus GAN-generated data: while the performance are not identical between GAN and real data (GAN data are easier to discriminate), the *theoretical performance predicted by random matrix theory on real data* are indeed a systematically accurate match to the actual performance.

From these observations, a careless conclusion may be to claim that Gaussian (mixture) vectors are accurate models for real data. In a way, this hasty conclusion is not necessarily inappropriate: it all depends on what is meant by “an accurate model”. If a model is appropriate because a human observer (or a machine) cannot distinguish real data from the model (as GANs have been designed to do), then Gaussian models are not accurate: Figure 8.1 evidences this fact by comparing digits from the MNIST database to their Gaussian model counterpart.

But, if an “accurate model” is defined as correctly *testifying of the performance of a given data processing method* on real data, then, as we already saw and will see next in more detail, the Gaussian model is quite accurate when studying a host of classification and regression problems in machine learning.

The conclusion here is quite fundamental to the vision of machine learning methods for (not so) large dimensional data: the conservative approach according to which real data need be appropriately modeled from a human eye standpoint to be worth theoretical analysis, reducing Gaussian vectors to “toy examples”, is strikingly disrupted in large dimensions. For large data, Gaussian models are often more than enough to account for the behavior of statistical learning mechanisms.




Figure 8.1: Images of digit “1” and “2” from the MNIST database (**left**) and random Gaussian generation from a model with the same mean and covariance (**right**), empirically estimated from all digits from the entire MNIST database. [\[Link to code\]](#)

8.1.2 A study of GAN data

8.1.2.1 Reminders on deep neural networks and GANs

The field of computer vision has recently experienced two successive tidal waves that brushed aside (i) years of conventional mathematical research in image classification with the arrival of deep convolutional networks [Krizhevsky et al., 2017], the performance of which is now near superhuman in some tasks (while previous, e.g., wavelet-based approaches, were far below human performance), and (ii) the conventional thinking that modern computers could not generate arbitrary samples of deceptively realistic images, here with the construction of generative adversarial networks (GANs) [Goodfellow et al., 2014], which are merely two competing instances of deep (convolutional) networks.

For a reminder, a neural network is a succession of L linear and entry-wise nonlinear maps, associating input datum $\mathbf{x} \in \mathbb{R}^p$ to an output $\mathbf{z} = \phi(\mathbf{x}) \in \mathbb{R}^q$ as

$$\mathbf{z} = \phi(\mathbf{x}) = \sigma_L(\mathbf{W}_L \sigma_{L-1}(\mathbf{W}_{L-1} \dots \sigma_1(\mathbf{W}_1 \mathbf{x}) \dots)) \quad (8.1)$$

with $\mathbf{W}_i \in \mathbb{R}^{l_i \times l_{i-1}}$ the linear maps (sometimes with additional bias terms $\mathbf{b}_i \in \mathbb{R}^{l_i}$) and $\sigma_i : \mathbb{R} \rightarrow \mathbb{R}$ the nonlinear maps applied entry-wise. Based on a (usually quite numerous) sequence of known input-output pairs $(\mathbf{x}_i, \mathbf{z}_i)$ and from a random initialization of the weights $\mathbf{W}_1, \dots, \mathbf{W}_L$, neural networks adapt these weights (for fixed σ_i) by running gradient descent to minimize some loss function of the type

$$\frac{1}{n} \sum_{i=1}^n \ell(\sigma_L(\mathbf{W}_L \sigma_{L-1}(\mathbf{W}_{L-1} \dots \sigma_1(\mathbf{W}_1 \mathbf{x}_i) \dots)), \mathbf{z}_i).$$

When the gradient vanishes, the algorithm stops and the weights $\mathbf{W}_1, \dots, \mathbf{W}_L$ ideally correspond to a (not too bad) local minimum of the above loss function.

Convolutional neural networks are simply more structured versions of this generic neural network for which the weight matrices \mathbf{W}_i have a block Toeplitz

structure (so to enforce local filtering of the data). State-of-the-art methods also use more elaborate optimization methods, add extra tricks to the general architecture, but are essentially based on this elementary model. They are called “deep” whenever both the number of “layers” L and the number of “neurons” per layer l_i ’s are large.

Generative adversarial networks (GANs) are the combination of two such neural networks: (i) a so-called *generator* which generates, from Gaussian input vectors $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, the output “data” vectors $\mathbf{z} \in \mathbb{R}^q$, which are then compared by (ii) the *discriminator* to real data. The objective of the generator is to maximize the loss function of the discriminator (hence the “adversarial” name), which aims to maximally discriminate genuine data from the generated ones. Upon convergence of this adversarial game, the expected output is that the discriminator, while having become skillful at discriminating fake from real data, can no longer distinguish them: the GAN (precisely the generator) has learned to generate fake but extremely realistic data.

The top display of Figure 8.2 schematically depicts the diagram of a GAN.




Figure 8.2: Schematics of modern data generation and representation frameworks: GANs (**top**) and CNNs (**bottom**).

8.1.2.2 GAN-induced data are concentrated random vectors

It is generally assumed that the weight matrices \mathbf{W}_i in neural networks have bounded operator norms (with respect to the data dimensions and numbers, which is one of the key ingredients for good network performance [Bartlett et al., 2017, Miyato et al., 2018]). Similarly, the functions σ_i are restricted to

be 1-Lipschitz (typical functions are the ReLU function $\sigma(x) = \max(x, 0)$, the sign function, or sigmoid functions).

As such, since the input of GANs are random Gaussian vectors $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ and that the successive operations $\mathbf{x} \mapsto \mathbf{W}_i \mathbf{x}$ and $\mathbf{x} \mapsto \sigma_i(\mathbf{x})$ are all bounded Lipschitz operations, the output of a GAN is, by definition, a bounded Lipschitz function of a Gaussian vector.

From the Lipschitz stability of concentrated random vectors (recall (2.45)) and the fact that $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ is concentrated, it then comes that the output of the GAN generator is a concentrated random vector with head and tail parameters of order $O(1)$ (i.e., the same as for \mathbf{x}). In practice, other operations are performed on neural networks, such as pooling operations, random or deterministic dropouts, various connectivity matrix normalization procedures, etc. All these, sometimes precisely designed to avoid the “explosion” of the norm of the weight matrices, can be shown to also consist in Lipschitz operations with $O(1)$ Lipschitz constants [Seddik et al., 2020]. This thus extends our previous statement on the concentration of GAN outputs to state-of-the-art deep neural networks, and in particular to the very popular convolutional neural networks (CNNs).

While being concentrated vectors, GAN-generated data (say fake images of dogs and cats) do not necessarily “cluster” in their ambient space as a well-separated mixture model of concentrated random vectors. This is even obviously far from being the case: well-performing GANs must have a large variance (or entropy) in ambient space so to avoid generating systematically similar data. Images of dogs (class \mathcal{C}_1) versus images of cats (class \mathcal{C}_2) differ by the fact that they are generated by different neural network maps $\mathbf{x} \mapsto \phi_1(\mathbf{x})$ and $\mathbf{x} \mapsto \phi_2(\mathbf{x})$ having differing statistical means $\boldsymbol{\mu}_a = \mathbb{E}[\phi_a(\mathbf{x})]$ and covariances $\mathbf{C}_a = \mathbb{E}[\phi_a(\mathbf{x})\phi_a(\mathbf{x})^\top] - \boldsymbol{\mu}_a\boldsymbol{\mu}_a^\top$ (or alternatively by a conditional GAN [Mirza and Osindero, 2014] having the same effect). Yet, they are clearly not linearly separable (as are real images), therefore implying that $\|\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2\|$, $\|\mathbf{C}_1 - \mathbf{C}_2\|$ are likely quite small (when compared to the typical values of $\|\boldsymbol{\mu}_a\|$ and $\|\mathbf{C}_a\|$) and thus not prone to immediate classification by, e.g., standard clustering methods. Feature extraction methods (from simple histograms of oriented gradients (HOG) [Dalal and Triggs, 2005] to modern CNNs such as VGG, ResNet, etc.) precisely aim at “increasing” these distances by further transforming $\phi_a(\mathbf{x})$ into some $\psi(\phi_a(\mathbf{x}))$ for which distances in means and covariances are much larger.

8.1.2.3 From GAN-data to CNN-features to GMM

State-of-the-art feature extractors in modern machine learning are based on deep neural networks, and specifically for multimedia data on convolutional neural networks. These networks, such as the popular VGG nets [Simonyan and Zisserman, 2014] or ResNets [He et al., 2016], have been pre-trained on huge collections of independent databases and are thus fixed, independent functions of the dataset of interest to the experimenter. The associated feature extractor, say $\psi : \mathbb{R}^p \rightarrow \mathbb{R}^q$, is then usually taken to be the function that maps the data to the second-to-last layer of the trained deep network, with the very last layer

(in general a fully connected “decision” layer from $\mathbb{R}^q \rightarrow \mathbb{R}^d$, with d the number of classes that the deep network was trained to classify) discarded, and only the mapping from input to q -dimensional output is maintained to form ψ .

Being a neural network map, ψ is naturally a Lipschitz function with well-controlled and bounded Lipschitz parameter [Bartlett et al., 2017, Miyato et al., 2018]. The features $\psi(\mathbf{z}_i) \in \mathbb{R}^q$ obtained from the machine learning algorithm are therefore some bounded Lipschitz images of the raw data $\mathbf{z}_i \in \mathbb{R}^p$. When these raw “data” \mathbf{z}_i are themselves the (close to realistic multimedia data) output from a GAN, i.e., $\mathbf{z}_i = \phi_a(\mathbf{x}_i)$ with our previous notations, we obtain that the second-to-last layer network features $\tilde{\mathbf{x}}_i$ to be classified are of the form $\tilde{\mathbf{x}}_i = \psi(\phi_a(\mathbf{x}_i))$ with $\mathbf{x}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$, which by definition are concentrated random vectors (since $\psi \circ \phi$ is Lipschitz with bounded parameter).

As a consequence, the dataset $\{\tilde{\mathbf{x}}_i\}_{i=1}^n$ in which each $\tilde{\mathbf{x}}_i$ is of the form $\tilde{\mathbf{x}}_i = \psi(\phi_a(\mathbf{x}_i))$, for some $a \in \{1, \dots, k\}$ identifying the class of $\tilde{\mathbf{x}}_i$, is a *mixture of concentrated random vectors*.

As such, to treat more realistic data models than the “toy” Gaussian mixture models, the results presented in the previous chapters should be updated to data $\{\tilde{\mathbf{x}}_i = \psi(\phi_a(\mathbf{x}_i))\}_{i=1}^n$ for $\mathbf{x}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ (where $a \in \{1, \dots, k\}$ implicitly depends on the class of \mathbf{x}_i) arising from a *mixture of concentrated random vectors*. This being said, from a purely mathematical concentration-theoretic standpoint, it is not formally necessary to specify the concentration origin of $\tilde{\mathbf{x}}_i$ and we may, in all generality, simply ask for the data to be *generic concentrated random vectors from a mixture model*.

Therefore, in the following, we will assume that the data (be they the raw random data or any kind of “representations” of the raw data), which we redefine as $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^p$, are simply issued from any mixture of concentrated random vectors as follows:

$$\mathbf{x}_1, \dots, \mathbf{x}_{n_1} \sim \mathcal{L}_1, \quad \dots, \quad \mathbf{x}_{n-n_k}, \dots, \mathbf{x}_n \sim \mathcal{L}_k$$

where \mathcal{L}_a is the law of a concentrated random vector of dimension p . We further denote, as usual, the statistics means and covariances of the law \mathcal{L}_a as $\boldsymbol{\mu}_a$ and \mathbf{C}_a . For technical reason, it is also necessary to demand that the (joint) data matrix $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{p \times n}$ also be concentrated.

The fundamental result and message of this section are the following: from Theorem 2.18, it appears, in a single-class setting ($k = 1$), that the resolvent $\mathbf{Q}(z) = (\frac{1}{n} \mathbf{XX}^\top - z \mathbf{I}_n)^{-1}$ of $\frac{1}{n} \mathbf{XX}^\top$, which is at the core of most of the machine learning algorithms studied thus far, admits a deterministic equivalent $\bar{\mathbf{Q}}(z)$ which *only* depends on $\mathbb{E}[\mathbf{x}_i \mathbf{x}_i^\top] = \boldsymbol{\mu}_1 \boldsymbol{\mu}_1^\top + \mathbf{C}_1$ for $\mathbf{x}_i \sim \mathcal{L}_1$, and thus on the *first two order statistics* of the law \mathcal{L}_1 .

It is thus reasonable to infer from Theorem 2.18 that, for a multi-class setting ($1 < k \ll n$), the same will hold. Besides, from Theorem 4.1 and the discussion preceding it, it is likely that kernel matrices with the standard normalization, e.g., $\mathbf{K} = \{f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)\}_{i,j=1}^n$ or $\mathbf{K} = \{f(\mathbf{x}_i^\top \mathbf{x}_j/p)\}_{i,j=1}^n$, and their spectral properties, which asymptotically essentially depend on the behavior of a low-

rank perturbation of $\mathbf{X}^\top \mathbf{X}$, will also depend on the first two order statistics of the data distribution.

8.1.2.4 Kernel asymptotics and GAN-generated data

The above intuition on kernel matrices turns out to be correct, at least to some extent. It is shown in [Seddik et al., 2019, Theorem 1] that Theorem 4.1 indeed holds identically for (a mixture of) concentrated random vectors. Precisely, it is shown (for technical simplicity) that $\|\mathbf{P}(\mathbf{K} - \tilde{\mathbf{K}})\mathbf{P}\| \xrightarrow{a.s.} 0$ where $\mathbf{P} = \mathbf{I}_n - \frac{1}{n}\mathbf{1}_n\mathbf{1}_n^\top$: this, as in Corollary 4.1, discards several terms in the expansion of $\tilde{\mathbf{K}}$ without affecting the practical relevance of the result (for instance, all terms of the form $\mathbf{v}\mathbf{1}_n^\top, \mathbf{1}_n\mathbf{v}^\top$ or $\mathbf{1}_n\mathbf{1}_n^\top$ disappear). This fact notwithstanding, the random $\tilde{\mathbf{K}}$ has the *same* expression under a Gaussian or “concentrated” mixture model. However, $\tilde{\mathbf{K}}$ does not solely depend on the first and second order moments of the data distribution, due to the presence of the vector $\psi = \{\|\mathbf{w}_i\|^2/p - \mathbb{E}[\|\mathbf{w}_i\|^2/p]\}_{i=1}^n$, the entries of which have mean zero but variance function of the fourth-order moment of \mathbf{x}_i (precisely of $\mathbf{w}_i = \mathbf{C}_a^{-1/2}(\mathbf{x}_i - \boldsymbol{\mu}_a)$ for $\mathbf{x}_i \sim \mathcal{L}_a$). Yet, this vector ψ (i) appears in the low-rank part of the expansion of $\tilde{\mathbf{K}}$ and thus does not asymptotically affects the limiting spectrum of \mathbf{K} , and (ii) does not affect the component $\mathbf{A}_{1,11}$ which essentially rules the *informative* isolated spectrum behavior of \mathbf{K} (position of isolated eigenvalues and content of the associated eigenvectors). As a consequence, various machine learning algorithms based on \mathbf{K} , such as the unsupervised extraction of “informative” eigenvectors and the use within a semi-supervised or supervised learning filter discussed in Section 4.5, are essentially *universal* with respect to the laws $\mathcal{L}_a = \mathcal{L}_a(\boldsymbol{\mu}_a, \mathbf{C}_a)$ in that they only depend on the first two order statistics $\boldsymbol{\mu}_a$ and \mathbf{C}_a .

The immediate outcome of this discussion is that most results above, defined for machine learning algorithms based on \mathbf{K} , *provably identically hold for realistic (GAN-generated) data and for their Gaussian mixture model equivalent* (i.e., for the Gaussian mixture model with same first order statistics).

This is visually confirmed in Figure 8.4 which provides a concrete comparison of the finite-dimensional spectrum (eigenvalues and two dominant eigenvectors) of $\mathbf{K} = \{\exp(-\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)\}_{i,j=1}^n$ for \mathbf{x}_i CNN-features of real images, or of GAN images (arising from the training of a GAN on the same real images), versus a Gaussian mixture with same empirical (mean and covariance) statistics as those CNN-features. The visual match, which we now know to be theoretically asymptotically perfect in the GAN-data case, is extremely accurate in this finite-dimensional case (p is of order a few thousands), even on the real data for which no guarantee can of course be claimed (so long that the theoretical relation between real data and their GAN-generated counterpart is not elucidated).



Figure 8.3: Images produced by the BigGAN model [Brock et al., 2019] for three data classes (“hamburger”, “mushroom”, “pizza”) (**top**) versus the dataset of real images used to learn the GAN, the ImageNet dataset [Krizhevsky et al., 2017] (**bottom**).

8.1.2.5 Beyond “classical” kernels

The previous section discussed the universality of the kernel matrices of the type $\mathbf{K} = \{f(\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)\}_{i,j=1}^n$ with respect to the concentrated random vector statistics of \mathbf{x}_i . The main reason follows from the fact that the higher-than-two order moments of \mathbf{x}_i play a marginal role, if not no role, in the asymptotics of \mathbf{K} (as we saw, only through the vector ψ which has asymptotically no impact on the relevant eigenvectors and low-rank informative terms in \mathbf{K}).

This may no longer be the case for more elaborate kernels, such as the α - β kernel and the properly scaling kernel, discussed in Section 4.3 and 4.4, respectively.

For the α - β kernel (in Theorem 4.2), the entries of the “second-order noise” matrix Φ are related to $(\mathbf{w}_i^\top \mathbf{w}_j)^2$: from the independence of \mathbf{w}_i and \mathbf{w}_j , the variance of this term depends on the fourth order moments of the (independent) entries of \mathbf{w}_i and \mathbf{w}_j , which impacts the overall spectrum of Φ . (Note that since the diagonal terms $(\mathbf{w}_i^\top \mathbf{w}_i)^2$ are discarded, the up-to-eighth order moments are not accounted for.) The universality thus holds, in this case, only up to the fourth order moments: the Gaussian mixture model likely becomes insufficient to properly model the behavior of such \mathbf{K} on concentrated random vectors and thus on realistic datasets.




Figure 8.4: Borrowed from [Seddik et al., 2020]: eigenvalue distribution and two dominant eigenvectors of $\mathbf{K} = \{\exp(-\|\mathbf{x}_i - \mathbf{x}_j\|^2/p)\}_{i,j=1}^n$ for CNN features of different deep networks (from left to right: ResNet-50 [He et al., 2016] with $p = 2048$ features, the VGG-16 [Simonyan and Zisserman, 2014] with $p = 4096$ features and DenseNet-201 [Huang et al., 2017] with $p = 1920$) of the dataset of original images in Figure 8.3. Comparison of the results obtained for the GAN-generated data (**top**) versus the real data (**bottom**), empirically on the dataset and on independent Gaussian vectors following the same statistical (means and covariances) mixture.

As for the more involved properly scaling kernels (in Theorem 4.6), such as the kernel $\mathbf{K} = \{f(\mathbf{x}_i^\top \mathbf{x}_j / \sqrt{p})\}_{i,j=1}^n$, recall that its asymptotics are inherently related to the Gaussian asymptotics (central limit) of $\mathbf{x}_i^\top \mathbf{x}_j / \sqrt{p}$ for independent $\mathbf{x}_i, \mathbf{x}_j$. This central limit must be preserved in concentrated random vectors for random matrix universality to hold. Yet, this is far from obvious and demands additional constraints on the laws \mathcal{L}_a (since, for instance, $\mathbf{x}_i^\top \mathbf{x}_j / \sqrt{p}$ does not necessarily rewrites as the sum of independent variables for concentrated $\mathbf{x}_i, \mathbf{x}_j$). In this setting, it is quite possible that significant deviations from universality will be observed. In the specific case of GAN-data, which arise from deep neural network learning, one is tempted to assume some sort of an inherent “isotropic” nature of the successive layers of the large trained network, which

may smooth-out the concentrated random vectors in a way to make them “more Gaussian-like”, so that one may still be confident that the Gaussian mixture modeling may be somewhat relevant.

8.2 Wide-sense universality in large dimensional machine learning

The example of GANs in the previous section underlies a seemingly more fundamental aspect of real (large dimensional) data processing: so long that real data can be assumed to be inherently constituted of a large number of degrees of freedom (or of randomness), a dual phenomenon arises:

- these degrees of freedom tend to regularize and induce robustness into machine learning algorithms: this is in particular at the very source of well-behaved deep neural networks (based on numerous data and numerous randomly-initialized neurons) versus ill-behaved small and shallow perceptrons;
- the machine learning algorithms essentially extract basic “small” dimensional statistics (scalar comparisons of first order moments and deterministic patterns) from the data, thereby completely “eliminating” the noise, irrespective of its nature (higher order moments of the distribution).

This suggests that, beyond images and sounds (that can be adequately modeled by GAN-generated data), data representations that are sufficiently rich in “degrees of freedom” should be similarly handled in a robust and theoretically tractable manner by standard machine learning methods. The recent success of word embeddings (such as the word2vec approach [Mikolov et al., 2013]) that aim to represent words, sentences and other structures in the complex field of natural language processing via vectors in a rather large dimensional space, confirms this intuition: these representations are sufficiently rich and diverse to perform theoretical analysis by means of Gaussian (or concentration-type) mixture approximations [Couillet et al., 2020]. A typical counterexample in this very field of natural language processing is the so-far exploited “bag-of-words” representation [Manning et al., 2008] which consists in large dimensional but *extremely sparse* dictionary vectors (each sentence being represented by one such vector counting the number of instances of each dictionary word in the sentence): being very sparse, these vectors *do not concentrate*, thereby hardly contributing to adding degrees of freedom to stabilize the machine learning algorithms.

Figure 8.5 compares the popular Gaussian kernel matrix structures observed when evaluating the CNN features for real images (of dimension $p = 1024$) in two classes, versus the word2vec embeddings for words (of dimensions $p = 300$) in two categories. The colormap strongly suggests the aforementioned concentration effect arising in real data, in both computer vision and natural language processing contexts.




Figure 8.5: Kernel matrices \mathbf{K} and the second dominant eigenvectors \mathbf{v}_2 for (**left**) VGG-16 [Simonyan and Zisserman, 2014] features of CIFAR-10 data (“airplane” versus “bird”) and (**right**) word2vec [Mikolov et al., 2013] features of GoogleNews-vectors data (“sports” versus “sales”), with $\mathbf{x}_1, \dots, \mathbf{x}_{n/2} \in \mathcal{C}_1$ and $\mathbf{x}_{n/2+1}, \dots, \mathbf{x}_n \in \mathcal{C}_2$. [\[Link to code\]](#)

This being said, we must note that the validity of all aforementioned random matrix predictions and improvements fundamentally relies on the *existence* of convenient data representations. Aside from subspace or manifold-based algorithms (such as PCA [Wold et al., 1987] or principle Hessian directions [Li, 1992]), the field of random matrix theory does not in itself propose such elaborate representations. If anything, it would naturally suggest to operate random (linear or nonlinear) projections on the data so to “generate” more randomness, and thus more predictability and robustness. However, mere random projections are a rather poor elementary representation technique which does not account for the data context and structure (as opposed to deep convolutional neural nets which intrinsically exploit the locality and multi-class nature of the data).

Recalling that machine learning can be seen as the elegant combination of “representation + decision” [Domingos, 2012], random matrix theory is so far merely able to operate on the “decision” aspect, assuming that the data representation is given and rather convenient. Better understanding and contributing to the “representation” part of machine learning would require to add supplementary data-related contextual ingredients to random matrix theory, so most likely more complex random and deterministic structures. Alternatively, empirically witnessing the powerful capability of deep neural networks to design appropriate representations of data, random matrix theory could also contribute to a better theoretical understanding of the deep learning mechanisms. This would mean characterization the dynamics of learning in the multi-layer, nonlinear,

and non-convex setting of deep networks, which raises multiple (so far unsurpassable) technical difficulties:

- several works predict that, despite the highly non-convex nature of the underlying optimization procedure, deep network learning owes its stability to the simultaneously large data dimension, number as well as network depth and width. This has not been formally proven but related problems in random Gaussian fields (formally not a neural network but that could resemble one [Choromanska et al., 2015]) show that in the large dimensional regime, while the number of local minima increases exponentially with the model size, these minima tend to locate at the same (loss) level, thereby ensuring that almost all initialization points reach the approximately same (good) performance upon convergence. These results however do not say much more: what are the performance reached? How do they relate to the data statistics? How could they be improved? Is there a training-test mismatch in the associated loss “landscape” (e.g., is reaching or getting close to the global minima a necessary and sufficient condition for good performance)? Besides, the setting of Gaussian random fields remains formally far from actual deep networks and notably ignores some of its key features, such as the nonlinear activations, their structure in multiple layers, their convolutive nature for convolutional nets, etc;
- by considering the infinite-width limit (that is, the limit of *infinitely many neurons per layer*), Jacot et al. [2018] proposed the so-called neural tangent kernel (NTK) as the key object to study the limiting behavior of deep neural networks. The NTK depends on both the data and the network random initialization, and most importantly, remains *unchanged* in the infinite-width limit during gradient descent [Lee et al., 2020] under some (rather restrictive) assumptions. In this NTK regime, as a consequence of the “constant” NTK that does not evolve during training, the resulting deep network behaves, in the infinite-width limit, as a kernel regression model, yielding performance that is closer to classical kernel methods than to modern deep networks [Chizat et al., 2019, Arora et al., 2019b]. In this vein, random matrix theory may come into play, as in the example of sample covariance and the Marčenko-Pastur law, to account for a non-trivial ratio between the input dimension/sample size and the network width, as well as for the different widths of each layer, thereby allowing for a “layer-by-layer” characterization of the network statistical behavior.

While it has been empirically observed that modern deep networks yield significantly better performance than the limiting NTKs [Chizat et al., 2019, Arora et al., 2019b], it remains true that neural networks and kernels are intimately related, so that further studies and explorations of the latter may help improve the understanding of the former.

8.3 Discussions and conclusions

Concentration of measure theory provides a powerful tool, quite complementary to random matrix theory, to analyze the performance of statistical learning algorithms applied to a host of realistic data models. According to our previous discussions, one is tempted to state that the existence of “good concentrated vector modeling” of real data, such as the images produced by GANs, fully justifies (through the proofs of universality) the further development of random matrix theory for the performance analysis and improvement of Gaussian mixture-based algorithms. However, claiming that a model is “good” is a fairly subjective statement (e.g., do GANs produce all images of a class one could possibly think of or do they over-produce a limited set of images?), and if ever a measure of goodness was appropriately designed, confirming that the analyses made on Gaussian mixtures are robust to deviations in the data models which would include real data is likely difficult. On this point, only extensive simulations can be used as a measure of faith.

This is the case of images, which differ from text, language, and some complex signals, in their not requiring “long-term memory”: correlations in images are indeed mostly “localized” and may thus be produced by (convolutional) feed-forward networks. Textual contents are rather produced by recurrent networks, and most particularly by long short-term memory (LSTM) networks [Hochreiter and Schmidhuber, 1997]. A possible parallel path to proving that learning in natural language processing can be theoretically analyzed by random matrix and concentration of measure methods would then consist in showing that LSTM networks are also Lipschitz mappings (for instance from and into some word embedding space). Similar to deep feedforward networks, understanding the behavior of LSTMs is difficult (as they use the same gradient descent learning) but the study of simpler networks, such as echo state networks, already capable of faithful predictions, may help improve our present lack of understanding.

In spirit though, without having to formally prove that “real data learning” is amenable to random matrix analysis, the validity of the random matrix approximation mostly holds on two complementary pillars: (i) a “concentration-like behavior” of the data representation, and (ii) a good “Lipschitz mixing behavior” of the studied algorithm. That is: (i) the data representation under study should resemble a vector with *rather independent and delocalized entries*, so as to exploit most of the degrees of freedom offered by its ambient space, in the manner of Gaussian random vectors: this ensures that their scalar Lipschitz observations have a (more or less) predictable behavior; and (ii) the learning algorithm maintains the “delocalized” behavior or data, if not reinforces it: this avoids the creation of outlying (thus difficult to predict) behavior and most well-performing algorithms tend to satisfy this rule (activation functions in neural networks are Lipschitz, their weight matrices are normalized, kernels are rarely more than quadratic to remain stable, etc.). In a sense, even data that would not be concentrated per se may be appropriately mixed (Gaussianized one may say) by the learning algorithm so to stabilize the performance. Conversely,

well-concentrated data may suffer the effects of non-strictly Lipschitz transformations (so to extract exotic or marginal features for instance) while remaining stable under random matrix analysis. This in essence justifies the wide applicability and robustness of the various random matrix analyses presented in the course of the monograph on various real data.

Some data, however, are clearly not concentrated: this is notably the case of sparse vectors and sparse graphs. A typical example in natural language processing is that of the bag-of-words or tf-idf (for term frequency-inverse document frequency) methods [Manning et al., 2008], which use large dictionary vectors (typically of size in the hundreds or thousands of words) filled with the number of occurrence or frequency of each word in a paragraph or text. These vectors are naturally quite “empty” and clearly do not adhere to the concentration of distance phenomenon observed for Gaussian-like vectors in Figure 8.5. The adjacency matrix of sparse graphs (that have n nodes with $O(1)$ neighbors per node) also loses key concentration properties to random matrix analysis: the norm or inner product between arbitrary rows or columns of the adjacency matrix do not converge and this makes most classical random matrix tools (starting with the trace lemma, Lemma 2.11) collapse at once. When additional statistical symmetries are assumed, for instance if the entries of the adjacency matrix are i.i.d., stable asymptotic behavior (of the eigenspectrum in particular) is empirically observed, but is to date not theoretically tractable to random matrix theory. If ever possible, the observed behavior would at any rate not be universal and thus quite dependent on the detailed model statistics: this raises a major issue in practice since, Gaussian approximations no longer being valid, the data models must be extremely accurate for the theoretical analysis to be of any value. With difficult-to-understand real data, this severely reduces the interest of large dimensional statistical results.

Nonetheless, where random matrix theory fails to work, ingredients of statistical physics (despite their sometimes lack of mathematical rigor) can be exploited. Informal techniques such as the replica method, the various linearizations and approximations of the belief-propagation algorithm using statistical physics concepts (of free energy, Hamiltonian, Bethe-Hessian approximation, etc.) [Mézard and Montanari, 2009] have provided tremendous advances in large dimensional statistical learning, precisely under scenarios where random matrices still lag behind. This is particularly the case of sparse graph mining, where heuristic but powerful new algorithms were designed out of statistical physics ideas [Krzakala et al., 2013, Saade et al., 2014]. Mathematicians are only recently managing to formally prove some of the fundamental predictions proposed in these articles. So long that no formal unified theory of sparse random matrices exists, the future of large dimensional statistical learning may in part lie in this two-stage process where physicists come first with intuitive ideas and new algorithm proposals, before random matrix experts formalize and mathematically push further these ideas.

Chapter 9

Solutions to exercises in Section 2.9

Exercise 1 (Stieltjes transform and moments). *Show that the Stieltjes transform $m_\mu(z)$, of a probability measure μ with bounded support (and thus finite moments), is a moment generating function in the sense that, for all $z \in \mathbb{C}$ such that $|z| > \max\{|\inf(\text{supp}(\mu))|, |\sup(\text{supp}(\mu))|\}$,*

$$m_\mu(z) = -\frac{1}{z} \sum_{k=0}^{\infty} M_k z^{-k}$$

where $M_k = \int t^k \mu(dt)$.

From this formulation, propose a method to evaluate the successive moments of μ .

Correction 1 (Stieltjes transform and moments). Note that

$$\begin{aligned} m_\mu(z) &= \int \frac{1}{t-z} \mu(dt) = -\frac{1}{z} \int \frac{1}{1-t/z} \mu(dt) \\ &= -\frac{1}{z} \sum_{k=0}^{\infty} \int \frac{t^k}{z^k} \mu(dt) = -\frac{1}{z} \sum_{k=0}^{\infty} M_k z^{-k}. \end{aligned}$$

Thus, differentiating successively $\delta(z) \equiv -\frac{1}{z} m_\mu(z^{-1}) = \int \frac{1}{1-zt} \mu(dt)$, we find

$$\frac{d^\ell}{dz^\ell} \delta(z) = \sum_{k \geq \ell} \frac{k!}{(k-\ell)!} M_k z^{k-\ell}.$$

In particular, letting $z = 0$ in the expression above, we obtain

$$\frac{d^\ell}{dz^\ell} \delta(0) = \ell! \cdot M_\ell.$$

Exercise 2 (Non-immediate Stieltjes transforms). Let $\mathbf{X} \in \mathbb{R}^{n \times n}$ be a Hermitian matrix and $\mathbf{Q}(z) = (\mathbf{X} - z\mathbf{I}_n)^{-1}$ its resolvent. Show that, for any $\mathbf{u} \in \mathbb{R}^n$ of unit norm $\|\mathbf{u}\| = 1$ and any \mathbf{A} nonnegative definite and such that $\text{tr } \mathbf{A} = 1$, the quantities $\mathbf{u}^\top \mathbf{Q}(z)\mathbf{u}$ and $\text{tr } \mathbf{A}\mathbf{Q}(z)$ are also Stieltjes transform of probability measures.

What are these measures and what are their supports?

Correction 2 (Non-immediate Stieltjes transforms). Denote $\mathbf{X} = \mathbf{V}\Lambda\mathbf{V}^\top$ the spectral decomposition of \mathbf{X} , with $\mathbf{V} = [\mathbf{v}_1, \dots, \mathbf{v}_n] \in \mathbb{R}^{n \times n}$ and $\Lambda = \text{diag}\{\lambda_i\}_{i=1}^n$. It then suffices to observe that, for $\mathbf{w} = \mathbf{V}^\top \mathbf{u}$,

$$\mathbf{u}^\top \mathbf{Q}(z)\mathbf{u} = \mathbf{u}^\top \mathbf{V} \text{diag} \left\{ \frac{1}{\lambda_i - z} \right\}_{i=1}^n \mathbf{V}^\top \mathbf{u} = \sum_{i=1}^n \frac{[\mathbf{w}]_i^2}{\lambda_i - z} = \int \frac{1}{t - z} \mu(dt)$$

where $\mu = \sum_{i=1}^n [\mathbf{w}]_i^2 \delta_{\lambda_i}$. In particular,

$$\int \mu(dt) = \sum_{i=1}^n [\mathbf{w}]_i^2 = \mathbf{w}^\top \mathbf{w} = \mathbf{u}^\top \mathbf{V} \mathbf{V}^\top \mathbf{u} = 1$$

so that, since $[\mathbf{w}]_i^2 \delta_{\lambda_i}(t) \geq 0$, μ is a probability measure.

Letting similarly $\mathbf{B} = \mathbf{V}^\top \mathbf{A} \mathbf{V}$,

$$\text{tr } \mathbf{A}\mathbf{Q} = \text{tr } \mathbf{B} \text{diag} \left\{ \frac{1}{\lambda_i - z} \right\}_{i=1}^n = \sum_{i=1}^n \frac{[\mathbf{B}]_{ii}}{\lambda_i - z} = \int \frac{1}{t - z} \nu(dt)$$

where $\nu = \sum_{i=1}^n [\mathbf{B}]_{ii} \delta_{\lambda_i}$, with $[\mathbf{B}]_{ii} \geq 0$ (to see this, write $\mathbf{A} = \mathbf{V}' \Lambda' (\mathbf{V}')^\top$, so that $[\mathbf{B}]_{ii} = [\mathbf{V}'^\top \mathbf{V}']_{i,i} \Lambda' [(\mathbf{V}')^\top \mathbf{V}]_{i,i} \geq 0$ since Λ' is a diagonal of non-negative entries) and $\int \nu(dt) = \text{tr } \mathbf{B} = \text{tr } \mathbf{A} = 1$.

The support of both measures μ and ν is included in the support of the original empirical spectral measure $\sum_{i=1}^n \delta_{\lambda_i}$ (i.e., $\{\lambda_1, \dots, \lambda_n\}$). They however only coincide with the support if $[\mathbf{w}]_i^2 \neq 0$ for all i 's associated to a common λ_j (respectively, $[\mathbf{B}]_{ii} \neq 0$).

Exercise 3 (Stieltjes transform and singular values). Let μ be a probability measure on \mathbb{R}^+ and ν, ν' be the measures defined by

$$\begin{aligned} \int f(t) \nu(dt) &= \int f(\sqrt{t}) \mu(dt) \\ \int f(t) \nu'(dt) &= \frac{1}{2} \left(\int f(t) \nu(dt) + \int f(-t) \nu(dt) \right) \end{aligned}$$

for all bounded continuous f .

What are ν and ν' when $\mu = \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i}$ for some $\lambda_1, \dots, \lambda_n \geq 0$? Show that the Stieltjes transform $m_{\nu'}$ of ν' satisfies

$$m_{\nu'}(z) = zm_\mu(z^2). \quad (9.1)$$

Letting $\mathbf{X} \in \mathbb{R}^{n \times p}$ and μ be the empirical spectral measure of \mathbf{XX}^\top (see Definition 2), relate the Stieltjes transform of the empirical spectral measure of the matrix

$$\Gamma = \begin{bmatrix} \mathbf{0}_{n \times n} & \mathbf{X} \\ \mathbf{X}^\top & \mathbf{0}_{p \times p} \end{bmatrix} \in \mathbb{R}^{(n+p) \times (n+p)} \quad (9.2)$$

to that of the measure μ , and conclude on the nature of this Stieltjes transform for $n = p$.

Correction 3 (Stieltjes transform and singular values). For $\mu = \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i}$, we have $\nu = \frac{1}{n} \sum_{i=1}^n \delta_{\sqrt{\lambda_i}}$ and $\nu' = \frac{1}{2n} \sum_{i=1}^n (\delta_{\sqrt{\lambda_i}} + \delta_{-\sqrt{\lambda_i}})$.

We thus have

$$\begin{aligned} m_{\nu'}(z) &= \frac{1}{2} \int \frac{1}{t-z} \nu(dt) + \frac{1}{2} \int \frac{1}{-t-z} \nu(dt) \\ &= \frac{1}{2} \int \frac{1}{\sqrt{t}-z} \mu(dt) + \frac{1}{2} \int \frac{1}{-\sqrt{t}-z} \mu(dt) \\ &= z \int \frac{1}{t-z^2} \mu(dt) = zm_\mu(z^2). \end{aligned}$$

The Stieltjes transform of the empirical spectral measure of Γ is given by

$$\begin{aligned} \frac{1}{n+p} \text{tr} \begin{pmatrix} -z\mathbf{I}_n & \mathbf{X} \\ \mathbf{X}^\top & -z\mathbf{I}_p \end{pmatrix}^{-1} &= \frac{z}{n+p} \text{tr}(\mathbf{XX}^\top - z^2\mathbf{I}_n)^{-1} + \frac{z}{n+p} \text{tr}(\mathbf{X}^\top \mathbf{X} - z^2\mathbf{I}_p)^{-1} \\ &= \frac{2z}{n+p} \text{tr}(\mathbf{XX}^\top - z^2\mathbf{I}_n)^{-1} + \frac{n-p}{n+p} \frac{1}{z} \end{aligned}$$

where we used the block-matrix inverse lemma, Lemma 2.5, and then Lemma 2.4. This Stieltjes transform has its singularities for z the singular values of \mathbf{X} (i.e., the square-root of the eigenvalues of \mathbf{XX}^\top) and thus relates to the singular spectrum of \mathbf{X} , or for $z = 0$ when $n \neq p$. In particular, taking $n = p$ (so that \mathbf{X} is a square matrix), we recover the relation in (9.1).

Exercise 4 (Proof of Lemma 2.9: special case). For $\mathbf{A}, \mathbf{M} \in \mathbb{R}^{p \times p}$ symmetric nonnegative definite matrices, $\mathbf{u} \in \mathbb{R}^p$, $\tau > 0$ and $z < 0$, show that

$$\left| \text{tr } \mathbf{A} (\mathbf{M} + \tau \mathbf{uu}^\top - z \mathbf{I}_p)^{-1} - \text{tr } \mathbf{A} (\mathbf{M} - z \mathbf{I}_p)^{-1} \right| \leq \frac{\|\mathbf{A}\|}{|z|}.$$

Correction 4 (Proof of Lemma 2.9: special case). Using the resolvent identity, Lemma 2.1, and Sherman–Morrison formula, Lemma 2.8, we write

$$\begin{aligned} &\left| \text{tr } \mathbf{A} (\mathbf{M} + \tau \mathbf{uu}^\top - z \mathbf{I}_p)^{-1} - \text{tr } \mathbf{A} (\mathbf{M} - z \mathbf{I}_p)^{-1} \right| \\ &= \left| \text{tr } \mathbf{A} (\mathbf{M} + \tau \mathbf{uu}^\top - z \mathbf{I}_p)^{-1} \tau \mathbf{uu}^\top (\mathbf{M} - z \mathbf{I}_p)^{-1} \right| \\ &= \frac{\left| \text{tr } \mathbf{A} (\mathbf{M} - z \mathbf{I}_p)^{-1} \tau \mathbf{uu}^\top (\mathbf{M} - z \mathbf{I}_p)^{-1} \right|}{1 + \tau \mathbf{u}^\top (\mathbf{M} - z \mathbf{I}_p)^{-1} \mathbf{u}} \\ &= \frac{\left| \tau \mathbf{u}^\top (\mathbf{M} - z \mathbf{I}_p)^{-1} \mathbf{A} (\mathbf{M} - z \mathbf{I}_p)^{-1} \mathbf{u} \right|}{1 + \tau \mathbf{u}^\top (\mathbf{M} - z \mathbf{I}_p)^{-1} \mathbf{u}}. \end{aligned}$$

At this point, we use the fact that $|\mathbf{x}^\top \mathbf{B}\mathbf{x}| \leq \|\mathbf{B}\| \cdot \|\mathbf{x}\|^2$ with $\mathbf{B} = (\mathbf{M} - z\mathbf{I}_p)^{-\frac{1}{2}} \mathbf{A} (\mathbf{M} - z\mathbf{I}_p)^{-\frac{1}{2}}$ (note the important for z to be real negative to ensure that \mathbf{B} is symmetric and nonnegative definite) and $\mathbf{x} = (\mathbf{M} - z\mathbf{I}_p)^{-\frac{1}{2}} \mathbf{u}$, so to obtain

$$\begin{aligned} & \left| \operatorname{tr} \mathbf{A} (\mathbf{M} + \tau \mathbf{u} \mathbf{u}^\top - z\mathbf{I}_p)^{-1} - \operatorname{tr} \mathbf{A} (\mathbf{M} - z\mathbf{I}_p)^{-1} \right| \\ & \leq \|(\mathbf{M} - z\mathbf{I}_p)^{-\frac{1}{2}} \mathbf{A} (\mathbf{M} - z\mathbf{I}_p)^{-\frac{1}{2}}\| \cdot \frac{\tau \mathbf{u}^\top (\mathbf{M} - z\mathbf{I}_n)^{-1} \mathbf{u}}{1 + \tau \mathbf{u}^\top (\mathbf{M} - z\mathbf{I}_n)^{-1} \mathbf{u}}. \end{aligned}$$

Using finally $\|(\mathbf{M} - z\mathbf{I}_p)^{-\frac{1}{2}} \mathbf{A} (\mathbf{M} - z\mathbf{I}_p)^{-\frac{1}{2}}\| \leq \|\mathbf{A}\| \cdot \|(\mathbf{M} - z\mathbf{I}_n)^{-1}\|$, $\|(\mathbf{M} - z\mathbf{I}_n)^{-1}\| \leq |z|^{-1}$ and $\frac{x}{1+x} \leq 1$ for $x > 0$, we conclude the proof.¹

Exercise 5 (The $\sqrt{|x - E|}$ behavior of the edges). Show that both the semicircle law (Theorem 2.5) and the Marčenko-Pastur law (Theorem 2.4, for $c \neq 1$) have a local $\sqrt{|x - E|}$ behavior at each of the edges E of their support.

Conclude on the typical number of eigenvalues of the Wishart matrix $\frac{1}{n} \mathbf{X} \mathbf{X}^\top \in \mathbb{R}^{p \times p}$ (with $\mathbf{X}_{ij} \sim \mathcal{N}(0, 1)$ independent) and the Wigner $\frac{1}{\sqrt{n}} \mathbf{X} \in \mathbb{R}^{n \times n}$ (with $\mathbf{X}_{ij} = \mathbf{X}_{ji} \sim \mathcal{N}(0, 1)$ independent up to symmetry) found near the edges of their respective supports.

Relate this finding to the fluctuations of the Tracy-Widom distribution of the largest and smallest eigenvalues in Theorem 2.15.

What happens for the left-edge of the support of the Marčenko-Pastur law and to the associated smallest eigenvalues of Wishart matrices when $\lim p/n = c = 1$? How many eigenvalues are then found close to the left edge in this so-called “hard-edge” setting? Conclude on the typical fluctuations of these eigenvalues and confirm numerically.

Correction 5 (The $\sqrt{|x - E|}$ behavior of the edges). The Marčenko-Pastur density can be approximated around, say, the right-edge $E_+ = (1 + \sqrt{c})^2$ (i.e., $x = (1 + \sqrt{c})^2 - \epsilon$, $\epsilon > 0$ small) of its support as

$$\mu(dx) \underset{x=E_+-0}{\simeq} \frac{\sqrt{(1 + \sqrt{c})^2 - (1 - \sqrt{c})^2}}{2\pi c(1 + \sqrt{c})^2} \sqrt{|x - (1 + \sqrt{c})^2|} dx.$$

Similarly, around $E_+ = 2$, the semicircle distribution has approximate density

$$\mu(dx) = \frac{1}{2\pi} \sqrt{(x^2 - 4)^+} dx \underset{x \sim E_+-0}{\simeq} \frac{1}{\pi} \sqrt{|x - 2|} dx.$$

As a consequence, for both models, within the interval $[E_+ - \epsilon, E_+]$ of length ϵ small, the typical integral of the density is of order

$$\int_{E_+-\epsilon}^{E_+} \mu(dx) \underset{\epsilon \rightarrow 0}{\simeq} C \int_{E_+-\epsilon}^{E_+} \sqrt{E_+ - x} dx = C \int_0^\epsilon \sqrt{y} dy = \frac{2C}{3} \epsilon^{\frac{3}{2}}$$

¹The proof of the result for z complex with positive imaginary part is available in the original article [Silverstein and Bai, 1995].

with $y = E_+ - x$ and $C > 0$ containing the constants provided above.

As a consequence, the number of eigenvalues of the corresponding matrix (of size $p \times p$ or $n \times n$) falling within an interval of small (but $O(1)$ with respect to n, p) size near E_+ is of order $O(n^{2/3})$. This is particularly interesting as, elsewhere in the support, the typical number of eigenvalues within a length of size $O(1)$ is naturally of order $O(n)$ (i.e., a non-trivial proportion of the total number of eigenvalues). As a consequence, the typical “spacing” between adjacent eigenvalues becomes larger near the edge and is precisely of order $1/n^{2/3} = n^{-2/3}$, which is (not surprisingly) also the typical fluctuation of the largest eigenvalues according to the Tracy-Widom theorem, Theorem 2.15.

When $c = 1$ in the Marčenko-Pastur limit though, due to the presence of the term x in the denominator of the density $\mu(dx)$, we obtain instead that, for x near the left edge $E_- = 0$,

$$\mu(dx) \simeq_{x=0+} \frac{1}{\pi\sqrt{x}} dx$$

so that the density diverges at zero and the integral of the density on $[0, \epsilon]$ is instead of order $O(\sqrt{\epsilon})$: consequently, the typical number of eigenvalues is here extremely large, of order n^2 (extremely locally of course).

Exercise 6 (The $\sqrt{|x - E|}$ behavior in elaborate models). We here seek to extend the results in Exercise 5 to the sample covariance matrix model $\frac{1}{n}\mathbf{X}\mathbf{X}^\top$ where $\mathbf{X} = \mathbf{C}^{1/2}\mathbf{Z}$ with \mathbf{Z} having independent entries of zero mean, unit variance and \mathbf{C} having a bounded limiting spectral measure ν with fast decaying tails. We denote $\tilde{m}(z)$ the Stieltjes transform of the limiting spectral measure $\tilde{\mu}$ of $\frac{1}{n}\mathbf{X}^\top\mathbf{X}$.

Using Figure 2.5 as a reference and recalling the formulation for functional inverse

$$x(\tilde{m}) = -\frac{1}{\tilde{m}} + c \int \frac{t\nu(dt)}{1+t\tilde{m}} \quad (9.3)$$

extensively discussed in Section 2.3.1, visually justify that $x''(\tilde{m})$ can be (complex) analytically extended in the neighborhood of the local extrema of $x(\tilde{m})$ (that is, each point \tilde{m} where $x'(\tilde{m}) = 0$) into a function $z(\tilde{m})$ which must locally coincide with the inverse Stieltjes transform of $\tilde{m}(z)$.

Deduce that $\tilde{m}(z)$ must be of the form $\sqrt{z - E}$ near an edge E and conclude.

Correction 6 (The $\sqrt{|x - E|}$ behavior in elaborate models). We observe visually that $x''(\tilde{m}) \neq 0$ near the edges of the support (so that $x(\tilde{m})$ has a quadratic local behavior per Taylor expansion). As such, the function $z(\tilde{m})$, defined similar to $x(\tilde{m})$ but extended for $\tilde{m} \in \mathbb{C} \setminus \{-1/\text{supp}(\nu)\}$, is analytic and locally invertible into a function $M(z)$. This function $M(z)$ coincides with the Stieltjes transform $\tilde{m}(z)$ of $\tilde{\mu}$ for all z real outside the support. By analyticity (in particular of all its derivatives), for z near the edges of the support (say for $z = x + i\varepsilon$ with x inside the support and $\varepsilon \ll 1$), $\tilde{m}(z)$ and its inverse $z(\tilde{m})$ must enjoy the same functional behavior: the latter is in particular locally quadratic, and thus, as its inverse, the former has a local square-root behavior. Since the density equals the imaginary part of $\tilde{m}(z)$ for z near the real axis, this square-root behavior propagates to the density which must then be of the form $\sqrt{z - E}$.

In passing, note that we implicitly assumed the existence of asymptotes for $-1/\tilde{m} \in \text{supp}(\nu)$ and the local extrema \tilde{m}_0 such that $x'(\tilde{m}_0) = 0$ (as in Figure 2.5), which, depending on the limiting spectral measure ν of \mathbf{C} , may not always be the case. In fact, unlike in Figure 2.5, $x(\tilde{m})$ may not have asymptotes on the edges of its domain of definition (i.e., the set of \tilde{m} such that $-1/\tilde{m} \notin \text{supp}(\nu)$) when ν has slow decaying tails near an edge. This is what occurs in Figure 9.1 where \mathbf{C} is of the form $\mathbf{C} = \frac{1}{n}\mathbf{D}^{\frac{1}{2}}\mathbf{Z}\mathbf{Z}^T\mathbf{D}^{\frac{1}{2}}$ for \mathbf{D} having a discrete limiting spectral measure and \mathbf{Z} has i.i.d. $\mathcal{N}(0, 1)$ entries; this example will be discussed in more detail in Exercise 7.




Figure 9.1: Functional inverse of $\tilde{m}(z)$, the Stieltjes transform of the limiting spectral measure of $\frac{1}{n}\mathbf{C}^{\frac{1}{2}}\mathbf{X}\mathbf{X}^T\mathbf{C}^{\frac{1}{2}}$ with $\mathbf{C} = \frac{1}{n}\mathbf{D}^{\frac{1}{2}}\mathbf{Z}\mathbf{Z}^T\mathbf{D}^{\frac{1}{2}}$ for $\mathbf{D} = \text{diag}[1_{p/3}, 3 \cdot 1_{p/3}, 10 \cdot 1_{p/3}]$, independent Gaussian $\mathbf{X}, \mathbf{Z} \in \mathbb{R}^{p \times n}$ and $p/n = 1/10$. Link to code: [\[Matlab\]](#) and [\[Python\]](#).

Exercise 7 (Further results on $x(\tilde{m})$). We aim in this exercise to justify some of the visual observations in Figure 2.5 with the help of

$$x(\tilde{m}) = -\frac{1}{\tilde{m}} + c \int \frac{t\nu(dt)}{1+t\tilde{m}}. \quad (9.4)$$

Show that, for $\tilde{m}_1 \neq \tilde{m}_2$ such that $x'(\tilde{m}_1), x'(\tilde{m}_2) > 0$, we cannot have $x(\tilde{m}_1) = x(\tilde{m}_2)$: that is, the increasing segments of $x(\tilde{m})$ never “overlap”.

Besides, show that, if $\tilde{m}_1 < \tilde{m}_2$ are both of the same sign, and $x'(\tilde{m}_1), x'(\tilde{m}_2) > 0$, then $x(\tilde{m}_1) < x(\tilde{m}_2)$: that is, the increasing segments of $x(\tilde{m})$ never “swap”. To this end, we may prove the intermediary result

$$(\tilde{m}_1 - \tilde{m}_2) \left(1 - \int \frac{c\tilde{m}_1 \tilde{m}_2 t^2 \nu(dt)}{(1+t\tilde{m}_1)(1+t\tilde{m}_2)} \right) = \tilde{m}_1 \tilde{m}_2 (x(\tilde{m}_1) - x(\tilde{m}_2)) \quad (9.5)$$

and use the Cauchy-Schwarz inequality to control the left-hand side parenthesis.

Finally show that, if ν has bounded support, then $x(\tilde{m}) \rightarrow 0$ as $\tilde{m} \rightarrow \pm\infty$.

As a final remark, note that the only important observation about Figure 2.5 which we have not shown here is the fact that the points \tilde{m} where $x'(\tilde{m}) = 0$

must exist. In fact, this is not always the case and heavily depends on the nature of the tails of the measure ν . Justify in particular that, for some ν , there may be no asymptote on the edges of the domain of definition of $x(\cdot)$ (as opposed to what is seen in Figure 2.5).

Correction 7 (Further results on $x(\tilde{m})$). If $x(\tilde{m}_1) = x(\tilde{m}_2)$ for \tilde{m}_1, \tilde{m}_2 such that $x'(\tilde{m}_1), x'(\tilde{m}_2) > 0$, then $x(\tilde{m})$ is locally invertible around both \tilde{m}_1 and \tilde{m}_2 with inverse the Stieltjes transform $\tilde{m}(\cdot)$ of $\tilde{\mu}$. In particular, $\tilde{m}(x(\tilde{m}_1))$ and $\tilde{m}(x(\tilde{m}_2))$ are both uniquely defined (since the Stieltjes transform of a valid z outside the support is unique) and equal to $\tilde{m}_1 \neq \tilde{m}_2$, respectively. It is thus impossible that $x(\tilde{m}_1) = x(\tilde{m}_2)$ if $\tilde{m}_1 \neq \tilde{m}_2$.

If $\tilde{m}_1 < \tilde{m}_2$ and $x'(\tilde{m}_1), x'(\tilde{m}_2) > 0$, then

$$x(\tilde{m}_1) - x(\tilde{m}_2) = (\tilde{m}_1 - \tilde{m}_2) \left[\frac{1}{\tilde{m}_1 \tilde{m}_2} - c \int \frac{t^2 \nu(dt)}{(1 + t\tilde{m}_1)(1 + t\tilde{m}_2)} \right]$$

or equivalently

$$(x(\tilde{m}_1) - x(\tilde{m}_2)) \tilde{m}_1 \tilde{m}_2 = (\tilde{m}_1 - \tilde{m}_2) \left[1 - c \int \frac{t^2 \tilde{m}_1 \tilde{m}_2 \nu(dt)}{(1 + t\tilde{m}_1)(1 + t\tilde{m}_2)} \right].$$

In addition, observe that, taking the limit where $\tilde{m}_1 - \tilde{m}_2 \rightarrow 0$, we have the derivative formulation

$$x'(\tilde{m}) \tilde{m}^2 = 1 - c \int \frac{t^2 \tilde{m}^2 \nu(dt)}{(1 + t\tilde{m})^2}$$

which is positive for both $\tilde{m} = \tilde{m}_1$ and $\tilde{m} = \tilde{m}_2$. Using the Cauchy-Schwarz inequality, we thus find that

$$\left| c \int \frac{t^2 \tilde{m}_1 \tilde{m}_2 \nu(dt)}{(1 + t\tilde{m}_1)(1 + t\tilde{m}_2)} \right| \leq \sqrt{c \int \frac{t^2 \tilde{m}_1^2 \nu(dt)}{(1 + t\tilde{m}_1)^2} \cdot c \int \frac{t^2 \tilde{m}_2^2 \nu(dt)}{(1 + t\tilde{m}_2)^2}} < 1.$$

As a consequence,

$$(x(\tilde{m}_1) - x(\tilde{m}_2)) \frac{\tilde{m}_1 \tilde{m}_2}{\tilde{m}_1 - \tilde{m}_2} > 0.$$

Since $\tilde{m}_1 \tilde{m}_2 > 0$, we conclude that $x(\tilde{m}_1) - x(\tilde{m}_2)$ is of the same sign as $\tilde{m}_1 - \tilde{m}_2$, as requested.

If ν has bounded support, by the dominated convergence theorem, it easily follows that $x(\tilde{m}) \rightarrow 0$ when $\tilde{m} \rightarrow \pm\infty$.

Denote \tilde{m}_0 an edge of the domain of definition of $x(\cdot)$, to see the asymptotes as in Figure 2.5 we need to have the integral $\int \frac{t\nu(dt)}{1+t\tilde{m}} \rightarrow \infty$ diverge as $\tilde{m} \rightarrow \tilde{m}_0$. However, if $\nu(dt)$ behaves locally like $(-1/\tilde{m}_0 - t)^\alpha$ around $t = -1/\tilde{m}_0$, the integral $\int \frac{t\nu(dt)}{1+t\tilde{m}}$ may not diverge for a sufficiently large α , and therefore no such asymptote can be observed. Figure 9.1 provides an illustrating example of this peculiar behavior, here for edges of order $\sqrt{-1/\tilde{m}_0 - t}$.

Exercise 8 (Alternative estimates of $\frac{1}{p} \text{tr}(\frac{1}{n} \mathbf{X}\mathbf{X}^\top)^2$). Let $\mathbf{X} = \mathbf{C}^{\frac{1}{2}}\mathbf{Z}$ for $\mathbf{Z} \in \mathbb{R}^{p \times n}$ with independent standard Gaussian entries, and \mathbf{C} deterministic symmetric nonnegative definite, of bounded operator norm, and limiting spectral measure ν .

Determine the limit, as $n, p \rightarrow \infty$ and $p/n \rightarrow c \in (0, \infty)$ of the (empirical) second order moment

$$M_2 = \frac{1}{p} \text{tr} \left(\frac{1}{n} \mathbf{X}\mathbf{X}^\top \right)^2$$

as a function of the moments of ν .

Retrieve the same result from the results of Exercise 1 along with the expression of the Stieltjes transform $m(z)$ of the limiting spectrum μ of $\frac{1}{n} \mathbf{X}\mathbf{X}^\top$. It may be useful to first show that $m(z)$ is also solution to

$$m(z) = \int \frac{\nu(dt)}{-z(1 + ctm(z)) + (1 - c)t}. \quad (9.6)$$

Correction 8 (Alternative estimates of $\frac{1}{p} \text{tr}(\frac{1}{n} \mathbf{X}\mathbf{X}^\top)^2$). A direct estimate is easily obtained by merely evaluating the expectation $\mathbb{E}[\text{tr}(\mathbf{C}^{\frac{1}{2}}\mathbf{Z}\mathbf{Z}^\top\mathbf{C}^{\frac{1}{2}}/n)^2]$. By the unitary invariance of the Gaussian matrix \mathbf{Z} (i.e., $\mathbf{U}\mathbf{Z} \sim \mathbf{Z}$ in law for all orthogonal matrix \mathbf{U}), we may assume $\mathbf{C} = \text{diag}\{[\mathbf{C}]_{ii}\}_{i=1}^p$ is diagonal. Therefore,

$$\frac{1}{p} \mathbb{E} \left[\text{tr} \left(\frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \mathbf{Z}^\top \mathbf{C}^{\frac{1}{2}} \right)^2 \right] = \frac{1}{n^2 p} \sum_{i,i'=1}^p \sum_{j,j'=1}^n \mathbf{C}_{ii} \mathbf{C}_{i'i'} \mathbb{E}[\mathbf{Z}_{ij} \mathbf{Z}_{i'j} \mathbf{Z}_{i'j'} \mathbf{Z}_{ij'}].$$

Of these sums, the case $i = i'$ and $j = j'$ simultaneously brings $\mathbb{E}[\mathbf{Z}_{ij} \mathbf{Z}_{i'j} \mathbf{Z}_{i'j'} \mathbf{Z}_{ij'}] = \mathbb{E}[\mathbf{Z}_{ij}^4] = 3$; the other non-trivial case is when $i = i'$ with $j \neq j'$, or $j = j'$ with $i \neq i'$, which both yield $\mathbb{E}[\mathbf{Z}_{ij} \mathbf{Z}_{i'j} \mathbf{Z}_{i'j'} \mathbf{Z}_{ij'}] = 1$. As a consequence,

$$\begin{aligned} \mathbb{E} \left[\frac{1}{p} \text{tr} \left(\frac{1}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{Z} \mathbf{Z}^\top \mathbf{C}^{\frac{1}{2}} \right)^2 \right] &= \frac{3n \text{tr } \mathbf{C}^2 + n(n-1) \text{tr } \mathbf{C}^2 + n[(\text{tr } \mathbf{C})^2 - \text{tr } \mathbf{C}^2]}{n^2 p} \\ &= \frac{1}{p} \text{tr } \mathbf{C}^2 + c \left(\frac{1}{p} \text{tr } \mathbf{C} \right)^2 + O(n^{-1}) \end{aligned}$$

where we recall that $c = \lim p/n$. In the large n, p limit, all empirical moments converge almost surely (this may be ensured by a control on their variances), and we thus find that

$$M_2 = \frac{1}{p} \text{tr } \mathbf{C}^2 + c \left(\frac{1}{p} \text{tr } \mathbf{C} \right)^2 + O(n^{-1})$$

almost surely. It is in particular interesting to note that, as $c \rightarrow 0$, we correctly recover the standard $\text{tr } \mathbf{C}^2/p$ estimate. In the large dimensional regime though, the additional non-negligible term $c(\text{tr } \mathbf{C}/p)^2$ contributes.

We have, from Exercise 1 that

$$M_2 = \frac{1}{2} \frac{d^2\delta}{dz^2}(0) + o(1)$$

almost surely, where $\delta(z) = -\frac{1}{z}m(z^{-1}) = \int \frac{\mu(dt)}{1-tz}$. Using $m(z) = \frac{1}{c}\tilde{m}(z) + \frac{1-c}{cz}$ and $\tilde{m}(z) = (-z + c \int \frac{t\nu(dt)}{1+t\tilde{m}(z)})^{-1}$, we may verify that

$$m(z) = \int \frac{\nu(dt)}{-z(1+ctm(z)) + (1-c)t}$$

(it suffices to replace $m(z)$ by $\frac{1}{c}\tilde{m}(z) + \frac{1-c}{cz}$ in this equation to fall back on the defining equation for $\tilde{m}(z)$). From this expression, we find that

$$\delta(z) = -\frac{1}{z}m(z^{-1}) = \int \frac{\nu(dt)}{1-ctz\delta(z) - (1-c)tz}$$

the derivative of which is given by

$$\delta'(z) = \int \frac{(1-c)t + ct\delta(z) + ctz\delta'(z)}{(1-ctz\delta(z) - (1-c)tz)^2} \nu(dt).$$

Remark that $\delta(z) = \int \frac{\mu(dt)}{1-tz}$ leads directly to $\delta(0) = 1$ and thus $\delta'(0) = \int t\nu(dt)$, as expected from Exercise 1. Differentiating a second time, we have

$$\delta''(z) = \int \left[\frac{2ct\delta'(z) + ctz\delta''(z)}{(1-ctz\delta(z) - (1-c)tz)^2} + 2 \frac{((1-c)t + ct\delta(z) + ctz\delta'(z))^2}{(1-ctz\delta(z) - (1-c)tz)^3} \right] \nu(dt)$$

which, after setting $z = 0$, gives

$$\delta''(0) = 2c \left(\int t\nu(dt) \right)^2 + 2 \int t^2 \nu(dt).$$

Consequently, we obtain

$$M_2 = c \left(\int t\nu(dt) \right)^2 + \int t^2 \nu(dt) + o(1) = c \left(\frac{1}{p} \text{tr } \mathbf{C} \right)^2 + \frac{1}{p} \text{tr } \mathbf{C}^2 + o(1)$$

almost surely, as requested.

Exercise 9 (Location of the zeros of $\tilde{m}(z)$). Figure 2.7 and Remark 2.12 both show that the zeros η_1, \dots, η_n of $m_{\mathbf{X}}(z)$, the Stieltjes transform of a symmetric matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$, are interlaced with the eigenvalues $\lambda_1, \dots, \lambda_n$ of \mathbf{X} .

In the sample covariance matrix case $\mathbf{X} = \frac{1}{n} \mathbf{Z}^T \mathbf{C} \mathbf{Z}$ with $\mathbf{Z} \in \mathbb{R}^{p \times n}$ having independent standard Gaussian entries and \mathbf{C} with limited spectral measure ν of bounded and connected support, this means that (up to zero eigenvalues) the roots η_i of $m_{\frac{1}{n} \mathbf{Z}^T \mathbf{C} \mathbf{Z}}(z)$ are all found in the limiting support of the empirical spectral measure $\tilde{\mu}$ of $\frac{1}{n} \mathbf{Z}^T \mathbf{C} \mathbf{Z}$, at the possible exception of the leftmost η_1 .

Using a change of variable involving $\tilde{m}(z)$ of the formula

$$0 = \frac{1}{2\pi i} \oint_{\Gamma} \frac{dw}{w}$$

for all Γ not enclosing zero, then the approximation $\tilde{m}(z) = m_{\frac{1}{n}} \mathbf{Z}^T \mathbf{C} \mathbf{Z}(z) + o(1)$ and finally the residue theorem, show that no zero of $m_{\frac{1}{n}} \mathbf{Z}^T \mathbf{C} \mathbf{Z}(z)$ can be found at macroscopic distance from the limiting support of $\tilde{\mu}$.

Correction 9 (Location of the zeros of $\tilde{m}(z)$). Letting $w = 1/\tilde{m}(z)$, we find that, upon the validity of the change of variable for the contour Γ ,

$$\frac{1}{2\pi i} \oint_{\Gamma} \frac{dw}{w} = -\frac{1}{2\pi i} \oint_{\tilde{\Gamma}} \frac{\tilde{m}'(z)}{\tilde{m}(z)} dz.$$

The idea is then to take $\tilde{\Gamma}$ any complex contour with leftmost real crossing slightly larger than zero and rightmost real crossing slightly smaller than the left edge of the support of $\tilde{\mu}$ (excluding the possible mass at 0 of course). From Figure 2.5 (and the discussions in that section), it appears that, for such a $\tilde{\Gamma}$, the associated Γ does not circle around zero, and thus the integral is null.

As such, in the large n, p limit, we have

$$0 = \frac{1}{2\pi i} \oint_{\tilde{\Gamma}} \frac{\tilde{m}'_{\frac{1}{n}} \mathbf{Z}^T \mathbf{C} \mathbf{Z}(z)}{\tilde{m}_{\frac{1}{n}} \mathbf{Z}^T \mathbf{C} \mathbf{Z}(z)} dz + o(1)$$

almost surely. By residue calculus, the right-hand side integral equals the cardinality of the η_i 's found inside $\tilde{\Gamma}$. As this is an integer and that, for all n, p large $o(1) < 1$, this cardinal must be zero: we therefore conclude that the smallest η_1 must also be found at an asymptotically vanishing distance from λ_1 (the smallest nonzero eigenvalue of $\frac{1}{n} \mathbf{Z}^T \mathbf{C} \mathbf{Z}$) almost surely.

Exercise 10 (Additive spiked model). Similar to Theorem 2.13, show the phase transition threshold for the additive model $\frac{1}{n} \mathbf{X} \mathbf{X}^T + \mathbf{P}$ for \mathbf{X} having i.i.d. entries of zero mean, unit variance and low rank $\mathbf{P} = \sum_{i=1}^k \ell_i \mathbf{u}_i \mathbf{u}_i^T$ with $\ell_1 > \dots > \ell_k > 0$, is determined by the condition

$$\ell_i > \sqrt{c}(1 + \sqrt{c})$$

with $c = \lim p/n$ as $p, n \rightarrow \infty$. Under this condition, show that the (almost sure) limiting value of the corresponding isolated eigenvalue $\hat{\lambda}_i$ of $\frac{1}{n} \mathbf{X} \mathbf{X}^T + \mathbf{P}$ is given by

$$\hat{\lambda}_i \xrightarrow{a.s.} \lambda_i = 1 + \ell_i + \frac{c}{\ell_i - c}.$$

Further show that, letting $\hat{\mathbf{u}}_i$ be the eigenvector associated with $\hat{\lambda}_i$, we have

$$|\hat{\mathbf{u}}_i^T \mathbf{u}_i|^2 \xrightarrow{a.s.} 1 - \frac{c}{(\ell_i - c)^2}.$$

Correction 10 (Additive spiked model). *The additive spiked model is in fact simpler to handle than the model in Theorem 2.13. We may answer the three questions (determination of the phase transition threshold, spike position and asymptotic eigenvector alignment) at once by writing*

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 = -\frac{1}{2\pi i} \oint_{\Gamma_{\lambda_i}} \mathbf{u}_i^\top \left(\frac{1}{n} \mathbf{X} \mathbf{X}^\top + \mathbf{P} - z \mathbf{I}_p \right)^{-1} \mathbf{u}_i dz$$

for all large n, p almost surely, where Γ_{λ_i} positively surrounds λ_i only. By Woodbury's identity, Lemma 2.7, we obtain

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 = -\frac{1}{2\pi i} \oint_{\Gamma_{\lambda_i}} \mathbf{u}_i^\top \mathbf{Q}(z) \mathbf{U} \mathbf{L} (\mathbf{I}_k + \mathbf{U}^\top \mathbf{Q}(z) \mathbf{U} \mathbf{L})^{-1} \mathbf{U}^\top \mathbf{Q}(z) \mathbf{u}_i dz \quad (9.7)$$

with $\mathbf{P} = \mathbf{U} \mathbf{L} \mathbf{U}^\top$ and $\mathbf{Q}(z) = (\frac{1}{n} \mathbf{X} \mathbf{X}^\top - z \mathbf{I}_p)^{-1}$ the resolvent of $\frac{1}{n} \mathbf{X} \mathbf{X}^\top$ that has asymptotically no residue inside Γ_{λ_i} . Since $\mathbf{U}^\top \mathbf{Q}(z) \mathbf{U} \xrightarrow{a.s.} m(z) \mathbf{I}_k$ by Theorem 2.4, we find that,

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 \xrightarrow{a.s.} -\frac{1}{2\pi i} \oint_{\Gamma_{\lambda_i}} \frac{\ell_i m^2(z)}{1 + \ell_i m(z)} dz. \quad (9.8)$$

By residue calculus, this term is null unless $1 + \ell_i m(\lambda_i) = 0$: this is the defining equation for λ_i . Since $m(x)$ increases on $x \in ((1 + \sqrt{c})^2, \infty)$ with image $(-1/(c + \sqrt{c}), 0)$, this equation has a solution if and only if $\ell_i > c + \sqrt{c}$; this provides the phase transition threshold. Using again the definition of $m(z)$, we find in this case that $\lambda_i = 1 + \ell_i + c/(\ell_i - c)$ as requested.

Exploiting the expression

$$m'(z) = \frac{m^2(z)}{1 - \frac{cm^2(z)}{(1+cm(z))^2}} \quad (9.9)$$

the residue calculus finally gives, by l'Hôpital's rule (i.e., by Taylor expanding around λ_i the numerator and denominator appearing in the residue calculus)

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 \xrightarrow{a.s.} \frac{m^2(\lambda_i)}{m'(\lambda_i)} = 1 - \frac{c}{(\ell_i - c)^2}$$

which concludes the proof.

Exercise 11 (Additive spiked model: the Wigner case). Let \mathbf{X} be symmetric with $[\mathbf{X}]_{ij}$, $i \geq j$, i.i.d. with zero mean and unit variance. As in Exercise 10, show that the “spiked” phase transition threshold for the model $\frac{1}{\sqrt{n}} \mathbf{X} + \mathbf{P}$ with $\mathbf{P} = \sum_{i=1}^k \ell_i \mathbf{u}_i \mathbf{u}_i^\top$, with $\ell_1 > \dots > \ell_k > 0$, is determined by the condition

$$\ell_i > 1$$

and that, under this condition, the isolated eigenvalue $\hat{\lambda}_i$ of $\frac{1}{\sqrt{n}} \mathbf{X} + \mathbf{P}$ associated with ℓ_i satisfies

$$\hat{\lambda}_i \xrightarrow{a.s.} \lambda_i = \ell_i + \frac{1}{\ell_i}. \quad (9.10)$$

Show finally that, for $\hat{\mathbf{u}}_i$ the eigenvector associated with $\hat{\lambda}_i$, we have

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 \xrightarrow{a.s.} 1 - \frac{1}{\ell_i^2}. \quad (9.11)$$

Correction 11 (Additive spiked model: the Wigner case). *The elements in the proof of Exercise 10 can be extensively reused, merely by replacing the definition of $m(z)$ by that of the Stieltjes transform of the semicircle law. In particular, here, $m(z) = \frac{-1}{z+m(z)}$ so that $m'(z) = \frac{m^2(z)}{1-m^2(z)}$.*

The asymptotic eigenvector alignment $|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2$ is again given by

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 \xrightarrow{a.s.} -\frac{1}{2\pi i} \oint_{\Gamma_{\lambda_i}} \frac{\ell_i m^2(z)}{1 + \ell_i m(z)} dz \quad (9.12)$$

where the result is non-trivial only if $1 + \ell_i m(\lambda_i) = 0$. On the set $(2, \infty)$ (outside the support of the semicircle law), $m(x)$ is increasing with $m(x=2+0) = -1$. The phase transition condition is thus $\ell_i > 1$, with $\lambda_i = \ell_i + 1/\ell_i$, as requested.

The residue of the above integral is $\frac{m^2(\lambda_i)}{m'(\lambda_i)}$ (again, as in Exercise 10) and we thus find that

$$|\hat{\mathbf{u}}_i^\top \mathbf{u}_i|^2 \xrightarrow{a.s.} 1 - \frac{1}{\ell_i^2} \quad (9.13)$$

as requested.

Exercise 12 (Sketch of proof of Theorem 2.17). *Inspired by the (sketch of) proof of Theorem 2.6, prove Theorem 2.17 using*

1. the trace lemma adapted to Haar random matrices, Lemma 2.16; and
2. Stein's lemma adapted to Haar random matrices, Lemma 2.17.

Correction 12 (Sketch of proof of Theorem 2.17). We first provide the derivation of Theorem 2.17 with the Haar trace lemma, Lemma 2.16. Before delving into the proof, we first proceed to a convenient change of variable: from the result of Theorem 2.17, for $\mathbf{Q}(z) = (\frac{p}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{U}^\top \mathbf{C}^{\frac{1}{2}} - z \mathbf{I}_p)^{-1}$ with $\mathbf{U} \in \mathbb{R}^{p \times n}$ having $n < p$ columns from a $p \times p$ Haar random matrix, we have, for $z < 0$, the deterministic equivalent

$$\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z) = \left(\frac{p/n}{1 + \delta(z)} \mathbf{C} - z \mathbf{I}_p \right)^{-1}$$

where $\delta(z)$ is the unique positive solution to

$$\delta(z) = \frac{p}{n} - 1 + \frac{p}{n} \frac{\delta(z)}{1 + \delta(z)} \frac{1}{n} \text{tr } \mathbf{C} \bar{\mathbf{Q}}(z).$$

This is indeed equivalent to the expected result by taking $\delta(z) = -\frac{p}{n} \frac{1}{z \tilde{m}_p(z)} - 1 > 0$ for $0 < -z \tilde{m}_p(z) < p/n \in (1, \infty)$. We now proceed to the proof by following the steps of the proof of Theorem 2.6.

For $\mathbf{u}_i \in \mathbb{R}^p$ the i -th column of \mathbf{U} , we have

$$\begin{aligned}\mathbf{I}_p + z\mathbb{E}[\mathbf{Q}] &= \mathbb{E}\left[\mathbf{Q} \cdot \frac{p}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{U}^\top \mathbf{C}^{\frac{1}{2}}\right] = \frac{p}{n} \sum_{i=1}^n \mathbb{E}[\mathbf{Q} \mathbf{C}^{\frac{1}{2}} \mathbf{u}_i \mathbf{u}_i^\top \mathbf{C}^{\frac{1}{2}}] \\ &= \frac{p}{n} \sum_{i=1}^n \mathbb{E}\left[\frac{\mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} \mathbf{u}_i \mathbf{u}_i^\top \mathbf{C}^{\frac{1}{2}}}{1 + \frac{p}{n} \mathbf{u}_i^\top \mathbf{C}^{\frac{1}{2}} \mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} \mathbf{u}_i}\right]\end{aligned}$$

where we denote $\mathbf{Q}_{-i} = (\frac{p}{n} \mathbf{C}^{\frac{1}{2}} \mathbf{U}_{-i} \mathbf{U}_{-i}^\top \mathbf{C}^{\frac{1}{2}} - z \mathbf{I}_p)^{-1}$ for $\mathbf{U}_{-i} \mathbf{U}_{-i}^\top = \mathbf{U} \mathbf{U}^\top - \mathbf{u}_i \mathbf{u}_i^\top$. At this point, note that, unlike in the proof of Theorem 2.6, here \mathbf{Q}_{-i} is no longer independent of \mathbf{u}_i and the usual “trace lemma”, Lemma 2.11, cannot be used. Instead, with Lemma 2.16, we have

$$\begin{aligned}\frac{p}{n} \mathbb{E}[\mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} \mathbf{u}_i \mathbf{u}_i^\top \mathbf{C}^{\frac{1}{2}}] &\simeq \frac{p}{n} \frac{1}{p-n} \mathbb{E}[\mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} (\mathbf{I}_p - \mathbf{U}_{-i} \mathbf{U}_{-i}^\top) \mathbf{C}^{\frac{1}{2}}] \\ &= \frac{1}{p-n} \mathbb{E}\left[\frac{p}{n} \mathbf{Q}_{-i} \mathbf{C} - (\mathbf{I}_p + z \mathbf{Q}_{-i})\right]\end{aligned}$$

as well as

$$\begin{aligned}\frac{p}{n} \mathbf{u}_i^\top \mathbf{C}^{\frac{1}{2}} \mathbf{Q}_{-i} \mathbf{C}^{\frac{1}{2}} \mathbf{u}_i &\simeq \frac{1}{p-n} \left(\frac{p}{n} \operatorname{tr} \mathbf{C} \mathbf{Q}_{-i} - (p+z \operatorname{tr} \mathbf{Q}_{-i}) \right) \\ &\simeq \frac{1}{p-n} \left(\frac{p}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}} - (p+z \operatorname{tr} \bar{\mathbf{Q}}) \right)\end{aligned}$$

with $\frac{1}{p} \operatorname{tr} \mathbf{C} \mathbf{Q}_{-i} \simeq \frac{1}{p} \operatorname{tr} \mathbf{C} \mathbf{Q} \simeq \frac{1}{p} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}}$ for some $\bar{\mathbf{Q}} \leftrightarrow \mathbf{Q}$ to be defined. Denoting

$$\delta(z) \equiv \frac{p-n}{n} + \frac{p}{n} \frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}} - \left(\frac{p}{n} + \frac{z}{n} \operatorname{tr} \bar{\mathbf{Q}} \right) \quad (9.14)$$

we thus have

$$\mathbf{I}_p + z\mathbb{E}[\mathbf{Q}] \simeq \frac{1}{n} \frac{\sum_{i=1}^n \mathbb{E}[\mathbf{Q}_{-i}] (\frac{p}{n} \mathbf{C} - z \mathbf{I}_p) - \mathbf{I}_p}{\delta(z)} \simeq \frac{\mathbb{E}[\mathbf{Q}] (\frac{p}{n} \mathbf{C} - z \mathbf{I}_p) - \mathbf{I}_p}{\delta(z)}$$

so that

$$\mathbb{E}[\mathbf{Q}(z)] \simeq \left(\frac{p/n}{1 + \delta(z)} \mathbf{C} - z \mathbf{I}_p \right)^{-1} \equiv \bar{\mathbf{Q}}(z).$$

Plugging back into (9.14) we obtain $\delta(z) = \frac{p}{n} - 1 + \frac{p}{n} \frac{\delta(z)}{1 + \delta(z)} \frac{1}{n} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}}(z)$ which concludes the asymptotic approximation of the expectation $\mathbb{E}[\mathbf{Q}] \simeq \bar{\mathbf{Q}}$. With additional concentration argument (for instance the Nash-Poincaré inequality for Haar matrices, Lemma 2.18) we conclude the proof.

We now present the derivation of Theorem 2.17 with Stein’s lemma for Haar matrices, Lemma 2.17, in the complex case for simplicity (see Remark 2.5). Since Lemma 2.17 only applies to square Haar random matrices, we may rewrite the resolvent as

$$\mathbf{Q}(z) = \left(\mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{D} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} - z \mathbf{I}_p \right)^{-1} \leftrightarrow \bar{\mathbf{Q}}(z)$$

for $\mathbf{U} \in \mathbb{C}^{p \times p}$ a random Haar matrix and diagonal $\mathbf{D} \in \mathbb{C}^{p \times p}$ with $\mathbf{D}_{ii} = d_i = \delta_{i \leq n}$ and some $\mathbf{Q}(z)$ to be specified. We start again from the relation $\mathbb{E}[\mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{D} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q}] = \mathbf{I}_p + z \mathbb{E}[\mathbf{Q}]$. Since \mathbf{U} is unitary, it suffices to consider the case of diagonal \mathbf{C} with $[\mathbf{C}]_{ii} = \alpha_i$. As such, defining

$$f(\mathbf{U}) = \sqrt{\alpha_q} \mathbf{U}_{qk} d_l [\mathbf{U}^*]_{lm} \sqrt{\alpha_m} \mathbf{Q}_{mq} \in \mathbb{C}$$

we have

$$\begin{aligned} \frac{\partial \mathbf{U}^*}{\partial \mathbf{U}_{ij}} &= -\mathbf{U}^* \mathbf{E}_{ij} \mathbf{U}^* \\ \frac{\partial \mathbf{Q}}{\partial \mathbf{U}_{ij}} &= -\mathbf{Q} \mathbf{C}^{\frac{1}{2}} \mathbf{E}_{ij} \mathbf{D} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q} + \mathbf{Q} \mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{D} \mathbf{U}^* \mathbf{E}_{ij} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q} \end{aligned}$$

and

$$\begin{aligned} f'_{ij}(\mathbf{U}) \mathbf{U}_{ij'} &= \sqrt{\alpha_q} \delta_{qi} \delta_{kj} d_l [\mathbf{U}^*]_{lm} \sqrt{\alpha_m} \mathbf{Q}_{mq} \mathbf{U}_{ij'} \\ &\quad - \sqrt{\alpha_q} \mathbf{U}_{qk} d_l [\mathbf{U}^*]_{li} [\mathbf{U}^*]_{jm} \sqrt{\alpha_m} \mathbf{Q}_{mq} \mathbf{U}_{ij'} \\ &\quad - \sqrt{\alpha_q} \mathbf{U}_{qk} d_l [\mathbf{U}^*]_{lm} \sqrt{\alpha_m} [\mathbf{Q} \mathbf{C}^{\frac{1}{2}}]_{mi} [\mathbf{D} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q}]_{jq} \mathbf{U}_{ij'} \\ &\quad + \sqrt{\alpha_q} \mathbf{U}_{qk} d_l [\mathbf{U}^*]_{lm} \sqrt{\alpha_m} [\mathbf{Q} \mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{D} \mathbf{U}^*]_{mi} [\mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q}]_{jq} \mathbf{U}_{ij'} \end{aligned}$$

so that with Lemma 2.17 for $j = k$ and $j' = l$,

$$\begin{aligned} 0 &= \mathbb{E} \left[\sum_{i=1}^p f'_{ij}(\mathbf{U}) \mathbf{U}_{ij'} \right] = \mathbb{E}[\sqrt{\alpha_q} \mathbf{U}_{ql} d_l [\mathbf{U}^*]_{lm} \sqrt{\alpha_m} \mathbf{Q}_{mq}] \\ &\quad - \mathbb{E}[\sqrt{\alpha_q} \mathbf{U}_{qk} [\mathbf{U}^*]_{km} \sqrt{\alpha_m} \mathbf{Q}_{mq} d_l] \\ &\quad - \mathbb{E}[\sqrt{\alpha_q} \mathbf{U}_{qk} [\mathbf{D} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q}]_{kq} d_l [\mathbf{U}^*]_{lm} \sqrt{\alpha_m} [\mathbf{Q} \mathbf{C}^{\frac{1}{2}} \mathbf{U}]_{ml}] \\ &\quad + \mathbb{E}[\sqrt{\alpha_q} \mathbf{U}_{qk} [\mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q}]_{kq} d_l [\mathbf{U}^*]_{lm} \sqrt{\alpha_m} [\mathbf{Q} \mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{D}]_{ml}] \end{aligned}$$

where we use $\mathbf{U}^* \mathbf{U} = \mathbf{I}_p$. Summing over k, l and m we get

$$\begin{aligned} 0 &= p \mathbb{E}[\alpha_q \mathbf{Q}_{qq}] - p \mathbb{E}[\alpha_q \mathbf{Q}_{qq}] - \mathbb{E}[[\mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{D} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q}]_{qq} \text{tr}(\mathbf{Q} \mathbf{C})] \\ &\quad + \mathbb{E}[\alpha_q [\mathbf{Q}]_{qq} \text{tr}(\mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q} \mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{D})] \end{aligned}$$

and with a concentration argument for the traces we have

$$\begin{aligned} \mathbb{E}[\mathbf{Q}]_{qq} &\simeq \frac{\text{tr}(\mathbb{E}[\mathbf{Q}] \mathbf{C})}{\alpha_q \mathbb{E}[\text{tr}(\mathbf{C}^{\frac{1}{2}} \mathbf{U} \mathbf{D} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q})] - z \text{tr}(\mathbb{E}[\mathbf{Q}] \mathbf{C})} \\ &\simeq \frac{\frac{1}{p} \text{tr} \bar{\mathbf{Q}} \mathbf{C}}{\alpha_q + \alpha_q \frac{z}{p} \text{tr} \bar{\mathbf{Q}} - \frac{z}{p} \text{tr} \bar{\mathbf{Q}} \mathbf{C}} \end{aligned}$$

so that $\bar{\mathbf{Q}}$ should take the form

$$\bar{\mathbf{Q}} = \left(\frac{p + z \text{tr} \bar{\mathbf{Q}}}{\text{tr} \mathbf{C} \bar{\mathbf{Q}}} \mathbf{C} - z \mathbf{I}_p \right)^{-1}$$

and therefore

$$\frac{1}{p} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}} \simeq \frac{1}{p} \operatorname{tr} \mathbf{C} \left(\frac{1 + \frac{z}{p} \operatorname{tr} \bar{\mathbf{Q}}}{\frac{1}{p} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}}} \mathbf{C} - z \mathbf{I}_p \right)^{-1}. \quad (9.15)$$

Similarly, for $\tilde{\mathbf{Q}}(z) = (\mathbf{D}^{\frac{1}{2}} \mathbf{U}^* \mathbf{C} \mathbf{U} \mathbf{D}^{\frac{1}{2}} - z \mathbf{I}_p)^{-1}$ we have

$$\frac{1}{p} \operatorname{tr} \mathbf{D} \tilde{\mathbf{Q}} \simeq \frac{1}{p} \operatorname{tr} \mathbf{D} \left(\frac{1 + \frac{z}{p} \operatorname{tr} \bar{\mathbf{Q}}}{\frac{1}{p} \operatorname{tr} \mathbf{D} \tilde{\mathbf{Q}}} \mathbf{D} - z \mathbf{I}_p \right)^{-1}. \quad (9.16)$$

To connect (9.15) to (9.16) and close the loop, it suffices to note that

$$\frac{1}{p} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}} \cdot \frac{1}{p} \operatorname{tr} \mathbf{D} \tilde{\mathbf{Q}} \simeq \left(1 + \frac{z}{p} \operatorname{tr} \bar{\mathbf{Q}} \right) \cdot \frac{1}{p} \operatorname{tr} \bar{\mathbf{Q}},$$

where we use the fact that $\mathbf{D}^{\frac{1}{2}} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}} \mathbf{Q} = \tilde{\mathbf{Q}} \mathbf{D}^{\frac{1}{2}} \mathbf{U}^* \mathbf{C}^{\frac{1}{2}}$. As a consequence, denoting $\tilde{m}_p(z) = -\frac{1}{z} \frac{p+z \operatorname{tr} \bar{\mathbf{Q}}}{\operatorname{tr} \mathbf{C} \bar{\mathbf{Q}}}$ we get

$$\tilde{m}_p(z) = \left(-z - (zc + z^2 \tilde{m}_p(z)) \frac{1}{p} \operatorname{tr} \mathbf{C} \bar{\mathbf{Q}} \right)^{-1}, \quad \bar{\mathbf{Q}}(z) = -\frac{1}{z} (\mathbf{I}_p + \tilde{m}_p(z) \mathbf{C})^{-1}$$

which, together with additional concentration arguments, concludes the proof.

Exercise 13 (Higher-order deterministic equivalent). *Theorem 2.4 provides a deterministic equivalent for the resolvent $\mathbf{Q} = (\frac{1}{n} \mathbf{X} \mathbf{X}^\top - z \mathbf{I}_p)^{-1}$ for $\mathbf{X} \in \mathbb{R}^{p \times n}$ having i.i.d. zero mean and unit variance entries, which, according to Notation 1, provides access to the asymptotic behavior of $\mathbf{a}^\top \mathbf{Q} \mathbf{b}$. In many machine learning applications, however, the object of natural interest (e.g., the mean squared error in a regression context and the variance in a classification context) often involves the asymptotic behavior of $\mathbf{a}^\top \mathbf{Q} \mathbf{A} \mathbf{Q} \mathbf{b}$ which requires a deterministic equivalent for random matrices of the type $\mathbf{Q} \mathbf{A} \mathbf{Q}$, for some \mathbf{A} independent of \mathbf{Q} . In particular, for $\mathbf{Q} \leftrightarrow \bar{\mathbf{Q}}$ (such that $\|\mathbb{E}[\mathbf{Q}] - \bar{\mathbf{Q}}\| \rightarrow 0$), $\bar{\mathbf{Q}} \mathbf{A} \bar{\mathbf{Q}}$ is in general not a deterministic equivalent for $\mathbf{Q} \mathbf{A} \mathbf{Q}$. This is due to the fact that*

$$\mathbb{E}[\mathbf{Q} \mathbf{A} \mathbf{Q}] \not\simeq \mathbb{E}[\mathbf{Q}] \mathbf{A} \mathbb{E}[\mathbf{Q}].$$

Instead, prove that, in the setting of Theorem 2.4, one has

$$\mathbf{Q}(z) \mathbf{A} \mathbf{Q}(z) \leftrightarrow m^2(z) \mathbf{A} + \frac{1}{n} \operatorname{tr} \mathbf{A} \cdot \frac{m'(z)m^2(z)}{(1 + cm(z))^2} \mathbf{I}_p. \quad (9.17)$$

As a sanity check, using the fact that $\partial \mathbf{Q}(z)/\partial z = \mathbf{Q}^2(z)$ and taking $\mathbf{A} = \mathbf{I}_p$ in the equation above, confirm that

$$\mathbf{Q}^2(z) \leftrightarrow m'(z) \mathbf{I}_p$$

for $m'(z) = \frac{m^2(z)}{1 - \frac{cm^2(z)}{(1+cm(z))^2}}$ obtained from differentiating the Marčenko–Pastur equation (2.9).

Correction 13 (Higher-order deterministic equivalent). *First recall from Theorem 2.4 that*

$$\mathbf{Q}(z) \leftrightarrow \bar{\mathbf{Q}}(z) = m(z)\mathbf{I}_p = \left(\frac{1}{1+cm(z)} - z \right)^{-1} \mathbf{I}_p, \quad m(z) = \frac{1}{-z + \frac{1}{1+cm(z)}}.$$

We then have

$$\begin{aligned} \mathbb{E}[\mathbf{Q}\mathbf{A}\mathbf{Q}] &= \mathbb{E}[\mathbf{Q}\mathbf{A}\bar{\mathbf{Q}}] + \mathbb{E}[\mathbf{Q}\mathbf{A}(\mathbf{Q} - \bar{\mathbf{Q}})] \\ &\simeq \bar{\mathbf{Q}}\mathbf{A}\bar{\mathbf{Q}} + \mathbb{E}\left[\mathbf{Q}\mathbf{A}\mathbf{Q}\left(\frac{\mathbf{I}_p}{1+\frac{1}{n}\text{tr }\bar{\mathbf{Q}}} - \frac{1}{n}\mathbf{X}\mathbf{X}^\top\right)\right]\bar{\mathbf{Q}} \\ &= \bar{\mathbf{Q}}\mathbf{A}\bar{\mathbf{Q}} + \frac{\mathbb{E}[\mathbf{Q}\mathbf{A}\mathbf{Q}]\bar{\mathbf{Q}}}{1+\frac{1}{n}\text{tr }\bar{\mathbf{Q}}} - \frac{1}{n}\sum_{i=1}^n \mathbb{E}\left[\frac{\mathbf{Q}\mathbf{A}\mathbf{Q}_{-i}\mathbf{x}_i\mathbf{x}_i^\top}{1+\frac{1}{n}\mathbf{x}_i^\top\mathbf{Q}_{-i}\mathbf{x}_i}\right]\bar{\mathbf{Q}} \\ &= \bar{\mathbf{Q}}\mathbf{A}\bar{\mathbf{Q}} + \frac{\mathbb{E}[\mathbf{Q}\mathbf{A}\mathbf{Q}]\bar{\mathbf{Q}}}{1+\frac{1}{n}\text{tr }\bar{\mathbf{Q}}} - \frac{1}{n}\sum_{i=1}^n \frac{\mathbb{E}[\mathbf{Q}_{-i}\mathbf{A}\mathbf{Q}_{-i}\mathbf{x}_i\mathbf{x}_i^\top]}{1+\frac{1}{n}\text{tr }\bar{\mathbf{Q}}}\bar{\mathbf{Q}} \\ &\quad + \frac{1}{n}\sum_{i=1}^n \mathbb{E}\left[\frac{\mathbf{Q}_{-i}\mathbf{x}_i\frac{1}{n}\mathbf{x}_i^\top\mathbf{Q}_{-i}\mathbf{A}\mathbf{Q}_{-i}\mathbf{x}_i\mathbf{x}_i^\top}{(1+\frac{1}{n}\text{tr }\bar{\mathbf{Q}})^2}\right]\bar{\mathbf{Q}} \\ &\simeq \bar{\mathbf{Q}}\mathbf{A}\bar{\mathbf{Q}} + \frac{1}{n}\sum_{i=1}^n \mathbb{E}\left[\mathbf{Q}_{-i}\frac{\frac{1}{n}\text{tr}(\mathbf{Q}_{-i}\mathbf{A}\mathbf{Q}_{-i})}{(1+\frac{1}{n}\text{tr }\bar{\mathbf{Q}})^2}\right]\bar{\mathbf{Q}} \simeq \bar{\mathbf{Q}}\mathbf{A}\bar{\mathbf{Q}} + \frac{\frac{1}{n}\text{tr}(\mathbb{E}[\mathbf{Q}\mathbf{A}\mathbf{Q}])}{(1+\frac{1}{n}\text{tr }\bar{\mathbf{Q}})^2}\bar{\mathbf{Q}}^2 \end{aligned}$$

where we apply Lemma 2.8 twice and use the fact that $\mathbb{E}[\mathbf{Q}] \simeq \bar{\mathbf{Q}}$ and $\mathbb{E}[\mathbf{Q}\mathbf{A}\mathbf{Q}] \simeq \mathbb{E}[\mathbf{Q}_{-i}\mathbf{A}\mathbf{Q}_{-i}]$ for \mathbf{A} of bounded norm. Taking the (normalized) trace and gathering the terms proportional to $\text{tr}(\mathbb{E}[\mathbf{Q}\mathbf{A}\mathbf{Q}])$, we obtain

$$\begin{aligned} \frac{1}{n}\text{tr}(\mathbb{E}[\mathbf{Q}\mathbf{A}\mathbf{Q}]) &\simeq \left(1 - \frac{\frac{1}{n}\text{tr }\bar{\mathbf{Q}}^2}{(1+\frac{1}{n}\text{tr }\bar{\mathbf{Q}})^2}\right)^{-1} \frac{1}{n}\text{tr }\bar{\mathbf{Q}}\mathbf{A}\bar{\mathbf{Q}} = \frac{m^2(z)\frac{1}{n}\text{tr }\mathbf{A}}{1 - \frac{cm^2(z)}{(1+cm(z))^2}} \\ &= m'(z)\frac{1}{n}\text{tr }\mathbf{A} \end{aligned}$$

where we recall that $m'(z) = \frac{m^2(z)}{1 - \frac{cm^2(z)}{(1+cm(z))^2}}$ by differentiating (2.9) and

$$\mathbf{Q}\mathbf{A}\mathbf{Q} \leftrightarrow \bar{\mathbf{Q}}\mathbf{A}\bar{\mathbf{Q}} + \frac{1}{n}\text{tr }\mathbf{A} \cdot \frac{m'(z)}{(1+cm(z))^2}\bar{\mathbf{Q}}^2 = m^2(z)\mathbf{A} + \frac{1}{n}\text{tr }\mathbf{A} \cdot \frac{m'(z)m^2(z)}{(1+cm(z))^2}\mathbf{I}_p.$$

In particular, with $\mathbf{A} = \mathbf{I}_p$, we obtain $\mathbf{Q}(z)\mathbf{A}\mathbf{Q}(z) \leftrightarrow m'(z)\mathbf{I}_p$.

Exercise 14 (Concentration of matrix quadratic forms). *Recalling the definitions and notations of Section 2.7, let $\mathbf{X} \in \mathbb{R}^{p \times n}$ be a random matrix satisfying*

$$\mathbf{X} \propto Ce^{c^2}, \text{ and } \|\mathbb{E}[\mathbf{X}]\| \leq K$$

for some $K, C, c > 0$. Given $\mathbf{A} \in \mathbb{R}^{p \times p}$ deterministic, we aim to prove the linear concentration of $\mathbf{X}^\top\mathbf{A}\mathbf{X}$ in $(\mathbb{R}^{n \times n}, \|\cdot\|_F)$. To this end, we consider a

deterministic matrix $\mathbf{B} \in \mathbb{R}^{n \times n}$ such that $\|\mathbf{B}\|_F \leq 1$ and study the behavior of $\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})$. Consider first the singular value decomposition

$$\mathbf{A} = \mathbf{U}_\mathbf{A} \Lambda_\mathbf{A} \mathbf{V}_\mathbf{A}^\top, \quad \mathbf{B} = \mathbf{U}_\mathbf{B} \Lambda_\mathbf{B} \mathbf{V}_\mathbf{B}^\top,$$

with $\mathbf{U}_\mathbf{A}, \mathbf{V}_\mathbf{A} \in \mathbb{R}^{p \times p}$ and $\mathbf{U}_\mathbf{B}, \mathbf{V}_\mathbf{B} \in \mathbb{R}^{n \times n}$ orthogonal matrices, $\Lambda_\mathbf{A} \in \mathbb{R}^{p \times p}$, $\Lambda_\mathbf{B} \in \mathbb{R}^{n \times n}$ diagonal matrices, and define $\tilde{\mathbf{X}}_1 = \mathbf{U}_\mathbf{A}^\top \mathbf{X} \mathbf{V}_\mathbf{B}$, $\tilde{\mathbf{X}}_2 = \mathbf{V}_\mathbf{A}^\top \mathbf{X} \mathbf{U}_\mathbf{B} \in \mathbb{R}^{p \times n}$. In the sequel, the constants $K', C', c' > 0$ are understood only depending on K, C, c and might change from line to line.

First show that there exist $K', C', c' > 0$ such that, for $t > K' \sqrt{\log(np)}$ and $\tilde{\mathbf{X}} \in \{\tilde{\mathbf{X}}_1, \tilde{\mathbf{X}}_2\}$

$$\mathbb{P}(\|\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}]\|_\infty \geq t) \leq C' e^{-c't^2/\log(np)}.$$

Deduce from the bound $\|\mathbb{E}[\mathbf{X}]\| \leq K$ that there exists a constant $K' > 0$ depending only on K, C, c such that

$$\mathbb{E}[\|\tilde{\mathbf{X}}\|_\infty] \leq K' \sqrt{\log(np)}.$$

This established, introduce the set $\mathcal{A}_\theta = \{\mathbf{X} \in \mathbb{R}^{p \times n}, \max\{\|\tilde{\mathbf{X}}_1\|_\infty, \|\tilde{\mathbf{X}}_2\|_\infty\} \leq \theta\} \subset \mathbb{R}^{p \times n}$ and show that for all $\theta \geq K' \sqrt{\log(np)}$ with $K' > 1$, we have

$$\mathbb{P}(\mathbf{X} \in \mathcal{A}_\theta^c) \leq C' e^{-c'\theta^2}$$

and that the mapping $\mathbf{X} \mapsto \text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})$ is $\theta\|\mathbf{A}\|_F$ -Lipschitz on \mathcal{A}_θ .

Introduce M , a median of $\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})$, and note that

$$\mathbb{P}(|\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X}) - M| \geq t, \mathbf{X} \in \mathcal{A}_\theta) \leq C' e^{-c't^2/(\theta\|\mathbf{A}\|_F)^2}.$$

Conclude by carefully choosing the parameter $\theta \geq K' \sqrt{\log(np)}$ and showing that

$$\mathbf{X}^\top \mathbf{A}\mathbf{X} \in \mathbb{E}[\mathbf{X}^\top \mathbf{A}\mathbf{X}] \pm C' e^{-c'.2/(\log(np)\|\mathbf{A}\|_F^2)} + C' e^{-c'./\|\mathbf{A}\|_F}.$$

Correction 14 (Concentration of matrix quadratic forms). First note that for $\|\mathbf{U}_\mathbf{A}\| = \|\mathbf{U}_\mathbf{B}\| = \|\mathbf{V}_\mathbf{A}\| = \|\mathbf{V}_\mathbf{B}\| = 1$, it follows from $\mathbf{X} \propto Ce^{c\cdot^2}$ that both $\tilde{\mathbf{X}}_1 = \mathbf{U}_\mathbf{A}^\top \mathbf{X} \mathbf{V}_\mathbf{B} \propto Ce^{c\cdot^2}$ and $\tilde{\mathbf{X}}_2 = \mathbf{V}_\mathbf{A}^\top \mathbf{X} \mathbf{U}_\mathbf{B} \propto Ce^{c\cdot^2}$ (linear and thus Lipschitz with constant one). Denote $\mathbf{e}_i \in \mathbb{R}^n$ the canonical basis vector of \mathbb{R}^n such that $[\mathbf{e}_i]_j = \delta_{ij}$, we thus have for $\mathbf{X} \in \{\tilde{\mathbf{X}}_1, \tilde{\mathbf{X}}_2\}$

$$\mathbb{P}(\|\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}]\|_\infty \geq t) \leq \mathbb{P}\left(\sup_{\substack{1 \leq i \leq p \\ 1 \leq j \leq n}} |\mathbf{e}_i^\top (\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}]) \mathbf{e}_j| \geq t\right) \leq np \cdot 2Ce^{-ct^2}.$$

Since $Ce^{\log(np)}e^{-ct^2} \leq C'e^{-c't^2/\log(np)}$ for $t \geq K' \sqrt{\log(np)}$ for some constants K', C', c' depending only on K, C, c , we have

$$\mathbb{P}(\|\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}]\|_\infty \geq t) \leq C' e^{-c't^2/\log(np)}$$

for $t \geq K'\sqrt{\log(np)}$ and therefore

$$\begin{aligned}\mathbb{E}[\|\tilde{\mathbf{X}}\|_\infty] &= \mathbb{E}\left[\|\mathbb{E}[\tilde{\mathbf{X}}] + (\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}])\|_\infty\right] \\ &\leq \|\mathbb{E}[\tilde{\mathbf{X}}]\|_\infty + \mathbb{E}[\|\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}]\|_\infty] \\ &\leq \|\mathbb{E}[\tilde{\mathbf{X}}]\| + \int_0^\infty \mathbb{P}\left(\|\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}]\|_\infty \geq t\right) dt \\ &\leq K + K'\sqrt{\log(np)} + \int_{K'\log(np)}^\infty \mathbb{P}\left(\|\tilde{\mathbf{X}} - \mathbb{E}[\tilde{\mathbf{X}}]\|_\infty \geq t\right) dt \\ &\leq K'\sqrt{\log(np)} + C'\sqrt{\log(np)} \int_0^\infty e^{-c't^2} dt \leq K'\sqrt{\log(np)}.\end{aligned}$$

Next, note that for $\theta \geq 2K'\sqrt{\log(np)}$, we have $\theta \geq \frac{\theta}{2} + \mathbb{E}[\|\tilde{\mathbf{X}}\|_\infty]$. Since $\tilde{\mathbf{X}} \propto Ce^{-c\cdot^2}$, we have $\|\tilde{\mathbf{X}}\|_\infty = \sup_{ij} |\mathbf{e}_i^\top \tilde{\mathbf{X}} \mathbf{e}_j| \propto Ce^{-c\cdot^2}$ and

$$\begin{aligned}\mathbb{P}(\mathbf{X} \in \mathcal{A}_\theta^c) &\leq \mathbb{P}\left(\|\tilde{\mathbf{X}}\|_\infty \geq \theta\right) \leq \mathbb{P}\left(\left|\|\tilde{\mathbf{X}}\|_\infty - \mathbb{E}[\|\tilde{\mathbf{X}}\|_\infty]\right| \geq \frac{\theta}{2}\right) \\ &\leq np \cdot 2Ce^{-c\theta^2/4} \leq C'e^{-c'\theta^2}\end{aligned}$$

for $\mathcal{A}_\theta = \{\mathbf{X} \in \mathbb{R}^{p \times n}, \max\{\|\tilde{\mathbf{X}}_1\|_\infty, \|\tilde{\mathbf{X}}_2\|_\infty\} \leq \theta\}$ and $K' > 1$. Besides, with $\Lambda_{\mathbf{A}} = \text{diag}\{\lambda_i(\mathbf{A})\}_{i=1}^p$ and $\Lambda_{\mathbf{B}} = \text{diag}\{\lambda_i(\mathbf{B})\}_{i=1}^n$, we find that for any $\mathbf{X} \in \mathcal{A}_\theta$,

$$\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X}) = \text{tr}(\Lambda_{\mathbf{B}} \tilde{\mathbf{X}}_1^\top \Lambda_{\mathbf{A}} \tilde{\mathbf{X}}_2) = \text{diag}(\Lambda_{\mathbf{A}})^\top (\tilde{\mathbf{X}}_1 \odot \tilde{\mathbf{X}}_2) \text{diag}(\Lambda_{\mathbf{B}})$$

where \odot is the Hadamard entry-wise product. Therefore

$$|\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})| \leq \|\mathbf{A}\|_F \cdot \|\mathbf{B}\|_F \cdot \|\tilde{\mathbf{X}}\|_F \cdot \|\tilde{\mathbf{X}}\|_\infty \leq \theta \|\mathbf{A}\|_F \cdot \|\mathbf{X}\|_F$$

which means that, for $\|\mathbf{B}\|_F \leq 1$, the map $\mathbf{X} \mapsto \text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})$ is $\theta \|\mathbf{A}\|_F$ -Lipschitz on \mathcal{A}_θ . As a consequence, we can bound

$$\begin{aligned}&\mathbb{P}(|\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X}) - \mathbb{E}[\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})]| \geq t) \\ &\leq \mathbb{P}(|\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X}) - M| \geq t, \mathbf{X} \in \mathcal{A}_\theta) + \mathbb{P}(\mathbf{X} \in \mathcal{A}_\theta^c) \\ &\leq C'e^{-c't^2/(\theta \|\mathbf{A}\|_F)^2} + C'e^{-c'\theta^2}\end{aligned}$$

where we used, in the last inequality, the Lipschitz nature of the map $\mathbf{X} \mapsto \text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})$ on the set \mathcal{A}_θ . Finally, by choosing $\theta = \max(K'\sqrt{\log(np)}, \sqrt{t/\|\mathbf{A}\|_F})$, we can bound

$$\begin{aligned}&\mathbb{P}(|\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X}) - \mathbb{E}[\text{tr}(\mathbf{B}\mathbf{X}^\top \mathbf{A}\mathbf{X})]| \geq t) \\ &\leq C'e^{-c't^2/(K'\sqrt{\log(np)} \|\mathbf{A}\|_F)^2} + C'e^{-c't/\|\mathbf{A}\|_F},\end{aligned}$$

from which we conclude on the linear concentration of $\mathbf{X}^\top \mathbf{A}\mathbf{X}$.

Exercise 15 (Towards spiked models in random tensors). Let $\mathcal{Y} \in \mathbb{R}^{n \times n \times n}$ be a three-way symmetric tensor, i.e., such that $[\mathcal{Y}]_{ijk}$ is constant to exchanges of its indexes, defined by

$$\mathcal{Y} = \ell \mathbf{x} \otimes \mathbf{x} \otimes \mathbf{x} + \frac{1}{\sqrt{n}} \mathcal{W}$$

where $\mathcal{W} \in \mathbb{R}^{n \times n \times n}$ has independent $\mathcal{N}(0, 1)$ entries up to symmetry, deterministic $\mathbf{x} \in \mathbb{R}^n$ of unit norm, and $[\mathbf{a} \otimes \mathbf{b} \otimes \mathbf{c}]_{ijk} = a_i b_j c_k$.

A possible definition of the “eigenvalue-eigenvector” pair $(\hat{\lambda}, \hat{\mathbf{u}})$ (without loss of generality such that $\hat{\lambda} \geq 0$ and $\|\hat{\mathbf{u}}\| = 1$) of a symmetric tensor \mathcal{Y} is the solution to [Lim, 2005]

$$\mathcal{Y} \cdot \hat{\mathbf{u}} \cdot \hat{\mathbf{u}} = \hat{\lambda} \hat{\mathbf{u}} \quad (9.18)$$

where $\mathcal{A} \cdot \mathbf{a} \cdot \mathbf{b} = \sum_{ij} [\mathcal{A}]_{ij} a_i b_j \in \mathbb{R}^n$ is the contraction of the tensor \mathcal{A} on the vectors $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$. The objective here is to characterize the (possible) spike $\hat{\lambda}$ as well as the associated eigenvector alignment $|\hat{\mathbf{u}}^\top \mathbf{x}|$ between the dominant eigenvector and \mathbf{x} .

Show first that the matrix $\mathbf{Y}_x = \mathcal{Y} \cdot \mathbf{x} = \sum_{i=1}^n \mathcal{Y}_{i..} x_i \in \mathbb{R}^{n \times n}$ takes the form

$$\mathbf{Y}_x = \ell \mathbf{x} \mathbf{x}^\top + \frac{1}{\sqrt{n}} \sum_{i=1}^n x_i \mathbf{W}_i \quad (9.19)$$

where $\mathbf{W}_i \in \mathbb{R}^{n \times n}$ is the i -th “layer” matrix of the tensor \mathcal{W} such that $[\mathbf{W}_i]_{ab} = \mathcal{W}_{iab}$.

Using the Gaussian method discussed in Section 2.2.2.2, show that the limiting spectral measure of \mathbf{Y}_x is the semicircle law supported on $[-2, 2]$ (we may discard the rank-one matrix $\ell \mathbf{x} \mathbf{x}^\top$ to retrieve this result). Then, using a spiked model analysis as in Section 2.5, show that

- for all $\ell > 0$, there must exist an isolated eigenvalue $\hat{\lambda}_x$ of \mathbf{Y}_x (thus no phase transition) asymptotically equal to (with high probability)

$$\hat{\lambda}_x \rightarrow \lambda_x = \sqrt{\ell^2 + 4};$$

- the eigenvector $\hat{\mathbf{u}}_x$ associated with $\hat{\lambda}_x$ satisfies (again with high probability)

$$|\hat{\mathbf{u}}_x^\top \mathbf{x}|^2 \rightarrow \frac{\ell}{\sqrt{\ell^2 + 4}}.$$

Conclude on an asymptotic upper bound for the quantity $\hat{\lambda} |\hat{\mathbf{u}}^\top \mathbf{x}|$.

Correction 15 (Towards spiked models in random tensors). After establishing (9.19), the main technical difficulty to obtain the semicircle limit lies in handling the multi-way symmetry of the tensor \mathcal{W} . Denoting the resolvent

$\mathbf{Q} = (\sum_{i=1}^n x_i \mathbf{W}_i / \sqrt{n} - z \mathbf{I}_n)^{-1}$ of $\frac{1}{\sqrt{n}} \sum_{i=1}^n x_i \mathbf{W}_i$, we write, with the help of Stein's lemma, Lemma 2.13, that

$$\frac{1}{\sqrt{n}} \sum_{l=1}^n x_l \mathbb{E}[\mathbf{W}_l \mathbf{Q}]_{ij} = \frac{1}{\sqrt{n}} \sum_{m,l} x_l \mathbb{E}[\mathcal{W}_{lim}[\mathbf{Q}]_{mj}] = \frac{1}{\sqrt{n}} \sum_{m,l} x_l \mathbb{E} \left[\frac{\partial [\mathbf{Q}]_{mj}}{\partial \mathcal{W}_{lim}} \right]. \quad (9.20)$$

At this point, note that by symmetry $\mathcal{W}_{abc} = \mathcal{W}_{bac} = \mathcal{W}_{cab} = \mathcal{W}_{cba} = \mathcal{W}_{bca} = \mathcal{W}_{acb}$ and therefore

$$\begin{aligned} \frac{\partial \mathbf{Q}}{\partial \mathcal{W}_{abc}} &= -\frac{1}{\sqrt{n}} \sum_{k=1}^n x_k \mathbf{Q} \frac{\partial \mathbf{W}_k}{\partial \mathcal{W}_{abc}} \mathbf{Q} \\ &= -\frac{1}{\sqrt{n}} \sum_{k=1}^n x_k \mathbf{Q} [(\mathbf{E}_{ab} + \mathbf{E}_{ba})\delta_{kc} + (\mathbf{E}_{ca} + \mathbf{E}_{ac})\delta_{kb} + (\mathbf{E}_{bc} + \mathbf{E}_{cb})\delta_{ka}] \mathbf{Q} \end{aligned}$$

with $\mathbf{E}_{ab} \in \mathbb{R}^{n \times n}$ the indicator matrix such that $[\mathbf{E}_{ab}]_{a'b'} = \delta_{aa'}\delta_{bb'}$. Plugging the above expression into (9.20) gives

$$\begin{aligned} \frac{1}{\sqrt{n}} \sum_{l=1}^n x_l \mathbb{E}[\mathbf{W}_l \mathbf{Q}]_{ij} &= -\frac{1}{n} \mathbb{E}[\mathbf{Q}^2]_{ij} - \frac{1}{n} \mathbb{E}[\text{tr } \mathbf{Q} \cdot [\mathbf{Q}]_{ij}] - \frac{1}{n} \mathbb{E}[\mathbf{x}^\top \mathbf{Q} \mathbf{x} \cdot [\mathbf{Q}]_{ij}] \\ &\quad - \frac{1}{n} \mathbb{E}[[\mathbf{Q} \mathbf{x}]_i [\mathbf{Q} \mathbf{x}]_j] - \frac{1}{n} \mathbb{E}[\text{tr } \mathbf{Q} \cdot [\mathbf{Q} \mathbf{x}]_j] x_i - \frac{1}{n} \mathbb{E}[\mathbf{Q}^2 \mathbf{x}]_j x_i \end{aligned}$$

where we used the fact that $\|\mathbf{x}\| = 1$. Using $\mathbf{Q} = -\frac{1}{z} \mathbf{I}_n + \frac{1}{z} \frac{1}{\sqrt{n}} \sum_{i=1}^n x_i \mathbf{W}_i \mathbf{Q}$, we then find

$$\begin{aligned} \mathbb{E}[\mathbf{Q}]_{ij} &= -\frac{1}{z} \delta_{ij} - \frac{1}{z} \frac{1}{n} \mathbb{E}[\mathbf{Q}^2]_{ij} - \frac{1}{z} \frac{1}{n} \mathbb{E}[\text{tr } \mathbf{Q} \cdot [\mathbf{Q}]_{ij}] - \frac{1}{z} \frac{1}{n} \mathbb{E}[\mathbf{x}^\top \mathbf{Q} \mathbf{x} \cdot [\mathbf{Q}]_{ij}] \\ &\quad - \frac{1}{z} \frac{1}{n} \mathbb{E}[[\mathbf{Q} \mathbf{x}]_i [\mathbf{Q} \mathbf{x}]_j] - \frac{1}{z} \frac{1}{n} \mathbb{E}[\text{tr } \mathbf{Q} \cdot [\mathbf{Q} \mathbf{x}]_j] x_i - \frac{1}{z} \frac{1}{n} \mathbb{E}[\mathbf{Q}^2 \mathbf{x}]_j x_i. \end{aligned}$$

This leads, with a concentration argument (using for instance the Nash-Poincaré inequality, Lemma 2.14, to show that the variances of $\frac{1}{n} \text{tr } \mathbf{Q}$ and $\mathbf{x}^\top \mathbf{Q} \mathbf{x}$ vanish as $n \rightarrow \infty$), to

$$\begin{aligned} \frac{1}{n} \text{tr } \mathbf{Q} &= -\frac{1}{z} - \frac{1}{z} \left(\frac{1}{n} \text{tr } \mathbf{Q} \right)^2 + O(n^{-1}) \\ \mathbf{x}^\top \mathbf{Q} \mathbf{x} &= -\frac{1}{z} - \frac{2}{z} \mathbf{x}^\top \mathbf{Q} \mathbf{x} \cdot \frac{1}{n} \text{tr } \mathbf{Q} + O(n^{-1}). \end{aligned}$$

Since $\frac{1}{n} \text{tr } \mathbf{Q}$ satisfies in the $n \rightarrow \infty$ limit the fixed-point equation of the Wigner semicircle law (i.e., $m^2(z) + zm(z) + 1 = 0$), the first part of the result unfolds. An interesting consequence of the results above is that, due to the existence of \mathbf{x} in the expression of \mathbf{Y}_x , $\mathbf{x}^\top \mathbf{Q} \mathbf{x}$ does not satisfy the same equation as $\frac{1}{n} \text{tr } \mathbf{Q}$. Precisely, we find that

$$\begin{aligned} \frac{1}{n} \text{tr } \mathbf{Q} &= -\frac{z}{2} + \frac{1}{2} \sqrt{z^2 - 4} + O(n^{-1}) \\ \mathbf{x}^\top \mathbf{Q} \mathbf{x} &= \frac{1}{\sqrt{z^2 - 4}} + O(n^{-1}) \end{aligned}$$

where the branch of the square root is chosen so that the right-hand side terms are Stieltjes transforms of (probability) measures. For further use, note that

$$\frac{\partial}{\partial z} \mathbf{x}^\top \mathbf{Q}(z) \mathbf{x} = -\frac{z}{(z^2 - 4)^{\frac{3}{2}}} + O(n^{-1}) \quad (9.21)$$

with the same convention on the square root.

As in Section 2.5 for spiked models, we write, for $(\hat{\lambda}_x, \hat{\mathbf{u}}_x)$ an eigenvalue-eigenvector pair of \mathbf{Y}_x ,

$$\begin{aligned} |\hat{\mathbf{u}}_x^\top \mathbf{x}|^2 &= -\frac{1}{2\pi i} \oint_{\Gamma} \mathbf{x}^\top (\mathbf{Y}_x - z\mathbf{I}_n)^{-1} \mathbf{x} dz \\ &= -\frac{1}{2\pi i} \oint_{\Gamma} \frac{\mathbf{x}^\top \mathbf{Q} \mathbf{x}}{1 + \ell \mathbf{x}^\top \mathbf{Q} \mathbf{x}} dz \end{aligned}$$

which holds for all large n with high probability over a positive contour Γ surrounding the limiting isolated eigenvalue λ_x (upon existence). By residue calculus, we find that this integral can only be non-vanishing if λ_x is a limiting root of $1 + \ell \mathbf{x}^\top \mathbf{Q}(z) \mathbf{x} = 0$, that is, if

$$\lambda_x = \sqrt{\ell^2 + 4}$$

with (using for instance l'Hôpital's rule to evaluate the residue)

$$|\hat{\mathbf{u}}_x^\top \mathbf{x}|^2 = -\frac{\mathbf{x}^\top \mathbf{Q}(\lambda_x) \mathbf{x}}{\ell \mathbf{x}^\top \mathbf{Q}'(\lambda_x) \mathbf{x}} + o(1) = \frac{\ell}{\sqrt{\ell^2 + 4}} + o(1)$$

as requested.

The proof is then concluded by remarking that, for $(\hat{\lambda}, \hat{\mathbf{u}})$ an “eigenvalue-eigenvector” pair of the tensor \mathcal{Y} (as defined in (9.18)),

$$\begin{aligned} \hat{\lambda} |\hat{\mathbf{u}}^\top \mathbf{x}| &= |\mathcal{Y} \cdot \hat{\mathbf{u}} \cdot \hat{\mathbf{u}} \cdot \mathbf{x}| \leq \max_{\|\mathbf{v}\|=1} |\mathcal{Y} \cdot \mathbf{v} \cdot \mathbf{v} \cdot \mathbf{x}| \\ &= \max_{\|\mathbf{v}\|=1} \mathbf{v}^\top \mathbf{Y}_x \mathbf{v} = \hat{\lambda}_x = \lambda_x + o(1) \end{aligned}$$

which, in the limit, provides the upper bound

$$\hat{\lambda} |\hat{\mathbf{u}}^\top \mathbf{x}| \leq \sqrt{\ell^2 + 4}.$$

Bibliography

- Radosław Adamczak. On the Marchenko-Pastur and Circular Laws for some Classes of Random Matrices with Dependent Entries. *Electronic Journal of Probability*, 16(0):1065–1095, 2011. ISSN 1083-6489. doi: 10.1214/ejp.v16-899.
- Lada A. Adamic and Natalie Glance. The political blogosphere and the 2004 U.S. election: divided they blog. In *Proceedings of the 3rd international workshop on Link discovery*, LinkKDD’05, pages 36–43. ACM, 2005. ISBN 9781595932151. doi: 10.1145/1134271.1134277.
- Ben Adlam and Jeffrey Pennington. The Neural Tangent Kernel in High Dimensions: Triple Descent and a Multi-Scale Theory of Generalization. In *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 74–84. PMLR, 2020. URL <http://proceedings.mlr.press/v119/adlam20a.html>.
- Ben Adlam, Jake Levinson, and Jeffrey Pennington. A Random Matrix Perspective on Mixtures of Nonlinearities for Deep Learning. *arXiv*, 2019. URL <https://arxiv.org/abs/1912.00827>.
- Madhu S. Advani, Andrew M. Saxe, and Haim Sompolinsky. High-dimensional dynamics of generalization error in neural networks. *Neural Networks*, 132: 428–446, 2020. ISSN 0893-6080. doi: 10.1016/j.neunet.2020.08.022.
- Oskari Ajanki, László Erdős, and Torben Krüger. Quadratic Vector Equations On Complex Upper Half-Plane. *Memoirs of the American Mathematical Society*, 261(1261):0–0, 2019. ISSN 0065-9266. doi: 10.1090/memo/1261.
- Naum Ilich Akhiezer and Izrail Markovich Glazman. *Theory of Linear Operators in Hilbert Space*. Dover Books on Mathematics. Dover Publications, 2013. ISBN 9780486677484. URL <http://cds.cern.ch/record/2009887>.
- Alnur Ali, J. Zico Kolter, and Ryan J. Tibshirani. A Continuous-Time View of Early Stopping for Least Squares Regression. In *Proceedings of Machine Learning Research*, volume 89 of *Proceedings of Machine Learning Research*, pages 1370–1378. PMLR, 2019. URL <http://proceedings.mlr.press/v89/ali19a.html>.

- Hafiz Tiomoko Ali and Romain Couillet. Improved spectral community detection in large heterogeneous networks. *Journal of Machine Learning Research*, 18(225):1–49, 2018. URL <http://jmlr.org/papers/v18/17-247.html>.
- Zeyuan Allen-Zhu, Yuanzhi Li, and Yingyu Liang. Learning and Generalization in Overparameterized Neural Networks, Going Beyond Two Layers. In *Advances in Neural Information Processing Systems*, volume 32 of *NIPS’19*, pages 6158–6169. Curran Associates, Inc., 2019. URL <https://proceedings.neurips.cc/paper/2019/file/62dad6e273d32235ae02b7d321578ee8-Paper.pdf>.
- Arash A. Amini, Aiyou Chen, Peter J. Bickel, and Elizaveta Levina. Pseudo-likelihood methods for community detection in large sparse networks. *The Annals of Statistics*, 41(4):2097–2122, 2013. ISSN 0090-5364. doi: 10.1214/13-aos1138.
- Greg W. Anderson, Alice Guionnet, and Ofer Zeitouni. *An Introduction to Random Matrices*, volume 118 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2010. ISBN 9780511801334. doi: 10.1017/cbo9780511801334.
- Theodore Wilbur Anderson et al. Asymptotic theory for principal component analysis. *Annals of Mathematical Statistics*, 34(1):122–148, 1963.
- Ralph G. Andrzejak, Klaus Lehnertz, Florian Mormann, Christoph Rieke, Peter David, and Christian E. Elger. Indications of nonlinear deterministic and finite-dimensional structures in time series of brain electrical activity: Dependence on recording region and brain state. *Physical Review E*, 64(6):061907, 2001. ISSN 1539-3755. doi: 10.1103/physreve.64.061907.
- Ludwig Arnold, Volker Matthias Gundlach, and Lloyd Demetrius. Evolutionary Formalism for Products of Positive Random Matrices. *The Annals of Applied Probability*, 4(3):859–901, 1994. ISSN 1050-5164. doi: 10.1214/aoap/1177004975.
- Sanjeev Arora, Rong Ge, Tengyu Ma, and Ankur Moitra. Simple, Efficient, and Neural Algorithms for Sparse Coding. In *Proceedings of The 28th Conference on Learning Theory*, volume 40 of *Proceedings of Machine Learning Research*, pages 113–149, Paris, France, 2015. PMLR. URL <http://proceedings.mlr.press/v40/Arora15.html>.
- Sanjeev Arora, Simon Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. Fine-Grained Analysis of Optimization and Generalization for Overparameterized Two-Layer Neural Networks. In *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 322–332. PMLR, 2019a. URL <http://proceedings.mlr.press/v97/arora19a.html>.

- Sanjeev Arora, Simon S. Du, Wei Hu, Zhiyuan Li, Russ R. Salakhutdinov, and Ruosong Wang. On Exact Computation with an Infinitely Wide Neural Net. In *Advances in Neural Information Processing Systems*, volume 32 of *NIPS'19*, pages 8141–8150. Curran Associates, Inc., 2019b. URL <https://proceedings.neurips.cc/paper/2019/file/dbc4d84bfcfe2284ba11beffb853a8c4-Paper.pdf>.
- Gérard Ben Arous and Sandrine Péché. Universality of local eigenvalue statistics for some sample covariance matrices. *Communications on Pure and Applied Mathematics*, 58(10):1316–1357, 2005. ISSN 1097-0312. doi: 10.1002/cpa.20070.
- Benson Au, Guillaume Cébron, Antoine Dahlqvist, Franck Gabriel, and Camille Male. Large permutation invariant random matrices are asymptotically free over the diagonal. *arXiv*, 2018. URL <https://arxiv.org/abs/1805.07045>.
- Nicolas Auguin, David Morales-Jimenez, Matthew R. McKay, and Romain Couillet. Large-dimensional Behavior of Regularized Maronna's M-estimators of Covariance Matrices. *IEEE Transactions on Signal Processing*, 66(13):3529–3542, 2018. ISSN 1053-587X. doi: 10.1109/tsp.2018.2831629.
- Konstantin Avrachenkov, Alexey Mishenin, Paulo Gonçalves, and Marina Sokol. Generalized Optimization Framework for Graph-based Semi-supervised Learning. In *Proceedings of the 2012 SIAM International Conference on Data Mining*, SDM'12, pages 966–974. SIAM, 2012. ISBN 9781611972320. doi: 10.1137/1.9781611972825.83.
- Zhidong Bai and Jack W. Silverstein. No eigenvalues outside the support of the limiting spectral distribution of large-dimensional sample covariance matrices. *The Annals of Probability*, 26(1):316–345, 1998. ISSN 0091-1798. doi: 10.1214/aop/1022855421.
- Zhidong Bai and Jack W. Silverstein. Exact Separation of Eigenvalues of Large Dimensional Sample Covariance Matrices. *The Annals of Probability*, 27(3):1536–1555, 1999. ISSN 0091-1798. doi: 10.1214/aop/1022677458.
- Zhidong Bai and Jack W. Silverstein. CLT for linear spectral statistics of large-dimensional sample covariance matrices. *The Annals of Probability*, 32(1A):553–605, 2004. ISSN 0091-1798. doi: 10.1214/aop/1078415845.
- Zhidong Bai and Jack W. Silverstein. *Spectral Analysis of Large Dimensional Random Matrices*, volume 20 of *Springer Series in Statistics*. Springer-Verlag New York, 2 edition, 2010. ISBN 9781441906601. doi: 10.1007/978-1-4419-0661-8.
- Zhidong Bai and Jian-feng Yao. Central limit theorems for eigenvalues in a spiked population model. *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, 44(3):447–474, 2008. ISSN 0246-0203. doi: 10.1214/07-aihp118.

- Zhidong Bai, Jack W. Silverstein, and Y. Q. Yin. A note on the largest eigenvalue of a large dimensional sample covariance matrix. *Journal of Multivariate Analysis*, 26(2):166–168, 1988. ISSN 0047-259X. doi: 10.1016/0047-259x(88)90078-4.
- Jinho Baik and Jack W. Silverstein. Eigenvalues of large sample covariance matrices of spiked population models. *Journal of Multivariate Analysis*, 97(6):1382–1408, 2006. ISSN 0047-259X. doi: 10.1016/j.jmva.2005.08.003.
- Jinho Baik, Gérard Ben Arous, and Sandrine Péché. Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. *The Annals of Probability*, 33(5):1643–1697, 2005. ISSN 0091-1798. doi: 10.1214/091179050000000233.
- Pierre Baldi, Peter Sadowski, and Zhiqin Lu. Learning in the machine: Random backpropagation and the deep learning channel. *Artificial Intelligence*, 260:1–35, 2018. ISSN 0004-3702. doi: 10.1016/j.artint.2018.03.003.
- Afonso S. Bandeira, Asad Lodhia, and Philippe Rigollet. Marchenko-Pastur law for Kendall’s tau. *Electronic Communications in Probability*, 22(0), 2017. ISSN 1083-589X. doi: 10.1214/17-ecp59.
- Richard G. Baraniuk. Compressive Sensing. *IEEE Signal Processing Magazine*, 24(4):118–121, 2007. ISSN 1053-5888. doi: 10.1109/msp.2007.4286571.
- Peter L. Bartlett, Dylan J. Foster, and Matus J. Telgarsky. Spectrally-normalized margin bounds for neural networks. In *Advances in Neural Information Processing Systems*, volume 30 of *NIPS’17*, pages 6240–6249. Curran Associates, Inc., 2017. URL <https://proceedings.neurips.cc/paper/2017/file/b22b257ad0519d4500539da3c8bcf4dd-Paper.pdf>.
- Heinz H Bauschke and Patrick L Combettes. *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. Number 2 in CMS Books in Mathematics. Springer International Publishing, 2 edition, 2017. ISBN 9783319483108. doi: 10.1007/978-3-319-48311-5.
- Amine Bejaoui, Khalil Elkhailil, Abla Kammoun, Mohamed Slim Alouini, and Tarek Al-Naffouri. Improved Design of Quadratic Discriminant Analysis Classifier in Unbalanced Settings. *arXiv*, 2020. URL <https://arxiv.org/abs/2006.06355>.
- Mikhail Belkin and Partha Niyogi. Semi-Supervised Learning on Riemannian Manifolds. *Machine Learning*, 56(1-3):209–239, 2004. ISSN 0885-6125. doi: 10.1023/b:mach.0000033120.25363.1e.
- Mikhail Belkin, Irina Matveeva, and Partha Niyogi. Regularization and Semi-supervised Learning on Large Graphs. In *International Conference on Computational Learning Theory (COLT)*, COLT’04, pages 624–638. Springer, 2004. doi: 10.1007/978-3-540-27819-1_43.

- Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine-learning practice and the classical bias–variance trade-off. *Proceedings of the National Academy of Sciences*, 116(32):15849–15854, 2019. ISSN 0027-8424. doi: 10.1073/pnas.1903070116.
- Florent Benaych-Georges and Romain Couillet. Spectral analysis of the Gram matrix of mixture models. *ESAIM: Probability and Statistics*, 20:217–237, 2016. ISSN 1292-8100. doi: 10.1051/ps/2016007.
- Florent Benaych-Georges and Raj Rao Nadakuditi. The eigenvalues and eigenvectors of finite, low rank perturbations of large random matrices. *Advances in Mathematics*, 227(1):494–521, 2011. ISSN 0001-8708. doi: 10.1016/j.aim.2011.02.007.
- Florent Benaych-Georges and Raj Rao Nadakuditi. The singular values and vectors of low rank perturbations of large rectangular random matrices. *Journal of Multivariate Analysis*, 111:120–135, 2012. ISSN 0047-259X. doi: 10.1016/j.jmva.2012.04.019.
- Lucas Benigni and Sandrine Péché. Eigenvalue distribution of nonlinear models of random matrices. *arXiv*, 2019. URL <https://arxiv.org/abs/1904.03090>.
- Pascal Bianchi, Mérourane Debbah, and Jamal Najim. Asymptotic Independence in the Spectrum of the Gaussian Unitary Ensemble. *Electronic Communications in Probability*, 15(0):376–395, 2010. ISSN 1083-589X. doi: 10.1214/ecp.v15-1568.
- Pascal Bianchi, Mérourane Debbah, Mylene Maida, and Jamal Najim. Performance of Statistical Tests for Single-Source Detection using Random Matrix Theory. *IEEE Transactions on Information Theory*, 57(4):2400–2419, 2011. ISSN 0018-9448. doi: 10.1109/tit.2011.2111710.
- Philippe Biane. Free probability for probabilists. *arXiv*, 1998. URL <https://arxiv.org/abs/math/9809193>.
- Alberto Bietti and Julien Mairal. On the Inductive Bias of Neural Tangent Kernels. In *Advances in Neural Information Processing Systems*, volume 32 of *NIPS’19*, pages 12893–12904. Curran Associates, Inc., 2019. URL <https://proceedings.neurips.cc/paper/2019/file/c4ef9c39b300931b69a36fb3dbb8d60e-Paper.pdf>.
- Patrick Billingsley. *Probability and Measure*. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd, 3 edition, 2012. ISBN 9781118122372. URL <https://www.wiley.com/en-us/Probability+and+Measure%2C+Anniversary+Edition-p-9781118122372>.
- Charles Bordenave. Eigenvalues of Euclidean random matrices. *Random Structures & Algorithms*, 33(4):515–532, 2008. ISSN 1098-2418. doi: 10.1002/rsa.20228.

- Charles Bordenave and Djamil Chafaï. Modern Aspects of Random Matrix Theory. *Proceedings of Symposia in Applied Mathematics*, pages 1–34, 2014. ISSN 0160-7634. doi: 10.1090/psapm/072/00617.
- Charles Bordenave and Marc Lelarge. Resolvent of large random graphs. *Random Structures & Algorithms*, 37(3):332–352, 2010. ISSN 1098-2418. doi: 10.1002/rsa.20313.
- Charles Bordenave, Marc Lelarge, and Justin Salez. The rank of diluted random graphs. *The Annals of Probability*, 39(3):1097–1121, 2011. ISSN 0091-1798. doi: 10.1214/10-aop567.
- Christian Borgs, Jennifer T Chayes, Henry Cohn, and Yufei Zhao. An $\$Lp\$$ theory of sparse graph convergence I: Limits, sparse random graph models, and power law distributions. *Transactions of the American Mathematical Society*, 372(5):3019–3062, 2019. ISSN 0002-9947. doi: 10.1090/tran/7543.
- Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013. ISBN 9780199535255. doi: 10.1093/acprof:oso/9780199535255.001.0001.
- Alan J. Bray and David S. Dean. Statistics of Critical Points of Gaussian Fields on Large-Dimensional Spaces. *Physical Review Letters*, 98(15):150201, 2007. ISSN 0031-9007. doi: 10.1103/physrevlett.98.150201.
- Andrew Brock, Jeff Donahue, and Karen Simonyan. Large Scale GAN Training for High Fidelity Natural Image Synthesis. In *International Conference on Learning Representations*, ICLR’19, 2019. URL <https://openreview.net/forum?id=B1xsqj09Fm>.
- Joël Bun, Jean-Philippe Bouchaud, and Marc Potters. Cleaning large correlation matrices: Tools from Random Matrix Theory. *Physics Reports*, 666:1–109, 2017. ISSN 0370-1573. doi: 10.1016/j.physrep.2016.10.005.
- Daniel M. Canaday. *Modeling and Control of Dynamical Systems with Reservoir Computing*. PhD thesis, 2019.
- Emmanuel J Candès and Terence Tao. Decoding by linear programming. *IEEE transactions on information theory*, 51(12):4203–4215, 2005.
- Emmanuel J. Candès. The restricted isometry property and its implications for compressed sensing. *Comptes Rendus Mathematique*, 346(9-10):589–592, 2008. ISSN 1631-073X. doi: 10.1016/j.crma.2008.03.014.
- Emmanuel J. Candès and Pragya Sur. The phase transition for the existence of the maximum likelihood estimate in high-dimensional logistic regression. *The Annals of Statistics*, 48(1):27–42, 2020. ISSN 0090-5364. doi: 10.1214/18-aos1789. URL <https://doi.org/10.1214/18-AOS1789>.

- Emmanuel J. Candès, Xiaodong Li, and Mahdi Soltanolkotabi. Phase Retrieval via Wirtinger Flow: Theory and Algorithms. *IEEE Transactions on Information Theory*, 61(4):1985–2007, 2015. ISSN 0018-9448. doi: 10.1109/tit.2015.2399924.
- Mireille Capitaine. Exact separation phenomenon for the eigenvalues of large Information-Plus-Noise type matrices. Application to spiked models. *Indiana University Mathematics Journal*, 63(6):1875–1910, 2014. ISSN 0022-2518. doi: 10.1512/iumj.2014.63.5432.
- Minmin Chen, Jeffrey Pennington, and Samuel Schoenholz. Dynamical Isometry and a Mean Field Theory of RNNs: Gating Enables Signal Propagation in Recurrent Neural Networks. In *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 873–882, Stockholm, Sweden, 2018. PMLR. URL <http://proceedings.mlr.press/v80/chen18i.html>.
- Yuxin Chen and Emmanuel J. Candès. Solving Random Quadratic Systems of Equations Is Nearly as Easy as Solving Linear Systems. *Communications on Pure and Applied Mathematics*, 70(5):822–883, 2017. ISSN 1097-0312. doi: 10.1002/cpa.21638.
- Xiuyuan Cheng and Amit Singer. The Spectrum of Random Inner-product Kernel Matrices. *Random Matrices: Theory and Applications*, 02(04):1350010, 2013. ISSN 2010-3263. doi: 10.1142/s201032631350010x.
- Marco Chiani. Distribution of the largest eigenvalue for real Wishart and Gaussian random matrices and a simple approximation for the Tracy–Widom distribution. *Journal of Multivariate Analysis*, 129:69–81, 2014. ISSN 0047-259X. doi: 10.1016/j.jmva.2014.04.002.
- Lénaïc Chizat, Edouard Oyallon, and Francis Bach. On Lazy Training in Differentiable Programming. In *Advances in Neural Information Processing Systems*, volume 32 of *NIPS’19*, pages 2937–2947. Curran Associates, Inc., 2019. URL <https://proceedings.neurips.cc/paper/2019/file/ae614c557843b1df326cb29c57225459-Paper.pdf>.
- Anna Choromanska, Mikaël Henaff, Michael Mathieu, Gerard Ben Arous, and Yann LeCun. The Loss Surfaces of Multilayer Networks. In *Proceedings of the Eighteenth International Conference on Artificial Intelligence and Statistics*, volume 38 of *Proceedings of Machine Learning Research*, pages 192–204, San Diego, California, USA, 2015. PMLR. URL <http://proceedings.mlr.press/v38/choromanska15.html>.
- Fan R. K. Chung. Spectral Graph Theory. *CBMS Regional Conference Series in Mathematics*, 1996. ISSN 0160-7642. doi: 10.1090/cbms/092.
- Tarin Clanuwat, Mikel Bober-Irizar, Asanobu Kitamoto, Alex Lamb, Kazuaki Yamamoto, and David Ha. Deep Learning for Classical Japanese Literature.

- arXiv*, 2018. doi: 10.20676/00000341. URL <https://arxiv.org/abs/1812.01718>.
- Amin Coja-Oghlan and André Lanka. Finding Planted Partitions in Random Graphs with General Degree Distributions. *SIAM Journal on Discrete Mathematics*, 23(4):1682–1714, 2010. ISSN 0895-4801. doi: 10.1137/070699354.
- Simon Coste and Yizhe Zhu. Eigenvalues of the non-backtracking operator detached from the bulk. *Random Matrices: Theory and Applications*, page 2150028, 2020. ISSN 2010-3263. doi: 10.1142/s2010326321500283.
- Romain Couillet. Robust spiked random matrices and a robust G-MUSIC estimator. *Journal of Multivariate Analysis*, 140:139–161, 2015. ISSN 0047-259X. doi: 10.1016/j.jmva.2015.05.009.
- Romain Couillet and Florent Benaych-Georges. Kernel spectral clustering of large dimensional data. *Electronic Journal of Statistics*, 10(1):1393–1454, 2016. ISSN 1935-7524. doi: 10.1214/16-ejs1144.
- Romain Couillet and Mérouane Debbah. *Random Matrix Methods for Wireless Communications*. Cambridge University Press, 2011. ISBN 9780511994746. doi: 10.1017/cbo9780511994746.
- Romain Couillet and Walid Hachem. Fluctuations of Spiked Random Matrix Models and Failure Diagnosis in Sensor Networks. *IEEE Transactions on Information Theory*, 59(1):509–525, 2013. ISSN 0018-9448. doi: 10.1109/tit.2012.2218572.
- Romain Couillet and Walid Hachem. Analysis of the limiting spectral measure of large random matrices of the separable covariance type. *Random Matrices: Theory and Applications*, 03(04):1450016, 2014. ISSN 2010-3263. doi: 10.1142/s2010326314500166.
- Romain Couillet and Abla Kammoun. Random Matrix Improved Subspace Clustering. In *2016 50th Asilomar Conference on Signals, Systems and Computers*, 2016 50th Asilomar Conference on Signals, Systems and Computers, pages 90–94. IEEE, 2016. ISBN 9781538639559. doi: 10.1109/acssc.2016.7869000.
- Romain Couillet and Matthew McKay. Large dimensional analysis and optimization of robust shrinkage covariance matrix estimators. *Journal of Multivariate Analysis*, 131:99–120, 2014. ISSN 0047-259X. doi: 10.1016/j.jmva.2014.06.018.
- Romain Couillet, Mérouane Debbah, and Jack W. Silverstein. A Deterministic Equivalent for the Analysis of Correlated MIMO Multiple Access Channels. *IEEE Transactions on Information Theory*, 57(6):3493–3514, 2011. ISSN 0018-9448. doi: 10.1109/tit.2011.2133151.

- Romain Couillet, Jakob Hoydis, and Mérouane Debbah. Random Beamforming over Quasi-Static and Fading Channels: A Deterministic Equivalent Approach. *IEEE Transactions on Information Theory*, 58(10):6392–6425, 2012. ISSN 0018-9448. doi: 10.1109/tit.2012.2201913.
- Romain Couillet, Frédéric Pascal, and Jack W. Silverstein. The random matrix regime of Maronna’s M-estimator with elliptically distributed samples. *Journal of Multivariate Analysis*, 139:56–78, 2015. ISSN 0047-259X. doi: 10.1016/j.jmva.2015.02.020.
- Romain Couillet, Abla Kammoun, and Frédéric Pascal. Second order statistics of robust estimators of scatter. Application to GLRT detection for elliptical signals. *Journal of Multivariate Analysis*, 143:249–274, 2016a. ISSN 0047-259X. doi: 10.1016/j.jmva.2015.08.021.
- Romain Couillet, Gilles Wainrib, Harry Sevi, and Hafiz Tiomoko Ali. The Asymptotic Performance of Linear Echo State Neural Networks. *Journal of Machine Learning Research*, 17(178):1–35, 2016b. URL <http://jmlr.org/papers/v17/16-076.html>.
- Romain Couillet, Malik Tiomoko, Steeve Zozor, and Eric Moisan. Random matrix-improved estimation of covariance matrix distances. *Journal of Multivariate Analysis*, 174:104531, 2019. ISSN 0047-259X. doi: 10.1016/j.jmva.2019.06.009.
- Romain Couillet, Yagmur Gizem Cinar, Eric Gaussier, and Muhammad Imran. Word Representations Concentrate and This is Good News! In *Proceedings of the 24th Conference on Computational Natural Language Learning*, CoNLL’20, pages 325–334. Association for Computational Linguistics, 2020. doi: 10.18653/v1/2020.conll-1.25. URL <https://www.aclweb.org/anthology/2020.conll-1.25>.
- Navneet Dalal and Bill Triggs. Histograms of Oriented Gradients for Human Detection. *2005 IEEE Computer Society Conference on Computer Vision and Pattern Recognition (CVPR’05)*, 1:886–893, 2005. doi: 10.1109/cvpr.2005.177.
- Lorenzo Dall’Amico, Romain Couillet, and Nicolas Tremblay. Revisiting the Bethe-Hessian: Improved Community Detection in Sparse Heterogeneous Graphs. In *Advances in Neural Information Processing Systems*, volume 32 of *NIPS’19*, pages 4037–4047. Curran Associates, Inc., 2019. URL <https://proceedings.neurips.cc/paper/2019/file/3e6260b81898beacda3d16db379ed329-Paper.pdf>.
- Yann N Dauphin, Razvan Pascanu, Caglar Gulcehre, Kyunghyun Cho, Surya Ganguli, and Yoshua Bengio. Identifying and attacking the saddle point problem in high-dimensional non-convex optimization. In *Advances in Neural Information Processing Systems*, volume 27 of *NIPS’14*, pages 2933–2941.

- Curran Associates, Inc., 2014. URL <https://proceedings.neurips.cc/paper/2014/file/17e23e50bedc63b4095e3d8204ce063b-Paper.pdf>.
- Chandler Davis. All convex invariant functions of hermitian matrices. *Archiv der Mathematik*, 8(4):276–278, 1957. ISSN 0003-889X. doi: 10.1007/bf01898787. URL <https://doi.org/10.1007/BF01898787>.
- Mérouane Debbah, Walid Hachem, Philippe Loubaton, and Marc De Courville. MMSE Analysis of Certain Large Isometric Random Precoded Systems. *IEEE Transactions on Information Theory*, 49(5):1293, 2003. ISSN 0018-9448. doi: 10.1109/tit.2003.810641.
- Aurelien Decelle, Florent Krzakala, Cristopher Moore, and Lenka Zdeborová. Inference and Phase Transitions in the Detection of Modules in Sparse Networks. *Physical Review Letters*, 107(6):065701, 2011. ISSN 0031-9007. doi: 10.1103/physrevlett.107.065701.
- Jia Deng, Wei Dong, Richard Socher, Li-Jia Li, Kai Li, and Li Fei-Fei. ImageNet: A large-scale hierarchical image database. *2009 IEEE Conference on Computer Vision and Pattern Recognition*, pages 248–255, 2009. ISSN 1063-6919. doi: 10.1109/cvpr.2009.5206848.
- Zeyu Deng, Abla Kammoun, and Christos Thrampoulidis. A Model of Double Descent for High-dimensional Binary Linear Classification. *arXiv*, 2019. URL <https://arxiv.org/abs/1911.05822>.
- Yen Do and Van Vu. The Spectrum of Random Kernel Matrices: Universality Results for Rough and Varying Kernels. *Random Matrices: Theory and Applications*, 02(03):1350005, 2013. ISSN 2010-3263. doi: 10.1142/s2010326313500056.
- Ivan Dokmanic, Reza Parhizkar, Juri Ranieri, and Martin Vetterli. Euclidean Distance Matrices: Essential theory, algorithms, and applications. *IEEE Signal Processing Magazine*, 32(6):12–30, 2015. ISSN 1053-5888. doi: 10.1109/msp.2015.2398954.
- Pedro Domingos. A Few Useful Things to Know about Machine Learning. *Communications of the ACM*, 55(10):78–87, 2012. ISSN 0001-0782. doi: 10.1145/2347736.2347755.
- David Donoho and Andrea Montanari. High dimensional robust M-estimation: asymptotic variance via approximate message passing. *Probability Theory and Related Fields*, 166(3-4):935–969, 2016. ISSN 0178-8051. doi: 10.1007/s00440-015-0675-z.
- David Donoho, Matan Gavish, and Iain M. Johnstone. Optimal shrinkage of eigenvalues in the spiked covariance model. *The Annals of Statistics*, 46(4):1742–1778, 2018. ISSN 0090-5364. doi: 10.1214/17-aos1601.

- David L. Donoho. Compressed Sensing. *IEEE Transactions on Information Theory*, 52(4):1289–1306, 2006. ISSN 0018-9448. doi: 10.1109/tit.2006.871582.
- R. Brent Dozier and Jack W. Silverstein. On the empirical distribution of eigenvalues of large dimensional information-plus-noise-type matrices. *Journal of Multivariate Analysis*, 98(4):678–694, 2007. ISSN 0047-259X. doi: 10.1016/j.jmva.2006.09.006.
- Simon Du, Jason Lee, Haochuan Li, Liwei Wang, and Xiyu Zhai. Gradient Descent Finds Global Minima of Deep Neural Networks. In *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 1675–1685. PMLR, 2019. URL <http://proceedings.mlr.press/v97/du19c.html>.
- Julien Dumont, Walid Hachem, Samson Lasaulce, Philippe Loubaton, and Jamal Najim. On the Capacity Achieving Covariance Matrix for Rician MIMO Channels: An Asymptotic Approach. *IEEE Transactions on Information Theory*, 56(3):1048–1069, 2010. ISSN 0018-9448. doi: 10.1109/tit.2009.2039063.
- Noureddine El Karoui. Spectrum estimation for large dimensional covariance matrices using random matrix theory. *The Annals of Statistics*, 36(6):2757–2790, 2008. ISSN 0090-5364. doi: 10.1214/07-aos581.
- Noureddine El Karoui. Concentration of measure and spectra of random matrices: Applications to correlation matrices, elliptical distributions and beyond. *The Annals of Applied Probability*, 19(6):2362–2405, 2009. ISSN 1050-5164. doi: 10.1214/08-aap548.
- Noureddine El Karoui, Derek Bean, Peter J. Bickel, Chinghway Lim, and Bin Yu. On robust regression with high-dimensional predictors. *Proceedings of the National Academy of Sciences*, 110(36):14557–14562, 2013. ISSN 0027-8424. doi: 10.1073/pnas.1307842110.
- Khalil Elkhailil, Abla Kammoun, Xiangliang Zhang, Mohamed-Slim Alouini, and Tareq Al-Naffouri. Risk Convergence of Centered Kernel Ridge Regression With Large Dimensional Data. *IEEE Transactions on Signal Processing*, 68: 1574–1588, 2019. ISSN 1053-587X. doi: 10.1109/tsp.2020.2975939.
- Khalil Elkhailil, Abla Kammoun, Romain Couillet, Tareq Y Al-Naffouri, and Mohamed-Slim Alouini. A Large Dimensional Study of Regularized Discriminant Analysis. *IEEE Transactions on Signal Processing*, 68:2464–2479, 2020. ISSN 1053-587X. doi: 10.1109/tsp.2020.2984160.
- Laszlo Erdős. Universality of Wigner random matrices: a survey of recent results. *Russian Mathematical Surveys*, 66(3):507–626, 2011. ISSN 0036-0279. doi: 10.1070/rm2011v06n03abeh004749.

- László Erdős, Sandrine Péché, José A. Ramírez, Benjamin Schlein, and Horng-Tzer Yau. Bulk universality for Wigner matrices. *Communications on Pure and Applied Mathematics*, 63(7):895–925, 2010. ISSN 1097-0312. doi: 10.1002/cpa.20317.
- Zhou Fan and Andrea Montanari. The spectral norm of random inner-product kernel matrices. *Probability Theory and Related Fields*, 173(1-2):27–85, 2019. ISSN 0178-8051. doi: 10.1007/s00440-018-0830-4.
- Zhou Fan and Zhichao Wang. Spectra of the Conjugate Kernel and Neural Tangent Kernel for linear-width neural networks. *arXiv*, 2020. URL <https://arxiv.org/abs/2005.11879>.
- James R. Fienup. Phase retrieval algorithms: a comparison. *Applied Optics*, 21(15):2758, 1982. ISSN 1539-4522. doi: 10.1364/ao.21.002758.
- Evelyn Fix and J. L. Hodges. Discriminatory Analysis. Nonparametric Discrimination: Consistency Properties. *International Statistical Review / Revue Internationale de Statistique*, 57(3):238–247, 1989. ISSN 0306-7734. doi: 10.2307/1403797. URL <http://www.jstor.org/stable/1403797>.
- Jonathan Frankle and Michael Carbin. The Lottery Ticket Hypothesis: Finding Sparse, Trainable Neural Networks. In *International Conference on Learning Representations*, ICLR’19, 2019. URL <https://openreview.net/forum?id=rJ1-b3RcF7>.
- Charlotte Frenkel, Martin Lefebvre, and David Bol. Learning without feedback: Direct random target projection as a feedback-alignment algorithm with layerwise feedforward training. *arXiv*, 2019. URL <https://arxiv.org/abs/1909.01311>.
- Jerome Friedman, Trevor Hastie, and Robert Tibshirani. *The Elements of Statistical Learning*, volume 1 of *Springer Series in Statistics*. Springer-Verlag New York, 1 edition, 2001. ISBN 9781489905192. doi: 10.1007/978-0-387-21606-5.
- Surya Ganguli, Dongsung Huh, and Haim Sompolinsky. Memory traces in dynamical systems. *Proceedings of the National Academy of Sciences*, 105(48):18970–18975, 2008. ISSN 0027-8424. doi: 10.1073/pnas.0804451105.
- Erol Gelenbe. Learning in the Recurrent Random Neural Network. *Neural Computation*, 5(1):154–164, 1993. ISSN 0899-7667. doi: 10.1162/neco.1993.5.1.154.
- Dar Gilboa, Bo Chang, Minmin Chen, Greg Yang, Samuel S Schoenholz, Ed H Chi, and Jeffrey Pennington. Dynamical Isometry and a Mean Field Theory of LSTMs and GRUs. *arXiv*, 2019. URL <https://arxiv.org/abs/1901.08987>.

- Vyacheslav L. Girko. *Theory of Stochastic Canonical Equations*, volume 535 of *Mathematics and Its Applications*. Springer Netherlands, 1 edition, 2001. ISBN 978-94-010-0989-8. doi: 10.1007/978-94-010-0989-8. URL <https://www.springer.com/cn/book/9781402000751>.
- Leon Glass and Michael C. Mackey. A simple model for phase locking of biological oscillators. *Journal of Mathematical Biology*, 7(4):339–352, 1979. ISSN 0303-6812. doi: 10.1007/bf00275153. URL <https://doi.org/10.1007/BF00275153>.
- Xavier Glorot and Yoshua Bengio. Understanding the difficulty of training deep feedforward neural networks. In *Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics*, volume 9 of *Proceedings of Machine Learning Research*, pages 249–256. JMLR Workshop and Conference Proceedings, 2010. URL <http://proceedings.mlr.press/v9/glorot10a.html>.
- Andrew Goldberg, Xiaojin Zhu, Aarti Singh, Zhiting Xu, and Robert Nowak. Multi-Manifold Semi-Supervised Learning. In *Proceedings of the Twelfth International Conference on Artificial Intelligence and Statistics*, volume 5 of *Proceedings of Machine Learning Research*, pages 169–176, Hilton Clearwater Beach Resort, Clearwater Beach, Florida USA, 2009. PMLR. URL <http://proceedings.mlr.press/v5/goldberg09a.html>.
- Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative Adversarial Nets. In *Advances in Neural Information Processing Systems*, volume 27 of *NIPS’14*, pages 2672–2680. Curran Associates, Inc., 2014. URL <https://proceedings.neurips.cc/paper/2014/file/5ca3e9b122f61f8f06494c97b1afccf3-Paper.pdf>.
- Yehoram Gordon. Some inequalities for Gaussian processes and applications. *Israel Journal of Mathematics*, 50(4):265–289, 1985. ISSN 0021-2172. doi: 10.1007/bf02759761.
- Robert M Gray. Toeplitz and Circulant Matrices: A Review. *Foundations and Trends® in Communications and Information Theory*, 2:155–239, 2006. ISSN 1567-2190. doi: 10.1561/0100000006. URL <http://dx.doi.org/10.1561/0100000006>.
- Lennart Gulikers, Marc Lelarge, and Laurent Massoulie. A spectral method for community detection in moderately sparse degree-corrected stochastic block models. *Advances in Applied Probability*, 49(3):686–721, 2017. ISSN 0001-8678. doi: 10.1017/apr.2017.18.
- Bernard Haasdonk. Feature Space Interpretation of SVMs with Indefinite Kernels. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 27(4):482–492, 2005. ISSN 0162-8828. doi: 10.1109/tpami.2005.78.

- Walid Hachem, Philippe Loubaton, and Jamal Najim. Deterministic equivalents for certain functionals of large random matrices. *The Annals of Applied Probability*, 17(3):875–930, 2007. ISSN 1050-5164. doi: 10.1214/105051606000000925.
- Walid Hachem, Philippe Loubaton, and Jamal Najim. A CLT for information-theoretic statistics of Gram random matrices with a given variance profile. *Annals of Applied Probability*, 18(6):2071–2130, 2008. ISSN 1050-5164. doi: 10.1214/08-aap515.
- Walid Hachem, Aris Moustakas, and Leonid A Pastur. The shannon’s mutual information of a multiple antenna time and frequency dependent channel: An ergodic operator approach. *Journal of Mathematical Physics*, 56(11):113501, 2015.
- James Douglas Hamilton. *Time Series Analysis*. Princeton University Press, Princeton, 1994. ISBN 978-0-691-21863-2. doi: 10.1515/9780691218632. URL <https://www.degruyter.com/princetonup/view/title/592052>.
- Donghyeon Han, Jinsu Lee, Jinmook Lee, and Hoi-Jun Yoo. A 1.32 TOPS/W Energy Efficient Deep Neural Network Learning Processor with Direct Feedback Alignment based Heterogeneous Core Architecture. In *2019 Symposium on VLSI Circuits*, volume 00 of *2019 Symposium on VLSI Circuits*, pages C304–C305, 2019. ISBN 9781728109145. doi: 10.23919/vlsic.2019.8778006.
- Trevor Hastie, Andrea Montanari, Saharon Rosset, and Ryan J Tibshirani. Surprises in High-Dimensional Ridgeless Least Squares Interpolation. *arXiv*, 2019. URL <https://arxiv.org/abs/1903.08560>.
- Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep Residual Learning for Image Recognition. In *2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR), pages 770–778. IEEE, 2016. ISBN 9781467388528. doi: 10.1109/cvpr.2016.90.
- Fumio Hiai and Dénes Petz. *The Semicircle Law, Free Random Variables and Entropy*. Mathematical Surveys and Monographs. American Mathematical Society, 2006. ISBN 9780821841358. doi: 10.1090/surv/077.
- Geoffrey E. Hinton and Sam Roweis. Stochastic Neighbor Embedding. In *Advances in Neural Information Processing Systems*, volume 15 of *NIPS’03*, pages 857–864. MIT Press, 2003. URL <https://proceedings.neurips.cc/paper/2002/file/6150ccc6069bea6b5716254057a194ef-Paper.pdf>.
- Sepp Hochreiter and Jurgen Schmidhuber. Long Short-Term Memory. *Neural Computation*, 9(8):1735–1780, 1997. ISSN 0899-7667. doi: 10.1162/neco.1997.9.8.1735.

- Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, 2 edition, 2012. ISBN 9780521548236. URL <http://www.cambridge.org/9780521548236>.
- Sebastian Houben, Johannes Stallkamp, Jan Salmen, Marc Schlipsing, and Christian Igel. Detection of Traffic Signs in Real-World Images: The German Traffic Sign Detection Benchmark. In *The 2013 International Joint Conference on Neural Networks (IJCNN)*, pages 1–8, 2013. doi: 10.1109/ijcnn.2013.6706807.
- Gao Huang, Zhuang Liu, Laurens van der Maaten, and Kilian Q. Weinberger. Densely Connected Convolutional Networks. In *2017 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2017 IEEE Conference on Computer Vision and Pattern Recognition (CVPR), pages 2261–2269. IEEE, 2017. ISBN 9781538604588. doi: 10.1109/cvpr.2017.243.
- Guang-Bin Huang, Hongming Zhou, Xiaojian Ding, and Rui Zhang. Extreme Learning Machine for Regression and Multiclass Classification. *IEEE Transactions on Systems, Man, and Cybernetics—Part B: Cybernetics*, 42(2):513–529, 2012. ISSN 1083-4419. doi: 10.1109/tsmcb.2011.2168604.
- Peter J. Huber. *Robust Statistics*. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd, 2011. ISBN 9780471725251. doi: 10.1002/0471725250.
- Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural Tangent Kernel: Convergence and Generalization in Neural Networks. In *Advances in Neural Information Processing Systems*, volume 31 of *NIPS’18*, pages 8571–8580. Curran Associates, Inc., 2018. URL <https://proceedings.neurips.cc/paper/2018/file/5a4be1fa34e62bb8a6ec6b91d2462f5a-Paper.pdf>.
- Herbert Jaeger. The “echo state” approach to analysing and training recurrent neural networks – with an Erratum note. Technical report, 2001. URL https://www.researchgate.net/profile/Herbert_Jaeger3/publication/215385037_The_echo_state_approach_to_analysing_and_training_recurrent_neural_networks-with_an_erratum_note/links/566a003508ae62b05f027be3/The-echo-state-approach-to-analysing-and-training-recurrent-neural-networks-with-an-erratum-note.pdf.
- Prateek Jain, Praneeth Netrapalli, and Sujay Sanghavi. Low-Rank Matrix Completion Using Alternating Minimization. In *Proceedings of the Forty-Fifth Annual ACM Symposium on Theory of Computing*, STOC ’13, page 665–674, New York, NY, USA, 2013. Association for Computing Machinery. ISBN 9781450320290. doi: 10.1145/2488608.2488693. URL <https://doi.org/10.1145/2488608.2488693>.
- Thorsten Joachims. Transductive Learning via Spectral Graph Partitioning. In *Proceedings of the Twentieth International Conference on Machine Learning*,

- volume 3 of *ICML'03*, pages 290–297. AAAI Press, 2003. URL <http://www.aaai.org/Library/ICML/2003/icml03-040.php>.
- Iain M. Johnstone. On the distribution of the largest eigenvalue in principal components analysis. *The Annals of Statistics*, 29(2):295–327, 2001. ISSN 0090-5364. doi: 10.1214/aos/1009210544.
- Iain M. Johnstone. Multivariate analysis and Jacobi ensembles: Largest eigenvalue, Tracy-Widom limits and rates of convergence. *The Annals of Statistics*, 36(6):2638–2716, 2008. ISSN 0090-5364. doi: 10.1214/08-aos605.
- Arun Kadavankandy and Romain Couillet. Asymptotic Gaussian Fluctuations of Spectral Clustering Eigenvectors. In *2019 IEEE 8th International Workshop on Computational Advances in Multi-Sensor Adaptive Processing (CAMSAP)*, volume 00 of *2019 IEEE 8th International Workshop on Computational Advances in Multi-Sensor Adaptive Processing (CAMSAP)*, pages 694–698. IEEE, 2019. ISBN 9781728155500. doi: 10.1109/camsap45676.2019.9022474.
- Abla Kammoun and Mohamed Alouini. On the Precise Error Analysis of Support Vector Machines. *IEEE Open Journal of Signal Processing*, pages 1–1, 2021. doi: 10.1109/ojsp.2021.3051849.
- Abla Kammoun and Romain Couillet. Subspace kernel spectral clustering of large dimensional data. *arxiv*, 2017. URL http://www.laneas.com/sites/default/files/attachments-186/paper_kernel.pdf.
- Abla Kammoun, Malika Kharouf, Walid Hachem, and Jamal Najim. A Central Limit Theorem for the SINR at the LMMSE Estimator Output for Large-Dimensional Signals. *IEEE Transactions on Information Theory*, 55(11):5048–5063, 2009. ISSN 0018-9448. doi: 10.1109/tit.2009.2030463.
- Abla Kammoun, Romain Couillet, Frédéric Pascal, and Mohamed-Slim Alouini. Optimal Design of the Adaptive Normalized Matched Filter Detector Using Regularized Tyler Estimators. *IEEE Transactions on Aerospace and Electronic Systems*, 54(2):755–769, 2017. ISSN 0018-9251. doi: 10.1109/taes.2017.2766538.
- Purushottam Kar and Harish Karnick. Random Feature Maps for Dot Product Kernels. In *Proceedings of the Fifteenth International Conference on Artificial Intelligence and Statistics*, volume 22 of *Proceedings of Machine Learning Research*, pages 583–591, La Palma, Canary Islands, 2012. PMLR. URL <http://proceedings.mlr.press/v22/kar12.html>.
- Brian Karrer and Mark E. J. Newman. Stochastic blockmodels and community structure in networks. *Physical Review E*, 83(1):016107, 2011. ISSN 1539-3755. doi: 10.1103/physreve.83.016107.
- Maurice G. Kendall. A New Measure of Rank Correlation. *Biometrika*, 30(1/2):81–93, 1938. ISSN 0006-3444. doi: 10.2307/2332226.

- Raghunandan H. Keshavan, Andrea Montanari, and Sewoong Oh. Matrix Completion from a Few Entries. *IEEE Transactions on Information Theory*, 56(6):2980–2998, 2010. ISSN 0018-9448. doi: 10.1109/tit.2010.2046205.
- Alexei M. Khorunzhy and Leonid A. Pastur. On the eigenvalue distribution of the deformed Wigner ensemble of random matrices. *Spectral Operator Theory and Related Topics*, pages 97–127, 1994. ISSN 1051-8037. doi: 10.1090/advsov/019/05.
- Alex Krizhevsky, Ilya Sutskever, and Geoffrey E. Hinton. ImageNet classification with deep convolutional neural networks. *Communications of the ACM*, 60(6):84–90, 2017. ISSN 0001-0782. doi: 10.1145/3065386.
- Florent Krzakala, Cristopher Moore, Elchanan Mossel, Joe Neeman, Allan Sly, Lenka Zdeborová, and Pan Zhang. Spectral redemption in clustering sparse networks. *Proceedings of the National Academy of Sciences*, 110(52):20935–20940, 2013. ISSN 0027-8424. doi: 10.1073/pnas.1312486110.
- Laurent Laloux, Pierre Cizeau, Marc Potters, and Jean-Philippe Bouchaud. Random matrix theory and financial correlations. *International Journal of Theoretical and Applied Finance*, 03(03):391–397, 2000. ISSN 0219-0249. doi: 10.1142/s0219024900000255.
- Yann LeCun, Leon Bottou, Yoshua Bengio, and Patrick Haffner. Gradient-based learning applied to document recognition. *Proceedings of the IEEE*, 86(11):2278–2324, 1998. ISSN 0018-9219. doi: 10.1109/5.726791.
- Olivier Ledoit and Sandrine Péché. Eigenvectors of some large sample covariance matrix ensembles. *Probability Theory and Related Fields*, 151(1-2):233–264, 2011. ISSN 0178-8051. doi: 10.1007/s00440-010-0298-3.
- Olivier Ledoit and Michael Wolf. Nonlinear shrinkage estimation of large-dimensional covariance matrices. *The Annals of Statistics*, 40(2):1024–1060, 2012. ISSN 0090-5364. doi: 10.1214/12-aos989.
- Michel Ledoux. *The Concentration of Measure Phenomenon*. Mathematical Surveys and Monographs. 2005. ISBN 9780821837924. doi: 10.1090/surv/089.
- Jaehoon Lee, Lechao Xiao, Samuel S Schoenholz, Yasaman Bahri, Roman Novak, Jascha Sohl-Dickstein, and Jeffrey Pennington. Wide neural networks of any depth evolve as linear models under gradient descent. *Journal of Statistical Mechanics: Theory and Experiment*, 2020(12):124002, 2020. doi: 10.1088/1742-5468/abc62b.
- Kiryung Lee, Yanjun Li, Marius Junge, and Yoram Bresler. Blind Recovery of Sparse Signals From Subsampled Convolution. *IEEE Transactions on Information Theory*, 63(2):802–821, 2017. ISSN 0018-9448. doi: 10.1109/tit.2016.2636204.

- Jure Leskovec and Andrej Krevl. SNAP Datasets: Stanford Large Network Dataset Collection, 2014. URL <http://snap.stanford.edu/data>.
- Jian Li and Petre Stoica. MIMO Radar with Colocated Antennas. *IEEE Signal Processing Magazine*, 24(5):106–114, 2007. ISSN 1053-5888. doi: 10.1109/msp.2007.904812.
- Ker-Chau Li. On Principal Hessian Directions for Data Visualization and Dimension Reduction: Another Application of Stein’s Lemma. *Journal of the American Statistical Association*, 87(420):1025–1039, 1992. ISSN 0162-1459. doi: 10.1080/01621459.1992.10476258.
- Zhenyu Liao and Romain Couillet. The Dynamics of Learning: A Random Matrix Approach. In *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 3072–3081, Stockholm, Sweden, 2018a. PMLR. URL <http://proceedings.mlr.press/v80/liao18b.html>.
- Zhenyu Liao and Romain Couillet. On the Spectrum of Random Features Maps of High Dimensional Data. In *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 3063–3071, Stockholm, Sweden, 2018b. PMLR. URL <http://proceedings.mlr.press/v80/liao18a.html>.
- Zhenyu Liao and Romain Couillet. Inner-product Kernels are Asymptotically Equivalent to Binary Discrete Kernels. *arXiv*, 2019a. URL <https://arxiv.org/abs/1909.06788>.
- Zhenyu Liao and Romain Couillet. A Large Dimensional Analysis of Least Squares Support Vector Machines. *IEEE Transactions on Signal Processing*, 67(4):1065–1074, 2019b. ISSN 1053-587X. doi: 10.1109/tsp.2018.2889954.
- Timothy P. Lillicrap, Daniel Cownden, Douglas B. Tweed, and Colin J. Akerman. Random synaptic feedback weights support error backpropagation for deep learning. *Nature Communications*, 7(1):13276, 2016. doi: 10.1038/ncomms13276.
- Lek-Heng Lim. Singular values and eigenvalues of tensors: a variational approach. In *1st IEEE International Workshop on Computational Advances in Multi-Sensor Adaptive Processing, 2005.*, 2005 IEEE International Workshop on Computational Advances in Multi-Sensor Adaptive Processing (CAMSAP), pages 129–132. IEEE, 2005. ISBN 9780780393228. doi: 10.1109/camap.2005.1574201.
- Zenan Ling and Robert C. Qiu. Spectrum Concentration in Deep Residual Learning: A Free Probability Approach. *IEEE Access*, 7:105212–105223, 2019. ISSN 2169-3536. doi: 10.1109/access.2019.2931991.

- Cosme Louart and Romain Couillet. Concentration of Measure and Large Random Matrices with an application to Sample Covariance Matrices. *arXiv*, 2018. URL <https://arxiv.org/pdf/1805.08295.pdf>.
- Cosme Louart, Zhenyu Liao, and Romain Couillet. A random matrix approach to neural networks. *Annals of Applied Probability*, 28(2):1190–1248, 2018. ISSN 1050-5164. doi: 10.1214/17-aap1328. URL <https://doi.org/10.1214/17-AAP1328>.
- Philippe Loubaton and Pascal Vallet. Almost Sure Localization of the Eigenvalues in a Gaussian Information Plus Noise Model. Application to the Spiked Models. *Electronic Journal of Probability*, 16(70):1934–1959, 2011. ISSN 1083-6489. doi: 10.1214/ejp.v16-943. URL <https://doi.org/10.1214/EJP.v16-943>.
- Daniel W. Lozier. NIST Digital Library of Mathematical Functions. *Annals of Mathematics and Artificial Intelligence*, 38(1-3):105–119, 2003. ISSN 1012-2443. doi: 10.1023/a:1022915830921.
- Lu Lu, Geoffrey Ye Li, A Lee Swindlehurst, Alexei Ashikhmin, and Rui Zhang. An Overview of Massive MIMO: Benefits and Challenges. *IEEE Journal of Selected Topics in Signal Processing*, 8(5):742–758, 2014. ISSN 1932-4553. doi: 10.1109/jstsp.2014.2317671.
- Yue M Lu and Gen Li. Phase transitions of spectral initialization for high-dimensional non-convex estimation. *Information and Inference: A Journal of the IMA*, 9(3):507–541, 2019. ISSN 2049-8772. doi: 10.1093/imaiai/iaz020.
- Ronny Luss and Alexandre D’aspremont. Support Vector Machine Classification with Indefinite Kernels. In *Advances in Neural Information Processing Systems*, volume 20 of *NIPS’08*, pages 953–960. Curran Associates, Inc., 2008. URL <https://proceedings.neurips.cc/paper/2007/file/c0c7c76d30bd3dcaefc96f40275bdc0a-Paper.pdf>.
- Ulrike Von Luxburg. A tutorial on spectral clustering. *Statistics and Computing*, 17(4):395–416, 2007. ISSN 0960-3174. doi: 10.1007/s11222-007-9033-z. URL <https://doi.org/10.1007/s11222-007-9033-z>.
- Ulrike von Luxburg, Mikhail Belkin, and Olivier Bousquet. Consistency of spectral clustering. *The Annals of Statistics*, 36(2):555–586, 2008. ISSN 0090-5364. doi: 10.1214/009053607000000640.
- Anna Lytova and Leonid Pastur. Central limit theorem for linear eigenvalue statistics of random matrices with independent entries. *The Annals of Probability*, 37(5):1778–1840, 2009. ISSN 0091-1798. doi: 10.1214/09-aop452.
- Zongming Ma. Accuracy of the Tracy-Widom limits for the extreme eigenvalues in white Wishart matrices. *Bernoulli*, 18(1):322–359, 2012. ISSN 1350-7265. doi: 10.3150/10-bej334. URL <https://doi.org/10.3150/10-BEJ334>.

- Andrew L. Maas, Awni Y. Hannun, and Andrew Y. Ng. Rectifier Nonlinearities Improve Neural Network Acoustic Models. In *ICML Workshop on Deep Learning for Audio, Speech and Language Processing*, ICML Workshop, page 3, 2013.
- Laurens van der Maaten and Geoffrey Hinton. Visualizing Data using t-SNE. *Journal of Machine Learning Research*, 9:2579–2605, 2008. URL <http://jmlr.org/papers/v9/vandermaaten08a.html>.
- Xiaoyi Mai. *Methods of random matrices for large dimensional statistical learning*. PhD thesis, 2019.
- Xiaoyi Mai and Romain Couillet. A Random Matrix Analysis and Improvement of Semi-Supervised Learning for Large Dimensional Data. *Journal of Machine Learning Research*, 19(79):1–27, 2018. URL <http://jmlr.org/papers/v19/17-421.html>.
- Xiaoyi Mai and Romain Couillet. Consistent Semi-Supervised Graph Regularization for High Dimensional Data. *arXiv*, 2020. URL <https://arxiv.org/abs/2006.07575>.
- Xiaoyi Mai and Zhenyu Liao. High Dimensional Classification via Regularized and Unregularized Empirical Risk Minimization: Precise Error and Optimal Loss. *arXiv*, 2019. URL <https://arxiv.org/abs/1905.13742>.
- Christopher D Manning, Hinrich Schütze, and Prabhakar Raghavan. *Introduction to Information Retrieval*. Cambridge University Press, 2008. ISBN 9780511809071. doi: 10.1017/cbo9780511809071.
- Vladimir A Marenko and Leonid Andreevich Pastur. Distribution of eigenvalues for some sets of random matrices. *Mathematics of the USSR-Sbornik*, 1(4):457, 1967. ISSN 0025-5734. doi: 10.1070/sm1967v001n04abeh001994.
- Ricardo A Maronna, R Douglas Martin, Victor J Yohai, and Matias Salibian-Barrera. *Robust Statistics: Theory and Methods (with R)*. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd, 2 edition, 2018. ISBN 9781119214656. doi: 10.1002/9781119214656.
- Ricardo Antonio Maronna. Robust M-Estimators of Multivariate Location and Scatter. *The Annals of Statistics*, 4(1):51–67, 1976. ISSN 0090-5364. doi: 10.1214/aos/1176343347.
- Laurent Massoulié. Community Detection Thresholds and the Weak Ramanujan Property. In *Proceedings of the Forty-Sixth Annual ACM Symposium on Theory of Computing*, STOC’14, page 694–703, New York, NY, USA, 2014. Association for Computing Machinery. ISBN 9781450327107. doi: 10.1145/2591796.2591857. URL <https://doi.org/10.1145/2591796.2591857>.

- Madan Lal Mehta and Michel Gaudin. On the density of Eigenvalues of a random matrix. *Nuclear Physics*, 18:420–427, 1960. ISSN 0029-5582. doi: 10.1016/0029-5582(60)90414-4.
- Song Mei and Andrea Montanari. The generalization error of random features regression: Precise asymptotics and double descent curve. *arXiv*, 2019. URL <https://arxiv.org/abs/1908.05355>.
- Xavier Mestre. Improved Estimation of Eigenvalues and Eigenvectors of Covariance Matrices Using Their Sample Estimates. *IEEE Transactions on Information Theory*, 54(11):5113–5129, 2008. ISSN 0018-9448. doi: 10.1109/tit.2008.929938.
- Xavier Mestre and Miguel Ángel Lagunas. Modified Subspace Algorithms for DoA Estimation with Large Arrays. *IEEE Transactions on Signal Processing*, 56(2):598–614, 2008. ISSN 1053-587X. doi: 10.1109/tsp.2007.907884.
- Sebastian Mika, Gunnar Ratsch, Jason Weston, Bernhard Scholkopf, and Klaus-Robert Mullers. Fisher discriminant analysis with kernels. In *Neural Networks for Signal Processing IX: Proceedings of the 1999 IEEE Signal Processing Society Workshop*, Neural Networks for Signal Processing IX: Proceedings of the 1999 IEEE Signal Processing Society Workshop (Cat. No.98TH8468), pages 41–48, 1999. ISBN 978078035673X. doi: 10.1109/nns.1999.788121.
- Tomas Mikolov, Kai Chen, Greg Corrado, and Jeffrey Dean. Efficient Estimation of Word Representations in Vector Space. *arXiv*, 2013. URL <https://arxiv.org/abs/1301.3781>.
- James A Mingo and Roland Speicher. *Free Probability and Random Matrices*, volume 35 of *Fields Institute Monographs*. Springer-Verlag New York, 1 edition, 2017. ISBN 9781493969418. doi: 10.1007/978-1-4939-6942-5.
- Mehdi Mirza and Simon Osindero. Conditional Generative Adversarial Nets. *arXiv*, 2014. URL <https://arxiv.org/abs/1411.1784>.
- Takeru Miyato, Toshiki Kataoka, Masanori Koyama, and Yuichi Yoshida. Spectral Normalization for Generative Adversarial Networks. In *International Conference on Learning Representations*, ICLR’18, 2018. URL <https://openreview.net/forum?id=B1QRgziT->.
- Marco Mondelli and Andrea Montanari. Fundamental Limits of Weak Recovery with Applications to Phase Retrieval. *Foundations of Computational Mathematics*, 19(3):703–773, 2019. ISSN 1615-3375. doi: 10.1007/s10208-018-9395-y.
- David Morales-Jimenez, Romain Couillet, and Matthew R. McKay. Large Dimensional Analysis of Robust M-Estimators of Covariance with Outliers. *IEEE Transactions on Signal Processing*, 63(21):5784–5797, 2015. ISSN 1053-587X. doi: 10.1109/tsp.2015.2460225.

- Amit Moscovich, Ariel Jaffe, and Nadler Boaz. Minimax-optimal semi-supervised regression on unknown manifolds. In *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics*, volume 54 of *Proceedings of Machine Learning Research*, pages 933–942. PMLR, 2016. URL <http://proceedings.mlr.press/v54/moscovich17a.html>.
- Elchanan Mossel, Joe Neeman, and Allan Sly. Reconstruction and estimation in the planted partition model. *Probability Theory and Related Fields*, 162(3-4):431–461, 2015. ISSN 0178-8051. doi: 10.1007/s00440-014-0576-6.
- Marc Mézard and Andrea Montanari. *Information, Physics, and Computation*. Oxford University Press, 2009. ISBN 9780198570837. doi: 10.1093/acprof:oso/9780198570837.001.0001.
- Jamal Najim and Jianfeng Yao. Gaussian fluctuations for linear spectral statistics of large random covariance matrices. *The Annals of Applied Probability*, 26(3):1837–1887, 2016. ISSN 1050-5164. doi: 10.1214/15-aap1135.
- Preetum Nakkiran, Gal Kaplun, Yamini Bansal, Tristan Yang, Boaz Barak, and Ilya Sutskever. Deep double descent: Where bigger models and more data hurt. In *International Conference on Learning Representations*, ICLR’19, 2020. URL <https://openreview.net/forum?id=B1g5sA4twr>.
- John Ashworth Nelder and R. W. M. Wedderburn. Generalized Linear Models. *Journal of the Royal Statistical Society: Series A (General)*, 135(3):370–384, 1972. ISSN 0035-9238. doi: 10.2307/2344614.
- Mark E. J. Newman. Modularity and community structure in networks. *Proceedings of the National Academy of Sciences*, 103(23):8577–8582, 2006. ISSN 0027-8424. doi: 10.1073/pnas.0601602103.
- Andrew Ng, Michael I. Jordan, and Yair Weiss. On spectral clustering: Analysis and an algorithm. In *Advances in Neural Information Processing Systems*, volume 14 of *NIPS’02*, pages 849–856. MIT Press, 2002. URL <https://proceedings.neurips.cc/paper/2001/file/801272ee79cfde7fa5960571fee36b9b-Paper.pdf>.
- Alexandru Nica and Roland Speicher. *Lectures on the Combinatorics of Free Probability*, volume 13 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, 2006. ISBN 9780511735127. doi: 10.1017/cbo9780511735127.
- Roman Novak, Lechao Xiao, Yasaman Bahri, Jaehoon Lee, Greg Yang, Daniel A. Abolafia, Jeffrey Pennington, and Jascha Sohl-dickstein. Bayesian Deep Convolutional Networks with Many Channels are Gaussian Processes. In *International Conference on Learning Representations*, ICLR’19, 2019. URL <https://openreview.net/forum?id=B1g30j0qF7>.

- Arild Nøkland. Direct Feedback Alignment Provides Learning in Deep Neural Networks. In *Advances in Neural Information Processing Systems*, volume 29 of *NIPS'16*, pages 1037–1045. Curran Associates, Inc., 2016. URL <https://proceedings.neurips.cc/paper/2016/file/d490d7b4576290fa60eb31b5fc917ad1-Paper.pdf>. backprop-less NN.
- Chapelle Olivier, Schölkopf Bernhard, and Zien Alexander. *Semi-Supervised Learning*. The MIT Press, 2006. ISBN 9780262033589. doi: 10.7551/mitpress/9780262033589.001.0001.
- Sean O'Rourke. A note on the Marchenko-Pastur law for a class of random matrices with dependent entries. *Electronic Communications in Probability*, 17(0):13, 2012. ISSN 1083-589X. doi: 10.1214/ecp.v17-2020.
- Alain Pajor and Lenoid Pastur. On the limiting empirical measure of eigenvalues of the sum of rank one matrices with log-concave distribution. *Studia Mathematica*, 195(1):11–29, 2009. ISSN 0039-3223. doi: 10.4064/sm195-1-2.
- Anastasios K. Papazafeiropoulos and Tharmalingam Ratnarajah. Deterministic Equivalent Performance Analysis of Time-Varying Massive MIMO Systems. *IEEE Transactions on Wireless Communications*, 14(10):5795–5809, 2015. ISSN 1536-1276. doi: 10.1109/twc.2015.2443040.
- Leonid Pastur. On Random Matrices Arising in Deep Neural Networks. Gaussian Case. *arXiv*, 2020. URL <https://arxiv.org/abs/2001.06188>.
- Leonid Pastur and Alexander Figotin. Spectra of random and almost-periodic operators. 1992.
- Leonid Pastur and Victor Slavin. On Random Matrices Arising in Deep Neural Networks: General I.I.D. Case. *arXiv*, 2020. URL <https://arxiv.org/abs/2011.11439>.
- Leonid A Pastur. A Simple Approach to the Global Regime of Gaussian Ensembles of Random Matrices. *Ukrainian Mathematical Journal*, 57(6):936–966, 2005. ISSN 0041-5995. doi: 10.1007/s11253-005-0241-4. URL <https://doi.org/10.1007/s11253-005-0241-4>.
- Leonid Andreevich Pastur and Mariya Shcherbina. *Eigenvalue Distribution of Large Random Matrices*, volume 171 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2011. doi: 10.1090/surv/1.
- Debashis Paul. Asymptotics of Sample Eigenstructure for a Large Dimensional Spiked Covariance Model. *Statistica Sinica*, 17(4):1617–1642, 2007. URL <http://www.jstor.org/stable/24307692>.
- Debashis Paul and Jack W. Silverstein. No eigenvalues outside the support of the limiting empirical spectral distribution of a separable covariance matrix. *Journal of Multivariate Analysis*, 100(1):37–57, 2009. ISSN 0047-259X. doi: 10.1016/j.jmva.2008.03.010.

- Judea Pearl. Fusion, propagation, and structuring in belief networks. *Artificial Intelligence*, 29(3):241–288, 1986. ISSN 0004-3702. doi: 10.1016/0004-3702(86)90072-x.
- Jeffrey Pennington and Yasaman Bahri. Geometry of Neural Network Loss Surfaces via Random Matrix Theory. In *Proceedings of the 34th International Conference on Machine Learning*, volume 70 of *Proceedings of Machine Learning Research*, pages 2798–2806, International Convention Centre, Sydney, Australia, 2017. PMLR. URL <http://proceedings.mlr.press/v70/pennington17a.html>.
- Jeffrey Pennington and Pratik Worah. Nonlinear random matrix theory for deep learning. In *Advances in Neural Information Processing Systems*, volume 30 of *NIPS’17*, pages 2637–2646. Curran Associates, Inc., 2017. URL <https://proceedings.neurips.cc/paper/2017/file/0f3d014eead934bbdbacb62a01dc4831-Paper.pdf>.
- Jeffrey Pennington, Samuel Schoenholz, and Surya Ganguli. Resurrecting the sigmoid in deep learning through dynamical isometry: theory and practice. In *Advances in Neural Information Processing Systems*, volume 30 of *NIPS’17*, pages 4785–4795. Curran Associates, Inc., 2017. URL <https://proceedings.neurips.cc/paper/2017/file/d9fc0cdb67638d50f411432d0d41d0ba-Paper.pdf>.
- Vinay Uday Prabhu. Kannada-MNIST: A new handwritten digits dataset for the Kannada language. *arXiv*, 2019. URL <https://arxiv.org/abs/1908.01242>.
- Tai Qin and Karl Rohe. Regularized Spectral Clustering under the Degree-Corrected Stochastic Blockmodel. In *Advances in Neural Information Processing Systems*, volume 26 of *NIPS’13*, pages 3120–3128. Curran Associates, Inc., 2013. URL <https://proceedings.neurips.cc/paper/2013/file/0ed9422357395a0d4879191c66f4faa2-Paper.pdf>.
- Ali Rahimi and Benjamin Recht. Random Features for Large-Scale Kernel Machines. In *Advances in Neural Information Processing Systems*, volume 20 of *NIPS’08*, pages 1177–1184. Curran Associates, Inc., 2008. URL <https://proceedings.neurips.cc/paper/2007/file/013a006f03dbc5392effeb8f18fda755-Paper.pdf>.
- Lorenzo Rosasco, Ernesto De Vito, Andrea Caponnetto, Michele Piana, and Alessandro Verri. Are Loss Functions All the Same? *Neural Computation*, 16(5):1063–1076, 2004. ISSN 0899-7667. doi: 10.1162/089976604773135104. URL <https://doi.org/10.1162/089976604773135104>.
- Frank Rosenblatt. The perceptron: A probabilistic model for information storage and organization in the brain. *Psychological Review*, 65(6):386–408, 1958. ISSN 0033-295X. doi: 10.1037/h0042519.

- Yu. A. Rozanov. *Stationary Random Processes*. Holden-Day series in time series analysis. Holden-Day, San Francisco, 1967. URL <https://openlibrary.org/books/OL21849368M/>.
- Walter Rudin. *Principles of Mathematical Analysis*, volume 3 of *International Series in Pure and Applied Mathematics*. McGraw-Hill Education, 3 edition, 1964. ISBN 9780070542358. URL <https://www.mheducation.com/highered/product/principles-mathematical-analysis-rudin/M9780070542358.html>.
- Alaa Saade, Florent Krzakala, and Lenka Zdeborová. Spectral Clustering of graphs with the Bethe Hessian. In *Advances in Neural Information Processing Systems*, volume 27 of *NIPS'14*, pages 406–414. Curran Associates, Inc., 2014. URL <https://proceedings.neurips.cc/paper/2014/file/63923f49e5241343aa7acb6a06a751e7-Paper.pdf>.
- Justin Salez. *Some implications of local weak convergence for sparse random graphs*. PhD thesis, 2011.
- Justin Salez. Spectral atoms of unimodular random trees. *Journal of the European Mathematical Society*, 22(2):345–363, 2019. ISSN 1435-9855. doi: 10.4171/jems/923.
- Simone Scardapane and Dianhui Wang. Randomness in neural networks: an overview. *Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery*, 7(2):e1200, 2017. ISSN 1942-4787. doi: 10.1002/widm.1200.
- Robert E. Schapire. A brief introduction to boosting. In *Proceedings of the Sixteenth International Joint Conference on Artificial Intelligence, IJCAI-99*, volume 99 of *IJCAI'99*, pages 1401–1406. International Joint Conferences on Artificial Intelligence Organization, 1999. URL <https://www.ijcai.org/Proceedings/99-2/Papers/103.pdf>.
- Ralph Schmidt. Multiple emitter location and signal parameter estimation. *IEEE Transactions on Antennas and Propagation*, 34(3):276–280, 1986. ISSN 0018-926X. doi: 10.1109/tap.1986.1143830.
- Wouter Schmidt, Martin Kraaijveld, and Robert Duin. Feedforward neural networks with random weights. In *11th IAPR International Conference on Pattern Recognition*, volume 1 of *ICPR*, pages 1–4. IEEE, 1992. doi: 10.1109/icpr.1992.201708. URL <https://doi.ieee.org/10.1109/ICPR.1992.201708>.
- Bernhard Schölkopf and Alexander J. Smola. *Learning with Kernels: Support Vector Machines, Regularization, Optimization, and Beyond*. The MIT Press, 2018. ISBN 9780262256933. doi: 10.7551/mitpress/4175.001.0001.
- Mohamed El Amine Seddik, Mohamed Tamaazousti, and Romain Couillet. Kernel Random Matrices of Large Concentrated Data: the Example of GAN-Generated Images. In *2019 IEEE International Conference on Acoustics,*

Speech and Signal Processing (ICASSP 2019), 2019 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP 2019), pages 7480–7484. IEEE, 2019. ISBN 9781479981328. doi: 10.1109/icassp.2019.8683333.

Mohamed El Amine Seddik, Cosme Louart, Mohamed Tamaazousti, and Romain Couillet. Random Matrix Theory Proves that Deep Learning Representations of GAN-data Behave as Gaussian Mixtures. In *Proceedings of the 37th International Conference on Machine Learning*, Proceedings of Machine Learning Research, pages 8573–8582. PMLR, 2020. URL <http://proceedings.mlr.press/v119/seddik20a.html>.

Jack W. Silverstein. The Limiting Eigenvalue Distribution of a Multivariate F Matrix. *SIAM Journal on Mathematical Analysis*, 16(3):641–646, 1985. ISSN 0036-1410. doi: 10.1137/0516047.

Jack W. Silverstein and Zhidong Bai. On the Empirical Distribution of Eigenvalues of a Class of Large Dimensional Random Matrices. *Journal of Multivariate Analysis*, 54(2):175–192, 1995. ISSN 0047-259X. doi: 10.1006/jmva.1995.1051.

Jack W. Silverstein and Sang-Il Choi. Analysis of the Limiting Spectral Distribution of Large Dimensional Random Matrices. *Journal of Multivariate Analysis*, 54(2):295–309, 1995. ISSN 0047-259X. doi: 10.1006/jmva.1995.1058.

Karen Simonyan and Andrew Zisserman. Very deep convolutional networks for large-scale image recognition. In *International Conference on Learning Representations*, ICLR’14, 2014. URL <http://arxiv.org/abs/1409.1556>.

Alexander Soshnikov. Universality at the Edge of the Spectrum in Wigner Random Matrices. *Communications in Mathematical Physics*, 207(3):697–733, 1999. ISSN 0010-3616. doi: 10.1007/s002200050743.

Alexander B. Soshnikov. Gaussian Fluctuation for the Number of Particles in Airy, Bessel, Sine, and Other Determinantal Random Point Fields. *Journal of Statistical Physics*, 100(3-4):491–522, 2000. ISSN 0022-4715. doi: 10.1023/a:1018672622921.

Daniel Soudry, Elad Hoffer, Mor Shpigel Nacson, Suriya Gunasekar, and Nathan Srebro. The Implicit Bias of Gradient Descent on Separable Data. *Journal of Machine Learning Research*, 19(70):1–57, 2018. URL <http://jmlr.org/papers/v19/18-188.html>.

Charles Spearman. The Proof and Measurement of Association between Two Things. *The American Journal of Psychology*, 100(3/4):441, 1987. ISSN 0002-9556. doi: 10.2307/1422689.

Roland Speicher and Carlos Vargas. Free deterministic equivalents, rectangular random matrix models, and operator-valued free probability theory. *Random*

- Matrices: Theory and Applications*, 01(02):1150008, 2012. ISSN 2010-3263. doi: 10.1142/s2010326311500080.
- Nitish Srivastava, Geoffrey Hinton, Alex Krizhevsky, Ilya Sutskever, and Ruslan Salakhutdinov. Dropout: A Simple Way to Prevent Neural Networks from Overfitting. *Journal of Machine Learning Research*, 15(56):1929–1958, 2014. URL <http://jmlr.org/papers/v15/srivastava14a.html>.
- Charles M. Stein. Estimation of the Mean of a Multivariate Normal Distribution. *The Annals of Statistics*, 9(6):1135–1151, 1981. ISSN 0090-5364. doi: 10.1214/aos/1176345632.
- Elias M. Stein and Rami Shakarchi. *Complex Analysis*, volume 2 of *Princeton lectures in analysis*. Princeton University Press, 2003. ISBN 9780691113852.
- Arun Suggala, Adarsh Prasad, and Pradeep K Ravikumar. Connecting Optimization and Regularization Paths. In *Advances in Neural Information Processing Systems*, volume 31 of *NIPS’18*, pages 10608–10619. Curran Associates, Inc., 2018. URL <https://proceedings.neurips.cc/paper/2018/file/6459257ddab7b85bf4b57845e875e4d4-Paper.pdf>.
- Pragya Sur and Emmanuel J Candes. A modern maximum-likelihood theory for high-dimensional logistic regression. *Proceedings of the National Academy of Sciences*, 116(29):14516–14525, 2019. URL <https://www.pnas.org/content/116/29/14516>.
- Johan A.K. Suykens and Joos Vandewalle. Least Squares Support Vector Machine Classifiers. *Neural Processing Letters*, 9(3):293–300, 1999. ISSN 1370-4621. doi: 10.1023/a:1018628609742.
- Martin Szummer and Tommi Jaakkola. Partially labeled classification with Markov random walks. In *Advances in Neural Information Processing Systems*, volume 14 of *NIPS’02*, pages 945–952. MIT Press, 2002. URL <https://proceedings.neurips.cc/paper/2001/file/a82d922b133be19c1171534e6594f754-Paper.pdf>.
- Hossein Taheri, Ramtin Pedarsani, and Christos Thrampoulidis. Sharp Guarantees for Solving Random Equations with One-Bit Information. In *57th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, volume 00 of *The 57th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 765–772, 2019. ISBN 9781728131528. doi: 10.1109/allerton.2019.8919905.
- Hossein Taheri, Ramtin Pedarsani, and Christos Thrampoulidis. Fundamental Limits of Ridge-Regularized Empirical Risk Minimization in High Dimensions. *arXiv*, 2020a. URL <https://arxiv.org/abs/2006.08917>.
- Hossein Taheri, Ramtin Pedarsani, and Christos Thrampoulidis. Optimality of Least-squares for Classification in Gaussian-Mixture Models. *2020 IEEE International Symposium on Information Theory (ISIT)*, 00:2515–2520, 2020b. doi: 10.1109/isit44484.2020.9174267.

- Hossein Taheri, Ramtin Pedarsani, and Christos Thrampoulidis. Sharp Asymptotics and Optimal Performance for Inference in Binary Models. *arXiv*, 2020c. URL <https://arxiv.org/abs/2002.07284>.
- Michel Talagrand. Concentration of measure and isoperimetric inequalities in product spaces. *Publications Mathématiques de l'Institut des Hautes Études Scientifiques*, 81(1):73–205, 1995. ISSN 0073-8301. doi: 10.1007/bf02699376.
- Terence Tao. *Topics in Random Matrix Theory*, volume 132 of *Graduate Studies in Mathematics*. 2012. ISBN 9780821874301. doi: 10.1090/gsm/132. URL <http://www.ams.org/books/gsm/132/>.
- Terence Tao and Van Vu. Random matrices: the circular law. *Communications in Contemporary Mathematics*, 10(02):261–307, 2008. ISSN 0219-1997. doi: 10.1142/s0219199708002788.
- Christos Thrampoulidis, Ehsan Abbasi, and Babak Hassibi. Precise Error Analysis of Regularized M-Estimators in High Dimensions. *IEEE Transactions on Information Theory*, 64(8):5592–5628, 2018. ISSN 0018-9448. doi: 10.1109/tit.2018.2840720.
- Malik Tiomoko and Romain Couillet. Random Matrix-Improved Estimation of the Wasserstein Distance between two Centered Gaussian Distributions. In *2019 27th European Signal Processing Conference (EUSIPCO)*, volume 00 of *2019 27th European Signal Processing Conference (EUSIPCO)*, pages 1–5, 2019a. ISBN 9781538673003. doi: 10.23919/eusipco.2019.8902795.
- Malik Tiomoko and Romain Couillet. Estimation of covariance matrix distances in the high dimension low sample size regime. In *2019 IEEE 8th International Workshop on Computational Advances in Multi-Sensor Adaptive Processing (CAMSAP)*, pages 341–345. IEEE, 2019b.
- Malik Tiomoko, Hafiz Tiomoko Ali, and Romain Couillet. Deciphering and Optimizing Multi-Task Learning: a Random Matrix Approach. In *International Conference on Learning Representations*, ICLR’21, 2021. URL <https://openreview.net/forum?id=Cri3xz59ga>.
- Hafiz Tiomoko Ali, Abla Kammoun, and Romain Couillet. Random Matrix-Improved Kernels For Large Dimensional Spectral Clustering. In *2018 IEEE Statistical Signal Processing Workshop (SSP)*, 2018 IEEE Statistical Signal Processing Workshop (SSP), pages 453–457. IEEE, 2018. ISBN 9781538615720. doi: 10.1109/ssp.2018.8450705.
- E. C. Titchmarsh. *The Theory of Functions*. Oxford University Press, New York, NY, USA, 1939.
- Craig A. Tracy and Harold Widom. On orthogonal and symplectic matrix ensembles. *Communications in Mathematical Physics*, 177(3):727–754, 1996. ISSN 0010-3616. doi: 10.1007/bf02099545.

- Craig A. Tracy and Harold Widom. The Distribution of the Largest Eigenvalue in the Gaussian Ensembles: $\beta = 1, 2, 4$. In *Calogero-Moser- Sutherland Models*, pages 461–472. Springer New York, 2000. ISBN 978-1-4612-1206-5. doi: 10.1007/978-1-4612-1206-5__29. URL https://doi.org/10.1007/978-1-4612-1206-5_29.
- Joel A. Tropp. An Introduction to Matrix Concentration Inequalities. *Foundations and Trends® in Machine Learning*, 8(1-2):1–230, 2015. ISSN 1935-8237. doi: 10.1561/2200000048.
- Antonia M. Tulino and Sergio Verdú. Random Matrix Theory and Wireless Communications. *Foundations and Trends® in Communications and Information Theory*, 1(1):1–182, 2004. ISSN 1567-2190. doi: 10.1561/0100000001.
- David E. Tyler. Robustness and Efficiency Properties of Scatter Matrices. *Biometrika*, 70(2):411, 1983. ISSN 0006-3444. doi: 10.2307/2335555.
- Aad W. Van der Vaart. *Asymptotic Statistics*, volume 3 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, 2000. ISBN 9780521784504. doi: 10.1017/cbo9780511802256.
- Pascal Vallet, Xavier Mestre, and Philippe Loubaton. Performance analysis of an improved music doa estimator. *IEEE transactions on signal processing*, 63(23):6407–6422, 2015.
- Vladimir Vapnik. Principles of Risk Minimization for Learning Theory. In *Advances in Neural Information Processing Systems*, volume 4 of *NIPS'92*, pages 831–838. Morgan-Kaufmann, 1992. URL <https://proceedings.neurips.cc/paper/1991/file/ff4d5fbafdf976cfdc032e3bde78de5-Paper.pdf>.
- Andrea Vedaldi and Andrew Zisserman. Efficient Additive Kernels via Explicit Feature Maps. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 34(3):480–492, 2012. ISSN 0162-8828. doi: 10.1109/tpami.2011.153.
- Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. In Yonina C. Eldar and Gitta Editors Kutyniok, editors, *Compressed Sensing: Theory and Applications*, page 210–268. Cambridge University Press, 2012. doi: 10.1017/cbo9780511794308.006.
- Roman Vershynin. *High-Dimensional Probability: An Introduction with Applications in Data Science*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2018. ISBN 9781108415194. doi: 10.1017/9781108231596.001.
- Dan Voiculescu, Kenneth Dykema, and Alexandru Nica. Free Random Variables. *CRM Monograph Series*, pages 55–66, 1992. ISSN 1065-8599. doi: 10.1090/crmm/001/05.

- Sebastian Wagner, Romain Couillet, Mérouane Debbah, and Dirk T. M. Slock. Large System Analysis of Linear Precoding in Correlated MISO Broadcast Channels Under Limited Feedback. *IEEE Transactions on Information Theory*, 58(7):4509–4537, 2012. ISSN 0018-9448. doi: 10.1109/tit.2012.2191700.
- Mati Wax and Thomas Kailath. Detection of signals by information theoretic criteria. *IEEE Transactions on acoustics, speech, and signal processing*, 33(2):387–392, 1985.
- Chao-Kai Wen, Guangming Pan, Kai-Kit Wong, Meihui Guo, and Jung-Chieh Chen. A Deterministic Equivalent for the Analysis of Non-Gaussian Correlated MIMO Multiple Access Channels. *IEEE Transactions on Information Theory*, 59(1):329–352, 2013. ISSN 0018-9448. doi: 10.1109/tit.2012.2218571.
- Eugene P. Wigner. Characteristic Vectors of Bordered Matrices with Infinite Dimensions. *The Annals of Mathematics*, 62(3):548, 1955. ISSN 0003-486X. doi: 10.2307/1970079.
- Christopher Williams. Computing with Infinite Networks. In *Advances in Neural Information Processing Systems*, volume 9 of *NIPS’97*, pages 295–301. MIT Press, 1997. URL <https://proceedings.neurips.cc/paper/1996/file/ae5e3ce40e0404a45ecacaaf05e5f735-Paper.pdf>.
- John Wishart. The Generalised Product Moment Distribution in Samples from a Normal Multivariate Population. *Biometrika*, 20A(1/2):32–52, 1928. ISSN 0006-3444. doi: 10.2307/2331939.
- Svante Wold, Kim Esbensen, and Paul Geladi. Principal component analysis. *Chemometrics and Intelligent Laboratory Systems*, 2(1-3):37–52, 1987. ISSN 0169-7439. doi: 10.1016/0169-7439(87)80084-9.
- Han Xiao, Kashif Rasul, and Roland Vollgraf. Fashion-MNIST: a Novel Image Dataset for Benchmarking Machine Learning Algorithms. *arXiv*, 2017. URL <https://arxiv.org/abs/1708.07747>.
- Lechao Xiao, Yasaman Bahri, Jascha Sohl-Dickstein, Samuel Schoenholz, and Jeffrey Pennington. Dynamical Isometry and a Mean Field Theory of CNNs: How to Train 10,000-Layer Vanilla Convolutional Neural Networks. In *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 5393–5402, Stockholm Sweden, 2018. PMLR. URL <http://proceedings.mlr.press/v80/xiao18a.html>.
- Liusha Yang, Romain Couillet, and Matthew R McKay. A Robust Statistics Approach to Minimum Variance Portfolio Optimization. *IEEE Transactions on Signal Processing*, 63(24):6684–6697, 2015. ISSN 1053-587X. doi: 10.1109/tsp.2015.2474298.

- Jianfeng Yao, Romain Couillet, Jamal Najim, and Mérouane Debbah. Fluctuations of an Improved Population Eigenvalue Estimator in Sample Covariance Matrix Models. *IEEE Transactions on Information Theory*, 59(2):1149–1163, 2013. ISSN 0018-9448. doi: 10.1109/tit.2012.2222862.
- Yuan Yao, Lorenzo Rosasco, and Andrea Caponnetto. On Early Stopping in Gradient Descent Learning. *Constructive Approximation*, 26(2):289–315, 2007. ISSN 0176-4276. doi: 10.1007/s00365-006-0663-2.
- Tayeb Zarrouk, Romain Couillet, Florent Chatelain, and Nicolas Le Bihan. Performance-Complexity Trade-Off in Large Dimensional Statistics. In *2020 IEEE 30th International Workshop on Machine Learning for Signal Processing (MLSP)*, volume 00 of *2020 IEEE 30th International Workshop on Machine Learning for Signal Processing (MLSP)*, pages 1–6. IEEE, 2020. ISBN 9781728166636. doi: 10.1109/mlsp49062.2020.9231568.
- Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. In *5th International Conference on Learning Representations*, ICLR’16, 2016. URL <https://openreview.net/forum?id=Sy8gdB9xx>.
- Teng Zhang, Xiuyuan Cheng, and Amit Singer. Marchenko-Pastur Law for Tyler’s M-estimator. *arXiv*, 2014. URL <https://arxiv.org/abs/1401.3424>.
- Shurong Zheng, Zhidong Bai, and Jianfeng Yao. CLT for eigenvalue statistics of large-dimensional general Fisher matrices with applications. *Bernoulli*, 23(2):1130–1178, 2017. ISSN 1350-7265. doi: 10.3150/15-bej772.
- Dengyong Zhou, Olivier Bousquet, Thomas Navin Lal, Jason Weston, and Bernhard Scholkopf. Learning with Local and Global Consistency. In *Advances in Neural Information Processing Systems*, volume 16 of *NIPS’04*, pages 321–328. MIT Press, 2004. URL <https://proceedings.neurips.cc/paper/2003/file/87682805257e619d49b8e0dfdc14affa-Paper.pdf>.
- Xueyuan Zhou and Mikhail Belkin. Semi-supervised Learning by Higher Order Regularization. In *Proceedings of the Fourteenth International Conference on Artificial Intelligence and Statistics*, volume 15 of *Proceedings of Machine Learning Research*, pages 892–900, Fort Lauderdale, FL, USA, 2011. JMLR Workshop and Conference Proceedings. URL <http://proceedings.mlr.press/v15/zhou11b.html>.
- Xiaojin Zhu. Semi-Supervised Learning Literature Survey. Technical report, University of Wisconsin-Madison Department of Computer Sciences, 9 2005. URL <https://minds.wisconsin.edu/bitstream/handle/1793/60444/TR1530.pdf>.
- Xiaojin Zhu and Zoubin Ghahramani. Learning from labeled and unlabeled data with label propagation. Technical report, Citeseer, 2002.

URL <http://citeseerx.ist.psu.edu/viewdoc/download;jsessionid=46D8FA6E43437BB5FAE59DF68862E758?doi=10.1.1.13.8280&rep=rep1&type=pdf>.

Xiaojin Zhu, Zoubin Ghahramani, and John Lafferty. Semi-Supervised Learning Using Gaussian Fields and Harmonic Functions. In *Proceedings of the Twentieth International Conference on Machine Learning (ICML-2003)*, volume 3 of *ICML '03*, pages 912–919. AAAI Press, 2003. URL <https://www.aaai.org/Library/ICML/2003/icml03-118.php>.