

# **SHAMBA v 1.1**

## **Methodology**

The Small-Holder Agriculture Mitigation Benefit Assessment model for estimation of greenhouse gas emission reductions and removals that result from smallholder farmers using Climate Smart Agriculture and/or tree planting in the tropics

Woollen, E., Berry, N., Cross, A., Hagdorn, M., Hughes, M., Wells, G.J., Ryan, C.M.

The University of Edinburgh

Tropical land use team

School of GeoSciences

Edinburgh, EH9 3JN, UK

## Contents

<b>Contents .....</b>	<b>2</b>
<b>Definitions .....</b>	<b>4</b>
<b>1. Model summary .....</b>	<b>5</b>
<b>2. Overview of GHG accounting approach .....</b>	<b>5</b>
<b>3. SHAMBA model overview .....</b>	<b>7</b>
<b>4. Modelling changes in woody biomass .....</b>	<b>8</b>
4.1 Biomass model description .....	8
4.2 Biomass model parameterisation .....	9
4.3 Biomass growth .....	11
4.4 Biomass loss .....	13
4.5 Biomass change .....	14
4.6 Stand density change .....	15
4.7 Tree-soil inputs .....	15
<b>5. Crop model .....</b>	<b>16</b>
5.1 Crop model description .....	16
5.2 Crop model parameterisation .....	17
5.3 Crop model calculations .....	17
<b>6. External organic soil inputs .....</b>	<b>19</b>
6.1 External organic soil input description .....	19
6.2 External organic input parameterisation .....	20
6.3 External organic input calculations .....	20
<b>7. Emissions due to biomass burning .....</b>	<b>20</b>
7.1 Biomass burning model description .....	21
7.2 Biomass burning model parameterisation .....	21
7.3 Biomass burning model calculations .....	22
7.4 Soil inputs reduced by fire .....	23
<b>8. Modelling changes in soil organic carbon .....</b>	<b>24</b>
8.1 SOC model overview .....	24
8.2 SOC model parameterisation .....	24
8.3 Model structure and SOC partitioning .....	25
8.4 SOC model initialisation .....	27
8.3 SOC changes .....	29
8.4 SOC model limitations .....	30
<b>9. Emissions due to nitrogen inputs from plants .....</b>	<b>30</b>
9.1 Plant nitrogen input emissions description .....	30
9.2 Plant nitrogen input emission parameterisation .....	30
9.3 Plant nitrogen input emission calculations .....	31

<b>10. Emissions due to fertiliser use.....</b>	<b>31</b>
<i>10.1 Emissions from fertiliser use description .....</i>	<i>31</i>
<i>10.2 Emissions from fertiliser use parameterisation .....</i>	<i>32</i>
<i>10.3 Emissions from fertiliser use calculations.....</i>	<i>32</i>
<b>11. Future model developments .....</b>	<b>33</b>
<b>References.....</b>	<b>35</b>
<b>Appendix 1: Model default values, sources, justification and applicability .....</b>	<b>37</b>

## Definitions

**Accounting period:** length of time over which greenhouse gas emissions and removals are quantified

**Activity:** a component of an intervention, such as agroforestry of a specific type

**Agroforestry:** agriculture incorporating the cultivation of trees

**Baseline:** land use/management practice(s) without any intervention(s) (i.e. “business as usual”)

**Climate Smart Agriculture (CSA):** using conservation agriculture and agroforestry techniques for improved GHG mitigation in agriculture

**Conservation agriculture:** uses agricultural methods (minimal soil disturbance, permanent soil cover and crop rotations) to achieve sustainable and profitable agriculture

**Intervention:** change in land use/management practice from baseline

**Project:** a collection of interventions

**Scenario:** a set of unique land use/management practices

**Small-holding/holders:** farms supporting single families with a mixture of cash crops and or subsistence farming

**User(s):** person(s) using the model

## 1. Model summary

The SHAMBA (Small-Holder Agriculture Mitigation Benefit Assessment) model estimates greenhouse gas (GHG) emissions or removals resulting from a change in land management practices. SHAMBA is designed to model a **baseline scenario** (where land management activities continue as business as usual) and an **intervention scenario** consisting of **activities** that can be described as Climate Smart Agricultural practices (CSA) including, conservation agriculture, agroforestry and other tree planting. SHAMBA models the changes in carbon stocks in soils and woody biomass, and the GHG emissions from biomass burning, plant nitrogen inputs to soils, and fertiliser use over the **accounting period** for baseline and intervention activities. Net emissions and removals are calculated on a yearly basis for the length of the accounting period, in units of tonnes (t) of carbon dioxide equivalent (CO<sub>2e</sub>) per hectare (ha). Version one of the SHAMBA model is designed to work with **smallholder** systems in the tropics.

This document describes the science of the SHAMBA model, and details the calculations and parameters, as well as outlining the data requirements to use the model. For a full description of how to use the model for carbon accounting, please see the SHAMBA methodology (<http://shambatool.wordpress.com/>). The methodology also outlines **applicability conditions** for using the SHAMBA tool to estimate GHG emissions and removals for carbon accounting projects in the tropics.

## 2. Overview of GHG accounting approach

This model was developed for the purpose of accounting for changes in soil and woody biomass carbon stocks and GHG emissions due to changing agricultural practices and tree planting. Soil carbon and woody biomass changes are modelled with simple quasi-process-based approaches, whilst emissions from other sources (e.g. biomass burning, the use of fertilisers) are accounted for using simpler (IPCC Tier 1-type) approaches. The model consists of three sub-models, one for soil, crops and woody biomass, working on a hectare basis.

Soil organic carbon is modelled using RothC, one of the most widely used soil carbon models. It has been used to model soil carbon dynamics globally, and is freely available to download<sup>1</sup>. RothC was originally developed and parameterised to model turnover of organic carbon in arable surface soils from the Rothamsted UK long-term field experiments, but has since been applied at a range of different scales and systems (Milne et al. 2007). It has been widely used to model the effects of agricultural (Farage et al. 2007, Traoré et al. 2008, Nakamura et al. 2010) and agroforestry practices (Diels et al. 2004, Kaonga and Coleman 2008), on soil carbon in the tropics.

However, several key effects of CSA and other land management practices are not represented in RothC. Therefore in the SHAMBA model the woody biomass growth and carbon inputs to the soil pool from agroforestry and tree planting activities are estimated with a new stock and flow biomass model

---

<sup>1</sup> [http://www.rothamsted.ac.uk/aen/carbon/mod26\\_3\\_win.pdf](http://www.rothamsted.ac.uk/aen/carbon/mod26_3_win.pdf)

driven by assumed tree growth rates. The carbon inputs to the soil pool from crops are estimated using IPCC (2006) guidelines. External organic inputs, such as additions of litter from outside sources, to the field are also included. Emissions from biomass burning, the volatilisation of nitrogen (N) in soils from plant inputs, and N fertiliser use are estimated through the use of simple equations and IPCC<sup>2</sup> default values. The following equations define the sources and sinks of GHGs included in this model.

For a baseline scenario, GHG emissions or removals per hectare in year  $y$  are calculated as:

$$BE_y = BE_{BB_y} + BE_{NI_y} + BE_{NF_y} + BE_{SO_y} + BE_{WB_y} \quad (\text{Equation 1})$$

For the intervention scenario, the calculation is identical:

$$PE_y = PE_{BB_y} + PE_{NI_y} + PE_{NF_y} + PE_{SO_y} + PE_{WB_y} \quad (\text{Equation 2})$$

Where for baseline (variables starting with  $B$ ) and intervention (variables starting with  $P$ ) emissions:

$E_y$  is the GHG emissions under the scenario for year  $y$  (tCO<sub>2e</sub>/ha);

$E_{BB_y}$  is the emissions from biomass burning in year  $y$  of the scenario (tCO<sub>2e</sub>/ha);

$E_{NI_y}$  is the emissions resulting from the nitrogen inputs to soils from plants in year  $y$  of the scenario (tCO<sub>2e</sub>/ha);

$E_{NF_y}$  is the direct emissions resulting from the use of N fertilisers in year  $y$  of the scenario (tCO<sub>2e</sub>/ha);

$E_{SO_y}$  is the emissions from change in soil organic carbon stocks in year  $y$  of the scenario (tCO<sub>2e</sub>/ha); and

$E_{WB_y}$  is the emissions from change in woody biomass of trees planted through scenario activities in year  $y$  of the scenario (tCO<sub>2e</sub>/ha).

For all emissions, positive numbers refer to emissions and negative numbers refer to removals. The units of all terms are tCO<sub>2e</sub>/ha, obtained for carbon stocks by multiplying by the ratio of the molecular weights of CO<sub>2</sub> and C (44/12), and for non-CO<sub>2</sub> fluxes, the appropriate global warming potential (GWPs).

Total emissions/removals for each scenario are given by summing over the years  $y = 1$  to  $y = d$ , where  $d$  is the accounting period:

$$BE = \sum_{y=1}^d BE_y \quad (\text{Equation 3})$$

$$PE = \sum_{y=1}^d PE_y \quad (\text{Equation 4})$$

Where:

$BE$  is the total emissions for the baseline scenario (tCO<sub>2e</sub>/ha) over the accounting period

$PE$  is the total emissions for the intervention scenario (tCO<sub>2e</sub>/ha) over the accounting period

The overall net impact of the intervention is given by the difference between the total baseline and intervention emissions:

$$E_{net} = PE - BE \quad (\text{Equation 5})$$

Where:

$E_{net}$  is the total net emissions/removals as a result of the intervention (tCO<sub>2e</sub>/ha)

Again, all terms are in units of tCO<sub>2e</sub>/ha, and negative numbers represent removals. Note that this formulation implies that all mitigation is considered to have an equal weight, regardless of the year in which it occurs.

### 3. SHAMBA model overview

Each section (Table 1) of this document details how to calculate the terms in Eq. 1 and Eq. 2, such that the modelled intervention and baseline emissions can be calculated (Eq. 3, 4), and the net impact estimated (Eq. 5). All calculations are performed on a per hectare basis on a yearly time step.

**Table 1: Summary of main sections of the model description and the origin of the approach**


Modelled Effect	Parameter in Eq 1 & 2	Section	Source of approach
Woody biomass carbon changes	$E_{WB}$	4	UoE tropical land use team
Crop residues and resultant soil inputs		5	IPCC <sup>2</sup>
External organic inputs		6	UoE tropical land use team
Emissions from biomass burning	$E_{BB}$	7	IPCC <sup>2</sup> and SALM <sup>3</sup>
Soil organic carbon changes	$E_{SO}$	8	RothC and UoE tropical land use team
Emissions from plant nitrogen inputs to soils	$E_{NI}$	9	IPCC and SALM
Emissions from N fertiliser use	$E_{NF}$	10	SALM and CDM <sup>4</sup> tool

SHAMBA models each component (Table 1) based on input data, using outputs from some components as inputs to others. A diagram of the basic structure of the model and links between components is shown below (Fig. 1).

<sup>2</sup> IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan

<sup>3</sup> <http://v-c-s.org/sites/v-c-s.org/files/VM0017%20SALM%20Methodolgy%20v1.0.pdf>

<sup>4</sup> <http://cdm.unfccc.int/methodologies/ARmethodologies/tools/ar-am-tool-07-v1.pdf>



**Fig. 1:** SHAMBA model showing each modelled component and the flow of information between them.

## 4. Modelling changes in woody biomass

This section details the parameterisation and use of the SHAMBA biomass model, which is used to estimate woody biomass growth and tree inputs to the soil for agroforestry and tree planting activities. The following section details the method.

### 4.1 Biomass model description


The SHAMBA biomass model (Fig. 2) estimates changes to woody biomass carbon pools over the project duration using a mass balance approach. It keeps track of the following pools; stem, branches, leaves (the sum of these three = AGB), fine roots, coarse roots (= BGB). Biomass flows into all the pools, based on a simple allocation of each year's net primary productivity (NPP), estimated from tree growth rates and an allometric equation (see below). Biomass flows out of each pool via a litter flux, determined by the turnover rate. Each pool is defined as the biomass in that pool per tree multiplied by the stand density per hectare, in units of tC/ha. Stand density, (also known as stocking density), is a function of initial planting density, tree mortality and thinning. As such the biomass model must be parameterised and run for a cohort of trees, where the cohort is planted at the same time and is made up of the same species (or generic tree type, see below). Thinned and dead biomass can be removed from the system, or can be left, in which case it is added to the litter flux. The litter flux drives the biomass burning model (section 7), the soil carbon model (section 8) and the emissions from plant N input model (section 9). Each pool is updated yearly, with fluxes calculated based on the previous year's pool sizes.

The biomass model assumes:

- The allometric equation is appropriate for the planted tree species



- If pruning occurs the allometric equation is appropriate for pruning (pruning will change the tree allometry)
- All trees planted in a cohort have similar growth rates and allocation parameters



**Fig. 2:** Graphical representation of the biomass model where growth of biomass (NPP) is allocated to each of the five biomass pools; leaves, branches, stem = above ground biomass (AGB), coarse roots, fine roots = below ground biomass (BGB). Biomass in each pool is lost due to turnover, thinning and mortality, entering the litter flux or removal flux.

## 4.2 Biomass model parameterisation

The data needed to parameterise and drive the model for a single tree species and/or cohort are shown in Table 2. The parameters for the biomass model should either be entered from user data, or defined by defaults if applicable (default parameter values are given in Appendix 1).

**Table 2:** Model parameters required for each tree species/cohort planted

Parameter	Symbol	Units	Default value	Notes
Expected growth rate of aboveground biomass of a single tree in year $y$	$B_{inc,y}$	ABG kg C/ year	<i>Users must provide this, or calculate it from the DBH increment</i>	See 4.3.1
Initial stand or planting density	$SD_{y=0}$	trees/ ha	<i>Users must provide this</i>	
Fraction of stand density thinned in year $y$	$th_y$	stems removed / total SD	<i>Users must provide this</i>	See 4.6
Fraction of thinned stems left in the field	$thf_{stem}$	t C stems left/ total t C of thinned stems	<i>Users must provide this</i>	See 4.4
Fraction of thinned branches left in the field	$thf_{branch}$	t C branches left / total t C thinned branches	<i>Users must provide this</i>	See 4.4

Fraction of thinned leaves left in the field	$thf_{leaf}$	t C leaves left / total t C thinned leaves	1	See 4.4
Fraction of thinned coarse roots left in the field	$thf_{croot}$	t C coarse roots left / total t C thinned coarse roots	1	See 4.4
Fraction of thinned fine roots left in the field	$thf_{froot}$	t C fine roots left / total t C thinned fine roots	1	See 4.4
Tree mortality in year y	$tm_y$	trees that die in year y/ total SD in year y	<i>Users must provide this</i>	See 4.4
Fraction of dead stems left in the field	$tmf_{stem}$	t C stems left/ total t C of dead stems	<i>Users must provide this</i>	See 4.4
Fraction of dead branches left in the field	$tmf_{branch}$	t C branches left / total t C dead branches	<i>Users must provide this</i>	See 4.4
Fraction of dead leaves left in the field	$tmf_{leaf}$	t C leaves left/ total t C of dead leaves	1	See 4.4
Fraction of dead coarse roots left in the field	$tmf_{croot}$	t C coarse roots left / total t C dead coarse roots	1	See 4.4
Fraction of dead fine roots left in the field	$tmf_{froot}$	t C fine roots left / total t C dead fine roots	1	See 4.4
NPP allocation to stem	$al_{stem}$	t C to stem/ total t C NPP	0.69	See 4.3.2
NPP allocation to branches	$al_{branch}$	t C to branches/ total t C NPP	0.31	See 4.3.2
NPP allocation to leaves	$al_{leaf}$	t C to leaves/ total t C NPP	0.10	See 4.3.2
NPP allocation to fine roots	$al_{froot}$	t C in fine roots/ total t C NPP	0.10	See 4.3.2
NPP allocation to coarse roots	$al_{croot}$	t C to roots/ total t C NPP	Equals product of root:shoot ratio and $al_{stem}$	See 4.3.2
Turnover rate of stem	$to_{stem}$	t C stem turned over /total t C stem	0	See 4.4
Turnover rate of branches	$to_{branch}$	t C branches turned over / total t C branches	0.05	See 4.4
Turnover rate of leaves	$to_{leaf}$	t C leaves turned over/ total t C leaves	1	See 4.4
Turnover rate of fine roots	$to_{froot}$	t C fine roots turned over / total t C fine roots	0.8	See 4.4
Turnover rate of coarse roots	$to_{croot}$	t C coarse roots turned over / total t C coarse roots	0	See 4.4
Root:shoot ratio	$t_{rs}$	t C BGB / t C AGB	0.26	
Fraction of roots in the top 0-30 cm of soil	$t_{30}$	t C roots in top 0-30 / total t C roots	0.7	
Wood density	$t_d$	g/ cm <sup>3</sup>	0.60	Only needed if using the Chave 2009 allometric eqs.
Stem carbon content	$tc_{stem}$	g C/ g DM	0.5	See 4.7
Branch carbon content	$tc_{branch}$	g C/ g DM	0.5	See 4.7
Leaf carbon content	$tc_{leaf}$	g C/ g DM	0.5	See 4.7
Fine root carbon content	$tc_{froot}$	g C/ g DM	0.5	See 4.7
Coarse root carbon content	$tc_{croot}$	g C/ g DM	0.5	See 4.7
Stem wood nitrogen content	$tn_{stem}$	g N/ g DM	0.0015	See 4.7
Branch wood nitrogen content	$tn_{branch}$	g N/ g DM	Assumed same as stem	See 4.7
Leaf litter nitrogen content	$tn_{leaf}$	g N/ g DM	0.01 if non-legume 0.02 if legume	See 4.7

Fine root nitrogen content	$tn_{root}$	g N/ g DM	0.0113	See 4.7
Coarse root nitrogen content	$tn_{croot}$	g N/ g DM	Assumed same as stem	See 4.7

## 4.3 Biomass growth

### 4.3.1 Net primary productivity

The SHAMBA biomass model uses growth models to estimate the net primary productivity (NPP) of tree growth. Given a dataset of age (years) and stem diameter (cm) of the planted tree species or generic type, the stem diameter data is converted to AGB using an allometric equation appropriate to the planted tree species/type, giving AGB (stem and branches) in units of kg C. The model provides several default allometric equations which are appropriate for generic tree types of dry tropical species, moist tropical species or wet tropical species (Chave et al. 2005), miombo tree species (Ryan et al. 2011), or tree species of *Markhamia lutea*, *Grevillea robusta*, and *Maesopsis eminii* (Tumwebaze et al. 2013). If none of the allometric equations are appropriate to the planted tree, user defined allometric equations can be used instead.

The biomass data is plotted against age of tree (Fig. 3), and several growth models are fitted to the data using optimisation methods:

$$agb = ax \quad (\text{Linear, Equation 6.1})$$

$$agb = (1 + a)^x - 1 \quad (\text{Exponential, Equation 6.2})$$

$$agb = a(1 - e^{-bx}) \quad (\text{Hyperbolic, Equation 6.3})$$

$$agb = \frac{a}{1 + e^{-b(x-c)}} \quad (\text{Logistic, Equation 6.4})$$

which when differentiated give:

$$\frac{dagb}{dx} = a \quad (\text{Equation 7.1})$$

$$\frac{dagb}{dx} = (1 + a)^x \cdot \log(1 + a) \quad (\text{Equation 7.2})$$

$$\frac{dagb}{dx} = a \cdot b \cdot e^{-bx} \quad (\text{Equation 7.3})$$

$$\frac{dagb}{dx} = \frac{a \cdot b \cdot e^{-b(x-c)}}{(e^{-b(x-c)} + 1)^2} \quad (\text{Equation 7.4})$$

Where:  $agb$  is the modelled tree AGB (kg C)

$x$  is the age of the tree


$a, b, c$  are fitted parameters

$dagb$  is the growth in AGB (kg C for the period  $dx$ )

The derivative of the best-fit equation is used to calculate the growth of AGB as a function of last year's AGB. This growth represents the NPP of above ground biomass of a single tree in year  $y$ :

$$NPP_{agb} = f(agb_{y-1}) \quad (\text{Equation 8})$$

Where:  $NPP_{agb,y}$  is the AGB growth of a single tree in year  $y$  (kg C)



**Fig. 3:** Growth curves (Eqs. 6.1-6.4) fitted to example data of age and biomass for a single tree species/type.

#### 4.3.2 Allocation to biomass pools

The NPP of a single tree is then converted to tonnes and scaled to the hectare by multiplying by the stand density, to obtain total NPP (t C/ha), assuming all trees of a particular species or cohort are of equal size and grow at similar rates. Total NPP is allocated to each of the five pools by the allocation parameters, giving biomass growth in each pool,  $i = \{\text{stem, leaf, branch, coarse roots, fine roots}\}$ :

$$NPP_{i,y} = (NPP_{agb,y}/1000) \cdot SD_{y-1} \cdot al_i \quad (\text{Equation 9})$$

Where:  $NPP_{i,y}$  is the NPP in pool  $i$  in year  $y$  (t C/ha)

$SD_{y-1}$  is the stand density in the year before  $y$  (trees/ha)

$al_i$  is the allocation of AGB NPP to pool  $i$

The default above-ground allocation parameters ( $al_{stem}$ ,  $al_{branch}$ ,  $al_{leaf}$ ) were determined using measurements taken from tree species planted as part of agroforestry activities (Tumwebaze et al. 2013), and several miombo woodland tree species (Ryan et al. 2011) (Table 3).

**Table 3:** Allocation patterns for several tree species, where allocation is a ratio to total above-ground biomass (AGB = stems and branches). nd indicates no data was available.

Species	DBH	Allocation to stems	Allocation to branches	Allocation to leaves	Reference
---------	-----	---------------------	------------------------	----------------------	-----------

<i>Units</i>	<i>cm</i>	<i>mass stems/ total AGB</i>	<i>mass branches/ total AGB</i>	<i>mass leaves/ total AGB</i>	
<i>Grevillea robusta</i>	31.08	0.63	0.37	0.08	Tumwebaze <i>et al.</i> (2013)
<i>Maesopsis eminii</i>	29.28	0.68	0.32	0.09	Tumwebaze <i>et al.</i> (2013)
<i>Markhamia lutea</i>	23.93	0.72	0.28	0.12	Tumwebaze <i>et al.</i> (2013)
<i>Julbernardia globiflora</i>	30.0	0.70	0.30	nd	Ryan <i>et al.</i> (2011)
<i>Brachystegia spiciformis</i>	32.3	0.69	0.31	nd	Ryan <i>et al.</i> (2011)
<i>Brachystegia bohemii</i>	28.0	0.69	0.31	nd	Ryan <i>et al.</i> (2011)
<b>Mean</b>	<b>29.10</b>	<b>0.69</b>	<b>0.31</b>	<b>0.10</b>	

The allocation patterns are fairly consistent between species (Table 3), and the default values for allocation of above-ground pools were therefore designated as the means. The default root allocation ( $al_{root}$ ) parameter is determined from root:shoot ratios. Due to a paucity of data for agroforestry trees, we use the average root:shoot ratio as reported in a global meta-analysis of tree root:shoot ratios (Cairns *et al.* 1997), which was within the range reported in Mokany *et al.* (2006) for tropical forest trees. We further assume that fine roots have the same allocation ratio as leaves (i.e. fine root productivity is the same as leaves). Other studies in temperate forests using a range of methods have found a wide range of fine root:leaf productivity ratios, but with a mean ( $\pm$ SD) of  $1.08 \pm 1.35$  (Hendricks *et al.* 2006), suggesting our assumption is reasonable. Finally, the default allocation parameters chosen for stems, leaves and roots agreed closely with a meta-analysis of biomass allocation (Poorter *et al.* 2012), where the mean allocation for natural tropical forests and woodlands ranged from 0.8-0.6 for stems, 0.02-0.06 for leaves, and 0.16-0.36 for roots.

The default allocation values, as used in the SHAMBA biomass model (Table 2), are therefore generally applicable across most tree species. However, as demonstrated for *Casuarina equisetifolia* (Tumwebaze *et al.* 2013), some trees may have a differing allometry to the 'generic' tree. Therefore, if local project data is available, or species specific data from peer reviewed literature is available, more specific allocation parameters should be used in the model.

#### 4.4 Biomass loss

From the growth model, biomass C stocks (t C/ha) are increased in each of the five pools over time. Some of the biomass from each pool is lost every year due to turnover, mortality and thinning. The flux of biomass from each pool is determined by the rate of loss:

$$BL_{i,y} = B_{i,y-1} \cdot to_i \quad (\text{Equation 10})$$

$$BT_{i,y} = B_{i,y-1} \cdot th_y \quad (\text{Equation 11})$$

$$BM_{i,y} = B_{i,y-1} \cdot tm_y \quad (\text{Equation 12})$$

$$B_{loss,i,y} = BL_{i,y} + BT_{i,y} + BM_{i,y} \quad (\text{Equation 13})$$

Where:  $BL_{i,y}$  is the biomass turned over in pool  $i$  in year  $y$  (t C/ha)

$B_{i,y-1}$  is the biomass in pool  $i$  in the year before  $y$  (t C/ha)

$to_i$  is the turnover rate of pool  $i$  (t C turned over/ total t C)

$BT_{i,y}$  is the biomass thinned in pool  $i$  in year  $y$  (t C/ha)

$th_y$  is the thinning fraction in year  $y$  (thinned trees/ total stand density)

$BM_{i,y}$  is the dead biomass in pool  $i$  in year  $y$  (t C/ha)

$tm_y$  is the mortality rate in year  $y$  (dead trees/ total stand density)

$B_{loss,i,y}$  is the total biomass lost in pool  $i$  in year  $y$  (t C/ha)

The default values for turnover (table 2) of each pool is based on the assumption that stems and coarse roots will not have significant losses of biomass as litter annually, and that trees are fully deciduous (see appendix 1). If planted trees are not deciduous (or have a leaf life span > 1 year), or have significant inputs from stem bark shedding or branches, the turnover rates should be specified by the user.

Thinning is defined as the removal of whole trees in this model, and not just pruning of branches or other parts. Mortality and thinning of trees can occur in any year, with different fractions dead or thinned in every year, if applicable. Biomass from the thinned ( $BT_i$ ) and dead ( $BM_i$ ) pools can be left in the field or removed. The fraction of dead and thinned biomass (e.g.  $tmf_{stem}$ ,  $thfs_{stem}$ ) which is left in the field determines the amount of thinned and dead biomass which is added to the litter flux. All coarse roots, fine roots and leaves of dead and felled trees are assumed to remain in the field in default settings. Thereby, biomass from the thinned and dead trees which is removed or left in the field is determined:

$$BT_{on,i,y} = BT_{i,y} \cdot thf_i \quad (\text{Equation 14})$$

$$BM_{on,i,y} = BM_{i,y} \cdot tmf_i \quad (\text{Equation 15})$$

$$BT_{off,i,y} = BT_{i,y}(1 - thf_i) \quad (\text{Equation 16})$$

$$BM_{off,i,y} = BM_{i,y}(1 - tmf_i) \quad (\text{Equation 17})$$

Where:  $BT_{on,i,y}$  is the biomass from pool  $i$  which is left in the field from thinned trees in year  $y$  (tC/ha)

$thf_i$  is the fraction of thinned biomass in pool  $i$  which is left in the field (t C left/ total t C thinned)

$BM_{on,i,y}$  is the biomass from pool  $i$  which is left in the field from dead trees in year  $y$  (tC/ha)

$tmf_i$  is the fraction of dead biomass in pool  $i$  which is left in the field (t C left/ total t C dead)

$BT_{off,i,y}$  is the biomass from pool  $i$  which is removed from the field from thinned trees in year  $y$  (tC/ha)

$BM_{off,i,y}$  is the biomass from pool  $i$  which is removed from the field from dead trees in  $y$  (tC/ha)

## 4.5 Biomass change

Biomass in each pool is thereby a function of the biomass growth and the biomass loss in each year:

$$B_{i,y} = B_{i,y-1} + NPP_{i,y} - B_{loss,i,y} \quad (\text{Equation 18})$$

Where:  $B_{i,y}$  is biomass in pool  $i$  in year  $y$  (t C/ha)

Thereby, total biomass carbon stocks (stems, leaves, branches, coarse roots and fine roots) which remains each year is used to calculate the total emissions from woody biomass (Eq. 1-2) due to tree planting (negative values are an uptake):

$$B_{total,y} = \sum_{i=1}^5 B_{i,y} \quad (\text{Equation 19})$$

$$E_{WB,y} = (B_{total,y-1} - B_{total,y}) \cdot mw_{CO_2} \quad (\text{Equation 20})$$

Where:  $B_{total,y}$  is the total woody biomass in year  $y$  (t C/ha)

$E_{WB,y}$  is the emissions from woody biomass change in year  $y$  (tCO<sub>2</sub>e/ha)

$B_{total,y-1}$  is the total woody biomass in the year before  $y$  (t C/ha)

$mw_{CO_2}$  is the molecular ratio of CO<sub>2</sub> and C (44/12)

From this equation the biomass removed due to thinning and mortality are included as an emission. Biomass removed from the field is assumed to be short lived and converted to CO<sub>2</sub> in the atmosphere immediately. Long-lived timber products are not included in this version of the model.

#### 4.6 Stand density change

The model accounts for tree mortality and thinning events by reducing the stand density:

$$SD_y = SD_{y-1} (1 - (tm_y + th_y)) \quad (\text{Equation 21})$$

Where:  $SD_y$  is the stocking or stand density in year  $y$  (trees/ha)

#### 4.7 Tree-soil inputs

The carbon inputs to the soils (t C/ha) from trees are the sum of all the inputs from turned over biomass, thinned biomass and dead biomass left in the field from each of the five pools:

$$B_{CI,i,y} = BL_{i,y} + BT_{on,i,y} + BM_{on,i,y} \quad (\text{Equation 22})$$

Where:  $B_{CI,i,y}$  is the C inputs from pool  $i$  in year  $y$  (t C/ha)

These inputs can be used to determine dry mass and soil N inputs of each pool as well, using the C and N content of each pool:

$$B_{DM,i,y} = \frac{B_{CI,i,y}}{tc_i} \quad (\text{Equation 23})$$

$$B_{NI,i,y} = B_{DM,i,y} \cdot tn_i \quad (\text{Equation 24})$$

Where:  $B_{DM,i,y}$  is the dry biomass inputs of pool  $i$  in year  $y$  (t DM/ha)

$tc_i$  is the carbon content of pool  $i$  (g C/g DM)

$B_{NI,i,y}$  is the N inputs of pool  $i$  in year  $y$  (t N/ha)

$tn_i$  is the N content of pool  $i$  (g N/g DM)

Total above and below ground tree inputs are thus:

$$T_{Clag,y} = B_{Cl,stem,y} + B_{Cl,branch,y} + B_{Cl,leaf,y} \quad (\text{Equation 25})$$

$$T_{Clbg,y} = (B_{Cl,root,y} + B_{Cl,froot,y}) \cdot t_{30} \quad (\text{Equation 26})$$

$$T_{Nlag,y} = B_{NI,stem,y} + B_{NI,branch,y} + B_{NI,leaf,y} \quad (\text{Equation 27})$$

$$T_{Nlbg,y} = (B_{NI,root,y} + B_{NI,froot,y}) \cdot t_{30} \quad (\text{Equation 28})$$

$$T_{DMag,y} = B_{DM,stem,y} + B_{DM,branch,y} + B_{DM,leaf,y} \quad (\text{Equation 29})$$

$$T_{DMbg,y} = (B_{DM,root,y} + B_{DM,froot,y}) \cdot t_{30} \quad (\text{Equation 30})$$

Where:  $T_{Clag,y}$  is the total tree carbon inputs from above ground pools in year  $y$  (t C/ha)

$T_{Clbg,y}$  is the total tree carbon inputs from below ground pools in year  $y$  (t C/ha)

$t_{30}$  is the below ground biomass found in the top 0-30 cm of soil (t C BGB in 0-30 cm/ total t C BGB)

$T_{Nlag,y}$  is the total tree nitrogen inputs from above ground pools in year  $y$  (t N/ha)

$T_{Nlbg,y}$  is the total tree nitrogen inputs from below ground pools in year  $y$  (t N/ha)

$T_{DMag,y}$  is the total tree dry matter inputs from above ground pools in year  $y$  (t DM/ha)

$T_{DMbg,y}$  is the total tree dry matter inputs from below ground pools in year  $y$  (t DM/ha)

These inputs are then passed to the biomass burning model (section 7) before being passed to the soil model (section 8) and the emissions from plant N input model (Section 9).

## 5. Crop model

This section details the parameterisation and use of the SHAMBA crop model, which is used to estimate annual crop residues and inputs to the soil. The following section details the method.

### 5.1 Crop model description

Crop inputs to the soil are difficult to measure and rarely directly observed. Instead, they may be estimated using more readily available data such as crop yields. The SHAMBA crop model uses an IPCC Tier-1 type approach (equations as outlined in table 11.2 of the IPCC<sup>5</sup> guidelines) to estimate crop above and below ground residues for various crop types. The model calculates the annual total above and below ground crop residues (t DM/ha), and the resultant above and below ground crop C (t C/ha) and N (t N/ha) inputs to the soil, on a hectare basis. The total nitrogen inputs to soils are estimated using the values given by the IPCC for nitrogen content of above and below ground residues. Carbon inputs are similarly calculated using peer reviewed values for crop carbon content.

As several different crops are often planted in one field over the course of a year, the crop model is parameterised and run for each crop type planted, summing together the results on a hectare basis to estimate total annual crop residues. The crop types planted and yields are assumed to be the same

<sup>5</sup> [http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4\\_Volume4/V4\\_11\\_Ch11\\_N2O&CO2.pdf](http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_11_Ch11_N2O&CO2.pdf)



for every year of the model run. Therefore, it is important that the model is parameterised with common crop types and mean yields for a typical year for each scenario.

The model assumes:

- Planted crop(s) are definable as one of the IPCC listed crop types or species
- The type or species of crops planted, and crop yields, do not vary between years. Different yields in each year can be accounted for in the model, but not as currently implemented.
- Crop yields are known for each planted crop type/species

## 5.2 Crop model parameterisation

The data required to parameterise the crop model are outlined in Table 4. For full details on default parameters and applicability see Appendix 1.

**Table 4:** Parameters required for each crop species or type planted in each scenario

Parameter	Symbol	Units	Default value
Crop type or species	$c_s$	<i>category</i>	<i>User must provide this</i>
Annual mean crop yield	$c_{yield}$	t DM/ha	<i>User must provide this</i>
Fraction of AG crop residues removed from the field post-harvest	$c_r$	t DM removed ha <sup>-1</sup> / total t DM ha <sup>-1</sup>	<i>User must provide this</i>
Crop root:shoot ratio	$c_{rs}$	t BG DM/ t AG DM	IPCC default value for crop type
Crop AG residue C content	$c_{ac}$	g C /g DM	0.4
Crop BG residue C content	$c_{bc}$	g C /g DM	0.4
Crop AG residue N content	$c_{an}$	g N /g DM	IPCC default value for crop type
Crop BG residue N content	$c_{bn}$	g N /g DM	IPCC default value for crop type
Fraction of crop BG residues in the top 0-30 cm of soil	$c_{30}$	t BG residues in 0-30/ total t BG residues	0.7

\*AG is above-ground, BG is below ground, DM is dry matter, C is carbon, N is nitrogen

## 5.3 Crop model calculations

Yield data should be gathered from local measurements for a typical year for each crop type, or alternatively country specific estimates can be obtained from the FAOSTAT<sup>6</sup> for specified crop species. The yield data can be used to calculate the mass of above-ground crop residues for each crop type ( $i$ ):

$$C_{ag,i} = c_{yield,i} \cdot a_i + b_i \quad (\text{Equation 31})$$

Where:  $C_{ag,i}$  is the mass of total crop above-ground residues for crop type  $i$  (t DM/ha)

$c_{yield,i}$  is annual mean crop dry matter yield for crop type  $i$  (t DM/ha)

$a_i$  and  $b_i$  are model parameters defined for specific crop types  $i$  by the IPCC

<sup>6</sup> <http://faostat.fao.org/site/567/DesktopDefault.aspx?PageID=567#ancor>

Users must provide information on crop residue management, such as if crop residues are removed from the field post-harvest or not. If residues are removed from the field post-harvest, the amount of crop residue which is taken off field can be calculated by:

$$C_{ag-off,i} = C_{ag,i} \cdot c_{f,i} \quad (\text{Equation 32})$$

Where:  $C_{ag-off,i}$  is the mass of above-ground residues that are removed from the field post-harvest for crop type  $i$  (t DM/ha)

$c_{f,i}$  is the fraction of total above-ground residues which are removed from the field post-harvest for crop type  $i$  (1 if all crop residues are removed, 0 if none are removed)

This will determine the amount of above-ground crop residue biomass that is left on the field post-harvest:

$$C_{ag-on} = C_{ag} - C_{ag-off} \quad (\text{Equation 33})$$

Where:  $C_{ag-on}$  is the mass of above-ground residues that are left in the field post-harvest (t DM/ ha)

Using default parameters of root-to-shoot ratios, the model estimates the total below ground crop residues for each crop type planted based on the total above ground biomass:

$$C_{bg,i} = (C_{yield,i} + C_{ag,i}) \cdot c_{rs,i} \quad (\text{Equation 34})$$

Where:  $C_{bg,i}$  is the mass of below ground residues for crop type  $i$  (t DM/ha)

$c_{rs,i}$  is the crop root:shoot ratio for crop type  $i$  (t below ground biomass/t above ground biomass)

Using further default parameters, the above and below ground C and N inputs to the soil from crop residues are estimated for each crop type ( $i$ ) planted:

$$C_{NIag,i} = C_{ag-on,i} \cdot c_{an,i} \quad (\text{Equation 35})$$

$$C_{NIbg,i} = C_{bg,i} \cdot c_{bn,i} \cdot c_{30,i} \quad (\text{Equation 36})$$

$$C_{CIag,i} = C_{ag-on,i} \cdot c_{ac,i} \quad (\text{Equation 37})$$

$$C_{CIbg,i} = C_{bg,i} \cdot c_{bc,i} \cdot c_{30,i} \quad (\text{Equation 38})$$

Where:  $C_{NIag,i}$  is the mass of above ground crop N inputs for crop type  $i$  (t N/ha)

$c_{an,i}$  is the crop above-ground residue N content for crop type  $i$  (g N/g DM)

$C_{NIbg,i}$  is the mass of below ground crop N inputs for crop type  $i$  (t N/ha)

$c_{bn,i}$  is the crop below-ground residue N content for crop type  $i$  (g N/g DM)

$c_{30,i}$  is the fraction of below-ground residues which can be found in the top 0-30 cm of soil for crop type  $i$  (t  $C_{bg}$  0-30 cm/ total t  $C_{bg}$ )

$C_{CIag,i}$  is the mass of above ground crop C inputs for crop type  $i$  (t C/ha)

$c_{ac,i}$  is the crop above-ground residue C content for crop type  $i$  (g C/g DM)

$C_{CIbg,i}$  is the mass of below ground crop C inputs for crop type  $i$  (t C/ha)

$c_{bc,i}$  is the crop below-ground residue C content for crop type  $i$  (g C/g DM)

Finally, the crop on and off farm above ground residues, below ground residues, above and below ground C and N inputs are summed for all planted crop types/species ( $i$ ):

$$C_{ag-off} = \sum_{i=1}^n C_{ag-off,i} \quad (\text{Equation 39})$$

$$C_{ag-on} = \sum_{i=1}^n C_{ag-on,i} \quad (\text{Equation 40})$$

$$C_{bg} = \sum_{i=1}^n C_{bg,i} \quad (\text{Equation 41})$$

$$C_{NIag} = \sum_{i=1}^n C_{NIag,i} \quad (\text{Equation 42})$$

$$C_{NIbg} = \sum_{i=1}^n C_{NIbg,i} \quad (\text{Equation 43})$$

$$C_{CIag} = \sum_{i=1}^n C_{CIag,i} \quad (\text{Equation 44})$$

$$C_{CIbg} = \sum_{i=1}^n C_{CIbg,i} \quad (\text{Equation 45})$$

These annual totals are assumed to be the same for every year of the model run, and are passed to the biomass burning model (section 7) before they are passed to the soil model (Section 8) and the emissions from plant N input model (section 9).

## 6. External organic soil inputs

### 6.1 External organic soil input description

Organic inputs such as litter, mulch or manure originating from outside of the boundaries of the intervention area can be added to fields or plots for additional soil inputs or as fertiliser. If applicable, the additional biomass, carbon, and nitrogen inputs to soils from external organic inputs are calculated using a Tier-1 type approach. External inputs are included in the model, but are not included in C accounting for projects (see the SHAMBA methodology), as external inputs to the soils reduce inputs outside of the project boundary and are therefore a source of leakage. For project accounting, external inputs are omitted from further calculations. The SHAMBA model has maintained external soil inputs here for added functionality.

To include external organic inputs in the model calculations, data on the amount of added material, and the C and N content are required (Table 5). The default values provided assume organic inputs are sourced from the surrounding forests/woodlands, or originate from woody tree/shrubs (i.e. tree litter). Other inputs, such as manure or mulch, can be included if data are available. However, further data on the decomposability of the organic matter is required (see section 8), and the emission and combustions factors when fire occurs (see section 7). The years when external organic inputs are added to fields can be estimated using a frequency interval (where a frequency interval of 2 would

mean addition of external inputs occurred every other year), or specified in known years of addition. The external organic soil inputs are incorporated into the emissions from biomass burning model (section 7), the soil C model (Section 8), and the emissions from fertiliser use (section 10).

The model assumes:

- The external inputs are organic
- The C and N content of the inputs are known, and do not differ between years

## 6.2 External organic input parameterisation

The data required to calculate the total external inputs of C and N to the field are outlined in table 5.

**Table 5:** Parameter requirements to calculate external inputs to soils for each organic input type

Parameter	Symbol	Units	Default value
Addition of external organic input to field in year $y$ ?	$a_{f,y}$	1 or 0 (Yes/No)	<i>User must provide this</i>
Mass of external input added in year $y$	$A_y$	t DM/ ha	<i>User must provide this</i>
N content of input	$a_n$	g N/ g DM	0.018
C content of input	$a_c$	g C/ g DM	0.5

## 6.3 External organic input calculations

Using data as outlined in table 5, external organic C and N inputs are calculated on a yearly time step for each organic input type ( $i$ ):

$$A_{CI,i,y} = A_{i,y} \cdot a_{f,i,y} \cdot a_{c,i} \quad (\text{Equation 46})$$

$$A_{NI,i,y} = A_{i,y} \cdot a_{f,i,y} \cdot a_{n,i} \quad (\text{Equation 47})$$

Where:  $A_{CI,i,y}$  is the mass of external C inputs for input type  $i$  in year  $y$  (t C/ha)

$A_{NI,i,y}$  is the mass of external N inputs for input type  $i$  in year  $y$  (t N/ha)

$A_{i,y}$  is the dry mass of external input added to the field for input type  $i$  in year  $y$  (t DM/ha)

$a_{f,i,y}$  is if external inputs type  $i$  are added in year  $y$  (1 if yes, 0 if no)

$a_{c,i}$  is the C content for input type  $i$  (g C/ g DM)

$a_{n,i}$  is the N content for input type  $i$  (g N/ g DM)

Total C and N inputs from all organic input types ( $i$ ) are then the sum of all inputs in each year:

$$A_{CI,y} = \sum_{i=1}^n A_{CI,y,i} \quad (\text{Equation 48})$$

$$A_{NI,y} = \sum_{i=1}^n A_{NI,y,i} \quad (\text{Equation 49})$$

Where:  $A_{CI,y}$  is the total mass of external C inputs in year  $y$  (t C/ha)

$A_{NI,y}$  is the mass of external N inputs in year  $y$  (t N/ha)

## 7. Emissions due to biomass burning

The following section details the calculation of emissions from biomass burning.

## 7.1 Biomass burning model description

Non-CO<sub>2</sub> emissions from burning of biomass can be estimated using a Tier-1 type approach, as outlined by IPCC (2006) and SALM<sup>7</sup>. The model accounts for emissions from burning of above ground crop residues, on and off the field, above ground tree litter and external organic input biomass on the field. Standing live trees, below ground biomass, or litter from previous years are not burnt during fires. The amount of crop biomass available to burn on and off the field (t DM/ha) is calculated from the crop model (section 5). The amount of biomass from tree litter available to burn on the field is calculated from the woody biomass model (section 4), and the biomass available to burn on the field from external inputs are calculated (section 6). Using default values of combustion factors and emission factors given by the IPCC (2006) for burning of crop residues and tree litter, we can calculate the emissions from burning of biomass from each of these sources. Combustion and emission factors for external organic inputs will need to be specified if not similar to tree litter.

In order to assess when fires will occur in the field, if at all, fire occurrence is either estimated using a fire return interval (where a fire return interval of 5 causes fields to be burnt every 5 years), or fire occurrence can be specified to particular years when fire was known to occur. When a fire occurs the biomass from above-ground sources on the field are combusted by an amount equivalent to the combustion factors, creating emissions from biomass burning and reducing the C inputs to the soil carbon model (section 8), the N inputs for the calculation of emissions from plant N inputs (section 9), and reducing N inputs from organic fertilisers and associated emissions (section 10). Emissions from biomass burning (tCO<sub>2e</sub>/ha) are calculated for baseline and intervention scenarios separately.

The model assumes:

- Only above-ground crop residues, tree litter and external organic inputs are burnt
- Only biomass added in the year of a field fire are burnt
- A field fire occurs at the end of the year, or post-harvest
- Removed crop residues are burnt annually post-harvest
- Live trees are not killed in fires, and live tree biomass is not burnt
- Soil organic C stocks are not directly affected by fire

## 7.2 Biomass burning model parameterisation

The data requirements for the biomass burning model are outlined in table 6.

**Table 6:** Parameter requirements to calculate emissions due to biomass burning for each scenario

Parameter	Symbol	Units	Default value
Fire occurrence in field in year $y$	$ff_y$	1 or 0 (Yes/No)	<i>User must provide this</i>
Are removed crop residues burnt else-where?	$C_{burn}$	1 or 0 (Yes/No)	<i>User must provide this</i>
Mass of above-ground crop residues removed from the field in year $y$	$C_{ag-off,y}$	t DM/ha	<i>Calculated in section 5</i>

<sup>7</sup> <http://v-c-s.org/sites/v-c-s.org/files/VM0017%20SALM%20Methodolgy%20v1.0.pdf>

Mass of above-ground crop residues available for fire in the field in year $y$	$C_{ag-on,y}$	t DM/ha	<i>Calculated in section 5</i>
Mass of above-ground tree litter available for fire in the field in year $y$	$T_{DMag,y}$	t DM/ha	<i>Calculated in section 4</i>
Mass of external organic input available for fire in the field in year $y$	$A_y$	t DM/ha	<i>Calculated in section 6</i>
Combustion factor for crop residues burned	$cf_c$	unitless	0.8
Combustion factor for tree litter burned	$cf_f$	unitless	0.74
Combustion factor for external input type $i$ burned	$cf_a$	unitless	Assumed the same as for tree litter
Emission factor for the production of methane for crop residues burned	$ef_{CH_4,c}$	g CH <sub>4</sub> /kg	2.7
Emission factor for the production of methane for tree litter burned	$ef_{CH_4,f}$	g CH <sub>4</sub> /kg	6.8
Emission factor for the production of methane for external input burned	$ef_{CH_4,a}$	g CH <sub>4</sub> /kg	Assumed the same as for tree litter
Emission factor for the production of nitrous oxide for crop residues burned	$ef_{N_2O,c}$	g N <sub>2</sub> O/kg	0.07
Emission factor for the production of nitrous oxide for tree litter burned	$ef_{N_2O,f}$	g N <sub>2</sub> O/kg	0.20
Emission factor for the production of nitrous oxide for external input type $i$ burned	$ef_{N_2O,a}$	g N <sub>2</sub> O/kg	Assumed the same as for tree litter
Global warming potential of methane for 100 years accounting period	$gwp_{CH_4}$	t CO <sub>2e</sub> /t gas	21
Global warming potential of nitrous oxide for 100 years accounting period	$gwp_{N_2O}$	t CO <sub>2e</sub> /t gas	310

### 7.3 Biomass burning model calculations

To calculate the emissions from burning of biomass the following equation is used:

$$E_{BB,y} = \left[ \left( (C_{ag-off} \cdot c_{burn}) + (C_{ag-on,y} \cdot ff_y) \right) \cdot cf_c \cdot (ef_{CH_4,c} \cdot gwp_{CH_4} + ef_{N_2O,c} \cdot gwp_{N_2O}) + \right. \\ \left. (T_{DMag,y} \cdot ff_y) \cdot cf_f \cdot (ef_{CH_4,f} \cdot gwp_{CH_4} + ef_{N_2O,f} \cdot gwp_{N_2O}) + \right. \\ \left. \sum_{i=0}^n (A_{i,y} \cdot ff_y) \cdot cf_{a,i} \cdot (ef_{CH_4,a,i} \cdot gwp_{CH_4} + ef_{N_2O,a,i} \cdot gwp_{N_2O}) \right] 10^{-3}$$

(Equation 50)

Where:  $E_{BB,y}$  is the emissions from burning above-ground biomass in year  $y$  (tCO<sub>2e</sub>/ha)

$C_{ag-off}$  is the mass of above ground crop residues removed from the field annually (t DM/ha)

$c_{burn}$  is if crop residues removed from the field are burnt annually (1 if yes, 0 if no)

$C_{ag-on,y}$  is the mass of above-ground crop residues available to burn in the field in year  $y$  (t DM/ha)

$ff_y$  is if fire occurs in the field in year  $y$  (1 for fire, 0 for no fire)

$cf_c$  is the combustion factor appropriate for burning crop residues (IPCC, 2006)

$ef_{CH_4,c}$  and  $ef_{N_2O,c}$  are the emission factors for the production of methane and nitrous oxide when burning crop residues (g CH<sub>4</sub>/kg and g N<sub>2</sub>O/kg, respectively) (IPCC, 2006)

$gwp_{CH_4}$  and  $gwp_{N_2O}$  are the global warming potentials (t CO<sub>2e</sub>/t gas) of CH<sub>4</sub> and N<sub>2</sub>O, respectively (IPCC, 2006)

$T_{DMag,y}$  is the mass of above ground tree litter available to burn in the field in year  $y$  (t DM/ha)

$cf_f$  is the combustion factor appropriate for burning litter from forests/woodlands (IPCC, 2006)

$ef_{CH_4,f}$  and  $ef_{N_2O,f}$  are the emission factors for the production of methane and nitrous oxide when burning tree litter (g CH<sub>4</sub>/kg and g N<sub>2</sub>O/kg, respectively) (IPCC, 2006)

$A_{i,y}$  is the mass of external organic input type  $i$  available to burn in the field in year  $y$  (t DM/ha)

$cf_{a,i}$  is the combustion factor appropriate for burning external biomass input type  $i$

$ef_{CH_4,a,i}$  and  $ef_{N_2O,a,i}$  are the emission factors for the production of methane and nitrous oxide when burning external input type  $i$  (g CH<sub>4</sub>/kg and g N<sub>2</sub>O/kg, respectively)

## 7.4 Soil inputs reduced by fire

When fire occurs biomass is combusted, decreasing the above ground C and N inputs to the soils in that year. Therefore, soil inputs need to be corrected for effects of fire before they can pass to the soil model (section 8), the emissions from plant N input model (section 9) and the fertiliser emissions model (section 10).

Using the outputs from the woody biomass model ( $T_{NIag,y}$ ,  $T_{NIbg,y}$ ,  $T_{CIag,y}$ ,  $T_{CIbg,y}$ ), the crop model ( $C_{NIag}$ ,  $C_{NIbg}$ ,  $C_{CIag}$ ,  $C_{CIbg}$ ), and external organic inputs ( $A_{NI,y}$ ,  $A_{CI,y}$ ), the soil inputs are corrected for losses to fire in the above-ground fraction using the respective combustion factors:

$$T_{NItotal,y} = \left( T_{NIag,y} - (T_{NIag,y} \cdot ff_y \cdot cf_f) \right) + T_{NIbg,y} \quad (\text{Equation 51})$$

$$T_{CItotal,y} = \left( T_{CIag,y} - (T_{CIag,y} \cdot ff_y \cdot cf_f) \right) + T_{CIbg,y} \quad (\text{Equation 52})$$

$$C_{NItotal,y} = \left( C_{NIag} - (C_{NIag} \cdot ff_y \cdot cf_c) \right) + C_{NIbg} \quad (\text{Equation 53})$$

$$C_{CItotal,y} = \left( C_{CIag} - (C_{CIag} \cdot ff_y \cdot cf_c) \right) + C_{CIbg} \quad (\text{Equation 54})$$

$$A_{NItotal,y} = \sum \left( A_{NI,i,y} - (A_{NI,i,y} \cdot ff_y \cdot cf_{a,i}) \right) \quad (\text{Equation 55})$$

$$A_{CItotal,y} = \sum \left( A_{CI,i,y} - (A_{CI,i,y} \cdot ff_y \cdot cf_{a,i}) \right) \quad (\text{Equation 56})$$

Where:  $T_{NItotal,y}$  is the total N inputs to soils from trees in year  $y$  (t N/ha)

$T_{CItotal,y}$  is the total C inputs to soils from trees in year  $y$  (t C/ha)

$C_{NItotal,y}$  is the total N inputs to soils from crops in year  $y$  (t N/ha)

$C_{CItotal,y}$  is the total C inputs to soils from crops in year  $y$  (t C/ha)

$A_{NItotal,y}$  is the total N inputs to soils from all external organic inputs in year  $y$  (t N/ha)

$A_{CItotal,y}$  is the total C inputs to soils from all external organic inputs in year  $y$  (t C/ha)

## 8. Modelling changes in soil organic carbon

This section details the parameterisation and use of the RothC model in SHAMBA, which is used to estimate soil organic carbon changes for baseline and intervention scenarios. The following sections describe the method.

### 8.1 SOC model overview

Soil organic carbon (SOC) stocks and changes, under baseline and intervention scenarios, are calculated using the RothC soil carbon model (Coleman and Jenkinson 1999), and implemented in Python (v 2.6.6). In SHAMBA, the RothC model runs on a yearly time step for the top 0-30 cm of soil, using annual C inputs (calculated in sections 4-7), modelling the turnover of organic carbon allowing for the effects of soil type, temperature, moisture content and plant cover on the turnover process.

As the equations for RothC are outlined in the user manual<sup>8</sup>, we describe the use of RothC in SHAMBA and show equations where they differ from RothC. Full model description and assumptions are described in the following section.

### 8.2 SOC model parameterisation

Some of the data needed to parameterise and drive the soil carbon model in SHAMBA are shown in Table 7. The data can either be entered from local measurements or other relevant data sources, or use default values (see Appendix 1).

**Table 7: Parameters and drivers required by the SHAMBA soil carbon model for each scenario**

Parameter/ driver	Symbol	Units	Default value
Monthly mean temperature	$t$	Degrees C	CRU TS 3.10 global dataset <sup>9</sup> , based on location
Monthly evapotranspiration*	$et$	mm	
Monthly rainfall	$p$	mm	
Soil carbon content at equilibrium	$soc_f$	t C/ha	Assumed to be 25 % higher than initial SOC stocks
Soil carbon content at start of intervention (initial)	$soc_{y=0}$	t C/ha	Harmonized World Soil Database <sup>10</sup> , based on location
Soil clay content at start of intervention (initial)	$clay_{y=0}$	%	
Is land covered or bare in each month of the year?	$sc$	Yes/No	<i>User must provide this</i>
Decomposable fraction of crop residue plant material	$dpm_c$	decomposable mass/ total mass	0.59
Resistant fraction of crop residue plant material	$rpm_c$	resistant mass/ total mass	0.41
Decomposable fraction of tree litter plant material	$dpm_r$	decomposable mass/ total mass	0.20
Resistant fraction of tree litter plant material	$rpm_r$	resistant mass/ total mass	0.80

<sup>8</sup> [http://www.rothamsted.ac.uk/aen/carbon/mod26\\_3\\_win.pdf](http://www.rothamsted.ac.uk/aen/carbon/mod26_3_win.pdf)

<sup>9</sup> University of East Anglia Climatic Research Unit (CRU). [Phil Jones, Ian Harris]. CRU TS3.10: Climatic Research Unit (CRU) Time-Series (TS) Version 3.10 of High Resolution Gridded Data of Month-by-month Variation in Climate (Jan. 1901 - Dec. 2009). Available at: [http://badc.nerc.ac.uk/view/badc.nerc.ac.uk\\_\\_ATOM\\_\\_ACTIVITY\\_fe67d66a-5b02-11e0-88c9-00e081470265](http://badc.nerc.ac.uk/view/badc.nerc.ac.uk__ATOM__ACTIVITY_fe67d66a-5b02-11e0-88c9-00e081470265)

<sup>10</sup> FAO/IIASA/ISRIC/ISSCAS/JRC, 2009. *Harmonized World Soil Database (version 1.1)*. Food and Agriculture Organization of the United Nations, Rome, Italy and IIASA, Laxenburg, Austria. Available at: <http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/index.html>




Decomposable fraction of the un-disturbed plant inputs	$dpm_e$	decomposable mass/ total mass	0.2 (Assumed the same as $dpm_i$ )
Resistant fraction of the un-disturbed plant inputs	$rpm_e$	resistant mass/ total mass	0.8 (Assumed the same as $rpm_i$ )
Decomposable fraction of external organic input	$dpm_a$	decomposable mass/ total mass	0.2 (Assumed the same as $dpm_i$ )
Resistant fraction of external organic input	$rpm_a$	resistant mass/ total mass	0.8 (Assumed the same as $rpm_i$ )
Humified fraction of external organic input	$hum_a$	humified mass/ total mass	0

\*If evapotranspiration is not available, pan evaporation (e) can be used instead

### 8.3 Model structure and SOC partitioning

The structure of RothC (Fig. 4) is such that the organic inputs (t C/ha) to the soil are split into four active pools; the decomposable plant material (DPM), resistant plant material (RPM), microbial biomass (BIO) and humified organic matter (HUM). Each pool then decomposes with its own characteristic rate. A small amount of inert organic matter (IOM) is present, but remains resistant to decomposition. SOC is defined as the total of all the organic carbon pools (DPM, RPM, BIO, HUM and IOM).



**Fig.4:** Structure of RothC, showing the partitioning of organic inputs to soil into the four active compartments, each decomposing at a specific rate (figure modified from Coleman & Jenkinson (1999), p.8).

The plant inputs or organic matter inputs to the soil pool are modelled for crop ( $C_{Ctotal}$ ) and tree ( $T_{Ctotal}$ ) inputs, and calculated for each external organic input type ( $A_{Ctotal}$ ) mediated by losses due to fire (section 7.4). Each year the incoming inputs are split between DPM and RPM fractions, depending on a DPM/RPM ratio. This ratio is based on the default values given by RothC for agricultural crop residues (59% DPM, 41% RPM) and tropical deciduous woodland litter (20% DPM, 80% RPM), and uses the DPM/RPM ratio provided by the user for each external organic input type. RothC provides a default value for manure (49% DPM, 49% RPM, 2% HUM), including a humified

fraction to account for the fact that manure is more decomposed than plant material is. Any other external organic inputs which are not from woodland litter or manure needs to have a specified DPM/RPM and HUM ratio in order to be incorporated into the soil model (Table 7).

The model uses a weighted DPM/RPM (and HUM) ratio calculated from the fraction of total inputs originating from crops, trees and external organic inputs:

$$dpm_y = (dpm_c \cdot f_{c,y}) + (dpm_f \cdot f_{f,y}) + \sum_{i=0}^n (dpm_{a,i} \cdot f_{a,i,y}) \quad (\text{Equation 57})$$

$$rpm_y = (rpm_c \cdot f_{c,y}) + (rpm_f \cdot f_{f,y}) + \sum_{i=0}^n (rpm_{a,i} \cdot f_{a,i,y}) \quad (\text{Equation 58})$$

$$hum_y = \sum_{i=0}^n (hum_{a,i} \cdot f_{a,i,y}) \quad (\text{Equation 59})$$

Where:  $dpm_y$  is the weighted fraction of the total inputs which are decomposable in year  $y$  (t decomposable/ total t input)

$dpm_c$  is the decomposable fraction of crop residue inputs (t decomposable crop input/ total t crop input)

$f_{c,y}$  is the fraction of total inputs which are from crops in year  $y$  (t crop inputs/ total t inputs)

$dpm_f$  is the decomposable fraction of tree litter inputs (t decomposable tree input/ total t tree input)

$f_{f,y}$  is the fraction of total inputs which are from trees in year  $y$  (t tree inputs/ total t inputs)

$dpm_{a,i}$  is the decomposable fraction of external organic inputs of type  $i$  (t decomposable organic input/ total t organic input)

$f_{a,i,y}$  is the fraction of total inputs which are from external organic input of type  $i$  in year  $y$  (t organic input/ total t inputs)

$rpm_y$  is the weighted fraction of the total inputs which are resistant in year  $y$  (t resistant/ total t input)

$rpm_c$  is the resistant fraction of crop residue inputs (t resistant crop input/ total t crop input)

$rpm_f$  is the resistant fraction of tree litter inputs (t resistant tree input/ total t tree input)

$rpm_{a,i}$  is the resistant fraction of external organic inputs of type  $i$  (t resistant organic input/ total t organic input)

$hum_y$  is the weighted fraction of total inputs which are humified in year  $y$  (t humified/ total t input)

$hum_{a,i}$  is the humified fraction of external organic input of type  $i$  (t humified organic input/ total t organic input)

Once the inputs have been split between decomposable and resistant fractions, the DPM and RPM fractions decompose further to form BIO, HUM and CO<sub>2</sub>. The proportion that goes to CO<sub>2</sub> and to BIO+HUM is determined by the soil clay content. The BIO+HUM is then split between BIO and HUM

using set partitioning coefficients (46% to BIO, 54% to HUM). BIO and HUM decompose to form more CO<sub>2</sub>, BIO and HUM, and so on (Fig. 4).

The inert organic matter fraction (IOM) remains constant, as no decomposition occurs in this fraction. IOM is calculated using the equation from Falloon et al. (1998):

$$IOM = 0.049 \cdot soc_f^{1.139} \quad (\text{Equation 60})$$

Where: *IOM* is the inert organic matter fraction (t C/ha)

*soc<sub>f</sub>* is the total organic carbon (t C/ha) of soils when at equilibrium (see sections 8.1.2)

Each active compartment of soil organic carbon (DPM, RPM, BIO, HUM) decays at a rate determined by individual decomposition rate constants (*k*) and a rate modifying factor (*r*) determined by temperature, soil moisture and soil cover (see the RothC<sup>11</sup> description for full details of calculations). The rate modifying factor (*r*) is calculated using data on soil cover and monthly climate variables (Table 2). The model default uses a global climate dataset to extract climate variables based on a specified geographical location. The climate dataset is the CRU TS 3.10<sup>12</sup> high resolution (0.5°) month-by-month global climate dataset (Harris et al. 2013), where climate variable are monthly means calculated from 1960-2009. If local measurements of monthly climate variables are available, users can enter these values into the model instead of using the defaults. The model currently assumes climate variables do not change between years or scenarios. Therefore, the rate modifier (*r*) will only change if the monthly soil cover changes (i.e. if soil is bare or covered in each month) between years or baseline and intervention scenarios.

## 8.4 SOC model initialisation

Before we can model baseline and intervention SOC changes, the soil model needs to be initialised to the *y=0* soil conditions if it is to accurately simulate future changes to soil carbon. Soils take decades to reach a steady state after changes to inputs or output fluxes of carbon. Therefore, unless land management has been consistent over ~30 years, which is rare and usually unknown, the soils are unlikely to be in equilibrium and may be losing or gaining carbon. The SHAMBA model allows the simulation of situations where SOC is changing rapidly, as it often is in situations where interventions are implemented. This means there is no assumption that the SOC is in equilibrium at the start of the interventions.

To initialise the SOC model, data on SOC stocks at equilibrium (*soc<sub>i</sub>*) and SOC stocks and clay content at the start of intervention activities (*soc<sub>y=0</sub>*, *clay<sub>y=0</sub>*) are required (Table 7). The model default uses the Harmonized World Soil Database<sup>13</sup> (HWSD) (FAO/IIASA/ISRIC/ISSCAS/JRC 2009) to estimate initial SOC stocks and clay content at the start of interventions, based on a geographical location. We assume the value given by HWSD is appropriate for disturbed soils, such as agricultural

<sup>11</sup> [http://www.rothamsted.ac.uk/aen/carbon/mod26\\_3\\_win.pdf](http://www.rothamsted.ac.uk/aen/carbon/mod26_3_win.pdf)

<sup>12</sup> [http://badc.nerc.ac.uk/view/badc.nerc.ac.uk\\_\\_ATOM\\_\\_ACTIVITY\\_fe67d66a-5b02-11e0-88c9-00e081470265](http://badc.nerc.ac.uk/view/badc.nerc.ac.uk__ATOM__ACTIVITY_fe67d66a-5b02-11e0-88c9-00e081470265)

<sup>13</sup> <http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/>

fields, based on the assumption that soil measurements used in the HWSD are more likely to come from disturbed soils. The HWSD provides soil characteristics for several different soil types at any one location. Therefore, the initial SOC stock and clay content values are calculated based on a weighted mean at the given location. The model default assumes SOC under equilibrium conditions are 25 % higher than the value given by HWSD (based on Guo and Gifford 2002, Don et al. 2011). The assumption is that the land was wooded before disturbance and that woodland or forest cover represent a pre-disturbance state where SOC was in equilibrium. If defaults are not applicable, local data on equilibrium and initial SOC stocks and soil clay content should be entered into the model instead.


The following procedure (illustrated in Figure 5) allows the simulation of non-steady state conditions:

1. SOC levels at steady state are simulated to estimate the pre-disturbance state of the SOC pools (i.e. the DPM, RPM, BIO, HUM and IOM) at equilibrium.
2. The change from this undisturbed state to the current state is modelled by imposing the baseline land management activities upon the undisturbed state until initial SOC levels observed in e.g. the HWSD are reached.
3. The distribution of soil carbon in the different modelled pools (i.e. the DPM, RPM, BIO and HUM) at the initial state can then be used to initialise the model prior to modelling of the baseline and intervention scenarios.

Step 1 requires an estimate of SOC stock at equilibrium, appropriate for the undisturbed soils (see above). The model uses a parameter search to find the mass of annual plant inputs to the soils, under the assumed forest/woodland cover, which allows SOC to reach the equilibrium value over 10,000 years. A DPM/RPM ratio of 0.25 is used as default and is appropriate for most deciduous and tropical woodlands (Coleman and Jenkinson 1999). IOM is based on SOC at equilibrium (Eq. 60), and remains constant throughout the model run.

Step 2 is to model the soil inputs to simulate the baseline scenario until the size of the total SOC pool equals initial SOC stocks (i.e SOC stocks at the start of interventions), in order to simulate the change in SOC pools following disturbance.

Step 3, at the time point where initial SOC stocks are reached, the relevant values for the carbon pools (DPM, RPM, BIO, HUM) are extracted and used as initial conditions for baseline and intervention scenarios.



**Fig. 5:** Schematic of soil model initialisation. The soil model is optimised to equilibrium conditions under woodland or forest conditions before modelling changes in SOC following conversion to baseline land use. The effects of the relevant interventions can then be modelled using initial soil carbon pool values derived at  $y = 0$ .

### 8.3 SOC changes

The soil organic carbon model initialises the SOC pools using data on equilibrium and initial SOC stocks and clay content. Once initialised, the baseline and intervention scenarios are modelled separately with modelled organic inputs, weighted DPM/RPM ratios, and climate and soil cover data. Total SOC stocks (i.e. DPM, RPM, HUM, BIO and IOM) in each year are modelled in units of t C/ha. Emissions from changes in total SOC stocks for each year (Eq.1 & 2) are calculated by SHAMBA for baseline and intervention scenarios:

$$E_{SOy} = (SOC_{y-1} - SOC_y) \cdot mw_{CO2} \quad (\text{Equation 61})$$

Where:  $E_{SOy}$  is the emissions from changes in SOC stocks (tCO<sub>2e</sub>/ha), where a negative is an uptake

$SOC_{y-1}$  is the SOC stocks in the year before  $y$  (t C/ha)

$SOC_y$  is the SOC stocks in the year  $y$  (t C/ha)

$mw_{CO2}$  is the ratio of molecular weights of CO<sub>2</sub> and C (44/12)

## 8.4 SOC model limitations

The model has a few limitations in that it cannot: model the effects of tillage on SOC stocks, include the impacts of fire on SOC, or incorporate other organic inputs such as charcoal from fire or bio-char. Furthermore, it assumes that the initialisation process is representative of past conditions. If soils have experienced several disturbance and recovery periods since equilibrium, the model initialisation may not be appropriate.

The model assumes:

- Soils are not waterlogged or seasonally flooded
- Only plant and organic inputs to the soil enter the soil carbon pool
- Impacts of fire and tillage on SOC are not considered
- Soils are at least 30 cm deep
- SOC losses through erosion or leakage are minimal
- Climate does not change over the model run

Due to the use of a yearly time step in the soil model, and not a monthly one as per RothC, the total modelled SOC stocks differ slightly between SHAMBA and the Rothamsted Carbon Model (v 26.3)<sup>14</sup>. In a test, based on a range of scenarios the difference in SOC at the end of the simulations between RothC (v 26.3) and the annual timestep implementation in SHAMBA never differed by more than 0.5 t C/ha.

## 9. Emissions due to nitrogen inputs from plants

This section details the calculations for estimating the emissions from plant nitrogen inputs to the soils.

### 9.1 Plant nitrogen input emissions description

Direct nitrous oxide emissions from plant nitrogen inputs to soils can be estimated using a Tier-1 type approach, as described in the SALM methodology. As all plants can fix nitrogen and contain some N in their biomass, we calculate emissions from all crop and tree N inputs to the soil, adopting a conservative approach to the calculation of emissions from this source. This differs to other approaches (e.g. SALM), which only account for emissions from the planting of N-fixing plants. The emissions from plant N inputs do not include N inputs from external organic inputs (section 6), as they are not always from plant sources, and are instead included in emissions from fertilisers (section 10).

The model assumes:

- All plant N inputs from crops and trees are included in emissions calculations

### 9.2 Plant nitrogen input emission parameterisation

The data required to estimate N emissions from plant inputs are outlined in Table 8.

<sup>14</sup> <http://www.rothamsted.ac.uk/sustainable-soils-and-grassland-systems/rothamsted-carbon-model-rothc>

**Table 8:** Parameters required for calculating emissions due to plant N inputs for each scenario

Parameter	Symbol	Units	Default value
Mass of total crop N inputs in year $y$	$C_{Ntotal,y}$	t N/ha	Calculated in section 5 and 7.4
Mass of total tree N inputs in year $y$	$T_{Ntotal,y}$	t N/ha	Calculated in section 4 and 7.4
Emission factor for emissions of N <sub>2</sub> O-N from N inputs	$ef_N$	t N <sub>2</sub> O-N/ t N input	0.01
The ratio of molecular weights of N <sub>2</sub> O and N <sub>2</sub>	$mw_{N2O}$	unitless	44/28
Global warming potential for N <sub>2</sub> O for 100 years accounting period	$gwp_{N2O}$	t CO <sub>2e</sub> /t N <sub>2</sub> O	310

### 9.3 Plant nitrogen input emission calculations

To calculate the emissions from plant N inputs the following method, based on the SALM methodology, was used:

$$E_{NI,y} = (C_{Ntotal,y} + T_{Ntotal,y}) \cdot ef_N \cdot mw_{N2O} \cdot gwp_{N2O} \quad (\text{Equation 62})$$

Where:  $E_{NI,y}$  is the emissions due to the N inputs to soils from plants in year  $y$  (tCO<sub>2e</sub>/ha)

$C_{Ntotal,y}$  is the total crop nitrogen inputs in year  $y$  (t N/ha)

$T_{Ntotal,y}$  is the total tree nitrogen inputs in year  $y$  (t N/ha)

$ef_N$  is the emission factor for emissions of N<sub>2</sub>O-N from N inputs (t N<sub>2</sub>O-N/t N input) (IPCC, 2006)

$mw_{N2O}$  is the ratio of molecular weights of N<sub>2</sub>O and N (44/28)

$gwp_{N2O}$  is the global warming potential for N<sub>2</sub>O over 100 years accounting period (tCO<sub>2e</sub>/ tN<sub>2</sub>O) (IPCC, 2006)

## 10. Emissions due to fertiliser use

This section details how emissions from fertiliser use are calculated.

### 10.1 Emissions from fertiliser use description

If applicable, emissions from the use of synthetic and/or organic nitrogen fertilisers are calculated using the CDM A/R Working Group Tool<sup>15</sup> *Estimation of direct nitrous oxide emission from nitrogen fertilisation (version 01)*. Using emission factors and volatilisation values as provided by the IPCC Guidelines (2006), emissions from organic and synthetic fertilisers are calculated. All external organic N inputs (section 6) are included as organic N fertilisers, and are assumed to have the same emission and volatilisation values as defined by the IPCC (2006) for organic fertilisers.

The model assumes:

- All external organic inputs (section 6) can be classified as organic fertilisers

<sup>15</sup> <http://cdm.unfccc.int/methodologies/ARmethodologies/tools/ar-am-tool-07-v1.pdf>

- All external organic inputs have similar emission and volatilisation parameters

## 10.2 Emissions from fertiliser use parameterisation

The data required to calculate the emissions from fertiliser use are outlined in table 9.

**Table 9:** Parameter requirements to calculate emissions due to fertiliser use

Parameter	Symbol	Units	Default value
Synthetic fertiliser application in year $y$	$sf_y$	1 or 0 (Yes/No)	<i>User must provide this</i>
Mass of synthetic fertiliser applied in year $y$	$S_y$	t/ha	<i>User must provide this</i>
Nitrogen content of synthetic fertiliser	$s_n$	g N/ g fertiliser	<i>User must provide this</i>
Mass of N inputs from external organic inputs in year $y$	$A_{Ntotal,y}$	t N/ha	Calculated in section 6 and 7.4
Emission factor for emissions of N <sub>2</sub> O-N from N inputs	$ef_N$	t N <sub>2</sub> O-N/t N input	0.01
Fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for synthetic fertilisers	$v_s$	(t NH <sub>3</sub> -N + NO <sub>x</sub> -N)/ t N applied	0.1
Fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for organic fertilisers	$v_o$	(t NH <sub>3</sub> -N + NO <sub>x</sub> -N)/ t N applied	0.2
Ratio of molecular weights of N <sub>2</sub> O and N	$mw_{N2O}$	unitless	44/28
Global warming potential of N <sub>2</sub> O for 100 years accounting period	$gwp_{N2O}$	t CO <sub>2e</sub> /t gas	310

## 10.3 Emissions from fertiliser use calculations

The direct nitrous oxide emissions from N fertilisation can be estimated as follows:

$$E_{NF,y} = \left[ (S_y \cdot sf_y \cdot s_n \cdot (1 - v_s)) + (A_{Ntotal,y} \cdot (1 - v_o)) \right] ef_N \cdot mw_{N2O} \cdot gwp_{N2O} \quad (\text{Equation 63})$$

Where:  $E_{NF,y}$  is the direct N<sub>2</sub>O emission as a result of nitrogen application in year  $y$  (tCO<sub>2e</sub>/ha)

$S_y$  is the mass of synthetic nitrogen fertiliser applied in year  $y$  (t /ha)

$sf_y$  is if synthetic fertiliser was applied in year  $y$  (1 if yes, 0 if no)

$s_n$  is the N content of synthetic fertiliser (g N/ g fertiliser)

$v_s$  is the fraction that volatilises as NH<sub>3</sub> and NO<sub>x</sub> for synthetic fertilisers (IPCC, 2006)

$A_{Ntotal,y}$  is the total mass of N inputs from external organic inputs (t N/ha)

$v_o$  is the fraction that volatilises as NH<sub>3</sub> and NO<sub>x</sub> for organic fertilisers (IPCC, 2006)

$ef_N$  is the emission factor for emissions of N<sub>2</sub>O-N from N inputs (t N<sub>2</sub>O-N /t N input) (IPCC, 2006)

$mw_{N2O}$  is the ratio of molecular weights of N<sub>2</sub>O and N

$gwp_{N2O}$  is the global warming potential of N<sub>2</sub>O (IPCC, 2006)



## 11. Future model developments

Here we suggest ways the model could be improved and further developed in subsequent versions, and outline some of the limitations of the SHAMBA model v1.0. Specific improvements are outlined in context of each model section.

### *SOC model*

- The SOC model runs on an annual time step, but RothC is designed to run on a monthly time step, causing some small errors in total SOC changes. Future versions should run RothC on a monthly time step, using annual plant inputs divided evenly over the year, or using plant inputs calculated for every month. This will require more data inputs, but the accuracy may be increased as a result.

### *Biomass model*

- The growth model chooses the best-fit model based on optimisation methods, which will bias those models with a greater number of fitted parameters (i.e. logistic model). To allow a more flexible growth model choice, users should be able to choose the best model to describe the growth of their trees.
- The biomass model is currently a mass-balance model and could be improved by using a process-based model of NPP, which would allow the effects of competition and nutrients to be modelled as well.
- Allocation parameters are static in time. Allocation changes with tree size/age and a more dynamic approach to allocation would be more realistic. Species specific allocations based on tree growth form would also increase the accuracy of the biomass model.
- If the default parameters are not applicable, a set of species specific parameters should be available for a range of commonly planted tree species to decrease the need for users to parameterise the model for each planted tree species themselves.

### *Crop model*

- The crop model assumes crops and yields do not change between years. It would be useful to allow crops planted and yields to change annually.
- If crops are not one of the default species listed in the IPCC, it should be possible for the user to parameterise the model for a different crop species, allowing a greater number of crop types (e.g. banana's, coffee).

### *External organic inputs*

- External organic inputs cannot easily be modelled if the N content, DPM/RPM ratio and combustion and emission factors are not known. These parameters are difficult to determine for various organic inputs. Default values for a range of typical organic inputs would make it easier for users to include these in calculations.

### *Emissions from biomass burning*

- The model does not take into account the emissions from burning of woody biomass taken off-farm, such as fire wood. The inclusion of emissions from burning of woody biomass removed from the fields would increase the accuracy of the emissions from this source.
- Fires do not volatilise SOC in this model or deposit carbon to the soil C pool. Emissions from these sources and impacts on these carbon stores should be considered in future versions, as fire can have a significant impact on SOC.
- Fire does not impact on living biomass in fields, including live trees. Fires only burn litter and crop residues, and assumes fire occur post-harvest. If a fire occurs before harvest, or burns whole standing trees, the emissions would be underestimated in the current model.

### *Emissions from plant N inputs*

- The N content of different plants and plant components can vary widely. Default values should be available for a range of different plant growth forms and/or N-fixing abilities to increase the accuracy of the model and decrease user data requirements.

### *Emissions from fertiliser use*

- Users are required to specify the N content of all applied synthetic and organic fertilisers. This process would be simplified if a set of default values were provided for various synthetic and organic fertiliser types, minimising the data requirements from users.

### *General*

- Increase the time steps from annual to monthly resolutions to capture the complexity of land management and subsequent emissions and removals within a year.

## References

- Cairns, M., S. Brown, E. Helmer, and G. Baumgardner. 1997. Root biomass allocation in the world's upland forests. *Oecologia* 111:1–11.
- Chave, J., C. Andalo, S. Brown, M. a Cairns, J. Q. Chambers, D. Eamus, H. Fölster, F. Fromard, N. Higuchi, T. Kira, J.-P. Lescure, B. W. Nelson, H. Ogawa, H. Puig, B. Riéra, and T. Yamakura. 2005. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia* 145:87–99.
- Coleman, K., and D. Jenkinson. 1999. RothC-26.3—A model for the turnover of carbon in soil: model description and windows users guide. Lawes Agricultural Trust, Harpenden.
- Diels, J., B. Vanlauwe, M. K. Van der Meersch, N. Sanginga, and R. Merckx. 2004. Long-term soil organic carbon dynamics in a subhumid tropical climate:  $^{13}\text{C}$  data in mixed C3/C4 cropping and modeling with RothC. *Soil Biology and Biochemistry* 36:1739–1750.
- Don, A., J. Schumacher, and A. FREIBAUER. 2011. Impact of tropical land-use change on soil organic carbon stocks - a meta-analysis. *Global Change Biology* 17:1658–1670.
- Falloon, P., P. Smith, K. Coleman, and S. Marshall. 1998. Estimating the size of the inert organic matter pool from total soil organic carbon content for use in the Rothamsted carbon model. *Soil Biology and ...* 30:1207–1211.
- FAO/IIASA/ISRIC/ISSCAS/JRC. 2009. *Harmonized World Soil Database (version 1.1)*. Food and Agriculture Organization of the United Nations, Rome, Italy and IIASA, Laxenburg, Austria. Available at: <http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/index.html>
- Farage, P., J. Ardo, L. Olsson, E. Rienzi, a Ball, and J. Pretty. 2007. The potential for soil carbon sequestration in three tropical dryland farming systems of Africa and Latin America: A modelling approach. *Soil and Tillage Research* 94:457–472.
- Guo, L., and R. Gifford. 2002. Soil carbon stocks and land use change: a meta analysis. *Global Change Biology* 8:345–360.
- Harris, I., P. D. Jones, T. J. Osborn, and D. H. Lister. 2013. Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology* 34:623–642.
- Hendricks, J. J., R. L. Hendrick, C. a. Wilson, R. J. Mitchell, S. D. Pecot, and D. Guo. 2006. Assessing the patterns and controls of fine root dynamics: an empirical test and methodological review. *Journal of Ecology* 94:40–57.
- IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds).
- Kaonga, M. L., and K. Coleman. 2008. Modelling soil organic carbon turnover in improved fallows in eastern Zambia using the RothC-26.3 model. *Forest Ecology and Management* 256:1160–1166.
- Milne, E., R. Al Adamat, N. H. Batjes, M. Bernoux, T. Bhattacharyya, C. C. Cerri, C. E. P. Cerri, K. Coleman, M. Easter, P. Falloon, C. Feller, P. Gicheru, P. Kamoni, K. Killian, D. K. Pal, K. Paustian, D. S. Powlson, Z. Rawajfih, M. Sessay, S. Williams, and S. Wokabi. 2007. National and sub-national assessments of soil organic carbon stocks and changes: The GEFSOC modelling system. *Agriculture, Ecosystems & Environment* 122:3–12.

- Mokany, K., R. J. Raison, and A. S. Prokushkin. 2006. Critical analysis of root : shoot ratios in terrestrial biomes. *Global Change Biology* 12:84–96.
- Nakamura, S., K. Hayashi, H. Omae, T. Ramadjita, F. Dougbedji, H. Shinjo, A. K. Saidou, and S. Tobita. 2010. Validation of soil organic carbon dynamics model in the semi-arid tropics in Niger, West Africa. *Nutrient Cycling in Agroecosystems* 89:375–385.
- Poorter, H., K. J. Niklas, P. B. Reich, J. Oleksyn, P. Poot, and L. Mommer. 2012. Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and environmental control. *The New phytologist* 193:30–50.
- Ryan, C. M., M. Williams, and J. Grace. 2011. Above- and Belowground Carbon Stocks in a Miombo Woodland Landscape of Mozambique. *Biotropica* 43:423–432.
- Traoré, P. C. S., W. M. Bostick, J. W. Jones, J. Koo, K. Goïta, and B. V Bado. 2008. A simple soil organic-matter model for biomass data assimilation in community-level carbon contracts. *Ecological applications : a publication of the Ecological Society of America* 18:624–36.
- Tumwebaze, S. B., E. Bevilacqua, R. Briggs, and T. Volk. 2013. Allometric biomass equations for tree species used in agroforestry systems in Uganda. *Agroforestry Systems* 87:781–795.

## Appendix 1: Model default values, sources, justification and applicability

This appendix outlines the default parameters used in the SHAMBA model. Their values, sources, justification and applicability are discussed for each parameter or set of parameters. These default values can be used by model users if applicability criteria are met. If they are not met, appropriate values must be found either through local measurements, appropriate databases, peer reviewed literature search, or other means.

Model component	Soil carbon model
Parameter	<b><i>t, et, p</i></b>
Description	Monthly climatic variables
Value	Depends on geographical location
Source of data	CRU TS 3.10 monthly global dataset
Reference	University of East Anglia Climatic Research Unit (CRU). [Phil Jones, Ian Harris]. CRU TS3.10: Climatic Research Unit (CRU) Time-Series (TS) Version 3.10 of High Resolution Gridded Data of Month-by-month Variation in Climate (Jan. 1901 - Dec. 2009),]. Available from: <a href="http://badc.nerc.ac.uk/view/badc.nerc.ac.uk__ATOM__ACTIVITY_fe67d66a-5b02-11e0-88c9-00e081470265">http://badc.nerc.ac.uk/view/badc.nerc.ac.uk__ATOM__ACTIVITY_fe67d66a-5b02-11e0-88c9-00e081470265</a>
Justification of choice of data or description of measurement methods and procedures applied	The dataset is based on weather station measurements worldwide, and has been peer reviewed. The dataset has been widely used in research and peer reviewed literature.
Applicability criteria	Climate data are representative of mean monthly climate conditions at the specified location
Comments	NA

Model component	Soil carbon model
Parameter	<b><math>soc_{y=0}</math>, <math>clay_{y=0}</math></b>
Description	SOC and clay content of soil in the top 0-30 cm at the start of the intervention
Value	<i>Depends on geographical location</i>
Source of data	Harmonized World Soil Database
Reference	FAO/IIASA/ISRIC/ISSCAS/JRC (2009) <i>Harmonized World Soil Database (version 1.1)</i> . Food and Agriculture Organization of the United Nations, Rome, Italy and IIASA, Laxenburg, Austria. Available at: <a href="http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/index.html">http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/index.html</a>
Justification of choice of data or description of measurement methods and procedures applied	This global dataset provides worldwide coverage of soil characteristics based on soil measurements. The HWSD has been peer reviewed, and has been widely used in research and peer reviewed literature.
Applicability criteria	The soil characteristics are representative of soils at the specified location, and represent conditions at the start of the intervention.
Comments	This dataset is used to define soil characteristics at the start of the intervention (i.e. disturbed soils or soils under agriculture). It is assumed that the HWSD is a suitable measure of this, as soil measurements taken in disturbed sites are far more common than in pristine habitats. Therefore, the HWSD is more likely to be based on disturbed soil measurements than undisturbed.

Model component	Soil carbon model
Parameter	<b><math>soc_r</math></b>
Description	SOC in the top 0-30 cm prior to conversion to agriculture (i.e. undisturbed soils)
Value	25 % higher than $soc_{y=0}$
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Don, A., J. Schumacher, and A. FREIBAUER. 2011. Impact of tropical land-use change on soil organic carbon stocks - a meta-analysis. <i>Global Change Biology</i> 17:1658–1670.</li> <li>- Guo, L., and R. Gifford. 2002. Soil carbon stocks and land use change: a meta analysis. <i>Global Change Biology</i> 8:345–360.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	The effects of land use change on soil organic carbon are poorly quantified, and lack consensus. We assume that SOC is lost when soils are disturbed. When forests are converted to agriculture, we expect an average loss of 25 % of SOC stocks. This value is based on peer reviewed meta-analyses of SOC stock changes in land-use change studies of the tropics (Don <i>et al.</i> , 2011), and elsewhere (Guo and Gifford, 2002)
Applicability criteria	Land was primary forest or woodland prior to disturbance, when it was converted to agriculture.
Comments	For other SOC loss rates, or conversion from non-primary forests, see Don <i>et al.</i> (2011) paper for other values.

Model component	Soil carbon model
Parameter	<i>dpm<sub>c</sub>, rpm<sub>c</sub>, dpm<sub>t</sub>, rpm<sub>t</sub></i>
Description	Decomposable and resistant fraction of crop and tree inputs
Value	0.59, 0.41, 0.20, 0.80
Source of data	Rothamsted Research
Reference	RothC v 26.6 (2008) A model for the turnover of carbon in soil. Available at: <a href="http://www.rothamsted.ac.uk/aen/carbon/mod26_3_win.pdf">http://www.rothamsted.ac.uk/aen/carbon/mod26_3_win.pdf</a>
Justification of choice of data or description of measurement methods and procedures applied	The RothC model provides default values for the DPM and RPM fractions of plant material from crop residues and from tropical woodlands. These values have been used by several peer reviewed publications using the RothC model.
Applicability criteria	Land was primary forest or woodland prior to disturbance, when it was converted to agriculture.
Comments	For other SOC loss rates, or conversion from non-primary forests, see Don <i>et al.</i> (2011) paper for other values.

Model component	Biomass model
Parameter	<i>al<sub>stem</sub>, al<sub>branch</sub>, al<sub>leaf</sub></i>
Description	Single tree allocation of NPP to the stem, branches and leaves
Value	0.69, 0.31, 0.10
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Poorter, H. et al., 2012. Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and environmental control. <i>The New phytologist</i>, 193(1), pp.30–50.</li> <li>- Ryan, C.M., Williams, M. &amp; Grace, J., 2011. Above- and Belowground Carbon Stocks in a Miombo Woodland Landscape of Mozambique. <i>Biotropica</i>, 43(4), pp.423–432.</li> <li>- Tumwebaze, S.B. <i>et al.</i>, 2013. Allometric biomass equations for tree species used in agroforestry systems in Uganda. <i>Agroforestry Systems</i>, 87(4), pp.781–795.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	Allocation of biomass to the stem, branches and leaves will vary depending on a trees allometry. However, similar allocation patterns were found between mature trees of agroforestry species (Tumwebaze <i>et al.</i> , 2013), and natural miombo woodland trees (Ryan <i>et al.</i> , 2011). These values were used as the default values, and were within the range reported in a meta-analysis of biomass allocation for tropical forests and woodlands (Poorter <i>et al.</i> , 2012).
Applicability criteria	Trees must have a similar allometry to species such as <i>G. robusta</i> , <i>M. lutea</i> , <i>M. eminii</i> , <i>J. globiflora</i> , and <i>B. spiciformis</i> (see section 4.4), as used in the studies. No pruning of branches or other biomass may occur, as this will change the allometry of trees.
Comments	If planted tree species have a differing allometry, species specific allocation parameters must be sought in peer reviewed literature.

Model component	Biomass model
Parameter	<i>al<sub>root</sub></i>
Description	NPP allocation to fine roots
Value	0.10 ( <i>same as AL<sub>leaf</sub></i> )
Source of data	Peer reviewed literature
Reference	Hendricks, J.J. et al., 2006. Assessing the patterns and controls of fine root dynamics: an empirical test and methodological review. <i>Journal of Ecology</i> , 94(1), pp.40–57.
Justification of choice of data or description of measurement methods and procedures applied	We assume that fine roots have the same allocation ratio as leaves (i.e. fine root productivity is the same as leaves). Other studies done in temperate forests have found a wide range of fine root:leaf productivity ratios, but with a mean of $1.26 \pm 1.44$ ( $\pm$ SD) (Hendricks et al., 2006). Therefore, the assumption that fine root allocation is the same as leaves agrees with the average of several measured values.
Applicability criteria	Fine root productivity is similar or equal to leaf productivity
Comments	If species specific or local measurements are available for fine root allocation patterns, these values should be used instead.

Model component	Biomass model
Parameter	<i>al<sub>crroot</sub></i>
Description	NPP allocation to coarse roots
Value	0.26 ( <i>same as root:shoot ratio</i> )
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Cairns, M. et al., 1997. Root biomass allocation in the world's upland forests. <i>Oecologia</i>, 111(1), pp.1–11.</li> <li>- Mokany, K., Raison, R.J. &amp; Prokushkin, A.S., 2006. Critical analysis of root : shoot ratios in terrestrial biomes. <i>Global Change Biology</i>, 12(1), pp.84–96.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	Coarse root allocation is assumed to be the same as the root:shoot ratio. Due to a paucity of data for agroforestry trees, we use the average root:shoot ratio as reported in a meta-analyses of tree root:shoot ratios (Cairns et al., 1997), which was within the range of values reported in Mokany et al. (2006) for tropical forest trees.
Applicability criteria	Allocation to coarse roots is the same as the root:shoot ratio
Comments	If species specific or local measurements are available for coarse root allocation, or root:shoot ratios, these values should be used instead.



Model component	Biomass model
Parameter	$to_{stem}$
Description	Annual turnover rate of stems
Value	0
Source of data	Conservative estimate
Reference	NA
Justification of choice of data or description of measurement methods and procedures applied	The model assumes as default that tree stems have no turnover of biomass annually. This assumes that there are no significant biomass losses in stems, such as bark shedding.
Applicability criteria	Stems do not shed their bark or lose biomass
Comments	If tree species planted shed a significant amount of stem biomass annually, such as bark, the turnover value should be > 0

Model component	Biomass model
Parameter	$to_{branch}$
Description	Annual turnover rate of branches
Value	0.05
Source of data	Conservative estimate
Reference	NA
Justification of choice of data or description of measurement methods and procedures applied	The model assumes as default that tree branches lose 5 % of their branch mass as litter annually. This is a conservative estimate, and might be a reasonable estimate if branches are broken off during storms or other events.
Applicability criteria	Trees lose approx. 5 % of their branch mass as litter annually
Comments	If tree species planted lose or shed a significant amount of branch biomass annually, such as through natural pruning, the turnover value should be adjusted accordingly.

Model component	Biomass model
Parameter	<b><i>to<sub>leaf</sub></i></b>
Description	Annual turnover rate of leaves
Value	1
Source of data	Estimate
Reference	NA
Justification of choice of data or description of measurement methods and procedures applied	The default assumes the trees lose all their leaves annually as litter, and that trees are either deciduous or have a leaf life span $\leq 1$ . This is set as the default as it was the simplest assumption to make.
Applicability criteria	Trees are deciduous or have a leaf life span $\leq 1$ year
Comments	If trees are evergreen, or have a leaf life span $>1$ , the leaf turnover rate can be estimated as follows: $1 / \text{leaf lifespan} = \text{turnover rate per year}$

Model component	Biomass model
Parameter	<b><i>to<sub>root</sub></i></b>
Description	Annual turnover rate of fine roots
Value	0.8
Source of data	Peer reviewed literature
Reference	Gill, R. & Jackson, R., 2000. Global patterns of root turnover for terrestrial ecosystems. <i>New Phytologist</i> , pp.13–31.
Justification of choice of data or description of measurement methods and procedures applied	Fine root turnover is difficult to measure and rarely reported in the literature. We base our default on a peer reviewed global meta-analysis of root turnover (Gill & Jackson, 2000), using the mean turnover rate reported for tropical tree fine roots.
Applicability criteria	NA
Comments	Local or species specific values for fine root turnover should be used where possible

Model component	Biomass model
Parameter	$t_{croot}$
Description	Annual turnover rate of coarse roots
Value	0
Source of data	Conservative estimate
Reference	NA
Justification of choice of data or description of measurement methods and procedures applied	Turnover rates of coarse roots are assumed to be zero as the default. This represents a conservative estimate, given the relatively low rates reported and the high uncertainty of this pool in the literature.
Applicability criteria	Coarse roots are not expected to have a high turnover rate
Comments	Local or species specific values for coarse root turnover should be used where possible

Model component	Biomass model
Parameter	$t_{rs}$
Description	Tree root:shoot ratio
Value	0.26
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Cairns, M. et al., 1997. Root biomass allocation in the world's upland forests. <i>Oecologia</i>, 111(1), pp.1–11.</li> <li>- Mokany, K., Raison, R.J. &amp; Prokushkin, A.S., 2006. Critical analysis of root : shoot ratios in terrestrial biomes. <i>Global Change Biology</i>, 12(1), pp.84–96.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	Root:shoot ratios can vary widely for different trees, and are dependent on several factors. Due to a paucity of data for agroforestry trees, we use the average root:shoot ratio as reported in a meta-analyses of tree root:shoot ratios (Cairns et al., 1997), which was within the range of values reported in Mokany et al. (2006) for tropical forest trees.
Applicability criteria	Trees have approx. 26 % of their biomass below-ground
Comments	If species specific or local measurements are available for root:shoot ratios, these values should be used

Model component	Biomass model
Parameter	<i>t<sub>30</sub></i>
Description	Fraction of tree roots in the top 0-30 cm of soil
Value	0.70
Source of data	Peer reviewed literature
Reference	Jackson, R.B. et al., 1996. A global analysis of root distributions for terrestrial biomes. <i>Oecologia</i> , 108(3), pp.389–411.
Justification of choice of data or description of measurement methods and procedures applied	Root distributions in the soil profile vary depending on vegetation type, soil characteristics and other factors. Therefore, a mean value for tropical forest biomes was used as the best available estimate. The default value was derived from a peer reviewed meta-analysis of root distributions for terrestrial biomes (Jackson et al., 1996), which has been widely cited in the literature.
Applicability criteria	Total tree roots biomass is mostly found in the top 0-30 cm of soil
Comments	Local or species specific values for root distributions should be used where possible

Model component	Biomass model
Parameter	<i>t<sub>Cstem</sub></i> , <i>t<sub>Cbranch</sub></i> , <i>t<sub>Cleaf</sub></i> , <i>t<sub>Croot</sub></i> , <i>t<sub>Cfruit</sub></i>
Description	Tree biomass C content
Value	0.5
Source of data	Estimate
Reference	<ul style="list-style-type: none"> <li>- Chave, J. et al., 2005. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. <i>Oecologia</i>, 145(1), pp.87–99.</li> <li>- Brown, S. &amp; Lugo, A.E., 1984. Biomass of tropical forests: a new estimate based on forest volumes. <i>Science</i> (New York, N.Y.), 223, pp.1290–1293.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	Woody biomass C content can vary between tree species, but it is traditionally assumed that the carbon content of dry biomass of a tree is 50 %.
Applicability criteria	NA
Comments	NA

Model component	Biomass model ( <i>Chave allometric equations only</i> )
Parameter	$t_d$
Description	Wood density
Value	0.60
Source of data	Peer reviewed literature
Reference	Chave, J. et al., 2009. Towards a worldwide wood economics spectrum. Ecology letters, 12(4), pp.351–66.
Justification of choice of data or description of measurement methods and procedures applied	Wood density can vary widely between tree species. Therefore, to estimate the wood density of planted trees we use an average for tropical African trees, as determined by a peer reviewed meta-analysis of 2482 tree species from this region (Chave et al., 2009).
Applicability criteria	If one of the Chave allometric equations is selected, and the planted trees are similar to a tropical African tree
Comments	Species specific values for wood density should be used where possible, and can be found at the following websites: <ul style="list-style-type: none"> <li>• <a href="http://db.worldagroforestry.org/wd">http://db.worldagroforestry.org/wd</a></li> <li>• <a href="https://datadryad.org/repo/handle/10255/dryad.235">https://datadryad.org/repo/handle/10255/dryad.235</a></li> </ul>

Model component	Biomass model
Parameter	$tn_{stem}$
Description	Nitrogen content of tree stems
Value	0.0015
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Chave, J. et al., 2009. Towards a worldwide wood economics spectrum. Ecology letters, 12(4), pp.351–66.</li> <li>- Weedon, J.T. et al., 2009. Global meta-analysis of wood decomposition rates: a role for trait variation among tree species? Ecology letters, 12(1), pp.45–56.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	Nitrogen content of wood can vary widely depending on location, tree species and other factors. Furthermore, few studies report wood N content for agroforestry trees. Therefore, a mean for woody debris N content is used, based on values reported in global meta-analyses of woody traits from peer reviewed literature.
Applicability criteria	NA
Comments	The default value is a very rough estimate of wood N content, and it is recommended that species specific values for wood N content are used where possible, especially if trees are N fixing or leguminous.

Model component	Biomass model
Parameter	<b><i>tn<sub>branch</sub></i></b> , <b><i>tn<sub>croot</sub></i></b>
Description	Nitrogen content of branches and coarse roots
Value	0.0015 ( <i>same as N<sub>stem</sub></i> )
Source of data	NA
Reference	NA
Justification of choice of data or description of measurement methods and procedures applied	Branches and coarse root N content are rarely reported in the literature. Therefore, we assume the N content of branches and coarse roots would be the same as for woody stems, as branches and coarse roots are also largely woody biomass.
Applicability criteria	NA
Comments	

Model component	Biomass model
Parameter	<b><i>tn<sub>leaf</sub></i></b>
Description	Nitrogen content of leaf litter from agroforestry trees (not fresh green leaves)
Value	0.01 if non-legume, 0.02 if legume
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Constantinides, M. &amp; Fownes, J., 1994. Nitrogen mineralization from leaves and litter of tropical plants: relationship to nitrogen, lignin and soluble polyphenol concentrations. <i>Soil Biology and Biochemistry</i>, 26(1), pp.49–55.</li> <li>- Ratnam, J. et al., 2008. Nutrient resorption patterns of plant functional groups in a tropical savanna: variation and functional significance. <i>Oecologia</i>, 157(1), pp.141–51.</li> <li>- Vitousek, P., 1984. Litterfall, nutrient cycling, and nutrient limitation in tropical forests. <i>Ecology</i>, 65(1), pp.285–298.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	Nitrogen content of leaf litter can vary widely depending on location, tree N-fixing ability, tree age and other factors. Therefore, a conservative value for leaf litter N content is used for leguminous trees and non-leguminous trees based on a study of several agroforestry trees (Constantinides et al., 1994). These values show close agreement to other studies of leaf litter N content from tropical Africa (Vitousek, 1984), and South Africa (Ratnam, 2008).
Applicability criteria	NA
Comments	This mean value is an estimate of leaf litter N content, and it is recommended that species specific values for leaf litter N content are used where possible, especially if trees are N-fixing or leguminous.

Model component	Biomass model
Parameter	<i>tn<sub>root</sub></i>
Description	Nitrogen content of agroforestry fine roots
Value	0.0113
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Gordon, W. &amp; Jackson, R., 2000. Nutrient concentrations in fine roots. <i>Ecology</i>, 81(January), pp.275–280.</li> <li>- Jackson, R.B., Mooney, H. a &amp; Schulze, E.D., 1997. A global budget for fine root biomass, surface area, and nutrient contents. <i>PNAS</i>, 94(14), pp.7362–6.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	Fine root nutrient contents are dynamic and can change widely depending on local conditions, tree species, and other factors. Therefore, we use a mean value of fine root nitrogen content as reported by a global meta-analysis, using the mean value reported for fine roots of broadleaf and coniferous trees (Gordon & Jackson, 2000), in close agreement with other global meta analyses (Jackson <i>et al.</i> , 1997).
Applicability criteria	NA
Comments	Local or species specific values should be used where possible

Model component	Crop model
Parameter	<i>Crs, Can, Cbn</i>
Description	Crop root:shoot ratio, crop residue above-ground N content and below-ground N content
Value	<i>Depends on crop species/type</i>
Source of data	IPCC, Table 11.2
Reference	<p>IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds).</p> <p>Available at: <a href="http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_11_Ch11_N2O&amp;CO2.pdf">http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_11_Ch11_N2O&amp;CO2.pdf</a></p>
Justification of choice of data or description of measurement methods and procedures applied	The IPCC Guidelines for National Greenhouse Gas Inventories is internationally recognised and the data provided in the guidelines is peer reviewed.
Applicability criteria	Crops must be one of the IPCC listed crop species or types
Comments	NA

Model component	Crop model
Parameter	<b>C<sub>ac</sub>, C<sub>bc</sub></b>
Description	Crop residue above-ground C content and below-ground C content
Value	0.40
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Johnson, J.M.-F., Allmaras, R.R. &amp; Reicosky, D.C., 2006. Estimating Source Carbon from Crop Residues, Roots and Rhizodeposits Using the National Grain-Yield Database. <i>Agronomy Journal</i>, 98(3), p.622.</li> <li>- Latshaw, W. &amp; Miller, E., 1924. Elemental composition of the corn plant. <i>Journal of Agricultural Research</i>, XXVII(11).</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	A default value of 0.4 was chosen as other studies have used this estimated mean C content for crop residues in shoots and roots (Johnson et al., 2006), and it agrees with the C content values reported for maize plants (Latshaw & Miller, 1924).
Applicability criteria	NA
Comments	Local or species specific values should be used where possible

Model component	Crop model
Parameter	<b>C<sub>30</sub></b>
Description	Fraction of crop roots (residues) in the top 0-30 cm of soil
Value	0.70
Source of data	Peer reviewed literature
Reference	Jackson, R.B. et al., 1996. A global analysis of root distributions for terrestrial biomes. <i>Oecologia</i> , 108(3), pp.389–411.
Justification of choice of data or description of measurement methods and procedures applied	Crop root distributions in the soil profile vary depending on crop type, soil characteristics and other factors. Therefore, a mean value for crop root distributions was used as the best available estimate, derived from a peer reviewed meta-analysis of root distributions for terrestrial biomes.
Applicability criteria	NA
Comments	Local or species specific values should be used where possible



Model component	External organic inputs
Parameter	$a_n$
Description	External organic input N content
Value	0.018
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Mtambanengwe, F. &amp; Kirchmann, H., 1995. Litter from a tropical savanna woodland (Miombo): chemical composition and C and N mineralization. <i>Soil Biology and Biochemistry</i>, 27(12), pp.1639–1651.</li> <li>- Musvoto, C., Campbell, B.M. &amp; Kirchmann, H., 2000. Decomposition and nutrient release from mango and miombo woodland litter in Zimbabwe. <i>Soil Biology and Biochemistry</i>, 32(8-9), pp.1111–1119.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	External organic inputs originating from surrounding woodland/forest is assumed to be a mixture of leaf, twig, bark and other woody litter. We assume forest litter is 1.8 % N, which agrees closely with values reported for miombo woodland mixed tree litter C content (Mtambanengwe et al., 1995; Musvoto et al., 2000).
Applicability criteria	External organic inputs must originate from forest/woodland litter (miombo), or similar woody trees/shrubs litter.
Comments	NA

Model component	External organic inputs
Parameter	$a_c$
Description	External organic input C content
Value	0.50
Source of data	Peer reviewed literature
Reference	<ul style="list-style-type: none"> <li>- Mtambanengwe, F. &amp; Kirchmann, H., 1995. Litter from a tropical savanna woodland (Miombo): chemical composition and C and N mineralization. <i>Soil Biology and Biochemistry</i>, 27(12), pp.1639–1651.</li> <li>- Musvoto, C., Campbell, B.M. &amp; Kirchmann, H., 2000. Decomposition and nutrient release from mango and miombo woodland litter in Zimbabwe. <i>Soil Biology and Biochemistry</i>, 32(8-9), pp.1111–1119.</li> </ul>
Justification of choice of data or description of measurement methods and procedures applied	External organic inputs originating from surround woodland/forest is assumed to be a mixture of leaf, twig, bark and other woody litter. We assume plant litter is 50 % C, which agrees closely with values reported for miombo woodland and mango mixed tree litter C content (Mtambanengwe et al., 1995; Musvoto et al., 2000).
Applicability criteria	External organic inputs must originate from forest/woodland litter, or woody trees/shrubs litter.
Comments	NA

<b>Model component</b>	Biomass burning, emissions from plant N inputs, emission from fertiliser use
<b>Parameter</b>	<i>cf, ef, gwp, v</i>
<b>Description</b>	Combustion factors, emissions factors, global warming potentials of greenhouse gases, and volatisation fraction
<b>Value</b>	<i>See section 7 (Table 6), section 9 (Table 8), section 10 (Table 9) for values</i>
<b>Source of data</b>	IPCC
<b>Reference</b>	IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds).
<b>Justification of choice of data or description of measurement methods and procedures applied</b>	The IPCC Guidelines for National Greenhouse Gas Inventories is internationally recognised and the data provided in the guidelines is peer reviewed.
<b>Applicability criteria</b>	Values are appropriate to biomass type (e.g. crop residues, tree litter), and gas emitted
<b>Comments</b>	