

Clustering in a hyperbolic model of complex networks.

Nikolaos Fountoulakis^{*1}, Pim van der Hoorn^{†2}, Tobias Müller^{‡3}, and Markus Schepers^{§3}

¹ School of Mathematics, University of Birmingham, United Kingdom.

²Department of Physics, Northeastern University, United States.

³Bernoulli Institute, University of Groningen, The Netherlands.

January 2, 2020

Abstract

In this paper we consider the clustering coefficient, and clustering function in a random graph model proposed by Krioukov et al. in 2010. In this model, nodes are chosen randomly inside a disk in the hyperbolic plane and two nodes are connected if they are at most a certain hyperbolic distance from each other. It has been previously shown that this model has various properties associated with complex networks, including a power-law degree distribution, “short distances” and a non-vanishing clustering coefficient. The model is specified using three parameters: the number of nodes n , which we think of as going to infinity, and $\alpha, \nu > 0$, which we think of as constant. Roughly speaking, the parameter α controls the power law exponent of the degree sequence and ν the average degree.

Here we show that the clustering coefficient tends in probability to a constant γ that we give explicitly as a closed form expression in terms of α, ν and certain special functions. This improves earlier work by Gugelmann et al., who proved that the clustering coefficient remains bounded away from zero with high probability, but left open the issue of convergence to a limiting constant. Similarly, we are able to show that $c(k)$, the average clustering coefficient over all vertices of degree exactly k , tends in probability to a limit $\gamma(k)$ which we give explicitly as a closed form expression in terms of α, ν and certain special functions. We are able to extend this last result also to sequences $(k_n)_n$ where k_n grows as a function of n . Our results show that $\gamma(k)$ scales differently, as k grows, for different ranges of α . More precisely, there exists constants $Q_{\alpha,\nu}$ depending on α and ν , such that as $k \rightarrow \infty$, $\gamma(k) \sim Q_{\alpha,\nu} k^{2-4\alpha}$ if $\frac{1}{2} < \alpha < \frac{3}{4}$, $\gamma(k) \sim Q_{\alpha,\nu} \log(k) k^{-1}$ if $\alpha = \frac{3}{4}$ and $\gamma = Q_{\alpha,\nu} k^{-1}$ when $\alpha > \frac{3}{4}$. These results contradict a prediction of Krioukov et al., which stated that the limiting values $\gamma(k)$ should always scale with k^{-1} as we let k grow.

^{*}Research partially supported by the Alan Turing Institute, grant no. EP/N510129/1.

[†]Research partially supported by ARO Grant W911NF-16-1-0391 and W911NF-17-1-0491

[‡]Research partially supported by NWO grants 639.032.529 and 612.001.409.

[§]Research partially supported by NWO grant 639.032.529.

Contents

1	Introduction and main results	4
1.1	KPKVB model	4
1.2	Main results	7
1.2.1	Degrees	7
1.2.2	The clustering coefficient	8
1.2.3	The clustering function	9
1.2.4	Scaling of $\gamma(k)$	10
1.3	Observations	10
1.4	Outline of the paper	11
2	Preliminaries	11
2.1	The infinite limit model G_∞	11
2.2	The finite box model G_{box}	12
2.3	The Poissonized KPKVB model G_{Po}	12
2.4	Coupling G_{Po} and G_{box}	12
2.5	The Campbell-Mecke formula	13
2.6	Concentration of heights	14
3	Degrees in the KPKVK model: proof of Theorem 1.1	14
3.1	Proof overview	15
3.2	The degree of the typical point in G_∞	15
3.3	Expected degrees in G_{box} and G_{Po}	16
3.4	Joint degrees in G_{box} and G_{Po}	20
3.5	Factorial moments of degrees	28
3.6	Coupling G_n to G_{Po}	33
3.7	Proof of Theorem 1.1	36
4	Clustering and the degree of the typical point in G_∞	38
4.1	The expected clustering coefficient and function of the typical point	38
4.1.1	Determining $P(y_0, y_1, y_2)$	40
4.1.2	Integrating over y_1, y_2	42
4.1.3	Computing γ and $\gamma(k)$	47
4.1.4	Explicit expressions for $\gamma, \gamma(k)$ when $\alpha = 1$.	51
4.2	The proof of Proposition 1.5	55
5	Proofs of Theorem 1.2 and Theorem 1.3	59
5.1	Clustering function, proving Theorem 1.3	59
5.1.1	Poissonization	59
5.1.2	Truncation	60
5.1.3	Coupling	62
5.1.4	Deriving the limit integral expression	62
5.1.5	Putting everything together	63
5.2	Overall clustering coefficient, proving Theorem 1.2	64
6	Overview of the proof strategy for $k \rightarrow \infty$	64
6.1	Adjusted clustering and the Poissonized KPKVB model	65
6.2	Coupling of local clustering between G_{Po} and G_{box}	65
6.3	From the finite box to the infinite model	67
6.4	Proof of the main results	67
7	From G_{box} to G_∞ (Proving Proposition 6.5)	68
7.1	Comparing triangles between G_∞ and G_{box}	70
7.2	Counting missing triangles	73
8	Concentration for $c(k; G_{\text{box}})$ (Proving Proposition 6.4)	82

8.1	The main contribution of triangles	82
8.2	Joint degrees in G_{box}	84
8.3	Concentration result for main triangle contribution	86
9	Equivalence for local clustering in G_{Po} and G_{box}	91
9.1	Some results on the hyperbolic geometric graph	91
9.2	Equivalence clustering G_{Po} and G_{box}	93
9.3	Coupling G_n to G_{Po}	111
A	Meijer's G-function	114
B	Incomplete Beta function	114
C	Some results on functions	116
D	Some results for random variables	116
E	Concentration of heights for vertices with degree k	119
E.1	Concentration of heights argument for the infinite model	120
E.2	Concentration of heights for the KPKVB and finite box model	122
F	Derivative of $\mu_{Po}(y)$	124
G	Code for the simulations	128

1 Introduction and main results

In this paper, we will consider clustering in a model of random graphs that involves points taken randomly in the hyperbolic plane. This model was introduced by Krioukov, Papadopoulos, Kitsak, Vahdat and Boguñá [25] in 2010 - we abbreviate it as *the KPKVB model*. We should however note that the model also goes by several other names in the literature, including *hyperbolic random geometric graphs* and *random hyperbolic graphs*. Krioukov et al. suggested this model as a suitable model for complex networks. It exhibits the three main characteristics usually associated with complex networks: a power-law degree distribution, a non-vanishing clustering coefficient and short graph distances.

1.1 KPKVB model

We start with the definition of the model. As mentioned, its nodes are situated in the hyperbolic plane \mathbb{H} , which is a surface with constant Gaussian curvature -1 . This surface has several convenient representations (i.e. coordinate maps), such as the Poincaré halfplane model, the Poincaré disk model and the Klein disk model. A gentle introduction to Gaussian curvature, hyperbolic geometry and these representations of the hyperbolic plane can be found in [34]. Throughout this paper we will be working with a representation of the hyperbolic plane using *hyperbolic polar coordinates*, sometimes called the *native representation*. That is, a point $u \in \mathbb{H}$ is represented as (r, θ) , where r is the hyperbolic distance between u and the origin O and θ as the angle between the line segment Ou and the positive x -axis. Here, when mentioning “the origin” and the angle between the line segment and the positive x -axis, we think of \mathbb{H} embedded as the Poincaré disk in the ordinary euclidean plane.

The KPKVB model has three parameters: the number of vertices n , which we think of as going to infinity, and $\alpha > \frac{1}{2}$, $\nu > 0$ which we think of as fixed. Given n, α, ν we define $R = 2 \log(n/\nu)$. Then the hyperbolic random graph $G(n; \alpha, \nu)$ is defined as follows:

- The vertex set is given by n i.i.d. points u_1, \dots, u_n denoted in polar coordinates $u_i = (r_i, \theta_i)$, where the angular coordinate θ is chosen uniformly from $(-\pi, \pi]$ while the radial coordinate r is sampled independently according to the cumulative distribution function

$$F_{\alpha, R}(r) = \begin{cases} 0 & \text{if } r < 0 \\ \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1} & \text{if } 0 \leq r \leq R \\ 1 & \text{if } r > R \end{cases} \quad (1)$$

- Any two vertices $u_i = (r_i, \theta_i)$ and $u_j = (r_j, \theta_j)$ are adjacent if and only if $d_{\mathbb{H}}(u_i, u_j) \leq R$, where $d_{\mathbb{H}}$ denotes the distance in the hyperbolic plane. We will frequently be using that, by the hyperbolic law of cosines, $d_{\mathbb{H}}(u_i, u_j) \leq R$ is equivalent to

$$\cosh(r_i) \cosh(r_j) - \sinh(r_i) \sinh(r_j) \cos(|\theta_i - \theta_j|_{2\pi}) \leq \cosh(R),$$

where $|a|_b = \min(|a|, b - |a|)$ for $-b \leq a \leq b$.

Figure 1 shows a computer simulation of $G(n; \alpha, \nu)$.

As observed by Krioukov et al. [25], and proved rigorously by Gugelmann et al. [21], the degree sequence of the KPKVB model follows a power-law with exponent $2\alpha + 1$. Gugelmann et al. [21] also showed that the average degree converges in probability to the constant $8\nu\alpha^2/\pi(2\alpha - 1)^2$, and they showed that the (local) clustering coefficient is non-vanishing in the sense that it is bounded below by a positive constant a.a.s. Here, and in the rest of the paper, for a sequence $(E_n)_n$ of events, E_n *asymptotically almost surely* (a.a.s.) means that $\mathbb{P}(E_n) \rightarrow 1$ as $n \rightarrow \infty$. Here the events E_n might be defined in different probability spaces, in which case \mathbb{P} will be the corresponding probability measure.

Apart from the degree sequence and clustering, the third main characteristic associated with complex networks, “short distances”, has also been established in the literature. In [2] it is shown




Figure 1: Simulation $G(n; \alpha, \nu)$ with $\alpha = 0.9$, $\nu = 0.2$ and $n = 5000$.

that for $\alpha < 1$ the largest component is what is called an *ultra-small world*: if we randomly sample two vertices of the graph then, a.a.s., conditional on them being in the same component, their graph distance is of order $\log \log n$. In [22] and [19] a.a.s. polylogarithmic upper and lower bounds on the graph diameter of the largest component are shown, and in [30], these were sharpened to show that $\log n$ is the correct order of the diameter.

Earlier work of the first and third authors with Bode [8] and of the first and third authors [17] has established the “threshold for a giant component”: if $\alpha < 1$ then there is a unique component of size linear in n no matter how small ν (i.e. the average degree); if $\alpha > 1$ all components are sublinear no matter the value of ν ; and if $\alpha = 1$ then there is a critical value ν_c such that for $\nu < \nu_c$ all components are sublinear and for $\nu > \nu_c$ there is a unique linearly sized component (all of these statements holding a.a.s.). Whether or not there is a giant component if $\alpha = 1$ and $\nu = \nu_c$ remains an open problem. In [22] and [23], Kiwi and Mitsche considered the size of the second largest component and showed that for $\alpha \in (1/2, 1)$, a.a.s., the second largest component has polylogarithmic order with exponent $1/(\alpha - 1/2)$.

In another paper of the first and third authors with Bode [9] it was shown that $\alpha = 1/2$ is the threshold for connectivity: for $\alpha < 1/2$ the graph is a.a.s. connected, for $\alpha > 1/2$ the graph is a.a.s. disconnected and when $\alpha = 1/2$ the probability of being connected tends to a continuous, nondecreasing function of ν which is identically one for $\nu \geq \pi$ and strictly less than one for $\nu < \pi$. Friedrich and Krophmer [6] studied the size of the largest clique as well as the number of cliques of a given size. Boguñá et al. [10] and Bläsius et al. [7] considered fitting the KPKVB model to data using maximum likelihood estimation. Kiwi and Mitsche [24] studied the spectral gap and related properties, and Bläsius et al. [5] considered the tree-width and related parameters of the KPKVB model. Recently Owada and Yogeshwaran [31] considered subgraph counts, and in particular established a central limit theorem for the number of copies of a fixed tree T in $G(n; \alpha, \nu)$, subject to some restrictions on the parameter α .

Clustering

In this work we study the clustering coefficient in the KPKVB model. In the literature there are unfortunately two distinct, rival definitions of the *clustering coefficient*. One of those, sometimes called the *global* clustering coefficient, is defined as three times the ratio of the number of triangles to the number of paths of length two in the graph. Results for this version of the clustering coefficient in the KPKVB model were obtained by Candellero and the first author [11] and for the

evolution of graphs on more general spaces with negative curvature by the first author in [16].

We will study the other notion of clustering, the one which is also considered by Krioukov et al. [25] and Gugelmann et al. [21]. It is sometimes called the *local* clustering coefficient, although we should point out that Gugelmann et al. actually call it the global clustering coefficient in their paper. For a graph G and a vertex $v \in V(G)$ we define the clustering coefficient of v as:

$$c(v) := \begin{cases} \frac{1}{\binom{\deg(v)}{2}} \sum_{u, w \sim v} 1_{\{uw \in E(G)\}}, & \text{if } \deg(v) \geq 2, \\ 0, & \text{otherwise,} \end{cases}$$

where $E(G)$ denotes the edge set of G and $\deg(v)$ is the degree of vertex v . That is, provided v has degree at least two, $c(v)$ equals the number of edges that are actually present between the neighbours of v divided by the number of edges that could possibly be present between the neighbours given the degree of v . The clustering coefficient of G is now defined as the average of $c(v)$ over all vertices v :

$$c(G) := \frac{1}{|V(G)|} \sum_{v \in V(G)} c(v).$$

As mentioned above, Gugelmann et al. [21], have established that $c(G(n; \alpha, \nu))$ is non-vanishing a.s., but they left open the question of convergence. Theorem 1.2 below establishes that the clustering coefficient indeed converges in probability to a constant γ that we give explicitly as a closed form expression involving α, ν and several classical special functions.

In addition to the clustering coefficient, we shall also be interested in the *clustering function*. This assigns to each non-negative integer k the value

$$c(k; G) := \begin{cases} \frac{1}{N(k)} \sum_{\substack{v \in V(G), \\ \deg(v) = k}} c(v), & \text{if } N(k) \geq 1, \\ 0, & \text{else,} \end{cases} \quad (2)$$

where $N(k)$ denotes the number of vertices of degree exactly k in G . In other words, the clustering function assigns to the integer k the average of the local clustering coefficient over all vertices of degree k . We remark that, while it might seem natural to consider $c(k)$ to be “undefined” when $N(k) = 0$, we prefer to use the above definition for technical convenience. This way $c(k; G(n; \alpha, \nu))$ is a plain vanilla random variable and we can for instance compute its moments without any issues.

A general expression of the clustering function for KPKVB random graphs is given in [25, Equation (59)]. The authors conjecture that as k tends to infinity, the clustering function decays as k^{-1} . They based this prediction on observations (Figure 8 in [25]) in experiments on the infrastructure of the Internet obtained in [13]. Despite these interesting observations and the attention the KPKVB model has generated since then, the behaviour of the clustering function in KPKVB random graphs had not been completely determined. In particular it has not been established whether it converges as $n \rightarrow \infty$ to some suitable limit function, nor how $c(k; G(n; \alpha, \nu))$ scales with k . Theorems 1.3, 1.4 and Proposition 1.5 below settle these questions. Theorem 1.3 shows that for each fixed k , the value $c(k; G(n; \alpha, \nu))$ converges in probability to a constant $\gamma(k)$ that we again give explicitly as a closed form expression involving α, ν and several classical special functions. Theorem 1.4 extends this result to growing sequences satisfying $k \ll n^{1/(2\alpha+1)}$. Proposition 1.5 clarifies the asymptotic behavior of the limiting function $\gamma(k)$, as $k \rightarrow \infty$. This depends on the parameter α , and $\gamma(k)$ only scales with k^{-1} when $\alpha > 3/4$, which corresponds to the exponent of the degree distribution exceeding $5/2$. So in particular our findings disprove the abovementioned conjecture of Krioukov et al. [25].

Notation

In the statement of our main results, and throughout the rest of the paper, we will use the following notations. We set

$$\xi := \frac{4\alpha\nu}{\pi(2\alpha-1)}.$$

We write $\Gamma(z) := \int_0^\infty t^{z-1}e^{-t}dt$ for the gamma function, $\Gamma^+(a, b) := \int_b^\infty t^{a-1}e^{-t}dt$ for the upper incomplete gamma function, $B(a, b) := \int_0^1 u^{a-1}(1-u)^{b-1}du = \Gamma(a)\Gamma(b)/\Gamma(a+b)$ for the beta function and $B^-(x; a, b) := \int_0^x u^{a-1}(1-u)^{b-1}du$ for the lower incomplete beta function. We write $U(a, b, z)$ for the hypergeometric U-function (also called Tricomi's confluent hypergeometric function), which has the integral representation

$$U(a, b, z) = \frac{1}{\Gamma(a)} \int_0^\infty e^{-zt} t^{a-1} (1+t)^{b-a-1} dt,$$

see [15, p.255 Equation (2)], and let $G_{p,q}^{m,\ell} \left(z \left| \begin{smallmatrix} \mathbf{a} \\ \mathbf{b} \end{smallmatrix} \right. \right)$ denote Meijer's G-Function [28], see Appendix A for more details.

For a sequence $(X_n)_n$ of random variables, we write $X_n \xrightarrow[n \rightarrow \infty]{\mathbb{P}} X$ to denote that X_n converges in probability to X , as $n \rightarrow \infty$.

1.2 Main results

Here we summarize our main results regarding the degrees and clustering in the KPKVB model.

1.2.1 Degrees

The first rigorous result on the degree distribution in the KPKVB model was given in [21], where they analyzed the fraction of nodes with degree k . Although their results allowed $k = k_n$ to scale with n as $k_n = n^\delta$, it did not cover the full range $1 \leq k_n \leq n-1$. Our main result for the degrees gives this full characterization of the degree distribution.

Theorem 1.1. *Let $\alpha > \frac{1}{2}$, denote by $N_n(k)$ the number of vertices with degree k in the KPKVB model $G(n; \alpha, \nu)$ and consider a sequence of integers k_n with $0 \leq k_n \leq n-1$.*

1. *If $k_n = o\left(n^{\frac{1}{2\alpha+1}}\right)$ as $n \rightarrow \infty$, then a.a.s.*

$$N_n(k_n) = (1 + o(1))np_{k_n},$$

where

$$p_{k_n} = \int_0^\infty \mathbb{P}(\text{Po}(\xi e^{\frac{y}{2}}) = k_n) \alpha e^{-\alpha y} dy,$$

$$\text{and } p_{k_n} = \frac{2\alpha \xi^{2\alpha} \Gamma^+(k_n - 2\alpha, \xi)}{k_n!} = (1 + o(1))2\alpha \xi^{2\alpha} k_n^{-(2\alpha+1)} \text{ as } k_n \rightarrow \infty.$$

2. *If $k_n = (1 + o(1))cn^{\frac{1}{2\alpha+1}}$ for some fixed $c > 0$, then*

$$N_n(k_n) \xrightarrow[n \rightarrow \infty]{d} \text{Po}(2\alpha \xi^{2\alpha} c^{-(2\alpha+1)}).$$

3. *If $k_n \gg n^{\frac{1}{2\alpha+1}}$, then a.a.s. $N_n(k_n) = 0$.*

The first result of Theorem 1.1 states that the fraction of nodes of degree k_n converges to p_{k_n} . Not surprisingly, the limit distribution is the same as found in [21]. However, their this result was proved only for $k_n \leq n^{\delta'}$, where $\delta' < \min \left\{ \frac{2\alpha-1}{4\alpha(2\alpha+1)}, \frac{2(2\alpha-1)}{5(2\alpha+1)} \right\} < \frac{1}{2\alpha+1}$. Therefore, our result extends theirs by showing that this converges holds all the way till $n^{\frac{1}{2\alpha+1}}$. Moreover, we establish what happens at the boundary, where k_n scales as $n^{\frac{1}{2\alpha+1}}$ and show that this scaling is indeed the bound for the distribution since $N_n(k_n) = 0$ a.a.s. when k_n grows faster than $n^{\frac{1}{2\alpha+1}}$.

1.2.2 The clustering coefficient

Our first result on the clustering shows the convergence of the local clustering coefficient in the KPKVB model and establishes the limit exactly.

Theorem 1.2. *Let $\alpha > \frac{1}{2}$, $\nu > 0$ be fixed. Writing $G_n := G(n; \alpha, \nu)$, we have*

$$c(G_n) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \gamma,$$

where γ is defined for $\alpha \neq 1$ as

$$\begin{aligned} \gamma = & \frac{2 + 4\alpha + 13\alpha^2 - 34\alpha^3 - 12\alpha^4 + 24\alpha^5}{16(\alpha - 1)^2\alpha(\alpha + 1)(2\alpha + 1)} + \frac{2^{-1-4\alpha}}{(\alpha - 1)^2} \\ & + \frac{(\alpha - 1/2)(B(2\alpha, 2\alpha + 1) + B^-(1/2; 1 + 2\alpha, -2 + 2\alpha))}{2(\alpha - 1)(3\alpha - 1)} \\ & + \frac{\xi^{2\alpha}(\Gamma^+(1 - 2\alpha, \xi) + \Gamma^+(-2\alpha, \xi))}{4(\alpha - 1)} \\ & + \frac{\xi^{2\alpha+2}\alpha(\alpha - 1/2)^2(\Gamma^+(-2\alpha - 1, \xi) + \Gamma^+(-2\alpha - 2, \xi))}{2(\alpha - 1)^2} \\ & - \frac{\xi^{2\alpha+1}\alpha(2\alpha - 1)(\Gamma^+(-2\alpha, \xi) + \Gamma^+(-2\alpha - 1, \xi))}{(\alpha - 1)} \\ & - \frac{\xi^{6\alpha-2}2^{-4\alpha}(3\alpha - 1)(\Gamma^+(-6\alpha + 3, \xi) + \Gamma^+(-6\alpha + 2, \xi))}{(\alpha - 1)^2} \\ & - \frac{\xi^{6\alpha-2}(\alpha - 1/2)B^-(1/2; 1 + 2\alpha, -2 + 2\alpha)(\Gamma^+(-6\alpha + 3, \xi) + \Gamma^+(-6\alpha + 2, \xi))}{(\alpha - 1)} \\ & - \frac{e^{-\xi}\Gamma(2\alpha + 1)(U(2\alpha + 1, 1 - 2\alpha, \xi) + U(2\alpha + 1, 2 - 2\alpha, \xi))}{4(\alpha - 1)} \\ & + \frac{\xi^{6\alpha-2}\Gamma(2\alpha + 1)\left(G_{2,3}^{3,0}\left(\xi \left| \begin{smallmatrix} 1, 3 - 2\alpha \\ 3 - 4\alpha, -6\alpha + 2, 0 \end{smallmatrix} \right. \right) + G_{2,3}^{3,0}\left(\xi \left| \begin{smallmatrix} 1, 3 - 2\alpha \\ 3 - 4\alpha, -6\alpha + 3, 0 \end{smallmatrix} \right. \right)\right)}{4(\alpha - 1)}, \end{aligned}$$

and for $\alpha = 1$ as the $\alpha \rightarrow 1$ limit of the above expression.

A plot of γ can be found in Figure 2.




Figure 2: Plot of γ for α varying from 0.5 to 5 on the horizontal axis and for $\nu = \frac{1}{2}$ (blue), $\nu = 1$ (purple), $\nu = 2$ (green); simulations (squares in corresponding colour) with $n = 10000$ and 100 repetitions.

In the above expression for γ , a factor $\alpha - 1$ occurs in the denominator of each term, but we will see that this corresponds to a removable singularity. We have not been able to find a closed

form expression in terms of known functions in the case when $\alpha = 1$, but in Section 4.1.4 we do provide an explicit expression involving integrals.

1.2.3 The clustering function

Our second result is on the clustering function for constant k .

Theorem 1.3. *Let $\alpha > \frac{1}{2}$, $\nu > 0$ and $k \geq 2$ be fixed. Writing $G_n := G(n; \alpha, \nu)$, we have*

$$c(k; G_n) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \gamma(k),$$

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\left| \frac{c(k; G_n)}{\gamma(k)} - 1 \right| \right] = 0.$$

where $\gamma(k)$ is defined for $\alpha \neq 1$ as

$$\begin{aligned} \gamma(k) = & \frac{1}{8\alpha(\alpha-1)\Gamma^+(k-2\alpha, \xi)} \left(-\Gamma^+(k-2\alpha, \xi) - 2 \frac{\alpha(\alpha-1/2)^2 \xi^2 \Gamma^+(k-2\alpha-2, \xi)}{(\alpha-1)} \right. \\ & + 8\alpha(\alpha-1/2) \xi \Gamma^+(k-2\alpha-1, \xi) \\ & + 4\xi^{4\alpha-2} \Gamma^+(k-6\alpha+2, \xi) \left(\frac{2^{-4\alpha}(3\alpha-1)}{(\alpha-1)} + (\alpha-1/2) B^-(1/2; 1+2\alpha, -2+2\alpha) \right) \\ & + \xi^{k-2\alpha} \Gamma(2\alpha+1) e^{-\xi} U(2\alpha+1, 1+k-2\alpha, \xi) \\ & \left. - \xi^{4\alpha-2} \Gamma(2\alpha+1) G_{2,3}^{3,0} \left(\xi \left| \begin{matrix} 1, 3-2\alpha \\ 3-4\alpha, -6\alpha+k+2, 0 \end{matrix} \right. \right) \right) \end{aligned}$$

and for $\alpha = 1$ as the $\alpha \rightarrow 1$ limit of the above expression.

A plot of $\gamma(k)$ can be found in Figure 3. Again, we remark that the above expression for $\gamma(k)$




Figure 3: Plot $\gamma(k)$ for k varying from 2 to 25 on the horizontal axis, for $\alpha = 0.8$ and $\nu = \frac{1}{2}$ (blue), $\nu = 1$ (purple), $\nu = 2$ (green); simulations (squares in corresponding colour) with $n = 10000$ and 100 repetitions.

appears to have a singularity at $\alpha = 1$, but this will turn out to be a removable singularity. Again, we have not been able to find a closed form expression in terms of known functions in the case when $\alpha = 1$, but in Section 4.1.4 we do provide an explicit expression involving integrals.

Theorem 1.3 in fact generalises to increasing sequences $(k_n)_{n \geq 1}$.

Theorem 1.4. *Let $\alpha > \frac{1}{2}$, $\nu > 0$ be fixed and let k_n be a sequence of non-negative integers satisfying $1 \ll k_n \ll n^{1/(2\alpha+1)}$. Then, writing $G_n := G(n; \alpha, \nu)$, we have*

$$\mathbb{E}[|c(k_n; G_n) - \gamma(k_n)|] = o(\gamma(k_n)),$$

as $n \rightarrow \infty$, where $\gamma(\cdot)$ is as in Theorem 1.3. In particular

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\left| \frac{c(k_n; G_n)}{\gamma(k_n)} - 1 \right| \right] = 0.$$

1.2.4 Scaling of $\gamma(k)$

To clarify the scaling behaviour of $\gamma(k)$ with k we offer the following result.

Proposition 1.5. *As $k \rightarrow \infty$, we have*

$$\gamma(k) = (1 + o(1)) \cdot \begin{cases} \frac{8\alpha\nu}{\pi(4\alpha-3)} \cdot k^{-1} & \text{if } \alpha > \frac{3}{4}, \\ \frac{6\nu}{\pi} \cdot \frac{\log(k)}{k} & \text{if } \alpha = \frac{3}{4}, \\ c_\alpha \cdot k^{2-4\alpha} & \text{if } \frac{1}{2} < \alpha < \frac{3}{4}, \end{cases},$$

$$\text{where } c_\alpha := \left(\frac{3\alpha-1}{2^{4\alpha+1}\alpha(\alpha-1)^2} + \frac{(\alpha-\frac{1}{2})B^-(\frac{1}{2}, 2\alpha+1, 2\alpha-2)}{2(\alpha-1)\alpha} - \frac{B(2\alpha, 3\alpha-4)}{4(\alpha-1)} \right) \cdot \xi^{4\alpha-2}.$$

Note that Theorem 1.4 implies that the clustering function of the KPKVB model scales as $\gamma(k)$ as k grows, whose scaling is given in the above result. In particular, this contradicts the scaling conjectured in [25] for $\alpha \leq \frac{3}{4}$, and confirms it only for $\alpha > \frac{3}{4}$.

We remark that simultaneously and independently Stegehuis, van der Hofstad and van Leeuwen [33] used a completely different technique to obtain a similar, though less detailed, result on the $k \rightarrow \infty$ scaling of the clustering function in the KPKVB model.

1.3 Observations

There are a few other observation to be made regarding our results on clustering.

Uniform convergence. Our results for the local clustering function in particular imply uniform convergence of $c(k; G_n)$ for all $2 \leq k \leq a_n$ where $a_n \ll n^{\frac{1}{2\alpha+1}}$. To see this let

$$b_n = \arg \max_{2 \leq k \leq a_n} \mathbb{E} \left[\left| \frac{c(k; G_n)}{\gamma(k)} - 1 \right| \right].$$

Then $b_n \leq a_n \ll n^{\frac{1}{2\alpha+1}}$ and therefore by Theorem 1.4

$$\lim_{n \rightarrow \infty} \max_{2 \leq k \leq a_n} \mathbb{E} \left[\left| \frac{c(k; G_n)}{\gamma(k)} - 1 \right| \right] = \lim_{n \rightarrow \infty} \mathbb{E} \left[\left| \frac{c(b_n; G_n)}{\gamma(b_n)} - 1 \right| \right] = 0.$$

Near maximum scaling for k_n . Our results for the clustering function in the KPKVB model are valid for any sequence of non-negative integers $k_n \rightarrow \infty$ such that $k_n \ll n^{\frac{1}{2\alpha+1}}$. Although one would like to have results for any sequence $k_n \leq n-1$, it turns out that $n^{1/(2\alpha+1)}$ is the near maximum scaling for which Theorem 1.2 can be true. To see why this is the case note that by definition of the local clustering function (2) we have that $c(k_n; G_n) = 0$ if $N_n(k_n) = 0$, where $N_n(k_n)$ denotes the number of vertices in a KPKVB graph $G(n; \alpha, \nu)$ with degree k_n . It follows by Markov's inequality that for any positive function $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$

$$\begin{aligned} \mathbb{E} \left[\left| \frac{c(k_n; G_n)}{f(k_n)} - 1 \right| \right] &\geq \mathbb{E} \left[\left| \frac{c(k_n; G_n)}{f(k_n)} - 1 \right| \mathbb{1}_{\{N_n(k_n)=0\}} \right] \\ &\geq \mathbb{P}(N_n(k_n) = 0) \geq 1 - \mathbb{E}[N_n(k_n)]. \end{aligned}$$

We shall later establish (see Lemma 3.12) that $\mathbb{E}[N_n(k_n)]$ scales as $nk_n^{-(2\alpha+1)}$. Therefore if k_n is such that $k_n^{-(2\alpha+1)}n$ tends to zero as $n \rightarrow \infty$ we have that

$$\lim_{n \rightarrow \infty} \mathbb{E}[N_n(k_n)] = 0$$

and hence

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\left| \frac{c(k_n; G_n)}{f(k_n)} - 1 \right| \right] \geq 1 - \lim_{n \rightarrow \infty} \mathbb{E}[N_n(k_n)] = 1 \neq 0,$$

for any positive function f . This implies that we cannot expect a result like that of Theorem 1.4 to hold as soon as $k_n \gg n^{\frac{1}{2\alpha+1}}$.

Transition in scaling at $\alpha = 3/4$. It follows from Proposition 1.5 that there is a transition in the scaling of the local clustering function at $\alpha = 3/4$. This corresponds to an exponent $5/2$ for the probability mass function of the degree distribution. This transition is different from those often observed for networks with scale-free degree distributions, where transitions occur at integer values of the exponent. At this point, it is unclear what the underlying reason is for the appearance of the transition at this half exponent. Interestingly, a similar transition point has also been observed for both majority vote models [12] and flocking dynamics [29] on networks with scale-free degree distributions.

1.4 Outline of the paper

In the next section we will recall some useful tools from the literature and define a series of auxiliary random graph models that will be used in the proofs. In particular, we relate in a series of steps the KPKVB model to an infinite percolation model G_∞ that was used in previous work of the first and third authors [17] on the largest component of the KPKVB model. The value of the limiting constant γ , respectively limiting clustering function $\gamma(k)$, correspond to the probability that two randomly chosen neighbours of a “typical point” in this infinite model are themselves neighbours, respectively the probability of this event conditional on the typical point having exactly k neighbours. These probabilities can be expressed as certain integrals, which we solve explicitly in Section 4. In the same section we also prove Proposition 1.5, on the asymptotics of $\gamma(k)$. We then proceed to prove Theorems 1.2 and 1.3 by relating said probabilities for the typical point of the infinite model to the corresponding clustering coefficient/function in the original KPKVB random graph, using the Campbell-Mecke formula and some other, relatively straightforward considerations.

The remaining sections are devoted to the proof of Theorem 1.4, which turns out to be a lot more involved. The main reason for this is that we push the possible scaling of k_n to its maximum and hence a great deal of work is needed to properly control the arising error terms and make sure these are of smaller order than $\gamma(k_n)$.

Finally, the Appendix includes some technical results on Meijer’s G-function, Chernoff bounds for Poisson and Binomial random variables and the code used for simulations.

2 Preliminaries

In this section we recall some definitions and tools that we will need in our proofs.

2.1 The infinite limit model G_∞

We start by recalling the definition of the infinite limit model from [17]. Let $\mathcal{P} = \mathcal{P}_{\alpha, \nu}$ be a Poisson point process on \mathbb{R}^2 with intensity function $f = f_{\alpha, \nu}$ given by

$$f(x, y) = \frac{\alpha\nu}{\pi} e^{-\alpha y} \cdot \mathbb{1}_{\{y>0\}}. \quad (3)$$

The *infinite limit model* $G_\infty = G_\infty(\alpha, \nu)$ has vertex set \mathcal{P} and edge set such that

$$pp' \in E(G_\infty) \iff |x - x'| \leq e^{\frac{y+y'}{2}},$$

for $p = (x, y), p' = (x', y') \in \mathcal{P}$.

For any point $p \in \mathbb{R} \times (0, \infty)$, we write $\mathcal{B}_\infty(p)$ to denote the *ball* around p , i.e.

$$\mathcal{B}_\infty(p) = \{p' \in \mathbb{R} \times (0, \infty) : |x - x'| \leq e^{\frac{y+y'}{2}}\}. \quad (4)$$

With this notation we then have that $\mathcal{B}_\infty(p) \cap \mathcal{P}$ denotes the set of neighbours of a vertex $p \in G_\infty$. We will denote the intensity measure of the Poisson process \mathcal{P} by $\mu = \mu_{\alpha, \nu}$, i.e. for every Borel-measurable subset $S \subseteq \mathbb{R}^2$ we have $\mu(S) = \int_S f(x, y) dx dy$.

2.2 The finite box model G_{box}

For the definition of the finite graph, recall that in the definition of the KPKVB model we set $R = 2 \log(n/\nu)$. We consider the box $\mathcal{R} = (-\frac{\pi}{2}e^{R/2}, \frac{\pi}{2}e^{R/2}] \times (0, R]$ in \mathbb{R}^2 . Then the *finite box model* $G_{\text{box}} := G_{\text{box}}(n; \alpha, \nu)$ has vertex set $V_{\text{box}} := \mathcal{P} \cap \mathcal{R}$ and edge set such that

$$pp' \in E(G_{\text{box}}(n; \alpha, \nu)) \iff |x - x'|_{\pi e^{R/2}} \leq e^{\frac{y+y'}{2}},$$

where $|x|_r = \min(|x|, r - |x|)$ for $-r \leq x \leq r$. Using $|\cdot|_{\pi e^{R/2}}$ instead of $|\cdot|$ results in the left and right boundaries of the box \mathcal{R} getting identified, which in particular makes the model invariant under horizontal shifts and reflections in vertical lines. The graph G_{box} can thus be seen as a subgraph of G_∞ induced on V_{box} , with some additional edges caused by the identification of the boundaries.

Similar to the infinite graph, for a point $p \in \mathcal{R}$ we define the ball $\mathcal{B}_{\text{box}}(p)$ as

$$\mathcal{B}_{\text{box}}(p) = \left\{p' \in \mathcal{R} : |x - x'|_{\pi e^{R/2}} \leq e^{\frac{y+y'}{2}}\right\}. \quad (5)$$

2.3 The Poissonized KPKVB model G_{Po}

Imagine that we have an infinite supply of i.i.d. points u_1, u_2, \dots in the hyperbolic plane \mathbb{H} chosen according to the (α, R) -quasi uniform distribution. In the standard KPKVB random graph $G(n; \alpha, \nu)$ we take u_1, \dots, u_n as our vertex set and add edges between points at hyperbolic distance at most $R = 2 \log(n/\nu)$. In the *Poissonized* KPKVB random graph $G_{\text{Po}} := G_{\text{Po}}(n; \alpha, \nu)$, we instead take $N \stackrel{\text{d}}{=} \text{Po}(n)$, a Poisson random variable with mean n , independent of our i.i.d. sequence of points and let the vertex set be u_1, \dots, u_N and add edges according to the same rule as before. Equivalently, we could say that the vertex set consists of the points of a Poisson point process with intensity function ng , where g denotes the probability density of the (α, R) -quasi uniform distribution. That is,

$$g(r, \theta) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)} \cdot 1_{\{0 \leq r \leq R, -\pi < \theta \leq \pi\}}. \quad (6)$$

Working with the Poissonized model has the advantage that when we take two disjoint regions A, B then the number of points in A and the number of points in B are independent Poisson-distributed random variables. As we will see, and as is to be expected, switching to the Poissonized model does not significantly alter the limiting behaviour of the clustering coefficient and function.

2.4 Coupling G_{Po} and G_{box}

The following lemmas from [17] establish a useful coupling between the Poissonized KPKVB random graph and the finite box model and relate the edge sets of the two graphs.

Lemma 2.1 ([17, Lemma 27]). *Let V_{Po} denote the vertex set of $G_{\text{Po}}(n; \alpha, \nu)$ and V_{box} the vertex set of $G_{\text{box}}(n; \alpha, \nu)$. Define the map $\Psi : [0, R] \times (-\pi, \pi] \rightarrow \mathcal{R}$ by*

$$\Psi(r, \theta) = \left(\theta \frac{e^{R/2}}{2}, R - r\right). \quad (7)$$

Then there exists a coupling such that, a.a.s., $V_{\text{box}} = \Psi[V_{\text{Po}}]$.

In the remainder of this paper we will write $\mathcal{B}(p)$ to denote the image under Ψ of the ball of hyperbolic radius R around the point $\Psi^{-1}(p)$ for $p \in \mathcal{R}$, i.e.

$$\mathcal{B}(p) := \Psi \left[\{u \in \mathbb{H} : d_{\mathbb{H}}(\Psi^{-1}(p), u), d_{\mathbb{H}}(O, u) \leq R\} \right] \subset \mathcal{R}.$$

Under the map Ψ , a point $p = (x, y) \in \mathcal{R}$ corresponds to $u := \Psi^{-1}(p) = (2e^{-R/2}x, R - y)$.

By the hyperbolic rule of cosines, for two points $p = (x, y) = \Psi((r, \theta)), p' = (x', y') = \Psi((r', \theta')) \in \mathcal{R}$ we have that $p' \in \mathcal{B}(p)$ iff. either $r + r' \leq R$ or $r + r' > R$ and

$$\cosh r \cosh r' - \sinh r \sinh r' \cos(|\theta - \theta'|_{2\pi}) \leq \cosh(R),$$

This can be rephrased as $p' \in \mathcal{B}(p)$ iff. either $y + y' \geq R$ or $y + y' < R$ and

$$|x - x'|_{\pi e^{R/2}} \leq \Phi(y, y') := \frac{1}{2}e^{R/2} \arccos \left(\frac{\cosh(R - y) \cosh(R - y') - \cosh R}{\sinh(R - y) \sinh(R - y')} \right). \quad (8)$$

The following lemma provides useful bounds on the function $\Phi(r, r')$. Note that in [17] the function Φ is written in terms of where $r := R - y, r' := R - y'$.

Lemma 2.2 ([17, Lemma 28]). *There exists a constant $K > 0$ such that, for every $\varepsilon > 0$ and for R sufficiently large, the following holds. For every $r, r' \in [\varepsilon R, R]$ with $y + y' < R$ we have that*

$$e^{\frac{1}{2}(y+y')} - Ke^{\frac{3}{2}(y+y')-R} \leq \Phi(y, y') \leq e^{\frac{1}{2}(y+y')} + Ke^{\frac{3}{2}(y+y')-R}, \quad (9)$$

Moreover:

$$\Phi(y, y') \geq e^{\frac{1}{2}(y+y')} \quad \text{if} \quad y, y' > K. \quad (10)$$

A key consequence of Lemma 2.2 is that the coupling from Lemma 2.1 preserves edges between points whose heights are not too large.

Lemma 2.3 ([17, Lemma 30]). *On the coupling space of Lemma 2.1 the following holds a.a.s.:*

1. for any two points $p, p' \in V_{\text{box}}$ with $y, y' \leq R/2$, we have

$$pp' \in E(G_{\text{box}}) \Rightarrow \Psi^{-1}(p)\Psi^{-1}(p') \in E(G_{\text{Po}}),$$

2. for any two points $p, p' \in V_{\text{box}}$ with $y, y' \leq R/4$, we have that

$$pp' \in E(G_{\text{box}}) \iff \Psi^{-1}(p)\Psi^{-1}(p') \in E(G_{\text{Po}}).$$

Remark 2.1 (Notion convention for points). *We will often be working with the finite box graph G_{Po} or the infinite graph G_{∞} , whose nodes are points in $\mathbb{R} \times \mathbb{R}_+$. For any point $p \in \mathbb{R} \times \mathbb{R}_+$ we will always use $p = (x, y)$. When considering different points $p, p' \in \mathbb{R} \times \mathbb{R}_+$, we will use primed coordinates to refer to p' , i.e. $p' = (x', y')$, and similar with subscripts, i.e. $p_i = (x_i, y_i)$.*

2.5 The Campbell-Mecke formula

A very useful tool for analyzing subgraph counts, and their generalizations, in the setting of the Poissonized random geometric graphs, and in particular the Poissonized KPKVB model and the box model is the *Campbell-Mecke formula*. We use a specific incarnation, which follows from the Palm theory of Poisson point processes on metric spaces, see [26]. For this consider a Poisson point process \mathcal{P} on some metric space \mathcal{M} with density μ and let \mathcal{N} denote the set of all possible point configurations in \mathcal{M} , equipped with the sigma algebra of the process \mathcal{P} . Then, for any natural number k and measurable function $h : \mathbb{R}^k \times \mathcal{N} \rightarrow \mathbb{R}$,

$$\mathbb{E} \left[\sum_{p_1, \dots, p_k \in \mathcal{P}}^{\neq} h(p_1, \dots, p_k, \mathcal{P}) \right] = \int_{\mathcal{M}} \cdots \int_{\mathcal{M}} \mathbb{E} [h(x_1, \dots, x_k, \mathcal{P})] d\mu(x_1) \cdots d\mu(x_k), \quad (11)$$

where the sum is over all distinct point $p_1, \dots, p_k \in \mathcal{P}$.

2.6 Concentration of heights

When analyzing degrees and clustering in the Poissonized KPKVB and related models we often encounter expressions of the form

$$\int_0^R \mathbb{P}(\text{Po}(\hat{\mu}(y)) = k_n) h(y) e^{-\alpha y} dy, \quad (12)$$

where $h(y)$ is some function and $\hat{\mu}(y)$ is $\mu(\mathcal{B}(y))$, $\mu(\mathcal{B}_{\text{box}}(y))$ or $\mu(\mathcal{B}_{\infty}(y))$. We will often have to either bound the behavior of such integrals as $k_n \rightarrow \infty$ or establish their asymptotic behavior. For this we will utilize that Poisson random variables are well concentrated around their mean.

In particular, if $\text{Po}(\lambda)$ denotes a Poisson random variable with mean λ , we have the following Chernoff bound (c.f. [32, Lemma 1.2])

$$\mathbb{P}(|\text{Po}(\lambda) - \lambda| \geq x) \leq 2e^{-\frac{x^2}{2(\lambda+x)}}. \quad (13)$$

In particular, if $\lambda = \lambda_n \rightarrow \infty$, then for any $C > 0$,

$$\mathbb{P}(|\text{Po}(\lambda_n) - \lambda_n| \geq C\sqrt{\lambda_n \log(\lambda_n)}) \leq O\left(\lambda_n^{-\frac{C^2}{2}}\right). \quad (14)$$

For our application these Chernoff bounds imply that if y is such that $\hat{\mu}(y)$ is far from k_n then $\mathbb{P}(\text{Po}(\hat{\mu}(y)) = k_n)$ becomes very small. To be more specific, we define for any $k \geq 0$ and $C > 0$,

$$y_{k,C}^{\pm} = 2 \log \left(\frac{k \pm C\sqrt{k \log(k)}}{\xi} \right), \quad (15)$$

where we set $y_{k,C}^{-} = 0$ if $k - C\sqrt{k \log(k)} < \xi$. For convenience we write $\mathcal{K}_C(k_n) := [y_{k_n,C}^{-}, y_{k_n,C}^{+}]$. Then we can show that for all y outside $\mathcal{K}_C(k_n)$

$$\mathbb{P}(\text{Po}(\hat{\mu}(y)) = k_n) \leq O\left(k_n^{-\frac{C^2}{2}}\right). \quad (16)$$

Since we can select C to be as big as we want we can make this error as small as needed. This implies that then the main contribution to the integral (12) comes from those “heights” y that are in the interval $\mathcal{K}_C(k_n)$. In other words, the main contribution is concentrated around the heights y for which $\mu(y) = k_n$. We thus refer to this as the concentration of heights result. More precisely, we prove the following.

Proposition 2.4 (Concentration of heights). *Let $\alpha > \frac{1}{2}$, $\nu > 0$, $(k_n)_{n \geq 1}$ be any positive sequence such that $k_n \rightarrow \infty$ and $k_n = o(n)$. Furthermore, let $\hat{\mu}(y)$ denoting either $\mu(\mathcal{B}(y))$, $\mu(\mathcal{B}_{\text{box}}(y))$ or $\mu(\mathcal{B}_{\infty}(y))$. Then for any continuous function $h : \mathbb{R}_+ \rightarrow \mathbb{R}$, such that $h(y) = O(e^{\beta y})$ as $y \rightarrow \infty$ for some $\beta < \alpha$, it holds that*

$$\int_0^{\infty} h(y) \mathbb{P}(\text{Po}(\hat{\mu}(y)) = k_n) \alpha e^{-\alpha y} dy \sim \int_{\mathcal{K}_C(k_n)} h(y) \mathbb{P}(\text{Po}(\hat{\mu}(y)) = k_n) \alpha e^{-\alpha y} dy, \quad (17)$$

as $n \rightarrow \infty$.

The key implication of Proposition 2.4 is that if the function $h(y)$ does not increase too fast, then we can restrict integration to the interval $\mathcal{K}_C(y)$. The full details associated with these concentration of heights and the proof of Proposition 2.4 can be found in the Section E of the Appendix.

3 Degrees in the KPKVK model: proof of Theorem 1.1

Since the new contribution of Theorem 1.1 concerns the cases where the degree $k_n \rightarrow \infty$, we will assume that this holds throughout this section.

3.1 Proof overview

We start by using the Campbell-Mecke formula to compare the degree distribution in G_{Po} with that of a *typical point* in G_∞ . For this we will be considering the Poisson process \mathcal{P} to which we add one additional point $(0, y)$ on the y -axis. In some computations the height y will be fixed, but eventually we shall take it exponentially distributed with parameter α , and independent of \mathcal{P} . We refer to $(0, y)$ as “the typical point”.

To provide some intuition for this definition and name, note that we can alternatively view \mathcal{P} as follows. We take a constant intensity Poisson process on \mathbb{R} corresponding to the x -coordinates, and to each point we attach a random “mark”, corresponding to the y -coordinate, where the marks are i.i.d. exponentially distributed with parameter α . By translation invariance of Poisson point process, each point will behave as $(0, y)$ with y exponentially distributed with parameter α , the typical point of the process.

We first compute the degree distribution of the typical point in G_∞ and show that this equals

$$p_k = \int_0^\infty \rho(y, k) \alpha e^{-\alpha y} dy.$$

We then relate this to the Poissonized KPKVB model G_{Po} . More precisely, let $N_{\text{Po}}(k)$ denote the set of degree k vertices in G_{Po} . We then show in Section 3.3 that for any $1 \ll k_n \leq n-1$ with $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$, $\mathbb{E}[N_{\text{Po}}(k_n)] = (1 + o(1))np_{k_n}$ and more generally,

$$\mathbb{E}\left[\binom{N_{\text{Po}}(k_n)}{r}\right] = (1 + o(1)) \frac{\mathbb{E}[N_{\text{Po}}(k_n)]^r}{r!}, \quad (18)$$

for any integer $r \geq 1$ in Section 3.5. The latter result requires us to analyze the joint degree distribution in G_{Po} , which we do in Section 3.4. The above result in particular implies concentration of $N_{\text{Po}}(k_n)$ from which the result on the degree distribution in G_{Po} follows for $k_n = o\left(n^{\frac{1}{2\alpha+1}}\right)$.

When $k_n = (1 + o(1))cn^{\frac{1}{2\alpha+1}}$ we use the above result to show that the fraction of degree k_n nodes in G_{Po} converges to a Poisson distribution.

To extend these results to G_n we couple the construction of the KPKVB model to that of the Poissonized version G_{Po} in Section 3.6 to show that a.a.s, $N_n(k_n) = (1 + o(1))N_{\text{Po}}(k_n)$. With these results we then prove Theorem 1.1 in Section 3.7.

We will also establish all the above mention results for the finite box model G_{box} , since the proofs only require small alterations and we will need these results later on when analyzing the clustering coefficient and function.

3.2 The degree of the typical point in G_∞

We start with investigating the distribution of the degree of the typical point. (Considered in the graph obtained from G_∞ by adding the typical point to its vertex set.)

If $p = (x, y) \in \mathbb{R} \times [0, \infty)$ is a point, not necessarily part of the Poisson process, then we will write

$$\mu(y) = \mu(p) := \mu(\mathcal{B}_\infty(p)).$$

Integrating the intensity function of \mathcal{P} over $\mathcal{B}_\infty(p)$ gives, using $\alpha > \frac{1}{2}$,

$$\mu(y) = \int_0^\infty \int_{-e^{(y+y')/2}}^{e^{(y+y')/2}} \frac{\alpha \nu}{\pi} e^{-\alpha y'} dx' dy' = \frac{2\alpha \nu e^{y/2}}{\pi} \int_0^\infty e^{(\frac{1}{2}-\alpha)y'} dy' = \frac{2\alpha \nu e^{y/2}}{\pi(\alpha - \frac{1}{2})} = \xi e^{y/2}.$$

For $p = (x, y) \in \mathbb{R} \times [0, \infty)$ we define

$$\rho(p, k) := \mathbb{P}(\text{Po}(\mu(p)) = k), \quad (19)$$

where $\text{Po}(\lambda)$ denotes a Poisson random variable with expectation λ . When $p = (0, y)$ we often write $\rho(y, k)$ instead of $\rho(p, k)$.

Let the random variable D denote the degree of the typical point. Since the typical point has a height that is independent of the Poisson process and exponential(α)-distributed, for $k \in \mathbb{N}_0$:

$$p_k := \mathbb{P}(D = k) = \int_0^\infty \rho(y, k) \alpha e^{-\alpha y} dy. \quad (20)$$

Using the transformation of variables $z = \xi e^{\frac{y}{2}}$ (so $dy = \frac{2}{z} dz$), we compute

$$\begin{aligned} p_k &= \frac{1}{k!} \int_0^\infty \left(\xi e^{\frac{y}{2}} \right)^k e^{-\xi e^{\frac{y}{2}}} \alpha e^{-\alpha y} dy = \frac{\alpha \xi^{2\alpha}}{k!} \int_0^\infty \left(\xi e^{\frac{y}{2}} \right)^{k-2\alpha} e^{-\xi e^{\frac{y}{2}}} dy \\ &= \frac{2\alpha \xi^{2\alpha}}{k!} \int_\xi^\infty z^{k-2\alpha-1} e^{-z} dz = \frac{2\alpha \xi^{2\alpha} \Gamma^+(k-2\alpha, \xi)}{k!}, \end{aligned}$$

where we recall that Γ^+ denotes the upper incomplete Gamma-function. Note that, unsurprisingly, this is identical to the expression Gugelmann et al. [21] gave for the limiting degree distribution of $G(n; \alpha, \nu)$. Using Stirling's approximation to the gamma function, we find that

$$p_k \sim 2\alpha \xi^{2\alpha} k^{-(2\alpha+1)} \quad \text{as } k \rightarrow \infty. \quad (21)$$

By a similar computation we have the following result, which will be useful later on. For any $\beta > 0$, as $k \rightarrow \infty$

$$\int_0^\infty e^{\beta y} \rho(y, k) \alpha e^{-\alpha y} dy \sim 2\alpha \xi^{2(\beta+\alpha)} k^{-2(\beta+\alpha)-1}. \quad (22)$$

3.3 Expected degrees in G_{box} and G_{Po}

We proceed with the expected degrees in the finite box and Poissonized KPKVB model. Recall the definition of the neighbourhood balls $\mathcal{B}_{\text{box}}(y)$ and $\mathcal{B}(y)$ of a point $(0, y)$ in, respectively G_{box} and G_{Po} . We introduce the short hand notation

$$\mu_{\text{Po}}(y) := \mu(\mathcal{B}(y)) \quad \text{and} \quad \mu_{\text{box}}(y) := \mu(\mathcal{B}_{\text{box}}(y)).$$

Our first results relate these measures to the measure $\mu(y)$, of the ball $\mathcal{B}_\infty(y)$ in the infinite model G_∞ .

Lemma 3.1 (Expected degree given height in G_{Po}). *Let $\alpha > \frac{1}{2}$, $\varepsilon > 0$ and $0 \leq y \leq (1 - \varepsilon)R$. Then as $n \rightarrow \infty$, uniformly in y ,*

$$\mu_{\text{Po}}(y) = (1 + o(1))\mu(y).$$

Proof. Recall that when $y' \geq R - y$ then $p' \in \mathcal{B}(y)$ while for $y' < R - y$ this is true when (8) holds. We split the integral for $\mu_{\text{Po},n}(y)$ accordingly, into two integrals I_1 and I_2 ,

$$\mu_{\text{Po}}(y) = \int_0^{R-y} 2\Phi(y, y_1) \frac{\alpha \nu}{\pi} e^{-\alpha y_1} dy_1 + \int_{R-y}^R \frac{\pi n}{\nu} \frac{\alpha \nu}{\pi} e^{-\alpha y_1} dy_1 =: I_1 + I_2.$$

Firstly, we will show that the second integral $I_2 = o(\mu(y))$ and then we will show that $I_1 = (1 + o(1))\mu(y)$ (both with convergence uniform in $0 \leq y \leq (1 - \varepsilon)R$).

For the second integral I_2 , we compute

$$I_2 = \int_{R-y}^R \frac{\pi n}{\nu} \frac{\alpha \nu}{\pi} e^{-\alpha y_1} dy_1 = n(e^{-\alpha(R-y)} - e^{-\alpha R}) = n e^{-\alpha R} (e^{\alpha y} - 1) = n^{1-2\alpha} \nu^{2\alpha} (e^{\alpha y} - 1).$$

To see that $n^{1-2\alpha} \nu^{2\alpha} (e^{\alpha y} - 1) = o(\mu(y))$, recall that $\mu(y) = \xi e^{\frac{y}{2}}$. So, we need to show that

$$\frac{n^{1-2\alpha} \nu^{2\alpha} (e^{\alpha y} - 1)}{\xi e^{\frac{y}{2}}} = o(1)$$

or equivalently that

$$\frac{e^{\alpha y} - 1}{\xi e^{\frac{y}{2}}} = o(n^{2\alpha-1}).$$

For this, note that

$$\frac{e^{\alpha y} - 1}{\xi e^{\frac{y}{2}}} = O\left(e^{\alpha y - \frac{y}{2}}\right) = O\left(e^{(\alpha - \frac{1}{2})y}\right).$$

As $y \leq (1 - \epsilon)R = (1 - \epsilon)2\ln \frac{n}{\nu}$ and $\alpha > \frac{1}{2}$, we have

$$e^{(\alpha - \frac{1}{2})y} \leq e^{(\alpha - \frac{1}{2})(1 - \epsilon)R} = \left(\frac{n}{\nu}\right)^{2(\alpha - \frac{1}{2})(1 - \epsilon)} = o(n^{2\alpha-1}),$$

where the convergence is uniform in y , $0 \leq y \leq (1 - \epsilon)R$, as the last upper bound does not depend on y .

For the first integral I_1 , we first recall from Lemma 2.2 that there is a positive constant K such that for any $\varepsilon > 0$, for all $y_1, y_2 \in [0, (1 - \varepsilon)R]$, $y_1 + y_2 < R$, we have

$$e^{\frac{1}{2}(y_1 + y_2)} - K e^{\frac{3}{2}(y_1 + y_2) - R} \leq \Phi(y_1, y_2) \leq e^{\frac{1}{2}(y_1 + y_2)} + K e^{\frac{3}{2}(y_1 + y_2) - R}.$$

We thus define the main and error term of I_1 as

$$\begin{aligned} I_{1,main} &= \int_0^{R-y} 2e^{\frac{y+y_1}{2}} \frac{\alpha\nu}{\pi} e^{-\alpha y_1} dy_1, \\ I_{1,error} &= \int_0^{R-y} 2K e^{\frac{3}{2}(y+y_1) - R} \frac{\alpha\nu}{\pi} e^{-\alpha y_1} dy_1. \end{aligned}$$

From the error bounds for Φ as given in Lemma 2.2, it follows that

$$I_{1,main} - I_{1,error} \leq I_1 \leq I_{1,main} + I_{1,error}.$$

We will firstly show that $I_{1,main} = (1 + o(1))\mu(y)$ and then that $I_{1,error} = o(\mu(y))$.

For the main term, we obtain, as $R - y \geq \epsilon R \rightarrow \infty$, uniformly in $0 \leq y \leq (1 - \epsilon)R$,

$$\begin{aligned} I_{1,main} &= \int_0^{R-y} 2e^{\frac{y+y_1}{2}} \frac{\alpha\nu}{\pi} e^{-\alpha y_1} dy_1 = \frac{2\alpha\nu}{\pi} e^{\frac{y}{2}} \int_0^{R-y} e^{(\frac{1}{2} - \alpha)y_1} dy_1 \\ &= \frac{2\alpha\nu}{\pi(\alpha - \frac{1}{2})} e^{\frac{y}{2}} \left(1 - e^{(\frac{1}{2} - \alpha)(R-y)}\right) = (1 + o(1))\xi e^{\frac{y}{2}} = (1 + o(1))\mu(y). \end{aligned}$$

For the error term, we obtain, for $\alpha \neq \frac{3}{2}$, uniformly in $0 \leq y \leq (1 - \epsilon)R$,

$$\begin{aligned} I_{1,error} &= \int_0^{R-y} 2K e^{\frac{3}{2}(y+y_1) - R} \frac{\alpha\nu}{\pi} e^{-\alpha y_1} dy_1 = 2K \frac{\alpha\nu}{\pi} e^{\frac{3}{2}y - R} \int_0^{R-y} e^{(\frac{3}{2} - \alpha)y_1} dy_1 \\ &= \frac{2K\alpha\nu}{\pi(\frac{3}{2} - \alpha)} e^{\frac{3}{2}y - R} \left(e^{(\frac{3}{2} - \alpha)(R-y)} - 1\right) \\ &= \frac{2K\alpha\nu}{\pi(\frac{3}{2} - \alpha)} e^{\frac{1}{2}y} \left(e^{(\frac{1}{2} - \alpha)(R-y)} - e^{-(R-y)}\right) = o\left(\xi e^{\frac{y}{2}}\right). \end{aligned}$$

For the error term with $\alpha = \frac{3}{2}$, uniformly in $0 \leq y \leq (1 - \epsilon)R$,

$$\int_0^{R-y} 3K e^{\frac{3}{2}(y+y_1) - R} \frac{\nu}{\pi} e^{-\frac{3}{2}y_1} dy_1 = 3K \frac{\nu}{\pi} e^{\frac{3}{2}y - R} \int_0^{R-y} dy_1 = \frac{3K\nu}{\pi} e^{\frac{3}{2}y - R} (R - y) = o\left(\xi e^{\frac{y}{2}}\right).$$

We conclude that $I_{1,error} = o(\mu(y))$ and hence $I_{1,main} \pm I_{1,error} = (1 + o(1))\mu(y)$, which finishes the proof. \square

Lemma 3.2 (Expected degree given height in G_{box}). *Let $\alpha > \frac{1}{2}$, $\varepsilon > 0$ and $0 \leq y \leq (1 - \varepsilon)R$. Then as $n \rightarrow \infty$, uniformly in y ,*

$$\mu_{\text{box}}(y) = (1 + o(1))\mu(y).$$

Proof. First note that since we have identified the boundaries of $[-\frac{\pi}{2}e^{\frac{R}{2}}, \frac{\pi}{2}e^{\frac{R}{2}}]$ we can assume, without loss of generality, that $p = (0, y)$. We then have that the boundaries of $\mathcal{B}_{\text{box}}(p)$ are given by the equations $x' = \pm e^{\frac{y+y'}{2}}$, which intersect the left and right boundaries of $[-\frac{\pi}{2}e^{\frac{R}{2}}, \frac{\pi}{2}e^{\frac{R}{2}}]$ at height

$$h(y) = R + 2 \log\left(\frac{\pi}{2}\right) - y.$$

Therefore, if $y \leq 2 \log(\pi/2)$ this intersection occurs above the height R of the box \mathcal{R} while in the other case the full region of the box above $h(y)$ is connected to p .

We will first consider the case where $y \leq 2 \log(\pi/2)$. Here we have

$$\begin{aligned} \mu(\mathcal{B}_{\text{box}}(p)) &= \int_0^R \int_{-e^{\frac{y+y'}{2}}}^{e^{\frac{y+y'}{2}}} f_{\alpha, \nu}(x', y') dx' dy' \\ &= \frac{2\alpha\nu}{\pi} e^{\frac{y}{2}} \int_0^R e^{-(\alpha - \frac{1}{2})y'} dy' \\ &= \mu(y) \left(1 - e^{-(\alpha - \frac{1}{2})R}\right), \end{aligned}$$

where the error term is $o(1)$, uniformly in y .

Now let $y > 2 \log(\pi/2)$ and recall that $\mu(y) = \xi e^{\frac{y}{2}}$ where $\xi = \frac{4\alpha\nu}{(2\alpha-1)\pi}$. Then, after some simple algebra, we have that

$$\begin{aligned} \mu_{\text{box}}(y) &= \int_0^{h(y)} \int_{-\frac{\pi}{2}e^{\frac{R}{2}}}^{\frac{\pi}{2}e^{\frac{R}{2}}} \mathbb{1}_{\{|x'| \leq e^{\frac{y+y'}{2}}\}} f_{\alpha, \nu}(x', y') dx' dy' + \int_{h(y)}^R \int_{-\frac{\pi}{2}e^{\frac{R}{2}}}^{\frac{\pi}{2}e^{\frac{R}{2}}} f_{\alpha, \nu}(x', y') dx' dy' \\ &= \frac{2\alpha\nu}{\pi} e^{\frac{y}{2}} \int_0^{h(y)} e^{-(\alpha - \frac{1}{2})y'} dy' + \alpha\nu e^{\frac{R}{2}} \int_{h(y)}^R e^{-\alpha y'} dy' \\ &= \xi e^{\frac{y}{2}} \left(1 - \left(\frac{\pi}{2}\right)^{-(2\alpha-1)} e^{-(\alpha - \frac{1}{2})(R-y)}\right) + \nu e^{\frac{R}{2}} \left(\left(\frac{\pi}{2}\right)^{-2\alpha} e^{-\alpha(R-y)} - e^{-\alpha R}\right) \\ &= \mu(y) (1 - \phi_n(y)). \end{aligned}$$

where

$$\phi_n(y) := \left(\frac{\pi}{2}\right)^{-(2\alpha-1)} e^{-(\alpha - \frac{1}{2})(R-y)} + \frac{\nu}{\xi} e^{-(\alpha - \frac{1}{2})R - \frac{y}{2}} - \frac{\nu}{\xi} \left(\frac{\pi}{2}\right)^{-2\alpha} e^{-(\alpha - \frac{1}{2})(R-y)}.$$

Since $R - y \geq \varepsilon R$ we have that $|\phi_n(y)|$ is uniformly bounded by $O\left(e^{-(\alpha - \frac{1}{2})\varepsilon R}\right)$, which is $o(1)$ for $\alpha > \frac{1}{2}$. \square

We can now use a concentration of heights argument to show that the integration of the Poisson probabilities $\mathbb{P}(\text{Po}(\mu_{\text{Po}}(y)) = k_n)$ over $0 \leq y \leq (1 - \varepsilon)R$ is asymptotically equivalent to p_{k_n} . And the same holds if we instead consider $\mu_{\text{box}}(y)$. The proof contains some technical elements that are contained in the Appendix to not hinder the flow of the argument.

Lemma 3.3. *Let $0 < \varepsilon < 1$. Then for all $0 \leq k_n \leq n - 1$, as $n \rightarrow \infty$,*

$$\int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\mu(\mathcal{B}(y)) = k_n) \alpha e^{-\alpha y} dy = (1 + o(1))p_{k_n}, \quad (23)$$

where

$$p_{k_n} = \int_0^\infty \mathbb{P}(\text{Po}(\xi e^{\frac{y}{2}}) = k_n) \alpha e^{-\alpha y} dy. \quad (24)$$

Moreover, the same holds if we replace $\mu(\mathcal{B}(y))$ with $\mu(\mathcal{B}_{\text{box}}(y))$ in (23).

Proof. We will show that

$$\int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\mu_{\text{Po}}(y)) = k_n) \alpha e^{-\alpha y} dy = (1 + o(1)) \int_0^\infty \mathbb{P}(\text{Po}(\xi e^{\frac{z}{2}}) = k_n) \alpha e^{-\alpha z} dz.$$

This implies the result because the last integral equals $(1 + o(1))2\alpha\xi^{2\alpha}k_n^{-(2\alpha+1)} = (1 + o(1))p_{k_n}$.

Define the function $z(y) = 2 \ln \frac{\mu_{\text{Po}}(y)}{\xi}$ (note that $z(y)$ is well-defined as $\mu_{\text{Po},n}(y) \geq 0$ and that $z(y)$ is bijective because $\mu_{\text{Po}}(y)$ is strictly monotone increasing and continuous, see Lemma 3.3. in [21]). By rearranging, we have that

$$\mu_{\text{Po}}(y) = \xi e^{\frac{z(y)}{2}}.$$

From Lemma 3.1, it follows that uniformly for all $0 \leq y \leq (1 - \varepsilon)R$, $\xi e^{\frac{y}{2}} = (1 + o(1))\mu_{\text{Po}}(y) = (1 + o(1))\xi e^{\frac{z(y)}{2}}$, and hence that

$$e^{-\alpha y} = (1 + o(1))e^{-\alpha z(y)}.$$

Next we need a similar result regarding the derivative of $\mu_{\text{Po}}(y)$, i.e.

$$\mu'_{\text{Po}}(y) = (1 + o(1))\frac{1}{2}\mu(y) = (1 + o(1))\frac{1}{2}\mu_{\text{Po}}(y) = (1 + o(1))\mu'(y),$$

uniformly for $0 \leq y \leq (1 - \varepsilon)R$. This result is given by Lemma F.1 in the Appendix. The lemma is placed there since the proof is a straightforward though cumbersome use of function analysis and we do not want to break the flow of the argument.

We now have that

$$z'(y) = \frac{2\mu'_{\text{Po}}(y)}{\mu_{\text{Po}}(y)} = 1 + o(1).$$

which implies that

$$\int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\mu_{\text{Po}}(y)) = k_n) \alpha e^{-\alpha y} dy = (1 + o(1)) \int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\xi e^{\frac{z(y)}{2}}) = k_n) \alpha e^{-\alpha z(y)} z'(y) dy.$$

We now apply integration by substitution to the integral, i.e. use the new variable $z = z(y)$, to obtain

$$\int_{z(0)}^{z((1-\varepsilon)R)} \mathbb{P}(\text{Po}(\xi e^{\frac{z}{2}}) = k_n) \alpha e^{-\alpha z} dz = \int_{z(0)}^{z((1-\varepsilon)R)} \mathbb{P}(\text{Po}(\mu(z)) = k_n) \alpha e^{-\alpha z} dz.$$

Note that since the function $y \mapsto 2 \ln \frac{y}{\xi}$ is monotone increasing it follows that for large enough n , $\mathcal{K}_C(k_n) \subset [z(0), z((1 - \varepsilon)R)]$. Therefore, by a concentration of heights argument (Proposition 2.4) it follows that

$$\int_{z(0)}^{z((1-\varepsilon)R)} \mathbb{P}(\text{Po}(\mu(z)) = k_n) \alpha e^{-\alpha z} dz = (1 + o(1)) \int_0^\infty \mathbb{P}(\text{Po}(\mu(z)) = k_n) \alpha e^{-\alpha z} dz.$$

and hence

$$\int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\mu_{\text{Po}}(y)) = k_n) \alpha e^{-\alpha y} dy = (1 + o(1)) \int_0^\infty \mathbb{P}(\text{Po}(\mu(z)) = k_n) \alpha e^{-\alpha z} dz,$$

which finishes the proof for $\mu_{\text{Po}}(y)$.

The proof for $\mu_{\text{box}}(y)$ follows similar arguments. First, we define $z(y) = 2 \log(\mu_{\text{box}}(y)/\xi)$ and use Lemma 3.2 instead of Lemma 3.1 to establish that $e^{-\alpha y} = (1 + o(1))e^{-\alpha z(y)}$. For the derivative $z'(y)$ we recall from the proof of Lemma 3.2 that $\mu_{\text{box}}(y) = (1 + \phi_n(y))\mu(y)$ with $\phi_n(y)$ given by (24). The derivative of $\phi_n(y)$ can be uniformly bounded by $o(1)$ for $0 \leq y \leq (1 - \varepsilon)R$. Hence we get $\mu'_{\text{box}}(y) = (1 + o(1))\mu'(y)$ and thus $z'(y) = 1 + o(1)$ uniformly for $0 \leq y \leq (1 - \varepsilon)R$. We can now apply the same change of variables and a concentration of heights argument to arrive at the required statement. \square

The main result of this section now follows almost immediately.

Lemma 3.4 (First moment of number of degree k vertices). *Let $N_{Po}(k)$ and $N_{box}(k)$ denote the number of vertices with degree k vertices in the poissonized KPKVB model G_{Po} and the finite box model G_{box} , respectively. Consider a sequence of integers $k_n \rightarrow \infty$ with $0 \leq k_n \leq n-1$. If $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$, then as $n \rightarrow \infty$,*

$$\mathbb{E}[N_{Po}(k_n)] = (1 + o(1))np_{k_n} \quad \text{and} \quad \mathbb{E}[N_{box}(k_n)] = (1 + o(1))np_{k_n}.$$

Moreover, if $k_n \gg n^{\frac{1}{2\alpha+1}}$ then

$$\mathbb{E}[N_{Po}(k_n)] = o(1) \quad \text{and} \quad \mathbb{E}[N_{box}(k_n)] = o(1)$$

Proof. We shall consider G_{Po} . The proof of the statements for G_{box} follows using the same arguments and we omit it here.

Let $D_{Po}(p)$ denote the degree of a node $p \in G_{Po}$. Then since $N_{Po}(k_n) = \sum_{v \in V(G_{Po})} \mathbb{1}_{\{D_{Po}(v)=k_n\}}$ and $\mathbb{P}(D_{Po}(p) = k)$ is invariant under translations in the x -axis, we can apply the Campbell-Mecke formula to obtain

$$\begin{aligned} \mathbb{E}[N_{Po}(k_n)] &= \int_0^R \mathbb{P}(\text{Po}(\mu_{Po}(y)) = k_n) \frac{\pi n}{\nu} \frac{\alpha \nu}{\pi} e^{-\alpha y} dy \\ &= n \int_0^R \mathbb{P}(\text{Po}(\mu_{Po}(y)) = k_n) \alpha e^{-\alpha y} dy \\ &= n \left(\int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\mu_{Po}(y)) = k_n) \alpha e^{-\alpha y} dy + \int_{(1-\varepsilon)R}^R \mathbb{P}(\text{Po}(\mu_{Po}(y)) = k_n) \alpha e^{-\alpha y} dy \right), \end{aligned}$$

where $0 < \varepsilon < 1$ is a constant to be chosen later. Note that

$$\int_{(1-\varepsilon)R}^R \mathbb{P}(\text{Po}(\mu_{Po}(y)) = k_n) \alpha e^{-\alpha y} dy \leq \int_{(1-\varepsilon)R}^R \alpha e^{-\alpha y} dy = \Theta(e^{-\alpha(1-\varepsilon)R}) = \Theta(n^{-2\alpha(1-\varepsilon)}).$$

Now first consider the case where $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$. Then, for $\alpha > \frac{1}{2}$ and $0 < \varepsilon < \frac{1}{2\alpha} - 1$, we have $2\alpha(1-\varepsilon) > 1$. Therefore, $\frac{2\alpha(1-\varepsilon)}{2\alpha+1} > \frac{1}{2\alpha+1}$ and thus $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right) = o\left(n^{\frac{2\alpha(1-\varepsilon)}{2\alpha+1}}\right)$. This implies that $k_n^{-(2\alpha+1)} \gg n^{-2\alpha(1-\varepsilon)}$ or, stated differently, $\Theta(n^{-2\alpha(1-\varepsilon)}) = o\left(k_n^{-(2\alpha+1)}\right)$. The first statement of the lemma now follows from Lemma 3.3.

When $k_n \gg n^{\frac{1}{2\alpha+1}}$, Lemma 3.3 implies that

$$n \int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\mu_{Po}(y)) = k_n) \alpha e^{-\alpha y} dy = (1 + o(1))np_{k_n} = O\left(nk_n^{-(2\alpha+1)}\right) = o(1).$$

On the other hand,

$$n \int_{(1-\varepsilon)R}^R \mathbb{P}(\text{Po}(\mu_{Po}(y)) = k_n) \alpha e^{-\alpha y} dy = O\left(n^{1-2\alpha(1-\varepsilon)}\right).$$

which is $o(1)$ since by our choice $2\alpha(1-\varepsilon) > 1$. Thus the second claim of the lemma follows. \square

3.4 Joint degrees in G_{box} and G_{Po}

To prove the factorization of higher moments of $N_{Po}(k_n)$ and $N_{box}(k_n)$ as in (18), we first have to understand the joint degree distribution in G_{Po} and G_{box} , respectively. This subsequently requires us to analyze the joint neighbourhoods of two points p, p' in these models. To explaining the proof strategy we will use the finite box model, since the formulas there are slightly easier. The results for the Poissonized KPKVB model G_{Po} follow the same idea.




Figure 4: Schematic representation of the neighbourhoods of p and p' in G_{box} when $|x - x'| > e^{\frac{y}{2}} + e^{\frac{y'}{2}}$ used for the proof of Lemma 3.5. Note that although here $p' \notin \mathcal{B}_{\text{box}}(p)$, this is not true in general. This situation was merely chosen to improve readability of the figure.

For $r \in \mathbb{N}$ and $p_1, \dots, p_r \in \mathcal{R}$, we write $G_{\text{box}} \cup \{p_1, \dots, p_r\}$ for the finite box model obtained by adding p_1, \dots, p_r to the vertex set of the graph and adding all corresponding edges according to the connection rule. Then we define, for any positive integer s and $V \subset \{p_1, \dots, p_s\}$,

$$\varphi_{\text{box}}(V, k; p_1, \dots, p_s) = \mathbb{P}(\text{every } p \in V \text{ has degree } k \text{ in } G_{\text{box}} \cup \{p_1, \dots, p_s\}). \quad (25)$$

In particular, for two points $p, p' \in \mathcal{R}$ and $V = \{p, p'\}$, $\varphi_{\text{box}}(V, k; p, p')$ is the joint degree distribution of p, p' in G_{box} . We will use similar notation for the Poissonized KPKVB model. That is, $G_{\text{Po}} \cup \{p_1, \dots, p_r\}$ denotes the Poissonized KPKVB model obtained by adding p_1, \dots, p_r to the vertex set of the graph and adding all corresponding edges and $\varphi_{\text{Po}}(V, k; p_1, \dots, p_s)$ the corresponding joint degree function.

If we define,

$$\begin{aligned} X_1(p, p') &:= \text{Po}(\mu(\mathcal{B}_{\text{box}}(p) \setminus \mathcal{B}_{\text{box}}(p'))), \\ X_2(p, p') &:= \text{Po}(\mu(\mathcal{B}_{\text{box}}(p') \setminus \mathcal{B}_{\text{box}}(p))), \\ Y(p, p') &:= \text{Po}(\mu(\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p'))) \end{aligned}$$

then each of these are independent Poisson random variables, while

$$\varphi_{\text{box}}(\{p, p'\}, k; p, p') = \mathbb{P}(X_1(p, p') + Y(p, p') = k, X_2(p, p') + Y(p, p') = k_n).$$

Recall the definition of $y_{k_n, C}^\pm$ from equation (15). We will show (see Lemma 3.7) that for any two points p, p' whose y -coordinate is in $\mathcal{K}_C(k_n)$ and whose x -coordinates are sufficiently separated, it holds that $\mu(\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')) = O(k_n^{1-\varepsilon})$. Since the mean of X_1 and X_2 for such two points is k_n , the contribution of the Poisson random variable $Y(p, p')$ to their degrees becomes negligible as $k_n \rightarrow \infty$ and hence the joint degree distribution will factorizes on this set. The main idea is that if p and p' are sufficiently separated in the x -direction, then the overlap of their neighbourhoods $\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')$ is of smaller order than $\mu(\mathcal{B}_{\text{box}}(p)) + \mu(\mathcal{B}_{\text{box}}(p'))$. We now proceed with analyzing these joint neighbourhoods.

Let $p, p' \in \mathcal{R}$ and denote by $\mathcal{N}_{\text{box}}(p, p')$ the number of common neighbours of p and p' in $G_{\text{box}} \cup \{p, p'\}$. We shall establish an upper bound on the expected number of joint neighbours when p and p' are sufficiently separated. Observe that $\mathbb{E}[\mathcal{N}_{\text{box}}(p, p')] = \mu(\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p'))$.

We start by analyzing the shape of the joint neighbourhood. Due to symmetry and the fact that we have identified the left and right boundaries of the box \mathcal{R} , we can, without loss of generality, assume that $p = (0, y)$ and $p' = (x', y')$ with $x' > 0$. To understand the computation it is helpful to have a picture. Figure 4 shows such an example. There are several different quantities that are important. The first are the heights where the left and right boundaries of the ball $\mathcal{B}_{\text{box}}(p)$ hit the boundaries of the box \mathcal{R} . Since $x = 0$ these heights are the same and we denote their common

value by $h(y)$. We also need to know the coordinates $\hat{y}_{\text{right}}(p, p')$ and $\hat{x}_{\text{right}}(p, p')$ of the intersection of the right boundary of the neighbourhood of p with the left boundary of the neighbourhood of p' and those for the intersection of the left boundary of the neighbourhood of p with the right boundary of the neighbourhood of p' , which we denote by $\hat{y}_{\text{left}}(p, p')$ and $\hat{x}_{\text{left}}(p, p')$. Finally we will denote by $d(p, p')$ the distance between the lower right boundary of $\mathcal{B}_{\text{box}}(p)$ and the lower left of $\mathcal{B}_{\text{box}}(p')$, which is positive only when the bottom parts of both neighbourhoods do not intersect, as is the case in Figure 4. The condition $d(p, p') > 0$ is exactly the right notion for p and p' being sufficiently separated.

We start by deriving expressions for these important coordinates. For this we introduce some notation. For any $p = (x, y) \in \mathcal{R}$ we will define the left and right boundary functions as, respectively,

$$b_p^-(z) = \begin{cases} 2 \log(x - z) - y & \text{if } -\frac{\pi}{2}e^{R/2} \leq z \leq x - e^{y/2} \\ 2 \log(\pi e^{R/2} + x - z) - y & \text{if } x - e^{(y+R)/2} + \pi e^{R/2} \leq z \leq \frac{\pi}{2}e^{R/2} \\ 0 & \text{otherwise} \end{cases} \quad (26)$$

$$b_p^+(z) = \begin{cases} 2 \log(z - x) - y & \text{if } x + e^{y/2} \leq z \leq \frac{\pi}{2}e^{R/2} \\ 2 \log(\pi e^{R/2} + z - x) - y & \text{if } -\frac{\pi}{2}e^{R/2} \leq z \leq x + e^{(y+R)/2} - \pi e^{R/2} \\ 0 & \text{otherwise} \end{cases} \quad (27)$$

Note that these functions describe the boundaries of the ball $\mathcal{B}_{\text{box}}(p)$. In particular, $p' = (x', y') \in \mathcal{B}_{\text{box}}(p)$ if and only if $y' \geq \min\{b_p^-(x'), b_p^+(x')\}$.

We shall derive the expressions for the point $(\hat{x}_{\text{left}}(p, p'), \hat{y}_{\text{left}}(p, p'))$. The x -coordinate $\hat{x}_{\text{left}}(p, p')$ is the solution to the equation $b_p^+(z) = b_{p'}^-(z)$ for $-\frac{\pi}{2}e^{R/2} \leq z \leq +e^{y'/2}$. This equation becomes

$$2 \log(\pi e^{R/2} + z - x') - y' = 2 \log(x' - z) - y',$$

whose solution is $\frac{x' - \pi e^{R/2}}{1 + e^{(y' - y)/2}}$. Plugging this into either the left or right hand side of the above equation yields the y -coordinate $\hat{y}_{\text{left}}(p, p') = 2 \log\left(\frac{\pi e^{R/2} - x'}{e^{y/2} + e^{y'/2}}\right)$. The expressions for $\hat{x}_{\text{right}}(p, p')$ and $\hat{y}_{\text{right}}(p, p')$ are derived in a similar way. The expression for $d(p, p')$ follows as the difference $b_{p'}^-(x' - e^{y'/2}) - b_p^+(e^{y/2})$.

The full expressions of all coordinates are given below for further reference.

$$h(y) = R - y + 2 \log\left(\frac{\pi}{2}\right) \quad (28)$$

$$\hat{y}_{\text{right}}(p, p') = 2 \log\left(\frac{x'}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}}\right) \quad (29)$$

$$\hat{x}_{\text{right}}(p, p') = \frac{x'}{1 + e^{\frac{y' - y}{2}}}, \quad (30)$$

$$\hat{y}_{\text{left}}(p, p') = 2 \log\left(\frac{\pi e^{R/2} - x'}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}}\right), \quad (31)$$

$$\hat{x}_{\text{left}}(p, p') = \frac{x' - \pi e^{R/2}}{1 + e^{\frac{y' - y}{2}}}, \quad (32)$$

$$d(p, p') = |x - x'|_n - \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}}\right). \quad (33)$$

The following result shows that if $d(p, p') > 0$, then the expected number of common neighbours is $o(\mu(\mathcal{B}_{\text{box}}(p)) + \mu(\mathcal{B}_{\text{box}}(p')))$.

Lemma 3.5. *Let $p, p' \in \mathcal{R}$. Then, whenever $|x - x'|_n > \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}}\right)$,*

$$\mathbb{E}[\mathcal{N}_{\text{box}}(p, p')] \leq \mu(\mathcal{B}_{\text{box}}(p)) \left(\left(\frac{|x - x'|}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}} \right)^{-(2\alpha-1)} + \frac{\nu}{\xi} e^{-(\alpha-\frac{1}{2})(R-y)} \right).$$

Proof. Again, without loss of generality we assume that $p = p_0 = (0, y)$ and $p' = (x', y')$ with $0 \leq x' \leq \frac{\pi}{2}e^{R/2}$. Note that since $0 < x' \leq \frac{\pi}{2}e^{R/2}$, $\hat{y}_{\text{right}}(p, p') \leq \hat{y}_{\text{left}}(p, p')$. We write \hat{y} for $\hat{y}_{\text{right}}(p, p')$ and observe that below \hat{y} the balls $\mathcal{B}_{\text{box}}(p)$ and $\mathcal{B}_{\text{box}}(p')$ are disjoint. Therefore, if we define $A := \{p_1 = (x_1, y_1) \in \mathcal{R} \cap \mathcal{B}_{\text{box}}(p) : y_1 \geq \hat{y}\}$. Then

$$\mathbb{E}[\mathcal{N}_{\text{box}}(p, p')] \leq \mu(A).$$

We proceed with computing the right hand side

$$\begin{aligned} \mu(A) &= \int_{\hat{y}}^{h(y)} \int_{-e^{\frac{y+y_1}{2}}}^{e^{\frac{y'+y_1}{2}}} f(x_1, y_1) dx_1 dy_1 + \int_{h(y)}^R \int_{-\frac{\pi}{2}e^{R/2}}^{\frac{\pi}{2}e^{R/2}} f(x_1, y_1) dx_1 dy_1 \\ &= \frac{2\alpha\nu}{\pi} e^{\frac{y}{2}} \int_{\hat{y}}^{h(y)} e^{-(\alpha-\frac{1}{2})y_1} dy_1 + \alpha\nu e^{R/2} \int_{h(y)}^R e^{-\alpha y_1} dy_1 \\ &\leq \xi \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}} \right) e^{-(\alpha-\frac{1}{2})\hat{y}} + \nu e^{R/2} e^{-\alpha h(y)} \\ &= \mu(\mathcal{B}_{\text{box}}(p)) \left(e^{-(\alpha-\frac{1}{2})\hat{y}} + \frac{\nu}{\xi} e^{-(\alpha-\frac{1}{2})(R-y)} \right). \end{aligned}$$

The result follows by plugging in

$$\hat{y} := \hat{y}_{\text{right}}(p, p') = 2 \log \left(\frac{x'}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}} \right),$$

and noting that x' is the same as $|x - x'|$, by our generalization step. \square

We can also prove a similar result for the Poissonized KPKVB model G_{Po} , denoting by $\mathcal{N}_{\text{Po}}(p, p')$ the number of joint neighbours in $G_{\text{Po}} \cup \{p, p'\}$.

Lemma 3.6. *Let $0 < \varepsilon < 1$, $p, p' \in \mathcal{R}$ with $y, y' \leq (1 - \varepsilon)R$, denote by $\mathcal{N}_{\text{Po}}(p, p')$ the number of joint neighbours of p, p' in G_{Po} and let K be the constant from Lemma 2.2. Then, whenever $|x - x'|_n > \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}} \right) \left(1 + \frac{\pi^2 K}{4} \right)$,*

$$\mathbb{E}[\mathcal{N}_{\text{Po}}(p, p')] \leq \mu(\mathcal{B}(p)) \left(e^{(2\alpha-1)\lambda} \left(\frac{|x - x'|}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}} \right)^{-(2\alpha-1)} + \frac{\nu}{\xi} e^{-(\alpha-\frac{1}{2})(R-y)} \right),$$

where

$$\lambda = \log \left(1 + \frac{\pi^2 K}{4} \right).$$

Proof. We will proceed in a similar fashion as for Lemma 3.5. That is, we will bound the expected number of common neighbours by the number of neighbors of p whose y -coordinate is above the intersection of the right boundary of $\mathcal{B}(p)$ and the left boundary of $\mathcal{B}(p')$. Denote by \hat{y} the height of this intersection point. Then

$$\mathbb{E}[\mathcal{N}_{\text{Po}}(p, p')] \leq \frac{2\alpha\nu}{\pi} \int_{\hat{y}}^{R-y} \Phi(y, y_1) e^{-\alpha y_1} dy_1 + \alpha\nu e^{R/2} \int_{R-y}^R e^{-\alpha y_1} dy_1.$$

The second integral is bounded by $\frac{\nu}{\xi} \mu(\mathcal{B}_{\text{box}}(y)) e^{-(\alpha-\frac{1}{2})(R-y)}$. We bound the first integral using Lemma 2.2 as

$$\begin{aligned} \frac{2\alpha\nu}{\pi} \int_{\hat{y}}^{R-y} \Phi(y, y_1) e^{-\alpha y_1} dy_1 &\leq \frac{2\alpha\nu}{\pi} \int_{\hat{y}}^{R-y} \left(e^{\frac{y+y_1}{2}} + K e^{\frac{3}{2}(y+y_1)-R} \right) e^{-\alpha y_1} dy_1 \\ &\leq \frac{2\alpha\nu}{\pi} (1 + K) e^{\frac{y}{2}} \int_{\hat{y}}^{R-y} e^{-(\alpha-\frac{1}{2})y_1} dy_1 \end{aligned}$$

$$\leq (1 + K)\mu(\mathcal{B}_{\text{box}}(p))e^{-(\alpha - \frac{1}{2})\hat{y}},$$

where we used that $\frac{3y_1}{2} \leq R - y + \frac{y_1}{2}$ for all $y_1 \leq R - y$ for the second line.

It remains to compute \hat{y} , for which we will establish the following bound

$$\hat{y} \geq 2 \log \left(\frac{x'}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}} \right) - 2\lambda. \quad (34)$$

To show (34) we note that for any point $y_1 \geq \hat{y}$, the corresponding x -coordinate of the left boundary of $\mathcal{B}(p')$ must be to the left of that of the ball $\mathcal{B}(p)$, i.e. $x' - \Phi(y', y_1) \leq \Phi(y, y_1)$. Therefore it is enough to show that for all

$$y_1 \leq 2 \log \left(\frac{x'}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}} \right) - 2\lambda$$

it holds that $\Phi(y, y_1) \leq x' - \Phi(y', y_1)$, with λ as defined in the statement of the lemma. Note that by assumption on $x' := |x - x'|_n$ (since we can take $x = 0$) the above upper bound is non-negative. Using Lemma 2.2 it suffices to prove that for all such y_1 ,

$$e^{\frac{y+y_1}{2}} + Ke^{\frac{3}{2}(y+y_1)-R} \leq x' - e^{\frac{y'+y_1}{2}} - Ke^{\frac{3}{2}(y'+y_1)-R},$$

which is equivalent to

$$\left(e^{\frac{y}{2}} + e^{\frac{y'}{2}} \right) e^{\frac{y_1}{2}} + Ke^{-R} \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}} \right)^3 e^{\frac{3y_1}{2}} \leq x'.$$

Plugging the upper bound for y_1 into the left hand side and using that $(e^{y/2} + e^{y'/2})^3 \geq e^{3y/2} + e^{3y'/2}$, we obtain

$$\begin{aligned} \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}} \right) e^{\frac{y_1}{2}} + Ke^{-R} \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}} \right)^3 e^{\frac{3y_1}{2}} &\leq x'e^{-\lambda} + Ke^{-R}(x')^3 e^{-3\lambda} \leq x' \left(e^{-\lambda} + \frac{\pi^2 K}{4} e^{-3\lambda} \right) \\ &\leq x'e^{-\lambda} \left(1 + \frac{\pi^2 K}{4} \right) = x', \end{aligned}$$

where we also used that $x' \leq \frac{\pi}{2}e^{-R/2}$. □

Let us now define the *stripe*

$$\mathcal{S}_{k_n, C} = \mathcal{R} \cap (\mathbb{R}_+ \times \mathcal{K}_C(k_n)). \quad (35)$$

and in addition define, for any $0 < \varepsilon < 1$, the following set

$$\mathcal{E}_\varepsilon(k_n) = \{(p, p') \in \mathcal{S}_{k_n, C} : |x - x'|_n > k_n^{1+\varepsilon}\}, \quad (36)$$

where $|x|_n = \min\{|x|, \pi e^{R/2} - |x|\}$ denotes the norm on the finite box \mathcal{R} where the left and right boundaries are identified. Then for any two points $p, p' \in \mathcal{E}_\varepsilon(k_n)$ the expect number of joint neighbours is $o(k_n)$.

Lemma 3.7. *Fix $0 < \varepsilon < 1$ and let $\varepsilon' = \min\{\varepsilon(2\alpha - 1), \varepsilon\}$. Then for all $(p, p') \in \mathcal{E}_\varepsilon(k_n)$, as $n \rightarrow \infty$,*

$$\mu(\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')) = O(k_n^{1-\varepsilon'}).$$

Proof. Since for all $(p, p') \in \mathcal{E}_\varepsilon(k_n)$ we have $\mu(\mathcal{B}_{\text{box}}(p)), \mu(\mathcal{B}_{\text{box}}(p')) = \Theta(k_n)$, Lemma 3.5 implies that

$$\mu(\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')) \leq O(k_n) \phi_n(p, p'),$$

where

$$\phi_n(p, p') = 2 \left(\frac{|x - x'|_n}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}} \right)^{-(2\alpha-1)} + \frac{3\nu^{2\alpha+1} e^{-(\alpha-\frac{1}{2})R} e^{\alpha y}}{2\pi^{2\alpha} \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}} \right)} + \frac{\nu e^{-(\alpha-\frac{1}{2})R}}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}}.$$

We thus need to show that $\phi_n(p, p') = O(k_n^{-\varepsilon'})$. For $(p, p') \in \mathcal{E}_\varepsilon(k_n)$, it holds that $e^{y/2}, e^{y'/2} = \Theta(k_n)$ and $|x - x'|_n > k_n^{1+\varepsilon}$ and hence

$$2 \left(\frac{|x - x'|_n}{e^{\frac{y}{2}} + e^{\frac{y'}{2}}} \right)^{-(2\alpha-1)} = O(k_n^{-\varepsilon(2\alpha-1)}).$$

For the second term in $\phi_n(p, p')$ we use that $e^{\alpha y^*} = \Theta(k_n^{2\alpha})$ and $e^R = \Theta(n^2)$ to obtain

$$\frac{3\nu^{2\alpha+1} e^{-(\alpha-\frac{1}{2})R} e^{\alpha y}}{2\pi^{2\alpha} \left(e^{\frac{y}{2}} + e^{\frac{y'}{2}} \right)} = O(1) n^{-(2\alpha-1)} k_n^{2\alpha-1} = O(n^{-(\alpha-\frac{1}{2})}).$$

Finally, the third term satisfies $O(n^{-(2\alpha-1)} k_n^{-1})$, and we conclude that

$$\phi_n(p, p') = O(k_n^{-\varepsilon(2\alpha-1)} + n^{-(\alpha-\frac{1}{2})} + n^{-(2\alpha-1)} k_n^{-1}) = O(k_n^{-\varepsilon'}),$$

where we used that $\varepsilon' = \min\{\varepsilon(2\alpha-1), \varepsilon\}$. \square

It is clear that using Lemma 3.6 instead of Lemma 3.5, the above proof applies to the Poissonized KPKVB model, yielding the following result.

Lemma 3.8. *Fix $0 < \varepsilon < 1$ and let $\varepsilon' = \min\{\varepsilon(2\alpha-1), \varepsilon\}$. Then for all $(p, p') \in \mathcal{E}_\varepsilon(k_n)$, as $n \rightarrow \infty$,*

$$\mu(\mathcal{B}(p) \cap \mathcal{B}(p')) = O(k_n^{1-\varepsilon'}).$$

Recall the definition of the stripe $\mathcal{S}_{k_n, C} = \mathcal{R} \cap (\mathbb{R}_+ \times [y_{k_n, C}^-, y_{k_n, C}^+])$ from (35). Consider a fixed number of points p_1, \dots, p_r . Then, if their x -coordinates are sufficient far apart and their y coordinates lie within the stripe $\mathcal{S}_{k_n, C}$, their degrees are asymptotically independent.

Lemma 3.9 (Asymptotic factorization of degree probabilities). *Let $C, C' > 0$ and r, s be positive integers with $r+1 \leq s$. Fix $0 < \varepsilon < 1$. Then, it holds uniformly for all $(v_1, \dots, v_s) \in (\mathcal{S}_{k_n, C})^s = \mathcal{S}_{k_n, C} \times \dots \times \mathcal{S}_{k_n, C}$, satisfying $|x_{v_i} - x_{v_{r+1}}| \geq k_n^{1+\varepsilon}$ for all $1 \leq i \leq r$, that*

$$\varphi(\{v_1, \dots, v_{r+1}\}; v_1, \dots, v_s) = (1+o(1))\varphi(\{v_1, \dots, v_r\}; v_1, \dots, v_s)\varphi(\{v_{r+1}\}; v_1, \dots, v_s) + O(k_n^{-C'}),$$

where φ is either φ_{Po} or φ_{box} . (Here uniformity means that the $o(1)$ and $O(k_n)$ terms do not depend on v_1, \dots, v_s .)

Proof. Let $H = G_{\text{Po}} \cup \{p_1, \dots, p_s\}$ or $H = G_{\text{box}} \cup \{p_1, \dots, p_s\}$ and $1 \leq r \leq s$. For $1 \leq j \leq r$, let Y_j be the number of vertices of H which are adjacent to both p_j and p_{r+1} . Let X_j be the number of vertices of H which are adjacent to p_j , but not to p_{r+1} . Then, $X_j + Y_j = D_H(p_j)$ is the degree of p_j in H .

Now let X_{r+1} be the number of vertices of H which are adjacent to p_{r+1} , but to none of p_1, \dots, p_r . Let Y_{r+1} be the number of vertices of H which are adjacent to p_{r+1} , and at least one of p_1, \dots, p_r . Then, $X_{r+1} + Y_{r+1} = D_H(p_{r+1})$ is the degree of p_{r+1} in H . By definition, we therefore have

$$\begin{aligned} \varphi(\{p_1, \dots, p_{r+1}\}; p_1, \dots, p_s) &= \mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + Y_{r+1} = k_n), \\ \varphi(\{p_1, \dots, p_r\}; p_1, \dots, p_s) &= \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n), \\ \varphi(\{p_{r+1}\}; p_1, \dots, p_s) &= \mathbb{P}(X_{r+1} + Y_{r+1} = k_n), \end{aligned}$$

and the claim of the lemma is that

$$\begin{aligned} & \mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + Y_{r+1} = k_n) \\ &= (1 + o(1)) \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n) \mathbb{P}(X_{r+1} + Y_{r+1} = k_n) + O(k_n^{-C}). \end{aligned} \quad (37)$$

To prove (37) let $\varepsilon' = \min(\varepsilon, \varepsilon(2\alpha - 1)) \in (0, 1)$. Since for $1 \leq i \leq r$, it is given that $|x_{p_i} - x_{p_{r+1}}| \geq k_n^{1+\varepsilon}$, then in the case where $H = G_{\text{Po}} \cup \{p_1, \dots, p_s\}$ it follows from Lemma 3.6 that $\mathbb{E}[Y_j] = O(k_n^{1-\varepsilon'})$. When $H = G_{\text{box}} \cup \{p_1, \dots, p_s\}$ we get the same result using Lemma 3.5. The rest of the proof is independent of which of the two models we consider and only uses that $\mathbb{E}[Y_j] = O(k_n^{1-\varepsilon'})$.

As $Y_{r+1} \leq Y_1 + \dots + Y_r$, we also have $\mu := \mathbb{E}[Y_{r+1}] = O(k_n^{1-\varepsilon'})$. In particular, there is $c_0 > 0$ such that $c_0 \sqrt{k_n^{1-\varepsilon'} \ln k_n^{1-\varepsilon'}} \geq c_1 \sqrt{\mu \ln \mu}$ (where $c_1 = \sqrt{\frac{2C}{1-\varepsilon'}}$, which is well-defined because $1 - \varepsilon' > 0$). Now define

$$A_n = [\mu - c_0 \sqrt{k_n^{1-\varepsilon'} \ln k_n^{1-\varepsilon'}}, \mu + c_0 \sqrt{k_n^{1-\varepsilon'} \ln k_n^{1-\varepsilon'}}] \cap \mathbb{N}_0.$$

By equation (130), we have

$$\begin{aligned} \mathbb{P}(Y_{r+1} \notin A_n) &= \mathbb{P}\left(|Y_{r+1} - \mu| \geq c_0 \sqrt{k_n^{1-\varepsilon'} \ln k_n^{1-\varepsilon'}}\right) \\ &\leq \mathbb{P}\left(|Y_{r+1} - \mu| \geq c_1 \sqrt{\mu \ln \mu}\right) = O\left(k_n^{-\frac{(1-\varepsilon')c_1^2}{2}}\right). \end{aligned}$$

As by definition c_1 satisfies $\frac{(1-\varepsilon')c_1^2}{2} = C$, this implies that for the event $S_n = \{Y_{r+1} \in A_n\}$,

$$\mathbb{P}(S_n^c) = O(k_n^{-C}).$$

Beginning with the left-hand side of the claim of the lemma, the law of total probability applied to the events $\{Y_{r+1} = y_{r+1}\}$, for all $y_{r+1} \in A_n$, and S_n^c implies that

$$\begin{aligned} & \mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + Y_{r+1} = k_n) \\ &= \sum_{y_{r+1} \in A_n} \mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + y_{r+1} = k_n | Y_{r+1} = y_{r+1}) \mathbb{P}(Y_{r+1} = y_{r+1}) \\ &\quad + \mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + Y_{r+1} = k_n | S_n^c) \mathbb{P}(S_n^c) \\ &= \sum_{y_{r+1} \in A_n} \mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + y_{r+1} = k_n | Y_{r+1} = y_{r+1}) \mathbb{P}(Y_{r+1} = y_{r+1}) + O(k_n^{-C}). \end{aligned}$$

As X_{r+1} is independent of $X_1, \dots, X_r, Y_1, \dots, Y_r$ by the properties of a Poisson process (as X_{r+1} counts the number of points in a set which is disjoint of $X_1, \dots, X_r, Y_1, \dots, Y_r$), it follows that

$$\begin{aligned} & \mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + Y_{r+1} = k_n) \\ &= \sum_{y_{r+1} \in A_n} \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n | Y_{r+1} = y_{r+1}) \mathbb{P}(X_{r+1} + y_{r+1} = k_n) \mathbb{P}(Y_{r+1} = y_{r+1}) \\ &\quad + O(k_n^{-C}). \end{aligned}$$

We will now show that uniformly for all $y_{r+1}, s \in A_n$, it holds that,

$$\mathbb{P}(X_{r+1} = k_n - y_{r+1}) = (1 + o(1)) \mathbb{P}(X_{r+1} = k_n - s). \quad (38)$$

To see this, observe that for all $y_{r+1}, s \in A_n$, we have that $|y_{r+1} - s| \leq 2c \sqrt{k_n^{1-\varepsilon'} \ln k_n^{1-\varepsilon'}}$. Denote the expectation of X_{r+1} by λ , write $\delta_n = k_n - y_{r+1} - \lambda$ and note that

$$\frac{\mathbb{P}(X_{r+1} = k_n - y_{r+1})}{\mathbb{P}(X_{r+1} = k_n - s)} = \frac{\mathbb{P}(X_{r+1} = k_n - y_{r+1})}{\mathbb{P}(X_{r+1} = k_n - y_{r+1} + (y_{r+1} - s))}$$

$$= \frac{(k_n - y_{r+1} + (y_{r+1} - s))!}{(k_n - y_{r+1})!} \lambda^{s - y_{r+1}}$$

We will now use that $\frac{(a+b)!}{a!} = (1 + o(1))(a+b)^b$ for $b^2 = o(a)$, applied to $a = k_n - y_{r+1}$ and $b = y_{r+1} - s$. To see this auxiliary fact, note that by Stirling's approximation to the factorial, it follows that

$$\frac{(a+b)!}{a!} = (1 + o(1)) \frac{(a+b)^{a+b+\frac{1}{2}} e^{-a-b}}{a^{a+\frac{1}{2}} e^{-a}} = (1 + o(1)) \left(1 + \frac{b}{a}\right)^{a+\frac{1}{2}} (a+b)^b e^{-b}.$$

Since $1 + \frac{b}{a} \leq e^{\frac{b}{a}}$, it holds that $(1 + \frac{b}{a})^a \leq e^b$. Furthermore, as $\ln(1+x) \geq x - \frac{x^2}{2}$ (for $x \in (-1, 1)$), we have that $(1 + \frac{b}{a})^a = e^{a \ln(1+\frac{b}{a})} \geq e^{a(\frac{b}{a} - \frac{b^2}{2a})} = e^{b - \frac{b^2}{2a}} = (1 + o(1))e^b$ because $b^2 = o(a)$. Finally, $b^2 = o(a)$ also implies that $(1 + \frac{b}{a})^{\frac{1}{2}} = 1 + o(1)$. This finishes the proof of the auxiliary fact and we can continue with

$$\begin{aligned} \frac{\mathbb{P}(X_{r+1} = k_n - y_{r+1})}{\mathbb{P}(X_{r+1} = k_n - s)} &= (1 + o(1)) (k_n - y_{r+1} + (y_{r+1} - s))^{y_{r+1} - s} \lambda^{s - y_{r+1}} \\ &= (1 + o(1)) (\lambda + \delta_n + (y_{r+1} - s))^{y_{r+1} - s} \lambda^{s - y_{r+1}} \\ &= (1 + o(1)) \left(1 + \frac{\delta_n + (y_{r+1} - s)}{\lambda}\right)^{y_{r+1} - s} \\ &= (1 + o(1)) e^{\frac{\delta_n (y_{r+1} - s)}{\lambda}} e^{\frac{(y_{r+1} - s)^2}{\lambda}} = 1 + o(1), \end{aligned}$$

where the last line follows since $\delta_n, |y_{r+1} - s| \leq 2c_0 \sqrt{k_n^{1-\varepsilon'} \ln k_n^{1-\varepsilon'}}$ and $\lambda = \Theta(k_n)$ and therefore, with convergence uniform in y_{r+1}, s ,

$$\frac{\delta_n (y_{r+1} - s)}{\lambda}, \frac{(y_{r+1} - s)^2}{\lambda} \leq \frac{4c_0^2 k_n^{1-\varepsilon'} \ln k_n^{1-\varepsilon'}}{\lambda} \rightarrow 0.$$

As $\mathbb{P}(S_n^c) = O(k_n^{-C})$, we have

$$1 = \sum_{s \in A_n} \mathbb{P}(Y_{r+1} = s) + O(k_n^{-C}).$$

From (38), it then follows that

$$\begin{aligned} \mathbb{P}(X_{r+1} = k_n - y_{r+1}) &= \mathbb{P}(X_{r+1} = k_n - y_{r+1}) \sum_{s \in A_n} \mathbb{P}(Y_{r+1} = s) + O(k_n^{-C}) \\ &= (1 + o(1)) \sum_{s \in A_n} \mathbb{P}(Y_{r+1} = s) \mathbb{P}(X_{r+1} = k_n - s) + O(k_n^{-C}) \\ &= (1 + o(1)) \sum_{s \in A_n} \mathbb{P}(X_{r+1} + Y_{r+1} = k_n, Y_{r+1} = s) + O(k_n^{-C}) \\ &= (1 + o(1)) \mathbb{P}(X_{r+1} + Y_{r+1} = k_n, Y_{r+1} \in A_n) + O(k_n^{-C}) \\ &= (1 + o(1)) (\mathbb{P}(X_{r+1} + Y_{r+1} = k_n) - \mathbb{P}(X_{r+1} + Y_{r+1} = k_n, Y_{r+1} \notin A_n)) + O(k_n^{-C}) \\ &= (1 + o(1)) \mathbb{P}(X_{r+1} + Y_{r+1} = k_n) + O(k_n^{-C}). \end{aligned}$$

Note that $\mathbb{P}(X_{r+1} + Y_{r+1} = k_n)$ no longer depends on y_{r+1} and neither does the $O(k_n^{-C})$ error term. Therefore we have

$$\begin{aligned} \mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + Y_{r+1} = k_n) \\ \sum_{y_{r+1} \in A_n} \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n | Y_{r+1} = y_{r+1}) \mathbb{P}(X_{r+1} + y_{r+1} = k_n) \mathbb{P}(Y_{r+1} = y_{r+1}) + O(k_n^{-C}) \end{aligned}$$

$$\begin{aligned}
&= (1 + o(1)) \mathbb{P}(X_{r+1} + Y_{r+1} = k_n) \sum_{y_{r+1} \in A_n} \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n | Y_{r+1} = y_{r+1}) \mathbb{P}(Y_{r+1} = y_{r+1}) \\
&\quad + O(k_n^{-C}).
\end{aligned}$$

For the last summation we have

$$\begin{aligned}
&\sum_{y_{r+1} \in A_n} \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n | Y_{r+1} = y_{r+1}) \mathbb{P}(Y_{r+1} = y_{r+1}) \\
&= \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n, Y_{r+1} \in A_n) \\
&= (1 + o(1)) \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n) + O(k_n^{-C}).
\end{aligned}$$

Finally, plugging this into the previous step gives

$$\begin{aligned}
&\mathbb{P}(X_1 + Y_1 = \dots = X_{r+1} + Y_{r+1} = k_n) \\
&= (1 + o(1)) \mathbb{P}(X_{r+1} + Y_{r+1} = k_n) \sum_{y_{r+1} \in A_n} \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n | Y_{r+1} = y_{r+1}) \mathbb{P}(Y_{r+1} = y_{r+1}) \\
&\quad + O(k_n^{-C}) \\
&= (1 + o(1)) \mathbb{P}(X_{r+1} + Y_{r+1} = k_n) \mathbb{P}(X_1 + Y_1 = \dots = X_r + Y_r = k_n) + O(k_n^{-C}).
\end{aligned}$$

which establishes (37) and thus the claim of the lemma. \square

3.5 Factorial moments of degrees

Now that we have analyzed the joint neighbourhoods and degree distributions in both the Poissonized KPKVB and finite box model, we can show convergence of the factorial moments of the number of nodes of degree k in both models.

Lemma 3.10 (Factorial moments). *Recall that $N_{Po}(k)$ denotes the number of degree k vertices in G_{Po} . Then, for any positive integer r and $k_n \rightarrow \infty$, it holds that*

$$\mathbb{E} \left[\binom{N_{Po}(k_n)}{r} \right] = (1 + o(1)) \frac{(\mathbb{E}[N_{Po}(k_n)])^r}{r!}.$$

The proof of this result requires the following technical lemma which states that the integration of the joint degree distribution can be factorized.

Lemma 3.11 (Factorization of degrees). *Let φ be either φ_{box} or φ_{Po} . Then we have that*

$$\begin{aligned}
&\int_{\mathcal{R}} \dots \int_{\mathcal{R}} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_r) f(x_1, y_1) dx_1 dy_1 \dots f(x_r, y_r) dx_r dy_r \\
&= (1 + o(1)) \left(\int_{\mathcal{R}} \varphi(\{p_1\}, k_n; p_1) f(x, y) dx dy \right)^r.
\end{aligned}$$

Proof. Let $C > r(2\alpha + 1)$ and define the set $A = (\mathcal{R} \times \dots \times \mathcal{R}) \setminus (\mathcal{S}_{k_n, C} \times \dots \times \mathcal{S}_{k_n, C})$. We will first show that the contribution of the integration of φ over this range is negligible.

For $(p_1, \dots, p_r) \in (\mathcal{R} \times \dots \times \mathcal{R}) \setminus (\mathcal{S}_{k_n, C} \times \dots \times \mathcal{S}_{k_n, C})$, there is a j , $1 \leq j \leq r$, such that $y_j \notin [y_{k_n, C}^-, y_{k_n, C}^+]$, so that the Chernoff bound (see in 16) yields that $\mathbb{P}(D_{G_{Po}}(v_j) = k_n) = O(k_n^{-C})$. As, for $1 \leq j \leq r$, the event $\{D_{G_{Po}}(p_1) = \dots = D_{G_{Po}}(p_r) = k_n\}$ implies the event $\{D_{G_{Po}}(v_j) = k_n\}$, it follows that $\mathbb{P}(D_{G_{Po}}(p_1) = \dots = D_{G_{Po}}(p_r) = k_n) = O(k_n^{-C})$ and hence,

$$\int_A \mathbb{P}(D_{G_{Po}}(p_1) = \dots = D_{G_{Po}}(p_r) = k_n) f(x_1, y_1) dx_1 dy_1 \dots f(x_r, y_r) dx_r dy_r = O(n^r k_n^{-C}).$$

Next we note that by the concentration of heights (Proposition 2.4) it follows that

$$\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) f(x, y) dx dy = (1 + o(1)) \int_{\mathcal{R}} \varphi(\{p_1\}, k_n; p_1) f(x, y) dx dy$$

$$= (1 + o(1))2\alpha\xi^{2\alpha}nk_n^{-(2\alpha+1)} \quad (39)$$

Since for $C > r(2\alpha + 1)$ it holds that $n^r k_n^{-C} = o\left(\left(nk_n^{-(2\alpha+1)}\right)^r\right)$ it now suffices to show that

$$\begin{aligned} & \int_{\mathcal{S}_{k_n, C}} \cdots \int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_r) f(x_1, y_1) dx_1 dy_1 \cdots f(x_r, y_r) dx_r dy_r \\ &= (1 + o(1)) \left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1 \right)^r. \end{aligned}$$

We will prove this using induction. More precisely, for every fixed positive integer s , we will show by induction on r , $1 \leq r \leq s$, that for all $p_{r+1}, \dots, p_s \in \mathcal{S}_{k_n, C}$, it holds that

$$\begin{aligned} & \int_{\mathcal{S}_{k_n, C}} \cdots \int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_r, y_r) dx_r dy_r \\ &= (1 + o(1)) \left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1 \right)^r. \end{aligned}$$

Note that for $r = s$ this is the claim of the lemma. Throughout the proof H is either $G_{\text{Po}} \cup \{p_1, \dots, p_s\}$ or $G_{\text{box}} \cup \{p_1, \dots, p_s\}$, depending on whether φ is, respectively, φ_{Po} or φ_{box} .

For $r = 1$, we only need to show that uniformly for $p_1 \in \mathcal{S}_{k_n, C}$,

$$\varphi(\{p_1\}, k_n; p_1, \dots, p_s) = (1 + o(1))\varphi(\{p_1\}, k_n; p_1).$$

To see this, note that as $p_1 \in \mathcal{S}_{k_n, C}$, the expected degree of p_1 in G_{Po} is $\Theta(k_n)$. Assume that p_1 is adjacent to $s' < s$ many vertices among p_2, \dots, p_s . Then, as $s' < s$ is bounded and $k_n \rightarrow \infty$ as $n \rightarrow \infty$, we have that

$$\mathbb{P}(D_{G_{\text{Po}}}(p_1) = k_n - s') = (1 + o(1))\mathbb{P}(D_{G_{\text{Po}}}(p_1) = k_n).$$

So, we have the base case of the induction,

$$\begin{aligned} \varphi(\{p_1\}, k_n; p_1, \dots, p_s) &= \mathbb{P}(D_H(p_1) = k_n) = \mathbb{P}(D_{G_{\text{Po}}}(p_1) = k_n - s') \\ &= (1 + o(1))\mathbb{P}(D_{G_{\text{Po}}}(p_1) = k_n) = (1 + o(1))\varphi(\{p_1\}, k_n; p_1). \end{aligned}$$

Assuming the claim holds for integer $r < s$, we will show that it holds for $r + 1$.

Let $p_{r+2}, \dots, p_s \in \mathcal{S}_{k_n, C}$ (if $r + 2 > s$, this definition is void and the corresponding points will never be used in the proof). Fix $0 < \epsilon < 1$ small enough s.t. $\frac{1}{2} + \epsilon < \alpha$. Define the region that the $(r + 1)$ -th vertex p_{r+1} is far apart from all other vertices horizontally,

$$\mathcal{F}_\epsilon(k_n) = \{(p_1, \dots, p_{r+1}) \in (\mathcal{S}_{k_n, C})^{r+1} : \forall 1 \leq i \leq r : |x_{p_i} - x_{p_{r+1}}| \geq k_n^{1+\epsilon}\}.$$

We will split the integration into this region and its complement $\mathcal{F}_\epsilon(k_n)^c = (\mathcal{S}_{k_n, C})^{r+1} \setminus \mathcal{F}_\epsilon(k_n)$. Firstly, we derive an upper bound for the complement $\mathcal{F}_\epsilon(k_n)^c$. Note that

$$\varphi(\{p_1, \dots, p_{r+1}\}, k_n; p_1, \dots, p_s) \leq \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s),$$

and so,

$$\begin{aligned} & \int \int_{\mathcal{F}_\epsilon(k_n)^c} \varphi(\{p_1, \dots, p_{r+1}\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1} \\ & \leq \int \int_{\mathcal{F}_\epsilon(k_n)^c} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1}. \end{aligned}$$

For $(p_1, \dots, p_{r+1}) \in \mathcal{F}_\epsilon(k_n)^c$, we have that $(p_1, \dots, p_r) \in (\mathcal{S}_{k_n, C})^r$ and $p_{r+1} = (x_{r+1}, y_{r+1})$ satisfies $y_{k_n, c}^- \leq y_{r+1} \leq y_{k_n, c}^+$ and x_{r+1} falls into an interval I_n of length $2k_n^{1+\epsilon}$. As the integrand

$\varphi(\{p_1, \dots, p_r\}; p_1, \dots, p_s)$ is constant in x_{r+1} , we can upper bound the corresponding integration as follows,

$$\begin{aligned} & \int_{\{(x_{r+1}, y_{r+1}) \in \mathcal{S}_{k_n, C} : x_{r+1} \in I_n\}} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1} \\ &= \int_{y_{k_n, c}^-}^{y_{k_n, c}^+} \int_{I_n} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) \frac{\alpha \nu}{\pi} e^{-\alpha y_{r+1}} dx_{r+1} dy_{r+1} \\ &\leq 2k_n^{1+\epsilon} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) \int_{y_{k_n, c}^-}^{y_{k_n, c}^+} \frac{\alpha \nu}{\pi} e^{-\alpha y_{r+1}} dy_{r+1}. \end{aligned}$$

Furthermore, we have that

$$\int_{y_{k_n, c}^-}^{y_{k_n, c}^+} e^{-\alpha y_{r+1}} dy_{r+1} \leq \int_{y_{k_n, c}^-}^{\infty} e^{-\alpha y_{r+1}} dy_{r+1} = O\left(e^{-\alpha y_{k_n, c}^-}\right) = O\left(\left(\frac{k_n - c\sqrt{k_n \ln k_n}}{\xi}\right)^{-2\alpha}\right) = O(k_n^{-2\alpha}).$$

We have thus established that

$$\begin{aligned} & \int \int_{\mathcal{F}_\epsilon(k_n)^c} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1} \\ &\leq O(k_n^{1+\epsilon} k_n^{-2\alpha}) \int_{\mathcal{S}_{k_n, C}} \cdots \int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_r, y_r) dx_r dy_r. \end{aligned}$$

The r -fold integral can be bounded from above using the induction hypothesis as

$$O(k_n^{1+\epsilon-2\alpha}) \left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}; p_1) f(x_1, y_1) dx_1 dy_1 \right)^r. \quad (40)$$

Finally, we use that $k_n^{1+\epsilon-2\alpha} = o\left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}; p_1) \mu_n(dp_1)\right)$. To see this, we will firstly show that $k_n^{1+\epsilon-2\alpha} = o\left(n k_n^{-(2\alpha+1)}\right)$ and then apply (39), which says that

$$\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}; p_1) f(x_1, y_1) dx_1 dy_1 = \Theta\left(n k_n^{-(2\alpha+1)}\right).$$

By our choice of ϵ , we have that $\frac{1}{2} + \epsilon < \alpha$, which implies that $\frac{2+\epsilon}{1+2\alpha} < 1$, and hence, using $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$,

$$\frac{k_n^{1+\epsilon-2\alpha}}{n k_n^{-(2\alpha+1)}} = n^{-1} k_n^{2+\epsilon} = O\left(n^{-1} n^{\frac{2+\epsilon}{1+2\alpha}}\right) = o(n^{-1} n) = o(1).$$

Having shown that $k_n^{1+\epsilon-2\alpha} = o\left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}; p_1) \mu_n(dp_1)\right)$, we therefore conclude from (40) that

$$\begin{aligned} & \int \int_{\mathcal{F}_\epsilon(k_n)^c} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1} \\ &= o\left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) \mu_n(dp_1)\right) \left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1\right)^r \\ &= o\left(\left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1\right)^{r+1}\right). \end{aligned}$$

For the integration over $\mathcal{F}_\epsilon(k_n)$, recall that by Lemma 3.9,

$$\varphi(\{p_1, \dots, p_{r+1}\}, k_n; p_1, \dots, p_s) = (1+o(1))\varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s)\varphi(\{p_{r+1}\}, k_n; p_1, \dots, p_s) + O(k_n^{-C}).$$

This implies that

$$\begin{aligned}
& \int \int_{\mathcal{F}_\epsilon(k_n)} \varphi(\{p_1, \dots, p_{r+1}\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1} \\
&= (1 + o(1)) \int \int_{\mathcal{F}_\epsilon(k_n)} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) \varphi(\{p_{r+1}\}; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots \\
&\quad \times f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1} \\
&\quad + O(k_n^{-C}) \int \int_{\mathcal{F}_\epsilon(k_n)} f(x_1, y_1) dx_1 dy_1 \cdots f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1} =: M + E.
\end{aligned}$$

To finish the induction step we have to show that

$$M = (1 + o(1)) \left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) \mu_n(dp_1) \right)^{r+1}$$

and

$$E = o \left(\left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) \mu_n(dp_1) \right)^{r+1} \right).$$

For M , note that we can factorize to integration over p_1, \dots, p_r and over p_{r+1} .

Furthermore, we note that the condition $(p_1, \dots, p_{r+1}) \in \mathcal{F}_\epsilon(k_n)$ implies that (writing $p_{r+1} = (x_{r+1}, y_{r+1})$) the horizontal coordinate x_{r+1} falls into an interval I_n of length L_n , satisfying

$$\frac{\pi n}{\nu} - 2rk_n^{1+\epsilon} \leq L_n \leq \frac{\pi n}{\nu} - 2k_n^{1+\epsilon}.$$

As $k_n^{1+\epsilon} = O(n^{\frac{1+\epsilon}{2\alpha+1}}) = o(n)$ for $\epsilon < 1$ and $\alpha > \frac{1}{2}$, this shows that the length of the integration range in x_{r+1} satisfies $L_n = (1 + o(1)) \frac{\pi n}{\nu}$. Thus, we have that

$$\begin{aligned}
M &= (1 + o(1)) \int_{I_n} \int_{y_{k_n, c}^-}^{y_{k_n, c}^+} \varphi(\{p_{r+1}\}, k_n; p_1, \dots, p_s) \frac{\alpha \nu}{\pi} e^{-\alpha y_{r+1}} dy_{r+1} dx_{r+1} \\
&\quad \times \int_{\mathcal{S}_{k_n, C}} \cdots \int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_r, y_r) dx_r dy_r \\
&= (1 + o(1)) n \int_{y_{k_n, c}^-}^{y_{k_n, c}^+} \varphi(\{p_{r+1}\}, k_n; p_1, \dots, p_s) \alpha e^{-\alpha y_{r+1}} dy_{r+1} \\
&\quad \times \int_{\mathcal{S}_{k_n, C}} \cdots \int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1, \dots, p_r\}; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_r, y_r) dx_r dy_r \\
&= (1 + o(1)) \int_{\mathcal{S}_{k_n, C}} \varphi(\{p_{r+1}\}, k_n; p_1, \dots, p_s) \mu_n(dp_1) \\
&\quad \times \int_{\mathcal{S}_{k_n, C}} \cdots \int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_s) f(x_1, y_1) dx_1 dy_1 \cdots f(x_r, y_r) dx_r dy_r.
\end{aligned}$$

By applying the base case of the induction to the first factor and the induction hypothesis to the second one, we have derived that

$$M = (1 + o(1)) \left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1 \right)^{r+1}.$$

Finally, for E we observe that

$$E = O(k_n^{-C}) \int \int_{\mathcal{F}_\epsilon(k_n)} f(x_1, y_1) dx_1 dy_1 \cdots f(x_{r+1}, y_{r+1}) dx_{r+1} dy_{r+1} = O(k_n^{-C}) (1 + o(1)) n^{r+1}.$$

Recall that again by (39),

$$\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1 = \Theta \left(n k_n^{-(2\alpha+1)} \right),$$

which implies that

$$\left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1 \right)^{r+1} = \Theta \left(n^{r+1} k_n^{-r(2\alpha+1)} \right).$$

For $C > r(2\alpha + 1)$, we can conclude that indeed

$$E = O \left(n^{r+1} k_n^{-C} \right) = o \left(n^{r+1} k_n^{-r(2\alpha+1)} \right) = o \left(\left(\int_{\mathcal{S}_{k_n, C}} \varphi(\{p_1\}, k_n; p_1) \mu_n(dp_1) \right)^{r+1} \right).$$

□

We now proof the result for the factorial moments.

Proof of Lemma 3.10. We give the proof for the Poissonized KPKVB model. The proof for the finite box model G_{box} follows using similar arguments.

First of all, we observe that

$$\binom{N_{\text{Po}}(k_n)}{r} = \frac{1}{r!} \sum_{p_1, \dots, p_r \in V(G_{\text{Po}})}^{\neq} \mathbb{1}_{\{D_{G_{\text{Po}}}(p_1) = \dots = D_{G_{\text{Po}}}(p_r) = k_n\}}.$$

This can be seen by induction on r . For $r = 1$, the claim is clear. Assuming it holds for $r \geq 1$, by the induction hypothesis,

$$\begin{aligned} \binom{N_{\text{Po}}(k_n)}{r+1} &= \binom{N_{\text{Po}}(k_n)}{r} \frac{N_{\text{Po}}(k_n) - r}{r+1} \\ &= \frac{1}{(r+1)!} \sum_{p_1, \dots, p_r \in V(G_{\text{Po}})}^{\neq} \mathbb{1}_{\{D_{G_{\text{Po}}}(p_1) = \dots = D_{G_{\text{Po}}}(p_r) = k_n\}} (N_{\text{Po}}(k_n) - r). \end{aligned}$$

Now, we can write

$$N_{\text{Po}}(k_n) = \sum_{\substack{p_{r+1} \in V(G_{\text{Po}}), \\ p_{r+1} \notin \{p_1, \dots, p_r\}}} \mathbb{1}_{\{D_{G_{\text{Po}}}(p_{r+1}) = k_n\}} + \sum_{p_{r+1} \in \{p_1, \dots, p_r\}} \mathbb{1}_{\{D_{G_{\text{Po}}}(p_{r+1}) = k_n\}}.$$

The first sum leads to the right-hand side of the claim for $r + 1$, whereas the second sum will cancel with the $-r$.

By the Campbell-Mecke formula

$$\begin{aligned} \mathbb{E} \left[\binom{N_{\text{Po}}(k_n)}{r} \right] &= \frac{1}{r!} \mathbb{E} \left[\sum_{p_1, \dots, p_r \in V(G_{\text{Po}})}^{\neq} \mathbb{1}_{\{D_{G_{\text{Po}}}(p_1) = \dots = D_{G_{\text{Po}}}(p_r) = k_n\}} \right] \\ &= \frac{1}{r!} \int_{\mathcal{R}} \dots \int_{\mathcal{R}} \varphi_{\text{Po}}(\{p_1, \dots, p_r\}, p_1, \dots, p_r) f(x_1, y_1) dx_1 dy_1 \dots f(x_r, y_r) dx_r dy_r, \end{aligned}$$

where we integrate over r additional points which we can think of as being added independently and with the same distribution as the vertices of the poissonized KPKVB model G_{Po} in the upper half-plane coordinates.

With $r = 1$, it follows that

$$\mathbb{E}[N_{\text{Po}}(k_n)] = \int_{\mathcal{R}} \phi_{\text{Po}}(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1,$$

which yields that the right-hand side of the claim of the lemma can be rewritten as

$$\frac{1}{r!} (\mathbb{E}[N_{\text{Po}}(k_n)])^r = \frac{1}{r!} \left(\int_{\mathcal{R}} \phi_{\text{Po}}(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1, \right)^r.$$

Using Lemma 3.11, we conclude that

$$\begin{aligned} & \mathbb{E} \left[\binom{N_{\text{Po}}(k_n)}{r} \right] \\ &= \frac{1}{r!} \int_{\mathcal{R}} \cdots \int_{\mathcal{R}} \varphi_{\text{Po}}(\{p_1, \dots, p_r\}, k_n; p_1, \dots, p_r) f(x_1, y_1) dx_1 dy_1 \cdots f(x_r, y_r) dx_r dy_r \\ &= (1 + o(1)) \frac{1}{r!} \left(\int_{\mathcal{R}} \varphi_{\text{Po}}(\{p_1\}, k_n; p_1) f(x_1, y_1) dx_1 dy_1 \right)^r \\ &= (1 + o(1)) \frac{1}{r!} (\mathbb{E}[N_{\text{Po}}(k_n)])^r. \end{aligned}$$

□

3.6 Coupling G_n to G_{Po}

In the previous sections we have established results for the degrees and the factorial moments of the degree k_n nodes in the Poissonized KPKVB and finite box model. Our intended result, however, was for the degree distribution in the original KPKVB model. In order to extend the result for the Poissonized KPKVB model to the original model we will use a coupling argument to show that the expected difference between the number of degree k_n nodes is negligible.

Lemma 3.12. *As $n \rightarrow \infty$, it holds that for $0 \leq k_n \leq n - 1$,*

$$\mathbb{E}[|N_n(k_n) - N_{\text{Po}}(k_n)|] = o(\mathbb{E}[N_{\text{Po}}(k_n)]).$$

Proof. We couple both models by taking an infinite supply of i.i.d. points u_1, u_2, \dots chosen according to the (α, R) -quasi uniform distribution and letting the vertices of $G(n; \alpha, \nu)$ be u_1, \dots, u_n and the vertices of $G_{\text{Po}}(n; \alpha, \nu)$ be u_1, \dots, u_N with $N \stackrel{\text{d}}{=} \text{Po}(n)$ independently of u_1, u_2, \dots . Thus, under this coupling, the only difference between $G_n = G(n; \alpha, \nu)$ and $G_{\text{Po}} = G_{\text{Po}}(n; \alpha, \nu)$ is the number of points. Note that since N is Poisson with mean n , it follows from the Chernoff bound (see also equation (135) in the paper) that we may assume that $n - C\sqrt{n \log n} \leq N \leq n + C\sqrt{n \log n}$. To keep notation simple we will suppress this conditioning in the derivations.

Clearly, if $N = n$ the graphs are the same. So we will consider the two cases $n - C\sqrt{n \log n} \leq N < n$ and $n < N \leq n + C\sqrt{n \log n}$. We will prove the latter case. The other case uses similar arguments and hence we omit the details here.

If $n < N \leq n + C\sqrt{n \log n}$ then the G_n has less vertices than G_{Po} . Write $V_n(k_n)$ and $V_{\text{Po}}(k_n)$ to denote the set of vertices that have degree k_n in G_n and G_{Po} , respectively. Then since the vertices u_{n+1}, \dots, u_N are not present in G_n ,

$$\begin{aligned} |N_n(k_n) - N_{\text{Po}}(k_n)| &= \sum_{i=1}^N \mathbb{1}_{\{u_i \in V_n(k_n) \Delta V_{\text{Po}}(k_n)\}} \\ &= \sum_{i=1}^n \mathbb{1}_{\{u_i \in V_n(k_n) \Delta V_{\text{Po}}(k_n)\}} + \sum_{i=n+1}^N \mathbb{1}_{\{u_i \in V_{\text{Po}}(k_n)\}}. \end{aligned}$$

Let D_{Po} denote the degree in the Poissonized KPKVB model of a point u placed according to the (α, R) -quasi uniform distribution. Then, using the Campbell-Mecke formula, the expectation of the second summation equals

$$\begin{aligned} (N - n)\mathbb{P}(D_{\text{Po}} = k_n) &= (N - n) \int_0^R \mathbb{P}(\text{Po}(\mu_{\text{Po},n}(y)) = k_n) \alpha e^{-\alpha y} dy \\ &\leq (1 + o(1))C\sqrt{n \log np_{k_n}} = o(\mathbb{E}[N_{\text{Po}}(k_n)]). \end{aligned}$$

Therefore it remains to consider the first summation.

Let $D_n(u)$ and $D_{\text{Po}}(u)$ denote the degree of a point u in G_n and G_{Po} , respectively. Then there are two scenarios to consider: 1) either $D_n(u_i) = k_n$ and $D_{\text{Po}}(u_i) \neq k_n$ or 2) $D_n(u_i) \neq k_n$ and $D_{\text{Po}}(u_i) = k_n$. In the first case, since u_i is present in both graphs it follows that $D_{\text{Po}}(u_i) > k_n$. Similarly, for the second case it must hold that $D_n(u_i) < k_n$. Hence we have

$$\begin{aligned} \sum_{i=1}^n \mathbb{1}_{\{u_i \in V_n(k_n) \Delta V_{\text{Po}}(k_n)\}} \\ = \sum_{i=1}^n \mathbb{1}_{\{D_n(u_i) = k_n, D_{\text{Po}}(u_i) > k_n\}} + \sum_{i=1}^n \mathbb{1}_{\{D_n(u_i) < k_n, D_{\text{Po}}(u_i) = k_n\}}. \end{aligned}$$

Let us first consider the second summation, i.e. the case where the node has degree smaller than k_n in G_n . Taking the expectation gives $n\mathbb{P}(D_n < k_n, D_{\text{Po}} = k_n)$, where D_n denotes the degree in the KPKVB model of a point u placed according to the (α, R) -quasi uniform distribution. We now observe that because the points u_1, \dots, u_N used to couple the graphs are independent, we can view the graph G_n as being obtained from G_{Po} by removing $N - n$ points, uniformly at random. Therefore if a point has degree k_n in G_{Po} but smaller degree in G_n , this means that at least one of its neighbors was removed. Denote by $Z(n)$ a random variable with a Hypergeometric distribution, for taking $N - n$ draws from a population of size N , where there are k_n good objects. That is, $Z(n)$ denote the number of removed neighbors of a node u with degree k_n in G_{Po} . We then have

$$\begin{aligned} \mathbb{P}(D_n < k_n, D_{\text{Po}} = k_n) &= \mathbb{P}(Z(n) > 1) \mathbb{P}(D_{\text{Po}} = k_n) \\ &\leq \mathbb{E}[Z(n)] \mathbb{P}(D_{\text{Po}} = k_n) = \frac{(N - n)k_n}{N} \mathbb{P}(D_{\text{Po}} = k_n). \end{aligned}$$

Because $\alpha > 1/2$ and $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$ it holds that $k_n = o\left(\sqrt{\frac{n}{\log n}}\right)$. Since $N = \Theta(n)$ and $N - n \leq O(\sqrt{n \log n})$ it then follows that $\frac{(N - n)k_n}{N} = o(1)$, from which we conclude that

$$\begin{aligned} \mathbb{E}\left[\sum_{i=1}^n \mathbb{1}_{\{D_n(u_i) < k_n, D_{\text{Po}}(u_i) = k_n\}}\right] &= n\mathbb{P}(D_n < k_n, D_{\text{Po}} = k_n) \\ &\leq o(1) n\mathbb{P}(D_{\text{Po}} = k_n) = o(\mathbb{E}[N_{\text{Po}}(k_n)]). \end{aligned}$$

We now proceed with the other summation, for the case where a vertex has degree k_n in G_n but larger degree in G_{Po} . Since the degree of u in G_{Po} can be at most $N - n$ larger we have

$$\mathbb{P}(D_n(u) = k_n, D_{\text{Po}}(u) > k_n) = \sum_{t=1}^{N-n} \mathbb{P}(D_n(u) = k_n, D_{\text{Po}}(u) = k_n + t).$$

Using that the graph G_n can be seen as being obtained from G_{Po} by removing $N - n$ points uniformly at random, a point with degree $k_n + t$ in G_{Po} can only have degree k_n in G_n if exactly t of its neighbors were removed. Let us therefore denote by $Z(n, t)$ a random variable with a Hypergeometric distribution, for taking $N - n$ draws from a population of size N , where there are $k_n + t$ good objects. Then

$$\mathbb{P}(D_n(u) = k_n, D_{\text{Po}}(u) = k_n + t) = \mathbb{P}(Z(n, t) = t) \mathbb{P}(D_{\text{Po}} = k_n + t).$$

Recall that, for any $0 < \varepsilon < 1$

$$\begin{aligned}\mathbb{P}(D_{\text{Po}} = k_n + t) &= \int_0^R \mathbb{P}(\text{Po}(\mu_{\text{Po},n}(y)) = k_n + t) \alpha e^{-\alpha y} dy \\ &= \int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\mu_{\text{Po},n}(y)) = k_n + t) \alpha e^{-\alpha y} dy \\ &\quad + \int_{(1-\varepsilon)R}^R \mathbb{P}(\text{Po}(\mu_{\text{Po},n}(y)) = k_n + t) \alpha e^{-\alpha y} dy\end{aligned}$$

By Lemma 3.3 the first part is $(1 + o(1))p_{k_n+t}$ while the second part is $O(n^{-2\alpha(1-\varepsilon)})$ and hence

$$\mathbb{P}(D_{\text{Po}} = k_n + t) \leq O(1) \left(p_{k_n+t} + n^{-2\alpha(1-\varepsilon)} \right).$$

In addition we have that $\mathbb{P}(Z(n, t) = t) \leq O(1) \frac{\mathbb{E}[Z(n, t)]}{t}$. We thus obtain

$$\begin{aligned}\sum_{t=1}^{N-n} \mathbb{P}(Z(n, t) = t) \mathbb{P}(D_{\text{Po}} = k_n + t) &\leq O(1) \sum_{t=1}^{N-n} \frac{\mathbb{E}[Z(n, t)]}{t} \left(p_{k_n+t} + n^{-2\alpha(1-\varepsilon)} \right) \\ &= O(1) \sum_{t=1}^{N-n} \frac{(N-m)(k_n+t)}{N} \left(p_{k_n+t} + n^{-2\alpha(1-\varepsilon)} \right) \\ &= O\left(\sqrt{\frac{\log n}{n}}\right) \sum_{t=1}^{N-n} \frac{k_n}{t} \left(p_{k_n+t} + n^{-2\alpha(1-\varepsilon)} \right) \\ &\quad + O\left(\sqrt{\frac{\log n}{n}}\right) \sum_{t=1}^{N-n} \left(p_{k_n+t} + n^{-2\alpha(1-\varepsilon)} \right),\end{aligned}$$

where we used that $\frac{N-n}{N} = O\left(\sqrt{\frac{\log n}{n}}\right)$.

We will show that both summations are $o(p_{k_n})$. For the first summation we recall that $\frac{k_n(\log n)^{3/2}}{\sqrt{n}} = o(1)$ for $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$, while for $\varepsilon > 0$ small enough $n^{-2\alpha(1-\varepsilon)} = o\left(k_n^{-(2\alpha+1)}\right) = o(p_{k_n})$. Hence, since $p_{k_n+t} \leq p_{k_n}$,

$$\begin{aligned}O\left(\sqrt{\frac{\log n}{n}}\right) \sum_{t=1}^{N-n} \frac{k_n}{t} \left(p_{k_n+t} + n^{-2\alpha(1-\varepsilon)} \right) &\leq O\left(k_n \sqrt{\frac{\log n}{n}} p_{k_n}\right) \sum_{t=1}^{N-n} \frac{1}{t} \\ &= O\left(\frac{k_n(\log n)^{3/2}}{\sqrt{n}}\right) p_{k_n} = o(p_{k_n}).\end{aligned}$$

For the other summation we use that

$$\sum_{t=1}^{N-n} p_{k_n+t} \leq \sum_{t=1}^{\infty} p_{k_n+t} \leq O(k_n^{-2\alpha}) = O(k_n p_{k_n}),$$

together with the fact that for ε small enough, $\log(n)n^{-2\alpha(1-\varepsilon)} = o\left(k_n^{-(2\alpha+1)}\right) = o(p_{k_n})$. This implies that

$$\begin{aligned}O\left(\sqrt{\frac{\log n}{n}}\right) \sum_{t=1}^{N-n} \left(p_{k_n+t} + n^{-2\alpha(1-\varepsilon)} \right) &\leq O\left(\sqrt{\frac{\log n}{n}} k_n p_{k_n}\right) + O\left((N-n) \sqrt{\frac{\log n}{n}} n^{-2\alpha(1-\varepsilon)}\right) \\ &\leq o(p_{k_n}) + O\left((\log n) n^{-2\alpha(1-\varepsilon)}\right) = o(p_{k_n}).\end{aligned}$$

It now follows that

$$\begin{aligned}
n\mathbb{P}(D_n(u) = k_n, D_{Po}(u) > k_n) &= \sum_{t=1}^{N-n} \mathbb{P}(D_n = k_n, D_{Po} = k_n + t) \\
&= n \sum_{t=1}^{N-n} \mathbb{P}(Z(n, t) = t) \mathbb{P}(D_{Po} = k_n + t) \\
&= n o(\mathbb{P}(D_{Po} = k_n)) = o(\mathbb{E}[N_{Po}(k_n)]),
\end{aligned}$$

which finishes the proof for the case where $N > n$. \square

3.7 Proof of Theorem 1.1

We now have all necessary ingredient to proof the main result on the degrees, Theorem 1.1.

Proof of Theorem 1.1. Recall that we shall only give the proof for the case where $k_n \rightarrow \infty$, since result (i) for fixed $k = O(1)$ follows from [21].

- (i) First we recall that the statements regarding p_k and its asymptotic behavior follow from Equation 20 and Equation 21.

By Lemma 3.4,

$$\mathbb{E}[N_{Po}(k_n)] = (1 + o(1))np_{k_n}.$$

Using Lemma 3.10 with $r = 2$ we have that $\mathbb{E}\left[\binom{N_{Po}(k_n)}{2}\right] = (1 + o(1))\frac{(\mathbb{E}[N_{Po}(k_n)])^2}{2}$, which implies $\mathbb{E}[N_{Po}(k_n)^2] = 2\mathbb{E}\left[\binom{N_{Po}(k_n)}{2}\right] + \mathbb{E}[N_{Po}(k_n)] = (1 + o(1))(\mathbb{E}[N_{Po}(k_n)]^2 + o((\mathbb{E}[N_{Po}(k_n)])^2))$ (because $\mathbb{E}[N_{Po}(k_n)] = (1 + o(1))np_{k_n} \rightarrow \infty$). Hence, by Chebychev for any $\epsilon > 0$,

$$\mathbb{P}(|N_{Po}(k_n) - \mathbb{E}[N_{Po}(k_n)]| \geq \epsilon \mathbb{E}[N_{Po}(k_n)]) \leq \frac{\text{Var}(N_{Po}(k_n))}{\epsilon^2 (\mathbb{E}[N_{Po}(k_n)])^2} = o(1).$$

As $N_n(k_n) = N_{Po}(k_n) + N_n(k_n) - N_{Po}(k_n) = N_{Po}(k_n) \pm |N_n(k_n) - N_{Po}(k_n)|$ (where the sign depends on whether $N_n(k_n) > N_{Po}(k_n)$ or not), due to Lemma 3.12, we have that

$$\mathbb{E}[N_n(k_n)] = \mathbb{E}[N_{Po}(k_n)] \pm \mathbb{E}[|N_n(k_n) - N_{Po}(k_n)|] = (1 + o(1))\mathbb{E}[N_{Po}(k_n)] = (1 + o(1))np_{k_n}.$$

- (ii) Let $\zeta = 2\alpha\xi^{2\alpha}c^{-(2\alpha+1)} \in \mathbb{R}$. The proof consists of showing that $\mathbb{E}\left[\binom{N_{Po}(k_n)}{r}\right] \rightarrow \frac{\zeta^r}{r!}$ for every positive integer r and then applying [3, Theorem 8.3.1].

If $k_n = (1 + o(1))cn^{\frac{1}{2\alpha+1}}$, then by Lemma 3.4,

$$\begin{aligned}
\mathbb{E}[N_{Po}(k_n)] &= (1 + o(1))2\alpha\xi^{2\alpha}n(1 + o(1))^{-(2\alpha+1)}c^{-(2\alpha+1)}n^{-1} \\
&= (1 + o(1))2\alpha\xi^{2\alpha}c^{-(2\alpha+1)} = (1 + o(1))\zeta,
\end{aligned}$$

which implies $\mathbb{E}[N_{Po}(k_n)] \rightarrow \zeta$ (as ζ is a positive constant). From Lemma 3.10, it then follows that $\mathbb{E}\left[\binom{N_{Po}(k_n)}{r}\right] = (1 + o(1))\frac{(\mathbb{E}[N_{Po}(k_n)])^r}{r!} \rightarrow \frac{\zeta^r}{r!}$. Thus, it follows from [3, Theorem 8.3.1] that $N_{Po}(k_n) \xrightarrow{d} Po(\zeta)$.

Finally, since $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$, by Lemma 3.12, $\mathbb{E}[|N_n(k_n) - N_{Po}(k_n)|] = o(\mathbb{E}[N_{Po}(k_n)]) = o(\zeta)$, from which it follows that $\mathbb{P}(|N_n(k_n) - N_{Po}(k_n)| \geq 1) \leq \mathbb{E}[|N_n(k_n) - N_{Po}(k_n)|] = o(\zeta)$. Hence, it also holds that $N_n(k_n) \xrightarrow{d} Po(\zeta)$.

(iii) We will show that in this case $\mathbb{E}[N_n(k_n)] = o(1)$. This then implies, by Markov's inequality,

$$\mathbb{P}(N_n(k_n) > 0) \leq \mathbb{E}[N_n(k_n)] = o(1).$$

First we observe that as the poissonized KPKVB model G_{Po} has the same intensity measure as the original KPKVB model with a fixed number n of points, the expected degree of a vertex of the KPKVB model with radial coordinate $r = R - y$ is given by $\mu_{\text{Po}}(y)$ and hence,

$$\mathbb{E}[N_n(k_n)] = n \int_0^R \mathbb{P}(\text{Bin}(n-1, \mu_{\text{Po}}(y)/n) = k_n) \frac{\alpha \sinh(\alpha(R-y))}{\cosh(\alpha R) - 1} dy.$$

Fix $0 < \varepsilon < \frac{4\alpha-1}{4\alpha+2} \wedge \frac{2\alpha-1}{2\alpha}$. We first show that we only need to consider integration up to $y \leq (1-\varepsilon)R$. By our choice of ε , $2\alpha(1-\varepsilon) > 1$, so that

$$\frac{\cosh(\alpha\varepsilon R) - 1}{\cosh(\alpha R) - 1} = O\left(n^{-2\alpha(1-\varepsilon)}\right) = o\left(n^{-1}\right).$$

This implies

$$n \int_{(1-\varepsilon)R}^R \mathbb{P}(\text{Bin}(n-1, \mu_{\text{Po}}(y)/n) = k_n) \frac{\alpha \sinh(\alpha(R-y))}{\cosh(\alpha R) - 1} dy \leq n \frac{\cosh(\alpha\varepsilon R)}{\cosh(\alpha R) - 1} = o(1),$$

and thus it is enough to show that

$$n \int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Bin}(n-1, \mu_{\text{Po}}(y)) = k_n) \frac{\alpha \sinh(\alpha(R-y))}{\cosh(\alpha R) - 1} dy = o(1).$$

Note that for all $0 \leq y \leq (1-\varepsilon)R$ we have

$$\frac{\alpha \sinh(\alpha(R-y))}{\cosh(\alpha R) - 1} = (1 + o(1))\alpha e^{-\alpha y}.$$

Hence, by bounding the Binomial probability (see Lemma D.3)

$$\begin{aligned} & n \int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Bin}(n-1, \mu_{\text{Po}}(y)) = k_n) \frac{\alpha \sinh(\alpha(R-y))}{\cosh(\alpha R) - 1} dy \\ & \leq (1 + o(1)) \frac{e}{\sqrt{2\pi}} \sqrt{\frac{n}{n-k_n}} n \int_0^{(1-\varepsilon)R} \mathbb{P}(\text{Po}(\mu_{\text{Po}}(y)) = k_n) \alpha e^{-\alpha y} dy \\ & \leq (1 + o(1)) \frac{e}{\sqrt{2\pi}} \sqrt{\frac{n}{n-k_n}} n p_{k_n} = O\left(\sqrt{\frac{n}{n-k_n}} n k_n^{-(2\alpha+1)}\right). \end{aligned}$$

We shall now consider two cases: $n^{\frac{1}{2\alpha+1}} \ll k_n < n^{1-\varepsilon}$ and $n^{1-\varepsilon} \leq k_n \leq n-1$.

If $n^{\frac{1}{2\alpha+1}} \ll k_n < n^{1-\varepsilon}$ then $\sqrt{\frac{n}{n-k_n}} = 1 + o(1)$ and hence

$$\sqrt{\frac{n}{n-k_n}} n k_n^{-(2\alpha+1)} = O\left(n k_n^{-(2\alpha+1)}\right) = o(1).$$

For $k_n \geq n^{1-\varepsilon}$ we have, by our choice of ε , that $\frac{3}{2} - (2\alpha+1)(1-\varepsilon) < 0$, and thus

$$\sqrt{\frac{n}{n-k_n}} n k_n^{-(2\alpha+1)} = O\left(n^{\frac{3}{2}} k_n^{-(2\alpha+1)}\right) = O\left(n^{\frac{3}{2} - (2\alpha+1)(1-\varepsilon)}\right) = o(1).$$

□

4 Clustering of the typical point in G_∞

Similar to the proof for the degrees we plan to make use of the Campbell-Mecke formula for comparing the clustering coefficient and function of $G_{\mathcal{P}_0}$ with certain quantities associated with G_∞ . Recall that a *typical point* is a point $p_0 = (0, y)$, where in some computations the height y will be fixed, but eventually we shall take it exponentially distributed with parameter α , and independent of \mathcal{P} .

Since $c(G)$ is defined as an average over all vertices of the graph, it is not immediately obvious how to meaningfully define a corresponding notion for infinite graphs, and similarly for the clustering function, the degree sequence, etc. We can however without any issues speak of the (expected) clustering coefficient of the typical point, or the expected clustering coefficient given that it has degree k , or the distribution of the degree of the typical point. (All considered in the graph obtained from G_∞ by adding the typical point to its vertex set.)

If $p = (x, y) \in \mathbb{R} \times [0, \infty)$ is a point, not necessarily part of the Poisson process, then we will write

$$\mu(y) = \mu(p) := \mu(\mathcal{B}_\infty(p)).$$

Integrating the intensity function of \mathcal{P} over $\mathcal{B}_\infty(p)$ gives, using $\alpha > \frac{1}{2}$,

$$\begin{aligned} \mu(y) &= \int_{\mathcal{B}_\infty(p)} f(x', y') dx' dy' = \int_0^\infty \int_{-e^{(y+y')/2}}^{e^{(y+y')/2}} \frac{\alpha\nu}{\pi} e^{-\alpha y'} dx' dy' \\ &= \int_0^\infty 2e^{(y+y')/2} \frac{\alpha\nu}{\pi} e^{-\alpha y'} dy' = \frac{2\alpha\nu e^{y/2}}{\pi} \int_0^\infty e^{(\frac{1}{2}-\alpha)y'} dy' \\ &= \frac{2\alpha\nu e^{y/2}}{\pi(\alpha - \frac{1}{2})} = \xi e^{y/2}. \end{aligned}$$

4.1 The expected clustering coefficient and function of the typical point

Let the random variable C denote the clustering coefficient of the typical point $(0, y)$, in the graph obtained from G_∞ by adding $(0, y)$. We now *define*

$$\gamma := \mathbb{E}[C], \quad \gamma(k) := \mathbb{E}[C | D = k].$$

(Where we take the expectation over both the Poisson point process \mathcal{P} and $y \stackrel{d}{=} \exp(\alpha)$, independently of the Poisson process \mathcal{P} .) We shall show shortly that these take on the values stated in Theorem 1.2 and 1.3.

For any fixed value $y_0 > 0$, the set of points inside $\mathcal{B}_\infty((0, y_0))$ is a Poisson process with intensity $f \cdot 1_{\mathcal{B}_\infty((0, y_0))}$. As $\mu(\mathcal{B}_\infty((0, y_0))) = \mu(y_0) = \xi e^{y_0/2} < \infty$, this can be described alternatively by first picking $N \stackrel{d}{=} \text{Po}(\mu(y_0))$ and then taking N i.i.d. points in $\mathcal{B}_\infty((0, y_0))$ according to the probability density $f \cdot 1_{\mathcal{B}_\infty((0, y_0))} / \mu(y_0)$. (That is, the intensity function of the Poisson point process, but set to zero outside of $\mathcal{B}_\infty((0, y_0))$ and re-normalized in such a way that it integrates to one.) Hence, if we condition on the event that y takes on some fixed value y_0 and that there are exactly k points of \mathcal{P} inside $\mathcal{B}_\infty((0, y_0))$, then those k points behave like k i.i.d. points in $\mathcal{B}_\infty((0, y_0))$ chosen according to the mentioned re-normalized probability density function. This shows that, for every $k \geq 2$:

$$\mathbb{E}[C | D = k, y = y_0] = \frac{1}{\binom{k}{2}} \mathbb{E} \left(\sum_{1 \leq i < j \leq k} \mathbb{1}_{\{u_i \in \mathcal{B}_\infty(u_j)\}} \right) = \mathbb{E} [\mathbb{1}_{\{u_1 \in \mathcal{B}_\infty(u_2)\}}],$$

where u_1, \dots, u_n are i.i.d. points in $\mathcal{B}_\infty((0, y_0))$ with the above mentioned density. Note that this does not depend on the value of k . For notational convenience, we will write

$$P(y_0) := \mathbb{E} [\mathbb{1}_{\{u_1 \in \mathcal{B}_\infty(u_2)\}}],$$

with u_1, u_2 as above.

We now observe that

$$\gamma(k) = \mathbb{E}[C|D=k] = \int_0^\infty \mathbb{E}[C|D=k, y=y_0] g_k(y_0) dy_0,$$

where g_k denotes the density of y conditional on $D=k$. That is,

$$g_k(y_0) = \frac{\rho(y_0, k) \alpha e^{-\alpha y_0}}{\int_0^\infty \rho(t, k) \alpha e^{-\alpha t} dt} = \frac{1}{p_k} \cdot \rho(y_0, k) \alpha e^{-\alpha y_0}.$$

Hence,

$$\gamma(k) = \frac{1}{p_k} \cdot \int_0^\infty P(y_0) \rho(y_0, k) \alpha e^{-\alpha y_0} dy_0. \quad (41)$$

This also gives

$$\begin{aligned} \gamma &= \mathbb{E}[C] = \sum_{k \geq 2} \mathbb{E}[C|D=k] \mathbb{P}(D=k) \\ &= \int_0^\infty P(y_0) \left(\sum_{k=2}^\infty \rho(y_0, k) \right) \alpha e^{-\alpha y_0} dy_0 \\ &= \int_0^\infty P(y_0) (1 - \rho(y_0, 0) - \rho(y_0, 1)) \alpha e^{-\alpha y_0} dy_0. \end{aligned} \quad (42)$$

A key step is to derive the following explicit expression for $P(y)$.

Lemma 4.1. *If $\alpha \neq 1$, then*

$$\begin{aligned} P(y) &= -\frac{1}{8(\alpha-1)\alpha} + \frac{(\alpha-1/2)e^{-\frac{1}{2}y}}{\alpha-1} - \frac{(\alpha-1/2)^2 e^{-y}}{4(\alpha-1)^2} \\ &\quad + (e^{-\frac{1}{2}y})^{4\alpha-2} \left(\frac{2^{-4\alpha-1}(3\alpha-1)}{\alpha(\alpha-1)^2} + \frac{(\alpha-1/2)B^-(1/2; 1+2\alpha, -2+2\alpha)}{2(\alpha-1)\alpha} \right) \\ &\quad + \frac{(1-e^{-\frac{1}{2}y})^{2\alpha}}{8(\alpha-1)\alpha} - \frac{(e^{-\frac{1}{2}y})^{4\alpha-2} B^-(1-e^{-\frac{1}{2}y}; 2\alpha, 3-4\alpha)}{4(\alpha-1)} \end{aligned}$$

We will prove this lemma in a sequence of steps.

Recall that $P(y_0)$ is the probability that $u_1 = (x_1, y_1), u_2 = (x_2, y_2)$ are neighbours in G_∞ , where u_1, u_2 are i.i.d. with probability density $f \cdot \mathbb{1}_{\mathcal{B}_\infty((0, y_0))} / \mu(y_0)$. In particular

$$\begin{aligned} \mathbb{P}(y_i > t) &= \frac{\nu\alpha}{\pi\mu(y_0)} \int_t^\infty \int_{-e^{(y+y_0)/2}}^{e^{(y+y_0)/2}} e^{-\alpha y} dx dy = \frac{\nu\alpha}{\pi\mu(y_0)} \int_t^\infty 2e^{(y+y_0)/2} \cdot e^{-\alpha y} dy \\ &= \frac{2\nu\alpha e^{y_0/2}}{\pi\xi e^{y_0/2}(\alpha - \frac{1}{2})} \cdot e^{(\frac{1}{2}-\alpha)t} = e^{(\frac{1}{2}-\alpha)t}, \end{aligned}$$

using that $\mu(y_0) = \xi e^{y_0/2} = \left(\frac{2\nu\alpha}{\pi(\alpha - \frac{1}{2})} \right) e^{y_0/2}$. Thus, y_1, y_2 are exponentially distributed with parameter $\alpha - \frac{1}{2}$. Now note that, for each $t > 0$, the probability density $f \cdot \mathbb{1}_{\mathcal{B}_\infty((0, y_0))} / \mu(y_0)$ is constant on $[-e^{(t+y_0)/2}, e^{(t+y_0)/2}] \times \{t\}$ and it vanishes on $(-\infty, -e^{(t+y_0)/2}) \times \{t\} \cup (e^{(t+y_0)/2}, \infty) \times \{t\}$.

Hence, given the height y_i of u_i , the x -coordinate of u_i is uniform in $[-e^{\frac{1}{2}(y+y_i)}, e^{\frac{1}{2}(y+y_i)}]$. With this in mind we define $P(y_0, y_1, y_2)$ to be the probability that $(0, y_0), (x_1, y_1), (x_2, y_2)$ satisfy $|x_1 - x_2| \leq e^{(y_1+y_2)/2}$, where x_1 and x_2 are independent uniform random variables in the intervals $[-e^{\frac{1}{2}(y_0+y_1)}, e^{\frac{1}{2}(y_0+y_1)}]$ and $[-e^{\frac{1}{2}(y_0+y_2)}, e^{\frac{1}{2}(y_0+y_2)}]$, respectively. We have that

$$P(y_0) = (\alpha - 1/2)^2 \int_0^\infty \int_0^\infty P(y_0, y_1, y_2) e^{-(\alpha-1/2)(y_1+y_2)} dy_2 dy_1. \quad (43)$$




Figure 5: Situation for the intersections of the connection intervals considered in Lemma 4.3, with $y_0 < y_1 < y_2$ fixed and for different cases of $0 \leq x_1 \leq e^{(y_0+y_1)/2}$. The top figure shows the case where $0 \leq x_1 \leq e^{(y_1+y_2)/2} - e^{(y_0+y_2)/2}$, while the bottom one shows the case $x_1 > e^{(y_1+y_2)/2} - e^{(y_0+y_2)/2}$. The solid red line indicates the range for x_2 such that the points p_0 , p_1 and p_2 form a triangle. The boundaries of their neighbourhoods are shown in, respectively, black, blue and green.

4.1.1 Determining $P(y_0, y_1, y_2)$

To compute the integral (43) it will be convenient to use the change of variable $z_i = e^{-y_i/2}$, for $i = 0, 1, 2$. We will write $y_i(z_i)$ to stress the dependence between y_i and z_i . The following result completely characterizes $P(y_0, y_1, y_2)$.

Lemma 4.2.

$$P(y_0(z_0), y_1(z_1), y_2(z_2)) = \begin{cases} 1, & \text{if } z_0 \geq z_1 + z_2, z_0 > z_1 > z_2, \\ 1 - G(z_0, z_1, z_2), & \text{if } z_0 < z_1 + z_2, z_0 > z_1 > z_2, \\ \frac{z_0}{z_1}, & \text{if } z_1 \geq z_0 + z_2, z_1 > \max(z_0, z_2), \\ \frac{z_0}{z_1} (1 - G(z_1, z_0, z_2)), & \text{if } z_1 < z_0 + z_2, z_1 > \max(z_0, z_2), \end{cases}$$

where

$$G(a, b, c) = \frac{1}{4} (b^{-1}c + bc^{-1} + a^2b^{-1}c^{-1} + 2 - 2ab^{-1} - 2ac^{-1}).$$

We split the proof of this lemma into a couple of smaller pieces. We begin with the following lemma.

Lemma 4.3. Let $z_i = e^{-y_i/2}$, $i = 0, 1, 2$. If $y_0 < y_1 < y_2$ (or equivalently $z_0 > z_1 > z_2$), then

$$P(y_0(z_0), y_1(z_1), y_2(z_2)) = \begin{cases} 1, & \text{if } z_0 \geq z_1 + z_2, \\ 1 - G(z_0, z_1, z_2), & \text{if } z_0 < z_1 + z_2 \end{cases}$$

Proof. Note that $P(y_0, y_1, y_2)$ is the probability that x_2 falls into the interval $[x_1 - e^{(y_1+y_2)/2}, x_1 + e^{(y_1+y_2)/2}]$, as well as into the interval $[-e^{(y_0+y_2)/2}, e^{(y_0+y_2)/2}]$. By symmetry considerations, we can take x_1 uniformly at random from $[0, e^{y_0/2+y_1/2}]$ as opposed to $[-e^{y_0/2+y_1/2}, e^{y_0/2+y_1/2}]$. Figure 5 shows the intersection of the intervals (red line) for two different cases for $x_1 \leq e^{(y_0+y_1)/2}$.

Since $y_0 < y_1 < y_2$ we have that $e^{(y_1+y_2)/2} > e^{(y_0+y_2)/2}$ and so, when $x_1 \geq 0$, the “right half” of the interval $[-e^{(y_0+y_2)/2}, e^{(y_0+y_2)/2}]$ is always covered by the interval $[x_1 - e^{(y_1+y_2)/2}, x_1 + e^{(y_1+y_2)/2}]$. If $e^{(y_1+y_2)/2} - e^{(y_0+y_1)/2} \geq e^{(y_0+y_2)/2}$ then the “left half” is always covered as well. In other words:

$$e^{(y_1+y_2)/2} - e^{(y_0+y_1)/2} \geq e^{(y_0+y_2)/2} \Rightarrow P(y_0, y_1, y_2) = 1.$$

Now consider the case where $e^{(y_1+y_2)/2} - e^{(y_0+y_1)/2} < e^{(y_0+y_2)/2}$. Then, if $x_1 \in [0, e^{(y_1+y_2)/2} - e^{(y_0+y_2)/2}]$ the whole interval $[-e^{(y_0+y_2)/2}, e^{(y_0+y_2)/2}]$ is still covered so that p_0, p_1 and p_2 form a triangle. If, on the other hand $e^{(y_1+y_2)/2} - e^{(y_0+y_2)/2} < x_1 \leq e^{(y_0+y_1)/2}$ then the probability that $|x_2 - x_1| \leq e^{(y_1+y_2)/2}$ equals

$$1 - \frac{x_1 - (e^{(y_1+y_2)/2} - e^{(y_0+y_2)/2})}{2e^{(y_0+y_2)/2}}.$$

Hence, when $e^{(y_1+y_2)/2} - e^{(y_0+y_1)/2} < e^{(y_0+y_2)/2}$ we have

$$\begin{aligned} P(y_0, y_1, y_2) &= \frac{e^{(y_1+y_2)/2} - e^{(y_0+y_2)/2}}{e^{(y_0+y_1)/2}} \\ &\quad + \int_{e^{(y_1+y_2)/2} - e^{(y_0+y_2)/2}}^{e^{(y_0+y_1)/2}} \left(1 - \frac{x_1 - (e^{(y_1+y_2)/2} - e^{(y_0+y_2)/2})}{2e^{(y_0+y_2)/2}}\right) \cdot \frac{1}{e^{(y_0+y_1)/2}} dx_1 \\ &= 1 - \frac{1}{2e^{y_0+y_1/2+y_2/2}} \int_0^{e^{(y_0+y_1)/2} + e^{(y_0+y_2)/2} - e^{(y_1+y_2)/2}} x_1 dx_1 \\ &= 1 - \frac{(e^{(y_0+y_1)/2} + e^{(y_0+y_2)/2} - e^{(y_1+y_2)/2})^2}{4e^{y_0+y_1/2+y_2/2}}. \end{aligned}$$

At this point it is convenient to rewrite everything in terms of $z_i := e^{-y_i/2}$. Note that $y_0 < y_1 < y_2$ if and only if $z_0 > z_1 > z_2$ while the condition $e^{(y_1+y_2)/2} - e^{(y_0+y_1)/2} < e^{(y_0+y_2)/2}$ becomes

$$e^{(y_1+y_2)/2} - e^{(y_0+y_1)/2} < e^{(y_0+y_2)/2} \Leftrightarrow z_1^{-1} z_2^{-1} < z_0^{-1} z_1^{-1} + z_0^{-1} z_2^{-1} \Leftrightarrow z_0 < z_1 + z_2.$$

We now conclude that

$$P(y_0(z_0), y_1(z_1), y_2(z_2)) = 1 \quad \text{if } z_0 > z_1 > z_2 \text{ and } z_0 \geq z_1 + z_2$$

while for $z_0 > z_1 > z_2$ and $z_0 < z_1 + z_2$

$$\begin{aligned} P(y_0(z_0), y_1(z_1), y_2(z_2)) &= 1 - \frac{z_0^2 z_1 z_2}{4} \cdot (z_0^{-1} z_1^{-1} + z_0^{-1} z_2^{-1} - z_1^{-1} z_2^{-1})^2 \\ &= 1 - \frac{1}{4} (z_1^{-1} z_2 + z_1 z_2^{-1} + z_0^2 z_1^{-1} z_2^{-1} + 2 - 2z_0 z_1^{-1} - 2z_0 z_2^{-1}), \end{aligned}$$

which finishes the proof. \square

The previous lemma covers the case when $y_0 < y_1 < y_2$. We now leverage it to take care of the other cases as well.

Proof of Lemma 4.2. Let $y_i > 0$ and $z_i = e^{-y_i/2}$, $i = 0, 1, 2$. Lemma 4.3 gives the expression for $P(y_0(z_0), y_1(z_1), y_2(z_2))$ in the case $y_0 < y_1 < y_2$, or equivalently $z_0 > z_1 > z_2$, i.e. the first two lines in the claim of Lemma 4.2. To analyze the other cases we shall express $P(y_1, y_0, y_2)$ and $P(y_1, y_2, y_0)$ in terms of $P(y_0, y_1, y_2)$ and z_i . For this we note that we can view $P(y_0, y_1, y_2)$ as a 2-fold integral of the indicator function

$$h(x_0, x_1, x_2) := \mathbb{1}_{\{|x_0 - x_1| < e^{(y_0+y_1)/2}, |x_0 - x_2| < e^{(y_0+y_2)/2}, |x_1 - x_2| < e^{(y_1+y_2)/2}\}},$$

where x_0 was set to zero, without loss of generality, and the other two x_i are uniform random variables on $[-e^{(y_0+y_i)/2}, e^{(y_0+y_i)/2}]$. When we consider the probability $P(y_1, y_0, y_2)$, this is the 2-fold integral of $h(x_0, 0, x_2)$ so that

$$\begin{aligned} P(y_1, y_0, y_2) &= \frac{1}{2e^{(y_1+y_0)/2}} \cdot \frac{1}{2e^{(y_1+y_2)/2}} \iint_{\mathbb{R}} h(x_0, 0, x_2) dx_0 dx_2 \\ &= \frac{e^{y_0/2}}{e^{y_1/2}} \frac{1}{2e^{(y_0+y_1)/2}} \frac{1}{2e^{(y_0+y_2)/2}} \iint_{\mathbb{R}} h(0, x_1, x_2) dx_1 dx_2 \\ &= \frac{e^{y_0/2}}{e^{y_1/2}} P(y_0, y_1, y_2) = \frac{z_1}{z_0} P(y_0, y_1, y_2). \end{aligned}$$

Finally we note that $h(x_0, 0, x_2) = h(x_2, 0, x_0)$ from which we conclude that

$$P(y_0, y_1, y_2) = (z_0/z_1) P(y_1, y_0, y_2) = (z_0/z_1) P(y_1, y_2, y_0). \quad (44)$$

To complete the proof for the other cases we note that since $P(y_0, y_1, y_2)$ is symmetric in y_1 and y_2 , we can assume, without loss of generality, that $y_1 < y_2$. Then, there are two more orderings of y_0, y_1, y_2 , namely $y_1 < y_0 < y_2$ and $y_1 < y_2 < y_0$, which can be summarized as $y_1 < \min(y_0, y_2)$, or equivalently $z_1 > \max(z_0, z_2)$. For $y_1 < y_0 < y_2$ and $y_1 < y_2 < y_0$ we can apply Lemma 4.3 to obtain $P(y_1, y_0, y_2) = P(y_1, y_2, y_0)$ which happen to agree due to the symmetry in the last two arguments of the expression found in Lemma 4.3. The expression for $P(y_0, y_1, y_2)$ then follows from (44). \square

4.1.2 Integrating over y_1, y_2

Now that we have established the expression for $P(y_0, y_1, y_2)$ we can proceed to compute $P(y_0)$ by integrating over y_1, y_2 . We however start with the following observation.

Lemma 4.4. *The function $\alpha \mapsto P_\alpha(y_0)$ is continuous for all $\alpha > \frac{1}{2}$.*

Proof. This follows from the theorem of dominated convergence: Let $\alpha > \frac{1}{2}$ and $(\alpha_n)_{n \in \mathbb{N}}$ a sequence of real numbers converging to α , so we can assume $|\alpha_n - \alpha| < \epsilon := \frac{\alpha - 1/2}{2}$. This means that $-\epsilon < \alpha_n - \alpha < \epsilon$, i.e. $\frac{\alpha - 1/2}{2} < \alpha_n - 1/2 < \frac{3\alpha - 3/2}{2}$. Define

$$f_n(y_1, y_2) = P(y_0, y_1, y_2)(\alpha_n - 1/2)^2 e^{-(\alpha_n - 1/2)(y_1 + y_2)}.$$

As the function $x \mapsto x^2$ is increasing in x for $x > 0$ and the function $x \mapsto e^{-(y_1 + y_2)x}$ is decreasing in x and $P(y_0, y_1, y_2) \in [0, 1]$, it holds that

$$|f_n(y_1, y_2)| \leq \left(\frac{3\alpha - 3/2}{2} \right)^2 e^{-(y_1 + y_2) \frac{\alpha - 1/2}{2}}$$

which is integrable over $\mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}$ (with integral equalling $(6\alpha - 3)^2 / (2\alpha - 1)^2$). Application of the theorem of dominated convergence yields that $P_{\alpha_n}(y_0) \rightarrow P_\alpha(y_0)$ which gives the claim as the sequence $(\alpha_n)_n$ was arbitrary. \square

Due to this lemma we can first assume $\alpha \notin \{\frac{3}{4}, 1\}$, compute $P(y_0)$ and then obtain the values of $P(y_0)$ at the remaining two points by taking the corresponding limit in α . This strategy is executed below. It involves the computation of several integrals which are involved and will take up a few pages. The proof is structured using headers, to aid the reader.

Note that when writing $P(y_0)$ as an integral, see equation (43), by symmetry in the integration variables y_1 and y_2 , we can assume that $y_1 < y_2$ in which case either y_0 or y_1 is the smallest height. This gives half the value of $P(y_0)$ and hence

$$P(y_0) = 2(I_1(y_0) + I_2(y_0)),$$

where I_1 and I_2 are given by:

$$I_1(y_0) := \int_{0 < y_0 < y_1 < y_2} P(y_0, y_1, y_2) \cdot (\alpha - 1/2)^2 e^{-(\alpha-1/2)(y_1+y_2)} dy_2 dy_1$$

$$I_2(y_0) := \int_{0 < y_1 < y_0, y_2} P(y_0, y_1, y_2) \cdot (\alpha - 1/2)^2 e^{-(\alpha-1/2)(y_1+y_2)} dy_2 dy_1$$

We proceed with computing each of these two integrals, each of which is split in two parts. The final expressions of those four integrals can be found in (45), (50), (51) and (53).

Computing $I_1(y_0)$ Applying the change of variables $z_i := e^{-y_i/2}$, $i = 1, 2$, and Lemma 4.2 gives

$$\begin{aligned} I_1(y_0) &= 4(\alpha - 1/2)^2 \cdot \int_{z_0 > z_1 > z_2 > 0} P(y_0, y_1(z), y_2(z)) z_1^{2\alpha-2} z_2^{2\alpha-2} dz_2 dz_1 \\ &= 4(\alpha - 1/2)^2 \cdot \left(\int_{z_0 > z_1 > z_2 > 0} 1 \cdot z_1^{2\alpha-2} z_2^{2\alpha-2} dz_2 dz_1 \right. \\ &\quad \left. - \int_{\substack{z_0 > z_1 > z_2 > 0, \\ z_0 < z_1 + z_2}} G(z_0, z_1, z_2) \cdot z_1^{2\alpha-2} z_2^{2\alpha-2} dz_2 dz_1 \right) \\ &=: 4(\alpha - 1/2)^2 (I_{11}(y_0) - I_{12}(y_0)). \end{aligned}$$

The integral $I_{11}(y_0)$ is easily obtained:

$$\begin{aligned} I_{11}(y_0) &= \int_0^{z_0} \int_0^{z_1} z_1^{2\alpha-2} z_2^{2\alpha-2} dz_2 dz_1 = \int_0^{z_0} z_1^{2\alpha-2} \left[\frac{z_2^{2\alpha-1}}{2\alpha-1} \right]_0^{z_1} dz_1 \\ &= \frac{1}{2\alpha-1} \cdot \int_0^{z_0} z_1^{4\alpha-3} dz_1 = \frac{1}{2(2\alpha-1)^2} \cdot z_0^{4\alpha-2}. \end{aligned} \quad (45)$$

To deal with I_{12} we note that $G(z_0, z_1, z_2)$ is a linear combination of monomials of the form $z_0^a z_1^b z_2^c$ with $a, b, c \in \{-1, 0, 1, 2\}$ and $a + b + c = 0$. Let us consider the integral $J_{(a,b,c)}(z_0)$ defined by

$$J_{a,b,c}(z_0) := z_0^a \int_{\substack{z_0 > z_1 > z_2 > 0, \\ z_0 < z_1 + z_2}} z_1^{b+2\alpha-2} z_2^{c+2\alpha-2} dz_2 dz_1. \quad (46)$$

and note that

$$I_{1,2}(y_0) = \frac{1}{4} (J_{0,-1,1}(z_0) + J_{0,1,-1}(z_0) + J_{2,-1,-1}(z_0) + 2J_{0,0,0}(z_0) - 2J_{1,-1,0}(z_0) - 2J_{1,0,-1}(z_0)). \quad (47)$$

Next we compute $J_{a,b,c}(z_0)$.

$$\begin{aligned} J_{a,b,c}(z_0) &= z_0^a \int_{z_0/2}^{z_0} \int_{z_0-z_1}^{z_1} z_1^{b+2\alpha-2} z_2^{c+2\alpha-2} dz_2 dz_1 = z_0^a \int_{z_0/2}^{z_0} z_1^{b+2\alpha-2} \left[\frac{z_2^{c+2\alpha-1}}{c+2\alpha-1} \right]_{z_0-z_1}^{z_1} dz_1 \\ &= \frac{z_0^a}{c+2\alpha-1} \cdot \left(\int_{z_0/2}^{z_0} z_1^{b+c+4\alpha-3} dz_1 - \int_{z_0/2}^{z_0} z_1^{b+2\alpha-2} (z_0 - z_1)^{c+2\alpha-1} dz_1 \right) \\ &= \frac{z_0^{a+b+c+4\alpha-2} (1 - (1/2)^{b+c+4\alpha-2})}{(c+2\alpha-1)(b+c+4\alpha-2)} \\ &\quad - \frac{z_0^{a+b+c+4\alpha-3}}{c+2\alpha-1} \int_{z_0/2}^{z_0} (z_1/z_0)^{b+2\alpha-2} (1 - (z_1/z_0))^{c+2\alpha-1} dz_1 \end{aligned}$$

$$\begin{aligned}
&= \frac{z_0^{4\alpha-2}(1 - (1/2)^{b+c+4\alpha-2})}{(c+2\alpha-1)(b+c+4\alpha-2)} - \frac{z_0^{4\alpha-2}}{c+2\alpha-1} \int_{1/2}^1 u^{b+2\alpha-2}(1-u)^{c+2\alpha-1} du \\
&= \frac{z_0^{4\alpha-2}(1 - (1/2)^{b+c+4\alpha-2})}{(c+2\alpha-1)(b+c+4\alpha-2)} - \frac{z_0^{4\alpha-2}}{c+2\alpha-1} B^-(1/2; c+2\alpha, b+2\alpha-1),
\end{aligned}$$

where we have used the substitution $u := z_1/z_0$ giving $z_0 du = dz_1$ in the penultimate line and B^- denotes the (lower) incomplete beta function. Note that since $c \geq -1$, $-a \in \{0, -1, -2\}$ and by our assumption $\alpha \notin \{\frac{3}{4}, 1\}$, the denominators that occur during the integration are all non-zero.

Plugging this back into (47) gives

$$\begin{aligned}
I_{1,2}(y_0) &= \frac{z_0^{4\alpha-2}(1 - (1/2)^{4\alpha-2})}{32\alpha(\alpha-1/2)} - \frac{z_0^{4\alpha-2}}{8\alpha} B^-(1/2; 1+2\alpha, 2\alpha-2) \\
&+ \frac{z_0^{4\alpha-2}(1 - (1/2)^{4\alpha-2})}{32(\alpha-1)(\alpha-1/2)} - \frac{z_0^{4\alpha-2}}{4(2\alpha-2)} B^-(1/2; 2\alpha-1, 2\alpha) \\
&+ \frac{z_0^{4\alpha-2}(1 - (1/2)^{4\alpha-4})}{32(\alpha-1)^2} - \frac{z_0^{4\alpha-2}}{4(2\alpha-2)} B^-(1/2; -1+2\alpha, 2\alpha-2) \\
&+ \frac{z_0^{4\alpha-2}(1 - (1/2)^{4\alpha-2})}{16(\alpha-1/2)^2} - \frac{z_0^{4\alpha-2}}{2(2\alpha-1)} B^-(1/2; 2\alpha, 2\alpha-1) \\
&- \frac{z_0^{4\alpha-2}(1 - (1/2)^{4\alpha-3})}{16(\alpha-1/2)(\alpha-3/4)} + \frac{z_0^{4\alpha-2}}{2(2\alpha-1)} B^-(1/2; 2\alpha, 2\alpha-2) \\
&- \frac{z_0^{4\alpha-2}(1 - (1/2)^{4\alpha-3})}{16(\alpha-1)(\alpha-3/4)} + \frac{z_0^{4\alpha-2}}{2(2\alpha-2)} B^-(1/2; -1+2\alpha, 2\alpha-1) \\
&= \frac{\left(\frac{3}{64} - \frac{3}{16}2^{-4\alpha} + \alpha\left(-\frac{41}{128} + \frac{13}{16}2^{-4\alpha}\right) + \alpha^2\left(\frac{5}{8} - \frac{3}{4}2^{-4\alpha}\right) - \frac{15}{32}\alpha^3 + \frac{1}{8}\alpha^4\right) z_0^{4\alpha-2}}{4(\alpha-1/2)^2(\alpha-1)^2(\alpha-3/4)\alpha} \\
&+ \frac{z_0^{4\alpha-2}}{8(\alpha-1)\alpha(2\alpha-1)} (4(\alpha-1)\alpha(B^-(1/2; 2\alpha, 2\alpha-2) - B^-(1/2; 2\alpha, 2\alpha-1)) \\
&- (2\alpha-1)\alpha(B^-(1/2; 2\alpha-1, 2\alpha-2) + B^-(1/2; 2\alpha-1, 2\alpha) - 2B^-(1/2; 2\alpha-1, 2\alpha-1)) \\
&- (2\alpha-1)(\alpha-1)B^-(1/2; 1+2\alpha, 2\alpha-2)) \\
&= \frac{\left(\frac{3}{64} - \frac{3}{16}2^{-4\alpha} + \alpha\left(-\frac{41}{128} + \frac{13}{16}2^{-4\alpha}\right) + \alpha^2\left(\frac{5}{8} - \frac{3}{4}2^{-4\alpha}\right) - \frac{15}{32}\alpha^3 + \frac{1}{8}\alpha^4\right) z_0^{4\alpha-2}}{4(\alpha-1/2)^2(\alpha-1)^2(\alpha-3/4)\alpha} \\
&+ \frac{z_0^{4\alpha-2}}{8(\alpha-1)\alpha(2\alpha-1)} (4(\alpha-1)\alpha B^-(1/2; 2\alpha+1, 2\alpha-2) \\
&- (2\alpha-1)\alpha B^-(1/2; 2\alpha+1, 2\alpha-2) \\
&- (2\alpha-1)(\alpha-1)B^-(1/2; 2\alpha+1, 2\alpha-2)).
\end{aligned}$$

For the last step we use the identities

$$B^-(z; a, b) - B^-(z; a, b+1) = B^-(z; a+1, b), \quad (48)$$

$$B^-(z; a, b) + B^-(z; a, b+2) - 2B^-(z; a, b+1) = B^-(z; a+2, b). \quad (49)$$

to obtain

$$\begin{aligned}
I_{1,2}(y_0) &= \frac{\left(\frac{3}{64} - \frac{3}{16}2^{-4\alpha} + \alpha\left(-\frac{41}{128} + \frac{13}{16}2^{-4\alpha}\right) + \alpha^2\left(\frac{5}{8} - \frac{3}{4}2^{-4\alpha}\right) - \frac{15}{32}\alpha^3 + \frac{1}{8}\alpha^4\right) z_0^{4\alpha-2}}{4(\alpha-1/2)^2(\alpha-1)^2(\alpha-3/4)\alpha} \\
&- \frac{z_0^{4\alpha-2} B^-(1/2; 2\alpha+1, 2\alpha-2)}{8(\alpha-1)\alpha(2\alpha-1)}
\end{aligned} \quad (50)$$

Computing $I_2(y_0)$ We will follow a similar strategy as for $I_1(y_0)$. First, using the change of variables $z_i := e^{-y_i/2}$, $i = 1, 2$, we get

$$\begin{aligned} I_2(y_0) &= 4(\alpha - 1/2)^2 \cdot \int_{1 > z_1 > z_2, z_0 > 0} P(y_0, y_1(z_1), y_2(z_2)) z_1^{2\alpha-2} z_2^{2\alpha-2} dz_2 dz_1 \\ &= 4(\alpha - 1/2)^2 \cdot \left(\int_{1 > z_1 > z_0, z_2 > 0} z_0 z_1^{2\alpha-3} z_2^{2\alpha-2} dz_2 dz_1 \right. \\ &\quad \left. - \int_{\substack{1 > z_1 > z_0, z_2 > 0 \\ z_1 < z_0 + z_2}} G(z_1, z_0, z_2) z_0 z_1^{2\alpha-3} z_2^{2\alpha-2} dz_2 dz_1 \right) \\ &=: 4(\alpha - 1/2)^2 (I_{21}(y_0) - I_{22}(y_0)). \end{aligned}$$

We start with the easy integral:

$$\begin{aligned} I_{21}(y_0) &= z_0 \int_{1 > z_1 > \max(z_2, z_0); z_0, z_2 > 0} z_1^{2\alpha-3} z_2^{2\alpha-2} dz_2 dz_1 = z_0 \int_{z_0}^1 \int_0^{z_1} z_1^{2\alpha-3} z_2^{2\alpha-2} dz_2 dz_1 \\ &= z_0 \int_{z_0}^1 \left[\frac{z_2^{2\alpha-1}}{2\alpha-1} \right]_0^{z_1} z_1^{2\alpha-3} dz_1 = \frac{z_0}{2\alpha-1} \int_{z_0}^1 z_1^{4\alpha-4} dz_1 = \frac{z_0 - z_0^{4\alpha-2}}{(4\alpha-3)(2\alpha-1)}. \end{aligned} \quad (51)$$

We note that the denominators above are non-zero as $\alpha > \frac{1}{2}$ and $\alpha \neq \frac{3}{4}$.

To deal with $I_{22}(y_0)$ we consider the function

$$J'_{a,b,c}(z_0) := z_0^a \int_{\substack{1 > z_1 > \max(z_0, z_2); z_0, z_2 > 0 \\ z_1 < z_0 + z_2}} z_1^{b+2\alpha-2} z_2^{c+2\alpha-2} dz_2 dz_1$$

and write

$$\begin{aligned} I_{2,2}(y_0) &= \frac{1}{4} (J'_{0,-1,1}(z_0) + J'_{2,-1,-1}(z_0) + J'_{0,1,-1}(z_0)) \\ &\quad + \frac{1}{2} (J'_{1,-1,0}(z_0) - J'_{0,0,0}(z_0) - J'_{1,0,-1}(z_0)). \end{aligned} \quad (52)$$

We now compute $J'_{a,b,c}(z_0)$

$$\begin{aligned} J'_{a,b,c}(z_0) &= z_0^a \int_{z_0}^1 \int_{z_1-z_0}^{z_1} z_1^{b+2\alpha-2} z_2^{c+2\alpha-2} dz_2 dz_1 \\ &= z_0^a \int_{z_0}^1 \frac{1}{c+2\alpha-1} z_1^{b+2\alpha-2} (z_1^{c+2\alpha-1} - (z_1 - z_0)^{c+2\alpha-1}) dz_1 \\ &= z_0^a \int_{z_0}^1 \frac{1}{c+2\alpha-1} z_1^{b+c+4\alpha-3} dz_1 - z_0^a \int_{z_0}^1 \frac{1}{c+2\alpha-1} z_1^{b+2\alpha-2} (z_1 - z_0)^{c+2\alpha-1} dz_1 \\ &= z_0^a \frac{1}{(c+2\alpha-1)(b+c+4\alpha-2)} (1 - z_0^{b+c+4\alpha-2}) \\ &\quad - \frac{z_0^a}{c+2\alpha-1} z_0^{b+c+4\alpha-2} B^-(1 - z_0; c+2\alpha, -b-c-4\alpha+2) \\ &= \frac{z_0^a - z_0^{4\alpha-2}}{(c+2\alpha-1)(b+c+4\alpha-2)} - \frac{z_0^{4\alpha-2} B^-(1 - z_0; c+2\alpha, -b-c-4\alpha+2)}{c+2\alpha-1}. \end{aligned}$$

Here we used that for $x \in \mathbb{R}, y > -1$ (note that as $c \geq -1$, it holds that $c+2\alpha-1 > -1$):

$$\begin{aligned} \int_{z_0}^1 z_1^x (z_1 - z_0)^y dz_1 &= \int_0^{1-z_0} (s + z_0)^x s^y ds \\ &= z_0^{x+y} \int_0^{1-z_0} ((s/z_0) + 1)^x (s/z_0)^y ds \end{aligned}$$

$$\begin{aligned}
&= z_0^{x+y+1} \int_0^{1/z_0-1} (t+1)^x t^y dt \\
&= z_0^{x+y+1} \int_0^{1-z_0} u^y (1-u)^{-(x+y+2)} du \\
&= z_0^{x+y+1} B^-(1-z_0; y+1, -x-y-1).
\end{aligned}$$

As $c \geq -1$ and $-a \in \{0, -1, -2\}$ and by our assumption $\alpha \notin \{\frac{3}{4}\}$, the denominators that occur during the computations above are non-zero.

Plugging the expression for $J'_{a,b,c}(z_0)$ back into (52) we get,

$$\begin{aligned}
I_{2,2}(y_0) &= \frac{1 - z_0^{4\alpha-2}}{32\alpha(\alpha-1/2)} - \frac{z_0^{4\alpha-2} B^-(1-z_0; 1+2\alpha, -4\alpha+2)}{8\alpha} \\
&+ \frac{z_0^2 - z_0^{4\alpha-2}}{32(\alpha-1)^2} - \frac{z_0^{4\alpha-2} B^-(1-z_0; -1+2\alpha, -4\alpha+4)}{8(\alpha-1)} \\
&+ \frac{1 - z_0^{4\alpha-2}}{32(\alpha-1)(\alpha-1/2)} - \frac{z_0^{4\alpha-2} B^-(1-z_0; -1+2\alpha, -4\alpha+2)}{8(\alpha-1)} \\
&+ \frac{z_0 - z_0^{4\alpha-2}}{16(\alpha-1/2)(\alpha-3/4)} - \frac{z_0^{4\alpha-2} B^-(1-z_0; 2\alpha, -4\alpha+3)}{4(\alpha-1/2)} \\
&- \frac{1 - z_0^{4\alpha-2}}{16(\alpha-1/2)^2} + \frac{z_0^{4\alpha-2} B^-(1-z_0; 2\alpha, -4\alpha+2)}{4(\alpha-1/2)} \\
&- \frac{z_0 - z_0^{4\alpha-2}}{16(\alpha-1)(\alpha-3/4)} + \frac{z_0^{4\alpha-2} B^-(1-z_0; -1+2\alpha, -4\alpha+3)}{4(\alpha-1)}.
\end{aligned}$$

Using some algebra and the identities (48) and (49) this can be reduced to

$$\begin{aligned}
I_{2,2}(y_0) &= \frac{1}{64\alpha(\alpha-1/2)^2(\alpha-1)} - \frac{(1-z_0)^{2\alpha}}{64\alpha(\alpha-1/2)^2(\alpha-1)} - \frac{z_0}{8(\alpha-1/2)(\alpha-1)(4\alpha-3)} \\
&+ \frac{z_0^2}{32(\alpha-1)^2} + \frac{(-6+25\alpha-48\alpha^2+44\alpha^3-16\alpha^4)z_0^{4\alpha-2}}{512\alpha(\alpha-1/2)^2(\alpha-1)^2(\alpha-3/4)} \\
&+ \frac{z_0^{4\alpha-2} B^-(1-z_0; 2\alpha, 3-4\alpha)}{32(\alpha-1)(\alpha-1/2)^2}.
\end{aligned} \tag{53}$$

Combining the results for $I_1(y_0)$ and $I_2(y_0)$ Combining the results for $I_{11}(y_0)$, $I_{12}(y_0)$, $I_{21}(y_0)$ and $I_{22}(y_0)$ we get, after some algebra, an explicit expression for $P(y_0)$ as a linear combination of terms of the form z_0^u , $(1-z_0)^u$ and $z_0^u B^-(1-z_0; a, b)$:

$$\begin{aligned}
P(y_0) &= 2(I_1 + I_2) = 8(\alpha-1/2)^2(I_{1,1} - I_{1,2} + I_{2,1} - I_{2,2}) \\
&= 8(\alpha-1/2)^2 \left(\frac{1}{2(2\alpha-1)^2} z_0^{4\alpha-2} \right. \\
&\quad - \frac{(\frac{3}{64} - \frac{3}{16} 2^{-4\alpha} + \alpha(-\frac{41}{128} + \frac{13}{16} 2^{-4\alpha}) + \alpha^2(\frac{5}{8} - \frac{3}{4} 2^{-4\alpha}) - \frac{15}{32} \alpha^3 + \frac{1}{8} \alpha^4) z_0^{4\alpha-2}}{4(\alpha-1/2)^2(\alpha-1)^2(\alpha-3/4)\alpha} \\
&\quad + \frac{z_0^{4\alpha-2} B^-(1/2; 2\alpha+1, 2\alpha-2)}{8(\alpha-1)\alpha(2\alpha-1)} + \frac{z_0 - z_0^{4\alpha-2}}{(4\alpha-3)(2\alpha-1)} \\
&\quad - \frac{1}{64\alpha(\alpha-1/2)^2(\alpha-1)} + \frac{(1-z_0)^{2\alpha}}{64\alpha(\alpha-1/2)^2(\alpha-1)} + \frac{z_0}{8(\alpha-1/2)(\alpha-1)(4\alpha-3)} \\
&\quad - \frac{z_0^2}{32(\alpha-1)^2} - \frac{(-6+25\alpha-48\alpha^2+44\alpha^3-16\alpha^4)z_0^{4\alpha-2}}{512\alpha(\alpha-1/2)^2(\alpha-1)^2(\alpha-3/4)} \\
&\quad \left. - \frac{z_0^{4\alpha-2} B^-(1-z_0; 2\alpha, 3-4\alpha)}{32(\alpha-1)(\alpha-1/2)^2} \right)
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8(\alpha-1)\alpha} + \frac{(\alpha-1/2)z_0}{\alpha-1} - \frac{(\alpha-1/2)^2 z_0^2}{4(\alpha-1)^2} \\
&\quad + z_0^{-2+4\alpha} \left(\frac{2^{-4\alpha-1}(3\alpha-1)}{\alpha(\alpha-1)^2} + \frac{(\alpha-1/2)B^-(1/2; 1+2\alpha, -2+2\alpha)}{2(\alpha-1)\alpha} \right) \\
&\quad + \frac{(1-z_0)^{2\alpha}}{8(\alpha-1)\alpha} - \frac{z_0^{4\alpha-2}B^-(1-z_0; 2\alpha, 3-4\alpha)}{4(\alpha-1)}
\end{aligned}$$

Observe that the above expression only contains terms of the form $\alpha-1$ in the denominator. The only expression of the form $\alpha-3/4$ is in the lower incomplete beta-function $B^-(1-z_0; 2\alpha, 3-4\alpha)$ which appears twice in the expression for $P(y_0)$.

The case of $\alpha = 3/4$

Note that the factor $\alpha - \frac{3}{4}$ does not occur in any denominator of the previously obtained expression. For the lower incomplete beta function, the last argument $3-4\alpha$ is zero for $\alpha = \frac{3}{4}$, however as $z_0 < 1$ the integration domain of the lower incomplete beta function does not touch the singularity at $t = 1$ (note $B^-(1-z_0; 2\alpha, 3-4\alpha) = \int_0^{1-z_0} t^{2\alpha-1}(1-t)^{2-4\alpha} dt$). Therefore, the previous expression holds for this case as well.

4.1.3 Computing γ and $\gamma(k)$

Now that we have an expression for $P(y_0)$ we can compute $\gamma, \gamma(k)$ by integrating over y_0 and prove that they equal the expressions given in, respectively, Theorem 1.2 and Theorem 1.3.

We define

$$I^{(k)} := \int_0^\infty P(y) \alpha e^{-\alpha y} \rho(y, k) dy = \int_0^\infty P(y) \alpha e^{-\alpha y} \frac{(\xi e^{y/2})^k}{k!} e^{-\xi e^{y/2}} dy$$

and

$$J := \int_0^\infty P(y) \alpha e^{-\alpha y} dy.$$

Then, recalling (42) and (41), we have

$$\gamma = J - I^{(1)} - I^{(2)} \quad \text{and} \quad \gamma(k) = \frac{I^{(k)}}{p_k}.$$

We will thus compute J and $I^{(k)}$. It will be helpful to change coordinates to $z := e^{-y/2}$. This yields

$$J = 2\alpha \int_0^1 P(y) z^{2\alpha-1} dz,$$

and

$$I^{(k)} = \frac{2\alpha \xi^k}{k!} \cdot \int_0^1 P(y(z)) \cdot z^{2\alpha-(k+1)} e^{-\xi z^{-1}} dz.$$

We shall be assuming $\alpha \neq 1$. We observe from Lemma 4.1 that for $\alpha \neq 1$, $P(y(z))$ is in fact a linear combination of terms of the form z^u , $(1-z)^u$ and $z^u B^-(1-z; v, w)$.

To compute J we observe that, by integration by parts,

$$\begin{aligned}
\int_0^1 z^{u+2\alpha-1} B^-(1-z; v, w) dz &= \left[\frac{z^{u+2\alpha}}{u+2\alpha} B^-(1-z; v, w) \right]_0^1 + \frac{1}{u+2\alpha} \int_0^1 z^{u+2\alpha+w-1} (1-z)^{v-1} dz \\
&= \frac{1}{u+2\alpha} B(u+w+2\alpha, v)
\end{aligned}$$

where we used that $\frac{\partial}{\partial z} B^-(1-z; v, w) = -z^{w-1}(1-z)^{v-1}$. This takes care of the two integrands involving the beta function in $P(y)$. The other integrals are easily computed and yield the following expression for J (note that it only depends on α but not on ν)

$$J = \frac{2 + 4\alpha + 13\alpha^2 - 34\alpha^3 - 12\alpha^4 + 24\alpha^5}{16(\alpha - 1)^2\alpha(\alpha + 1)(2\alpha + 1)} + \frac{2^{-1-4\alpha}}{(\alpha - 1)^2} + \frac{(\alpha - 1/2)(B(2\alpha, 2\alpha + 1) + B^-(1/2; 1 + 2\alpha, -2 + 2\alpha))}{2(\alpha - 1)(3\alpha - 1)}.$$

We proceed to work out $I^{(k)}$. For this we will compute the integrals involving terms in $P(y(z))$ of the form z^u , $(1-z)^u$ and $B(1-z, v, w)$ separately. We first point out that for any $0 \leq a < b \leq 1$

$$\begin{aligned} \int_a^b z^{u+2\alpha-(k+1)} e^{-\xi z^{-1}} dz &= \xi^{u+2\alpha-k} \int_{\xi/b}^{\xi/a} t^{k-1-2\alpha-u} e^{-t} dt \\ &= \xi^{u+2\alpha-k} (\Gamma^+(k-2\alpha-u, \xi/b) - \Gamma^+(k-2\alpha-u, \xi/a)). \end{aligned}$$

In particular

$$\int_0^1 z^{u+2\alpha-k-1} e^{-\xi z^{-1}} dz = \xi^{u+2\alpha-k} \Gamma^+(k-2\alpha-u, \xi). \quad (54)$$

where Γ^+ denotes the (upper) incomplete gamma function, and we have used the substitution $t = \xi/z$ which gives $dz = -\xi t^{-2} dt$. (And of course it is understood that $\xi/0 = \infty$). This takes care of the integrals of all terms in $P(y(z))$ of the form z^u .

Next we will consider the integrals over the terms in $P(y(z))$ of the form $(1-z)^u$. For this we need the hypergeometric U-function (also called Tricomi's confluent hypergeometric function), which has the integral representation

$$U(a, b, z) = \frac{1}{\Gamma(a)} \int_0^\infty e^{-zt} t^{a-1} (1+t)^{b-a-1} dt.$$

which holds for $a, b, z \in \mathbb{C}$, $b \notin \mathbb{Z}_{\leq 0}$, $\operatorname{Re}(a), \operatorname{Re}(z) > 0$, see [15, p.255]. Applying the change of variables $t = \frac{1-s}{s}$ (i.e. $dt = -s^{-2} ds$ and $s = \frac{1}{t+1}$) yields

$$U(a, b, z) = \frac{e^z}{\Gamma(a)} \int_0^1 s^{-b} (1-s)^{a-1} e^{-z/s} ds$$

Setting $a = 2\alpha + 1 > 0$, $b = -2\alpha + k + 1$, $z = \xi > 0$, then gives

$$\int_0^1 z_0^{2\alpha-k-1} e^{-\xi/z_0} (1-z_0)^{2\alpha} dz_0 = \Gamma(2\alpha+1) e^{-\xi} U(2\alpha+1, 1+k-2\alpha, \xi). \quad (55)$$

Finally we need to deal with the terms in $P(y(z))$ that involve the incomplete beta function. Let $a, c \in \mathbb{R}$, $\xi, b > 0$ positive real numbers. Using the integral definition of the incomplete beta function, the change of variables $s = 1-t$ gives:

$$\begin{aligned} \int_0^1 z^a e^{-\xi/z} B^-(1-z; b, c) dz &= \int_0^1 z^a e^{-\xi/z} \int_0^{1-z} t^{b-1} (1-t)^{c-1} dt dz \\ &= \int_0^1 z^a e^{-\xi/z} \int_z^1 s^{c-1} (1-s)^{b-1} ds dz. \end{aligned}$$

Then changing the order of integration and using the substitution $u = \xi/z$ and recognizing the upper incomplete gamma function yields

$$\int_0^1 z^a e^{-\xi/z} \int_z^1 s^{c-1} (1-s)^{b-1} ds dz$$

$$\begin{aligned}
&= \int_0^1 \int_0^s z^a e^{-\xi/z} dz s^{c-1} (1-s)^{b-1} ds \\
&= \int_0^1 \int_{\xi/s}^\infty \xi^{a+1} u^{-a-2} e^{-u} du s^{c-1} (1-s)^{b-1} ds \\
&= \xi^{a+1} \int_0^1 \Gamma^+(-a-1, \xi/s) s^{c-1} (1-s)^{b-1} ds.
\end{aligned} \tag{56}$$

To compute this last integral we make use of the fact that the incomplete Γ -function has a representation in terms of Meijer's G -function (see Lemma A.1 in Appendix A)

$$\Gamma^+(-a-1, \xi/s) = G_{1,2}^{2,0} \left(\frac{\xi}{s} \middle| \begin{matrix} 1 \\ -a-1, 0 \end{matrix} \right),$$

which holds for any $a \in \mathbb{R}$ and $s > 0$ (that for a fixed second argument, the upper incomplete gamma function is entire in the first argument, see [20, pp. 899, 1032ff.]). We can now evaluate the integral in (56) using several identities for Meijer's G -function. First, inserting the expression for the incomplete Gamma-function into (56) gives

$$\xi^{a+1} \int_0^1 s^{c-1} (1-s)^{b-1} G_{1,2}^{2,0} \left(\frac{\xi}{s} \middle| \begin{matrix} 1 \\ -a-1, 0 \end{matrix} \right) ds.$$

Next we apply the inversion identity for Meijer's G -function (see [15, p. 209, 5.3.1.(9)]) to get

$$\xi^{a+1} \int_0^1 s^{c-1} (1-s)^{b-1} G_{2,1}^{0,2} \left(\frac{s}{\xi} \middle| \begin{matrix} 2+a, 1 \\ 0 \end{matrix} \right) ds.$$

This expression is actually the Euler transform of Meijer's G -function (see [15, p. 214, 5.5.2.(5)]) and (as the conditions $2+1 < 2(0+2)$ and $|\arg(\xi^{-1})| < \frac{\pi}{2}$ (as $\xi > 0$) and $1-c-b < 1-c$ (as $b > 0$) are satisfied) it equals

$$\xi^{a+1} \Gamma(b) G_{3,2}^{0,3} \left(\xi^{-1} \middle| \begin{matrix} 1-c, 2+a, 1 \\ 0, 1-c-b \end{matrix} \right).$$

Using again the inversion identity for Meijer's G -function we now get

$$\xi^{a+1} \Gamma(b) G_{2,3}^{3,0} \left(\xi \middle| \begin{matrix} 1, b+c \\ c, -1-a, 0 \end{matrix} \right).$$

Finally, plugging in $a = 6\alpha - k - 3$, $b = 2\alpha$, $c = 3 - 4\alpha$ we obtain

$$\int_0^1 z^a e^{-\xi/z} B^-(1-z; b, c) dz = \xi^{6\alpha-k-2} \Gamma(2\alpha) G_{2,3}^{3,0} \left(\xi \middle| \begin{matrix} 1, 3-2\alpha \\ 3-4\alpha, -6\alpha+k+2, 0 \end{matrix} \right). \tag{57}$$

Using equation (54), (55) and (57) we get

$$\begin{aligned}
I^{(k)} &= \frac{\xi^{2\alpha}}{4k!(\alpha-1)} \left(-\Gamma^+(k-2\alpha, \xi) - 2 \frac{\alpha(\alpha-1/2)^2 \xi^2 \Gamma^+(k-2\alpha-2, \xi)}{(\alpha-1)} \right. \\
&\quad + 8\alpha(\alpha-1/2) \xi \Gamma^+(k-2\alpha-1, \xi) \\
&\quad + 4\xi^{4\alpha-2} \Gamma^+(k-6\alpha+2, \xi) \left(\frac{2^{-4\alpha}(3\alpha-1)}{(\alpha-1)} + (\alpha-1/2) B^-(1/2; 1+2\alpha, -2+2\alpha) \right) \\
&\quad + \xi^{k-2\alpha} \Gamma(2\alpha+1) e^{-\xi} U(2\alpha+1, 1+k-2\alpha, \xi) \\
&\quad \left. - \xi^{4\alpha-2} \Gamma(2\alpha+1) G_{2,3}^{3,0} \left(\xi \middle| \begin{matrix} 1, 3-2\alpha \\ 3-4\alpha, -6\alpha+k+2, 0 \end{matrix} \right) \right)
\end{aligned}$$

With the expressions for J and $I^{(k)}$ and using $\Gamma^*(q, z) = \Gamma^+(q+1, z) + \Gamma^+(q, z)$ we now obtain, after some algebra, the expression for γ

$$\begin{aligned}
\gamma &= J - I^{(0)} - I^{(1)} \\
&= \frac{2 + 4\alpha + 13\alpha^2 - 34\alpha^3 - 12\alpha^4 + 24\alpha^5}{16(\alpha - 1)^2\alpha(\alpha + 1)(2\alpha + 1)} + \frac{2^{-1-4\alpha}}{(\alpha - 1)^2} \\
&\quad + \frac{(\alpha - 1/2)(B(2\alpha, 2\alpha + 1) + B^-(1/2; 1 + 2\alpha, -2 + 2\alpha))}{2(\alpha - 1)(3\alpha - 1)} \\
&\quad - \frac{\xi^{2\alpha}}{4(\alpha - 1)} \left(-\Gamma^+(-2\alpha, \xi) - 2 \frac{\alpha(\alpha - 1/2)^2 \xi^2 \Gamma^+(-2\alpha - 2, \xi)}{(\alpha - 1)} \right. \\
&\quad \left. + 8\alpha(\alpha - 1/2) \xi \Gamma^+(-2\alpha - 1, \xi) \right. \\
&\quad \left. + 4\xi^{4\alpha-2} \Gamma^+(-6\alpha + 2, \xi) \left(\frac{2^{-4\alpha}(3\alpha - 1)}{(\alpha - 1)} + (\alpha - 1/2) B^-(1/2; 1 + 2\alpha, -2 + 2\alpha) \right) \right. \\
&\quad \left. + \xi^{-2\alpha} \Gamma(2\alpha + 1) e^{-\xi} U(2\alpha + 1, 1 - 2\alpha, \xi) \right. \\
&\quad \left. - \xi^{4\alpha-2} \Gamma(2\alpha + 1) G_{2,3}^{3,0} \left(\xi \left| \begin{matrix} 1, 3 - 2\alpha \\ 3 - 4\alpha, -6\alpha + 2, 0 \end{matrix} \right. \right) \right) \\
&\quad - \frac{\xi^{2\alpha}}{4(\alpha - 1)} \left(-\Gamma^+(1 - 2\alpha, \xi) - 2 \frac{\alpha(\alpha - 1/2)^2 \xi^2 \Gamma^+(-2\alpha - 1, \xi)}{(\alpha - 1)} \right. \\
&\quad \left. + 8\alpha(\alpha - 1/2) \xi \Gamma^+(1 - 2\alpha - 1, \xi) \right. \\
&\quad \left. + 4\xi^{4\alpha-2} \Gamma^+(1 - 6\alpha + 2, \xi) \left(\frac{2^{-4\alpha}(3\alpha - 1)}{(\alpha - 1)} + (\alpha - 1/2) B^-(1/2; 1 + 2\alpha, -2 + 2\alpha) \right) \right. \\
&\quad \left. + \xi^{1-2\alpha} \Gamma(2\alpha + 1) e^{-\xi} U(2\alpha + 1, 2 - 2\alpha, \xi) \right. \\
&\quad \left. - \xi^{4\alpha-2} \Gamma(2\alpha + 1) G_{2,3}^{3,0} \left(\xi \left| \begin{matrix} 1, 3 - 2\alpha \\ 3 - 4\alpha, -6\alpha + 3, 0 \end{matrix} \right. \right) \right) \\
&= \frac{2 + 4\alpha + 13\alpha^2 - 34\alpha^3 - 12\alpha^4 + 24\alpha^5}{16(\alpha - 1)^2\alpha(\alpha + 1)(2\alpha + 1)} + \frac{2^{-1-4\alpha}}{(\alpha - 1)^2} \\
&\quad + \frac{(\alpha - 1/2)(B(2\alpha, 2\alpha + 1) + B^-(1/2; 1 + 2\alpha, -2 + 2\alpha))}{2(\alpha - 1)(3\alpha - 1)} \\
&\quad + \frac{\xi^{2\alpha} \Gamma^*(-2\alpha, \xi)}{4(\alpha - 1)} + \frac{\xi^{2\alpha+2} \alpha(\alpha - 1/2)^2 \Gamma^*(-2\alpha - 2, \xi)}{2(\alpha - 1)^2} \\
&\quad - \frac{\xi^{2\alpha+1} \alpha(2\alpha - 1) \Gamma^*(-2\alpha - 1, \xi)}{(\alpha - 1)} - \frac{\xi^{6\alpha-2} 2^{-4\alpha} (3\alpha - 1) \Gamma^*(-6\alpha + 2, \xi)}{(\alpha - 1)^2} \\
&\quad - \frac{\xi^{6\alpha-2} (\alpha - 1/2) B^-(1/2; 1 + 2\alpha, -2 + 2\alpha) \Gamma^*(-6\alpha + 2, \xi)}{(\alpha - 1)} \\
&\quad - \frac{e^{-\xi} \Gamma(2\alpha + 1) (U(2\alpha + 1, 1 - 2\alpha, \xi) + U(2\alpha + 1, 2 - 2\alpha, \xi))}{4(\alpha - 1)} \\
&\quad + \frac{\xi^{6\alpha-2} \Gamma(2\alpha + 1) \left(G_{2,3}^{3,0} \left(\xi \left| \begin{matrix} 1, 3 - 2\alpha \\ 3 - 4\alpha, -6\alpha + 2, 0 \end{matrix} \right. \right) + G_{2,3}^{3,0} \left(\xi \left| \begin{matrix} 1, 3 - 2\alpha \\ 3 - 4\alpha, -6\alpha + 3, 0 \end{matrix} \right. \right) \right)}{4(\alpha - 1)}.
\end{aligned}$$

which is the expression in Theorem 1.2.

Similarly, we get

$$\begin{aligned}
\gamma(k) &= \frac{I^{(k)}}{p_k} \\
&= \frac{1}{8\alpha(\alpha - 1) \Gamma^+(k - 2\alpha, \xi)} \left(-\Gamma^+(k - 2\alpha, \xi) - 2 \frac{\alpha(\alpha - 1/2)^2 \xi^2 \Gamma^+(k - 2\alpha - 2, \xi)}{(\alpha - 1)} \right)
\end{aligned}$$

$$\begin{aligned}
& +8\alpha(\alpha - 1/2)\xi\Gamma^+(k - 2\alpha - 1, \xi) \\
& +4\xi^{4\alpha-2}\Gamma^+(k - 6\alpha + 2, \xi) \left(\frac{2^{-4\alpha}(3\alpha - 1)}{(\alpha - 1)} + (\alpha - 1/2)B^-(1/2; 1 + 2\alpha, -2 + 2\alpha) \right) \\
& +\xi^{k-2\alpha}\Gamma(2\alpha + 1)e^{-\xi}U(2\alpha + 1, 1 + k - 2\alpha, \xi) \\
& -\xi^{4\alpha-2}\Gamma(2\alpha + 1)G_{2,3}^{3,0} \left(\xi \left| \begin{matrix} 1, 3 - 2\alpha \\ 3 - 4\alpha, -6\alpha + k + 2, 0 \end{matrix} \right. \right),
\end{aligned}$$

which equals the expression in Theorem 1.3.

4.1.4 Explicit expressions for $\gamma, \gamma(k)$ when $\alpha = 1$.

Although we've already established that $\gamma, \gamma(k)$ can be obtained at $\alpha = 1$ by taking the $\alpha \rightarrow 1$ limit of the expression obtained for $\alpha = 1$, it is still helpful to derive an alternative, more explicit expression. This is what we will do in the current section. We will prove

Proposition 4.5. *If $\alpha = 1$ then*

$$\begin{aligned}
\gamma &= \frac{575 - 12\pi^2}{576} + \frac{\eta^4(7 + \pi^2)\Gamma^*(-4, \eta)}{4} \\
& - \frac{1}{2} \int_0^1 (1 - 4z + 3z^3) \log(1 - z)(z + \eta)e^{-\eta/z} dz \\
& - \int_0^1 \text{Li}_2(z)(z^3 + \eta z^2)e^{-\eta/z} dz,
\end{aligned}$$

and

$$\begin{aligned}
\gamma(k) &= \frac{9\eta^3}{2k!}\Gamma^+(k - 3, \eta) - \frac{\xi^4}{k!}\frac{7 + \pi^2}{4}\Gamma^+(k - 4, \eta) \\
& + \frac{\eta^k}{2k!} \int_0^1 (1 - 4z + 3z^2) \ln(1 - z)z^{1-k}e^{-\eta/z} dz \\
& + \frac{\eta^k}{k!} \int_0^1 z^{3-k} \text{Li}_2(z)e^{-\eta/z} dz,
\end{aligned}$$

with $\eta = 4\nu/\pi$ and $\text{Li}_2(z) = \sum_{t=1}^{\infty} z^t/t^2$, the dilogarithm function.

Naturally, the proof proceeds by proving the analogue of Lemma 4.1:

Lemma 4.6. *If $\alpha = 1$, then*

$$P(y) = \frac{9}{4}e^{-\frac{1}{2}y} + \frac{1 - 4e^{-\frac{1}{2}y} + 3e^{-y}}{4} \ln(1 - e^{-\frac{1}{2}y}) - \frac{7 + \pi^2}{8}e^{-y} + \frac{1}{2}e^{-y} \text{Li}_2(e^{-y})$$

where $\text{Li}_2(z) = \int_0^z \frac{\ln(1-t)}{t} dt$ is the dipolylogarithm function.

Proof. **Tobias: Warning!** this has not been properly proofread by me, or anyone else. We want to compute the limit $\lim_{\alpha \rightarrow 1} P_\alpha(y_0(z_0))$. For $\alpha \neq 1$, we label the terms as follows:

$$\begin{aligned}
& P_\alpha(y_0(z_0)) \\
&= \frac{1}{\alpha - 1} \left(s_1(\alpha, z_0) + s_2(\alpha, z_0) + \frac{1}{\alpha - 1} (s_3(\alpha, z_0) + s_4(\alpha, z_0)) + s_5(\alpha, z_0) + s_6(\alpha, z_0) + s_7(\alpha, z_0) \right)
\end{aligned}$$

where

$$\begin{aligned}
s_1(\alpha, z_0) &= -\frac{1}{8\alpha} \\
s_2(\alpha, z_0) &= (\alpha - 1/2)z_0
\end{aligned}$$

$$\begin{aligned}
s_3(\alpha, z_0) &= -\frac{(\alpha - 1/2)^2 z_0^2}{4} \\
s_4(\alpha, z_0) &= z_0^{-2+4\alpha} \frac{2^{-4\alpha-1}(3\alpha - 1)}{\alpha} \\
s_5(\alpha, z_0) &= z_0^{-2+4\alpha} \frac{(\alpha - 1/2)B^-(1/2; 1 + 2\alpha, -2 + 2\alpha)}{2\alpha} \\
s_6(\alpha, z_0) &= \frac{(1 - z_0)^{2\alpha}}{8\alpha} \\
s_7(\alpha, z_0) &= -\frac{z_0^{4\alpha-2}B^-(1 - z_0; 2\alpha, 3 - 4\alpha)}{4}
\end{aligned}$$

Now, we consider the functions $s_i(\alpha) = s_i(\alpha, z_0)$ as functions of α only and compute their Taylor expansion at $\alpha = 1$, for $i \in \{1, 2, 5, 6, 7\}$ up to linear and for $i \in \{3, 4\}$ up to quadratic order, i.e. we write $s_i(\alpha) = s_i(1) + s'_i(1)(\alpha - 1) + o(\alpha - 1)$ for $i \in \{1, 2, 5, 6, 7\}$ and $s_i(\alpha) = s_i(1) + s'_i(1)(\alpha - 1) + \frac{s''_i(1)}{2}(\alpha - 1)^2 + o((\alpha - 1)^2)$ for $i \in \{3, 4\}$. Using these expansions, we can rewrite

$$\begin{aligned}
P(y_0(z_0)) &= \frac{1}{\alpha - 1} \left(\sum_{i \in \{1, 2, 5, 6, 7\}} s_i(1) + \sum_{i \in \{1, 2, 5, 6, 7\}} s'_i(1)(\alpha - 1) + o(\alpha - 1) \right. \\
&\quad \left. + \frac{s_3(1) + s_4(1)}{\alpha - 1} + s'_3(1) + s'_4(1) + \frac{1}{2}(s''_3(1) + s''_4(1))(\alpha - 1) + o((\alpha - 1)) \right)
\end{aligned}$$

In order to continue, we compute:

$$\begin{aligned}
s_1(\alpha) &= -\frac{1}{8} + \frac{1}{8}(\alpha - 1) + o(\alpha - 1) \\
s_2(\alpha) &= \frac{1}{2}z_0 + z_0(\alpha - 1) + o(\alpha - 1) \\
s_3(\alpha) &= -\frac{1}{16}z_0^2 - \frac{1}{4}z_0^2(\alpha - 1) - \frac{1}{2}z_0^2(\alpha - 1)^2 + o((\alpha - 1)^2) \\
s_4(\alpha) &= \frac{1}{16}z_0^2 + \frac{z_0^2}{4} \left(\frac{1}{8} + \ln \frac{z_0}{2} \right) (\alpha - 1) \\
&\quad + \frac{z_0^2}{8} \left(8 \left(\ln \frac{z_0}{2} \right)^2 + 2 \ln \frac{z_0}{2} - \frac{1}{2} \right) (\alpha - 1)^2 + o((\alpha - 1)^2) \\
s_5(\alpha) &= \frac{z_0^2}{4} B^-(1/2; 3, 0) + o(\alpha - 1) \\
&\quad + z_0^2 \left(\left(\ln(z_0) + \frac{1}{4} \right) B^-(1/2; 3, 0) + 1/2 \int_0^{\frac{1}{2}} \ln(t(1-t)) t^2 (1-t)^{-1} dt \right) (\alpha - 1) + o(\alpha - 1) \\
s_6(\alpha) &= \frac{(1 - z_0)^2}{8} + \frac{(1 - z_0)^2}{4} (\ln(1 - z_0) - 1/2)(\alpha - 1 + o(\alpha - 1)) \\
s_7(\alpha) &= -\frac{z_0^2}{4} B^-(1 - z_0; 2, -1) + o(\alpha - 1) \\
&\quad - z_0^2 \left(\ln(z_0) B^-(1 - z_0; 2, -1) + \int_0^{1-z_0} t(1-t)^{-2} \ln \left(\frac{\sqrt{t}}{1-t} \right) t(1-t)^{-2} dt \right) (\alpha - 1).
\end{aligned}$$

Based on this we see that

$$s_3(1) + s_4(1) = -\frac{1}{16}z_0^2 + \frac{1}{16}z_0^2 = 0$$

and

$$\sum_{i \in \{1, 2, 5, 6, 7\}} s_i(1) + s'_3(1) + s'_4(1)$$

$$\begin{aligned}
&= -\frac{1}{8} + \frac{1}{2}z_0 - \frac{1}{4}z_0^2 + \frac{z_0^2}{32} + \frac{z_0^2}{4}\ln\left(\frac{z_0}{2}\right) + \frac{z_0^2}{4}B^-(1/2; 3, 0) + \frac{(1-z_0)^2}{8} - \frac{z_0^2}{4}B^-(1-z_0; 2, -1) \\
&= -\frac{1}{8} + \frac{1}{2}z_0 - \frac{1}{4}z_0^2 + \frac{z_0^2}{32} + \left(\frac{z_0^2}{4}\ln(z_0) - \frac{z_0^2}{4}\ln 2\right) + \left(-\frac{5z_0^2}{32} + \frac{z_0^2}{4}\ln 2\right) \\
&\quad + \left(\frac{1}{8} - \frac{z_0}{4} + \frac{z_0^2}{8}\right) + \left(\frac{z_0^2}{4} - \frac{z_0}{4} - \frac{z_0^2}{4}\ln z_0\right) = 0,
\end{aligned}$$

using that

$$\begin{aligned}
B^-\left(\frac{1}{2}; 3, 0\right) &= \int_0^{\frac{1}{2}} t^2(1-t)^{-1} dt = \int_{\frac{1}{2}}^1 (1-s)^2 s^{-1} ds \\
&= \int_{\frac{1}{2}}^1 s^{-1} - 2 + s ds = -2 + \frac{1}{2} - \ln \frac{1}{2} + 1 - \frac{1}{8} = -\frac{5}{8} + \ln 2
\end{aligned}$$

and

$$\begin{aligned}
B^-(1-z_0; 2, -1) &= \int_0^{1-z_0} t(1-t)^{-2} dt = \int_{z_0}^1 (1-s)s^{-2} ds \\
&= \int_{z_0}^1 s^{-2} - s^{-1} ds = -1 + z_0^{-1} + \ln z_0.
\end{aligned}$$

Finally, it follows that as $\alpha \rightarrow 1$,

$$P(y_0(z_0)) = \sum_{i \in \{1, 2, 5, 6, 7\}} s'_i(1) + \frac{1}{2}(s''_3(1) + s''_4(1)) + o(1)$$

Therefore, the desired value of $\lim_{\alpha \rightarrow 1} P(y_0(z_0))$ is given by

$$\begin{aligned}
&\sum_{i \in \{1, 2, 5, 6, 7\}} s'_i(1) + \frac{1}{2}(s''_3(1) + s''_4(1)) \\
&= \frac{1}{8} + z_0 - \frac{z_0^2}{4} + \frac{z_0^2}{8}(4(\ln \frac{z_0}{2})^2 + \ln \frac{z_0}{2} - \frac{1}{4}) + \frac{(1-z_0)^2}{4}(\ln(1-z_0) - 1/2) \\
&\quad + z_0^2 \left(\left(\ln(z_0) + \frac{1}{4} \right) B^-(1/2; 3, 0) + \frac{1}{2} \int_0^{\frac{1}{2}} \ln(t(1-t)) t^2(1-t)^{-1} dt \right) \\
&\quad - z_0^2 \left(\ln(z_0) B^-(1-z_0; 2, -1) + \int_0^{1-z_0} \ln\left(\frac{\sqrt{t}}{1-t}\right) t(1-t)^{-2} dt \right) \\
&= \frac{1}{8} + z_0 - \frac{z_0^2}{4} + \frac{z_0^2}{2}(\ln \frac{z_0}{2})^2 + \frac{z_0^2}{8} \ln \frac{z_0}{2} - \frac{z_0^2}{32} \\
&\quad - \frac{5}{8} z_0^2 \ln(z_0) + z_0^2 \ln(z_0) \ln 2 - \frac{5z_0^2}{32} + \frac{z_0^2 \ln 2}{4} \\
&\quad + z_0^2/2 \int_0^{\frac{1}{2}} \ln(t(1-t)) t^2(1-t)^{-1} dt \\
&\quad + \frac{(1-z_0)^2}{4} \ln(1-z_0) - \frac{1}{8} + \frac{z_0}{4} - \frac{z_0^2}{8} \\
&\quad + z_0^2 \ln(z_0) - z_0 \ln z_0 - z_0^2(\ln z_0)^2 - z_0^2 \int_0^{1-z_0} \ln\left(\frac{\sqrt{t}}{1-t}\right) t(1-t)^{-2} dt \\
&= \frac{5}{4} z_0 - \frac{9}{16} z_0^2 + \frac{z_0^2}{2}(\ln \frac{z_0}{2})^2 + \frac{z_0^2}{8} \ln \frac{z_0}{2} + \frac{(1-z_0)^2}{4} \ln(1-z_0) \\
&\quad + \frac{3}{8} z_0^2 \ln(z_0) + z_0^2 \ln(z_0) \ln 2 + \frac{z_0^2 \ln 2}{4} + z_0^2/2 \int_0^{\frac{1}{2}} \ln(t(1-t)) t^2(1-t)^{-1} dt
\end{aligned}$$

$$\begin{aligned}
& -z_0 \ln z_0 - z_0^2 (\ln z_0)^2 - z_0^2 \int_0^{1-z_0} \ln \left(\frac{\sqrt{t}}{1-t} \right) t(1-t)^{-2} dt \\
&= \frac{5}{4} z_0 - \frac{9}{16} z_0^2 + \frac{z_0^2}{2} (\ln \frac{z_0}{2})^2 + \frac{z_0^2}{8} \ln \frac{z_0}{2} + \frac{(1-z_0)^2}{4} \ln(1-z_0) \\
& \quad + \frac{3}{8} z_0^2 \ln(z_0) + z_0^2 \ln(z_0) \ln 2 + \frac{z_0^2 \ln 2}{4} + z_0^2/2(11/8 - 1/4 \ln 2 - 3/2 \ln(2)^2 - \text{Li}_2(1/2)) \\
& \quad - z_0 \ln z_0 - z_0^2 (\ln z_0)^2 + z_0(1 + \frac{1}{2}(2-z_0) \ln(z_0) + \frac{1}{2} z_0 \ln(z_0)^2 - \frac{1}{2}(1-z_0) \ln(1-z_0) \\
& \quad + \frac{1}{2} z_0 \text{Li}_2(z_0)) - z_0^2 - \frac{1}{2} z_0^2 \text{Li}_2(1) \\
&= \frac{9}{4} z_0 - \frac{25}{16} z_0^2 + \frac{z_0^2}{2} (\ln \frac{z_0}{2})^2 + \frac{z_0^2}{8} \ln \frac{z_0}{2} + \frac{(1-z_0)^2}{4} \ln(1-z_0) \\
& \quad - \frac{1}{8} z_0^2 \ln(z_0) + z_0^2 \ln(z_0) \ln 2 + \frac{z_0^2 \ln 2}{4} + z_0^2/2(11/8 - 1/4 \ln 2 - 3/2 \ln(2)^2 \\
& \quad - \text{Li}_2(1/2) - \text{Li}_2(1) + \text{Li}_2(z_0)) - \frac{1}{2} z_0^2 (\ln z_0)^2 - \frac{1}{2} z_0(1-z_0) \ln(1-z_0)
\end{aligned}$$

where we used that

$$z_0^2/2 \int_0^{\frac{1}{2}} \ln(t) t^2 (1-t)^{-1} + \ln(1-t) t^2 (1-t)^{-1} dt = 11/8 - 1/4 \ln 2 - 3/2 \ln(2)^2 - \text{Li}_2(1/2),$$

and

$$\begin{aligned}
& z_0^2 \int_0^{1-z_0} 1/2 \ln(t) t(1-t)^{-2} - t \ln(1-t) (1-t)^{-2} dt \\
&= -\frac{1}{z_0} (1 + \frac{1}{2}(2-z_0) \ln(z_0) + \frac{1}{2} z_0 \ln(z_0)^2 - \frac{1}{2}(1-z_0) \ln(1-z_0) + \frac{1}{2} z_0 \text{Li}_2(z_0)) + 1 + \frac{1}{2} \text{Li}_2(1).
\end{aligned}$$

By expanding the squares and collecting terms, the last expression can be simplified to

$$\begin{aligned}
& \frac{9}{4} z_0 + \frac{1-4z_0+3z_0^2}{4} \ln(1-z_0) + z_0^2 \left(-7/8 - \frac{\ln(2)^2 + 2 \text{Li}_2(1/2) + 2 \text{Li}_2(1)}{4} \right) + \frac{1}{2} z_0^2 \text{Li}_2(z) \\
&= \frac{9}{4} z_0 + \frac{1-4z_0+3z_0^2}{4} \ln(1-z_0) - \frac{7+\pi^2}{8} z_0^2 + \frac{1}{2} z_0^2 \text{Li}_2(z)
\end{aligned}$$

which finishes the computation. \square

Proof of Proposition 4.5: It suffices to find the value of J and $I^{(k)}$ at $\alpha = 1$. We can do this by computing the integrals with the expression for $P(y)$ that we found for $\alpha = 1$, i.e.

$$\begin{aligned}
J &= 2\alpha \int_0^1 \left(\frac{9}{4} z + \frac{1-4z+3z^2}{4} \ln(1-z) - \frac{7+\pi^2}{8} z^2 + \frac{1}{2} z^2 \text{Li}_2(z) \right) z^{2\alpha-1} dz \\
&= \frac{575-12\pi^2}{576}
\end{aligned}$$

and

$$\begin{aligned}
I^{(k)} &= \frac{2\alpha \xi^k}{k!} \int_0^1 \left(\frac{9}{4} z + \frac{1-4z+3z^2}{4} \ln(1-z) - \frac{7+\pi^2}{8} z^2 + \frac{1}{2} z^2 \text{Li}_2(z) \right) z^{2\alpha-k-1} e^{-\xi/z} dz \\
&= \frac{2\eta^k}{k!} \int_0^1 \left(\frac{9}{4} z + \frac{1-4z+3z^2}{4} \ln(1-z) - \frac{7+\pi^2}{8} z^2 + \frac{1}{2} z^2 \text{Li}_2(z) \right) z^{1-k} e^{-\eta/z} dz \\
&= \frac{9\eta^k}{2k!} \eta^{3-k} \Gamma^+(k-3, \eta) - \frac{\eta^k}{k!} \frac{7+\pi^2}{4} \eta^{4-k} \Gamma^+(k-4, \eta)
\end{aligned}$$

$$\begin{aligned}
& + \frac{\eta^k}{2k!} \int_0^1 (1-4z+3z^2) \ln(1-z) z^{1-k} e^{-\eta/z} dz + \frac{\eta^k}{k!} \int_0^1 z^{3-k} \text{Li}_2(z) e^{-\eta/z} dz \\
& = \frac{9\eta^3}{2k!} \Gamma^+(k-3, \eta) - \frac{\eta^4}{k!} \frac{7+\pi^2}{4} \Gamma^+(k-4, \eta) \\
& + \frac{\eta^k}{2k!} \int_0^1 (1-4z+3z^2) \ln(1-z) z^{1-k} e^{-\eta/z} dz + \frac{\eta^k}{k!} \int_0^1 z^{3-k} \text{Li}_2(z) e^{-\eta/z} dz
\end{aligned}$$

where $\eta = \frac{4\nu}{\pi}$ and $\text{Li}_2(z) = \sum_{t=1}^{\infty} z^t/t^2$, the dilogarithm function. Plugging this into (42) and (41) yields the expressions in the statement of the proposition. \blacksquare

4.2 The proof of Proposition 1.5

Instead of extracting the scaling of $\gamma(k)$ from its explicit expression, it turns out to be more convenient to derive it directly. Recall that

$$\gamma(k) = \frac{\int_0^{\infty} \rho(y, k) P(y) \alpha e^{-\alpha y} dy}{\int_0^{\infty} \rho(y, k) \alpha e^{-\alpha y} dy}.$$

The asymptotic behavior for the denominator follows from (21). Hence, the main term to consider is the numerator

$$\int_0^{\infty} P(y) \rho(y, k) \alpha e^{-\alpha y} dy,$$

and in particular the function $P(y)$. We therefore start with establishing the asymptotic behavior of the latter. First we combine (21) and (22) to obtain the following scaling result

$$\frac{\int_0^{\infty} e^{-\beta y} \rho(y, k) \alpha e^{-\alpha y} dy}{\int_0^{\infty} \rho(y, k) \alpha e^{-\alpha y} dy} \sim \xi^{2\beta} k^{-2\beta}. \quad (58)$$

Proposition 4.7 (Asymptotic behavior of $P(y)$). *Let $\alpha > \frac{1}{2}$, $\nu > 0$ and c_α as defined in Proposition 1.5. Then, as $y \rightarrow \infty$,*

1. *for $\frac{1}{2} < \alpha < \frac{3}{4}$,*

$$P(y) \sim e^{-\frac{y}{2}(4\alpha-2)} c_\alpha \xi^{4\alpha-2},$$

2. *for $\alpha = \frac{3}{4}$,*

$$P(y) \sim \frac{y}{2} e^{-\frac{y}{2}},$$

3. *and for $\alpha > \frac{3}{4}$,*

$$P(y) \sim e^{-\frac{y}{2}} \frac{\alpha - \frac{1}{2}}{\alpha - \frac{3}{4}}.$$

Proof. We shall deal with each of the three cases for α separately.

Proof for $1/2 < \alpha < 3/4$ By Lemma 4.1 we get that

$$\begin{aligned}
e^{(4\alpha-2)\frac{y}{2}} P(y) &= \frac{2^{-4\alpha-1}(3\alpha-1)}{\alpha(\alpha-1)^2} + \frac{(\alpha-\frac{1}{2})B^-(\frac{1}{2}; 1+2\alpha, -2+2\alpha)}{2(\alpha-1)\alpha} - \frac{B^-(1-e^{-\frac{y}{2}}; 2\alpha, 3-4\alpha)}{4(\alpha-1)} \\
&+ \frac{e^{(4\alpha-2)\frac{y}{2}}}{8(\alpha-1)\alpha} \left((1-e^{-\frac{y}{2}})^{2\alpha} - 1 \right) + \frac{\alpha-\frac{1}{2}}{\alpha-1} e^{(4\alpha-3)\frac{y}{2}} - \frac{(\alpha-\frac{1}{2})^2}{4(\alpha-1)^2} e^{4(\alpha-1)\frac{y}{2}}.
\end{aligned}$$

Because for any $b < 1$, $B^-(1-z; a, b)$ converges to $B(a, b) < \infty$ as $z \rightarrow 0$, we get that as $y \rightarrow \infty$, the first line is asymptotically equivalent to

$$\frac{3\alpha-1}{2^{4\alpha+1}\alpha(\alpha-1)^2} + \frac{(\alpha-1/2)B^-(1/2; 1+2\alpha, -2+2\alpha)}{2(\alpha-1)\alpha} - \frac{B(2\alpha, 3-4\alpha)}{4(\alpha-1)} = c_\alpha \xi^{-(4\alpha-2)},$$

with c_α as defined in Proposition (1.5). The proof now follows since for $1/2 < \alpha < 3/4$, the remaining three terms go to zero as $y \rightarrow \infty$.

Proof for $\alpha = 3/4$ Similar to the previous case we use Lemma 4.1 to obtain (evaluating the expressions for $\alpha = 3/4$)

$$\begin{aligned} \frac{2}{y} e^{\frac{y}{2}} P(y) &= \frac{2}{y} B^-(1 - e^{-\frac{y}{2}}; 3/2, 0) - \frac{4}{y} \frac{e^{\frac{y}{2}} ((1 - e^{-\frac{y}{2}})^{3/2} - 1)}{3} - \frac{1}{y} - \frac{e^{-\frac{y}{2}}}{4y} \\ &\quad + \frac{2}{y} \left(\frac{5}{3} - \frac{2B^-(\frac{1}{2}; 5/2, -1/2)}{3} \right) \end{aligned}$$

First we note that as $y \rightarrow \infty$,

$$e^{\frac{y}{2}} \left((1 - e^{-\frac{y}{2}})^{3/2} - 1 \right) \sim -\frac{3}{2}, \quad (59)$$

which implies that

$$\lim_{y \rightarrow \infty} \frac{4}{y} \frac{e^{\frac{y}{2}} ((1 - e^{-\frac{y}{2}})^{3/2} - 1)}{3} = 0.$$

We can now conclude that all terms in $\frac{2}{y} e^{\frac{y}{2}} P(y)$ except the first one are $o(1)$ as $y \rightarrow \infty$. By writing $z = e^{-\frac{y}{2}}$ we can rewrite the first term as

$$\frac{2}{y} B^-(1 - e^{-\frac{y}{2}}; 3/2, 0) = -\frac{B^-(1 - z; 3/2, 0)}{\log(z)}.$$

Since $B^-(1 - z, 3/2, 0) \sim -\log(z)$ as $z \rightarrow 0$, see Lemma B.1, it now follows that for $\alpha = 3/4$,

$$\lim_{y \rightarrow \infty} \frac{2}{y} B^-(1 - e^{-\frac{y}{2}}; 3/2, 0) = \lim_{z \rightarrow 0} -\frac{1}{\log(z)} B^-(1 - z; 3/2, 0) = 1.$$

We therefore conclude that

$$P(y) \sim \frac{y}{2} e^{-\frac{y}{2}},$$

as $y \rightarrow \infty$.

Proof for $\alpha > 3/4$ We first deal with the case $\alpha = 1$. Here it follows from Proposition 4.5 that

$$\begin{aligned} e^{y/2} P(y) &= \frac{9}{4} + \frac{e^{y/2} \log(1 - e^{-y/2})}{4} \\ &\quad - \log(1 - e^{-y/2}) + e^{-y/2} \left(\frac{3}{4} \log(1 - e^{-y/2}) - \frac{7 + \pi^2}{8} + \frac{1}{2} \text{Li}_2(e^{-y}) \right) \\ &= 2 + \left(\frac{e^{y/2} \log(1 - e^{-y/2})}{4} + 1 \right) \\ &\quad - \log(1 - e^{-y/2}) + e^{-y/2} \left(\frac{3}{4} \log(1 - e^{-y/2}) - \frac{7 + \pi^2}{8} + \frac{1}{2} \text{Li}_2(e^{-y}) \right) \end{aligned}$$

The last two terms are $o(1)$ as $y \rightarrow \infty$, while $2 = (\alpha - 1/2)/(\alpha - 3/4)$ for $\alpha = 1$.

Now we will deal with the case $\alpha > 3/4$ and $\alpha \neq 1$. For simplicity we write

$$Q_\alpha := \frac{2^{-4\alpha-1}(3\alpha-1)}{\alpha(\alpha-1)^2} + \frac{(\alpha-1/2)B^-(1/2; 1+2\alpha, -2+2\alpha)}{2(\alpha-1)\alpha}.$$

Then, by Lemma 4.1 we get

$$\begin{aligned} e^{y/2} P(y) &= \frac{\alpha - \frac{1}{2}}{\alpha - 1} + \frac{e^{\frac{y}{2}}}{8(\alpha-1)\alpha} \left((1 - e^{-\frac{y}{2}})^{2\alpha} - 1 \right) \\ &\quad - e^{-(4\alpha-3)\frac{y}{2}} \frac{B^-(1 - e^{-\frac{1}{2}y}; 2\alpha, 3-4\alpha)}{4(\alpha-1)} \end{aligned}$$

$$+ e^{-(4\alpha-3)\frac{y}{2}} Q_\alpha + \frac{(\alpha - \frac{1}{2})^2}{4(\alpha-1)^2} e^{-\frac{y}{2}}.$$

The first term is constant while the last two terms go to zero as $y \rightarrow \infty$. We will therefore focus on the remaining two terms. For the first we have, see (59)

$$\frac{e^{\frac{y}{2}}}{8(\alpha-1)\alpha} \left(\left(1 - e^{-\frac{y}{2}}\right)^{2\alpha} - 1 \right) \sim \frac{-2\alpha}{8(\alpha-1)\alpha} = -\frac{1}{4(\alpha-1)},$$

as $y \rightarrow \infty$. Finally, writing $z = e^{-\frac{y}{2}}$ we get that

$$e^{-(4\alpha-3)\frac{y}{2}} B^-(1 - e^{-\frac{1}{2}y}; 2\alpha, 3 - 4\alpha) = z^{4\alpha-3} B^-(1 - z, 2\alpha, 3 - 4\alpha).$$

Therefore it follows, see Lemma B.1, that

$$\begin{aligned} \lim_{y \rightarrow \infty} -e^{-(4\alpha-3)\frac{y}{2}} \frac{B^-(1 - e^{-\frac{1}{2}y}; 2\alpha, 3 - 4\alpha)}{4(\alpha-1)} &= \lim_{z \rightarrow 0} z^{4\alpha-3} \frac{B^-(1 - z, 2\alpha, 3 - 4\alpha)}{4(\alpha-1)} \\ &= \frac{1}{4(\alpha-1)(4\alpha-3)}. \end{aligned}$$

We conclude that as $y \rightarrow \infty$

$$e^{y/2} P(y) \sim \frac{\alpha - \frac{1}{2}}{\alpha - 1} - \frac{1}{4(\alpha-1)} - \frac{1}{4(\alpha-1)(4\alpha-3)} = \frac{1 - 3\alpha + 2\alpha^2}{(\alpha-1)(\alpha - \frac{3}{4})} = \frac{\alpha - \frac{1}{2}}{\alpha - \frac{3}{4}},$$

which finishes the proof. \square

With the asymptotic behavior of $P(y)$ we are almost ready to prove Proposition 1.5. First we will prove a result that will allow us to limit the values of y , when performing the integration. For this, recall that for any $C > 0$ we defined

$$y_{k,C}^\pm = 2 \log \left(\frac{k \pm C \sqrt{k \log(k)}}{\xi} \vee 1 \right).$$

Since $P(y) \leq 1$ by the concentration of heights results (Proposition 2.4) we have that We will show that, as $k \rightarrow \infty$,

$$\int_0^\infty P(y) \rho(y, k) \alpha e^{-\alpha y} dy = (1 + o(1)) \int_{y_{k,C}^-}^{y_{k,C}^+} P(y) \rho(y, k) \alpha e^{-\alpha y} dy. \quad (60)$$

Note that this implies that if $P(y) = h(y)(1 + o(1))$, uniformly in y , as $y \rightarrow \infty$, then

$$\int_0^\infty P(y) \rho(y, k) \alpha e^{-\alpha y} dy \sim \int_0^\infty h(y) \rho(y, k) \alpha e^{-\alpha y} dy, \quad (61)$$

as $y \rightarrow \infty$.

We now proceed with the proof of Proposition 1.5, which is split over the different cases for α .

Proof when $1/2 < \alpha < 3/4$ By Proposition 4.7 and (61) it follows that as $k \rightarrow \infty$,

$$\gamma(k) \sim c_\alpha \xi^{-(4\alpha-2)} \frac{\int_0^\infty e^{-(4\alpha-2)y/2} \rho(y, k) \alpha e^{-\alpha y} dy}{\int_0^\infty \rho_y(k) \alpha e^{-\alpha y} dy} \sim c_\alpha k^{-4\alpha+2}.$$

where the last line is due to (58) with $\beta = 2\alpha - 1$.

Proof when $\alpha = 3/4$ Similar to the previous case Proposition 4.7 and (61) imply that as $k \rightarrow \infty$

$$\gamma(k) = \frac{\int_0^\infty P(y)\rho(y, k)\alpha e^{-\alpha y} dy}{\int_0^\infty \rho_y(k)\alpha e^{-\alpha y} dy} \sim \frac{\int_0^\infty \frac{y}{2}e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy}{\int_0^\infty \rho_y(k)\alpha e^{-\alpha y} dy}.$$

However, the final step does not follow immediately from (58) because of the additional logarithmic term. To prove the result we first show that

$$\int_{a^-(k)}^{a^+(k)} P(y)\rho(y, k)\alpha e^{-\alpha y} dy \sim \int_{a^-(k)}^{a^+(k)} \frac{y}{2}e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy. \quad (62)$$

For this we establish an upper bound for the left hand side

$$\int_{a^-(k)}^{a^+(k)} \frac{y}{2}e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy \leq \frac{a^+(k)}{2} \int_{a^-(k)}^{a^+(k)} e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy$$

and similarly, a lower bound

$$\int_{a^-(k)}^{a^+(k)} \frac{y}{2}e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy \geq \frac{a^-(k)}{2} \int_{a^-(k)}^{a^+(k)} e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy$$

Now observe that as $k \rightarrow \infty$,

$$\frac{a^\pm(k)}{2} = \log \left(\frac{k \pm \sqrt{k \log(k)}}{\xi_{\alpha, \nu}} \right) \sim \log(k)$$

and therefore it follows that

$$\limsup_{k \rightarrow \infty} \frac{\int_{a^-(k)}^{a^+(k)} \frac{y}{2}e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy}{\log(k) \int_{a^-(k)}^{a^+(k)} e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy} \leq 1.$$

and

$$\liminf_{k \rightarrow \infty} \frac{\int_{a^-(k)}^{a^+(k)} \frac{y}{2}e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy}{\log(k) \int_{a^-(k)}^{a^+(k)} e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy} \geq 1.$$

This proves (62).

Next we note that by (58) with $\beta = 1/2$ we have

$$\frac{\int_0^\infty e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy}{\int_0^\infty \rho(y, k)\alpha e^{-\alpha y} dy} \sim \xi k^{-1}.$$

Therefore, since by (60),

$$\int_0^\infty P(y)\rho(y, k)\alpha e^{-\alpha y} dy \sim \int_{a^-(k)}^{a^+(k)} P(y)\rho(y, k)\alpha e^{-\alpha y} dy$$

it follows from (62) that as $k \rightarrow \infty$,

$$\begin{aligned} \gamma(k) &\sim \frac{\int_0^\infty \frac{y}{2}e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy}{\int_0^\infty \rho(y, k)\alpha e^{-\alpha y} dy} \\ &\sim \log(k) \frac{\int_0^\infty e^{-y/2}\rho(y, k)\alpha e^{-\alpha y} dy}{\int_0^\infty \rho(y, k)\alpha e^{-\alpha y} dy} \sim \xi \log(k) k^{-1} = \frac{6\nu}{\pi} \log(k) k^{-1}, \end{aligned}$$

when $\alpha = 3/4$.

Proof when $\alpha > 3/4$ Again, by Proposition 4.7, equation (61) and (58) with $\beta = 1/2$, it follows that as $k \rightarrow \infty$,

$$\gamma(k) \sim \frac{\alpha - \frac{1}{2}}{\alpha - \frac{3}{4}} \frac{\int_0^\infty e^{-y/2} \rho(y, k) \alpha e^{-\alpha y} dy}{\int_0^\infty \rho(y, k) \alpha e^{-\alpha y} dy} \sim \frac{\alpha - \frac{1}{2}}{\alpha - \frac{3}{4}} \xi k^{-1} = \frac{8\alpha\nu}{\pi(4\alpha - 3)}.$$

5 Proofs of Theorem 1.2 and Theorem 1.3

We will first derive Theorem 1.3. It will turn out that Theorem 1.2 has a quick derivation assuming Theorem 1.3.

5.1 Clustering function, proving Theorem 1.3

In this subsection, we want to show that the clustering function of the KPKVB model $c_n(k) \xrightarrow{\mathbb{P}} \gamma(k)$ for a fixed k . The key idea is that for the poissonized KPKVB model the coupling with the box model is guaranteed to be exact (in the sense that it also preserves edges) for all vertices up to height $H = \frac{R}{4}$ (which is all we need to show the convergence of the clustering function for constant k).

The overall proof proceeds in four steps: firstly poissonizing the KPKVB model, secondly truncating the poissonized KPKVB model at height $H = H_n \rightarrow \infty$, thirdly using that the coupling between the poissonized KPKVB and box model is exact when truncated at this height H and finally showing that the clustering function in the truncated box model agrees with the integral for $\gamma(k)$.

We will encounter the following models: we will start with the KPKVB model $G_n = G(n; \alpha, \nu)$, then the Poissonized KPKVB model $G_{\text{Po}} = G_{\text{Po}}(n; \alpha, \nu)$, moving to the truncated Poissonized KPKVB model $G_{\text{Po}, H}$ which is the subgraph of G_{Po} induced on all vertices with height at most H resp. radial coordinate at least $R - H$, then the coupling will bring us to the truncated box model $G_{\text{box}, H}$ which is the subgraph of the box model G_{box} induced by all vertices with height at most H .

5.1.1 Poissonization

We first show we can restrict attention to the Poissonized instead of the standard KPKVB model.

Lemma 5.1. *Let $\alpha > \frac{1}{2}, \nu > 0, k \geq 2$ be fixed. Then*

$$c(k; G_n) - c(k; G_{\text{Po}}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0.$$

Proof. We consider the coupling where we have an infinite supply of i.i.d. points u_1, u_2, \dots chosen according to the (α, R) -quasi uniform distribution, the vertices of $G(n; \alpha, \nu)$ are u_1, \dots, u_n and the vertices of $G_{\text{Po}}(n; \alpha, \nu)$ are u_1, \dots, u_N with $N \stackrel{d}{=} \text{Po}(n)$ independently of u_1, u_2, \dots .

Due to the coupling, the KPKVB graph or its poissonization is contained in the other, i.e. we can use new graph labels $G_1 \subset G_2$; e.g. if $N \geq n$, then $G_1 \stackrel{d}{=} G(n; \alpha, \nu)$ and $G_2 \stackrel{d}{=} G_{\text{Po}}(n; \alpha, \nu)$. We write $V_{1,k}$ for the set of vertices of G_1 which have k neighbours in G_1 and $V_{2,k}$ for the set of vertices of G_2 which have k neighbours in G_2 . Write $N_{1,k} = |V_{1,k}|$ and $N_{2,k} = |V_{2,k}|$.

Write W_1 for the set of all vertices of G_1 which have k neighbours in G_1 but no neighbour in $G_2 \setminus G_1$. Write W_2 for the set of all vertices of G_1 which have k neighbours in G_1 and at least one neighbour in $G_2 \setminus G_1$. Write W_3 for the set of all vertices of G_2 which have strictly less than k neighbours in G_1 , but k neighbours in G_2 .

With this notation, we can rewrite:

$$|c(k; G_n) - c(k; G_{\text{Po}})| = \left| \sum_{v \in V_{1,k}} \frac{c_{G_1}(v)}{N_{1,k}} - \sum_{v \in V_{2,k}} \frac{c_{G_2}(v)}{N_{2,k}} \right|$$

$$= \left| \sum_{v \in W_1} \frac{c_{G_1}(v)}{N_{1,k}} + \sum_{v \in W_2} \frac{c_{G_1}(v)}{N_{1,k}} - \sum_{v \in W_1} \frac{c_{G_2}(v)}{N_{2,k}} - \sum_{v \in W_3} \frac{c_{G_2}(v)}{N_{2,k}} \right|.$$

For every vertex of W_1 , the clustering coefficient in G_1 and in G_2 have the same value. Hence, we can simplify:

$$|c(k; G_n) - c(k; G_{Po})| = \left| \sum_{v \in W_1} c_{G_1}(v) \frac{N_{2,k} - N_{1,k}}{N_{1,k} N_{2,k}} + \sum_{v \in W_2} \frac{c_{G_1}(v)}{N_{1,k}} - \sum_{v \in W_3} \frac{c_{G_2}(v)}{N_{2,k}} \right|.$$

Using the triangle inequality for the absolute value and the fact that $|c_G(v)| \leq 1$,

$$|c(k; G_n) - c(k; G_{Po})| \leq \frac{|W_1| |N_{2,k} - N_{1,k}|}{N_{1,k} N_{2,k}} + \frac{|W_2|}{N_{1,k}} + \frac{|W_3|}{N_{2,k}}.$$

Using that $|W_1| \leq N_{1,k}$,

$$|c(k; G_n) - c(k; G_{Po})| \leq \frac{|N_{2,k} - N_{1,k}|}{N_{2,k}} + \frac{|W_2|}{N_{1,k}} + \frac{|W_3|}{N_{2,k}}.$$

From the Chernoff bound for Poisson random variables, it follows that for any $c > 0$, $N \in [n - c\sqrt{n \log n}, n + c\sqrt{n \log n}]$ a.a.s.. This implies that a.a.s. the vertex sets of G_1 and G_2 differ by at most $c\sqrt{n \log n}$ many vertices. In particular also $|N_{1,k} - N_{2,k}|, |W_2|, |W_3| = O(\sqrt{n \log n})$ a.a.s.

By the paper by Gugelmann et al. [21] it holds that $N_{1,k}$ or $N_{2,k}$ is $\Theta(n)$ a.a.s. for fixed degree k and as $|N_{1,k} - N_{2,k}| = O(\sqrt{n \log n})$ a.a.s., it follows that both $N_{1,k}, N_{2,k} = \Theta(n)$ a.a.s. Putting everything together yields the claim:

$$|c(k; G_n) - c(k; G_{Po})| \leq \frac{|N_{2,k} - N_{1,k}|}{N_{2,k}} + \frac{|W_2|}{N_{1,k}} + \frac{|W_3|}{N_{2,k}} \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0.$$

□

5.1.2 Truncation

Consider the truncated Poissonized KPKVB model $G_{Po,H}$ with $H = H_n = \frac{R}{4}$.

Lemma 5.2. *Let $N_1(H)$ denote the number of vertices with radial coordinate at most $R - H \in [0, R]$ in the Poissonized KPKVB model G_{Po} . Then $\mathbb{E}[N_1(H)] = o(n)$.*

Proof. By the Poisson distribution of the number of points inside the disk with radius $R - H$, by the asymptotic behaviour of cosh and using that all $R, R - H, H \rightarrow \infty$,

$$\mathbb{E}[N_1(H)] = n \frac{\cosh(\alpha(R - H)) - 1}{\cosh(\alpha R) - 1} = (1 + o(1)) n e^{-\alpha H} = o(n)$$

□

Lemma 5.3. *Let $N_2(H)$ denote the number of vertices with at least one neighbour with radial coordinate at most $R - H \in [0, R]$ in the Poissonized KPKVB model. If $H \rightarrow \infty$, then $\mathbb{E}[N_2(H)] = o(n)$.*

Proof. Among the vertices with at least one neighbour with radial coordinate at most $R - H$, we distinguish between those that themselves have radial coordinate at most $R - H_0$ and those that themselves have radial coordinate at least $R - H_0$ where $H_0 = \log H$. We will show that the expectation of both numbers is sublinear in n .

Since $H \rightarrow \infty$ we also have that $H_0 = \log H \rightarrow \infty$. Hence by Lemma 5.2, the expected number of vertices with radial coordinate at most $R - H_0$ (in particular those vertices among them with at least one neighbour with radial coordinate at most $R - H$), $\mathbb{E}[N_1(H_0)] = o(n)$ is sublinear. For a vertex with radial coordinate at least $R - H_0$, the probability that it has a neighbour with radial coordinate at most $R - H$, tends to zero. Therefore, by the Campbell-Mecke formula the number of such vertices is sublinear. □

Lemma 5.4. Let $\alpha > \frac{1}{2}$, $\nu > 0$ and $k \geq 2$ be fixed. Let G_{Po} denote the Poissonized KPKVB model and $G_{\text{Po},H}$ denote the truncated Poissonized KPKVB model. Then

$$c(k; G_{\text{Po}}) - c(k; G_{\text{Po},H}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0.$$

Proof. Let $V_n(k)$ denote the set of all vertices with degree k in the Poissonized KPKVB model G_{Po} and $V_{n,H}(k)$ the set of all vertices with degree k in the truncated Poissonized KPKVB model $G_{\text{Po},H}$. Note that a vertex may have a smaller degree and different clustering coefficient in G_{Po} and $G_{\text{Po},H}$ (even if it is a vertex of both graphs). We write $c_n(v)$ for the clustering coefficient of a vertex v in G_{Po} and $c_{n,H}(v)$ for the clustering coefficient in $G_{\text{Po},H}$. Let $N_n(k) = |V_n(k)|$ denote the number of degree k vertices in G_{Po} . Then as $N_n(k) = \Theta(nk^{-(2\alpha+1)})$ a.a.s., there is a constant $a_1(k) > 0$ such that a.a.s. $N_n(k) \geq a_1(k)n$. Let $N_{n,H}(k) = |V_{n,H}(k)|$ be the number of degree k vertices in $G_{\text{Po},H}$.

Using a telescoping sum, we obtain that

$$\begin{aligned} |c(k; G_{\text{Po}}) - c(k; G_{\text{Po},H})| &\leq \left| c(k; G_{\text{Po}}) - \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} c_n(v) \right| \\ &\quad + \left| \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} c_n(v) - \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} c_{n,H}(v) \right| \\ &\quad + \left| \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} c_{n,H}(v) - c(k; G_{\text{Po},H}) \right|. \end{aligned}$$

Now we use $X_1 := |V_n(k) \setminus V_{n,H}(k)|$, $X_2 := \sum_{v \in V_{n,H}(k)} |c_n(v) - c_{n,H}(v)|$ and $X_3 := |N_{n,H}(k) - N_n(k)|$ to bound the three differences. In the first difference, after factorizing $N_n(k)$, all terms in $V_{n,H}(k)$ cancel and the remaining terms are all bounded by one:

$$\left| c(k; G_{\text{Po}}) - \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} c_n(v) \right| = \left| \frac{1}{N_n(k)} \sum_{v \in V_n \setminus V_{n,H}(k)} c_n(v) \right| \leq \frac{X_1}{N_n(k)}.$$

In the second difference, the sums can be aligned and then the triangle inequality for the absolute value applied:

$$\left| \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} c_n(v) - \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} c_{n,H}(v) \right| \leq \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} |c_n(v) - c_{n,H}(v)| = \frac{X_2}{N_n(k)}.$$

In the third difference, we can simplify:

$$\begin{aligned} \left| \frac{1}{N_n(k)} \sum_{v \in V_{n,H}(k)} c_{n,H}(v) - c(k; G_{\text{Po},H}) \right| &= \left| \sum_{v \in V_{n,H}(k)} c_{n,H}(v) \left(\frac{1}{N_n(k)} - \frac{1}{N_{n,H}(k)} \right) \right| \\ &= \frac{|N_{n,H}(k) - N_n(k)|}{N_n(k)N_{n,H}(k)} \sum_{v \in V_{n,H}(k)} c_{n,H}(v). \end{aligned}$$

Thus, we end up with the upper bound, using that $c_{n,H}(k) \leq 1$ and $N_{n,H}(k) = |V_{n,H}(k)|$,

$$|c(k; G_{\text{Po}}) - c(k; G_{\text{Po},H})| \leq \frac{X_1}{N_n(k)} + \frac{X_2}{N_n(k)} + \frac{X_3}{N_n(k)} \frac{\sum_{v \in V_{n,H}(k)} c_{n,H}(v)}{N_{n,H}(k)} \leq \frac{X_1 + X_2 + X_3}{N_n(k)}.$$

Now using that a.a.s. $N_n(k) \geq a_1(k)n$ gives that a.a.s.

$$|c(k; G_{\text{Po}}) - c(k; G_{\text{Po},H})| \leq \frac{X_1 + X_2 + X_3}{a_1(k)n}.$$

By Lemma 5.2, we have that $\mathbb{E}[X_1] = \mathbb{E}[X_3] = o(n)$ and by Lemma 5.3, we have that $\mathbb{E}[X_2] = o(n)$. We conclude that $\mathbb{E}|c(k; G_{Po}) - c(k; G_{Po,H})| = o(1)$. From this, convergence in probability follows by Markov's inequality: for $\epsilon > 0$,

$$\mathbb{P}(|c(k; G_{Po}) - c(k; G_{Po,H})| \geq \epsilon) \leq \frac{\mathbb{E}|c(k; G_{Po}) - c(k; G_{Po,H})|}{\epsilon} = o(1).$$

□

5.1.3 Coupling

Lemma 5.5. *The clustering function in the truncated Poissonized KPKVB model $G_{Po,H}$ and in the truncated box model $G_{box,H}$ agree, i.e. $c(k; G_{Po,H}) = c(k; G_{box,H})$.*

Proof. As all vertices involved have height $\leq H$ resp. radial coordinate $\geq R - H$ and as $H \leq \frac{R}{4}$, all vertices involved (in the disk) will have radial coordinate $\geq \frac{3}{4}R$, therefore by Lemma 2.3, edges exist between such vertices in the disk if and only if they exist in the box. So the two graphs are isomorphic and hence the clustering functions agree. □

5.1.4 Deriving the limit integral expression

Lemma 5.6. *Let $\alpha > \frac{1}{2}$, $\nu > 0$ and $k \geq 2$ be fixed. Then*

$$c(k; G_{box,H}) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \gamma(k).$$

Let $\mathcal{R}_H = \mathcal{R}([0, H])$ denote the truncated box. For a vertex v of the vertex set $V = V(G_{box,H})$, writing $\Gamma(v)$ for the set of all neighbours of vertex v , let

$$h(V, v) = \binom{k}{2}^{-1} \sum_{uv \in \binom{\Gamma(v)}{2}} \mathbb{1}_{\{uw \in E(G_{box,H})\}} \mathbb{1}_{\{D_{G_{box,H}}(v)=k\}}.$$

Let $X = \frac{1}{n} \sum_{v \in V(G_{box,H})} h(V(G_{box,H}), v)$. Let $Z = \frac{N_{box,H}(k)}{n} = \frac{1}{n} \sum_{v \in G_{box,H}} \mathbb{1}_{\{D_{G_{box,H}}(v)=k\}}$. Then we have that $c(k; G_{box,H}) = \frac{X}{Z}$. We will show separately that numerator X and denominator Z converge in probability (where the limit of Z is a positive constant). From this, the claim of the lemma will follow as convergence in probability is closed under taking quotients.

By the Campbell-Mecke formula, we have

$$\mathbb{E}[X] = \frac{1}{n} \int_{\mathcal{R}_H} \mathbb{E}h(V, v) \mu(dv) = \int_0^H \mathbb{E}h(V, (0, y)) \alpha e^{-\alpha y} dy.$$

Then, we condition on the additional vertex $(0, y) \in \mathcal{R}_H$ having degree k ,

$$\mathbb{E}[X] = \int_0^H \mathbb{E}[h(V, (0, y)) | D_{G_{box,H}}(0, y) = k] \mathbb{P}(D_{G_{box,H}}(0, y) = k) \alpha e^{-\alpha y} dy.$$

We observe that for $H \rightarrow \infty$, the integrand converges point-wise for $y \in (0, \infty)$ to $P(y)\rho(y, k)\alpha e^{-\alpha y}$ and is always bounded by the integrable function $\alpha e^{-\alpha y}$. Therefore, by the theorem of dominated convergence, as $H \rightarrow \infty$,

$$\mathbb{E}[X] \rightarrow \int_0^\infty P(y)\rho(y, k)\alpha e^{-\alpha y} dy.$$

Note that for X^2 , the summation can be split into off-diagonal (with $v_1 \neq v_2$) and diagonal (with $v_1 = v_2$) terms, i.e. write $X^2 = \frac{1}{n^2}(Y_1 + Y_2)$ where

$$Y_1 = \sum_{v_1 \neq v_2 \in V(G_{box,H})} h(V(G_{box,H}), v_1) h(V(G_{box,H}), v_2),$$

and $Y_2 = \sum_{v_1 \in V(G_{box,H})} h(V(G_{box,H}), v_1)^2$. As $h \leq 1$, $Y_2 \leq nX \leq n$ and so, $\mathbb{E}Y_2 \leq n$, we have by the Campbell-Mecke formula,

$$\mathbb{E}[Y_1] = \int_{\mathcal{R}_H} \int_{\mathcal{R}_H} \mathbb{E}[h(V(G_{box,H}), v_1)h(V(G_{box,H}), v_2)]\mu(dv_1)\mu(dv_2).$$

If $v_1 = (x_1, y_1)$ and $v_2 = (x_2, y_2)$ have horizontal distance $|x_1 - x_2| > 2e^H$, their neighbourhood balls are disjoint w.r.t. the connection rule of the truncated box model $G_{box,H}$. To see this, assume that $v_3 = (x_3, y_3)$ is adjacent to both v_1 and v_2 . Then $|x_1 - x_3| \leq e^{\frac{1}{2}(y_1+y_3)}$ and $|x_2 - x_3| \leq e^{\frac{1}{2}(y_2+y_3)}$. As $y_1, y_2, y_3 \leq H$, it follows that $|x_1 - x_3| \leq e^H$ and $|x_2 - x_3| \leq e^H$, which implies that $|x_1 - x_2| \leq |x_1 - x_3| + |x_2 - x_3| \leq 2e^H$, contradiction.

Due to the disjointedness of the neighbourhood balls and the properties of the Poisson process $h(V, v_1)$ and $h(V, v_2)$ are independent. The horizontal distance is uniform in $[0, \frac{\pi n}{2\nu}]$. Therefore, the probability that two vertices have horizontal distance $\leq 2e^H$ is $\leq \frac{4\nu e^H}{\pi n} = o(1)$. Writing d_H for the horizontal distance, we can continue the calculation of the second moment,

$$\begin{aligned} \mathbb{E}[Y_1] &= \iint_{\substack{\mathcal{R}_H \times \mathcal{R}_H: \\ |x_1 - x_2| > 2e^H}} \mathbb{E}[h(V, v_1)]\mathbb{E}[h(V, v_2)]\mu(dv_1)\mu(dv_2) \\ &\quad + \iint_{\substack{\mathcal{R}_H \times \mathcal{R}_H: \\ |x_1 - x_2| \leq 2e^H}} \mathbb{E}[h(V, v_1)]\mathbb{E}[h(V, v_2)]\mu(dv_1)\mu(dv_2) \\ &= (1 + o(1)) \iint_{\mathcal{R}_H \times \mathcal{R}_H} \mathbb{E}[h(V, v_1)]\mathbb{E}[h(V, v_2)]\mu(dv_1)\mu(dv_2) \\ &= (1 + o(1))(n\mathbb{E}X)^2. \end{aligned}$$

It follows that $\mathbb{E}[X^2] = (1 + o(1))\mathbb{E}[X]^2$, which implies $Var(X) \rightarrow 0$ using that $\mathbb{E}[X]$ tends to a constant (see above). We conclude

$$X \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \int_0^\infty P(y)\rho(y, k)\alpha e^{-\alpha y} dy.$$

In a similar way, by the Campbell-Mecke formula,

$$\mathbb{E}[Z] = \frac{1}{n} \int_{\mathcal{R}_H} \mathbb{P}(D_{G_{box,H}}(v) = k)\mu(dv) = \int_0^H \mathbb{P}(D_{G_{box,H}}(0, y) = k)\alpha e^{-\alpha y} dy.$$

By the theorem of dominated convergence, it follows that

$$\mathbb{E}[Z] \rightarrow \int_0^\infty \rho(y, k)\alpha e^{-\alpha y} dy.$$

The statement that $Var(Z) \rightarrow 0$ follows by the same steps as for X , just using $h(V, v) = \mathbb{1}_{\{D_{G_{box,H}}(v)=k\}}$ instead. Hence,

$$\mathbb{E}[Z] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \int_0^\infty \rho(y, k)\alpha e^{-\alpha y} dy.$$

Finally note that

$$\gamma(k) = \frac{\int_0^\infty P(y)\rho(y, k)\alpha e^{-\alpha y} dy}{\int_0^\infty \rho(y, k)\alpha e^{-\alpha y} dy}.$$

5.1.5 Putting everything together

Proof of Theorem 1.3: The overall statement follows from the lemmas together with the fact that convergence in probability is closed under addition. Write

$$c(k; G_n) - \gamma(k) = c(k; G_n) - c(k; G_{P_0}) + c(k; G_{P_0}) - c(k; G_{P_0, H}) + c(k; G_{P_0, H}) - \gamma(k).$$

The first difference converges in probability to zero due to the Poissonization Lemma 5.1, the second difference due to the truncation Lemma 5.4, then for the third difference we use $c(k; G_{P_{o,H}}) = c(k; G_{box,H})$ from Lemma 5.5 and then Lemma 5.6 on the derivation of the limit integral. ■

5.2 Overall clustering coefficient, proving Theorem 1.2

In this subsection, we want to infer that the clustering coefficient of the KPKVB random graph $c(G_n) \xrightarrow{\mathbb{P}} \gamma$ by using that $c(k; G_n) \xrightarrow{\mathbb{P}} \gamma(k)$ for fixed k .

Proof of Theorem 1.2: Let $\epsilon > 2\delta > 0$. As the p_k form the probability mass function of an \mathbb{N}_0 -valued random variable, we have that $\sum_{k=0}^{\infty} p_k = 1$, which implies that there is $K_\delta > 0$ such that $1 - \delta < \sum_{k=0}^{K_\delta} p_k \leq 1$. From Equations (20), (41) and (42), we know that $\gamma = \sum_{k=2}^{\infty} \gamma(k) p_k$. From this, it follows that $(1 - \delta)\gamma < \sum_{k=2}^{K_\delta} \gamma(k) p_k \leq 1$ and hence that $|\gamma - \sum_{k=2}^{K_\delta} \gamma(k) p_k| \leq \delta$. Write $p(k; G_n) = \frac{N_n(k)}{n}$ for the fraction of degree k vertices in the KPKVB model. From the definition of the clustering coefficient and function, it follows immediately that $c(G_n) = \sum_{k=2}^{\infty} c(k; G_n) p(k; G_n)$. By the initial observation, it follows again that $(1 - \delta)c(k; G_n) < \sum_{k=2}^{K_\delta} c(k; G_n) p(k; G_n) \leq 1$, and hence, $|c(G_n) - \sum_{k=2}^{K_\delta} c(k; G_n) p(k; G_n)| \leq \delta$. Putting these observations together:

$$\begin{aligned} |c(G_n) - \gamma| &\leq \left| c(G_n) - \sum_{k=2}^{K_\delta} c(k; G_n) p(k; G_n) \right| + \left| \sum_{k=2}^{K_\delta} c(k; G_n) p(k; G_n) - \sum_{k=2}^{K_\delta} \gamma(k) p_k \right| + \left| \sum_{k=2}^{K_\delta} \gamma(k) p_k - \gamma \right| \\ &\leq \delta + \left| \sum_{k=2}^{K_\delta} c(k; G_n) p(k; G_n) - \sum_{k=2}^{K_\delta} \gamma(k) p_k \right| + \delta. \end{aligned}$$

Therefore,

$$\mathbb{P}(|c(G_n) - \gamma| \geq \epsilon) \leq \mathbb{P}\left(\left| \sum_{k=2}^{K_\delta} c(k; G_n) p(k; G_n) - \sum_{k=2}^{K_\delta} \gamma(k) p_k \right| \geq \epsilon - 2\delta\right) \rightarrow 0,$$

because $c(k; G_n) \xrightarrow{\mathbb{P}} \gamma(k)$ for $k = 2, \dots, K_\delta$ and $p(k; G_n) \xrightarrow{\mathbb{P}} p_k$ together imply that

$$\sum_{k=2}^{K_\delta} c(k; G_n) p(k; G_n) \xrightarrow{\mathbb{P}} \sum_{k=2}^{K_\delta} \gamma(k) p_k.$$

■

6 Overview of the proof strategy for $k \rightarrow \infty$

The proof of Theorem 1.4 follows the same strategy as outlined in Section 2 and executed in Section 5. However, the fact that $k = k_n \rightarrow \infty$ as $n \rightarrow \infty$, introduces significant technical challenges, especially for k_n close to the maximum scale $n^{\frac{1}{2\alpha+1}}$. For example, the coupling between G_{P_o} and G_{box} we use becomes less exact so that we can no longer use Lemma 2.3 to conclude that triangle counts in G_{P_o} and G_{box} are asymptotically equivalent. Moreover, since we are ultimately interested in recovering the scaling of $c(k_n; G_n)$, which Theorem 1.4 claims is $\gamma(k_n)$, we need to show that each step in the strategy outlined in Section 2 only introduces error terms that are of smaller order, i.e. that are $o(\gamma(k_n))$. This will turn out to require a great deal of care in bounding all error terms we encounter.

In this section we explain the challenges associated with each step and give a detailed overview of the structure for the proof of Theorem 1.4 using intermediate results for each of the steps. To this end we define the scaling function

$$s(k) = \begin{cases} k^{-(4\alpha-2)} & \text{if } \frac{1}{2} < \alpha < \frac{3}{4}, \\ \log(k) k^{-1} & \text{if } \alpha = \frac{3}{4}, \\ k^{-1} & \text{if } \alpha > \frac{3}{4}, \end{cases} \quad (63)$$

so that $\gamma(k) = \Theta(s(k))$ as $k \rightarrow \infty$. We will end this section with the proof of Theorem 1.4, based on the intermediate results.

Remark 6.1 (Diverging k_n). *Throughout the remainder of this paper, unless stated otherwise, $\{k_n\}_{n \geq 1}$ will always denote a sequence of non-negative integers satisfying $k_n \rightarrow \infty$ and $k_n = o\left(n^{\frac{1}{2\alpha+1}}\right)$, as $n \rightarrow \infty$.*

We start with introducing a slightly modified version of the local clustering function, which will be convenient for computations later,

$$c^*(k; G) = \frac{1}{\mathbb{E}[N_k]} \sum_{\substack{v \in V(G) \\ \deg(v)=k}} c(v). \quad (64)$$

Notice that the only difference between $c(k; G)$ and $c^*(k; G)$ is that we replace N_k by its expectation $\mathbb{E}[N_k]$. The advantage is that now, the only randomness is in the formation of triangles. In addition, note that since $\mathbb{E}[N_k] > 0$ a case distinction for N_k is no longer needed for $c^*(k; G)$. It is however still relevant since we are eventually interested in $c(k; G)$. Following the notational convention, throughout the remainder of this paper we write $c^*(k; G_{\text{Po}})$ and $c^*(k; G_{\text{box}})$ to denote the modified local clustering function in G_{Po} and $G_{\mathcal{P},n}(\alpha, \nu)$, respectively.

Figure 6 shows a schematic overview of the proof of Theorem 1.4 based on the different propositions described below, plus the sections in which these propositions are proved. Observe that the order in which the intermediate results are proved is reversed with respect to the natural order of reasoning. This does not create any circular logic, since each intermediate result is independent of the others. We choose this order because results proved in the later stages are helpful to deal with error terms coming up in proofs at earlier stages and hence help streamline those proofs. Below briefly describe each of the intermediate steps leading up to the proof of Theorem 1.4. We start with an observation on the dependence between the degree of a point $p = (x, y)$ and its height y .

6.1 Adjusted clustering and the Poissonized KPKVB model

Recall that the first step for the fixed k case was to show that the transition from the KPKVB graph $G_n = G(n; \alpha, \nu)$ to the Poissonized version G_{Po} did not influence clustering. Here we first make a transition from the local clustering function $c(k_n; G_n)$ to the adjusted version $c^*(k_n; G_n)$. The following lemma justifies working with this modified version. The proof uses a concentration result for $N_n(k_n)$ and full details can be found in Section 9.3.

Lemma 6.1. *As $n \rightarrow \infty$,*

$$\mathbb{E}[|c(k_n; G_n) - c^*(k_n; G_n)|] = o(s(k_n)).$$

We then establish that the modified local clustering function for KPKVB graphs G_n behaves similarly to that in the G_{Po} . The proof, found in Section 9.3, is based on a standard coupling between a Binomial Point Process and Poisson Point Process.

Proposition 6.2. *As $n \rightarrow \infty$,*

$$\mathbb{E}[|c^*(k_n; G_n) - c^*(k_n; G_{\text{Po}})|] = o(s(k_n)).$$

6.2 Coupling of local clustering between G_{Po} and G_{box}

The next step is to show that the modified clustering is preserved under the coupling described in Section 2.4. The proof can be found in Section 9.2. This step is one of the key technical challenges we face in proving Theorem 1.4.

To understand why, recall that the degree k of a node is related to its height y , roughly speaking, by $k \approx \xi e^{y/2}$. Therefore, when k is fixed we have that the heights of nodes with that




Figure 6: Overview of the proof strategy for Theorem 1.3. The left column denote the models in which the true hyperbolic balls are used while the right column contains the models that use an approximation of these. The most important part is the transition between these to setting which is accomplished by Proposition 6.3.

degree are also fixed, in particular $y < R/4$ for large enough n . In addition, the main contribution of triangles would also come from nodes with heights $y' < R/4$. This allowed us to use Lemma 2.3 and conclude that the triangles present in the graph G_{Po} were exactly those present in G_{box} and therefore the local clustering function was the same in both models. When $k_n \rightarrow \infty$ this is no longer true in general. For instance, suppose $k_n = n^{\frac{1-\varepsilon}{2\alpha+1}}$, for some small $0 < \varepsilon < 1$. Then the relation $k_n \approx \xi e^{y_n/2}$ implies that $y_n \approx \frac{2(1-\varepsilon)}{2\alpha+1} \log(n) - 2 \log(\xi)$. Since $R/4 = \frac{1}{2} \log(n) - \frac{1}{2} \log(\nu)$ we get that $R/4 = o(y_n)$ for all $\alpha > (3 - 4\varepsilon)/2$ and hence $y_n > R/4$ for large enough n , violating the conditions of Lemma 2.3. However, by carefully analyzing the difference between the adjusted local clustering function in both models we can still make the same conclusion. This is summarized in the following proposition whose proof is found in Section 9.2.

Proposition 6.3 (Coupling result for adjusted clustering function). *As $n \rightarrow \infty$,*

$$\mathbb{E}[|c^*(k_n; G_{Po}) - c^*(k_n; G_{box})|] = o(s(k_n)).$$

Together, the three results described so far imply that the difference between the clustering function for a KPKVB graph and the adjusted clustering function for the finite box graph G_{box} converges to zero faster than the proposed scaling $\gamma(k_n)$ in Theorem 1.4. Hence, to prove this theorem it is enough to prove it for $c^*(k; G_{box})$.

6.3 From the finite box to the infinite model

To compute the limit of the adjusted clustering function $c^*(k; G_{\text{box}})$ we first prove in Section 8 that it is concentrated around its mean $\mathbb{E}[c^*(k_n; G_{\text{box}})]$.

Proposition 6.4 (Concentration for adjusted clustering function in G_{box}). *As $n \rightarrow \infty$,*

$$\mathbb{E}[|c^*(k_n; G_{\text{box}}) - \mathbb{E}[c^*(k_n; G_{\text{box}})]|] = o(s(k_n)).$$

This result represents another technical challenge we face when considering $k_n \rightarrow \infty$. For the proof, we first identify the specific range of heights that give the main contribution to the triangle count, showing that the triangles coming from nodes with heights outside this range is of smaller order. Then we prove a concentration result for the main term, by carefully analyzing the joint neighbourhoods of two nodes whose heights fall into the identified range. The full details are found in Section 8.

Assuming this concentration result, we are left to compute expectation $\mathbb{E}[c^*(k_n; G_{\text{box}})]$ and show that it is asymptotically equivalent to $\gamma(k_n)$ as $n \rightarrow \infty$. To accomplish this we move to the infinite limit model G_∞ and show that the difference between the expected value of $c^*(k; G_{\text{box}})$ and $\gamma(k_n)$ goes to zero faster than the proposed scaling in Theorem 1.3.

Proposition 6.5 (Transition to the infinite limit model). *As $n \rightarrow \infty$,*

$$|\mathbb{E}[c^*(k_n; G_{\text{box}})] - \gamma(k_n)| = o(s(k_n)).$$

Recall that for the finite box model the left and right boundaries of \mathcal{R}_n were identified, so that graph G_{box} contains some additional edge with respect to the induced subgraph of G_∞ on \mathcal{R}_n . The proof of Proposition 6.5 therefore relies on analyzing the number of triangles coming from these additional edges and showing that their contribution to the clustering function are of negligible order, see Section 7.

Remark 6.2 (Notations for different graphs). *We will use the subscripts n , Po, box and ∞ to identify properties of, respectively, the KPKVB mode G_n , the Poisson version G_{Po} , the finite box model G_{box} and the infinite model G_∞ . For example $N_{\text{Po}}(k)$ denotes number of nodes with degree k in G_{Po} and $\rho_{\text{box}}(y, k) = \mathbb{P}(\text{Po}(\mu(\mathcal{B}_{\text{box}}(y))) = k)$, i.e. the degree distribution in G_{box} for a point $p = (x, y)$.*

6.4 Proof of the main results

We are now ready to prove Theorem 1.4, using the results stated in the previous sections.

Proof of Theorem 1.4. Note that the second of the theorem follows immediately from the first.

To prove the first statement, we rewrite $c(k_n; G_n)$ as

$$\begin{aligned} c(k_n; G_n) - \gamma(k_n) &= (c(k_n; G_n) - c^*(k_n; G_n)) + (c^*(k_n; G_n) - c^*(k_n; G_{\text{Po}})) \\ &\quad + (c^*(k_n; G_{\text{Po}}) - c^*(k_n; G_{\text{box}})) + (c^*(k_n; G_{\text{box}}) - \mathbb{E}c^*(k_n; G_{\text{box}})) \\ &\quad + \mathbb{E}c^*(k_n; G_{\text{box}}) - \gamma(k_n) \end{aligned}$$

Then, we take absolute values and apply the triangle inequality. By monotonicity of expectation, we can apply it to both sides and obtain

$$\begin{aligned} \mathbb{E}[|c(k_n; G_n) - \gamma(k_n)|] &\leq \mathbb{E}[|c(k_n; G_n) - c^*(k_n; G_n)|] + \mathbb{E}[|c^*(k_n; G_n) - c^*(k_n; G_{\text{Po}})|] \\ &\quad + \mathbb{E}[|c^*(k_n; G_{\text{Po}}) - c^*(k_n; G_{\text{box}})|] + \mathbb{E}[|c^*(k_n; G_{\text{box}}) - \mathbb{E}c^*(k_n; G_{\text{box}})|] \\ &\quad + |\mathbb{E}[c^*(k_n; G_{\text{box}})] - \gamma(k_n)| \end{aligned}$$

At this point, the lemmas and propositions presented above in this section can be applied in order to show that all summands are $o(\gamma(k_n))$: Lemma 6.1 for the transition to the modified clustering function in the first term, Proposition 6.2 for the Poissonization in the second term,

Proposition 6.3 for the coupling between the Poissonized KPKVB and the finite box model in the third term, Proposition 6.4 for the concentration in the fourth term and finally Proposition 6.5 for the transition to the infinite limit model.

All of this together yields that:

$$\mathbb{E}[|c(k_n; G_n) - \gamma(k_n)|] = o(s(k_n)) = o(\gamma(k_n)),$$

which establishes the first statement of the theorem and finishes the proof. \square

7 From G_{box} to G_∞ (Proving Proposition 6.5)

In this section we shall relate the clustering in the finite box model G_{box} to that of the infinite model. The main goal is to prove Proposition 6.5 which states that

$$|\mathbb{E}[c^*(k_n; G_{\text{box}})] - \gamma(k_n)| = o(s(k_n)).$$

Recall that G_{box} is obtained by restricting the Poisson point process \mathcal{P} to the box $\mathcal{R} = (-I_n, I_n] \times (0, R]$, with $I_n = \frac{\pi}{2}e^{R/2}$ and connecting two points $p_1, p_2 \in \mathcal{R}$ if and only if $|x_1 - x_2|_{\pi e^{R/2}} \leq e^{(y_1 + y_2)/2}$. We also recall that by definition of the norm $|\cdot|_{\pi e^{R/2}}$ the left and right boundaries of \mathcal{R} are identified. See Section 2.2 for more details. Due to this identification of the boundaries some triples of nodes that form a triangle in the finite box model do not form a triangle in the infinite model. Therefore, to establish the required result we need to compute the asymptotic difference between triangle counts in both models. To keep notation concise we write $|\cdot|_n$ for the norm $|\cdot|_{\pi e^{R/2}}$.

For any $p \in \mathbb{R} \times \mathbb{R}_+$ we define for the finite box model,

$$T_{\text{box}}(p) = \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus \{p\}}} T_{\text{box}}(p, p_1, p_2)$$

where the sum is over all distinct pairs in $\mathcal{P} \setminus p$ and

$$T_{\text{box}}(p, p_1, p_2) = \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{box}}(p)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{box}}(p)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{box}}(p_1)\}}.$$

Similarly, for the infinite model we define

$$T_\infty(y) = \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus (0, y)}} T_\infty(y, p_1, p_2),$$

where

$$T_\infty(y, p_1, p_2) = \mathbb{1}_{\{p_1 \in \mathcal{B}_\infty(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_\infty(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_\infty(p_1)\}}.$$

Here we slightly abuse notation and write $\mathcal{B}_\infty(y)$ for $\mathcal{B}_\infty((0, y))$. We will adopt this notational convention throughout the remainder of this section, to keep notation concise. We also recall that $N_{\text{box}}(k)$ denotes the number of vertices with degree k in G_{box} .

We will first relate $\gamma(k_n)$ to an integral expression involving $T_\infty(y)$ and $\mathbb{E}[c^*(k_n; G_{\text{box}})]$ to one involving $\mathbb{E}[T_{\text{box}}(y)]$. Recall the definition of $y_{k,C}^\pm$ from (15) and the interval $\mathcal{K}_C(k_n) = [y_{k_n,C}^-, y_{k_n,C}^+]$. Note that for any $y \in \mathcal{K}_C(k_n)$ it holds that as

$$\frac{k_n - C\sqrt{k_n \log(k_n)}}{\xi} \leq e^{\frac{y}{2}} \leq \frac{k_n + C\sqrt{k_n \log(k_n)}}{\xi} \quad (65)$$

and thus $k_n - C\sqrt{k_n \log(k_n)} \leq \mu(y) \leq k_n + C\sqrt{k_n \log(k_n)}$.

Lemma 7.1. *Let $\gamma(k_n)$ be defined as in (41). Then as $n \rightarrow \infty$*

$$\gamma(k_n) = (1 + o(1)) \frac{1}{k_n^2 p_{k_n}} \int_{\mathcal{K}_C(k_n)} \mathbb{E}[T_\infty(y)] \rho(y, k) \alpha e^{-\alpha y} dy. \quad (66)$$

Moreover,

$$\mathbb{E}[c^*(k_n; G_{\text{box}})] = (1 + o(1)) \frac{1}{k_n^2 p_{k_n}} \int_{\mathcal{K}_C(k_n)} \mathbb{E}[T_{\text{box}}(y)] \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy \quad (67)$$

as $n \rightarrow \infty$,

Proof. Recall that

$$P(y) = \mathbb{E}[\mathbb{1}_{\{u_1 \in \mathcal{B}_\infty(u_2)\}}],$$

where u_1 and u_2 are independent and distributed according to the probability density $\mu(\mathcal{B}_\infty(y))^{-1} \mathbb{1}_{\{u_i \in \mathcal{B}_\infty(y)\}} f(x_i, y_i)$. It then follows from the Campbell-Mecke formula that

$$\begin{aligned} \mathbb{E}[T_\infty(y)] &= \int \mathbb{1}_{\{p_1 \in \mathcal{B}_\infty(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_\infty(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_\infty(p_1)\}} f(x_1, y_1) f(x_2, y_2) dx_1 dx_2 dy_1 dy_2 \\ &= \mu(\mathcal{B}_\infty(y))^2 P(y). \end{aligned}$$

It then follows that,

$$\gamma(k_n) = \frac{1}{p_{k_n}} \cdot \int_0^\infty P(y) \rho(y, k) \alpha e^{-\alpha y} dy = \frac{1}{p_{k_n}} \int_0^\infty \mathbb{E}[T_\infty(y)] \mu(\mathcal{B}_\infty(y))^{-2} \rho(y, k) \alpha e^{-\alpha y} dy.$$

Because $\mathbb{E}[T_\infty(y)] \mu(\mathcal{B}_\infty(y))^{-2} = O(1)$, uniformly in $y \in \mathbb{R}_+$ and $\mu(y) = (1 + o(1))k_n$ uniformly for $y \in \mathcal{K}_C(k_n)$, by the concentration of heights (Proposition 2.4)

$$\begin{aligned} &\frac{1}{p_{k_n}} \int_0^\infty \mathbb{E}[T_\infty(y)] \mu(\mathcal{B}_\infty(y))^{-2} \rho(y, k) \alpha e^{-\alpha y} dy \\ &= (1 + o(1)) \frac{1}{p_{k_n}} \int_{\mathcal{K}_C(k_n)} \mathbb{E}[T_\infty(y)] \mu(\mathcal{B}_\infty(y))^{-2} \rho(y, k) \alpha e^{-\alpha y} dy \\ &= (1 + o(1)) \frac{1}{k_n^2 p_{k_n}} \int_{\mathcal{K}_C(k_n)} \mathbb{E}[T_\infty(y)] \rho(y, k) \alpha e^{-\alpha y} dy. \end{aligned}$$

For (67) we recall that

$$c^*(k_n; G_{\text{box}}) = \frac{1}{\mathbb{E}[N_{\text{box}}(k_n)]} \sum_{p \in \mathcal{P}} c_{\text{box}}(p) \mathbb{1}_{\{D_{G_{\text{box}}}(p) = k_n\}},$$

where $c_{\text{box}}(p)$ can be expressed as

$$c_{\text{box}}(p) = \frac{1}{\binom{D_{G_{\text{box}}}(p)}{2}} \sum_{p_1, p_2 \in \mathcal{P} \setminus p}^{\neq} T_{\text{box}}(p, p_1, p_2) = \frac{T_{\text{box}}(p)}{\binom{D_{G_{\text{box}}}(p)}{2}}.$$

By the Campbell-Mecke formula

$$\begin{aligned} \mathbb{E}[c^*(k_n; G_{\text{box}})] &= \frac{1}{\mathbb{E}[N_{\text{box}}(k_n)]} \int_{\mathcal{R}} \mathbb{E}[c_{\text{box}}(p) \mathbb{1}_{\{D_{G_{\text{box}}}(p) = k_n\}}] f(x, y) dx dy \\ &= \frac{1}{\mathbb{E}[N_{\text{box}}(k_n)]} \int_{\mathcal{R}} \mathbb{E}[c_{\text{box}}(p) | D_{G_{\text{box}}}(p) = k_n] \rho_{\text{box}}(p, k_n) f(x, y) dx dy \\ &= (1 + o(1)) \frac{n}{\mathbb{E}[N_{\text{box}}(k_n)]} \int_{\mathcal{K}_C(k_n)} \mathbb{E}[c_{\text{box}}(y) | D_{G_{\text{box}}}(y) = k_n] \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy, \end{aligned}$$

where the last line follows from the concentration of heights, for which we used the upper bound $\mathbb{E}[c_{\text{box}}(y) | D_{G_{\text{box}}}(y) = k_n] \leq 1$.

To analyze the conditional expectation we observe that, similar to the analysis of $\gamma(k_n)$, conditioned on there being k_n points in $\mathcal{B}_{\text{box}}(y)$, each point $u_i = (x_i, y_i)$ is independently distributed according to the probability density $\mu(\mathcal{B}_{\text{box}}(y))^{-1} \mathbb{1}_{\{u_i \in \mathcal{B}_{\text{box}}(y)\}} f(x_i, y_i)$. Therefore,

$$\begin{aligned} \mathbb{E}[c_{\text{box}}(y) | D_{G_{\text{box}}}(y) = k_n] &= \binom{k_n}{2}^{-1} \mathbb{E} \left[\sum_{1 \leq i < j \leq k_n} \mathbb{1}_{\{u_i \in \mathcal{B}_{\text{box}}(u_j)\}} \right] \\ &= \mathbb{E}[u_1 \in \mathcal{B}_{\text{box}}(u_2)] \\ &= \mu(\mathcal{B}_{\text{box}}(y))^{-2} \iint T_{\text{box}}(y, p_1, p_2) f(x_1, y_1) f(x_2, y_2) dx_1 dy_1 dx_2 dy_2 \\ &= \mu(\mathcal{B}_{\text{box}}(y))^{-2} \mathbb{E}[T_{\text{box}}(y)]. \end{aligned}$$

and thus, by applying a concentration of heights argument on $\mu(\mathcal{B}_{\text{box}}(y))^{-2}$,

$$\mathbb{E}[c^*(k_n; G_{\text{box}})] = (1 + o(1)) \frac{n\mu(\mathcal{B}_{\text{box}}(2\log(k_n/\xi)))^{-2}}{\mathbb{E}[N_{\text{box}}(k_n)]} \int_{\mathcal{K}_C(k_n)} \mathbb{E}[T_{\text{box}}(y)] \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy.$$

To finish the argument, we first note that $\mu(\mathcal{B}_{\text{box}}(2\log(k_n/\xi)))^{-2} = (1 + o(1))k_n^2$, while

$$\mathbb{E}[N_{\text{box}}(k_n)] = (1 + o(1))np_{k_n}$$

by Lemma 3.2. We therefore conclude that

$$\mathbb{E}[c^*(k_n; G_{\text{box}})] = (1 + o(1)) \frac{1}{k_n^2 p_{k_n}} \int_{\mathcal{K}_C(k_n)} \mathbb{E}[T_{\text{box}}(y)] \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy.$$

□

As a result of this lemma we only need to compare the difference in triangles between both models for height in the interval $\mathcal{K}_C(k_n)$. This will significantly help the analysis.

7.1 Comparing triangles between G_∞ and G_{box}

To analyze $T_{\text{box}}(y_0) - T_\infty(y_0)$ we first reiterate that the difference between the indicator $\mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{box}}(p)\}}$ in the finite box model and $\mathbb{1}_{\{p_1 \in \mathcal{B}_\infty(p)\}}$ is that in G_{box} we identified the boundaries of the interval $[-\frac{\pi}{2}e^{R/2}, \frac{\pi}{2}e^{R/2}]$ and we stop at height $y = R$. This induces a difference in triangle counts between both models. To see this, note that for any $p = (x, y)$ with $0 \leq y \leq R$ we have that $\mathcal{B}_{\text{box}}(p) = \mathcal{B}_\infty(p) \cap \mathcal{R}$. This means that if $p', p_2 \in \mathcal{B}_{\text{box}}(p)$ and $p_2 \in \mathcal{B}_\infty(p') \cap \mathcal{R}$ then $p_2 \in \mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')$ and hence (p, p', p_2) form a triangle both in G_{box} and G_∞ . However, it could happen that there are points in the intersection $\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')$ that are not in $\mathcal{B}_\infty(p) \cap \mathcal{B}_\infty(p')$. Let us denote this region by $\mathcal{T}(p, p')$, see Figure 7 for an example of this region. Then, any $p_2 \in \mathcal{T}(p, p')$ creates a triangle with p and p' in G_{box} that is not present in G_∞ . Finally, any point $p_2 \in \mathcal{B}_\infty(p) \cap \mathcal{B}_\infty(p')$ with height $y_2 > R$ creates a triangle with p, p' in G_∞ but not in G_{box} .

Let us now define the following triangle count function

$$\tilde{T}_{\text{box}}(p_0) = \sum_{(p_1, p_2) \in \mathcal{P} \setminus \{p_0\}}^{\neq} \tilde{T}_{\text{box}}(p_0, p_1, p_2).$$

where

$$\tilde{T}_{\text{box}}(p_0, p_1, p_2) = \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{box}}(p_0)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{box}}(p_0)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_\infty(p_1) \cap \mathcal{R}\}}.$$

Then $\tilde{T}_{\text{box}}(p_0)$ only counts those triangles attached to p_0 that exist in both G_{box} and G_∞ and thus, by definition of the region $\mathcal{T}(p_0, p_1)$,

$$T_{\text{box}}(p_0) - \tilde{T}_{\text{box}}(p_0) = \sum_{p_1, p_2 \in \mathcal{P} \setminus \{p_0\}}^{\neq} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{box}}(p_0)\}} \mathbb{1}_{\{p_2 \in \mathcal{T}(p_0, p_1)\}}.$$

The next result, which is crucial for the proof of Proposition 6.5, computes the expected measure of $\mathcal{T}(p, p')$ with respect to p' .

Lemma 7.2. *Let $p_0 = (0, y)$ with $y \in \mathcal{K}_C(k_n)$. Then as $n \rightarrow \infty$,*

$$\mathbb{E} \left[\left| T_{\text{box}}(p_0) - \tilde{T}_{\text{box}}(p_0) \right| \right] = y O \left(n^{-(2\alpha-1)} \right) + e^y O \left(n^{-(4\alpha-2)} \right).$$

The proof of the lemma is not difficult but cumbersome, since it involves computing many different integrals. We postpone this proof till the end of this section and proceed with the main goal, proving Proposition 6.5. First we state a small lemma about the scaling of $s(k_n)$ that will be very useful.

Lemma 7.3. *Let $s(k_n)$ be as defined in (63). Then for any $k_n = o \left(n^{\frac{1}{2\alpha+1}} \right)$, as $n \rightarrow \infty$,*

$$n^{-(2\alpha-1)} = o \left(s(k_n) \right).$$

Proof. First let $\frac{1}{2} < \alpha < \frac{3}{4}$. Then

$$n^{-(2\alpha-1)} s(k_n)^{-1} = n^{-(2\alpha-1)} k_n^{4\alpha-2} = o \left(n^{-(2\alpha-1) + \frac{4\alpha-2}{2\alpha+1}} \right) = o \left(n^{-\frac{4\alpha^2-4\alpha+1}{2\alpha+1}} \right) = o(1),$$

since $4\alpha^2 - 4\alpha + 1 > 0$ for all $\alpha > \frac{1}{2}$. Similarly, for $\alpha \geq \frac{3}{4}$ we have that $4\alpha^2 > 2$ and hence,

$$n^{-(2\alpha-1)} s_\alpha(k_n) = o \left(n^{-(2\alpha-1)} k_n \right) = o \left(n^{-\frac{4\alpha^2-2}{2\alpha+1}} \right) = o(1).$$

□

The first key implication of Lemma 7.2 is that the triangle count in the finite box model is equivalent to $k_n^2 P(y)$, where $P(y)$ is defined in [??].

Lemma 7.4. *Let $p_0 = (0, y)$. Then uniformly for $y \in \mathcal{K}_C(k_n)$,*

$$\mathbb{E} [T_{\text{box}}(p_0)] = (1 + o(1)) k_n^2 P(y) + o \left(s(k_n) k_n^2 \right),$$

as $n \rightarrow \infty$.

Proof. Recall that $\mathbb{E} [T_\infty(y)] = \mu(y)^2 P(y) = (1 + o(1)) k_n^2 P(y)$ on $\mathcal{K}_C(k_n)$. We will show that

$$\mathbb{E} [|\mathbb{E} [T_{\text{box}}(p_0)] - T_\infty(y)|] = o \left(s(k_n) k_n^2 \right),$$

which implies the result.

Define $\mathcal{R}' := (\mathbb{R} \times \mathbb{R}_+) \setminus \mathcal{R}$. Then we can write

$$\begin{aligned} |T_{\text{box}}(p_0) - T_\infty(y)| &\leq \left| T_{\text{box}}(p_0) - \tilde{T}_{\text{box}}(p_0) \right| + \left| \tilde{T}_{\text{box}}(p_0) - T_\infty(y) \right| \\ &= \left| T_{\text{box}}(p_0) - \tilde{T}_{\text{box}}(p_0) \right| + \sum_{p_1, p_2 \in \mathcal{P} \cap \mathcal{R}'}^{\neq} T_\infty(p_0, p_1, p_2). \end{aligned}$$

Then by the Campbell-Mecke formula

$$|\mathbb{E} [T_{\text{box}}(p_0) - T_\infty(p_0)]| \leq \mathbb{E} \left[\left| T_{\text{box}}(p_0) - \tilde{T}_{\text{box}}(p_0) \right| \right]$$

$$+ \int_{\mathcal{R}'} \int_{\mathcal{R}'} T_{\infty}(p_0, p_1, p_2) f(x_1, y_1) f(x_2, y_2) dx_2 dy_2 dx_1 dy_1.$$

The first part is taken care of by Lemma 7.2. For the other integral we have

$$\begin{aligned} & \iint_{\mathcal{R}'} T_{\infty}(p_0, p_1, p_2) f(x_1, y_1) f(x_2, y_2) dx_2 dy_2 dx_1 dy_1 \\ & \leq \left(\int_{\mathcal{R}'} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\infty}(p_0)\}} f(x_1, y_1) dx_1 dy_1 \right)^2 = O \left(\left(e^{y/2} \int_R^{\infty} e^{-(\alpha - \frac{1}{2})y_1} dy_1 \right)^2 \right) \\ & = O \left(e^y e^{-(2\alpha-1)R} \right) = O \left(e^y n^{-(4\alpha-2)} \right). \end{aligned}$$

Thus we conclude, using Lemma 7.2, that,

$$|\mathbb{E}[T_{\text{box}}(p_0) - T_{\infty}(p_0)]| = O \left(y n^{-(2\alpha-1)} + n^{-(4\alpha-2)} e^y \right). \quad (68)$$

Therefore, on $\mathcal{K}_C(k_n)$,

$$\begin{aligned} |\mathbb{E}[T_{\text{box}}(p_0) - T_{\infty}(p_0)]| &= O \left(\log(k_n) n^{-(2\alpha-1)} + k_n^2 n^{-(4\alpha-2)} \right) \\ &= O \left(\log(k_n) n^{-(2\alpha-1)} + k_n^2 n^{-(4\alpha-2)} \right) = o \left(s(k_n) k_n^2 \right), \end{aligned}$$

where the last part follows from Lemma 7.3 and the fact that $s(k_n)^2 = o(s(k_n))$. \square

We can now prove the main result of this section.

Proof of Proposition 6.5. First, by Lemma 7.4 and (67) we have

$$\begin{aligned} \mathbb{E}[c^*(k_n; G_{\text{box}})] &= (1 + o(1)) \frac{1}{p_{k_n}} \int_{\mathcal{K}_C(k_n)} P(y) \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy \\ &\quad + o(s(k_n)) \frac{1}{p_{k_n}} \int_{\mathcal{K}_C(k_n)} P(y) \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy. \end{aligned}$$

By Lemma 3.3 the integral in the second term is $(1 + o(1))p_{k_n}$ and thus the second term is $o(s(k_n)) = o(\gamma(k_n))$. Hence it remain to prove that the first term is $(1 + o(1))\gamma(y)$. Using (66) it is enough to show that

$$\int_{\mathcal{K}_C(k_n)} P(y) \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy = (1 + o(1)) \int_{\mathcal{K}_C(k_n)} P(y) \rho(y, k_n) \alpha e^{-\alpha y} dy.$$

Now recall the substitution of variables from the proof of Lemma 3.3: $z(y) = 2 \log(\mu_{\text{box}}(y)/\xi)$. We will apply this change of variables to the right hand side in the above equation. This yields

$$\begin{aligned} & \int_{\mathcal{K}_C(k_n)} P(y) \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy \\ &= (1 + o(1)) \int_{\mathcal{K}_C(k_n)} P(z(y) - 2 \log(1 + o(1))) \rho(z(y), k_n) z'(y) \alpha e^{-\alpha z(y)} dy \\ &= (1 + o(1)) \int_{\mathcal{K}_C(k_n)} P(z - 2 \log(1 + o(1))) \rho(z, k_n) \alpha e^{-\alpha z} dz. \end{aligned}$$

Next we recall that by Proposition 4.7

$$P(y) \sim \begin{cases} e^{-\frac{y}{2}(4\alpha-2)} c_{\alpha} \xi^{4\alpha-2} & \text{if } \frac{1}{2} < \alpha < \frac{3}{4} \\ \frac{y}{2} e^{-\frac{y}{2}} & \text{if } \alpha = \frac{3}{4} \\ e^{-\frac{y}{2} \frac{\alpha-\frac{1}{2}}{\alpha-\frac{3}{4}}} & \text{if } \alpha > \frac{3}{4}. \end{cases}$$




Figure 7: Example configuration of two points p and p' for which $\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')$ is not a subset of $\mathcal{B}_{\infty}(p) \cap \mathcal{B}_{\infty}(p')$. The red region indicates the area belonging to $\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')$ but not to $\mathcal{B}_{\infty}(p) \cap \mathcal{B}_{\infty}(p')$.

In particular, this implies that $P(z - 2\log(1 + o(1))) = (1 + o(1))P(z)$, uniformly on $\mathcal{K}_C(k_n)$. We therefore conclude that

$$\int_{\mathcal{K}_C(k_n)} P(y) \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy = (1 + o(1)) \int_{\mathcal{K}_C(k_n)} P(z) \rho(z, k_n) \alpha e^{-\alpha z} dz,$$

which finishes the proof. \square

From the proof of Proposition 6.5 we immediately obtain the following useful corollary, which will be used in Section 8.

Corollary 7.5. *Let $p_0 = (0, y)$. Then, as $n \rightarrow \infty$,*

$$\int_{-I_n}^{I_n} \int_{\mathcal{K}_C(k_n)} \rho_{\text{box}}(y, k_n) \mathbb{E} [\tilde{T}_{\text{box}}(p_0)] f(x, y) dx dy = (1 + o(1)) n k_n^2 \int_0^\infty P(y) \rho(y, k_n) \alpha e^{-\alpha y} dy.$$

In particular,

$$\int_{\mathcal{K}_C(k_n)} \rho_{\text{box}}(y, k_n) \mathbb{E} [\tilde{T}_{\text{box}}(p_0)] f(x, y) dx dy = \Theta \left(n k_n^{-(2\alpha-1)} s(k_n) \right).$$

7.2 Counting missing triangles

We now come back to computing the expected number of triangles attached to a node at height y in G_{box} that are not present in G_{∞} .

Recall that $\mathcal{T}(p, p')$ denotes the region of points which form triangles with p and p' in G_{box} but not in G_{∞} . Figure 7 shows an example of a configuration where $\mathcal{T}(p, p') \neq \emptyset$. We observe that $\mathcal{T}(p, p') \neq \emptyset$ because the right boundary of the ball $\mathcal{B}_{\text{box}}(p')$ exits the right boundary of the box \mathcal{R} and then, since we identified the boundaries, continues from the left so that $\mathcal{B}_{\text{box}}(p')$ covers part of the ball $\mathcal{B}_{\text{box}}(p)$ which would not be covered in the infinite limit model.

To further analyze this, let us introduce some notation. For any $p = (x, y) \in \mathcal{R}$ we will define the left and right boundary functions as, respectively,

$$b_p^-(z) = \begin{cases} 2 \log(x - z) - y & \text{if } -\frac{\pi}{2} e^{R/2} \leq z \leq x - e^{y/2} \\ 2 \log(\pi e^{R/2} + x - z) - y & \text{if } x - e^{(y+R)/2} + \pi e^{R/2} \leq z \leq \frac{\pi}{2} e^{R/2} \\ 0 & \text{otherwise} \end{cases} \quad (69)$$

$$b_p^+(z) = \begin{cases} 2 \log(z - x) - y & \text{if } x + e^{y/2} \leq z \leq \frac{\pi}{2} e^{R/2} \\ 2 \log(\pi e^{R/2} + z - x) - y & \text{if } -\frac{\pi}{2} e^{R/2} \leq z \leq x + e^{(y+R)/2} - \pi e^{R/2} \\ 0 & \text{otherwise} \end{cases} \quad (70)$$

Note that these functions describe the boundaries of the ball $\mathcal{B}_{\text{box}}(p)$. In particular, $p' = (x', y') \in \mathcal{B}_{\text{box}}(p)$ if and only if $y' \geq \min \{b_p^-(x'), b_p^+(x')\}$.

Since we have identified the left and right boundary of \mathcal{R} we can assume, without loss of generality that $x = 0$. Due to symmetry it is then enough to restrict the analysis to the case where $x' > 0$. For this case there are two important points in the box \mathcal{R} . These are the intersection between the left boundary of p' and the right boundary of p' , as it continues from the left side of the box, and the left boundary of p . We denote by $(x^*(p'), y^*(p'))$ the intersection between the left and right boundary of p' and by $(\hat{x}(p, p'), \hat{y}(p, p'))$ the intersection between the left boundary of p and the right boundary of p' , see Figure 7.

Let us derive the expressions for the coordinates of these two points, starting with $(x^*(p'), y^*(p'))$. The x -coordinate $x^*(p')$ is the solution to the equation $b_{p'}^+(z) = b_p^-(z)$ for $-\frac{\pi}{2}e^{R/2} \leq z \leq x + e^{(y+R)/2} - \pi e^{R/2}$. This equation becomes

$$2 \log \left(\pi e^{R/2} + z - x' \right) - y' = 2 \log (x' - z) - y',$$

whose solution is $x^*(p') := x' - \frac{\pi}{2}e^{R/2}$. Plugging this into either the left or right hand side of the above equation yields the y -coordinate $y^*(p') = 2 \log \left(\frac{\pi}{2}e^{R/2} \right) - y'$. In a similar way, the x -coordinate $\hat{x}(p, p')$ is the solution to the equation $b_{p'}^+(z) = b_p^-(z)$ for $-\frac{\pi}{2}e^{R/2} \leq z \leq x + e^{(y+R)/2} - \pi e^{R/2}$, i.e.

$$2 \log \left(\pi e^{R/2} + z - x' \right) - y' = 2 \log (x - z) - y.$$

This solution is $\frac{x' - \pi e^{R/2}}{1 + e^{(y' - y)/2}}$ and again $\hat{y}(p, p')$ is obtained by plugging the solution into either the left or right hand side of the equation, yielding $\hat{y}(p, p') = 2 \log \left(\frac{\pi e^{R/2} - x'}{e^{y/2} + e^{y'/2}} \right)$.

To summarize we have:

$$\begin{aligned} x^*(p') &= x' - \frac{\pi}{2}e^{R/2} \\ y^*(p') &= 2 \log \left(\frac{\pi}{2}e^{R/2} \right) - y' \\ \hat{x}(p, p') &= \frac{x' - \pi e^{R/2}}{1 + e^{(y' - y)/2}} \\ \hat{y}(p, p') &= 2 \log \left(\frac{\pi e^{R/2} - x'}{e^{y/2} + e^{y'/2}} \right) \end{aligned}$$

The crucial observation is that $\mathcal{T}(p, p') = \emptyset$ as long as the point $(x^*(p'), y^*(p'))$ is above the left boundary of p . This happens exactly when $y^*(p') > b_p^-(x^*(p'))$. Therefore the boundary of this event is given by the equation $y^*(p') = b_p^-(x^*(p'))$ which reads

$$2 \log \left(\frac{\pi}{2}e^{R/2} \right) - y' = 2 \log \left(\frac{\pi}{2}e^{R/2} - x' \right) - y.$$

Solving this equation gives us the function

$$b_p^*(z) = y - 2 \log \left(1 - \frac{z}{\frac{\pi}{2}e^{R/2}} \right), \quad (71)$$

which is displayed by the red curve in Figure 8. It holds that $y^*(p') > b_p^-(x^*(p'))$ if and only if $y' < b_p^*(x')$ and hence we have that $\mathcal{T}(p, p') = \emptyset$ for all $p' \in \mathcal{R}$ for which $y' \geq b_p^*(x')$. We also note that when $y' = b_p^*(x')$ the two points $(x^*(p'), y^*(p'))$ and $(\hat{x}(p, p'), \hat{y}(p, p'))$ coincide.

This analysis allows us to compute the expected difference in the number of triangles for the finite box model and the infinite model, for a typical node with height y , i.e. prove Lemma 7.2.

Proof of Lemma 7.2. Due to symmetry it is enough to show that

$$\int_0^R \int_0^{I_n} \mu(\mathcal{T}(p, p_1)) f(x_1, y_1) dx_1 dy_1 = O \left(y n^{-(2\alpha-1)} + n^{-(2\alpha-1)} e^y \right) \quad (72)$$




Figure 8: Example for a given p of the boundary function $x' \mapsto b_p^*(x')$, given by the red curve, which determines whether $\mathcal{T}(p, p') = \emptyset$. We see that when $y' = b_p^*(x')$ then $(\hat{x}(p, p'), \hat{y}(p, p')) = (x^*(p'), y^*(p'))$.




Figure 9: Three different areas $B_n^{(i)}$ used in the proof of Lemma 7.2.

The proof goes in two stages. First we compute $\mu(\mathcal{T}(p, p_1))$ by splitting it over three disjoint regimes with respect to p_1 , with $x_1 \geq 0$. Then we do the integration with respect to p_1 .

Computing $\mu(\mathcal{T}(p, p_1))$

Recall that $I_n = \frac{\pi}{2}e^{R/2}$ and define the sets

$$\begin{aligned} A_n^{(1)} &= \{p_1 = (x_1, y_1) \in \mathcal{R} : 0 \leq y_1 \leq y - 2 \log(I_n/(I_n - x_1))\}, \\ A_n^{(2)} &= \left\{p_1 = (x_1, y_1) \in \mathcal{R} : y - 2 \log(I_n/(I_n - x_1)) < y_1 \leq y + 2 \log\left(1 + \frac{x_1}{I_n}\right)\right\}, \\ A_n^{(3)} &= \left\{p_1 = (x_1, y_1) \in \mathcal{R} : y + 2 \log\left(1 + \frac{x_1}{I_n}\right) < y_1 \leq y + 2 \log\left(\frac{I_n}{I_n - x_1}\right)\right\}, \end{aligned}$$

and let $B_n^{(i)} = \mathcal{B}_{\text{box}}(p) \cap A_n^{(i)}$, for $i = 1, 2, 3$, see Figure 9. Here the heights of the two intersections are given by

$$h_*(y) = y + 2 \log\left(\frac{I_n}{I_n + e^y}\right) \quad (73)$$

$$h^*(y) = y + 2 \log\left(\frac{I_n}{I_n - e^y}\right). \quad (74)$$

With these definitions we have that the union $B_n := \bigcup_{i=1}^n B_n^{(i)}$ denotes the area under the red curve in Figure 8 and hence, for all $p_1 \in \mathcal{R} \setminus B_n$ with $x_1 \geq 0$ we have that $\mathcal{T}(p, p_1) = \emptyset$. So we only need to consider $p_1 \in B_n$. We shall establish the following result:

$$\mu(\mathcal{T}(p, p_1)) = \begin{cases} O(I_n^{-2\alpha} e^{\alpha y_1}) & \text{if } p_1 \in B_n^{(1)} \\ O(I_n^{-2\alpha} e^{\alpha y}) & \text{if } p_1 \in B_n^{(2)} \cup B_n^{(3)} \end{cases} \quad (75)$$

Depending on which set p_1 belongs to, the set $\mathcal{T}(p, p_1)$ has a different shape. We displayed these shapes in Figure 10 as a visual aid to follow the computations below.

Case $p_1 \in B_n^{(1)}$: $0 \leq y_1 \leq y - 2\log(I_n/(I_n - x_1))$ In this case the integral over p_2 splits into two parts

$$\begin{aligned} \mathcal{I}_n^{(1)}(p_1) &:= \int_{h_2(p_1)}^{y^*(p_1)} \int_{-I_n}^{x_1 + e^{(y_1+y_2)/2} - 2I_n} e^{-\alpha y_2} dx_2 dy_2 \\ \mathcal{I}_n^{(2)}(p_1) &:= \int_{y^*(p_1)}^{h_1(p_1)} \int_{x^*(p_1)}^{x_1 - e^{(y_1+y_2)/2}} e^{-\alpha y_2} dx_2 dy_2. \end{aligned}$$

We first compute $\mathcal{I}_n^{(1)}$.

$$\begin{aligned} \mathcal{I}_n^{(1)}(p_1) &= \int_{h_2(p_1)}^{y^*(p_1)} \left(x_1 + e^{(y_1+y_2)/2} - I_n \right) e^{-\alpha y_2} dy_2 \\ &\leq e^{y_1/2} \int_{h_2(p_1)}^{y^*(p_1)} e^{-(\alpha - \frac{1}{2})y_2} dy_2 \\ &= \frac{2e^{y_1/2}}{2\alpha - 1} \left(e^{-(\alpha - \frac{1}{2})h_2(p_1)} - e^{-(\alpha - \frac{1}{2})y^*(p_1)} \right) \\ &= \frac{2e^{\alpha y_1}}{2\alpha - 1} I_n^{-(2\alpha-1)} \left(\left(1 - \frac{x_1}{I_n} \right)^{-(2\alpha-1)} - 1 \right) \\ &= O(I_n^{-2\alpha} x_1 e^{\alpha y_1}), \end{aligned}$$

where we used that $x_1 \leq e^{(y+y_1)/2} = o(I_n)$ for all $y_1 \leq y$ and $y \in \mathcal{K}_C(k_n)$ so that

$$\left(\left(1 - \frac{x_1}{I_n} \right)^{-(2\alpha-1)} - 1 \right) = O\left(\frac{x_1}{I_n} \right) \quad \text{as } n \rightarrow \infty.$$

For $\mathcal{I}_n^{(2)}(p_1)$ we have

$$\begin{aligned} \mathcal{I}_n^{(2)}(p_1) &= \int_{y^*(p_1)}^{h_1(p_1)} \left(I_n + x_1 - e^{(y_1+y_2)/2} \right) e^{-\alpha y_2} dy_2 \\ &\leq 2I_n \int_{y^*(p_1)}^{h_1(p_1)} e^{-\alpha y_2} dx_2 dy_2 \\ &= \frac{2}{\alpha} I_n \left(I_n^{-2\alpha} e^{\alpha y_1} - (I_n + x_1)^{-2\alpha} e^{-\alpha y_1} \right) \\ &= O(I_n^{-2\alpha} x_1 e^{\alpha y_1}) = O(I_n^{-(2\alpha-1)} e^{\alpha y_1}). \end{aligned}$$

We conclude that for $p_1 \in B_n^{(1)}$:

$$\mu(\mathcal{T}(p, p_1)) = O(I_n^{-2\alpha} x_1 e^{\alpha y_1}),$$

which establishes the first part of (75).




Figure 10: The different shapes of $\mathcal{T}(p, p_1)$ depending on the regime to which p_1 belongs. The top figure is for $p_1 \in B_n^{(1)}$, the middle for $p_1 \in B_n^{(2)}$ and the bottom one for $p_1 \in B_n^{(3)}$.

Case $p_1 \in B_n^{(2)}$: $y - 2 \log(I_n/(I_n - x_1)) < y_1 \leq y + 2 \log\left(1 + \frac{x_1}{I_n}\right)$ Here we split the integration into two parts (see Figure 10). Recall that $x^*(p, p_1) = x_1 - I_n$. Then, for the first part we have

$$\begin{aligned}
\mathcal{I}_n^{(1)}(p, p_1) &\leq \int_{h(y)}^{h_1(p_1)} \int_{-I_n}^{x^*(p, p_1)} f(x_2, y_2) dx_2 dy_2 \\
&= O\left(x_1 \left(e^{-\alpha h(y)} - e^{-\alpha h_1(p_1)}\right)\right) \\
&= O\left(x_1 I_n^{-2\alpha} \left(e^{\alpha y} - e^{\alpha y_1} \left(1 + \frac{x_1}{I_n}\right)^{-2\alpha}\right)\right) \\
&= O\left(I_n^{-2\alpha} x_1 e^{\alpha y_1} \left(\left(1 - \frac{x_1}{I_n}\right)^{-2\alpha} - \left(1 + \frac{x_1}{I_n}\right)^{-2\alpha}\right)\right) \\
&= O\left(I_n^{-2\alpha} x_1 e^{\alpha y_1}\right) = O\left(I_n^{-(2\alpha-1)} e^{\alpha y}\right),
\end{aligned}$$

where we used that $y \leq y_1 + 2 \log(I_n/(I_n - x_1))$ for $p_1 \in B_n^{(2)}$ for the third line and

$$\left(1 - \frac{x_1}{I_n}\right)^{-2\alpha} - \left(1 + \frac{x_1}{I_n}\right)^{-2\alpha} = O\left(\frac{x_1}{I_n}\right) = O(1),$$

for the last line.

For the second part we first use the upper bound on y_1 to compute that

$$\begin{aligned}
x_1 + e^{(y_1+y_2)/2} - 2I_n + e^{(y+y_2)/2} &\leq \left(e^{y/2} + e^{y_1/2}\right) e^{y_2/2} \\
&\leq e^{y/2} \left(2 + \frac{x_1}{I_n}\right) e^{y_2/2} = O\left(e^{(y+y_2)/2}\right),
\end{aligned}$$

since $|x_1| \leq I_n$. Then we have

$$\begin{aligned}
\mathcal{I}_n^{(2)} &= \int_{\hat{y}(p, p_1)}^{h(y)} \int_{-e^{(y+y_2)/2}}^{x_1 + e^{(y+y_1)/2} - 2I_n} f(x_2, y_2) dx_2 dy_2 \\
&= O\left(e^{y/2} \int_{\hat{y}(p, p_1)}^{h(y)} e^{-(\alpha - \frac{1}{2})y_2} dy_2\right) \\
&= O\left(e^{y/2} \left(e^{-(\alpha - \frac{1}{2})\hat{y}(p, p_1)} - e^{-(\alpha - \frac{1}{2})h(y)}\right)\right) \\
&= O\left(e^{y/2} \left(\left(\frac{2I_n - x_1}{e^{y/2} + e^{y_1/2}}\right)^{-(2\alpha-1)} - I_n^{-(2\alpha-1)} e^{(\alpha - \frac{1}{2})y}\right)\right) \\
&= O\left(I_n^{-(2\alpha-1)} e^{\alpha y}\right),
\end{aligned}$$

where for the last line we first used that $(2I_n - x_1)^{-(2\alpha-1)} \leq I_n^{-(2\alpha-1)}$ and then

$$\left(\left(e^{y/2} + e^{y_1/2}\right)^{2\alpha-1} - e^{(\alpha - \frac{1}{2})y}\right) \leq e^{(\alpha - \frac{1}{2})y} \left(\left(1 + \sqrt{1 + \frac{x_1}{I_n}}\right)^{2\alpha-1} - 1\right) = O\left(e^{(\alpha - \frac{1}{2})y}\right).$$

It then follows that for $p_1 \in B_n^{(2)}$

$$\mu(\mathcal{T}(p, p_1)) = O\left(I_n^{-(2\alpha-1)} e^{\alpha y}\right).$$

Case $p_1 \in B_n^{(3)}$: $y + 2 \log(1 + x_1/I_n) < y_1 \leq y + 2 \log(I_n/(I_n - x_1))$

$$\begin{aligned} \mathcal{I}_n^{(1)} &= \int_{y^*}^{\tilde{y}} \int_{-e^{(y+y_2)/2}}^{x_1 - e^{(y_1+y_2)/2}} f(x_2, y_2) dx_2 dy_2 \\ &= O \left(\int_{y^*}^{\tilde{y}} x_1 e^{-\alpha y_2} - \left(e^{y_1/2} - e^{y/2} \right) e^{-(\alpha - \frac{1}{2})y_2} dy_2 \right) \\ &= O \left(x_1 \int_{y^*}^{\tilde{y}} e^{-\alpha y_2} dy_2 \right). \end{aligned}$$

Now

$$\begin{aligned} \int_{y^*}^{\tilde{y}} e^{-\alpha y_2} dy_2 &= \frac{1}{\alpha} \left(e^{-\alpha y^*} - e^{-\alpha \tilde{y}} \right) = \frac{1}{\alpha} \left(I_n^{-2\alpha} e^{\alpha y_1} - \left(\frac{x_1}{e^{y_1/2} - e^{y/2}} \right)^{-2\alpha} \right) \\ &= \frac{I_n^{-2\alpha} e^{\alpha y_1}}{\alpha} \left(1 - \left(1 - e^{(y-y_1)/2} \right)^{2\alpha} \left(\frac{x_1}{I_n} \right)^{-2\alpha} \right) = O \left(I_n^{-2\alpha} e^{\alpha y_1} \right), \end{aligned}$$

and hence we have

$$\mathcal{I}_n^{(1)} = O \left(I_n^{-2\alpha} x_1 e^{\alpha y_1} \right).$$

For the second integral we have, using that $y \leq y_1$ for $p_1 \in B_n^{(3)}$,

$$\begin{aligned} \mathcal{I}_n^{(2)} &= \int_{\hat{y}}^{y^*} \int_{-e^{(y+y_2)/2}}^{e^{(y_1+y_2)/2} + x_1 - 2I_n} f(x_2, y_2) dx_2 dy_2 \\ &= O \left(\int_{\hat{y}}^{y^*} \left(e^{y/2} + e^{y_1/2} \right) e^{-(\alpha - \frac{1}{2})y_2} dy_2 \right) \\ &= O \left(e^{y_1/2} \int_{\hat{y}}^{y^*} e^{-(\alpha - \frac{1}{2})y_2} dy_2 \right). \end{aligned}$$

For the integral we have

$$\begin{aligned} \int_{\hat{y}}^{y^*} e^{-(\alpha - \frac{1}{2})y_2} dy_2 &= \frac{2}{2\alpha - 1} \left(e^{-(\alpha - \frac{1}{2})\hat{y}} - e^{-(\alpha - \frac{1}{2})y^*} \right) \\ &= \frac{2}{2\alpha - 1} \left(\left(\frac{2I_n - x_1}{e^{y/2} + e^{y_1/2}} \right)^{-(2\alpha - 1)} - I_n^{-(2\alpha - 1)} e^{(\alpha - \frac{1}{2})y_1} \right) = O \left(I_n^{-(2\alpha - 1)} e^{(\alpha - \frac{1}{2})y_1} \right), \end{aligned}$$

where we used the upper bound on y_1 and the fact that $2I_n - x_1 = \Theta(I_n)$ for all $x_1 \in [-I_n, I_n]$. We conclude that

$$\mathcal{I}_n^{(2)} = O \left(I_n^{-(2\alpha - 1)} x_1 e^{\alpha y} \right)$$

and hence for $p_1 \in B_n^{(3)}$

$$\mu(\mathcal{T}(p, p_1)) = O \left(I_n^{-2\alpha} x_1 e^{\alpha y} \right) = O \left(I_n^{-(2\alpha - 1)} e^{\alpha y} \right).$$

Integration $\mu(\mathcal{T}(p, p_1))$ with respect to p_1

We now proceed with the second part of the computation leading to (72). Here we will integrate $\mu(\mathcal{T}(p, p'))(p, p_1)$ over the region $B_n := B_n^{(1)} \cup B_n^{(2)} \cup B_n^{(3)}$, see Figure 9. Let us first identify the boundaries of these areas.

The area $B_n^{(1)}$ is bounded from above by the line given by the equation

$$y_1 = y - 2 \log \left(\frac{I_n}{I_n - x_1} \right).$$

Solving this for x_1 yields $x_1 = I_n (1 - e^{(y_1 - y)/2})$ and hence the area $B_n^{(1)}$ is given by

$$B_n^{(1)} = \left\{ (x_1, y_1) : 0 \leq y_1 \leq y, \quad 0 \leq x_1 \leq I_n \left(1 - e^{(y_1 - y)/2} \right) \wedge e^{(y + y_1)/2} \right\}.$$

In a similar way we have that $B_n^{(2)}$ is bounded from above by line

$$y_1 = y + 2 \log \left(\frac{I_n}{I_n + x_1} \right),$$

which yields $x_1 = I_n (e^{(y_1 - y)/2} - 1)$. The lower red boundary is the upper boundary of $B_n^{(2)}$ and hence we have

$$B_n^{(2)} = \left\{ (x_1, y_1) : h_*(y) \leq y_1 \leq h^*(y), \quad I_n \left(1 - e^{(y_1 - y)/2} \right) \vee I_n \left(e^{(y_1 - y)/2} - 1 \right) \leq x_1 \leq e^{(y + y_1)/2} \right\}.$$

We continue in the same way for $B_n^{(3)}$

$$B_n^{(3)} = \left\{ (x_1, y_1) : y \leq y_1 \leq R, \quad I_n \left(1 - e^{(y - y_1)/2} \right) \leq x_1 \leq I_n \left(e^{(y_1 - y)/2} - 1 \right) \wedge e^{(y + y_1)/2} \wedge I_n \right\}.$$

We these characterizations of the areas we now integrate $\mu(\mathcal{T}(p, p_1))$ over B_n , splitting the computations over the three different areas.

Integration over $B_n^{(1)}$: We use that $I_n (1 - e^{(y_1 - y)/2}) \wedge e^{(y + y_1)/2} \leq I_n (1 - e^{(y_1 - y)/2})$ so that

$$\begin{aligned} & \int_{B_n^{(1)}} \mu(\mathcal{T}(p, p_1)) f(x_1, y_1) dx_1 dy_1 \\ & \leq \int_0^y \int_0^{I_n (1 - e^{(y_1 - y)/2})} \mu(\mathcal{T}(p, p_1)) f(x_1, y_1) dx_1 dy_1 \\ & = O \left(I_n^{-2\alpha} \int_0^y \int_0^{e^{(y + y_1)/2}} x_1 dx_1 dy_1 \right) \\ & = O \left(I_n^{-(2\alpha - 1)} \int_0^y \left(1 - e^{(y_1 - y)/2} \right)^2 dy_1 \right) \\ & = O \left(I_n^{-(2\alpha - 1)} y \right) = O \left(y n^{-(2\alpha - 1)} \right). \end{aligned}$$

Integration over $B_n^{(2)}$: We will show that

$$\mu(B_n^{(2)}) = O \left(I_n^{-1} e^{(2 - \alpha)y} \right), \quad (76)$$

which together with (75) yields

$$\begin{aligned} \int_{B_n^{(2)}} \mu(\mathcal{T}(p, p_1)) f(x_1, y_1) dx_1 dy_1 & = O \left(\mu(B_n^{(2)}) I_n^{-(2\alpha - 1)} e^{\alpha y} \right) \\ & = O \left(I_n^{-2\alpha} e^{2y} \right). \end{aligned}$$

The integration is split into two parts determined by $I_n (1 - e^{(y_1 - y)/2}) \vee I_n (e^{(y_1 - y)/2} - 1)$:

$$\mu(B_n^{(3)}) = \int_{h_*(y)}^y \int_{I_n (1 - e^{(y_1 - y)/2})}^{e^{(y + y_1)/2}} f(x_1, y_1) dx_1 dy_1$$

$$+ \int_y^{h^*(y)} \int_{I_n(e^{(y_1-y)/2}-1)}^{e^{(y+y_1)/2}} f(x_1, y_1) dx_1 dy_1.$$

For the first integral we use that $e^{(y+y_1)/2} - I_n(1 - e^{(y_1-y)/2}) \leq e^{y_1/2} (e^{y/2} + e^{-y/2})$ to obtain

$$\begin{aligned} & \int_{h_*(y)}^y \int_{I_n(1-e^{(y_1-y)/2})}^{e^{(y+y_1)/2}} f(x_1, y_1) dx_1 dy_1 \\ &= O \left(e^{y/2} \int_{h_*(y)}^y e^{-(\alpha-\frac{1}{2})y_1} dy_1 \right) \\ &= O \left(e^{y/2} \left(e^{-(\alpha-\frac{1}{2})y} - e^{-(\alpha-\frac{1}{2})y} \left(\frac{I_n}{I_n + e^y} \right)^{-(2\alpha-1)} \right) \right) \\ &= O \left(I_n^{-1} e^{(2-\alpha)y} \right). \end{aligned}$$

For the second integral note that $e^{(y+y_1)/2} - I_n(e^{(y_1-y)/2} - 1) \leq e^{(y+y_1)/2}$ and hence

$$\begin{aligned} & \int_y^{h^*(y)} \int_{I_n(e^{(y_1-y)/2}-1)}^{e^{(y+y_1)/2}} f(x_1, y_1) dx_1 dy_1 \\ &= O \left(e^{y/2} \int_y^{h^*(y)} e^{-(\alpha-\frac{1}{2})y_1} dy_1 \right) \\ &= O \left(e^{y/2} \left(e^{-(\alpha-\frac{1}{2})y} - e^{-(\alpha-\frac{1}{2})y} \left(\frac{I_n}{I_n - e^y} \right)^{-(2\alpha-1)} \right) \right) \\ &= O \left(I_n^{-1} e^{(2-\alpha)y} \right), \end{aligned}$$

so that (76) follows.

Integration over $B_n^{(3)}$: For this case we show that

$$\mu(B_n^{(3)}) = O \left(e^{(1-\alpha)y} \right) \quad (77)$$

so that

$$\begin{aligned} \int_{B_n^{(3)}} \mu(\mathcal{T}(p, p_1)) f(x_1, y_1) dx_1 dy_1 &= O \left(\mu(B_n^{(2)}) I_n^{-(2\alpha-1)} e^{\alpha y} \right) \\ &= O \left(I_n^{-(2\alpha-1)} e^y \right). \end{aligned}$$

Here the integral is split into three parts:

$$\begin{aligned} \mu(B_n^{(3)}) &= \int_y^{h^*(y)} \int_{I_n(1-e^{(y_1-y)/2})}^{I_n(e^{(y_1-y)/2}-1)} f(x_1, y_1) dx_1 dy_1 \\ &+ \int_{h^*(y)}^{h(y)} \int_{I_n(1-e^{(y_1-y)/2})}^{e^{(y+y_1)/2}} f(x_1, y_1) dx_1 dy_1 \\ &+ \int_{h(y)}^R \int_{I_n(1-e^{(y_1-y)/2})}^{I_n} f(x_1, y_1) dx_1 dy_1. \end{aligned}$$

Let us first focus on the first integral. Since $I_n(e^{(y_1-y)/2} - 1) - I_n(1 - e^{(y_1-y)/2}) \leq I_n e^{(y_1-y)/2}$ we get, using similar arguments as above

$$\int_y^{h^*(y)} \int_{I_n(1-e^{(y_1-y)/2})}^{I_n(e^{(y_1-y)/2}-1)} f(x_1, y_1) dx_1 dy_1 = O \left(I_n e^{-y/2} \int_y^{h^*(y)} e^{-(\alpha-\frac{1}{2})y_1} dy_1 \right)$$

$$\begin{aligned}
&= O \left(I_n e^{-\alpha y} \left(1 - \left(\frac{I_n}{I_n - e^y} \right)^{-(2\alpha-1)} \right) \right) \\
&= O \left(e^{(1-\alpha)y} \right).
\end{aligned}$$

Proceeding to the second integral, we first note that $e^{(y+y_1)/2} - I_n(1-e^{(y-y_1)/2}) = O(I_n e^{(y_1-y)/2})$ so that similar calculations as before yield

$$\int_{h^*(y)}^{h(y)} \int_{I_n(1-e^{(y-y_1)/2})}^{e^{(y+y_1)/2}} f(x_1, y_1) dx_1 dy_1 = O \left(I_n e^{-y/2} \int_{h^*(y)}^{h(y)} e^{-(\alpha-\frac{1}{2})y_1} dy_1 \right) = O \left(e^{(1-\alpha)y} \right).$$

□

8 Concentration for $c(k; G_{\text{box}})$ (Proving Proposition 6.4)

In this section we establish a concentration result for the local clustering function $c^*(k; G_{\text{box}})$ in the finite box model G_{box} . Similar to the previous section we will focus on typical points $p = (0, y)$ with $y \in \mathcal{K}_C(k_n)$.

8.1 The main contribution of triangles

Recall that $N_{\text{box}}(k_n)$ denotes the number of vertices in G_{box} with degree k_n . We first write

$$c^*(k_n; G_{\text{box}}) = \frac{T_{\text{box}}(k_n)}{\binom{k_n}{2} \mathbb{E}[N_{\text{box}}(k_n)]},$$

where

$$T_{\text{box}}(k_n) = \sum_{p \in \mathcal{P}} \mathbb{1}_{\{D_{G_{\text{box}}}(p)=k_n\}} \sum_{(p_1, p_2) \in \mathcal{P} \setminus \{p\}} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{box}}(p)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{box}}(p)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{box}}(p_1)\}}$$

In particular, the variance of $c^*(k_n; G_{\text{box}})$ is determined by the variance of $T_{\text{box}}(k_n)$.

Next, recall the adjusted triangle count function

$$\tilde{T}_{\text{box}}(p_0) = \sum_{(p_1, p_2) \in \mathcal{P} \setminus \{p_0\}}^{\neq} \tilde{T}_{\text{box}}(p_0, p_1, p_2).$$

where

$$\tilde{T}_{\text{box}}(p_0, p_1, p_2) = \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{box}}(p_0)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{box}}(p_0)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\infty}(p_1) \cap \mathcal{R}\}},$$

as well as the definition of $\mathcal{K}_C(k_n)$

$$\mathcal{K}_C(k_n) = \left\{ y \in \mathbb{R}_+ : \frac{k_n - C\sqrt{k_n \log(k_n)}}{\xi} \vee 1 \leq e^{\frac{y}{2}} \leq \frac{k_n + C\sqrt{k_n \log(k_n)}}{\xi} \right\},$$

and write $\mathcal{R}(k_n, C) = [-I_n, I_n] \times \mathcal{K}_C(k_n)$ for the part of the box \mathcal{R} with heights in $\mathcal{K}_C(k_n)$. Slightly abusing notation, we will define the corresponding triangle degree function

$$\tilde{T}_{\text{box}}(k_n, C) = \sum_{p \in \mathcal{P} \cap \mathcal{R}(k_n, C)} \mathbb{1}_{\{\deg_{\text{box}}(p)=k_n\}} \tilde{T}_{\text{box}}(p). \tag{78}$$

and with that a different clustering function.

$$\tilde{c}_{\text{box}}(k_n) = \frac{\tilde{T}_{\text{box}}(k_n, C)}{\binom{k_n}{2} \mathbb{E}[N_{\text{box}}(k_n)]}. \tag{79}$$

The idea is that the main contribution of triangles of degree k_n to the triangle count $T_{\text{box}}(k_n)$ is given by $\tilde{T}_{\text{box}}(k_n, C)$. Therefore, in order to prove Proposition 6.4 it suffices to show that $\tilde{T}_{\text{box}}(k_n, C)$ is sufficiently concentrated around its mean. This last part is done in the following proposition.

Proposition 8.1 (Concentration $\tilde{T}_{\text{box}}(k_n, C)$). *Let $\alpha > \frac{1}{2}$, $\nu > 0$ and let $(k_n)_{n \geq 1}$ be any positive sequence satisfying $k_n = o\left(n^{\frac{1}{2\alpha+1}}\right)$. Then for any $C > 0$, as $n \rightarrow \infty$,*

$$\mathbb{E} \left[\tilde{T}_{\text{box}}(k_n, C)^2 \right] = (1 + o(1)) \mathbb{E} \left[\tilde{T}_{\text{box}}(k_n, C) \right]^2.$$

We first use this result to prove Proposition 6.4. The remainder of this section is devoted to the proof of Proposition 8.1. The final proof can be found in Section 8.3.

Proof of Proposition 6.4. We bound the expectation as follows,

$$\begin{aligned} \mathbb{E} [|c^*(k_n; G_{\text{box}}) - \mathbb{E}[c^*(k_n; G_{\text{box}})]|] &\leq \frac{\mathbb{E} \left[\left| \tilde{T}_{\text{box}}(k_n, C) - \mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)] \right| \right]}{\binom{k_n}{2} \mathbb{E}[N_{\text{box}}(k_n)]} \\ &\quad + 2\mathbb{E} [|c^*(k_n; G_{\text{box}}) - \tilde{c}_{\text{box}}(k_n)|]. \end{aligned}$$

We will show that both terms are $o(s(k_n))$.

First we note that $\mathbb{1}_{\{p_2 \in \mathcal{B}_{\infty}(p_1) \cap \mathcal{R}\}} \leq \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{box}}(p_1)\}}$ and hence $\tilde{T}_{\text{box}}(p) \leq T_{\text{box}}(p)$. This implies that

$$\tilde{c}_{\text{box}}(k_n) = \frac{\tilde{T}_{\text{box}}(k_n, C)}{\binom{k_n}{2} \mathbb{E}[N_{\text{box}}(k_n)]} \leq c^*(k_n; G_{\text{box}}).$$

and therefore

$$\mathbb{E} [|c^*(k_n; G_{\text{box}}) - \tilde{c}_{\text{box}}(k_n)|] = \mathbb{E} [c^*(k_n; G_{\text{box}})] - \mathbb{E} [\tilde{c}_{\text{box}}(k_n)].$$

For the expectation of $\tilde{T}_{\text{box}}(k_n, C)$ we use that

$$\mathbb{E} \left[\tilde{T}_{\text{box}}(p) \mid D_{G_{\text{box}}}(p) = k_n \right] = \binom{k_n}{2} \mu(\mathcal{B}_{\text{box}}(y))^{-2} \mathbb{E} [\tilde{T}_{\text{box}}(p)],$$

to get

$$\begin{aligned} \mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)] &= \int_{\mathcal{R}(k_n, C)} \mathbb{E} [\tilde{T}_{\text{box}}(p) \mid D_{G_{\text{box}}}(p) = k_n] \rho_{\text{box}}(y, k_n) f(x, y) \, dx \, dy \\ &= (1 + o(1)) \binom{k_n}{2} \int_{\mathcal{R}(k_n, C)} \mu(\mathcal{B}_{\text{box}}(y))^{-2} \mathbb{E} [\tilde{T}_{\text{box}}(y)] \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} \, dy \\ &= (1 + o(1)) \frac{1}{2} \int_{\mathcal{R}(k_n, C)} \mathbb{E} [\tilde{T}_{\text{box}}(y)] \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} \, dy \\ &= (1 + o(1)) n \binom{k_n}{2} \int_0^\infty P(y) \rho(y, k_n) \alpha e^{-\alpha y} \, dy, \end{aligned}$$

where the last line is due to Corollary 7.5. In particular, since the last integral is $\Theta\left(k_n^{-(2\alpha+1)} s(k_n)\right)$ we conclude that

$$\mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)] = \Theta\left(n k_n^{-(2\alpha-1)} s(k_n)\right). \quad (80)$$

Since $\mathbb{E}[N_{\text{box}}(k_n)] = (1 + o(1)) n p_{k_n}$ it follows that

$$\tilde{c}_{\text{box}}(k_n) = \frac{\mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)]}{\binom{k_n}{2} \mathbb{E}[N_{\text{box}}(k_n)]} = (1 + o(1)) \frac{\int_0^\infty P(y) \alpha e^{-\alpha y} \, dy}{p_{k_n}} = (1 + o(1)) \gamma(k_n).$$

On the other hand, Proposition 6.5 implies that $\mathbb{E}[c^*(k_n; G_{\text{box}})] = (1 + o(1))\gamma(k_n)$ and thus we conclude that

$$2\mathbb{E}[|c^*(k_n; G_{\text{box}}) - \tilde{c}_{\text{box}}(k_n)|] = o(\gamma(k_n)) = o(s(k_n)).$$

For the remaining term we use Hölder's inequality and Proposition 8.1 to obtain

$$\begin{aligned} \mathbb{E}\left[\left|\tilde{T}_{\text{box}}(k_n, C) - \mathbb{E}\left[\tilde{T}_{\text{box}}(k_n, C)\right]\right|\right] &\leq \left(\mathbb{E}\left[\tilde{T}_{\text{box}}(k_n, C)^2\right] - \mathbb{E}\left[\tilde{T}_{\text{box}}(k_n, C)\right]^2\right)^{\frac{1}{2}} \\ &= o\left(\mathbb{E}\left[\tilde{T}_{\text{box}}(k_n, C)\right]\right). \end{aligned}$$

This implies

$$\frac{\mathbb{E}\left[\left|\tilde{T}_{\text{box}}(k_n, C) - \mathbb{E}\left[\tilde{T}_{\text{box}}(k_n, C)\right]\right|\right]}{\binom{k_n}{2}\mathbb{E}[N_{\text{box}}(k_n)]} = o\left(\frac{\mathbb{E}\left[\tilde{T}_{\text{box}}(k_n, C)\right]}{\binom{k_n}{2}\mathbb{E}[N_{\text{box}}(k_n)]}\right) = o(s(k_n)),$$

which finishes the proof. \square

We note that the above proof establishes the following important result

Corollary 8.2. *Let $k_n \rightarrow \infty$. Then, as $n \rightarrow \infty$,*

$$\mathbb{E}[|c^*(k_n; G_{\text{box}}) - \tilde{c}_{\text{box}}(k_n)|] = o(s(k_n)).$$

8.2 Joint degrees in G_{box}

To prove Proposition 8.1 will use results from Section 3.4 regarding the joint degree distribution in G_{box} . For any two points $p, p' \in \mathcal{R}$ we will denote by

$$\rho_{\text{box}}(p, p', k, k') := \mathbb{P}(\text{Po}(\mu(\mathcal{B}_{\text{box}}(p))) = k, \text{Po}(\mu(\mathcal{B}_{\text{box}}(p'))) = k'). \quad (81)$$

the joint degree distribution.

Recall the definition of $\mathcal{E}_\varepsilon(k_n)$ from Section 3.4,

$$\mathcal{E}_\varepsilon(k_n) = \left\{(p, p') \in \mathcal{R} \times \mathcal{R} : y, y' \in [y_{k_n, C}^-, y_{k_n, C}^+] \text{ and } |x - x'|_n > k_n^{1+\varepsilon}\right\},$$

where

$$y_{k, C}^\pm = 2 \log \left(\frac{k \pm C\sqrt{k \log(k)}}{\xi} \right),$$

as defined in (15). Furthermore, we recall that by Lemma 3.9 the joint degree distribution of two point $p, p' \in \mathcal{E}_\varepsilon(k_n)$ factorizes, i.e. on the set $\mathcal{E}_\varepsilon(k_n)$ the joint degree distribution in G_{box} is asymptotically equivalent to the product of the degree distributions. We shall now prove a slightly stronger result (Lemma 8.4) which also takes care of bounded shifts in the joint degree distribution $\rho_{\text{box}}(p, p', k_n - t, k_n - t')$, for some uniformly bounded $t, t' \in \mathbb{Z}$. For this we first need the following simple result for Poisson distributions.

Lemma 8.3. *Let $k_n \rightarrow \infty$ be a sequence of non-negative integers and $X = \text{Po}(\lambda_n)$ be a Poisson random variable with mean λ_n satisfying*

$$k_n - C\sqrt{k_n \log(k_n)} \leq \lambda_n \leq k_n + C\sqrt{k_n \log(k_n)}$$

for some $C > 0$. Then, for any $t_n, s_n = O(1)$, as $n \rightarrow \infty$,

$$\mathbb{P}(X = k_n - t_n) \sim \mathbb{P}(X = k_n - s_n).$$

Proof. Note that $k_n > t_n, s_n$ for large enough n . Hence, using Stirling's formula, as $n \rightarrow \infty$,

$$\begin{aligned} \frac{\mathbb{P}(X = k_n - t_n)}{\mathbb{P}(X = k_n - s_n)} &= \frac{(k_n - t_n - (s_n - t_n))!}{(k_n - t_n)!} \lambda_n^{s_n - t_n} \\ &\sim \sqrt{\frac{k_n - s_n}{k_n - t_n}} \frac{(k_n - s_n)^{k_n - s_n}}{(k_n - t_n)^{k_n - t_n}} e^{t_n - s_n} \lambda_n^{s_n - t_n} \\ &= \sqrt{\ell_n} (\ell_n)^{k_n - t_n} e^{t_n - s_n} (k_n - s_n)^{t_n - s_n} \lambda_n^{s_n - t_n} \\ &= \sqrt{\ell_n} e^{(k_n - t_n) \log(\ell_n) + t_n - s_n} \left(\frac{k_n - s_n}{\lambda_n} \right)^{t_n - s_n} \end{aligned}$$

where we wrote $\ell_n = (k_n - s_n)/(k_n - t_n)$. Note that $\ell_n \rightarrow 1$ and hence $\sqrt{\ell_n} \rightarrow 1$. Moreover, since $(k_n - s_n)/\lambda_n \rightarrow 1$ and $|s_n - t_n| = O(1)$ we have that $\left(\frac{k_n - s_n}{\lambda_n} \right)^{t_n - s_n} \sim 1$. Therefore it remains to show that

$$\lim_{n \rightarrow \infty} e^{(k_n - t_n) \log(\ell_n) + t_n - s_n} = 1.$$

For this we note that for any x , such that $|x| \leq 1/2$, we have

$$x - x^2 \leq \log(1 + x) \leq x.$$

Write $x_n = \ell_n - 1 = \frac{t_n - s_n}{k_n - t_n}$. Then by the assumptions of the lemma, $x_n \rightarrow 0$, and thus, for n large enough,

$$t_n - s_n - \frac{(t_n - s_n)^2}{k_n - t_n} \leq (k_n - t_n) \log(\ell_n) \leq t_n - s_n.$$

In particular

$$e^{-\frac{(t_n - s_n)^2}{k_n - t_n}} \leq e^{(k_n - t_n) \log(\ell_n) + t_n - s_n} \leq 1,$$

and the result follows since $\frac{(t_n - s_n)^2}{k_n - t_n} \rightarrow 0$. \square

We can now prove the main result of this section.

Lemma 8.4. *Let $0 < \varepsilon < 1$, $k_n \rightarrow \infty$ and let $t_n, t'_n, s_n, s'_n \in \mathbb{Z}$ be uniformly bounded. Then for any $(p, p') \in \mathcal{E}_\varepsilon(k_n)$, as $n \rightarrow \infty$,*

$$\rho_{\text{box}}(p, p', k_n - t_n, k_n - t'_n) = (1 + o(1)) \rho_{\text{box}}(p, k_n - s_n) \rho_{\text{box}}(p', k_n - s'_n).$$

Proof. Define the random variables

$$\begin{aligned} X_1(p, p') &:= \text{Po}(\mu(\mathcal{B}_{\text{box}}(p) \setminus \mathcal{B}_{\text{box}}(p'))), \\ X_2(p, p') &:= \text{Po}(\mu(\mathcal{B}_{\text{box}}(p') \setminus \mathcal{B}_{\text{box}}(p))), \\ Y(p, p') &:= \text{Po}(\mu(\mathcal{B}_{\text{box}}(p) \cup \mathcal{B}_{\text{box}}(p'))), \end{aligned}$$

so that

$$\rho_{\text{box}}(p, p', k_n - t_n, k_n - t'_n) = \mathbb{P}(X_1(p, p') + Y(p, p') = k_n - t_n, X_2(p, p') + Y(p, p') = k_n - t'_n).$$

Since by Lemma 3.7 $\mu(\mathcal{B}_{\text{box}}(p) \cap \mathcal{B}_{\text{box}}(p')) = O(k_n^{1-\varepsilon'})$, it follows from Lemma 3.9 that

$$\rho_{\text{box}}(p, p', k_n - t_n, k_n - t'_n) = (1 + o(1)) \rho_{\text{box}}(p, k_n - t_n) \rho_{\text{box}}(p', k_n - t'_n).$$

The result then follows by applying Lemma 8.3 twice. \square

8.3 Concentration result for main triangle contribution

We now turn to Proposition 8.1. Before we dive into the proof let us first give a high level overview of the strategy and the flow of the arguments.

Recall (see (78)) that for any $C > 0$

$$\tilde{T}_{\text{box}}(k_n, C) = \sum_{p \in \mathcal{P}_n \cap \mathcal{K}_{C,n}(k_n)} \mathbb{1}_{\{\deg_{\text{box}}(p)=k\}} \tilde{T}_{\text{box}}(p)$$

Then we have

$$\tilde{T}_{\text{box}}(k_n, C)^2 = \sum_{p, p' \in \mathcal{P}_n \cap \mathcal{K}_C(k_n)} \mathbb{1}_{\{D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p')=k_n\}} \sum_{\substack{\neq \\ (p_1, p_2), (p'_1, p'_2) \in \mathcal{P}_n}} \tilde{T}_{\mathcal{P}}(p, p_1, p_2) \tilde{T}_{\mathcal{P}}(p', p'_1, p'_2),$$

This expression can be written as the sum of several terms, depending on how $\{p, p_1, p_2\}$ and $\{p', p'_1, p'_2\}$ intersect. To this end we define, for $a \in \{0, 1\}$ and $b \in \{0, 1, 2\}$,

$$I_{a,b} = \sum_{\substack{p, p' \in \mathcal{P}_n \cap \mathcal{K}_C(k) \\ |\{p\} \cap \{p'\}|=a}} \mathbb{1}_{\{D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p')=k_n\}} J_b(p, p'),$$

where

$$J_b(p, p') = \sum_{\substack{\neq \\ p_1, p_2, p'_1, p'_2 \in \mathcal{P}_n \\ |\{p_1, p_2\} \cap \{p'_1, p'_2\}|=b}} T_{\mathcal{P},n}(p, p_1, p_2) T_{\mathcal{P},n}(p', p'_1, p'_2),$$

with the sum taken over all two distinct pairs (p_1, p_2) and (p'_1, p'_2) . Then we have

$$\tilde{T}_{\text{box}}(k, C)^2 = \sum_{a=0}^1 \sum_{b=0}^2 I_{a,b}.$$

To prove Proposition 8.1 we will deal with each of the $I_{a,b}$ separately, showing that

$$\mathbb{E}[I_{0,0}] = (1 + o(1)) \mathbb{E} \left[\tilde{T}_{\text{box}}(k_n, C) \right]^2 \quad (82)$$

and for all other combinations

$$\mathbb{E}[I_{a,b}] = o \left(\mathbb{E} \left[\tilde{T}_{\text{box}}(k_n, C) \right]^2 \right). \quad (83)$$

Note $I_{1,2} = \tilde{T}_{\text{box}}(k_n, C)$ and since (80) implies that $\mathbb{E} \left[\tilde{T}_{\text{box}}(k_n, C) \right] \rightarrow \infty$, it follows that (83) holds for $I_{1,2}$.

Recall that $\mathcal{R}(k_n, C) = [-I_n, I_n] \times \mathcal{K}_C(k_n)$ and (36)

$$\mathcal{E}_\varepsilon(k_n) = \{(p, p') \in \mathcal{R} \times \mathcal{R} : y, y' \in \mathcal{K}_C(k_n) \text{ and } |x - x'|_n > k_n^{1+\varepsilon}\}.$$

Let $\mathcal{E}_\varepsilon(k_n)^c$ be the same set but with $|x - x'|_n \leq k_n^{1+\varepsilon}$ and denote by $I_{a,b}^*$ the the part of $I_{a,b}$ where $(p, p') \in \mathcal{E}_\varepsilon(k_n)$. Will split the analysis between $I_{a,b}^*$ and $I_{a,b} - I_{a,b}^*$. The idea for these two cases is that by Lemma 8.4 it follows that on the set $\mathcal{E}_\varepsilon(k_n)$ and for any uniformly bounded $t, t' \in \mathbb{Z}$, the joint degree distribution factorizes,

$$\rho_{\text{box}}(p, p', k_n + t, k_n + t') = (1 + o(1)) \rho_{\text{box}}(p, k_n) \rho_{\text{box}}(p', k_n).$$

In particular this allows us to prove that $\mathbb{E}[I_{0,0}^*] = (1 + o(1)) \mathbb{E} \left[\tilde{T}_{\text{box}}(k_n, C) \right]^2$. On the other hand, the expected number of points in $\mathcal{E}_\varepsilon(k_n)^c$ is $O(k_n^{1+\varepsilon} k_n^{-2\alpha} \mathbb{E}[N_{\text{box}}(k_n)]) = o \left(\mathbb{E}[N_{\text{box}}(k_n)]^2 \right)$, where the latter is the expected number of points in $\mathcal{R}(k_n, C) \times \mathcal{R}(k_n, C)$. Hence we expect the contributions coming from $\mathcal{E}_\varepsilon(k_n)^c$ to be negligible.

Proof of Proposition 8.1. Throughout this proof we set $i = |\{p', p_1, p_2, p'_1, p'_2\} \cap \mathcal{B}_{\text{box}}(p)|$, $j = |\{p'\} \cap \mathcal{B}_{\text{box}}(p)|$ and define i', j' in a similar way by interchanging the primed and non-primed variables. In addition, we write $\tilde{D}_{\text{box}}(p, p', k, \ell)$ to denote the indicator that $|\mathcal{B}_{\text{box}}(p) \cap (\mathcal{P} \setminus \{p, p', p_1, p_2, p'_1, p'_2\})| = k$ and $|\mathcal{B}_{\text{box}}(p') \cap (\mathcal{P} \setminus \{p, p', p_1, p_2, p'_1, p'_2\})| = \ell$. Note that this also depend on $\{p_1, p_2, p'_1, p'_2\}$ but we suppressed this to keep notation concise. Similarly we write $D_{\text{box}}(p, p', k, \ell)$ to denote the indicator that $|\mathcal{B}_{\text{box}}(p) \cap (\mathcal{P} \setminus \{p, p'\})| = k$ and $|\mathcal{B}_{\text{box}}(p') \cap (\mathcal{P} \setminus \{p, p'\})| = \ell$, which now only depends on p and p' . Then, by the Campbell-Mecke formula

$$\begin{aligned} & \mathbb{E} \left[\mathbb{1}_{\{D_{G_{\text{box}}}(p)=k_n, D_{G_{\text{box}}}(p')=k_n\}} J_b(p, p') \right] \\ &= \mathbb{E} \left[\sum_{\substack{p_1, p_2, p'_1, p'_2 \in \mathcal{P}_n \\ |\{p_1, p_2\} \cap \{p'_1, p'_2\}|=b}}^{\neq} \tilde{D}_{\text{box}}(p, p', k_n - i, k_n - i') \tilde{T}_{\text{box}}(p, p_1, p_2) \tilde{T}_{\text{box}}(p', p'_1, p'_2) \right] \end{aligned}$$

where the sum is over all distinct pairs (p_1, p_2) and (p'_1, p'_2) . We also know that

$$\mathbb{E}[T_{\mathcal{P}}(k_n)] = \Theta \left(n k_n^{-(2\alpha-1)} s_{\alpha}(k_n) \right).$$

We will now proceed to establish (82) and (83).

Computing $I_{0,0}$ We first show that

$$\mathbb{E}[I_{0,0} - I_{0,0}^*] = o \left(\mathbb{E}[T_{\text{box}}(k_n, C)]^2 \right), \quad (84)$$

so that for the remainder of the proof we only need to consider $p, p' \in \mathcal{E}_{\varepsilon}(k_n)$ and hence, we can apply Lemma 8.4.

For J_0 we have, using Lemma 8.4

$$\begin{aligned} & \mathbb{E} \left[\mathbb{1}_{\{D_{G_{\text{box}}}(p)=k_n, D_{G_{\text{box}}}(p')=k_n\}} J_0(p, p') \right] \\ &= \mathbb{E} \left[\sum_{\substack{p_1, p_2, p'_1, p'_2 \in \mathcal{P} \setminus \{p, p'\} \\ |\{p_1, p_2\} \cap \{p'_1, p'_2\}|=0}}^{\neq} \tilde{D}_{\text{box}}(p, p', k_n - i, k_n - i') \tilde{T}_{\text{box}}(p, p_1, p_2) \tilde{T}_{\text{box}}(p', p'_1, p'_2) \right] \\ &= \mathbb{E} \left[D_{\text{box}}(p, p', k_n - j - 2, k_n - j' - 2) \sum_{p_1, p_2 \in \mathcal{P} \setminus p}^{\neq} \tilde{T}_{\text{box}}(p, p_1, p_2) \sum_{p'_1, p'_2 \in \mathcal{P} \setminus p'}^{\neq} \tilde{T}_{\text{box}}(p', p'_1, p'_2) \right] \\ &= (1 + o(1)) \rho_{\text{box}}(p, p', k_n, k_n) \mathbb{E} \left[\tilde{T}_{\text{box}}(p) \middle| D_{G_{\text{box}}}(p) = k_n \right] \mathbb{E} \left[\tilde{T}_{\text{box}}(p') \middle| D_{G_{\text{box}}}(p') = k_n \right], \end{aligned}$$

Next we recall that for all $y' \in \mathcal{K}_C(k_n)$ (see (65)),

$$\mathbb{E} \left[\tilde{T}_{\text{box}}(p') \middle| D_{G_{\text{box}}}(p') = k_n \right] = \binom{k_n}{2} \mu(\mathcal{B}_{\text{box}}(p'))^{-2} \mathbb{E} \left[\tilde{T}_{\text{box}}(p') \right] = O(1) k_n^2 P(y'),$$

where $p' = (x', y')$ and we used that $\mathbb{E} \left[\tilde{T}_{\text{box}}(p') \right] = (1 + o(1)) k_n^2 P(y')$, for all $y' \in \mathcal{K}_C(k_n)$. Therefore, using that $\rho_{\text{box}}(p, p', k_n, k_n) \leq \rho_{\text{box}}(p, k_n)$,

$$\begin{aligned} & \mathbb{E} \left[\mathbb{1}_{\{D_{G_{\text{box}}}(p)=k_n, D_{G_{\text{box}}}(p')=k_n\}} J_0(p, p') \right] \\ & \leq O(k_n^2) \rho_{\text{box}}(p, k_n) \mathbb{E} \left[\tilde{T}_{\text{box}}(p) \middle| D_{G_{\text{box}}}(p) = k_n \right] P(y') \end{aligned}$$

and thus

$$\mathbb{E} [I_{0,0} - I_{0,0}^*]$$

$$\begin{aligned}
&= \int_{\mathcal{E}_\varepsilon(k_n)^c} \mathbb{E} \left[\mathbb{1}_{\{D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p')=k_n\}} J_0(p, p') \right] f(x, y) f(x', y') dx' dx dy' dy \\
&\leq O(k_n^2) k_n^{1+\varepsilon} \left(\int_{a_n^-}^{a_n^+} P(y') e^{-\alpha y'} dy' \right) \mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)] \\
&= O(k_n^{3+\varepsilon-2\alpha} s_\alpha(k_n) \mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)]) \\
&= o(n k_n^{-(2\alpha-1)} s_\alpha(k_n) \mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)]) = o\left(\mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)]^2\right),
\end{aligned}$$

which proves (84). Here we used that $k_n^{2+\varepsilon} = o(n)$ and

$$\mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)] = \Theta\left(\mathbb{E} [\tilde{T}_{\text{box}}(k_n)]\right) = \Theta\left(n k_n^{-(2\alpha-1)} s_\alpha(k_n)\right)$$

for the last line.

We will now show that

$$\mathbb{E} [I_{0,0}^*] = (1 + o(1)) \mathbb{E} [\mathbb{E} [T_{\text{box}}(k_n, C)]^2].$$

Recall the result from Lemma 8.4, that for $(p, p') \in \mathcal{E}_\varepsilon(k_n)$ and any two uniformly bounded $t, t' \in \mathbb{Z}$,

$$\rho_{\text{box}}(p, p', k_n + t, k_n + t') = (1 + o(1)) \rho_{\text{box}}(p, k_n) \rho_{\text{box}}(p', k_n).$$

Therefore, by defining $h(y) = \mathbb{E} [\tilde{T}_{\text{box}}(y) | D_{G_{\text{box}}}(y) = k_n]$

$$\mathbb{E} [I_{0,0}^*] = (1 + o(1)) \int_{\mathcal{E}_\varepsilon(k_n)} \rho_{\text{box}}(p, k_n) \rho_{\text{box}}(p', k_n) h(y) h(y') f(x, y) f(x', y') dx' dx dy' dy.$$

The difference with $\mathbb{E} [\mathbb{E} [T_{\text{box}}(k_n, C)]^2]$ is in that the above integral is over $\mathcal{E}_\varepsilon(k_n)$ instead of $\mathcal{R}(k_n, C) \times \mathcal{R}(k_n, C)$. Since the difference between the two sets is $\mathcal{E}_\varepsilon(k_n)^c$ and $n k_n^{1+\varepsilon} = o(n^2)$ it follows that

$$\begin{aligned}
&\mathbb{E} [\mathbb{E} [T_{\text{box}}(k_n, C)]^2] - \int_{\mathcal{E}_\varepsilon(k_n)} \rho_{\text{box}}(p, k_n) \rho_{\text{box}}(p', k_n) h(y) h(y') f(x, y) f(x', y') dx' dx dy' dy \\
&= \int_{\mathcal{E}_\varepsilon(k_n)^c} \rho_{\text{box}}(p, k_n) \rho_{\text{box}}(p', k_n) h(y) h(y') f(x, y) f(x', y') dx' dx dy' dy \\
&= O(k_n^{1+\varepsilon} n) \left(\int_{\mathcal{K}_C(k_n)} h(y) \rho_{\text{box}}(y, k_n) \alpha e^{-\alpha y} dy \right)^2 = o\left(\mathbb{E} [\mathbb{E} [T_{\text{box}}(k_n, C)]^2]\right).
\end{aligned}$$

Thus we conclude that $\mathbb{E} [I_{0,0}^*] = (1 + o(1)) \mathbb{E} [\mathbb{E} [T_{\text{box}}(k_n, C)]^2]$, which finishes the proof of (82).

Computing $\mathbb{E} [I_{0,1}]$ We first write

$$\mathbb{E} [\mathbb{1}_{\{D_{G_{\text{box}}}(p)=k_n, D_{G_{\text{box}}}(p')=k_n\}} J_1] \leq O(1) k_n \rho_{\text{box}}(p, p', k_n, k_n) \mathbb{E} [\tilde{T}_{\text{box}}(p) | D_{G_{\text{box}}}(p) = k_n] \quad (85)$$

Then, using that $\rho_{\text{box}}(p, p', k_n, k_n) \leq \rho_{\text{box}}(p, k_n)$,

$$\begin{aligned}
&\mathbb{E} [I_{0,1} - I_{0,1}^*] \\
&\int_{\mathcal{E}_\varepsilon(k_n)^c} \mathbb{E} [\mathbb{1}_{\{D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p')=k_n\}} J_1(p, p')] f(x, y) f(x', y') dx' dx dy' dy \\
&= k_n \int_{\mathcal{K}_C(k_n)^2} \mathbb{1}_{\{|x-x'| \leq k_n^{1+\varepsilon}\}} \rho_{\text{box}}(p, k_n) \mathbb{E} [\tilde{T}_{\text{box}}(p) | D_{G_{\text{box}}}(p) = k_n] f(x, y) f(x', y') dx' dx dy' dy
\end{aligned}$$

$$\begin{aligned} &\leq O(k_n^{2+\varepsilon}) \left(\int_{a_n^-}^{a_n^+} e^{-\alpha y'} dy' \right) \mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)] \\ &= O(k_n^{2+\varepsilon-2\alpha} \mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)]). \end{aligned}$$

Recall that $\mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)] = \Theta(nk_n^{-(2\alpha-1)} s(k_n))$. Therefore to show that $\mathbb{E} [I_{0,1} - I_{0,1}^*] = o\left(\mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)]^2\right)$ it suffices to show that $k_n^{2+\varepsilon-2\alpha} = o(nk_n^{-(2\alpha-1)} s(k_n))$. When $\frac{1}{2} < \alpha \leq \frac{3}{4}$ we have

$$\frac{4\alpha - 1 + \varepsilon}{2\alpha + 1} < 1,$$

for ε small enough. Hence

$$n^{-1} k_n^{2\alpha-1} s(k_n)^{-1} k_n^{2+\varepsilon-2\alpha} = n^{-1} k_n^{4\alpha-1+\varepsilon} = o\left(n^{-1} n^{\frac{4\alpha-1+\varepsilon}{2\alpha+1}}\right) = o(1)$$

When $\alpha \geq \frac{3}{4}$,

$$n^{-1} k_n^{2\alpha-1} s(k_n)^{-1} k_n^{2+\varepsilon-2\alpha} = O(\log(k_n)) n^{-1} k_n^{2+\varepsilon} = o(1),$$

for ε small enough.

For $(p, p') \in \mathcal{E}_\varepsilon(k_n)$ we assume without loss of generality that $p'_1 = p_1 = (x_1, y_1)$, i.e.

$$J_{0,1} = \sum_{(p_1, p_2) \in \mathcal{P} \setminus \{p\}}^{\neq} \tilde{T}_{\text{box}}(p, p_1, p_2) \sum_{p'_2 \in \mathcal{P} \setminus \{p', p_1\}} \tilde{T}_{\text{box}}(p', p_1, p'_2).$$

Now let $Z_{0,1}$ denote the part of $J_{0,1}$ where $y_1 \leq 4\log(k_n)$ and $y_2, y'_2 \leq \varepsilon \log(k_n)$.

We first analyze $\mathbb{E} [Z_{0,1} | D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p') = k_n]$. When $y_1 \leq 4\log(k_n)$ and both $y_2, y'_2 \leq \varepsilon \log(k_n)$ we have that

$$|x_2 - x'_2| \leq |x_1 - x_2| + |x_1 - x'_2| \leq e^{\frac{y_1}{2}} \left(e^{\frac{y_2}{2}} + e^{\frac{y'_2}{2}} \right) \leq 2k_n^{2+\varepsilon}$$

whenever $\tilde{T}_{\text{box}}(p, p_1, p_2) \tilde{T}_{\text{box}}(p', p_1, p'_2) > 0$ while both $|x - x_2|, |x' - x'_2| = O(k_n^{1+\varepsilon})$. Hence it follows that $\tilde{T}_{\text{box}}(p, p_1, p_2) \tilde{T}_{\text{box}}(p', p_1, p'_2) > 0$ implies that

$$|x - x'| \leq |x - x_2| + |x_2 - x'_2| + |x'_2 - x'| = O(k_n^{2+\varepsilon}).$$

Next, by integrating only over x'_2 and y'_2 we get

$$\begin{aligned} \mathbb{E} [Z_{0,1} | D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p') = k_n] &= O\left(e^{\frac{y'_1}{2}} \mathbb{1}_{\{|x-x'| \leq O(1)k_n^{2+\varepsilon}\}}\right) \mathbb{E} \left[\tilde{T}_{\text{box}}(p) \middle| D_{G_{\text{box}}}(p) = k_n \right] \\ &= O\left(k_n \mathbb{E} \left[\tilde{T}_{\text{box}}(p) \middle| D_{G_{\text{box}}}(p) = k_n \right]\right). \end{aligned}$$

Thus

$$\begin{aligned} &\int_{\mathcal{E}_\varepsilon(k_n)} \rho_{\text{box}}(p, p', k_n, k_n) \mathbb{E} [Z_{0,1} | D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p') = k_n] f(x, y) f(x', y') dx dy dx' dy' \\ &= O(k_n^{3+\varepsilon}) \mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)] \int_{\mathcal{K}_C(k_n)} \rho_{\text{box}}(y', k_n) e^{-\alpha y'} dy' \\ &= O\left(k_n^{2+\varepsilon} k_n^{-2\alpha} \mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)]\right) = o\left(\mathbb{E} [\tilde{T}_{\text{box}}(k_n, C)]^2\right), \end{aligned}$$

where the last line follows from the analysis done for $\mathbb{E} [I_{0,0} - I_{0,0}^*]$.

It now remains to consider $J_{0,1} - Z_{0,1} := Z_{0,1}^*$. We will show that

$$\mathbb{E} [Z_{0,1}^* | D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p') = k_n] = o(k_n^4 s(k_n)^2). \quad (86)$$

Using that the joint degree distribution factorizes on $\mathcal{E}_\varepsilon(k_n)$ this then implies that

$$\begin{aligned}\mathbb{E}[I_{0,1}^*] &= o(k_n^4 s(k_n)^2) \left(\int_{\mathcal{R}(k_n, C)} \rho_{\text{box}}(y, k_n) f(x, y) \, dx \, dy \right)^2 \\ &= o\left((ns(k_n)k_n^{-2\alpha+1})^2\right) = o\left(\mathbb{E}\left[\tilde{T}_{\text{box}}(k_n, C)\right]^2\right),\end{aligned}$$

which finished the proof of (83) for $a = 0, b = 1$.

We first consider the part with $y_1 > 4\log(k_n)$. Since the integration over x_1, x_2 and x'_2 of $\mathbb{E}[Z_{0,1}^* | D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p') = k_n]$ is bounded by $O(e^y e^{\frac{y'}{2}})$ we get that the contribution to $\mathbb{E}[Z_{0,1}^* | D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p') = k_n]$ with $y > 4\log(k_n)$ and $(p, p') \in \mathcal{E}_\varepsilon(k_n)$ is

$$\begin{aligned}O\left(e^y e^{\frac{y'}{2}} \int_{4\log(k_n)}^R e^{-(\alpha-\frac{1}{2})y_1} \, dy_1\right) &= O\left(k_n^3 \int_{4\log(k_n)}^R e^{-(\alpha-\frac{1}{2})y_1} \, dy_1\right) \\ &= O\left(k_n^{3-(4\alpha-2)}\right) = o(k_n^4 s_\alpha(k_n)^2).\end{aligned}$$

Here the last step follows since for $\frac{1}{2} < \alpha < \frac{3}{4}$

$$k_n^{3-(4\alpha-2)-4} s(k_n)^{-2} = k_n^{3-(4\alpha-2)-4+2(4\alpha-2)} = k_n^{-5+4\alpha} = o(1),$$

while for $\alpha = \frac{3}{4}$

$$k_n^{3-(4\alpha-2)-4} s(k_n)^{-2} = O(\log(k_n)^{-2}) k_n^{3-(4\alpha-2)-2} = O(\log(k_n)^{-2}) = o(1),$$

and for $\alpha > \frac{3}{4}$

$$k_n^{3-(4\alpha-2)-4} s(k_n)^{-2} = k_n^{3-(4\alpha-2)-2} = o(1).$$

Next we consider the case where $y_1 \leq 4\log(k_n)$ and at least one of y_2, y'_2 is larger than $\varepsilon \log(k_n)$. Due to symmetry it is enough to consider the case with $y_2 > \varepsilon \log(k_n)$. Here the contribution to $\mathbb{E}[Z_{0,1}^* | D_{G_{\text{box}}}(p), D_{G_{\text{box}}}(p') = k_n]$ is

$$\begin{aligned}\mathbb{E}[\tilde{T}_{\text{box}}(p)] O\left(e^{\frac{y'}{2}} \int_{\varepsilon \log(k_n)}^R e^{-(\alpha-\frac{1}{2})y_2} \, dy_2\right) &= O\left(k_n^{1-\varepsilon(\alpha-\frac{1}{2})}\right) \mathbb{E}[\tilde{T}_{\text{box}}(p)] \\ &= O\left(k_n^{3-\varepsilon(\alpha-\frac{1}{2})} s(k_n)\right) = o(k_n^4 s(k_n)^2).\end{aligned}$$

The last line follows since $k_n^{-1} = o(s(k_n))$ for $\frac{1}{2} < \alpha < \frac{3}{4}$ and $k_n^{-1} = O(s(k_n))$ for $\alpha \geq \frac{3}{4}$.

Computing $\mathbb{E}[I_{0,2}]$ In this case we have

$$\mathbb{E}\left[\mathbb{1}_{\{D_{G_{\text{box}}}(p)=k_n, D_{G_{\text{box}}}(p')=k_n\}} J_2\right] = (1 + o(1)) \rho_{\text{box}}(p, p', k_n, k_n) \mathbb{E}\left[\tilde{T}_{\text{box}}(p) \mid D_{G_{\text{box}}}(p) = k_n\right].$$

We then use that $\rho_{\text{box}}(p, p', k_n, k_n) \leq \rho_{\text{box}}(p, k_n)$ to obtain

$$\begin{aligned}\mathbb{E}[I_{0,2} - I_{0,2}^*] &= O(k_n^{1+\varepsilon}) \left(\int_{\mathcal{K}_C(k_n)} e^{-\alpha y'} \, dy' \right) \mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)] \\ &= O\left(k_n^{\varepsilon-(2\alpha-1)}\right) \mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)] = o\left(\mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)]\right)\end{aligned}$$

where the last line follows since $\mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)] = \Theta(nk_n^{-(2\alpha-1)} s(k_n))$ and $k_n^\varepsilon n^{-1} = o(s(k_n))$.

For the other term we use the fact that the degree distribution factorizes;

$$\begin{aligned}\mathbb{E}[I_{0,2}^*] &= O(1) \left(\int_{\mathcal{R}(k_n, C)} \rho_{\text{box}}(y', k_n) f(x', y') dx' dy' \right) \mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)] \\ &= O(nk_n^{-(2\alpha+1)}) \mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)] = o\left(\mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)]^2\right),\end{aligned}$$

where we also used that $k_n^{-2} = o(s(k_n))$.

Computing $\mathbb{E}[I_{1,1}]$ Using (86) we get

$$\begin{aligned}\mathbb{E}[I_{1,1}] &= O(k_n) \int_{\mathcal{R}(k_n, C)} \rho_{\text{box}}(y, k_n) \mathbb{E}[\tilde{T}_{\text{box}} | D_{G_{\text{box}}}(p) = k_n] f(x, y) dx dy \\ &= O(k_n) \mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)].\end{aligned}$$

Now observe that for $\frac{1}{2} < \alpha < \frac{3}{4}$

$$k_n n^{-1} k_n^{(2\alpha-1)} s(k_n)^{-1} = k_n^{6\alpha-2} n^{-1} = O\left(n^{\frac{4\alpha-3}{2\alpha+1}}\right) = o(1),$$

while for $\alpha \geq \frac{3}{4}$

$$k_n n^{-1} k_n^{(2\alpha-1)} s(k_n)^{-1} = O\left(n^{-1} k_n^{-(2\alpha-1)}\right) = o(1).$$

We conclude that $k_n = o(nk_n^{-(2\alpha-1)} s(k_n))$ and hence $\mathbb{E}[I_{1,1}] = o\left(\mathbb{E}[\tilde{T}_{\text{box}}(k_n, C)]^2\right)$. □

9 Equivalence for local clustering in G_{Po} and G_{box}

In this section we establish the equivalence between $c^*(k; G_n)$ and $c^*(k; G_{\text{box}})$ as expressed in Proposition 6.3, using the coupling procedure explained in Section 2.4. As in the previous section we write $|\cdot|_n$ for the norm $|\cdot|_{\pi_{e^{R/2}}}$.

Recall the map Ψ from (7)

$$\Psi(r, \theta) = \left(\theta \frac{e^{R/2}}{2}, R - r \right),$$

and that $\mathcal{B}(p)$ denotes the image under Ψ of the ball of hyperbolic radius R around the point $\Psi^{-1}(p)$. Under the coupling between the hyperbolic random graph and the finite box model, described in Section 2.4, two points $p = (x, y)$ and $p' = (x', y')$ are connected if and only if

$$|x - x'|_n \leq \Phi(y, y') = \frac{1}{2} e^{R/2} \arccos \left(\frac{\cosh(R - y) \cosh(R - y') - \cosh R}{\sinh(R - y) \sinh(R - y')} \right),$$

see (8). We will often use the result from Lemma 2.2 to approximate the function Φ , for $y + y' < R$, by

$$e^{\frac{1}{2}(y+y')} - K e^{\frac{3}{2}(y+y')-R} \leq \Phi(R - y, R - y') \leq e^{\frac{1}{2}(y+y')} + K e^{\frac{3}{2}(y+y')-R},$$

where K is a constant determined by the lemma.

9.1 Some results on the hyperbolic geometric graph

We start with some basic results for the hyperbolic random geometric graph. Recall that $\mathcal{B}_\infty(p) = \{p' \in \mathbb{R} \times \mathbb{R}_+ : |x - x'| \leq e^{(y+y')/2}\}$ and observe that (10) from Lemma 2.2 implies the following.

Corollary 9.1. *For sufficiently large n and $p \in \mathcal{R}$,*

$$\mathcal{B}_\infty(p) \cap \mathcal{R}([K, R]) \subseteq \mathcal{B}(p) \cap \mathcal{R}([K, R]),$$

where K is the constant from Lemma 2.2.

Furthermore, Lemma 2.2 enables us to determine the measure of a ball around a given point $p = (0, y)$ - this is will be fairly useful in our subsequent analysis.

Let $p \in \mathcal{R}$. Then we can see that the curve $x' = e^{\frac{1}{2}(y+y')}$ with $x' \geq 0$ meets the right boundary of \mathcal{R} , that is, the line $x' = \frac{\pi}{2}e^{R/2}$ at $y' = R - y + 2 \ln \frac{\pi}{2}$. Hence, any point $p' \in \mathcal{R}([R - y + 2 \ln \frac{\pi}{2}, R])$ is included in $\mathcal{B}_\infty(p)$. In other words,

$$\mathcal{B}_\infty(p) \cap \mathcal{R}([R - y + 2 \ln \frac{\pi}{2}, R]) = \mathcal{R}([R - y + 2 \ln \frac{\pi}{2}, R]).$$

This together with the fact that for any $u' = (r', \theta')$,

$$r' < y = R - r \Rightarrow d_{\mathbb{H}}(\Psi^{-1}(p), u') \leq R$$

implies that

$$(\mathcal{B}(p) \triangle \mathcal{B}_\infty(p)) \cap \mathcal{R}([R - y + 2 \ln \frac{\pi}{2}, R]) = \emptyset, \quad (87)$$

where $A \triangle B$ denotes the symmetric difference of the sets A and B . We can now compute the expected number of points in $\mathcal{B}(p) \triangle \mathcal{B}_\infty(p)$, i.e. those that belong are a neighbor of p in only one of the two models.

Lemma 9.2. *Let $0 \leq y_n < R$ be such that $R - y_n \rightarrow \infty$ and write $p_n = (x_n, y_n)$. Then we have, as $n \rightarrow \infty$,*

$$\mu(\mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n)) = \Theta(1) \cdot \begin{cases} e^{(1/2-\alpha)R+\alpha y_n}, & \text{if } \alpha < 3/2 \\ (R - y_n)e^{3y_n/2-R}, & \text{if } \alpha = 3/2 \\ e^{3y_n/2-R}, & \text{if } \alpha > 3/2 \end{cases}.$$

Proof. Let $r_n := R - y$. Lemma 2.2 implies that for such a p_n , if a point p belongs to $\mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n) \cap \mathcal{R}([0, r_n])$ then

$$|x_n - x| = \Theta(1) \cdot e^{\frac{3}{2}(y_n+y)-R}.$$

Now, if $p \in [r_n, r_n + 2 \ln \frac{\pi}{2}]$ and also $p \in \mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n)$, then

$$|x_n - x|_n = \frac{\pi}{2}e^{R/2} - e^{\frac{1}{2}(y_n+y)}.$$

Finally, (87) implies that no point in $\mathcal{R}([r_n + 2 \ln \frac{\pi}{2}, R])$ belongs to $\mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n)$. We first compute the expected number of points in $p \in \mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n)$ that have $R - y \leq r_n$. The result depends on the value of α , yielding the following three cases

$$\begin{aligned} \mu(\mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n) \cap \mathcal{R}([0, r_n])) &= \Theta(1) \cdot e^{3y_n/2-R} \int_0^{r_n} e^{(3/2-\alpha)y} dy \\ &= \Theta(1) \cdot \begin{cases} e^{(1/2-\alpha)R+\alpha y_n}, & \text{if } \alpha < 3/2 \\ (R - y_n)e^{3y_n/2-R}, & \text{if } \alpha = 3/2 \\ e^{3y_n/2-R}, & \text{if } \alpha > 3/2 \end{cases} \end{aligned}$$

Next we compute the number of remaining points in $\mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n)$,

$$\begin{aligned} \mu(\mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n) \cap \mathcal{R}([r_n, R])) &= \frac{\nu\alpha}{\pi} \int_{r_n}^{r_n+2 \ln \frac{\pi}{2}} \left(\frac{\pi}{2}e^{R/2} - e^{\frac{1}{2}(y_n+y)} \right) e^{-\alpha y} dy \\ &= O(1) \cdot e^{R/2} \int_{r_n}^{r_n+2 \ln \frac{\pi}{2}} e^{-\alpha y} dy = O(1) \cdot e^{R/2} e^{-\alpha r_n} \end{aligned}$$

$$= O(1) \cdot e^{(1/2-\alpha)R+\alpha y_n}.$$

Now note that for any $\alpha > 3/2$, we have

$$((1/2 - \alpha)R + \alpha y_n) - (3y_n/2 - R) = (3/2 - \alpha)(R - y_n) \rightarrow -\infty,$$

by our assumption on y_n . For $\alpha = 3/2$, these two quantities are equal. From these observations, we deduce that

$$\mu(\mathcal{B}(p_n) \triangle \mathcal{B}_\infty(p_n)) = \Theta(1) \cdot \begin{cases} e^{(1/2-\alpha)R+\alpha y_n}, & \text{if } \alpha < 3/2 \\ r_n e^{3y_n/2-R}, & \text{if } \alpha = 3/2 \\ e^{3y_n/2-R}, & \text{if } \alpha > 3/2 \end{cases}.$$

□

9.2 Equivalence clustering G_{Po} and G_{box}

Here we prove Proposition 6.3. We first note that Lemma 3.1 and Lemma 3.2 imply the following

$$\mathbb{E}[N_{\text{Po}}(k_n)] = \Theta(1) n k_n^{-(2\alpha+1)}, \quad (88)$$

and

$$\mathbb{E}[N_{\text{box}}(k_n)] = \Theta(1) n k_n^{-(2\alpha+1)}. \quad (89)$$

Moreover,

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}[N_{\text{Po}}(k_n)]}{\mathbb{E}[N_{\text{box}}(k_n)]} = 1. \quad (90)$$

Recall that Proposition 6.3 states

$$\lim_{n \rightarrow \infty} s(k_n)^{-1} \mathbb{E}[|c^*(k_n; G_{\text{Po}}) - c^*(k_n; G_{\text{box}})|] = 0.$$

Next recall the definition of $\mathcal{K}_C(k_n)$

$$\mathcal{K}_C(k_n) = \left\{ y \in \mathbb{R}_+ : \frac{k_n - C\sqrt{k_n \log(k_n)}}{\xi} \vee 0 \leq e^{\frac{y}{2}} \leq \frac{k_n + C\sqrt{k_n \log(k_n)}}{\xi} \wedge e^{R/2} \right\},$$

and (79)

$$\tilde{c}_{\text{box}}(k_n) = \frac{\tilde{T}_{\text{box}}(k_n, C)}{\binom{k_n}{2} \mathbb{E}[N_{\text{box}}(k_n)]},$$

where $\tilde{T}_{\text{box}}(k_n, C)$ counts for all nodes $p = (x, y)$ with $y \in \mathcal{K}_C(k_n)$ the pairs (p_1, p_2) that form a triangle with p , with the exception that it considers $p_2 \in \mathcal{B}_\infty(p_1) \cap \mathcal{R}$ instead of $\mathcal{B}_{\text{box}}(p_1)$. Then using Corollary 8.2 we get

$$\mathbb{E}[|c^*(k_n; G_{\text{Po}}) - c^*(k_n; G_{\text{box}})|] \leq \mathbb{E}[|c^*(k_n; G_{\text{Po}}) - \tilde{c}_{\text{box}}(k_n)|] + o(s(k_n)),$$

and hence it is enough to prove that

$$\lim_{n \rightarrow \infty} s(k_n)^{-1} \mathbb{E}[|c^*(k_n; G_{\text{Po}}) - \tilde{c}_{\text{box}}(k_n)|] = 0.$$

The following lemma will be frequently used in the proof of Proposition 6.3.

Lemma 9.3. *Let $t, r \in \mathbb{R}$ be fixed and let $\hat{\rho}(y, k)$ be any of the three probability functions $\rho_{\text{Po}}(y, k)$, $\rho_{\text{box}}(y, k)$ or $\rho(y, k)$. Then for any sequence k_n of non-negative integers with $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$ and $C > 0$ large enough,*

$$\int_{\mathcal{K}_C} e^{ty} \hat{\rho}_n(y, k_n - r) e^{-\alpha y} dy = O(1) k_n^{-2\alpha-1+2t}$$

as $n \rightarrow \infty$.

Proof. Note that on $\mathcal{K}_C(k_n)$ we have that $e^{ty} = \Theta(k_n^{2t})$. Hence, by Lemma 3.3

$$\begin{aligned} \int_{\mathcal{K}_C} e^{ty} \hat{\rho}_n(y, k_n - r) e^{-\alpha y} dy &= \Theta(k_n^{2t}) \int_{\mathcal{K}_C} \hat{\rho}_n(y, k_n - r) e^{-\alpha y} dy \\ &= O(k_n^{2t}) (k_n - r)^{-(2\alpha+1)} = O(1) k_n^{-2\alpha-1+2t}. \end{aligned}$$

□

Proof of Proposition 6.3. To keep notation concise we abbreviate $\mathbb{E}[N_{\text{Po}}(k_n)]$ and $\mathbb{E}[N_{\text{box}}(k_n)]$ by $\bar{n}_{\text{Po}}(k_n)$ and $\bar{n}_{\text{box}}(k_n)$, respectively. We will also suppress the subscript n in most expressions regarding the graphs G_{Po} and G_{box} . Finally we will write

$$T_{\text{Po}}(p) = \sum_{(p_1, p_2) \in \mathcal{P} \setminus \{p\}}^{\neq} T_{\text{Po}}(p, p_1, p_2),$$

with

$$T_{\text{Po}}(p, p_1, p_2) = \mathbb{1}_{\{p_1 \in \mathcal{B}(p)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(p)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(p_1)\}}$$

to denote the triangle count function for p in G_{Po} . Then we have

$$\begin{aligned} \mathbb{E}[|c^*(k_n; G_{\text{Po}}) - \tilde{c}_{\text{box}}(k_n)|] &= \binom{k_n}{2}^{-1} \mathbb{E} \left[\left| \sum_{p \in \mathcal{P}} \frac{\mathbb{1}_{\{\deg_{\text{Po}}(p)=k_n\}}}{\bar{n}_{\text{Po}}(k_n)} T_{\text{Po}}(p) - \frac{\mathbb{1}_{\{\deg_{\text{box}}(p)=k_n\}}}{\bar{n}_{\text{box}}(k_n)} \tilde{T}_{\text{box}}(p) \right| \right] \\ &\leq \binom{k_n}{2}^{-1} \bar{n}_{\text{Po}}(k_n)^{-1} \mathbb{E} \left[\left| \sum_{p \in \mathcal{P}} \mathbb{1}_{\{\deg_{\text{Po}}(p)=k_n\}} T_{\text{Po}}(p) - \mathbb{1}_{\{\deg_{\text{box}}(p)=k_n\}} \tilde{T}_{\text{box}}(p) \right| \right] \\ &\quad + \binom{k_n}{2}^{-1} \left| \frac{1}{\bar{n}_{\text{Po}}(k_n)} - \frac{1}{\bar{n}_{\text{box}}(k_n)} \right| \mathbb{E} \left[\sum_{p \in \mathcal{P}} \mathbb{1}_{\{\deg_{\text{box}}(p)=k_n\}} \tilde{T}_{\text{box}}(p) \right] \end{aligned}$$

The last term can be rewritten as

$$\left| 1 - \frac{\bar{n}_{\text{Po}}(k_n)}{\bar{n}_{\text{box}}(k_n)} \right| \mathbb{E}[\tilde{c}_{\text{box}}(k_n)] = \left| 1 - \frac{\bar{n}_{\text{Po}}(k_n)}{\bar{n}_{\text{box}}(k_n)} \right| \gamma(k_n)(1 + o(1)),$$

where we used Proposition 6.5 (See Section 7). The first term in this product converges to zero by Lemma ?? while the second term scales as $s(k_n)$. Hence

$$\left| 1 - \frac{\bar{n}_{\text{Po}}(k_n)}{\bar{n}_{\text{box}}(k_n)} \right| \mathbb{E}[\tilde{c}_{\text{box}}(k_n)] = o(s(k_n)),$$

and therefore we are left to analyze the other term. By the Campbell-Mecke formula we have that

$$\begin{aligned} &\mathbb{E} \left[\left| \sum_{p \in \mathcal{P}} \mathbb{1}_{\{\deg_{\text{Po}}(p)=k_n\}} T_{\text{Po}}(p) - \mathbb{1}_{\{\deg_{\text{box}}(p)=k_n\}} \tilde{T}_{\text{box}}(p) \right| \right] \\ &= \int_{\mathcal{R}} \mathbb{E} \left[\left| \mathbb{1}_{\{\deg_{\text{Po}}(y)=k_n\}} T_{\text{Po}}(y) - \mathbb{1}_{\{\deg_{\text{box}}(y)=k_n\}} \tilde{T}_{\text{box}}(y) \right| \right] f(x, y) dy dx. \end{aligned}$$

Since

$$\begin{aligned} \mathbb{E} \left[\frac{\mathbb{1}_{\{\deg_{\text{Po}}(y)=k_n\}}}{\bar{n}_{\text{Po}}(k_n)} T_{\text{Po}}(y) \right] &\leq \binom{k_n}{2} \rho_{\text{Po}}(y, k_n) \bar{n}_{\text{Po}}(k_n)^{-1} \\ &= \binom{k_n}{2} \rho_{\text{Po}}(y, k_n) \Theta(\bar{n}_{\text{box}}(k_n)^{-1}) \end{aligned}$$

$$= \Theta(n^{-1}k_n^{2\alpha+3}) \rho_{\text{Po}}(y, k_n)$$

and similar for the other term, it follows that

$$\begin{aligned} \mathbb{E} \left[\left| \frac{\mathbb{1}_{\{\deg_{\text{Po}}(y)=k_n\}}}{\bar{n}_{\text{Po}}(k_n)} T_{\text{Po}}(y) - \frac{\mathbb{1}_{\{\deg_{\text{box}}(y)=k_n\}}}{\bar{n}_{\text{Po}}(k_n)} \tilde{T}_{\text{box}}(y) \right| \right] \\ \leq \Theta(n^{-1}k_n^{2\alpha+3}) (\rho_{\text{Po}}(y, k_n) + \rho_{\text{box}}(y, k_n)). \end{aligned}$$

Therefore, by a concentration of heights argument (c.f. first statement of Lemma ??), it is enough to consider the integral

$$n \int_{\mathcal{K}_C(k_n)} \mathbb{E} \left[\left| \mathbb{1}_{\{\deg_{\text{Po}}(y)=k_n\}} T_{\text{Po}}(y) - \mathbb{1}_{\{\deg_{\text{box}}(y)=k_n\}} \tilde{T}_{\text{box}}(y) \right| \right] e^{-\alpha y} dy, \quad (91)$$

where we also used that $f(x, y)$ is simply a constant multiple of the function $e^{-\alpha y}$. Since $\binom{k_n}{2} \bar{n}_{\text{Po}}(k_n) = \Theta(nk_n^{-(2\alpha-1)})$ we have to show that

$$\lim_{n \rightarrow \infty} k_n^{2\alpha-1} s(k_n)^{-1} \int_{\mathcal{K}_C(k_n)} \mathbb{E} \left[\left| \mathbb{1}_{\{\deg_{\text{Po}}(y)=k_n\}} T_{\text{Po}}(y) - \mathbb{1}_{\{\deg_{\infty}(y)=k_n\}} \tilde{T}_{\text{box}}(y) \right| \right] e^{-\alpha y} dy = 0.$$

For $\alpha > 3/4$, $s_{3/4}(k_n) = \log(k_n)^{-1} s_{\alpha}(k_n) = o(s_{\alpha}(k_n))$ and thus it suffices to prove the following two cases:

1. if $1/2 < \alpha \leq 3/4$, then

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} \int_{\mathcal{K}_C(k_n)} \mathbb{E} \left[\left| \mathbb{1}_{\{\deg_{\text{Po}}(y)=k_n\}} T_{\text{Po}}(y) - \mathbb{1}_{\{\deg_{\text{box}}(y)=k_n\}} \tilde{T}_{\text{box}}(y) \right| \right] e^{-\alpha y} dy = 0,$$

2. if $3/4 < \alpha$, then

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \int_{\mathcal{K}_C(k_n)} \mathbb{E} \left[\left| \mathbb{1}_{\{\deg_{\text{Po}}(y)=k_n\}} T_{\text{Po}}(y) - \mathbb{1}_{\{\deg_{\text{box}}(y)=k_n\}} \tilde{T}_{\text{box}}(y) \right| \right] e^{-\alpha y} dy = 0.$$

We shall proceed by expanding the integrand and analyzing the individual terms. With a slight abuse of notation we shall write y instead of $(0, y)$ in an expression such as $\mathcal{B}(y)$. In addition we write $D_{\text{Po}}(y, k_n; \mathcal{P})$ for the indicator which is equal to 1 if and only if $\mathcal{B}(y)$ contains k_n points from $\mathcal{P} \setminus \{(0, y)\}$. We define $D_{\text{box}}(y, k_n; \mathcal{P})$ analogously for the ball $\mathcal{B}_{\text{box}}(y)$. It is important to note that for any $p' \in \mathcal{R}$ it holds that $p' \in \mathcal{B}_{\text{box}}(y) \iff p' \in \mathcal{B}_{\infty}(y)$.

We need to split the integrand over several terms and then analyze each of these separately. Applying the Campbell-Mecke formula yields

$$\begin{aligned} \mathbb{E} \left[\left| \mathbb{1}_{\{\deg_{\text{Po}}(y)=k_n\}} P_{\text{Po}}(y) - \mathbb{1}_{\{\deg_{\infty}(y)=k_n\}} \tilde{T}_{\text{box}}(y) \right| \right] \leq \\ \mathbb{E} \left[\sum_{(p_1, p_2) \in \mathcal{P} \setminus \{(0, y)\}}^{\neq} |D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) T_{\text{Po}}(y, p_1, p_2) \right. \\ \left. - D_{\text{box}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \tilde{T}_{\text{box}}(y, p_1, p_2) \right| \right], \end{aligned}$$

where the sum ranges over all distinct pairs of points in $\mathcal{P} \setminus \{(0, y)\}$. In what follows, we will set $\mathcal{B}_{\text{Po} \triangle \infty}(p') = \mathcal{B}(p') \triangle (\mathcal{B}_{\infty}(p') \cap \mathcal{R})$ and $\mathcal{B}_{\text{Po} \cap \text{box}}(p') = \mathcal{B}(p') \cap \mathcal{B}_{\text{box}}(p')$ and observe that $\mathcal{B}_{\text{Po} \cap \text{box}}(y) = \mathcal{B}(y) \cap \mathcal{B}_{\infty}(y)$. We will now bound the sum that is inside the expectation. We will split the sum into different parts, depending on combinations of $p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\}$ for which only one of the two terms of the difference is non-zero. Clearly, for this we need that either $p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)$ and $p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_1)$ or $p_1 \in \mathcal{B}_{\text{Po} \triangle \infty}(y)$ and $p_2 \in \mathcal{B}_{\text{Po} \cap \text{box}}(p_1)$. We will consider the following four cases:

1. $p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)$ and $p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_1)$
 - (a) $y_1, y_2 < (1 - \varepsilon)R \wedge (R - y)$
 - (b) $y_1 \geq (1 - \varepsilon)R \wedge (R - y)$
2. $p_1 \in \mathcal{B}(y) \setminus \mathcal{B}_\infty(y)$ with $y_1 < K$ and $p_2 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)$.
3. $p_1 \in \mathcal{B}_{\text{Po} \triangle \infty}(y)$ with $y_1 \geq K$ and $p_2 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)$,

where K in the last two cases is the constant from Lemma 2.2.

Observe that when $y_1 < (1 - \varepsilon)R \wedge (R - y)$ and $y_2 \geq (1 - \varepsilon)R \wedge (R - y)$ it follows from Corollary 9.1 that $p_2 \in \mathcal{B}_{\text{Po} \cap \text{box}}(p_1)$ and thus we do not have to consider this case when $p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)$ and $p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_1)$. Similarly, when $y_1 \geq K$ and $p_1 \in \mathcal{B}_{\text{Po} \triangle \infty}(y)$ Corollary 9.1 implies that $p_1 \in \mathcal{B}(y) \setminus \mathcal{B}_\infty(y)$ which explains the setting of case 2.

We can now bound the sum by the following expression:

$$\sum_{\substack{\neq \\ (p_1, p_2) \in \mathcal{P} \setminus \{(0, y)\}}} |D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) T_{\text{Po}}(y, p_1, p_2) - D_{\text{box}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \tilde{T}_{\text{box}}(y, p_1, p_2)|$$

$$\leq \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y_1, y_2 < (1 - \varepsilon)R \wedge (R - y)}} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_1)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \quad (92)$$

$$+ \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y_1, y_2 < (1 - \varepsilon)R \wedge (R - y)}} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_1)\}} D_{\text{box}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \quad (93)$$

$$+ \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y_1 \geq (1 - \varepsilon)R \wedge (R - y)}} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_1) \cap \mathcal{B}(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \quad (94)$$

$$+ \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y_1 \geq (1 - \varepsilon)R \wedge (R - y)}} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_1) \cap \mathcal{B}(y)\}} D_{\text{box}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \quad (95)$$

$$+ \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y(p_1) \geq K}} \mathbb{1}_{\{p_1 \in \mathcal{B}(y) \setminus \mathcal{B}_\infty(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \quad (96)$$

$$+ \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y(p_1) \geq K}} \mathbb{1}_{\{p_1 \in \mathcal{B}(y) \setminus \mathcal{B}_\infty(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(y)\}} D_{\text{box}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \quad (97)$$

$$+ \sum_{\substack{\neq \\ p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y(p_1) < K}} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \triangle \infty}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(y)\}}. \quad (98)$$

In the following paragraphs we will give upper bounds on the expected values of each one of these partial sums.

The sums (92) and (93) We will analyze (92). The analysis of the other sum (93) is similar. Note first that for any two points p_1, p_2 the following holds: $p_1 \in \mathcal{B}(y)$ and $p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_1) \cap \mathcal{B}(y)$, then $p_2 \in \mathcal{B}(y)$ and $p_1 \in \mathcal{B}_{\text{Po} \triangle \infty}(p_2) \cap \mathcal{B}(y)$. Using this symmetry, it suffices to consider distinct pairs $(p_1, p_2) \in \mathcal{P} \setminus \{(0, y)\}$ with $0 \leq y_2 \leq y_1 \leq R - y$. Let \mathcal{D} denote the set of these pairs.

We are going to consider several sub-cases and, thereby, split the domain \mathcal{D} into the corresponding sub-domains. Let $\omega = \omega(n) \rightarrow \infty$ as $n \rightarrow \infty$ be a slowly growing function and set $y_\omega := y + \omega$. We let

$$\begin{aligned}\mathcal{D}_1 &= \{(p_1, p_2) \in \mathcal{D} \cap \mathcal{P} : y \leq y_1 \leq R/2, y_\omega \leq y_2 \leq y_1\}, \\ \mathcal{D}_2 &= \{(p_1, p_2) \in \mathcal{D} \cap \mathcal{P} : y_1 \leq R/2, y_2 \leq y_\omega\} \text{ and} \\ \mathcal{D}_3 &= \{(p_1, p_2) \in \mathcal{D} \cap \mathcal{P} : R/2 < y_1 \leq R - y, y_2 \leq y_1\}.\end{aligned}$$

Note that $\mathcal{D} \subseteq \mathcal{D}_1 \cup \mathcal{D}_2 \cup \mathcal{D}_3$. Hence, we can write

$$\begin{aligned}\mathbb{E} &\left[\sum_{\substack{p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y_1, y_2 \leq (1-\varepsilon)R \wedge (R-y)}} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\ &\leq \sum_{i=1}^3 \mathbb{E} \left[\sum_{(p_1, p_2) \in \mathcal{D}_i} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} \cdot D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right].\end{aligned}\tag{99}$$

We bound each one of the above three summands as follows:

$$\begin{aligned}\mathbb{E} &\left[\sum_{(p_1, p_2) \in \mathcal{D}_1} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\ &\leq \mathbb{E} \left[\sum_{(p_1, p_2) \in \mathcal{D}_1} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] := \mathcal{I}_n^{(1)}(y),\end{aligned}\tag{100}$$

$$\begin{aligned}\mathbb{E} &\left[\sum_{(p_1, p_2) \in \mathcal{D}_2} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\ &\leq \mathbb{E} \left[\sum_{(p_1, p_2) \in \mathcal{D}_2} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] := \mathcal{I}_n^{(2)}(y)\end{aligned}\tag{101}$$

and

$$\begin{aligned}\mathbb{E} &\left[\sum_{(p_1, p_2) \in \mathcal{D}_3} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\ &\leq \mathbb{E} \left[\sum_{(p_1, p_2) \in \mathcal{D}_3} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] := \mathcal{I}_n^{(3)}(y).\end{aligned}\tag{102}$$

We will bound each term using the Campbell-Mecke formula and show for $i = 1, 2, 3$ that for $1/2 < \alpha < 3/4$

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(i)}(y) e^{-\alpha} dy = 0,\tag{103}$$

and for $\alpha \geq 3/4$

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(i)}(y) e^{-\alpha} dy = 0.\tag{104}$$

For the first term (100), we note that

$$\mathbb{E}[D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\})] = \rho_{\text{Po}}(y, k_n - 2).$$

and hence $\mathcal{I}_n^{(1)}(y)$ becomes

$$\rho_{\text{Po}}(y, k_n - 2) \int_{-I_n}^{I_n} \int_y^{R/2} \int_{-I_n}^{I_n} \int_{y_\omega}^{y_1} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y)\}} e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1. \quad (105)$$

Next, Lemma 2.2 implies that for $y' \leq R - y$, we have that if $(x', y') \in \mathcal{B}(y)$, then $|x'| < (1 + K)e^{y/2+y'/2}$, where $K > 0$ is as in Lemma 2.2. Using these observations, we obtain:

$$\begin{aligned} \mathbb{E} \left[\sum_{p_1, p_2 \in \mathcal{D}_1} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}((0, y))\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}(y)\}} \cdot D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\ = \rho_{\text{Po}}(y, k_n - 2) e^y \int_y^{R/2} e^{y_1/2} \int_{y_\omega}^{y_1} e^{y_2/2} e^{-\alpha y_2} \cdot e^{-\alpha y_1} dy_2 dy_1. \end{aligned}$$

Now, the double integral becomes

$$\begin{aligned} \int_y^{R/2} e^{y_1/2} \int_{y_\omega}^{y_1} e^{y_2/2} e^{-\alpha y_2} \cdot e^{-\alpha y_1} dy_2 dy_1 &= \\ O(1) \cdot \int_y^{R/2} e^{y_1/2-\alpha y_1} \cdot e^{(1/2-\alpha)y_\omega} dy_1 & \\ = O(1) \cdot e^{(1/2-\alpha)y_\omega} \cdot \int_y^{R/2} e^{y_1/2-\alpha y_1} dy_1 & \\ = O(1) \cdot e^{(1/2-\alpha)y_\omega + (1/2-\alpha)y} & \\ \ll e^{(1-2\alpha)y}, & \end{aligned} \quad (106)$$

since $y_\omega = y + \omega$ and $\omega \rightarrow \infty$. We then deduce that

$$\begin{aligned} \mathbb{E} \left[\sum_{p_1, p_2 \in \mathcal{D}_1} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \cap \text{box}}((0, y))\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}(y)\}} \cdot D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\ \ll \rho_{\text{Po}}(y, k_n - 2) e^{(1-2\alpha)y}. \end{aligned} \quad (107)$$

We now integrate this with respect to y and determine its contribution to (91);

$$\begin{aligned} \int_{\mathcal{K}_C(k_n)} \rho_{\text{Po}}(y, k_n - 2) e^{(1-2\alpha)y} e^{-\alpha y} dy dx \\ = O(k_n^{-6\alpha+1}) \end{aligned}$$

where we used Lemma 9.3 with $t = 1 - 2\alpha$.

Since $1 - 6\alpha + \min\{6\alpha - 3, 2\alpha\} < 0$ for all $\alpha > 1/2$ we deduce that for $1/2 < \alpha < 3/4$

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(1)}(y) e^{-\alpha y} dy = 0,$$

while for $\alpha \geq 3/4$

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(1)}(y) e^{-\alpha y} dy = 0.$$

We will now bound the term in (101). Using similar observations as for the previous term we get that $\mathcal{I}_n^{(2)}(y)$ equals

$$\rho_{\text{Po}}(y, k_n - 2) \int_{-I_n}^{I_n} \int_0^{R/2} \int_{-I_n}^{I_n} \int_0^{y_\omega} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po} \triangle \infty}((0, y))\}} e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1.$$

Now, Lemma 2.2 implies that for $y_2 \leq R - y_1$, we have that if $(x_2, y_2) \in \mathcal{B}_{\text{Po}\Delta\infty}((x_1, y_1))$, then x_2 lies in an interval of length $Ke^{3y/2+3y'/2-R}$, where $K > 0$ is again the constant in Lemma 2.2. Using these observations we obtain:

$$\mathcal{I}_n^{(2)}(y) = \rho_{\text{Po}}(y, k_n - 2)e^{y/2} \int_0^{R/2} e^{y_1/2+3y_1/2} \int_0^{y_\omega} e^{3y_2/2-R} e^{-\alpha y_2} \cdot e^{-\alpha y_1} dy_2 dy_1. \quad (108)$$

The integrals satisfy

$$\begin{aligned} & e^{-R} \left(\int_0^{R/2} e^{(2-\alpha)y_1} dy_1 \right) \left(\int_0^{y_\omega} e^{(3/2-\alpha)y_2} dy_2 \right) \\ &= O(1) e^{-R} \left(\begin{cases} e^{(1-\alpha/2)R} & \text{if } \frac{1}{2} < \alpha < 2 \\ R & \text{if } \alpha \geq 2 \end{cases} \right) \left(\begin{cases} e^{(3/2-\alpha)y_\omega} & \text{if } \frac{1}{2} < \alpha < \frac{3}{2} \\ y & \text{if } \alpha \geq \frac{3}{2} \end{cases} \right) \\ &= O(1) \begin{cases} e^{-\frac{\alpha}{2}R} e^{(3/2-\alpha)y} & \text{if } \frac{1}{2} < \alpha < \frac{3}{2} \\ (y + \omega(n))e^{-\frac{\alpha}{2}R} & \text{if } \frac{3}{2} \leq \alpha < 2 \\ (y + \omega(n))Re^{-R} & \text{if } \alpha \geq 2 \end{cases}. \end{aligned}$$

Since $y_\omega := y + \omega(n) \leq R = O(\log(n))$ we conclude that on $\mathcal{K}_C(k_n)$

$$\mathcal{I}_n^{(2)}(y) = O(1) \rho_{\text{Po}}(y, k_n - 2) \begin{cases} n^{-\alpha} k_n^{3-2\alpha} & \text{if } \frac{1}{2} < \alpha < \frac{3}{2} \\ n^{-\alpha} \log(n) & \text{if } \frac{3}{2} \leq \alpha < 2 \\ n^{-2} \log(n)^2 & \text{if } \alpha \geq 2 \end{cases},$$

and hence

$$\begin{aligned} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(2)}(y) e^{-\alpha y} dy &= O(1) k_n^{-(2\alpha+1)} \begin{cases} n^{-\alpha} k_n^{3-2\alpha} & \text{if } \frac{1}{2} < \alpha < \frac{3}{2} \\ n^{-\alpha} \log(n) & \text{if } \frac{3}{2} \leq \alpha < 2 \\ n^{-2} \log(n)^2 & \text{if } \alpha \geq 2 \end{cases} \\ &= O(1) \begin{cases} n^{-\alpha} k_n^{2-4\alpha} & \text{if } \frac{1}{2} < \alpha < \frac{3}{2} \\ n^{-\alpha} \log(n) k_n^{-(2\alpha+1)} & \text{if } \frac{3}{2} \leq \alpha < 2 \\ n^{-2} \log(n)^2 k_n^{-(2\alpha+1)} & \text{if } \alpha \geq 2 \end{cases}. \end{aligned}$$

Now for $1/2 < \alpha < 3/4$ it holds that $4\alpha^2 - \alpha + 1 > 0$. Hence since $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right)$, we have

$$k_n^{6\alpha-3} n^{-\alpha} k_n^{2-4\alpha} = n^{-\alpha} k_n^{2\alpha-1} = O\left(n^{-\alpha+\frac{2\alpha-1}{2\alpha+1}}\right) = O\left(k_n^{-\frac{4\alpha^2-\alpha+1}{2\alpha+1}}\right) = o(1),$$

from which we deduce that

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(2)}(y) e^{-\alpha y} dy = 0.$$

For $\alpha \geq 3/4$ we have that both $n^{-\alpha} \log(n) k_n^{-1}$ and $n^{-2} \log(n)^2 k_n^{-1}$ converge to zero as $n \rightarrow \infty$ and hence in this case

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(2)}(y) e^{-\alpha y} dy = 0.$$

We will now consider the term in (102). Recall that \mathcal{D}_3 consists of all pairs $(p_1, p_2) \in \mathcal{D}$ such that $R/2 < y_1 \leq (1-\varepsilon)R \wedge (R-y)$ and $y_1 \leq y_\omega$ with the property that $p_1 \in \mathcal{B}(y)$ and $p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)$. So, in particular, $p_2 \in (\mathcal{B}(p_1) \cup \mathcal{B}_\infty(p_1)) \cap \mathcal{B}(y)$.

We will consider this intersection more closely. We use Lemma 2.2 to define a ball around p_1 that contains both $\mathcal{B}(p_1)$ and $\mathcal{B}_\infty(p_1)$. For $K > 0$, we define, for any point $p_1 = (x_1, y_1) \in \mathbb{R} \times \mathbb{R}_+$,

$$\check{\mathcal{B}}_{\text{Po}}(p_1) := \{(x', y') : y' < R - y_1, |x_1 - x'| < (1+K)e^{\frac{1}{2}(y_1+y')}\}. \quad (109)$$

It is an implication of Lemma 2.2 that

$$(\mathcal{B}(p_1) \cup \mathcal{B}_\infty(p_1)) \cap \mathcal{R}([0, R - y_1]) \subseteq \check{\mathcal{B}}_{\text{Po}}(p_1)$$

Therefore, any point $p_2 = (x_2, y_2) \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)$ with $y_2 \leq R - y_1$ must belong to $\check{\mathcal{B}}_{\text{Po}}(p_1) \cap \check{\mathcal{B}}_{\text{Po}}(y)$.

We will use this in order to derive a lower bound on y_2 as a function of x_1, y_1 . Let us suppose without loss of generality that $x_1 < 0$. The left boundary of $\check{\mathcal{B}}_{\text{Po}}((0, y))$ is given by the equation $x' = (1 - K)e^{\frac{1}{2}(y+y')}$ whereas the right boundary of $\check{\mathcal{B}}_{\text{Po}}(p_1)$ is given by the curve having equation $x' = x_1 + (1 + K)e^{\frac{1}{2}(y_1+y')}$. The equation that determines the intersection point (\hat{x}, \hat{y}) of these curves is

$$x_1 + (1 + K)e^{(y_1+\hat{y})/2} = (1 - K)e^{(y+\hat{y})/2}.$$

We can solve the above for \hat{y}

$$|x_1| = (1 + K)e^{\hat{y}/2} \left(e^{y_1/2} + e^{y/2} \right).$$

But $y_1 > R/2$ and since $y \in \mathcal{K}_C(k_n)$, it follows that for sufficiently large n , $y \leq (1 + \varepsilon)R/(2\alpha + 1)$. So if ε is small enough depending on α , we have

$$|x_1| = (1 + K)e^{\hat{y}/2} \left(e^{y_1/2} + e^{y/2} \right) = (1 + K + o(1))e^{\hat{y}/2+y_1/2}.$$

Let c_K^2 denote the multiplicative term $1 + K + o(1)$, which appears in the above. The above yields

$$\hat{y} = \left(2 \log(|x_1|e^{-y_1/2}) - \log c_K \right) \vee 0 := \hat{y}(x_1, y_1). \quad (110)$$

In particular, note that $\hat{y} = 0$ if and only if $|x_1| \leq c_K e^{y_1/2}$. Moreover, since $p_1 \in \mathcal{B}(y)$ and $x_1 \leq R - y$, we also have that $|x_1| \leq e^{(y+y_1)/2}(1 + o(1))$. This upper bound on $|x_1|$ together with (110), imply that for n sufficiently large, we have $\hat{y} \leq y$. This observation will be used below, where we integrate over y_2 , thus ensuring that the integrals are non-zero.

We conclude that

$$p' \in \check{\mathcal{B}}_{\text{Po}}(y) \cap \check{\mathcal{B}}_{\text{Po}}((x_1, y_1)) \Rightarrow y' \geq \hat{y}(x_1, y_1),$$

which implies

$$\mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} \leq \mathbb{1}_{\{y_2 \geq \hat{y}(x_1, y_1), p_2 \in \check{\mathcal{B}}_{\text{Po}}((0, y))\}}. \quad (111)$$

If we integrate this over x_2, y_2 we get

$$\begin{aligned} \int_{-I_n}^{I_n} \int_0^{y_1} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} e^{-\alpha y_2} dy_2 dx_2 &\leq \int_{-I_n}^{I_n} \int_0^{y_1} \mathbb{1}_{\{y_2 \geq \hat{y}(x_1, y_1), p_2 \in \check{\mathcal{B}}_{\text{Po}}(y)\}} e^{-\alpha y_2} dy_2 dx_2 \\ &\leq (1 + K) \cdot e^{y/2} \int_{\hat{y}(x_1, y_1)}^{y_1} e^{y_2/2 - \alpha y_2} dy_2 \\ &= O(1) \cdot e^{y/2 + (1/2 - \alpha)\hat{y}(x_1, y_1)}. \end{aligned}$$

Note also that

$$\mathbb{E}[D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\})] = \rho_{\text{Po}}(y, k_n - 2),$$

uniformly over all $(p_1, p_2) \in \mathcal{D}_3$. Hence the Campbell-Mecke formula yields that $\mathcal{I}_n^{(3)}(y)$ equals:

$$\begin{aligned} O(1) \rho_{\text{Po}}(y, k_n - 2) e^{y/2} \int_{-I_n}^{I_n} \int_{R/2}^{(R-y) \wedge (1-\varepsilon)R} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} e^{(1/2 - \alpha)\hat{y}(x_1, y_1) - \alpha y_1} dy_1 dx_1 \\ = O(1) \rho_{\text{Po}}(y, k_n - 2) e^{y/2} \int_{-I_n}^{I_n} \int_{R/2}^{(R-y) \wedge (1-\varepsilon)R} \mathbb{1}_{\{p_1 \in \check{\mathcal{B}}_{\text{Po}}(y)\}} e^{(1/2 - \alpha)\hat{y}(x_1, y_1) - \alpha y_1} dy_1 dx_1. \end{aligned}$$

Due to the symmetry of $\check{\mathcal{B}}_{\text{Po}}(y)$, the integration over x_1 is:

$$O(1) \cdot e^{y/2} \cdot \int_0^{(1+K)e^{y/2+y_1/2}} e^{\hat{y}(x_1, y_1)(1/2-\alpha)} dx_1$$

We will split this integral into two parts according to the value of $\hat{y}(x_1, y_1)$:

$$\int_0^{(1+K)e^{y/2+y_1/2}} e^{\hat{y}(x_1, y_1)(1/2-\alpha)} dx_1 = \int_{c_K e^{y_1/2}}^{(1+K)e^{y/2+y_1/2}} e^{\hat{y}(x_1, y_1)(1/2-\alpha)} dx_1 + \int_0^{c_K e^{y_1/2}} dx_1.$$

The first integral becomes:

$$\begin{aligned} & \int_{c_K e^{y_1/2}}^{(1+K)e^{y/2+y_1/2}} e^{\hat{y}(x_1, y_1)(1/2-\alpha)} dx_1 = \int_{c_K e^{y_1/2}}^{(1+K)e^{y/2+y_1/2}} e^{\hat{y}(x_1, y_1)/2(1-2\alpha)} dx_1 \\ &= O(1) \cdot \int_{c_K e^{y_1/2}}^{(1+K)e^{y/2+y_1/2}} x_1^{1-2\alpha} e^{-\frac{y_1}{2}(1-2\alpha)} dx_1 \\ &= O(1) \cdot e^{-y_1/2+\alpha y_1} \cdot e^{\frac{(y+y_1)}{2}2(1-\alpha)} \\ &= O(1) \cdot e^{y_1/2+y(1-\alpha)}. \end{aligned}$$

The second integral trivially gives:

$$\int_0^{c_K e^{y_1/2}} dx_1 = O(1) \cdot e^{y_1/2} = O(1) \cdot e^{y_1/2+y(1-\alpha)}.$$

We conclude that

$$e^{y/2} \cdot \int_0^{(1+K)e^{y/2+y_1/2}} e^{\hat{y}(x_1, y_1)(1/2-\alpha)} dx_1 = O(1) \cdot e^{y_1/2+y(3/2-\alpha)}.$$

Now, we integrate this with respect to y_1 and get

$$e^{y(3/2-\alpha)} \int_{R/2}^{R-y} e^{(1/2-\alpha)y_1} dy_1 = O(1) \cdot e^{y(3/2-\alpha)} e^{(1/2-\alpha)R/2} = O(1) \cdot n^{1/2-\alpha} \cdot e^{y(3/2-\alpha)},$$

from which we deduce

$$\mathcal{I}_n^{(3)}(y) = O(1) \cdot n^{1/2-\alpha} e^{y(3/2-\alpha)} \rho_{\text{Po}}(y, k_n - 2). \quad (112)$$

We now apply Lemma 9.3 with $t = \frac{3}{2} - \alpha$ and get

$$\begin{aligned} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(3)}(y) e^{-\alpha y} dy &= O(1) n^{-(\alpha-\frac{1}{2})} \int_{\mathcal{K}_C(k_n)} e^{(3/2-\alpha)y} \rho_{\text{Po}}(y, k_n - 2) e^{-\alpha y} dy \\ &= O\left(n^{-(\alpha-\frac{1}{2})} k_n^{2-4\alpha}\right). \end{aligned}$$

Since for $\alpha > 1/2$, $k_n = O\left(n^{\frac{1}{2\alpha+1}}\right) = o\left(n^{1/2}\right)$ we have that $k_n^{6\alpha-3} k_n^{2-4\alpha} n^{-(\alpha-1/2)} = o(1)$ and hence for $1/2 < \alpha < 3/4$.

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(3)}(y) e^{-\alpha y} dx dy = 0,$$

For $\alpha \geq 3/4$ we observe that $2\alpha^2 + 2\alpha - 5/2 > 0$. Hence,

$$k_n^{2\alpha} n^{-(\alpha-\frac{1}{2})} k_n^{2-4\alpha} = O\left(n^{-(\alpha-1/2)} n^{\frac{2-2\alpha}{2\alpha+1}}\right) = O\left(n^{-\frac{2\alpha^2+2\alpha-5/2}{2\alpha+1}}\right) = o(1).$$

and we get for $\alpha \geq 3/4$

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(3)}(y) e^{-\alpha y} dx dy = 0.$$

The sums (94) and (95) Again, we will only consider (94) since the analysis for the other term is similar. Recall that in this case, we consider pairs (p_1, p_2) , with $p_1 = (x_1, y_1)$ satisfying $y_1 \geq (R - y) \wedge (1 - \varepsilon)R$, and $p_1 \in \mathcal{B}(y)$, $p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)$. We split this into three sub-domains: i) $y_2 \geq R - y$; ii) $R - y_1 \leq y_2 \leq R - y$ and iii) $y_2 < R - y_1$. Similar to the analysis above we define

$$\begin{aligned}\mathcal{D}_1 &:= \{(p_1, p_2) : p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\}, y_1 \geq (1 - \varepsilon)R \wedge (R - y), R - y \leq y_2 \leq R\} \\ \mathcal{D}_2 &:= \{(p_1, p_2) : p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\}, y_1 \geq (1 - \varepsilon)R \wedge (R - y), R - y_1 \leq y_2 \leq R - y\} \\ \mathcal{D}_3 &:= \{(p_1, p_2) : p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\}, y_1 \geq (1 - \varepsilon)R \wedge (R - y), y_2 \leq R - y_1\}\end{aligned}$$

and write, for $i = 1, 2, 3$,

$$\mathcal{I}_n^{(i)}(y) := \mathbb{E} \left[\sum_{(p_1, p_2) \in \mathcal{D}_i} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} \cdot D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right].$$

In the first case, note that for $y \in \mathcal{K}_C(k_n)$ we have, for small enough ε and sufficiently large n , $2y \leq 2(1 + \varepsilon)\frac{R}{2\alpha+1} = o(R)$. Thus $y_1 + y_2 \geq 2(R - y) = \Omega(R)$ and thus $p_2 \in \mathcal{B}(p_1)$ for large enough n . Furthermore, $y_2 > R - y_1 + 2\ln(\pi/2)$, which implies that $p_2 \in \mathcal{B}_\infty(p_1)$ too. Hence, the contribution from these pairs is zero.

The Campbell-Mecke formula yields that:

$$\begin{aligned}\mathcal{I}_n^{(1)}(y) &= O(1) \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \times \\ &\quad \int_{-I_n}^{I_n} \int_{R-y}^R \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} \rho_{\text{Po}}(y, k_n - 2) \cdot e^{-\alpha(y_2+y_1)} dy_2 dx_2 dy_1 dx_1.\end{aligned}$$

We proceed to bound the integral:

$$\begin{aligned}&\int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \int_{-I_n}^{I_n} \int_{R-y}^R \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1 \\ &\leq \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \int_{-I_n}^{I_n} \int_{R-y}^R e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1 \\ &= \left(\int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R e^{-\alpha y_1} dy_1 dx_1 \right) \left(\int_{-I_n}^{I_n} \int_{R-y}^R e^{-\alpha y_2} dy_2 dx_2 \right).\end{aligned}$$

We evaluate

$$\int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R e^{-\alpha y_1} dy_1 dx_1 = O(1) \cdot n \cdot e^{-\alpha R + ((\varepsilon R) \vee y)\alpha} = O(1) \cdot n \cdot e^{-\alpha R + \alpha y + \alpha \varepsilon R}$$

and

$$\int_{-I_n}^{I_n} \int_{R-y}^R e^{-\alpha y_2} dy_2 dx_2 = O(1) \cdot n \cdot e^{-\alpha R + \alpha y}.$$

Also, $n \cdot e^{-\alpha R} = O(1) \cdot e^{(1/2-\alpha)R}$, whereby we deduce that

$$\begin{aligned}&\int_{\mathcal{D}_1} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1 \\ &= O(1) \cdot e^{(1-2\alpha)R + 2\alpha y + \alpha \varepsilon R} = O(1) \cdot n^{2(1-2\alpha) + 2\alpha \varepsilon} \cdot e^{2\alpha y}.\end{aligned}$$

With these computations we obtain

$$\int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(1)}(y) e^{-\alpha y} dx dy = O(1) n^{2(1-2\alpha) + 2\alpha \varepsilon} \int_{\mathcal{K}_C(k_n)} e^{2\alpha y} \rho_{\text{Po}}(y, k_n - 2) e^{-\alpha y} dy dx$$

$$= O(1)n^{2(1-2\alpha)+2\alpha\varepsilon} k_n^{2\alpha-1}.$$

Thus, for $1/2 < \alpha < 3/4$, we have

$$k_n^{6\alpha-3} n^{2(1-2\alpha)+2\alpha\varepsilon} k_n^{2\alpha-1} = n^{2\alpha\varepsilon} \left(\frac{k_n^2}{n} \right)^{2(2\alpha-1)} = o(1),$$

provided that $\varepsilon = \varepsilon(\alpha) > 0$ is small enough, and hence for such ε

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(1)}(y) e^{-\alpha y} dx dy = 0.$$

When $\alpha \geq 3/4$ we have $2(1-2\alpha) < 1/2(4\alpha-1)$ and we get

$$k_n^{2\alpha} n^{2(1-2\alpha)+2\alpha\varepsilon} \cdot k_n^{2\alpha-1} \leq k_n^{4\alpha-1} n^{2(1-2\alpha)} n^{2\alpha\varepsilon} = o(1),$$

provided that ε is small enough, depending on α , so that

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(1)}(y) e^{-\alpha y} dx dy = 0.$$

We now consider the second sub-domain \mathcal{D}_2 . The Campbell-Mecke formula yields that:

$$\begin{aligned} \mathcal{I}_n^{(2)}(y) &= \mathbb{E} \left[\sum_{(p_1, p_2) \in \mathcal{D}_2} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1\}) \right] \\ &= O(1) \rho_{\text{Po}}(y, k_n - 2) \cdot \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \times \\ &\quad \int_{-I_n}^{I_n} \int_{R-y_1}^{R-y} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1. \end{aligned}$$

We bound the integral as follows:

$$\begin{aligned} &\int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \int_{-I_n}^{I_n} \int_{R-y_1}^{R-y} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1 \\ &\leq \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \int_{-I_n}^{I_n} \int_{R-y_1}^{R-y} \mathbb{1}_{\{p_2 \in \mathcal{B}(y)\}} e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1. \end{aligned}$$

Now, by Lemma 2.2,

$$\begin{aligned} &\int_{-I_n}^{I_n} \int_{R-y_1}^{R-y} \mathbb{1}_{\{p_2 \in \mathcal{B}(y)\}} \cdot e^{-\alpha y_2} dy_2 dx_2 = O(1) \cdot e^{y/2} \int_{R-y_1}^{R-y} e^{(1/2-\alpha)y_2} dy_2 \\ &= O(1) \cdot e^{y/2+(1/2-\alpha)(R-y_1)}. \end{aligned}$$

We then integrate with respect to y_1 :

$$\begin{aligned} &O(1) \cdot e^{y/2} \cdot \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} e^{(1/2-\alpha)(R-y_1)} e^{-\alpha y_1} dy_1 dx_1 \\ &\leq O(1) \cdot e^{y/2+(1/2-\alpha)R} \cdot \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R e^{(\alpha-1/2)y_1} e^{-\alpha y_1} dy_1 dx_1 \\ &= O(1) \cdot e^{y/2+(1-\alpha)R-((1-\varepsilon)R \wedge (R-y))/2} \\ &= O(1) \cdot e^{y/2+(1/2-\alpha)R+((\varepsilon R) \vee y)/2} \\ &= O(1) \cdot e^{y+(1/2-\alpha)R+\varepsilon R} = O(1) \cdot n^{1-2\alpha+\varepsilon} \cdot e^y. \end{aligned}$$

Therefore we get

$$\begin{aligned}
& \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(2)}(y) e^{-\alpha y} dx dy \\
&= O(n^{1-2\alpha+\varepsilon}) \int_{\mathcal{K}_C(k_n)} \rho_{\text{Po}}(y, k_n - 2) e^y e^{-\alpha y} dx dy \\
&= O(1) n^{1-2\alpha+\varepsilon} k_n^{-2\alpha+1},
\end{aligned}$$

where we used Lemma 9.3 with $t = 1$.

For $1/2 < \alpha < 3/4$, we have

$$k_n^{4\alpha-2} \cdot n^{1-2\alpha+\varepsilon} = n^\varepsilon \left(\frac{k_n^2}{n} \right)^{2\alpha-1} = o(1),$$

provided that $\varepsilon = \varepsilon(\alpha) > 0$ is small enough, yielding

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(2)}(y) e^{-\alpha y} dx dy = 0.$$

Similarly, for $\alpha > 3/4$ we have $2\alpha - 1 > 1/2$ and we get

$$k_n \cdot n^{1-2\alpha+\varepsilon} \ll n^{-1/2+\varepsilon} \cdot k_n = o(1),$$

provided that ε is small enough, so that

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(2)}(y) e^{-\alpha y} dx dy = 0.$$

For the third sub-domain \mathcal{D}_3 we shall use (111) which states that if $p_2 = (x_2, y_2) \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)$ and $y_2 \leq R - y_1$, then $y_2 \geq \hat{y}(x_1, y_1)$, where $\hat{y}(x_1, y_1) = (2 \log(|x_1| e^{-y_1/2}) - \log c_K) \vee 0$ (cf. (115)). Moreover, $p_2 \in \check{\mathcal{B}}_{\text{Po}}(p_1)$.

Again, we will use the Campbell-Mecke formula:

$$\begin{aligned}
\mathcal{I}_n^{(3)}(y) &= \mathbb{E} \left[\sum_{(p_1, p_2) \in \mathcal{D}_3} \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \cdot \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} \cdot D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\
&= O(1) \rho_{\text{Po}}(y, k_n - 2) \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \times \\
&\quad \int_{-I_n}^{I_n} \int_0^{R-y_1} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1
\end{aligned}$$

The inner integral with respect to $p_2 := (x_2, y_2)$ is

$$\begin{aligned}
& \int_{-I_n}^{I_n} \int_0^{R-y_1} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} e^{-\alpha y_2} dy_2 dx_2 \\
& \leq \int_{-I_n}^{I_n} \int_0^{R-y_1} \mathbb{1}_{\{y_2 \geq \hat{y}(x_1, y_1), p_2 \in \check{\mathcal{B}}_{\text{Po}}((0, y))\}} e^{-\alpha y_2} dy_2 dx_2 \\
& = O(1) e^{y/2} \int_{\hat{y}(x_1, y_1)}^{R-y_1} e^{y_2/2 - \alpha y_2} dy_2 \\
& = O(1) e^{y/2 + (1/2 - \alpha)\hat{y}(x_1, y_1)}.
\end{aligned}$$

Thus, we get

$$\begin{aligned} & \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R \mathbb{1}_{\{p_1 \in \mathcal{B}(y)\}} \int_{-I_n}^{I_n} \int_0^{R-y_1} \mathbb{1}_{\{p_2 \in \mathcal{B}_{\text{Po}\Delta\infty}(p_1) \cap \mathcal{B}(y)\}} \times \\ & \quad e^{-\alpha(y_1+y_2)} dy_2 dx_2 dy_1 dx_1 \\ & \leq O(1) \int_{-I_n}^{I_n} \int_{(1-\varepsilon)R \wedge (R-y)}^R e^{y/2+(1/2-\alpha)\hat{y}(x_1, y_1)} e^{-\alpha y_1} dy_1 dx_1. \end{aligned}$$

Due to symmetry, to bound the integral it is enough to integrate this with respect to x_1 from 0 to I_n . We will split this integral into two parts according to the value of $c(x_1, y_1)$:

$$\int_0^{I_n} e^{\hat{y}(x_1, y_1)(1/2-\alpha)} dx_1 = \int_{c_K e^{y_1/2}}^{I_n} e^{c(x_1, y_1)(1/2-\alpha)} dx_1 + \int_0^{c_K e^{y_1/2}} dx_1.$$

The first integral becomes:

$$\begin{aligned} & \int_{c_K e^{y_1/2}}^{I_n} e^{\hat{y}(x_1, y_1)(1/2-\alpha)} dx_1 = O(1) \cdot \int_{c_K e^{y_1/2}}^{I_n} x_1^{1-2\alpha} e^{-\frac{y_1}{2}(1-2\alpha)} dx_1 \\ & = \begin{cases} O(R) \cdot e^{-y_1/2+\alpha y_1} \cdot e^{\frac{R}{2}2(1-\alpha)} & \text{if } \alpha \leq 1 \\ O(1) \cdot e^{-y_1/2+\alpha y_1+2(1-\alpha)y_1/2} & \text{if } \alpha > 1 \end{cases} \\ & = \begin{cases} O(R) \cdot e^{(\alpha-1/2)y_1} \cdot n^{2(1-\alpha)} & \text{if } \alpha \leq 1 \\ O(1) \cdot e^{y_1/2} & \text{if } \alpha > 1 \end{cases}. \end{aligned}$$

The second integral trivially gives:

$$\int_0^{c_K e^{y_1/2}} dx_1 = O(1) \cdot e^{y_1/2}.$$

Putting these two together we conclude that

$$e^{y/2} \cdot \int_0^{I_n} e^{\hat{y}(x_1, y_1)(1/2-\alpha)} dx_1 = O(1) \cdot e^{y_1/2+y(3/2-\alpha)}.$$

Now, we integrate these with respect to y_1 :

$$\begin{aligned} & n^{2(1-\alpha)} \cdot \int_{(1-\varepsilon)R \wedge (R-y)}^R e^{(\alpha-1/2)y_1-\alpha y_1} dy_1 = O(1) \cdot n^{2(1-\alpha)} \cdot e^{-R/2+\varepsilon R/2+y/2} \\ & = O(1) \cdot n^{1-2\alpha+\varepsilon} \cdot e^{y/2}. \end{aligned}$$

Therefore, we conclude that

$$\mathcal{I}_n^{(3)}(y) = O(R) n^{1-2\alpha+\varepsilon(2\alpha-1)} e^{y/2} \rho_{\text{Po}}(y, k_n - 2)$$

and hence, using again Lemma 9.3,

$$\begin{aligned} & \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(3)}(y) e^{-\alpha y} dx dy = O(R) n^{1-2\alpha+\varepsilon(2\alpha-1)} \int_{\mathcal{K}_C(k_n)} e^{y/2} \rho_{\text{Po}}(y, k_n - 2) e^{-\alpha y} dx dy \\ & = O(R) n^{1-2\alpha+\varepsilon(2\alpha-1)} k_n^{-2\alpha+1}. \end{aligned}$$

It follows that for $\varepsilon = \varepsilon(\alpha)$ small enough

$$k_n^{6\alpha-3} R n^{1-2\alpha+\varepsilon(2\alpha-1)} k_n^{-2\alpha+1} = R n^{\varepsilon(2\alpha-1)} \left(\frac{k_n^2}{n} \right)^{2\alpha-1} = o(1)$$

and hence for $\alpha > 1/2$,

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(3)}(y) e^{-\alpha y} dx dy = 0.$$

Since $2\alpha - 1 \geq 1/2$ when $\alpha \geq 3/4$ it immediately follows that

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(3)}(y) e^{-\alpha y} dx dy = 0.$$

The sums (96) and (97) Again, the analysis for both terms are similar and we shall analyze (96). Let us set $p = (0, y)$. Recall that $\mathcal{B}_{\text{Po}\Delta\infty}(y) \cap \mathcal{R}([R - y + 2 \log(\frac{\pi}{2}), R]) = \emptyset$. Thus, the summand in (96) is equal to 0, when $y_1 > R - y + 2 \log(\pi/2)$.

Recall the definition of the extended ball $\check{\mathcal{B}}_{\text{Po}}(p)$ around p (109) that contains both $\mathcal{B}(p)$ and $\mathcal{B}_\infty(p)$

$$\check{\mathcal{B}}_{\text{Po}}(y) := \{p' : y' < R - y, |x'| < (1 + K)e^{\frac{1}{2}(y+y')}\},$$

and that we have $\mathbb{E}[D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\})] = \rho_{\text{Po}}(y, k_n - 2)$.

Further, observe that,

$$\mathcal{B}(y) \cap \mathcal{R}([0, R - y]) \subseteq \check{\mathcal{B}}_{\text{Po}}(y)$$

and

$$\mathcal{B}(y) \cap \mathcal{R}([R - y, R]) = \mathcal{R}([R - y, R]).$$

We thus conclude that

$$\mathcal{B}(y) \subseteq \check{\mathcal{B}}_{\text{Po}}(y) \cup \mathcal{R}([R - y, R]). \quad (113)$$

Hence, if we set

$$h_y(p_1) := \mathbb{1}_{\{p_1 \in \mathcal{B}(p) \setminus \mathcal{B}_\infty(y)\}} \cdot (\mu(\check{\mathcal{B}}_{\text{Po}}(p_1) \cap \check{\mathcal{B}}_{\text{Po}}(y)) + \mu(\mathcal{R}([R - y, R]))) ,$$

then

$$\begin{aligned} & \mathbb{1}_{\{p_1 \in \mathcal{B}(p) \setminus \mathcal{B}_\infty(y)\}} \cdot \mathbb{E} \left[\left(\sum_{p_2 \in \mathcal{P} \setminus \{p, p_1\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(p_1)\}} \right) \cdot D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\ &= O(1) \cdot \mathbb{1}_{\{p_1 \in \mathcal{B}(y) \setminus \mathcal{B}_\infty(y)\}} \cdot \mu(\mathcal{B}(y) \cap \mathcal{B}(p_1)) \rho_{\text{Po}}(y, k_n - 2) \\ &\leq O(1) \cdot h_y(p_1) \rho_{\text{Po}}(y, k_n - 2). \end{aligned}$$

To calculate the expectation of the above function we need to approximate the intersection of the two balls $\check{\mathcal{B}}_{\text{Po}}(y)$ and $\check{\mathcal{B}}_{\text{Po}}(p_1)$, where $p_1 = (x_1, y_1)$. Let us assume without loss of generality that $x_1 > 0$. The right boundary of $\check{\mathcal{B}}_{\text{Po}}(y)$ is given by the equation $x = x(y') = (1 + K)e^{\frac{1}{2}(y+y')}$ whereas the left boundary of $\check{\mathcal{B}}_{\text{Po}}(p_1)$ is given by the curve $x = x(y') = x_1 - (1 + K)e^{\frac{1}{2}(y_1+y')}$.

The equation that determines the intersecting point of the two curves is

$$x_1 - (1 + K)e^{(\hat{y}+y_1)/2} = (1 + K)e^{(\hat{y}+y)/2},$$

where \hat{y} is the y -coordinate of the intersecting point. We can solve the above for \hat{y}

$$x_1 = (1 + K)e^{\hat{y}/2} \left(e^{y/2} + e^{y_1/2} \right).$$

But since $p_1 = (x_1, y_1) \in \mathcal{B}_{\text{Po}\Delta\infty}(p)$, we also have $x_1 > e^{\frac{y+y_1}{2}}$. Therefore,

$$e^{\hat{y}/2} > \frac{1}{1 + K} \frac{e^{\frac{y+y_1}{2}}}{e^{y/2} + e^{y_1/2}} \geq \frac{1}{2(1 + K)} \frac{e^{\frac{y_1+y}{2}}}{e^{(y \vee y_1)/2}} > \frac{1}{2(1 + K)} e^{(y \wedge y_1)/2}. \quad (114)$$

The above yields

$$\hat{y} > (y \wedge y_1) - 2 \log(2(1 + K)) := \hat{y}(y_1, y). \quad (115)$$

which, in turn, implies the following

$$p \in \check{\mathcal{B}}_{\text{Po}}((0, y)) \cap \check{\mathcal{B}}_{\text{Po}}(p_1) \Rightarrow y(p) \geq \hat{y}(y_1, y). \quad (116)$$

We thus conclude that

$$\mathcal{B}(p_1) \cap \mathcal{B}(p) \subseteq (\check{\mathcal{B}}_{\text{Po}}(p) \cap \mathcal{R}([\hat{y}(y_1, y), R])) \cup \mathcal{R}([R - y, R]),$$

which in turn implies that

$$\mu(\check{\mathcal{B}}_{\text{Po}}(p_1) \cap \mathcal{B}(p)) \leq \mu(\check{\mathcal{B}}_{\text{Po}}(p) \cap \mathcal{R}([\hat{y}(y_1, y), R])) + \mu(\mathcal{R}([R - y, R])).$$

Therefore,

$$\begin{aligned} h_y(p_1, \mathcal{P}) &\leq \mathbb{1}_{\{p_1 \in \mathcal{B}(p) \setminus \mathcal{B}_\infty(p)\}} \mu(\check{\mathcal{B}}_{\text{Po}}(p) \cap \mathcal{R}([\hat{y}(y_1, y), R])) \\ &\quad + \mathbb{1}_{\{p_1 \in \mathcal{B}(p) \setminus \mathcal{B}_\infty(p)\}} \mu(\mathcal{R}([R - y, R])). \end{aligned}$$

Now, the Campbell-Mecke formula gives

$$\begin{aligned} &\mathbb{E} \left[\sum_{\substack{p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y(p_1) \geq K}} \mathbb{1}_{\{p_1 \in \mathcal{B}(y) \setminus \mathcal{B}_\infty(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(y)\}} D_{\text{Po}}(y, k_n - 2; \mathcal{P} \setminus \{p_1, p_2\}) \right] \\ &\leq \mathbb{E} \left[\left(\sum_{p_1 \in \mathcal{P}} h_y(p_1, \mathcal{P} \setminus \{p_1\}) \right) \right] \\ &= \frac{\nu\alpha}{\pi} \int_{\mathcal{R}} \mathbb{E}[h_y(p_1, \mathcal{P} \setminus \{p_1\})] e^{-\alpha y_1} dx_1 dy_1 \\ &\leq \frac{\nu\alpha}{\pi} \int_{\mathcal{R}} \mathbb{1}_{\{p_1 \in \mathcal{B}(p) \setminus \mathcal{B}_\infty(p)\}} \mu(\check{\mathcal{B}}_{\text{Po}}(p) \cap \mathcal{R}([\hat{y}(y_1, y), R])) e^{-\alpha y_1} dx_1 dy_1 \quad (117) \\ &\quad + \frac{\nu\alpha}{\pi} \int_{\mathcal{R}} \mathbb{1}_{\{p_1 \in \mathcal{B}(p) \setminus \mathcal{B}_\infty(p)\}} \mu(\mathcal{R}([R - y, R])) e^{-\alpha y_1} dx_1 dy_1. \quad (118) \end{aligned}$$

Recall that $(\mathcal{B}_{\text{Po} \triangle \infty}((0, y))) \cap \mathcal{R}([R - y + 2 \log(\frac{\pi}{2}), R]) = \emptyset$. We will first calculate the measures μ appearing in (117) and (118). The first one is:

$$\begin{aligned} \mu(\check{\mathcal{B}}_{\text{Po}}(y) \cap \mathcal{R}([c(y_1, y), R])) &\leq (1 + K) \frac{\nu\alpha}{\pi} \cdot e^{y/2} \int_{\hat{y}(y_1, y)}^R e^{-(\alpha - \frac{1}{2})y'} dy' \\ &= O\left(e^{\frac{y}{2} - (\alpha - \frac{1}{2})(y \wedge y_1)}\right). \end{aligned}$$

The second term is:

$$\mu(\mathcal{R}([R - y, R])) = \frac{\nu\alpha}{\pi} \int_{R-y}^R \pi e^{\frac{R}{2}} e^{-\alpha y'} dy' = O\left(e^{\frac{R}{2}} e^{-\alpha(R-y)}\right) = O\left(e^{\alpha y - (\alpha - \frac{1}{2})R}\right).$$

Using these, we get

$$\begin{aligned} &\int_{\mathcal{R}([0, R - y_n + 2 \ln \frac{\pi}{2}])} \mathbb{E}[h_y(p_1, \mathcal{P} \setminus \{p_1\})] e^{-\alpha y_1} dx_1 dy_1 \\ &= O(1) \int_{\mathcal{R}([0, R - y + 2 \ln \frac{\pi}{2}])} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \triangle \infty}(p)\}} e^{\frac{y}{2} - (\alpha - \frac{1}{2})(y \wedge y_1) - \alpha y_1} dx_1 dy_1 \quad (119) \end{aligned}$$

$$+ O(1) \int_{\mathcal{R}([0, R - y + 2 \ln \frac{\pi}{2}])} \mathbb{1}_{\{p_1 \in \mathcal{B}((0, y))\}} e^{\alpha y - (\alpha - \frac{1}{2})R - \alpha y_1} dx_1 dy_1. \quad (120)$$

Now, Lemma 2.2 implies that for any $y_1 \in [0, R - y + 2 \ln \frac{\pi}{2}]$, we have

$$\int_{-I_n}^{I_n} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \triangle \infty}(y)\}} dx_1 \leq 2K e^{\frac{3}{2}(y_1+y)-R}.$$

Therefore, (119) is

$$\begin{aligned} & O(1) \cdot e^{2y-R} \int_0^{R-y+2 \ln \frac{\pi}{2}} e^{\frac{3y_1}{2} - (\alpha - \frac{1}{2})(y_1 \wedge y) - \alpha y_1} dy_1 \\ &= O(1) \cdot e^{2y-R} \left(\int_0^y e^{\frac{3y_1}{2} - (2\alpha - \frac{1}{2})y_1} dy_1 + e^{-(\alpha - \frac{1}{2})y} \int_y^{R-y+2 \ln \frac{\pi}{2}} e^{(\frac{3}{2} - \alpha)y_1} dy_1 \right) \\ &= O(1) \left(\begin{cases} e^{(4-2\alpha)y-R}, & \text{if } \alpha < 1 \\ R \cdot e^{2y-R}, & \text{if } \alpha \geq 1 \end{cases} + \begin{cases} e^{-(\alpha - \frac{1}{2})R+y}, & \text{if } \alpha < 3/2 \\ R \cdot e^{2(2-\alpha)y-R}, & \text{if } \alpha \geq 3/2 \end{cases} \right). \end{aligned}$$

Similarly, for (120) we have

$$\begin{aligned} & \int_{\mathcal{R}([0, R-y+2 \ln \frac{\pi}{2}])} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po} \triangle \infty}((0, y))\}} e^{\alpha y - (\alpha - \frac{1}{2})R - \alpha y_1} dx_1 dy_1 \\ &= e^{\frac{3y}{2} - R + \alpha y - (\alpha - \frac{1}{2})R} \cdot \int_0^{R-y+2 \ln \frac{\pi}{2}} e^{\frac{3y_1}{2} - \alpha y_1} dy_1 \\ &= O(1) \cdot \begin{cases} e^{\frac{3y}{2} - R + \alpha y - (\alpha - \frac{1}{2})R + (\frac{3}{2} - \alpha)(R-y)}, & \text{if } \alpha < 3/2 \\ R \cdot e^{(\frac{3}{2} + \alpha)y - (\alpha + \frac{1}{2})R}, & \text{if } \alpha \geq 3/2 \end{cases} \\ &= O(1) \cdot \begin{cases} e^{-(2\alpha-1)R+2\alpha y}, & \text{if } \alpha < 3/2 \\ R \cdot e^{(\frac{3}{2} + \alpha)y - (\alpha + \frac{1}{2})R}, & \text{if } \alpha \geq 3/2 \end{cases}. \end{aligned}$$

We thus conclude, using $2(2 - \alpha)y \leq y$ for $\alpha > 3/2$, that

$$\mathbb{E} \left[\left(\sum_{p_1 \in \mathcal{P} \setminus \{p\}} h_y(p_1) \right) \right] \leq O(1) \cdot \left(\mathcal{I}_n^{(1)}(y) + \mathcal{I}_n^{(2)}(y) + \mathcal{I}_n^{(3)}(y) \right), \quad (121)$$

where

$$\begin{aligned} \mathcal{I}_n^{(1)}(y) &= \begin{cases} e^{(4-2\alpha)y-R}, & \text{if } \alpha < 1 \\ R \cdot e^{2y-R}, & \text{if } \alpha \geq 1 \end{cases}, \\ \mathcal{I}_n^{(2)}(y) &= \begin{cases} e^{-(\alpha - \frac{1}{2})R+y}, & \text{if } \alpha < 3/2 \\ R \cdot e^{y-R}, & \text{if } \alpha \geq 3/2 \end{cases} \\ \mathcal{I}_n^{(3)}(y) &= \begin{cases} e^{-(2\alpha-1)R+2\alpha y}, & \text{if } \alpha < 3/2 \\ R \cdot e^{(\frac{3}{2} + \alpha)y - (\alpha + \frac{1}{2})R}, & \text{if } \alpha \geq 3/2 \end{cases}. \end{aligned}$$

We proceed to calculate:

$$\int_{\mathcal{K}_C(k_n)} \mathbb{E} \left[\left(\sum_{p_1 \in \mathcal{P}} h_y(p_1, \mathcal{P} \setminus \{p_1\}) \right) \right] \cdot \rho_{\text{Po}}(y, k_n - 2) e^{-\alpha y} dy.$$

For this we define

$$M_i = \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(i)}(y) \rho_{\text{Po}}(y, k_n - 1) e^{-\alpha y} dy$$

so that

$$\int_{\mathcal{K}_C(k_n)} \mathbb{E} \left[\left(\sum_{p_1 \in \mathcal{P} \setminus \{(0,y)\}} h_y(p_1) \right) \right] \rho_{\text{Po}}(y, k_n - 1) e^{-\alpha y} dy = O(M_1 + M_2 + M_3).$$

Computing each of the integral separately we obtain, using Lemma 9.3 and the fact that $n = \nu e^{R/2}$,

$$M_1 := \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(1)}(y) \rho_{\text{Po}}(y, k_n - 1) e^{-\alpha y} dy = O(1) \cdot \begin{cases} \frac{k_n^{7-6\alpha}}{n^2}, & \text{if } \alpha < 1 \\ R \frac{k_n^{3-2\alpha}}{n^2}, & \text{if } \alpha \geq 1 \end{cases}.$$

$$M_2 := \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(2)}(y) \rho_{\text{Po}}(y, k_n - 1) e^{-\alpha y} dy = O(1) \cdot \begin{cases} \frac{k_n^{1-2\alpha}}{n^{2\alpha-1}}, & \text{if } \alpha < 3/2 \\ R \frac{k_n^{1-2\alpha}}{n^2}, & \text{if } \alpha \geq 3/2 \end{cases}$$

and finally

$$M_3 := \int_{\mathcal{K}_C(k_n)} \mathcal{I}_n^{(3)}(y) \rho_{\text{Po}}(y, k_n - 1) e^{-\alpha y} dy = O(1) \cdot \begin{cases} \frac{k_n^{2\alpha-1}}{n^{4\alpha-2}}, & \text{if } \alpha < 3/2 \\ R \cdot \frac{k_n^2}{n^{2\alpha+1}}, & \text{if } \alpha \geq 3/2 \end{cases}.$$

Now, we will consider the two cases according to the value of α . First we note that $R = O(\log(n))$ and since $k_n = O(n^{\frac{1}{2\alpha+1}})$ and $\alpha > 1/2$ we have that $R k_n^2 n^{-1} = o(1)$. Assume first that $1/2 < \alpha < 3/4$. In this case, we want to show that

$$\lim_{n \rightarrow \infty} k_n^{6\alpha-3} (M_1 + M_2 + M_3) = 0. \quad (122)$$

Using the above expression for M_i , we have

$$k_n^{6\alpha-3} (M_1 + M_2 + M_3) = O(1) \cdot k_n^{6\alpha-3} \left(\frac{k_n^{7-6\alpha}}{n^2} + \frac{k_n^{1-2\alpha}}{n^{2\alpha-1}} + \frac{k_n^{2\alpha-1}}{n^{4\alpha-3}} \right)$$

We wish to show that each one of the above three terms is $o(1)$ for $k_n = O(n^{\frac{1}{2\alpha+1}})$. For the first one we have

$$k_n^{6\alpha-3} \frac{k_n^{7-6\alpha}}{n^2} = \left(\frac{k_n^2}{n} \right)^2 = o(1).$$

The second term yields:

$$k_n^{6\alpha-3} \frac{k_n^{-2\alpha+1}}{n^{2\alpha-1}} = \left(\frac{k_n^2}{n} \right)^{2\alpha-1} = o(1).$$

Finally, the third one yields:

$$k_n^{6\alpha-3} \cdot \frac{k_n^{2\alpha-1}}{n^{4\alpha-2}} = \left(\frac{k_n^2}{n} \right)^{4\alpha-2} = o(1).$$

For $\alpha \geq 3/4$, we would like to show that

$$\lim_{n \rightarrow \infty} k_n^{2\alpha} \cdot (M_1 + M_2 + M_3) = 0. \quad (123)$$

Firstly, we note that each M_i is as above if $3/4 < \alpha < 1$. Therefore, since for this range $2\alpha < 6\alpha - 3$ the result follows from the above analysis. Next we consider the case $1 \leq \alpha < 3/2$. Here, only the value of M_1 changes and we compute that

$$k_n^{6\alpha-3} M_1 = O(1) \log(n) n^{-2} k_n^{4\alpha} \leq O(\log(n)) \left(\frac{k_n^2}{n} \right)^2 = o(1),$$

so that (123) holds for $3/4 < \alpha < 1$.

Proceeding with the case $\alpha \geq 3/2$, it is only M_2 and M_3 that change values. In particular, for any $\alpha \geq 3/2$ we have

$$\frac{k_n}{n} M_2 = O(1) R \frac{k_n}{n^2} = o(1).$$

Also,

$$k_n^{2\alpha} M_3 = O(1) R \frac{k_n^{2\alpha+2}}{n^{2\alpha+1}} = R o\left(\frac{n^{\alpha+1}}{n^{2\alpha+1}}\right) = o(1),$$

since $k_n = o(n^{1/2})$ and hence (123) holds. This finished the proof for (96).

The sum of (98) Using the Campbell-Mecke formula, we write

$$\begin{aligned} \mathbb{E} & \left[\sum_{\substack{p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y_1 < K}}^{\neq} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po}\Delta\infty}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(y)\}} \right] \\ & \leq \int_0^K \int_{-I_n}^{I_n} \int_0^R \int_{-I_n}^{I_n} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po}\Delta\infty}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(y)\}} e^{-\alpha y_2} e^{-\alpha y_1} dx_2 dy_2 dx_1 dy_1 \\ & \leq \mu(\mathcal{B}(y)) \cdot \int_{-I_n}^{I_n} \int_0^K \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po}\Delta\infty}(y)\}} e^{-\alpha y_1} dx_1 dy_1. \end{aligned}$$

Recall that by Lemma ??, $\mu(\mathcal{B}(y)) = O(1)e^{y/2}$. We bound the integral using Lemma 2.2. In particular, (9) implies that if $p_1 = (x_1, y_1) \in \mathcal{B}_{\text{Po}\Delta\infty}(y)$, then because $y_1 < K$

$$|x_1 - e^{(y+y_1)/2}| \leq e^{(y+y_1)/2} \cdot K e^{y+y_1-R} = O(1) e^{(y+y_1)/2} \cdot e^{y-R}.$$

Therefore,

$$\int_{-I_n}^{I_n} \int_0^K \mathbb{1}_{\{(x_1, y_1) \in \mathcal{B}_{\text{Po}\Delta\infty}((0, y))\}} e^{-\alpha y_1} dx_1 dy_1 = O(1) \cdot e^{y-R} \cdot \int_0^K e^{(y+y_1)/2} e^{-\alpha y_1} dy_1 = O(1) \cdot e^{3y/2-R},$$

and hence

$$\mathbb{E} \left[\sum_{\substack{p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y_1 < K}}^{\neq} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po}\Delta\infty}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(y)\}} \right] = O(1) \cdot e^{2y-R}.$$

Now, we integrate this over y to obtain that

$$\begin{aligned} & \int_{\mathcal{K}_C(k_n)} \mathbb{E} \left[\sum_{\substack{p_1, p_2 \in \mathcal{P} \setminus \{(0, y)\} \\ y_1 < K}}^{\neq} \mathbb{1}_{\{p_1 \in \mathcal{B}_{\text{Po}\Delta\infty}(y)\}} \mathbb{1}_{\{p_2 \in \mathcal{B}(y) \cap \mathcal{B}_\infty(y)\}} \right] e^{-\alpha y} dy \\ & = O(1) e^{-R} \int_{\mathcal{K}_C(k_n)} e^{2y-\alpha y} dy = O(1) n^{-2} \begin{cases} k_n^{4-2\alpha}, & \text{if } \alpha < 2 \\ \log k_n, & \text{if } \alpha = 2 \\ 1, & \text{if } \alpha > 2 \end{cases}. \end{aligned}$$

To finish the argument assume first that $1/2 < \alpha \leq 3/4$. In this case,

$$k_n^{6\alpha-3} n^{-2} k_n^{4-2\alpha} = n^{-2} \cdot k_n^{4\alpha+1} = o(1).$$

For $3/4 \leq \alpha < 2$ we use that $2\alpha < 6\alpha - 3$, so that $k_n^{2\alpha} n^{-2} k_n^{4-2\alpha} = o(1)$. Finally, when $\alpha \geq 2$, we have that

$$k_n^{2\alpha} (\log(k_n) \wedge 1) n^{-2} \leq k_n^{2\alpha+1} n^{-2} = O(n^{-1}) = o(1).$$

which completes the proof for (98) and thus the proof of Proposition 6.3. \square

9.3 Coupling G_n to G_{Po}

Now that we have established the equivalence of the clustering function between the Poissonized KPKVB graph G_{Po} and the finite box graph G_{box} the final step is to relate the clustering function in G_{Po} to the KPKVB graph G_n . As mentioned in Section 6.1, this is done by moving from $c(k_n; G_n)$ to the adjusted clustering function $c^*(k_n; G_n)$ (Lemma 6.1) and then to $c^*(k_n; G_{\text{Po}})$ (Proposition 6.2). For this we will use the coupling result (Lemma 3.12) from Section 3.6. We first give the proof of Proposition 6.2 and after that we prove Lemma 6.1. Recall that Proposition 6.2 states

$$\lim_{n \rightarrow \infty} s(k_n) \mathbb{E} [|c^*(k_n; G_n) - c^*(k_n; G_{\text{Po}})|] = 0.$$

Proof of Proposition 6.2. First we note that Proposition 6.3, 6.4 and 6.5 together imply that

$$\mathbb{E} [c^*(k_n; G_{\text{Po}})] = (1 + o(1))s(k_n)$$

Therefore it suffices to show that

$$\mathbb{E} [|c^*(k_n; G_n) - c^*(k_n; G_{\text{Po}})|] = o(\mathbb{E} [c^*(k_n; G_{\text{Po}})]).$$

For this we observe that we are looking at the modified clustering coefficient, where we divide by the expected number of degree k_n vertices. As the expected numbers of degree k_n vertices in G_{Po} and G_n are asymptotically equivalent (see Lemma 3.12), it is therefore sufficient to consider the sum of the clustering coefficients of all vertices of degree k_n . Given again the standard coupling between the binomial and Poisson process (as used in the proof of Lemma 3.12), we again denote by $V_n(k_n)$ the set of degree k_n vertices in G_n and by $V_{\text{Po}}(k_n)$ the set of degree k_n vertices in G_{Po} . If a vertex is contained in both sets, it must have the same degree in both the Poisson and KPKVB graph, and given the nature of the coupling, the neighbourhoods are therefore the same and hence also their clustering coefficients agree.

The difference of the sum of the clustering coefficients therefore comes from all the clustering coefficients of the symmetric difference $V_n(k_n) \Delta V_{\text{Po}}(k_n)$. By Lemma 3.12 the expected number vertices in this set is $\mathbb{E} [|N_n(k_n) - N_{\text{Po}}(k_n)|] = o(\mathbb{E} [N_{\text{Po}}(k_n)])$. Therefore we have that

$$\mathbb{E} [|c^*(k_n; G_n) - c^*(k_n; G_{\text{Po}})|] \leq \frac{\mathbb{E} [|N_n(k_n) - N_{\text{Po}}(k_n)|]}{(1 + o(1))\mathbb{E} [N_{\text{Po}}(k_n)]} \mathbb{E} [c^*(k_n; G_{\text{Po}})] = o(1) \mathbb{E} [c^*(k_n; G_{\text{Po}})],$$

which finishes the proof. \square

Finally we prove Lemma 6.1, whose statement is

$$\mathbb{E} [|c^*(k_n; G_n) - c(k_n; G_n)|] = o(s(k_n)).$$

Proof of Lemma 6.1. Let $0 < \delta < 1$ and define the event

$$A_n = \left\{ |N_n(k_n) - \mathbb{E} [N_n(k_n)]| \leq \mathbb{E} [N_n(k_n)]^{\frac{1+\delta}{2}} \right\}.$$

Since $N_n(k_n) = \sum_{i=1}^n \mathbb{1}_{\{\deg_n(u_i)=k_n\}}$, with u_i sampled according to (1), it follows from Lemma D.1, with $c = \mathbb{E} [N_n(k_n)]^{-\frac{1-\delta}{2}}$, that

$$\mathbb{P}(A_n) \geq 1 - O\left(e^{-\frac{\mathbb{E} [N_n(k_n)]^\delta}{2}}\right) = 1 - O\left(e^{-\frac{n^\delta k_n^{-\delta(2\alpha+1)}}{2}}\right), \quad (124)$$

where for the last part we used that $\mathbb{E} [N_n(k_n)] = \Theta\left(nk_n^{-(2\alpha+1)}\right)$ (cf. Lemma 3.12).

On the event A_n

$$\left| \frac{\mathbb{E} [N_n(k_n)]}{N_n(k_n)} - 1 \right| \leq \frac{\mathbb{E} [N_n(k_n)]^{\frac{1+\delta}{2}}}{\mathbb{E} [N_n(k_n)] + \mathbb{E} [N_n(k_n)]^{\frac{1+\delta}{2}}} \leq \mathbb{E} [N_n(k_n)]^{-\frac{1-\delta}{2}}.$$

Therefore we have

$$\begin{aligned}
\mathbb{E}[|c^*(k_n; G_n) - c(k_n; G_n)|] &\leq \mathbb{E}[|c^*(k_n; G_n) - c(k_n; G_n)| \mathbb{1}_{\{A_n\}}] + O(1 - \mathbb{P}(A_n)) \\
&= \mathbb{E}\left[c^*(k_n; G_n) \left| \frac{\mathbb{E}[N_n(k_n)]}{N_n(k_n)} - 1 \right| \mathbb{1}_{\{A_n\}}\right] + O\left(e^{-\frac{n^\delta k_n^{-\delta(2\alpha+1)}}{2}}\right) \\
&\leq \mathbb{E}[c^*(k_n; G_n)] \mathbb{E}[N_n(k_n)]^{-\frac{1-\delta}{2}} + O\left(e^{-\frac{n^\delta k_n^{-\delta(2\alpha+1)}}{2}}\right).
\end{aligned}$$

The second term is clearly $o(s(k_n))$. The first term is clearly $o(\mathbb{E}[c^*(k_n; G_n)])$ and since Propositions 6.2-6.5 imply that

$$\mathbb{E}[c^*(k_n; G_n)] = O(s(k_n)).$$

the first term is also $o(s(k_n))$. □

Tobias: Maybe a “conclusion and further work” section here, before the references and appendix?

Also, appendix usually comes after references in my experience.

Acknowledgments We are grateful to Dmitri Krioukov for pointing us to the problem of local clustering in this model and his helpful insights during discussions of the topic.

References

- [1] Nist digital library of mathematical functions. Release 1.0.22 of 2019-03-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller and B. V. Saunders, eds.
- [2] Mohammed Amin Abdullah, Nikolaos Fountoulakis, and Michel Bode. Typical distances in a geometric model for complex networks. *Internet Mathematics*, aug 2017.
- [3] Noga Alon and Joel H Spencer. *The probabilistic method*. John Wiley & Sons, 2016.
- [4] George E Andrews, Richard Askey, and Ranjan Roy. *Special functions*, volume 71. Cambridge university press, 2000.
- [5] Thomas Bläsius, Tobias Friedrich, and Anton Krohmer. Hyperbolic random graphs: Separators and treewidth. In *24th Annual European Symposium on Algorithms (ESA 2016)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2016.
- [6] Thomas Bläsius, Tobias Friedrich, and Anton Krohmer. Cliques in hyperbolic random graphs. *Algorithmica*, 80(8):2324–2344, 2018.
- [7] Thomas Bläsius, Tobias Friedrich, Anton Krohmer, and Sören Laue. Efficient embedding of scale-free graphs in the hyperbolic plane. *IEEE/ACM Transactions on Networking*, 26(2):920–933, 2018.
- [8] Michel Bode, Nikolaos Fountoulakis, and Tobias Müller. On the largest component of a hyperbolic model of complex networks. *The Electronic Journal of Combinatorics*, 22(3):3–24, 2015.
- [9] Michel Bode, Nikolaos Fountoulakis, and Tobias Müller. The probability of connectivity in a hyperbolic model of complex networks. *Random Structures & Algorithms*, 49(1):65–94, 2016.
- [10] Marián Boguná, Fragkiskos Papadopoulos, and Dmitri Krioukov. Sustaining the internet with hyperbolic mapping. *Nature communications*, 1:62, 2010.
- [11] Elisabetta Candellero and Nikolaos Fountoulakis. Clustering and the hyperbolic geometry of complex networks. *Internet Mathematics*, 12(1-2):2–53, 2016.

- [12] Hanshuang Chen, Chuansheng Shen, Gang He, Haifeng Zhang, and Zhonghuai Hou. Critical noise of majority-vote model on complex networks. *Physical Review E*, 91(2):022816, 2015.
- [13] Kimberly Claffy, Young Hyun, Ken Keys, Marina Fomenkov, and Dmitri Krioukov. Internet mapping: from art to science. In *Conference For Homeland Security, 2009. CATCH'09. Cybersecurity Applications & Technology*, pages 205–211. IEEE, 2009.
- [14] Brian Davies. *Integral transforms and their applications*, volume 41. Springer Science & Business Media, 2012.
- [15] Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, Francesco G Tricomi, et al. Higher transcendental functions, vol. 1, 1953.
- [16] Nikolaos Fountoulakis. On the evolution of random graphs on spaces of negative curvature. *arXiv*, 2012.
- [17] Nikolaos Fountoulakis and Tobias Müller. Law of large numbers for the largest component in a hyperbolic model of complex networks. *The Annals of Applied Probability*, 28(1):607–650, 2018.
- [18] David Freedman. Another note on the borel–cantelli lemma and the strong law, with the poisson approximation as a by-product. *The Annals of Probability*, pages 910–925, 1973.
- [19] Tobias Friedrich and Anton Krohmer. On the diameter of hyperbolic random graphs. *SIAM Journal on Discrete Mathematics*, 32(2):1314–1334, 2018.
- [20] Izrail Solomonovich Gradshteyn and Iosif Moiseevich Ryzhik. *Table of Integrals, Series, and Products*. Elsevier, 2015.
- [21] Luca Gugelmann, Konstantinos Panagiotou, and Ueli Peter. Random hyperbolic graphs: degree sequence and clustering. In *International Colloquium on Automata, Languages, and Programming*, pages 573–585. Springer, 2012.
- [22] Marcos Kiwi and Dieter Mitsche. A bound for the diameter of random hyperbolic graphs. In *Proceedings of the Meeting on Analytic Algorithmics and Combinatorics*, pages 26–39. Society for Industrial and Applied Mathematics, 2015.
- [23] Marcos Kiwi and Dieter Mitsche. On the second largest component of random hyperbolic graphs. *arXiv*, 2017.
- [24] Marcos Kiwi and Dieter Mitsche. Spectral gap of random hyperbolic graphs and related parameters. *The Annals of Applied Probability*, 28(2):941–989, 2018.
- [25] Dmitri Krioukov, Fragkiskos Papadopoulos, Maksim Kitsak, Amin Vahdat, and Marián Boguná. Hyperbolic geometry of complex networks. *Physical Review E*, 82(3):036106, 2010.
- [26] Günter Last and Mathew Penrose. *Lectures on the Poisson process*, volume 7. Cambridge University Press, 2017.
- [27] Yudell L Luke. *Mathematical functions and their approximations*. Academic Press, 2014.
- [28] Cornelis Simon Meijer. On the G-function. I–VIII. In *Nederl. Akad. Wetensch., Proc.*, volume 49, 1946.
- [29] M Carmen Miguel, Jack T Parley, and Romualdo Pastor-Satorras. Effects of heterogeneous social interactions on flocking dynamics. *Physical review letters*, 120(6):068303, 2018.
- [30] Tobias Müller and Merlijn Staps. The diameter of kpkvb random graphs. *arXiv*, 2017.
- [31] Takashi Owada and D Yogeshwaran. Sub-tree counts on hyperbolic random geometric graphs. *arXiv*, 2018.

- [32] Mathew Penrose. *Random geometric graphs*. Number 5. Oxford university press, 2003.
- [33] Clara Stegehuis, Remco van der Hofstad, and Johan SH van Leeuwaarden. Scale-free network clustering in hyperbolic and other random graphs. *Journal of Physics A: Mathematical and Theoretical*, 2019.
- [34] John Stillwell. *Geometry of surfaces*. Springer Science & Business Media, 2012.
- [35] Nico M Temme. *Special functions: An introduction to the classical functions of mathematical physics*. John Wiley & Sons, 2011.

A Meijer's G-function

Recall that $\Gamma(z)$ denotes the Gamma function. Let p, q, m, ℓ be four integers satisfying $0 \leq m \leq q$ and $0 \leq \ell \leq p$ and consider two sequences $\mathbf{a}_p = \{a_1, \dots, a_p\}$ and $\mathbf{b}_q = \{b_1, \dots, b_q\}$ of reals such that $a_i - b_j$ is not a positive integer for all $1 \leq i \leq p$ and $1 \leq j \leq q$ and $a_i - a_j$ is not an integer for all distinct indices $1 \leq i, j \leq p$. Then, with ι denoting the complex unit, Meijer's G-Function [28] is defined as

$$G_{p,q}^{m,\ell} \left(z \left| \begin{matrix} \mathbf{a} \\ \mathbf{b} \end{matrix} \right. \right) = \frac{1}{2\pi\iota} \int_L \frac{\prod_{j=1}^m \Gamma(b_j - t) \prod_{j=1}^{\ell} \Gamma(1 - a_j + t)}{\prod_{j=m+1}^q \Gamma(1 - b_j + t) \prod_{j=\ell+1}^p \Gamma(a_j - t)} z^t dt, \quad (125)$$

where the path L is an upward oriented loop contour which separates the poles of the function $\prod_{j=1}^m \Gamma(b_j - t)$ from those of $\prod_{j=1}^p \Gamma(1 - a_j + t)$ and begins and ends at $+\infty$ or $-\infty$.

The Meijer's G-Function is of very general nature and has relation to many known special functions such as the Gamma function and the generalized hypergeometric function. For more details, such as many identities for $G_{p,q}^{m,\ell} \left(z \left| \begin{matrix} \mathbf{a} \\ \mathbf{b} \end{matrix} \right. \right)$ see [20, 27].

For our purpose we need the following identity which follows from an Mellin transform operation.

Lemma A.1. *For any $a \in \mathbb{R}$ and $\xi, s > 0$,*

$$\Gamma^+(-a-1, \xi/s) = G_{1,2}^{2,0} \left(\frac{\xi}{s} \left| \begin{matrix} 1 \\ -a-1, 0 \end{matrix} \right. \right)$$

Proof. Let $x > 0$ and $q \in \mathbb{R}$ and note that as the Γ -function is the Mellin transform of e^{-x} , by the inverse Mellin transform formula, we have $e^{-x} = \frac{1}{2\pi\iota} \int_{c-\iota\infty}^{c+\iota\infty} \Gamma(p) x^{-p} dp$ for $c > 0$ (see [14, p.196]). Applying the change of variable $p(r) = q - r$ yields $e^{-x} = \frac{1}{2\pi\iota} \int_{c+q-\iota\infty}^{c+q+\iota\infty} \Gamma(q-r) x^{r-q} dr$, then multiplying both sides with $-x^{q-1}$ gives $-x^{q-1} e^{-x} = -\frac{1}{2\pi\iota} \int_{c+q-\iota\infty}^{c+q+\iota\infty} \Gamma(q-r) x^{r-1} dr$. Now, integrating both sides gives $\int_x^\infty t^{q-1} e^{-t} dt = \frac{1}{2\pi\iota} \int_{c+q-\iota\infty}^{c+q+\iota\infty} \frac{\Gamma(q-r)}{-r} x^r dr$. On the left-hand side is the incomplete gamma function and on the right-hand side with using $-r = \frac{\Gamma(1-r)}{\Gamma(-r)}$ is the Meijer G-function, i.e. $\Gamma^+(q, x) = G_{1,2}^{2,0} \left(x \left| \begin{matrix} 1 \\ q, 0 \end{matrix} \right. \right)$. The claim follows by plugging in $q = -a-1$ and $x = \frac{\xi}{s}$. \square

B Incomplete Beta function

Here we derive the asymptotic behavior for the function $B^-(1-z; 2\alpha, 3-4\alpha)$ as $z \rightarrow 0$, which is used to analyze the asymptotic behavior of $P(y)$, see Section 4.2.

Lemma B.1. *We have the following asymptotic results for $B^-(1-z; 2\alpha, 3-4\alpha)$*

1. For $1/2 < \alpha < 3/4$

$$\lim_{z \rightarrow 0} B^-(1-z, 2\alpha, 3-4\alpha) = B(2\alpha, 3-4\alpha).$$

2. When $\alpha = 3/4$,

$$\lim_{z \rightarrow 0} \frac{B^-(1-z, 2\alpha, 3-4\alpha)}{\log(z)} = -1.$$

3. For $\alpha > 3/4$,

$$\lim_{z \rightarrow 0} z^{4\alpha-3} B^-(1-z, 2\alpha, 3-4\alpha) = \frac{1}{4\alpha-3}.$$

Proof. We use the hypergeometric representation of the incomplete Beta function,

$$B^-(x, a, b) = \frac{x^a}{2a} F(a, 1-b, a+1, x),$$

where F denote the hypergeometric function [35] (or see [1, Section 8.17 (ii)]). In particular we have that

$$B^-(1-z; 2\alpha, 3-4\alpha) = \frac{(1-z)^{2\alpha}}{2\alpha} F(2\alpha, 4\alpha-2, 2\alpha+1, 1-z).$$

The behavior of $F(a, b, c, 1-z)$ as $z \rightarrow 0$ depend on the real part of the sum of $c-a-b$ and whether $c = a+b$ [4] (or see [1, Section 15.4(ii)]). Since in our case a, b, c will be real it only depends on the sum of $c-a-b$. For $c-a-b > 0$ we have

$$\lim_{z \rightarrow 0} F(a, b, c, 1-z) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}, \quad (126)$$

if $c = a+b$ then

$$\lim_{z \rightarrow 0} \frac{F(a, b, c, 1-z)}{\log(z)} = -\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)}, \quad (127)$$

and finally, when $c-a-b < 0$

$$\lim_{z \rightarrow 0} \frac{F(a, b, c, 1-z)}{z^{c-a-b}} = \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)}. \quad (128)$$

In our case we have,

$$B^-(1-z; 2\alpha, 3-4\alpha) = \frac{(1-z)^{2\alpha}}{2\alpha} F(a, b, c, 1-z),$$

with $a := 2\alpha$, $b := 4\alpha-2$ and $c := 2\alpha+1$. Therefore,

$$c-a-b = 2\alpha+1-2\alpha-(4\alpha-2) = 3-4\alpha.$$

Now if $\alpha < 3/4$ then $c-a-b > 0$ and hence

$$\lim_{z \rightarrow 0} B^-(1-z; 2\alpha, 3-4\alpha) = \frac{1}{2\alpha} \frac{\Gamma(2\alpha+1)\Gamma(3-4\alpha)}{\Gamma(1)\Gamma(3-2\alpha)} = \frac{\Gamma(2\alpha)\Gamma(3-4\alpha)}{\Gamma(3-2\alpha)} = B(2\alpha, 3-4\alpha),$$

where we used that $\Gamma(2\alpha+1) = 2\alpha\Gamma(2\alpha)$.

When $\alpha = 3/4$ then $c-a-b = 0$ and therefore (127), together with the fact that $(1-z)^{3/2} \sim 1$ as $z \rightarrow 0$, implies that

$$\lim_{z \rightarrow 0} \frac{B^-(1-z; 2\alpha, 3-4\alpha)}{\log(z)} = -\frac{1}{2\alpha} \frac{\Gamma(6\alpha-2)}{\Gamma(2\alpha)\Gamma(4\alpha-2)} = -\frac{\Gamma(5/2)}{\frac{3}{2}\Gamma(3/2)} = -1.$$

Finally, when $\alpha > 3/4$, $c-a-b = 3-4\alpha < 0$ and using (128) we get

$$\lim_{z \rightarrow 0} z^{4\alpha-3} B^-(1-z, 2\alpha, 3-4\alpha) = \frac{1}{2\alpha} \frac{\Gamma(2\alpha+1)\Gamma(4\alpha-3)}{\Gamma(2\alpha)\Gamma(4\alpha-2)} = \frac{\Gamma(4\alpha-3)}{\Gamma(4\alpha-2)} = \frac{1}{4\alpha-3}.$$

□

C Some results on functions

Lemma C.1. *For any $0 < \lambda < 1$ there exists a $K > 0$, such that for all $0 < x \leq (1 - \lambda)2$*

$$\frac{1}{2} \arccos(1 - x) \leq \frac{x}{\sqrt{1 - (1 - x)^2}} \leq \frac{1}{2} \arccos(1 - x) (1 + x).$$

In particular, as $x \rightarrow 0$,

$$\frac{x}{\sqrt{1 - (1 - x)^2}} \sim \frac{1}{2} \arccos(1 - x).$$

Proof. First we observe that for all $0 < x < 2$

$$0 < \sqrt{2x} \left(1 - \frac{x}{\sqrt{8}}\right) \leq \arccos(1 - x) \leq \sqrt{2x} \left(1 + \frac{x}{\sqrt{8}}\right)$$

while for every $0 < \lambda < 1$, there exists a $K > 0$ such that for all $0 < x \leq (1 - \lambda)2$,

$$0 < \frac{1}{\sqrt{2x}} \left(1 - \frac{x}{2}\right) \leq \frac{1}{\sqrt{1 - (1 - x)^2}} \leq \frac{1}{\sqrt{2x}} (1 + Kx).$$

It then follows that for all $0 < x \leq (1 - \lambda)2$,

$$\frac{x}{\sqrt{1 - (1 - x)^2}} \leq \frac{1}{2} \sqrt{2x} \left(1 + K \frac{x}{\sqrt{2}}\right) \leq \frac{1}{2} \arccos(1 - x) \frac{1 + Kx}{1 - \frac{x}{\sqrt{8}}} \leq \frac{1}{2} \arccos(1 - x) \left(1 + \frac{(K + 1)x}{1 - x}\right),$$

and

$$\frac{x}{\sqrt{1 - (1 - x)^2}} \geq \frac{1}{2} \sqrt{2x} \left(1 - \frac{x}{2}\right) \geq \frac{1}{2} \arccos(1 - x) \frac{1 - \frac{x}{2}}{1 + \frac{x}{\sqrt{8}}} \geq \frac{1}{2} \arccos(1 - x) \left(1 - \frac{(1 + \sqrt{2})x}{1 + x}\right),$$

which finishes the proof. \square

D Some results for random variables

Here we summarize several known results for random variables and provide one technical lemma for Binomial random variables.

We start with the following concentration result which follows from [18, Theorem 4], together with the note directly after it.

Lemma D.1. *Let X_n be a sum of n , possibly dependent, indicators and $c > 0$. Then*

$$\mathbb{P}(|X_n - \mathbb{E}[X_n]| > c\mathbb{E}[X_n]) \leq 2e^{-\frac{c^2\mathbb{E}[X_n]^2}{2}}.$$

Next we recall two Chernoff bound for Poisson and Binomial random variables. The following Chernoff bound can be found in [32, Lemma 1.2].

Lemma D.2. *Let $\text{Po}(\lambda)$ denote a Poisson random variable with mean λ and let $H(x) = x \log(x) - x + 1$. Then*

$$\begin{aligned} \mathbb{P}(\text{Po}(\lambda) \geq k) &\leq e^{-\lambda H(k/\lambda)} \quad \text{for all } k \geq \lambda \\ \mathbb{P}(\text{Po}(\lambda) \leq k) &\leq e^{-\lambda H(k/\lambda)} \quad \text{for all } k \leq \lambda. \end{aligned}$$

It follows from the above lemma that

$$\mathbb{P}(|\text{Po}(\lambda) - \lambda| \geq x) \leq 2e^{-\frac{x^2}{2(\lambda + x)}}. \tag{129}$$

In particular, if $\lambda_n \rightarrow \infty$, then, for any $C > 0$,

$$\mathbb{P}\left(|\text{Po}(\lambda_n) - \lambda_n| \geq C\sqrt{\lambda_n \log(\lambda_n)}\right) \leq 2e^{-\frac{C^2 \lambda_n \log(\lambda_n)}{2(\lambda_n + C\sqrt{\lambda_n \log(\lambda_n)})}} = O\left(\lambda_n^{-\frac{C^2}{2}}\right). \quad (130)$$

Let $\text{Bin}(n, p)$ denote a Binomial random variable with n trials and success probability p , and $0 < \delta < 1$. Then we have the following well-known Chernoff bound.

$$\mathbb{P}(|\text{Bin}(n, p) - np| > \delta np) \leq e^{-\frac{\delta^2 np}{3}}. \quad (131)$$

The following lemma gives an upper bound on the Binomial distribution for $p = \lambda/n$ in terms of a Poisson distribution with mean λ .

Lemma D.3. *Let $n \geq 1$, $0 < \lambda < n$. Then, for any integer $0 \leq k \leq n - 1$,*

$$\mathbb{P}(\text{Bin}(n, \lambda/n) = k) \leq \frac{e}{\sqrt{2\pi}} \sqrt{\frac{n}{n-k}} \mathbb{P}(\text{Po}(\lambda) = k).$$

Proof. Using Stirling's bounds

$$\sqrt{2\pi} \sqrt{ss^s} e^{-s} \leq s! \leq e \sqrt{ss^s} e^{-s},$$

we have

$$\begin{aligned} \mathbb{P}(\text{Bin}(n, \lambda/n) = k) &= \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \\ &\leq \frac{e}{\sqrt{2\pi}} \sqrt{\frac{n}{n-k}} \frac{n^n}{k!} (n-k)^{-(n-k)} e^{-k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \\ &= \frac{e}{\sqrt{2\pi}} \sqrt{\frac{n}{n-k}} \frac{\lambda^k e^{-\lambda}}{k!} \left(\frac{n-\lambda}{n-k}\right)^{n-k} e^{\lambda-k} \\ &= \frac{e}{\sqrt{2\pi}} \sqrt{\frac{n}{n-k}} \mathbb{P}(\text{Po}(\lambda) = k) \left(\frac{n-\lambda}{n-k}\right)^{n-k} e^{\lambda-k}. \end{aligned}$$

The result then follows by observing that $\left(\frac{n-\lambda}{n-k}\right)^{n-k} e^{\lambda-k} \leq 1$ for all $0 < \lambda < n$ and $0 \leq k \leq n - 1$. \square

The following technical lemma establishes two results that are important in Section 9.

Lemma D.4. *Let $\text{Bin}(m, p)$ denote a Binomial random variable with m trials and success probability p , let $k_n \rightarrow \infty$ be a sequence of integers such that $k_n = o(n)$ and fix some $C > 0$. Then the following holds for any $s, t > 0$:*

1. *for any $0 < p_n < 1$ such that $|np_n - k_n| > C\sqrt{k_n \log(k_n)}$,*

$$\mathbb{P}(\text{Bin}(n - t, p_n) = k_n) = O\left(k_n^{-\frac{C^2}{3}}\right).$$

2. *for any $0 < p_n < 1$ such that $|np_n - k_n| \leq C\sqrt{k_n \log(k_n)}$ and any sequence of integers m_n such that $|m_n - n| \leq C\sqrt{n \log(n)}$*

$$\mathbb{P}(\text{Bin}(n - t, p_n) = k_n) = (1 + o(1)) \mathbb{P}(\text{Bin}(m_n - s, p_n) = k_n).$$

Proof. First we observe that

$$\frac{\partial}{\partial x} \mathbb{P}(\text{Bin}(m, x) = k) = \binom{m}{k} (kx^{k-1}(1-x)^{m-k} - (m-k)x^k(1-x)^{m-k-1}).$$

Hence, the function $x \mapsto \mathbb{P}(\text{Bin}(m, x) = k)$ attains its maximum at $x = k/m$ and is strictly increasing on $(0, k/m]$ and strictly decreasing on $[k/m, 1)$.

We proceed with proving the first statement. Consider the case where $np_n < k_n - C\sqrt{k_n \log(k_n)}$. Define

$$q_n = \frac{k_n - C\sqrt{k_n \log(k_n)}}{n - t}$$

and set $Y_n = \text{Bin}(n - t, q_n)$. Then, since $q_n < k_n/(n - t)$ we get

$$\mathbb{P}(\text{Bin}(n - t, p_n) = k_n) \leq \mathbb{P}(Y_n = k_n) \leq \mathbb{P}(Y_n > k_n - 1) = \mathbb{P}(Y_n > (1 + \delta_n)(n - t)q_n),$$

with

$$\delta_n = \frac{k_n - 1 - nq_n}{(n - t)q_n} = \frac{C\sqrt{k_n \log(k_n)} - 1}{k_n - C\sqrt{k_n \log(k_n)}}.$$

By a Chernoff bound we get

$$\mathbb{P}(\text{Bin}(n - t, p_n) = k_n) \leq \mathbb{P}(Y_n > (1 + \delta_n)nq_n) \leq e^{-\frac{\delta_n^2 nq_n}{2}} \leq e^{-\frac{\delta_n^2 nq_n}{3}}.$$

The result now follows by observing that $\delta_n^2 nq_n = \Omega(C^2 \log(k_n))$.

For the case $np_n > k_n - C\sqrt{k_n \log(k_n)}$ we set $q_n = (k_n + C\sqrt{k_n \log(k_n)})/n > k_n/n$ and

$$\delta_n = \frac{nq_n - (k_n + 1)}{nq_n} = \frac{C\sqrt{k_n \log(k_n)} - 1}{k_n + C\sqrt{k_n \log(k_n)}}.$$

Then, $p_n > q_n > k_n/n$ and $\delta_n^2 nq_n = \Omega(C^2 \log(k_n))$. Thus by another Chernoff bound

$$\begin{aligned} \mathbb{P}(\text{Bin}(n - t, p_n) = k_n) &\leq \mathbb{P}(Y_n = k_n) \leq \mathbb{P}(Y_n < k_n + 1) \\ &= \mathbb{P}(Y_n < (1 - \delta_n)nq_n) \leq e^{-\frac{\delta_n^2 nq_n}{3}} = O\left(k_n^{-\frac{C^2}{3}}\right). \end{aligned}$$

Now let us prove the second statement. For this we first note that by Stirling's formula

$$\binom{n - t}{k_n} \sim (2\pi k_n)^{-1/2} \left(\frac{k_n}{n - t}\right)^{-k_n} \left(1 - \frac{k_n}{n - t}\right)^{k_n - (n - t)},$$

and similarly for $\binom{m_n - s}{k_n}$. Therefore

$$\binom{n - t}{k_n} \binom{m_n - s}{k_n}^{-1} = (1 + o(1)) \left(\frac{n - t}{m_n - s}\right)^{k_n} \left(1 - \frac{k_n}{n - t}\right)^{k_n - (n - t)} \left(1 - \frac{k_n}{m_n - s}\right)^{m_n - s - k_n}$$

and thus

$$\frac{\mathbb{P}(\text{Bin}(n - t, p_n) = k_n)}{\mathbb{P}(\text{Bin}(m_n - s, p_n) = k_n)} = (1 + o(1)) \left(\frac{n - t - k_n}{n - t - p_n}\right)^{k_n - (n - t)} \left(\frac{m_n - s - k_n}{m_n - s - p_n}\right)^{k_n - (m_n - s)}$$

Let us rewrite the first multiplicative term as

$$\left(\frac{n - t - k_n}{n - t - p_n}\right)^{k_n - (n - t)} = \left(1 - \frac{k_n - p_n}{n - t - p_n}\right)^{k_n - (n - t)} := (1 - x_n)^{k_n - (n - t)}.$$

Then, using that for all $-1/2 \leq x \leq 1/2$, $-x - x^2 \leq \log(1 - x) \leq -x$, we get

$$e^{-(k_n - (n-t))(x_n + x_n^2)} \leq \left(\frac{n - t - k_n}{n - t - p_n} \right)^{k_n - (n-t)} \leq e^{-(k_n - (n-t))x_n}.$$

Similarly,

$$e^{-(m_n - s - k_n)(y_n + y_n^2)} \leq \left(\frac{m_n - s - k_n}{m_n - s - p_n} \right)^{m_n - s - k_n} \leq e^{-(m_n - s - k_n)y_n},$$

where

$$y_n := \frac{k_n - p_n}{m_n - s - p_n}.$$

We first show that

$$\lim_{n \rightarrow \infty} e^{-(k_n - (n-t))x_n} e^{-(m_n - s - k_n)y_n} = 1. \quad (132)$$

With some algebra we get

$$\begin{aligned} & - (k_n - (n-t))x_n - (m_n - s - k_n)y_n \\ &= \left(1 - \frac{k_n - p_n}{n - t - p_n} \right) (k_n - p_n) - \left(1 - \frac{k_n - p_n}{m_n - s - p_n} \right) (k_n - p_n) \\ &= (k_n - p_n)^2 \left(\frac{1}{m_n - s - p_n} - \frac{1}{n - t - p_n} \right) \\ &= (k_n - p_n)^2 \frac{n - m_n - t + s}{(m_n - s - p_n)(n - t - p_n)}. \end{aligned}$$

Now by our assumptions

$$(k_n - p_n)^2 = \Theta(k_n \log(k_n))$$

while

$$m_n - s - p_n = \Theta(n - t - p_n) = \Theta(n)$$

and

$$-C\sqrt{n \log(n)} \leq n - m_n \leq C\sqrt{n \log(n)}.$$

Therefore we conclude that

$$(k_n - p_n)^2 \frac{n - m_n - t + s}{(m_n - s - p_n)(n - t - p_n)} = \Theta\left(\frac{k_n \log(k_n) \sqrt{n \log(n)}}{n^2} \right)$$

from which (132) follows.

In a similar way it follows that

$$\lim_{n \rightarrow \infty} e^{-(k_n - (n-t))x_n^2} e^{-(m_n - s - k_n)y_n^2} = 1,$$

which implies that

$$\lim_{n \rightarrow \infty} \frac{\mathbb{P}(\text{Bin}(n - t, p_n) = k_n)}{\mathbb{P}(\text{Bin}(m_n - s, p_n) = k_n)} = 1,$$

as thus finishes the proof. \square

E Concentration of heights for vertices with degree k

Here we will prove Proposition 2.4. We start by considering integration with respect to the function $\rho(y, k_n) = \mathbb{P}(\text{Po}(\mu(y)) = k_n)$ (the degree distribution of a typical point in G_∞). Here we show that we may restrict integration with respect to the *height* y to the interval $\mathcal{K}_C(k_n) = [y_{k_n, C}^-, y_{k_n, C}^+]$ on which $\mu(y) = \Theta(k_n)$. Next we show that if we consider any other measure $\hat{\mu}_n(y)$ that is

sufficiently equivalent to $\mu(y)$ on this interval (which will be made precise later), then we may replace $\hat{\rho}_n(y, k_n) := \mathbb{P}(\text{Po}(\hat{\mu}_n(y)) = k_n)$ in integrals with $\rho(y, k_n)$. This then implies that we can also restrict integration to the interval $\mathcal{K}_C(k_n)$. We will refer to such results as a *concentration of heights* result.

We start with a concentration of heights result for the infinite model G_∞ (Lemma E.1). We then present a generalization of this result (Lemma E.1) and use this to establish concentration of heights results for the Poissonized KPKVB G_{Po} and finite box model G_{box} .

Finally we provide a general result that allow to substitute $\hat{\rho}_n(y, k_n)$ in the integrand with $\rho(y, k_n)$ and show that this holds in particular for the degree distributions in G_{Po} and G_{box} , given by, respectively $\rho_{\text{Po}}(y, k_n) := \mathbb{P}(\text{Po}(\mu_{\text{Po}}(y)) = k_n)$ and $\rho_{\text{box}}(y, k_n) := \mathbb{P}(\text{Po}(\mu_{\text{box}}(y)) = k_n)$.

E.1 Concentration of heights argument for the infinite model

The next lemma states that for a large class of functions $h(y)$ and $k_n \rightarrow \infty$, to compute the integral

$$\int_0^\infty \rho(y, k_n) h(y) e^{-\alpha y} dy$$

it is enough to consider integration over a small interval on which $e^{y/2} \approx k_n$, instead of \mathbb{R}_+ .

Lemma E.1. *Let $\alpha > \frac{1}{2}$, $\nu > 0$, $(k_n)_{n \geq 1}$ be any positive sequence such that $k_n \rightarrow \infty$ and $k_n = o(n)$. Then the following holds.*

For any continuous function $h : \mathbb{R}_+ \rightarrow \mathbb{R}$, such that $h(y) = O(e^{\beta y})$ as $y \rightarrow \infty$ for some $\beta < \alpha$,

$$\int_{\mathbb{R}_+ \setminus \mathcal{K}_C(k_n)} \rho(y, k_n) h(y) \alpha e^{-\alpha y} dy = O(k_n^{-C^2/2}), \quad (133)$$

as $n \rightarrow \infty$.

Proof. Since $\mu'(y) = \mu(y)/2$, we get that

$$\frac{\partial \rho(y, k)}{\partial y} = \frac{1}{2} (k - \mu(y)) \rho(y, k),$$

which implies that $\rho(y, k)$ attains its maximum at $\mu(y) = k$. Moreover we see that the derivative is strictly positive when $\mu(y) < k$ and strictly negative when $\mu(y) > k$. Since $\mu(y_{k,C}^-) < k$ and $\mu(y_{k,C}^+) > k$, we conclude that $\rho(y, k)$, as a function of y , is strictly increasing on $[0, y_{k,C}^-]$ and strictly decreasing on $[y_{k,C}^+, \infty)$.

Therefore, by our assumption on $h(y)$,

$$\begin{aligned} & \int_{\mathbb{R}_+ \setminus \mathcal{K}_C(k_n)} h(y) \rho(y, k_n) \alpha e^{-\alpha y} dy \\ &= O(1) \int_0^{y_{k_n,C}^-} e^{\beta y} \rho(y, k_n) \alpha e^{-\alpha y} dy + O(1) \int_{y_{k_n,C}^+}^\infty e^{\beta y} \rho(y, k_n) \alpha e^{-\alpha y} dy \\ &= O(1) \int_0^{y_{k_n,C}^-} \rho(y, k_n) e^{-(\alpha-\beta)y} dy + O(1) \int_{y_{k_n,C}^+}^\infty \rho(y, k_n) e^{-(\alpha-\beta)y} dy \\ &\leq O(1) \rho(y_{k_n,C}^-, k_n) \int_0^{y_{k_n,C}^-} e^{-(\alpha-\beta)y} dy + O(1) \rho(y_{k_n,C}^+, k_n) \int_{y_{k_n,C}^+}^\infty e^{-(\alpha-\beta)y} dy. \end{aligned}$$

Since $\alpha - \beta > 0$, we conclude that

$$\int_{\mathbb{R}_+ \setminus \mathcal{K}_C(k_n)} h(y) \rho(y, k_n) \alpha e^{-\alpha y} dy = O(1) \left(\rho(y_{k_n,C}^-, k_n) + \rho(y_{k_n,C}^+, k_n) \right). \quad (134)$$

We shall now bound the terms $\rho(y_{k_n, C}^\pm, k_n)$. We explicitly show the bound for $\rho(y_{k_n, C}^+, k_n)$, the computation for $\rho(y_{k_n, C}^-, k_n)$ is similar. First note that $\mu(y_{k_n, C}^+) = k_n + C\sqrt{\frac{\log(k_n)}{k_n}}$. Hence we can write

$$\begin{aligned}\rho(y_{k_n, C}^+, k_n) &= \mathbb{P}\left(\text{Po}(\mu(y_{k_n, C}^+)) = k_n\right) \leq \mathbb{P}\left(\text{Po}(\mu(y_{k_n, C}^+)) \geq k_n\right) \\ &\leq \mathbb{P}\left(\left|\text{Po}(\mu(y_{k_n, C}^+)) - \mu(y_{k_n, C}^+)\right| \geq C\sqrt{\frac{\log(k_n)}{k_n}}\right).\end{aligned}$$

Apply the Chernoff bound (130) then yields

$$\rho(y_{k_n, C}^+, k_n) = O\left(k_n^{-C^2/2}\right). \quad (135)$$

A similar analysis yields

$$\rho(y_{k_n, C}^-, k_n) \leq O\left(k_n^{-C^2/2}\right). \quad (136)$$

Plugging (136) and (135) into (134) yields the result. \square

Note that we can tune the error in (133) by selecting an appropriately large $C > 0$, i.e. by restricting the function $h(y)$ inside the integral to an appropriate interval around $2\log(k_n/\xi)$. This makes Lemma E.1 very powerful. As an example we give the following corollary, which allows us to bound integrals of functions $h_n(y)$ by considering their maximum of $\mathcal{K}_C(k_n)$.

Corollary E.2. *Let $h_n : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a sequence of continuous functions which such that for some $s \in \mathbb{R}$ and $\beta < \alpha$, as $n \rightarrow \infty$, $h_n(y) = O(k_n^s e^{\beta y})$ and $h_n(y) = \Omega(1)$, uniformly on $0 \leq y \leq (1-\varepsilon)R$ for some $0 < \varepsilon < 1$. Then for large enough $C > 0$, as $n \rightarrow \infty$,*

$$\int_{\mathbb{R}_+} h_n(y) \rho(y, k_n) e^{-\alpha y} dy = (1 + o(1)) \int_{\mathcal{K}_C(k_n)} h_n(y) \rho(y, k_n) \alpha e^{-\alpha y} dy.$$

In particular,

$$\int_{\mathbb{R}_+} h_n(y) \rho(y, k_n) e^{-\alpha y} dy = O(1) k_n^{-(2\alpha+1)} \max_{y \in \mathcal{K}_C(k_n)} h_n(y),$$

as $n \rightarrow \infty$.

Proof. The second result follows immediately from the first. For the first result we note that by Lemma E.1

$$\begin{aligned}\int_{\mathbb{R}_+ \setminus \mathcal{K}_C(k_n)} h_n(y) \rho(y, k_n) e^{-\alpha y} dy &\leq O(k_n^s) \int_{\mathbb{R}_+ \setminus \mathcal{K}_C(k_n)} e^{\beta y} \rho(y, k_n) e^{-\alpha y} dy \\ &= O\left(k_n^{s-C^2/2}\right).\end{aligned}$$

By assumption on $h_n(y)$,

$$\int_{\mathcal{K}_C(k_n)} h_n(y) \rho(y, k_n) e^{-\alpha y} dy = O(k_n^{s+2\beta}) \int_{\mathcal{K}_C(k_n)} \rho(y, k_n) e^{-\alpha y} dy = O\left(k_n^{s+2\beta-(2\alpha+1)}\right),$$

and

$$\int_{\mathcal{K}_C(k_n)} h_n(y) \rho(y, k_n) e^{-\alpha y} dy = \Omega(1) \int_{\mathcal{K}_C(k_n)} \rho(y, k_n) e^{-\alpha y} dy = \Omega(k_n^{-(2\alpha+1)}).$$

Hence, by taking $C > 0$ such that $C^2/2 > \max\{2\alpha + 1 + s, 2\alpha + 1 - \beta\}$ we get that

$$\int_{\mathbb{R}_+ \setminus \mathcal{K}_C(k_n)} h_n(y) \rho(y, k_n) e^{-\alpha y} dy = o(1) \int_{\mathcal{K}_C(k_n)} h_n(y) \rho(y, k_n) e^{-\alpha y} dy.$$

\square

E.2 Concentration of heights for the KPKVB and finite box model

Although powerful, the current versions of the concentration of heights argument is only valid for the function $\rho(y, k_n) := \mathbb{P}(\text{Po}(\mu(\mathcal{B}_\infty(y))) = k_n)$. We want to extend this to the Poissonized KPKVB model G_{Po} and the finite box model G_{box} . To be more precise, recall that $\mu_{\text{Po}}(y) = \mu(\mathcal{B}(y))$ and $\mu_{\text{box}}(y) = \mu(\mathcal{B}_{\text{box}}(y))$ and let us define

$$\rho_{\text{Po}}(y, k) = \mathbb{P}(\text{Po}(\mu_{\text{Po}}(y)) = k)$$

and

$$\rho_{\text{box}}(y, k) = \mathbb{P}(\text{Po}(\mu_{\text{box}}(y)) = k).$$

Then we want when Lemma E.1 to remain true if we replace $\rho(y, k_n)$ with either the function $\rho_{\text{Po}}(y, k_n)$ or $\rho_{\text{box}}(y, k_n)$. To establish this result we first prove the following technical lemma.

Lemma E.3. *Let $0 < \delta < 1$ and $k_n \rightarrow \infty$ be such that $k_n = O(n^{1-\delta})$. Let $\hat{\mu}_n(y)$ be a monotone increasing differentiable function such for some $0 < \varepsilon < 1$, $\hat{\mu}_n(y) = (1 + o(1))\mu(y)$ holds uniformly for $0 \leq y \leq (1 - \varepsilon)R$. Furthermore, let $h : \mathbb{R}_+ \rightarrow \mathbb{R}$, be a continuous function such that $h(y) = O(e^{\beta y})$ as $y \rightarrow \infty$ for some $\beta < \alpha$. Then, for $C > 0$ large enough*

$$\int_0^\infty h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy \sim \int_{\mathcal{K}_C(k_n)} h(y) \rho(y, k_n) e^{-\alpha y} dy,$$

as $n \rightarrow \infty$.

Proof. Take any $0 < \eta < \min\{\delta, \varepsilon\}$. We first show that we can restrict to integration to the interval $[0, (1 - \eta)R]$. By construction $\eta < \varepsilon$, and hence by the assumption on $\hat{\mu}$ we have that $\hat{\mu}_n((1 - \eta)R) = \Theta(\mu((1 - \eta)R)) = \Theta(n^{1-\delta})$. Therefore, since $\eta < \delta$ and $k_n = O(n^{1-\delta})$, it follows that $\hat{\mu}_n((1 - \eta)R)/k_n = \omega(n^{\delta-\eta}) = \omega(1)$ as $n \rightarrow \infty$. Hence $\hat{\rho}_n(y, k_n) \leq \hat{\rho}((1 - \eta)R, k_n)$ for all $y \geq (1 - \eta)R$. It now follow that

$$\begin{aligned} \int_{(1-\eta)R}^R h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy &= O(1) \hat{\rho}_n((1 - \eta)R, k_n) e^{-(\alpha-\beta)(1-\eta)R} \\ &= O\left(\hat{\rho}_n((1 - \eta)R, k_n) n^{-2(\alpha-\beta)(1-\eta)}\right), \end{aligned}$$

where we used that that $h(y) = O(e^{\beta y})$. Next we use Stirling's bound $k! \geq \sqrt{2\pi} k^{k+\frac{1}{2}} e^{-k}$, to bound $\hat{\rho}_n((1 - \eta)R, k_n)$,

$$\begin{aligned} \hat{\rho}_n((1 - \eta)R, k_n) &= \mathbb{P}(\text{Po}(\hat{\mu}_n((1 - \eta)R)) = k_n) \\ &= \frac{\hat{\mu}_n((1 - \eta)R)^{k_n}}{k_n!} e^{-\hat{\mu}_n((1 - \eta)R)} \\ &= O(1) k_n^{-1/2} \left(\frac{\hat{\mu}_n((1 - \eta)R)}{k_n} \right)^{k_n} e^{k_n - \hat{\mu}_n((1 - \eta)R)} \\ &= O(1) k_n^{-1/2} e^{k_n \left(1 - \frac{\hat{\mu}_n((1 - \eta)R)}{k_n} + \log\left(\frac{\hat{\mu}_n((1 - \eta)R)}{k_n}\right)\right)} \\ &\leq O(1) k_n^{-1/2} e^{-\hat{\mu}_n((1 - \eta)R)/2}, \end{aligned}$$

where the last line follows since $\hat{\mu}_n((1 - \eta)R)/k_n \rightarrow \infty$ and $1 - x + \log(x) \leq -x/2$ for large enough x . Because $\hat{\mu}_n((1 - \eta)R) = \Theta(n^{1-\delta})$ we conclude that for any $C > 0$

$$\int_{(1-\delta)R}^R h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy = O\left(k_n^{-1/2} n^{-2(\beta-\alpha)(1-\delta)} e^{-n^{(1-\delta)/2}}\right) = O\left(k_n^{-C^2/2}\right).$$

It thus remains to prove that

$$\int_{y_{k_n, C}^+}^{(1-\eta)R} h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy \leq O\left(k_n^{-C^2/8}\right)$$

and

$$\int_0^{y_{k_n, C}^-} h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy \leq O\left(k_n^{-C^2/8}\right).$$

Define \hat{y}_n^\pm to be such that $\hat{\mu}_n(y_n^\pm) = k_n \pm C\sqrt{k_n \log(k_n)}$. Then by assumption on $\hat{\mu}_n$ we have that

$$k_n \pm C\sqrt{k_n \log(k_n)} = \hat{\mu}_n(\hat{y}_n^\pm) = (1 + o(1))\mu(\hat{y}_n^\pm) = (1 + o(1))\xi e^{\hat{y}_n^\pm/2},$$

and hence

$$\hat{y}_n^\pm = 2 \log\left(\frac{k_n \pm C\sqrt{k_n \log(k_n)}}{\xi}\right) - 2 \log(1 + o(1)) = y_{k_n, C}^\pm - 2 \log(1 + o(1)) := y_{k_n, C}^\pm - \epsilon_n,$$

with $\epsilon_n \rightarrow 0$. Recall that $\hat{\mu}_n(y)$ is monotonic increasing. Now let n be large enough such that $\hat{\mu}_n(\hat{y}_n^+ - \epsilon_n) > k_n + \frac{C}{2}\sqrt{k_n \log(n)}$. Then

$$\begin{aligned} \int_{y_{k_n, C}^+}^{(1-\eta)R} h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy &\leq \int_{\hat{y}_n^+ - \epsilon_n}^{(1-\eta)R} h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy \\ &\leq \hat{\rho}_n(\hat{y}_n^+ - \epsilon_n, k_n) \int_{\hat{y}_n^+ - \epsilon_n}^{(1-\eta)R} h(y) e^{-\alpha y} dy \\ &\leq O(1) \hat{\rho}_n(\hat{y}_n^+ - \epsilon_n, k_n), \end{aligned}$$

where we used that $\hat{\mu}_n(y)$ is monotonic increasing and $\hat{\rho}_n(y, k_n)$ is decreasing for all $y \geq \hat{y}_n^+$. Write $\lambda_n = \hat{\mu}_n(\hat{y}_n^+ - \epsilon_n)$. Then, similar to the proof of Lemma E.1, we have that

$$\hat{\rho}_n(\hat{y}_n^+ - \epsilon_n, k_n) \leq \mathbb{P}\left(|\text{Po}(\lambda_n) - \lambda_n| \geq \frac{C}{2}\sqrt{k_n \log(k_n)}\right) \leq O\left(k_n^{-C^2/8}\right),$$

where the last step follows from the Chernoff bound (130) with $C = C/2$.

In a similar fashion we can let n be large enough such that $\hat{\mu}_n(\hat{y}_n^- + \epsilon_n) < k_n - \frac{C}{2}\sqrt{k_n \log(n)}$ can show that

$$\begin{aligned} \int_0^{y_{k_n, C}^-} h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy &\leq \int_0^{\hat{y}_n^- + \epsilon_n} h(y) \hat{\rho}_n(y, k_n) e^{-\alpha y} dy \\ &\leq O(1) \hat{\rho}_n(\hat{y}_n^- + \epsilon_n, k_n) = O\left(k_n^{-C^2/8}\right). \end{aligned}$$

□

The conclusion of Lemma E.3 is that as long as $\mu_{Po}(y)$ and $\mu_{\text{box}}(y)$ are $(1 + o(1))\mu(y)$, uniformly on $[0, (1 - \varepsilon)R]$, then indeed the concentration of height result (Lemma E.1) also holds in both G_{Po} and G_{box} . This was proven in Lemma 3.1 and Lemma 3.2, respectively. For completeness we give the proof of Proposition 2.4.

Proof of Proposition 2.4. The proof for $\hat{\mu}(y) = \mu(y)$ follows directly from Lemma E.1.

Now consider the case $\hat{\mu}(y) = \mu_{Po}(y)$. Then by Lemma 3.1 $\hat{\mu}(y) = (1 + o(1))\mu(y)$, uniformly on $[0, (1 - \varepsilon)R]$ and thus in particular on $\mathcal{K}_C(k_n)$. Finally we note that by Lemma 3.3. in [21] $\hat{\mu}(y)$ is monotonic increasing. The statement then follows by applying Lemma E.3.

Finally, for $\hat{\mu}(y) = \mu_{\text{box}}(y)$ we recall that by Lemma 3.2 $\hat{\mu}(y) = (1 + o(1))\mu(y)$. More precisely,

$$\hat{\mu}(y) = \mu(y)(1 - \phi_n(y)) = \begin{cases} \mu(y) \left(1 - e^{-(\alpha - \frac{1}{2})R}\right) & \text{if } 0 \leq y \leq 2 \log(\pi/2) \\ \mu(y) (1 - \phi_n(y)) & \text{if } 2 \log(\pi/2) < y \leq (1 - \varepsilon)R, \end{cases}$$

where

$$\phi_n(y) := \left(\frac{\pi}{2}\right)^{-(2\alpha-1)} e^{-(\alpha - \frac{1}{2})(R-y)} + \frac{\nu}{\xi} e^{-(\alpha - \frac{1}{2})R - \frac{y}{2}} - \frac{\nu}{\xi} \left(\frac{\pi}{2}\right)^{-2\alpha} e^{-(\alpha - \frac{1}{2})(R-y)}.$$

Note that $\phi_n(2\log(\pi/2)) = e^{-(\alpha-\frac{1}{2})R}$. In addition, since $|\phi_n(y)| \leq O\left(e^{-(\alpha-\frac{1}{2})\varepsilon R}\right)$ for $0 \leq y \leq (1-\varepsilon)R$ we have that $\hat{\mu}(y)$ is monotonic increasing for large enough n . The statement now follows by applying Lemma E.3. \square

F Derivative of $\mu_{Po}(y)$

Recall that $\mu_{Po}(y) = \mu(\mathcal{B}(y))$ denote the measure of the ball at height y in the KPKVB model and $\mu(y) = \xi e^{\frac{y}{2}}$ denotes the measure of a ball at height h in the infinite model G_∞ . In this section we will show that $\mu'_{Po}(y) = (1 + o(1))\mu'(y)$, uniformly on $[0, (1-\varepsilon)R]$, for some $0 < \varepsilon < 1$. This is a technical result that is needed in the proof of Lemma 3.3 in Section 3.3.

First we note that it follows from Lemma 3.1 that $\mu_{Po}(y) = \mu(y)(1 + \phi_n(y))$, where $\phi_n(y) := \mu_{Po}(y)/\mu(y) - 1$. Taking the derivative we have

$$\mu'_{Po}(y) = \mu'(y)(1 + \phi_n(y)) + \mu(y)\phi'_n(y) = \mu'(y)(1 + \phi_n(y) + 2\phi'_n(y)),$$

where we used that $\frac{\partial}{\partial y}\mu(y) = \frac{1}{2}\mu(y)$. Hence, to show the result for we thus need to show that $\phi'_n(y) = o(1)$, uniformly on $[0, (1-\varepsilon)R]$.

Writing out the derivative we have

$$\phi'_n(y) = \mu_{Po}(y)^{-1} \frac{\partial}{\partial y} \mu_{Po}(y) - \frac{1}{2} \frac{\mu_{Po}(y)}{\mu(y)},$$

where we used again that $\frac{\partial}{\partial y}\mu(y) = \frac{1}{2}\mu(y)$. For the second term Lemma 3.1 implies that $\frac{1}{2} \frac{\mu_{Po}(y)}{\mu(y)} = (1 + o(1))\frac{1}{2}$ uniformly on $[0, (1-\varepsilon)R]$. The following lemma shows that the same holds for the first term from which we conclude that $\phi'_n(y) = o(1)$ and hence $\mu'_{Po}(y) = (1 + o(1))\mu'(y)$, uniformly on $[0, (1-\varepsilon)R]$.

Lemma F.1. *For any $0 < \varepsilon < 1$,*

$$\lim_{n \rightarrow \infty} \sup_{0 \leq y \leq (1-\varepsilon)R} \left| \mu(y)^{-1} \frac{\partial}{\partial y} \mu_{Po}(y) - \frac{1}{2} \right| = 0.$$

Proof. We again split $\mu_{Po}(y)$ over the top and bottom part,

$$\mu_{Po}(y) = \mu(\mathcal{B}(y) \cap \mathcal{R}([0, R-y])) + \mu(\mathcal{B}(y) \cap \mathcal{R}([R-y, R])),$$

where

$$\mu(\mathcal{B}(y) \cap \mathcal{R}([0, R-y])) = \frac{2\alpha\nu}{\pi} \int_0^{R-y} \Phi(y, y') e^{-\alpha y'} dy',$$

with $\Phi(y, y')$ defined as in (8). For the second term we have

$$\mu(\mathcal{B}(y) \cap \mathcal{R}([R-y, R])) = \int_{R-y}^R \int_{I_n} f(x', y') dx' dy' = \xi e^{y/2} \frac{2\alpha-1}{4\pi} \left(e^{-(\alpha-\frac{1}{2})(R-y)} - e^{-(\alpha-\frac{1}{2})R-y/2} \right).$$

Taking the derivative of the last expression gives

$$\begin{aligned} & \frac{\partial}{\partial y} \mu(\mathcal{B}(y) \cap \mathcal{R}([R-y, R])) \\ &= \frac{1}{2} \mu(\mathcal{B}(y) \cap \mathcal{R}([R-y, R])) + \xi e^{y/2} \frac{2\alpha-1}{4\pi} \left(\left(\alpha - \frac{1}{2} \right) e^{-(\alpha-\frac{1}{2})(R-y)} + \frac{1}{2} e^{-(\alpha-\frac{1}{2})R-y/2} \right) \\ &= \frac{1}{2} \mu(\mathcal{B}(y) \cap \mathcal{R}([R-y, R])) \left(1 + \frac{(2\alpha-1)e^{-(\alpha-\frac{1}{2})(R-y)} + e^{-(\alpha-\frac{1}{2})R-y/2}}{e^{-(\alpha-\frac{1}{2})(R-y)} - e^{-(\alpha-\frac{1}{2})R-y/2}} \right). \end{aligned}$$

Since, $\lim_{n \rightarrow \infty} \sup_{0 < y \leq (1-\varepsilon)R} \mu(\mathcal{B}_\infty(y))^{-1} \mu(\mathcal{B}(y) \cap \mathcal{R}([R-y, R])) = 0$, we are left to show that

$$\lim_{n \rightarrow \infty} \sup_{0 < y \leq (1-\varepsilon)R} \left| \mu(\mathcal{B}_\infty(y))^{-1} \frac{2\alpha\nu}{\pi} \frac{\partial}{\partial y} \int_0^{R-y} \Phi(y, y') e^{-\alpha y'} dy' - \frac{1}{2} \right| = 0. \quad (137)$$

We start with some preliminary computations. For convenience we define

$$\Xi(y, y') = 1 - \frac{\cosh(R-y) \cosh(R-y') - \cosh(R)}{\sinh(R-y) \sinh(R-y')},$$

so that

$$\Phi(y, y') = \frac{1}{2} e^{R/2} \arccos(1 - \Xi(y, y')).$$

Next, following the same calculation as in the proof of [17, Lemma 28], we write

$$\begin{aligned} \Xi(y, y') &= 2e^{-(R-y-y')} \frac{(1 - e^{y'-y-R}) (1 - e^{y-y'-R})}{(1 - e^{-2(R-y')}) (1 - e^{-2(R-y)})} \\ &:= 2e^{-(R-y-y')} \frac{h_1(y) h_2(y)}{h_3(y') h_3(y)}, \end{aligned}$$

with

$$h_1(y) = 1 - e^{y'-y-R}, \quad h_2(y) = 1 - e^{y-y'-R} \quad \text{and} \quad h_3(y) = 1 - e^{-2(R-y)}.$$

We suppressed the dependence on n and, in some cases, on y' for notation convenience.

We make two important observations. First, $\Xi(y, y')$ is an increasing function in both arguments, for $y, y' < R$ and $y + y' < R$. Second, for all $y + y' < R$, $h_1(y) \leq h_3(y')$ and $h_2(y) \leq h_3(y)$, while $h_3(y), h_3(y') < 1$, so that

$$2e^{-(R-y-y')} h_1(y) h_2(y) \leq \Xi(y, y') \leq 2e^{-(R-y-y')}. \quad (138)$$

In particular, since $R-y$ is an increasing function of n uniformly on $0 < y < (1-\varepsilon)R$, there exists a $0 < \delta < 1$ such that $1/2 \leq \Xi(y, y') < 2$ for all $y + y' < R$ and $(1-\delta)(R-y) < y' < R$ and n large enough.

Next, taking the derivative of $\Xi(y, y')$ yields,

$$\begin{aligned} \frac{\partial}{\partial y} \Xi(y, y') &= \Xi(y, y') + 2e^{-(R-y-y')} \left(\frac{h'_1(y) h_2(y)}{h_3(y') h_3(y)} + \frac{h_1(y) h'_2(y)}{h_3(y') h_3(y)} - \frac{h_1(y) h_2(y) h'_3(y)}{h_3(y') h_3(y)^2} \right) \\ &= \Xi(y, y') \left(1 + \frac{h'_1(y)}{h_1(y)} + \frac{h'_2(y)}{h_2(y)} - \frac{h'_3(y)}{h_3(y)} \right) \\ &:= \Xi(y, y') (1 + \varphi_n(y, y')), \end{aligned}$$

with

$$\varphi_n(y, y') = \frac{e^{y'-y-R}}{1 - e^{y'-y-R}} - \frac{e^{y-y'-R}}{1 - e^{y-y'-R}} - \frac{2e^{-2(R-y)}}{1 - e^{-2(R-y)}}.$$

Therefore, by the chain rule,

$$\begin{aligned} \frac{\partial}{\partial y} \Phi(y, y') &= \frac{1}{2} e^{R/2} \frac{1}{\sqrt{1 - (1 - \Xi(y, y'))^2}} \frac{\partial}{\partial y} \Xi(y, y') \\ &= \frac{\frac{1}{2} e^{R/2} \Xi(y, y')}{\sqrt{1 - (1 - \Xi(y, y'))^2}} (1 + \varphi_n(y, y')). \end{aligned} \quad (139)$$

Applying the Leibniz's rule we then get

$$\frac{\partial}{\partial y} \int_0^{R-y} \Phi(y, y') e^{-\alpha y'} dy'$$

$$\begin{aligned}
&= -\Phi(y, R-y)e^{-\alpha(R-y)} + \int_0^{R-y} \frac{\partial}{\partial y} \Phi(y, y') e^{-\alpha y'} dy' \\
&= -\frac{1}{2}e^{-(\alpha-\frac{1}{2})R+\alpha y} + \int_0^{R-y} \frac{\frac{1}{2}e^{R/2}\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} (1+\varphi_n(y, y')) e^{-\alpha y'} dy' \\
&= -\frac{1}{2}e^{-(\alpha-\frac{1}{2})R+\alpha y} + \int_0^{(1-\delta)(R-y)} \frac{\frac{1}{2}e^{R/2}\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} (1+\varphi_n(y, y')) e^{-\alpha y'} dy' \\
&\quad + \int_{(1-\delta)(R-y)}^{R-y} \frac{\frac{1}{2}e^{R/2}\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} (1+\varphi_n(y, y')) e^{-\alpha y'} dy' \\
&:= -I_1(y) + I_2(y) + I_3(y),
\end{aligned}$$

with $0 \leq \delta < 1$ such that $0 < \Xi(y, y') < 2$ for all $0 < y < R$ and $(1-\delta)(R-y) < y' < R$.

We proceed by showing that

$$\lim_{n \rightarrow \infty} \sup_{0 < y \leq (1-\varepsilon)R} \left| \frac{I_t(y)}{\mu(\mathcal{B}_\infty(y))} \right| = 0, \quad \text{for } t = 1, 3 \quad (140)$$

while

$$\lim_{n \rightarrow \infty} \sup_{0 \leq y \leq (1-\varepsilon)R} \left| \frac{2\nu\alpha}{\pi\mu(\mathcal{B}_\infty(y))} I_2(y) - \frac{1}{2} \right| = 0. \quad (141)$$

This then implies (137) and finishes the proof.

For $I_1(y)$ we have

$$\lim_{n \rightarrow \infty} \sup_{0 < y \leq (1-\varepsilon)R} \mu(\mathcal{B}_\infty(y))^{-1} I_1(y) \leq \lim_{n \rightarrow \infty} \sup_{0 < y \leq (1-\varepsilon)R} \frac{1}{2\xi} e^{-(\alpha-\frac{1}{2})(R-y)} = 0.$$

For $I_3(y)$ we first use that $y' < R-y$ to bound $\varphi(y, y')$ as follows,

$$\varphi_n(y, y') \leq \frac{e^{y'-y-R}}{1-e^{y'-y-R}} \leq \frac{e^{-2y}}{1-e^{-2y}}.$$

This then yields that

$$I_3(y) \leq \frac{1}{2} \left(1 + \frac{e^{-2y}}{1-e^{-2y}} \right) e^{R/2} \int_{(1-\delta)(R-y)}^{R-y} \frac{\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} e^{-\alpha y'} dy'.$$

To bound the integral we recall that $0 < \Xi(y, y') \leq 2e^{-(R-y-y')} < 2$ and for all $1/2 \leq x < 2$,

$$\frac{1}{\sqrt{1-(1-x)^2}} \leq \frac{2}{\sqrt{2-x}},$$

where the right hand side is a monotonic increasing function. Therefore

$$\begin{aligned}
\int_{(1-\delta)(R-y)}^{R-y} \frac{\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} e^{-\alpha y'} dy' &\leq 2 \int_{(1-\delta)(R-y)}^{R-y} \frac{\Xi(y, y')}{\sqrt{(2-\Xi(y, y'))}} e^{-\alpha y'} dy' \\
&\leq \sqrt{2} e^{-\alpha(R-y)} \int_{(1-\delta)(R-y)}^{R-y} \frac{e^{-(R-y-y')}}{\sqrt{1-e^{-(R-y-y')}}} e^{\alpha(R-y-y')} dy',
\end{aligned}$$

Making the change of variables $z = e^{-(R-y-y')}$ ($dy' = z^{-1} dz$) we get that

$$\sqrt{2} e^{-\alpha(R-y)} \int_{(1-\delta)(R-y)}^{R-y} \frac{e^{-(R-y-y')}}{\sqrt{1-e^{-(R-y-y')}}} e^{\alpha(R-y-y')} dy'$$

$$= \sqrt{2}e^{-\alpha(R-y)} \int_{e^{-\delta(R-y)}}^1 \frac{z^{-\alpha}}{\sqrt{1-z}} dz \leq \sqrt{2}e^{-\alpha(R-y)} \sqrt{1-e^{-\delta(R-y)}} \leq \sqrt{2}e^{-\alpha(R-y)}.$$

We therefore conclude that

$$I_3(y) \leq \frac{1}{\sqrt{2}} \left(1 + \frac{e^{-2y}}{1-e^{-2y}} \right) e^{-(\alpha-\frac{1}{2})R+\alpha y}.$$

which implies (140) for $t = 3$.

Finally, to show (141) we first write

$$\begin{aligned} \left| \frac{2\alpha\nu}{\pi\mu(\mathcal{B}_\infty(y))} I_2(y) - \frac{1}{2} \right| &\leq \left| \frac{2\alpha\nu}{\pi\mu(\mathcal{B}_\infty(y))} \int_0^{(1-\delta)(R-y)} \frac{\Phi(y, y')}{2} e^{-\alpha y'} dy' - \frac{1}{2} \right| \\ &\quad + \frac{2\alpha\nu}{\pi\mu(\mathcal{B}_\infty(y))} \left| \int_0^{(1-\delta)(R-y)} \frac{\Phi(y, y')}{2} e^{-\alpha y'} dy' - I_2(y) \right|. \end{aligned}$$

By Lemma ??,

$$\lim_{n \rightarrow \infty} \sup_{0 < y \leq (1-\varepsilon)R} \left| \frac{2\nu\alpha}{\pi\mu(\mathcal{B}_\infty(y))} \int_0^{(1-\delta)(R-y)} \Phi(y, y') \alpha e^{-\alpha y'} dy' - 1 \right| = 0,$$

and thus it suffices to show that the $\lim_{n \rightarrow \infty} \sup_{0 < y \leq (1-\varepsilon)R}$ of the second term goes to zero.

Recalling the definition of $I_2(y)$ we have

$$\begin{aligned} &\left| \int_0^{(1-\delta)(R-y)} \frac{\Phi(y, y')}{2} e^{-\alpha y'} dy' - I_2(y) \right| \\ &\leq \int_0^{(1-\delta)(R-y)} \left| \frac{\Phi(y, y')}{2} - \frac{\frac{1}{2}e^{R/2}\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} (1 + \varphi_n(y, y')) \right| e^{-\alpha y'} dy'. \end{aligned} \quad (142)$$

We will proceed to bound the term inside the integral. For this we first note that for $0 \leq y' \leq (1-\delta)(R-y)$,

$$\varphi_n(y, y') \leq \frac{e^{-\delta(R-y)}}{1-e^{-\delta(R-y)}}.$$

and recall that $\Xi(y, y') \leq 2e^{-(R-y-y')}$. Moreover, since $x/\sqrt{1-(1-x)^2} = x/\sqrt{2x-x^2}$ is an increasing function and $e^{-(R-y-y')} \leq e^{-\delta(R-y)}$ for $0 < y' < (1-\delta)(R-y)$,

$$\frac{\frac{1}{2}e^{R/2}\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} \leq e^{R/2} \frac{e^{-\delta(R-y)}}{\sqrt{2e^{-\delta(R-y)} - e^{-2\delta(R-y)}}}.$$

Next, recall that $\Phi(y, y') = \frac{1}{2}e^{R/2} \arccos(1 - \Xi(y, y'))$. Then, since $\Xi(y, y') < 1$ for all $y' < (1-\delta)(R-y)$, $y < R$ and n large enough, we have (see Lemma C.1),

$$\left| \frac{1}{2}\Phi(y, y') - \frac{\frac{1}{2}e^{R/2}\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} \right| \leq \frac{1}{2}\Phi(y, y')\Xi(y, y').$$

for all $y' < (1-\delta)(R-y)$ and $y < R$. Together these facts imply that for n large enough

$$\left| \frac{\Phi(y, y')}{2} - \frac{\frac{1}{2}e^{R/2}\Xi(y, y')}{\sqrt{1-(1-\Xi(y, y'))^2}} (1 + \varphi_n(y, y')) \right|$$

$$\begin{aligned}
&\leq \frac{\Phi(y, y')\Xi(y, y')}{2} + \frac{\frac{1}{2}e^{R/2}\Xi(y, y')\varphi_n(y, y')}{\sqrt{1 - (1 - \Xi(y, y'))^2}} \\
&\leq e^{-\delta(R-y)}\Phi(y, y') + \frac{e^{-\delta(R-y)}}{1 - e^{-\delta(R-y)}} \frac{e^{R/2}e^{-\delta(R-y)}}{\sqrt{2e^{-\delta(R-y)} - e^{-2\delta(R-y)}}} \\
&\leq e^{-\delta(R-y)}\Phi(y, y') + \frac{e^{\frac{R}{2}}e^{-\frac{3}{2}\delta(R-y)}}{(1 - e^{-\delta(R-y)})^{3/2}}
\end{aligned}$$

Plugging this into (142) yields

$$\begin{aligned}
&\left| \int_0^{(1-\delta)(R-y)} \frac{\Phi(y, y')}{2} e^{-\alpha y'} dy' - I_2(y) \right| \\
&\leq \int_0^{(1-\delta)(R-y)} \left(e^{-\delta(R-y)}\Phi(y, y') + \frac{e^{\frac{R}{2}}e^{-\frac{3}{2}\delta(R-y)}}{(1 - e^{-\delta(R-y)})^{3/2}} \right) e^{-\alpha y'} dy' \\
&\leq e^{-\delta(R-y)}\mu(\mathcal{B}_\infty(y)) + e^{\frac{y}{2}} \frac{e^{-(\alpha - \frac{1}{2} - (\alpha - \frac{3}{2})\delta)(R-y)}}{\alpha (1 - e^{-\delta(R-y)})^{3/2}}
\end{aligned}$$

To finish the argument we note that $R - y > 0$ for all $0 < y \leq (1 - \varepsilon)R$ and observe that $\delta < 1$ implies that $(\alpha - \frac{1}{2} - (\alpha - \frac{3}{2})\delta) > 1$. Since $\mu(\mathcal{B}_\infty(y)) = \Theta(e^{\frac{y}{2}})$ it then follows that

$$\lim_{n \rightarrow \infty} \sup_{0 < y \leq (1-\varepsilon)R} \frac{2\alpha\nu}{\pi\mu(\mathcal{B}_\infty(y))} \left| \int_0^{(1-\delta)(R-y)} \frac{\Phi(y, y')}{2} e^{-\alpha y'} dy' - I_2(y) \right| = 0,$$

which completes the proof. \square

G Code for the simulations

The simulations of the clustering coefficient and function in the KPKVB model were done using Wolfram Mathematica 11.1. The simulation dots for the clustering coefficient in Figure 2 were generated by the following code (where in the second line, the entire script was also run for the values `nu=1` and `nu=0.5`):

```

1 n=10000;
2 nu=2;
3 R=2*Log[n/nu];
4 plotpoints=20;
5 reps=100;
6 Plottingdataalpha = ConstantArray[0,{plotpoints,2}];
7 SeedRandom[1];
8 For[z=1,z<=plotpoints,z++,a=0.4+z (4.6/plotpoints); sum=0;
9   For[r=1,r<=reps,r++,V = ConstantArray[0,{n,2}];
10     For[i=1,i<=n,i++,
11       V[[i,1]]=RandomReal[{ -Pi ,Pi }];
12       V[[i,2]]=ArcCosh[RandomReal[{0,1}]](Cosh[a*R]-1)+1/a];
13     A= ConstantArray[0,{n,n}];
14     For[i=1,i<=n,i++,
15       For[j=1,j<=n,j++,
16         If[Cosh[V[[i,2]]]Cosh[V[[j,2]]] - Sinh[V[[i,2]]]Sinh[V[[j,2]]]
17           Cos[Abs[V[[i,1]]-V[[j,1]]]] <= Cosh[R] && i
18             != j,A[[i,j]]=1,A[[i,j]]=0]];
19     g = AdjacencyGraph[A];

```



```

18      sum=sum+MeanClusteringCoefficient [g] ];
19      Plotingdataalpha [[z,1]]=a;
20      Plotingdataalpha [[z,2]]=1.0*sum/ reps ;]
21 Print [Plotingdataalpha]

```

The simulation dots for the clustering function in Figure 3 were generated by the following code (where in the third line, the entire script was also run for the values `nu=1` and `nu=0.5`):

```

1  n=10000;
2  a=0.8;
3  nu=2;
4  R=2*Log [n/nu] ;
5  plotpoints=24;
6  reps=100;
7  Plotingdatak = ConstantArray[0,{reps,plotpoints,2}];
8  SeedRandom [1];
9  For [r=1,r<=reps,r++,V = ConstantArray[0,{n,2}];
10     For [i=1,i<=n,i++,
11         V[[i,1]]=RandomReal[{-Pi,Pi }];
12         V[[i,2]]=ArcCosh [RandomReal[{0,1}](Cosh [a*R]-1)+1]/a];
13  A= ConstantArray[0,{n,n}];
14  For [i=1,i<=n,i++,
15     For [j=1,j<=n,j++,
16         If [Cosh [V[[i,2]]] Cosh [V[[j,2]]] - Sinh [V[[i,2]]] Sinh [V[[j,2]]]
17             Cos [Abs[V[[i,1]]-V[[j,1]]]] <= Cosh [R] && i != j,A
18             [[i,j]]=1,A[[i,j]]=0]];
19  g = AdjacencyGraph[A];
20  For [k=1,k<=plotpoints,k++,
21     sum=0;
22     result=0;
23     nrdegk=0;
24     For [v =1,v<=n,v++,
25         If [VertexDegree [g,v]==k+1,
26             result=result+LocalClusteringCoefficient [g,v];nrdegk
27             ++]];
28     Plotingdatak [[r,k,1]]=k+1;
29     If [nrdegk>0,Plotingdatak [[r,k,2]]=1.0*result/nrdegk];]
30 Print [Mean[Plotingdatak]];

```
