

A Search for Sterile Neutrinos at the NO ν A Far Detector

A DISSERTATION PRESENTED
BY
GARETH KAFKA
TO
THE DEPARTMENT OF PHYSICS

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY
IN THE SUBJECT OF
PHYSICS

HARVARD UNIVERSITY
CAMBRIDGE, MASSACHUSETTS
MAY 2016

©2016 – GARETH KAFKA
ALL RIGHTS RESERVED.

A Search for Sterile Neutrinos at the NO ν A Far Detector

ABSTRACT

We measured things. And searched for other things. Here is what we found, please let me graduate.

Contents

ABSTRACT	iii
LIST OF FIGURES	vi
LIST OF TABLES	vii
DEDICATION	vii
ACKNOWLEDGMENTS	ix
1 A BRIEF HISTORY OF NEUTRINOS	1
1.1 Introduction	1
1.2 First Detection of Neutrinos	2
1.3 First of Evidence of Oscillations	2
1.4 Possible Evidence of Sterile Neutrinos	2
2 THEORY OF NEUTRINO OSCILLATIONS	4
2.1 Standard 3-Flavor Oscillations	4
2.2 Matter Effects	4
2.3 Sterile Neutrinos	4
2.4 Current Measurements	4
3 THE $\text{NO}\nu\text{A}$ EXPERIMENT	5
3.1 Introduction	5
3.2 The NuMI Beam	5
3.3 The $\text{NO}\nu\text{A}$ Detectors	5
3.3.1 Near Detector	5
3.3.2 Far Detector	5
4 EXPERIMENT SIMULATION	6
4.1 Introduction	6
4.2 Flux Simulation	6
4.3 Detector Simulation	6
5 EVENT RECONSTRUCTION	7
5.1 Reconstruction Chain	7
5.2 Calibration	7

6	NEUTRAL CURRENT EVENT SELECTION	8
6.1	Preselection	8
6.2	CVN Based Selection	8
6.3	Standard PID Cross Check	8
7	NEUTRAL CURRENT DISAPPEARANCE ANALYSIS	9
7.1	The Analysis Chain	9
7.2	Near Detector Decomposition	9
7.3	Extrapolation	9
7.4	Far Detector Prediction	9
8	ANALYSIS RESULTS AND SYSTEMATIC ERRORS	10
8.1	Fitting Method	10
8.2	Systematic Errors	10
8.3	Results	10
9	CONCLUSIONS AND FUTURE IMPROVEMENTS	11
9.1	Conclusions	11
9.2	Future Improvements	11
	REFERENCES	12

List of Figures

List of Tables

THIS IS THE DEDICATION.

Acknowledgments

These people were cool.

1

A Brief History of Neutrinos

1.1 INTRODUCTION

The neutrino was first postulated by Wolfgang Pauli as a possible explanation for the continuous spectrum of electrons emitted from nuclear β decay [1]. This decay was originally thought to be the emission of an electron from an atom, resulting in a different nucleus, via the process,

$$N \rightarrow N' + e \tag{1.1}$$

where N and N' are the parent and daughter nuclei, respectively. In a two body decay such as this, the momenta and energies of the outgoing particles are exactly constrained. Pauli's new particle explained the continuous spectrum of electron energy via a modified decay process:

$$N \rightarrow N' + e + \nu \tag{1.2}$$

where ν is the outgoing neutral particle. Pauli's original proposal called the new particle the neutron, but this name was later used to name the massive neutral nucleon discovered by Chadwick in 1932 [2]. Three years after Pauli's idea, Fermi proposed a model for nuclear β decay that included the new particle, which he coined the neutrino, or little neutral one [3].

1.2 FIRST DETECTION OF NEUTRINOS

Twenty years passed from Fermi's model proposal before neutrinos were discovered experimentally. Reines and Cowan made the discovery by placing a detector near a nuclear reactor as a source of neutrinos and observing inverse β decay [4, 5]. The neutrinos observed were anti-electron neutrinos, thus the following was the observed process.

$$p + \bar{\nu}_e \rightarrow n + e^+. \quad (1.3)$$

Fred Reines earned the Nobel Prize in Physics in 1995 for the detection of the neutrino.

In 1962, the muon neutrino was discovered at Brookhaven National Laboratory using the first neutrino beam [6] in a scheme still used in neutrino experiments today. The beam was generated by colliding protons with a target, producing pions that decayed into muons and muon neutrinos. The resultant beam then passed through thick steel, absorbing everything but the neutrinos. Leon Lederman, Melvin Schwartz, and Jack Steinberger won the Nobel Prize in Physics in 1988 for the discovery of the muon neutrino.

The last generation of neutrino, the tau neutrino, was discovered at Fermilab by the DONUT collaboration in 2000 [7].

1.3 FIRST OF EVIDENCE OF OSCILLATIONS

1.4 POSSIBLE EVIDENCE OF STERILE NEUTRINOS

The number of active neutrinos is constrained by measurements of the width of the Z boson. LEP has measured the number of active neutrinos to be 2.984 ± 0.008 [8], so the discoveries of the ν_e , ν_μ , and ν_τ

leave no room for new active neutrinos.

2

Theory of Neutrino Oscillations

2.1 STANDARD 3-FLAVOR OSCILLATIONS

2.2 MATTER EFFECTS

2.3 STERILE NEUTRINOS

2.4 CURRENT MEASUREMENTS

3

The NO ν A Experiment

3.1 INTRODUCTION

3.2 THE NUMI BEAM

3.3 THE NO ν A DETECTORS

3.3.1 NEAR DETECTOR

3.3.2 FAR DETECTOR

4

Experiment Simulation

4.1 INTRODUCTION

4.2 FLUX SIMULATION

4.3 DETECTOR SIMULATION

5

Event Reconstruction

5.1 RECONSTRUCTION CHAIN

5.2 CALIBRATION

6

Neutral Current Event Selection

6.1 PRESELECTION

6.2 CVN BASED SELECTION

6.3 STANDARD PID CROSS CHECK

7

Neutral Current Disappearance Analysis

7.1 THE ANALYSIS CHAIN

7.2 NEAR DETECTOR DECOMPOSITION

7.3 EXTRAPOLATION

7.4 FAR DETECTOR PREDICTION

8

Analysis Results and Systematic Errors

8.1 FITTING METHOD

8.2 SYSTEMATIC ERRORS

8.3 RESULTS

9

Conclusions and Future Improvements

9.1 CONCLUSIONS

The results of this analysis are consistent with no sterile neutrinos.

9.2 FUTURE IMPROVEMENTS

References

- [1] W. Pauli. Letter to a physicists' gathering at tubingen, 1930.
- [2] J. Chadwick. Possible existence of a neutron. *Nature*, 192:312, 1932.
- [3] E. Fermi. Versuch einer theorie der β -strahlen. *Zeitschrift für Physik*, 88:161–177, 1934.
- [4] F. Reines and C. L. Cowan. Detection of the free neutrino. *Physical Review*, 92:830–831, 1953.
- [5] C. L. Cowan, F. Reines, F. B. Harrison, H. W. Kruse, and A. D. McGuire. Detection of the free neutrino: A confirmation. *Science*, 124:103–104, 1956.
- [6] G. Danby et al. Observation of high-energy neutrino reactions and the existence of two kinds of neutrinos. *Physical Review Letters*, 9:36–44, 1962.
- [7] DONUT Collaboration, K. Kodama et al. Observation of tau neutrino interactions. *Physics Letters B*, 504:218–224, 2001.
- [8] ALEPH Collaboration, DELPHI Collaboration, L3 Collaboration, OPAL Collaboration, SID Collaboration, LEP Electroweak Working Group, SID Electroweak Group, Heavy Flavour Group. Precision electroweak measurements on the Z resonance. *Physics Reports*, 427:218–224, 2006.



THIS THESIS WAS TYPESET using \LaTeX , originally developed by Leslie Lamport and based on Donald Knuth's \TeX .

The body text is set in 11 point Egenolff-Berner Garamond, a revival of Claude Garamont's humanist typeface. The above illustration, *Science Experiment 02*, was created by Ben Schlitter and released under [CC BY-NC-ND 3.0](#). A template that can be used to format a PhD dissertation with this look & feel has been released under the permissive AGPL license, and can be found online at github.com/asm-products/Dissertate or from its lead author, Jordan Suchow, at suchow@post.harvard.edu.