A Search for Sterile Neutrinos at the $NO\nu A$ Far Detector

A DISSERTATION PRESENTED
BY
GARETH KAFKA
TO
THE DEPARTMENT OF PHYSICS

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS

FOR THE DEGREE OF

DOCTOR OF PHILOSOPHY

IN THE SUBJECT OF

PHYSICS

Harvard University Cambridge, Massachusetts May 2016 ©2016 – Gareth Kafka all rights reserved.

A Search for Sterile Neutrinos at the NO ν A Far Detector

Abstract

We measured things. And searched for other things. Here is what we found, please let me graduate.

Contents

A:	STRACT	iii
Lı	t of Figures	vi
Lı	t of Tables	vii
D	DICATION	vii
A	KNOWLEDGMENTS	ix
I	A Brief History of Neutrinos	I
	I.I Introduction	I
	1.2 First Detection of Neutrinos	2
	1.3 Evidence of Neutrino Oscillations	2
	Possible Evidence of Sterile Neutrinos	3
2	Theory of Neutrino Oscillations	4
	2.1 Standard 3-Flavor Oscillations	4
	2.2 Matter Effects	4
	2.3 Sterile Neutrinos	4
	Current Measurements	4
3	The NO $ u$ A Experiment	5
	3.1 Introduction	5
	3.2 The NuMI Beam	5
	The NO ν A Detectors	5
	3.3.1 Near Detector	5
	3.3.2 Far Detector	5
4	Experiment Simulation	6
	4.1 Introduction	6
	4.2 Flux Simulation	6
	4.3 Detector Simulation	6
5	Event Reconstruction	7
	Reconstruction Chain	7
	s 2. Calibration	7

6	Neutral Current Event Selection	8
	6.1 Preselection	8
	6.2 CVN Based Selection	8
	6.3 Standard PID Cross Check	8
7	Neutral Current Disappearance Analysis	9
	7.1 The Analysis Chain	9
	7.2 Near Detector Decomposition	9
	7.3 Extrapolation	9
	7.4 Far Detector Prediction	9
8	Analysis Results and Systematic Errors	IO
	8.1 Fitting Method	Ю
	8.2 Systematic Errors	Ю
	Results	
9	Conclusions and Future Improvements	II
	9.1 Conclusions	II
	9.2 Future Improvements	
Ri	FERENCES	12

List of Figures

List of Tables

This is the dedication.

Acknowledgments

These people were cool.

A Brief History of Neutrinos

I.I INTRODUCTION

The neutrino was first postulated by Wolfgang Pauli as a possible explanation for the continuous spectrum of electrons emitted from nuclear β decay [1]. This decay was originally thought to be the emission of an electron from an atom, resulting in a different nucleus, via the process,

$$N \to N' + e \tag{1.1}$$

where N and N' are the parent and daughter nuclei, respectively. In a two body decay such as this, the momenta and energies of the outgoing particles are exactly constrained. Pauli's new particle explained the continuous spectrum of electron energy via a modified decay process:

$$N \to N' + e + \nu \tag{1.2}$$

where ν is the outgoing neutral particle. Pauli's original proposal called the new particle the neutron, but this name was later used to name the massive neutral nucleon discovered by Chadwick in 1932 [2]. Three years after Pauli's idea, Fermi proposed a model for nuclear β decay that included the new particle, which he coined the neutrino, or little neutral one [3].

1.2 First Detection of Neutrinos

Twenty years passed from Fermi's model proposal before neutrinos were discovered experimentally. Fred Reines and Clyde Cowan made the discovery by placing a detector near a nuclear reactor as a source of neutrinos and observing inverse β decay [4, 5]. The neutrinos observed were anti-electron neutrinos, thus the following was the observed process.

$$p + \bar{\nu}_e \to n + e^+ \tag{1.3}$$

Reines earned the Nobel Prize in Physics in 1995 for the detection of the neutrino.

In 1962, the muon neutrino was discovered at Brookhaven National Laboratory using the first neutrino beam [6] in a scheme still used in neutrino experiments today. The beam was generated by colliding protons with a target, producing pions that decayed into muons and muon neutrinos. The resultant beam then passed through thick steel, absorbing everything but the neutrinos. Leon Lederman, Melvin Schwartz, and Jack Steinberger won the Nobel Prize in Physics in 1988 for the discovery of the muon neutrino.

The last generation of neutrino, the tau neutrino, was discovered at Fermilab by the DONUT collaboration in 2000 [7].

1.3 EVIDENCE OF NEUTRINO OSCILLATIONS

Pontecorvo first postulated neutrino oscillations between neutrinos and anti-neutrinos, analogous to K^0/\bar{K}^0 oscillations, in 1957 [8]. Nothing came of the proposal immediately, but the idea was later revived and modified to solve the solar neutrino problem. The physics community initially viewed neutrino oscillations with skepticism and believed the experiments to be flawed, but over time oscillations

have become an unmistakable and accepted phenomenon.

The solar neutrino problem was born from a large discrepancy between the theoretical and observed number of neutrinos produced by the sun. Neutrinos were used as a study for solar models because photons take a thousand years to escape the dense nuclear plasma to the surface of the sun, but neutrinos are unimpeded. The models, which have been confirmed today, describe a somewhat complicated chain of nuclear reactions, many of which produce neutrinos. Each individual process contributes a neutrinos in a different energy spectrum, but all of the neutrinos are created as electron neutrinos.

The experimental observations and theoretical predictions were both published in 1968. Ray Davis designed an experiment underground in the South Dakota Homestake mine consisting of a tank of an ultra pure chlorine cleaning solution capable of neutrino capture via the process

$$\nu_e + {}^{37}\text{Cl} \rightarrow {}^{37}\text{Ar} + e$$
 (1.4)

The argon atoms could be collected and counted for a direct measurement of the neutrino flux [9]. Meanwhile, John Bahcall precisely calculated the expected neutrino flux [10], and the observed rate was found to be about one third of the predicted rate. Pontecorvo revived his theory with the modification of allowing ν_e to ν_μ oscillations [11], but the idea was still not taken seriously and it was another 20 years before the solar neutrino problem was confirmed.

1.4 Possible Evidence of Sterile Neutrinos

There exists some evidence of more than three neutrinos, but the number of active neutrinos is constrained by measurements of the width of the Z boson. LEP has measured the number of active neutrinos to be 2.984 ± 0.008 [12], so the discoveries of the ν_e , ν_μ , and ν_τ leave no room for new active neutrinos. (Strictly speaking, there could be other active neutrinos if they had mass greater than half the mass of the Z boson so the Z could not decay to them, but the evidence that does exist suggests a mass splitting from the other neutrino states much smaller than this.)

Theory of Neutrino Oscillations

- 2.1 STANDARD 3-FLAVOR OSCILLATIONS
- 2.2 MATTER EFFECTS
- 2.3 STERILE NEUTRINOS
- 2.4 Current Measurements

The NOuA Experiment

- 3.1 Introduction
- 3.2 THE NUMI BEAM
- 3.3 The NO ν A Detectors
- 3.3.1 NEAR DETECTOR
- 3.3.2 FAR DETECTOR

Experiment Simulation

- 4.1 Introduction
- 4.2 FLUX SIMULATION
- 4.3 Detector Simulation

Event Reconstruction

- 5.1 RECONSTRUCTION CHAIN
- 5.2 Calibration

Neutral Current Event Selection

- 6.1 Preselection
- 6.2 CVN BASED SELECTION
- 6.3 STANDARD PID CROSS CHECK

Neutral Current Disappearance Analysis

- 7.1 THE ANALYSIS CHAIN
- 7.2 NEAR DETECTOR DECOMPOSITION
- 7.3 EXTRAPOLATION
- 7.4 FAR DETECTOR PREDICTION

Analysis Results and Systematic Errors

- 8.1 FITTING METHOD
- 8.2 Systematic Errors
- 8.3 RESULTS

Conclusions and Future Improvements

9.1 Conclusions

The results of this analysis are consistent with no sterile neutrinos.

9.2 Future Improvements

References

- [1] W. Pauli. Letter to a physicists' gathering at tubingen, 1930.
- [2] J. Chadwick. Possible existence of a neutron. *Nature*, 192:312, 1932.
- [3] E. Fermi. Versuch einer theorie der β -strahlen. Zeitschrift für Physik, 88:161–177, 1934.
- [4] F. Reines and C. L. Cowan. Detection of the free neutrino. *Physical Review*, 92:830-831, 1953.
- [5] C. L. Cowan, F. Reines, F. B. Harrison, H. W. Kruse, and A. D. McGuire. Detection of the free neutrino: A confirmation. *Science*, 124:103–104, 1956.
- [6] G. Danby et al. Observation of high-energy neutrino reactions and the existence of two kinds of neutrinos. *Physical Review Letters*, 9:36–44, 1962.
- [7] DONUT Collaboration, K. Kodama et al. Observation of tau neutrino interactions. *Physics Letters B*, 504:218–224, 2001.
- [8] B. Pontecorvo. Mesonium and anti-mesonium. Soviet Journal of Experimental and Theoretical Physics, 6:429, 1957.
- [9] R. Davis, Jr., D. S. Harmer, and K. C. Hoffman. Search for neutrinos from the sun. *Physical Review Letters*, 20:1205–1209, 1968.
- [10] J. N. Bahcall, N. A. Bahcall, and G. Shaviv. Present status of the theoretical predictions for the ³⁷cl solar-neutrino experiment. *Physical Review Letters*, 20:1209–1212, 1968.
- [11] V. Gribov and B. Pontecorvo. Neutrino astronomy and lepton charge. *Physics Letters B*, 28:493–496, 1969.
- [12] ALEPH Collaboration, DELPHI Collaboration, L3 Collaboration, OPAL Collaboration, SID Collaboration, LEP Electroweak Working Group, SID Electroweak Group, Heavy Flavour Group. Precision electroweak measurements on the Z resonance. *Physics Reports*, 427:218–224, 2006.



originally developed by Leslie Lamport and based on Donald Knuth's TEX.

The body text is set in 11 point Egenolff-Berner Garamond, a revival of Claude Garamont's humanist typeface. The above illustration, *Science Experiment 02*, was created by Ben Schlitter and released under CC BY-NC-ND 3.0. A template that can be used to format a PhD dissertation with this look & feel has been released under the permissive AGPL license, and can be found online at github.com/asm-products/Dissertate or from its lead author, Jordan Suchow, at suchow@post.harvard.edu.