PHYC 3590 - Advanced Classical Mechanics Assignment 4

Gavin Kerr B00801584 2023-02-07

Problem 4.2

(a)

$$W = \int_0^P F \cdot dr$$

$$= \int_0^Q F \cdot dr + \int_Q^P F \cdot dr$$

$$= \int_0^1 F_x dx + \int_Q^P F_y dy$$

$$= \int_0^1 x^2 dx + \int_Q^P 2xy dy$$

$$= \left[\frac{x^3}{3}\right]_0^1 + \int_Q^P 2xy dy$$

$$= \frac{1}{3} + \int_Q^P 2xy dy$$

On the path from Q to P, x is equal to 1.

$$= \frac{1}{3} + \int_0^1 2y \, dy$$

$$= \frac{1}{3} + [y^2]_0^1$$

$$W = \frac{1}{3} + 1$$

$$W = \boxed{\frac{4}{3}}$$

(b)

$$W = \int_{0}^{P} F \cdot dr$$

$$= \int_{0}^{P} F_{x} dx + \int_{0}^{P} F_{y} dy$$

$$= \int_{0}^{P} x^{2} dx + \int_{0}^{P} 2xy dy$$

$$y = x^{2}, dy = 2x dx$$

$$W = \int_{0}^{P} x^{2} dx + \int_{0}^{P} 4x^{4} dx$$

$$= \int_{0}^{1} x^{2} dx + \int_{0}^{1} 4x^{4} dx$$

$$= \left[\frac{x^{3}}{3}\right]_{0}^{1} + \left[\frac{4x^{5}}{5}\right]_{0}^{1}$$

$$= \frac{1}{3} + \frac{4}{5}$$

$$W = \boxed{\frac{17}{15}}$$

(c)

$$W = \int_{0}^{x=1} x^{2} dx + \int_{0}^{y=1} 2xy dy$$

$$W = \int_{0}^{t=1} t^{6} dx + \int_{0}^{t=1} 2t^{3}t^{2} dy$$

$$dx = 3t^{2} dt, dy = 2t dt$$

$$W = \int_{0}^{t=1} t^{6} \cdot 3t^{2} dt + \int_{0}^{t=1} 2t^{5} \cdot 2t dt$$

$$W = \int_{0}^{1} 3t^{8} dt + \int_{0}^{1} 4t^{6} dt$$

$$W = \left[\frac{2x^{9}}{9}\right]_{0}^{1} + \left[\frac{4x^{7}}{7}\right]_{0}^{1}$$

$$W = \frac{3}{9} + \frac{4}{7}$$

$$W = \frac{19}{21}$$

Problem 4.8

Problem 4.22