

Functional Analysis

Georges Khater

American University of Beirut, Math 305

September 25, 2026

Contents

1 Metric Spaces.	1
1.1 Topological Spaces	1
1.2 Compactness	8
1.3 The Arzela-Ascoli theorem	12

1 Metric Spaces.

1.1 Topological Spaces

Definition 1.1. Given a set X , a topology τ on X is a subset of $\mathcal{P}(X)$ s.t.

1. $\emptyset, X \in \tau$
2. Closed under union: $E_\alpha \in \tau \forall \alpha \in I \implies \bigcup_{\alpha \in I} E_\alpha \in \tau$
3. Closed under finite intersection: $E_1, E_2, \dots, E_n \in \tau \implies \bigcap_{i=1}^n E_i \in \tau$

We say that (X, τ) is a topological space.

Definition 1.2. Let (X, τ) a topological space and $A \subseteq X$

- A open $\iff A \in \tau$.
- A is closed $\iff X \setminus A$ is open.

Definition 1.3. (X, τ) topological space and $x \in X$. A neighborhood of x is an open set $U_x \ni x$.

Definition 1.4. (X, τ) topological space, and $A \subseteq X$. Then x is an interior point of $A \iff \exists U_x$ an open neighborhood of x s.t. $U_x \subseteq A$.

We denote by $\mathring{A}$ the set of interior points of A . Notice $\mathring{A} \subseteq A$.

Now x is a limit point of $A \iff \forall U_x$ a neighborhood of x , we have that $U_x \setminus \{x\} \cap A \neq \emptyset$.
 A' denotes the set of all limit points of A and $\bar{A} := A \cup A'$ denotes the closure of A .

Proposition 1.5. 1. $\mathring{A}$ is an open set. In fact, A is open $\iff A = \mathring{A}$.

2. $\bar{A}$ is closed. In fact, A is closed $\iff A' \subseteq A \iff \bar{A} = A$.

Consequence, A is open iff $\forall x \in A, \exists U_x \ni x$ an open neighborhood s.t. $U_x \subseteq A$.

Proof. Easy. □

Definition 1.6. Let (X, τ_x) , (Y, τ_y) topological spaces, and $f: X \rightarrow Y$ and $x_0 \in X$. Then f is continuous at $x_0 \iff \forall V$ neighborhood of $f(x_0)$ there exists a neighborhood U of x_0 s.t. $f(U) \subseteq V$.

f is said to be continuous on X iff f is continuous at every $x_0 \in X$.

Proposition 1.7. $f: X \rightarrow Y$ continuous iff $f^{-1}(G)$ is open in X for every G open in Y .

Definition 1.8. (X, τ) topological spaces, and x_n a sequence in X . We say that x_n converges to $x_0 \in X$ iff for every U_{x_0} neighborhood of x_0 , $\exists N \in \mathbb{N}$ s.t. $x_n \in U_{x_0} \forall n \geq N$.

Example 1.9. $x_n = \frac{1}{n}$.

- In $(\mathbb{R}, \tau_{\text{trivial}})$, this sequence converges to every point in $\mathbb{R}$.
- In $(\mathbb{R}, \tau_{\text{disc}})$, this sequence doesn't converge to any point (we say that it diverges).

Proposition 1.10. (X, τ) topological space. $A \subseteq X$, x_n sequence in A that converges to x_0 . Then $x_0 \in \bar{A}$.

Proposition 1.11. $f: (X, \tau_x) \rightarrow (Y, \tau_y)$ and $x_0 \in X$ with f continuous at x_0 . If x_n converges to x_0 , then $f(x_n)$ converges to $f(x_0)$.

Definition 1.12. (X, τ) topological space, $A \subseteq X$. We say that A is dense in X iff $\bar{A} = X$. We say that X is separable iff it has a countable dense subset.

Example 1.13. $(\mathbb{R}^n, |\cdot|)$ is separable. $(\mathbb{R}, \tau_{\text{disc}})$ is not separable. $(X, \tau_{\text{trivial}})$ is always separable.

Definition 1.14. (X, τ) is a Hausdorff space iff $\forall x, y \in X$ $x \neq y \exists U_x, U_y$ neighborhoods of x, y respectively s.t. $U_x \cap U_y = \emptyset$.

Proposition 1.15. The limit of a convergent sequence in a Hausdorff space is unique.

Proof. Assume x_n converges to x, y with $x \neq y$, since Hausdorff we have U_x, U_y neighborhoods with $U_x \cap U_y = \emptyset$. Now for n big enough $x_n \in U_x$ for $n \geq N$ then $x_n \notin U_y$ for these $n \geq N$, contradiction. □

Definition 1.16. X a set, then a metric on X is a function $d: X \times X \rightarrow \mathbb{R}$ s.t.

1. $d(x, y) \geq 0$, $\forall x, y \in X$ with equality iff $x = y$.
2. $d(x, y) = d(y, x) \forall x, y \in X$.
3. $d(x, z) \leq d(x, y) + d(y, z) \forall x, y, z \in X$

We say that (X, d) is a metric space.

Definition 1.17. $x \in X$ and $r \geq 0$ then

$$B(x, r) = \{y \in X : d(x, y) < r\}$$

Now define

$$\tau_d \{U \in \tau_d \iff \forall x \in U \exists r \geq 0 \text{ s.t. } B(x, r) \subseteq U\}$$

Proposition 1.18. *the $B(x, r)$. τ_d is a topology on X , called the metric topology. Then (X, τ_d) is always Hausdorff and τ_d is the smallest topology containing all*

Proposition 1.19. *Take (X, d) metric space and x_n converges to x_0 iff $\forall \varepsilon > 0, \exists N \in \mathbb{N}$ s.t. $d(x_n, x_0) < \varepsilon \forall n \geq N$.*

Proposition 1.20. *(X, d) metric space and $E \subseteq X$, then $x \in \overline{E}$ iff $\exists x_n \in E$ a convergent sequence with $x_n \rightarrow x$*

Proof ($\Leftarrow$) Any topological space.

($\Rightarrow$) Take $x \in \overline{E} = E \cup E'$, if $x \in E$ take $x_n = x$. Now if $x \in E'$, then for every $n \in \mathbb{N}$, $\exists x_n \in B^*(x, 1/n) \cap E$ then $d(x_n, x) < 1/n$, by Archimedean property we get that $x_n \rightarrow x$. □

Proposition 1.21. *(X, d) metric space with $E \subseteq X$, then E is closed iff for every convergent sequence $x_n \in E$ the limit as $n \rightarrow \infty$ of $x_n \in E$.*

Definition 1.22. *(X, d) metric space, then $E \subseteq X$ is bounded iff $E \subseteq B(x_0, R)$ for some $x_0 \in x$ and $R > 0$. In particular, a sequence x_n is bounded iff $\exists M$ s.t. $|x_n| \leq M \forall n \in \mathbb{N}$.*

Example 1.23 (Some examples of some separable or not separable metric spaces).

- $(\mathbb{R}, |\cdot|)$ separable with countable dense set $\mathbb{Q}^n$.
- The space $C([a, b])$ of real valued continuous functions with $d_\infty(f, g) = \max_{[a, b]} |f(x) - g(x)|$. By Stone-Weierstrass, for every $f \in C([a, b])$ there exists a sequence of polynomials P_n s.t. $\lim_{n \rightarrow \infty} P_n = f$ uniformly. Let's call $P(\mathbb{R})$ the space of polynomials and fix $\varepsilon > 0$ and $f \in C([a, b])$. Then $\exists P \in P(\mathbb{R})$ s.t. $d_\infty(P, f) < \varepsilon/2$ with

$$p(x) = c_k x^k + c_{k-1} x^{k-1} + \cdots + c_1 x + c_0$$

Now for each $c_i \in q_i \in \mathbb{Q}$ s.t. $|c_i - q_i| < \frac{\varepsilon}{2M}$ where $M \geq \sum_{i=0}^k |x|^i, \forall x \in [a, b]$. Now let

$$q(x) = q_k x^k + \cdots + q_1 x + q_0$$

So we get that

$$\begin{aligned} |p(x) - q(x)| &\leq \sum_{i=0}^k \sum_{q_i - c_i} |x|^i \\ &\leq \frac{\varepsilon}{2M} \sum_{i=0}^k |x|^i \end{aligned}$$

So we get that

$$d_\infty(q, f) < \varepsilon$$

Hence $C([a, b])$ is separable.

- Let $\ell^\infty(\mathbb{R})$ be the space of bounded sequences. Now define

$$x = (x_1, x_2, \cdots), \quad y = (y_1, y_2, y_3, \cdots)$$

and

$$d_\infty(x, y) = \sup_{i \in \mathbb{N}} |x_i - y_i|$$

Assume it's separable, let M be the countable dense subset. Let $A = \{0, 1\}^{\mathbb{N}}$ be the space of all sequences with entries 0 or 1, clearly A is uncountable. But if $x, y \in A$ with $x \neq y$ then $d_\infty(x, y) = 1$. For every $a \in A$, construct $B(a, 1/2)$, let m_a be an element of $B(a, 1/2) \cap M$ (uses AOC and density of m_a). Clearly if $a \neq b$, we have $B(a, 1/2) \cap B(b, 1/2) = \emptyset$, then $a \rightarrow m_a$ is injective, hence M is uncountable. Contradiction.

Definition 1.24. Let (X, d) a metric space, given a sequence x_n in X , we say that x_n is Cauchy iff $\forall \varepsilon > 0, \exists N$ s.t.

$$d(x_n, x_m) < \varepsilon, \forall n, m \geq N$$

Proposition 1.25. Every Cauchy sequence is bounded, every convergent sequence is Cauchy.

Definition 1.26. (X, d) is complete iff every Cauchy Sequence converges.

Example 1.27. $(\mathbb{R}^n, |\cdot|)$ is complete.

Some properties of metric spaces.

- i) if x_n converges, then x_n is bounded.
- ii) Let C be the space of convergent sequences, then $C \subseteq \ell^\infty$. We want to show that C is closed in ℓ^∞ . Let $x^1 = (x_{1,1}, x_{1,2}, \dots), x^2 = (x_{2,1}, x_{2,2}, \dots), x^3 = (x_{3,1}, x_{3,2}, \dots), \dots$ be a sequence in C that converges to some sequence $x = (x_1, x_2, \dots) \in \ell^\infty$. Let $\varepsilon > 0$, since x^n converges to x in ℓ^∞ , $\exists N$ s.t $d_\infty(x^n, x) < \varepsilon/3 \forall n \geq N$. In particular for $n = N$ we have

$$|x_{N,i} - x_i| < \varepsilon/3, \quad \forall i = 1, 2, 3, \dots$$

Now x^N is Cauchy, then $\exists i_0$ s.t.

$$|x_{N,i} - x_{N,j}| < \varepsilon/3, \quad \forall i, j \geq i_0$$

Hence

$$\begin{aligned} |x_i - x_j| &\leq |x_i - x_{N,i}| + |x_{N,i} - x_{N,j}| + |x_{N,j} - x_j| \\ &< \varepsilon \end{aligned}$$

Then the sequence x is Cauchy, since $\mathbb{R}$ is complete we have that $x \in C$. So C is closed.

- iii) Is $\ell^\infty(\mathbb{R})$ complete. Let $x^1 = (x_{1,1}, x_{2,1}, \dots), x^2 = (x_{1,2}, x_{2,2}, \dots), \dots$ be a Cauchy sequence in $\ell^\infty(\mathbb{R})$. We define

$$x_i := (x_i^1, x_i^2, x_i^3, \dots)$$

Clearly we have that for each $i \in \mathbb{N}$,

$$d_\infty(x^k, x^l) \geq |x_{k,i} - x_{l,i}|$$

Hence it must be the case that all the x_i 's are Cauchy, and hence converge to some x_i^* (since $\mathbb{R}$ is complete). Define

$$x^* := (x_1^*, x_2^*, \dots)$$

Now by Cauchy we know that $\exists N \in \mathbb{N}$ s.t.

$$d_\infty(x_n, x_m) < \varepsilon/2, \quad \forall n, m \geq N$$

Therefore $\forall i \in \mathbb{N}$ we have that

$$|x_i^n - x_i^m| < \varepsilon/2$$

But we know that $(x_i^m)_i \rightarrow x_i^*$, by continuity of $|\cdot|$, we have

$$|x_i^n - x_i^*| \leq \varepsilon/2, \quad \forall i \in \mathbb{N}, \forall n \geq N$$

Note that N is fixed over all i 's hence

$$d_\infty(x^n, x^*) = \sup_{i \in \mathbb{N}} |x_i^n - x_i^*| \leq \varepsilon/2 < \varepsilon$$

Hence $\ell^\infty(\mathbb{R})$ is complete.

Note that x^* is bounded because each x^n is bounded in ℓ^∞ , therefore $|x_{n,i}| \leq M$, $\forall n, i$, taking $n \rightarrow \infty$ we get that $|x_i| \leq M$, $\forall i$.

- iv) Is $C([a, b])$ complete. This is just uniformly Cauchy iff uniformly convergent and the uniform limit of continuous functions is continuous.
- v) Is $P([a, b])$ closed in $C([a, b])$. No, this is because $P([a, b])$ is dense with nonempty complement. So if its complement was open, it should contain a nonempty open set, contradicting density of the polynomials.

Proposition 1.28. Let (X, d_x) , (Y, d_y) metric spaces and $f: X \rightarrow Y$ and $x_0 \in X$ then f is continuous at x_0 iff $\forall \varepsilon > 0$, $\exists \delta > 0$ s.t.

$$d_y(f(x), f(x_0)) < \varepsilon, \quad x \text{ s.t. } d_x(x, x_0) < \delta$$

iff for every sequence $x_n \rightarrow x_0$ we have $f(x_n) \rightarrow f(x_0)$.

Proposition 1.29. (X, d) metric space, $E \subseteq X$. If E is complete, then E is closed.

Proof. Let x_n be a sequence in E that converges to some $x \in X$, but x_n is Cauchy in E so it converges to $x \in E$ (by uniqueness of the limit). $\square$

Example 1.30. Note that the converse is false, $(\mathbb{Q}, |\cdot|)$ is closed in itself but is not Cauchy.

Proposition 1.31. Every closed subset of a complete metric space is complete.

Proof. Take (X, d) complete and $E \subseteq X$ closed. Take a Cauchy sequence $x_n \in E$, since $E \subseteq X$ it converges to some $x \in X$. Now E is closed so we must have $x \in E$. $\square$

Example 1.32. $C \subseteq \ell^\infty(\mathbb{R})$ is complete.

Remark 1.33. (X, d) metric space:

$$|d(x, y) - d(x', y')| \leq d(x, x') + d(y, y')$$

This is because

$$\begin{aligned} d(x, y) &\leq d(x, x') + d(x', y') + d(y', y) \\ d(x, y) - d(x', y') &\leq d(x, x') + d(y, y') \end{aligned}$$

Similarly,

$$d(x', y) - d(x, y) \leq d(x', x) + d(y', y)$$

So we get the desired inequality.

Corollary 1.34. (X, d) metric space and x_n, y_n Cauchy sequences, then $d(x_n, y_n)$ converge in $\mathbb{R}$.

Proof. Take $d_n = d(x_n, y_n) \in \mathbb{R}$, we show that d_n is Cauchy. Now

$$\begin{aligned} |d_n - d_m| &= |d(x_n, y_n) - d(x_m, y_m)| \\ &\leq d(x_n, x_m) + d(y_n, y_m) \end{aligned}$$

Now since x_n, y_n are Cauchy then d_n is Cauchy in $\mathbb{R}$. Which by completeness gives that d_n converges. $\square$

Lemma 1.35. (X, d) metric space, and M a dense subset of X s.t. every Cauchy sequence in M converges in X . Then X is complete.

Proof. Let x_n be a Cauchy sequence in X , for every $n \in \mathbb{N}$, $\exists y_n \in M$ s.t.

$$d(x_n, y_n) < \frac{1}{n}$$

Let $\varepsilon > 0$, now

$$\begin{aligned} d(y_n, y_m) &\leq d(y_n, x_n) + d(x_n, x_m) + d(x_m, y_m) \\ &< \frac{1}{n} + d(x_n, x_m) + \frac{1}{m} \end{aligned}$$

Now $\exists N$ s.t. $n, m \geq N$ we have

$$d(y_n, y_m) < \varepsilon$$

This is because

$$1/n, 1/m < \varepsilon/3, \quad d(x_n, x_m) < \varepsilon/3$$

Now $y_n \in M$ is Cauchy, so y_n converges to some $y_0 \in X$. Now let ε

$$\begin{aligned} d(x_n, y_0) &\leq d(x_n, y_n) + d(y_n, y_0) \\ &< \frac{1}{n} + d(y_n, y_0) < \varepsilon \end{aligned}$$

for $n \geq N$ for some large enough index N . Then $x_n \rightarrow y_0$ so X is complete. $\square$

Definition 1.36. $\varphi: (X, d_X) \rightarrow (Y, d_Y)$ is an isometry iff

$$d_Y(\varphi(x), \varphi(y)) = d_X(x, y)$$

We say that A is isometric to B iff there exists a bijective isometry $\varphi: A \rightarrow B$.

Theorem 1.37 (Completeness theorem). (X, d) metric space, then there exists a complete metric space $(\hat{X}, \hat{d})$ and a dense subset M of $\hat{X}$ s.t. M is isometric to X .

In fact, the completion is "unique" (up to isometry), we call it the completion of X .

Proof. Let C_X be the space of Cauchy sequences in X . We define the relation $\sim$ on C_X as follows

$$x_n \sim y_n \iff \lim_{n \rightarrow \infty} d(x_n, y_n) = 0$$

Not that the RHS exists by the above. Clearly $\sim$ is an equivalence relation. Let $\hat{X} = C_X / \sim$ (the set of equivalence classes), i.e.

$$\hat{X} = \{[x_n] : x_n \in C_X\}$$

We define $\hat{d}([x_n], [y_n]) = \lim_{n \rightarrow \infty} d(x_n, y_n)$ (this exists by the corollary above). Note that this is well-defined, indeed let $x_n \sim x'_n$ and $y_n \sim y'_n$ so we have that

$$|d(x_n, y_n) - d(x'_n, y'_n)| \leq d(x_n, x'_n) + d(y_n, y'_n) \rightarrow 0$$

Therefore

$$\lim_{n \rightarrow \infty} d(x_n, y_n) = \lim_{n \rightarrow \infty} d(x'_n, y'_n)$$

We now show that $\hat{d}$ is a metric (clear since d is a metric):

- Clearly $\hat{d}([x_n], [y_n]) \geq 0$ and equal to zero iff $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0 \iff x_n \sim y_n \iff [x_n] = [y_n]$.
- Symmetry follows from Symmetry of d .
- Triangle inequality follows from triangle inequality on d .

Define $\varphi: X \rightarrow \hat{X}$ as follows,

$$\varphi(x) = [(x, x, x, \dots)], \quad (x, x, x, \dots) \in C_x$$

This is clearly an isometry, we denote by $M := \varphi(X) = \{[(x, x, x, \dots)] : x \in X\}$. We now show that $\overline{M} = \hat{X}$. Consider an element $[x_n] = [(x_1, x_2, \dots)] \in \hat{X}$. Let $\varepsilon > 0$, $\exists N \in \mathbb{N}$ s.t. $\forall m, n \geq N$

$$d(x_n, x_m) < \varepsilon/2$$

Consider the sequence $P_n = (x_N, x_N, \dots)$, clearly

$$d(x_n, P_n) = d(x_n, x_N) < \varepsilon/2, \quad \forall n \geq N$$

Then

$$\hat{d}([x_n], [P_n]) = \lim_{n \rightarrow \infty} d(x_n, P_n) \leq \varepsilon/2 < \varepsilon$$

With

$$[P_n] = [(x_N, x_N, \dots)] \in M$$

Therefore $\overline{M} = \hat{X}$. It remains to show that $\hat{X}$ is complete, we use the lemma. Take a Cauchy sequence in M ,

$$P_{x_1} = [(x_1, x_1, \dots)], P_{x_2} = [(x_2, x_2, \dots)], \dots, P_{x_k} = [(x_k, x_k, x_k, \dots)]$$

Construct the sequence $p = (x_1, x_2, x_3, \dots)$, note that

$$d(x_n, x_m) = \hat{d}([(x_n)], [(x_m)]) = \hat{d}(P_{x_n}, P_{x_m})$$

But P_{x_n} is Cauchy in $\hat{X}$, hence p is Cauchy in X , so $P := [p] \in \hat{X}$. Take $\varepsilon > 0$, then $\exists N \in \mathbb{N}$ s.t.

$$d(x_n, x_k) \leq \varepsilon/2, \quad \forall n, k \geq N$$

So we get that for $k \geq N$

$$\begin{aligned} \hat{d}(P, P_{x_k}) &= \lim_{n \rightarrow \infty} d(x_n, x_k) \\ &\leq \varepsilon/2 \\ &< \varepsilon \end{aligned}$$

Hence

$$\lim_{k \rightarrow \infty} P_{x_k} = P \in \hat{X}$$

□

1.2 Compactness

Definition 1.38. (X, τ) topological spaces and $K \subseteq X$. Then K is compact iff every open cover $\mathcal{G} = \{G_\alpha\}_{\alpha \in I}$ of K has a finite subcover.

Example 1.39. $[0, 1]$ is not compact in $(\mathbb{R}, |\cdot|)$, $\mathcal{G} = \{(-1, 1 - 1/n) : n \in \mathbb{N}\}$ has no finite subcover. Finite sets are clearly compact.

Proposition 1.40. (X, τ) topological space, then every closed subset of a compact set is compact.

Proof. Let K be compact, and $E \subseteq K$ is closed. Let $\mathcal{G} = \{G_\alpha\}_{\alpha \in I}$ open cover of E , so

$$E \subseteq \bigcup_{\alpha \in I} G_\alpha$$

But E^c is open with

$$K \subseteq \left(\bigcup_{\alpha \in I} G_\alpha \right) \cup E^c$$

Hence $\{G_\alpha\} \cup \{E^c\}$ is an open cover of K , so it has a finite subcover

$$K \subseteq G_{\alpha_1} \cup \dots \cup G_{\alpha_N} \cup E^c$$

Then clearly

$$E = E \cap K \subseteq G_{\alpha_1} \cup \dots \cup G_{\alpha_N}$$

so E is compact. □

Proposition 1.41. *Let (X, τ) be a Hausdorff space, then K compact $\implies K$ is closed.*

Proof. Take $q \in K^c$, for every $p \in K$ by Hausdorff $\exists U_p, V_p^q$ open neighborhoods of p, q resp. s.t. $U_p \cap V_p^q = \emptyset$. Take $\mathcal{G} = \{U_p\}_{p \in K}$ an open cover of K , so it has a finite subcover

$$K \subseteq U_{p_1} \cup \dots \cup U_{p_N}$$

Now define

$$V^q = V_{p_1}^q \cap \dots \cap V_{p_N}^q$$

open set with

$$V^q \cap K = \emptyset$$

So K is closed. □

Proposition 1.42. *(X, d) metric space, $K \subseteq X$ is compact $\implies K$ is closed and bounded.*

Proof. Ex. □

Theorem 1.43 (Heine Borel). *The converse is true for $(\mathbb{R}^n, |\cdot|)$ or more generally f.d. vector spaces.*

Example 1.44. $\ell^\infty(\mathbb{R})$ $d_\infty(x, y) = \sup_{x_i - y_i}$ and $A = \{0, 1\}^\mathbb{N}$. Note that A is bounded, since if $x_0 = (0, 0, 0, 0, \dots)$, notice that $d_\infty(x, x_0) \leq 1$, so A is bounded. Moreover, every convergent sequence in A is eventually a constant and so belongs to A so A is closed. We could also say that A^c is open, take $x \notin A$, then $\exists x_i \neq 0, x_i \neq 1$. So $\exists \varepsilon > 0$ s.t. for all $y \in (x_i - \varepsilon, x_i + \varepsilon)$ $y_0 \neq 0, y \neq 1$. take $z \in B(x, \varepsilon)$ so

$$|z_i - x_i| \leq d_\infty(z, x) < \varepsilon$$

i.e. $z_i \neq 0$ and $z_i \neq 1$ so $z \in A^c$.

Now A is not compact. Take $\mathcal{G} = \{B(x, 1/2)\}_{x \in A}$, now $B(x, 1/2) \cap A = \{x\}$ but A is infinite (uncountable even), then $\mathcal{G}$ does not have a finite subcover.

Definition 1.45. (X, d) metric space and $K \subseteq X$, then K is totally bounded iff $\forall \varepsilon > 0, \exists x_1, x_2, \dots, x_n \in K$ s.t. $K \subseteq B(x_1, \varepsilon) \cup B(x_2, \varepsilon) \cup \dots \cup B(x_n, \varepsilon)$.

Example 1.46. $\{0, 1\}^\mathbb{N}$ is not totally bounded in $\ell^\infty(\mathbb{R})$ (although it is bounded).

Proposition 1.47. *K is totally bounded $\implies K$ is bounded.*

Proof. easy. □

Proposition 1.48. *A subset of a totally bounded set is totally bounded.*

Proof. Let K be totally bounded and $E \subseteq K$, fix $\varepsilon > 0$ then we have $x_1, x_2, \dots, x_n \in K$ s.t.

$$K \subseteq B(x_1, \varepsilon/2) \cup \dots \cup B(x_n, \varepsilon/2)$$

Note that $E \subseteq K$, so pick $y_1, \dots, y_k \in B(x_i, \varepsilon/2)$ (in case of empty just skip). Then we have that

$$E \subseteq B(x_1, \varepsilon) \cup \dots \cup B(x_k, \varepsilon)$$

□

Lemma 1.49. *Assume that x_n is a Cauchy sequence in (X, d) with a subsequence that converges to x_0 , then $x_n \rightarrow x_0$.*

Proof. easy. □

Remark 1.50. In $\mathbb{R}^n$ (finite dimensional vector spaces), K is totally bounded iff K is bounded.

Proof. Let K be bounded, then $\exists Q$ a closed cube in $\mathbb{R}^n$ s.t. $K \subseteq Q$. Now Q is compact (closed and bounded), so given $\varepsilon > 0$ it can be covered by finitely many balls of radius ε and center in Q . So the cube is totally bounded, and hence K is a subset of a totally bounded set, hence is totally bounded. □

Theorem 1.51. *(X, d) metric space and $K \subseteq Q$, then the following are equivalent:*

- i) K is compact
- ii) Every infinite subset E of K has a limit point in K .
- iii) Every sequence in K has a convergent subsequence (with its limit in K).
- iv) K is complete and totally bounded.

(Proof $\Rightarrow$ (ii)) Let $E \subset K$ be infinite s.t. E has no limit points in K . Then for each $x \in K$, $x \notin E'$ so $\exists \varepsilon_x$ s.t.

$$B^*(x, \varepsilon_x) \cap E = \emptyset$$

Now take $\mathcal{G} = \{B(x, \varepsilon_x)\}_{x \in K}$, this is an open cover of K so it has a finite subcover. We get

$$K \subseteq B(x_1, \varepsilon_{x_1}) \cup \dots \cup B(x_N, \varepsilon_{x_N})$$

But $E \subseteq K$ so

$$E = E \cap K \subseteq \bigcup_{i=1}^N E \cap B(x_i, \varepsilon_{x_i}) \subseteq \{x_1, \dots, x_N\}$$

But E is infinite, contradiction.

(ii) $\Rightarrow$ (iii) Let x_n be a sequence in K and define

$$E = \{x_n : n \in \mathbb{N}\}$$

If E is finite, by pigeonhole principle $\exists n_1 < n_2 < \dots$ and $x_0 \in E$ s.t. $x_{n_k} = x_0 \in E$ so we have found a convergent subsequence. If not then by the hypothesis E has a limit point $x_0 \in K$. We know that

$$B(x_0, 1) \cap E \neq \emptyset$$

So let n_1 be such that $x_{n_1} \in B(x_0, 1) \cap E$. In fact if $x \in E'$ we have that $B(x, \varepsilon) \cap E$ is infinite (since we are in a metric space). Hence for each k ,

$$B^*(x_0, 1/k) \cap E \text{ is infinite}$$

So we can find some $n_k > n_{k-1}$ with $d(x_{n_k}, x_0) < 1/k$, we have that

$$\lim_{k \rightarrow \infty} x_{n_k} = x_0$$

(iii) $\implies$ (iv) Let x_n be a Cauchy sequence in K . But K is sequentially compact, then x_n has a convergent subsequence $x_{n_k} \rightarrow x_0 \in K$. Now by the lemma, x_n converges to x_0 and so K is complete. Assume K is not totally bounded, then $\exists \varepsilon_0 > 0$ s.t. K cannot be covered by finitely many ε_0 -balls with center in K . take $x_1 \in K$, then take $x_2 \in K \setminus B(x_1, \varepsilon_0)$, then $x_3 \in K \setminus (B(x_1, \varepsilon_0) \cup B(x_2, \varepsilon_0))$. Similarly take

$$x_k \in K \setminus (B(x_1, \varepsilon_0) \cup \dots \cup B(x_{k-1}, \varepsilon_0))$$

Consider the sequence $\{x_n\}$, then for $n \neq m$ we have

$$d(x_n, x_m) \geq \varepsilon_0$$

So x_n has no convergent subsequence, contradiction.

(iv) $\implies$ (i) Assume K is not compact, then $\exists \mathcal{G} = \{G_\alpha\}_{\alpha \in I}$ an open cover of K that does not have a finite subcover. Take $\varepsilon = 1$, so K can be covered by finitely many 1-balls. So $\exists x_1 \in K$ and $B(x_1, 1)$ s.t. $K_1 := B(x_1, 1) \cap K$ cannot be covered by a finite subcover of $\mathcal{G}$. Now take $\varepsilon = 1/2$, since $K_1 \subseteq K$ is totally bounded we can similarly find $x_2 \in K_1$ with $K_2 := B(x_2, 1/2) \cap K_1$ cannot be covered by finitely many G_α 's. Recursively, we construct

$$K_0 \subseteq K_1 \subseteq K_2 \subseteq \dots$$

with $K_n := B(x_n, 1/n) \cap K_{n-1}$ with $x_n \in K_{n-1}$ and K_n cannot be covered by finitely many elements in $\mathcal{G}$. We show that x_n is Cauchy. Fix $\varepsilon > 0$, take N s.t. $2/N < \varepsilon$, then for $n, m > N$ then

$$x_n \in K_{n-1} \subseteq K_N, \quad x_m \in K_{m-1} \subseteq K_N \implies x_n, x_m \in K_N = B(x_n, 1/N) \cap K_{N-1}$$

So

$$\begin{aligned} d(x_n, x_m) &\leq d(x_n, x_N) + d(x_N, x_m) \\ &< 2/N \\ &< \varepsilon \end{aligned}$$

So x_n is Cauchy.

Therefore by completeness, x_n converges to some $x_0 \in K$. But we must have that $\exists \alpha_0 \in I$ s.t. $x_0 \in G_{\alpha_0}$ which is open, so $\exists \delta_0 > 0$ s.t. $B(x_0, \delta_0) \subseteq G_{\alpha_0}$. Since $x_n \rightarrow x_0$, then $\exists N$ s.t. $1/N < \delta_0/2$ and $d(x_n, x_0) < \delta_0/2$. So take $y \in B(x_n, 1/N)$ so

$$\begin{aligned} d(y, x_0) &\leq d(y, x_n) + d(x_n, x_0) \\ &< 1/N + d(x_n, x_0) \\ &< \delta_0 \end{aligned}$$

So

$$B(x_n, 1/N) \subseteq B(x_0, \delta_0)$$

We get that $K_N = B(x_N, 1/N) \cap K_{N-1} \subseteq B(x_0, \delta_0) \subseteq G_{\alpha_0}$. □

Proposition 1.52. $f: (X, \tau_X) \rightarrow (Y, \tau_Y)$ and $K \subseteq X$ compact, then $f(K)$ is compact.

Proof. easy. □

1.3 The Arzela-Ascoli theorem

Definition 1.53. Let X be a compact metric space, we denote by $C(X)$ the space of real-valued continuous functions on X . We define

$$d_\infty(f, g) = \sup_{x \in X} |f(x) - g(x)|$$

Proposition 1.54. $C(X)$ is complete and separable.

Proof. Uniformly Cauchy in $\mathbb{R} \implies$ uniformly continuous, follows from the general Stone-Weierstrass theorem. □

Remark 1.55. Since X is compact, then every continuous function is uniformly continuous. I.e. $\forall \varepsilon > 0, \exists \delta > 0$ s.t.

$$d_Y(f(x), f(x')) < \varepsilon, \quad \forall d(x, x') < \delta$$

Definition 1.56. $\mathcal{F} \subseteq C(X)$ is said to be equicontinuous iff $\forall \varepsilon > 0, \exists \delta > 0$ s.t.

$$d(f(x), f(x')) < \varepsilon, \quad \forall x, x' \text{ } d(x, x') < \delta, \forall f \in \mathcal{F}$$

Theorem 1.57 (Arzela-Ascoli). $\mathcal{F} \subseteq C(X)$ is totally bounded $\iff \mathcal{F}$ is bounded and equicontinuous.

Proof. $\implies \mathcal{F}$ is bounded since it is totally bounded. Let $\varepsilon > 0$, since $\mathcal{F}$ is totally bounded, there exist $f_1, f_2, \dots, f_N \in \mathcal{F}$ s.t.

$$\mathcal{F} \subseteq B(f_1, \varepsilon/3) \cup \dots \cup B(f_N, \varepsilon/3)$$

Now $f_i \in C(X)$, so uniformly continuous, and so $\exists \delta_i > 0$ s.t.

$$|f_i(x) - f_i(y)| < \varepsilon/3, \quad \forall d(x, y) < \delta_i$$

Let $\delta = \min_i \delta_i$. Now take $f \in \mathcal{F}$, and $x, y \in X$ with $d(x, y) < \delta$, now $\exists i_0 \in \{1, \dots, N\}$ s.t. $f \in B(f_{i_0}, \varepsilon/3)$. Hence

$$\begin{aligned} |f(x) - f(y)| &\leq |f(x) - f_{i_0}(x)| + |f_{i_0}(x) - f_{i_0}(y)| + |f_{i_0}(y) - f(y)| \\ &\leq 2 \times d_\infty(f, f_{i_0}) + \varepsilon/3 \\ &< \varepsilon \end{aligned}$$

Therefore $\mathcal{F}$ is equicontinuous.

$\Leftarrow$ Let $\varepsilon > 0$, then $\exists \delta > 0$ s.t. $\forall f \in \mathcal{F}$ we have

$$|f(x) - f(y)| < \varepsilon/3, \quad \forall x, y \in X \text{ s.t. } d(x, y) < \delta$$

Now X is compact, then we can cover X with finitely many δ -balls, i.e.

$$X = B(x_1, \delta) \cup \dots \cup B(x_N, \delta)$$

Define a map $\varphi: \mathcal{F} \rightarrow \mathbb{R}^N$,

$$\varphi(f) = (f(x_1), \dots, f(x_N))$$

We want to show that $\varphi(\mathcal{F})$ is bounded in $\mathbb{R}^N$. We know that $\mathcal{F}$ is bounded in $C(X)$, so $\exists M > 0$ s.t.

$$|f(x)| \leq M, \quad \forall f \in \mathcal{F}, \forall x \in X$$

Therefore

$$|\varphi(f)| = \sqrt{f(x_1)^2 + \dots + f(x_N)^2} \leq \sqrt{N}M$$

Therefore $\varphi(\mathcal{F})$ is bounded in $\mathbb{R}^N$. Therefore $\varphi(\mathcal{F})$ is totally bounded. Then there exists $f_1, f_2, \dots, f_K \in \mathcal{F}$ s.t.

$$\varphi(\mathcal{F}) \subseteq B(\varphi(f_1), \varepsilon/3) \cup \dots \cup B(\varphi(f_K), \varepsilon/3)$$

We want to show that

$$\mathcal{F} \subseteq B(f_1, \varepsilon) \cup \dots \cup B(f_K, \varepsilon)$$

Let $f \in \mathcal{F}$, then $\varphi(f) \in \varphi(\mathcal{F})$ i.e. $\exists i_0$ s.t. $\varphi(f) \in B(\varphi(f_{i_0}), \varepsilon/3)$. Recall that

$$\varphi(f) = (f(x_1), \dots, f(x_N))$$

But notice that for each j , we have

$$|f(x_j) - f_{i_0}(x_j)| < \varepsilon/3, \quad \forall j = 1, \dots, N$$

Let $x \in X$, then $x \in B(x_{j_0}, \delta)$ for some

$$\begin{aligned} |f(x) - f_{i_0}(x)| &\leq |f(x) - f(x_{j_0})| + |f(x_{j_0}) - f_{i_0}(x_{j_0})| + |f_{i_0}(x_{j_0}) - f_{i_0}(x)| \\ &< \varepsilon \end{aligned}$$

Since this holds for each $x \in X$ we have

$$|f(x) - f_{i_0}(x)| < \varepsilon, \quad \forall x \in X$$

Then since X is compact

$$\sup |f(x) - f_{i_0}(x)| = \max |f(x) - f_{i_0}(x)| < \varepsilon$$

Hence $f \in B(f_{i_0}, \varepsilon)$, therefore $\mathcal{F}$ is totally bounded.

□

Corollary 1.58. $\mathcal{F} \subseteq C(X)$ is compact iff $\mathcal{F}$ is closed and bounded and equicontinuous.

Lemma 1.59. $\mathcal{F} \subseteq C(X)$ is equicontinuous, then $\overline{\mathcal{F}}$ is equicontinuous.

Proof. Take $\varepsilon > 0$ and $f \in \overline{\mathcal{F}}$, then $\exists f_n \in \mathcal{F}$ s.t.

$$\sup_X |f_n - f|$$

but $\mathcal{F}$ is equicontinuous, so $\exists \delta > 0$ s.t. $\forall h \in \mathcal{F}$

$$|h(x) - h(x')| < \varepsilon/3, \quad \forall x, x' \text{ s.t. } d(x, x') < \delta$$

for $d(x, x') < \delta$

$$\begin{aligned} |f(x) - f(x')| &\leq |f(x) - f_n(x)| + |f_n(x) - f_n(x')| + |f_n(x') - f(x')| \\ &\leq 2 \sup_X |f - f_n| + \varepsilon/3 \end{aligned}$$

But $\exists N$ s.t. for $n \geq N$ $\sup |f - f_n| < \varepsilon/3$ and

$$|f(x) - f(x')| < \varepsilon$$

□

Theorem 1.60 (Sequential Arzela-Ascoli). *Let f_n be a sequence in $C(X)$ s.t.*

- i) f_n is uniformly bounded, i.e. $|f_n(x)| \leq M, \forall x \in X, \forall n \in \mathbb{N}$.
- ii) f_n is equicontinuous, i.e. $\forall \varepsilon > 0, \exists \delta$ s.t.

$$|f_n(x) - f_n(x')| < \varepsilon, \forall x, x' \text{ s.t. } d(x, x') < \delta, \forall n \in \mathbb{N}$$

Then f_n has a subsequence that converges uniformly to a function $f \in C(X)$.

Proof. $\mathcal{F} = \{f_n \in \mathbb{N}\}$, then $\overline{\mathcal{F}}$ is closed, bounded and equicontinuous. Then $\overline{\mathcal{F}}$ is compact and hence is sequentially compact. $\square$

Example 1.61. In $C([0, 1])$, let $\mathcal{F} = \{f \in \text{Lip}_1([0, 1]) : f(0) = \alpha\}$ where

$$\text{Lip}_1([0, 1]) = \{f : |f(x) - f(y)| \leq |x - y|, \forall x, y \in X\}$$

Now $\mathcal{F}$ is compact in $C(X)$.

- **Closed.** Take $f_n \in \mathcal{F}$ s.t. $f_n \rightarrow f$ in $C([0, 1])$. Clearly $f_n(0) = \alpha$ hence $f(0) = \alpha$. Now

$$|f_n(x) - f_n(y)| \leq |x - y|$$

let $n \rightarrow \infty$ we have

$$|f(x) - f(y)| \leq |x - y|$$

Hence $f \in \mathcal{F}$.

- **Bounded.** Now let $f \in \mathcal{F}$, then $|f(x) - f(0)| \leq |x - 0| \leq 1 + \alpha$, therefore for any $f \in \mathcal{F}$ and $x \in X$ we have

$$|f(x)| \leq |f(x) - f(0)| + |f(0)| \leq 1 + 2\alpha$$

Therefore

$$\sup |f(x)| \leq 1 + 2\alpha$$

So $\mathcal{F}$ is bounded.

- **Equicontinuous.** Let $\varepsilon > 0$, take $\delta = \varepsilon$, then for $|x - y| < \delta = \varepsilon$ we have

$$|f(x) - f(y)| \leq |x - y| < \varepsilon$$

By Arzela-Ascoli, we get that $\mathcal{F}$ is compact.

Example 1.62. Now fix $f \in C([0, 1])$, and look at $m = \inf_g \{d_\infty(f, g) \in \text{Lip}_1([0, 1]) : g(0) = f(0)\}$, want to show that this is attained. Indeed, let g_n be a sequence in $\text{Lip}_1([0, 1])$ $g_n(0) = f(0)$ with $d_\infty(f, g_n) \rightarrow m$. By compactness, g_n has convergent subsequence g_{n_k} in $C(X)$ to a function $g \in \text{Lip}_1([0, 1])$. Now $d_\infty(f, g) = \lim_{k \rightarrow \infty} d_\infty(f, g_{n_k}) = m$.

1.4 Baire Category Theorem

Definition 1.63. (X, τ) topological space, $A \subseteq X$. We say that A is nowhere dense in X iff $\overline{A}$ has empty interior.

Example 1.64. In $\mathbb{R}$, all singletons (or discrete sets) are nowhere dense in $\mathbb{R}$. But $\mathbb{Q}$ is not nowhere dense in $\mathbb{R}$.

Proposition 1.65. A is nowhere dense in $X \iff X \setminus \bar{A}$ is dense.

Proof. $\implies$ take $x \in X$ and $U_x \ni x$ open neighborhood, but x is not an interior point of $\bar{A}$ because $\overset{\circ}{\bar{A}} = \emptyset$, therefore $U_x \cap \bar{A} \neq \emptyset$.

$\impliedby$ Same.

□

Remark 1.66. A closed set is nowhere dense iff $\overset{\circ}{A} = \emptyset$.

Definition 1.67. (X, τ) topological space and $A \subseteq X$. We say that A is first Category or meagre in X iff A can be written as a countable union of nowhere dense subsets of X . Otherwise we say that A is second category (or non meagre).

Example 1.68. $\mathbb{Q}$ is first category in $\mathbb{R}$.

Remark 1.69. If a set A is second Category with $A = \bigcup_{n=1}^{\infty} F_n$ with F_n closed, then $\exists n_0$ with F_{n_0} has nonempty interior.

Theorem 1.70 (Baire Category Theorem). (X, d) complete metric space, then if $G_1, G_2, \dots$ are open dense subsets of X then $\bigcap_{n=1}^{\infty} G_n$ is dense in X . Equivalently, every nonempty open subset of X is second Category. In particular, if $X = \bigcup_{n=1}^{\infty} F_n$ with F_n closed, then $\exists n_0$ with F_{n_0} has nonempty interior.

Why equivalently?

$\implies$ Let G open in X , suppose that we can write $G = \bigcup_{n=1}^{\infty} A_n$ where A_n is nowhere dense. Take $G_i = X \setminus \bar{A}_i$, note that G_i is open and dense in X , therefore $\bigcap_{n=1}^{\infty} G_n$ is dense in X . But

$$\begin{aligned} \bigcap_{n=1}^{\infty} G_n &= \bigcap_{n=1}^{\infty} (\bar{A}_n)^c \\ &= \left(\bigcup_{n=1}^{\infty} \bar{A}_n \right)^c \end{aligned}$$

Which is dense in X . But we have that

$$\bigcup_{n=1}^{\infty} \bar{A}_n \supseteq G$$

with G open, contradiction.

$\impliedby$ Let $G_1, G_2, \dots$ be open dense subsets of X . Note that G_i is second Category, and suppose $\bigcap_{n=1}^{\infty} G_n$ is not dense, so $\exists H \subseteq (\bigcap_{n=1}^{\infty} G_n)^c = \bigcup_{n=1}^{\infty} G_n^c$ with H open. But this means that

$$H = \bigcup_{n=1}^{\infty} G_n^c \cap H$$

with H second category, then $\exists n_0 \in \mathbb{N}$ and an open set $U \subseteq G_{n_0}^c \cap H \subseteq G_{n_0}^c$, contradiction.

Proof of Baire Category Theorem. Take $x_0 \in X$ and $\delta > 0$, take $B_0 = B(x_0, \delta)$. Now B_0 intersects G_1 , and $\exists x_1 \in B_0 \cap G_1$ and $\delta_1 < \delta/2$ s.t.

$$B_1 = B(x_1, \delta_1) \subseteq \overline{B(x_1, \delta_1)} \subseteq B_0 \cap G_1$$

Now B_1 intersects G_2 , so $\exists x_2 \in B_1 \cap G_2$ and $\delta_2 < \delta/2^2$ with

$$B_2 = B(x_2, \delta_2) \subseteq \overline{B(x_2, \delta_2)} \subseteq B_1 \cap G_2 \subseteq B_0 \cap G_1 \cap G_2$$

Continue this recursively to get $x_n \in B_{n-1} \cap G_n$ and $\delta_n < \delta/2^n$ s.t.

$$B_n = B(x_n, \delta_n) \subseteq \overline{B(x_n, \delta_n)} \subseteq B_0 \cap G_1 \cap \dots \cap G_n$$

We show that x_n is a Cauchy sequence in X . Indeed, let $n, m > N$ for some $N \in \mathbb{N}$ then $x_n, x_m \in B_N = B(x_N, \delta_N)$, so

$$d(x_n, x_m) < 2\delta/2^N = \delta/2^{N-1}$$

Now let $\varepsilon > 0$, choose N s.t. $\delta/2^{N-1} < \varepsilon$, then

$$d(x_n, x_m) < \varepsilon$$

so x_n is Cauchy. By completeness, x_n converges to some $z \in X$. We show that $z \in (\bigcap_{n=1}^{\infty} G_n) \cap B_0$. Fix m , we have

$$x_n \in \overline{B_m}, \quad \forall n > m$$

Therefore $z \in \overline{B_m} \subseteq B_0 \cap (\bigcap_{k=1}^m G_k)$, since this holds for every m , we have

$$z \in B_0 \cap \left(\bigcap_{k=1}^{\infty} G_k \right)$$

□

2 Normed Spaces

Definition 2.1. Given a vector space V over $\mathbb{F}$, a norm is a map

$$\|\cdot\|: V \rightarrow \mathbb{R}$$

satisfying

- i) $\|x\| \geq 0$ with equality iff $x = 0$.
- ii) $\|\alpha x\| = |\alpha| \cdot \|x\|$, $\alpha \in \mathbb{F}$, $x \in V$.
- iii) $\|x + y\| \leq \|x\| + \|y\|$, $\forall x, y \in V$.

Example 2.2. $\ell^\infty(\mathbb{R})$ (set of bounded sequences) under coordinate-wise addition and scalar multiplication.

$$\|x\|_\infty = \sup_{i \in \mathbb{N}} |x_i| = d_\infty(x, 0)$$

$C(X)$ with X compact

$$\|f\|_\infty = \sup_{x \in X} |f(x)|$$

Proposition 2.3. Every norm induces a metric

$$d(x, y) = \|x - y\|$$

This allows us to deal with normed space as metric spaces w.r.t. induced metric.

Definition 2.4. V is a vector space and $\|\cdot\|_a, \|\cdot\|_b$ norms on V , we say that $\|\cdot\|_a, \|\cdot\|_b$ are equivalent iff they induce the same topology.

Proposition 2.5. V vector space, $\|\cdot\|_a, \|\cdot\|_b$ norms on V . The following are equivalent:

- $\|\cdot\|_a, \|\cdot\|_b$ are equivalent.
- $\exists c_1, c_2 > 0$ s.t.

$$c_2\|x\|_b \leq \|x\|_a \leq c_1\|x\|_b, \quad \forall x \in V$$

- $x_n \rightarrow x_0$ in $(V, \|\cdot\|_a) \iff x_n \rightarrow x_0$ in $(V, \|\cdot\|_b)$.

(Proof $\Rightarrow$ (ii) $B_{\|\cdot\|_a}(0, 1) = \{x \in V, \|x\|_a < 1\}$ is open in $(V, \|\cdot\|_a)$ then $B_{\|\cdot\|_a}(0, 1)$ is open in $(V, \|\cdot\|_b)$. Therefore 0 is an interior point of $B_{\|\cdot\|_a}(0, 1)$ w.r.t. the $\|\cdot\|_b$ norm. Therefore $\exists \delta > 0$ s.t.

$$B_{\|\cdot\|_b}(0, \delta) \subseteq B_{\|\cdot\|_a}(0, 1)$$

Now take $x \in V, x \neq 0$, then

$$\frac{\delta}{2} \frac{x}{\|x\|_b} \in B_{\|\cdot\|_b}(0, \delta) \subseteq B_{\|\cdot\|_a}(0, 1)$$

Hence

$$\left| \frac{\delta}{2} \frac{x}{\|x\|_b} \right|_a < 1 \implies \|x\|_b < \frac{2}{\delta} \|x\|_a$$

do the same by switching $\|\cdot\|_a$ and $\|\cdot\|_b$.

(ii) $\implies$ (iii) We have that

$$c_1\|x_n - x_0\|_b \leq \|x_n - x_0\|_a \leq c_1\|x_n - x_0\|_b$$

(iii) $\implies$ (i) Closed in $(V, \|\cdot\|_a) \iff$ Closed in $(V, \|\cdot\|_b)$ by sequential definition.

□

Definition 2.6. Take V finite dimensional vector space and $S = \{v_1, \dots, v_n\}$ a basis of V . Take $x \in V$, then $\exists! r_1, r_2, \dots, r_n$ s.t.

$$x = r_1 v_1 + \dots + r_n v_n$$

We define

$$\|x\|_\infty := \max_{1 \leq i \leq n} |r_i|$$

We then have the following:

Proposition 2.7. *V is a f.d. vector space and $S = \{v_1, \dots, v_n\}$ a basis:*

1. $(V, \|\cdot\|_\infty)$ is complete.
2. $\{x \in V : \|x\|_\infty \leq M\}$ is compact.
3. $K \subseteq (V, \|\cdot\|_\infty)$ is compact iff K is closed and bounded.

Proof. 1. Let x_k be a Cauchy sequence in V , then

$$x_k = r_{k1}v_1 + \dots + r_{kn}v_n$$

Then

$$|r_{ki} - r_{li}| \leq \|x_k - x_l\|_\infty$$

Then the sequence $(r_{ki})_k$ is Cauchy in $\mathbb{R}$ for every i , then r_{ki} converges to some r_i . Take $x = r_1v_1 + \dots + r_nv_n$, we get that

$$\begin{aligned} \|x_k - x\|_\infty &= \|(r_{k1} - r_1)v_1 + \dots + (r_{kn} - r_n)v_n\|_\infty \\ &\leq \|r_{k1} - r_1\| \|v_1\|_\infty + \dots + \|r_{kn} - r_n\| \|v_n\|_\infty \\ &\rightarrow 0 \end{aligned}$$

2. Take a sequence x s.t. $\|x_k\|_\infty \leq M$. Apply Bolzano Weierstrass n times and take subsequence of the subsequence.
3. $(\implies)$ is always true. $(\impliedby)$ since K is bounded, then $\exists M$ s.t.

$$K \subseteq \{x \in V : \|x\|_\infty \leq M\} =: Q$$

But K is closed Q is compact, then K is compact.

□

Theorem 2.8. *V is a finite dimensional vector space, then all norms on V are equivalent.*

Corollary 2.9. *V finite dimensional and $\|\cdot\|$ is a norm on V then $(V, \|\cdot\|)$ has the properties of the theorem above.*

Moreover, if V is a normed space and M is a finite dimensional subspace of V .

Remark 2.10. Note that the converse is not true in general, take $P([a, b])$ as a subspace of $C([a, b])$.

Proof. Take $S = \{v_1, \dots, v_n\}$ a basis of V , define

$$\|x\|_\infty = \max_{1 \leq i \leq N} |r_i|$$

Let $\|\cdot\|_a$ be a norm on V , we will show that $\|\cdot\|_a$ is equivalent to $\|\cdot\|$.

$$\begin{aligned}\|x\|_a &= \|r_1 v_1 + \cdots + r_n v_n\|_a \\ &\leq |r_1| \|v_1\|_a + \cdots + |r_n| \|v_n\|_a \\ &\leq (\|v_1\|_a + \cdots + \|v_n\|_a) \|x\|_\infty\end{aligned}$$

Let $\varphi: (V, \|\cdot\|_\infty) \rightarrow \mathbb{R}$, note that this is continuous since if $x_n \rightarrow x_0$ in $\|\cdot\|_\infty$, then

$$\begin{aligned}|\varphi(x_n) - \varphi(x_0)| &= (\|x_n\|_a - \|x_0\|_a) \\ &\leq \|x_n - x_0\|_a \\ &\leq c_1 \|x_n - x_0\|_\infty \\ &\rightarrow 0\end{aligned}$$

Now let $K = \{x \in V: \|x\|_\infty = 1\}$, note that K is compact (closed and bounded), so $\varphi(K)$ is compact in $\mathbb{R}$ so $m = \inf \varphi(K)$ is attained at some $x_0 \in K$, $x_0 \neq 0$. Then $m = \|x_0\|_a > 0$. So $\forall x \in V$ s.t. $\|x\|_\infty = 1$ we have

$$\|x\|_a \geq m$$

Take $v \in V$ s.t. $x \neq 0$ we have that

$$\begin{aligned}\left\| \frac{x}{\|x\|_\infty} \right\|_a &\geq m \\ \|x\|_a &\geq m \|x\|_\infty\end{aligned}$$

□

Lemma 2.11 (Riesz Lemma). *$(V, \|\cdot\|)$ is a normed v.s. and M a proper closed subspace of V . Then for every $0 < \alpha < 1$ there exists $x \in V$ s.t. $\|x\| = 1$ and*

$$d(x, M) \geq \alpha$$

Remark 2.12. If x is s.t. $\|x\| = 1$, then $d(x, M) \leq \|x - 0\| \leq 1$.

Proof. Take $x_0 \notin M$, $d(x_0, M) > 0$ since M is closed. Fix $0 < \alpha < 1$, there exists $m_0 \in M$ s.t.

$$d(x_0, M) \leq \|x_0 - m_0\| < \frac{d(x_0, M)}{\alpha}$$

Define

$$x = \frac{x_0 - m_0}{\|x_0 - m_0\|}, \quad \|x\| = 1$$

Let $m \in M$, then

$$\begin{aligned}\|x - m\| &= \left\| \frac{x_0 - m_0}{\|x_0 - m_0\|} - m \right\| \\ &= \frac{\|x_0 - (m_0 + m\|x_0 - m_0\|)\|}{\|x_0 - m_0\|}\end{aligned}$$

But $m_0 + m\|x_0 - m_0\| \in M$, so

$$\|x - m\| \geq \frac{d(x_0, M)}{\|x_0 - m_0\|} > \alpha$$

Therefore

$$d(x, M) = \inf_{m \in M} \|x - m\| \geq \alpha$$

□

Theorem 2.13. $(V, \|\cdot\|)$ normed space, the closed unit sphere in V is compact iff V is finite dimensional.

Proof. $\Leftarrow$ ok.

$\Rightarrow$ Assume that V is infinite dimensional, then we can construct

$$E_1 = \text{Span}\{v_1\}, E_2 = \text{Span}\{v_1, v_2\}, \dots$$

With E_i finite dimensional and $\{v_i\}$ linearly independent. Then

$$E_1 \subsetneq E_2 \subsetneq E_3 \subsetneq \dots$$

Fix some $u_1 \in E_1$ with $\|u_1\| = 1$, now by Riesz' lemma $\exists u_2 \in E_2$ s.t. $\|u_2\| = 1$ and $d(u_2, E_1) \geq 1/2$. Now $\exists u_3 \in E_3$ with $\|u_3\| = 1$ and $d(u_3, E_2) \geq 1/2$, recursively we can construct $u_n \in E_n$ s.t. $\|u_n\| = 1$ and $d(u_n, E_{n-1}) \geq 1/2$. Now notice that for $m \neq n$ we have

$$\|u_m - u_n\| \geq 1/2$$

Therefore u_n is a sequence which belongs to the unit sphere that has no Cauchy (and hence convergent) subsequence. So the unit sphere is not sequentially compact, and hence not compact.

□

Definition 2.14. A complete normed space is called a Banach space.

Example 2.15. Some Banach spaces: $\mathbb{R}^N$. $\ell^\infty(\mathbb{R})$ (which is the set of bounded sequences). $\ell^1(\mathbb{R}) = \{x: (x_1, x_2, \dots) : \sum |x_i| < \infty\}$. $C(X)$ when X is compact.

$P(\mathbb{R})$ is not Banach.

ℓ^p Spaces.

Lemma 2.16. $1 < p, q < \infty$ s.t. $\frac{1}{p} + \frac{1}{q} = 1$ and $x, y > 0$ s.t. $xy = 1$ then

$$\frac{x^p}{p} + \frac{y^q}{q} \geq 1$$

Proof. $f(x, y) = \frac{x^p}{p} + \frac{y^q}{q}$, we want to minimize $f(x, y)$ on $g(x, y) = xy = 1$. Note that this minimum exists since this is a closed set and $f(x, y) \geq 0$ for all x, y s.t. $xy = 1$. Using Lagrange multipliers, we solve

$$\begin{cases} \nabla f &= \lambda \nabla g \\ xy &= 1 \end{cases}$$

i.e.

$$\begin{cases} x^{p-1} = \lambda y \\ y^{q-1} = \lambda x \\ xy = 1 \end{cases}$$

Then we get

$$x^p = \lambda xy = \lambda, \quad y^q = \lambda$$

Then $x = \lambda^{1/p}, y = \lambda^{1/q}$, then

$$\lambda^{\frac{1}{p} + \frac{1}{q}} = 1 \implies \lambda = 1$$

Then $x = y = 1$. So the minimum is $f(1, 1) = 1$. □

Corollary 2.17. $\frac{1}{p} + \frac{1}{q} = 1, x, y \geq 0$ then

$$\frac{x^p}{p} + \frac{y^q}{q} \geq xy$$

Proof. Take $X = \frac{x}{(xy)^{1/p}}$ and $Y = \frac{y}{(xy)^{1/q}}$ then

$$XY = \frac{xy}{(xy)^{1/p + 1/q}} = 1$$

Then apply the lemma, we get

$$\begin{aligned} \frac{X^p}{p} + \frac{Y^q}{q} &\geq 1 \\ \frac{x^p}{(xy)p} + \frac{y^q}{(xy)q} &\geq 1 \\ \frac{x^p}{p} + \frac{y^q}{q} &\geq xy \end{aligned}$$

□

Theorem 2.18 (Young's inequality). Let $x, y \in \mathbb{R}$ then

$$|xy| \leq \frac{|x|^p}{p} + \frac{|y|^q}{q}$$

with $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 2.19 (Holder inequality). Let $x_1, x_2, \dots, x_n \in \mathbb{R}$ and $y_1, y_2, \dots, y_n \in \mathbb{R}$ then

$$\sum_{i=1}^n |x_i y_i| \leq \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \left(\sum_{i=1}^n |y_i|^q \right)^{1/q}$$

With $\frac{1}{p} + \frac{1}{q} = 1$.

Letting $n \rightarrow \infty$ we have

$$\sum_{n=1}^{\infty} |x_n y_n| \leq \left(\sum_{n=1}^{\infty} |x_n|^p \right)^{1/p} \left(\sum_{n=1}^{\infty} |y_n|^q \right)^{1/q}$$

Proof. Wlog suppose not all zeroes. Let $X_i = \frac{|x_i|}{(\sum_{k=1}^n |x_k|^p)^{1/p}}$ and $y_i = \frac{|y_i|}{(\sum_{k=1}^n |y_k|^q)^{1/q}}$. From Young's inequality, we have that

$$\begin{aligned} X_i Y_i &= \frac{|x_i| |y_i|}{(\sum_{k=1}^n |x_k|^p)^{1/p} (\sum_{k=1}^n |y_k|^q)^{1/q}} \\ &\leq \frac{X_i^p}{p} + \frac{Y_i^q}{q} \\ &= \frac{|x_i|^p}{(\sum_{k=1}^n |x_k|^p) p} + \frac{|y_i|^q}{(\sum_{k=1}^n |y_k|^q) q} \end{aligned}$$

Therefore

$$\begin{aligned} \sum_{i=1}^n X_i Y_i &\leq \frac{1}{p} + \frac{1}{q} \\ &= 1 \end{aligned}$$

Substituting we get the desired inequality. $\square$

Theorem 2.20 (Minkowski inequality). $(|x_i + y_i|)^p \leq (\sum_{i=1}^n |x_i|^p)^{1/p} + (\sum_{i=1}^n |y_i|^p)^{1/p}$. In the infinite case we get

$$\left(\sum_{n=1}^{\infty} |x_n + y_n|^p \right)^{1/p} \leq \left(\sum_{n=1}^{\infty} |x_n|^p \right)^{1/p} + \left(\sum_{n=1}^{\infty} |y_n|^p \right)^{1/p}$$

Proof. If $p = 1$, trivial. Take $p > 1$ and let q be s.t. $\frac{1}{p} + \frac{1}{q} = 1$. Then

$$\begin{aligned} \sum_{i=1}^n |x_i + y_i|^p &= \sum_{i=1}^n |x_i + y_i| \cdot |x_i + y_i|^{p-1} \\ &\leq \sum_{i=1}^n (|x_i| + |y_i|) |x_i + y_i|^{p-1} \\ &= \sum_{i=1}^n |x_i| |x_i + y_i|^{p-1} + \sum_{i=1}^n |y_i| |x_i + y_i|^{p-1} \\ &\leq \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \left(\sum_{i=1}^n |x_i + y_i|^{(p-1)q} \right)^{1/q} \\ &\quad + \left(\sum_{i=1}^n |y_i|^p \right)^{1/p} \left(\sum_{i=1}^n |x_i + y_i|^{(p-1)q} \right)^{1/q} \end{aligned}$$

We have that $\frac{1}{p} + \frac{1}{q} = 1 \implies p + q = pq \implies p = q(p-1)$, therefore

$$\begin{aligned} \sum_{i=1}^n |x_i + y_i|^p &\leq \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{1/q} \left(\left(\sum_{i=1}^n |x_i|^p \right)^{1/p} + \left(\sum_{i=1}^n |y_i|^p \right)^{1/p} \right) \\ \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{1-1/q=1/p} &\leq \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} + \left(\sum_{i=1}^n |y_i|^p \right)^{1/p} \end{aligned}$$

as desired. $\square$

Definition 2.21. Let $p \geq 1$, we define the space $\ell^p(\mathbb{R}) = \{x = (x_1, \dots) : \sum_{n=1}^{\infty} |x_n|^p < \infty\}$ with the following norm $\|x\|_{\ell^p} = (\sum_{n=1}^{\infty} |x_n|^p)^{1/p}$.

Proposition 2.22. • $\ell^p(\mathbb{R})$ is complete $\forall 1 \leq p \leq \infty$.

- $\ell^p(\mathbb{R})$ is separable $\forall 1 \leq p < \infty$ (recall that $\ell^\infty(\mathbb{R})$ is not separable). Since $c_{00}(\mathbb{Q})$ is dense in $\ell^p(\mathbb{R})$.
- $K \subseteq \ell^p(\mathbb{R})$ ($p \leq 1 < \infty$) is compact iff it is closed and bounded and

$$\lim_{N \rightarrow \infty} \sup_{x \in K} \sum_{n=1}^{\infty} |x_n|^p = 0$$

- $1 \leq r \leq s < \infty$, then

$$\ell^r(\mathbb{R}) \subseteq \ell^s(\mathbb{R})$$

Hence

$$\ell^1(\mathbb{R}) \subseteq \ell^r(\mathbb{R}) \subseteq \ell^s(\mathbb{R}) \subseteq \ell^\infty(\mathbb{R})$$

- If $1 \leq r < s < \infty$ then $\ell^r(\mathbb{R}) \subsetneq \ell^s(\mathbb{R})$

$$x_n = \frac{1}{n^{1/r}}$$

Then $\sum |x_n| < \infty$ but $\sum |x_n|^r = \infty$ so $x_n \in \ell^s(\mathbb{R})$ and $x_n \notin \ell^r(\mathbb{R})$.

2.1 Bounded Linear Transformation

Definition 2.23. V, W vector spaces and $\mathcal{L}(V, W)$ is the space of linear transformations $T: V \rightarrow W$, this is a vector space.

Remark 2.24. If V, W are finite dimensional then $\mathcal{L}(V, W)$ is finite dimensional with

$$\dim \mathcal{L}(V, W) = \dim(V) \cdot \dim(W)$$

Definition 2.25. Take $(V, \|\cdot\|_V)$ and $(W, \|\cdot\|_W)$ and $T \in \mathcal{L}(V, W)$ then we say that T is bounded iff $\exists M > 0$ s.t.

$$\|Tx\|_W \leq M\|x\|_V$$

Equivalently $\exists M$ s.t.

$$\|Tx\| \leq M, \quad \forall x \text{ s.t. } \|x\|_V = 1$$

Proposition 2.26. $(V, \|\cdot\|_V)$ and $(W, \|\cdot\|_W)$ normed spaces and $T \in \mathcal{L}(V, W)$, then

$$T \text{ bounded} \iff T \text{ is continuous} \iff T \text{ is continuous at } 0$$

(Proof $\Rightarrow$) (ii) Then $\exists M$ s.t. $\|Tx\| \leq M\|x\|$. Then

$$\begin{aligned} \|Tx - Ty\| &= \|T(x - y)\| \\ &\leq M\|x - y\| \end{aligned}$$

Then T is Lipschitz so it is continuous.

(ii) $\implies$ (iii) ok.

(iii) $\implies$ (i) Let T be continuous at 0, then $\exists \delta > 0$ s.t. $\|Tx - T0\| < 1$ for $\|x - 0\| < \delta$ but this gives

$$\|Tx\| \leq 1$$

for $\|x\| < \delta$. Now take $x \neq 0$, then

$$\|T\left(\frac{\delta}{\|x\|}x\right)\| \leq 1$$

So we get

$$\|Tx\| \leq \frac{\delta}{2}\|x\|$$

□

Remark 2.27. If V is finite dimensional, and $T \in \mathcal{L}(V, W)$, then T is continuous. Take $x \in V$, and $S = \{v_1, \dots, v_n\}$ basis of V . Then

$$x = r_1v_1 + \dots + r_nv_n \implies \|Tx\| \leq r_1\|Tv_1\| + \dots + r_n\|Tv_n\| \leq (\|Tv_1\| + \dots + \|Tv_n\|)\|x\|_\infty$$

But all norms on V are equivalent, then

$$\|Tx\|_W \leq M\|x\|_V$$

Then T is bounded.

Definition 2.28. $(V, \|\cdot\|)$, $(W, \|\cdot\|)$ normed spaces, we define $B(V, W)$ to be the space of bounded linear transformations with the following norm

$$\|T\|_{B(V,W)} = \sup_{\substack{x \in V \\ \|x\|=1}} \|Tx\|_W = \sup_{\substack{x \in V \\ x \neq 0}} \frac{\|Tx\|}{\|x\|}$$

Check that this is subspace of $\ell(V, W)$ and $\|\cdot\|_{B(V,W)}$ is a norm.

Remark 2.29. Notice that for $x \in V$ $x \neq 0$ we have that

$$\left|T \frac{x}{\|x\|}\right| \leq \|T\|$$

$$\|Tx\| \leq \|T\| \cdot \|x\|$$

Proposition 2.30. If W is Banach, then $B(V, W)$ is Banach.

Proof. Let T_n be a Cauchy sequence in $B(V, W)$, note that

$$\|T_n x - T_m x\| \leq \|T_n - T_m\| \cdot \|x\|$$

But T_n is Cauchy in $B(V, W)$, therefore $T_n(x)$ is Cauchy in W (which is Banach), then it converges. So we define

$$T: V \rightarrow W, \quad T(x) = \lim_{n \rightarrow \infty} T_n(x)$$

We want to show T is linear, this is trivial by properties of limits and that T_n are linear. Moreover $\{T_n\}$ is Cauchy in $B(V, W)$, then it is bounded i.e. $\exists M > 0$ s.t. $\|T_n\| \leq M, \forall n$. Now

$$\|T_n x\| \leq \|T_n\| \cdot \|x\|$$

$$\|T_n x\| \leq M \|x\|$$

Letting $n \rightarrow \infty$ we get that $\|Tx\| \leq M\|x\|$ so $T \in B(V, W)$.

We want to show that $T_n \rightarrow T$ in $B(V, W)$. Given $\varepsilon > 0$, since T_n is Cauchy, so $\exists N_1$ s.t.

$$\|T_n - T_m\| < \varepsilon/2, \quad \forall n, m \geq N_1$$

Now fix $x \in V$, since $T_n x \rightarrow Tx$ in W we have that $\exists N_2(x) > N_1$ s.t.

$$\|T_{N_2} x - Tx\| < \varepsilon/2$$

Now for $n \geq N_1$ and $\|x\| = 1$ we have

$$\begin{aligned} \|T_n x - Tx\| &< \|T_n x - T_{N_2} x\| + \|T_{N_2} x - Tx\| \\ &\leq \|T_n - T_{N_2}\| \cdot \|x\| + \frac{\varepsilon}{2} \\ &< \varepsilon \end{aligned}$$

Therefore given $\varepsilon > 0, \exists N_1$ s.t. $\forall x \in V$ with $\|x\| = 1$ then

$$\|T_n x - Tx\| < \varepsilon, \quad \forall n \geq N_1$$

Therefore

$$\sup_{\substack{x \in V \\ \|x\|=1}} \|T_n x - Tx\| \leq \varepsilon, \quad \forall n \geq N_1$$

Hence

$$T_n \rightarrow T \in B(V, W)$$

□

Example 2.31. Define

$$T: (C([0, 1]), \|\cdot\|_\infty) \rightarrow \mathbb{R}, \quad T(f) = \int_0^1 f(x) dx$$

Show that $T \in B(C([0, 1]), \mathbb{R})$ and find $\|T\|$.

Solution: Clearly T is linear because integration is linear. Now

$$\begin{aligned} |Tf| &= \left| \int_0^1 f(x) dx \right| \\ &\leq \int_0^1 |f(x)| dx \\ &\leq \|f\|_\infty \end{aligned}$$

Take $T \in B(C([0, 1]), \mathbb{R})$, note that

$$\sup_{\|f\|_\infty} \|Tf\| \leq 1$$

moreover if we take $f = 1$ we have that $\|Tf\| = 1$. Therefore

$$\|T\| = 1$$

2.2 Duality

Definition 2.32. Let V normed space over $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$. Then the dual set $V^* = B(V, \mathbb{F})$, which is the set of all bounded linear maps from $V \rightarrow \mathbb{F}$. In particular, the dual norm

$$\|f\|_* = \sup_{\substack{x \in V \\ \|x\|=1}} \|f(x)\| = \sup_{\substack{x \in V \\ x \neq 0}} \frac{|f(x)|}{\|x\|}$$

Remark 2.33. If V is finite dimensional, then $V^* = B(V, \mathbb{F}) = \mathcal{L}(V, \mathbb{F})$, then

$$\dim V^* = \dim V < \infty$$

Therefore V^* and V are isomorphic.

notation. Given $(V, \|\cdot\|)$, $(W, \|\cdot\|)$ we write $V \cong W$ iff V and W are isometrically isomorphic. i.e. $\exists \varphi: V \rightarrow W$ and isomorphism and isometry.

Example 2.34. $(\ell^1(\mathbb{R}))^* \cong \ell^\infty(\mathbb{R})$, more particularly if $1 < p < q < \infty$ and $1/p + 1/q = 1$, then

$$(\ell^p)^* \cong \ell^q$$

So for $p = 2$ we get that

$$(\ell^2)^* \cong \ell^2$$

Now if $1 < p < q < +\infty$ then

$$(\ell^p)^{**} \cong (\ell^q)^* \cong (\ell^p)^*$$

If $p = 1$, then

$$(\ell^1)^* \cong \ell^\infty \text{ but } (\ell^\infty)^* \neq \ell^1$$