QMWS Fall 2026: Introduction to Python for Data Analysis

Day Three: Applications

Gavin Klorfine
gklorfin@yorku.ca

Deborah Laze
laze@yorku.ca

York University

Today

1. matplotlib.pyplot
2. statsmodels

Section 1

matplotlib.pyplot

matplotlib.pyplot

- **matplotlib.pyplot** is a package for visualizing data in Python
- Fun fact: If you have asked ChatGPT for a graph, you likely received one that was made using this package


matplotlib.pyplot

Basic Histogram

```
import numpy as np
import matplotlib.pyplot as plt

x = np.random.normal(0, 1, 1000)
plt.hist(x)
plt.show()
```

Basic Histogram




- A few ways to modify the histogram are illustrated below

Modifying the Plot

```
plt.hist(
    x,                          # Data
    color = "green",            # Bar Colour
    edgecolor = "black",        # Outline Colour
    bins = np.arange(-3, 3.5, 0.5) # Bins of 0.5 from -3 to +3
)
plt.show()
```

Modifying the Plot




More Ways to Modify

```
plt.hist(
    x, color = "green", edgecolor = "black",
    bins = np.arange(-3, 3.5, 0.5)
)

plt.xticks(np.arange(-3, 4, 1)) # x-axis ticks spaced by 1
plt.title("Sample from N ~ (0, 1)")
plt.xlabel("Value")
plt.ylabel("Count")
plt.show()
```

More Ways to Modify




- The below is how data was generated for subsequent scatterplot examples

Simulated Data for a Simple Regression

```
errors = np.random.normal(0, 2, 10) # Normal errors
nums_x = np.arange(0, 10, 1) # Nums 0 to 9
nums_y = nums_x + errors # y = x + errors
```

Scatterplot

```
# scatter(x, y, colour, transparency)
plt.scatter(nums_x, nums_y, color = "red", alpha = 0.6)
plt.show()
```




- To colour the dots by a **categorical** variable (e.g., group), call `plt.scatter()` once per group
- Give each call its own `color` and `label`, then add `plt.legend()`

Colouring by Group (Categorical)

```
groups = np.array(["A", "B"] * 5)     # Group label for each po

plt.scatter(nums_x[groups == "A"], nums_y[groups == "A"],
    color = "blue", label = "Group A")
plt.scatter(nums_x[groups == "B"], nums_y[groups == "B"],
    color = "red", label = "Group B")
plt.legend()
plt.show()
```

Colouring by Group (Categorical)




- `groups == "A"` selects only the points in Group A
- `plt.legend()` uses each call's `label` to name the groups

- For a **continuous** variable, pass the numbers to `c` and choose a colour map with `cmap`
- `plt.colorbar()` acts as the legend, showing which colour matches which value

Colouring with a Colour Map (Continuous)

```
plt.scatter(
    nums_x, nums_y,
    c = errors,           # Values that determine each dot's co
    cmap = "coolwarm"     # Colour map: low = blue, high = red
)
plt.colorbar(label = "Error")
plt.show()
```

Colouring with a Colour Map (Continuous)



- Redder dots have larger errors (further above the line $y = x$); bluer dots have smaller errors (further below it)
- Other colour maps include `"viridis"`, `"plasma"`, and `"Greys"`

`color` vs. `c` + `cmap`

- `color` + `label`: one colour per group, one `plt.scatter()` call per group; legend with `plt.legend()` (categorical)
- `c` + `cmap`: a list of numbers that matplotlib converts to colours; legend with `plt.colorbar()` (continuous)

Note

- `cmap` only works with `c` — matplotlib needs numbers to map onto a colour scale

matplotlib.pyplot


- You can add another plot layer by calling another plotting function before `plt.show()`
- This is useful for adding:
  - reference lines
  - fitted lines
  - multiple groups

Adding a Line to a Scatterplot

```
# Observed data as red dots
plt.scatter(nums_x, nums_y, color = "red")
# Reference line: black solid line
plt.plot(nums_x, nums_x, '-k')
plt.show()
```

- `plt.scatter()` draws dots; `plt.plot()` with `'-k'` draws a line

Adding Labels and a Legend



- `label` gives each layer a name
- `plt.legend()` displays the labels on the plot

Adding Labels and a Legend, continued

- `plt.title()`, `plt.xlabel()`, and `plt.ylabel()` make the plot easier to interpret

Section 2

statsmodels

statsmodels

- **statsmodels** is a package for statistical testing and modelling
- It is useful when you want:
  - formal hypothesis tests
  - regression models
  - p-values and confidence intervals
  - detailed statistical output

Importing **statsmodels** Functions

```
import statsmodels.formula.api as smf
from statsmodels.stats.weightstats import ttest_ind
```

statsmodels

- **statsmodels.formula.api** contains functions that use formula notation
- It is commonly renamed `smf`
- `ttest_ind` is a function for independent-samples t-tests

Import Structure

```
import statsmodels.formula.api as smf
#      package.module          alias

from statsmodels.stats.weightstats import ttest_ind
#    package.module.submodule        function
```

Statistical Testing

- Statistical tests help us use a **sample** to make inferences about a larger population
- A test usually starts with a **null hypothesis**
  - Often "no difference" or "no relationship"
- We compare the observed data to what would be expected if the null hypothesis were true

Note

- The code runs the test, but the researcher chooses whether the test makes sense

statsmodels

General Workflow

1. State the research question
2. Identify the level of measurements and study design
3. Summarize and visualize the data
4. Run the appropriate test
5. Interpret the estimate, confidence interval, and p-value

Example Data

- We will use simulated scores from two independent groups
- Each row represents one participant
- The group labels are stored in a pandas DataFrame

Creating the Data

```
import pandas as pd

np.random.seed(42)

df = pd.DataFrame({
    "group": ["control"] * 30 + ["treatment"] * 30,
    "score": np.concatenate([
        np.random.normal(70, 10, 30),
        np.random.normal(76, 10, 30)
    ])
})
```

Inspecting the Data

```
print(df.head())    # First five rows
print(df.shape)     # (number of rows, number of columns)
```

```
     group      score
0  control  74.967142
1  control  68.617357
2  control  76.476885
3  control  85.230299
4  control  67.658466
(60, 2)
```

- Always inspect the data before running a test

Independent-Samples t-test

- Used to compare the means of **two independent groups**
- Do not use this test if the same participants appear in both groups
- Two versions exist depending on whether the groups' variances are assumed equal — both use `ttest_ind()`, just with a different `usevar` argument
- We'll work through this test using our general workflow

Research Question:

- Do treatment and control groups have different mean scores?

Null hypothesis:

$$H_0 : \mu_{\text{treatment}} - \mu_{\text{control}} = 0$$

Independent-Samples t-test: Level of Measurement

- `group`: categorical (nominal), two independent levels — grouping variable
  - Study design: independent (between-subjects) groups
- `score`: continuous (interval/ratio) — outcome variable

Independent-Samples t-test: Inspecting the Data

```
print(df.groupby("group")["score"].describe().round(2))
```

group	count	mean	std	min	25%	50%	75%	max
control	30.0	68.12	9.00	50.87	64.09	67.66	73.60	85.79
treatment	30.0	74.79	9.31	56.40	68.91	75.35	81.45	94.52

- Always re-check the data immediately before running a test
- `groupby("group")` summarizes the control and treatment rows separately
- `.describe()` gives common summary statistics for each group

Selecting Each Group

```
control = df.loc[
    df["group"] == "control",  # Rows to include
    "score"                    # Column to return
]

treatment = df.loc[
    df["group"] == "treatment",
    "score"
]
```

- `ttest_ind()` needs the two groups as separate variables

Independent-Samples t-test: The `usevar` Argument

```
ttest_ind(
    x1,                  # First group's values
    x2,                  # Second group's values
    alternative = "two-sided",
    usevar = "pooled",   # or "unequal" (Welch's)
    value = 0            # Null-hypothesis difference
)
```

- `usevar = "pooled"` assumes equal population variances (the regular independent-samples t-test)
- `usevar = "unequal"` does **not** assume equal population variances (**Welch's t-test**) — a useful default when in doubt

Independent-Samples t-test: The `usevar` Argument

- Use `alternative = "two-sided"` when either group could have the larger mean
- Use `value = 0` when the null hypothesis is no mean difference

Independent-Samples t-test: Running the Test

```
from statsmodels.stats.weightstats import ttest_ind

pool_result = ttest_ind(
    treatment, control, usevar = "pooled"
)
welch_result = ttest_ind(
    treatment, control, usevar = "unequal"
)
```

- Only the `usevar` argument changes between the two tests

Independent-Samples t-test: Results

- `ttest_ind()` returns three values: the t-statistic, the p-value, and the degrees of freedom.

```
print(pool_result)
print(welch_result)
```

```
(np.float64(2.8210747685177098), np.float64(0.006542565964576
(np.float64(2.82107476851771), np.float64(0.006544738502602256
```

```
# You can also index these results:
print(f"{pool_result[0]},"
f"{pool_result[1]}, {pool_result[2]}")
print(f"{welch_result[0]},"
f"{welch_result[1]}, {welch_result[2]}")
```

```
2.8210747685177098,0.006542565964576848, 58.0
2.82107476851771,0.006544738502602256, 57.93320998252979
```

t-Test Interpretation

- The t-statistic describes the group difference relative to its uncertainty
- The p-value is the probability, assuming the null hypothesis is true, of observing a test statistic at least as extreme as the one observed
- The degrees of freedom help determine the t-distribution used by the test

Important

- A p-value is **not** the probability that the null hypothesis is true

Calculating the Mean Difference

```
mean_difference = treatment.mean() - control.mean()

print(f"Treatment mean: {treatment.mean():.2f}")
print(f"Control mean: {control.mean():.2f}")
print(f"Mean difference: {mean_difference:.2f}")
```

```
Treatment mean: 74.79
Control mean: 68.12
Mean difference: 6.67
```

- The t-test output should be interpreted alongside the observed group difference
- The mean difference itself does not depend on the variance assumption, so it is the same for both versions

Interpreting the Results: Equal Variances

```
print(
    f"Mean difference = {mean_difference:.2f}\n"
    f"t({pool_result[2]:.2f}) = {pool_result[0]:.2f}\n"
    f"p = {pool_result[1]:.3f}"
)
```

```
Mean difference = 6.67
t(58.00) = 2.82
p = 0.007
```

- Assumes the two groups have equal population variances

Interpreting the Results: Unequal Variances (Welch's)

```
print(
    f"Mean difference = {mean_difference:.2f}\n"
    f"t({welch_result[2]:.2f}) = {welch_result[0]:.2f}\n"
    f"p = {welch_result[1]:.3f}"
)
```

```
Mean difference = 6.67
t(57.93) = 2.82
p = 0.007
```

- Does **not** assume equal population variances
- Compare the two reports above — here the conclusion is the same either way, which is common when group variances are similar

Correlation

- A correlation describes the direction and strength of a **linear relationship**
- Pearson's correlation coefficient is represented by r
- It ranges from -1 to 1

Research Question:

- Is there a linear relationship between study hours and score?

! Important

- Correlation does not establish causation

Correlation: Level of Measurement

- `study_hours`: continuous (interval/ratio)
- `score`: continuous (interval/ratio)
- Study design: observational/correlational (no groups being compared)
- Both variables are continuous, so a Pearson correlation is appropriate

Adding a Second Variable

```
np.random.seed(7)


df["study_hours"] = (
    2
    + 0.08 * df["score"]
    + np.random.normal(0, 1.5, len(df))
)
```

Scatterplot

```
plt.scatter(df["study_hours"], df["score"], color = "blue")
plt.xlabel("Study Hours")
plt.ylabel("Score")
plt.show()
```

- Use a scatterplot before calculating a correlation — this is the *inspecting the data* step of our workflow

Scatterplot



.corr() Method

```
correlation = df["study_hours"].corr(
    df["score"]
)

print(f"r = {correlation:.2f}")
```

```
r = 0.53
```

- .corr() describes the observed sample relationship

Correlation: Running the Test

- `.corr()` gives the **estimate** (r) of the linear relationship, but not a p-value or confidence interval on its own
- In **simple linear regression** with one predictor, the slope's significance test is mathematically equivalent to testing whether r differs from zero
- We'll get the confidence interval and p-value for this same relationship next, using regression

Simple Linear Regression

- Simple linear regression examines the relationship between:
  - one continuous **outcome**
  - one continuous **predictor**
- Here, the outcome is `score` and the predictor is `study_hours`

Research Question (same as above):

- Is there a linear relationship between study hours and score?

$$\text{score} = \text{intercept} + (\text{slope} \times \text{study hours}) + \text{error}$$

Simple Linear Regression: Level of Measurement


- `study_hours`: continuous (interval/ratio) — predictor
- `score`: continuous (interval/ratio) — outcome
- Study design: observational/correlational, same as above

Simple Linear Regression: Inspecting the Data

```
plt.scatter(df["study_hours"], df["score"], color = "blue")
plt.xlabel("Study Hours")
plt.ylabel("Score")
plt.show()
```

- Same scatterplot as before — always inspect the data before fitting a model

Simple Linear Regression: Inspecting the Data



Formula Notation: Formula Structure

```
"score ~ study_hours"
# outcome ~ predictor
```

- The outcome goes on the left of ~
- The predictor goes on the right of ~
- Column names must match the DataFrame

Creating a Model with `smf.ols()`

```
model_definition = smf.ols(
    formula = "score ~ study_hours",
    data = df
)

print(type(model_definition))
```

```
<class 'statsmodels.regression.linear_model.OLS'>
```

- `ols` stands for **ordinary least squares**
- `smf.ols()` defines the model but does not estimate its coefficients yet

Fitting a Model with `.fit()`

```
model = model_definition.fit()

print(type(model))
```

- `.fit()` estimates the intercept, slope, and other model statistics
- Store the fitted result so its attributes can be inspected

Method Chaining: Defining and Fitting in One Step

```
model = smf.ols(
    "score ~ study_hours",
    data = df
).fit()
```

- This is a common way to define and fit a model

Model Parameters with `.params`

```
print("Intercept:", round(model.params["Intercept"], 2))
print("study_hours:", round(model.params["study_hours"], 2))
```

```
Intercept: 50.43
study_hours: 2.75
```

- `Intercept`: predicted score when study hours equals zero
- `study_hours`: predicted score change for one additional study hour

Accessing One Parameter

```
intercept = model.params["Intercept"]
slope = model.params["study_hours"]

print(f"Intercept: {intercept:.2f}")
print(f"Slope: {slope:.2f}")
```

```
Intercept: 50.43
Slope: 2.75
```

- Use the parameter name to extract a specific estimate

Testing the Slope with .pvalues

```
slope_p_value = model.pvalues["study_hours"]

print(f"study_hours p-value: {slope_p_value:.3f}")
```

```
study_hours p-value: 0.000
```

Null hypothesis:

$$H_0 : \text{population slope} = 0$$

Confidence Intervals with `.conf_int()`

```
slope_ci = model.conf_int().loc["study_hours"]

print(f"Lower: {slope_ci[0]:.2f}")
print(f"Upper: {slope_ci[1]:.2f}")
```

```
Lower: 1.59
Upper: 3.90
```

- A confidence interval gives a range of values compatible with the model and data

Model Overview: `.summary().tables[0]`

```
summary = model.summary()
print(summary.tables[0])
```

```
                            OLS Regression Results
==============================================================================
Dep. Variable:                  score   R-squared:                       0.280
Model:                            OLS   Adj. R-squared:                  0.268
Method:                 Least Squares   F-statistic:                     22.57
Date:                Tue, 03 Nov 2026   Prob (F-statistic):           1.37e-05
Time:                        13:52:24   Log-Likelihood:                -210.99
No. Observations:                  60   AIC:                             426.0
Df Residuals:                      58   BIC:                             430.2
Df Model:                           1
Covariance Type:            nonrobust
==============================================================================
```

- Table 0 summarizes the model setup and overall fit
- Useful for checking the outcome, sample size, R^2, and model type

Model Overview: `.summary().tables[1]`

```
print(summary.tables[1])
```

```
==============================================================================
                 coef    std err          t      P>|t|      [0.025      0.975]
------------------------------------------------------------------------------
Intercept      50.4278      4.553     11.076      0.000      41.314      59.541
study_hours     2.7466      0.578      4.751      0.000       1.589       3.904
==============================================================================
```

- Table 1 gives the estimates for each model term
- Focus first on the coefficient, p-value, and confidence interval columns

Model Overview: `.summary().tables[2]`

```
print(summary.tables[2])
```

```
==============================================================================
Omnibus:                        0.940   Durbin-Watson:                   1.864
Prob(Omnibus):                  0.625   Jarque-Bera (JB):                0.843
Skew:                          -0.281   Prob(JB):                        0.656
Kurtosis:                       2.852   Cond. No.                         34.0
==============================================================================
```

- Table 2 contains diagnostic and assumption-related information
- Useful later when checking whether the model is behaving reasonably

.rsquared

- R^2 is the proportion of outcome variance described by the fitted model
- It ranges from 0 to 1
- A larger R^2 does not prove that the model is correct or causal

```
print(f"$R^2$: {model.rsquared:.3f}")
```

```
$R^2$: 0.280
```

Predicted Outcomes with .predict()

```
df["predicted_score"] = model.predict(df)

for i in range(3):
    print(
        f"Observed: {df['score'][i]:.2f}; "
        f"Predicted: {df['predicted_score'][i]:.2f}"
    )
```

```
Observed: 74.97; Predicted: 79.36
Observed: 68.62; Predicted: 69.08
Observed: 76.48; Predicted: 72.86
```


- .predict() calculates the fitted outcome for each row

Plotting Model Predictions: Observed and Predicted Values

```
ordered = df.sort_values("study_hours")

plt.scatter(df["study_hours"], df["score"], color = "blue")
plt.plot(
    ordered["study_hours"], ordered["predicted_score"], "k-"
)
plt.xlabel("Study Hours")
plt.ylabel("Score")
plt.show()
```

Fitted Line



- Blue circles are observed values
- The black line contains model-predicted values

Reporting Regression Results: Formatting the Result

```
slope_ci = model.conf_int().loc["study_hours"]

print(f"b = {model.params['study_hours']:.2f}, "
    f"95% CI [{slope_ci[0]:.2f}, {slope_ci[1]:.2f}], "
    f"$p$ = {model.pvalues['study_hours']:.3f}")
```

```
b = 2.75, 95% CI [1.59, 3.90], $p$ = 0.000
```

- Report the estimated slope, confidence interval, and p-value together
- Use **associated with**, rather than **caused**
- This matches the significance test for the Pearson correlation computed earlier — the slope's p-value is mathematically identical to the correlation's p-value

Multiple Regression

- Multiple regression extends simple regression to **more than one predictor**
- Predictors can be continuous or categorical — categorical predictors are automatically dummy-coded in formula notation
- Here we add `group` as a second predictor alongside `study_hours`

Research Question:

- Does study hours predict score, after accounting for group — and does the effect of study hours depend on group?

$$\text{score} = \text{intercept} + b_1(\text{study hours}) + b_2(\text{group}) + \text{error}$$

Multiple Regression: Level of Measurement

- `study_hours`: continuous (interval/ratio) — predictor
- `group`: categorical (nominal), two levels — predictor (dummy-coded)
- `score`: continuous (interval/ratio) — outcome


Multiple Regression: Inspecting the Data

```
control_rows = df[df["group"] == "control"]
treatment_rows = df[df["group"] == "treatment"]

plt.scatter(
    control_rows["study_hours"], control_rows["score"],
    color = "blue", label = "Control")
plt.scatter(
    treatment_rows["study_hours"], treatment_rows["score"],
    color = "red", label = "Treatment")
plt.xlabel("Study Hours")
plt.ylabel("Score")
plt.legend()
plt.show()
```

- Same approach as before: one `plt.scatter()` call per group, each with its own `color` and `label`, so `plt.legend()` can name the

Multiple Regression: Inspecting the Data



Multiple Regression: Formula Notation

```
"score ~ study_hours + group"
# Main effects only

"score ~ study_hours * group"
# Main effects AND their interaction; equivalent to:
"score ~ study_hours + group + study_hours:group"
```

- `+` includes each predictor's **main effect** only, holding the other predictor constant
- `*` includes the main effects **and** the interaction between them

Multiple Regression: Running the Test

```
main_effects_model = smf.ols(
    "score ~ study_hours + group", data = df
).fit()

interaction_model = smf.ols(
    "score ~ study_hours * group", data = df
).fit()
```

- Only the formula string changes between the two models

Multiple Regression: Interpreting the Results (Main Effects)

```
main_effects_model.summary()
```

```
Intercept: b = 49.68, 95% CI [40.91, 58.45], p = 0.000
group[T.treatment]: b = 5.09, 95% CI [0.91, 9.26], p = 0.018
study_hours: b = 2.51, 95% CI [1.38, 3.64], p = 0.000
```

Multiple Regression: Interpreting the Results: (Main Effects)

- Each coefficient is interpreted *holding the other predictor constant*

Multiple Regression: Interpreting the Results (Interaction Effects)

```
interaction_model.summary()
```

```
Intercept: b = 49.73, 95% CI [36.69, 62.77], p = 0.000
group[T.treatment]: b = 4.99, 95% CI [-
13.00, 22.98], p = 0.581
study_hours: b = 2.51, 95% CI [0.77, 4.24], p = 0.005
study_hours:group[T.treatment]: b = 0.01, 95% CI [-
2.29, 2.31], p = 0.991
```

Multiple Regression: Interpreting the Results (Interaction Effects)

- Switching + to * added exactly one new term: `study_hours:group[T.treatment]`
- It tests whether the `study_hours` slope differs between `control` and `treatment`
- Here it is far from significant and barely shifts the other coefficients — evidence the effect of `study_hours` does **not** depend on `group`

Comparing the Two Models: Running the Test

```
from statsmodels.stats.anova import anova_lm

comparison_table = anova_lm(
    main_effects_model, interaction_model
)
print(comparison_table)
```

```
   df_resid           ssr  df_diff  ss_diff         F    Pr(>F)
0      57.0  3604.470329      0.0      NaN       NaN       NaN
1      56.0  3604.462629      1.0   0.0077   0.00012  0.991312
```

Comparing the Two Models: Interpreting the Results

- Tests whether adding the interaction explains significantly more variance in `score` than the main-effects-only model
- Use this p-value, not the individual interaction term's p-value, to judge whether the interaction belongs in the model
- Here, p is far from significant — the interaction does not improve the model

Multiple Regression: Two Continuous Predictors

- Multiple regression also works when **all** predictors are continuous
- Here we add `sleep_hours` (hours slept the night before the exam) as a second predictor alongside `study_hours`

Research Question:

- Does study hours predict score, after accounting for sleep hours?

$$\text{score} = \text{intercept} + b_1(\text{study hours}) + b_2(\text{sleep hours}) + \text{error}$$

Two Continuous Predictors: Level of Measurement

- `study_hours`: continuous (interval/ratio) — predictor
- `sleep_hours`: continuous (interval/ratio) — predictor
- `score`: continuous (interval/ratio) — outcome
- Study design: observational/correlational, same as above

Adding a Third Variable

```
np.random.seed(21)

df["sleep_hours"] = (
    4
    + 0.06 * df["score"]
    - 0.25 * df["study_hours"]
    + np.random.normal(0, 0.6, len(df))
)
```


- Students who study more tend to sleep a little less

Two Continuous Predictors: Inspecting the Data

```
plt.scatter(
    df["study_hours"], df["score"],
    c = df["sleep_hours"],   # Colour each dot by sleep hours
    cmap = "viridis"
)
plt.colorbar(label = "Sleep Hours")
plt.xlabel("Study Hours")
plt.ylabel("Score")
plt.show()
```

- With a second continuous predictor, use `c` and `cmap` instead of one colour per group

Two Continuous Predictors: Inspecting the Data



- At any given number of study hours, lighter (more sleep) dots tend to sit higher

Two Continuous Predictors: Correlations Between Variables

```
print(
    df[["score", "study_hours", "sleep_hours"]].corr().round(2
)
```

```
              score  study_hours  sleep_hours
score          1.00         0.53         0.47
study_hours    0.53         1.00        -0.23
sleep_hours    0.47        -0.23         1.00
```

- Both predictors correlate positively with `score`
- The predictors are slightly **negatively** correlated with each other
- Very strong correlations between predictors (**multicollinearity**) make coefficients hard to interpret

Two Continuous Predictors: Formula Notation

```
"score ~ study_hours + sleep_hours"
# Main effects only

"score ~ study_hours * sleep_hours"
# Main effects AND their interaction
```

- The formula works exactly as it did with `group` — the predictor types do not change the syntax
- `*` would test whether the effect of study hours depends on sleep hours; we focus on the main-effects model here

Two Continuous Predictors: Running the Test

```
two_cont_model = smf.ols(
    "score ~ study_hours + sleep_hours", data = df
).fit()
```

- Same `smf.ols()` and `.fit()` steps as before; only the formula string changes

Two Continuous Predictors: Interpreting the Results

```
two_cont_model.summary()
```

```
Intercept: b = -1.94, 95% CI [-16.92, 13.05], p = 0.797
study_hours: b = 3.49, 95% CI [2.65, 4.33], p = 0.000
sleep_hours: b = 7.29, 95% CI [5.41, 9.18], p = 0.000
```

Two Continuous Predictors: Interpreting the Results

- `study_hours`: predicted score change for one additional study hour, *holding sleep hours constant*
- `sleep_hours`: predicted score change for one additional hour of sleep, *holding study hours constant*
- `Intercept`: predicted score at zero study hours **and** zero sleep hours — not meaningful here, since no one in the data slept zero hours

Comparing to Simple Regression

```
print(f"Simple: b = {model.params['study_hours']:.2f}, "
    f"R^2 = {model.rsquared:.3f}")
print(f"Multiple: b = {two_cont_model.params['study_hours']:.2
    f"R^2 = {two_cont_model.rsquared:.3f}")
```

```
Simple: b = 2.75, R^2 = 0.280
Multiple: b = 3.49, R^2 = 0.649
```

- The `study_hours` slope grows once sleep is held constant — heavy studiers sleep less, which had been masking part of the study effect
- R^2 increases: the two predictors together describe more variance in `score` than study hours alone