

# Deuring Correspondence and Public Key Cryptography

Oral Comprehensive Examination

---

Gaurish Korpai

December 07, 2023

The University of Arizona

# Table of contents

Appetizer: Discrete logarithm

Entrée: Supersingular isogeny graph


Dessert: Quantum-safe signature

## Appetizer: Discrete logarithm

---

# Public Key Cryptography

You are securely connected to this website. **Key exchange** allows two parties to agree on a common secret using only publicly exchanged information. **Digital signature** allows parties to authenticate themselves.



# Public Key Cryptography

*Let's Encrypt* is the world's largest certificate authority with over 2.53 billion certificates issued.



The screenshot shows a web browser window with the URL <https://letsencrypt.org/docs/glossary/>. The page features the Let's Encrypt logo and a navigation menu. A pink arrow points from the text on the left to the ECDSA entry. A pink oval highlights the text: "Let's Encrypt supports ECDSA for end-entity or leaf certificates, but not yet for the entire chain: /upcoming-features".

**ECDSA** (Elliptic Curve Digital Signature Algorithm) : A variant of the Digital Signature Algorithm (DSA) which uses elliptic curve cryptography. Wikipedia Let's Encrypt supports ECDSA for end-entity or leaf certificates, but not yet for the entire chain: /upcoming-features

**Ed25519** : A specific type of EdDSA, along with Ed448.

**EdDSA** (Edwards-curve Digital Signature Algorithm) : A modern public-key signature system based on elliptic curves, designed to solve several common implementation issues with elliptic curve cryptography. Certificate Authorities like Let's Encrypt can't provide EdDSA certificates yet. Wikipedia

**Elliptic Curve Cryptography** (ECC) : An type of public-key cryptography based on elliptic curves. ECC uses smaller keys compared to non-EC cryptography while providing equivalent security. Cloudflare - Wikipedia

# Public Key Cryptography

For 128-bit security,  
DSA (based on DLP)  
needs 4096-bit keys,  
but ECDSA (based on  
ECDLP) only needs  
256-bit key.



Glossary - Let's Encrypt X + Private browsing

https://letsencrypt.org/docs/glossary/

**Let's Encrypt**

ISRG celebrates 10 years of helping build a brighter Internet →

Glossary

**ECDSA** (Elliptic Curve Digital Signature Algorithm) : A variant of the Digital Signature Algorithm (DSA) which uses elliptic curve cryptography. [Wikipedia](#). Let's Encrypt supports ECDSA for [end-entity or leaf certificates](#), but not yet for the entire [chain](#): [/upcoming-features](#)


**Ed25519** : A specific type of [EdDSA](#), along with Ed448.

**EdDSA** (Edwards-curve Digital Signature Algorithm) : A modern public-key signature system based on elliptic curves, designed to solve several common [implementation issues](#) with elliptic curve cryptography. Certificate Authorities like [Let's Encrypt](#) can't provide EdDSA certificates yet. [Wikipedia](#)

**Elliptic Curve Cryptography** (ECC) : An type of public-key cryptography based on elliptic curves. [ECC uses smaller keys compared to non-EC cryptography while providing equivalent security.](#) [Cloudflare](#) - [Wikipedia](#)

# Public Key Cryptography

EdDSA is not ECDSA over a different curve. Rather, it is a *Schnorr signature* implemented for the Ed25519 Edwards curve.



Glossary - Let's Encrypt X + Private browsing

https://letsencrypt.org/docs/glossary/

**Let's Encrypt**

ISRG celebrates 10 years of helping build a brighter Internet →

Glossary

**ECDSA** (Elliptic Curve Digital Signature Algorithm) : A variant of the Digital Signature Algorithm (DSA) which uses elliptic curve cryptography. [Wikipedia](#). Let's Encrypt supports ECDSA for [end-entity or leaf certificates](#), but not yet for the entire [chain](#): [/upcoming-features](#)

**Ed25519** : A specific type of [EdDSA](#), along with Ed448.

**EdDSA** (Edwards-curve Digital Signature Algorithm) : A modern public-key signature system based on elliptic curves, designed to solve several common [implementation issues](#) with elliptic curve cryptography. Certificate Authorities like Let's Encrypt can't provide EdDSA certificates yet. [Wikipedia](#)

**Elliptic Curve Cryptography** (ECC) : An type of public-key cryptography based on elliptic curves. ECC uses smaller keys compared to non-EC cryptography while providing equivalent security. [Cloudflare](#) - [Wikipedia](#)

# Preparation for Schnorr

Acrobat Enterprise

Acrobat Desktop Digital Signatures

Search docs

Supported Standards

Signature algorithms

- Message digest creation algorithms
- Message digest encryption algorithms

Standards

APIs and miscellaneous

Enterprise Toolkit Console Help Forum Mobile

» Supported Standards

Previous

*Supported algorithms for creating the signature*

Product version	PDF version	Supported encryption algorithms
11.x and later	PDF 1.7	<ul style="list-style-type: none"><li>• RSA and DSA SHA1 up to 4096-bit</li><li>• ECDSA elliptic curve P256 with digest algorithm SHA256</li><li>• ECDSA elliptic curve P384 with digest algorithm SHA384</li><li>• ECDSA elliptic curve P512 with digest algorithm SHA512</li></ul>

Most of us have used Adobe Acrobat Sign to digitally sign PDF documents.



# Preparation for Schnorr

Acrobat Enterprise

Enterprise Toolkit Console Help Forum Mobile

Acrobat Desktop Digital Signatures

Search docs

Supported Standards

Signature algorithms

Message digest creation algorithms

Message digest encryption algorithms

Standards

APIs and miscellaneous

Supported Standards

Supported algorithms for creating the signature

Product version	PDF version	Supported encryption algorithms
11.x and later	PDF 1.7	<ul style="list-style-type: none"><li>• RSA and <b>DSA</b> SHA1 up to 4096-bit</li><li>• ECDSA elliptic curve P256 with digest algorithm SHA256</li><li>• ECDSA elliptic curve P384 with digest algorithm SHA384</li><li>• ECDSA elliptic curve P512 with digest algorithm SHA512</li></ul>

Let  $(\mathbb{G}, \cdot)$  be a finite abelian group of prime order  $\ell$ . The *discrete logarithm problem* (DLP) in  $\mathbb{G}$  is: given  $\langle g \rangle = \mathbb{G}$  and  $h \in \mathbb{G}$ , find an integer  $k \in \{0, \dots, \ell - 1\}$  such that  $g^k = h$ .

Number of operations for generic  $\mathbb{G}$  is  $\sqrt{\#\mathbb{G}}$ .

# Preparation for Schnorr

Acrobat Enterprise

Enterprise Toolkit Console Help Forum Mobile

Acrobat Desktop Digital Signatures

Search docs

Supported Standards

Signature algorithms

- Message digest creation algorithms
- Message digest encryption algorithms
- Standards
- APIs and miscellaneous

Supported Standards

Previous

Supported algorithms for creating the signature

Product version	PDF version	Supported encryption algorithms
11.x and later	PDF 1.7	<ul style="list-style-type: none"><li>RSA and DSA SHA1 up to 4096-bit</li><li>ECDSA elliptic curve P256 with digest algorithm SHA256</li><li>ECDSA elliptic curve P384 with digest algorithm SHA384</li><li>ECDSA elliptic curve P512 with digest algorithm SHA512</li></ul>

Let  $p$  be a prime larger than 3 and  $q = p^n$  for  $n > 0$ . An elliptic curve  $E$  over finite field  $\mathbb{F}_q$  can be written as  $E : y^2 = x^3 + ax + b$  where  $a, b \in \mathbb{F}_q$  and  $4a^3 + 27b^2 \neq 0$ , along with an extra point  $\mathcal{O}_E$ . Points on  $E$  form a group with  $\mathcal{O}_E$  as the neutral element.  $P$  be a point on  $E$  of prime order  $\ell$ , then  $\mathbb{G} = \langle P \rangle$  with the exponentiation replaced with scalar point multiplication, we get ECDLP.

# Preparation for Schnorr

Acrobat Enterprise

Enterprise Toolkit Console Help Forum Mobile

Acrobat Desktop Digital Signatures

Search docs

Supported Standards

Signature algorithms

Message digest creation algorithms

Message digest encryption algorithms

Standards

APIs and miscellaneous

» Supported Standards

Previous

Supported algorithms for creating the signature

Product version	PDF version	Supported encryption algorithms
11.x and later	PDF 1.7	<ul style="list-style-type: none"><li>• RSA and DSA SHA1 up to 4096-bit</li><li>• ECDSA elliptic curve P256 with digest algorithm <u>SHA256</u></li><li>• ECDSA elliptic curve P384 with digest algorithm <u>SHA384</u></li><li>• ECDSA elliptic curve P512 with digest algorithm <u>SHA512</u></li></ul>

A cryptographic hash function  $H$  takes arbitrary length bit strings as input and produces a fixed-length bit string as output, such that it is preimage resistant (can't find input of given output), second preimage resistant (can't find a different input leading to given output), and collision resistant (can't find two inputs with same output).

## Schnorr signature, Step 1: $\Sigma$ -protocol

Let  $\mathbb{G} = \langle g \rangle$  where  $g \in (\mathbb{Z}/p\mathbb{Z})^\times$  is an element of prime order  $\ell$ .

## Schnorr signature, Step 1: $\Sigma$ -protocol

Let  $\mathbb{G} = \langle g \rangle$  where  $g \in (\mathbb{Z}/p\mathbb{Z})^\times$  is an element of prime order  $\ell$ .  
The prover  $P$  randomly chooses (secret)  $k \in \{0, \dots, \ell - 1\}$  and publishes  $h = g^k \pmod{p}$ .

## Schnorr signature, Step 1: $\Sigma$ -protocol

Let  $\mathbb{G} = \langle g \rangle$  where  $g \in (\mathbb{Z}/p\mathbb{Z})^\times$  is an element of prime order  $\ell$ . The prover  $P$  randomly chooses (secret)  $k \in \{0, \dots, \ell - 1\}$  and publishes  $h = g^k \pmod{p}$ . Now,  $P$  can prove “knowledge” of a discrete logarithm  $k$  to a verifier  $V$ :

$P$  -----  $\mathbb{G}, p, h$  -----  $V$

$$t \xleftarrow{\$} \{0, \dots, \ell - 1\}$$
$$e \leftarrow g^t \pmod{p}$$

commitment:  $e$


$$r \leftarrow \{0, \dots, \ell - 1\}$$

$$s \leftarrow t + rk \pmod{\ell}$$

$$g^s \stackrel{?}{=} e h^r \pmod{p}$$


# Schnorr signature, Step 1: $\Sigma$ -protocol

Let  $\mathbb{G} = \langle g \rangle$  where  $g \in (\mathbb{Z}/p\mathbb{Z})^\times$  is an element of prime order  $\ell$ . The prover  $P$  randomly chooses (secret)  $k \in \{0, \dots, \ell - 1\}$  and publishes  $h = g^k \pmod{p}$ . Now,  $P$  can prove “knowledge” of a discrete logarithm  $k$  to a verifier  $V$ :



## Schnorr signature, Step 1: $\Sigma$ -protocol

Let  $\mathbb{G} = \langle g \rangle$  where  $g \in (\mathbb{Z}/p\mathbb{Z})^\times$  is an element of prime order  $\ell$ . The prover  $P$  randomly chooses (secret)  $k \in \{0, \dots, \ell - 1\}$  and publishes  $h = g^k \pmod{p}$ . Now,  $P$  can prove “knowledge” of a discrete logarithm  $k$  to a verifier  $V$ :





## Schnorr signature, Step 2: Fiat-Shamir transformation

Choose a *random oracle* cryptographic hash function  $H$  with appropriate domain and codomain.

## Schnorr signature, Step 2: Fiat-Shamir transformation

Choose a *random oracle* cryptographic hash function  $H$  with appropriate domain and codomain. The key generation algorithm  $G$  outputs a pair  $(k, h)$  such that  $h = g^k \pmod{p}$ , where  $k$  is the *secret signing key* and  $h$  is the *public verification key*.

## Schnorr signature, Step 2: Fiat-Shamir transformation

Choose a *random oracle* cryptographic hash function  $H$  with appropriate domain and codomain. The key generation algorithm  $G$  outputs a pair  $(k, h)$  such that  $h = g^k \pmod{p}$ , where  $k$  is the *secret signing key* and  $h$  is the *public verification key*.

### Signing $(\mathbb{G}, g, k, H, m)$

1.  $t \xleftarrow{\$} \{1, \dots, \ell - 1\}$
2.  $e \leftarrow g^t \pmod{p}$
3.  $r \leftarrow H(m \| e)$
4.  $s \leftarrow t + rk \pmod{\ell}$
5. return  $\sigma := (e, s)$

## Schnorr signature, Step 2: Fiat-Shamir transformation

Choose a *random oracle* cryptographic hash function  $H$  with appropriate domain and codomain. The key generation algorithm  $G$  outputs a pair  $(k, h)$  such that  $h = g^k \pmod{p}$ , where  $k$  is the *secret signing key* and  $h$  is the *public verification key*.

### Signing $(\mathbb{G}, g, k, H, m)$

1.  $t \xleftarrow{\$} \{1, \dots, \ell - 1\}$
2.  $e \leftarrow g^t \pmod{p}$
3.  $r \leftarrow H(m \| e)$
4.  $s \leftarrow t + rk \pmod{\ell}$
5. return  $\sigma := (e, s)$

### Verification $(\mathbb{G}, g, h, H, m, \sigma)$

1.  $r \leftarrow H(m \| e)$
2. return  $g^s \stackrel{?}{=} eh^r \pmod{p}$

On Dec 20, 2016,  
NIST initiated the  
process.

Neal Koblitiz | University of Washington  
Alfred Menezes | University of Waterloo

In August 2015, the NSA released a major policy statement on the need for postquantum cryptography. Certain peculiarities in its wording and timing have puzzled many people and given rise to speculation concerning the NSA, elliptic curve cryptography, and quantum-safe cryptography. Of the various theories that have been proposed, some seem more plausible than others, but a definitive explanation is elusive.

"It is a riddle wrapped in a mystery inside an enigma; but perhaps there is a key." —Winston Churchill, 1939  
(in reference to the Soviet Union)

In August 2015, the US government's NSA released a major policy statement on the need to develop standards for postquantum cryptography (PQC).<sup>1</sup> The NSA, like many other organizations, believes that the time is right to make a major push to design public-key cryptographic protocols whose security depends on hard problems that can't be solved efficiently by a quantum computer. Ever since Peter Shor's pioneering work more than 20 years ago,<sup>2</sup> it has been known that both the integer factorization problem, upon which RSA is based, and the elliptic curve discrete logarithm problem (ECDLP), upon which elliptic curve cryptography (ECC) is based, can be solved in polynomial time by a quantum computer.

The NSA announcement will give a tremendous boost to efforts to develop, standardize, and

commercialize quantum-safe cryptography. While standards for new postquantum algorithms are several years away, in the immediate future the NSA is encouraging vendors to add quantum resistance to existing protocols by means of conventional symmetric-key tools such as the Advanced Encryption Standard (AES). Given the NSA's strong interest in PQC, the demand for quantum-safe cryptographic solutions by governments and industry will likely grow dramatically in the coming years.

Most of the NSA statement was unexceptionable. However, one passage was puzzling and unexpected:<sup>1</sup>

For those partners and vendors that have not yet made the transition to Suite B algorithms, we recommend not making a significant expenditure to do so at this point but instead to prepare for the upcoming quantum resistant algorithm transition.... Unfortunately, the growth of elliptic curve use has bumped up against the fact of continued progress in the research on quantum computing, necessitating a re-evaluation of our cryptographic strategy.

In 1994, quantum algorithm for solving the DLP. Unfortunately, that is the hard-problem used by state-of-the-art digital signatures.

Neal Koblitz | University of Washington  
Alfred Menezes | University of Waterloo

In August 2015, the NSA released a major policy statement on the need for postquantum cryptography. Certain peculiarities in its wording and timing have puzzled many people and given rise to speculation concerning the NSA, elliptic curve cryptography, and quantum-safe cryptography. Of the various theories that have been proposed, some seem more plausible than others, but a definitive explanation is elusive.

"It is a riddle wrapped in a mystery inside an enigma; but perhaps there is a key." —Winston Churchill, 1939  
(in reference to the Soviet Union)

In August 2015, the US government's NSA released a major policy statement on the need to develop standards for postquantum cryptography (PQC).<sup>1</sup> The NSA, like many other organizations, believes that the time is right to make a major push to design public-key cryptographic protocols whose security depends on hard problems that can't be solved efficiently by a quantum computer. Ever since Peter Shor's pioneering work more than 20 years ago,<sup>2</sup> it has been known that both the integer factorization problem, upon which RSA is based, and the elliptic curve discrete logarithm problem (ECDLP), upon which elliptic curve cryptography (ECC) is based, can be solved in polynomial time by a quantum computer.

The NSA announcement will give a tremendous boost to efforts to develop, standardize, and

commercialize quantum-safe cryptography. While standards for new postquantum algorithms are several years away, in the immediate future the NSA is encouraging vendors to add quantum resistance to existing protocols by means of conventional symmetric-key tools such as the Advanced Encryption Standard (AES). Given the NSA's strong interest in PQC, the demand for quantum-safe cryptographic solutions by governments and industry will likely grow dramatically in the coming years.

Most of the NSA statement was unexceptionable. However, one passage was puzzling and unexpected:<sup>1</sup>

For those partners and vendors that have not yet made the transition to Suite B algorithms, we recommend not making a significant expenditure to do so at this point but instead to prepare for the upcoming quantum resistant algorithm transition.... Unfortunately, the growth of elliptic curve use has bumped up against the fact of continued progress in the research on quantum computing, necessitating a re-evaluation of our cryptographic strategy.

Entrée: Supersingular isogeny graph

---

# Supersingular elliptic curves

Recall,  $p$  is a prime larger than 3 and  $q = p^n$  for  $n > 0$ . For  $E/\mathbb{F}_q$  we have  $\#E(\mathbb{F}_q) = q + 1 - t$ , where  $|t| \leq 2\sqrt{q}$ .



# Supersingular elliptic curves

Recall,  $p$  is a prime larger than 3 and  $q = p^n$  for  $n > 0$ . For  $E/\mathbb{F}_q$  we have  $\#E(\mathbb{F}_q) = q + 1 - t$ , where  $|t| \leq 2\sqrt{q}$ .

## Supersingular elliptic curve

An elliptic curve over  $\mathbb{F}_q$  is called *supersingular* if  $p \mid t$ .

# Supersingular elliptic curves

Recall,  $p$  is a prime larger than 3 and  $q = p^n$  for  $n > 0$ . For  $E/\mathbb{F}_q$  we have  $\#E(\mathbb{F}_q) = q + 1 - t$ , where  $|t| \leq 2\sqrt{q}$ .

## Supersingular elliptic curve

An elliptic curve over  $\mathbb{F}_q$  is called *supersingular* if  $p \mid t$ .

## Example

$E_1 : y^2 = x^3 + x$  over  $\mathbb{F}_{23}$  is a supersingular elliptic curve because  $\#E(\mathbb{F}_{23}) = 24$  and  $t = 0$ .

## Isogeny

An isogeny between two elliptic curves  $E/\mathbb{F}_q$  and  $E'/\mathbb{F}_q$  is a non-constant rational function that maps points from  $E$  to points on  $E'$  and is compatible with the group law.

# Isogeny

## Isogeny

An isogeny between two elliptic curves  $E/\mathbb{F}_q$  and  $E'/\mathbb{F}_q$  is a non-constant rational function that maps points from  $E$  to points on  $E'$  and is compatible with the group law. In fact, an isogeny  $\phi : E \rightarrow E'$  exists iff  $\#E(\mathbb{F}_q) = \#E'(\mathbb{F}_q)$ .

# Isogeny

## Isogeny

An isogeny between two elliptic curves  $E/\mathbb{F}_q$  and  $E'/\mathbb{F}_q$  is a non-constant rational function that maps points from  $E$  to points on  $E'$  and is compatible with the group law. In fact, an isogeny  $\phi : E \rightarrow E'$  exists iff  $\#E(\mathbb{F}_q) = \#E'(\mathbb{F}_q)$ .

## Example

For  $E_1 : y^2 = x^3 + x$  and  $E_2 : y^2 = x^3 + 19x$  over  $\mathbb{F}_{23}$  we have

$$\phi : E_1 \rightarrow E_2$$

$$(x, y) \mapsto \left( \frac{x^2 + 1}{x}, \frac{x^2 y - y}{x^2} \right)$$

# Degree of isogeny

## Degree of (separable) isogeny

The degree of a (separable) isogeny  $\phi : E \rightarrow E'$  over  $\mathbb{F}_q$ , is the number of points on  $E$ , taken over any extension field of  $\mathbb{F}_q$ , mapping to  $\mathcal{O}_{E'}$ .

# Degree of isogeny

## Degree of (separable) isogeny

The degree of a (separable) isogeny  $\phi : E \rightarrow E'$  over  $\mathbb{F}_q$ , is the number of points on  $E$ , taken over any extension field of  $\mathbb{F}_q$ , mapping to  $\mathcal{O}_{E'}$ .

## Example

The degree of  $\phi : E_1 \rightarrow E_2$  defined above is 2, because  $(0, 0)$  and  $\mathcal{O}_{E_1}$  are the only two points mapping to  $\mathcal{O}_{E_2}$ .

# Degree of isogeny

## Degree of (separable) isogeny

The degree of a (separable) isogeny  $\phi : E \rightarrow E'$  over  $\mathbb{F}_q$ , is the number of points on  $E$ , taken over any extension field of  $\mathbb{F}_q$ , mapping to  $\mathcal{O}_{E'}$ .

## Example

The degree of  $\phi : E_1 \rightarrow E_2$  defined above is 2, because  $(0, 0)$  and  $\mathcal{O}_{E_1}$  are the only two points mapping to  $\mathcal{O}_{E_2}$ .

Degree is multiplicative:  $\deg(\phi \circ \psi) = \deg(\phi) \deg(\psi)$



# Isomorphism

## Isomorphism

An isogeny of degree 1 is called an isomorphism.

# Isomorphism

## Isomorphism

An isogeny of degree 1 is called an isomorphism. That is, two elliptic curves  $E/\mathbb{F}_q$  and  $E'/\mathbb{F}_q$  are isomorphic over  $\mathbb{F}_q$  if there exists a polynomial map over  $\mathbb{F}_q$  that maps points on  $E$  to points on  $E'$  in a one-to-one way which is compatible with the group operation.

# Isomorphism

## Isomorphism

An isogeny of degree 1 is called an isomorphism. That is, two elliptic curves  $E/\mathbb{F}_q$  and  $E'/\mathbb{F}_q$  are isomorphic over  $\mathbb{F}_q$  if there exists a polynomial map over  $\mathbb{F}_q$  that maps points on  $E$  to points on  $E'$  in a one-to-one way which is compatible with the group operation.

## Example

For  $E_1 : y^2 = x^3 + x$  and  $E_3 : y^2 = x^3 + 2x$  over  $\mathbb{F}_{23}$  we have

$$\tau : E_1 \rightarrow E_3$$

$$(x, y) \mapsto (-5x, -6y)$$

# Isomorphism class label

## $j$ -invariant

The  $j$ -invariant uniquely describes isomorphism classes over an algebraic closure of  $\mathbb{F}_q$ .

# Isomorphism class label

## $j$ -invariant

The  $j$ -invariant uniquely describes isomorphism classes over an algebraic closure of  $\mathbb{F}_q$ . For  $E : y^2 = x^3 + ax + b$  we have

$$j(E) = 1728 \frac{4a^3}{(4a^3 + 27b^2)}$$

# Isomorphism class label

## $j$ -invariant

The  $j$ -invariant uniquely describes isomorphism classes over an algebraic closure of  $\mathbb{F}_q$ . For  $E : y^2 = x^3 + ax + b$  we have

$$j(E) = 1728 \frac{4a^3}{(4a^3 + 27b^2)}$$

Note that two curves having the same  $j$ -invariant need not be isomorphic over  $\mathbb{F}_q$ .

# Isomorphism class label

## $j$ -invariant

The  $j$ -invariant uniquely describes isomorphism classes over an algebraic closure of  $\mathbb{F}_q$ . For  $E : y^2 = x^3 + ax + b$  we have

$$j(E) = 1728 \frac{4a^3}{(4a^3 + 27b^2)}$$

Note that two curves having the same  $j$ -invariant need not be isomorphic over  $\mathbb{F}_q$ .

## Example

We have  $j(E_1) = j(E_2) = j(E_3) = 1728 \pmod{23} = 3$ . But  $E_1$  and  $E_2$  are not isomorphic over  $\mathbb{F}_{23}$ .

# Isomorphism class label

## $j$ -invariant

The  $j$ -invariant uniquely describes isomorphism classes over an algebraic closure of  $\mathbb{F}_q$ . For  $E : y^2 = x^3 + ax + b$  we have

$$j(E) = 1728 \frac{4a^3}{(4a^3 + 27b^2)}$$

Note that two curves having the same  $j$ -invariant need not be isomorphic over  $\mathbb{F}_q$ .

## Example

We have  $j(E_1) = j(E_2) = j(E_3) = 1728 \pmod{23} = 3$ . But  $E_1$  and  $E_2$  are not isomorphic over  $\mathbb{F}_{23}$ .

If  $E$  is supersingular, then we can replace “algebraic closure of  $\mathbb{F}_q$ ” with  $\mathbb{F}_{p^2}$ .



# Supersingular isomorphism classes

The number of supersingular isomorphism classes over an algebraic closure of  $\mathbb{F}_p$ , with representative curves defined over  $\mathbb{F}_{p^2}$ , is  $S_p := \left\lfloor \frac{p}{12} \right\rfloor + \epsilon$  where  $\epsilon \in \{0, 1, 2\}$ .

# Supersingular isomorphism classes

The number of supersingular isomorphism classes over an algebraic closure of  $\mathbb{F}_p$ , with representative curves defined over  $\mathbb{F}_{p^2}$ , is  $S_p := \left\lfloor \frac{p}{12} \right\rfloor + \epsilon$  where  $\epsilon \in \{0, 1, 2\}$ .

## Example

$S_{23} = 3$  with the classes represented by the  $j$ -invariants 0, 3, 19.

## Supersingular $\ell$ -isogeny graph

Let  $\ell$  be a prime different from  $p$ . The supersingular  $\ell$ -isogeny graph over an algebraic closure of  $\mathbb{F}_q$  is the directed multigraph  $G_\ell(p)$  whose vertices belong to the set of isomorphism classes of supersingular elliptic curves  $\{j(E_1), \dots, j(E_s)\}$  with  $s = S_p$  and  $E_i/\mathbb{F}_{p^2}$ ; there is a directed edge  $[E_i, E_{i'}]$  for each equivalence class (same kernel) of  $\ell$ -isogenies from  $E_i$  to  $E_{i'}$ .


# Supersingular $\ell$ -isogeny graph

Let  $\ell$  be a prime different from  $p$ . The supersingular  $\ell$ -isogeny graph over an algebraic closure of  $\mathbb{F}_q$  is the directed multigraph  $G_\ell(p)$  whose vertices belong to the set of isomorphism classes of supersingular elliptic curves  $\{j(E_1), \dots, j(E_s)\}$  with  $s = S_p$  and  $E_i/\mathbb{F}_{p^2}$ ; there is a directed edge  $[E_i, E_{i'}]$  for each equivalence class (same kernel) of  $\ell$ -isogenies from  $E_i$  to  $E_{i'}$ . Hence the out-degree of each vertex is  $\ell + 1$ .

# Supersingular $\ell$ -isogeny graph

Let  $\ell$  be a prime different from  $p$ . The supersingular  $\ell$ -isogeny graph over an algebraic closure of  $\mathbb{F}_q$  is the directed multigraph  $G_\ell(p)$  whose vertices belong to the set of isomorphism classes of supersingular elliptic curves  $\{j(E_1), \dots, j(E_s)\}$  with  $s = S_p$  and  $E_i/\mathbb{F}_{p^2}$ ; there is a directed edge  $[E_i, E_{i'}]$  for each equivalence class (same kernel) of  $\ell$ -isogenies from  $E_i$  to  $E_{i'}$ . Hence the out-degree of each vertex is  $\ell + 1$ .

Example:  $G_2(23)$



# Endomorphism ring

## Endomorphism ring

The endomorphism ring of  $E$ ,  $\text{End}_{\mathbb{F}_q}(E)$ , is the set of  $\mathbb{F}_q$ -isogenies from  $E$  to itself, together with the zero map  $[0] : E \rightarrow E$  given  $[0](P) = \mathcal{O}_E$ .

# Endomorphism ring

## Endomorphism ring

The endomorphism ring of  $E$ ,  $\text{End}_{\mathbb{F}_q}(E)$ , is the set of  $\mathbb{F}_q$ -isogenies from  $E$  to itself, together with the zero map  $[0] : E \rightarrow E$  given  $[0](P) = \mathcal{O}_E$ . In particular, for isogenies over an algebraic closure of  $\mathbb{F}_q$ , we write  $\text{End}(E)$ .

# Endomorphism ring

## Endomorphism ring

The endomorphism ring of  $E$ ,  $\text{End}_{\mathbb{F}_q}(E)$ , is the set of  $\mathbb{F}_q$ -isogenies from  $E$  to itself, together with the zero map  $[0] : E \rightarrow E$  given  $[0](P) = \mathcal{O}_E$ . In particular, for isogenies over an algebraic closure of  $\mathbb{F}_q$ , we write  $\text{End}(E)$ .

## Example

For  $E_1 : y^2 = x^3 + x$  over  $\mathbb{F}_{23}$ , we have

$$\text{End}(E_1) = \mathbb{Z} \text{id} + \mathbb{Z} \iota + \mathbb{Z} \frac{\iota + \pi}{2} + \mathbb{Z} \frac{\text{id} + \iota \circ \pi}{2}$$

where  $\pi, \iota \in \text{End}(E_1)$  such that  $\pi(x, y) = (x^{23}, y^{23})$  and  $\iota(x, y) = (-x, \alpha y)$  is an isomorphism over  $\mathbb{F}_{23^2} = \mathbb{F}_{23}(\alpha)$  with  $\alpha^2 + 1 = 0$ .



# Endomorphism ring

## Endomorphism ring

The endomorphism ring of  $E$ ,  $\text{End}_{\mathbb{F}_q}(E)$ , is the set of  $\mathbb{F}_q$ -isogenies from  $E$  to itself, together with the zero map  $[0] : E \rightarrow E$  given  $[0](P) = \mathcal{O}_E$ . In particular, for isogenies over an algebraic closure of  $\mathbb{F}_q$ , we write  $\text{End}(E)$ .

## Example

For  $E_1 : y^2 = x^3 + x$  over  $\mathbb{F}_{23}$ , we have

$$\text{End}(E_1) = \mathbb{Z} \text{id} + \mathbb{Z} \iota + \mathbb{Z} \frac{\iota + \pi}{2} + \mathbb{Z} \frac{\text{id} + \iota \circ \pi}{2}$$

where  $\pi, \iota \in \text{End}(E_1)$  such that  $\pi(x, y) = (x^{23}, y^{23})$  and  $\iota(x, y) = (-x, \alpha y)$  is an isomorphism over  $\mathbb{F}_{23^2} = \mathbb{F}_{23}(\alpha)$  with  $\alpha^2 + 1 = 0$ . Moreover,  $\iota \circ \iota = [-1]$ ,  $\pi \circ \pi = [-23]$ , and  $\iota \circ \pi = -\pi \circ \iota$ ; i.e.  $\text{End}(E_1)$  is a non-commutative ring.

## Deuring correspondence - I

$E$  is a supersingular elliptic curve over  $\mathbb{F}_q$  if and only if  $\text{End}(E)$  is isomorphic to a maximal order in the quaternion algebra  $B_{p,\infty}$ .

# Maximal orders

## Deuring correspondence - I

$E$  is a supersingular elliptic curve over  $\mathbb{F}_q$  if and only if  $\text{End}(E)$  is isomorphic to a maximal order in the quaternion algebra  $B_{p,\infty}$ .

## Quaternion algebra

A quaternion algebra over  $\mathbb{Q}$  is of the form  $\mathbb{Q}\langle i, j \rangle = \mathbb{Q} + \mathbb{Q}i + \mathbb{Q}j + \mathbb{Q}ij$ , where  $i^2, j^2 \in \mathbb{Q}^\times$ , and  $ij = -ji$ . In particular, we have

$$B_{p,\infty} = \begin{cases} i^2 = -1, j^2 = -1 & \text{if } p = 2 \\ i^2 = -1, j^2 = -p & \text{if } p \equiv 3 \pmod{4} \\ i^2 = -2, j^2 = -p & \text{if } p \equiv 5 \pmod{8} \\ i^2 = -\ell, j^2 = -p & \text{if } p \equiv 1 \pmod{8} \end{cases}$$

where  $\ell \equiv 3 \pmod{4}$  is a prime quadratic non-residue mod  $p$ .

# Maximal orders

## Deuring correspondence - I

$E$  is a supersingular elliptic curve over  $\mathbb{F}_q$  if and only if  $\text{End}(E)$  is isomorphic to a maximal order in the quaternion algebra  $B_{p,\infty}$ .

## Quaternion (maximal) order

$O \subseteq \mathbb{Q}\langle i, j \rangle$  is called an *order* if  $O$  is a ring whose elements are integral,  $\mathbb{Z} \subseteq O$ , and contains a basis for  $\mathbb{Q}\langle i, j \rangle$  as  $\mathbb{Q}$ -vector space. Moreover, an order  $O \subsetneq B$  is called *maximal* if it is not properly contained in another order.

# Maximal orders

## Deuring correspondence - I

$E$  is a supersingular elliptic curve over  $\mathbb{F}_q$  if and only if  $\text{End}(E)$  is isomorphic to a maximal order in the quaternion algebra  $B_{p,\infty}$ .

## Example

In  $B_{23,\infty} = \langle i, j \mid i^2 = -1, j^2 = -23, ij = -ji \rangle$ , two examples of maximal orders are

$$O_1 = \mathbb{Z} + \mathbb{Z}i + \mathbb{Z}\frac{i+j}{2} + \mathbb{Z}\frac{1+ij}{2}; \text{ and}$$

$$O_2 = \mathbb{Z} + \mathbb{Z}i + \mathbb{Z}\frac{1+j}{2} + \mathbb{Z}\frac{i(1+j)}{2}$$

Note that  $O_1$  is isomorphic to  $\text{End}(E_1)$  we saw above.

# Maximal orders

## Deuring correspondence - I

$E$  is a supersingular elliptic curve over  $\mathbb{F}_q$  if and only if  $\text{End}(E)$  is isomorphic to a maximal order in the quaternion algebra  $B_{p,\infty}$ .

## Mestre-Oesterle-Ribet

$$\left\{ \begin{array}{l} \text{isomorphism classes} \\ \text{of supersingular} \\ \text{elliptic curves over } \overline{\mathbb{F}}_p \end{array} \right\} / \text{Gal}(\overline{\mathbb{F}}_p / \mathbb{F}_p) \longleftrightarrow \left\{ \begin{array}{l} \text{maximal orders} \\ \text{of } B_{p,\infty} \end{array} \right\} / \cong$$

That is, there is one-to-one correspondence if  $j(E) \in \mathbb{F}_p$  and two-to-one correspondence if  $j(E) \in \mathbb{F}_{p^2} \setminus \mathbb{F}_p$ .

## Deuring correspondence - II

Fix,  $E$ , a supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  with  $\text{End}(E) \cong O \subseteq B_{p,\infty}$ . There is a bijection between isomorphism classes over  $\overline{\mathbb{F}}_p$  and the left class set  $\text{Cls}_L(O)$ .

## Deuring correspondence - II

Fix,  $E$ , a supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  with  $\text{End}(E) \cong O \subseteq B_{p,\infty}$ . There is a bijection between isomorphism classes over  $\overline{\mathbb{F}}_p$  and the left class set  $\text{Cls}_L(O)$ .

## Quaternion (left) $O$ -ideal

$I \subseteq \mathbb{Q}\langle i, j \rangle$  is called an *ideal* if  $I$  is a  $\mathbb{Z}$ -module that contains a basis for  $\mathbb{Q}\langle i, j \rangle$  as  $\mathbb{Q}$ -vector space. Furthermore, given an order  $O$  of  $\mathbb{Q}\langle i, j \rangle$ ,  $I$  is called a *left  $O$ -ideal* if  $\alpha I \subseteq I$  for all  $\alpha \in O$ .



## Deuring correspondence - II

Fix,  $E$ , a supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  with  $\text{End}(E) \cong O \subseteq B_{p,\infty}$ . There is a bijection between isomorphism classes over  $\overline{\mathbb{F}}_p$  and the left class set  $\text{Cls}_L(O)$ .

## Left-ideal class set

We say ideals  $I, J$  are in the *same left class*,  $I \sim_L J$ , if there exists  $\alpha \in B^\times$  such that  $I\alpha = J$ . Furthermore, the left equivalence class is denoted by  $[I]_L$ . In particular, we have

$$\text{Cls}_L(O) := \{[I]_L \mid I \text{ is an invertible left } O\text{-ideal}\}$$

$\text{Cls}_L(O)$  has the structure of a *pointed set* with distinguished element  $[O]_L \in \text{Cls}_L(O)$ .

## Deuring correspondence - II

Fix,  $E$ , a supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  with  $\text{End}(E) \cong O \subseteq B_{p,\infty}$ . There is a bijection between isomorphism classes over  $\overline{\mathbb{F}}_p$  and the left class set  $\text{Cls}_L(O)$ .

### Example

Let  $O_2 \subset B_{23,\infty}$  as above. Then we have  $\text{Cls}_L(O_2) = \{[l_1]_L, [l_2]_L, [l_3]_L\}$  with

$$l_1 = 2\mathbb{Z}(1 + j) + 2\mathbb{Z}i(1 + j) + 4\mathbb{Z}j + 4\mathbb{Z}ij,$$

$$l_2 = 2\mathbb{Z}(1 + 3j) + 2\mathbb{Z}i(1 + 3j) + 8\mathbb{Z}j + 8\mathbb{Z}ij, \text{ and}$$

$$l_3 = 2\mathbb{Z}(1 + 3j + 4ij) + 2\mathbb{Z}(i + 4j + 3ij) + 16\mathbb{Z}j + 16\mathbb{Z}ij$$

Here  $[l_1]_L$ ,  $[l_2]_L$ , and  $[l_3]_L$  correspond to the isomorphism classes of supersingular curves represented by  $j$ -invariants 3, 19, and 0, respectively.

## Deuring correspondence - II

Fix,  $E$ , a supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  with  $\text{End}(E) \cong \mathcal{O} \subseteq B_{p,\infty}$ . There is a bijection between isomorphism classes over  $\overline{\mathbb{F}}_p$  and the left class set  $\text{Cls}_L(\mathcal{O})$ .

## Waterhouse

$E[I] := \{P \in E(\overline{\mathbb{F}}_p) \mid \phi(P) = 0 \ \forall \text{ separable } \phi \in I\}$ , where  $I$  is a nonzero left  $\text{End}(E)$ -ideal.  $\phi_I : E \rightarrow E/E[I]$  with  $\deg(\phi_I) = \#E[I]$ .

$I(H) := \{\phi \in \text{End}(E) \mid \phi(P) = 0 \text{ for all } P \in H\}$ , where  $H \leq E(\overline{\mathbb{F}}_p)$  is finite. If  $\phi : E \rightarrow E'$  an isogeny, then  $I_\phi := I(\ker(\phi))$  a left  $\text{End}(E)$ -ideal and right  $\text{End}(E')$ -ideal (connecting ideal).

## Deuring correspondence - II

Fix,  $E$ , a supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  with  $\text{End}(E) \cong O \subseteq B_{p,\infty}$ . There is a bijection between isomorphism classes over  $\overline{\mathbb{F}}_p$  and the left class set  $\text{Cls}_L(O)$ .

## Waterhouse

- $E[I(H)] = H$  and  $I(E[I]) = I$  (overloaded notation).
- If  $I \sim_L J$  then  $E/E[I] \cong E/E[J]$ .
- $\phi_{I \cdot J} = \tau_J \circ \phi_I$  and  $I_{\tau \circ \phi} = I_\phi \cdot I_\tau$
- $\phi_{\bar{I}} = \widehat{\phi_I}$  (dual isogeny) and  $I_{\widehat{\phi}} = \overline{I_\phi}$
- $\deg(\phi_I) = \text{nrd}(I)$  and  $\text{nrd}(I_\phi) = \deg(\phi)$

## Deuring correspondence - III

$(E, C)$  is a pair of supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  and cyclic subgroup of order  $M$  with  $\gcd(p, M) = 1$  iff  $\text{End}(E, C) \cong O(M) \subseteq B_{p, \infty}$  an Eichler order of level  $M$ .

## Deuring correspondence - III

$(E, C)$  is a pair of supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  and cyclic subgroup of order  $M$  with  $\gcd(p, M) = 1$  iff  $\text{End}(E, C) \cong O(M) \subseteq B_{p, \infty}$  an Eichler order of level  $M$ .

## Eichler order

An *Eichler order*  $O \subset B$  is the intersection of two (not necessarily distinct) maximal orders. Therefore, maximal orders are also Eichler orders.

## Deuring correspondence - III

$(E, C)$  is a pair of supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  and cyclic subgroup of order  $M$  with  $\gcd(p, M) = 1$  iff  $\text{End}(E, C) \cong O(M) \subseteq B_{p, \infty}$  an Eichler order of level  $M$ .

## Level of an Eichler order

The level of an Eichler order  $O$ , is defined as the ratio of the reduced discriminant of order  $O$  and the discriminant of the quaternion algebra  $B = \mathbb{Q}\langle i, j \rangle$ .

$$\text{lev}(O) = \frac{\text{discrd}(O)}{\text{disc}(B)}$$

From the definition of (reduced) discriminants it follows that maximal orders are Eichler orders of level 1.

## Deuring correspondence - III

$(E, C)$  is a pair of supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  and cyclic subgroup of order  $M$  with  $\gcd(p, M) = 1$  iff  $\text{End}(E, C) \cong O(M) \subseteq B_{p, \infty}$  an Eichler order of level  $M$ .

## Example

$O_3 = O_1 \cap O_2 = \mathbb{Z} + \mathbb{Z}\mathbf{i} + \mathbb{Z}\mathbf{j} + \mathbb{Z}\frac{1 + \mathbf{i} + \mathbf{j} + \mathbf{ij}}{2}$  is an Eichler order of level 2, because  $\text{discrd}(O_3) = 46$  and  $\text{disc}(B_{23, \infty}) = 23$ . Therefore, if  $\phi \in \text{End}(E, C) \cong O_3$  then  $\phi \in \text{End}(E)$  such that  $\phi(C) = C$  with  $\#C = 2$ .



## Deuring correspondence - III

$(E, C)$  is a pair of supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  and cyclic subgroup of order  $M$  with  $\gcd(p, M) = 1$  iff  $\text{End}(E, C) \cong O(M) \subseteq B_{p, \infty}$  an Eichler order of level  $M$ .

## Kohel

Fix a base point  $(E_0, C_0)$ , where  $C_0 \leq E(\overline{\mathbb{F}}_p)$  is a cyclic subgroup of order  $M$ . Then  $\text{End}(E_0, C_0)$ , the subring of  $\text{End}(E_0)$  that maps  $C_0$  to itself, is an Eichler order of level  $M$  and reduced discriminant  $pM$ .

## Deuring correspondence - III

$(E, C)$  is a pair of supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  and cyclic subgroup of order  $M$  with  $\gcd(p, M) = 1$  iff  $\text{End}(E, C) \cong O(M) \subseteq B_{p, \infty}$  an Eichler order of level  $M$ .

## Kohel

Let  $\mathcal{S}_M$  be the category of supersingular elliptic curves over  $\overline{\mathbb{F}}_p$  equipped with a cyclic  $M$ -isogeny (under isogenies identifying the cyclic subgroups).

Let  $\mathcal{I}_M$  be the category of left  $\text{End}(E_0, C_0)$ -ideals (under module homomorphisms).

Then the functor  $\text{Hom}(-, (E_0, C_0))$  from  $\mathcal{S}_M$  to  $\mathcal{I}_M$  is an equivalence of categories.

## Spectral graph theory

Deuring correspondence lets us use the relationship between quaternion algebras and modular forms to study the eigenvalues of the adjacency matrix of  $G_\ell(p)$ .


Deuring correspondence lets us use the relationship between quaternion algebras and modular forms to study the eigenvalues of the adjacency matrix of  $G_\ell(p)$ .

1.  $G_\ell(p)$  is connected with diameter  $O(\log p)$ , where the constant in the bound is independent of  $\ell$ . That is, the largest number of vertices which must be traversed in order to travel from one vertex to another when paths which backtrack, detour, or loop are excluded from consideration is  $O(\log p)$ .

Deuring correspondence lets us use the relationship between quaternion algebras and modular forms to study the eigenvalues of the adjacency matrix of  $G_\ell(p)$ .

1.  $G_\ell(p)$  is connected with diameter  $O(\log p)$ , where the constant in the bound is independent of  $\ell$ . That is, the largest number of vertices which must be traversed in order to travel from one vertex to another when paths which backtrack, detour, or loop are excluded from consideration is  $O(\log p)$ .
2.  $G_\ell(p)$  is an *expander graph*, i.e. simultaneously sparse and highly connected. Therefore, the natural random walk on  $G_\ell(p)$  converges to its limiting distribution as rapidly as possible.

$G_2(431)$  with 37 vertices and diameter 7



# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .

## Easier problems

# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .

## Easier problems

1. Given maximal order  $O \subseteq B_{p,\infty}$ , find a supersingular  $j$ -invariant such that  $\text{End}(E(j)) \cong O$ .



# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .

## Easier problems

1. Given maximal order  $O \subseteq B_{p,\infty}$ , find a supersingular  $j$ -invariant such that  $\text{End}(E(j)) \cong O$ .

# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .

## Easier problems

1. Given maximal order  $O \subseteq B_{p,\infty}$ , find a supersingular  $j$ -invariant such that  $\text{End}(E(j)) \cong O$ .
2. Find all the maximal orders (up to isomorphism) of  $B_{p,\infty}$ .

# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .
3. Given  $j, j' \in \mathbb{F}_{p^2}$  find an isogeny  $\phi : E \rightarrow E'$  such that  $j(E) = j$  and  $j(E') = j'$ .

## Easier problems

1. Given maximal order  $O \subseteq B_{p,\infty}$ , find a supersingular  $j$ -invariant such that  $\text{End}(E(j)) \cong O$ .
2. Find all the maximal orders (up to isomorphism) of  $B_{p,\infty}$ .

# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .
3. Given  $j, j' \in \mathbb{F}_{p^2}$  find an isogeny  $\phi : E \rightarrow E'$  such that  $j(E) = j$  and  $j(E') = j'$ .

## Easier problems

1. Given maximal order  $O \subseteq B_{p,\infty}$ , find a supersingular  $j$ -invariant such that  $\text{End}(E(j)) \cong O$ .
2. Find all the maximal orders (up to isomorphism) of  $B_{p,\infty}$ .
3. Given maximal orders  $O, O' \subseteq B_{p,\infty}$  find an ideal  $I$  that is left  $O$ -ideal and right  $O'$ -ideal.

# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .
3. Given  $j, j' \in \mathbb{F}_{p^2}$  find an isogeny  $\phi : E \rightarrow E'$  such that  $j(E) = j$  and  $j(E') = j'$ .
4. Given a maximal order in  $B_{p,\infty}$ , determine the ideal class set.

## Easier problems

1. Given maximal order  $O \subseteq B_{p,\infty}$ , find a supersingular  $j$ -invariant such that  $\text{End}(E(j)) \cong O$ .
2. Find all the maximal orders (up to isomorphism) of  $B_{p,\infty}$ .
3. Given maximal orders  $O, O' \subseteq B_{p,\infty}$  find an ideal  $I$  that is left  $O$ -ideal and right  $O'$ -ideal.

# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .
3. Given  $j, j' \in \mathbb{F}_{p^2}$  find an isogeny  $\phi : E \rightarrow E'$  such that  $j(E) = j$  and  $j(E') = j'$ .
4. Given a maximal order in  $B_{p,\infty}$ , determine the ideal class set.

## Easier problems

1. Given maximal order  $O \subseteq B_{p,\infty}$ , find a supersingular  $j$ -invariant such that  $\text{End}(E(j)) \cong O$ .
2. Find all the maximal orders (up to isomorphism) of  $B_{p,\infty}$ .
3. Given maximal orders  $O, O' \subseteq B_{p,\infty}$  find an ideal  $I$  that is left  $O$ -ideal and right  $O'$ -ideal.
4. Given  $p$ , determine (all) supersingular  $j$ -invariants in  $\mathbb{F}_{p^2}$ .

# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .
3. Given  $j, j' \in \mathbb{F}_{p^2}$  find an isogeny  $\phi : E \rightarrow E'$  such that  $j(E) = j$  and  $j(E') = j'$ .

## Equivalent and quantum-safe

- All three problems are known to be equivalent.

# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .
3. Given  $j, j' \in \mathbb{F}_{p^2}$  find an isogeny  $\phi : E \rightarrow E'$  such that  $j(E) = j$  and  $j(E') = j'$ .

## Equivalent and quantum-safe

- All three problems are known to be equivalent.
- The fact that  $\text{End}(E)$  is non-commutative makes these problems resistant to known quantum algorithms.



# Problems about supersingular elliptic curves

## Difficult problems

1. Given  $E/\mathbb{F}_{p^2}$ , find a maximal order  $O \subseteq B_{p,\infty}$  such that  $O \cong \text{End}(E)$ .
2. Given  $E/\mathbb{F}_{p^2}$ , chosen uniformly at random, determine  $\text{End}(E)$ .
3. Given  $j, j' \in \mathbb{F}_{p^2}$  find an isogeny  $\phi : E \rightarrow E'$  such that  $j(E) = j$  and  $j(E') = j'$ .

## Equivalent and quantum-safe

- All three problems are known to be equivalent.
- The fact that  $\text{End}(E)$  is non-commutative makes these problems resistant to known quantum algorithms.
- We can rewrite these problems in terms of cyclic  $M$ -isogenies and Eichler orders of level  $M$ . For SQIsign, we assume that given  $E/\mathbb{F}_{p^2}$  it is *difficult* to find a (non-trivial) cyclic endomorphism of  $E$  of smooth degree.

## Dessert: Quantum-safe signature

---

In 2022, NIST selected two lattice-based signatures (*CRYSTALS-Dilithium* and *FALCON*) and one hash-based signature (*SPHINCS+*)

The screenshot shows the NIST Computer Security Resource Center website. The page title is "Post-Quantum Cryptography: Digital Signature Schemes". Below the title, there are links for "PROJECTS" and "PQC DIGITAL SIGNATURE SCHEMES". The "Overview" link is highlighted with a red circle, and a red arrow points from the text "CRYSTALS-Dilithium and FALCON" to it. The page also features a "Round 1 Additional Signatures" section with a table of submitters and comments.

## Post-Quantum Cryptography: Digital Signature Schemes

PROJECTS PQC DIGITAL SIGNATURE SCHEMES

Overview News & Updates

### Round 1 Additional Signatures


Official comments on the **First Round Signatures** should be submitted using the "Submit Comment" link for the appropriate algorithm. Comments from the [pqc-forum Google group subscribers](#) will also be forwarded to the [pqc-forum Google group list](#). We will periodically post and update the comments received to the appropriate algorithm.

All relevant comments will be posted in their entirety and should not include PII information in the body of the email message.

#### Isogeny Signatures

Algorithm	Algorithm Information	Submitters	Comments
SQIsign	<a href="#">Specification</a> <a href="#">JP Statements</a> <a href="#">Zip file (14 MB)</a> <a href="#">Website</a>	Jorge Chavez-Saab Maria Corte-Real Santos Luca De Feo Jonathan Komada Eriksen Basil Hess David Kohel Antonin Leroux Patrick Longa Michael Meyer Lorenz Panny Sikhar Patranabis Christophe Petit Francisco Rodríguez Henríquez Sina Schaeffler Benjamin Wesolowski	<a href="#">Submit Comment</a> <a href="#">View Comments</a>

Signature schemes  
with short signatures  
and fast verification;  
not based on struc-  
tured lattices.



**NIST**  
Information Technology Laboratory  
**COMPUTER SECURITY RESOURCE CENTER**

PROJECTS PQC DIGITAL SIGNATURE SCHEMES

## Post-Quantum Cryptography: Digital Signature Schemes

f t

PROJECT LINKS

- Overview
- News & Updates

### Round 1 Additional Signatures

Official comments on the **Round 1 Signatures** should be submitted using the "Submit Comment" link for the appropriate algorithm. Comments from the [pqc-forum Google group subscribers](#) will also be forwarded to the [pqc-forum Google group](#) list. We will periodically post and update the comments received to the appropriate algorithm.

All relevant comments will be posted in their entirety and should not include PII information in the body of the email message.

#### Isogeny Signatures

Algorithm	Algorithm Information	Submitters	Comments
<b>SQIsign</b>	<a href="#">Specification</a> <a href="#">JP Statements</a> <a href="#">Zip file (14 MB)</a> <a href="#">Website</a>	Jorge Chavez-Saab Maria Corte-Real Santos Luca De Feo Jonathan Komada Eriksen Basil Hess David Kohel Antonin Leroux Patrick Longa Michael Meyer Lorenz Panny Sikhar Patranabis Christophe Petit Francisco Rodriguez Henriquez Sina Schaeffler Benjamin Wesolowski	<a href="#">Submit Comment</a> <a href="#">View Comments</a>

Only submission  
based on isogeny;  
shortest signatures;  
fast verification; com-  
plex signing procedure



**NIST**  
Information Technology Laboratory  
**COMPUTER SECURITY RESOURCE CENTER**

[PROJECTS](#) [PQC DIGITAL SIGNATURE SCHEMES](#)

## Post-Quantum Cryptography: Digital Signature Schemes

[f](#) [t](#)

[PROJECT LINKS](#)

[Overview](#)  
[News & Updates](#)

### Round 1 Additional Signatures

Official comments on the **First Round Signatures** should be submitted using the "Submit Comment" link for the appropriate algorithm. Comments from the [pqc-forum Google group subscribers](#) will also be forwarded to the pqc-forum Google group list. We will periodically post and update the comments received to the appropriate algorithm.

All relevant comments will be posted in their entirety and should not include PII information in the body of the email message.

#### Isogeny Signatures

Algorithm	Algorithm Information	Submitters	Comments
<b>SQIsign</b>	<a href="#">Specification</a> <a href="#">IP Statements</a> <a href="#">Zip file (14 MB)</a> <a href="#">Website</a>	Jorge Chavez-Saab Maria Corte-Real Santos Luca De Feo Jonathan Komada Eriksen Basil Hess David Kohel Antonin Leroux Patrick Longa Michael Meyer Lorenz Panny Sikhar Patranabis Christophe Petit Francisco Rodriguez Henriquez Sina Schaeffler Benjamin Wesolowski	<a href="#">Submit Comment</a> <a href="#">View Comments</a>

# Preparation for SQIsign

Let  $\lambda$  be the security parameter.

- Fix a prime  $p \equiv 3 \pmod{4}$  with  $\log_2(p) \approx 2\lambda$ . such that the  $N2^f$ -torsion subgroup is defined over a small extension of  $\mathbb{F}_{p^2}$  for smooth number  $N \simeq p^{5/4}$  and  $f$  is as big as possible.

# Preparation for SQIsign

Let  $\lambda$  be the security parameter.

- Fix a prime  $p \equiv 3 \pmod{4}$  with  $\log_2(p) \approx 2\lambda$ . such that the  $N2^f$ -torsion subgroup is defined over a small extension of  $\mathbb{F}_{p^2}$  for smooth number  $N \simeq p^{5/4}$  and  $f$  is as big as possible.
- Let  $N2^f = MM'$  such that  $M$  is a  $\lambda$ -bit integer consisting all the smallest factors, and  $M'$  is a  $2\lambda$ -bit integer.

# Preparation for SQIsign

Let  $\lambda$  be the security parameter.

- Fix a prime  $p \equiv 3 \pmod{4}$  with  $\log_2(p) \approx 2\lambda$ . such that the  $N2^f$ -torsion subgroup is defined over a small extension of  $\mathbb{F}_{p^2}$  for smooth number  $N \simeq p^{5/4}$  and  $f$  is as big as possible.
- Let  $N2^f = MM'$  such that  $M$  is a  $\lambda$ -bit integer consisting all the smallest factors, and  $M'$  is a  $2\lambda$ -bit integer.
- Let  $L = 2^e \simeq p^{15/4}$ , where  $e$  is greater than the diameter of  $G_2(p)$ .



# Preparation for SQIsign

Let  $\lambda$  be the security parameter.

- Fix a prime  $p \equiv 3 \pmod{4}$  with  $\log_2(p) \approx 2\lambda$ . such that the  $N2^f$ -torsion subgroup is defined over a small extension of  $\mathbb{F}_{p^2}$  for smooth number  $N \simeq p^{5/4}$  and  $f$  is as big as possible.
- Let  $N2^f = MM'$  such that  $M$  is a  $\lambda$ -bit integer consisting all the smallest factors, and  $M'$  is a  $2\lambda$ -bit integer.
- Let  $L = 2^e \simeq p^{15/4}$ , where  $e$  is greater than the diameter of  $G_2(p)$ .
- Fix  $E_0 : y^2 = x^3 + x$  with known endomorphism ring  $O_0 := \text{End}(E_0)$ .

# SQIsign, Step 1: $\Sigma$ -protocol

The prover  $P$  chooses a random isogeny  $\phi : E_0 \rightarrow E_1$  such that  $\deg(\phi)$  is a prime smaller than  $2^{\lambda/2}$ , leading to a random elliptic curve  $E_1$ .  $P$  keeps  $\phi$  secret and publishes  $E_1$ . Now,  $P$  can prove “knowledge” of  $O_1 := \text{End}(E_1)$  to a verifier  $V$ :

$P$

$$\begin{aligned}\phi' &\xleftarrow{\$} \text{Hom}(E_0, -) \text{ of degree } M' \\ E'_1 &= \phi'(E_0)\end{aligned}$$

$V$

$$\begin{aligned}C &\leq E'_1(\overline{\mathbb{F}}_p), C \cong \mathbb{Z}/M\mathbb{Z} \\ \tau &\leftarrow \text{Hom}((E'_1, C), -)\end{aligned}$$

$$\eta : E_1 \rightarrow E'_2, \ker(\hat{\tau} \circ \eta) \text{ cyclic}$$

$$\eta \stackrel{?}{\in} \text{Hom}(E_1, E'_2), \ker(\hat{\tau} \circ \eta) \stackrel{?}{=} \text{cyclic}$$

# SQIsign, Step 1: $\Sigma$ -protocol

The prover  $P$  chooses a random isogeny  $\phi : E_0 \rightarrow E_1$  such that  $\deg(\phi)$  is a prime smaller than  $2^{\lambda/2}$ , leading to a random elliptic curve  $E_1$ .  $P$  keeps  $\phi$  secret and publishes  $E_1$ . Now,  $P$  can prove “knowledge” of  $O_1 := \text{End}(E_1)$  to a verifier  $V$ :

$P$  -----  $p, E_0, O_0, M, L, E_1$  -----  $V$

$$\begin{aligned} \phi' &\xleftarrow{\$} \text{Hom}(E_0, -) \text{ of degree } M' \\ E'_1 &= \phi'(E_0) \end{aligned}$$

$$\begin{aligned} C &\leq E'_1(\overline{\mathbb{F}}_p), C \cong \mathbb{Z}/M\mathbb{Z} \\ \tau &\leftarrow \text{Hom}((E'_1, C), -) \end{aligned}$$

$$\eta : E_1 \rightarrow E'_2, \ker(\hat{\tau} \circ \eta) \text{ cyclic}$$

$$\eta \stackrel{?}{\in} \text{Hom}(E_1, E'_2), \ker(\hat{\tau} \circ \eta) \stackrel{?}{=} \text{cyclic}$$

# SQIsign, Step 1: $\Sigma$ -protocol

The prover  $P$  chooses a random isogeny  $\phi : E_0 \rightarrow E_1$  such that  $\deg(\phi)$  is a prime smaller than  $2^{\lambda/2}$ , leading to a random elliptic curve  $E_1$ .  $P$  keeps  $\phi$  secret and publishes  $E_1$ . Now,  $P$  can prove “knowledge” of  $O_1 := \text{End}(E_1)$  to a verifier  $V$ :

$P$  -----  $p, E_0, O_0, M, L, E_1$  -----  $V$

$\phi' \xleftarrow{\$} \text{Hom}(E_0, -)$  of degree  $M'$   
 $E'_1 = \phi'(E_0)$

$E'_1$

$C \leq E'_1(\overline{\mathbb{F}}_p), C \cong \mathbb{Z}/M\mathbb{Z}$   
 $\tau \leftarrow \text{Hom}((E'_1, C), -)$

$\eta : E_1 \rightarrow E'_2, \ker(\hat{\tau} \circ \eta) \text{ cyclic}$

$\eta \stackrel{?}{\in} \text{Hom}(E_1, E'_2), \ker(\hat{\tau} \circ \eta) \stackrel{?}{=} \text{cyclic}$

# SQIsign, Step 1: $\Sigma$ -protocol

The prover  $P$  chooses a random isogeny  $\phi : E_0 \rightarrow E_1$  such that  $\deg(\phi)$  is a prime smaller than  $2^{\lambda/2}$ , leading to a random elliptic curve  $E_1$ .  $P$  keeps  $\phi$  secret and publishes  $E_1$ . Now,  $P$  can prove “knowledge” of  $O_1 := \text{End}(E_1)$  to a verifier  $V$ :

$P$  -----  $p, E_0, O_0, M, L, E_1$  -----  $V$

$\phi' \xleftarrow{\$} \text{Hom}(E_0, -)$  of degree  $M'$   
 $E'_1 = \phi'(E_0)$

$E'_1$

$C \leq E'_1(\overline{\mathbb{F}}_p), C \cong \mathbb{Z}/M\mathbb{Z}$   
 $\tau \leftarrow \text{Hom}((E'_1, C), -)$

$\tau : E'_1 \rightarrow E'_2$

$\eta : E_1 \rightarrow E'_2, \ker(\hat{\tau} \circ \eta) \text{ cyclic}$

$\eta \in \text{Hom}(E_1, E'_2), \ker(\hat{\tau} \circ \eta) \stackrel{?}{=} \text{cyclic}$

# SQIsign, Step 1: $\Sigma$ -protocol

The prover  $P$  chooses a random isogeny  $\phi : E_0 \rightarrow E_1$  such that  $\deg(\phi)$  is a prime smaller than  $2^{\lambda/2}$ , leading to a random elliptic curve  $E_1$ .  $P$  keeps  $\phi$  secret and publishes  $E_1$ . Now,  $P$  can prove “knowledge” of  $O_1 := \text{End}(E_1)$  to a verifier  $V$ :

$P$  -----  $p, E_0, O_0, M, L, E_1$  -----  $V$

$\phi' \xleftarrow{\$} \text{Hom}(E_0, -)$  of degree  $M'$   
 $E'_1 = \phi'(E_0)$

$E'_1$


$C \leq E'_1(\overline{\mathbb{F}}_p), C \cong \mathbb{Z}/M\mathbb{Z}$   
 $\tau \leftarrow \text{Hom}((E'_1, C), -)$

$\tau : E'_1 \rightarrow E'_2$

$\eta : E_1 \rightarrow E'_2, \ker(\hat{\tau} \circ \eta) \text{ cyclic} \xrightarrow{\eta} \eta \stackrel{?}{\in} \text{Hom}(E_1, E'_2), \ker(\hat{\tau} \circ \eta) \stackrel{?}{=} \text{cyclic}$

# SQIsign, Step 1: $\Sigma$ -protocol

The prover  $P$  chooses a random isogeny  $\phi : E_0 \rightarrow E_1$  such that  $\deg(\phi)$  is a prime smaller than  $2^{\lambda/2}$ , leading to a random elliptic curve  $E_1$ .  $P$  keeps  $\phi$  secret and publishes  $E_1$ . Now,  $P$  can prove “knowledge” of  $O_1 := \text{End}(E_1)$  to a verifier  $V$ :



## Computing $L$ -isogeny $\eta : E_1 \rightarrow E_2'$

1. Translate isogeny  $\tau \circ \phi' \circ \hat{\phi}$  to left  $O_1$ -ideal  $I := \bar{l}_\phi \cdot l_{\phi'} \cdot l_\tau$  (*isogeny-to-kernel-to-ideal*).
2. From  $I, l_\phi$  get  $J \in [I]_L$  with  $\text{nrd}(J) = L$ .
3. Translate left  $O_1$ -ideal  $J$  to  $\eta$  (*ideal-to-kernel-to-isogeny*)

## SQIsign, Step 2: Fiat-Shamir transformation

Choose a *random oracle* cryptographic hash function  $H$  with appropriate domain and codomain (based on a compression algorithm).



## SQIsign, Step 2: Fiat-Shamir transformation

Choose a *random oracle* cryptographic hash function  $H$  with appropriate domain and codomain (based on a compression algorithm). The key generation algorithm  $G$  outputs a pair  $(O_1, E_1)$  such that  $O_1 = \text{End}(E_1)$ , where  $O_1$  is the *secret signing key* and  $E_1$  is the *public verification key*.

## SQIsign, Step 2: Fiat-Shamir transformation

Choose a *random oracle* cryptographic hash function  $H$  with appropriate domain and codomain (based on a compression algorithm). The key generation algorithm  $G$  outputs a pair  $(O_1, E_1)$  such that  $O_1 = \text{End}(E_1)$ , where  $O_1$  is the *secret signing key* and  $E_1$  is the *public verification key*.

### Signing $(M', M, L, O_1, H, m)$

1.  $\phi' \xleftarrow{\$} \text{Hom}(E_0, -)$  of degree  $M'$
2.  $E'_1 = \phi'(E_0)$
3.  $b = H(m \| j(E'_1))$
4.  $\tau = \text{Decompress}(E'_1, b)$
5.  $\eta : E_1 \rightarrow E'_2$ ,  $\ker(\hat{\tau} \circ \eta)$  cyclic
6. return  $\sigma := (E'_1, \eta)$

## SQIsign, Step 2: Fiat-Shamir transformation

Choose a *random oracle* cryptographic hash function  $H$  with appropriate domain and codomain (based on a compression algorithm). The key generation algorithm  $G$  outputs a pair  $(O_1, E_1)$  such that  $O_1 = \text{End}(E_1)$ , where  $O_1$  is the *secret signing key* and  $E_1$  is the *public verification key*.

### Signing $(M', M, L, O_1, H, m)$

1.  $\phi' \xleftarrow{\$} \text{Hom}(E_0, -)$  of degree  $M'$
2.  $E'_1 = \phi'(E_0)$
3.  $b = H(m \parallel j(E'_1))$
4.  $\tau = \text{Decompress}(E'_1, b)$
5.  $\eta : E_1 \rightarrow E'_2$ ,  $\ker(\hat{\tau} \circ \eta)$  cyclic
6. return  $\sigma := (E'_1, \eta)$

### Verification $(M, L, E_1, H, m, \sigma)$

1.  $b = H(m \parallel j(E'_1))$
2.  $\tau = \text{Decompress}(E'_1, b)$
3. return  $\eta \stackrel{?}{\in} \text{Hom}(E_1, E'_2)$ ,  
 $\ker(\hat{\tau} \circ \eta) \stackrel{?}{=} \text{cyclic}$

SIDH (2011-2022) reached  
Round 4 of NIST's quantum-  
safe KEM list.

A screenshot of a Quanta Magazine article. The page has a white background with a dark header. The Quanta Magazine logo is in the top left, and navigation icons are in the top right. The article title is highlighted with a red rounded rectangle. The author's name and the date are below the title. A short paragraph of the article is visible below the title. The sub-header 'A New Twist on Old Mathematics' is in bold. The main text of the article is in a standard font, with some words underlined. The article discusses the breaking of a quantum-resistant encryption protocol.

Quanta magazine

CRYPTOGRAPHY

**'Post-Quantum' Cryptography Scheme Is Cracked on a Laptop**

By JORDANA CEPELEWICE

August 24, 2022

*Two researchers have broken an encryption protocol that many saw as a promising defense against the power of quantum computing.*


**A New Twist on Old Mathematics**





Thomas Decru didn't set out to break SIDH. He was trying to build on it — to generalize the method to enhance another type of cryptography. That didn't work out, but it sparked an idea: His approach might be useful for attacking SIDH. And so he approached [Wouter Castryck](#), his colleague at the Catholic University of Leuven in Belgium and one of his former doctoral advisers, and the two dived into the relevant literature.

They stumbled across a paper published by the mathematician [Ernst Kani](#) in 1997. In it was a theorem that “was almost immediately applicable to SIDH,” Castryck said. “I think once we realized that ... the attack came quite quickly, in one or two days.”

# Quantum-safe?

On August 5, 2022, Castryck and Decru posted a preprint outlining an efficient classical key recovery algorithm against SIDH.

 **Quanta** magazine

CRYPTOGRAPHY

## 'Post-Quantum' Cryptography Scheme Is Cracked on a Laptop

By JORDANA CEPELEWICZ  
August 24, 2022

*Two researchers have broken an encryption protocol that many saw as a promising defense against the power of quantum computing.*


### A New Twist on Old Mathematics

Thomas Decru didn't set out to break SIDH. He was trying to build on it — to generalize the method to enhance another type of cryptography. That didn't work out, but it sparked an idea: His approach might be useful for attacking SIDH. And so he approached [Wouter Castryck](#), his colleague at the Catholic University of Leuven in Belgium and one of his former doctoral advisers, and the two dived into the relevant literature.

They stumbled across a paper published by the mathematician [Ernst Kani](#) in 1997. In it was a theorem that “was almost immediately applicable to SIDH,” Castryck said. “I think once we realized that ... the attack came quite quickly, in one or two days.”

# Quantum-safe?

SQISignHD uses this constructively: easier to generate public parameters & simpler signing procedure; but needs efficient implementation of 4D isogeny.



Questions?

Choose a cryptographic hash function  $H$  with appropriate domain and codomain. The key generation algorithm  $G$  outputs a pair  $(k, Q)$  such that  $Q = [k]P$ , where  $k$  is the *secret signing key* and  $Q$  is the *public verification key*.

## Signing ( $\mathbb{G}, P, k, H, m$ )

1.  $t \xleftarrow{\$} \{1, \dots, \ell - 1\}$
2.  $R \leftarrow [t]P$
3.  $r \leftarrow x(R) \pmod{\ell}$
4. if  $r = 0$  then goto Step 1.
5.  $e \leftarrow H(m)$
6.  $s \leftarrow (e + kr)t^{-1} \pmod{\ell}$
7. if  $s = 0$  then goto Step 1.
8. return  $\sigma := (r, s)$

## Verification ( $\mathbb{G}, P, Q, H, m, \sigma$ )

1.  $e \leftarrow H(m)$
2.  $u_1 \leftarrow es^{-1} \pmod{\ell}$ ,  $u_2 \leftarrow rs^{-1} \pmod{\ell}$
3.  $T \leftarrow [u_1]P + [u_2]Q$
4. return  $r \stackrel{?}{=} x(T) \pmod{\ell}$



# EdDSA, a footnote

## Twisted Edwards model

A twisted Edwards curve defined over  $\mathbb{F}_q$  is the curve

$$C : ax^2 + y^2 = 1 + dx^2y^2, \quad a, d \in \mathbb{F}_q, \text{ and } ad(a - d) \neq 0$$

with two singular points. It is birationally equivalent to  $E : v^2 = u^3 + 2(a + d)u^2 + (a - d)^2u$  such that every point has order divisible by 4.

