







Robot Perception: Compute Projective Transformations

Kostas Daniilidis

A perspective projection of a plane
(like a camera image) is always a
projective transformation



Using the projective transformation the pose of a robot with respect to a planar pattern:



Projective Transformation

Definition

A **projective transformation** is any invertible matrix transformation $\mathbb{P}^2 \rightarrow \mathbb{P}^2$.

A projective transformation A maps p to $p' \sim Ap$.


Invertibility means that $\det(A) \neq 0$ and that there exists $\lambda \neq 0$ such that $\lambda p' = Ap$.

Observe that we will write either $p' \sim Ap$ or $\lambda p' = Ap$.

A projective transformation is also known as **collineation** or **homography**.

A projective transformation preserves incidence:

- Three collinear points are mapped to three collinear points.
- and three concurrent lines are mapped to three concurrent lines.



Projective transformation of lines

If A maps a point to Ap , then where does a line l map to?

Line equation in original plane

$$l^T p = 0$$

Line equation in image plane $p' \sim Ap$

$$l^T A^{-1} p' = 0$$

implies that $l' = A^{-T} l$.

Projective transformation of lines

If A maps a point to Ap , then where does a line l map to?

Line equation in original plane

$$l^T p = 0$$

Line equation in image plane $p' \sim Ap$

$$l^T A^{-1} p' = 0$$

implies that $l' = A^{-T} l$.

Projective transformation of lines

If A maps a point to Ap , then where does a line l map to?

Line equation in original plane


$$l^T p = 0$$

Line equation in image plane $p' \sim Ap$

$$l^T A^{-1} p' = 0$$

implies that $l' = A^{-T} l$.

How can we compute the projective transformation between




Floor tiles measured in [m]

and




Points in pixel coordinates


The result of such a transformation would map any point in one plane to the corresponding point in the other




How can we compute the projective transformation between



How can we compute the projective transformation between



How can we compute the projective transformation between




Assume that a mapping A maps the three points $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$ to the non-collinear points A, B, C

with coordinate vectors a, b and $c \in \mathbb{P}^2$. Then the following is a possible projective transformation:

$$(a \quad b \quad c) = (\alpha a \quad \beta b \quad \gamma c) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

with 3 degrees of freedoms α, β and γ . This means 3 points do not suffice to compute a projective transformation.

Let us introduce a 4th point D



Let us assume that the same A maps $(1, 1, 1)$ to the point d . Then, the following should hold:

$$\lambda d = \begin{pmatrix} \alpha a & \beta b & \gamma c \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix},$$

hence

$$\lambda d = \alpha a + \beta b + \gamma c.$$

There always exist such $\lambda, \alpha, \beta, \gamma$ because four elements of $\mathbb{R}^3 \setminus \{(0, 0, 0)\}$ are always linearly dependent.

Because a, b, c are not collinear, there exist unique $\alpha/\lambda, \beta/\lambda, \gamma/\lambda$ for writing this linear combination.

Since A is the same as A/λ we solve for α, β, γ such that $d = \alpha a + \beta b + \gamma c$, which can be written as a linear system

$$\begin{pmatrix} a & b & c \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} = d.$$

Since a, b, c are not collinear we can always find a unique triple α, β, γ . The resulting projective transformation is $A = \begin{pmatrix} \alpha a & \beta b & \gamma c \end{pmatrix}$.

There always exist such $\lambda, \alpha, \beta, \gamma$ because four elements of $\mathbb{R}^3 \setminus \{(0, 0, 0)\}$ are always linearly dependent.

Because a, b, c are not collinear, there exist unique $\alpha/\lambda, \beta/\lambda, \gamma/\lambda$ for writing this linear combination.

Since A is the same as A/λ we solve for α, β, γ such that $d = \alpha a + \beta b + \gamma c$, which can be written as a linear system

$$\begin{pmatrix} a & b & c \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} = d.$$

Since a, b, c are not collinear we can always find a unique triple α, β, γ . The resulting projective transformation is $A = \begin{pmatrix} \alpha a & \beta b & \gamma c \end{pmatrix}$.

There always exist such $\lambda, \alpha, \beta, \gamma$ because four elements of $\mathbb{R}^3 \setminus \{(0, 0, 0)\}$ are always linearly dependent.

Because a, b, c are not collinear, there exist unique $\alpha/\lambda, \beta/\lambda, \gamma/\lambda$ for writing this linear combination.

Since A is the same as A/λ we solve for α, β, γ such that $d = \alpha a + \beta b + \gamma c$, which can be written as a linear system

$$\begin{pmatrix} a & b & c \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} = d.$$

Since a, b, c are not collinear we can always find a unique triple α, β, γ . The resulting projective transformation is $A = \begin{pmatrix} \alpha a & \beta b & \gamma c \end{pmatrix}$.

Four points not three of them collinear suffice to recover unambiguously a projective transformation.

Knowledge of this projective transformation makes Virtual Billboards possible!



Microsoft Office Lens App



Office Lens

Microsoft Corporation Productivity


Everyone

Top Developer



★★★★★ 91,419

This app is compatible with all of your devices.




Installed



Office Lens



What happens when the original set of points is not a square?



Find projective transformation mapping $(a, b, c, d) \rightarrow (a', b', c', d')$:

To determine this mapping we go through the four canonical points.

We find the mapping from $(1, 0, 0)$, *etc* to (a, b, c, d) and we call it T :

$$a \sim T(1, 0, 0)^T, \text{ etc}$$

We find the mapping from $(1, 0, 0)$, *etc* to (a', b', c', d') and we call it T' :

$$a' \sim T'(1, 0, 0)^T, \text{ etc}$$

Then, back-substituting $(1, 0, 0)^T \sim T^{-1}a$, *etc* we obtain that

$$a' = T'T^{-1}a, \text{ etc}$$

To determine this mapping we go through the four canonical points.

We find the mapping from $(1, 0, 0)$, *etc* to (a, b, c, d) and we call it T :

$$a \sim T(1, 0, 0)^T, \text{ etc}$$

We find the mapping from $(1, 0, 0)$, *etc* to (a', b', c', d') and we call it T' :

$$a' \sim T'(1, 0, 0)^T, \text{ etc}$$

Then, back-substituting $(1, 0, 0)^T \sim T^{-1}a$, *etc* we obtain that

$$a' = T'T^{-1}a, \text{ etc}$$

To determine this mapping we go through the four canonical points.

We find the mapping from $(1, 0, 0)$, *etc* to (a, b, c, d) and we call it T :

$$a \sim T(1, 0, 0)^T, \text{ etc}$$

We find the mapping from $(1, 0, 0)$, *etc* to (a', b', c', d') and we call it T' :

$$a' \sim T'(1, 0, 0)^T, \text{ etc}$$

Then, back-substituting $(1, 0, 0)^T \sim T^{-1}a$, *etc* we obtain that

$$a' = T'T^{-1}a, \text{ etc}$$