

C3S-CDS marine in-situ data

Release r092019-000000

Annex to User Manual

Scottish Fishery Cruisers MARIDS (1930 code)


MetO MDB: Flatfile 1A (Flatfile plus cardimage data)

1950 to 1956 summary

091-205


Data source summary
Optimal data quality subset

Temporal extent of reports





Shaded area: number of reports with passed report_quality flag. Series: number observed parameter reports with passed quality_flag flag.

Spatial distribution of reports




Spatial distribution of reports. Reports included are those with passed report_quality flag.

Air Temperature




TOP. Series, left axis: latitudinal band aggregations of air temperature: minimum and maximum value (min/max) and mean (mean). The asterisks indicate where values fall outside of the plotting area; Shaded area, right axis: number of reports. BOTTOM. Spatial distribution on a 1x1 grid of number of reports (left), minimum value (center) and maximum value (right). Data values and counts extracted from the best data quality data subset only (report_quality and parameter_quality_flag passed).

Sea Surface Temperature



TOP. Series, left axis: latitudinal band aggregations of sea surface temperature: minimum and maximum value (min/max) and mean (mean). The asterisks indicate where values fall outside of the plotting area; Shaded area, right axis: number of reports. BOTTOM. Spatial distribution on a 1x1 grid of number of reports (left), minimum value (center) and maximum value (right). Data values and counts extracted from the best data quality subset only (report_quality and parameter quality_flag passed).


Wind Direction



unavailable

TOP. Series, left axis: latitudinal band aggregations of wind direction: minimum and maximum value (min/max) and mean (mean). The asterisks indicate where values fall outside of the plotting area; Shaded area, right axis: number of reports. BOTTOM. Spatial distribution on a 1x1 grid of number of reports (left), minimum value (center) and maximum value (right). Data values and counts extracted from the best data quality data subset only (report_quality and parameter_quality_flag passed).


Wind Speed



TOP. Series, left axis: latitudinal band aggregations of wind speed: minimum and maximum value (min/max) and mean (mean). The asterisks indicate where values fall outside of the plotting area; Shaded area, right axis: number of reports. BOTTOM. Spatial distribution on a 1x1 grid of number of reports (left), minimum value (center) and maximum value (right). Data values and counts extracted from the best data quality data subset only (report_quality and parameter_quality_flag passed).


Data source summary
All available data

IO flow from ICOADS to C3S




Series, left axis: input/output main flow of reports from source data (ICOADS) to dataset included in release. Shaded areas, right axis: percent of invalid data with respect to the initially selected reports (PT selection). Invalid status are: invalid data model (md-invalid), invalid datetime (dt-invalid) and invalid station ID (id-invalid)

Duplicate status of reports




Duplicate status of reports included in release. Series, left axis: number of reports; shaded areas, right axis: stacked percent. Duplicate status are: best duplicate (best), unique report (unique), duplicate (dup), worst duplicate (worst), duplicate status not checked (unchecked)

Report quality status




Report quality status for reports included in release. Series, left axis: number of reports; shaded areas, right axis: stacked percent. Quality status are: passed (pass), location check failed (loc-fail), quality control of all measured parameters failed while location check passed (param-failed), quality not checked (unchecked)

Temporal extent of reports



Shaded area: total number of reports included in release (all report_quality flags). Series: number reports from the observed parameter (all quality_flag flags).


Spatial distribution of reports



Spatial distribution of reports. Reports of all qualities are included.


Air Temperature

Air Temperature
1953 to 1953



TOP: Series, left axis: latitudinal band aggregations of air temperature: minimum and maximum value (min/max) and mean (mean); Shaded areas, right axis: stacked number of reports with different parameter quality status. Quality status are: report and parameter passed (qc passed), parameter qc failed (qc failed), location check failed (qcr failed), quality not checked (qcr unchecked). BOTTOM: Spatial distribution on a 1x1 grid of number of reports (left), minimum value (center) and maximum value (right). Data values and counts extracted from all the available data from the source. Axis represent the full extent of the data available.


Sea Surface Temperature



unavailable

TOP: Series, left axis: latitudinal band aggregations of sea surface temperature: minimum and maximum value (min/max) and mean (mean); Shaded areas, right axis: stacked number of reports with different parameter quality status. Quality status are: report and parameter passed (qc passed), parameter qc failed (qc failed), location check failed (qcr failed), quality not checked (qcr unchecked) . BOTTOM: Spatial distribution on a 1x1 grid of number of reports (left), minimum value (center) and maximum value (right). Data values and counts extracted from all the available data from the source. Axis represent the full extent of the data available.


Wind Direction



unavailable

TOP: Series, left axis: latitudinal band aggregations of wind direction: minimum and maximum value (min/max) and mean (mean); Shaded areas, right axis: stacked number of reports with different parameter quality status. Quality status are: report and parameter passed (qc passed), parameter qc failed (qc failed), location check failed (qcr failed), quality not checked (qcr unchecked). BOTTOM: Spatial distribution on a 1x1 grid of number of reports (left), minimum value (center) and maximum value (right). Data values and counts extracted from all the available data from the source. Axis represent the full extent of the data available.

Wind Speed



unavailable

TOP: Series, left axis: latitudinal band aggregations of wind speed: minimum and maximum value (min/max) and mean (mean); Shaded areas, right axis: stacked number of reports with different parameter quality status. Quality status are: report and parameter passed (qc passed), parameter qc failed (qc failed), location check failed (qcr failed), quality not checked (qcr unchecked) . BOTTOM: Spatial distribution on a 1x1 grid of number of reports (left), minimum value (center) and maximum value (right). Data values and counts extracted from all the available data from the source. Axis represent the full extent of the data available.