# Discrete differential geometry in homotopy type theory

Greg Langmead

Carnegie Mellon University

April 2025

- 1 Intro
- 2 Motivation
- 3 Discrete manifolds
- 4 Codomain
- 6 Results

Intro

- Form BG, study maps  $I:BG\to B\mathbb{R}$  (invariant polynomials
- $\bullet$  Fix a principal bundle  $P: M \to BG$

- 6 / guarantees independence of class on connection

Intro

#### Motivation

- Form BG, study maps  $I:BG\to B\mathbb{R}$  (invariant polynomials such as det, trace)
- ② Fix a principal bundle  $P: M \rightarrow BG$
- **3** Choose a connection A on P, compute curvature  $F_A$
- 4 Compose  $I \circ F_A$  (cohomology class)
- 6 / guarantees independence of class on connection

Intro

- Form BG, study maps  $I:BG\to B\mathbb{R}$  (invariant polynomials such as det, trace)
- Fix a principal bundle  $P: M \rightarrow BG$

- 6 / guarantees independence of class on connection

Intro

- Form BG, study maps  $I:BG\to B\mathbb{R}$  (invariant polynomials such as det, trace)
- **2** Fix a principal bundle  $P: M \rightarrow BG$
- **3** Choose a connection A on P, compute curvature  $F_{\Delta}$
- 6 / guarantees independence of class on connection

Intro

- Form BG, study maps  $I:BG\to B\mathbb{R}$  (invariant polynomials such as det, trace)
- Fix a principal bundle  $P: M \rightarrow BG$
- **3** Choose a connection A on P, compute curvature  $F_A$
- **4** Compose  $I ∘ F_A$  (cohomology class)
- 6 / guarantees independence of class on connection

Intro

- Form BG, study maps  $I:BG\to B\mathbb{R}$  (invariant polynomials such as det, trace)
- **2** Fix a principal bundle  $P: M \rightarrow BG$
- **3** Choose a connection A on P, compute curvature  $F_A$
- **4** Compose  $I ∘ F_A$  (cohomology class)
- 6 I guarantees independence of class on connection

Intro

- What is  $BG \to B\mathbb{R}$  in HoTT?
- What are bundles, connections, and curvature in HoTT?
- What is cohomology; what are exact forms?

## Questions

Where is the Bianchi identity? What is "the space of connections?"

Discrete manifolds

## Simplicial complexes


$$egin{aligned} extit{$M_1$} imes \partial \Delta^1 & \stackrel{\mathrm{pr}_1}{\longrightarrow} & extit{$M_1$} \ & \mathbb{A}_0 & \mathbb{M}_0 & \mathbb{M}_0 & \mathbb{M}_1 \end{aligned}$$

Discrete manifolds 000

## Simplicial complexes

$$P(2) imes \partial \Delta^1 \stackrel{r}{\longrightarrow} \partial P(2) \ egin{array}{ccc} \mathbb{A}_0 & & & \downarrow^*_{\partial \Delta} \ P(2)_0 & & & \partial \Delta^2 \end{array}$$


Discrete manifolds



## Codomain

 $\mathrm{EM}(\mathbb{Z},1)$ ,  $\mathrm{K}(\mathbb{Z},2)$ 

## Classical proof



**[26.2]** The difference  $\Re(\Delta) - 2\pi \Im_F(s)$  can be found by summing over the edges  $K_j$  the change  $\Phi(K_j)$  in the illustrated angle  $\angle Fw_{||}$ , i.e., the rotation of  $w_{||}$  relative to F.

Every aspect was informed by the classical proof, which is already discrete-flavored. The " $\angle Fw_{||}$ " looked a lot like a pathover.

Figure: Tristan Needham, Visual Differential Geometry and Forms