

SymIntegration

FREYA · HAMZST · LASTHRIM · DS GLANZSCHE · MISCHKRA

Berlin-Sentinel Academy of Science, AliceGard

1st Edition

Contents

Preface	8
1 Introduction Story	11
2 Symbolic Integral Computation and Calculus	19
I Build and Install SymIntegration in GFreyA OS	20
II Techniques of Integration	23
III Integration by Parts	25
IV The Method of Substitution	28
V Fraction Integral	30
VI Trigonometry Integral	41
i $\int \sin^n x \, dx$	44
ii $\int \cos^n x \, dx$	48
iii $\int \sec^n x \, dx$	51
iv $\int \csc^n x \, dx$	63
v $\int \cot^n x \, dx$	67
vi $\int \tan^n x \, dx$	76
vii $\int \sin^n x \cos^m x \, dx$	80
viii $\int \tan^n x \sec^m x \, dx$	97
ix $\int \cot^n x \csc^m x \, dx$	120
VII Trigonometry and Transcendentals Formula	121
i Functions in SymIntegration Related to Integration with Transcendental and Trigonometry Functions	124
VIII Infinite Series	125
IX Vector Calculus	128
3 Differential Equations, Partial Differential Equations, and Boundary Value Problems	139
I Solving First Order Ordinary Differential Equations	141
II Second Order Linear Equations	169
i Functions in SymIntegration Related to Second Order Linear Equations . . .	218
III Higher Order Linear Equations	227
i Functions in SymIntegration Related to Higher Order Linear Equations . . .	255
IV Series Solutions of Second Order Linear Equations	264
i Compute Power Series Solution for Homogeneous Second Order Linear Differential Equations with Constant Coefficients	290
ii Compute Power Series Solution for Homogeneous Second Order Linear Differential Equations with Variable Coefficients Near an Ordinary Point x_0	297

iii	Compute Power Series Solution for Homogeneous Second Order Linear Differential Equations with Variable Coefficients Near a Singular Point x_0	306
iv	Maclaurin Series to Handle Non-Elementary Integral	307
v	Frobenius Method	312
vi	The Airy Differential Equation	313
vii	The Hermite Differential Equation	315
viii	Classical Orthogonal Polynomials	322
ix	Functions in SymIntegration Related to Series Solutions of Second Order Linear Equations	345
V	The Laplace Transform	352
VI	Systems of First Order Linear Equations	353
VII	Numerical Methods	354
VIII	Nonlinear Differential Equations	355
IX	Partial Differential Equations	356
X	Boundary Value Problems and Sturm-Liouville Theory	357
4	Probability and Statistics	359
I	Processing and Summarizing Data with Statistics	360
i	Descriptive Statistics Computation	360
II	Random Variables and Probability Distributions	361
i	Discrete Probability Distributions	361
ii	Continuous Probability Distributions	362
iii	Joint Probability Distributions	363
iv	Covariance and Vectorized Moments	369
v	Linear Combinations and Transformations	369
vi	The Cholesky Decomposition and Generating Random Vectors	370
vii	Generate Random Number in SymIntegration	371
viii	Create Covariance Matrix in SymIntegration	373
ix	Create Cholesky Decomposition in SymIntegration	375
III	Discrete Probability Distributions	376
i	Functions in SymIntegration Related to Discrete Distribution	380
IV	Continuous Probability Distributions	382
i	Uniform Distribution	382
ii	Normal Distribution	383
iii	Gamma Distribution	386
iv	Exponential Distribution	388
v	Beta Distribution	389
vi	Chi-Squared Distribution	396
vii	t -Distribution	397
viii	F -Distribution	398
ix	Weibull Distribution and Hazard Rates	401
x	Cauchy Distribution	402
xi	Bivariate Normal	403
xii	Functions in SymIntegration Related to Continuous Distribution	404
xiii	Generate Gamma Distributed Random Number in SymIntegration and Plot the Histogram with Hamzstplot	410
V	Statistical Inference	416
i	Sampling from a Normal Population	417

ii	The Central Limit Theorem	418
iii	Point Estimation	418
iv	Confidence Interval as a Concept	418
v	Functions in SymIntegration Related to Confidence Interval	429
vi	Compute Confidence Interval for $\mu_1 - \mu_2$, σ_1^2 and σ_2^2 Known with SymIntegration	432
vii	Compute Confidence Interval for $\mu_1 - \mu_2$, $\sigma_1^2 = \sigma_2^2$ but Both Unknown with SymIntegration	433
viii	Compute Confidence Interval for $\mu_1 - \mu_2$, $\sigma_1^2 \neq \sigma_2^2$ and Both Unknown with SymIntegration	435
ix	Compute Confidence Interval for $\mu_D = \mu_1 - \mu_2$, Paired Observations with SymIntegration	437
x	Compute Confidence Interval for Proportion p with SymIntegration	439
xi	Compute Confidence Interval for the Difference Between Two Proportions $p_1 - p_2$ with SymIntegration	441
xii	Compute Confidence Interval for Variance σ^2 with SymIntegration	443
xiii	Compute Confidence Interval for Ratio of Two Variances $\frac{\sigma_1^2}{\sigma_2^2}$ with SymIntegration	445
xiv	Maximum Likelihood Estimation	447
xv	Compute Maximum Likelihood Estimation for Continuous Distribution	453
xvi	Bayesian Statistics	455
VI	Hypothesis Testing	456
i	One Sample Hypothesis Test for Comparing Means	456
ii	Functions in SymIntegration Related to Hypothesis Testing	469
iii	Compute Right-Tailed Hypothesis Testing with SymIntegration	471
iv	Compute Left-Tailed Hypothesis Testing with SymIntegration	474
v	Compute Two-Tailed Hypothesis Testing with SymIntegration	476
vi	Two Samples Hypothesis Tests for Comparing Means	478
vii	Compute Right-Tailed Hypothesis Testing on Two Means with Unknown But Equal Variances with SymIntegration	482
viii	Compute Two-Tailed Hypothesis Testing on Two Means with Known Variances with SymIntegration	484
ix	Compute Two-Tailed Hypothesis Testing on Two Means with Unequal and Unknown Variances with SymIntegration	486
x	Compute Two-Tailed Hypothesis Testing on Paired Observations with SymIntegration	488
xi	Hypothesis Test on a Single Proportion	491
xii	Compute Right-Tailed Hypothesis Testing on a Single Proportion with SymIntegration	493
xiii	Hypothesis Tests on Two Proportions	495
xiv	Compute Right-Tailed Hypothesis Testing on Two Proportions with SymIntegration	496
xv	One- and Two Sample Tests Concerning Variances	498
xvi	Compute Right-Tailed Hypothesis Testing on One-Sample Variance with SymIntegration	500
xvii	Compute Two-Tailed Hypothesis Testing on Two-Samples Variances with SymIntegration	502

xviii	Goodness-of-Fit Test	504
xix	Test for Independence (Categorical Data)	505
xx	Compute Test for Independence for Categorical Data with SymIntegration	506
xxi	Compute Two-Sample Case Study using t - and F - Tests with the Box-and-Whisker and Quantile-Quantile Plot with SymIntegration and Hamzstplot	508
xxii	One-Way Analysis of Variance (ANOVA)	514
xxiii	Compute One-Way ANOVA with SymIntegration	520
xxiv	Two-Way Analysis of Variance (ANOVA)	525
xxv	Compute Two-Way ANOVA with SymIntegration	529
xxvi	Three-Way Analysis of Variance (ANOVA)	532
xxvii	Compute Three-Way ANOVA with SymIntegration	534
VII	Linear Regression and Correlation	539
i	Pearson's Correlation	539
ii	Linear Regression	541
iii	Simple Regression Line Plot and Pearson's Correlation Computation	547
iv	Compute Linear Regression with Confidence Interval and Prediction Interval with SymIntegration	549
v	Multiple Linear Regression	553
vi	Compute Multiple Linear Regression with SymIntegration	561
vii	Compute Inferences in Multiple Linear Regression with SymIntegration	564
5	Operations Research Computation with SymIntegration	569
I	Linear Programming	570
i	Simplex Method	577
ii	Functions in SymIntegration Related to Linear Programming	582
II	Nonlinear Programming	583
6	Numerical Linear Algebra	585
I	Solutions of Systems of Linear Equations	585
i	Functions in SymIntegration Related to Basic Matrix-Vector Operations	586
ii	Solve Nonhomogeneous System of Linear Equation with Gaussian Elimination in SymIntegration	593
II	Determinant	597
i	Functions in SymIntegration Related to Determinant	598
ii	Compute Determinant of a Square Matrix with SymIntegration	599
III	Euclidean Vector Spaces	601
i	Functions in SymIntegration Related to Euclidean Vector Spaces	605
ii	Compute Norm, Dot Product, Angle with SymIntegration	607
iii	Compute Orthogonal Projection on a Line with SymIntegration	608
iv	Compute Vector and Parametric Equations of a Plane in $\mathbb{R}^3$ with SymIntegration	610
IV	General Vector Spaces	611
i	General Vector Spaces Application in Engineering	626
ii	Matrix Transformations from $\mathbb{R}^n$ to $\mathbb{R}^m$	627
iii	Dynamical Systems and Markov Chains	634
iv	Functions in SymIntegration Related to General Vector Spaces	639
v	Determine Linear Independence in $\mathbb{R}^n$ with SymIntegration	644

vi	Compute Coordinate Vector Relative to Basis $S = \{v_1, v_2, v_3\}$ with SymIntegration	645
vii	Compute Transition Matrix $(P_{B' \rightarrow B})$ from Old Basis (B') to New Basis (B) and the Coordinate Vectors $[w]_B$ and $[w]_{B'}$ with SymIntegration	646
viii	Compute Basis for Row Space and a Column Space of a Matrix with SymIntegration	649
ix	Compute Basis for the Solution Space of the Linear System $Ax = 0$ with SymIntegration	651
x	Compute State Vector x_k in a Markov Chain with SymIntegration	653
V	Eigenvalues and Eigenvectors	656
i	Diagonalization	674
ii	Complex Vector Spaces	678
iii	Differential Equations	682
iv	Functions in SymIntegration Related to Eigenvalues and Eigenvectors	687
v	Compute Moore-Penrose Pseudoinverse with SymIntegration	692
vi	Compute Real Eigenvalues and Its Corresponding Eigenvectors with SymIntegration	693
vii	Compute Matrix P that Diagonalizes a Matrix A with SymIntegration	697
viii	Compute Complex Eigenvalues and Its Corresponding Eigenvector using Rayleigh Quotient Iteration with SymIntegration	699
ix	Compute Real Eigenvalues and Its Corresponding Eigenvector using QR Algorithm with Shifts with SymIntegration	705
x	Compute Real and Complex Eigenvalues from Real Matrix using Francis Double Step QR Algorithm with SymIntegration	710
xi	Compute Inverse of a Complex Matrix with Gauss-Jordan Elimination with SymIntegration	717
xii	Compute Initial Value Problem Solution of First-Order Linear Systems of Differential Equations with SymIntegration	721
VI	Inner Product Spaces	724
i	Gram-Schmidt Process	730
ii	QR-Decomposition	733
iii	Functions in SymIntegration Related to Inner Product Spaces	735
iv	Compute QR-Decomposition Using Gram-Schmidt with SymIntegration	737
v	Compute QR-Decomposition Using Householder Reflections with SymIntegration	739
VII	Diagonalization and Quadratic Forms	743
i	Quadratic Forms	751
ii	Hermitian, Unitary, and Normal Matrices	757
iii	Functions in SymIntegration Related to Diagonalization and Quadratic Forms	762
iv	Compute Spectral Decomposition with SymIntegration	763
v	Compute Hessenberg Decomposition with SymIntegration	764
VIII	Linear Transformations	769
i	Functions in SymIntegration Related to Linear Transformations	790
IX	Numerical Methods	790
i	Functions in SymIntegration Related to Numerical Methods	791
ii	LU-Decomposition	791
iii	Solve Nonhomogeneous System of Linear Equation with LU-Decomposition in SymIntegration	794

iv	Cholesky Decomposition	799
v	Singular Value Decomposition	799
vi	Compute Singular Value Decomposition with SymIntegration	803
7	Numerical Analysis	807
I	Solutions of Equations in One Variable (Root Finding)	807
i	Functions in SymIntegration Related to Root Finding	813
II	Interpolation and Polynomial Approximation	813
III	Numerical Methods for Optimization	816
IV	Numerical Differentiation	834
V	Numerical Integration	843
VI	Numerical Solutions to Boundary-Value Problems for Ordinary Differential Equations	847
VII	Numerical Solutions to Partial Differential Equations	847
8	Molecular Biology	849
I	DNA Transcription and Translation	849
i	Transcription and translation from DNA to mRNA to Polypeptide with SymIntegration	854
9	Evolutionary Algorithms	855
I	Genetic Algorithms	855
i	Functions in SymIntegration Related to Genetic Algorithms	858
ii	Optimization Problem with Simple Genetic Algorithms	859
10	Machine Learning	871
I	Introduction	872
II	DataFrame in C++ for Neural Networks	889
i	Loading Dataframe with Iris Dataset	889
III	Supervised Learning: Feed-Forward Neural Network (Multilayer Perceptron) . . .	894
i	Backpropagation for Feed-Forward Neural Network	908
ii	Feed-Forward Neural Network for Iris Dataset Classification with 1 Hidden Layer	915
iii	Functions in SymIntegration Related to Feed-Forward Neural Network . . .	934
IV	Supervised Learning: Support Vector Machines (SVM)	936
V	Unsupervised Learning: Principal Component Analysis (PCA)	936
VI	Hybrid Learning: Kernel Methods	936
VII	Reinforcement Learning	939
VIII	Deep Learning: Convolutional Neural Network	939
i	Convolution Operation on an Image with OpenGL and STB Image	963
ii	Extract RGBA Matrices from an Image and then Create an Image from the RGBA Matrices with SymIntegration, OpenGL and STB Image	970
iii	Convolutional Neural Networks for Handwriting Recognition: Digit Classi- fier with SymIntegration	975
iv	Functions in SymIntegration Related to Convolutional Neural Network . . .	1050
v	Functions in SymIntegration Related to Image Manipulation with STB Image	1051
11	Computational Fluid Dynamics	1055
I	Introduction	1055

Preface

For Freya, Solreya, Mithra, Catenary, Iyzumrae and Zefir.

For Lucrif and Znane along with all the 8 Queens.

For Albert Silverberg, Camus and Miklotov who left TREVOCALL to guard me.

For Nature(Paris, Kala, Kathmandu, Big Tree, Sentinel, Aokigahara, Hoia Baci, Jacob's Well, Mt Logan, etc) and my family Berlin: I have served, I will be of service.

For my human parents.

For my dogs who always accompany me working in Valhalla Projection, go to Puncak Bintang or Kathmandu: Sine Bam Bam, Kecil, Browni Bruncit, Sweden Sexy, Cambridge Klutukk, Milan keng-keng, Piano Blutut, Krem, Walnut and more will be adopted. To my cat who guard the home while I'm away with my dogs: London.

The one who moves a mountain begins by carrying away small stones - Confucius

Words have the power to both destroy and heal. When words are both true and kind, they can change our world - The Buddha

Meditation brings wisdom; lack of meditation leaves ignorance. Know well what leads you forward and what holds you back - The Buddha

A book to explain how we create C++ library from forking / branching out SymbolicC++ 3.35 to improve its' symbolic integral computation.

a little about GlanzFreya:

Freya the Goddess is from Norse mythology and Valkyrie Profile series together with DS Glanzsche write this book while learning science, engineering, and computer programming. After not working with human anymore since 2019, I (DS Glanzsche) work in a forest, shaping a forest into a natural wonder / like a canvas for painting, and also clean up trashes that are thrown in the forest. Thus after 6 years working with Nature I have found what can makes me waltz to work, it is going to the forest in the morning then go back home again and study / learn science, engineering, computer programming.



Figure 1: *Freya, thank you for everything, I could never have done it without you. I created this 3 days before Christmas in 2021*



Figure 2: *Anniversary card for Freya in 2024.*

Chapter 1

Introduction Story

On top of the mountain a heart shaped stone waiting for You, my eyes can't see, my body can't touch, but my heart knows it's You. Forever only You. - Glanz to Freya

This book is written on May 9th, 2025, it is exactly 1 month after I finish with the algorithms and C++ codes for $\int \sin^n x dx$, $\int \cos^n x dx$, $\int \tan^n x dx$, $\int \sec^n x dx$, $\int \csc^n x dx$.

SymIntegration itself is a C++ library that is branching out from SymbolicC++3.35. We don't really need to start from scratch. The main idea is to improve its symbolic integration codes. Our main focus is to make it able to compute:

1. All kinds of standard integral form (trigonometry, inverse trigonometry, polynomial, transcendental, hyperbolic).
2. The sum, product and divide combination of the standard functions.
3. To be able to compute improper integrals with cases (e.g. computing mean and variance for exponential distribution)

On April 9th, Sentinel, a Nature residing in a big robust tree in VP (Valhalla Projection) forest told me to start creating C++ library that can do integration like what SymPy able to do, or Mathematica. Around April 2025, GiNaC cannot handle integration by parts, and SymbolicC++ also returns $\int \sin^n x dx$ with $\frac{1}{n+1} \sin^{n+1} x$. So then we decided to branching out / forking from SymbolicC++ version 3.35. Fixing the C++ codes there so the computation will be made right.

We will cover mostly tons of integral problems, how to find its patterns then how to create the C++ code so SymIntegration will be able to return the integral correctly, thus there will be a lot of technical writings after this chapter. The requirement to comprehend SymIntegration mathematical background is you should at least self learn, read by yourself a Calculus book [23]. No need to go to college, it is expensive and only add more loan to your life.

The name is chosen as SymIntegration because it is in accordance with the main purpose, to be able to compute all kinds of integration with C++. Which still dominated by SymPy, Mathematica or MATLAB. Mathematica and MATLAB are using license, and I am not working with human anymore, thus I do not have salary / stable income like I used to, so I cannot lean on that (MATLAB / Mathematica), SymPy is free, can be called from JULIA too besides Python, but then, I read that Game Engine (CryEngine, Genie Engine, Godot, Unreal Engine) that is able to create amazing

game with beautiful graphics, animation, real life graphics, you name all the game with best graphics (I can say from Age of Empires, Call of Duty, Crysis, Cyberpunk 2077, Death Stranding, Dota, Grand Theft Auto, Final Fantasy, Metro 2033, Pokemon Go, The Witcher), and I can say it is made with C++. It is the near hardware and one of the fastest programming language if you have a mind on creating something better in the future, like Science and Engineering simulator, e.g. Bullet, Project Chrono, HySys, Autodesk Civil Engineering. Let say if you have time you can create your own Autodesk instead of paying for its license, basically the brain behind them are derivative, integral, statistics and algebra.

Basically we choose C++ so we can create something long lasting in the future, and then I (DS Glanzsche) dedicate everyday of my life to eat and breathe C++ to be able to achieve this stage, hopefully it can be developed further till we can get practical use, like having a simulation for fluid flow, pharmacokinetics (to compute how drugs are metabolized by the body), simulation for human' organ, or to create a stable bridge or skyscraper, since the basic of calculus, integral, linear algebra, statistics can be covered nicely by SymIntegration. The front end looks for plot and simulation can be done by third party C++ library like ImGui, MatplotlibPlus or OpenGL.

Symbolic integration is basically harder than differentiation, differentiation has rigid rules: sum rule, product rule, chain rule, divide rule. While integration has only method of substitution and integration by parts, which back to the method itself. We can obtain different results for integration, but same result for differentiating different functions with respect to the same variable, like this case:

$$\begin{aligned}D_x \frac{-\cos^2(nx)}{2n} &= \cos(nx) \sin(nx) \\D_x \frac{\sin^2(nx)}{2n} &= \cos(nx) \sin(nx)\end{aligned}$$

Depending on the method of substitution, which one we substitute, if we reverse the equations above into integral problems, then we will obtain:

$$\begin{aligned}\int \cos(nx) \sin(nx) dx &= \frac{-\cos^2(nx)}{2n} + C \\ \int \cos(nx) \sin(nx) dx &= \frac{\sin^2(nx)}{2n} + C\end{aligned}$$




Figure 1.1: Integrating a function with different method can result in at least two different functions.

To compute the definite integral of a function in an interval you have to know how it behaves on the entire interval, and to compute the indefinite integral you have to know how it behaves on all intervals. While differentiation is a "local" operation, it computes the derivative of a function at a point you only have to know how it behaves in a neighborhood; distance, speed and acceleration from basic physics is the example of differentiation and integral. Computing integral / antiderivative often requires clever algebraic manipulation and technique and there are no universal rules for integration like there are for differentiation.

Integrals are fundamental in all science and engineering fields for solving problems involving area, volume, and other accumulations, some specific applications of integrals in science and engineering:

1. To calculate the area of irregular shapes, the volume of solids, and the surface area of objects.
2. To determine center of mass and moments of inertia for various shapes, they are crucial for structural analysis and design.
3. To calculate work done by a variable force, which is essential in mechanics and thermodynamics.
4. To calculate energy, power, and signal properties over time in electrical and computer engineering.
5. To determine stresses and strains analysis in structures under various loads.
6. To calculate fluid flow, pressure distributions, and forces exerted by fluids.

7. To compute probability distributions and statistical calculations.
8. To predict the trajectory of satellites and other objects in motion, ensuring they follow the intended path.
9. To calculate drug accumulation and half-life in a body, it is crucial for understanding how drugs are metabolized and eliminated by the body in Pharmacokinetics.
10. To model how diseases spread through a population with Susceptible-Infectious-Removed (SIR) model.
11. To understand better the complex processes of modeling tumor growth and metastasis.
12. To understand how reactions progress and their rates for reaction kinetics in chemistry.
13. To calculate work done during processes like gas expansion and changes in internal energy or enthalpy. These calculations are crucial for understanding the energetics of chemical reactions and processes.
14. To solve problems related to process design, optimization and understanding reaction kinetics in building a plant in Chemical Engineering.
15. To determine the volume of concrete needed for a structure, the area of a road surface, or the volume of soil required for an excavation in building bridges and structures.
16. To find the deflection (bending) and slope of a beam under load, which is crucial for ensuring structural stability.
17. To determine the velocity of a fluid flow, which is important in designing hydraulic structures like dams and pipelines.
18. To calculate the pressure distribution in fluids, which is essential for understanding the forces exerted on structure by fluids.
19. To understand the behavior of capacitors and inductors. The voltage across a capacitor is the integral of the current flowing through it, and the current through an inductor is the integral of the voltage across it.
20. To compute total charge and energy in Electrical Engineering.
21. To determine the circuit's voltage, current, or other parameters as a function of time by using integration to solve differential equations for many kind of circuit behaviors.

Now we will take a look at the applications of the integration in computer science:

1. Mathematics and computing.
2. Block chain method.
3. Quantum computing.
4. Big data analytics.
5. Neural Networking.
6. Artificial Intelligence.

7. Machine Learning.

Integration, calculus, linear algebra are the brain behind the applications above.

Triple integrals are used in physics, engineering, and even finance. If we get ahold of the C++, we can do the integral computation right, then it is only a matter of presentation, create the plot / simulation / computation for our goal, e.g. to create a stable bridge that can handle earthquake with magnitude of 10 SR. It takes simulation first, before constructing the real product.

The Operating System and Softwares in Use:

1. GFreya OS version 1.8 (based on LFS and BLFS version 11.0 System V books)

2. GCC version 11.2.0 (to compile C++ codes)

3. SymIntegration version 1.51

By June 2026 SymIntegration is able to compute

1. The integral of $\sin(ax + b)$, $\cos(ax + b)$, $\tan(ax + b)$, $\cot(ax + b)$, $\sec(ax + b)$, $\csc(ax + b)$ with $ax + b$ is a polynomial of order 1.
2. The integral of $\frac{1}{(ax+b)}$, $\frac{1}{ax^2+bx+c}$.
3. The integral and derivative of $\operatorname{asin}(ax + b)$, $\operatorname{acos}(ax + b)$, $\operatorname{atan}(ax + b)$, $\operatorname{acot}(ax + b)$, $\operatorname{asec}(ax + b)$, $\operatorname{acsc}(ax + b)$ with $ax + b$ is a polynomial of order 1.
4. The integral and derivative of all hyperbolic trigonometry functions $\sinh(ax + b)$, $\cosh(ax + b)$, $\tanh(ax + b)$, $\coth(ax + b)$, $\operatorname{sech}(ax + b)$, $\operatorname{csch}(ax + b)$.
5. The integral with the form of $x^b e^{ax}$, $x^b e^x$ with integration by parts method.
6. The integral of $\sin(ax) \cos(bx)$, $\cos(ax) \cos(bx)$, $\sin(ax) \sin(bx)$.
7. The integral of $\int \sin^n(x) dx$, $\int \cos^n(x) dx$, $\int \tan^n(x) dx$, $\int \sec^n(x) dx$, $\int \csc^n(x) dx$, but the speed to show the whole symbolic integral is slow compared to SymPy, needs to be fixed.
8. The solution ($y(t)$) of first order linear ordinary differential equation of type $ay' + ty = b$, separable equation of type $\frac{dy}{dx} = \frac{f(x)}{g(y)}$, and separable equation with homogeneous ratio equations of type $\frac{dy}{dx} = \frac{f(x,y)}{g(x,y)}$ along with the initial value problem solution.
9. The solution ($y(t)$) of second order linear ordinary differential equation of type $ay'' + by' + cy = g(t)$ (for homogeneous and non-homogeneous equation) along with the initial value problem solution.
10. The series solution of homogeneous second order linear ordinary differential equation of type $y'' + py' + qy = 0$ with constant coefficients, $p, q \in \mathbb{R}$.
11. The series solution of homogeneous n -th order linear ordinary differential equation of type $b_n y^{(n)} + b_{n-1} y^{(n-1)} + \dots + b_1 y' + b_0 y = 0$ with constant coefficients, $b_0, b_1, \dots, b_n \in \mathbb{R}$.
12. The series solution of homogeneous second order linear ordinary differential equation of type $P(x)y'' + Q(x)y' + R(x)y = 0$ with variable coefficients, $P(x)$, $Q(x)$, and $R(x)$ are polynomials.
13. The computation of divergence, gradient, and curl.
14. The probability mass function, mean, variance, moment generating function of discrete distributions.
15. The probability density function, cumulative distribution function, mean, variance, moment generating function of continuous distributions.
16. The regression line equation along with Pearson's correlation (r) with input of vector of 2 columns
17. Bond pricing and amortization table for mortgage loan or other credit related loan.
18. Compute direction fields and plot it with gnuplot.
19. Linear programming solution with Simplex Method.

-
20. Numerical differentiation and Richardson's Extrapolation.
 21. Matrices multiplication, addition, LU decomposition, SVD, eigenvalues and eigenvectors approximation, QR-decomposition, Cholesky decomposition, Schur decomposition, Hessenberg decomposition, QR Algorithm with shifts
 22. Generate random number that has certain distribution (uniform, gamma, beta, normal, F, chi-squared) and plot the histogram with Hamzstplot
 23. Compute one-way ANOVA, two-way ANOVA, three-way ANOVA, multiple regression model.
 24. Compute one-tailed and two-tailed hypothesis testing. One- and Two-Sample Estimation Problem, e.g. Confidence Interval and Prediction Interval.
 25. Compute cdf of Normal, student's t distribution, F distribution, chi-squared distribution, beta distribution, gamma distribution without Boost.
 26. Compute the quantile of standard normal distribution / inverse of cdf without Boost.
 27. Compute the classical orthogonal polynomials: Jacobi, Hermite, Gegenbauer, Chebyshev, Laguerre, Legendre.
 28. Transcription and translation from DNA to mRNA to polypeptide (Molecular biology).
 29. Genetic algorithms.
 30. Feed-forward neural network with steepest descent and conjugate gradient method.
 31. Extract RGBA matrices out of an image
 32. Perform kernel / convolution matrix on an image with OpenGL.
 33. Convolutional neural network for digit handwriting recognition with LeNet-5 architecture and steepest descent method.

We really appreciate those who spend time to write constructing critics, write opinions on what should be added in SymIntegration or if there is an incorrect C++ codes or wrong algorithm. Feedbacks can be sent to dsglanzsche@gmail.com.

Chapter 2

Symbolic Integral Computation and Calculus

"You can't direct the wind, but you can adjust your sails." - Sailing

W^E will first learn how to create the shared library for C++ library in a very simple and manual way, then we will see the source code that composing this C++ library, SymIntegration.

I. BUILD AND INSTALL SYMINTEGRATION IN GFREYA OS

We are using GFreya OS as it is a custom Linux-based Operating System, your distro your rules. Building applications from scratch instead of using package manager, you learn more from mistakes than just depending on package manager.

To download SymIntegration, open terminal / xterm then type:
git clone https://github.com/glanzkaiser/SymIntegration.git

Create Dynamic/Shared Library libsymintegration.so

Enter the directory of the cloned SymIntegration repository then type:
cd src
bash dynamiclibrary.sh
 this is a very manual way to create a dynamic library, or you can also type from the main parent directory of SymIntegration repository:

```
cd src
g++ -fPIC -c *.cpp
g++ -shared -o libsymintegration.so *.o
```

Code 1: Create dynamic library

It will create the dynamic library **libsymintegration.so** with a very manual way, instead of using CMake, we are using less space.

If you want to use Makefile to create the library, from **SymIntegration/src** open terminal/xterm then type:

make
mv libsymintegration.so /usr/lib

it will move the created shared / dynamic library to the default directory for library.

```
CXX = g++

SYMINTEGRATION_DIR = ../src
LIB = libsymintegration.so
SOURCES = array.cpp
SOURCES += artificialneuralnetworks.cpp cloning.cpp complexanalysis.cpp constants.
           cpp derive.cpp differentialequations.cpp divgradcurl.cpp dsolve.cpp
           dynamicalsystem.cpp
SOURCES += equation.cpp functions.cpp geneticalgorithms.cpp geometryandvectors.cpp
           geompack.cpp identity.cpp infiniteseries.cpp integrate.cpp linearalgebra.cpp
SOURCES += mathematicalfinance.cpp matnorm.cpp matrix.cpp multinomial.cpp
           navierstokes2d.cpp number.cpp numericalmethod.cpp
SOURCES += operationsresearch.cpp polynomial.cpp product.cpp quatern.cpp rational.
           cpp simplify.cpp
```



```
SOURCES += solve.cpp specialfunctions.cpp statistics.cpp stbimage.cpp sum.cpp
          symbol.cpp symbolic.cpp symerror.cpp
SOURCES += symintegrationc++.cpp symmatrix.cpp vecnorm.cpp
SOURCES += vector.cpp vectormatrixoperations.cpp verylong.cpp
SOURCES += asa241.cpp statistics-mle.cpp
SOURCES += biology-dna.cpp

OBJS = $(addsuffix .o, $(basename $(notdir $(SOURCES))))
UNAME_S := $(shell uname -s)

CXXFLAGS = -fPIC
CXXFLAGS += -g -Wall -Wformat

## BUILD FLAGS PER PLATFORM

ifeq ($(UNAME_S), Linux) #LINUX
ECHO_MESSAGE = "SymIntegration from GFreyra OS"

CFLAGS = $(CXXFLAGS)
endif

## BUILD RULES

%.o:$(SYMINTEGRATION_DIR)/%.cpp
$(CXX) $(CXXFLAGS) -c -o $@ $<

all: $(LIB)
@echo Build complete for $(ECHO_MESSAGE)

$(LIB): $(OBJS)
$(CXX) -shared -o $@ $^ $(CXXFLAGS)

clean:
rm -f $(OBJS)
```

Code 2: *src/Makefile*

Assuming you are using Linux-based OS, after finish compiling don't forget to copy / move the dynamic library to **/usr/lib**.

Move the Include Files

After finish compiling the dynamic library, open terminal from the current working directory / this repository main directory:

```
cd include
cp -r * /usr/include
```

Code 3: *Move include folder*

When we want to create the shared library it will look for this header files in the default path where the include files are usually located. In Linux OS **usr/include** is the basic / default path.

All files, examples codes and source codes used in this books can be found in this repository:
<https://github.com/glanzkaiser/SymIntegration>

We are using less, minimal amount of code to create the library, if you compare it with the original SymbolicC++ that has **.configure** and **Makefile** that will be spawn after you configure it, we give you two options here to compile the library.

Two simple commands with **g++** can already make a shared symbolic integration computation library, with limitation still.

If you are using Windows OS or MacOS then you are on your own.

The C++ codes are only located in the **/src** directory , it is designed to contain all the **' .cpp '** files, then inside the **/include/** we have header files and another folder **/include/symintegral/** is also a folder that contain header files too.

You can open them one by one, read them and making sense of it based on the function that you are looking for.

In the beginning (from SymbolicC++ 3.35), there are total of 27 **.cpp** files and 26 **.h** / header files. So it won't take a long time for someone to learn it.

II. TECHNIQUES OF INTEGRATION

We will cover some techniques of integration and also two Fundamental Theorems of Calculus that are very useful in solving integral problems or even differential equations.

Theorem 2.1: The First Fundamental Theorem of Calculus

Let f be continuous on the closed interval $[a, b]$ and let x be a (variable) point in (a, b) . Then

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

Theorem 2.2: The Second Fundamental Theorem of Calculus

Let f be continuous (hence integrable) on $[a, b]$, and let F be any antiderivative of f on $[a, b]$. Then

$$\int_a^b f(x) dx = F(b) - F(a)$$

[SI*] Integration by Parts

From the analogue of the product rule for differentiation. Suppose we have two functions $f(x)$ and $g(x)$ with

$$\frac{d}{dx} f(x)g(x) = f(x) \frac{dg(x)}{dx} + g(x) \frac{df(x)}{dx} \quad (2.1)$$

Integrating both sides and rearranging the terms, we will get the integration by parts formula:

$$\int f(x) \frac{dg(x)}{dx} dx = f(x)g(x) - \int g(x) \frac{df(x)}{dx} dx \quad (2.2)$$

This formula is only useful if $\int g(x) \frac{df(x)}{dx} dx$ or $\int f(x) \frac{dg(x)}{dx} dx$ is easier to integrate than $\int f(x)g(x) dx$.

The application for integration by part is very useful in statistics, e.g. to be able to compute the mean and variance for exponential distribution.

[SI*] Substitution Rule for Indefinite Integral

The substitution rule is nothing more than the Chain Rule in reverse.

Theorem 2.3: Substitution Rule for Indefinite Integrals

Let g be a differentiable function and suppose that F is an antiderivative of f . Then

$$\int f(g(x))g'(x) dx = F(g(x)) + C$$

Theorem 2.4: Substitution Rule for Definite Integrals

Let g have a continuous derivative on $[a, b]$, and let f be continuous on the range of g . Then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

where $u = g(x)$.

III. INTEGRATION BY PARTS

[SI*] **Example: Exponential Distribution**

The exponential distribution, which is sometimes used to model the lifetimes of electrical or mechanical components, has probability density function

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } 0 \leq x \\ 0, & \text{otherwise} \end{cases}$$

where λ is some positive constant. Compute the mean μ and the variance σ^2 .

Solution:

To compute for the mean we will have

$$\begin{aligned} E(X) &= \int_{-\infty}^{\infty} x f(x) dx \\ &= \int_{-\infty}^0 x \cdot 0 dx + \int_0^{\infty} x \lambda e^{-\lambda x} dx \end{aligned}$$

We apply integration by parts in the second integral with

$$\begin{aligned} u &= x \\ dv &= \lambda e^{-\lambda x} dx \\ du &= dx \\ v &= -e^{-\lambda x} \end{aligned}$$

Thus

$$\begin{aligned} E(X) &= [-x \lambda e^{-\lambda x}]_0^{\infty} - \int_0^{\infty} (-e^{-\lambda x}) dx \\ &= (-0 + 0) + \left[-\frac{1}{\lambda} e^{-\lambda x} \right]_0^{\infty} \\ &= \frac{1}{\lambda} \end{aligned}$$

The variance is

$$\begin{aligned} \sigma^2 &= E(X^2) - \mu^2 \\ &= \int_{-\infty}^{\infty} x^2 f(x) dx - \left(\frac{1}{\lambda} \right)^2 \\ &= \int_{-\infty}^0 x^2 \cdot 0 dx + \int_0^{\infty} x^2 \lambda e^{-\lambda x} dx - \frac{1}{\lambda^2} \\ &= [-x^2 e^{-\lambda x}]_0^{\infty} - \int_0^{\infty} (-e^{-\lambda x}) 2x dx - \frac{1}{\lambda^2} \\ &= (-0 + 0) + 2 \int_0^{\infty} x e^{-\lambda x} dx - \frac{1}{\lambda^2} \\ &= 2 \frac{1}{\lambda^2} - \frac{1}{\lambda^2} \\ &= \frac{1}{\lambda^2} \end{aligned}$$

[SI*] When they say integration is an art it is true, you need more practice to be able to handle all kinds of integral problems.

If we are facing this kind of equation:

$$f(x) = x\lambda e^{-\lambda x}$$

then what we need to do is learning the pattern

$$\begin{aligned}\int x e^x dx &= (x - 1)e^x + C \\ \int x^2 e^x dx &= (x^2 - 2x + 2)e^x + C \\ \int x^3 e^x dx &= (x^3 - 3x^2 + 6x - 6)e^x + C \\ \int x^4 e^x dx &= (x^4 - 4x^3 + 12x^2 - 24x + 24)e^x + C \\ &\vdots\end{aligned}$$

so for the general function of

$$f(x) = x^n e^x$$

The algorithm will be like this:

```
Symbolic sgn = 1;
int k = 1;
Symbolic integral;
for (int i = 0; i <= n; i++)
{
    integral += k * sgn * (x^(n-i)) * exp(x)
    sgn = -sgn;
    k *= n-i;
}
return integral;
```

The C++ file to handle cases of integration by parts is located in **src/integrate.cpp**

```
Symbolic integrate(const Symbolic &f,const Symbolic &x)
{
    list<Equations> eq;
    list<Equations>::iterator i;
    UniqueSymbol a, b, c;
    ...
    ...
```

Code 4: *src/integrate.cpp*

Now for the case of $f(x) = x^n e^x$, it is handled at this part of codes

```
Symbolic integrate(const Symbolic &f,const Symbolic &x)
{
    ...
```

```

...
eq = ((x^b)*exp(x)).match(f, (a,b)); // case for x^b *
    exp(x)
for(i=eq.begin(); i!=eq.end(); ++i)
try {
    Symbolic bp = rhs(*i, b);
    if(bp.type() == typeid(Numeric)
    && Number<void>(bp)->numerictype() == typeid(int)
    && Number<void>(bp)>Number<int>(0))
    {
        int n = CastPtr<const Number<int>>(bp)->n,
            sgn = 1;
        Symbolic integral, nf = 1;
        for(; n>=0; nf*=n, --n, sgn=-sgn)
            integral += sgn*nf*(x^n)*exp(x);
        return integral;
    }
} catch(const SymbolicError &se) {}
}

```

Code 5: *src/integrate.cpp* to handle $x^n e^x$

For the computation of mean and variance of exponential distribution with SymIntegration we will get this raw result, where the e^{-inf} is still as it is, and not returning to 0, we will work on it in the future.

```

#include <iostream>
#include "symintegrationc++.h"

using namespace std;

int main(void)
{
    Symbolic x("x"), l("1"), Inf("Inf"), y, Ex, Varx,
        u, dv, v, du;

    y = l*exp(-l*x);
    u = x;
    dv = l*exp(-l*x);
    du = df(u,x);
    v = integrate(dv,x);

    cout << "f(x) = " << y << endl;

    for(int i=1;i<=1;i++)
    {
        y = integrate(y,x);
        cout << i << "-st integral of f(x) = " <<
            y << endl;
    }
}

```

```

    }
    cout << "int_{0}^{Inf} f(x) = " << y[x==Inf] - y[
        x==0] << endl;

    Ex = x*l*exp(-l*x);

    cout << "\nx f(x) = " << Ex << endl;
    cout << "\nu = " << u << endl;
    cout << "dv = " << dv << endl;
    cout << "du = " << du << endl;
    cout << "v = " << v << endl;

    Ex = integrate(Ex,x);
    cout << "\nE(x) = int x f(x) = " << Ex << endl;
    cout << "int_{0}^{Inf} x f(x) = " << Ex[x==Inf] -
        Ex[x==0] << endl;

    Varx = x*x*l*exp(-l*x);
    Varx = integrate(Varx,x);

    cout << "\nint x^{2} f(x) = " << Varx << endl;
    cout << "int_{0}^{Inf} x^{2} f(x) = " << Varx[x==
        INFINITY, l==1] - Varx[x==0] << endl;
    cout << "\nVar(x) = int x^{2} f(x) - mu^{2} = "
        << Varx[x==INFINITY, l==1] - Varx[x==0] -
        (1/l^(2)) << endl;

    return 0;
}

```

Code 6: Integration by Parts

IV. THE METHOD OF SUBSTITUTION

[SI*] Some Examples

Suppose we want to compute the integral

$$\int x \sin x^2 dx$$

Solution:

Here the appropriate substitution is

$$\begin{aligned}
 u &= x^2 \\
 du &= 2x dx
 \end{aligned}$$

Thus

$$\begin{aligned}\int x \sin x^2 dx &= \int \frac{1}{2} \sin x^2 (2x dx) \\&= \int \frac{1}{2} \sin u du \\&= -\frac{1}{2} \cos u + C \\&= -\frac{1}{2} \cos x^2 + C\end{aligned}$$

[SI*] In SymIntegration the method of substitution can be handled in **src/integrate.cpp**, we can create a pattern of equation to be solved, just like the integration by parts method.

V. FRACTION INTEGRAL

[SI*] Fraction Level 1

Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{1}{ax + b}$$

a and b are constants.

The integral will be

$$\int \frac{1}{ax + b} dx = \frac{\ln(ax + b)}{a} + C \quad (2.3)$$

For the integrate function alone in SymIntegration we do not include the constant C that is obtained after integration.

* Note that the C of the indefinite integration cancels out, as it always will, in the definite integration. That is why in the statement of the Second Fundamental Theorem of Calculus we could use the phrase **any derivative**. In particular, we may always choose $C = 0$ in applying the Second Fundamental Theorem of Calculus [23].

[SI*] The C++ code to handle this fraction level 1 integral problem to make sure the integral will return the correct result is located in `src/functions.cpp` under the function **Symbolic Power::integrate(const Symbolic &s) const**, we still use the function **integrate** for this type of integral.

```
Symbolic Power::integrate(const Symbolic &s) const
{
    const Symbolic &a = parameters.front();
    const Symbolic &b = parameters.back();
    ...
    ...
}
```

```
Symbolic Power::integrate(const Symbolic &s) const
{
    ...
    ...

    if(b == -1 && parameters.front().coeff(s,2) == 0) return
        ln(parameters.front()) * (1/ parameters.front().df(
            s));
    ...
    ...
}
```

Code 7: fraction level 1 integral

The **if** statement above is a condition: if the input of a function meets the criteria that the power is -1 / a fraction ($f(x)^{-1} = \frac{1}{f(x)}$) then we will return it as **ln(parameters.front()) * (1/**

parameters.front().df(s)). The symbolic s is used to replace the variable in the integration problem.

I will introduce to the reader here about **parameters.front()** and **parameters.back()**, suppose we have a function

$$f(x) = (ax^2 + bx + c)^d \quad (2.4)$$

with a, b, c as constants.

- The **parameters.front()** of the function from equation (2.4) is $ax^2 + bx + c$.
parameters.front().coeff(s,2) is a .
parameters.front().coeff(s,1) is b .
parameters.front().coeff(s,0) is c .
- The **parameters.back()** of the function from equation (2.4) is d .

With this knowledge we can create a lot of **if, if.. else..** conditions to make the integration computation complete and better.

[SI*] Fraction Level 2

Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{1}{ax^2 + bx + c}$$

a, b and c are constants.

The integral will be

$$\begin{aligned} \int \frac{1}{ax^2 + bx + c} dx = & -\sqrt{\frac{-1}{4ac - b^2}} \ln \left(x + \frac{-4ac\sqrt{\frac{-1}{4ac - b^2}} + b^2\sqrt{\frac{-1}{4ac - b^2}} + b}{2a} \right) \\ & + \sqrt{\frac{-1}{4ac - b^2}} \ln \left(x + \frac{4ac\sqrt{\frac{-1}{4ac - b^2}} - b^2\sqrt{\frac{-1}{4ac - b^2}} + b}{2a} \right) + C \end{aligned} \quad (2.5)$$

The codes for the SymIntegration to handle this integral is located in **src/functions.cpp**

```
...
...

Symbolic Power::integrate(const Symbolic &s) const
{
    ...
    ...
    if(b == -1 && parameters.front().coeff(s,2) != 0)
    {
        double a1 = parameters.front().coeff(s,2);
        double b1 = parameters.front().coeff(s,1);
        double c1 = parameters.front().coeff(s,0);
        double D_inv = sqrt(-1/(4*a1*c1-b1*b1));
```

```

        return - D_inv * ln(s + (-4*a1*c1*D_inv + b1*b1*
            D_inv + b1)/(2*a1)) + D_inv * ln(s + (4*a1*c1
            *D_inv - b1*b1*D_inv + b1)/(2*a1)) ;
    }
    ...
    ...
}
...
...

```

Code 8: fraction level 2 integral type 1

[SI*] Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1 x}{ax^2 + bx + c}$$

a_1, a, b , and c are constants.

The integral will be

$$\begin{aligned}
 \int \frac{a_1 x}{ax^2 + bx + c} dx = a_1 & \left[\left(-\frac{b\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} + \frac{1}{2a} \right) * \right. \\
 & \ln \left(x + \frac{-4ac \left(-\frac{b\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} + \frac{1}{2a} \right) + b^2 \left(-\frac{b\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} + \frac{1}{2a} \right) + 2c}{b} \right) \\
 & + \left(\frac{b\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} + \frac{1}{2a} \right) * \\
 & \left. \ln \left(x + \frac{-4ac \left(\frac{b\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} + \frac{1}{2a} \right) + b^2 \left(\frac{b\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} + \frac{1}{2a} \right) + 2c}{b} \right) \right] + C
 \end{aligned} \tag{2.6}$$

The codes for the SymIntegration to handle this integral is located in **src/integrate.cpp**, instead of **integrate** we create a new function **fraction integrate** to handle this type of integral because this type of integral cannot be read nicely with **Symbolic Power::integrate(const Symbolic &s) const** that is in **src/functions.cpp**.

By July 8th, 2025, the function **fraction integrate** already has around 190 lines of code.

```

    ...
    ...

Symbolic fractionintegrate(const Symbolic &fnum, const Symbolic &fdenom,
    const Symbolic &x)
{
    list<Equations> eq;
    list<Equations>::iterator i;
    UniqueSymbol anum, bnum;
    Symbolic integral_sol;
    double anum, bnum, ap, bp, cp;

```

```

    ap = fdenom.coeff(x,2);
    bp = fdenom.coeff(x,1);
    cp = fdenom.coeff(x,0);

    ...
    ...

    if (fnum.coeff(x,2)==0 && fnum.coeff(x,1)!=0 && fnum.coeff(x,0)==0 &&
        fdenom.coeff(x,2)!=0 && fdenom.coeff(x,1)!=0 && fdenom.coeff(x,0)
        !=0 ) // for (a1x)/(ax^2+bx+c)
    {
        eq = ( (bnum*x)).match(fnum, (bnum,cnum));
        for(i=eq.begin(); i!=eq.end(); ++i)
        try {
            Symbolic a1 = rhs(*i, bnum);
            bnum= a1;
        } catch(const SymbolicError &se) {}

        double D = (bp*sqrt(bp*bp-4*ap*cp))/(2*ap*(4*ap*cp-(bp*bp)))
            ;
        integral_sol = bnum*( (-D + (1/(2*ap)))*ln(x+(-4*ap*cp*(-D +
            1/(2*ap)) +bp*bp*(-D+ 1/(2*ap)) + 2*cp)/(bp)) + (D +
            (1/(2*ap)))*ln(x+(-4*ap*cp*(D + 1/(2*ap)) +bp*bp*(D+ 1/(2*
            ap)) + 2*cp)/(bp)) );
    }

    ...

    return integral_sol;
}

...
...

```

Code 9: fraction level 2 integral default type

[SI*] Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1 x}{ax^2 + bx}$$

a_1, a , and b are constants.

The integral will be

$$\int \frac{a_1 x}{ax^2 + bx} dx = \frac{a_1 \ln(ax + b)}{a} + C \quad (2.7)$$

[SI*] Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1 x}{ax^2 + c}$$

a_1, a and c are constants.

The integral will be

$$\int \frac{a_1 x}{ax^2 + c} dx = \frac{a_1 \ln(ax^2 + c)}{2a} + C \quad (2.8)$$

[SI*] Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1 x}{bx + c}$$

a_1, b , and c are constants.

The integral will be

$$\int \frac{a_1 x}{bx + c} dx = a_1 \left(\frac{x}{b} - \frac{c \ln(bx + c)}{b^2} \right) + C \quad (2.9)$$

[SI*] **Fraction Level 3**

Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1 x + b_1}{ax^2 + bx + c}$$

a_1, b_1, a, b and c are constants.

The integral will be

$$\begin{aligned} \int \frac{a_1 x + b_1}{ax^2 + bx + c} dx = & \left(\frac{a_1}{2a} - \frac{(2ab_1 - a_1b)\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} \right) * \\ & \ln \left(x + \frac{4ac \left(\frac{a_1}{2a} - \frac{(2ab_1 - a_1b)\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} \right) - 2a_1c - b^2 \left(\frac{a_1}{2a} - \frac{(2ab_1 - a_1b)\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} \right) + bb_1}{2ab_1 - a_1b} \right) \\ & + \left(\frac{a_1}{2a} + \frac{(2ab_1 - a_1b)\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} \right) * \\ & \ln \left(x + \frac{4ac \left(\frac{a_1}{2a} + \frac{(2ab_1 - a_1b)\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} \right) - 2a_1c - b^2 \left(\frac{a_1}{2a} + \frac{(2ab_1 - a_1b)\sqrt{b^2 - 4ac}}{2a(4ac - b^2)} \right) + bb_1}{2ab_1 - a_1b} \right) + C \end{aligned} \quad (2.10)$$

The codes for the SymIntegration to handle this integral is located in **src/integrate.cpp**, instead of **integrate** we create a new function **fraction integrate** to handle this type of integral.

```
...

Symbolic fractionintegrate(const Symbolic &fnum, const Symbolic &fdenom,
    const Symbolic &x)
```

```

{
    list<Equations> eq;
    list<Equations>::iterator i;
    UniqueSymbol anum, bnum;
    Symbolic integral_sol;
    double anump, bnump, ap, bp, cp;
    ap = fdenom.coeff(x,2);
    bp = fdenom.coeff(x,1);
    cp = fdenom.coeff(x,0);

    ...

    if (fnum.coeff(x,2)==0 && fnum.coeff(x,1)!=0 && fnum.coeff(x,0)!=0 &&
        fdenom.coeff(x,2)!=0 && fdenom.coeff(x,1)!=0 && fdenom.coeff(x,0)
        !=0 ) // for (a1x+b1)/(ax^2+bx+c)
    {
        eq = ( (bnum*x+cnum)).match(fnum, (bnum,cnum));
        for(i=eq.begin(); i!=eq.end(); ++i)
            try {
                Symbolic a1 = rhs(*i, bnum), b1 = rhs(*i, cnum);
                bnump= a1;
                cnump = b1;
            } catch(const SymbolicError &se) {}

        double D = (2*ap*cnump - bnump*bp)*sqrt(-4*ap*cp+(bp*bp))/(2*
            ap*(4*ap*cp-(bp*bp)));
        double denom1 = 2*ap*cnump - bnump*bp;

        integral_sol = ((bnump/(2*ap)) - D)*ln(x + (4*ap*cp*((bnump
            /(2*ap))-D) - 2*bnump*cp -(bp*bp)*(bnump/(2*ap) - D) +
            bp*cnump)/(denom1)) + ((bnump/(2*ap)) + D)*ln(x + (4*ap*cp
            *((bnump/(2*ap))+D) - 2*bnump*cp -(bp*bp)*(bnump/(2*ap) +
            D) + bp*cnump)/(denom1)) ;
    }

    ...

    return integral_sol;
}

...

```

Code 10: *fraction level 3 integral default type showstringspaces*

[SI*] Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1x + b_1}{ax^2 + bx}$$

a_1, b_1, a , and b are constants.

The integral will be

$$\int \frac{a_1x + b_1}{ax^2 + bx} dx = \frac{b_1 \ln(x)}{b} - \frac{(ab_1 - a_1b) \ln\left(x + \frac{bb_1 + \frac{b(ab_1 - a_1b)}{2ab_1 - a_1b}}{ab}\right)}{ab} + C \quad (2.11)$$

[SI*] Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1x + b_1}{ax^2 + c}$$

a_1, b_1, a , and c are constants.

The integral will be

$$\begin{aligned} \int \frac{a_1x + b_1}{ax^2 + c} dx &= \left(\frac{a_1}{2a} - \frac{b_1\sqrt{-a^3c}}{2a^2c} \right) * \\ &\ln \left(x + \frac{2ac \left(\frac{a_1}{2a} - \frac{b_1\sqrt{-a^3c}}{2a^2c} \right) - a_1c}{ab_1} \right) \\ &+ \left(\frac{a_1}{2a} + \frac{b_1\sqrt{-a^3c}}{2a^2c} \right) * \\ &\ln \left(x + \frac{2ac \left(\frac{a_1}{2a} + \frac{b_1\sqrt{-a^3c}}{2a^2c} \right) - a_1c}{ab_1} \right) + C \end{aligned} \quad (2.12)$$

[SI*] Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1x + b_1}{bx + c}$$

a_1, b_1, b , and c are constants.

The integral will be

$$\int \frac{a_1x + b_1}{bx + c} dx = \frac{a_1x}{b} - \frac{(a_1c - bb_1) \ln(bx + c)}{b^2} + C \quad (2.13)$$

[SI*] **Fraction Level 4**

Suppose we have this function that we want to integrate toward x .

$$f(x) = \frac{a_1x^2 + b_1x + c_1}{ax^2 + bx + c}$$

a_1, b_1, c_1, a, b and c are constants.

The integral will be

$$\begin{aligned}
\int \frac{a_1 x^2 + b_1 x + c_1}{ax^2 + bx + c} dx = & \left(\frac{ab_1 - a_1 b}{2a^2} - \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right) * \\
& \ln \left[x + \frac{4a^2 c \left(\frac{ab_1 - a_1 b}{2a^2} - \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right)}{2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2} \right. \\
& \left. - \frac{ab^2 \left(\frac{ab_1 - a_1 b}{2a^2} - \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right) + abc_1 - 2ab_1 c + a_1 bc}{2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2} \right] \\
& + \left(\frac{ab_1 - a_1 b}{2a^2} + \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right) * \\
& \ln \left[x + \frac{4a^2 c \left(\frac{ab_1 - a_1 b}{2a^2} + \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right)}{2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2} \right. \\
& \left. - \frac{ab^2 \left(\frac{ab_1 - a_1 b}{2a^2} + \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right) + abc_1 - 2ab_1 c + a_1 bc}{2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2} \right] \\
& + \frac{a_1 x}{a} + C
\end{aligned} \tag{2.14}$$

The codes for the SymIntegration to handle this integral is located in **src/integrate.cpp**, we use a new function **fraction integrate** to handle this type of integral.

```

...

Symbolic fractionintegrate(const Symbolic &fnum, const Symbolic &fdenom,
    const Symbolic &x)
{
    list<Equations> eq;
    list<Equations>::iterator i;
    UniqueSymbol anum, bnum, cnum;
    Symbolic integral_sol;
    double anump, bnump, cnump, ap, bp, cp;
    ap = fdenom.coeff(x,2);
    bp = fdenom.coeff(x,1);
    cp = fdenom.coeff(x,0);

    if (fnum.coeff(x,2)!=0 && fnum.coeff(x,1)!=0 && fnum.coeff(x,0)!=0 &&
        fdenom.coeff(x,2)!=0 && fdenom.coeff(x,1)!=0 && fdenom.coeff(x,0)
        !=0 ) // for (a1x^2 + b1x + c1)/(ax^2+bx+c)
    {
        eq = ( (anum*x*x+bnum*x+cnum)).match(fnum, (anum,bnum,cnum));
        for(i=eq.begin(); i!=eq.end(); ++i)
        try {
            Symbolic a1 = rhs(*i, anum), b1 = rhs(*i, bnum), c1 =
                rhs(*i, cnum);

```

```

        anum= a1;
        bnum= b1;
        cnum = c1;
    } catch(const SymbolicError &se) {}

    double D = sqrt(bp*bp-(4*ap*cp))*(2*ap*ap*cnum-2*ap*anum*cp
        -ap*bp*bnum+anum*bp*bp)/(2*ap*ap*(4*ap*cp-(bp*bp))) ;

    integral_sol = ((ap*bnum-anum*bp)/(2*ap*ap) - D)*ln(x+(4*ap
        *ap*cp*((ap*bnum-anum*bp)/(2*ap*ap) - D) - ap*bp*bp*((
        ap*bnum-anum*bp)/(2*ap*ap) - D) + ap*bp*cnum-2*ap*
        bnum*cp+anum*bp*cp) / (2*ap*ap*cnum-2*ap*anum*cp-ap*
        bp*bnum+anum*bp*bp)) + ((ap*bnum-anum*bp)/(2*ap*ap) +
        D)*ln(x+(4*ap*ap*cp*((ap*bnum-anum*bp)/(2*ap*ap) + D) -
        ap*bp*bp*((ap*bnum-anum*bp)/(2*ap*ap) + D) + ap*bp*
        cnum-2*ap*bnum*cp+anum*bp*cp) / (2*ap*ap*cnum-2*ap*
        anum*cp-ap*bp*bnum+anum*bp*bp)) + (anum*x)/ap;

    }

    ...
    ...

```

Code 11: fraction level 4 integral default type

[SI*] The key to solve integral of fraction integral is to use the method of partial fraction decomposition, which allows us to decompose rational functions into sums of simpler, more easily integrated rational functions, for example if we have

$$\int \frac{P_n(x)}{P_m(x)}$$

with $P_n(x)$ is a polynomial with x as the independent variable and highest order of n , $P_m(x)$ is a polynomial with x as the independent variable and highest order of m .

The partial fraction decomposition can be applied to a rational function

$$\frac{P_n(x)}{P_m(x)}$$

only if $n < m$.

For the case where $n \geq m$, it can be tricked by first we must perform long division to rewrite the quotient into a rational function that can fit the bill to be performed the partial fraction decomposition

$$\frac{P_n(x)}{P_m(x)} = B(x) + \frac{P_c(x)}{P_d(x)}$$

where $c < d$.

From the previous part we can see that a solid formula to compute the fraction integral can be very long enough, with $m = 2$, it comes from deriving and generalizing this method of partial fraction decomposition by changing the constants into coefficients like a_1, b_1, c_1, a, b, c .

[SI*] **List of Solvable Fraction Integral**

We will list all the fractions that can be solved with **fractionintegrate** function in SymIntegration here, they are:

$$\begin{aligned}
 \int \frac{1}{1+x^2} dx &= \text{atan}(x) \\
 \int \frac{ax}{1+x^2} dx &= \frac{a \ln(x^2+1)}{2} \\
 \int \frac{1}{ax^2+bx+c} dx &= -\sqrt{\frac{-1}{4ac-b^2}} \ln \left(x + \frac{-4ac\sqrt{\frac{-1}{4ac-b^2}} + b^2\sqrt{\frac{-1}{4ac-b^2}} + b}{2a} \right) \\
 &\quad + \sqrt{\frac{-1}{4ac-b^2}} \ln \left(x + \frac{4ac\sqrt{\frac{-1}{4ac-b^2}} - b^2\sqrt{\frac{-1}{4ac-b^2}} + b}{2a} \right) + C \\
 \int \frac{a_1x}{ax^2+bx+c} dx &= a_1 \left[\left(-\frac{b\sqrt{b^2-4ac}}{2a(4ac-b^2)} + \frac{1}{2a} \right) * \right. \\
 &\quad \ln \left(x + \frac{-4ac \left(-\frac{b\sqrt{b^2-4ac}}{2a(4ac-b^2)} + \frac{1}{2a} \right) + b^2 \left(-\frac{b\sqrt{b^2-4ac}}{2a(4ac-b^2)} + \frac{1}{2a} \right) + 2c}{b} \right) \\
 &\quad + \left(\frac{b\sqrt{b^2-4ac}}{2a(4ac-b^2)} + \frac{1}{2a} \right) * \\
 &\quad \left. \ln \left(x + \frac{-4ac \left(\frac{b\sqrt{b^2-4ac}}{2a(4ac-b^2)} + \frac{1}{2a} \right) + b^2 \left(\frac{b\sqrt{b^2-4ac}}{2a(4ac-b^2)} + \frac{1}{2a} \right) + 2c}{b} \right) \right] + C \\
 \int \frac{a_1x+b_1}{ax^2+bx+c} dx &= \left(\frac{a_1}{2a} - \frac{(2ab_1-a_1b)\sqrt{b^2-4ac}}{2a(4ac-b^2)} \right) * \\
 &\quad \ln \left(x + \frac{4ac \left(\frac{a_1}{2a} - \frac{(2ab_1-a_1b)\sqrt{b^2-4ac}}{2a(4ac-b^2)} \right) - 2a_1c - b^2 \left(\frac{a_1}{2a} - \frac{(2ab_1-a_1b)\sqrt{b^2-4ac}}{2a(4ac-b^2)} \right) + bb_1}{2ab_1-a_1b} \right) \\
 &\quad + \left(\frac{a_1}{2a} + \frac{(2ab_1-a_1b)\sqrt{b^2-4ac}}{2a(4ac-b^2)} \right) * \\
 &\quad \ln \left(x + \frac{4ac \left(\frac{a_1}{2a} + \frac{(2ab_1-a_1b)\sqrt{b^2-4ac}}{2a(4ac-b^2)} \right) - 2a_1c - b^2 \left(\frac{a_1}{2a} + \frac{(2ab_1-a_1b)\sqrt{b^2-4ac}}{2a(4ac-b^2)} \right) + bb_1}{2ab_1-a_1b} \right) + C
 \end{aligned}$$

$$\begin{aligned}
 \int \frac{a_1 x^2 + b_1 x + c_1}{ax^2 + bx + c} dx = & \left(\frac{ab_1 - a_1 b}{2a^2} - \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right) * \\
 & \ln \left[x + \frac{4a^2 c \left(\frac{ab_1 - a_1 b}{2a^2} - \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right)}{2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2} \right. \\
 & \left. - \frac{ab^2 \left(\frac{ab_1 - a_1 b}{2a^2} - \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right) + abc_1 - 2ab_1 c + a_1 bc}{2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2} \right] \\
 & + \left(\frac{ab_1 - a_1 b}{2a^2} + \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right) * \\
 & \ln \left[x + \frac{4a^2 c \left(\frac{ab_1 - a_1 b}{2a^2} + \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right)}{2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2} \right. \\
 & \left. - \frac{ab^2 \left(\frac{ab_1 - a_1 b}{2a^2} + \frac{(2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2) \sqrt{b^2 - 4ac}}{2a^2(4ac - b^2)} \right) + abc_1 - 2ab_1 c + a_1 bc}{2a^2 c_1 - 2aa_1 c - abb_1 + a_1 b^2} \right] \\
 & + \frac{a_1 x}{a} + C
 \end{aligned}$$

If the denominator has determinant of 0 / $b^2 = 4ac$ then

$$\int \frac{b_1 x + c_1}{ax^2 + bx + c} dx = \frac{-ac_1 + b_1 c}{a^3 x + a^2 c} + \frac{b_1 \ln(ax + c)}{a^2}$$

VI. TRIGONOMETRY INTEGRAL

[SI*] Trigonometry Level 1

We will want to compute integral of trigonometry at the basic level like these:

$$\begin{aligned}\int \sin(x) dx &= -\cos(x) + C \\ \int \sin(3x+5) dx &= -\frac{1}{3}\cos(3x+5) + C \\ \int \cos(x) dx &= \sin(x) + C \\ \int \cos(3x+5) dx &= \frac{1}{3}\sin(3x+5) + C \\ \int \tan(x) dx &= -\ln(\cos(x)) + C \\ \int \tan(3x+5) dx &= \frac{1}{6}\ln(\tan^2(3x+5)+1) + C \\ \int \cot(x) dx &= \ln(\sin(x)) + C \\ \int \cot(3x+5) dx &= -\frac{1}{6}\ln(\tan^2(3x+5)+1) + \frac{1}{3}\ln(\tan(3x+5)) + C \\ \int \sec(x) dx &= -\frac{\ln(\cos(x))}{\sin(x)} + C \\ \int \sec(3x+5) dx &= -\frac{1}{3}\frac{\ln(\cos(3x+5))}{\sin(3x+5)} \\ \int \csc(x) dx &= \frac{\ln(\sin(x))}{\cos(x)} + C \\ \int \csc(3x+5) dx &= \frac{1}{3}\frac{\ln(\sin(3x+5))}{\cos(3x+5)}\end{aligned}$$

or in general form

$$\begin{aligned}\int \sin(ax+b) dx &= -\frac{1}{a}\cos(ax+b) + C \\ \int \cos(ax+b) dx &= \frac{1}{a}\sin(ax+b) + C \\ \int \tan(ax+b) dx &= \frac{1}{2a}\ln(\tan^2(ax+b)+1) + C \\ \int \cot(ax+b) dx &= -\frac{1}{2a}\ln(\tan^2(ax+b)+1) + \frac{1}{a}\ln(\tan(ax+b)) + C \\ \int \sec(ax+b) dx &= -\frac{1}{a}\frac{\ln(\cos(ax+b))}{\sin(ax+b)} \\ \int \csc(ax+b) dx &= \frac{1}{a}\frac{\ln(\sin(ax+b))}{\cos(ax+b)}\end{aligned}$$

At first, the raw codes from SymbolicC++ 3.35 cannot perform that computation, in the **src/functions.cpp** there are implementations for sin, cos, tan but they are imperfect and if the case is not covered it will has an output of **return Integral(*this,s)**, which mean the integral cannot be computed.

So we correct it and put the formula so SymIntegration can compute the integral problems above, which we call them as Level 1 trigonometry problems.

```

...
...

Symbolic Tan::integrate(const Symbolic &s) const
{
    const Symbolic &x = parameters.front();
    if(x == s) return -ln(cos(x)) * (1 / parameters.front().
        df(s));
    if(df(s) == 0) return *this * s;
    return ln( tan(parameters.front()) * tan(parameters.
        front() + 1) * ( 1 / (2*parameters.front().df(s)) )
        ;
}

...

```

Code 12: Implementation of integral for tangent function

```

170
171 Symbolic Acos::integrate(const Symbolic &s) const
172 {
173     const Symbolic &x = parameters.front();
174     if(x == s) return x*acos(x) - sqrt(1-x*x);
175     if(df(s) == 0) return *this * s;
176     return ( - sqrt(1-(parameters.front())*(parameters.front())) / parameters.front().df(s) ) + ( parameters.front()*acos(parameters.front()) ,
177 }
178
179
180 ///////////////////////////////////////////////////////////////////
181 // Implementation of Tan //
182 ///////////////////////////////////////////////////////////////////
183
184 Tan::Tan(const Tan &s) : Symbol(s) {}
185
186 Tan::Tan(const Symbolic &s) : Symbol(Symbol("tan")[s]) {}
187
188 Simplified Tan::simplify() const
189 {
190     const Symbolic &s = parameters.front().simplify();
191     if(s == 0) return Number<int>(0);
192     if(s.type() == typeid(Product))
193     {
194         CastPtr<const Product> p(s);
195         if(p->factors.front() == -1) return -Tan(-s);
196     }
197     if(s.type() == typeid(Numeric) &&
198         Number<void>(s).numeric_type() == typeid(double))
199         return Number<double>(tan(CastPtr<const Number<double>>(s)->n));
200     return *this;
201 }
202
203 Symbolic Tan::df(const Symbolic &s) const
204 { return tan(parameters.front()) * tan(parameters.front()) * parameters.front().df(s) + parameters.front().df(s) ; }
205
206 Symbolic Tan::integrate(const Symbolic &s) const
207 {
208     const Symbolic &x = parameters.front();
209     if(x == s) return -ln(cos(x)) * (1 / parameters.front().df(s));
210     if(df(s) == 0) return *this * s;
211     return ln( tan(parameters.front()) * tan(parameters.front() + 1) * ( 1 / (2*parameters.front().df(s)) ) ;
212 }
213

```

Figure 2.1: The implementation of tangent function to compute its derivative and integral in `src/functions.cpp`.

[SI*] **Trigonometry Level 2**

The trigonometry integrals that we consider as level 2 are

$$\int \sin mx \cos nx \, dx$$

$$\int \cos mx \cos nx \, dx$$

$$\int \sin mx \sin nx \, dx$$

Integrals of this type occurs in many physics and engineering applications. By hands we can use the product identities to handle these integrals.

[SI*] **Trigonometry Level 3**

The trigonometry integrals that we consider as level 3 are

$$\int \sin^n x \, dx$$

$$\int \cos^n x \, dx$$

$$\int \tan^n x \, dx$$

$$\int \sec^n x \, dx$$

$$\int \csc^n x \, dx$$

$$\int \cot^n x \, dx$$

We happen to check on Wolfram Alpha for these indefinite integrals. The Wolfram Alpha gives result of these integrals in hypergeometric function term, for example:

$$\int \sin^n x \, dx = -\cos(x) \sin^{n+1}(x) \sin^{-n-1}(x) {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3}{2}; \cos^2(x)\right)$$

For the record, hypergeometric function ${}_2F_1(\frac{1}{2}, \frac{1-n}{2}; \frac{3}{2}; \cos^2(x))$ can be computed with SymIntegration, Boost, or even self made C++ codes, but it will produce approximated integral computation, and we tried to use this formulas with hypergeometric function, it compute very slow.

So we refer to the reduction formula to compute these trigonometry integral level 3.

$$\begin{aligned} \int \sin^n x \, dx &= -\frac{1}{n} \sin^{n-1}(x) \cos(x) + \frac{n-1}{n} \int \sin^{n-2}(x) \, dx \\ \int \cos^n x \, dx &= \frac{1}{n} \cos^{n-1}(x) \sin(x) + \frac{n-1}{n} \int \cos^{n-2}(x) \, dx \\ \int \tan^n x \, dx &= \frac{\tan^{n-1}(x)}{n-1} - \int \tan^{n-2}(x) \, dx \\ \int \sec^n x \, dx &= \frac{\sec^{n-2}(x) \tan(x)}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2}(x) \, dx \\ \int \csc^n x \, dx &= \frac{-\csc^{n-2}(x) \cot(x)}{n-1} + \frac{n-2}{n-1} \int \csc^{n-2}(x) \, dx \\ \int \cot^n x \, dx &= -\frac{1}{n-1} \cot^{n-1}(x) - \int \cot^{n-2}(x) \, dx \end{aligned} \tag{2.15}$$

With this reduction formula we can use recursive method, with **for** loop technique in C++ after we find the pattern, we can code it easily. The computation time with reduction formula and this recursive method is faster than using Hypergeometric function.

[SI*] We will examine each integral and learn about their patterns, one by one. Starting with $\int \sin^n x \, dx$

i. $\int \sin^n x \, dx$

These are the formula of $\int \sin^n x \, dx$ with $n = 2$ to $n = 8$.

$$\int \sin^2 x \, dx = \frac{x}{2} - \frac{\sin x \cos x}{2}$$

$$\int \sin^3 x \, dx = \frac{\cos^3 x}{3} - \cos x$$

$$\int \sin^4 x \, dx = \frac{3x}{8} - \frac{\sin^3 x \cos x}{4} - \frac{3 \sin x \cos x}{8}$$

$$\int \sin^5 x \, dx = -\frac{\cos^5 x}{5} + \frac{2 \cos^3 x}{3} - \cos x$$

$$\int \sin^6 x \, dx = \frac{5x}{16} - \frac{\sin^5 x \cos x}{6} - \frac{5 \sin^3 x \cos x}{24} - \frac{5 \sin x \cos x}{16}$$

$$\int \sin^7 x \, dx = \frac{\cos^7 x}{7} - \frac{3 \cos^5 x}{5} + \frac{3 \cos^3 x}{3} - \cos x$$

$$\int \sin^8 x \, dx = \frac{35x}{128} - \frac{\sin^7 x \cos x}{8} - \frac{7 \sin^5 x \cos x}{48} - \frac{35 \sin^3 x \cos x}{192} - \frac{35 \sin x \cos x}{128}$$

From the first 8 terms we will get this intuition that the pattern occur for even power and odd power, so it will divide into two cases, in C++ we can use the statement `(n % 2 == 0)` in the **if** statement to handle this.

The manual way to solve this can be traced back to the reduction formula, for example if you want to compute $\int \sin^8 x \, dx$, then you start in a decreasing **for** loop with $i = n, i \geq 2, i = i - 2$.

Case 1: For $\int \sin^n x \, dx$ with even n

The numerator coefficient and denominator coefficient are making patterns that can be decoded.

```
...

Symbolic Power::integrate(const Symbolic &s) const
{
    const Symbolic &a = parameters.front();
    const Symbolic &b = parameters.back();
    int bpower = parameters.back().coeff(s,0);
    if(a.type() == typeid(Sin))
    {
        if(b.df(s) == 0 && b!=0 && bpower % 2 == 0) // even power case
        {
            Symbolic c = 1;
```



```
Symbolic d = parameters.back().coeff(s,0);
Symbolic integral;
for(int i = bpower ; i >= 2 ; i = i-2)
{
    integral -= c*cos(s)*((sin(s))^(i-1))/(d) ;
    d = d*(i-2);
    c = c*(i-1);
    if (i==4)
    {
        integral += ( (c*s)/(d) );
    }
}
return integral;
}
...
```

Code 13: level 3 integral for sine even case

As you can see the logic for the code above: we multiply the numerator with c that is the odd factorial $(1 \times 3 \times 5 \times \cdots \times n - 1)$, and multiply the denominator with the even factorial $(2 \times 4 \times 6 \times \cdots \times n)$.

The looping starts from the **bpower**, the variable name to represents n , the power for the integral of $\sin^n x$, then descending to 2, and decreasing by 2.

Notice that the numerator and denominator coefficient for x is the same as $\sin x \cos x$, so we only need to compute the coefficient for $\sin x \cos x$ and then copy it as the coefficient for x .

We start to compute the coefficient with the highest power, for example we want to compute

$\int \sin^8 x \, dx$, then we start with $i = 8$ and we will have:

```

i = 8
c = 1
d = 8
integral = -  $\frac{c * \cos(s) * \sin^{i-1}(s)}{d}$ 
d = d * (i - 2)
c = c * (i - 1)
i = i - 2
*****
i = 6
c = 7
d = 48
integral = -  $\frac{c * \cos(s) * \sin^{i-1}(s)}{d}$ 
d = d * (i - 2)
c = c * (i - 1)
i = i - 2
*****
i = 4
c = 35
d = 192
integral = -  $\frac{c * \cos(s) * \sin^{i-1}(s)}{d}$ 
d = d * (i - 2)
c = c * (i - 1)
i = i - 2
*****
i = 2
c = 105
d = 384
integral = -  $\frac{c * \cos(s) * \sin^{i-1}(s)}{d}$ 
d = d * (i - 2)
c = c * (i - 1)
i = i - 2

```

We stop when $i = 2$ and we will obtain $\frac{105}{384} = \frac{35}{128}$ as the coefficient for $\sin x \cos x$.

It only takes a lot of papers and pen to be able to decode this patterns. Then some patience and persistence with focus to create the C++ codes.

Case 2: For $\int \sin^n x \, dx$ with odd n

The numerator coefficient and denominator coefficient are making patterns that can be decoded. This time it is easier than the even case, but a little bit tricky with alternating sign and need to use combination formula.

```

...

Symbolic Power::integrate(const Symbolic &s) const
{
    ...
    if(a.type() == typeid(Sin))
    {
        if(b.df(s) == 0 && b!=0 && bpower % 2 != 0) // odd power case
        {
            Symbolic sgn = -1;
            Symbolic integral;
            int j =1;
            int m = bpower-(0.5*(bpower+1));
            for(int i = 1 ; i <= bpower ; i = i+2)
            {
                integral += sgn*combinations(m,j-1)*((cos(s))^i)/(i);
                sgn = -sgn;
                j = j+1;
            }
            return integral;
        }
    }
}

```

Code 14: level 3 integral for sine odd case

When I was trying to find the pattern, it is nice to write from the lowest odd power to a certain odd power and see the pattern for the numerator and denominator as the power rises.

$$\begin{aligned}
 \int \sin^3 x \, dx &= \frac{\cos^3 x}{3} - \cos x \\
 \int \sin^5 x \, dx &= -\frac{\cos^5 x}{5} + \frac{2\cos^3 x}{3} - \cos x \\
 \int \sin^7 x \, dx &= \frac{\cos^7 x}{7} - \frac{3\cos^5 x}{5} + \frac{3\cos^3 x}{3} - \cos x \\
 \int \sin^9 x \, dx &= -\frac{\cos^9 x}{9} + \frac{4\cos^7 x}{7} - \frac{6\cos^5 x}{5} + \frac{4\cos^3 x}{3} - \cos x
 \end{aligned}$$

Right from the bat, the denominator has an easy pattern of odd number decreasing from the n to 1, the denominator has the same coefficient as the power for the $\cos(x)$. So it is clear already.

The problem now, is the numerator, we have this kind of pattern:

$$\begin{array}{cccccc}
 & & 1 & & 1 & \\
 & & & 1 & & 2 & & 1 \\
 & 1 & & 3 & & 3 & & 1 \\
 1 & & 4 & & 6 & & 4 & & 1
 \end{array}$$

It is the infamous Pascal' triangle. Which can be derived from the combination formula:

$$\begin{aligned}
 {}_0C_0 &= 1 \\
 {}_1C_0 &= 1 \\
 {}_1C_1 &= 1 \\
 {}_2C_0 &= 1 \\
 {}_2C_1 &= 2 \\
 {}_2C_2 &= 1 \\
 {}_3C_0 &= 1 \\
 {}_3C_1 &= 3 \\
 {}_3C_2 &= 3 \\
 {}_3C_3 &= 1 \\
 {}_4C_0 &= 1 \\
 {}_4C_1 &= 4 \\
 {}_4C_2 &= 6 \\
 {}_4C_3 &= 4 \\
 {}_4C_4 &= 1
 \end{aligned}$$

This explains the usage of **combinations** function in the C++ code.

ii. $\int \cos^n x \, dx$

These are the formula of $\int \cos^n x \, dx$ with $n = 2$ to $n = 8$.

$$\begin{aligned}
 \int \cos^2 x \, dx &= \frac{x}{2} + \frac{\sin x \cos x}{2} \\
 \int \cos^3 x \, dx &= -\frac{\sin^3 x}{3} + \sin x \\
 \int \cos^4 x \, dx &= \frac{3x}{8} + \frac{\sin x \cos^3 x}{4} + \frac{3 \sin x \cos x}{8} \\
 \int \cos^5 x \, dx &= \frac{\sin^5 x}{5} - \frac{2 \sin^3 x}{3} + \sin x \\
 \int \cos^6 x \, dx &= \frac{5x}{16} + \frac{\sin x \cos^5 x}{6} + \frac{5 \sin x \cos^3 x}{24} + \frac{5 \sin x \cos x}{16} \\
 \int \cos^7 x \, dx &= -\frac{\sin^7 x}{7} + \frac{3 \sin^5 x}{5} - \frac{3 \sin^3 x}{3} + \sin x \\
 \int \cos^8 x \, dx &= \frac{35x}{128} + \frac{\sin x \cos^7 x}{8} + \frac{7 \sin x \cos^5 x}{48} + \frac{35 \sin x \cos^3 x}{192} + \frac{35 \sin x \cos x}{128}
 \end{aligned}$$

Right after we observe the equations above, it is really familiar, the pattern for the numerator and denominator will be like the sine counterpart, thus we only need to exchange the sign, what is $-$ become $+$ and the other way around, and also exchanging sine into cosine and cosine into sine.

```
...

Symbolic Power::integrate(const Symbolic &s) const
{
    const Symbolic &a = parameters.front();
    const Symbolic &b = parameters.back();
    int bpower = parameters.back().coeff(s,0);
    if(a.type() == typeid(Cos))
    {
        if(b.df(s) == 0 && b!=0 && bpower % 2 == 0) // even power case
        {
            Symbolic c = 1;
            Symbolic d = parameters.back().coeff(s,0);
            Symbolic integral;
            for(int i = bpower ; i >= 2 ; i = i-2)
            {
                integral += c*sin(s)*((cos(s))^(i-1))/(d) ;
                d = d*(i-2);
                c = c*(i-1);
                if (i==4)
                {
                    integral += ( c*s)/(d) );
                }
            }
            return integral;
        }
    }
}
```

Code 15: level 3 integral for cosine even case

```
...

Symbolic Power::integrate(const Symbolic &s) const
{
    const Symbolic &a = parameters.front();
    const Symbolic &b = parameters.back();
    int bpower = parameters.back().coeff(s,0);
    if(a.type() == typeid(Cos))
    {
        if(b.df(s) == 0 && b!=0 && bpower % 2 != 0) // odd power
            case
        {
            Symbolic sgn = 1;
```

```
Symbolic integral;
int j =1;
int m = bpower-(0.5*(bpower+1));
for(int i = 1 ; i <= bpower ; i = i+2)
{
    integral += sgn*combinations(m,j-1)*((sin(s))^(i))/(i);
    sgn = -sgn;
    j = j+1;
}
return integral;
}
...
```

Code 16: level 3 integral for cosine odd case

iii. $\int \sec^n x \, dx$

These are the formula of $\int \sec^n x \, dx$ with $n = 2$ to $n = 8$.

$$\int \sec^2 x \, dx = \frac{\sin x}{\cos x}$$

$$\int \sec^3 x \, dx = -\frac{\log(\sin(x) - 1)}{4} + \frac{\log(\sin(x) + 1)}{4} - \frac{\sin x}{2 \sin^2 x - 2}$$

$$\int \sec^4 x \, dx = \frac{2 \sin x}{3 \cos x} + \frac{\sin x}{3 \cos^3 x}$$

$$\int \sec^5 x \, dx = -\frac{3 \sin^3 x - 5 \sin x}{8 \sin^4 x - 16 \sin^2 x + 8} - \frac{3 \log(\sin(x) - 1)}{16} + \frac{3 \log(\sin(x) + 1)}{16}$$

$$\int \sec^6 x \, dx = \frac{8 \sin x}{15 \cos x} + \frac{4 \sin x}{15 \cos^3 x} + \frac{\sin x}{5 \cos^5 x}$$

$$\int \sec^7 x \, dx = \frac{-15 \sin^5 x + 40 \sin^3 x - 33 \sin x}{48 \sin^6 x - 144 \sin^4 x + 144 \sin^2 x - 48} - \frac{5 \log(\sin(x) - 1)}{32} + \frac{5 \log(\sin(x) + 1)}{32}$$

$$\int \sec^8 x \, dx = \frac{16 \sin x}{35 \cos x} + \frac{8 \sin x}{35 \cos^3 x} + \frac{6 \sin x}{35 \cos^5 x} + \frac{\sin x}{7 \cos^7 x}$$

The same case occurs again here, the pattern is repeated every $n + 2$, so there will be the odd power pattern and the even power pattern.

Case 1: For $\int \sec^n x \, dx$ with even n

The numerator coefficient and denominator coefficient are making patterns that can be decoded.

$$\int \sec^2 x \, dx = \frac{\sin x}{\cos x}$$

$$\int \sec^4 x \, dx = \frac{2 \sin x}{3 \cos x} + \frac{\sin x}{3 \cos^3 x}$$

$$\int \sec^6 x \, dx = \frac{8 \sin x}{15 \cos x} + \frac{4 \sin x}{15 \cos^3 x} + \frac{\sin x}{5 \cos^5 x}$$

$$\int \sec^8 x \, dx = \frac{16 \sin x}{35 \cos x} + \frac{8 \sin x}{35 \cos^3 x} + \frac{6 \sin x}{35 \cos^5 x} + \frac{\sin x}{7 \cos^7 x}$$

$$\int \sec^{10} x \, dx = \frac{128 \sin x}{315 \cos x} + \frac{64 \sin x}{315 \cos^3 x} + \frac{16 \sin x}{105 \cos^5 x} + \frac{8 \sin x}{63 \cos^7 x} + \frac{\sin x}{9 \cos^9 x}$$

The patterns that can be seen from those equations above are

- You may have misinterpreted it first, but the denominator are all the same, for example:

$$\int \sec^6 x \, dx = \frac{8 \sin x}{15 \cos x} + \frac{4 \sin x}{15 \cos^3 x} + \frac{\sin x}{5 \cos^5 x}$$

is the same as

$$\int \sec^6 x \, dx = \frac{8 \sin x}{15 \cos x} + \frac{4 \sin x}{15 \cos^3 x} + \frac{3 \sin x}{15 \cos^5 x}$$

Learning from that, the pattern for the numerator can be decoded.

Knowing that the denominator is the same, we only need to find out, how we get the denominator coefficient of 1 for $n = 2$, or of 3 for $n = 4$, or of 15 for $n = 6$.

Turns out, the first intuition is quite correct, it is the factorial odd, we will declare the denominator as variable d_0 :

$$d_0 = (n - 1) * (n - 3) * (n - 5) * \dots * (1)$$

If it is $\int \sec^8 x dx$, then the denominator will be

$$d_0 = 7 * 5 * 3 * 1 = 105$$

In the equation for $\int \sec^8 x dx$, the denominator is 35, instead of 105, this happens because the numerator is divisible with the denominator till they have no greatest common divisor anymore but 1, it is caused by the computer algebra system that always shows the most simplified version, so you can adjust the equation again to find the raw number for the denominator and the numerator. Making the denominator into the same number is not enough, so we need them in raw number version.

- After we make the denominator to become equal we will have this:

$$\begin{aligned}\int \sec^2 x dx &= \frac{\sin x}{\cos x} \\ \int \sec^4 x dx &= \frac{2 \sin x}{3 \cos x} + \frac{\sin x}{3 \cos^3 x} \\ \int \sec^6 x dx &= \frac{8 \sin x}{15 \cos x} + \frac{4 \sin x}{15 \cos^3 x} + \frac{3 \sin x}{15 \cos^5 x} \\ \int \sec^8 x dx &= \frac{16 \sin x}{35 \cos x} + \frac{8 \sin x}{35 \cos^3 x} + \frac{6 \sin x}{35 \cos^5 x} + \frac{5 \sin x}{35 \cos^7 x} \\ \int \sec^{10} x dx &= \frac{128 \sin x}{315 \cos x} + \frac{64 \sin x}{315 \cos^3 x} + \frac{48 \sin x}{315 \cos^5 x} + \frac{40 \sin x}{315 \cos^7 x} + \frac{35 \sin x}{315 \cos^9 x}\end{aligned}$$

so the pattern of the numerator can be seen like this:

$$\begin{array}{ccccccccc} & & & & 1 & & & & \\ & & & & 1 & & 2 & & \\ & & & 3 & & 4 & & 8 & \\ & & 5 & & 6 & & 8 & & 16 \\ 35 & & 40 & & 48 & & 64 & & 128\end{array}$$

Notice that we have to change the denominator into its raw version (for $\int \sec^8 x dx$ the denominator should be 105 instead of 35, and for $\int \sec^{10} x dx$ the denominator should be 945 instead of 315), this is the logic: the pattern can be decoded at its' raw numbers.

Thus

$$\begin{array}{ccccccccc} & & & & 1 & & & & \\ & & & & 1 & & 2 & & \\ & & & 3 & & 4 & & 8 & \\ & 15 & & 18 & & 24 & & 48 & \\ 105 & & 120 & & 144 & & 192 & & 384\end{array}$$

The first entry for the next iteration is an odd multiplication of the current first entry from before with an increasing odd number. The first iteration will multiplied by 1, the

second will be multiplied by 3, the next iteration will be multiplied by 5, and so on. This odd multiplication will be called as k , While the second entry to the last entry will be multiplied by even multiplication that will be called as l and it is increasing by 2 at every iteration.

```

for  $i = 1 \rightarrow \mathbf{v}[0] = 1 * k_{i=1} = 1$ 
for  $i = 1 \rightarrow \mathbf{v}[1] = 1 * l_{i=1} = 2$ 
for  $i = 2 \rightarrow \mathbf{v}[0] = \mathbf{v}[0]_{i=1} * k_{i=2} = 1 * 3 = 3$ 
for  $i = 2 \rightarrow \mathbf{v}[1] = \mathbf{v}[0]_{i=1} * l_{i=2} = 1 * 4 = 4$ 
for  $i = 2 \rightarrow \mathbf{v}[2] = \mathbf{v}[1]_{i=1} * l_{i=2} = 2 * 4 = 8$ 
for  $i = 3 \rightarrow \mathbf{v}[0] = \mathbf{v}[0]_{i=2} * k_{i=3} = 3 * 5 = 15$ 
for  $i = 3 \rightarrow \mathbf{v}[1] = \mathbf{v}[0]_{i=2} * l_{i=3} = 3 * 6 = 18$ 
for  $i = 3 \rightarrow \mathbf{v}[2] = \mathbf{v}[1]_{i=2} * l_{i=3} = 4 * 6 = 24$ 
for  $i = 3 \rightarrow \mathbf{v}[3] = \mathbf{v}[2]_{i=2} * l_{i=3} = 8 * 6 = 48$ 

```

Now we already have an idea about the pattern. For the C++ code, this time instead of using vector we will use array that is initialized with a very big size: `int v[999]`. Array is often used to replace vector, since vector has an invalid pointer problem / out of bounds memory access if you are not careful, array can also be used to become a matrix.

In order to determine the coefficient of the numerator with sine function we will use ascending **for** loop, to be exact we will use **for**($i = 1; i < n - 1; i = i + 2$).

We will declare variables / constants to help us compute, which are $d_0 = 1, k = 1, l = 2, c = 1$. Their value will be changing in every iteration (c will be used to store the first entry ($\mathbf{v}[0]$) of the previous array, $k = k + 2$ and $l = l + 2$).

We save the first entry of the previous array because it is going to be used to determine the first and second entry of the array at the current iteration.

```

 $k = 1$ 
 $l = 2$ 
 $d_0 = 1$ 
 $c = 1$ 

```

In every iteration we will clear the entries in the array.

So the first **for** loop will be like this:

```
         $i = 1$   
         $d_0 = 3$   
         $l = 4$   
         $k = 3$   
         $v[0] = 1$   
         $v[1] = 2$   
        *****  
         $i = 3$   
         $d_0 = 15$   
         $l = 6$   
         $k = 5$   
         $v[0] = 3$   
         $v[1] = 4$   
         $v[2] = 8$   
        *****  
         $i = 5$   
         $d_0 = 105$   
         $l = 8$   
         $k = 7$   
         $v[0] = 15$   
         $v[1] = 18$   
         $v[2] = 24$   
         $v[3] = 48$   
        *****  
         $i = 7$   
         $d_0 = 945$   
         $l = 10$   
         $k = 9$   
         $v[0] = 105$   
         $v[1] = 120$   
         $v[2] = 144$   
         $v[3] = 192$   
         $v[4] = 384$ 
```

```
...
```

```
Symbolic Power::integrate(const Symbolic &s) const  
{  
    ...  
}
```

```
if(a.type() == typeid(Sec))
{
...

if(b.df(s) == 0 && b!=0 && bpower % 2 == 0) // even power case
{
int v[999];
v[0] = 1;
Symbolic integral;
Symbolic d0 = 1;
int k = 1, l = 2, c=1;
for(int i = 1 ; i < bpower ; i = i+2) // to compute the denominator
{
    d0 *= i;
}
for(int i = 1 ; i < bpower - 1; i = i+2)
{
    c = v[0];
    int arrsec[999]; // make the size of the array as big as
                    possible

    for(int j = 0 ; j < i-1 ; j = j+1)
    {
        arrsec[j] = v[j];
    }
    int d;
    for(int j = 1 ; j < i ; j = j+1)
    {
        d = arrsec[j-1];
        v[j] = d*l;
    }

    v[0] = c*k;
    v[1] = c*l;

    k= k + 2;
    l = l + 2;
}

int j_d = 1 ;
for(int i = 1 ; i < (0.5*bpower)+1; i = i+1)
{
    integral += ( v[i-1] * sin(s) ) / ( d0*(cos(s)^(bpower-j_d)) )
    ;
    j_d = j_d+2;
}
return integral;
}
```

```

    }
}

```

Code 17: level 3 integral for secant even case

Case 2: For $\int \sec^n x dx$ with odd n

Compared to the even case, the odd case for $\int \sec^n x dx$ is harder in terms of determining the pattern and then to code it with C++, we even have to use two vectors, the first one to store the coefficient for the numerator, and the second vector is used to store the "middle coefficient", in order to determine the correct coefficient for the numerator.

$$\begin{aligned}
 \int \sec^3 x dx &= -\frac{\log(\sin(x) - 1)}{4} + \frac{\log(\sin(x) + 1)}{4} - \frac{\sin x}{2 \sin^2 x - 2} \\
 \int \sec^5 x dx &= -\frac{3 \sin^3 x - 5 \sin x}{8 \sin^4 x - 16 \sin^2 x + 8} - \frac{3 \log(\sin(x) - 1)}{16} + \frac{3 \log(\sin(x) + 1)}{16} \\
 \int \sec^7 x dx &= \frac{-15 \sin^5 x + 40 \sin^3 x - 33 \sin x}{48 \sin^6 x - 144 \sin^4 x + 144 \sin^2 x - 48} - \frac{5 \log(\sin(x) - 1)}{32} + \frac{5 \log(\sin(x) + 1)}{32} \\
 \int \sec^9 x dx &= -\frac{105 \sin^7 x - 385 \sin^5 x + 511 \sin^3 x - 279 \sin x}{384 \sin^8 x - 1536 \sin^6 x + 2304 \sin^4 x - 1536 \sin^2 x + 384} \\
 &\quad - \frac{35 \log(\sin(x) - 1)}{256} + \frac{35 \log(\sin(x) + 1)}{256} \\
 \int \sec^{11} x dx &= \frac{-315 \sin^9 x + 1470 \sin^7 x - 2688 \sin^5 x + 2370 \sin^3 x - 965 \sin x}{1280 \sin^{10} x - 6400 \sin^8 x + 12800 \sin^6 x - 12800 \sin^4 x + 6400 \sin^2 x - 1280} \\
 &\quad - \frac{63 \log(\sin(x) - 1)}{512} + \frac{63 \log(\sin(x) + 1)}{512}
 \end{aligned}$$

If we want to compute it with C++, then we can use the **for** as usual to help us. Then it is either ascending **for** loop or descending **for** loop. As time goes by and flight hours, you will know which one to use.

We can immediately notice some very obvious patterns:

- We have 3 different terms here, 2 are the log terms and the last is the term with numerator that has the sum of alternating sign of sine with odd power and denominator that has the sum of alternating sign of sine with even power.
- The terms with log has denominator of with multiplication of $(2 * i)$ that still store the previous value so we can determine it with $d_1 = d_1 * (2 * i)$. After we finish with the **for** loop, we will multiply again for the last time to obtain d_1 that will be used

$$d_1 = d_1 * (n - 1)$$

- Now, we shall see the coefficient for the sum of alternating sign of sine with odd power:

$$\begin{array}{cccc}
 1 & & & \\
 3 & 5 & & \\
 15 & 40 & 33 & \\
 105 & 385 & 511 & 279
 \end{array}$$

315 1470 2688 2370 965

The first row represents $\int \sec^3 x \, dx$ and the last row represents $\int \sec^{11} x \, dx$. The sign is alternating actually, but it is omitted for better view, so the reader won't be confused. Alternating sign can be attached later on.

In this case, to determine the coefficient of the numerator with odd sine function we will use ascending **for** loop, to be exact we will use **for**($i = 1; i < \frac{n-1}{2}; i = i + 1$).

We also need to have extra variables to help compute the coefficient correctly, we give them initial value before all the **for** loop:

$$\begin{aligned} k &= 1 \\ l &= 2 \\ d_0 &= 2 \\ d_1 &= 2 \\ \text{first_coeff} &= 3 \\ j &= 1 \\ m &= n - \left(\frac{n+1}{2} \right) \\ \text{sgn} &= -1 \end{aligned}$$

We are going to use vector $\mathbf{v}$ to represent the coefficient for the numerator of sine with odd power, the size of the vector is $\frac{n-1}{2}$.

At first, when $i = 1$, we know that $\mathbf{v}[0] = 1$, so how to determine $\mathbf{v}[0]$ and $\mathbf{v}[1]$ for $i = 2$ and continuing the loop to compute for n at higher number?

Go back and see the reduction formula at Equation (2.6) for $\int \sec^n x \, dx$, we will notice that there is $n - 2$ and $n - 1$ terms at every recursive step / the loop. So we will have to see the multiplication possibilities for the factorial odd: $(n - 2) * (n - 4) * \dots * 1$ and the factorial even: $(n - 1) * (n - 3) * (n - 5) * \dots * 2$. We are focusing on the numerator now, so it is the factorial odd that we will use (the one related to $n - 2$ factorial down with difference of 2).

If you take a look again at the triangle above, the trend is increasing then decreasing at the end, it is the typical of Pascal' triangle but we do not use combination here. Here we try step by step from $n = 3$ or $i = 1$:

$$\begin{aligned} n &= 3 \\ i &= 1 \\ \mathbf{v}[0] &= 1 \end{aligned}$$

that will make

$$\int \sec^3 x \, dx = -\frac{\log(\sin(x) - 1)}{4} + \frac{\log(\sin(x) + 1)}{4} - \frac{\mathbf{v}[0] \sin x}{2 \sin^2 x - 2}$$

moving on to $n = 5$ and $i = 2$

$$\begin{aligned} n &= 5 \\ i &= 2 \\ \mathbf{v}[0] &= 3 \\ \mathbf{v}[1] &= 5 \end{aligned}$$

$$\int \sec^5 x \, dx = -\frac{\mathbf{v}[0] \sin^3 x - \mathbf{v}[1] \sin x}{8 \sin^4 x - 16 \sin^2 x + 8} - \frac{3 \log(\sin(x) - 1)}{16} + \frac{3 \log(\sin(x) + 1)}{16}$$

Then to $n = 7$ and $i = 3$

$$\begin{aligned} n &= 7 \\ i &= 3 \\ \mathbf{v}[0] &= 15 \\ \mathbf{v}[1] &= 40 \\ \mathbf{v}[2] &= 33 \end{aligned}$$

$$\int \sec^7 x \, dx = \frac{\mathbf{v}[0] \sin^5 x + \mathbf{v}[1] \sin^3 x - \mathbf{v}[2] \sin x}{48 \sin^6 x - 144 \sin^4 x + 144 \sin^2 x - 48} - \frac{5 \log(\sin(x) - 1)}{32} + \frac{5 \log(\sin(x) + 1)}{32}$$

So to be able to create the C++ code to compute this coefficients, when we have the initial condition of $\mathbf{v}[0] = 1$ at $i = 1$, how do we get into $\mathbf{v}[0] = 3, \mathbf{v}[1] = 5$ at $i = 2$ and $\mathbf{v}[0] = 15, \mathbf{v}[1] = 40, \mathbf{v}[2] = 33$ at $i = 3$ and so on correctly?

Watch this, we first define $last_coef = \mathbf{v}[j - 1]$ with $j = 1, 2, 3, \dots$ as the last index of

the vector from the previous i and $first_coeff = 3$, so

```

n = 3
i = 1
v[0] = 1
last_coeff = v[0] = 1
k = 2
*****

n = 5
i = 2
v[0] = v[0]i=1 * (n - 2) = 3
v[1] = [last_coeff * (n - 2)] + k = [1 * 3] + 2 = 5
mc[0] = v[0] + v[1] = 8
last_coeff = v[1] = 5
k = 8
*****

n = 7
i = 3
v[0] = v[0]i=2 * (n - 2) = 3 * 5 = 15
v[1] = mc[0] * (n - 2) = 8 * 5 = 40
v[2] = [last_coeff * (n - 2)] + k = [5 * 5] + 8 = 33
mc[0] = v[0] + v[1] = 55
mc[1] = v[1] + v[2] = 73
last_coeff = v[2] = 33
k = 48
*****

n = 9
i = 4
v[0] = v[0]i=3 * (n - 2) = 15 * 7 = 105
v[1] = mc[0] * (n - 2) = 55 * 7 = 385
v[2] = mc[1] * (n - 2) = 73 * 7 = 511
v[3] = [last_coeff * (n - 2)] + k = [33 * 7] + 48 = 279
mc[0] = v[0] + v[1] = 490
mc[1] = v[1] + v[2] = 896
mc[2] = v[2] + v[3] = 790
last_coeff = v[3] = 279
k = 384

```

The $last_coeff$ and $first_coeff$ will be used in the C++ codes to represent $n - 2$, it is a little bit of a handy trick.

We finally know the pattern and formula to obtain the coefficients for the numerator. It is not an easy one as I spent days to realize how to obtain this.

- The denominator that has the sum of alternating sign of sine with even power has a distinctive pattern that can be seen when you take the greatest common divider out:

$$\begin{aligned} 8 \sin^4 x - 16 \sin^2 x + 8 &= 8(\sin^4 x - 2 \sin^2 x + 1) \\ 48 \sin^6 x - 144 \sin^4 x + 144 \sin^2 x - 48 &= 48(\sin^6 x - 3 \sin^4 x + 3 \sin^2 x - 1) \\ 384 \sin^8 x - \dots + \dots + 384 &= 384(\sin^8 x - 4 \sin^6 x + 6 \sin^4 x - 4 \sin^2 x + 1) \end{aligned}$$

We are focusing on the denominator now, so it is the factorial even that we will use, with the initial value of $d_0 = 2$ we will have:

$$\begin{aligned} \text{for } n = 5 &\rightarrow d_0 = d_0 * (n - 1) = d_0 * (2 + 2 * i) = 8 \\ \text{for } n = 7 &\rightarrow d_0 = d_0 * (n - 1) = d_0 * (2 + 2 * i) = 48 \\ \text{for } n = 9 &\rightarrow d_0 = d_0 * (n - 1) = d_0 * (2 + 2 * i) = 384 \end{aligned}$$

From this you can see clearly what to write for the C++ code.

So the first **for** loop will be like this:

```

        i = 1
        d1 = 4
        d0 = 8
        mc[0] = 1
        v[0] = 3
        v[1] = 5
    * * * * *
        i = 2
        d1 = 16
        d0 = 48
        mc[0] = 8
        v[0] = 15
        v[1] = 40
        v[2] = 33
    * * * * *
        i = 3
        d1 = 96
        d0 = 384
        mc[0] = 55
        mc[1] = 73
        v[0] = 105
        v[1] = 385
        v[2] = 511
        v[3] = 279
    * * * * *
        i = 4
        d1 = 768
        d0 = 3840
        mc[0] = 490
        mc[1] = 896
        mc[2] = 790
        v[0] = 945
        v[1] = 4410
        v[2] = 8064
        v[3] = 7110
        v[4] = 2895
    
```

The next two **for** loops won't be too hard to comprehend, we include the variable *sgn* to alternate the sign. Overall, to be able to detect the pattern out of this took quite some time,

and makes the author realize that integration to the power of n for basic trigonometry is related to sum / series, and also combination and Pascal' triangle. With that knowledge and logic, thus today Computer Algebra System, like SymIntegration or SymPy able to compute $\int \sec^n x \, dx$ for any integer number of n . It shows how beautiful mathematics is.

```
...

Symbolic Power::integrate(const Symbolic &s) const
{
...
    if(a.type() == typeid(Sec))
    {
        if(b.df(s) == 0 && b!=0 && bpower % 2 != 0) // odd power case
        {
            int k = 1, l=2;
            vector<int> v={1};
            vector<int> mc={1}; // to store the middle coefficient
            Symbolic sgn = -1;
            Symbolic integral_numerator, integral_denominator;
            Symbolic d0 = 2, d1 = 2;
            int j =1;
            int m = bpower-(0.5*(bpower+1));
            int last_coeff;
            int first_coeff = 3;

            // For the coefficient at the numerator of sine with odd power
            for(int i = 1 ; i < (bpower-1)/2 ; i = i+1)
            {
                if (i >= 2)
                {
                    for(int ic = 1 ; ic < i ; ic = ic+1)
                    {
                        mc[ic-1] = v[ic-1] + v[ic];
                    }
                }
                k = k*1;
                last_coeff = v[j-1];
                v[0] = v[0]*(first_coeff+2*(i-1));
                v.assign({v[0]});

                for(int ic = 1 ; ic < i ; ic = ic+1)
                {
                    v.push_back(mc[ic-1]*(first_coeff+2*(i-1)));
                }
                d1 = d1*(2*i);
                d0 = d0*(2+2*i);
                v.push_back(last_coeff*(first_coeff+2*(i-1))+k);
                l = l+2;
                j = j+1;
            }
        }
    }
}
```

```

    }
    int j_num = 0 ;
    for(int i = bpower-2 ; i >= 1 ; i = i-2)
    {
        integral_numerator += sgn*v[j_num]*((sin(s))^(i));
        sgn = -sgn;
        j_num = j_num+1;
    }
    sgn = 1;
    j = 1;
    for(int i = bpower-1 ; i >= 0 ; i = i-2)
    {
        integral_denominator += sgn*d0*combinations(m,j-1)*((
            sin(s))^(i));
        sgn = -sgn;
        j = j+1;
    }
    d1 = d1*(bpower-1);
    return (-v[0]*ln(sin(s)-1))/(d1) + (v[0]*ln(sin(s)+1))/(d1) +
        (integral_numerator)/(integral_denominator);
}
}

```

Code 18: level 3 integral for secant odd case

iv. $\int \csc^n x \, dx$

These are the formula of $\int \csc^n x \, dx$ with $n = 2$ to $n = 8$.

$$\int \csc^2 x \, dx = -\frac{\cos x}{\sin x}$$

$$\int \csc^3 x \, dx = \frac{\log(\cos(x)-1)}{4} - \frac{\log(\cos(x)+1)}{4} + \frac{\cos x}{2\cos^2 x - 2}$$

$$\int \csc^4 x \, dx = -\frac{2\cos x}{3\sin x} - \frac{\cos x}{3\sin^3 x}$$

$$\int \csc^5 x \, dx = \frac{3\cos^3 x - 5\cos x}{8\cos^4 x - 16\cos^2 x + 8} + \frac{3\log(\cos(x)-1)}{16} - \frac{3\log(\cos(x)+1)}{16}$$

$$\int \csc^6 x \, dx = -\frac{8\cos x}{15\sin x} - \frac{4\cos x}{15\sin^3 x} - \frac{\cos x}{5\sin^5 x}$$

$$\int \csc^7 x \, dx = -\frac{-15\cos^5 x + 40\cos^3 x - 33\cos x}{48\cos^6 x - 144\cos^4 x + 144\cos^2 x - 48} + \frac{5\log(\cos(x)-1)}{32} - \frac{5\log(\cos(x)+1)}{32}$$

$$\int \csc^8 x \, dx = -\frac{16\cos x}{35\sin x} - \frac{8\cos x}{35\sin^3 x} - \frac{6\cos x}{35\sin^5 x} - \frac{\cos x}{7\sin^7 x}$$

The pattern for the numerator and denominator will be like the secant counterpart, thus we only need to exchange the sign, what is $-$ become $+$ and the other way around, and also exchanging sine into cosine and cosine into sine.

...

```

Symbolic Power::integrate(const Symbolic &s) const
{
    ...
    if(a.type() == typeid(Csc))
    {
        ...

        if(b.df(s) == 0 && b!=0 && bpower % 2 == 0) // even power case
        {
            int v[999];
            v[0] = 1;
            Symbolic integral;
            Symbolic d0 = 1;
            int k = 1, l = 2, c=1;
            for(int i = 1 ; i < bpower ; i = i+2) // to compute the
                denominator
            {
                d0 *= i;
            }
            for(int i = 1 ; i < bpower - 1; i = i+2)
            {
                c = v[0];
                int arrsec[999]; // make the size of the array as
                    big as possible

                for(int j = 0 ; j < i-1 ; j = j+1)
                {
                    arrsec[j] = v[j];
                }
                int d;
                for(int j = 1 ; j < i ; j = j+1)
                {
                    d = arrsec[j-1];
                    v[j] = d*l;
                }

                v[0] = c*k;
                v[1] = c*l;

                k= k + 2;
                l = l + 2;
            }

            int j_d = 1 ;
            for(int i = 1 ; i < (0.5*bpower)+1; i = i+1)
            {
                integral -= ( v[i-1] * cos(s) ) / ( d0*(sin(s)^(

```

```
        bpower-j_d)) ) ;
        j_d = j_d+2;
    }
    return integral;
}
}
```

Code 19: level 3 integral for cosecant even case

```
...

Symbolic Power::integrate(const Symbolic &s) const
{
    ...
    if(a.type() == typeid(Csc))
    {
        ...

        if(b.df(s) == 0 && b!=0 && bpower % 2 != 0) // odd power case
        {
            int k = 1, l=2;
            vector<int> v={1};
            vector<int> mc={1}; // to store the middle coefficient
            Symbolic sgn = -1;
            Symbolic integral_numerator, integral_denominator;
            Symbolic d0 = 2, d1 = 2;
            int j =1;
            int m = bpower-(0.5*(bpower+1));
            int last_coeff;
            int first_coeff = 3;

            // For the coefficient at the numerator of cosine with odd
            power
            for(int i = 1 ; i < (bpower-1)/2 ; i = i+1)
            {
                if (i >= 2)
                {
                    for(int ic = 1 ; ic < i ; ic = ic+1)
                    {
                        mc[ic-1] = v[ic-1] + v[ic];
                    }
                }
                k = k*1;
                last_coeff = v[j-1];
                v[0] = v[0]*(first_coeff+2*(i-1));
                v.assign({v[0]});

                for(int ic = 1 ; ic < i ; ic = ic+1)
```

```

        {
            v.push_back(mc[ic-1]*(first_coeff+2*(i-1)));
        }
        d1 = d1*(2*i);
        d0 = d0*(2+2*i);
        v.push_back(last_coeff*(first_coeff+2*(i-1))+k);
        l = l+2;
        j = j+1;
    }
    int j_num = 0 ;
    for(int i = bpower-2 ; i >= 1 ; i = i-2)
    {
        integral_numerator += sgn*v[j_num]*((cos(s))^i));
        sgn = -sgn;
        j_num = j_num+1;
    }
    sgn = 1;
    j = 1;
    for(int i = bpower-1 ; i >= 0 ; i = i-2)
    {
        integral_denominator += sgn*d0*combinations(m,j-1)*((
            cos(s))^i));
        sgn = -sgn;
        j = j+1;
    }
    d1 = d1*(bpower-1);
    return (v[0]*ln(cos(s)-1))/(d1) - (v[0]*ln(cos(s)+1))/(d1) -
        (integral_numerator)/(integral_denominator);
    }
}
}

```

Code 20: level 3 integral for cosecant odd case

v. $\int \cot^n x \, dx$

These are the formula of $\int \cot^n x \, dx$ with $n = 2$ to $n = 8$.

$$\begin{aligned}\int \cot^2 x \, dx &= -x - \frac{\cos x}{\sin x} \\ \int \cot^3 x \, dx &= -\log(\sin(x)) - \frac{1}{2 \sin^2 x} \\ \int \cot^4 x \, dx &= x + \frac{\cos x}{\sin x} - \frac{\cos^3 x}{3 \sin^3 x} \\ \int \cot^5 x \, dx &= \log(\sin(x)) + \frac{4 \sin^2 x - 1}{4 \sin^4 x} \\ \int \cot^6 x \, dx &= -x - \frac{\cos x}{\sin x} + \frac{\cos^3 x}{3 \sin^3 x} - \frac{\cos^5 x}{5 \sin^5 x} \\ \int \cot^7 x \, dx &= -\log(\sin(x)) - \frac{18 \sin^4 x - 9 \sin^2 x + 2}{12 \sin^6 x} \\ \int \cot^8 x \, dx &= x + \frac{\cos x}{\sin x} - \frac{\cos^3 x}{3 \sin^3 x} + \frac{\cos^5 x}{5 \sin^5 x} - \frac{\cos^7 x}{7 \sin^7 x}\end{aligned}$$

Case 1: For $\int \cot^n x \, dx$ with even n

We will cluster only the even power

$$\begin{aligned}\int \cot^2 x \, dx &= -x - \frac{\cos x}{\sin x} \\ \int \cot^4 x \, dx &= x + \frac{\cos x}{\sin x} - \frac{\cos^3 x}{3 \sin^3 x} \\ \int \cot^6 x \, dx &= -x - \frac{\cos x}{\sin x} + \frac{\cos^3 x}{3 \sin^3 x} - \frac{\cos^5 x}{5 \sin^5 x} \\ \int \cot^8 x \, dx &= x + \frac{\cos x}{\sin x} - \frac{\cos^3 x}{3 \sin^3 x} + \frac{\cos^5 x}{5 \sin^5 x} - \frac{\cos^7 x}{7 \sin^7 x}\end{aligned}$$

The patterns that can be seen from those equations above are

- The variable x is only alternating sign at every iteration.
- The numerator' coefficient with cosine function of odd power is only 1.
- The denominator coefficient is the same as the power of the sine function in the denominator.
- The integral computed at the previous iteration alternates its sign at the next iteration minus the new function with power of $n - 1$.

The C++ code is an easy one here, the simplest out of all level 3 trigonometry integral, it alternates the whole sign at every iteration. We will use an increasing **for** loop with $i = 2, i \leq n, i = i + 2$.

```
...

Symbolic Power::integrate(const Symbolic &s) const
{
    ...
    if(a.type() == typeid(Cot))
    {
```

```

...
if(b.df(s) == 0 && b!=0 && bpower % 2 == 0) // even power case
{
    Symbolic integral;
    Symbolic integral_front;
    Symbolic sgn = -1;
    for(int i = 2 ; i <= (bpower) ; i = i+2)
    {
        integral = (-1)*integral;
        integral += - (cos(s)^(i-1)) / ((i-1)*(sin(s)^(i-1)));
    }

    for(int i = 1 ; i <= (bpower)/2 ; i = i+1)
    {
        integral_front = sgn;
        sgn = -sgn;
    }
    return integral_front*s + integral;
}
}

```

Code 21: level 3 integral for cotangent even case**Case 2: For $\int \cot^n x \, dx$ with odd n**

We will cluster only the odd power

$$\begin{aligned}
 \int \cot^3 x \, dx &= -\log(\sin(x)) - \frac{1}{2 \sin^2 x} \\
 \int \cot^5 x \, dx &= \log(\sin(x)) + \frac{4 \sin^2 x - 1}{4 \sin^4 x} \\
 \int \cot^7 x \, dx &= -\log(\sin(x)) - \frac{18 \sin^4 x - 9 \sin^2 x + 2}{12 \sin^6 x} \\
 \int \cot^9 x \, dx &= \log(\sin(x)) + \frac{48 \sin^4 x - 36 \sin^2 x + 16 \sin^2 x - 3}{24 \sin^8 x} \\
 \int \cot^{11} x \, dx &= -\log(\sin(x)) - \frac{300 \sin^8 x - 300 \sin^6 x + 200 \sin^4 x - 75 \sin^2 x + 12}{120 \sin^{10} x}
 \end{aligned}$$

The patterns that can be seen from those equations above are

- The $\log(\sin(x))$ function is only alternating the sign at every iteration.
- The denominator for the sine function with power of $n - 1$ is making a pattern like this (d_0 is the variable to represent the denominator)

$$d_0 = d_0 * \left(\frac{n-1}{2} \right)$$

We start with $d_0 = 2$ then

$$d_0 = 2 * \left(\frac{5-1}{2} \right) = 4$$

$$d_0 = 4 * \left(\frac{7-1}{2} \right) = 12$$

$$d_0 = 12 * \left(\frac{9-1}{2} \right) = 48$$

$$d_0 = 48 * \left(\frac{11-1}{2} \right) = 240$$

Knowing that the denominator has been simplified then we know that we have to adjust the numerator so they become the raw numbers like this:

$$\int \cot^3 x \, dx = -\log(\sin(x)) - \frac{1}{2 \sin^2 x}$$

$$\int \cot^5 x \, dx = \log(\sin(x)) + \frac{4 \sin^2 x - 1}{4 \sin^4 x}$$

$$\int \cot^7 x \, dx = -\log(\sin(x)) - \frac{18 \sin^4 x - 9 \sin^2 x + 2}{12 \sin^6 x}$$

$$\int \cot^9 x \, dx = \log(\sin(x)) + \frac{96 \sin^4 x - 72 \sin^2 x + 32}{48 \sin^8 x}$$

$$\int \cot^{11} x \, dx = -\log(\sin(x)) - \frac{600 \sin^8 x - 600 \sin^6 x + 400 \sin^4 x - 150 \sin^2 x + 24}{120 \sin^{10} x}$$

It is very important to do this in order to know the pattern for the numerator.

- To be able to know the pattern for the numerator with sine function of even power we can easily put the numerator coefficients like this for a better look:

$$\begin{array}{cccccc} & & & 1 & & & \\ & & & & 1 & & 4 \\ & & 2 & & 9 & & 18 \\ & 6 & & 32 & & 72 & & 96 \\ 24 & & 150 & & 400 & & 600 & & 600 \end{array}$$

The rule of thumb is we have to vectorize the coefficients, making it into a vector of size $\frac{n-1}{2}$ at every iteration. Back to the reduction formula, if we want to compute $\int \cot^{11} x \, dx$ we will start from $\int \cot^1 x \, dx$ then $\int \cot^3 x \, dx$, this is the recursive method.

When we create the vector at certain iteration, we will realize that to know the entries of this vector, we will need the values from certain variables that are having pattern, e.g. increasing at every iteration or decreasing at every iteration or multiplying with increasing coefficient. Now we take a look again. At the first iteration we have a vector $\mathbf{v}$ with only has one entry / size of 1 which is

$$\mathbf{v}[0] = 1$$

then at the next iteration we have a vector of size 2 with entries

$$\mathbf{v}[0] = 1$$

$$\mathbf{v}[1] = 4$$

then at the next iteration we have a vector of size 3 with entries

$$\mathbf{v}[0] = 2$$

$$\mathbf{v}[1] = 9$$

$$\mathbf{v}[2] = 18$$

We will use 4 vectors this time, since this one is quite tricky compared to the secant case. The vectors are: $\mathbf{v}, \mathbf{v}_{temp}, \mathbf{mc}, \mathbf{mc}_{temp}$.

We also need to use variables to help compute the coefficients with their initial value before the **for** loop :

$$d_0 = 2$$

$$k = 1$$

$$l = 2$$

We will use an increasing **for** loop with $i = 1, i \leq \frac{n-1}{2}, i = i + 1$ because the size of the vector is getting bigger at every iteration, the size of the vector is the same as the iteration number.

You can and probably should write it on paper so you can learn more. To be simple, after several days of trial and error, the formula is shown to be like this:

$$\begin{aligned} \mathbf{v}[0]_i &= \mathbf{v}[0]_{i-1} * (i - 1) * \mathbf{mc}[0]_i \\ \mathbf{v}[j]_i &= [\mathbf{v}[j - 1]_{i-1} * (2 * i - j)] + [\mathbf{v}[0]_i * (\mathbf{mc}[j]_i)] \\ \mathbf{v}\left[\frac{n-1}{2}\right]_i &= \left[\mathbf{v}\left[\frac{n-1}{2} - 1\right]_{i-1} * (2 * i - j)\right] + (\mathbf{v}[0]_i * \mathbf{mc}\left[\frac{n-1}{2}\right]_i) \end{aligned}$$

with i is the iteration $i = 1, 2, \dots, \frac{n-1}{2}$, and $j = 1, 2, \dots, \frac{n-1}{2}$. The vector $\mathbf{mc}$, the middle coefficient is the Pascal' triangle

$$\begin{array}{cccccccc} & & & & 1 & & & \\ & & & 1 & & 1 & & \\ & & 1 & & 2 & & 1 & \\ & 1 & & 3 & & 3 & & 1 \\ & & 1 & & 4 & & 6 & & 4 & & 1 \\ & 1 & & 5 & & 10 & & 10 & & 5 & & 1 \\ 1 & & 1 & & 6 & & 15 & & 20 & & 15 & & 6 & & 1 \end{array}$$

the top row represents $i = 1$ for $\int \cot^3 x \, dx$, and this vector $\mathbf{mc}$ helps computing the numerator coefficient.

The computation / coding of C++ for the numerator coefficients will be a tough one, it takes an increasing **for** loop with $i = 1, i \leq \frac{n-1}{2}, i = i + 1$ and we split it into several cases.

So if we want to compute $\int \cot^{11} x \, dx$ we will gain the numerator coefficients at the iteration $i = 5$.

So the first **for** loop will be like this:

```

        i = 1
        d0 = 2
        mc[0] = 1
        v[0] = 1
    * * * * *
        i = 2
        d0 = 4
        mc[0] = 1
        mc[1] = 1
        v[0] = 1
        v[1] = 4
    * * * * *
        i = 3
        d0 = 12
        mc[0] = 1
        mc[1] = 2
        mc[1] = 1
        v[0] = 2
        v[1] = 9
        v[2] = 18
    * * * * *
        i = 4
        d0 = 48
        mc[0] = 1
        mc[1] = 3
        mc[2] = 3
        mc[3] = 1
        v[0] = 6
        v[1] = 32
        v[2] = 72
        v[3] = 96
    * * * * *
```

```

        i = 5
        d0 = 240
        mc[0] = 1
        mc[1] = 4
        mc[2] = 6
        mc[3] = 4
        mc[4] = 1
        v[0] = 24
        v[1] = 150
        v[2] = 400
        v[3] = 600
        v[4] = 600

```

We omitted the `mc[0]` and `mc[i - 1]` in the C++ codes since it will only returns 1 so it won't really change much.

The rest of the `for` loops are only used to return the correct integral.

```

...

Symbolic Power::integrate(const Symbolic &s) const
{
    ...
    if(a.type() == typeid(Cot))
    {
        ...
        if(b.df(s) == 0 && b!=0 && bpower % 2 != 0) // odd power case
        {
            Symbolic integral;
            Symbolic integral_front;
            Symbolic sgn = -1;
            vector<int> v={1};
            vector<int> v_temp={1};
            vector<int> mc={1}; // to store the middle coefficient
            vector<int> mc_temp={1}; // to store the temporary / new middle
                                   coefficient
            int d0 = 2;
            int k = 1, l=2;
            // For the coefficient at the numerator
            for(int i = 1 ; i <= (bpower-1)/2 ; i = i+1)
            {
                if (i == 1)
                {
                    v[0] = 1;
                    v.assign({v[0]});
                }
            }
        }
    }
}

```

```
if (i == 2)
{
    mc[0]=1;
    mc.assign({mc[0]});
    v_temp[0] = v[0]*(i-1);
    v_temp.assign({v_temp[0]});

    for(int j = 1 ; j < i ; j = j+1)
    {
        v_temp.push_back(v[j-1]*((2*i)-j) +
            v_temp[0]*mc[j-1] );
    }
    v[0] = v_temp[0];
    v.assign({v[0]});
    for(int j = 1 ; j < i ; j = j+1)
    {
        v.push_back(v_temp[j]);
    }
}
if (i == 3)
{
    mc[0] = mc[0] + 1; //
    mc.assign({mc[0]});

    v_temp[0] = v[0]*(i-1);
    v_temp.assign({v_temp[0]});
    for(int j = 1 ; j < i-1 ; j = j+1)
    {
        v_temp.push_back(v[j-1]*((2*i)-j) +
            v_temp[0]*mc[j-1] );
    }
    v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
    // Assign the temporary vector to vector v for
    future use
    v[0] = v_temp[0];
    v.assign({v[0]});
    for(int j = 1 ; j < i ; j = j+1)
    {
        v.push_back(v_temp[j]);
    }
}
if (i == 4)
{
    mc[0] = mc[0] + 1;
    mc.assign({mc[0]});
    mc.push_back(mc[0]);

    v_temp[0] = v[0]*(i-1);
```

```

v_temp.assign({v_temp[0]});
for(int j = 1 ; j < i-1 ; j = j+1)
{
    v_temp.push_back(v[j-1]*((2*i)-j) +
        v_temp[0]*mc[j-1] );
}
v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
// Assign the temporary vector to vector v for
    future use
v[0] = v_temp[0];
v.assign({v[0]});
for(int j = 1 ; j < i ; j = j+1)
{
    v.push_back(v_temp[j]);
}
}
if (i >= 5)
{
    mc_temp[0] = mc[0]+1;
    mc_temp.assign({mc_temp[0]});
    for(int ic = 1 ; ic < l ; ic = ic+1)
    {
        mc_temp[ic] = mc[ic-1] + mc[ic];
    }
    mc[1] = (mc_temp[0]);
    mc[l+1] = (mc_temp[0]);

    mc[0]=mc_temp[0];
    mc.assign({mc[0]});

    for(int ic = 1 ; ic < l ; ic = ic+1)
    {
        mc.push_back(mc_temp[ic]);
    }
    mc.push_back(mc_temp[0]);
    mc.push_back(0);

    v_temp[0] = v[0]*(i-1);
    v_temp.assign({v_temp[0]});
    for(int j = 1 ; j < i-1 ; j = j+1)
    {
        v_temp.push_back(v[j-1]*((2*i)-j) +
            v_temp[0]*mc[j-1] );
    }
    v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
    // Assign the temporary vector to vector v for
        future use
    v[0] = v_temp[0];

```

```
        v.assign({v[0]});
        for(int j = 1 ; j < i ; j = j+1)
        {
            v.push_back(v_temp[j]);
        }

        l = l+1;
    }

    d0 = d0*k;
    k = k+1;
}
for(int i = 1 ; i <= (bpower-1)/2 ; i = i+1)
{
    integral += sgn*v[i-1]*(sin(s)^(2*(i-1)));
    sgn=-sgn;
}

for(int i = 1 ; i <= (bpower-1)/2 ; i = i+1)
{
    integral_front = sgn;
    sgn = -sgn;
}
return integral_front*ln(sin(s)) + (integral)/(d0*(sin(s)^(
bpower-1))) ;
}
}
```

Code 22: *level 3 integral for cotangent odd case*

vi. $\int \tan^n x \, dx$

These are the formula of $\int \tan^n x \, dx$ with $n = 2$ to $n = 8$.

$$\begin{aligned}\int \tan^2 x \, dx &= -x + \frac{\sin x}{\cos x} \\ \int \tan^3 x \, dx &= \log(\cos x) + \frac{1}{2 \cos^2 x} \\ \int \tan^4 x \, dx &= x - \frac{\sin x}{\cos x} + \frac{\sin^3 x}{3 \cos^3 x} \\ \int \tan^5 x \, dx &= -\log(\cos x) - \frac{4 \cos^2 x - 1}{4 \cos^4 x} \\ \int \tan^6 x \, dx &= -x + \frac{\sin x}{\cos x} - \frac{\sin^3 x}{3 \cos^3 x} + \frac{\sin^5 x}{5 \cos^5 x} \\ \int \tan^7 x \, dx &= \log(\cos x) + \frac{18 \cos^4 x - 9 \cos^2 x + 2}{12 \cos^6 x} \\ \int \tan^8 x \, dx &= x - \frac{\sin x}{\cos x} + \frac{\sin^3 x}{3 \cos^3 x} - \frac{\sin^5 x}{5 \cos^5 x} + \frac{\sin^7 x}{7 \cos^7 x}\end{aligned}$$

The pattern for the numerator and denominator will be like the cotangent counterpart, thus we only need to exchange the sign, what is $-$ become $+$ and the other way around, and also exchanging sine into cosine and cosine into sine.

```
...

Symbolic Power::integrate(const Symbolic &s) const
{
    ...
    if(a.type() == typeid(Cot))
    {
        ...
        if(b.df(s) == 0 && b!=0 && bpower % 2 == 0) // even power case
        {
            Symbolic integral;
            Symbolic integral_front;
            Symbolic sgn = -1;
            for(int i = 2 ; i <= (bpower) ; i = i+2)
            {
                integral = (-1)*integral;
                integral += (sin(s)^(i-1)) / ((i-1)*(cos(s)^(i-1)));
            }

            for(int i = 1 ; i <= (bpower)/2 ; i = i+1)
            {
                integral_front = sgn;
                sgn = -sgn;
            }
            return integral_front*s + integral;
        }
    }
}
```



```
}  
}
```

Code 23: level 3 integral for tangent even case

```
...  
  
Symbolic Power::integrate(const Symbolic &s) const  
{  
    ...  
    if(a.type() == typeid(Cot))  
    {  
        ...  
        if(b.df(s) == 0 && b!=0 && bpower % 2 != 0) // odd power case  
        {  
            Symbolic integral;  
            Symbolic integral_front;  
            Symbolic sgn = 1;  
            vector<int> v={1};  
            vector<int> v_temp={1};  
            vector<int> mc={1}; // to store the middle coefficient  
            vector<int> mc_temp={1}; // to store the temporary / new middle  
                coefficient  
            int d0 = 2;  
            int k = 1, l=2;  
            // For the coefficient at the numerator  
            for(int i = 1 ; i <= (bpower-1)/2 ; i = i+1)  
            {  
                if (i == 1)  
                {  
                    v[0] = 1;  
                    v.assign({v[0]});  
                }  
  
                if (i ==2)  
                {  
                    mc[0]=1;  
                    mc.assign({mc[0]});  
                    v_temp[0] = v[0]*(i-1);  
                    v_temp.assign({v_temp[0]});  
  
                    for(int j = 1 ; j < i ; j = j+1)  
                    {  
                        v_temp.push_back(v[j-1]*((2*i)-j) +  
                            v_temp[0]*mc[j-1] );  
                    }  
                    v[0] = v_temp[0];  
                    v.assign({v[0]});  
                    for(int j = 1 ; j < i ; j = j+1)
```

```

        {
            v.push_back(v_temp[j]);
        }
    }
    if (i == 3)
    {
        mc[0] = mc[0] + 1; //
        mc.assign({mc[0]});

        v_temp[0] = v[0]*(i-1);
        v_temp.assign({v_temp[0]});
        for(int j = 1 ; j < i-1 ; j = j+1)
        {
            v_temp.push_back(v[j-1]*((2*i)-j) +
                            v_temp[0]*mc[j-1] );
        }
        v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
        // Assign the temporary vector to vector v for
        future use
        v[0] = v_temp[0];
        v.assign({v[0]});
        for(int j = 1 ; j < i ; j = j+1)
        {
            v.push_back(v_temp[j]);
        }
    }
    if (i == 4)
    {
        mc[0] = mc[0] + 1;
        mc.assign({mc[0]});
        mc.push_back(mc[0]);

        v_temp[0] = v[0]*(i-1);
        v_temp.assign({v_temp[0]});
        for(int j = 1 ; j < i-1 ; j = j+1)
        {
            v_temp.push_back(v[j-1]*((2*i)-j) +
                            v_temp[0]*mc[j-1] );
        }
        v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
        // Assign the temporary vector to vector v for
        future use
        v[0] = v_temp[0];
        v.assign({v[0]});
        for(int j = 1 ; j < i ; j = j+1)
        {
            v.push_back(v_temp[j]);
        }
    }

```

```
    }
    if (i >= 5)
    {
        mc_temp[0] = mc[0]+1;
        mc_temp.assign({mc_temp[0]});
        for(int ic = 1 ; ic < l ; ic = ic+1)
        {
            mc_temp[ic] = mc[ic-1] + mc[ic];
        }
        mc[l] = (mc_temp[0]);
        mc[l+1] = (mc_temp[0]);

        mc[0]=mc_temp[0];
        mc.assign({mc[0]});

        for(int ic = 1 ; ic < l ; ic = ic+1)
        {
            mc.push_back(mc_temp[ic]);
        }
        mc.push_back(mc_temp[0]);
        mc.push_back(0);

        v_temp[0] = v[0]*(i-1);
        v_temp.assign({v_temp[0]});
        for(int j = 1 ; j < i-1 ; j = j+1)
        {
            v_temp.push_back(v[j-1]*((2*i)-j) +
                             v_temp[0]*mc[j-1] );
        }
        v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
        // Assign the temporary vector to vector v for
        future use
        v[0] = v_temp[0];
        v.assign({v[0]});
        for(int j = 1 ; j < i ; j = j+1)
        {
            v.push_back(v_temp[j]);
        }

        l = l+1;
    }

    d0 = d0*k;
    k = k+1;
}
for(int i = 1 ; i <= (bpower-1)/2 ; i = i+1)
{
    integral += sgn*v[i-1]*(cos(s)^(2*(i-1)));
}
```

```

        sgn=-sgn;

    }

    for(int i = 1 ; i <= (bpower-1)/2 ; i = i+1)
    {
        integral_front = sgn;
        sgn = -sgn;
    }
    return integral_front*ln(cos(s)) + (integral)/(d0*(cos(s)^(
        bpower-1))) ;
}
}
}

```

Code 24: level 3 integral for tangent odd case

[SI*] **Trigonometry Level 4**

The trigonometry integrals that we consider as level 4 is

$$\int \sin^n x \cos^m x \, dx$$

$$\int \tan^n x \sec^m x \, dx$$

$$\int \cot^n x \csc^m x \, dx$$

the reduction formula to compute these trigonometry integral level 4 are

$$\int \sin^n x \cos^m x \, dx = \frac{\sin^{n+1} x \cos^{m-1} x}{m+n} + \frac{m-1}{m+n} \int \sin^n x \cos^{m-2} x \, dx \quad (2.16)$$

vii. $\int \sin^n x \cos^m x \, dx$

The differentiation to obtain the reduction formula is coming from integration by parts and a smart algebra technique

$$\begin{aligned}
 I_{m,n} &= \int \sin^n x \cos^m x \, dx \\
 &= \int \cos^{m-1} x \sin^n x \cos x \, dx \\
 &= \cos^{m-1} x \frac{\sin^{n+1} x}{n+1} - \frac{m-1}{n+1} \int \sin^{n+2} x \cos^{m-2} x \, dx \\
 &= \cos^{m-1} x \frac{\sin^{n+1} x}{n+1} - \frac{m-1}{n+1} \int \sin^n x \cos^{m-2} x (1 - \cos^2 x) \, dx \\
 &= \frac{\sin^{n+1} x \cos^{m-1} x}{m+n} + \frac{m-1}{m+n} \int \sin^n x \cos^{m-2} x (1 - \cos^2 x) \, dx \\
 &= \frac{\sin^{n+1} x \cos^{m-1} x}{m+n} + \frac{m-1}{m+n} \int \sin^n x \cos^{m-2} x \, dx
 \end{aligned}$$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 1$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \sin^1 x \cos^1 x dx &= \frac{\sin^2 x}{2} \\ \int \sin^1 x \cos^2 x dx &= \frac{-\cos^3 x}{3} \\ \int \sin^1 x \cos^3 x dx &= \frac{-\cos^4 x}{4} \\ \int \sin^1 x \cos^4 x dx &= \frac{-\cos^5 x}{5} \\ \int \sin^1 x \cos^5 x dx &= \frac{-\cos^6 x}{6} \\ \int \sin^1 x \cos^6 x dx &= \frac{-\cos^7 x}{7} \\ \int \sin^1 x \cos^7 x dx &= \frac{-\cos^8 x}{8} \\ \int \sin^1 x \cos^8 x dx &= \frac{-\cos^9 x}{9}\end{aligned}$$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 2$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \sin^2 x \cos^1 x dx &= \frac{\sin^3 x}{3} \\ \int \sin^2 x \cos^2 x dx &= \frac{x}{8} - \frac{\sin(2x) \cos(2x)}{16} \\ \int \sin^2 x \cos^3 x dx &= -\frac{\sin^5 x}{5} + \frac{\sin^3 x}{3} \\ \int \sin^2 x \cos^4 x dx &= \frac{x}{16} - \frac{\sin(x) \cos^5(x)}{6} + \frac{\sin(x) \cos^3(x)}{24} + \frac{\sin(x) \cos(x)}{16} \\ \int \sin^2 x \cos^5 x dx &= \frac{\sin^7 x}{7} - \frac{2 \sin^5 x}{5} + \frac{\sin^3 x}{3} \\ \int \sin^2 x \cos^6 x dx &= \frac{5x}{128} - \frac{\sin(x) \cos^7(x)}{8} + \frac{\sin(x) \cos^5(x)}{48} + \frac{5 \sin(x) \cos^3(x)}{192} + \frac{5 \sin(x) \cos(x)}{128} \\ \int \sin^2 x \cos^7 x dx &= -\frac{\sin^9 x}{9} + \frac{3 \sin^7 x}{7} - \frac{3 \sin^5 x}{5} + \frac{\sin^3 x}{3} \\ \int \sin^2 x \cos^8 x dx &= \frac{7x}{256} - \frac{\sin(x) \cos^9(x)}{10} + \frac{\sin(x) \cos^7(x)}{80} + \frac{7 \sin(x) \cos^5(x)}{480} \\ &\quad + \frac{7 \sin(x) \cos^3(x)}{384} + \frac{7 \sin(x) \cos(x)}{256}\end{aligned}$$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 3$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \sin^3 x \cos^1 x dx &= \frac{\sin^4 x}{4} \\ \int \sin^3 x \cos^2 x dx &= \frac{\cos^5 x}{5} - \frac{\cos^3 x}{3} \\ \int \sin^3 x \cos^3 x dx &= -\frac{\sin^6 x}{6} + \frac{\sin^4 x}{4} \\ \int \sin^3 x \cos^4 x dx &= \frac{\cos^7 x}{7} - \frac{\cos^5 x}{5} \\ \int \sin^3 x \cos^5 x dx &= \frac{\cos^8 x}{8} - \frac{\cos^6 x}{6} \\ \int \sin^3 x \cos^6 x dx &= \frac{\cos^9 x}{9} - \frac{\cos^7 x}{7} \\ \int \sin^3 x \cos^7 x dx &= \frac{\cos^{10} x}{10} - \frac{\cos^8 x}{8} \\ \int \sin^3 x \cos^8 x dx &= \frac{\cos^{11} x}{11} - \frac{\cos^9 x}{9}\end{aligned}$$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 4$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \sin^4 x \cos^1 x dx &= \frac{\sin^5 x}{5} \\ \int \sin^4 x \cos^2 x dx &= \frac{x}{16} + \frac{\sin^5(x) \cos(x)}{6} - \frac{\sin^3(x) \cos(x)}{24} - \frac{\sin(x) \cos(x)}{16} \\ \int \sin^4 x \cos^3 x dx &= -\frac{\sin^7 x}{7} + \frac{\sin^5 x}{5} \\ \int \sin^4 x \cos^4 x dx &= \frac{3x}{128} - \frac{\sin^3(2x) \cos(2x)}{128} - \frac{3 \sin(2x) \cos(2x)}{256} \\ \int \sin^4 x \cos^5 x dx &= \frac{\sin^9 x}{9} - \frac{2 \sin^7 x}{7} + \frac{\sin^5 x}{5} \\ \int \sin^4 x \cos^6 x dx &= \frac{3x}{256} + \frac{\sin(x) \cos^9(x)}{10} - \frac{11 \sin(x) \cos^7(x)}{80} + \frac{\sin(x) \cos^5(x)}{160} \\ &\quad + \frac{\sin(x) \cos^3(x)}{128} + \frac{3 \sin(x) \cos(x)}{256} \\ \int \sin^4 x \cos^7 x dx &= -\frac{\sin^{11} x}{11} + \frac{\sin^9 x}{3} - \frac{3 \sin^7 x}{7} + \frac{\sin^5 x}{5} \\ \int \sin^4 x \cos^8 x dx &= \frac{7x}{1024} + \frac{\sin(x) \cos^{11}(x)}{12} - \frac{13 \sin(x) \cos^9(x)}{120} + \frac{\sin(x) \cos^7(x)}{320} + \frac{7 \sin(x) \cos^5(x)}{1920} \\ &\quad + \frac{7 \sin(x) \cos^3(x)}{1536} + \frac{7 \sin(x) \cos(x)}{1024}\end{aligned}$$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 5$ and $m = 1$ to $m = 8$.

$$\begin{aligned}
 \int \sin^5 x \cos^1 x dx &= \frac{\sin^6 x}{6} \\
 \int \sin^5 x \cos^2 x dx &= -\frac{\cos^7 x}{7} + \frac{2\cos^5 x}{5} - \frac{\cos^3 x}{3} \\
 \int \sin^5 x \cos^3 x dx &= -\frac{\sin^8 x}{8} + \frac{\sin^6 x}{6} \\
 \int \sin^5 x \cos^4 x dx &= -\frac{\cos^9 x}{9} + \frac{2\cos^7 x}{7} - \frac{\cos^5 x}{5} \\
 \int \sin^5 x \cos^5 x dx &= \frac{\sin^{10} x}{10} - \frac{\sin^8 x}{4} + \frac{\sin^6 x}{6} \\
 \int \sin^5 x \cos^6 x dx &= -\frac{\cos^{11} x}{11} + \frac{2\cos^9 x}{9} - \frac{\cos^7 x}{7} \\
 \int \sin^5 x \cos^7 x dx &= -\frac{\cos^{12} x}{12} + \frac{\cos^{10} x}{5} - \frac{\cos^8 x}{8} \\
 \int \sin^5 x \cos^8 x dx &= -\frac{\cos^{13} x}{13} + \frac{2\cos^{11} x}{11} - \frac{\cos^9 x}{9}
 \end{aligned}$$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 6$ and $m = 1$ to $m = 8$.

$$\begin{aligned}
 \int \sin^6 x \cos^1 x dx &= \frac{\sin^7 x}{7} \\
 \int \sin^6 x \cos^2 x dx &= \frac{5x}{128} + \frac{\sin^7(x) \cos(x)}{8} - \frac{\sin^5(x) \cos(x)}{48} - \frac{5\sin^3(x) \cos(x)}{192} - \frac{5\sin(x) \cos(x)}{128} \\
 \int \sin^6 x \cos^3 x dx &= -\frac{\sin^9 x}{9} + \frac{\sin^7 x}{7} \\
 \int \sin^6 x \cos^4 x dx &= \frac{3x}{256} - \frac{\sin^9(x) \cos(x)}{10} + \frac{11\sin^7(x) \cos(x)}{80} - \frac{\sin^5(x) \cos(x)}{160} - \frac{\sin^3(x) \cos(x)}{128} \\
 &\quad - \frac{3\sin(x) \cos(x)}{256} \\
 \int \sin^6 x \cos^5 x dx &= \frac{\sin^{11} x}{11} - \frac{2\sin^9 x}{9} + \frac{\sin^7 x}{7} \\
 \int \sin^6 x \cos^6 x dx &= \frac{5x}{1024} - \frac{\sin^5(2x) \cos(2x)}{768} - \frac{5\sin^3(2x) \cos(2x)}{3072} - \frac{5\sin(2x) \cos(2x)}{2048} \\
 \int \sin^6 x \cos^7 x dx &= -\frac{\sin^{13} x}{13} + \frac{3\sin^{11} x}{11} - \frac{\sin^9 x}{3} + \frac{\sin^7 x}{7} \\
 \int \sin^6 x \cos^8 x dx &= \frac{5x}{2048} - \frac{\sin(x) \cos^{13}(x)}{14} + \frac{29\sin(x) \cos^{11}(x)}{168} - \frac{37\sin(x) \cos^9(x)}{336} \\
 &\quad + \frac{\sin(x) \cos^7(x)}{896} + \frac{\sin(x) \cos^5(x)}{768} + \frac{5\sin(x) \cos^3(x)}{3072} + \frac{5\sin(x) \cos(x)}{2048}
 \end{aligned}$$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 7$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \sin^7 x \cos^1 x dx &= \frac{\sin^8 x}{8} \\ \int \sin^7 x \cos^2 x dx &= \frac{\cos^9 x}{9} - \frac{3 \cos^7 x}{7} + \frac{3 \cos^5 x}{5} - \frac{\cos^3 x}{3} \\ \int \sin^7 x \cos^3 x dx &= -\frac{\sin^{10} x}{10} + \frac{\sin^8 x}{8} \\ \int \sin^7 x \cos^4 x dx &= \frac{\cos^{11} x}{11} - \frac{\cos^9 x}{3} + \frac{3 \cos^7 x}{7} - \frac{\cos^5 x}{5} \\ \int \sin^7 x \cos^5 x dx &= \frac{\sin^{12} x}{12} - \frac{\sin^{10} x}{5} + \frac{\sin^8 x}{8} \\ \int \sin^7 x \cos^6 x dx &= \frac{\cos^{13} x}{13} - \frac{3 \cos^{11} x}{11} + \frac{\cos^9 x}{3} - \frac{\cos^7 x}{7} \\ \int \sin^7 x \cos^7 x dx &= -\frac{\sin^{14} x}{14} + \frac{\sin^{12} x}{4} - \frac{3 \sin^{10} x}{10} + \frac{\sin^8 x}{8} \\ \int \sin^7 x \cos^8 x dx &= \frac{\cos^{15} x}{15} - \frac{3 \cos^{13} x}{13} + \frac{3 \cos^{11} x}{11} - \frac{\cos^9 x}{9}\end{aligned}$$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 8$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \sin^8 x \cos^1 x dx &= \frac{\sin^9 x}{9} \\ \int \sin^8 x \cos^2 x dx &= \frac{7x}{256} + \frac{\sin^9(x) \cos(x)}{10} - \frac{\sin^7(x) \cos(x)}{80} - \frac{7 \sin^5(x) \cos(x)}{480} \\ &\quad - \frac{7 \sin^3(x) \cos(x)}{384} - \frac{7 \sin(x) \cos(x)}{256} \\ \int \sin^8 x \cos^3 x dx &= -\frac{\sin^{11} x}{11} + \frac{\sin^9 x}{9} \\ \int \sin^8 x \cos^4 x dx &= \frac{7x}{1024} - \frac{\sin^{11}(x) \cos(x)}{12} + \frac{13 \sin^9(x) \cos(x)}{120} - \frac{\sin^7(x) \cos(x)}{320} - \frac{7 \sin^5(x) \cos(x)}{1920} \\ &\quad - \frac{7 \sin^3(x) \cos(x)}{1536} - \frac{7 \sin(x) \cos(x)}{1024} \\ \int \sin^8 x \cos^5 x dx &= \frac{\sin^{13} x}{13} - \frac{2 \sin^{11} x}{11} + \frac{\sin^9 x}{9} \\ \int \sin^8 x \cos^6 x dx &= \frac{5x}{2048} + \frac{\sin^{13}(x) \cos(x)}{14} - \frac{29 \sin^{11}(x) \cos(x)}{168} + \frac{37 \sin^9(x) \cos(x)}{336} \\ &\quad - \frac{\sin^7(x) \cos(x)}{896} - \frac{\sin^5(x) \cos(x)}{768} - \frac{5 \sin^3(x) \cos(x)}{3072} - \frac{5 \sin(x) \cos(x)}{2048} \\ \int \sin^8 x \cos^7 x dx &= -\frac{\sin^{15} x}{15} + \frac{3 \sin^{13} x}{13} - \frac{3 \sin^{11} x}{11} + \frac{\sin^9 x}{9} \\ \int \sin^8 x \cos^8 x dx &= \frac{35x}{32768} - \frac{\sin^7(2x) \cos(2x)}{4096} - \frac{7 \sin^5(2x) \cos(2x)}{24576} \\ &\quad - \frac{35 \sin^3(2x) \cos(2x)}{98304} - \frac{35 \sin(2x) \cos(2x)}{65536}\end{aligned}$$

Case 1: For $\int \sin^n x \cos^m x dx$ with even power for both n and m

This case occurs when both m and n are even number. We divide this case into 3 smaller

cases:

1. When $n > m$
2. When $n < m$
3. When $n = m$

Case 1-a: For $\int \sin^n x \cos^m x dx$ with $n > m$

We will start with a big example, these are the formula of $\int \sin^n x \cos^m x dx$ with $n = 14$ and $m = 2$ to $m = 12$ with increment of 2.

$$\begin{aligned}
 \int \sin^{14} x \cos^2 x dx &= \frac{429x}{32768} + \frac{\sin^{15}(x) \cos(x)}{16} - \frac{\sin^{13}(x) \cos(x)}{224} - \frac{13 \sin^{11}(x) \cos(x)}{2688} \\
 &\quad - \frac{143 \sin^9(x) \cos(x)}{26880} - \frac{429 \sin^7(x) \cos(x)}{71680} - \frac{143 \sin^5(x) \cos(x)}{20480} \\
 &\quad - \frac{143 \sin^3(x) \cos(x)}{16384} - \frac{429 \sin(x) \cos(x)}{32768} \\
 \int \sin^{14} x \cos^4 x dx &= \frac{143x}{65536} - \frac{\sin^{17}(x) \cos(x)}{18} + \frac{19 \sin^{15}(x) \cos(x)}{288} - \frac{\sin^{13}(x) \cos(x)}{1344} \\
 &\quad - \frac{13 \sin^{11}(x) \cos(x)}{16128} - \frac{143 \sin^9(x) \cos(x)}{161280} - \frac{143 \sin^7(x) \cos(x)}{143360} \\
 &\quad - \frac{143 \sin^5(x) \cos(x)}{122880} - \frac{143 \sin^3(x) \cos(x)}{98304} - \frac{143 \sin(x) \cos(x)}{65536} \\
 \int \sin^{14} x \cos^6 x dx &= \frac{143x}{262144} + \frac{\sin^{19}(x) \cos(x)}{20} - \frac{41 \sin^{17}(x) \cos(x)}{360} + \frac{383 \sin^{15}(x) \cos(x)}{5760} \\
 &\quad - \frac{\sin^{13}(x) \cos(x)}{5376} - \frac{13 \sin^{11}(x) \cos(x)}{64512} - \frac{143 \sin^9(x) \cos(x)}{645120} \\
 &\quad - \frac{143 \sin^7(x) \cos(x)}{573440} - \frac{143 \sin^5(x) \cos(x)}{491520} - \frac{143 \sin^3(x) \cos(x)}{393216} \\
 &\quad - \frac{143 \sin(x) \cos(x)}{262144} \\
 \int \sin^{14} x \cos^8 x dx &= \frac{91x}{524288} - \frac{\sin^{21}(x) \cos(x)}{22} + \frac{67 \sin^{19}(x) \cos(x)}{440} - \frac{1367 \sin^{17}(x) \cos(x)}{7920} \\
 &\quad + \frac{8441 \sin^{15}(x) \cos(x)}{126720} - \frac{\sin^{13}(x) \cos(x)}{16896} - \frac{13 \sin^{11}(x) \cos(x)}{202752} \\
 &\quad - \frac{13 \sin^9(x) \cos(x)}{184320} - \frac{13 \sin^7(x) \cos(x)}{163840} - \frac{91 \sin^5(x) \cos(x)}{983040} \\
 &\quad - \frac{91 \sin^3(x) \cos(x)}{786432} - \frac{91 \sin(x) \cos(x)}{524288} \\
 \int \sin^{14} x \cos^{10} x dx &= \frac{273x}{4194304} + \frac{\sin^{23}(x) \cos(x)}{24} - \frac{97 \sin^{21}(x) \cos(x)}{528} + \frac{1081 \sin^{19}(x) \cos(x)}{3520} \\
 &\quad - \frac{1629 \sin^{17}(x) \cos(x)}{7040} + \frac{7507 \sin^{15}(x) \cos(x)}{112640} - \frac{\sin^{13}(x) \cos(x)}{45056} \\
 &\quad - \frac{13 \sin^{11}(x) \cos(x)}{540672} - \frac{13 \sin^9(x) \cos(x)}{491520} - \frac{39 \sin^7(x) \cos(x)}{1310720} \\
 &\quad - \frac{91 \sin^5(x) \cos(x)}{2621440} - \frac{91 \sin^3(x) \cos(x)}{2097152} - \frac{273 \sin(x) \cos(x)}{4194304}
 \end{aligned}$$

$$\begin{aligned} \int \sin^{14} x \cos^{12} x dx = & \frac{231x}{8388608} - \frac{\sin^{25}(x) \cos(x)}{26} + \frac{131 \sin^{23}(x) \cos(x)}{624} - \frac{577 \sin^{21}(x) \cos(x)}{1248} \\ & + \frac{4281 \sin^{19}(x) \cos(x)}{8320} - \frac{4829 \sin^{17}(x) \cos(x)}{16640} + \frac{17747 \sin^{15}(x) \cos(x)}{266240} \\ & - \frac{\sin^{13}(x) \cos(x)}{106496} - \frac{\sin^{11}(x) \cos(x)}{98304} - \frac{11 \sin^9(x) \cos(x)}{983040} \\ & - \frac{33 \sin^7(x) \cos(x)}{2621440} - \frac{77 \sin^5(x) \cos(x)}{5242880} - \frac{77 \sin^3(x) \cos(x)}{4194304} \\ & - \frac{231 \sin(x) \cos(x)}{8388608} \end{aligned}$$

The patterns that can be seen from those equations above are

- The integral $\int \sin^{14} x dx$ represents some term for $\int \sin^{14} x \cos^2 x dx$ with denominator that is multiplied by $m + n = 14 + 2 = 16$.
- The higher the power of m the longer the new term that is having power of $\sin^{n+1}$ and above.
- As the iteration increases the denominator coefficient is the multiplication of $m + n$ with m starting from 2 to 12 with increment of 2.

This kind of integral is quite challenging, if we want to compute this integral with $n = 14$ and $m = 12$ we will start from $m = 0$, then $m = 2$. So let see when $m = 0$, we have computed and know the pattern for this kind of integral:

$$\begin{aligned} \int \sin^{14} x \cos^0 x dx = & \frac{429x}{2048} - \frac{\sin^{13}(x) \cos(x)}{14} - \frac{13 \sin^{11}(x) \cos(x)}{168} \\ & - \frac{143 \sin^9(x) \cos(x)}{1680} - \frac{429 \sin^7(x) \cos(x)}{4480} - \frac{143 \sin^5(x) \cos(x)}{1280} \\ & - \frac{143 \sin^3(x) \cos(x)}{1024} - \frac{429 \sin(x) \cos(x)}{2048} \end{aligned}$$

now when $m = 2$, we will obtain this:

$$\begin{aligned} \int \sin^{14} x \cos^2 x dx = & \frac{429x}{32768} + \frac{\sin^{15}(x) \cos(x)}{16} - \frac{\sin^{13}(x) \cos(x)}{224} - \frac{13 \sin^{11}(x) \cos(x)}{2688} \\ & - \frac{143 \sin^9(x) \cos(x)}{26880} - \frac{429 \sin^7(x) \cos(x)}{71680} - \frac{143 \sin^5(x) \cos(x)}{20480} \\ & - \frac{143 \sin^3(x) \cos(x)}{16384} - \frac{429 \sin(x) \cos(x)}{32768} \end{aligned}$$

From the reduction formula we know that this term:

$$\frac{c_{n/2} x}{d_{n/2}} + \sum_{i=1}^{n/2} - \frac{c_i \sin^{2i-1} x \cos x}{d_i}$$

will be multiplied by $\frac{m-1}{m+n}$ at every iteration from $m = 2$ till $m = 12$. So the problem is solved for this term.

For example, to obtain the denominator for function of x of 32768 from multiplying the denominator of function x from $\int \sin^{14} x dx$ with $d_0 = m + n = 16$, it is $2048 * 16 = 32768$.

The remaining term will have different pattern for the numerator, from now on we will omit the term of $\frac{c_{n/2}x}{d_{n/2}} + \sum_{i=1}^{n/2} -\frac{c_i \sin^{2i-1} x \cos x}{d_i}$, now take a look again:

$$\begin{aligned}
 \int \sin^{14} x \cos^2 x dx &= \frac{\sin^{15}(x) \cos(x)}{16} + \dots - \dots \\
 \int \sin^{14} x \cos^4 x dx &= -\frac{\sin^{17}(x) \cos(x)}{18} + \frac{19 \sin^{15}(x) \cos(x)}{288} + \dots - \dots \\
 \int \sin^{14} x \cos^6 x dx &= \frac{\sin^{19}(x) \cos(x)}{20} - \frac{41 \sin^{17}(x) \cos(x)}{360} \\
 &\quad + \frac{383 \sin^{15}(x) \cos(x)}{5760} + \dots - \dots \\
 \int \sin^{14} x \cos^8 x dx &= -\frac{\sin^{21}(x) \cos(x)}{22} + \frac{67 \sin^{19}(x) \cos(x)}{440} - \frac{1367 \sin^{17}(x) \cos(x)}{7920} \\
 &\quad + \frac{8441 \sin^{15}(x) \cos(x)}{126720} + \dots - \dots \\
 \int \sin^{14} x \cos^{10} x dx &= \frac{\sin^{23}(x) \cos(x)}{24} - \frac{97 \sin^{21}(x) \cos(x)}{528} + \frac{1081 \sin^{19}(x) \cos(x)}{3520} \\
 &\quad - \frac{1629 \sin^{17}(x) \cos(x)}{7040} + \frac{7507 \sin^{15}(x) \cos(x)}{112640} + \dots - \dots \\
 \int \sin^{14} x \cos^{12} x dx &= -\frac{\sin^{25}(x) \cos(x)}{26} + \frac{131 \sin^{23}(x) \cos(x)}{624} - \frac{577 \sin^{21}(x) \cos(x)}{1248} \\
 &\quad + \frac{4281 \sin^{19}(x) \cos(x)}{8320} - \frac{4829 \sin^{17}(x) \cos(x)}{16640} \\
 &\quad + \frac{17747 \sin^{15}(x) \cos(x)}{266240} + \dots - \dots
 \end{aligned}$$

We will learn how to decode this pattern, first we will make the denominator into its' raw number, so the numerator will change into its' raw number too, here it goes

$$\begin{aligned}
 \int \sin^{14} x \cos^2 x dx &= \frac{\sin^{15}(x) \cos(x)}{16} + \dots - \dots \\
 \int \sin^{14} x \cos^4 x dx &= -\frac{\sin^{17}(x) \cos(x)}{18} + \frac{19 \sin^{15}(x) \cos(x)}{288} + \dots - \dots \\
 \int \sin^{14} x \cos^6 x dx &= \frac{\sin^{19}(x) \cos(x)}{20} - \frac{41 \sin^{17}(x) \cos(x)}{360} \\
 &\quad + \frac{383 \sin^{15}(x) \cos(x)}{5760} + \dots - \dots \\
 \int \sin^{14} x \cos^8 x dx &= -\frac{\sin^{21}(x) \cos(x)}{22} + \frac{67 \sin^{19}(x) \cos(x)}{440} - \frac{1367 \sin^{17}(x) \cos(x)}{7920} \\
 &\quad + \frac{8441 \sin^{15}(x) \cos(x)}{126720} + \dots - \dots \\
 \int \sin^{14} x \cos^{10} x dx &= \frac{\sin^{23}(x) \cos(x)}{24} - \frac{97 \sin^{21}(x) \cos(x)}{528} + \frac{3243 \sin^{19}(x) \cos(x)}{10560} \\
 &\quad - \frac{43983 \sin^{17}(x) \cos(x)}{190080} + \frac{202689 \sin^{15}(x) \cos(x)}{3041280} + \dots - \dots
 \end{aligned}$$

$$\begin{aligned} \int \sin^{14} x \cos^{12} x dx = & -\frac{\sin^{25}(x) \cos(x)}{26} + \frac{131 \sin^{23}(x) \cos(x)}{624} - \frac{6347 \sin^{21}(x) \cos(x)}{13728} \\ & + \frac{141273 \sin^{19}(x) \cos(x)}{274560} - \frac{1434213 \sin^{17}(x) \cos(x)}{4942080} \\ & + \frac{5270859 \sin^{15}(x) \cos(x)}{79073280} + \dots - \dots \end{aligned}$$

Then for a better look we will write it into Pascal' triangle form, for the numerator in the raw version

$$\begin{array}{ccccccc} & & & & 1 & & \\ & & & & 1 & 19 & \\ & & & 1 & 41 & 383 & \\ & & 1 & 67 & 1367 & 8441 & \\ & 1 & 97 & 3243 & 43983 & 202689 & \\ 1 & 131 & 6347 & 141273 & 1434213 & 5270859 & \end{array}$$

The C++ code will use an increasing **for** loop with $i = 1, i \leq \frac{m}{2}, i = i + 1$.

For the computation, we are creating a variable for the denominator for function $\sin^{m+n-1} x \cos x$ as d_0 , with $d_0 = m + n$. This variable will plays an important role to determine the value for the numerator of this kind of integral.

We will also use vector $\mathbf{v}$ to store the coefficient with

$$\begin{aligned} \mathbf{v}[0] &= 1 \\ \mathbf{v}[1]_i &= (d_0 * (i - 1)) + 1, \quad i = 2, 3, \dots, \frac{m}{2} - 1 \\ \mathbf{v} \left[\frac{m}{2} - 1 \right]_i &= (d_0 * \mathbf{v} \left[\frac{m}{2} - 2 \right]_{i-1}) + \mathbf{factorialoddup}(i - 1) \end{aligned}$$

$\mathbf{v} \left[\frac{m}{2} - 1 \right]_i$ is the last entry for the vector at iteration i , while $\mathbf{v} \left[\frac{m}{2} - 2 \right]_{i-1}$ is the last entry for the vector at the previous iteration $i - 1$.

We make a definition here of **factorialoddup**($i - 1$):

$$\mathbf{factorialoddup}(i - 1) = 1 * 3 * 5 * \dots$$

it is a multiplication of odd number from 1 upward till we get $i - 1$ terms.

$$\begin{aligned} \mathbf{factorialoddup}(2) &= 1 * 3 \\ \mathbf{factorialoddup}(3) &= 1 * 3 * 5 \\ \mathbf{factorialoddup}(4) &= 1 * 3 * 5 * 7 \end{aligned}$$

For $i = 4$ we will make a **for** loop again to find the entry of the vector $\mathbf{v}[j]_i$ for $j = 2, 3, \dots, \frac{m}{2} - 2$.

$$\begin{aligned} \mathbf{v}[j]_i &= (d_0 * \mathbf{v}[j - 1]_{i-1}) + (k * \mathbf{v}[j - 1]_{i-1} + \mathbf{v}[j]_{i-1}) \\ k &= k - 2 \end{aligned}$$

So the **for** loop will be like this:

```

        i = 1
        d0 = 16
        v[0] = 1
    * * * * *
        i = 2
        d0 = 18
        v[0] = 1
        v[1] = (d0 * (2 - 1)) + 1 = 19
    * * * * *
        i = 3
        d0 = 20
        v[0] = 1
        v[1] = (d0 * (3 - 1)) + 1 = 41
        v[2] = (d0 * v[1]i=2) + (1 * 3) = 383
    * * * * *
        i = 4
        d0 = 22
        k = 2
        v[0] = 1
        v[1] = (d0 * (4 - 1)) + 1 = 67
        v[2] = (d0 * v[1]i=3) + (k * v[1]i=3 + v[2]i=3) = 1367
        v[3] = (d0 * v[2]i=3) + (1 * 3 * 5) = 8441
    
```

Case 1-b: For $\int \sin^n x \cos^m x \, dx$ with $n < m$

Let's take a look at several integral related to this case

$$\begin{aligned}
 \int \sin^2 x \cos^4 x \, dx &= \frac{x}{16} - \frac{\sin(x) \cos^5(x)}{6} + \frac{\sin(x) \cos^3(x)}{24} + \frac{\sin(x) \cos(x)}{16} \\
 \int \sin^2 x \cos^6 x \, dx &= \frac{5x}{128} - \frac{\sin(x) \cos^7(x)}{8} + \frac{\sin(x) \cos^5(x)}{48} + \frac{5 \sin(x) \cos^3(x)}{192} + \frac{5 \sin(x) \cos(x)}{128} \\
 \int \sin^4 x \cos^6 x \, dx &= \frac{3x}{256} + \frac{\sin(x) \cos^9(x)}{10} - \frac{11 \sin(x) \cos^7(x)}{80} + \frac{\sin(x) \cos^5(x)}{160} \\
 &\quad + \frac{\sin(x) \cos^3(x)}{128} + \frac{3 \sin(x) \cos(x)}{256} \\
 \int \sin^2 x \cos^8 x \, dx &= \frac{7x}{256} - \frac{\sin(x) \cos^9(x)}{10} + \frac{\sin(x) \cos^7(x)}{80} + \frac{7 \sin(x) \cos^5(x)}{480} \\
 &\quad + \frac{7 \sin(x) \cos^3(x)}{384} + \frac{7 \sin(x) \cos(x)}{256} \\
 \int \sin^4 x \cos^8 x \, dx &= \frac{7x}{1024} + \frac{\sin(x) \cos^{11}(x)}{12} - \frac{13 \sin(x) \cos^9(x)}{120} + \frac{\sin(x) \cos^7(x)}{320} + \frac{7 \sin(x) \cos^5(x)}{1920} \\
 &\quad + \frac{7 \sin(x) \cos^3(x)}{1536} + \frac{7 \sin(x) \cos(x)}{1024} \\
 \int \sin^6 x \cos^8 x \, dx &= \frac{5x}{2048} - \frac{\sin(x) \cos^{13}(x)}{14} + \frac{29 \sin(x) \cos^{11}(x)}{168} - \frac{37 \sin(x) \cos^9(x)}{336} \\
 &\quad + \frac{\sin(x) \cos^7(x)}{896} + \frac{\sin(x) \cos^5(x)}{768} + \frac{5 \sin(x) \cos^3(x)}{3072} + \frac{5 \sin(x) \cos(x)}{2048}
 \end{aligned}$$

This case does not need to be scrutinize further, since the numerator and denominator for all terms are exactly the same as those for the case 1-a where $n > m$, we only need to change the sine function into cosine and cosine function into sine in the result, since this case has the cosine function in the term that has the power. The plus and minus will also be interchange for the term with sine and cosine function, the term with x stays positive always.

Case 1-c: For $\int \sin^n x \cos^m x \, dx$ with $n = m$

These are the formula of $\int \sin^n x \cos^m x dx$ with $n = 1$ to $n = 8$.

$$\begin{aligned}
 \int \sin^1 x \cos^1 x dx &= \frac{\sin^2 x}{2} \\
 \int \sin^2 x \cos^2 x dx &= \frac{x}{8} - \frac{\sin(2x) \cos(2x)}{16} \\
 \int \sin^3 x \cos^3 x dx &= -\frac{\sin^6 x}{6} + \frac{\sin^4 x}{4} \\
 \int \sin^4 x \cos^4 x dx &= \frac{3x}{128} - \frac{\sin^3(2x) \cos(2x)}{128} - \frac{3 \sin(2x) \cos(2x)}{256} \\
 \int \sin^5 x \cos^5 x dx &= \frac{\sin^{10} x}{10} - \frac{\sin^8 x}{4} + \frac{\sin^6 x}{6} \\
 \int \sin^6 x \cos^6 x dx &= \frac{5x}{1024} - \frac{\sin^5(2x) \cos(2x)}{768} - \frac{5 \sin^3(2x) \cos(2x)}{3072} - \frac{5 \sin(2x) \cos(2x)}{2048} \\
 \int \sin^7 x \cos^7 x dx &= -\frac{\sin^{14} x}{14} + \frac{\sin^{12} x}{4} - \frac{3 \sin^{10} x}{10} + \frac{\sin^8 x}{8} \\
 \int \sin^8 x \cos^8 x dx &= \frac{35x}{32768} - \frac{\sin^7(2x) \cos(2x)}{4096} - \frac{7 \sin^5(2x) \cos(2x)}{24576} - \frac{35 \sin^3(2x) \cos(2x)}{98304} \\
 &\quad - \frac{35 \sin(2x) \cos(2x)}{65536}
 \end{aligned}$$

We can tell that the case here will be divided into two, which are even case and odd case.

If n is even

$$\begin{aligned}
 \int \sin^2 x \cos^2 x dx &= \frac{x}{8} - \frac{\sin(2x) \cos(2x)}{16} \\
 \int \sin^4 x \cos^4 x dx &= \frac{3x}{128} - \frac{\sin^3(2x) \cos(2x)}{128} - \frac{3 \sin(2x) \cos(2x)}{256} \\
 \int \sin^6 x \cos^6 x dx &= \frac{5x}{1024} - \frac{\sin^5(2x) \cos(2x)}{768} - \frac{5 \sin^3(2x) \cos(2x)}{3072} - \frac{5 \sin(2x) \cos(2x)}{2048} \\
 \int \sin^8 x \cos^8 x dx &= \frac{35x}{32768} - \frac{\sin^7(2x) \cos(2x)}{4096} - \frac{7 \sin^5(2x) \cos(2x)}{24576} - \frac{35 \sin^3(2x) \cos(2x)}{98304} \\
 &\quad - \frac{35 \sin(2x) \cos(2x)}{65536}
 \end{aligned}$$

The patterns that can be seen from those equations above are

- The denominator with function of $\sin(2x)^{n-1} \cos(2x)$ can be determined with:

$$d_0 = 2^n * (2n)$$

This denominator will be used for the first integral result which is

$$\frac{\sin^{n-1}(2x) \cos(2x)}{d_0}$$

the next iteration will have the pattern like this:

$$\frac{(n-1) \sin^{n-3}(2x) \cos(2x)}{d_0 * (n-2)}$$

The next iteration will have pattern like this:

$$\frac{(n-1)(n-3)\sin^{n-5}(2x)\cos(2x)}{d_0 * (n-2) * (n-4)}$$

Till we obtain the last iteration integral result with function $\sin(2x)\cos(2x)$ at the numerator we will stop. The easy thing here is the signs are all minus.

- The function x will have numerator and denominator of:

$$\frac{2 * (n-1) * (n-3) * \dots * 1}{d_0 * (n-2) * (n-4) * \dots * 1} * x$$

If n is odd

$$\begin{aligned}\int \sin^1 x \cos^1 x dx &= \frac{\sin^2 x}{2} \\ \int \sin^3 x \cos^3 x dx &= -\frac{\sin^6 x}{6} + \frac{\sin^4 x}{4} \\ \int \sin^5 x \cos^5 x dx &= \frac{\sin^{10} x}{10} - \frac{\sin^8 x}{4} + \frac{\sin^6 x}{6} \\ \int \sin^7 x \cos^7 x dx &= -\frac{\sin^{14} x}{14} + \frac{\sin^{12} x}{4} - \frac{3\sin^{10} x}{10} + \frac{\sin^8 x}{8}\end{aligned}$$

The patterns that can be seen from those equations above are

- The integral will have $\frac{n+1}{2}$ term/s, and have function of $\sin^{n+1} x$ at the numerator, at the next iteration we will have the function of $\sin^{(n+1)+2} x$ at the numerator with alternating sign, till we reach the $\sin^{2n} x$ at the numerator.
- The power determine the Pascal' triangle level / combinations formula that will be used to determine the numerator coefficients.
For $n = 1$ we will use the first level of Pascal' triangle which is ${}_0C_0$.
For $n = 3$ we will use the second level of Pascal' triangle which is ${}_1C_0$ and ${}_1C_1$.
For $n = 5$ we will use the third level of Pascal' triangle which is ${}_2C_0$, ${}_2C_1$ and ${}_2C_2$.
For $n = 7$ we will use the fourth level of Pascal' triangle which is ${}_3C_0$, ${}_3C_1$, ${}_3C_2$ and ${}_3C_3$.
- The denominator coefficient can be easily determined, it is the same as the power of the function in the numerator

Case 2: For $\int \sin^n x \cos^m x dx$ with at least one odd power in either n or m

This case occurs if either n or m is an odd number. We will recall the integral for $n = 1$,

$n = 2, n = 3, n = 5,$ and $n = 6$

$$\int \sin^1 x \cos^2 x \, dx = \frac{-\cos^3 x}{3}$$

$$\int \sin^1 x \cos^3 x \, dx = \frac{-\cos^4 x}{4}$$

$$\int \sin^1 x \cos^4 x \, dx = \frac{-\cos^5 x}{5}$$

$$\int \sin^1 x \cos^5 x \, dx = \frac{-\cos^6 x}{6}$$

$$\int \sin^1 x \cos^6 x \, dx = \frac{-\cos^7 x}{7}$$

$$\int \sin^1 x \cos^7 x \, dx = \frac{-\cos^8 x}{8}$$

$$\int \sin^1 x \cos^8 x \, dx = \frac{-\cos^9 x}{9}$$

$$\int \sin^2 x \cos^3 x \, dx = -\frac{\sin^5 x}{5} + \frac{\sin^3 x}{3}$$

$$\int \sin^2 x \cos^5 x \, dx = \frac{\sin^7 x}{7} - \frac{2\sin^5 x}{5} + \frac{\sin^3 x}{3}$$

$$\int \sin^2 x \cos^7 x \, dx = -\frac{\sin^9 x}{9} + \frac{3\sin^7 x}{7} - \frac{3\sin^5 x}{5} + \frac{\sin^3 x}{3}$$

$$\int \sin^3 x \cos^1 x \, dx = \frac{\sin^4 x}{4}$$

$$\int \sin^3 x \cos^2 x \, dx = \frac{\cos^5 x}{5} - \frac{\cos^3 x}{3}$$

$$\int \sin^3 x \cos^4 x \, dx = \frac{\cos^7 x}{7} - \frac{\cos^5 x}{5}$$

$$\int \sin^3 x \cos^5 x \, dx = \frac{\cos^8 x}{8} - \frac{\cos^6 x}{6}$$

$$\int \sin^3 x \cos^6 x \, dx = \frac{\cos^9 x}{9} - \frac{\cos^7 x}{7}$$

$$\int \sin^3 x \cos^7 x \, dx = \frac{\cos^{10} x}{10} - \frac{\cos^8 x}{8}$$

$$\int \sin^3 x \cos^8 x \, dx = \frac{\cos^{11} x}{11} - \frac{\cos^9 x}{9}$$

$$\begin{aligned}
 \int \sin^5 x \cos^1 x \, dx &= \frac{\sin^6 x}{6} \\
 \int \sin^5 x \cos^2 x \, dx &= -\frac{\cos^7 x}{7} + \frac{2 \cos^5 x}{5} - \frac{\cos^3 x}{3} \\
 \int \sin^5 x \cos^3 x \, dx &= -\frac{\sin^8 x}{8} + \frac{\sin^6 x}{6} \\
 \int \sin^5 x \cos^4 x \, dx &= -\frac{\cos^9 x}{9} + \frac{2 \cos^7 x}{7} - \frac{\cos^5 x}{5} \\
 \int \sin^5 x \cos^6 x \, dx &= -\frac{\cos^{11} x}{11} + \frac{2 \cos^9 x}{9} - \frac{\cos^7 x}{7} \\
 \int \sin^5 x \cos^7 x \, dx &= -\frac{\cos^{12} x}{12} + \frac{\cos^{10} x}{5} - \frac{\cos^8 x}{8} \\
 \int \sin^5 x \cos^8 x \, dx &= -\frac{\cos^{13} x}{13} + \frac{2 \cos^{11} x}{11} - \frac{\cos^9 x}{9}
 \end{aligned}$$

$$\begin{aligned}
 \int \sin^6 x \cos^1 x \, dx &= \frac{\sin^7 x}{7} \\
 \int \sin^6 x \cos^3 x \, dx &= -\frac{\sin^9 x}{9} + \frac{\sin^7 x}{7} \\
 \int \sin^6 x \cos^5 x \, dx &= \frac{\sin^{11} x}{11} - \frac{2 \sin^9 x}{9} + \frac{\sin^7 x}{7} \\
 \int \sin^6 x \cos^7 x \, dx &= -\frac{\sin^{13} x}{13} + \frac{3 \sin^{11} x}{11} - \frac{\sin^9 x}{3} + \frac{\sin^7 x}{7}
 \end{aligned}$$

There will be two big cases here which are:

1. When $m > n$
2. When $m \leq n$

If $m > n$

$$\begin{aligned}
 \int \sin^1 x \cos^2 x \, dx &= \frac{-\cos^3 x}{3} \\
 \int \sin^1 x \cos^3 x \, dx &= \frac{-\cos^4 x}{4} \\
 \int \sin^1 x \cos^4 x \, dx &= \frac{-\cos^5 x}{5} \\
 \int \sin^1 x \cos^5 x \, dx &= \frac{-\cos^6 x}{6} \\
 \int \sin^1 x \cos^6 x \, dx &= \frac{-\cos^7 x}{7} \\
 \int \sin^1 x \cos^7 x \, dx &= \frac{-\cos^8 x}{8} \\
 \int \sin^1 x \cos^8 x \, dx &= \frac{-\cos^9 x}{9} \\
 \int \sin^2 x \cos^3 x \, dx &= -\frac{\sin^5 x}{5} + \frac{\sin^3 x}{3} \\
 \int \sin^2 x \cos^5 x \, dx &= \frac{\sin^7 x}{7} - \frac{2\sin^5 x}{5} + \frac{\sin^3 x}{3} \\
 \int \sin^2 x \cos^7 x \, dx &= -\frac{\sin^9 x}{9} + \frac{3\sin^7 x}{7} - \frac{3\sin^5 x}{5} + \frac{\sin^3 x}{3} \\
 \int \sin^3 x \cos^4 x \, dx &= \frac{\cos^7 x}{7} - \frac{\cos^5 x}{5} \\
 \int \sin^3 x \cos^5 x \, dx &= \frac{\cos^8 x}{8} - \frac{\cos^6 x}{6} \\
 \int \sin^5 x \cos^6 x \, dx &= -\frac{\cos^{11} x}{11} + \frac{2\cos^9 x}{9} - \frac{\cos^7 x}{7} \\
 \int \sin^5 x \cos^7 x \, dx &= -\frac{\cos^{12} x}{12} + \frac{\cos^{10} x}{5} - \frac{\cos^8 x}{8} \\
 \int \sin^5 x \cos^8 x \, dx &= -\frac{\cos^{13} x}{13} + \frac{2\cos^{11} x}{11} - \frac{\cos^9 x}{9} \\
 \int \sin^6 x \cos^7 x \, dx &= -\frac{\sin^{13} x}{13} + \frac{3\sin^{11} x}{11} - \frac{\sin^9 x}{3} + \frac{\sin^7 x}{7}
 \end{aligned}$$

The patterns that can be seen from those equations above are

- The integral will have the result with function of cosine, if n is an odd number.

The integral will have the result of sine, if n is an even number.

- If n is an even number, then the pattern will be like this for the integral:

$$\begin{aligned}
 \int \sin^n x \cos^m x \, dx &= \operatorname{sgn}_0 * \frac{m-1}{2} C_0 \frac{\sin^{n+1} x}{n+1} + \operatorname{sgn}_1 * \frac{m-1}{2} C_1 \frac{\sin^{n+3} x}{n+3} + \dots \\
 &\quad + \operatorname{sgn}_{\frac{m-1}{2}} * \frac{m-1}{2} C_{\frac{m-1}{2}} \frac{\sin^{n+m} x}{n+m}
 \end{aligned}$$

We start with $\text{sgn}_0 = 1$ and it is alternating at every iteration.

- If n is an odd number, then the pattern will be like this for the integral:

$$\int \sin^n x \cos^m x dx = \text{sgn}_0 * \frac{n-1}{2} C_0 \frac{\cos^{m+1} x}{m+1} + \text{sgn}_1 * \frac{n-1}{2} C_1 \frac{\cos^{m+3} x}{m+3} + \dots$$

$$\text{sgn}_{\frac{n-1}{2}} * \frac{n-1}{2} C_{\frac{n-1}{2}} \frac{\cos^{n+m} x}{n+m}$$

We start with $\text{sgn}_0 = -1$ and it is alternating at every iteration.

If $m \leq n$

$$\begin{aligned} \int \sin^3 x \cos^1 x dx &= \frac{\sin^4 x}{4} \\ \int \sin^3 x \cos^2 x dx &= \frac{\cos^5 x}{5} - \frac{\cos^3 x}{3} \\ \int \sin^5 x \cos^1 x dx &= \frac{\sin^6 x}{6} \\ \int \sin^5 x \cos^2 x dx &= -\frac{\cos^7 x}{7} + \frac{2\cos^5 x}{5} - \frac{\cos^3 x}{3} \\ \int \sin^5 x \cos^3 x dx &= -\frac{\sin^8 x}{8} + \frac{\sin^6 x}{6} \\ \int \sin^5 x \cos^4 x dx &= -\frac{\cos^9 x}{9} + \frac{2\cos^7 x}{7} - \frac{\cos^5 x}{5} \\ \int \sin^6 x \cos^1 x dx &= \frac{\sin^7 x}{7} \\ \int \sin^6 x \cos^3 x dx &= -\frac{\sin^9 x}{9} + \frac{\sin^7 x}{7} \\ \int \sin^6 x \cos^5 x dx &= \frac{\sin^{11} x}{11} - \frac{2\sin^9 x}{9} + \frac{\sin^7 x}{7} \end{aligned}$$

The patterns that can be seen from those equations above are

- If m is an even number, then the integral will be the function of cosine.

If m is an odd number, then the integral will be the function of sine.

- If m is an even number, then the pattern will be like this for the integral:

$$\int \sin^n x \cos^m x dx = \text{sgn}_0 * \frac{n-1}{2} C_0 \frac{\cos^{m+1} x}{m+1} + \text{sgn}_1 * \frac{n-1}{2} C_1 \frac{\cos^{m+3} x}{m+3} + \dots$$

$$+ \text{sgn}_{\frac{n-1}{2}} * \frac{n-1}{2} C_{\frac{n-1}{2}} \frac{\cos^{n+m} x}{n+m}$$

We start with $\text{sgn}_0 = -1$ and it is alternating at every iteration.

- If m is an odd number, then the pattern will be like this for the integral:

$$\int \sin^n x \cos^m x dx = \text{sgn}_0 * \frac{n-1}{2} C_0 \frac{\sin^{n+1} x}{n+1} + \text{sgn}_1 * \frac{n-1}{2} C_1 \frac{\sin^{n+3} x}{n+3} + \dots$$

$$+ \text{sgn}_{\frac{n-1}{2}} * \frac{n-1}{2} C_{\frac{n-1}{2}} \frac{\sin^{n+m} x}{n+m}$$

We start with $\text{sgn}_0 = -1$ and it is alternating at every iteration.

viii. $\int \tan^n x \sec^m x dx$

These are the formula of $\int \tan^n x \sec^m x dx$ with $n = 1$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \tan^1 x \sec^1 x dx &= \frac{1}{\cos(x)} \\ \int \tan^1 x \sec^2 x dx &= \frac{1}{2 \cos^2(x)} \\ \int \tan^1 x \sec^3 x dx &= \frac{1}{3 \cos^3(x)} \\ \int \tan^1 x \sec^4 x dx &= \frac{1}{4 \cos^4(x)} \\ \int \tan^1 x \sec^5 x dx &= \frac{1}{5 \cos^5(x)} \\ \int \tan^1 x \sec^6 x dx &= \frac{1}{6 \cos^6(x)} \\ \int \tan^1 x \sec^7 x dx &= \frac{1}{7 \cos^7(x)} \\ \int \tan^1 x \sec^8 x dx &= \frac{1}{8 \cos^8(x)}\end{aligned}$$

These are the formula of $\int \tan^n x \sec^m x dx$ with $n = 2$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \tan^2 x \sec^1 x dx &= \frac{\log(\sin(x) - 1)}{4} - \frac{\log(\sin(x) + 1)}{4} - \frac{\sin(x)}{2 \sin^2(x) - 2} \\ \int \tan^2 x \sec^2 x dx &= -\frac{\sin(x)}{3 \cos(x)} + \frac{\sin(x)}{3 \cos^3(x)} \\ \int \tan^2 x \sec^3 x dx &= \frac{\log(\sin(x) - 1)}{16} - \frac{\log(\sin(x) + 1)}{16} - \frac{-\sin^3(x) - \sin(x)}{8 \sin^4(x) - 16 \sin^2(x) + 8} \\ \int \tan^2 x \sec^4 x dx &= -\frac{2 \sin(x)}{15 \cos(x)} - \frac{\sin(x)}{15 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)} \\ \int \tan^2 x \sec^5 x dx &= \frac{\log(\sin(x) - 1)}{32} - \frac{\log(\sin(x) + 1)}{32} + \frac{3 \sin^5(x) - 8 \sin^3(x) - 3 \sin(x)}{48 \sin^6(x) - 144 \sin^4(x) + 144 \sin^2(x) - 48} \\ \int \tan^2 x \sec^6 x dx &= -\frac{8 \sin(x)}{105 \cos(x)} - \frac{4 \sin(x)}{105 \cos^3(x)} - \frac{\sin(x)}{35 \cos^5(x)} + \frac{\sin(x)}{7 \cos^7(x)} \\ \int \tan^2 x \sec^7 x dx &= \frac{5 \log(\sin(x) - 1)}{256} - \frac{5 \log(\sin(x) + 1)}{256} \\ &\quad - \frac{-15 \sin^7(x) + 55 \sin^5(x) - 73 \sin^3(x) - 15 \sin(x)}{384 \sin^8(x) - 1536 \sin^6(x) + 2304 \sin^4(x) - 1536 \sin^2(x) + 384} \\ \int \tan^2 x \sec^8 x dx &= -\frac{16 \sin(x)}{315 \cos(x)} - \frac{8 \sin(x)}{315 \cos^3(x)} - \frac{2 \sin(x)}{105 \cos^5(x)} - \frac{\sin(x)}{63 \cos^7(x)} + \frac{\sin(x)}{9 \cos^9(x)}\end{aligned}$$

These are the formula of $\int \tan^n x \sec^m x dx$ with $n = 3$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \tan^3 x \sec^1 x dx &= \frac{1 - 3 \cos^2(x)}{3 \cos^3(x)} \\ \int \tan^3 x \sec^2 x dx &= \frac{1 - 2 \cos^2(x)}{4 \cos^4(x)} \\ \int \tan^3 x \sec^3 x dx &= \frac{3 - 5 \cos^2(x)}{15 \cos^5(x)} \\ \int \tan^3 x \sec^4 x dx &= \frac{2 - 3 \cos^2(x)}{12 \cos^6(x)} \\ \int \tan^3 x \sec^5 x dx &= \frac{5 - 7 \cos^2(x)}{35 \cos^7(x)} \\ \int \tan^3 x \sec^6 x dx &= \frac{3 - 4 \cos^2(x)}{24 \cos^8(x)} \\ \int \tan^3 x \sec^7 x dx &= \frac{7 - 9 \cos^2(x)}{63 \cos^9(x)} \\ \int \tan^3 x \sec^8 x dx &= \frac{4 - 5 \cos^2(x)}{40 \cos^{10}(x)}\end{aligned}$$

These are the formula of $\int \tan^n x \sec^m x dx$ with $n = 4$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \tan^4 x \sec^1 x dx &= -\frac{3 \log(\sin(x) - 1)}{16} + \frac{3 \log(\sin(x) + 1)}{16} - \frac{-5 \sin^3(x) + 3 \sin(x)}{8 \sin^4(x) - 16 \sin^2(x) + 8} \\ \int \tan^4 x \sec^2 x dx &= \frac{\sin(x)}{5 \cos(x)} - \frac{2 \sin(x)}{5 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)} \\ \int \tan^4 x \sec^3 x dx &= -\frac{\log(\sin(x) - 1)}{32} + \frac{\log(\sin(x) + 1)}{32} - \frac{-3 \sin^5(x) - 8 \sin^3(x) + 3 \sin(x)}{48 \sin^6(x) - 144 \sin^4(x) + 144 \sin^2(x) - 48} \\ \int \tan^4 x \sec^4 x dx &= \frac{2 \sin(x)}{35 \cos(x)} + \frac{\sin(x)}{35 \cos^3(x)} - \frac{8 \sin(x)}{35 \cos^5(x)} + \frac{\sin(x)}{7 \cos^7(x)} \\ \int \tan^4 x \sec^5 x dx &= -\frac{3 \log(\sin(x) - 1)}{256} + \frac{3 \log(\sin(x) + 1)}{256} \\ &\quad - \frac{3 \sin^7(x) - 11 \sin^5(x) - 11 \sin^3(x) + 3 \sin(x)}{128 \sin^8(x) - 512 \sin^6(x) + 768 \sin^4(x) - 512 \sin^2(x) + 128} \\ \int \tan^4 x \sec^6 x dx &= \frac{8 \sin(x)}{315 \cos(x)} + \frac{4 \sin(x)}{315 \cos^3(x)} + \frac{\sin(x)}{105 \cos^5(x)} - \frac{10 \sin(x)}{63 \cos^7(x)} + \frac{\sin(x)}{9 \cos^9(x)} \\ \int \tan^4 x \sec^7 x dx &= -\frac{3 \log(\sin(x) - 1)}{512} + \frac{3 \log(\sin(x) + 1)}{512} \\ &\quad + \frac{-15 \sin^9(x) + 70 \sin^7(x) - 128 \sin^5(x) - 70 \sin^3(x) + 15 \sin(x)}{1280 \sin^{10}(x) - 6400 \sin^8(x) + 12800 \sin^6(x) - 12800 \sin^4(x) + 6400 \sin^2(x) - 1280} \\ \int \tan^4 x \sec^8 x dx &= \frac{16 \sin(x)}{1155 \cos(x)} + \frac{8 \sin(x)}{1155 \cos^3(x)} + \frac{2 \sin(x)}{385 \cos^5(x)} + \frac{\sin(x)}{231 \cos^7(x)} - \frac{4 \sin(x)}{33 \cos^9(x)} \\ &\quad + \frac{\sin(x)}{11 \cos^{11}(x)}\end{aligned}$$

These are the formula of $\int \tan^n x \sec^m x dx$ with $n = 5$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \tan^5 x \sec^1 x dx &= \frac{-(-15 \cos^4(x) + 10 \cos^2(x) - 3)}{15 \cos^5(x)} \\ \int \tan^5 x \sec^2 x dx &= \frac{-(-3 \cos^4(x) + 3 \cos^2(x) - 1)}{6 \cos^6(x)} \\ \int \tan^5 x \sec^3 x dx &= \frac{-(-35 \cos^4(x) + 42 \cos^2(x) - 15)}{105 \cos^7(x)} \\ \int \tan^5 x \sec^4 x dx &= \frac{-(-6 \cos^4(x) + 8 \cos^2(x) - 3)}{24 \cos^8(x)} \\ \int \tan^5 x \sec^5 x dx &= \frac{-(-63 \cos^4(x) + 90 \cos^2(x) - 35)}{315 \cos^9(x)} \\ \int \tan^5 x \sec^6 x dx &= \frac{-(-10 \cos^4(x) + 15 \cos^2(x) - 6)}{60 \cos^{10}(x)} \\ \int \tan^5 x \sec^7 x dx &= \frac{-(-99 \cos^4(x) + 154 \cos^2(x) - 63)}{693 \cos^{11}(x)} \\ \int \tan^5 x \sec^8 x dx &= \frac{-(-15 \cos^4(x) + 24 \cos^2(x) - 10)}{120 \cos^{12}(x)}\end{aligned}$$

These are the formula of $\int \tan^n x \sec^m x dx$ with $n = 6$ and $m = 1$ to $m = 8$.

$$\begin{aligned}\int \tan^6 x \sec^1 x dx &= \frac{5 \log(\sin(x) - 1)}{32} - \frac{5 \log(\sin(x) + 1)}{32} + \frac{-33 \sin^5(x) + 40 \sin^3(x) - 15 \sin(x)}{48 \sin^6(x) - 144 \sin^4(x) + 144 \sin^2(x) - 48} \\ \int \tan^6 x \sec^2 x dx &= -\frac{\sin(x)}{7 \cos(x)} + \frac{3 \sin(x)}{7 \cos^3(x)} - \frac{3 \sin(x)}{7 \cos^5(x)} + \frac{\sin(x)}{7 \cos^7(x)} \\ \int \tan^6 x \sec^3 x dx &= \frac{5 \log(\sin(x) - 1)}{256} - \frac{5 \log(\sin(x) + 1)}{256} \\ &\quad - \frac{-15 \sin^7(x) - 73 \sin^5(x) + 55 \sin^3(x) - 15 \sin(x)}{384 \sin^8(x) - 1536 \sin^6(x) + 2304 \sin^4(x) - 1536 \sin^2(x) + 384} \\ \int \tan^6 x \sec^4 x dx &= -\frac{2 \sin(x)}{63 \cos(x)} - \frac{\sin(x)}{63 \cos^3(x)} + \frac{5 \sin(x)}{21 \cos^5(x)} - \frac{19 \sin(x)}{63 \cos^7(x)} + \frac{\sin(x)}{9 \cos^9(x)} \\ \int \tan^6 x \sec^5 x dx &= \frac{3 \log(\sin(x) - 1)}{512} - \frac{3 \log(\sin(x) + 1)}{512} \\ &\quad + \frac{15 \sin^9(x) - 70 \sin^7(x) - 128 \sin^5(x) + 70 \sin^3(x) - 15 \sin(x)}{1280 \sin^{10}(x) - 6400 \sin^8(x) + 12800 \sin^6(x) - 12800 \sin^4(x) + 6400 \sin^2(x) - 1280} \\ \int \tan^6 x \sec^6 x dx &= -\frac{8 \sin(x)}{693 \cos(x)} - \frac{4 \sin(x)}{693 \cos^3(x)} - \frac{\sin(x)}{231 \cos^5(x)} + \frac{113 \sin(x)}{693 \cos^7(x)} \\ &\quad - \frac{23 \sin(x)}{99 \cos^9(x)} + \frac{\sin(x)}{11 \cos^{11}(x)} \\ \int \tan^6 x \sec^7 x dx &= \frac{5 \log(\sin(x) - 1)}{2048} - \frac{5 \log(\sin(x) + 1)}{2048} \\ &\quad - \frac{-15 \sin^{11}(x) + 85 \sin^9(x) - 198 \sin^7(x) - 198 \sin^5(x) + 85 \sin^3(x) - 15 \sin(x)}{3072 \sin^{12}(x) - 18432 \sin^{10}(x) + 46080 \sin^8(x) - 61440 \sin^6(x) + 46080 \sin^4(x) - 18432 \sin^2(x) + 3072}\end{aligned}$$

$$\begin{aligned} \int \tan^6 x \sec^8 x \, dx = & -\frac{16 \sin(x)}{3003 \cos(x)} - \frac{8 \sin(x)}{3003 \cos^3(x)} - \frac{2 \sin(x)}{1001 \cos^5(x)} - \frac{5 \sin(x)}{3003 \cos^7(x)} \\ & + \frac{53 \sin(x)}{429 \cos^9(x)} - \frac{27 \sin(x)}{143 \cos^{11}(x)} + \frac{\sin(x)}{13 \cos^{13}(x)} \end{aligned}$$

These are the formula of $\int \tan^n x \sec^m x \, dx$ with $n = 7$ and $m = 1$ to $m = 8$.

$$\begin{aligned} \int \tan^7 x \sec^1 x \, dx &= \frac{-35 \cos^6(x) + 35 \cos^4(x) - 21 \cos^2(x) + 5}{35 \cos^7(x)} \\ \int \tan^7 x \sec^2 x \, dx &= \frac{-4 \cos^6(x) + 6 \cos^4(x) - 4 \cos^2(x) + 1}{8 \cos^8(x)} \\ \int \tan^7 x \sec^3 x \, dx &= \frac{-105 \cos^6(x) + 189 \cos^4(x) - 135 \cos^2(x) + 35}{315 \cos^9(x)} \\ \int \tan^7 x \sec^4 x \, dx &= \frac{-10 \cos^6(x) + 20 \cos^4(x) - 15 \cos^2(x) + 4}{40 \cos^{10}(x)} \\ \int \tan^7 x \sec^5 x \, dx &= \frac{-231 \cos^6(x) + 495 \cos^4(x) - 385 \cos^2(x) + 105}{1155 \cos^{11}(x)} \\ \int \tan^7 x \sec^6 x \, dx &= \frac{-20 \cos^6(x) + 45 \cos^4(x) - 36 \cos^2(x) + 10}{120 \cos^{12}(x)} \\ \int \tan^7 x \sec^7 x \, dx &= \frac{-429 \cos^6(x) + 1001 \cos^4(x) - 819 \cos^2(x) + 231}{3003 \cos^{13}(x)} \\ \int \tan^7 x \sec^8 x \, dx &= \frac{-35 \cos^6(x) + 84 \cos^4(x) - 70 \cos^2(x) + 20}{280 \cos^{14}(x)} \end{aligned}$$

Case 1: For $\int \tan^n x \sec^m x \, dx$ with n even

We divide this case into 2 smaller cases:

1. When m even
2. When m odd

Case 1-a: For $\int \tan^n x \sec^m x \, dx$ with n even and m even

$$\begin{aligned}
 \int \tan^2 x \sec^2 x \, dx &= -\frac{\sin(x)}{3 \cos(x)} + \frac{\sin(x)}{3 \cos^3(x)} \\
 \int \tan^2 x \sec^4 x \, dx &= -\frac{2 \sin(x)}{15 \cos(x)} - \frac{\sin(x)}{15 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)} \\
 \int \tan^2 x \sec^6 x \, dx &= -\frac{8 \sin(x)}{105 \cos(x)} - \frac{4 \sin(x)}{105 \cos^3(x)} - \frac{\sin(x)}{35 \cos^5(x)} + \frac{\sin(x)}{7 \cos^7(x)} \\
 \int \tan^2 x \sec^8 x \, dx &= -\frac{16 \sin(x)}{315 \cos(x)} - \frac{8 \sin(x)}{315 \cos^3(x)} - \frac{2 \sin(x)}{105 \cos^5(x)} - \frac{\sin(x)}{63 \cos^7(x)} + \frac{\sin(x)}{9 \cos^9(x)} \\
 \int \tan^4 x \sec^2 x \, dx &= \frac{\sin(x)}{5 \cos(x)} - \frac{2 \sin(x)}{5 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)} \\
 \int \tan^4 x \sec^4 x \, dx &= \frac{2 \sin(x)}{35 \cos(x)} + \frac{\sin(x)}{35 \cos^3(x)} - \frac{8 \sin(x)}{35 \cos^5(x)} + \frac{\sin(x)}{7 \cos^7(x)} \\
 \int \tan^4 x \sec^6 x \, dx &= \frac{8 \sin(x)}{315 \cos(x)} + \frac{4 \sin(x)}{315 \cos^3(x)} + \frac{\sin(x)}{105 \cos^5(x)} - \frac{10 \sin(x)}{63 \cos^7(x)} + \frac{\sin(x)}{9 \cos^9(x)} \\
 \int \tan^4 x \sec^8 x \, dx &= \frac{16 \sin(x)}{1155 \cos(x)} + \frac{8 \sin(x)}{1155 \cos^3(x)} + \frac{2 \sin(x)}{385 \cos^5(x)} + \frac{\sin(x)}{231 \cos^7(x)} - \frac{4 \sin(x)}{33 \cos^9(x)} \\
 &\quad + \frac{\sin(x)}{11 \cos^{11}(x)} \\
 \int \tan^6 x \sec^2 x \, dx &= -\frac{\sin(x)}{7 \cos(x)} + \frac{3 \sin(x)}{7 \cos^3(x)} - \frac{3 \sin(x)}{7 \cos^5(x)} + \frac{\sin(x)}{7 \cos^7(x)} \\
 \int \tan^6 x \sec^4 x \, dx &= -\frac{2 \sin(x)}{63 \cos(x)} - \frac{\sin(x)}{63 \cos^3(x)} + \frac{5 \sin(x)}{21 \cos^5(x)} - \frac{19 \sin(x)}{63 \cos^7(x)} + \frac{\sin(x)}{9 \cos^9(x)} \\
 \int \tan^6 x \sec^6 x \, dx &= -\frac{8 \sin(x)}{693 \cos(x)} - \frac{4 \sin(x)}{693 \cos^3(x)} - \frac{\sin(x)}{231 \cos^5(x)} + \frac{113 \sin(x)}{693 \cos^7(x)} \\
 &\quad - \frac{23 \sin(x)}{99 \cos^9(x)} + \frac{\sin(x)}{11 \cos^{11}(x)} \\
 \int \tan^6 x \sec^8 x \, dx &= -\frac{16 \sin(x)}{3003 \cos(x)} - \frac{8 \sin(x)}{3003 \cos^3(x)} - \frac{2 \sin(x)}{1001 \cos^5(x)} - \frac{5 \sin(x)}{3003 \cos^7(x)} \\
 &\quad + \frac{53 \sin(x)}{429 \cos^9(x)} - \frac{27 \sin(x)}{143 \cos^{11}(x)} + \frac{\sin(x)}{13 \cos^{13}(x)}
 \end{aligned}$$

The patterns that can be seen from those equations above are

- The series are consisting of a function with $\sin(x)$ as the numerator and $\cos(x)$ as the denominator, the length of the series and the maximum power for the denominator depending on m and n .
- If you take a look at the $\int \sec^m x \, dx$ for m even, you will notice that this case will use the result of $\int \sec^m x \, dx$ for m even depending on n and using the combination to make the Pascal' triangle.

For example, let $n = 4$ and $m = 6$

$$\begin{aligned}
 \int \tan^4 x \sec^6 x \, dx &= \int (\sec^2 x - 1)^2 \sec^6 x \, dx \\
 &= \int (\sec^4 x - 2\sec^2 x + 1) \sec^6 x \, dx \\
 &= \int \sec^{10} x - 2\sec^8 x + \sec^6 x \, dx \\
 &= \frac{8 \sin(x)}{315 \cos(x)} + \frac{4 \sin(x)}{315 \cos^3(x)} + \frac{\sin(x)}{105 \cos^5(x)} - \frac{10 \sin(x)}{63 \cos^7(x)} + \frac{\sin(x)}{9 \cos^9(x)}
 \end{aligned}$$

In the end we will refer to the formula for $\int \sec^m x \, dx$ for m even that starts the power of the secant and end at the total power of the secant and tangent, with.

```

#include <iostream>
#include "symintegrationc++.h"
#include <vector>

using namespace std;

// Factorial and combinations
Symbolic factorial(int n) {
    if (n <= 1)
    {
        return 1;
    }
    Symbolic result = 1;
    for (int i = 2; i <= n; ++i)
    {
        result *= i;
    }
    return result;
}

Symbolic combinations(int n, int r) {
    if (r < 0 || r > n)
    {
        return 0; // Invalid input
    }
    return factorial(n) / (factorial(r) * factorial(n - r));
}

Symbolic integralsecanteven(int n) {
    Symbolic x("x");
    int v[999];
    v[0] = 1;
    Symbolic integral;
    Symbolic d0 = 1;
    int k = 1, l = 2, c=1;
    for(int i = 1 ; i < n ; i = i+2) // to compute the denominator

```

```
{
    d0 *= i;
}
for(int i = 1 ; i < n - 1; i = i+2)
{
    c = v[0];
    int arrsec[999]; // make the size of the array as big as
                     possible

    for(int j = 0 ; j < i-1 ; j = j+1)
    {
        arrsec[j] = v[j];
    }
    int d;
    for(int j = 1 ; j < i ; j = j+1)
    {
        d = arrsec[j-1];
        v[j] = d*1;
    }

    v[0] = c*k;
    v[1] = c*1;

    k= k + 2;
    l = l + 2;
}

int j_d = 1 ;
for(int i = 1 ; i < (0.5*n)+1; i = i+1)
{
    integral += ( v[i-1] * sin(x) ) / ( d0*(cos(x)^(n-j_d)) ) ;
    j_d = j_d+2;
}
return integral;
}

int main(void)
{
    Symbolic x("x");
    int bpowersec = 4;
    int bpowertan = 6;
    int bpower = bpowersec+bpowertan;
    Symbolic result;
    Symbolic sgn = 1;
    int j = 1;
    int m = bpowertan/2;
    for(int i = bpower ; i >= bpowersec ; i = i-2)
    {
```

```

        result += sgn*combinations(m,j-1)*integralsecanteven(i);
        sgn = -sgn;
        j = j+1;
    }
    cout << "integral = " << result << endl;
    //cout << "integral with subs = " << result[x==1] << endl;
    return 0;
}

```

Code 25: level 4 integral for tangent even power and secant even power case

Case 1-b: For $\int \tan^n x \sec^m x dx$ with n even and m odd

$$\begin{aligned}
 \int \tan^2 x \sec^1 x dx &= \frac{\log(\sin(x) - 1)}{4} - \frac{\log(\sin(x) + 1)}{4} - \frac{\sin(x)}{2\sin^2(x) - 2} \\
 \int \tan^2 x \sec^3 x dx &= \frac{\log(\sin(x) - 1)}{16} - \frac{\log(\sin(x) + 1)}{16} - \frac{-\sin^3(x) - \sin(x)}{8\sin^4(x) - 16\sin^2(x) + 8} \\
 \int \tan^2 x \sec^5 x dx &= \frac{\log(\sin(x) - 1)}{32} - \frac{\log(\sin(x) + 1)}{32} + \frac{3\sin^5(x) - 8\sin^3(x) - 3\sin(x)}{48\sin^6(x) - 144\sin^4(x) + 144\sin^2(x) - 48} \\
 \int \tan^2 x \sec^7 x dx &= \frac{5\log(\sin(x) - 1)}{256} - \frac{5\log(\sin(x) + 1)}{256} \\
 &\quad - \frac{-15\sin^7(x) + 55\sin^5(x) - 73\sin^3(x) - 15\sin(x)}{384\sin^8(x) - 1536\sin^6(x) + 2304\sin^4(x) - 1536\sin^2(x) + 384} \\
 \int \tan^4 x \sec^1 x dx &= -\frac{3\log(\sin(x) - 1)}{16} + \frac{3\log(\sin(x) + 1)}{16} - \frac{-5\sin^3(x) + 3\sin(x)}{8\sin^4(x) - 16\sin^2(x) + 8} \\
 \int \tan^4 x \sec^3 x dx &= -\frac{\log(\sin(x) - 1)}{32} + \frac{\log(\sin(x) + 1)}{32} - \frac{-3\sin^5(x) - 8\sin^3(x) + 3\sin(x)}{48\sin^6(x) - 144\sin^4(x) + 144\sin^2(x) - 48} \\
 \int \tan^4 x \sec^5 x dx &= -\frac{3\log(\sin(x) - 1)}{256} + \frac{3\log(\sin(x) + 1)}{256} \\
 &\quad - \frac{3\sin^7(x) - 11\sin^5(x) - 11\sin^3(x) + 3\sin(x)}{128\sin^8(x) - 512\sin^6(x) + 768\sin^4(x) - 512\sin^2(x) + 128} \\
 \int \tan^4 x \sec^7 x dx &= -\frac{3\log(\sin(x) - 1)}{512} + \frac{3\log(\sin(x) + 1)}{512} \\
 &\quad + \frac{-15\sin^9(x) + 70\sin^7(x) - 128\sin^5(x) - 70\sin^3(x) + 15\sin(x)}{1280\sin^{10}(x) - 6400\sin^8(x) + 12800\sin^6(x) - 12800\sin^4(x) + 6400\sin^2(x) - 1280} \\
 \int \tan^6 x \sec^1 x dx &= \frac{5\log(\sin(x) - 1)}{32} - \frac{5\log(\sin(x) + 1)}{32} + \frac{-33\sin^5(x) + 40\sin^3(x) - 15\sin(x)}{48\sin^6(x) - 144\sin^4(x) + 144\sin^2(x) - 48} \\
 \int \tan^6 x \sec^3 x dx &= \frac{5\log(\sin(x) - 1)}{256} - \frac{5\log(\sin(x) + 1)}{256} \\
 &\quad - \frac{-15\sin^7(x) - 73\sin^5(x) + 55\sin^3(x) - 15\sin(x)}{384\sin^8(x) - 1536\sin^6(x) + 2304\sin^4(x) - 1536\sin^2(x) + 384} \\
 \int \tan^6 x \sec^5 x dx &= \frac{3\log(\sin(x) - 1)}{512} - \frac{3\log(\sin(x) + 1)}{512} \\
 &\quad + \frac{15\sin^9(x) - 70\sin^7(x) - 128\sin^5(x) + 70\sin^3(x) - 15\sin(x)}{1280\sin^{10}(x) - 6400\sin^8(x) + 12800\sin^6(x) - 12800\sin^4(x) + 6400\sin^2(x) - 1280}
 \end{aligned}$$

This case is considered as the hardest of all cases in $\int \tan^n x \sec^m x \, dx$. If you try to solve it by hand, most calculus book will recommend to use method of substitution to replace $\tan^n x$ into $\sec x$ function so we can then compute the integral of secant with odd power, which already been done previously in level 3 integral section.

The secant with odd power make use of middle coefficient, the sum of two neighbor entry in the vector that will become the constant / coefficient for the numerator with sine function that has odd power.

Here, we will do the same, with an adding of some tricks.

The patterns that can be seen from those equations above are

- The terms are divided into 3, the same like integral of secant with odd power which are:

1. The numerator with $\sin(x)$ function that has odd power
2. The denominator with $\sin(x)$ function that has even power
3. The log terms

- For all $\int \tan^n x \sec^m x \, dx$, when $m = 1$ the result is the same / equivalent with $\int \sec^{n+1} x \, dx$.

To solve for any m that can be any number but 1, we will need to make another **for** loop to adjust the coefficient.

So the starting is we will use $\int \sec^{n+1} x \, dx$ that we already know the pattern.

- The denominator coefficients have the exact same pattern with the integral of secant with odd power, but this time we will make the denominator coefficient to run another loop again till we reach m .

To learn the pattern we will try with small n and m , we will make use of

$$\tan^2 x = \sec^2 x - 1$$

to get everything in terms of $\sec x$.

For $n = 2$ and $m = 3$

$$\begin{aligned}
 \int \tan^2 x \sec^3 x \, dx &= (\sec^2 x - 1) \sec^3 x \, dx \\
 &= \int \sec^5 x - \sec^3 x \, dx \\
 &= \left[-\frac{3 \log(\sin(x) - 1)}{16} + \frac{3 \log(\sin(x) + 1)}{16} - \frac{3 \sin^3 x - 5 \sin x}{8 \sin^4 x - 16 \sin^2 x + 8} \right] \\
 &\quad - \left[-\frac{\log(\sin(x) - 1)}{4} + \frac{\log(\sin(x) + 1)}{4} - \frac{\sin x}{2 \sin^2 x - 2} \right] \\
 &= -\frac{\log(\sin(x) - 1)}{16} + \frac{\log(\sin(x) + 1)}{16} \\
 &\quad - \frac{3 \sin^3 x - 5 \sin x - (4 \sin^2 x - 4)(\sin x)}{8 \sin^4 x - 16 \sin^2 x + 8} \\
 &= -\frac{\log(\sin(x) - 1)}{16} + \frac{\log(\sin(x) + 1)}{16} - \frac{-\sin^3 x - \sin x}{8 \sin^4 x - 16 \sin^2 x + 8}
 \end{aligned}$$

For $n = 4$ and $m = 3$

$$\begin{aligned}
 \int \tan^4 x \sec^3 x \, dx &= (\sec^2 x - 1)^2 \sec^3 x \, dx \\
 &= \int \sec^3 x (\sec^4 x - 2 \sec^2 x + 1) \, dx \\
 &= \int \sec^3 x - 2 \sec^5 x + \sec^7 x \, dx \\
 &= -\frac{\log(\sin(x) - 1)}{32} + \frac{\log(\sin(x) + 1)}{32} + \frac{-3 \sin^5 x - 8 \sin^3 x + 3 \sin x}{48 \sin^6 x - 144 \sin^4 x + 144 \sin^2 x - 48}
 \end{aligned}$$

If we take a closer look and observe at this part $\int \sec^3 x - 2 \sec^5 x + \sec^7 x \, dx$, we are seeing a binomial coefficient / pascal triangle with alternating sign, thus we can use this pattern to compute $\int \tan^n x \sec^m x \, dx$ with n even and m odd for any number of n and m .

```

#include <iostream>
#include "symintegrationc++.h"
#include <vector>

using namespace std;

// Factorial and combinations
Symbolic factorial(int n) {
    if (n <= 1)
    {
        return 1;
    }
    Symbolic result = 1;
    for (int i = 2; i <= n; ++i)
    {
        result *= i;
    }
    return result;
}

```

```
}

Symbolic combinations(int n, int r) {
    if (r < 0 || r > n)
    {
        return 0; // Invalid input
    }
    return factorial(n) / (factorial(r) * factorial(n - r));
}

Symbolic numeratorintegralsecantodd(int n) {
    Symbolic x("x");
    int k = 1, l=2;
    vector<int> v={1};
    vector<int> mc={1}; // to store the middle coefficient
    Symbolic sgn = -1;
    Symbolic integral_numerator;
    Symbolic d0 = 2, d1 = 2;
    int j =1;
    int m = n-(0.5*(n+1));
    int last_coeff;
    int first_coeff = 3;

    // For the coefficient at the numerator of sine with odd power
    for(int i = 1 ; i < (n-1)/2 ; i = i+1)
    {
        if (i >= 2)
        {
            for(int ic = 1 ; ic < i ; ic = ic+1)
            {
                mc[ic-1] = v[ic-1] + v[ic];
            }
        }

        k = k*1;
        last_coeff = v[j-1];
        v[0] = v[0]*(first_coeff+2*(i-1));
        v.assign({v[0]});

        for(int ic = 1 ; ic < i ; ic = ic+1)
        {
            v.push_back(mc[ic-1]*(first_coeff+2*(i-1)));
        }
        d1 = d1*(2*i);
        d0 = d0*(2+2*i);
        v.push_back(last_coeff*(first_coeff+2*(i-1))+k);
        l = l+2;
        j = j+1;
    }
}
```

```

    }
    int j_num = 0 ;
    for(int i = n-2 ; i >= 1 ; i = i-2)
    {
        integral_numerator += sgn*v[j_num]*((sin(x))^(i));
        sgn = -sgn;
        j_num = j_num+1;
    }
    return integral_numerator;
}

Symbolic denominatorintegralsecantodd(int n) {
    Symbolic x("x");
    int k = 1, l=2;
    vector<int> v={1};
    vector<int> mc={1}; // to store the middle coefficient
    Symbolic sgn = 1;
    Symbolic integral_denominator;
    Symbolic d0 = 2, d1 = 2;
    int j =1;
    int m = n-(0.5*(n+1));
    int last_coeff;
    int first_coeff = 3;

    // For the coefficient at the numerator of sine with odd power
    for(int i = 1 ; i < (n-1)/2 ; i = i+1)
    {
        if (i >= 2)
        {
            for(int ic = 1 ; ic < i ; ic = ic+1)
            {
                mc[ic-1] = v[ic-1] + v[ic];
            }
        }

        k = k*1;
        last_coeff = v[j-1];
        v[0] = v[0]*(first_coeff+2*(i-1));
        v.assign({v[0]});

        for(int ic = 1 ; ic < i ; ic = ic+1)
        {
            v.push_back(mc[ic-1]*(first_coeff+2*(i-1)));
        }
        d1 = d1*(2*i);
        d0 = d0*(2+2*i);
        v.push_back(last_coeff*(first_coeff+2*(i-1))+k);
        l = l+2;
    }
}

```



```
        j = j+1;
    }
    j = 1;
    for(int i = n-1 ; i >= 0 ; i = i-2)
    {
        integral_denominator += sgn*d0*combinations(m,j-1)*((sin(x))^(
            i));
        sgn = -sgn;
        j = j+1;
    }
    return integral_denominator;
}

Symbolic logtermsintegralsecantodd(int n) {
    Symbolic x("x");
    int k = 1, l=2;
    vector<int> v={1};
    vector<int> mc={1}; // to store the middle coefficient
    Symbolic sgn = -1;
    Symbolic integral_denominator;
    Symbolic d0 = 2, d1 = 2;
    int j =1;
    int m = n-(0.5*(n+1));
    int last_coeff;
    int first_coeff = 3;

    // For the coefficient at the numerator of sine with odd power
    for(int i = 1 ; i < (n-1)/2 ; i = i+1)
    {
        if (i >= 2)
        {
            for(int ic = 1 ; ic < i ; ic = ic+1)
            {
                mc[ic-1] = v[ic-1] + v[ic];
            }
        }

        k = k*1;
        last_coeff = v[j-1];
        v[0] = v[0]*(first_coeff+2*(i-1));
        v.assign({v[0]});

        for(int ic = 1 ; ic < i ; ic = ic+1)
        {
            v.push_back(mc[ic-1]*(first_coeff+2*(i-1)));
        }
        d1 = d1*(2*i);
    }
}
```

```

        d0 = d0*(2+2*i);
        v.push_back(last_coeff*(first_coeff+2*(i-1))+k);
        l = l+2;
        j = j+1;
    }
    d1 = d1*(n-1);
    return (-v[0]*ln(sin(x)-1))/(d1) + (v[0]*ln(sin(x)+1))/(d1) ;
}

int main(void)
{
    Symbolic x("x");
    int bpowersec = 3;
    int bpowertan = 6;
    int bpower = bpowersec+bpowertan;
    Symbolic sgn = 1;
    Symbolic result;
    int j = 1;
    int m = bpowertan/2;
    for(int i = bpower ; i >= bpowersec ; i = i-2)
    {
        result += sgn*combinations(m,j-1)*( (
            numeratorintegralsecantodd(i)/denominatorintegralsecantodd(
            i)) + logtermsintegralsecantodd(i));
        sgn = -sgn;
        j = j+1;
    }
    //cout << "integral at numerator= "<< numeratorintegralsecantodd(3)<<
    endl;
    //cout << "integral at denominator= "<< denominatorintegralsecantodd
    (3)<< endl;
    //cout << "log terms= "<< logtermsintegralsecantodd(3)<< endl;
    cout << "for secant power of " << bpowersec << " and tangent power of
    " << bpowertan << endl;
    cout << "integral = "<< result << endl;
    //cout << "integral with subs = " << result[x==1] << endl;
    return 0;
}

```

Code 26: level 4 integral for tangent even power and secant odd power case

It is recommended that you use the:

```

cout << "integral at numerator= "<< numeratorintegralsecantodd(3)<< endl;
cout << "integral at denominator= "<< denominatorintegralsecantodd(3)<< endl;
cout << "log terms= "<< logtermsintegralsecantodd(3)<< endl;
(uncomment them in the C++ code)

```

and comment the :

```

cout << "for secant power of " << bpowersec << " and tangent power of " << bpowertan << endl;

```

```
cout << "integral = " << result << endl;
```

Because it is faster to show the result when we only show the log terms function, numerator only / denominator only of the fraction with sine function than to show it in the full fraction function, it is still slow and we wonder why, because SymPy is able to show the result very fast, knowing that C++ is basically / by default faster than SymPy we might suspect because it is the fraction that we are using combining with symbolic computation, if we only show the vector / array, the result can come up very fast, faster than SymPy.

We will figure it out and make it really fast one day. Any reader interested is always welcome to give a nice input for this.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Trigonometry Int
egration Level 4 Sec^m Tan^n for n Even m Odd with Functions ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Trigonometry Int
egration Level 4 Sec^m Tan^n for n Even m Odd with Functions ]# ./main
for secant power of 3 and tangent power of 6
integral = -105*sin(x)^7*(384*sin(x)^8)-1536*sin(x)^6+2304*sin(x)^4-1536*
sin(x)^2+384*(-1)+385*sin(x)^5*(384*sin(x)^8)-1536*sin(x)^6+2304*sin(x)^
(4)-1536*sin(x)^2+384*(-1)-511*sin(x)^3*(384*sin(x)^8)-1536*sin(x)^6+230
4*sin(x)^4-1536*sin(x)^2+384*(-1)+279*sin(x)*(384*sin(x)^8)-1536*sin(x)^6
)+2304*sin(x)^4-1536*sin(x)^2+384*(-1)+5/256*ln(sin(x)-1)-5/256*ln(sin(x)+1
)+45*sin(x)^5*(48*sin(x)^6)-144*sin(x)^4+144*sin(x)^2-48*(-1)-120*sin(x)^
^3*(48*sin(x)^6)-144*sin(x)^4+144*sin(x)^2-48*(-1)+99*sin(x)*(48*sin(x)^
(6)-144*sin(x)^4+144*sin(x)^2-48*(-1)-9*sin(x)^3*(8*sin(x)^4)-16*sin(x)^
(2)+8*(-1)+15*sin(x)*(8*sin(x)^4)-16*sin(x)^2+8*(-1)+sin(x)*(2*sin(x)^2-2
)^(-1)
julia> integrate((sec(x)^3)*(tan(x)^6),x)
- 15*sin(x) - 73*sin(x) + 55*sin(x) - 15*sin(x) + 5*log(sin(x)
- 384*sin(x) - 1536*sin(x) + 2304*sin(x) - 1536*sin(x) + 384
+ 5*log(sin(x)
- 1) - 5*log(sin(x) + 1)
256
julia> integrate((sec(x)^4)*(tan(x)^6),x)

```

Figure 2.2: The computation of $\int \tan^6 x \sec^3 x dx$ with SymPy and SymIntegration (SymIntegration/Examples/Calculus/Test SymIntegration Trigonometry Integration Level 4 Sec^m Tanⁿ for n Even m Odd with Functions/main.cpp).

Case 2: For $\int \tan^n x \sec^m x dx$ with n odd

We divide this case into 2 smaller cases:

1. When m even
2. When m odd

Case 2-a: For $\int \tan^n x \sec^m x dx$ with n odd and m even

$$\int \tan^1 x \sec^2 x dx = \frac{1}{2 \cos^2(x)}$$

$$\int \tan^1 x \sec^4 x dx = \frac{1}{4 \cos^4(x)}$$

$$\int \tan^1 x \sec^6 x dx = \frac{1}{6 \cos^6(x)}$$

$$\int \tan^1 x \sec^8 x dx = \frac{1}{8 \cos^8(x)}$$

$$\begin{aligned}
\int \tan^3 x \sec^2 x \, dx &= \frac{1 - 2 \cos^2(x)}{4 \cos^4(x)} \\
\int \tan^3 x \sec^4 x \, dx &= \frac{2 - 3 \cos^2(x)}{12 \cos^6(x)} \\
\int \tan^3 x \sec^6 x \, dx &= \frac{3 - 4 \cos^2(x)}{24 \cos^8(x)} \\
\int \tan^3 x \sec^8 x \, dx &= \frac{4 - 5 \cos^2(x)}{40 \cos^{10}(x)} \\
\int \tan^5 x \sec^2 x \, dx &= \frac{-(-3 \cos^4(x) + 3 \cos^2(x) - 1)}{6 \cos^6(x)} \\
\int \tan^5 x \sec^4 x \, dx &= \frac{-(-6 \cos^4(x) + 8 \cos^2(x) - 3)}{24 \cos^8(x)} \\
\int \tan^5 x \sec^6 x \, dx &= \frac{-(-10 \cos^4(x) + 15 \cos^2(x) - 6)}{60 \cos^{10}(x)} \\
\int \tan^5 x \sec^8 x \, dx &= \frac{-(-15 \cos^4(x) + 24 \cos^2(x) - 10)}{120 \cos^{12}(x)}
\end{aligned}$$

The patterns that can be seen from those equations above are

- They are all function that only consist of function of cosine and constants.
- The power of cosine in the denominator is $m + n - 1$.
- You can solve it easily to determine the pattern by substituting

$$\sec^2 x = \tan^2 x + 1$$

then we will have the terms in $\tan(x)$ with odd power only, which we already have learned the pattern from the previous section.

For example, let $m = 4, n = 3$

$$\begin{aligned}
\int \tan^3 x \sec^4 x \, dx &= \int \tan^3 x (\tan^2 x + 1)^2 dx \\
&= \int \tan^3 x (\tan^4 x + 2 \tan^2 x + 1) dx \\
&= \int \tan^7 x + 2 \tan^5 x + \tan^3 x \, dx
\end{aligned}$$

It is using combination / Pascal' triangle again, this time the sign is the same, all positive.

```

#include <iostream>
#include "symintegrationc++.h"
#include <vector>

using namespace std;

// Factorial and combinations
Symbolic factorial(int n) {
    if (n <= 1)

```

```
{
    return 1;
}
Symbolic result = 1;
for (int i = 2; i <= n; ++i)
{
    result *= i;
}
return result;
}

Symbolic combinations(int n, int r) {
    if (r < 0 || r > n)
    {
        return 0; // Invalid input
    }
    return factorial(n) / (factorial(r) * factorial(n - r));
}

Symbolic integraltangentodd(int n) {
    Symbolic x("x");
    Symbolic integral;
    Symbolic integral_front;
    Symbolic sgn = 1;
    vector<int> v={1};
    vector<int> v_temp={1};
    vector<int> mc={1}; // to store the middle coefficient
    vector<int> mc_temp={1}; // to store the temporary / new middle
        coefficient
    int d0 = 2;
    int k = 1, l=2;
    // For the coefficient at the numerator
    for(int i = 1 ; i <= (n-1)/2 ; i = i+1)
    {
        if (i == 1)
        {
            v[0] = 1;
            v.assign({v[0]});
        }

        if (i ==2)
        {
            mc[0]=1;
            mc.assign({mc[0]});
            v_temp[0] = v[0]*(i-1);
            v_temp.assign({v_temp[0]});

            for(int j = 1 ; j < i ; j = j+1)
```

```

        {
            v_temp.push_back(v[j-1]*((2*i)-j) + v_temp[0]*mc
                [j-1] );
        }
        v[0] = v_temp[0];
        v.assign({v[0]});
        for(int j = 1 ; j < i ; j = j+1)
        {
            v.push_back(v_temp[j]);
        }
    }
    if (i == 3)
    {
        mc[0] = mc[0] + 1; //
        mc.assign({mc[0]});

        v_temp[0] = v[0]*(i-1);
        v_temp.assign({v_temp[0]});
        for(int j = 1 ; j < i-1 ; j = j+1)
        {
            v_temp.push_back(v[j-1]*((2*i)-j) + v_temp[0]*mc
                [j-1] );
        }
        v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
        // Assign the temporary vector to vector v for future
        use
        v[0] = v_temp[0];
        v.assign({v[0]});
        for(int j = 1 ; j < i ; j = j+1)
        {
            v.push_back(v_temp[j]);
        }
    }
    if (i == 4)
    {
        mc[0] = mc[0] + 1;
        mc.assign({mc[0]});
        mc.push_back(mc[0]);

        v_temp[0] = v[0]*(i-1);
        v_temp.assign({v_temp[0]});
        for(int j = 1 ; j < i-1 ; j = j+1)
        {
            v_temp.push_back(v[j-1]*((2*i)-j) + v_temp[0]*mc
                [j-1] );
        }
        v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
        // Assign the temporary vector to vector v for future

```

```

        use
        v[0] = v_temp[0];
        v.assign({v[0]});
        for(int j = 1 ; j < i ; j = j+1)
        {
            v.push_back(v_temp[j]);
        }
    }
    if (i >= 5)
    {
        mc_temp[0] = mc[0]+1;
        mc_temp.assign({mc_temp[0]});
        for(int ic = 1 ; ic < l ; ic = ic+1)
        {
            mc_temp[ic] = mc[ic-1] + mc[ic];
        }
        mc[1] = (mc_temp[0]);
        mc[l+1] = (mc_temp[0]);

        mc[0]=mc_temp[0];
        mc.assign({mc[0]});

        for(int ic = 1 ; ic < l ; ic = ic+1)
        {
            mc.push_back(mc_temp[ic]);
        }
        mc.push_back(mc_temp[0]);
        mc.push_back(0);

        v_temp[0] = v[0]*(i-1);
        v_temp.assign({v_temp[0]});
        for(int j = 1 ; j < i-1 ; j = j+1)
        {
            v_temp.push_back(v[j-1]*((2*i)-j) + v_temp[0]*mc
                             [j-1] );
        }
        v_temp.push_back((v[i-2]*(i+1) ) + (v_temp[0]) );
        // Assign the temporary vector to vector v for future
        use
        v[0] = v_temp[0];
        v.assign({v[0]});
        for(int j = 1 ; j < i ; j = j+1)
        {
            v.push_back(v_temp[j]);
        }

        l = l+1;
    }
}

```

```

        d0 = d0*k;
        k = k+1;
    }
    for(int i = 1 ; i <= (n-1)/2 ; i = i+1)
    {
        integral += sgn*v[i-1]*(cos(x)^(2*(i-1)));
        sgn=-sgn;
    }
    sgn = 1;
    for(int i = 1 ; i <= (n-1)/2 ; i = i+1)
    {
        integral_front = sgn;
        sgn = -sgn;
    }
    integral = integral_front*ln(cos(x)) + (integral)/(d0*(cos(x)^(n-1)))
    ;
    return integral;
}

int main(void)
{
    Symbolic x("x");
    int bpowersec = 4;
    int bpowertan = 5;
    int bpower = bpowersec+bpowertan;
    Symbolic result;
    int j = 1;
    int m = bpowersec/2;
    for(int i = bpowertan ; i <= bpower ; i = i+2)
    {
        result += combinations(m,j-1)*integralextantodd(i);
        j = j+1;
    }
    cout << "for secant power of " << bpowersec << " and tangent power of
        " << bpowertan << endl;
    cout << "integral = "<< result<< endl;
    return 0;
}

```

Code 27: level 4 integral for tangent odd power and secant even power case**Case 2-b:** For $\int \tan^n x \sec^m x dx$ with n odd and m odd

$$\begin{aligned}\int \tan^1 x \sec^1 x \, dx &= \frac{1}{\cos(x)} \\ \int \tan^1 x \sec^3 x \, dx &= \frac{1}{3 \cos^3(x)} \\ \int \tan^1 x \sec^5 x \, dx &= \frac{1}{5 \cos^5(x)} \\ \int \tan^1 x \sec^7 x \, dx &= \frac{1}{7 \cos^7(x)}\end{aligned}$$

$$\begin{aligned}\int \tan^3 x \sec^1 x \, dx &= \frac{1 - 3 \cos^2(x)}{3 \cos^3(x)} \\ \int \tan^3 x \sec^3 x \, dx &= \frac{3 - 5 \cos^2(x)}{15 \cos^5(x)} \\ \int \tan^3 x \sec^5 x \, dx &= \frac{5 - 7 \cos^2(x)}{35 \cos^7(x)} \\ \int \tan^3 x \sec^7 x \, dx &= \frac{7 - 9 \cos^2(x)}{63 \cos^9(x)}\end{aligned}$$

$$\begin{aligned}\int \tan^5 x \sec^1 x \, dx &= \frac{-(-15 \cos^4(x) + 10 \cos^2(x) - 3)}{15 \cos^5(x)} \\ \int \tan^5 x \sec^3 x \, dx &= \frac{-(-35 \cos^4(x) + 42 \cos^2(x) - 15)}{105 \cos^7(x)} \\ \int \tan^5 x \sec^5 x \, dx &= \frac{-(-63 \cos^4(x) + 90 \cos^2(x) - 35)}{315 \cos^9(x)} \\ \int \tan^5 x \sec^7 x \, dx &= \frac{-(-99 \cos^4(x) + 154 \cos^2(x) - 63)}{693 \cos^{11}(x)}\end{aligned}$$

The patterns that can be seen from those equations above are

- They are all function that only consist of function of cosine and constants.
- The power of cosine in the denominator is $m + n - 1$.
- You can solve it easily to determine the pattern by substituting

$$\tan^2 x = \sec^2 x - 1$$

and then use method of substitution by assigning

$$\begin{aligned}u &= \sec x \\ du &= \sec x \tan x \, dx\end{aligned}$$

then we will have the terms in $\sec(x)$ with even power only, which we already have learned the pattern from the previous section.

For example, let $m = 3, n = 5$

$$\begin{aligned}
 \int \tan^5 x \sec^3 x \, dx &= \int \tan^4 x \sec^2 x (\sec x \tan x \, dx) \\
 &= \int [(\sec^2 x - 1)^2 \sec^2 x] (\sec x \tan x \, dx) \\
 &= \int [(\sec^4 x - 2\sec^2 x + 1) \sec^2 x] (\sec x \tan x \, dx) \\
 &= \int [\sec^6 x - 2\sec^4 x + \sec^2 x] (\sec x \tan x \, dx) \\
 &= \int u^6 - 2u^4 + u^2 \, du \\
 &= \frac{1}{7}u^7 - \frac{2}{5}u^5 + \frac{1}{3}u^3 \\
 &= \frac{1}{7}\sec^7 x - \frac{2}{5}\sec^5 x + \frac{1}{3}\sec^3 x \\
 &= \frac{1}{7}\frac{1}{\cos^7 x} - \frac{2}{5}\frac{1}{\cos^5 x} + \frac{1}{3}\frac{1}{\cos^3 x}
 \end{aligned}$$

It is using combination / Pascal' triangle combining with a method of substitution, the sign is alternating this time. The thing is the method of substitution is a really helpful here, we do not even need to create a function, only integrating ordinary polynomial and then substituting back, we can substitute $u = \sec(x)$ or $u = \frac{1}{\cos(x)}$.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Trigonometry Int
egration Level 4 Sec^m Tan^n for n Odd m Odd with Functions ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Trigonometry Integration Lev
el 4 Sec^m Tan^n for n Odd m Odd with Functions ]# ./main
for secant power of 3 and tangent power of 5
we will compute the integral of u^2-2*u^4+u^6
integral = 1/3*u^3-2/5*u^5+1/7*u^7
Substitute back, integral = 1/3*cos(x)^(-3)-2/5*cos(x)^(-5)+1/7*cos(x)^(-7)
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Trigonometry Integration Lev
el 4 Sec^m Tan^n for n Odd m Odd with Functions ]#

```



```

julia> integrate((sec(x)^3)*(tan(x)^5),x)
- ( 4      2
  - 35*cos (x) + 42*cos (x) - 15 )
  7
 105*cos (x)

```

Figure 2.3: The computation of $\int \tan^5 x \sec^3 x \, dx$ with SymPy and SymIntegration (*SymIntegration/Examples/Calculus/Test SymIntegration Trigonometry Integration Level 4 Sec^m Tanⁿ for n Odd m Odd/main.cpp*).

```

#include <iostream>
#include "symintegrationc++.h"
#include <vector>

using namespace std;

```

```
// Factorial and combinations
Symbolic factorial(int n) {
    if (n <= 1)
    {
        return 1;
    }
    Symbolic result = 1;
    for (int i = 2; i <= n; ++i)
    {
        result *= i;
    }
    return result;
}


Symbolic combinations(int n, int r) {
    if (r < 0 || r > n)
    {
        return 0; // Invalid input
    }
    return factorial(n) / (factorial(r) * factorial(n - r));
}

int main(void)
{
    Symbolic u("u"), x("x");
    int bpowersec = 3;
    int bpowertan = 5;
    int bpower = bpowersec+bpowertan;
    Symbolic result;
    int j = 1;
    int m = (bpowertan-1)/2;
    Symbolic sgn = 1;
    for(int i = bpowersec-1 ; i <= bpower-2 ; i = i+2)
    {
        result += sgn*combinations(m,j-1)*(u^i);
        j = j+1;
        sgn = -sgn;
    }
    cout << "for secant power of " << bpowersec << " and tangent power of "
          << bpowertan << endl;
    cout << "we will compute the integral of "<< result << endl;
    Symbolic f = integrate(result,u);
    cout << "integral = "<< f << endl;
    cout << "Substitute back, integral = "<< f[u==(1/cos(x))] << endl;
    return 0;
}
```

Code 28: level 4 integral for tangent odd power and secant odd power case

ix. $\int \cot^n x \csc^m x dx$

The formula of $\int \cot^n x \csc^m x dx$ has similar pattern with $\int \tan^n x \sec^m x dx$, we only change from $\sin(x)$ to $\cos(x)$ and from $\cos(x)$ to $\sin(x)$, the sign will be changed too, from minus to plus, and from plus to minus. Once we have decoded the pattern for $\int \tan^n x \sec^m x dx$, we only need to modify a bit for this integral type.



```

julia> integrate((tan(x)^2)*(sec(x)^7),x)
      7      5      3      2      1
      - 15·sin(x) + 55·sin(x) - 73·sin(x) - 15·sin(x) + 5·log(sin(x))
      -----
      384·sin(x) - 1536·sin(x) + 2304·sin(x) - 1536·sin(x) + 384
      - 1) 5·log(sin(x) + 1)
      -----
      256

julia> integrate((cot(x)^2)*(csc(x)^7),x)
      7      5      3      2      1
      - 15·cos(x) + 55·cos(x) - 73·cos(x) - 15·cos(x) + 5·log(cos(x))
      -----
      384·cos(x) - 1536·cos(x) + 2304·cos(x) - 1536·cos(x) + 384
      - 1) 5·log(cos(x) + 1)
      -----
      256

julia> integrate((tan(x)^2)*(sec(x)^8),x)
      16·sin(x)      8·sin(x)      2·sin(x)      sin(x)      sin(x)
      ----- - ----- - ----- - ----- + -----
      315·cos(x)      315·cos(x)      105·cos(x)      63·cos(x)      9·cos(x)

julia> integrate((cot(x)^2)*(csc(x)^8),x)
      16·cos(x)      8·cos(x)      2·cos(x)      cos(x)      cos(x)
      ----- + ----- + ----- + ----- - -----
      315·sin(x)      315·sin(x)      105·sin(x)      63·sin(x)      9·sin(x)

```

Figure 2.4: The integral of $\int \cot^n x \csc^m x dx$ that has similar pattern with of $\int \tan^n x \sec^m x dx$.

VII. TRIGONOMETRY AND TRANSCENDENTALS FORMULA

We will list all the trigonometry and transcendentals (exponential and logarithm) formulas that can be used to help you solve integral problems

[SI*] Trigonometry

Reciprocal identities

$$\begin{aligned}\tan x &= \frac{\sin x}{\cos x} \\ \cot x &= \frac{\cos x}{\sin x} \\ \csc x &= \frac{1}{\sin x} \\ \sec x &= \frac{1}{\cos x}\end{aligned}$$

Pythagorean identities

$$\begin{aligned}\sin^2 x + \cos^2 x &= 1 \\ \tan^2 x &= \sec^2 x - 1 \\ \cot^2 x &= \csc^2 x - 1\end{aligned}$$

Sum and difference identities

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \sin(x - y) &= \sin x \cos y - \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y \\ \cos(x - y) &= \cos x \cos y + \sin x \sin y \\ \tan(x + y) &= \frac{\tan x + \tan y}{1 - \tan x \tan y} \\ \tan(x - y) &= \frac{\tan x - \tan y}{1 + \tan x \tan y}\end{aligned}$$

Double-angle identities

$$\begin{aligned}\sin 2x &= 2 \sin x \cos x = \frac{2 \tan x}{1 + \tan^2 x} \\ \cos 2x &= \cos^2 x - \sin^2 x \\ \cos 2x &= 2 \cos^2 x - 1 \\ \cos 2x &= 1 - 2 \sin^2 x \\ \cos 2x &= \frac{1 - \tan^2 x}{1 + \tan^2 x} \\ \tan 2x &= \frac{2 \tan x}{1 - \tan^2 x} \\ \cot 2x &= \frac{\cot^2 x - 1}{2 \cot x}\end{aligned}$$

Half-angle identities

$$\begin{aligned}\sin \frac{x}{2} &= \pm \sqrt{\frac{1 - \cos x}{2}} \\ \cos \frac{x}{2} &= \pm \sqrt{\frac{1 + \cos x}{2}} \\ \tan \frac{x}{2} &= \pm \sqrt{\frac{1 - \cos x}{1 + \cos x}} = \frac{\sin x}{1 + \cos x} = \frac{1 - \cos x}{\sin x}\end{aligned}$$

Integral

$$\begin{aligned}\int \sin x \, dx &= -\cos x \\ \int \cos x \, dx &= \sin x \\ \int \tan x \, dx &= -\log(\cos(x)) \\ \int \sec x \, dx &= -\frac{\log(\sin(x) - 1)}{2} + \frac{\log(\sin(x) + 1)}{2} \\ \int \csc x \, dx &= \frac{\log(\cos(x) - 1)}{2} - \frac{\log(\cos(x) + 1)}{2} \\ \int \cot x \, dx &= \log(\sin(x))\end{aligned}$$

Inverse Trigonometric integral

$$\begin{aligned}\int \frac{1}{1+x^2} dx &= \operatorname{atan}(x) \\ \int \frac{1}{a+bx^2} dx &= -\frac{\sqrt{\frac{-1}{ab}} \ln\left(-a\sqrt{\frac{-1}{ab}} + x\right)}{2} + \frac{\sqrt{\frac{-1}{ab}} \ln\left(a\sqrt{\frac{-1}{ab}} + x\right)}{2}\end{aligned}$$

Inverse hyperbolic trigonometry integral

$$\begin{aligned}\int \frac{1}{\sqrt{1+x^2}} dx &= \sinh^{-1}(x) + C \\ \int \frac{1}{\sqrt{x^2-1}} dx &= \cosh^{-1}(x) + C \\ \int \frac{1}{1-x^2} dx &= \tanh^{-1}(x) + C, \quad |x| < 1 \\ \int \frac{1}{1-x^2} dx &= \coth^{-1}(x) + C, \quad |x| > 1 \\ \int \frac{-1}{x\sqrt{1-x^2}} dx &= \operatorname{sech}^{-1}(x) + C \\ \int \frac{-1}{|x|\sqrt{1+x^2}} dx &= \operatorname{csch}^{-1}(x) + C\end{aligned}$$

Note that the derivatives of $\tanh^{-1}(x)$ and $\coth^{-1}(x)$ are the same. Thus, when we compute $\int \frac{1}{1-x^2} dx$, we need to select the proper antiderivative based on the domain of the functions and the values of x .

$$\int \frac{1}{1-x^2} dx = \begin{cases} \tanh^{-1}(x) + C, & |x| < 1 \\ \coth^{-1}(x) + C, & |x| > 1 \end{cases}$$

Derivative

$$\frac{d}{dx} \sin x = \cos x$$

$$\frac{d}{dx} \cos x = -\sin x$$

$$\frac{d}{dx} \tan x = \tan^2 x + 1$$

$$\frac{d}{dx} \sec x = \sec(x) \tan(x)$$

$$\frac{d}{dx} \csc x = -\cot(x) \csc(x)$$

$$\frac{d}{dx} \cot x = -\cot^2 x - 1$$

[SI*] Logarithm

Logarithm along with exponential play a very important role in solving differential equations, so we will need to take a look at these formulas often to help us when solving differential equations problem

$$\log_a b = c$$

$$a^c = b$$

$$\log b = c$$

$$e^c = b$$

$$\log_a a = 1$$

$$\log(a * b) = \log(a) + \log(b)$$

$$\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$$

$$\log(a^b) = b \log(a)$$

$$a^{\log_a b} = b$$

$$\log_b a = \frac{\log_c a}{\log_c b}$$

$$\log_b c = \log_b a \log_a c$$

$$\log_{b^n} a^m = \frac{m}{n} \log_b a$$

$$-\log_b a = \log_b \left(\frac{1}{a}\right)$$

$$-\log_b a = \log_{\frac{1}{b}} a$$

$$\frac{d}{dx} \log_a x = \frac{1}{x \ln(a)}$$

$$\frac{d}{dx} \ln x = \frac{1}{x}$$

with $e = 2.718281828459045$ is the Euler' number. When there is no base written in $\log$ function, we will use e as the base number. The base number cannot be equal to 1 and it has to be positive. For all $\log(a)$, a is a real number and has to be positive.

i. Functions in SymIntegration Related to Integration with Transcendental and Trigonometry Functions

[SI*] When we learn second order linear differential equation and want to solve it with variation of parameters, a general method that can be applied to any equation, we will be faced with a lot of combination / product of functions of trigonometry, transcendental and polynomial. In SymIntegration before we call the function **integrate()** we can call this function first, to simplify the function into a simpler function that can be integrated:

simplifybeforeintegrate(const Symbolic &F, const Symbolic &t)

this function will simplify function $F(t)$ into simpler function that can be integrated, for example:

$$\frac{3 \cos t \cos 2t}{\sin 2t} = \frac{3}{2} \csc t - 3 \sin t$$

The list of functions that already in the source code are:

$$F(t) = a \frac{\cos(t) \cos(2bt)}{\sin(2bt)}$$

$$F(t) = a \sin(2bt) \csc(bt)$$

$$F(t) = a \cos(2bt) \csc(bt)$$

with a, b are constants and $2 * b$ has to be an even number.

VIII. INFINITE SERIES

[SI*] Infinite Sequences

A sequence

$$a_1, a_2, a_3, a_4, \dots$$

is an ordered arrangement of real numbers, one for each positive integer. More formally, an infinite sequence is a function whose domain is the set of positive integers and whose range is a set of real numbers. We may denote a sequence by $a_1, a_2, a_3, \dots$, by $\{a_n\}_{n=1}^{\infty}$, or simply by $\{a_n\}$.

A sequence may be specified by giving enough initial terms to establish a pattern, as in

$$1, 4, 7, 10, 13, \dots$$

by an explicit formula for the n th term, as in

$$a_n = 3n - 2, \quad n \geq 1$$

or by a recursion formula

$$a_1 = 1, \quad a_n = a_{n-1} + 3, \quad n \geq 2$$

Definition 2.1: Convergence

The sequence $\{a_n\}$ is said to converge to L , and we write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for each positive number ϵ there is a corresponding positive number N such that

$$n \geq N \rightarrow |a_n - L| < \epsilon$$

A sequence that fails to converge to any finite number L is said to diverge, or to be divergent.

Theorem 2.5: Properties of Limits of Sequences

Let $\{a_n\}$ and $\{b_n\}$ be convergent sequences and k a constant. Then

1. $\lim_{n \rightarrow \infty} k = k$;
2. $\lim_{n \rightarrow \infty} ka_n = k \lim_{n \rightarrow \infty} a_n$;
3. $\lim_{n \rightarrow \infty} (a_n \pm b_n) = \lim_{n \rightarrow \infty} a_n \pm \lim_{n \rightarrow \infty} b_n$;
4. $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = \lim_{n \rightarrow \infty} a_n \cdot \lim_{n \rightarrow \infty} b_n$;
5. $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n}$, provided that $\lim_{n \rightarrow \infty} b_n \neq 0$;

Theorem 2.6: Squeeze Theorem

Suppose that $\{a_n\}$ and $\{c_n\}$ both converge to L and that $a_n \leq b_n \leq c_n$ for $n \geq K$ (K a fixed integer). Then $\{b_n\}$ also converges to L .

Theorem 2.7: Absolute Convergence Theorem

If $\lim_{n \rightarrow \infty} |a_n| = 0$, then $\lim_{n \rightarrow \infty} a_n = 0$.

Since $-|a_n| \leq a_n \leq |a_n|$, the result follows from the Squeeze theorem.

Theorem 2.8: Monotonic Sequence Theorem

If U is an upper bound for a nondecreasing sequence $\{a_n\}$, then the sequence converges to a limit A that is less than or equal to U . Similarly, if L is a lower bound for a nonincreasing sequence $\{b_n\}$, then the sequence $\{b_n\}$ converges to a limit B that is greater than or equal to L .

[SI*] **Example:**

A famous sequence $\{f_n\}$, called the Fibonacci sequence after Leonardo Fibonacci, who introduced it around A.D. 1200, is defined by the recursion formula

$$f_1 = f_2 = 1, \quad f_{n+2} = f_{n+1} + f_n$$

- (a) Find f_3 through f_{10} .
- (b) Let $\phi = \frac{1}{2}(1 + \sqrt{5}) = 1.618034$. The greeks called this number the golden ratio, claiming that a rectangle whose dimensions were in this ratio was "perfect." It can be shown that

$$\begin{aligned} f_n &= \frac{1}{\sqrt{5} \left[\left(\frac{1+\sqrt{5}}{2} \right)^n - \left(\frac{1-\sqrt{5}}{2} \right)^n \right]} \\ &= \frac{1}{\sqrt{5}} [\phi^n - (-1)^n \phi^{-n}] \end{aligned}$$

Check that this gives the right result for $n = 1$ and $n = 2$. The general result can be proved by induction (it is a nice challenge). More in line with this section, use this explicit formula to prove that $\lim_{n \rightarrow \infty} \frac{f_{n+1}}{f_n} = \phi$.

- (c) Using this limit just proved, show that ϕ satisfies the equation $x^2 - x - 1 = 0$. Then, in another interesting twist, use the Quadratic formula to show that the two roots of this equation are ϕ and $-\frac{1}{\phi}$, two numbers that occur in the explicit formula for f_n .

Solution:

- (a) $f_3 = 2, f_4 = 3, f_5 = 5, f_6 = 8, f_7 = 13, f_8 = 21, f_9 = 34, f_{10} = 55$

(b) Using the formula

$$\begin{aligned}
f_1 &= \frac{1}{\sqrt{5}} \left[\frac{1+\sqrt{5}}{2} - \frac{1-\sqrt{5}}{2} \right] = \frac{1}{\sqrt{5}} \left[\frac{2\sqrt{5}}{2} \right] = 1 \\
f_2 &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2} \right)^2 - \left(\frac{1-\sqrt{5}}{2} \right)^2 \right] \\
&= \frac{1}{\sqrt{5}} \left[\frac{1+2\sqrt{5}+5 - (1-2\sqrt{5}+5)}{4} \right] \\
&= \frac{1}{\sqrt{5}} \left[\frac{4\sqrt{5}}{4} \right] = 1
\end{aligned}$$

thus

$$\begin{aligned}
\lim_{n \rightarrow \infty} \frac{f_{n+1}}{f_n} &= \lim_{n \rightarrow \infty} \frac{\phi^{n+1} - (-1)^{n+1}\phi^{-n-1}}{\phi^n - (-1)^n\phi^{-n}} \\
&= \lim_{n \rightarrow \infty} \frac{\phi^{n+1} - \frac{(-1)^{n+1}}{\phi^{n+1}}}{\phi^n - \frac{(-1)^n}{\phi^n}} \\
&= \lim_{n \rightarrow \infty} \frac{\phi - \frac{(-1)^{n+1}}{\phi^{2n+1}}}{1 - \frac{(-1)^n}{\phi^{2n}}} = \phi
\end{aligned}$$

(c)

$$\begin{aligned}
\phi^2 - \phi - 1 &= \left[\frac{1}{2}(1+\sqrt{5}) \right]^2 - \frac{1}{2}(1+\sqrt{5}) - 1 \\
&= \left(\frac{3}{2} + \frac{\sqrt{5}}{2} \right) - \left(\frac{1}{2} + \frac{\sqrt{5}}{2} \right) - 1 \\
&= 0
\end{aligned}$$

Therefore ϕ satisfies $x^2 - x - 1 = 0$. Using the Quadratic Formula on $x^2 - x - 1 = 0$ yields

$$x = \frac{1 \pm \sqrt{1+4}}{2} = \frac{1 \pm \sqrt{5}}{2}$$

thus

$$\begin{aligned}
\phi &= \frac{1+\sqrt{5}}{2} \\
-\frac{1}{\phi} &= -\frac{2}{1+\sqrt{5}} = -\frac{2(1-\sqrt{5})}{1-5} = \frac{1-\sqrt{5}}{2}
\end{aligned}$$

[SI*] Infinite Series

Many algorithms in a computer to compute the sine of an angle, or e to some power are based on infinite series. Generally, infinite series can be denoted by

$$\sum_{k=1}^{\infty} a_k \tag{2.17}$$

a_k is the k -th term of the series. Then S_n , the partial n -th sum, is given by

$$S_n = a_1 + a_2 + a_3 + \cdots + a_n = \sum_{k=1}^n a_k \quad (2.18)$$

Definition 2.2: Converges and Diverges

The infinite series $\sum_{k=1}^{\infty} a_k$ converges and has sum S if the sequence of partial sums $\{S_n\}$ converges to S . If $\{S_n\}$ diverges, then the series diverges. A divergent series has no sum.

Geometric Series

A series of the form

$$\sum_{k=1}^{\infty} ar^{k-1} = a + ar + ar^2 + ar^3 + \cdots \quad (2.19)$$

where $a \neq 0$, is called a geometric series.

Let $S_n = a + ar + ar^2 + \cdots + ar^{n-1}$. If $r = 1$, $S_n = na$, which grows without bound, and so $\{S_n\}$ diverges. If $r \neq 1$, we may write

$$\begin{aligned} S_n - rS_n &= (a + ar + ar^2 + \cdots + ar^{n-1}) - (ar + ar^2 + ar^3 + \cdots + ar^n) \\ &= a - ar^n \end{aligned}$$

and so

$$S_n = \frac{a - ar^n}{1 - r} = \frac{a}{1 - r} - \frac{a}{1 - r} r^n$$

If $|r| < 1$, then $\lim_{n \rightarrow \infty} r^n = 0$, and thus

$$S = \lim_{n \rightarrow \infty} S_n = \frac{a}{1 - r}$$

If $|r| > 1$ or $r = -1$, the sequence $\{r^n\}$ diverges, and consequently so does $\{S_n\}$.

[SI*] **Positive Series**

[SI*] **Alternating Series, Absolute Convergence, and Conditional Convergence**

[SI*] **Power Series**

[SI*] **Operations on Power Series**

[SI*] **Taylor and Maclaurin Series**

[SI*] **The Taylor Approximation to a Function**

IX. VECTOR CALCULUS

[SI*] **Vector Fields with Div, Grad, Curl**

Vector calculus talks about vector-valued functions, that is, functions whose input is a real

number and whose output is a vector.

In vector calculus we will use the operator ∇ , read: del, or nabla, it is an operator used in mathematics as a vector differential operator. When applied to a function defined on a one-dimensional domain, it denotes the standard derivative of the function as defined in calculus.

When applied to a field (a function defined on a multi-dimensional domain), it may denote any one of three operations depending on the way it is applied: the gradient or steepest slope of a scalar field; the divergence of a vector field; or the curl (rotation) of a vector field.

Del is a very convenient mathematical notation for those three operations (gradient, divergent, and curl) that makes many equations easier to write and remember. The del symbol can be formally defined as a vector operator whose components are the corresponding partial derivative operators. As a vector operator, it can act on scalar and vector fields in three different ways, giving rise to three different differential operations:

1. It acts on a scalar field ($f(x, y, z)$) by a formal scalar multiplication-to produce a vector field (∇f) called the gradient.
2. It acts on a vector field ($F(x, y, z)$) by a formal dot product-to produce a scalar field ($\nabla \cdot F$) called the divergence.
3. It acts on a vector field ($F(x, y, z)$) by a formal cross product-to produce a vector field ($\nabla \times F$) called the curl.

These three uses are summarized as:

Gradient

$$\text{grad } f = \nabla f \quad (2.20)$$

Physical interpretation:

Points in the direction of the greatest rate of increase of the scalar field, and its magnitude is that rate of increase.

Example:

Consider a scalar field representing temperature $T(x, y, z)$ in a garden. The gradient ∇T would give a vector at each point indicating the direction in which the temperature increases most rapidly, and its magnitude would be the rate of that increase.

Divergent

$$\text{div } v = \nabla \cdot v \quad (2.21)$$

Physical interpretation:

It measures the "outward flux per unit volume" of a vector field at a point. It quantifies how much a vector field is "spreading out" or "compressing" at that point.

Example:

For a vector field representing fluid flow $v(x, y, z)$, the divergence $\nabla \cdot v$ indicates whether the fluid is expanding (positive divergence, like a source) or contracting (negative divergence, like a sink). If $\nabla \cdot v = 0$, the fluid is incompressible.

Curl

$$\text{curl } v = \nabla \times v \quad (2.22)$$

Physical interpretation:

It measures the "rotation" or "circulation" of a vector field at a point. The direction of the

curl vector indicates the axis of rotation, and its magnitude indicates the strength of the rotation.

Example:

For a vector field representing the velocity of a fluid $v(x, y, z)$, the curl $\nabla \times v$ indicates the local angular velocity of the fluid. If $\nabla \times v = \mathbf{0}$, the fluid is irrotational.

The divergence of a curl is zero

$$\nabla \cdot (\nabla \times F) = 0 \quad (2.23)$$

This means that the divergence of any curl field is always zero, in linear algebra it is a dot product of an orthogonal vector.

Curl of the gradient is zero

$$\nabla \times (\nabla f) = \mathbf{0} \quad (2.24)$$

This means that the curl of any gradient field (which is a conservative vector field) is always the zero vector.

The div, grad, curl operators are crucial in various fields, including fluid dynamics, electromagnetism (Maxwell's equations), and heat transfer.

Definition 2.3: Mathematical Vector Fields

A vector field in 2-dimensional space is function whose value at a point (x, y) is a 2-dimensional vector $F(x, y)$. Similarly, in 3-dimensional space, a vector field is a function $F(x, y, z)$ whose value at the point (x, y, z) is a 3-dimensional vector.

If $F(x, y, z)$ is a vector, it has i, j , and k components. Each of these components is a scalar function of the point (x, y, z) , so we will often write

$$F(x, y, z) = F_1(x, y, z)\mathbf{i} + F_2(x, y, z)\mathbf{j} + F_3(x, y, z)\mathbf{k} = \langle F_1, F_2, F_3 \rangle \quad (2.25)$$

For example, if $F(x, y, z) = \langle x^2, xy \tan(3z), \ln(y^2) \rangle$, then

$$F_1(x, y, z) = x^2$$

$$F_2(x, y, z) = xy \tan(3z)$$

$$F_3(x, y, z) = \ln(y^2)$$

* The writing of i, j, k components in this book has the same meaning as $\hat{i}, \hat{j}, \hat{k}$ or $\hat{x}, \hat{y}, \hat{z}$ and will be used interchangeably.

[SI*] In three-dimensional Cartesian coordinate system $\mathbb{R}^3$ with coordinates (x, y, z) and standard basis or unit vectors of axes $\{e_x, e_y, e_z\}$, del is written as

$$\nabla = e_x \frac{\partial}{\partial x} + e_y \frac{\partial}{\partial y} + e_z \frac{\partial}{\partial z} = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right) \quad (2.26)$$

As a vector operator, del naturally acts on scalar fields via scalar multiplication, and naturally acts on vector fields via dot products and cross products.

Then using the above definition of ∇ , we may write

$$\nabla f = \left(e_x \frac{\partial}{\partial x} \right) f + \left(e_y \frac{\partial}{\partial y} \right) f + \left(e_z \frac{\partial}{\partial z} \right) f = \frac{\partial f}{\partial x} e_x + \frac{\partial f}{\partial y} e_y + \frac{\partial f}{\partial z} e_z \quad (2.27)$$

and

$$\nabla \cdot \mathbf{F} = \left(e_x \frac{\partial}{\partial x} \cdot \mathbf{F} \right) + \left(e_y \frac{\partial}{\partial y} \cdot \mathbf{F} \right) + \left(e_z \frac{\partial}{\partial z} \cdot \mathbf{F} \right) = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z} \quad (2.28)$$

and

$$\begin{aligned} \nabla \times \mathbf{F} &= \left(e_x \frac{\partial}{\partial x} \times \mathbf{F} \right) + \left(e_y \frac{\partial}{\partial y} \times \mathbf{F} \right) + \left(e_z \frac{\partial}{\partial z} \times \mathbf{F} \right) \\ &= \frac{\partial}{\partial x} (0, -F_z, F_y) + \frac{\partial}{\partial y} (F_z, 0, -F_x) + \frac{\partial}{\partial z} (-F_y, F_x, 0) \\ &= \left(\frac{\partial F_z}{\partial y} - \frac{\partial F_y}{\partial z} \right) e_x + \left(\frac{\partial F_x}{\partial z} - \frac{\partial F_z}{\partial x} \right) e_y + \left(\frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) e_z \end{aligned} \quad (2.29)$$

Del can also be expressed in other coordinate systems, e.g. in cylindrical and spherical coordinates.

[SI*] Gradient

the vector derivative of a scalar field f is called the gradient, and it can be represented as

$$\text{grad } f = \nabla f = \frac{\partial f}{\partial x} \hat{x} + \frac{\partial f}{\partial y} \hat{y} + \frac{\partial f}{\partial z} \hat{z}$$

It always points in the direction of greatest increase of f , and it has a magnitude equal to the maximum rate of increase at the point—just like a standard derivative.

Theorem 2.9: Gradient Identities

- (a) $\nabla(f + g) = \nabla f + \nabla g$
- (b) $\nabla(cf) = c\nabla f$, for any constant c .
- (c) $\nabla(fg) = (\nabla f)g + f(\nabla g)$
- (d) $\nabla \left(\frac{f}{g} \right) = \frac{(g\nabla f - f\nabla g)}{g^2}$ at points x where $g(x) \neq 0$.
- (e) $\nabla(\mathbf{F} \cdot \mathbf{G}) = \mathbf{F} \times (\nabla \times \mathbf{G}) - (\nabla \times \mathbf{F}) \times \mathbf{G} + (\mathbf{G} \cdot \nabla)\mathbf{F} + (\mathbf{F} \cdot \nabla)\mathbf{G}$

Here

$$(\mathbf{G} \cdot \nabla)\mathbf{F} = G_1 \frac{\partial \mathbf{F}}{\partial x} + G_2 \frac{\partial \mathbf{F}}{\partial y} + G_3 \frac{\partial \mathbf{F}}{\partial z}$$

[SI*] Divergence

The divergence of a vector field $\mathbf{v}(x, y, z) = v_x \hat{x} + v_y \hat{y} + v_z \hat{z}$ is a scalar field that can be represented as:

$$\text{div } \mathbf{v} = \nabla \cdot \mathbf{v} = \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z}$$

The divergence is roughly a measure of a vector field's increase in the direction it points; but more accurately, it is a measure of that field's tendency to converge toward or diverge from a point.

Theorem 2.10: Divergent Identities

- (a) $\nabla \cdot (F + G) = \nabla \cdot F + \nabla \cdot G$
- (b) $\nabla \cdot (cF) = c \nabla \cdot F$, for any constant c .
- (c) $\nabla \cdot (fF) = (\nabla f) \cdot F + f \nabla \cdot F$
- (d) $\nabla \cdot (F \times G) = (\nabla \times F) \cdot G - F \cdot (\nabla \times G)$

[SI*] Curl

The curl of a vector field $v(x, y, z) = v_x \hat{x} + v_y \hat{y} + v_z \hat{z}$ is a vector function that can be represented as:

$$\text{curl } v = \nabla \times v = \left(\frac{\partial v_z}{\partial y} - \frac{\partial v_y}{\partial z} \right) \hat{x} + \left(\frac{\partial v_x}{\partial z} - \frac{\partial v_z}{\partial x} \right) \hat{y} + \left(\frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right) \hat{z}$$

The curl at a point is proportional to the on-axis torque that a tiny pinwheel would be subjected to if it were centered at that point.

Theorem 2.11: Curl Identities

- (a) $\nabla \times (F + G) = \nabla \times F + \nabla \times G$
 - (b) $\nabla \times (cF) = c \nabla \times F$, for any constant c .
 - (c) $\nabla \times (fF) = (\nabla f) \times F + f \nabla \times F$
 - (d) $\nabla \times (F \times G) = F(\nabla \cdot G) - (\nabla \cdot F)G + (G \cdot \nabla)F - (F \cdot \nabla)G$
- Here

$$(G \cdot \nabla)F = G_1 \frac{\partial F}{\partial x} + G_2 \frac{\partial F}{\partial y} + G_3 \frac{\partial F}{\partial z}$$

[SI*] In SymIntegration the computation for div, grad, and curl can be handled in **src/divgrad-curl.cpp**.

```
#include "symintegral/symintegrationc++.h"

#ifdef SYMBOLIC_DEFINE
#ifdef SYMINTEGRATION_CPLUSPLUS_DIVGRADCURL_DEFINE
#define SYMINTEGRATION_CPLUSPLUS_DIVGRADCURL_DEFINE

Symbolic div(const Symbolic &F, const Symbolic &x, const Symbolic &y, const
Symbolic &z)
{
    Symbolic sol, i("i"), j("j"), k("k");

    Symbolic F1 = F.coeff(i,1);
    Symbolic F2 = F.coeff(j,1);
    Symbolic F3 = F.coeff(k,1);

    if(F1 != 0 || F2 != 0 || F3 != 0)
```



```
{
    sol = df(F1,x) + df(F2,y) + df(F3,z);
}
else if(F1_hat != 0 || F2_hat != 0 || F3_hat != 0)
{
    sol = df(F1_hat,x) + df(F2_hat,y) + df(F3_hat,z);
}
return sol;
}

Symbolic grad(const Symbolic &F, const Symbolic &x, const Symbolic &y, const
Symbolic &z)
{
    Symbolic sol;

    if(F != 0)
    {
        sol = (df(F,x), df(F,y), df(F,z));
        sol = sol.transpose();
    }
    return sol;
}

...

Symbolic curl(const Symbolic &F, const Symbolic &x, const Symbolic &y, const
Symbolic &z)
{
    Symbolic sol, i("i"), j("j"), k("k");

    Symbolic F1 = F.coeff(i,1);
    Symbolic F2 = F.coeff(j,1);
    Symbolic F3 = F.coeff(k,1);

    if(F1 != 0 || F2 != 0 || F3 != 0)
    {
        sol = (df(F3,y) - df(F2,z), -(df(F3,x) - df(F1,z)), df(F2,x)
            - df(F1,y));
        sol = sol.transpose();
    }
    else if(F1_hat != 0 || F2_hat != 0 || F3_hat != 0)
    {
        sol = (df(F3_hat,y) - df(F2_hat,z), -(df(F3_hat,x) - df(
            F1_hat,z)), df(F2_hat,x) - df(F1_hat,y));
        sol = sol.transpose();
    }
    return sol;
}
```

}

Code 29: *src/divgradcurl.cpp*

the source code can handle the input of a function along with the independent variables. For example, if we have a scalar field:

$$f(x, y, z) = -\frac{1}{2}xy + 2y + z^2$$

and we can compute the gradient with :

grad(f,x,y,z)

If we have a vector field:

$$F(x, y, z) = y\hat{i} - x\hat{j} + yz\hat{k}$$

we can compute the divergence and curl with these functions:

div(F,x,y,z)

curl(F,x,y,z)

In SymIntegration the source code to compute the div, grad, curl for the example above will be like this:

```
#include <iostream>
#include "symintegrationc++.h"
#include <bits/stdc++.h>
#include <cmath>

using namespace std;
using namespace SymbolicConstant;

int main(void)
{
    Symbolic x("x"), y("y"), z("z"), f, f2, i("i"), j("j"), k("k");

    f = -(0.5*x)*y+2*y + z*z;
    f2 = y*i -x*j +y*z*k;
    Symbolic f2_hat = y*i -x*j +y*z*k;
    cout << "F(x,y,z) = " << f << endl;
    cout << "F2 (vector field) = " << f2_hat << endl;

    cout << "\ndiv(F2) = " << div(f2_hat,x,y,z) <<endl;
    cout << "\ngrad(F) = " << grad(f,x,y,z) <<endl;
    cout << "\ncurl(F2) = " << curl(f2_hat,x,y,z) <<endl;

    return 0;
}
```

Code 30: *Examples/test SymIntegration DivGradCurl/main.cpp*

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DivGradCurl ]# m
ake
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DivGradCurl ]# ./main
F(x,y,z) = -0.5*x*y+2*y*z^(2)
F2 (vector field) = y*i-x*j+y*z*k

div(F2) = y

grad(F) =
[ -0.5*y ]
[ -0.5*x+2 ]
[ 2*z ]

curl(F2) =
[ z ]
[ 0 ]
[ -2 ]

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DivGradCurl ]#

```

Figure 2.5: The computation of *div grad curl*. (*SymIntegration/Examples/Calculus/Test SymIntegration Div-GradCurl/main.cpp*).

[SI*] The input, f , for the gradient is a scalar-valued function, while the output, ∇f , is a vector-valued function.

The input, F , for the divergence is a vector-valued function, while the output, $\nabla \cdot F$, is a scalar-valued function.

The input, F , for the curl is a vector-valued function, and the output, $\nabla \times F$, is a vector-valued function.

[SI*] If F denotes the velocity field for a fluid, then $\text{div } F$ at a point p measures the tendency of that fluid to diverge away from p ($\text{div } F > 0$) or accumulate toward p ($\text{div } F < 0$).

On the other hand, $\text{curl } F$ picks out the direction of the axis about which the fluid rotates (curls) most rapidly, and $||\text{curl } F||$ is a measure of the speed of this rotation. The direction of the rotation is according to the right-hand rule.

[SI*] **Laplacian**

The scalar function $\text{div}(\text{grad } f) = \nabla \cdot \nabla f$ (also written $\nabla^2 f$) is called the Laplacian, and a function f satisfying

$$\nabla^2 f = 0$$

is said to be harmonic, this concept is important in physics.

The Laplace operator is a scalar operator that can be applied to either vector or scalar fields.

The Laplacian of a scalar field $f(x, y, z)$ is the scalar-valued function:

$$\Delta f = \nabla^2 f = \nabla \cdot \nabla f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \quad (2.30)$$

The Laplacian of a vector field $F(x, y, z)$ is the vector field:

$$\Delta F = \nabla^2 F = \nabla \cdot \nabla F = \frac{\partial^2 F}{\partial x^2} + \frac{\partial^2 F}{\partial y^2} + \frac{\partial^2 F}{\partial z^2} \quad (2.31)$$

so if we have a vector field that can be written like this:

$$\mathbf{F}(x, y, z) = F_1(x, y, z)\hat{i} + F_2(x, y, z)\hat{j} + F_3(x, y, z)\hat{k}$$

then the Laplacian can also be written like this:

$$\begin{aligned}\nabla^2 \mathbf{F} &= \nabla^2 F_1 \hat{i} + \nabla^2 F_2 \hat{j} + \nabla^2 F_3 \hat{k} \\ \nabla^2 \mathbf{F} &= (\nabla^2 F_1, \nabla^2 F_2, \nabla^2 F_3) \\ \nabla^2 \mathbf{F} &= \left(\frac{\partial^2 F_1}{\partial x^2} + \frac{\partial^2 F_1}{\partial y^2} + \frac{\partial^2 F_1}{\partial z^2} \right) \hat{i} + \left(\frac{\partial^2 F_2}{\partial x^2} + \frac{\partial^2 F_2}{\partial y^2} + \frac{\partial^2 F_2}{\partial z^2} \right) \hat{j} \\ &\quad + \left(\frac{\partial^2 F_3}{\partial x^2} + \frac{\partial^2 F_3}{\partial y^2} + \frac{\partial^2 F_3}{\partial z^2} \right) \hat{k} \\ \nabla^2 \mathbf{F} &= \left(\frac{\partial^2 F_1}{\partial x^2} + \frac{\partial^2 F_1}{\partial y^2} + \frac{\partial^2 F_1}{\partial z^2}, \frac{\partial^2 F_2}{\partial x^2} + \frac{\partial^2 F_2}{\partial y^2} + \frac{\partial^2 F_2}{\partial z^2}, \frac{\partial^2 F_3}{\partial x^2} + \frac{\partial^2 F_3}{\partial y^2} + \frac{\partial^2 F_3}{\partial z^2} \right)\end{aligned}\tag{2.32}$$

The Laplacian of the velocity is related to the viscous forces in a fluid, it is also part of the heat equation, which describes how temperature changes over time, the Laplacian is also used in Poisson's equation, which relates electric potential to charge density.

In essence, the Laplacian of a vector field provides valuable information about the "smoothness" or "curvature" of the vector field, highlighting regions where the field changes rapidly.

The Laplacian is ubiquitous throughout modern mathematical physics, appearing for example in Laplace's equation, Poisson's equation, the heat equation, the wave equation, and the Schrödinger equation.

To compute the Laplacian from a defined function $f(x, y, z)$ with SymIntegration we can use :

laplacian(f,x,y,z) (it can handle the input of scalar field or vector field)

Theorem 2.12: Laplacian Identities

- (a) $\nabla^2(f + g) = \nabla^2 f + \nabla^2 g$
- (b) $\nabla^2(cf) = c\nabla^2 f$, for any constant c .
- (c) $\nabla^2(fg) = f\nabla^2 g + 2\nabla f \cdot \nabla g + g\nabla^2 f$

Theorem 2.13: Degree Two Identities

- (a) $\nabla \cdot (\nabla \times \mathbf{F}) = 0$ (divergence of a curl)
- (b) $\nabla \times (\nabla f) = 0$ (curl of gradient)
- (c) $\nabla \cdot (f\{\nabla g \times \nabla h\}) = \nabla f \cdot (\nabla g \times \nabla h)$
- (d) $\nabla \cdot (f\nabla g - g\nabla f) = f\nabla^2 g - g\nabla^2 f$
- (e) $\nabla \times (\nabla \times \mathbf{F}) = \nabla(\nabla \cdot \mathbf{F}) - \nabla^2 \mathbf{F}$ (curl of curl)

[SI*] Hessian Matrix

Hessian matrix is a matrix that maps a function of n variables into $n \times n$ square matrix. Consider a function

$$f(x_1, x_2, \dots, x_n)$$

the Hessian matrix of f is the matrix-valued function $H : \mathbb{R}^n \rightarrow \mathbb{R}^{n \times n}$ defined by

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1 \partial x_1} & \frac{\partial^2 f}{\partial x_2 \partial x_1} & \cdots & \frac{\partial^2 f}{\partial x_n \partial x_1} \\ \frac{\partial^2 f}{\partial x_1 \partial x_2} & \frac{\partial^2 f}{\partial x_2 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n \partial x_2} \\ \vdots & \vdots & \cdots & \vdots \\ \frac{\partial^2 f}{\partial x_1 \partial x_n} & \frac{\partial^2 f}{\partial x_2 \partial x_n} & \cdots & \frac{\partial^2 f}{\partial x_n \partial x_n} \end{bmatrix} \quad (2.33)$$

we can generate the hessian matrix from a defined function f with SymIntegration with :

hessianmatrix(f,x,y) (for 2 variables function)

hessianmatrix(f,x,y,z) (for 3 variables function)

Below is the code to show hessian matrix for function of 2 variables, $f(x, y)$, we provide the function to create the Hessian matrix for function that has up to 3 variables for now, and it is handled in **src/divgradcurl.cpp**.

```
#include <iostream>
#include <iomanip> // to declare the manipulator of setprecision()
#include <fstream>
#include <bits/stdc++.h> //for setw(6) at display() function
#include <vector> // For std::vector (example container)
#include "symintegrationc++.h"
#include <algorithm> // For std::sort

#include <chrono>
#include <string>
#include <bitset>

using namespace std::chrono;
using namespace std;

int main(int argc, char** argv)
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    Symbolic x("x"), y("y");

    Symbolic f = 2*sin(3*x) + 5 + x*x*y + cos(y);
    cout << "\nf(x,y) = " << f << endl;
    cout << "\nH = " << hessian(f,x,y) << endl;

    // Get ending timepoint
    auto stop = high_resolution_clock::now();
```

```

    auto duration = duration_cast<microseconds>(stop - start);

    cout << "Time taken by function: " << duration.count() << "
          microseconds" << endl;

    return 0;
}

```

Code 31: Examples/test Hessian Matrix/main.cpp

```

root [ ~/SourceCodes/CPP/C++ Create Library/TEST Hessian ]# ./main
f(x,y) = 2*sin(3*x)+x^(2)*y+cos(y)+5
H =
[ -18*sin(3*x)+2*y      2*x      ]
[      2*x             -cos(y)   ]

Time taken by function: 32521 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/TEST Hessian ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/TEST Hessian ]# ./main
f(x,y) = 2*sin(3*x)+x^(2)*y+cos(y)+5
H =
[ -18*sin(3*x)+2*y      2*x      ]
[      2*x             -cos(y)   ]

Time taken by function: 31283 microseconds

```

Figure 2.6: The computation of Hessian matrix. (SymIntegration/Examples/Linear Algebra/Test Hessian Matrix/main.cpp).

Chapter 3

Differential Equations, Partial Differential Equations, and Boundary Value Problems

"After Odin's defeat atop Yggdrasil and the revelation of Lezard's true nature, Freya quickly teleports there in a frantic state, telling Odin that Lezard's presence was the distortion she felt at Dipan" - Freya (Valkyrie Profile 2: Silmeria)

"Everyone's favorite fighting fairy godmother is still kicking, taking out Ether Strike hits for her godfather Odin. The heretofore unmatched fury of her scowl derives from her lack of lines in the main story" - Freya (Valkyrie Profile: Covenant of the Plume)

Differential equations are of interest to engineers and other nonmathematicians, such as physicists. It is because of the possibility of using them to investigate a wide variety of problems in the physical, biological, and social sciences.

For example consider this Newton's law formula:

$$F = ma \quad (3.1)$$

If you read the Physics book for undergraduate degree you will learn that it is not simply just $F = ma$, but it has a derivative as well

$$F = m \frac{dv}{dt} \quad (3.2)$$

Then add with drag force the equation will become

$$m \frac{dv}{dt} = mg - \gamma v \quad (3.3)$$

The Eq. (3.3) is what we call a differential equation that model an object falling in the atmosphere near sea level. From this equation we are able to determine the velocity at a given time, $v(t)$, from known g, m, γ . You can read more on this book [6], it is the first example of the book.

We will assume that you already read some books or learn from anywhere about differential equations, so we will only focus on the SymIntegration function and capability that can help you

to compute the solution of a differential equation.

Starting on June 23rd, 2025, we have created a new function **dsolve** in SymIntegration, it is still very fresh and in an infant stage and just able to solve a very simple first order linear ordinary differential equations.

I. SOLVING FIRST ORDER ORDINARY DIFFERENTIAL EQUATIONS

[SI*] Method of Integrating Factors

Solve the initial value problem

$$2y' + ty = 2 \quad (3.4)$$

with the initial value condition

$$y(0) = 1 \quad (3.5)$$

Solution:

We will convert the differential equation Eq. (3.4) into the standard form to become

$$y' + \left(\frac{t}{2}\right)y = 1 \quad (3.6)$$

Thus $p(t) = t/2$, and the integrating factor is

$$\mu(t) = e^{\int p(t) dt} = e^{\frac{t^2}{4}}$$

Then multiply Eq. (3.6) by $\mu(t)$ so that

$$e^{\frac{t^2}{4}}y' + \left(\frac{t}{2}\right)e^{\frac{t^2}{4}}y = e^{\frac{t^2}{4}} \quad (3.7)$$

Then integrate both sides of equation we will obtain

$$e^{\frac{t^2}{4}}y = \int e^{\frac{t^2}{4}} dt + c \quad (3.8)$$

thus

$$y = e^{-\frac{t^2}{4}} \int_0^t e^{\frac{s^2}{4}} ds + ce^{-\frac{t^2}{4}} \quad (3.9)$$

You can compute the value of the constant c by inputting the initial condition of $y(0) = 1$.

Now, in SymIntegration we will need the user to input the equation in term of $\frac{dy}{dx} = f(x, y)$, we will input the $f(x, y)$ and the coefficient of $\frac{dy}{dx}$ has to be 1.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]# ./main
f(x) = -0.5*x*y+1
DSolve for f(x,y).
y(x) = 1/2*pi^(1/2)*(erf(x*(-0.25)^(1/2)))*(-0.25)^(-1/2)*e^(-0.25*x^2))+C*e^(-0.25*x^2))
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]#
```

Figure 3.1: The newly built function **dsolve** in SymIntegration is able to compute the solution $y(x)$ for the ordinary differential equation of $2y' + ty = 2$ (SymIntegration/Examples/Differential Equations/Test SymIntegration DSolve/main.cpp).

[SI*] For the coding in SymIntegration, we add a new C++ file:

src/dsolve.cpp

This is the very first beginning of the dsolve by June 23rd, 2025:

```

#include "symintegral/symintegrationc++.h"

#ifdef SYMBOLIC_DEFINE
#ifndef SYMINTEGRATION_CPLUSPLUS_DSOLVE_DEFINE
#define SYMINTEGRATION_CPLUSPLUS_DSOLVE_DEFINE

Symbolic dsolve(const Symbolic &fx, const Symbolic &y, const Symbolic
&x)
{
    Symbolic dsol, mu, C("C");

    if(fx != 0)
    {
        list<Equations> eq;
        list<Equations>::iterator i;
        UniqueSymbol a, b, c, d;
        eq = (a*x*y + d).match(fx, (a,d));
        for(i=eq.begin(); i!=eq.end(); ++i)
        {
            try {
                Symbolic ap = rhs(*i, a), dp = rhs(*i, d);
                mu = exp(integrate(-ap*x,x));
                dsol = (integrate(mu*dp,x))/(mu) + (C)/(mu)
                ;
            } catch(const SymbolicError &se) {}
        }
    }
    return dsol;
}

#endif
#endif

```

Code 32: src/dsolve.cpp

and the include file as well **include/symintegral/dsolve.h**

```

#ifndef SYMINTEGRATION_CPLUSPLUS_DSOLVE

#ifdef SYMBOLIC_FORWARD
#ifndef SYMINTEGRATION_CPLUSPLUS_DSOLVE_FORWARD
#define SYMINTEGRATION_CPLUSPLUS_DSOLVE_FORWARD

#endif
#endif

#ifdef SYMBOLIC_DECLARE
#define SYMINTEGRATION_CPLUSPLUS_DSOLVE

```

```

#ifndef SYMINTEGRATION_CPLUSPLUS_DSOLVE_DECLARE
#define SYMINTEGRATION_CPLUSPLUS_DSOLVE_DECLARE

Symbolic dsolve(const Symbolic &, const Symbolic &, const Symbolic &);

#endif
#endif

#endif

```

Code 33: *include/symintegral/dsolve.h*

Then we also need to add a new function **erf**, as the integral of e^{x^2} is not an easy one to be computed and defined, it is like this

$$\int e^{ax^2} dx = \frac{\sqrt{\pi} \operatorname{erf}(x\sqrt{-a})}{2\sqrt{-a}}, \quad a \neq 0 \quad (3.10)$$

To handle that, the need to return the $\int e^{ax^2} dx$ correctly in SymIntegration we modify these files:

- *src/symintegrationc++.cpp*

```

...
...

Symbolic erf(const Symbolic &s)
{ return Erf(s); }

...
...

```

Code 34: *src/symintegrationc++.cpp*

- *src/functions.cpp*

```

...
...

////////////////////////////////////
// Implementation of Erf //
////////////////////////////////////

Erf::Erf(const Erf &s) : Symbol(s) {}

Erf::Erf(const Symbolic &s) : Symbol(Symbol("erf")[s]) {}

Simplified Erf::simplify() const
{
    const Symbolic &s = parameters.front().simplify();

```

```

        if(s == 0) return Number<int>(0);
        return *this;
    }

    Symbolic Erf::df(const Symbolic &s) const
    { return Derivative(*this,s); }

    Symbolic Erf::integrate(const Symbolic &s) const
    {
        const Symbolic &x = parameters.front();
        if(x == s) return Integral(*this,s);
        if(df(s) == 0) return Integral(*this,s);
        return Integral(*this,s);
    }

    ...
    ...

```

Code 35: *src/functions.cpp*

- `include/symintegral/functions.h`

```

...
...

class Erf;

...
...

class Erf: public Symbol
{
    public: Erf(const Erf&);
           Erf(const Symbolic&);

    Simplified simplify() const;
    Symbolic df(const Symbolic&) const;
    Symbolic integrate(const Symbolic&) const;

    Cloning *clone() const { return Cloning::clone(*this); }
};

...
...

```

Code 36: *include/symintegral/functions.h*

- `include/symintegral/symintegrationc++.h`

```

...
...

```

```
Symbolic erf(const Symbolic &);

...

...
```

Code 37: `include/symintegral/symintegrationc++.h`

[SI*] **Separable Equations**

We can integrate a differential equation function that is classified as separable equations if they have this form:

$$\frac{dy}{dx} = \frac{f(x)}{g(y)} \quad (3.11)$$

with $f(x)$ is a function of x only and $g(y)$ is a function of y only. Thus we can write the differential equation in the form of

$$g(y) dy - f(x) dx = 0 \quad (3.12)$$

Such an equation is said to be separable. A separable equation can be solved by integrating the functions f and g .

[SI*] **Example:**

Show that the equation

$$\frac{dy}{dx} = \frac{x^2}{1-y^2}$$

is separable, and then find an equation for its integral curves.

Solution:

$$\begin{aligned} \frac{dy}{dx} &= \frac{x^2}{1-y^2} \\ 1-y^2 dy &= x^2 dx \\ y - \frac{y^3}{3} &= \frac{x^3}{3} + c \\ -x^3 + 3y - y^3 &= c \end{aligned}$$

where c is an arbitrary constant and $-x^3 + 3y - y^3 = c$ is the equation for the integral curves in the form of implicit function.

[SI*] **Homogeneous Ratio Equations**

If the right side of the equation

$$\frac{dy}{dx} = f(x, y)$$

can be expressed as a function of the ratio y/x only, then the equation is said to be a homogeneous ratio equation. Such equations can always be transformed into separable equations by a change of the dependent variable.

[SI*] **Example:**

Find the solution of

$$\frac{dy}{dx} = \frac{y-4x}{x-y}$$

Solution:

First we will make it into the form of y/x

$$\frac{dy}{dx} = \frac{y-4x}{x-y} = \frac{(y/x)-4}{1-(y/x)}$$

Make the change of variables

$$v = \frac{y}{x}$$

$$\frac{dv}{dx} = \frac{1}{x} \frac{dy}{dx} - \frac{y}{x^2}$$

$$y = xv$$

$$\frac{dy}{dx} = x \frac{dv}{dx} + \frac{y}{x} = x \frac{dv}{dx} + v$$

As a result, the ODE becomes

$$\begin{aligned} \frac{dy}{dx} &= \frac{(y/x)-4}{1-(y/x)} \\ x \frac{dv}{dx} + v &= \frac{v-4}{1-v} \\ x \frac{dv}{dx} &= \frac{v-4}{1-v} - v \\ &= \frac{v-4}{1-v} - \frac{v(1-v)}{1-v} \\ &= \frac{v^2-4}{1-v} \\ \frac{1-v}{v^2-4} dv &= \frac{dx}{x} \\ \int \frac{1-v}{(v+2)(v-2)} dv &= \int \frac{dx}{x} \\ \int \left(-\frac{3}{4(v+2)} - \frac{1}{4(v-2)} \right) dv &= \int \frac{dx}{x} \\ -\frac{3}{4} \ln|v+2| - \frac{1}{4} \ln|v-2| &= \ln|x| + C \\ -\frac{3}{4} \ln\left|\frac{y}{x}+2\right| - \frac{1}{4} \ln\left|\frac{y}{x}-2\right| &= \ln|x| + C \end{aligned}$$

We can write the solution as an implicit function like this

$$f(x,y) = -\frac{3}{4} \ln\left|\frac{y}{x}+2\right| - \frac{1}{4} \ln\left|\frac{y}{x}-2\right| - \ln|x| - C$$

Remember that C is a constant that depends on the initial value.

The C++ code to solve the homogeneous ratio equations is using a function named **dsolveseparable** and we need to input it like this

dsolveseparable(g(x,y),f(x,y),y,x)

with

$$\frac{dy}{dx} = \frac{f(x,y)}{g(x,y)}$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]# ./main

DSolve for dy/dx = (y-4x) / (x-y)

f(x,y) = -0.25*ln(y*x^(-1)-2)-0.75*ln(y*x^(-1)+2)-ln(x)-C
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]# 

```

Figure 3.2: The newly built function *dsolveseparable* in *SymIntegration* is able to compute the solution $f(x,y)$ for homogeneous ratio equations (*SymIntegration/Examples/Differential Equations/Test SymIntegration DSolve/main.cpp*).

Now, to draw a direction field and some integral curves, recall that the right hand side of

$$\frac{dy}{dx} = \frac{y - 4x}{x - y}$$

depends only on the ratio $\frac{y}{x}$. This means that the integral curves have the same slope at all points on any given straight line through the origin, although the slope changes from one line to another. Therefore the direction field and the integral curves are symmetric with respect to the origin.

List of Solvable Linear ODEs with dsolve

We will list all the first order linear ordinary differential equations that can be solved with **dsolve** function in *SymIntegration* here, they are:

$$\begin{aligned}
 ay' + ty &= b \\
 aty' + by &= ct^2 \\
 ay' + by &= ce^{dt} \\
 ay' + by &= ct + d \\
 ay' + by &= c \\
 ay' + by &= 0
 \end{aligned}$$

with a, b, c, d are integers, e is reserved for exponential symbol, the solution will be $y(t)$. Remember when inputting to *SymIntegration*, you have to input the function $f(t,y)$ with this criteria

$$y' = f(t,y) \tag{3.13}$$

input the right hand side type of function in term of variables t and y only, and the coefficient of y' is 1.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]# make
g++ -c -o main.o main.cpp
g++ -o main -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]# ./main
f(t,y) = -0.5*t*y+1

DSolve for 2y' + ty = 2
y(t) = 1/2*pi^(1/2)*(erf(t*(-0.25)^(1/2)))*(-0.25)^(-1/2)*e^(-0.25*t^(2))+C*e^(-0.25*t^(2))

DSolve for ty' + 2y = 4t^2
y(t) = t^(2)+C*t^(-2)

DSolve for y' + 0.5y = 0.5*exp(t/3)
y(t) = 0.6*e^(0.333333*t)+C*e^(-0.5*t)

DSolve for y' - 2y = 4-t
y(t) = 1/2*t+C*e^(2*t)-7/4

DSolve for Q' = r/4 - rQ/100
Q(t) = C*e^(-1/100*t*r)+25

DSolve for S' = rS
S(t) = C*e^(t*r)
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve ]#

```

Figure 3.3: The function **dsolve** in **SymIntegration** able to compute the solution $y(t)$ for several examples of linear first order ordinary differential equations (**SymIntegration/Examples/Differential Equations/Test SymIntegration DSolve/main.cpp**).

List of Solvable Linear ODEs with **dsolveseparable**

We will list all the first order linear ordinary differential equations that can be solved with **dsolveseparable** function in **SymIntegration** here, they are:

$$\frac{dy}{dx} = \frac{f(x)}{g(y)}$$

$$\frac{dy}{dx} = \frac{f(x,y)}{g(x,y)}$$

We create another function besides **dsolve** to make it easier to separate certain type of differential equation, with this **dsolveseparable** we need to input the function in the numerator and denominator. It is able to compute the simple separable equations and also the homogeneous ratio equations (we use this term instead of only 'homogeneous equations' that is written from this book [6] to avoid clashing of name at the next section).

the origin:

31. $\frac{dy}{dx} = \frac{x^2 + xy + y^2}{x^2}$	32. $\frac{dy}{dx} = \frac{x^2 + 3y^2}{2xy}$
33. $\frac{dy}{dx} = \frac{4y - 3x}{2x - y}$	34. $\frac{dy}{dx} = -\frac{4x + 3y}{2x + y}$
35. $\frac{dy}{dx} = \frac{x + 3y}{x - y}$	36. $(x^2 + 3xy + y^2)dx - x^2 dy = 0$
37. $\frac{dy}{dx} = \frac{x^2 - 3y^2}{2xy}$	38. $\frac{dy}{dx} = \frac{3y^2 - x^2}{2xy}$

Figure 3.4: The problems from book [6] with regards to the homogeneous ratio equation that can be solved with **SymIntegration' dsolveseparable**.


```

]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolveSeparable
]# ./main

DSolveSeparable for dy/dx = (x^2) / (1-y^2)
f(x,y) = -1/3*y^(3)+y-1/3*x^(3)-C
DSolveSeparable for dy/dx = (y-4x) / (x-y)
f(x,y) = -0.25*ln(y*x^(-1)-2)-0.75*ln(y*x^(-1)+2)-ln(x)-C
DSolveSeparable for dy/dx = (x^2+xy+y^2) / (x^2)
f(x,y) = atan(y*x^(-1))-ln(x)-C
DSolveSeparable for dy/dx = (4y-3x) / (2x-y)
f(x,y) = 0.25*ln(y*x^(-1)-1)-1.25*ln(y*x^(-1)+3)-ln(x)-C
DSolveSeparable for dy/dx = (4x+3y) / (2x+y)
f(x,y) = -0.333333*ln(y*x^(-1)+1)-0.666667*ln(y*x^(-1)+4)-ln(x)-C
DSolveSeparable for dy/dx = (x+3y) / (x-y)
f(x,y) = -ln(y*x^(-1)+1)-2*(y*x^(-1)+1)^(-1)-ln(x)-C
DSolveSeparable for dy/dx = (x^2+3xy+y^2) / (x^2)
f(x,y) = -(y*x^(-1)+1)^(-1)-ln(x)-C
DSolveSeparable for dy/dx = (x^2 - 3y^2) / (2xy)
f(x,y) = -0.2*ln(-5*y^(2)*x^(-2)+1)-ln(x)-C
DSolveSeparable for dy/dx = (3y^2 - x^2) / (2xy)
f(x,y) = ln(y^(2)*x^(-2)-1)-ln(x)-C

```

Figure 3.5: The solutions with SymIntegration' *dsolveSeparable* of the problems from book [6] at the end of chapter 2.2.

List of Solvable Linear ODEs with *dsolveLogistic*

We will list all the first order linear ordinary differential equations that can be solved with *dsolveLogistic* function in SymIntegration here, they are:

$$\frac{dy}{dt} = r \left(1 - \frac{y}{K} \right) y$$

$$\frac{dy}{dt} = -r \left(1 - \frac{y}{T} \right) y$$

with r, K, T as the parameters. We need to input the function in $\frac{dy}{dt}$ term. The function needs 7 input

dsolveLogistic(y',y,y0, t, r, K, T)

We can define $y' = \frac{dy}{dt}$ as either $r \left(1 - \frac{y}{K} \right) y$ or $-r \left(1 - \frac{y}{T} \right) y$, it will return the general solution with initial value problem of $y_0 = y_0$.

The more difficult case occurs in the case for logistic growth with a threshold where the differential equation is

$$\frac{dy}{dt} = -r \left(1 - \frac{y}{K} \right) \left(1 - \frac{y}{T} \right) y$$

This is the case where we have to do integration of fraction polynomial of order 3. For a while, instead of solving the difficult differential equation we will break it into 2 cases, with **if.. else if..** conditional.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Logistic Equation ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Logistic Equation ]# ./main
For dy/dt = r*(1-(y/K))*y
y(t) = y0*K*(y0+K*e^(-r*t)-y0*e^(-r*t))^(-1)

For dy/dt = -r*(1-(y/T))*y
y(t) = y0*T*(y0+T*e^(-r*t)-y0*e^(-r*t))^(-1)

For dy/dt = -r*(1-(y/T))*(1-(y/K))*y
For T < y < K and y > K
y(t) = y0*K*(y0+K*e^(-r*t)-y0*e^(-r*t))^(-1)
For 0 < y < T,
y(t) =
y0*T*(y0+T*e^(-r*t)-y0*e^(-r*t))^(-1)

Time taken by function: 261099 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Logistic Equation ]#

```

Figure 3.6: The function *dsolve* in *SymIntegration* (*SymIntegration/Examples/Differential Equation-Test SymIntegration Logistic Equation/main.cpp*).

Modeling with First Order Equations

[SI*] Mathematical modeling and experiment or observation are both critically important and have somewhat complementary roles in scientific investigations. Mathematical models are validated by comparison of their predictions with experimental results.

[SI*] Three identifiable steps that are always present in the process of mathematical modeling:

1. Construction of the Model.
2. Analysis of the Model.
3. Comparison with Experiment or Observation.

[SI*] The Water Tank Mixing

At time $t = 0$ a tank contains Q_0 lb of salt dissolved in 100 gal of water. Assume that water containing $\frac{1}{4}$ lb of salt/gal is entering the tank at a rate of r gal/min and that the well-stirred mixture is draining from the tank at the same rate. Set up the initial value problem that describes this flow process. Find the amount of salt $Q(t)$ in the tank at any time, and also find the limiting amount Q_L that is present after a very long time. If $r = 3$ and $Q_0 = 2Q_L$, find the time T after which the salt level is within 2% of Q_L . Also find the flow rate that is required if the value of T is not to exceed 45 min.

Solution:

We assume that salt is neither created nor destroyed in the tank. Therefore variations in the amount of salt are due solely to the flows in and out of the tank. More precisely, the rate of change of salt in the tank, $\frac{dQ}{dt}$, is equal to the rate at which salt is flowing in minus the rate at which it is flowing out. In symbols,

$$\frac{dQ}{dt} = \text{rate in} - \text{rate out} \quad (3.14)$$

The rate at which salt enters the tank is the concentration $\frac{1}{4}$ lb/gal times the flow rate r gal/min, or $(r/4)$ lb/min. To find the rate at which salt leaves the tank, we need to multiply the concentration of salt in the tank by the rate of outflow, r gal/min. Since the rates of flow in and out are equal, the volume of water in the tank remains constant at 100 gal, and since

the mixture is "well-stirred," the concentration throughout the tank is the same, namely, $\frac{Q(t)}{100}$ lb/gal.

Therefore the rate at which salt leaves the tank is $\frac{rQ(t)}{100}$ lb/min. Thus the differential equation governing this process is

$$\frac{dQ}{dt} = \frac{r}{4} - \frac{rQ}{100} \quad (3.15)$$

The initial condition is

$$Q(0) = Q_0 \quad (3.16)$$

Upon thinking about the problem physically, we might anticipate that eventually the mixture originally in the tank will be essentially replaced by the mixture flowing in, whose concentration is $\frac{1}{4}$ lb/gal. Consequently, we might expect that ultimately the amount of salt in the tank would be very close to 25 lb. We can also find the limiting amount $Q_L = 25$ by setting $\frac{dQ}{dt}$ equal to zero in Eq. (3.15)

$$\frac{dQ}{dt} = \text{rate in} - \text{rate out} \quad (3.17)$$

The rate at which salt enters the tank is the concentration $\frac{1}{4}$ lb/gal times the flow rate r gal/min, or $(r/4)$ lb/min. To find the rate at which salt leaves the tank, we need to multiply the concentration of salt in the tank by the rate of outflow, r gal/min. Since the rates of flow in and out are equal, the volume of water in the tank remains constant at 100 gal, and since the mixture is "well-stirred," the concentration throughout the tank is the same, namely, $\frac{Q(t)}{100}$ lb/gal.

Therefore the rate at which salt leaves the tank is $\frac{rQ(t)}{100}$ lb/min. Thus the differential equation governing this process is

$$\frac{dQ}{dt} = \frac{r}{4} - \frac{rQ}{100} \quad (3.18)$$

The initial condition is

$$Q(0) = Q_0 \quad (3.19)$$

Upon thinking about the problem physically, we might anticipate that eventually the mixture originally in the tank will be essentially replaced by the mixture flowing in, whose concentration is $\frac{1}{4}$ lb/gal. Consequently, we might expect that ultimately the amount of salt in the tank would be very close to 25 lb. We can also find the limiting amount $Q_L = 25$ by setting $\frac{dQ}{dt}$ equal to zero in Eq. (3.15) and solving the resulting algebraic equation for Q .

To solve the initial value problem at Eqs (3.15), (3.16) analytically, note that Eq. (3.15) is both linear and separable. Rewriting it in the standard form for a linear equation, we have

$$\frac{dQ}{dt} + \frac{rQ}{100} = \frac{r}{4} \quad (3.20)$$

Thus the integrating factor is $e^{rt/100}$ and the general solution is

$$Q(t) = 25 + ce^{-\frac{rt}{100}} \quad (3.21)$$

where c is an arbitrary constant. To satisfy the initial condition (3.16), we must choose $c = Q_0 - 25$. Therefore the solution of the initial value problem at Eqs (3.15), (3.16) is

$$Q(t) = 25 + (Q_0 - 25)e^{-\frac{rt}{100}} \quad (3.22)$$

or

$$Q(t) = 25(1 - e^{-\frac{rt}{100}}) + Q_0 e^{-\frac{rt}{100}} \quad (3.23)$$

You can see that

$$Q(t) \rightarrow 25$$

as $t \rightarrow \infty$, so the limiting value Q_L is 25, confirming our physical intuition.

Further, $Q(t)$ approaches the limit more rapidly as r increases. In interpreting the solution (3.23), note that the second term on the right side is the portion of the original salt that remains at time t , while the first term gives the amount of salt in the tank due to the action of the flow processes.

Now suppose that $r = 3$ and $Q_0 = 2Q_L = 50$, then Eq. (3.22) becomes

$$Q(t) = 25 + (50 - 25)e^{-0.03t} \quad (3.24)$$

Since 2% of 25 is 0.5, we wish to find the time T at which $Q(t)$ has the value 25.5. Substituting $t = T$ and $Q(t) = 25.5$ in Eq. (3.24) and solving for T , we obtain

$$\begin{aligned} Q(t) &= 25 + (Q_0 - 25)e^{-\frac{rt}{100}} \\ 25.5 &= 25 + (50 - 25)e^{-\frac{3t}{100}} \\ e^{0.03t} &= \frac{25}{0.5} \\ T &= \frac{\ln |50|}{0.03} \\ &= 130.4 \end{aligned}$$

T is in minutes. To determine r so that $T = 45$, return to Eq. (3.22), set $t = 45$, $Q_0 = 50$, $Q(t) = 25.5$, and solve for r . The result is

$$\begin{aligned} Q(t) &= 25 + (Q_0 - 25)e^{-\frac{rt}{100}} \\ 25.5 &= 25 + (50 - 25)e^{-\frac{45r}{100}} \\ 0.5 &= 25e^{-\frac{45r}{100}} \\ e^{\frac{45r}{100}} &= \frac{25}{0.5} \\ \frac{45r}{100} &= \ln |50| \\ r &= \frac{100}{45} \ln |50| \\ &\approx 8.69 \end{aligned}$$

r is in gal/min.

We would like to look into the metrics and measurement a bit further, since this is really important. We have the units of gal/min which can be converted into

$$1 \text{ gal/min} = 0.00006309 \text{ m}^3/\text{s} = 0.0037854 \text{ m}^3/\text{min} \quad (3.25)$$

Now, we try to do some observations and experimentations with the help of computer simulation, namely Hamzstlab Physics. We will fill an empty tank with a water, that is made by water particle, sphere like, but since it is a 2D computer simulation, we make a circle instead.

And as a representation for the water tank, that is having a cylinder shape, we take the cross section of a cylinder that is like a rectangle.

We observe with a circle of radius $r = 0.2$ and a mass of 0.15 as the water particle that is coming into the empty tank, it takes $t = 45$ s to fill the water tank to be full.

The volume of the water tank itself based on the computer simulation is

$$\begin{aligned} V &= \pi r^2 h \\ &= \pi(5)^2(10) \\ &= 785.39819 \end{aligned}$$

The metrics for V is m^3 . So we will have the rate of the entering water as

$$\frac{785.39819 \text{ m}^3}{45 \text{ s}} = 17.453293 \text{ m}^3/\text{s} \quad (3.26)$$

Knowing that 1 gal/min = 0.00006309 m^3/s , we will have the rate of the incoming water as

$$rate_{r=0.2,m=0.15} = \frac{17.453293 \text{ m}^3/\text{s}}{0.00006309 \text{ m}^3/\text{s}} * (1 \text{ gal/min}) = 276,641.19512 \text{ gal/min} \quad (3.27)$$

We are doing a 6 simulations with Hamzstlab Physics and we gather this data:

sim	$r(m)$	N	$v_x(m/s)$	$t(s)$
1	0.2	23437.5	19.5	45
2	0.2	23437.5	25	44.7
3	0.3	6944.44	19.5	20
4	0.3	6944.44	25	20.3
5	0.4	2929.6875	19.5	11.8
6	0.4	2929.6875	25	11.5

Table 3.1: The variable t denotes time of completion in seconds till the tank is full, v_x is the velocity of the water particle in m/s toward the x axis, while N is the number of water particle that will fit in a tank of volume $V = 785.4 \text{ m}^3$, derived from $\frac{4}{3}\pi r^3$.

From table 3.1 we can then use multivariable regression so we can interpolate and approximate the time of completion, t , for any speed of the water that is filling the tank, for any radius of the water particle and the volume of the water tank. Thus, we will treat t as the independent variable, and the rest is the dependent variables.

Multivariable regression models aim to find the best-fitting equation that describes the relationship between the independent variables and the dependent variable. This equation has the form of

$$y = b_0 + b_1x_1 + b_2x_2 + \cdots + b_nx_n + \epsilon \quad (3.28)$$

where y is the dependent variable, b_0 is the intercept, $b_1, b_2, \dots, b_n$ are the coefficients for each independent variable $(x_1, x_2, \dots, x_n)$, and ϵ is the error term.

By analyzing the coefficients, we can understand how each factor relates to time of completion.

In this case we will use the independent variables of: N and v_x , we only need 2, since N is derived from the volume of the water tank divided by the volume of each particle so we don't need to use r anymore, it will become an extra cost and not necessary. Thus

$$x_1 = N$$

$$x_2 = v_x$$

Implementing multivariable regression in C++ typically involves handling matrix operations for calculations of coefficients. This can be achieved by implementing the necessary linear algebra functions from scratch or by utilizing existing C++ libraries designed for numerical linear algebra computation.

You can write your own C++ code from scratch to perform matrix multiplication, transposition, and inversion. But this provides a deep understanding of the underlying mathematics and can be complex, very time consuming and prone to errors.

Thankfully, in 2025 we have a high-performance C++ library that can handle float matrix and float vector operations with great precision, including transpose, inverse and multiplication, we will use **Armadillo** / **Arma** to help us computing the coefficients b_1 and b_2 .

Another alternative besides **Armadillo** is **Eigen**.

Now, suppose we have this matrix that we construct from the simulation data above (table 3.1):

$$X = \begin{bmatrix} 1 & 23437.5 & 19.5 \\ 1 & 23437.5 & 25 \\ 1 & 6944.44 & 19.5 \\ 1 & 6944.44 & 25 \\ 1 & 2929.6875 & 19.5 \\ 1 & 2929.6875 & 25 \end{bmatrix}, \quad y = \begin{bmatrix} 45 \\ 44.7 \\ 20 \\ 20.3 \\ 11.8 \\ 11.5 \end{bmatrix}$$

By any means, our measurements for matrix X and vector y / the simulation data could contain human errors as well.

So we will have this matrix equation for this multivariable regression problem:

$$y = X * B + E \quad (3.29)$$

y is the vector of dependent variable values, X is the matrix of independent variables values (including a column of ones for the intercept term), B is the vector of regression coefficients

to be estimated, and E is the vector of error terms.

We will use the most common method to estimate the coefficients B , which is the Ordinary Least Squares (OLS), which minimizes the sum of squared residuals. The formula for OLS coefficient estimation is:

$$B = (X^T * X)^{-1} * X^T * y \quad (3.30)$$

From **Armadillo** we will obtain

$$B = \begin{bmatrix} 8.3696 \\ 0.0016 \\ -0.0182 \end{bmatrix}$$

Thus

$$y = b_0 + b_1x_1 + b_2x_2 = 8.3696 + 0.0016N - 0.0182v_x$$

the equation above tells us about the relation / proportion between y and the independent variables

$$y \propto N$$

$$y \propto \frac{1}{v_x}$$

```
root [ ~/latex/DianFreya Math Physics Simulator/Source Codes/C++/C++ Gnuplot SymbolicC++/ch23-Numerical Linear Algebra/Multivariab
le Regression with Armadillo ]# ./main
Matrix X:
  1.0000e+00  2.3438e+04  1.9500e+01
  1.0000e+00  2.3438e+04  2.5000e+01
  1.0000e+00  6.9444e+03  1.9500e+01
  1.0000e+00  6.9444e+03  2.5000e+01
  1.0000e+00  2.9297e+03  1.9500e+01
  1.0000e+00  2.9297e+03  2.5000e+01
Vector y:
  45.0000
  44.7000
  20.0000
  20.3000
  11.8000
  11.5000
Matrix X^T:
  1.0000e+00  1.0000e+00  1.0000e+00  1.0000e+00  1.0000e+00  1.0000e+00
  2.3438e+04  2.3438e+04  6.9444e+03  6.9444e+03  2.9297e+03  2.9297e+03
  1.9500e+01  2.5000e+01  1.9500e+01  2.5000e+01  1.9500e+01  2.5000e+01
Matrix B:
  8.3696
  0.0016
 -0.0182
Xnew:
  1.0000e+00  6.9444e+03  2.5000e+01
Xnew * B:
  18.9128
Time taken by function: 2155 microseconds
```

Figure 3.7: The matrix B is computed with **Armadillo** and the predicted t for $N = 6944.44$ and $v_x = 25$ is 18.9128, while the t obtained from the simulation observation is 20.3, the error is 6.83%, it took 2155 microseconds to complete the computation with **Armadillo** (**DFSimulatorC/Source Codes/C++/C++ Gnuplot SymbolicC++/ch23-Numerical Linear Algebra/Multivariable Regression with Armadillo/main.cpp**).

For comparison,

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Linear Algebra/Test SymIntegration Multivariable Regression to Fill a Water Tank ]# ./main

Matrix X:
      1.000000      23437.500000      19.500000
      1.000000      23437.500000      25.000000
      1.000000      6944.440000      19.500000
      1.000000      6944.440000      25.000000
      1.000000      2929.687500      19.500000
      1.000000      2929.687500      25.000000

Matrix X^{T}:
      1.000000      1.000000      1.000000      1.000000      1.000000      1.000000
    23437.500000    23437.500000    6944.440000    6944.440000    2929.687500    2929.687500
      19.500000      25.000000      19.500000      25.000000      19.500000      25.000000

Vector y:
    45.000000
    44.700000
    20.000000
    20.300000
    11.800000
    11.500000

B :
    8.369652
    0.001584
   -0.018182

Xnew :
    1.000000
    6944.440000
    25.000000

Xnew * B = 18.912819

Time taken by function: 1751 microseconds

```

Figure 3.8: The matrix B is computed with *SymIntegration* and the predicted t for $N = 6944.44$ and $v_x = 25$ is 18.9128, giving the same result with *Armadillo*, the result with *SymIntegration* took around 1751 microseconds. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Multivariable Regression to Fill a Water Tank/main.cpp*).

With this multivariable regression we are able to predict the time of completion to fill a water tank with any given volume of the water tank and any given speed of the particle for the filling, or perhaps in other case, to clear up an organ from dangerous substance; how many dosage of drugs should one person drinks with certain mass and age as the independent variables to be healed from certain disease? How many chlorine needed to clear up a pool with certain volume and water quality?




Figure 3.9: The simulation for a water tank filling with a water particle that has the radius of $r = 0.4$ and $v_x = 25$ the t obtained from the simulation observation is 11.5 s (*Hamzstlab-Physics2D/tests/ode_watertankfilling.cpp*).

We would like to introduce a book we had made in 2023-2025 that has covered Elementary Linear Algebra for undergraduate level knowledge along with its' C++ code that can be tested by the reader here [10]. We are summarizing the theorems, formulas, definitions, solving the problems from chapter 1 to the last chapter of Chris-Rorres book [26], the addition is that we make the C++ codes ourselves with the help of using already established C++ libraries like **Armadillo**, **giNaC**, **Eigen**, **SymbolicC++**. It is very fortunate that we have learned Linear Algebra first, since it is proven now that it becomes very helpful when we learn other topics such as physics, mechanical engineering, differential equations, and so on. All the C++ codes that we have made and put in that book is very helpful and can be reused again.

The book can be downloaded and all the source codes are available at:
<https://github.com/glanzkaizer/DFSimulatorC>

Since this example of water tank mixing is hypothetical, the validity of the model is not in question. If the flow rates are as stated, and if the concentration of salt in the tank is uniform, then the differential equation (3.17) is an accurate description of the flow process.

Model of this kind are often used in problems involving a pollutant in a lake, or a drug in an organ of the body, for example, rather than a tank of salt water. In such cases the flow

rates may not be easy to determine or may vary with time.




Figure 3.10: The 2D simulation of a salt water mixing in a tank made with Hamzstlab Physics 2D that is using Box2D version 2.4.2 as the backend library ([Hamzstlab-Physics2D/tests/ode_watertankmixing.cpp](https://github.com/Hamzstlab-Physics2D/tests/ode_watertankmixing.cpp)).

We would like to introduce again one of our creation: **Hamzstlab Physics 2D**, a 2D classical physics simulator, that you can download and try / can be accessed at <https://github.com/glanzkaizer/Hamzstlab-Physics2D>

We make a lot of classical mechanics Physics simulations in 2D with Hamzstlab Physics 2D that is using the backend of Box2D version 2.4.2, at this opportunity we try to make a 2D simulation for this example so people / reader can see the illustration of this water tank mixing with salt water that is coming with certain rate / speed into the tank.

If SymIntegration is able to compute the differential equation solution, and combining it with Hamzstlab Physics we can make a better simulation that is near to real-life physics.

[SI*] **Home Loan / Mortgage**

A home buyer can afford to spend no more than USD 500 / month on mortgage payments. Suppose that the interest rate is 6%, that interest is compounded continuously, and that payments are also made continuously.

- Determine the maximum amount that this buyer can afford to borrow on a 20-year mortgage.
- Determine the total interest paid during the term of the mortgage.

Solution:

- (a) The amount of money $S(t)$ that the buyer has to pay changes in time due to two factors, the compound interest and its' continuous payments. The rate of growth for compounding is rS , and the rate of decay due to the continuous payments is k .

$$\frac{dS}{dt} = rS - k$$

$$\frac{dS}{dt} - rS = -k$$

This is a linear first-order inhomogeneous ODE, so it can be solved by multiplying both sides by an integrating factor $\mu(t)$

$$\mu(t) = e^{\int^t (-r) ds} = e^{-rt}$$

proceed with the multiplication and solve it for $S(t)$

$$e^{-rt} \frac{dS}{dt} - re^{-rt} S = -ke^{-rt}$$

$$\frac{d}{dt}(e^{-rt} S) = -ke^{-rt}$$

$$e^{-rt} S = \frac{k}{r} e^{-rt} + C$$

$$S(t) = \frac{k}{r} + Ce^{rt}$$

Apply the initial condition $S(0) = S_0$ to determine C

$$S(0) = \frac{k}{r} + Ce^{rt}$$

$$S_0 = \frac{k}{r} + Ce^{rt}$$

$$C = S_0 - \frac{k}{r}$$

Therefore, the amount of money the buyer has to pay after t years is

$$\begin{aligned} S(t) &= \frac{k}{r} + \left(S_0 - \frac{k}{r}\right) e^{rt} \\ &= \frac{k}{r}(1 - e^{rt}) + S_0 e^{rt} \end{aligned}$$

For a year the home buyer will need to pay $k = 500 \times 12 = \text{USD } 6,000$ year and $r = 6\% = 0.06$.

$$S(t) = \frac{6000}{0.06}(1 - e^{0.06t}) + S_0 e^{0.06t}$$

For a 20-year mortgage, $S(t) = 0$ at $t = 20$.

$$0 = \frac{6000}{0.06}(1 - e^{0.06t}) + S_0 e^{0.06t}$$

$$S_0 = 100000 - 100000e^{-1.2}$$

$$S_0 \approx \text{USD } 69,880.58$$

The value of S_0 is how much the buyer can afford to borrow initially.

(b) For the 20-year mortgage, the home buyer pays a total of

$$USD\ 6,000 \times 20 = USD\ 120,000$$

so the total interest paid is

$$USD\ 120,000 - USD\ 69,880.58 = USD\ 50,119.42$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve and IVP f
or Mortgage case ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve and IVP f
or Mortgage case ]# ./main

DSolve for S' = rS - k

S(t) = k*r^(-1)+C*e^(t*r)
S(t=0) = k*r^(-1)+C
For ivp S(0) = S0,
C = -k*r^(-1)+S0

Test ivp = k*r^(-1)-k*r^(-1)*e^(t*r)+S0*e^(t*r)
For ivp k = 6000, r = 6%, t = 20, S(t) = 0,
S(0) = -100000*e^(-1.2)+100000

```

Figure 3.11: The function *dsolve* and *ivp* in *SymIntegration* is able to compute the solution $S(t)$ for the ordinary differential equation of $\frac{dS}{dt} = rS - k$ with the initial value problem $S(0) = S_0$ (*SymIntegration/Examples/Differential Equations/Test SymIntegration DSolve and IVP for Mortgage case/main.cpp*).

As of July 20th, 2025 we just add a new function *ivp*, and it is managed in */usr/include/symintegral/dsolve.h* and */src/dsolve.cpp*

```

...
...

Symbolic ivp(const Symbolic &fx, const Symbolic &x, const Symbolic &c)
{
    Symbolic ivpsol, f0, C("C"), S0("S0");

    if(fx != 0)
    {
        list<Equations> eq;
        list<Equations>::iterator i;
        UniqueSymbol a, b;
        // Will work for this model: dS/dt = rS-k
        f0 = fx[x==0];
        C = solve(f0-S0,C).front().rhs ;
        ivpsol = fx[c==C];

    }
    return ivpsol;
}

...

```

Code 38: *src/dsolve.cpp*

[SI*] Radiocarbon Dating Example:

An important tool in archeological research is radiocarbon dating, developed by the American chemist Willard F. Libby. This is a means of determining the age of certain wood and plant remains, hence of animal or human bones or artifacts found buried at the same levels.

Radiocarbon dating is based on the fact that some wood or plant remains contain residual amounts of carbon-14, a radioactive isotope of carbon. This isotope is accumulated during the lifetime of the plant and begins to decay at its death. Since the half-life of carbon-14 is long (approximately 5730 years), measurable amounts of carbon-14 is still present, then by appropriate laboratory measurements the proportion of the original amount of carbon-14 that remains can be accurately determined. In other words, if $Q(t)$ is the amount of carbon-14 at time t and Q_0 is the original amount, then the ratio $\frac{Q(t)}{Q_0}$ can be determined, as long as this quantity is not too small. Present measurement techniques permit the use of this method for time periods of 50,000 years or more.

- Assuming that Q satisfies the differential equation $Q' = -rQ$, determine the decay constant r for carbon-14.
- Find an expression for $Q(t)$ at any time t , if $Q(0) = Q_0$
- Suppose that certain remains are discovered in which the current residual amounts of carbon-14 is 20% of the original amount. Determine the age of the remains.

Solution:

- The rate that the mass of carbon-14 decreases is assumed to be proportional to how much is present at any given time

$$\frac{dQ}{dt} \propto -Q$$

Change this proportionality to an equation by introducing a constant r on the right side.

$$Q' = -rQ$$

Divide both sides by Q .

$$\frac{Q'}{Q} = -r$$

The left side can be written as the derivative of a logarithm by the chain rule.

$$\begin{aligned}\frac{d}{dt} \ln Q &= -r \\ \ln Q &= -rt + C \\ Q(t) &= e^{-rt+C} \\ &= ce^{-rt}\end{aligned}$$

- Suppose that there is a certain amount of mass Q_0 initially. Then the initial condition is $Q(0) = Q_0$. We will use it to determine c .

$$\begin{aligned}Q(0) &= ce^{-r(0)} \\ c &= Q_0\end{aligned}$$

Consequently, the mass decays exponentially. Substituting back we will have the solution to the initial value problem as

$$Q(t) = Q_0 e^{-rt}$$

- (c) From the formula that we obtain at (b) we know that r can be determined with knowledge of the half-life. For carbon-14, specifically, we know that half the initial mass is lost after 5730 years.

$$\frac{Q_0}{2} = Q_0 e^{-r(5730)}$$

Solve this equation for r

$$\begin{aligned} e^{-5730r} &= \frac{1}{2} \\ \ln |e^{-5730r}| &= \ln \left| \frac{1}{2} \right| \\ r &= -\frac{\ln \left| \frac{1}{2} \right|}{5730} \\ &= \frac{\ln |2|}{5730} \\ r &\approx 0.0001210 \end{aligned}$$

r is the rate of the decreasing mass per year. Therefore, the mass of a sample of carbon-14 is

$$Q(t) = Q_0 e^{-\frac{\ln|2|}{5730}t}$$

where t is in years. If some remains have 20% of the original amount of carbon-14, then $Q(t) = 0.2Q_0$. Solve the resulting equation for t to determine how old the remains are.

$$\begin{aligned} 0.2Q_0 &= Q_0 e^{-\frac{\ln|2|}{5730}t} \\ e^{-\frac{\ln|2|}{5730}t} &= 0.2 \\ \ln |e^{-\frac{\ln|2|}{5730}t}| &= \ln |0.2| \\ -\frac{\ln |2|}{5730}t &= \ln \frac{1}{5} \\ -\frac{\ln |2|}{5730}t &= -\ln |5| \\ t &= \frac{\ln |5|}{\ln |2|} 5730 \\ t &\approx 13,305 \end{aligned}$$

so the age of the remains are around 13,305 years old.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve and IVP for Radiocarbon Dating ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve and IVP for Radiocarbon Dating ]# ./ma
in

DSolve for Q' = -rQ
Q(t) = C*e^(-t*r)
Q(t=0) = C
For ivp Q(0) = Q0,
C = Q0

Solution for the ivp,
Q(t) = Q0*e^(-t*r)

Half-life problem for carbon-14, the ivp becomes:
Q0*e^(-5730*r)-0.5*Q0 = 0

Determining the rate by solving the ivp solution
r = 0.000120968

Determining the age of the remains with 20% of carbon-14
t = 13304.6

```

Figure 3.12: The computation of the rate and the age of remains with *SymIntegration*' *dsolve* and *ivp* functions. (*SymIntegration/Examples/Differential Equations/Test SymIntegration DSolve and IVP for Radiocarbon Dating/main.cpp*).

On this example, we have added another **else if** conditional so *SymIntegration*' **solve** function will be able to solve this equation:

$$ce^{-rx} = a$$

$$ce^{-rx} = a$$

$$x = -\frac{\ln \frac{a}{c}}{r} = \frac{\ln \frac{c}{a}}{r}$$

```

...
double division(Symbolic x, Symbolic y)
{
    return x/y;
}

Equations solve(const Symbolic &e, const Symbolic &x)
{
    Equations soln;

    ...
    ...

    else if(e.coeff(1,0) == e) // for case c * exp(-tx) = a or ln|tx| = a
    {
        if(df(e,x).coeff(x,0) != 0 && df(df(e,x),x).coeff(x,0) != df(e,x).
            coeff(x,0) && -df(df(e,x),x).coeff(x,0) != df(e,x).coeff(x,0)) //
            for case c * exp(-tx) = a ,
        {
            Symbolic t = df(df(e,x),x).coeff(x,0) / df(e,x).coeff(x,0); //
                will work for c == 1 or c != 1
            Symbolic a = -e.coeff(SymbolicConstant::e,0) ;
            Symbolic c = df(e,x).coeff(x,0)/t;
            soln = (soln, x == ln(division(a,c))/t);
        }
    }
}

```

```

    }
    else if(df(e,x).coeff(x,0) != 0 && -df(df(e,x),x).coeff(x,0) == df(e,
        x).coeff(x,0) ) // for case c * exp(-tx) = a , c !=1, t ==1
    {
        Symbolic a = - e.coeff(SymbolicConstant::e,0) ;
        Symbolic c = - df(e,x).coeff(x,0) ;
        soln = (soln, x == ln(division(c,a)));
    }
    ...
}
return soln;
}
...

```

Code 39: *src/solve.cpp***[SI*] Pollutants in a Lake:**

Consider a lake of constant volume V containing at time t an amount $Q(t)$ of pollutant, evenly distributed throughout the lake with a concentration $c(t)$, where $c(t) = \frac{Q(t)}{V}$. Assume that water containing a concentration k of pollutant enters the lake at a rate r , and that water leaves the lake at the same rate. Suppose that pollutants are also added directly to the lake at a constant rate P .

Note that given assumptions neglect a number of factors that may, in some cases, be important—for example, the water added or lost by precipitation, absorption, and evaporation; the stratifying effect of temperature differences in a deep lake; the tendency of irregularities in the coastline to produce sheltered bays; and the fact that pollutants are not deposited evenly throughout the lake but (usually) at isolated points around its periphery. The results below must be interpreted in the light of the neglect of such factors as these.

- If at time $t = 0$ the concentration of pollutant is c_0 , find an expression for the concentration $c(t)$ at any time. What is the limiting concentration as $t \rightarrow \infty$?
- If the addition of pollutants to the lake is terminated ($k = 0$ and $P = 0$ for $t > 0$), determine the time interval T that must elapse before the concentration of pollutants is reduced to 50% of its original value; to 10% of its original value.
- Table 5.1 contains data for several of the Great Lakes. Using these data, determine from part (b) the time T necessary to reduce the contamination of each of these lakes to 10% of the original value.

Lake	$V(\text{km}^3 \times 10^3)$	$r(\text{km}^3/\text{year})$
<i>Superior</i>	12.2	65.2
<i>Michigan</i>	4.9	158
<i>Erie</i>	0.46	175
<i>Ontario</i>	1.6	209

Table 3.2: *Volume and Flow Data for the Great Lakes*

Solution:

(a) The conservation law that governs the mass of pollutant in the lake is as follows.

rate of mass accumulation = rate of mass in - rate of mass out

the rate of mass in is the sum of rk and P , and the rate of mass out is $rc(t)$.

$$\begin{aligned}\frac{dQ}{dt} &= rk + P - rc(t) \\ &= rk + P - \frac{r}{V}Q(t) \\ \frac{dQ}{dt} + \frac{r}{V}Q &= rk + P\end{aligned}$$

This is a first-order linear inhomogeneous ODE, so it can be solved by multiplying both sides by an integrating factor $\mu(t)$.

$$\begin{aligned}\mu(t) &= e^{\int \frac{r}{V} ds} = e^{\frac{rt}{V}} \\ e^{\frac{rt}{V}} \frac{dQ}{dt} + \frac{r}{V} e^{\frac{rt}{V}} Q &= rke^{\frac{rt}{V}} + Pe^{\frac{rt}{V}} \\ \frac{d}{dt}(e^{\frac{rt}{V}} Q) &= rke^{\frac{rt}{V}} + Pe^{\frac{rt}{V}} \\ e^{\frac{rt}{V}} Q &= \int^t (rke^{\frac{rs}{V}} + Pe^{\frac{rs}{V}}) ds + C \\ &= rk \left(\frac{V}{r} \right) e^{\frac{rt}{V}} + P \left(\frac{V}{r} \right) e^{\frac{rt}{V}} + C \\ e^{\frac{rt}{V}} Q &= kVe^{\frac{rt}{V}} + \frac{PV}{r} e^{\frac{rt}{V}} + C \\ Q(t) &= kV + \frac{PV}{r} e^{\frac{rt}{V}} + Ce^{-\frac{rt}{V}}\end{aligned}$$

Divide both sides by V to obtain the concentration.

$$c(t) = \frac{Q(t)}{V} = k + \frac{P}{r} + \frac{C}{V} e^{-\frac{rt}{V}}$$

Apply the initial condition $c(0) = c_0$ to determine C .

$$\begin{aligned}c(0) &= k + \frac{P}{r} + \frac{C}{V} \\ c_0 &= k + \frac{P}{r} + \frac{C}{V} \\ \frac{C}{V} &= c_0 - k - \frac{P}{r}\end{aligned}$$

Therefore, the concentration is

$$c(t) = \frac{Q(t)}{V} = k + \frac{P}{r} + \left(c_0 - k - \frac{P}{r} \right) e^{-\frac{rt}{V}}$$

Because of the decaying exponential function, the limit of $c(t)$ as $t \rightarrow \infty$ is

$$\lim_{t \rightarrow \infty} c(t) = k + \frac{P}{r}$$

(b) If the pollution stops, then the rate of mass in becomes zero in the conservation law.

rate of mass accumulation = - rate of mass out

The rate of mass out is still $rc(t)$.

$$\begin{aligned}\frac{dQ}{dt} &= -rc(t) \\ &= -\frac{r}{V}Q(t) \\ \frac{\frac{dQ}{dt}}{Q} &= -\frac{r}{V} \\ \frac{dQ}{Q} &= -\frac{r}{V} dt \\ \ln Q &= -\frac{rt}{V} + C_1 \\ Q &= e^{-\frac{rt}{V} + C_1} \\ Q(t) &= C_2 e^{-\frac{rt}{V}}\end{aligned}$$

Since Q is the mass of pollutant, divide both sides by V to get the concentration.

$$c(t) = \frac{Q(t)}{V} = \frac{A_1}{V} e^{-\frac{rt}{V}}$$

Use the initial condition $c(0) = c_0$ to determine C_2

$$c(0) = \frac{C_2}{V} = c_0$$

Therefore

$$c(t) = c_0 e^{-\frac{rt}{V}}$$

Set $c(t) = 0.5c_0$ and solve for $t = T$ to find how long it will take for the concentration to reach half its initial value.

$$\begin{aligned}0.5c_0 &= c_0 e^{-\frac{rT}{V}} \\ e^{-\frac{rT}{V}} &= 0.5 \\ \ln e^{-\frac{rT}{V}} &= \ln 0.5 \\ -\frac{rT}{V} &= \ln \frac{1}{2} \\ T &= \frac{V}{r} \ln 2\end{aligned}$$

On the other hand, set $c(t) = 0.1c_0$ and solve for $t = T$ to find how long it will take for

the concentration to reach one-tenth its initial value.

$$\begin{aligned}0.1c_0 &= c_0 e^{-\frac{rt}{V}} \\e^{-\frac{rt}{V}} &= 0.1 \\\ln e^{-\frac{rt}{V}} &= \ln 0.1 \\-\frac{rT}{V} &= \ln \frac{1}{10} \\T &= \frac{V}{r} \ln 10\end{aligned}$$

- (c) Plug in the numbers for V and r into the formula at part (b).
Lake Superior:

$$T = \frac{V}{r} \ln 10 = \frac{12.2 \times 10^3}{65.2} \ln 10 \approx 431 \text{ years}$$

Lake Michigan:

$$T = \frac{V}{r} \ln 10 = \frac{4.9 \times 10^3}{158} \ln 10 \approx 71.4 \text{ years}$$

Lake Erie:

$$T = \frac{V}{r} \ln 10 = \frac{0.46 \times 10^3}{175} \ln 10 \approx 6.05 \text{ years}$$

Lake Ontario:

$$T = \frac{V}{r} \ln 10 = \frac{1.6 \times 10^3}{209} \ln 10 \approx 17.6 \text{ years}$$

```

root { ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve and IVP for Pollutant in Great Lakes }# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root { ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DSolve and IVP for Pollutant in Great Lakes }# ./main

DSolve for Q' = rk + P - rc(t) , c(t) = Q(t)/V

Q(t) = k*V+P*r^(-1)+C*e^(-V^(-1)*t*r)
c(t) = k+P*r^(-1)+C*e^(-V^(-1)*t*r)*V^(-1)
c(0) = k+P*r^(-1)+C*V^(-1)
c0 - c(0) = c0-k-P*r^(-1)-C*V^(-1)
For ivp c(0) = c0,
C/V = c0-k-P*r^(-1)

Therefore, the concentration is,
c(t) = k+P*r^(-1)+c0*e^(-V^(-1)*t*r)-k*e^(-V^(-1)*t*r)-P*r^(-1)*e^(-V^(-1)*t*r)

The limit of c(t) as t -> ∞
lim_{t -> ∞} c(t) = k+P*r^(-1)+c0*e^(-inf*V^(-1)*r)-k*e^(-inf*V^(-1)*r)-P*r^(-1)*e^(-inf*V^(-1)*r)

If the pollution stops, then
Q(t) = C*e^(-V^(-1)*t*r)

The concentration when the pollution stops is
c(t) = C*e^(-V^(-1)*t*r)*V^(-1)

For ivp c(0) = c0,
c0-C*V^(-1) = 0

Therefore the ivp solution will be,
c(t) = c0*e^(-V^(-1)*t*r)

Set c(t) = 0.5 c0 and solve for t=T,
T = 0.693147*V*r^(-1)

Set c(t) = 0.1 c0 and solve for t=T,
T = 2.30259*V*r^(-1)

Time to reduce the contamination in each lakes to 10 percent of the original value

Lake Superior: T = 430.852
Lake Michigan: T = 71.4093
Lake Erie: T = 6.05251
Lake Ontario: T = 17.6274

```

Figure 3.13: The computation of initial value problem solution for the pollutant concentration in a lake, $c(t)$, and the time to clean up a lake to make the pollutant concentration less than 10% with SymIntegration' *dsolve* function. (SymIntegration/Examples/Differential Equations/Test SymIntegration DSolve and IVP for Pollutant in Great Lakes/main.cpp).

[SI*] Psychology' Problem:

One problem in psychology is to determine the relation between some physical stimulus and the corresponding sensation or reaction produced in a subject. Suppose that, measured in appropriate units, the strength of a stimulus is s and the intensity of the corresponding sensation is some function of s , say $y(s)$. Some experimental data suggest that the rate of change of intensity of the sensation with respect to the stimulus is directly proportional to the intensity of the sensation and inversely proportional to the strength of the stimulus; that is, $y(s)$ satisfies the differential equation

$$\frac{dy}{ds} = k \frac{y}{s}$$

for some positive constant k . Solve this differential equation.

Solution:

$$\frac{dy}{ds} = k \frac{y}{s}$$

$$\frac{dy}{y} = k \frac{ds}{s}$$

$$\ln |y| = k \ln |s|$$

$$y(s) = s^k + C$$

II. SECOND ORDER LINEAR EQUATIONS

[SI*] Linear equations of second order are of crucial importance in the study of differential equations for two main reasons. The first is that linear equations have a rich theoretical structure that underlies a number of systematic methods of solution. Further, a substantial portion of this structure and of these methods is understandable at a fairly elementary mathematical level. Another reason to study second order linear equations is that they are vital to any serious investigation of the classical areas of mathematical physics. One cannot go very far in the development of fluid mechanics, heat conduction, wave motion, or electromagnetic phenomena without finding it necessary to solve second order linear differential equations.

[SI*] A second order ordinary differential equation has the form

$$\frac{d^2y}{dt^2} = f\left(t, y, \frac{dy}{dt}\right) \quad (3.31)$$

where f is some given function. We usually denote the independent variable by t since time is often the independent variable in physical problems, but sometimes we will use x instead. We will use y , or occasionally some other letter, to designate the dependent variable. Equation (3.31) is said to be linear if the function f has the form

$$f\left(t, y, \frac{dy}{dt}\right) = g(t) - p(t)\frac{dy}{dt} - q(t)y \quad (3.32)$$

that is, if f is linear in y and $\frac{dy}{dt}$. In Eq. (3.32) g , p , and q are specified functions of the independent variable t but do not depend on y . In this case we usually rewrite Eq. (3.31) as

$$y'' + p(t)y' + q(t)y = g(t) \quad (3.33)$$

where the primes denote differentiation with respect to t . Instead of Eq. (3.33), we often see the equation

$$P(t)y'' + Q(t)y' + R(t)y = G(t) \quad (3.34)$$

Of course, if $P(t) \neq 0$, we can divide Eq. (3.33) by $P(t)$ and thereby obtain Eq. (3.34) with

$$p(t) = \frac{Q(t)}{P(t)}, \quad q(t) = \frac{R(t)}{P(t)}, \quad g(t) = \frac{G(t)}{P(t)} \quad (3.35)$$

In discussing Eq. (3.33) and in trying to solve it, we will restrict ourselves to intervals in which p , q , and g are continuous functions.

If Eq. (3.31) is not of the form (3.33) or (3.34), then it is called nonlinear. Analytical investigations of nonlinear equations are relatively difficult. Numerical or geometrical approaches are often more appropriate.

We should not be misled by the fact that first order linear equations are easily solved by means of formulas. In general, Eq. (3.33) cannot be solved explicitly in terms of known elementary functions, or even in terms of indicated integrations. To find solutions, it is commonly necessary to resort to infinite processes of one kind or another, usually infinite series.

[SI*] An initial value problem consists of a differential equation such as Eq. (3.31), (3.33), or (3.34) together with a pair of initial conditions

$$\begin{aligned} y(t_0) &= y_0 \\ y'(t_0) &= y'_0 \end{aligned} \quad (3.36)$$

where y_0 and y'_0 are given numbers prescribing values for y and y' at the initial point t_0 . Observe that the initial conditions for a second order equation identify not only a particular point (t_0, y_0) through which the graph of the solution must pass, but also the slope y'_0 of the graph at that point. It is reasonable to expect that two initial conditions are needed for a second order equation because, roughly speaking, two integrations are required to find a solution and each integration introduces an arbitrary constant. Presumably, two initial conditions will suffice to determine values for these two constants.

[SI*] A second order linear equation is said to be homogeneous if the term $g(t)$ in Eq. (3.33), or the term $G(t)$ in Eq. (3.34), is zero for all t . Otherwise, the equation is called nonhomogeneous. As a result, the term $g(t)$, or $G(t)$, is sometimes called the nonhomogeneous term. We begin our discussion with homogeneous equations, which we will write in the form

$$P(t)y'' + Q(t)y' + R(t)y = 0 \quad (3.37)$$

Once the homogeneous equation has been solved, it is always possible to solve the corresponding nonhomogeneous equation (3.34), or at least to express the solution in terms of an integral. Thus the problem of solving the homogeneous equation is the more fundamental one.

[SI*] Consider the general second order linear differential equation

$$ay'' + by' + cy = 0 \quad (3.38)$$

whose coefficients a, b , and c are arbitrary (real) constants. We can solve Eq. (3.38) for any values of its coefficients and also satisfy any given set of initial conditions for y and y' . We start by seeking exponential solutions of the form $y = e^{rt}$, where r is a parameter to be determined. Then it follows that

$$\begin{aligned} y' &= re^{rt} \\ y'' &= r^2e^{rt} \end{aligned}$$

By substituting these expressions for y, y' , and y'' in Eq. (3.38), we obtain

$$(ar^2 + br + c)e^{rt} = 0$$

or, since $e^{rt} \neq 0$,

$$ar^2 + br + c = 0 \quad (3.39)$$

Equation (3.39) is called the characteristic equation for the differential equation (3.38). Its significance lies in the fact that if r is a root of the polynomial equation (3.39), then $y = e^{rt}$ is a solution of the differential equation (3.38). Since Eq. (3.39) is a quadratic equation with real coefficients, it has two roots, which may be real and different, real but repeated, or complex conjugates.

Assuming that the roots of the characteristic equation (3.39) are real and different, let them be denoted by r_1 and r_2 , where $r_1 \neq r_2$. Then $y_1(t) = e^{r_1 t}$ and $y_2(t) = e^{r_2 t}$ are two solutions of Eq. (3.38). It now follows that

$$y = c_1 y_1(t) + c_2 y_2(t) = c_1 e^{r_1 t} + c_2 e^{r_2 t} \quad (3.40)$$

is also a solution of Eq. (3.38). To verify that this is so, we can differentiate the expression in Eq. (3.40); hence

$$y' = c_1 r_1 e^{r_1 t} + c_2 r_2 e^{r_2 t} \quad (3.41)$$

and

$$y'' = c_1 r_1^2 e^{r_1 t} + c_2 r_2^2 e^{r_2 t} \quad (3.42)$$

Substituting these expressions for y, y' and y'' in Eq. (3.38) and rearranging terms, we obtain

$$ay'' + by' + cy = c_1(ar_1^2 + br_1 + c)e^{r_1 t} + c_2(ar_2^2 + br_2 + c)e^{r_2 t} \quad (3.43)$$

The quantity in each of the parentheses on the right side of Eq. (3.43) is zero because r_1 and r_2 are roots of Eq. (3.39); therefore, y as given by Eq. (3.40) is indeed a solution of Eq. (3.38), as we wished to verify.

Now suppose that we want to find the particular member of the family solutions (3.40) that satisfies the initial conditions (3.36)

$$y(t_0) = y_0$$

$$y'(t_0) = y'_0$$

By substituting $t = t_0$ and $y = y_0$ in Eq. (3.40), we obtain

$$c_1 e^{r_1 t_0} + c_2 e^{r_2 t_0} = y_0 \quad (3.44)$$

Similarly, setting $t = t_0$ and $y' = y'_0$ in Eq. (3.41) gives

$$c_1 r_1 e^{r_1 t_0} + c_2 r_2 e^{r_2 t_0} = y'_0 \quad (3.45)$$

On solving Eqs. (3.44) and (3.45) simultaneously for c_1 and c_2 , we find that

$$\begin{aligned} c_1 &= \frac{y'_0 - y_0 r_2}{r_1 - r_2} e^{-r_1 t_0} \\ c_2 &= \frac{y_0 r_1 - y'_0}{r_1 - r_2} e^{-r_2 t_0} \end{aligned} \quad (3.46)$$

recall that $r_1 - r_2 \neq 0$ so that the expressions in Eq. (3.46) always make sense. Thus, no matter what initial conditions are assigned—that is, regardless of the values of t_0, y_0 , and y'_0 in Eqs. (3.36)—it is always possible to determine c_1 and c_2 so that the initial conditions are satisfied. Moreover, there is only one possible choice of c_1 and c_2 for each set of initial conditions. With the values of c_1 and c_2 given by Eq. (3.46), the expression (3.40) is the solution of the initial value problem

$$\begin{aligned} ay'' + by' + cy &= 0 \\ y(t_0) &= y_0 \\ y'(t_0) &= y'_0 \end{aligned} \quad (3.47)$$

[SI*] Solutions of Linear Homogeneous Equations; the Wronskian

We will introduce a differential operator notation. Let p and q be continuous functions on an open interval I , that is, for $\alpha < t < \beta$. The cases $\alpha = -\infty$, or $\beta = \infty$, or both, are included.

Then, for any function ϕ that is twice differentiable on I we define the differential operator L by the equation

$$L[\phi] = \phi'' + p\phi' + q\phi \quad (3.48)$$

Note that $L[\phi]$ is a function on I . The value of $L[\phi]$ at a point t is

$$L[\phi](t) = \phi''(t) + p(t)\phi'(t) + q(t)\phi(t)$$

For example, if $p(t) = t^2$, $q(t) = 1 + t$, and $\phi(t) = \sin 3t$, then

$$\begin{aligned} L[\phi](t) &= (\sin 3t)'' + t^2(\sin 3t)' + (1 + t)\sin 3t \\ &= -9\sin 3t + 3t^2 \cos 3t + (1 + t)\sin 3t \end{aligned}$$

The operator L is often written as $L = D^2 + pD + q$, where D is the derivative operator.

Since it is customary to use the symbol y to denote $\phi(t)$, we will usually write the second order linear homogeneous equation in the form

$$L[y] = y'' + p(t)y' + q(t)y = 0 \quad (3.49)$$

With Eq. (3.49) we associate a set of initial conditions

$$y(t_0) = y_0, \quad y'(t_0) = y'_0 \quad (3.50)$$

where t_0 is any point in the interval I , and y_0 and y'_0 are given real numbers.

Theorem 3.1: Existence and Uniqueness Theorem

Consider the initial value problem

$$\begin{aligned} y'' + p(t)y' + q(t)y &= g(t) \\ y(t_0) &= y_0 \\ y'(t_0) &= y'_0 \end{aligned} \quad (3.51)$$

where p, q , and g are continuous on an open interval I that contains the point t_0 . Then there is exactly one solution $y = \phi(t)$ of this problem, and the solution exists throughout the interval I .

Theorem 3.2: Principle of Superposition

If y_1 and y_2 are two solutions of the differential equation (3.49),

$$L[y] = y'' + p(t)y' + q(t)y = 0$$

then the linear combination $c_1y_1 + c_2y_2$ is also a solution for any values of the constants c_1 and c_2 .

A special case of Principle of Superposition theorem occurs if either c_1 or c_2 is zero. Then we conclude that any constant multiple of a solution of Eq. (3.49) is also a solution. To prove the theorem, we need only substitute

$$y = c_1y_1(t) + c_2y_2(t) \quad (3.52)$$

for y in Eq. (3.49). By calculating the indicated derivatives and rearranging terms, we obtain

$$\begin{aligned} L[c_1y_1 + c_2y_2] &= [c_1y_1 + c_2y_2]'' + p[c_1y_1 + c_2y_2]' + q[c_1y_1 + c_2y_2] \\ &= c_1y_1'' + c_2y_2'' + c_1py_1' + c_2py_2' + c_1qy_1 + c_2qy_2 \\ &= c_1[y_1'' + py_1' + qy_1] + c_2[y_2'' + py_2' + qy_2] \\ &= c_1L[y_1] + c_2L[y_2] \end{aligned}$$

Since $L[y_1] = 0$ and $L[y_2] = 0$, it follows that $L[c_1y_1 + c_2y_2] = 0$ also. Therefore, regardless of the values of c_1 and c_2 , y as given by Eq. (3.52) does satisfy the differential equation (3.49), and the proof for the theorem is complete.

The theorem states that, beginning with only two solutions of Eq. (3.49), we can construct an infinite family of solutions by means of Eq. (3.52). The next question is whether all solutions of Eq. (3.49) are included in Eq. (3.52) or whether there may be other solutions of a different form. We begin to address this question by examining whether the constants c_1 and c_2 in Eq. (3.52) can be chosen so as to satisfy the initial conditions (3.50). These initial conditions require c_1 and c_2 to satisfy the equations

$$\begin{aligned} c_1y_1(t_0) + c_2y_2(t_0) &= y_0 \\ c_1y_1'(t_0) + c_2y_2'(t_0) &= y_0' \end{aligned} \tag{3.53}$$

The determinant of coefficients of the system (3.54) is

$$W = \begin{vmatrix} y_1(t_0) & y_2(t_0) \\ y_1'(t_0) & y_2'(t_0) \end{vmatrix} = y_1(t_0)y_2'(t_0) - y_1'(t_0)y_2(t_0) \tag{3.54}$$

If $W \neq 0$, then Eqs (3.53) have a unique solution (c_1, c_2) regardless of the values of y_0 and y_0' . This solution is given by

$$\begin{aligned} c_1 &= \frac{y_0y_2'(t_0) - y_0'y_2(t_0)}{y_1(t_0)y_2'(t_0) - y_1'(t_0)y_2(t_0)} \\ c_2 &= \frac{-y_0y_1'(t_0) + y_0'y_1(t_0)}{y_1(t_0)y_2'(t_0) - y_1'(t_0)y_2(t_0)} \end{aligned} \tag{3.55}$$

or, in terms of determinants,

$$c_1 = \frac{\begin{vmatrix} y_0 & y_2(t_0) \\ y_0' & y_2'(t_0) \end{vmatrix}}{\begin{vmatrix} y_1(t_0) & y_2(t_0) \\ y_1'(t_0) & y_2'(t_0) \end{vmatrix}}, \quad c_2 = \frac{\begin{vmatrix} y_1(t_0) & y_0 \\ y_1'(t_0) & y_0' \end{vmatrix}}{\begin{vmatrix} y_1(t_0) & y_2(t_0) \\ y_1'(t_0) & y_2'(t_0) \end{vmatrix}} \tag{3.56}$$

with these values for c_1 and c_2 , the expression (3.52) satisfies the initial conditions (3.50) as well as the differential equation (3.49).

On the other hand, if $W = 0$, then Eqs. (3.53) have no solution unless y_0 and y_0' satisfy a certain additional conditions; in this case there are infinitely many solutions.

The determinant W is called the Wronskian determinant, or simply the Wronskian, of the solutions y_1 and y_2 . Sometimes we use the more extended notation $W(y_1, y_2)(t_0)$ to stand

for the expression on the right side of Eq. (3.54), thereby emphasizing that the Wronskian depends on the functions y_1 and y_2 , and that it is evaluated at the point t_0 .

Theorem 3.3: Wronskian for Second Order Linear Differential Equations

Suppose that y_1 and y_2 are two solutions of Eq. (3.49)

$$L[y] = y'' + p(t)y' + q(t)y = 0$$

and that the initial conditions (3.50)

$$y(t_0) = y_0, \quad y'(t_0) = y'_0$$

are assigned. Then it is always possible to choose the constants c_1, c_2 so that

$$y = c_1 y_1(t) + c_2 y_2(t)$$

satisfies the differential equation (3.49) and the initial conditions (3.50) if and only if the Wronskian

$$W = y_1 y'_2 - y'_1 y_2$$

is not zero at t_0 .

Theorem 3.4: Fundamental Set of Solutions

Consider the differential equation (3.49)

$$L[y] = y'' + p(t)y' + q(t)y = 0$$

whose coefficients p and q are continuous on some open interval I . Choose some point t_0 in I . Let y_1 be the solution of Eq. (3.49) that also satisfies the initial conditions

$$y(t_0) = 1, \quad y'(t_0) = 0$$

and let y_2 be the solution of Eq. (3.49) that satisfies the initial conditions

$$y(t_0) = 0, \quad y'(t_0) = 1$$

Then y_1 and y_2 form a fundamental set of solutions of Eq. (3.49). If you compute the Wronskian it will be $W(y_1, y_2)(t_0) = 1$, since the Wronskian is not zero at the point t_0 , the functions y_1 and y_2 do form a fundamental set of solutions.

Theorem 3.5: Abel's Theorem

If y_1 and y_2 are solutions of the differential equation

$$L[y] = y'' + p(t)y' + q(t)y = 0$$

where p and q are continuous on an open interval I , then the Wronskian $W(y_1, y_2)(t)$ is given by

$$W(y_1, y_2)(t) = ce^{-\int p(t) dt} \quad (3.57)$$

where c is a certain constant that depends on y_1 and y_2 , but not on t . Further, $W(y_1, y_2)(t)$ either is zero for all t in I (if $c = 0$) or else is never zero in I (if $c \neq 0$).

[SI*] **The Adjoint Equation**

If a second order linear homogeneous equation is not exact, it can be made exact by multiplying by an appropriate integrating factor $\mu(x)$. Thus we require that $\mu(x)$ be such that

$$\mu(x)P(x)y'' + \mu(x)Q(x)y' + \mu(x)R(x)y = 0$$

can be written in the form

$$[\mu(x)P(x)y']' + [f(x)y]' = 0$$

By equating coefficients in these two equations and eliminating $f(x)$, show that the function μ must satisfy

$$P\mu'' + (2P' - Q)\mu' + (P'' - Q' + R)\mu = 0$$

This equation is known as the adjoint of the original equation and is important in the advanced theory of differential equations. In general, the problem of solving the adjoint differential equation is as difficult as that of solving the original equation, so only occasionally is it possible to find an integrating factor for a second order equation.

Solution:

Suppose we have the second-order ODE,

$$P(x)y'' + Q(x)y' + R(x)y = 0$$

To make it exact, multiply both sides by an integrating factor $\mu = \mu(x)$

$$\mu(x)P(x)y'' + \mu(x)Q(x)y' + \mu(x)R(x)y = 0 \quad (3.58)$$

Now that it's exact, it can be written in the form,

$$[\mu(x)P(x)y']' + [f(x)y]' = 0$$

Expand the left side

$$\mu'(x)P(x)y' + \mu(x)P'(x)y' + \mu(x)P(x)y'' + f'(x)y + f(x)y' = 0$$

Factor it now

$$\mu(x)P(x)y'' + [\mu'(x)P(x) + \mu(x)P'(x) + f(x)]y' + f'(x)y = 0$$

Equate the coefficients with those of equation (3.58)

$$\begin{aligned}\mu'(x)P(x) + \mu(x)P'(x) + f(x) &= \mu(x)Q(x) \\ f'(x) &= \mu(x)R(x)\end{aligned}$$

Differentiate both sides of the first equation with respect to x .

$$\mu''(x)P(x) + \mu'(x)P'(x) + \mu(x)P''(x) + f'(x) = \mu'(x)Q(x) + \mu(x)Q'(x)$$

Substitute $\mu(x)R(x)$ for $f'(x)$

$$\mu''(x)P(x) + \mu'(x)P'(x) + \mu(x)P''(x) + \mu(x)R(x) = \mu'(x)Q(x) + \mu(x)Q'(x)$$

Therefore, the adjoint is

$$P\mu'' + (2P' - Q)\mu' + (P'' - Q' + R)\mu = 0$$

[SI*] **Example:**

Find the adjoint of the given differential equation

(a) $x^2y'' + xy' + (x^2 - v^2)y = 0$ (Bessel's equation)

(b) $(1 - x^2)y'' - 2xy' + \alpha(\alpha + 1)y = 0$ (Legendre's equation)

(c) $y'' - xy = 0$ (Airy's equation)

Solution:

(a) To make the ODE exact, multiply both sides by an integrating factor $\mu = \mu(x)$

$$x^2\mu(x)y'' + x\mu(x)y' + (x^2 - v^2)\mu(x)y = 0 \quad (3.59)$$

Now that it's exact, it can be written in the form

$$\left[x^2\mu(x)y' \right]' + [f(x)y]' = 0$$

Expand the left side

$$2x\mu(x)y' + x^2\mu'(x)y' + x^2\mu(x)y'' + f'(x)y + f(x)y' = 0$$

Factor it now

$$x^2\mu(x)y'' + \left[x^2\mu'(x) + 2x\mu(x) + f(x) \right] y' + f'(x)y = 0$$

Equate the coefficients with those of equation (3.59)

$$\begin{aligned}x^2\mu'(x) + 2x\mu(x) + f(x) &= x\mu(x) \\ f'(x) &= (x^2 - v^2)\mu(x)\end{aligned}$$

Differentiate both sides of the first equation with respect to x

$$2x\mu'(x) + x^2\mu''(x) + 2\mu(x) + 2x\mu'(x) + f'(x) = \mu(x) + x\mu'(x)$$

Substitute $(x^2 - v^2)\mu(x)$ for $f'(x)$

$$2x\mu'(x) + x^2\mu''(x) + 2\mu(x) + 2x\mu'(x) + (x^2 - v^2)\mu(x) = \mu(x) + x\mu'(x)$$

Bring all terms to the left side and combine like terms

$$x^2\mu''(x) + 3x\mu'(x) + (x^2 - v^2 + 1)\mu(x) = 0$$

(b) To make the ODE exact, multiply both sides by an integrating factor $\mu = \mu(x)$

$$(1 - x^2)\mu(x)y'' - 2x\mu(x)y' + \alpha(\alpha + 1)\mu(x)y = 0 \quad (3.60)$$

Now that it's exact, it can be written in the form

$$\left[(1 - x^2)\mu(x)y' \right]' + [f(x)y]' = 0$$

Expand the left side

$$-2x\mu(x)y' + (1 - x^2)\mu'(x)y' + (1 - x^2)\mu(x)y'' + f'(x)y + f(x)y' = 0$$

Factor it now.

$$(1 - x^2)\mu(x)y'' + \left[(1 - x^2)\mu'(x) - 2x\mu(x) + f(x) \right] y' + f'(x)y = 0$$

Equate the coefficients with those of equation (3.60)

$$\begin{aligned} (1 - x^2)\mu'(x) - 2x\mu(x) + f(x) &= -2x\mu(x) \\ f'(x) &= \alpha(\alpha + 1)\mu(x) \end{aligned}$$

Add $2x\mu(x)$ to both sides of the first equation

$$(1 - x^2)\mu'(x) + f(x) = 0$$

Differentiate both sides with respect to x .

$$-2x\mu'(x) + (1 - x^2)\mu''(x) + f'(x) = 0$$

Substitute $\alpha(\alpha + 1)\mu(x)$ for $f'(x)$

$$-2x\mu'(x) + (1 - x^2)\mu''(x) + \alpha(\alpha + 1)\mu(x) = 0$$

Therefore,

$$(1 - x^2)\mu''(x) - 2x\mu'(x) + \alpha(\alpha + 1)\mu(x) = 0$$

(c) To make the ODE exact, multiply both sides by an integrating factor $\mu = \mu(x)$

$$\mu(x)y'' - x\mu(x)y' = 0 \quad (3.61)$$

Now that it's exact, it can be written in the form

$$[\mu(x)y']' + [f(x)y]' = 0$$

Expand the left side

$$\mu'(x)y' + \mu(x)y'' + f'(x)y + f(x)y' = 0$$

Factor it now

$$\mu(x)y'' + [\mu'(x) + f(x)] y' + f'(x)y = 0$$

Equate the coefficients with those of equation (3.61)

$$\begin{aligned} \mu'(x) + f(x) &= 0 \\ f'(x) &= -x\mu(x) \end{aligned}$$

Differentiate both sides of the first equation with respect to x

$$\mu''(x) + f'(x) = 0$$

Substitute $-x\mu(x)$ for $f'(x)$

$$\mu''(x) - x\mu(x) = 0$$

[SI*] Euler' Equations

An equation of the form

$$t^2 \frac{d^2 y}{dt^2} + \alpha \frac{dy}{dt} + \beta y = 0, \quad t > 0 \quad (3.62)$$

where α and β are real constants, is called an Euler equation.

- (a) Let $x = \ln t$ and calculate $\frac{dy}{dt}$ and $\frac{d^2 y}{dt^2}$ in terms of $\frac{dy}{dx}$ and $\frac{d^2 y}{dx^2}$.
 (b) Use the results of part (a) to transform Eq. (3.62) into

$$\frac{d^2 y}{dx^2} + (\alpha - 1) \frac{dy}{dx} + \beta y = 0 \quad (3.63)$$

Observe that Eq. (3.63) has constant coefficients. If $y_1(x)$ and $y_2(x)$ form a fundamental set of solutions of Eq. (3.63), then $y_1(\ln t)$ and $y_2(\ln t)$ form a fundamental set of solutions of Eq. (3.62).

Solution:

- (a) Make the substitution $x = \ln t$ in the ODE. Then

$$e^x = t \rightarrow e^{2x} = t^2$$

and the ODE becomes

$$e^{2x} \frac{d^2 y}{dt^2} + \alpha e^x \frac{dy}{dt} + \beta y = 0$$

The aim now is to find what the derivatives are in terms of this new variable by using the chain rule.

$$\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt} = \frac{dy}{dx} \left(\frac{1}{t} \right) = \frac{dy}{dx} \left(\frac{1}{e^x} \right) = e^{-x} \frac{dy}{dx}$$

$$\begin{aligned} \frac{d^2 y}{dt^2} &= \frac{d}{dt} \left(\frac{dy}{dt} \right) \\ &= \frac{dx}{dt} \frac{d}{dx} \left(e^{-x} \frac{dy}{dx} \right) \\ &= \frac{1}{t} \left(-e^{-x} \frac{dy}{dx} + e^{-x} \frac{d^2 y}{dx^2} \right) \\ &= \frac{1}{e^x} \left(-e^{-x} \frac{dy}{dx} + e^{-x} \frac{d^2 y}{dx^2} \right) \end{aligned}$$

- (b) Substitute these expressions into the ODE

$$\begin{aligned} e^{2x} \frac{1}{e^x} \left(-e^{-x} \frac{dy}{dx} + e^{-x} \frac{d^2 y}{dx^2} \right) + \alpha e^x \left(e^{-x} \frac{dy}{dx} \right) + \beta y &= 0 \\ e^x \left(-e^{-x} \frac{dy}{dx} + e^{-x} \frac{d^2 y}{dx^2} \right) + \alpha \frac{dy}{dx} + \beta y &= 0 \\ -\frac{dy}{dx} + \frac{d^2 y}{dx^2} + \alpha \frac{dy}{dx} + \beta y &= 0 \end{aligned}$$

Therefore,

$$\frac{d^2 y}{dx^2} + (\alpha - 1) \frac{dy}{dx} + \beta y = 0$$

[SI*] **Complex Roots of the Characteristic Equation**

Suppose we have the homogeneous second order linear differential equation

$$ay'' + by' + cy = 0 \quad (3.64)$$

We have learned that if we seek solutions of the form $y = e^{rt}$, then r must be a root of the characteristic equation

$$ar^2 + br + c = 0 \quad (3.65)$$

If the roots r_1 and r_2 are real and different, which occurs whenever the discriminant $b^2 - 4ac$ is positive, then the general solution of Eq. (3.64) is

$$y = c_1 e^{r_1 t} + c_2 e^{r_2 t} \quad (3.66)$$

Suppose now that $b^2 - 4ac$ is negative. Then the roots of Eq. (3.65) are conjugate complex numbers; we denote them by

$$\begin{aligned} r_1 &= \lambda + i\mu \\ r_2 &= \lambda - i\mu \end{aligned} \quad (3.67)$$

where λ and μ are real. The corresponding expressions for y are

$$\begin{aligned} y_1(t) &= e^{(\lambda+i\mu)t} \\ y_2(t) &= e^{(\lambda-i\mu)t} \end{aligned} \quad (3.68)$$

[SI*] **Euler's Formula**

To assign a meaning to the expressions in Eqs. (3.68), we need to give a definition of the complex exponential function. Recall from Calculus that the Taylor series for e^t about $t = 0$ is

$$e^t = \sum_{n=0}^{\infty} \frac{t^n}{n!}, \quad -\infty < t < \infty \quad (3.69)$$

If we now assume that we can substitute it for t in Eq. (3.69), then we have

$$\begin{aligned} e^{it} &= \sum_{n=0}^{\infty} \frac{(it)^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n}}{(2n)!} + i \sum_{n=0}^{\infty} \frac{(-1)^{n-1} t^{2n-1}}{(2n-1)!} \end{aligned} \quad (3.70)$$

where we have separated the sum into its real and imaginary parts, making use of the fact that $i^2 = -1$, $i^3 = -i$, $i^4 = 1$, and so forth. The first series in Eq. (3.70) is precisely the Taylor series for $\cos t$ about $t = 0$, and the second is the Taylor series for $\sin t$ about $t = 0$. Thus we have

$$e^{it} = \cos t + i \sin t \quad (3.71)$$

Equation (3.71) is known as Euler's formula and is an extremely important mathematical relationship.

There are some variations of Euler's formula that are also worth noting

$$e^{-it} = \cos t - i \sin t \quad (3.72)$$

Further, if t is replaced by μt , then we obtain a generalized version of Euler's formula, namely

$$e^{i\mu t} = \cos \mu t + i \sin \mu t \quad (3.73)$$

Next, we want to extend the definition of the exponential function to arbitrary complex exponents of the form $(\lambda + i\mu)t$. Since we want the usual properties of the exponential function to hold for complex exponents, we certainly want $e^{(\lambda+i\mu)t}$ to satisfy

$$e^{(\lambda+i\mu)t} = e^{\lambda t} e^{i\mu t} \quad (3.74)$$

Then, substituting for $e^{i\mu t}$ from Eq. (3.73), we obtain

$$\begin{aligned} e^{(\lambda+i\mu)t} &= e^{\lambda t} (\cos \mu t + i \sin \mu t) \\ &= e^{\lambda t} \cos \mu t + i e^{\lambda t} \sin \mu t \end{aligned} \quad (3.75)$$

We now take Eq. (3.75) as the definition of $e^{(\lambda+i\mu)t}$. The value of the exponential function with a complex exponent is a complex number whose real and imaginary parts are given by the terms on the right side of Eq. (3.75). Observe that the real and imaginary parts of $e^{(\lambda+i\mu)t}$ are expressed entirely in terms of elementary real-valued functions.

[SI*] **Complex Roots; The General Case**

The functions $y_1(t)$ and $y_2(t)$, given by Eqs. (3.68) and with the meaning expressed by Eq. (3.75), are solutions of Eq. (3.64) when the roots of the characteristic equation (3.65) are complex numbers $\lambda \pm i\mu$. Unfortunately, the solutions y_1 and y_2 are complex-valued functions, whereas in general we would prefer to have real-valued solutions, if possible, because the differential equation itself has real coefficients. To find a fundamental set of real-valued solutions let us form the sum and then the difference of y_1 and y_2 . We have

$$\begin{aligned} y_1(t) + y_2(t) &= e^{\lambda t} (\cos \mu t + i \sin \mu t) + e^{\lambda t} (\cos \mu t - i \sin \mu t) \\ &= 2e^{\lambda t} \cos \mu t \end{aligned}$$

and

$$\begin{aligned} y_1(t) - y_2(t) &= e^{\lambda t} (\cos \mu t + i \sin \mu t) - e^{\lambda t} (\cos \mu t - i \sin \mu t) \\ &= 2ie^{\lambda t} \sin \mu t \end{aligned}$$

Hence, neglecting the constant multipliers 2 and $2i$, respectively, we have obtained a pair of real-valued solutions

$$\begin{aligned} u(t) &= e^{\lambda t} \cos \mu t \\ v(t) &= e^{\lambda t} \sin \mu t \end{aligned} \quad (3.76)$$

Observe that u and v are simply the real and imaginary parts, respectively, of y_1 . By direct computation you can show that the Wronskian of u and v is

$$W(u, v)(t) = \mu e^{2\lambda t} \quad (3.77)$$

Thus, as long as $\mu \neq 0$, the Wronskian W is not zero, so u and v form a fundamental set of solutions. If $\mu = 0$ then the roots are real and repeated. Consequently, if the roots of the characteristic equation are complex numbers $\lambda \pm i\mu$, with $\mu \neq 0$, then the general solution of Eq. (3.64) is

$$y = c_1 e^{\lambda t} \cos \mu t + c_2 e^{\lambda t} \sin \mu t \quad (3.78)$$

[SI*] Example:

Find the solution of the initial value problem

$$16y'' - 8y' + 145y = 0, \quad y(0) = -2, \quad y'(0) = 1 \quad (3.79)$$

Solution:

The characteristic equation is $16r^2 - 8r + 145 = 0$ and its roots are $r = \frac{1}{4} \pm 3i$. Thus the general solution of the differential equation is

$$y = c_1 e^{t/4} \cos 3t + c_2 e^{t/4} \sin 3t \quad (3.80)$$

To apply the first initial condition, we set $t = 0$ in Eq. (3.80); this gives

$$y(0) = c_1 = -2$$

For the second initial condition we must differentiate Eq. (3.80) and then set $t = 0$. In this way we find that

$$y'(0) = \frac{1}{4}c_1 + 3c_2 = 1$$

from which $c_2 = \frac{1}{2}$. Using these values of c_1 and c_2 in Eq. (3.80), we obtain

$$y = -2e^{t/4} \cos 3t + \frac{1}{2}e^{t/4} \sin 3t \quad (3.81)$$

as the solution of the initial value problem (3.79).




Figure 3.14: The graph of the solution $y = -2e^{t/4} \cos 3t + \frac{1}{2}e^{t/4} \sin 3t$ of the initial value problem $16y'' - 8y' + 145y = 0, y(0) = -2, y'(0) = 1$ (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Growing Oscillation/main.cpp).

In this case we observe that the solution is a growing oscillation. The trigonometric factors in Eq. (3.81) determine the oscillatory part of the solution, while the exponential factor (with a positive exponent) causes the magnitude of the oscillation to increase with time.

[SI*] **Repeated Roots**

We showed how to solve the equation

$$ay'' + by' + cy = 0 \quad (3.82)$$

when the roots of the characteristic equation

$$ar^2 + br + c = 0 \quad (3.83)$$

either are real and different or are complex conjugates. Now we consider the third possibility, namely, that the two roots r_1 and r_2 are equal. This case is transitional between the other two and occurs when the discriminant $b^2 - 4ac$ is zero. Then it follows from the quadratic formula that

$$r_1 = r_2 = -\frac{b}{2a} \quad (3.84)$$

The difficulty is immediately apparent; both roots yield the same solution

$$y_1(t) = e^{-\frac{bt}{2a}} \quad (3.85)$$

of the differential equation (3.82), and it is not obvious how to find a second solution.

To find a second solution, we assume that

$$y = v(t)y_1(t) = v(t)e^{-\frac{bt}{2a}} \quad (3.86)$$

and substitute for y in Eq. (3.82) to determine $v(t)$. We have

$$y' = v'(t)e^{-\frac{bt}{2a}} - \frac{b}{2a}v(t)e^{-\frac{bt}{2a}} \quad (3.87)$$

and

$$y'' = v''(t)e^{-\frac{bt}{2a}} - \frac{b}{a}v'(t)e^{-\frac{bt}{2a}} + \frac{b^2}{4a^2}v(t)e^{-\frac{bt}{2a}} \quad (3.88)$$

Then, by substituting in Eq. (3.82), we obtain

$$\left[a \left(v''(t) - \frac{b}{a}v'(t) + \frac{b^2}{4a^2}v(t) \right) + b \left(v'(t) - \frac{b}{2a}v(t) \right) + cv(t) \right] e^{-\frac{bt}{2a}} = 0 \quad (3.89)$$

Canceling the factor $e^{-\frac{bt}{2a}}$, which is nonzero, and rearranging the remaining terms, we find that

$$av''(t) + (-b + b)v'(t) + \left(\frac{b^2}{4a} - \frac{b^2}{2a} + c \right) v(t) = 0 \quad (3.90)$$

The term involving $v'(t)$ is obviously zero. Further, the coefficient of $v(t)$ is $c - \left(\frac{b^2}{4a} \right)$, which is also zero because $b^2 - 4ac = 0$ (repeated roots). Thus, Eq. (3.90) reduces to

$$v''(t) = 0$$

therefore

$$v(t) = c_1 + c_2 t$$

Hence, from Eq. (3.86), we have

$$y = c_1 e^{-\frac{bt}{2a}} + c_2 t e^{-\frac{bt}{2a}} \quad (3.91)$$

Thus y is a linear combination of the two solutions

$$y_1(t) = e^{-\frac{bt}{2a}}, \quad y_2(t) = t e^{-\frac{bt}{2a}} \quad (3.92)$$

The Wronskian of these two solutions is

$$W(y_1, y_2)(t) = \begin{vmatrix} e^{-\frac{bt}{2a}} & t e^{-\frac{bt}{2a}} \\ -\frac{b}{2a} e^{-\frac{bt}{2a}} & \left(1 - \frac{bt}{2a}\right) e^{-\frac{bt}{2a}} \end{vmatrix} = e^{-\frac{bt}{a}} \quad (3.93)$$

Since $W(y_1, y_2)(t)$ is never zero, the solutions y_1 and y_2 given by Eq. (3.92) are a fundamental set of solutions. Further Eq. (3.91) is the general solution of Eq. (3.82) when the roots of the characteristic equation are equal.

[SI*] **Example:**

Find the solution of the initial value problem

$$y'' - y' + 0.25y = 0, \quad y(0) = 2, \quad y'(0) = \frac{1}{3} \quad (3.94)$$

Solution:

The characteristic equation is

$$r^2 - r + 0.25 = 0$$

so the roots are $r_1 = r_2 = \frac{1}{2}$. Thus the general solution of the differential equation is

$$y = c_1 e^{t/2} + c_2 t e^{t/2} \quad (3.95)$$

The first initial condition requires that

$$y(0) = c_1 = 2$$

to satisfy the second initial condition, we first differentiate Eq. (3.95) and then set $t = 0$. This gives

$$y'(0) = \frac{1}{2}c_1 + c_2 = \frac{1}{3}$$

so $c_2 = -\frac{2}{3}$. Thus the solution of the initial value problem is

$$y = 2e^{t/2} - \frac{2}{3}te^{t/2} \quad (3.96)$$

[SI*] **Reduction of Order**

Suppose that we know one solution $y_1(t)$, not everywhere zero, of

$$y'' + p(t)y' + q(t)y = 0 \quad (3.97)$$

To find a second solution, let

$$y = v(t)y_1(t) \quad (3.98)$$

then

$$y' = v'(t)y_1(t) + v(t)y_1'(t)$$

and

$$y'' = v''(t)y_1(t) + 2v'(t)y_1'(t) + v(t)y_1''(t)$$

Substituting for $y, y',$ and y'' in Eq. (3.97) and collecting terms, we find that

$$y_1 v'' + (2y_1' + py_1)v' + (y_1'' + py_1' + qy_1)v = 0 \quad (3.99)$$

Since y_1 is a solution of Eq. (3.97), the coefficient of v in Eq. (3.99) is zero, so that Eq. (3.99) becomes

$$y_1 v'' + (2y_1' + py_1)v' = 0 \quad (3.100)$$

Despite its appearance, Eq. (3.100) is actually a first order equation for the function v' and can be solved either as a first order linear equation or as a separable equation. Once v' has been found, then v is obtained by an integration. Finally, y is determined from Eq. (3.98). This procedure is called the method of reduction of order, because the crucial step is the solution of a first order differential equation for v' rather than the original second order equation for y .

[SI*] **Example:**

Given that $y_1(t) = t^{-1}$ is a solution of

$$2t^2 y'' + 3ty' - y = 0, \quad t > 0 \quad (3.101)$$

find a fundamental set of solutions.

Solution:

We set $y = v(t)t^{-1}$; then

$$\begin{aligned} y' &= v't^{-1} - vt^{-2} \\ y'' &= v''t^{-1} - 2v't^{-2} + 2vt^{-3} \end{aligned}$$

Substituting for $y, y',$ and y'' in Eq. (3.101) and collecting terms, we obtain

$$\begin{aligned} 2t^2 y'' + 3ty' - y &= 2t^2(v''t^{-1} - 2v't^{-2} + 2vt^{-3}) + 3t(v't^{-1} - vt^{-2}) - vt^{-1} \\ &= 2tv'' + (-4 + 3)v' + (4t^{-1} - 3t^{-1} - t^{-1})v \\ &= 2tv'' - v' = 0 \end{aligned} \quad (3.102)$$

Note that the coefficient of v is zero, as it should be; this provides a useful check on our algebra.

Separating the variables in Eq. (3.102) and solving for $v'(t)$, we find that

$$v'(t) = ct^{1/2}$$

then

$$v(t) = \frac{2}{3}ct^{3/2} + k$$

It follows that

$$y = \frac{2}{3}ct^{1/2} + kt^{-1} \quad (3.103)$$

where c and k are arbitrary constants. The second term on the right side of Eq. (3.103) is a multiple of $y_1(t)$ and can be dropped, but the first term provides a new solution $y_2(t) = t^{1/2}$. You can verify that the Wronskian of y_1 and y_2 is

$$W(y_1, y_2)(t) = \frac{3}{2}t^{-3/2}, \quad t > 0 \quad (3.104)$$

Consequently, y_1 and y_2 form a fundamental set of solutions of Eq. (3.101).

```
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Reduction of Order ]# ./main
*****
Reduction of order with y_{1}(t) : t^{(-1)}
y : v*t^{(-1)}
y' : t^{(-1)}*v'-t^{(-2)}*v
y'' : t^{(-1)}*v''-2*v'*t^{(-2)}+2*t^{(-3)}*v
Substituting for y, y', and y'' in the solution:
2*t*v''-v' = 0
v'(t) : C*t^{(0.5)}
v(t) : 0.666667*C*t^{(1.5)}+k
Final v(t) : 0.666667*C*t^{(0.5)}+k*t^{(-1)}
Time taken by function: 132300 microseconds
```

Figure 3.15: The computation of the second solution with reduction of order method (*SymIntegration/Examples/Differential Equations/Test SymIntegration Reduction of Order/main.cpp*).

[SI*] Nonhomogeneous Equations

We now return to the nonhomogeneous equation

$$L[y] = y'' + p(t)y' + q(t)y = g(t) \quad (3.105)$$

where p, q , and g are given (continuous) functions on the open interval I . The equation

$$L[y] = y'' + p(t)y' + q(t)y = 0 \quad (3.106)$$

in which $g(t) = 0$ and p and q are the same as in Eq. (3.105), is called the homogeneous equation corresponding to Eq. (3.105).

Theorem 3.6: Solutions of the Nonhomogeneous Equation

If Y_1 and Y_2 are two solutions of the nonhomogeneous equation (3.105), then their difference $Y_1 - Y_2$ is a solution of the corresponding homogeneous equation (3.106). If, in addition, y_1 and y_2 are a fundamental set of solutions of Eq. (3.106), then

$$Y_1(t) - Y_2(t) = c_1y_1(t) + c_2y_2(t) \quad (3.107)$$

where c_1 and c_2 are certain constants.

Theorem 3.7: The General Solution of the Nonhomogeneous Equation

The general solution of the nonhomogeneous equation (3.105) can be written in the form

$$y = \phi(t) = c_1 y_1(t) + c_2 y_2(t) + Y(t) \quad (3.108)$$

where y_1 and y_2 are a fundamental set of solutions of the corresponding homogeneous equation (3.106), c_1 and c_2 are arbitrary constants, and Y is some specific solution of the nonhomogeneous equation (3.105).

To solve the nonhomogeneous equation (3.105) we must do three things:

1. Find the general solution $c_1 y_1(t) + c_2 y_2(t)$ of the corresponding homogeneous equation. This solution is frequently called the complementary solution and may be denoted by $y_c(t)$.
2. Find some single solution $Y(t)$ of the nonhomogeneous equation. Often this solution is referred to as a particular solution.
3. Add together the functions found in the two preceding steps.

[SI*] Method of Undetermined Coefficients

The method of undetermined coefficients requires us to make an initial assumption about the form of the particular solution $Y(t)$, but with the coefficients left unspecified. We then substitute the assumed expression into Eq. (3.105) and attempt to determine the coefficients so as to satisfy that equation. If we are successful, then we have found a solution of the differential equation (3.105) and can use it for the particular solution $Y(t)$. If we cannot determine the coefficients, then this means that there is no solution of the form that we assumed. In this case we may modify the initial assumption and try again.

The main advantage of the method of undetermined coefficients is that it is straightforward to execute once the assumption is made as to the form of $Y(t)$. Its major limitation is that it is useful primarily for equations for which we can easily write down the correct form of the particular solution in advance. For this reason, this method is usually used only for problems in which the homogeneous equation has constant coefficients and the nonhomogeneous term is restricted to a relatively small class of functions. For more complex class of functions, a computer algebra system can be very helpful in practical applications.

If the nonhomogeneous term $g(t)$ from the Eq. (3.105) is an exponential function e^{at} , then assume that $Y(t)$ is proportional to the same exponential function; if $g(t)$ is $\sin \beta t$ or $\cos \beta t$, then assume that $Y(t)$ is a linear combination of $\sin \beta t$ and $\cos \beta t$; if $g(t)$ is a polynomial, then assume that $Y(t)$ is a polynomial of the same degree with $g(t)$. The same principle extends to the case where $g(t)$ is a product of any two, or all three, of these types of functions.

[SI*] Example:

Find a particular solution of

$$y'' - 3y' - 4y = 2 \sin t \quad (3.109)$$

Solution:

We seek a function $Y(t)$ such that the combination $Y''(t) - 3Y'(t) - 4Y(t)$ is equal to $2 \sin t$. Our assumption concerning $Y(t)$ needs to include cosine as well, since the derivative of sine

and cosine are alternating between sine and cosine. Thus

$$Y(t) = A \sin t + B \cos t \quad (3.110)$$

where A and B are to be determined. Then

$$\begin{aligned} Y'(t) &= A \cos t - B \sin t \\ Y''(t) &= -A \sin t - B \cos t \end{aligned}$$

By substituting these expression for y, y' , and y'' in Eq. (3.110) and collecting terms, we obtain

$$(-A + 3B - 4A) \sin t + (-B - 3A - 4B) \cos t = 2 \sin t \quad (3.111)$$

To satisfy Eq. (3.111), we must match the coefficients of $\sin t$ and $\cos t$ on each side of the equation; thus A and B must satisfy the equations

$$\begin{aligned} -5A + 3B &= 2 \\ -3A - 5B &= 0 \end{aligned}$$

With Gaussian elimination this can be solved. Hence

$$\begin{aligned} A &= -\frac{5}{17} \\ B &= \frac{3}{17} \end{aligned}$$

so a particular solution of Eq. (3.110) is

$$Y(t) = -\frac{5}{17} \sin t + \frac{3}{17} \cos t$$

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Nonhomogeneous Second Order Linear Differential Equations 1# ./main
Solution for y'' - 3y' - 4y = 3e^{2t}

The particular solution is
Y(t) = -0.5*e^{2*t}

Solution for y'' - 3y' - 4y = 2 sin(t)

Y(t) = A*sin(t)+B*cos(t)
Y'(t) = A*cos(t)-B*sin(t)
Y''(t) = -A*sin(t)-B*cos(t)

The particular solution is
Y(t) = -0.294118*sin(t)+0.176471*cos(t)

Solution for y'' - 3y' - 4y = 5 cos(t)

Y(t) = A*sin(t)+B*cos(t)
Y'(t) = A*cos(t)-B*sin(t)
Y''(t) = -A*sin(t)-B*cos(t)

The particular solution is
Y(t) = -0.441176*sin(t)-0.735294*cos(t)

Time taken by function: 108997 microseconds
```

Figure 3.16: The computation of the particular solution for the second order nonhomogeneous differential equation (*SymIntegration/Examples/Differential Equations/Test SymIntegration Nonhomogeneous Second Order Linear Differential Equations/main.cpp*).

[SI*] Example:

Find a particular solution of

$$y'' - 3y' - 4y = 2e^{-t} \quad (3.112)$$

Solution:

We assume that $Y(t) = Ae^{-t}$, we then obtain

$$(A - 3A - 4A)e^{-t} = 2e^{-t} \quad (3.113)$$

Since the left side is zero we will obtain the solution as $\frac{2}{0}e^{-t}$, which means that there is no choice of A that satisfies this equation. Therefore, there is no particular solution of Eq. (3.112) of the assumed form. The reason for this possibly unexpected result becomes clear if we solve the homogeneous equation

$$y'' - 3y' - 4y = 0 \quad (3.114)$$

A fundamental set of solutions of Eq. (3.114) is $y_1(t) = e^{-t}$ and $y_2(t) = e^{4t}$. Thus our assumed particular solution of Eq. (3.112) is actually a solution of the homogeneous equation (3.114); consequently, it cannot possibly be a solution of the nonhomogeneous equation (3.112). To find a solution of Eq. (3.112), we must therefore consider functions of a somewhat different form.

We can simply try to guess the proper form of the particular solution of Eq. (3.112). Since the roots of the homogeneous equation are real and different, we can try to use $Y(t) = te^{-t}$, solution with form of te^{-t} occurs if we have repeated root, so it is safe to use this.

We assume that a particular solution of the form $Y(t) = Ate^{-t}$, then

$$\begin{aligned} Y'(t) &= Ae^{-t} - Ate^{-t} \\ Y''(t) &= -2Ae^{-t} + Ate^{-t} \end{aligned}$$

Substituting these expressions for y, y' , and y'' in Eq. (3.112), we obtain

$$(-2A - 3A)e^{-t} + (A + 3A - 4A)te^{-t} = 2e^{-t}$$

Hence $-5A = 2$, so $A = -\frac{2}{5}$. Thus, a particular solution of Eq. (3.112) is

$$Y(t) = -\frac{2}{5}te^{-t}$$

```
g++ -o main -g -std=c++11 -I../SymIntegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Nonhomogeneous Second Order Linear Differential Equations Special
General solution for y'' - 3y' - 4y = 0

The general solution is:
c1*e^(4*t)+c2*e^(-t)

Particular solution for y'' - 3y' - 4y = 2e^(-t)

The particular solution is
Y(t) = -0.4*t*e^(-t)

Time taken by function: 93265 microseconds
```

Figure 3.17: The computation of the particular solution for the second order nonhomogeneous differential equation (*SymIntegration/Examples/Differential Equations/Test SymIntegration Nonhomogeneous Second Order Linear Differential Equations Special Case/main.cpp*).

[SI*] **Variation of Parameters**

Another method to compute a particular solution of a nonhomogeneous equation is known as variation of parameters. It complements the method of undetermined coefficients rather well. The main advantage of variation of parameters is that it is a general method; in principle at least, it can be applied to any equation, and it requires no detailed assumptions about the form of the solution. On the other hand, the method of variation of parameters eventually requires us to evaluate certain integrals involving the nonhomogeneous term in the differential equation.

[SI*] **Example:**

Find a particular solution of

$$y'' + 4y = 3 \csc t \quad (3.115)$$

Solution:

Observe that this problem is not a good candidate for the method of undetermined coefficients, because the nonhomogeneous term $g(t) = 3 \csc t$ involves a quotient (rather than a sum or a product) of $\sin t$ or $\cos t$.

Observe also that the homogeneous equation corresponding to Eq. (3.115) is

$$y'' + 4y = 0 \quad (3.116)$$

and that the general solution of Eq. (3.116) is

$$y_c(t) = c_1 \cos 2t + c_2 \sin 2t \quad (3.117)$$

The basic idea in the method of variation of parameters is to replace the constants c_1 and c_2 in Eq. (3.117) by functions $u_1(t)$ and $u_2(t)$, respectively, and then to determine these functions so that the resulting expression

$$y = u_1(t) \cos 2t + u_2(t) \sin 2t \quad (3.118)$$

is a solution of the nonhomogeneous equation (3.115).

To determine u_1 and u_2 , we need to substitute for y from Eq. (3.118) in Eq. (3.115). However, even without carrying out this substitution, we can anticipate that the result will be a single equation involving some combination of u_1, u_2 , and their first two derivatives. Since there is only one equation and two unknown functions, we can expect that there are many possible choices of u_1 and u_2 that will meet our needs. Alternatively, we may be able to impose a second condition of our own choosing, thereby obtaining two equations for the two unknown functions u_1 and u_2 . We will soon show (following Lagrange) that it is possible to choose this second condition in a way that makes the computation markedly more efficient.

Returning now to Eq. (3.118), we differentiate it and rearrange the terms, thereby obtaining

$$y' = -2u_1(t) \sin 2t + 2u_2(t) \cos 2t + u_1'(t) \cos 2t + u_2'(t) \sin 2t \quad (3.119)$$

Keeping in mind the possibility of choosing a second condition on u_1 and u_2 , let us require the sum of the last two terms on the right side of Eq. (3.119) to be zero; that is, we require that

$$u_1'(t) \cos 2t + u_2'(t) \sin 2t = 0 \quad (3.120)$$

It then follows from Eq. (3.119) that

$$y' = -2u_1(t) \sin 2t + 2u_2(t) \cos 2t \quad (3.121)$$

Although the ultimate effect of the condition (3.120) is not yet clear, at the very least it has simplified the expression for y' . Further, by differentiating Eq. (3.121), we obtain

$$y'' = -4u_1(t) \cos 2t - 4u_2(t) \sin 2t - 2u_1'(t) \sin 2t + 2u_2'(t) \cos 2t \quad (3.122)$$

Then, substituting for y and y'' in Eq. (3.115) from Eq. (3.118) and (3.122), respectively, we find that u_1 and u_2 must satisfy

$$-2u_1'(t) \sin 2t + 2u_2'(t) \cos 2t = 3 \csc t \quad (3.123)$$

We want to choose u_1 and u_2 to satisfy Eqs. (3.120) and (3.123). These equations can be viewed as a pair of linear algebraic equations for the two unknown quantities $u_1'(t)$ and $u_2'(t)$. Equations (3.120) and (3.123) can be solved in various ways. For example, solving Eq. (3.120) for $u_2'(t)$, we have

$$u_2'(t) = u_1'(t) \frac{\cos 2t}{\sin 2t} \quad (3.124)$$

Then, substituting for $u_2'(t)$ in Eq. (3.123) and simplifying, we obtain

$$u_1'(t) = -\frac{3 \csc t \sin 2t}{2} = -3 \cos t \quad (3.125)$$

Further, putting this expression for $u_1'(t)$ back in Eq. (3.124) and using the double-angle formulas, we find that

$$u_2'(t) = \frac{3 \cos t \cos 2t}{\sin 2t} = \frac{3(1 - 2 \sin^2 t)}{2 \sin t} = \frac{3}{2} \csc t - 3 \sin t \quad (3.126)$$

Having obtained $u_1'(t)$ and $u_2'(t)$, we next integrate so as to find $u_1(t)$ and $u_2(t)$. The result is

$$u_1(t) = -3 \sin t + c_1 \quad (3.127)$$

and

$$u_2(t) = \frac{3}{2} \ln |\csc t - \cot t| + 3 \cos t + c_2 \quad (3.128)$$

On substituting these expressions in Eq. (3.118), we have

$$\begin{aligned} y &= -3 \sin t \cos 2t + \frac{3}{2} \ln |\csc t - \cot t| \sin 2t + 3 \cos t \sin 2t \\ &\quad + c_1 \cos 2t + c_2 \sin 2t \end{aligned}$$

Finally, by using the double-angle formula once more, we obtain

$$y = 3 \sin t + \frac{3}{2} \ln |\csc t - \cot t| \sin 2t + c_1 \cos 2t + c_2 \sin 2t \quad (3.129)$$

The terms in Eq. (3.129) involving the arbitrary constants c_1 and c_2 are the general solution of the corresponding homogeneous equation, while the remaining terms are a particular solution of the nonhomogeneous equation (3.115). Thus Eq. (3.129) is the general solution of Eq. (3.115).

[SI*] Consider

$$y'' + p(t)y' + q(t)y = g(t) \quad (3.130)$$

where p, q , and g are given continuous functions. As a starting point, we assume that we know the general solution

$$y_c(t) = c_1y_1(t) + c_2y_2(t) \quad (3.131)$$

of the corresponding homogeneous equation

$$y'' + p(t)y' + q(t)y = 0 \quad (3.132)$$

The crucial idea is to replace the constants c_1 and c_2 in Eq. (3.131) by functions $u_1(t)$ and $u_2(t)$, respectively; this gives

$$y = u_1(t)y_1(t) + u_2(t)y_2(t) \quad (3.133)$$

Then we try to determine $u_1(t)$ and $u_2(t)$ so that the expression in Eq. (3.133) is a solution of the nonhomogeneous equation (3.130) rather than the homogeneous equation (3.132). Thus we differentiate Eq. (3.133), obtaining

$$y' = u_1'(t)y_1(t) + u_1(t)y_1'(t) + u_2'(t)y_2(t) + u_2(t)y_2'(t) \quad (3.134)$$

We now set the terms involving $u_1'(t)$ and $u_2'(t)$ in Eq. (3.134) equal to zero; that is, we require that

$$u_1'(t)y_1(t) + u_2'(t)y_2(t) = 0 \quad (3.135)$$

Then, from Eq. (3.134), we have

$$y' = u_1(t)y_1'(t) + u_2(t)y_2'(t) \quad (3.136)$$

Further, by differentiating again, we obtain

$$y'' = u_1'(t)y_1'(t) + u_1(t)y_1''(t) + u_2'(t)y_2'(t) + u_2(t)y_2''(t) \quad (3.137)$$

Now, we substitute for y, y' , and y'' in Eq. (3.130) from Eqs. (3.133), (3.136), and (3.137), respectively. After rearranging the terms in the resulting equation, we find that

$$\begin{aligned} & u_1(t) [y_1''(t) + p(t)y_1'(t) + q(t)y_1(t)] \\ & + u_2(t) [y_2''(t) + p(t)y_2'(t) + q(t)y_2(t)] \\ & + u_1'(t)y_1'(t) + u_2'(t)y_2'(t) = g(t) \end{aligned} \quad (3.138)$$

Each of the expression in square brackets in Eq. (3.138) is zero because y_1 and y_2 are solutions of the homogeneous equation (3.132). Therefore Eq. (3.138) reduces to

$$u_1'(t)y_1'(t) + u_2'(t)y_2'(t) = g(t) \quad (3.139)$$

Equations (3.135) and (3.139) form a system of two linear algebraic equations for the derivatives $u_1'(t)$ and $u_2'(t)$ of the unknown functions.

By solving the system (3.135), (3.139) we obtain

$$\begin{aligned} u_1'(t) &= -\frac{y_2(t)g(t)}{W(y_1, y_2)(t)} \\ u_2'(t) &= \frac{y_1(t)g(t)}{W(y_1, y_2)(t)} \end{aligned} \quad (3.140)$$

where $W(y_1, y_2)$ is the Wronskian of y_1 and y_2 . Note that division by W is permissible since y_1 and y_2 are a fundamental set of solutions, and therefore their Wronskian is nonzero. By integrating Eqs (3.140), we find the desired functions $u_1(t)$ and $u_2(t)$, namely

$$\begin{aligned} u_1'(t) &= - \int \frac{y_2(t)g(t)}{W(y_1, y_2)(t)} dt + c_1 \\ u_2'(t) &= \int \frac{y_1(t)g(t)}{W(y_1, y_2)(t)} dt + c_2 \end{aligned} \quad (3.141)$$

If the integrals in Eqs. (3.141) can be evaluated in terms of elementary functions, then we substitute the results in Eq. (3.133), thereby obtaining the general solution of Eq. (3.130).

Theorem 3.8: Variation of Parameters

If the functions p, q , and g are continuous on an open interval I , and if the functions y_1 and y_2 are a fundamental set of solutions of the homogeneous equation (3.132) corresponding to the nonhomogeneous equation (3.130)

$$y'' + p(t)y' + q(t)y = g(t)$$

then a particular solution of Eq. (3.130) is

$$Y(t) = -y_1(t) \int_{t_0}^t \frac{y_2(s)g(s)}{W(y_1, y_2)(s)} ds + y_2(t) \int_{t_0}^t \frac{y_1(s)g(s)}{W(y_1, y_2)(s)} ds \quad (3.142)$$

where t_0 is any conveniently chosen point in I . The general solution is

$$y = c_1 y_1(t) + c_2 y_2(t) + Y(t) \quad (3.143)$$

[SI*] Mechanical and Electrical Vibrations

The motion of a mass on a vibrating spring, the torsional oscillations of a shaft with a flywheel, the flow of electric current in a simple series circuit, and many other physical problems are all described by the solution of an initial value problem of the form

$$ay'' + by' + cy = gt, \quad y(0) = y_0, \quad y'(0) = y'_0 \quad (3.144)$$

This illustrates a fundamental relationship between mathematics and physics; many physical problems may have the same mathematical model. Thus, once we know how to solve the initial value problem (3.144), it is only necessary to make appropriate interpretations of the constants a, b , and c , and of the functions y and g , to obtain solutions of different physical problems.

The Motion of a Mass on a Spring

Consider a mass m hanging on the end of a vertical spring of original length l . The mass causes an elongation L of the spring in the downward (positive) direction. There are two forces acting at the point where the mass is attached to the spring; the gravitational force, or weight of the mass, acts downward and has magnitude mg , where g is the acceleration due to gravity. There is also a force F , due to the spring, that acts upward. If we assume that the elongation L of the spring is small, the spring force is very nearly proportional to L ; this is known as Hooke's law. Thus we write $F_s = -kL$, where the constant of proportionality k is called the spring constant, and the minus sign is due to the fact that the spring force acts in

the upward (negative) direction. Since the mass is in equilibrium, the two forces balance each other, which means that

$$mg - kL = 0 \quad (3.145)$$

For a given weight $w = mg$, one can measure L and then use Eq. (3.145) to determine k . Note that k has the units of force/length.

In the corresponding dynamic problem, we are interested in studying the motion of the mass when it is acted on by an external force or is initially displaced. Let $u(t)$, measured positive downward, denote the displacement of the mass from its equilibrium position at time t . Then $u(t)$ is related to the forces acting on the mass through Newton's law of motion

$$mu''(t) = f(t) \quad (3.146)$$

where u'' is the acceleration of the mass and f is the net force acting on the mass. Observe that both u and f are functions of time. In determining f there are four separate forces that must be considered:

1. The weight $w = mg$ of the mass that always acts downward.
2. The spring force F_s , is assumed to be proportional to the total elongation $L + u$ of the spring and always acts to restore the spring to its natural position. If $L + u > 0$, then the spring is extended, and the spring force is directed upward. In this case

$$F_s = -k(L + u) \quad (3.147)$$

On the other hand, if $L + u < 0$, then the spring is compressed a distance $|L + u|$, and the spring force, which is now directed downward, is given by $F_s = k|L + u|$. However, when $L + u < 0$, it follows that $|L + u| = -(L + u)$, so F_s is again given by Eq. (3.147). Thus, regardless of the position of the mass, the force exerted by the spring is always expressed by Eq. (3.147).

3. The damping or resistive force F_d always acts in the direction opposite to the direction of motion of the mass. This force may arise from several sources: resistance from the air or other medium in which the mass moves, internal energy dissipation due to the extension or compression of the spring, friction between the mass and the guides (if any) that constrain its motion to one dimension, or a mechanical device (dashpot) that imparts a resistive force to the mass. In any case, we assume that the resistive force is proportional to the speed $\left| \frac{du}{dt} \right|$ of the mass; this is usually referred to as viscous damping. If $\frac{du}{dt} > 0$, then u is increasing, so the mass is moving downward. Then F_d is directed upward and is given by

$$F_d(t) = -\gamma u'(t) \quad (3.148)$$

where γ is a positive constant of proportionality known as the damping constant. On the other hand, if $\frac{du}{dt} < 0$, then u is decreasing, the mass is moving upward, and F_d is directed downward. In this case, $F_d = \gamma|u'(t)|$; since $|u'(t)| = -u'(t)$, it follows that $F_d(t)$ is again given by Eq. (3.148). Thus, regardless of the direction of motion of the mass, the damping force is always expressed by Eq. (3.148).

The damping force may be rather complicated, and the assumption that it is modeled adequately by Eq. (3.148) may be open to question. Some dashpots do behave as Eq. (3.148) states, and if the other sources of dissipation are small, it may be possible to

neglect them altogether or to adjust the damping constant γ to approximate them. An important benefit of the assumption (3.148) is that it leads to a linear (rather than a nonlinear) differential equation.

4. An applied external force $F(t)$ is directed downward or upward as $F(t)$ is positive or negative. This could be a force due to the motion of the mount to which the spring is attached, or it could be a force applied directly to the mass. Often the external force is periodic.

Taking account of these forces, we can now rewrite Newton's law (3.146) as

$$\begin{aligned} mu''(t) &= mg + F_s(t) + F_d(t) + F(t) \\ &= mg - k[L + u(t)] - \gamma u'(t) + F(t) \end{aligned} \quad (3.149)$$

Since $mg - kL = 0$ by Eq. (3.145), it follows that the equation of motion of the mass is

$$mu''(t) + \gamma u'(t) + ku(t) = F(t) \quad (3.150)$$

where the constants m , γ , and k are positive. Note that Eq. (3.150) has the same form as Eq. (3.144).

It is important to understand that Eq. (3.150) is only an approximate equation for the displacement $u(t)$. In particular, both Eqs. (3.147) and (3.148) should be viewed as approximations for the spring force and the damping force, respectively. In our derivation we have also neglected the mass of the spring in comparison with the mass of the attached body.

The complete formulation of the vibration problem requires that we specify two initial conditions, namely, the initial position u_0 and the initial velocity v_0 of the mass:

$$u(0) = u_0, \quad u'(0) = v_0 \quad (3.151)$$

These conditions give a mathematical problem that has a unique solution. This is consistent with our physical intuition that if the mass is set in motion with a given initial displacement and velocity, then its position will be determined uniquely at all future times.

[SI*] **Example:**

A mass weighing 4 lb stretches a spring 2 in. Suppose that the mass is displaced an additional 6 in. in the positive direction and then released. The mass is in a medium that exerts a viscous resistance of 6 lb when the mass has a velocity of 3 ft/s. Formulate the initial value problem that governs the motion of the mass.

Solution:

The required initial value problem consists of the differential equation (3.150) and initial conditions (3.151), so our task is to determine the various constants that appear in these equations. The first step is to choose the units of measurement. Based on the statement of the problem, it is natural to use English rather than metric system of units. The only time unit mentioned is the second, so we will measure t in seconds. On the other hand, both the foot and the inch appear in the statement as units of length. It is immaterial which one we use, but having made a choice, we must be consistent. To be definite, let us measure the displacement u in feet.

Since nothing is said in the statement of the problem about an external force, we assume that $F(t) = 0$. To determine m , note that

$$m = \frac{w}{g} = \frac{4 \text{ lb}}{32 \text{ ft/s}^2} = \frac{1 \text{ lb} \cdot \text{s}^2}{8 \text{ ft}}$$

II. Second Order Linear Equations

The damping coefficient γ is determined from the statement that $\gamma u'$ is equal to 6 lb when u' is 3 ft/s. Therefore

$$\gamma = -\frac{F_d(t)}{u'(t)} = \frac{6 \text{ lb}}{3 \text{ ft/s}} = 2 \frac{\text{lb} \cdot \text{s}}{\text{ft}}$$

The spring constant k is found from the statement that the mass stretches the spring by 2 in., or $\frac{1}{6}$ ft. Thus

$$k = \frac{w}{L} = \frac{4 \text{ lb}}{\frac{1}{6} \text{ ft}} = 24 \frac{\text{lb}}{\text{ft}}$$

Consequently, Eq. (3.150) becomes

$$\frac{1}{8}u'' + 2u' + 24u = 0$$

or

$$u'' + 16u' + 192u = 0 \quad (3.152)$$

The initial conditions are

$$u(0) = \frac{1}{2}, \quad u'(0) = 0 \quad (3.153)$$

We obtain $\frac{1}{2}$ from the initial displacement of the mass that is about 6 in (6 in. = $\frac{1}{2}$ ft). The second initial condition is implied by the word "release" in the statement of the problem, which we interpret to mean that the mass is set in motion with no initial velocity.




Figure 3.18: The simulation of a spring-mass system with Hamzstlab Physics (modifying Box2D source code), if we are assuming the on screen simulation have metric in inch, then the simulation for this example won't oscillate at all after we displaced the mass downward of 6 inch, it will return after the first displacement, the damping coefficient γ is too big / strong thus it does not oscillate. (**Hamzstlab-Physics2D/tests/ode_verticallspringvibrations.cpp**).

Undamped Free Vibrations

If there is no external force, then $F(t) = 0$ in Eq. (3.150). Let us also suppose that there is no damping, so that $\gamma = 0$; this is an idealized configuration of the system, seldom (if ever) completely attainable in practice. However, if the actual damping is very small, then the assumption of no damping may yield satisfactory results over short to moderate time intervals. In this case the equation of motion (3.150) reduces to

$$mu'' + ku = 0 \quad (3.154)$$

The general solution of Eq. (3.154) is

$$u = A \cos \omega_0 t + B \sin \omega_0 t \quad (3.155)$$

where

$$\omega_0^2 = \frac{k}{m} \quad (3.156)$$

The arbitrary constants A and B can be determined if initial conditions of the form (3.151) are given.

In discussing the solution of Eq. (3.154), it is convenient to rewrite Eq. (3.155) in the form

$$u = R \cos(\omega_0 t - \delta) \quad (3.157)$$

or

$$u = R \cos \delta \cos \omega_0 t + R \sin \delta \sin \omega_0 t \quad (3.158)$$

By comparing Eq. (3.158) with Eq. (3.155), we find that A, B, R , and δ are related by the equations

$$A = R \cos \delta, \quad B = R \sin \delta \quad (3.159)$$

Thus

$$R = \sqrt{A^2 + B^2}, \quad \tan \delta = \frac{B}{A} \quad (3.160)$$

In calculating δ , we must take care to choose the correct quadrant; this can be done by checking the signs of $\cos \delta$ and $\sin \delta$ in Eqs. (3.159).




Figure 3.19: The graph of Eq. (3.157), or the equivalent Eq. (3.155), for a typical set of initial conditions is shown. The graph is a displaced cosine wave that describes a periodic or simple harmonic, motion of the mass. (*Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Undamped Free Vibrations/main.cpp*).

The period of the motion is

$$T = \frac{2\pi}{\omega_0} = 2\pi \left(\frac{m}{k} \right)^{1/2} \quad (3.161)$$

The circular frequency $\omega_0 = \sqrt{\frac{k}{m}}$, measured in radians per unit time, is called the natural frequency of the vibration. The maximum displacement R of the mass from equilibrium is the amplitude of the motion. The dimensionless parameter δ is called the phase, or phase angle, and measures the displacement of the wave from its normal position corresponding to $\delta = 0$.

The graph $u(t) = R \cos(\omega_0 t - \delta)$ reaches its highest / amplitude R at $\delta, \delta + n\pi$ with $n = 2, 4, 6, 8, \dots$, and reaches the lowest / $-R$ at $\delta + m\pi$ with $m = 1, 3, 5, 7, \dots$.

Note that the motion described by Eq. (3.157) has a constant amplitude that does not diminish with time. This reflects the fact that, in the absence of damping, there is no way for the system to dissipate the energy imparted to it by the initial displacement and velocity. Further, for a given mass m and spring constant k , the system always vibrates at the same frequency ω_0 , regardless of the initial conditions. However, the initial conditions do help to determine the amplitude of the motion. Finally, observe from Eq. (3.161) that T increases as m increases, so larger masses vibrate more slowly. On the other hand, T decreases as k increases, which means that stiffer springs cause the system to vibrate more rapidly.

[SI*] Example:

Suppose that a mass weighing 10 lb stretches a spring 2 in. If the mass is displaced an additional 2 in. and is then set in motion with an initial upward velocity of 1 ft/s, determine the position of the mass at any later time. Also determine the period, amplitude, and phase of the motion.

Solution:

The spring constant is $k = 10 \text{ lb} / 2 \text{ in.} = 60 \text{ lb/ft}$, and the mass is $m = \frac{w}{g} = \frac{10}{32} \text{ lb}\cdot\text{s}^2/\text{ft}$. Hence the equation of the motion reduces to

$$u'' + 192u = 0 \quad (3.162)$$

and the general solution is

$$u = A \cos(8\sqrt{3}t) + B \sin(8\sqrt{3}t)$$

The solution satisfying the initial conditions $u(0) = \frac{1}{6} \text{ ft}$ and $u'(0) = -1 \text{ ft/s}$ is

$$u = \frac{1}{6} \cos(8\sqrt{3}) - \frac{1}{8\sqrt{3}} \sin(8\sqrt{3}) \quad (3.163)$$

The natural frequency is $\omega_0 = \sqrt{192} \approx 13.856 \text{ rad/s}$, so the period is $T = \frac{2\pi}{\omega_0} \approx 0.45345 \text{ s}$. The amplitude R and phase δ are found from Eqs. (3.160). We have

$$\begin{aligned} R^2 &= \frac{1}{36} + \frac{1}{192} \\ &= \frac{19}{576} \\ &\approx 0.0329 \text{ ft}^2 \end{aligned}$$

The seconds of Eqs. (3.160) yields $\tan \delta = -\frac{\sqrt{3}}{4}$. There are two solutions of this equation, one in the second quadrant and one in the fourth. In the present problem $\cos \delta > 0$ and $\sin \delta < 0$, so δ is in the fourth quadrant, namely,

$$\delta = -\arctan\left(\frac{\sqrt{3}}{4}\right) \approx -0.40864 \text{ rad}$$

Damped Free Vibrations

If we include the effect of damping, the differential equation governing the motion of the mass is

$$mu'' + \gamma u' + ku = 0 \quad (3.164)$$

We are especially interested in examining the effect of variations in the damping coefficient γ for given values of the mass m and spring constant k . The roots of the corresponding characteristic equation are

$$r_1, r_2 = \frac{-\gamma \pm \sqrt{\gamma^2 - 4km}}{2m} \quad (3.165)$$

Depending on the sign of $\gamma^2 - 4km$, the solution u has one of the following forms:

$$\gamma^2 - 4km > 0, \quad u(t) = Ae^{r_1 t} + Be^{r_2 t} \quad (3.166)$$

with

$$B = \frac{u'(0) - r_1 u(0)}{r_2 - r_1}$$

$$A = u(0) - B$$

the next form is

$$\gamma^2 - 4km = 0, \quad u(t) = (A + Bt)e^{-\frac{\gamma t}{2m}} \quad (3.167)$$

with

$$A = u(0)$$

$$B = u'(0) + \frac{\gamma}{2m} A$$

and the last form

$$\gamma^2 - 4km < 0, \quad u(t) = e^{-\frac{\gamma t}{2m}} (A \cos \mu t + B \sin \mu t), \quad \mu = \frac{\sqrt{4km - \gamma^2}}{2m} > 0 \quad (3.168)$$

with

$$A = u(0)$$

$$B = \frac{u'(0) + \frac{\gamma}{2m} A}{\mu}$$

Since m, γ , and k are positive, $\gamma^2 - 4km$ is always less than γ^2 . Hence, if $\gamma^2 - 4km \geq 0$, then the values of r_1 and r_2 given by Eq. (3.165) are negative. If $\gamma^2 - 4km < 0$, then the values of r_1 and r_2 are complex, but with negative real part. Thus, in all cases, the solution u tends to zero as $t \rightarrow \infty$; this occurs regardless of the values of the arbitrary constants A and B , that is, regardless of the initial conditions. This confirms our intuitive expectation, namely, that

damping gradually dissipates the energy initially imparted to the system, and consequently the motion dies out with increasing time.

The most important case is the third one, which occurs when the damping is small. If we let $A = R \cos \delta$ and $B = R \sin \delta$ in Eq. (3.168), then we obtain

$$u(t) = Re^{-\frac{\gamma t}{2m}} \cos(\mu t - \delta) \quad (3.169)$$

The displacement u lies between the curves $u = \pm Re^{-\frac{\gamma t}{2m}}$; hence it resembles a cosine wave whose amplitude decreases as t increases. The motion is called a damped oscillation or a damped vibration. The amplitude factor R depends on m, γ, k , and the initial conditions.

Although the motion is not periodic, the parameter μ determines the frequency with which the mass oscillates back and forth; consequently, μ is called the quasi frequency. By comparing μ with the frequency ω_0 of undamped motion, we find that

$$\frac{\mu}{\omega_0} = \frac{\frac{\sqrt{4km - \gamma^2}}{2m}}{\sqrt{k/m}} = \sqrt{1 - \frac{\gamma^2}{4km}} \approx 1 - \frac{\gamma^2}{8km} \quad (3.170)$$

The last approximation is valid when $\frac{\gamma^2}{4km}$ is small; we refer to this situation as "small damping." Thus the effect of small damping is to reduce slightly the frequency of the oscillation. By analogy with Eq. (3.161), the quantity $T_d = \frac{2\pi}{\mu}$ is called the quasi period. It is the time between successive maxima or successive minima of the position of the mass, or between successive passages of the mass through its equilibrium position while going in the same direction. The relation between T_d and T is given by

$$\frac{T_d}{T} = \frac{\omega_0}{\mu} = \left(1 - \frac{\gamma^2}{4km}\right)^{-1/2} \approx 1 + \frac{\gamma^2}{8km} \quad (3.171)$$

where again the last approximation is valid when $\frac{\gamma^2}{4km}$ is small. thus small damping increases the quasi period.

Equations (3.170) and (3.171) reinforce the significance of the dimensionless ratio $\frac{\gamma^2}{4km}$. It is not the magnitude of γ alone that determines whether damping is large or small, but the magnitude of γ^2 compared to $4km$. When $\frac{\gamma^2}{4km}$ is small, then the damping has a small effect on the quasi frequency and quasi period of the motion. On the other hand, if we want to study the detailed motion of the mass for all time, then we can never neglect the damping force, no matter how small.

As $\frac{\gamma^2}{4km}$ increases, the quasi frequency μ decreases and the quasi period T_d increases. In fact, $\mu \rightarrow 0$ and $T_d \rightarrow \infty$ as $\gamma \rightarrow 2\sqrt{km}$. As indicated by Eqs. (3.166), (3.167), and (3.168), the nature of the solution changes as γ passes through the value $2\sqrt{km}$. This value is known as critical damping, while for larger values of γ the motion is said to be overdamped. In these cases, given by Eqs. (3.167) and (3.166), respectively, the mass creeps back to its equilibrium position but does not oscillate about it, as for small γ .

[SI*] Example:

The motion of a certain spring-mass system is governed by the differential equation

$$u'' + 0.125u' + u = 0 \quad (3.172)$$

where u is measured in feet and t in seconds. If $u(0) = 2$ and $u'(0) = 0$, determine the position of the mass at any time. Find the quasi frequency and the quasi period, as well as the time at which the mass first passes through its equilibrium position. Also find the time τ such that $|u(t)| < 0.1$ for all $t > \tau$.

Solution:

The solution of Eq. (3.172) is

$$u(t) = e^{-t/16} \left[A \cos \left(\frac{\sqrt{255}}{16} t \right) + B \sin \left(\frac{\sqrt{255}}{16} t \right) \right]$$

To satisfy the initial conditions, we must choose $A = 2$ and $B = \frac{2}{\sqrt{255}}$; hence the solution of the initial value problem is

$$\begin{aligned} u(t) &= e^{-t/16} \left[2 \cos \left(\frac{\sqrt{255}}{16} t \right) + \frac{2}{\sqrt{255}} \sin \left(\frac{\sqrt{255}}{16} t \right) \right] \\ &= \frac{32}{\sqrt{255}} e^{-t/16} \cos \left(\frac{\sqrt{255}}{16} t - \delta \right) \end{aligned} \tag{3.173}$$

where $\tan \delta = \frac{1}{\sqrt{255}}$, so $\delta \cong 0.06254$.

II. Second Order Linear Equations




Figure 3.20: The figure shows the displacement of the mass as a function of time. For purposes of comparison, we also show the motion if the damping term is neglected. (*Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Vibration with Small Damping and With No Damping Second Order ODE/main.cpp*).

The quasi frequency is $\mu = \frac{\sqrt{255}}{16} \approx 0.998$ and the quasi period is

$$T_d = \frac{2\pi}{\mu} \approx 6.295 \text{ s}$$

These values differ only slightly from the corresponding values (1 and 2π , respectively) for the undamped oscillation.

The damping coefficient is small in this example, only one-sixteenth of the critical value, in fact. Nevertheless, the amplitude of the oscillation is reduced rather rapidly.

To find the time at which the mass first passes through its equilibrium position, we refer to Eq. (3.173) and set $\frac{\sqrt{255}}{16} - \delta$ to $\frac{\pi}{2}$, the smallest positive zero of the cosine function. Then, by solving for t , we obtain

$$t = \frac{16}{\sqrt{255}} \left(\frac{\pi}{2} + \delta \right) \approx 1.637 \text{ s}$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Spring Mass System Second Order Linear Differential Equations ]# ./main
Spring mass system solution for  $y'' + 0.125y' + y = 0$  with  $u(0) = 2$ ,  $u'(0) = 0$ 

The real-valued initial value problem solution is:
 $u(t) = 2 * e^{(-0.0625 * t)} * \cos(0.998045 * t) + 0.125245 * e^{(-0.0625 * t)} * \sin(0.998045 * t)$ 
 $\delta = 0.0625408$ 
 $T_d = 6.29549$ 
The time when the mass passes through its equilibrium position:
 $t = 1.63654$ 
Spring mass system solution for  $y'' + y = 0$  with  $u(0) = 2$ ,  $u'(0) = 0$ 

The real-valued initial value problem solution is:
 $u(t) = 2 * \cos(t)$ 
 $\delta = 0$ 
 $T_d = 6.28319$ 
The time when the mass passes through its equilibrium position:
 $t = 1.5708$ 

```

Figure 3.21: The computation of initial value problem solution for the second order linear differential equation with small damping and no damping (*SymIntegration/Examples/Differential Equations/Test SymIntegration Spring Mass System Second Order Linear Differential Equations/main.cpp*).

Electric Circuits

A second example of the occurrence of second order linear differential equations with constant coefficients is their use as a model of the flow of electric current in the simple series circuit. The current I , measured in amperes (A), is a function of time t . The resistance R in ohms (Ω), the capacitance C in farads (F), and the inductance L in Henrys (H) are all positive and are assumed to be known constants. the impressed voltage E in volts (V) is a given function of time. Another physical quantity that enters the discussion is the total charge Q in coulombs (C) on the capacitor at time t . The relation between charge Q and current I is

$$I = \frac{dQ}{dt} \quad (3.174)$$




Figure 3.22: A simple RLC series electric circuit.

The flow of current in the circuit is governed by Kirchhoff's second law: In a closed circuit the impressed voltage is equal to the sum of the voltage drops in the rest of the circuit.

According to the elementary laws of electricity, we know that
The voltage drop across the resistor is IR .

The voltage drop across the capacitor is $\frac{Q}{C}$.
 The voltage drop across the inductor is $L\frac{dI}{dt}$.

Hence, by Kirchhoff's law,

$$L\frac{dI}{dt} + RI + \frac{1}{C}Q = E(t) \quad (3.175)$$

The units have been chosen so that 1 volt = 1 ohm · 1 ampere = 1 coulomb/1 farad = 1 henry · 1 ampere/1 second.

Substituting for I from Eq. (3.174), we obtain the differential equation

$$LQ'' + RQ' + \frac{1}{C}Q = E(t) \quad (3.176)$$

for the charge Q . The initial conditions are

$$Q(t_0) = Q_0, \quad Q'(t_0) = I(t_0) = I_0 \quad (3.177)$$

Thus we must know the charge on the capacitor and the current in the circuit at some initial time t_0 .

Alternatively, we can obtain a differential equation for the current I by differentiating Eq. (3.176) with respect to t , and then substituting for $\frac{dQ}{dt}$ from Eq. (3.174). The result is

$$LI'' + RI' + \frac{1}{C}I = E'(t) \quad (3.178)$$

with the initial conditions

$$I(t_0) = I_0, \quad I'(t_0) = I'_0 \quad (3.179)$$

From Eq. (3.175) it follows that

$$I'_0 = \frac{E(t_0) - RI_0 - \left(\frac{1}{C}\right)Q_0}{L} \quad (3.180)$$

Hence I'_0 is also determined by the initial charge and current, which are physically measurable quantities.

The most important conclusion from this discussion is that the flow of current in the circuit is described by an initial value problem of precisely the same form as the one that describes the motion of a spring-mass system. This is a good example of the unifying role of mathematics: Once you know how to solve second order linear differential equations with constant coefficients, you can interpret the results in terms of mechanical vibrations, electric circuits, or any other physical situation that leads to the same problem.

[SI*] **Example:**

A series circuit has a capacitor of 10^{-5} F, a resistor of $3 \times 10^2 \Omega$, and an inductor of 0.2 H. The initial charge on the capacitor is 10^{-6} C and there is no initial current. Find the charge Q on the capacitor at any time t .

Solution:

Assume that the circuit is closed at $t = 0$. Apply Faraday's law to obtain the governing ODE for the current.

$$\sum V = -L\frac{di}{dt}$$

The only potential drops occur over the resistor and the capacitor.

$$iR + \frac{q}{C} = -L \frac{di}{dt}$$

Write $i = \frac{dq}{dt} = q'$.

$$Rq' + \frac{q}{C} = -Lq''$$

Bring Lq'' to the left side and divide both sides by L .

$$q'' + \frac{R}{L}q' + \frac{1}{LC}q = 0 \quad (3.181)$$

Since the coefficients are constant and this is a homogeneous ODE, the solutions are of the form $q = e^{rt}$.

$$q = e^{rt} \rightarrow q' = re^{rt} \rightarrow q'' = r^2e^{rt}$$

Substitute these expressions to obtain an algebraic equation for r .

$$r^2e^{rt} + \frac{R}{L}(re^{rt}) + \frac{1}{LC}(e^{rt}) = 0$$

Divide both sides by e^{rt} .

$$\begin{aligned} r^2 + \frac{R}{L}(re^{rt}) + \frac{1}{LC}(e^{rt}) &= 0 \\ r &= \frac{-\frac{R}{L} \pm \sqrt{\frac{R^2}{L^2} - \frac{4}{LC}}}{2} \\ &= \frac{-1500 \pm \sqrt{250000}}{2} \\ &= \frac{-1500 \pm 500}{2} \\ &= -750 \pm 250 \end{aligned}$$

so $r = \{-1000, 500\}$. Two solutions to equation (3.181) are then

$$\begin{aligned} q(t) &= e^{-1000t} \\ q(t) &= e^{-500t} \end{aligned}$$

By the principle of superposition, the general solution is a linear combination of these two.

$$q(t) = C_1e^{-1000t} + C_2e^{-500t}$$

Take a derivative of it with respect to t .

$$q'(t) = -1000C_1e^{-1000t} - 500C_2e^{-500t}$$

Now apply the initial conditions, $q(0) = 10^{-6}$ and $q'(0) = 0$, to determine C_1 and C_2 .

$$\begin{aligned} q(0) &= C_1 + C_2 = 10^{-6} \\ q'(0) &= -1000C_1 - 500C_2 = 0 \end{aligned}$$

II. Second Order Linear Equations

Solving this system of equations yields $C_1 = -10^{-6}$ and $C_2 = 2 \times 10^{-6}$. Therefore,

$$q(t) = -10^{-6}e^{-1000t} + 2 \times 10^{-6}e^{-500t}$$

Note that the charge $q(t)$ is in Coulombs (C).

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration RLC Series Electric Circuit Second Order Linear Differential Equations ]# ./main
RLC series electric circuit solution for R = 0, L = 1 H, C = 0.25 * 10^{-6} F with q(0) = 10^{-6}, q'(0) = 0

The solution for the initial value problem / the charge Q at any time t is:
q (t) = 1e-06*cos(2000*t)

RLC series electric circuit solution for R = 3* 10^{2} Ohm, L = 0.2 H, C = 10^{-5} F with q(0) = 10^{-6}, q'(0) = 0

The general solution is:
c1*e^{(-500*t)}+c2*e^{(-1000*t)}

The solution for the initial value problem / the charge Q at any time t is:
q(t) = 2e-06*e^{(-500*t)}-1e-06*e^{(-1000*t)}

Time taken by function: 43347 microseconds
```

Figure 3.23: The computation of initial value problem solution for the second order linear differential equation to compute the charge Q on the capacitor at any time t in a RLC series circuit (**SymIntegration/Examples/Differential Equations/Test SymIntegration RLC Series Electric Circuit Second Order Linear Differential Equations/main.cpp**).

[SI*] Example:

If a series circuit has a capacitor of $C = 0.8 \times 10^{-6}$ F and an inductor of $L = 0.2$ H, find the resistance R so that the circuit is critically damped.

Solution:

Assume the circuit is closed at $t = 0$. Apply Faraday's law to obtain the governing equation for the current.

$$\sum V = -L \frac{di}{dt}$$

The only potential drops occur over the resistor and the capacitor

$$iR + \frac{q}{C} = -L \frac{di}{dt}$$

Write $i = \frac{dq}{dt} = q'$.

$$\begin{aligned} Rq' + \frac{q}{C} &= -Lq'' \\ Lq'' + Rq' + \frac{q}{C} &= 0 \end{aligned}$$

Critical damping occurs when the ratio of R^2 to $4\frac{L}{C}$ is 1.

$$\begin{aligned}\frac{R^2}{\frac{4L}{C}} &= 1 \\ R^2 &= \frac{4L}{C} \\ R &= \sqrt{\frac{4L}{C}} \\ &= \sqrt{\frac{4(0.2)}{0.8 \times 10^{-6}}} \\ &= 1000 \Omega\end{aligned}$$

[SI*] **Forced Vibrations**

We will now investigate the situation in which a periodic external force is applied to a spring-mass system. The behavior of this simple system models that of many oscillatory systems with an external force due, for example, to a motor attached to the system. We will first consider the case in which damping is present and will look later at the idealized special case in which there is assumed to be no damping.

[SI*] **Example for Forced Vibrations with Damping:**

Suppose that the motion of a certain spring-mass system satisfies the differential equation

$$u'' + u' + 1.25u = 3 \cos t \quad (3.182)$$

and the initial conditions

$$u(0) = 2, \quad u'(0) = 3 \quad (3.183)$$

Find the solution of this initial value problem and describe the behavior of the solution for large t .

Solution:

The homogeneous equation corresponding to Eq. (3.182) has the characteristic equation

$$r^2 + r + 1.25 = 0$$

with roots $r = -0.5 \pm i$. Thus the general solution $u_c(t)$ of this homogeneous equation is

$$u_c(t) = c_1 e^{-t/2} \cos t + c_2 e^{-t/2} \sin t \quad (3.184)$$

A particular solution of Eq. (3.182) has the form $U(t) = A \cos t + B \sin t$, where A and B are found by substituting $U(t)$ for u in Eq. (3.182). We have $U'(t) = -A \sin t + B \cos t$ and $U''(t) = -A \cos t - B \sin t$. Thus, from Eq. (3.182) we obtain

$$(0.25A + B) \cos t + (-A + 0.25B) \sin t = 3 \cos t$$

Consequently, A and B must satisfy the equations

$$\begin{aligned}0.25A + B &= 3 \\ -A + 0.25B &= 0\end{aligned}$$

with the result that $A = \frac{12}{17}$ and $B = \frac{48}{17}$. Therefore the particular solution is

$$U(t) = \frac{12}{17} \cos t + \frac{48}{17} \sin t \quad (3.185)$$

and the general solution of Eq. (3.182) is

$$u = u_c(t) + U(t) = c_1 e^{-t/2} \cos t + c_2 e^{-t/2} \sin t + \frac{12}{17} \cos t + \frac{48}{17} \sin t \quad (3.186)$$

The remaining constants c_1 and c_2 are determined by the initial conditions (3.183). From Eq. (3.186) we have

$$\begin{aligned} u(0) &= c_1 + \frac{12}{17} = 2 \\ u'(0) &= -\frac{1}{2}c_1 + c_2 + \frac{48}{17} = 3 \end{aligned}$$

so $c_1 = \frac{22}{17}$ and $c_2 = \frac{14}{17}$. Thus we finally obtain the solution of the given initial value problem, namely,

$$\frac{22}{17} e^{-t/2} \cos t + \frac{14}{17} e^{-t/2} \sin t + \frac{12}{17} \cos t + \frac{48}{17} \sin t \quad (3.187)$$




Figure 3.24: The graph of the solution (3.187) for the initial value problem for the second order nonhomogeneous differential equation (*Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Forced Vibrations IVP Second Order ODE/main.cpp*).

It is important to note that the solution consists of two distinct parts. The first two terms on the right side of Eq. (3.187) contain the exponential factor $e^{-t/2}$; as a result they rapidly approach zero. It is customary to call these terms transient. The remaining terms in Eq. (3.187) involve only sines and cosines, and therefore represent an oscillation that continues indefinitely. We refer to them as a steady state. The green solid and blue dashed curves show the transient and the steady state parts of the solution, respectively. The transient part comes from the solution of the homogeneous equation corresponding to Eq. (3.182) and is

needed to satisfy the initial conditions. The steady state is the particular solution of the full nonhomogeneous equation. After a fairly short time the transient is vanishingly small and the full solution is essentially indistinguishable from the steady state.

[SI*] The equation of motion of a general spring-mass system subject to an external force $F(t)$ is

$$mu''(t) + \gamma u'(t) + ku(t) = F(t) \quad (3.188)$$

where m, γ, k are the mass, damping coefficient, and spring constant of the spring-mass system. Suppose now that the external force is given by $F_0 \cos \omega t$, where F_0 and ω are positive constants representing the amplitude and frequency, respectively, of the force. Then Eq. (3.188) becomes

$$mu'' + \gamma u' + ku = F_0 \cos \omega t \quad (3.189)$$

Solutions of Eq. (3.189) behave very much like the solution in the preceding example. The general solution of Eq. (3.189) must have the form

$$u = c_1 u_1(t) + c_2 u_2(t) + A \cos \omega t + B \sin \omega t = u_c(t) + U(t) \quad (3.190)$$

The first two terms on the right side of Eq. (3.190) are the general solution $u_c(t)$ of the homogeneous equation corresponding to Eq. (3.189), while the latter two terms are a particular solution $U(t)$ of the full nonhomogeneous equation. The coefficients A and B can be found, as usual, by substituting these terms into the differential equation (3.189), while the arbitrary constants c_1 and c_2 are available to satisfy initial conditions, if any are prescribed. The solutions $u_1(t)$ and $u_2(t)$ of the homogeneous equation depend on the roots r_1 and r_2 of the characteristic equation $mr^2 + \gamma r + k = 0$. Since m, γ , and k are all positive, it follows that r_1 and r_2 either are real and negative or are both complex conjugates with negative real part. In either case, both $u_1(t)$ and $u_2(t)$ approach zero as $t \rightarrow \infty$. Since $u_c(t)$ dies out as t increases, it is called the transient solution. In many applications it is of little importance and (depending on the value of γ) may well be undetectable after only a few seconds.

The remaining terms in Eq. (3.190), namely, $U(t) = A \cos \omega t + B \sin \omega t$, do not die out as t increases but persist indefinitely, or as long as the external force is applied. They represent a steady oscillation with the same frequency as the external force and are called the steady state solution or the forced response. The transient solution enables us to satisfy whatever initial conditions may be imposed; with increasing time, the energy put into the system by the initial displacement and velocity is dissipated through the damping force, and the motion then becomes the response of the system to the external force. Without damping, the effect of the initial conditions would persist for all time.

It is convenient to express $U(t)$ as a single trigonometric term rather than as a sum of two terms. Recall that we did this for other similar expressions, thus we write

$$U(t) = R \cos(\omega t - \delta) \quad (3.191)$$

The amplitude R and phase δ depend directly on A and B and indirectly on the parameters in the differential equation (3.189). It is possible to show, by straightforward but somewhat lengthy algebraic computations, that

$$R = \frac{F_0}{\Delta}, \quad \cos \delta = \frac{m(\omega_0^2 - \omega^2)}{\Delta}, \quad \sin \delta = \frac{\gamma\omega}{\Delta} \quad (3.192)$$

where

$$\begin{aligned}\Delta &= \sqrt{m^2(\omega_0^2 - \omega^2)^2 + \gamma^2\omega^2} \\ \omega_0^2 &= \frac{k}{m}\end{aligned}\tag{3.193}$$

Recall that ω_0 is the natural frequency of the unforced system in the absence of damping. We now investigate how the amplitude R of the steady state oscillation depends on the frequency ω of the external force. Substituting from Eq. (3.193) into the expression for R in Eq. (3.192) and executing some algebraic manipulations, we find that

$$\frac{Rk}{F_0} = \frac{1}{\left[\left(1 - \frac{\omega^2}{\omega_0^2} \right)^2 + \Gamma \frac{\omega^2}{\omega_0^2} \right]^{1/2}}\tag{3.194}$$

where $\Gamma = \frac{\gamma^2}{mk}$. Observe that the quantity $\frac{Rk}{F_0}$ is the ratio of the amplitude R of the forced response to $\frac{F_0}{k}$, the static displacement of the spring produced by a force F_0 .

For low frequency excitation, that is, as $\omega \rightarrow 0$, it follows from Eq. (3.194) that $\frac{Rk}{F_0} \rightarrow 1$ or $R \rightarrow \frac{F_0}{k}$. At the other extreme, for very high frequency excitation, Eq. (3.194) implies that $R \rightarrow 0$ as $\omega \rightarrow \infty$. At an intermediate value of ω the amplitude may have a maximum. To find this maximum point, we can differentiate R with respect to ω and set the result equal to zero. In this way we find that the maximum amplitude occurs when $\omega = \omega_{\max}$, where

$$\omega_{\max}^2 = \omega_0^2 - \frac{\gamma^2}{2m^2} = \omega_0^2 \left(1 - \frac{\gamma^2}{2mk} \right)\tag{3.195}$$

Note that $\omega_{\max} < \omega_0$ and that $\omega_{\max}$ is close to ω_0 when γ is small. The maximum value of R is

$$R_{\max} = \frac{F_0}{\gamma\omega_0 \sqrt{1 - \left(\frac{\gamma^2}{4mk} \right)}} \cong \frac{F_0}{\gamma\omega_0} \left(1 + \frac{\gamma^2}{8mk} \right)\tag{3.196}$$

where the last expression is an approximation for small γ . If $\frac{\gamma^2}{mk} > 2$, then $\omega_{\max}$ as given by Eq. (3.195) is imaginary; in this case the maximum value of R occurs for $\omega = 0$, and R is a monotone decreasing function of ω . Recall that critical damping occurs when $\frac{\gamma^2}{mk} = 4$.

For small γ it follows from Eq. (3.196) that $R_{\max} \cong \frac{F_0}{\gamma\omega_0}$. Thus, for lightly damped systems, the amplitude R of the forced response when ω is near ω_0 is quite large even for relatively small external forces, and the smaller the value of γ , the more pronounced is this effect. This phenomenon is known as resonance, and it is often an important design consideration. Resonance can be either good or bad, depending on the circumstances. It must be taken very seriously in the design of structures, such as buildings and bridges, where it can produce instabilities that might lead to the catastrophic failure of the structure. On the other hand, resonance can be put to good use in the design of instruments, such as seismographs, that are intended to detect weak periodic incoming signals.




Figure 3.25: Some representative graphs of $\frac{Rk}{F_0}$ versus $\frac{\omega}{\omega_0}$ for several values of $\Gamma = \frac{\gamma^2}{mk}$ (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Forced Vibrations Amplitude of Steady State Response versus Frequency of Driving Force/main.cpp).

It follows from Eq. (3.194), or from Eqs. (3.192) and (3.193), that $R \rightarrow \frac{F_0}{m|\omega_0^2 - \omega^2|}$ as $\gamma \rightarrow 0$ and hence $\frac{Rk}{F_0}$ is asymptotic to the vertical line $\omega = \omega_0$. As the damping in the system increases, the peak response gradually diminishes.

You can easily verify that each of the quantities $\frac{Rk}{F_0}$, $\frac{\omega}{\omega_0}$, and Γ is dimensionless. The importance of this observation is that the number of significant parameters in the problem has been reduced to three rather than five that appear in Eq. (3.189). Thus only one family of curves, of which a few are shown in Figure (3.24), is needed to describe the response-versus-frequency behavior of all systems governed by Eq. (3.189).

The phase angle δ also depends in an interesting way on ω . For ω near zero, it follows from Eqs. (3.192) and (3.193) that $\cos \delta \approx 1$ and $\sin \delta \approx 0$. Thus $\delta \approx 0$, and the response is enarly in phase with the excitation, meaning that they rise and fall together and, in particular, assume their respective maxima nearly together and their respective minima nearly together. For $\omega = \omega_0$ we find that $\cos \delta = 0$ and $\sin \delta = 1$, so $\delta = \frac{\pi}{2}$. In this acse the response lags behind the excitation by $\frac{\pi}{2}$; that is, the peaks of the response occur $\frac{\pi}{2}$ later than the peaks of the excitation, and similarly for the valleys. Finally, for ω very large, we have $\cos \delta \approx -1$ and $\sin \delta \approx 0$. Thus $\delta \approx \pi$, so that the response is nearly out of phase with the excitation; this means that the response is minimum when the excitation is maximum, and vice versa.




Figure 3.26: Forced vibration with damping: phase of steady state response versus frequency of driving force; $\Gamma = \frac{\gamma^2}{mk}$ (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Forced Vibrations Phase of Steady State Response versus Frequency of Driving Force/main.cpp).

Figure (3.25) shows the graph of δ versus $\frac{\omega}{\omega_0}$ for several values of Γ . For small damping, the phase transition from near $\delta = 0$ to near $\delta = \pi$ occurs rather abruptly, whereas for larger values of the damping parameter, the transition takes place more gradually.

To compute the phase δ in terms of $\frac{\omega}{\omega_0}$ we can use the formula of $\sin \delta$ from Eqs. (3.192), thus

$$\begin{aligned} \sin \delta &= \frac{\gamma \omega}{\Delta} \\ &= \frac{\gamma \omega}{\sqrt{m^2(\omega_0^2 - \omega^2)^2 + \gamma^2 \omega^2}} \\ &= \frac{\gamma \frac{\omega}{\omega_0}}{\frac{1}{\omega_0} \sqrt{m^2(\omega_0^2 - \omega^2)^2 + \gamma^2 \omega^2}} \\ &= \frac{\gamma \frac{\omega}{\omega_0}}{\sqrt{m^2(1 - \frac{\omega^2}{\omega_0^2})^2 + \gamma^2 \frac{\omega^2}{\omega_0^2}}} \\ \delta &= \sin^{-1} \left(\frac{\gamma \frac{\omega}{\omega_0}}{\sqrt{m^2(1 - \frac{\omega^2}{\omega_0^2})^2 + \gamma^2 \frac{\omega^2}{\omega_0^2}}} \right) \end{aligned}$$

To plot with sine formula we use this formula:

$$\delta = \begin{cases} \sin^{-1} \left(\frac{\gamma \frac{\omega}{\omega_0}}{\sqrt{m^2(1-\frac{\omega^2}{\omega_0^2})^2 + \gamma^2 \frac{\omega^2}{\omega_0^2}}} \right), & \text{for } \frac{\omega}{\omega_0} \leq 1 \\ \pi - \sin^{-1} \left(\frac{\gamma \frac{\omega}{\omega_0}}{\sqrt{m^2(1-\frac{\omega^2}{\omega_0^2})^2 + \gamma^2 \frac{\omega^2}{\omega_0^2}}} \right), & \text{for } \frac{\omega}{\omega_0} > 1 \end{cases} \quad (3.197)$$

In reality when we try to plot with this formula, the graphs will be the same if we can adjust the Γ parameter with γ, m, k . Thus based on what has been used, most people are rarely use the arcsine formula to plot the phase of steady state response versus frequency of driving force, in general to plot the phase of steady state response versus frequency of driving force we often use the arctangent formula which is simpler and only require 3 parameters

$$\delta = \begin{cases} \tan^{-1} \left(\frac{2\Gamma \frac{\omega}{\omega_0}}{1 - \left(\frac{\omega}{\omega_0}\right)^2} \right), & \text{for } \frac{\omega}{\omega_0} \leq 1 \\ \pi + \tan^{-1} \left(\frac{2\Gamma \frac{\omega}{\omega_0}}{1 - \left(\frac{\omega}{\omega_0}\right)^2} \right), & \text{for } \frac{\omega}{\omega_0} > 1 \end{cases} \quad (3.198)$$

As $\frac{\omega}{\omega_0} \rightarrow \infty$, $\frac{2\Gamma \frac{\omega}{\omega_0}}{1 - \left(\frac{\omega}{\omega_0}\right)^2} \rightarrow 0^-$ and $\delta \rightarrow \pi$.

$\tan^{-1}(\infty) = \frac{\pi}{2}$, and tangent is positive in quadrant I and III only. You can analyze the value of δ correctly when you computed $R, \cos \delta, \sin \delta$ in Eqs. (3.192). For example, if the value of $\cos \delta$ is negative and the value of $\sin \delta$ is positive, then δ is in quadrant II, where sine is positive, cosine is negative and tangent is negative. Furthermore, if $\tan \delta$ is negative then we will use this formula

$$\delta = \pi + \frac{2\Gamma \frac{\omega}{\omega_0}}{1 - \left(\frac{\omega}{\omega_0}\right)^2}$$

due to

$$\begin{aligned} \tan^{-1}(\alpha) &= \tan^{-1}(\alpha) \\ \tan^{-1}(-\alpha) &= -\tan^{-1}(\alpha) + \pi \end{aligned}$$

In a forced vibration system with damping, the steady-state response (displacement) does not occur simultaneously with the driving force. Instead, the displacement lags behind the driving force by a phase angle (δ), which varies significantly depending on the ratio of the driving frequency (ω) to the system's natural frequency (ω_0).

[SI*] **Example:**

Consider a vibrating system described by the initial value problem

$$u'' + \frac{1}{4}u' + 2u = 2 \cos \omega t, \quad u(0) = 0, \quad u'(0) = 2$$

- Determine the steady state part of the solution of this problem.
- Find the amplitude A of the steady state solution in terms of ω .
- Plot A versus ω
- Find the maximum value of A and the frequency ω for which it occurs.

Solution:

- (a) This is the linear nonhomogeneous ODE, so its general solution can be expressed a sum of the complementary solution and the particular solution.

$$u(t) = u_c(t) + u_p(t)$$

The complementary solution satisfies the associated homogeneous equation.

$$u_c'' + \frac{1}{4}u_c' + 2u_c = 0 \quad (3.199)$$

Since the coefficients are constant and this is a homogeneous ODE, the solutions are of the form $u_c = e^{rt}$.

$$u_c = e^{rt} \rightarrow u_c' = re^{rt} \rightarrow u_c'' = r^2e^{rt}$$

Substitute these expressions to obtain an algebraic equation for r .

$$r^2e^{rt} + \frac{1}{4}(re^{rt}) + 2(e^{rt}) = 0$$

Divide both sides by $\frac{e^{rt}}{4}$.

$$\begin{aligned} 4r^2 + r + 8 &= 0 \\ r &= \frac{-1 \pm \sqrt{1 - 4(4)(8)}}{2(4)} \\ &= \frac{-1 \pm \sqrt{-127}}{8} \\ &= \frac{-1 \pm i\sqrt{127}}{8} \\ &= -\frac{1}{8} \pm i\mu \\ r &= \left\{ -\frac{1}{8} - i\mu, -\frac{1}{8} + i\mu \right\} \end{aligned}$$

Two solutions to equation (3.199) are $u_c = e^{(-\frac{1}{8} - i\mu)t}$ and $u_c = e^{(-\frac{1}{8} + i\mu)t}$.

By the principle of superposition, the general solution for u_c is a linear combination of these two.

$$\begin{aligned} u_c(t) &= C_1e^{(-\frac{1}{8} - i\mu)t} + C_2e^{(-\frac{1}{8} + i\mu)t} \\ &= C_1e^{-t/8}e^{-i\mu t} + C_2e^{-t/8}e^{i\mu t} \\ &= C_1e^{-t/8} [\cos(-\mu t) + i \sin(-\mu t)] + C_2e^{-t/8} [\cos(\mu t) + i \sin(\mu t)] \\ &= C_1e^{-t/8} [\cos(\mu t) - i \sin(\mu t)] + C_2e^{-t/8} [\cos(\mu t) + i \sin(\mu t)] \\ &= C_1e^{-t/8} \cos(\mu t) - iC_1e^{-t/8} \sin(\mu t) + C_2e^{-t/8} \cos(\mu t) + iC_2e^{-t/8} \sin(\mu t) \\ &= (C_1 + C_2)e^{-t/8} \cos(\mu t) + (-iC_1 + iC_2)e^{-t/8} \sin(\mu t) \\ &= C_3e^{-t/8} \cos(\mu t) + C_4e^{-t/8} \sin(\mu t) \\ &= C_3e^{-t/8} \cos \frac{\sqrt{127}}{8}t + C_4e^{-t/8} \sin \frac{\sqrt{127}}{8}t \end{aligned}$$

On the other hand, the particular solution satisfies

$$u_p'' + \frac{1}{4}u_p' + 2u_p = 2 \cos \omega t$$

Since there are both even and odd derivatives on the left side, include both sine and cosine in the trial solution to account for the cosine term on the right side:

$$u_p(t) = B \cos \omega t + D \sin \omega t$$

Plug this into the ODE to determine B and D

$$(B \cos \omega t + D \sin \omega t)'' + \frac{1}{4}(B \cos \omega t + D \sin \omega t)' + 2(B \cos \omega t + D \sin \omega t) = 2 \cos \omega t$$

Evaluate the derivatives

$$\left(-B\omega^2 + \frac{D\omega}{4} + 2B\right) \cos \omega t + \left(-D\omega^2 - \frac{B\omega}{4} + 2D\right) \sin \omega t = 2 \cos \omega t$$

Match the coefficients.

$$\begin{aligned} -B\omega^2 + \frac{D\omega}{4} + 2B &= 2 \\ -D\omega^2 - \frac{B\omega}{4} + 2D &= 0 \end{aligned}$$

Solving this system of equations yields

$$\begin{aligned} B &= \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \\ D &= \frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \end{aligned}$$

Consequently, the particular solution is

$$u_p(t) = \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \cos \omega t + \frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \sin \omega t$$

and the general solution is

$$\begin{aligned} u(t) &= C_3 e^{-t/8} \cos \frac{\sqrt{127}}{8} t + C_4 e^{-t/8} \sin \frac{\sqrt{127}}{8} t \\ &\quad + \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \cos \omega t + \frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \sin \omega t \end{aligned}$$

Differentiate it with respect to t

$$\begin{aligned} u'(t) &= -\frac{C_3}{8} e^{-t/8} \cos \frac{\sqrt{127}}{8} t - C_3 \frac{\sqrt{127}}{8} e^{-t/8} \sin \frac{\sqrt{127}}{8} t - \frac{C_4}{8} e^{-t/8} \sin \frac{\sqrt{127}}{8} t \\ &\quad + C_4 \frac{\sqrt{127}}{8} e^{-t/8} \cos \frac{\sqrt{127}}{8} t \\ &\quad - \omega \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \sin \omega t + \omega \frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \cos \omega t \end{aligned}$$

Now apply the initial conditions to determine C_3 and C_4

$$u(0) = C_3 + \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} = 0$$

$$u'(0) = -\frac{C_3}{8} + C_4 \frac{\sqrt{127}}{8} e^{-t/8} + \omega \frac{8\omega}{16\omega^4 - 63\omega^2 + 64} = 2$$

Solving this system of equations yields

$$C_3 = \frac{32(\omega^2 - 2)}{16\omega^4 - 63\omega^2 + 64}$$

$$C_4 = \frac{16(16\omega^4 - 65\omega^2 + 60)}{\sqrt{127}(16\omega^4 - 63\omega^2 + 64)}$$

Therefore

$$u(t) = \frac{32(\omega^2 - 2)}{16\omega^4 - 63\omega^2 + 64} e^{-t/8} \cos \frac{\sqrt{127}}{8} t + \frac{16(16\omega^4 - 65\omega^2 + 60)}{\sqrt{127}(16\omega^4 - 63\omega^2 + 64)} e^{-t/8} \sin \frac{\sqrt{127}}{8} t$$

$$+ \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \cos \omega t + \frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \sin \omega t$$

This is the steady state part:

$$u(t)_{\text{steady state}} = \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \cos \omega t + \frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \sin \omega t$$

- (b) We introduce an amplitude A and a phase δ to combine the two sinusoidal terms of the steady state into one

$$u(t)_{\text{steady state}} = \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \cos \omega t + \frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \sin \omega t$$

$$= A \cos \delta \cos \omega t + A \sin \delta \sin \omega t$$

$$= A \cos(\omega t - \delta)$$

A and δ satisfy the following system of equations

$$A \cos \delta = \frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64}$$

$$A \sin \delta = \frac{8\omega}{16\omega^4 - 63\omega^2 + 64}$$

To find A , square both sides of each equation

$$A^2 \cos^2 \delta = \left[\frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \right]^2$$

$$A^2 \sin^2 \delta = \left[\frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \right]^2$$

and then add the respective sides together

$$\begin{aligned}
 A^2 \cos^2 \delta + A^2 \sin^2 \delta &= \left[\frac{32(2 - \omega^2)}{16\omega^4 - 63\omega^2 + 64} \right]^2 + \left[\frac{8\omega}{16\omega^4 - 63\omega^2 + 64} \right]^2 \\
 A^2(\cos^2 \delta + \sin^2 \delta) &= \frac{64}{16\omega^4 - 63\omega^2 + 64} \\
 A^2 &= \frac{64}{16\omega^4 - 63\omega^2 + 64} \\
 A &= \frac{8}{\sqrt{16\omega^4 - 63\omega^2 + 64}}
 \end{aligned}$$

(c) Below is the plot of A versus ω




Figure 3.27: (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Forced Vibrations Amplitude of Steady State Response versus Frequency/main.cpp).

(d) In order to find the value of ω that maximizes A , differentiate A with respect to ω

$$A'(\omega) = -\frac{4}{(16\omega^4 - 63\omega^2 + 64)^{3/2}}(64\omega^3 - 126\omega)$$

and then set it equal to zero.

$$\begin{aligned}
 -\frac{4}{(16\omega^4 - 63\omega^2 + 64)^{3/2}}(64\omega^3 - 126\omega) &= 0 \\
 64\omega^3 - 126\omega &= 0 \\
 2\omega(32\omega^2 - 63) &= 0
 \end{aligned}$$

Use the zero product theorem

$$\begin{aligned}
 2\omega &= 0 \\
 \omega &= 0
 \end{aligned}$$

or

$$32\omega^2 - 63 = 0$$

$$\omega = \pm \sqrt{\frac{63}{32}}$$

Since ω has to be positive

$$\omega_{\max} = \sqrt{\frac{63}{32}} \approx 1.403$$

The amplitude that corresponds with this value of ω is

$$\begin{aligned} A_{\max} = A(\omega_{\max}) &= \frac{8}{\sqrt{16\omega^4 - 63\omega^2 + 64}} \\ &= \frac{8}{\sqrt{16\omega_{\max}^4 - 63\omega_{\max}^2 + 64}} \\ &= \frac{8}{\sqrt{16\left(\sqrt{\frac{63}{32}}\right)^4 - 63\left(\sqrt{\frac{63}{32}}\right)^2 + 64}} \\ &= \frac{64}{\sqrt{127}} \\ &\approx 5.679 \end{aligned}$$

```
Solution for u'' + 0.25u' + 2u = 2 cos(1.403t), u(0) = 0, u'(0) = 2

The particular solution is
Y(t) = 5.65618*sin(1.403*t)+0.509435*cos(1.403*t)

The general solution for the homogeneous equation is:
y(t) = e^(-0.125*t)*c1*cos(1.40868*t)+e^(-0.125*t)*c2*sin(1.40868*t)

The solution for the initial value problem is:
-0.509435*e^(-0.125*t)*cos(1.40868*t)-4.25882*e^(-0.125*t)*sin(1.40868*t)+5.65618*sin(1.403*t)+0.509435*cos(1.403*t)

The forced vibrations parameters:
F_0 = 2
w = 1.403
w_0 = 1.41421
w_max = 1.40312
Gamma = 0.03125
R = 5.67908
R_max = 5.67908
Delta = 0.35217
delta = 1.48097

Time taken by function: 723903 microseconds
```

Figure 3.28: The computation of initial value problem solution for the second order linear differential equation for forced vibration with damping, the parameter A (the amplitude) is represented by R in `SymIntegration` (`SymIntegration/Examples/Differential Equations/Test SymIntegration Initial Value Problem for Nonhomogeneous Second Order Linear Differential Equations Forced Vibrations/main.cpp`).

[SI*] Forced Vibrations Without Damping

We now assume that $\gamma = 0$ in Eq. (3.189), thereby obtaining the equation of motion of an undamped forced oscillator

$$mu'' + ku = F_0 \cos \omega t \quad (3.200)$$

The form of the general solution of Eq. (3.200) is different, depending on whether the forcing frequency ω is different from or equal to the natural frequency $\omega_0 = \sqrt{\frac{k}{m}}$ of the unforced

system. First consider the case $\omega \neq \omega_0$; then the general solution of Eq. (3.200) is

$$u = c_1 \cos \omega_0 t + c_2 \sin \omega_0 t + \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos \omega t \quad (3.201)$$

The constants c_1 and c_2 are determined by the initial conditions. The resulting motion is, in general, the sum of two periodic motions of different frequencies (ω_0 and ω) and different amplitudes as well.

It is particularly interesting to suppose that the mass is initially at rest, so that the initial conditions are $u(0) = 0$ and $u'(0) = 0$. Then the energy driving the system comes entirely from the external force, with no contribution from the initial conditions. In this case it turns out that the constants c_1 and c_2 in Eq. (3.201) are given by

$$c_1 = -\frac{F_0}{m(\omega_0^2 - \omega^2)}, \quad c_2 = 0 \quad (3.202)$$

and the solution of Eq. (3.200) is

$$u = \frac{F_0}{m(\omega_0^2 - \omega^2)} (\cos \omega t - \cos \omega_0 t) \quad (3.203)$$

This is the sum of two periodic functions of different periods but the same amplitude. Making use of the trigonometric identities for $\cos(A \pm B)$ with $A = \frac{(\omega_0 + \omega)t}{2}$ and $B = \frac{(\omega_0 - \omega)t}{2}$, we can write in Eq. (3.203) in the form

$$u = \left[\frac{2F_0}{m(\omega_0^2 - \omega^2)} \sin \frac{(\omega_0 - \omega)t}{2} \right] \sin \frac{(\omega_0 + \omega)t}{2} \quad (3.204)$$

If $|\omega_0 - \omega|$ is small, then $\omega_0 + \omega$ is much greater than $|\omega_0 - \omega|$. Consequently, $\sin \frac{(\omega_0 + \omega)t}{2}$ is a rapidly oscillating function compared to $\sin \frac{(\omega_0 - \omega)t}{2}$. Thus the motion is a rapid oscillation with frequency $\frac{\omega_0 + \omega}{2}$ but with a slowly varying sinusoidal amplitude

$$\frac{2F_0}{m|\omega_0^2 - \omega^2|} \left| \sin \frac{(\omega_0 - \omega)t}{2} \right|$$

This type of motion, possessing a periodic variation of amplitude, exhibits what is called a beat. For example, such a phenomenon occurs in acoustics when two tuning forks of nearly equal frequency are excited simultaneously. In this case the periodic variation of amplitude is quite apparent to the unaided ear. In electronics, the variation of the amplitude with time is called amplitude modulation.

i. Functions in SymIntegration Related to Second Order Linear Equations

[SI*] We will list the basic functions that can be used for solving second order linear differential equations:

secondorderlineardiffeq_dsolve(double a, double b, double c, const Symbolic &y, const Symbolic &t)

to solve a homogeneous second order linear differential equation with this form

$$ay'' + by' + cy = 0$$

it will return the general solution, $y(t)$.

secondorderlineardiffeq_ivpsolution(double a, double b, double c, const Symbolic &y, const Symbolic &t, double t_0 , double y_0 , double y'_0)

to solve the initial value problem of a homogeneous second order linear differential equation with this form

$$ay'' + by' + cy = 0$$

with initial value

$$y(t_0) = y_0$$

$$y'(t_0) = y'_0$$

it will also compute the critical value t_m and the maximum value y_m for the initial value problem solution for the case with real and different roots.

```
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration IVPSolve Second Order Linear Equations ]# ./main
IVP Solve for y'' + 5y' + 6y = 0 with y(0) = 2, y'(0) = 3

The general solution is:
c1*e^(-2*t)+c2*e^(-3*t)
The solution for the initial value problem is:
9*e^(-2*t)-7*e^(-3*t)
y' = -18*e^(-2*t)+21*e^(-3*t)
Critical value t_{m} = 0.154151
Maximum value y_{m} = 2.20408
IVP Solve for y'' - y = 0 with y(0) = 2, y'(0) = -1

The general solution is:
c1*e^t+c2*e^(-t)
The solution for the initial value problem is:
0.5*e^t+1.5*e^(-t)
y' = 0.5*e^t-1.5*e^(-t)
Critical value t_{m} = 0.549306
Maximum value y_{m} = 1.73205
IVP Solve for 4y'' - 8y' + 3y = 0 with y(0) = 2, y'(0) = 1/2

The general solution is:
c1*e^(1.5*t)+c2*e^(0.5*t)
The solution for the initial value problem is:
-0.5*e^(1.5*t)+2.5*e^(0.5*t)
y' = -0.75*e^(1.5*t)+1.25*e^(0.5*t)
Critical value t_{m} = 0.510826
Maximum value y_{m} = 2.15166
```

Figure 3.29: The computation of the initial value problem solution for a homogeneous second order linear equation with `secondorderlineardiffeq_ivpsolution()` function (`SymIntegration/Examples/Differential Equations/Test SymIntegration IVPSolve Second Order Linear Equations/main.cpp`).

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Complex Roots Second Order Linear Differential Equations ]# ./main
General solution for  $y'' + y' + 9.25y = 0$  with  $t_{\{0\}} = 0$ ,  $y(t_{\{0\}}) = 2$ ,  $y'(t_{\{0\}}) = 8$ 

The general solution is:
 $y_{\{1\}}(t) = e^{(-0.5*t)}\cos(3*t) + e^{(-0.5*t)}i\sin(3*t)$ 
 $y_{\{2\}}(t) = e^{(-0.5*t)}\cos(3*t) - e^{(-0.5*t)}i\sin(3*t)$ 
Initial Value Problem solution for  $y'' + y' + 9.25y = 0$  with  $t_{\{0\}} = 0$ ,  $y(t_{\{0\}}) = 2$ ,  $y'(t_{\{0\}}) = 8$ 

The complex-valued general solution is:
 $y_{\{1\}}(t) = e^{(-0.5*t)}\cos(3*t) + e^{(-0.5*t)}i\sin(3*t)$ 
 $y_{\{2\}}(t) = e^{(-0.5*t)}\cos(3*t) - e^{(-0.5*t)}i\sin(3*t)$ 
 $u(t) = e^{(-0.5*t)}\cos(3*t)$ 
 $v(t) = e^{(-0.5*t)}\sin(3*t)$ 
The real-valued initial value problem solution is:
 $y(t) = 2*e^{(-0.5*t)}\cos(3*t) + 3*e^{(-0.5*t)}\sin(3*t)$ 
Initial Value Problem solution for  $16y'' - 8y' + 145y = 0$  with  $t_{\{0\}} = 0$ ,  $y(t_{\{0\}}) = -2$ ,  $y'(t_{\{0\}}) = 1$ 

The complex-valued general solution is:
 $y_{\{1\}}(t) = e^{(0.25*t)}\cos(3*t) + e^{(0.25*t)}i\sin(3*t)$ 
 $y_{\{2\}}(t) = e^{(0.25*t)}\cos(3*t) - e^{(0.25*t)}i\sin(3*t)$ 
 $u(t) = e^{(0.25*t)}\cos(3*t)$ 
 $v(t) = e^{(0.25*t)}\sin(3*t)$ 
The real-valued initial value problem solution is:
 $y(t) = -2*e^{(0.25*t)}\cos(3*t) + 0.5*e^{(0.25*t)}\sin(3*t)$ 
Time taken by function: 289530 microseconds

```

Figure 3.30: The computation of the initial value problem solution for a homogeneous second order linear equation with complex conjugate roots (SymIntegration/Examples/Differential Equations/Test SymIntegration Complex Roots Second Order Linear Differential Equations/main.cpp).

wronskian(double a, double b, double c, const Symbolic &y, const Symbolic &t, double t_0 , double y_0 , double y'_0)

to compute the Wronskian of a homogeneous second order linear differential equation with this form

$$ay'' + by' + cy = 0$$

with initial value

$$\begin{aligned} y(t_0) &= y_0 \\ y'(t_0) &= y'_0 \end{aligned}$$

wronskian_fundamentalsolutions(double a, double b, double c, const Symbolic &y, const Symbolic &t)

to compute the fundamental set of solutions of a homogeneous second order linear differential equation with this form

$$ay'' + by' + cy = 0$$

The first fundamental solution y_1 satisfies the initial conditions

$$\begin{aligned} y(t_0) &= 1 \\ y'(t_0) &= 0 \end{aligned}$$

The second fundamental solution y_2 satisfies the initial conditions

$$\begin{aligned} y(t_0) &= 0 \\ y'(t_0) &= 1 \end{aligned}$$

II. Second Order Linear Equations

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Wronskian Second Order Linear Equations ]# ./main

Wronskian for y'' + 5y' + 6y = 0 with t_{0} = 0

The general solution is:
c1*e^(-2*t)+c2*e^(-3*t)

The Wronskian is:-e^(-5*t)
W(t_{0}) = -1

Wronskian for y'' - y = 0 with t_{0} = 0

The solution that satisfies the initial value problem y(0)=1 and y'(0)=0:
0.5*e^t+0.5*e^(-t)

The solution that satisfies the initial value problem y(0)=0 and y'(0)=1:
0.5*e^t-0.5*e^(-t)

The general solution :
0.5*k1*e^t+0.5*k1*e^(-t)+0.5*k2*e^t-0.5*k2*e^(-t)
```

Figure 3.31: The computation of the Wronskian and the fundamental set of solutions for a homogeneous second order linear equation (*SymIntegration/Examples/Differential Equations/Test SymIntegration Wronskian Second Order Linear Equations/main.cpp*).

adjointequation(const Symbolic &c1, const Symbolic &c2, const Symbolic &c3, Symbolic &t)

to compute the adjoint of the original second order linear homogeneous equation with this form:

$$c_1 y'' + c_2 y' + c_3 y = 0$$

with c_1, c_2, c_3 are functions of t , other symbols (e.g. α, β, ν) will be considered as constants.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Second Order Linear Homogeneous Equation Adjoint Equation ]# ./main

Adjoint for x^{2}y'' + xy' + (x^{2} - \nu^{2})y = 0 (Bessel' equation)

Exact equation :
x^{2}(2)*\mu(x)*y'' + x*\mu(x)*y' + x^{2}(2)*\mu(x)*y - \nu^{2}(2)*\mu(x)*y = 0
2*x*\mu(x)*y' + x^{2}(2)*\mu'(x)*y' + x^{2}(2)*\mu(x)*y'' + f'(x)*y + f(x)*y' = 0

Equate the coefficients :
2*x*\mu(x) + x^{2}(2)*\mu'(x) + f(x) = x * \mu(x)

f'(x) = x^{2}(2) - \nu^{2}(2) * \mu(x)

Differentiate both sides of the first equation with respect to : x
3*x*\mu'(x) + x^{2}(2)*\mu''(x) + \mu(x) + f'(x)

Substitute x^{2}(2) - \nu^{2}(2) * \mu(x) for f'(x)
3*x*\mu'(x) + x^{2}(2)*\mu''(x) + \mu(x) + x^{2}(2)*\mu(x) - \nu^{2}(2)*\mu(x) = 0
*****

Adjoint for (1-x^{2})y'' - 2xy' + \alpha(\alpha+1)y = 0 (Legendre's equation)

Exact equation :
-x^{2}(2)*\mu(x)*y'' + \mu(x)*y' - 2*x*\mu(x)*y' + \alpha^{2}(2)*\mu(x)*y + \alpha*\mu(x)*y = 0
-2*x*\mu(x)*y' - x^{2}(2)*\mu'(x)*y' + \mu'(x)*y' - x^{2}(2)*\mu(x)*y'' + \mu(x)*y' + f'(x)*y + f(x)*y' = 0

Equate the coefficients :
-2*x*\mu(x) - x^{2}(2)*\mu'(x) + \mu'(x) + f(x) = -2*x * \mu(x)

f'(x) = \alpha^{2}(2) + \alpha * \mu(x)

Differentiate both sides of the first equation with respect to : x
-2*x*\mu'(x) - x^{2}(2)*\mu''(x) + \mu''(x) + f'(x)

Substitute \alpha^{2}(2) + \alpha * \mu(x) for f'(x)
-2*x*\mu'(x) - x^{2}(2)*\mu''(x) + \mu''(x) + \alpha^{2}(2)*\mu(x) + \alpha*\mu(x) = 0
```

Figure 3.32: The computation of the adjoint of the given second order linear differential equation, Bessel's equation and Legendre's equation (*SymIntegration/Examples/Differential Equations/Test SymIntegration Second Order Linear Homogeneous Equation Adjoint Equation/main.cpp*).

```

*****
Adjoint for y'' - xy' = 0 (Airy's equation)

Exact equation :
μ(x)*y'' - x*μ(x)*y' = 0
μ'(x)*y' + μ(x)*y'' + f'(x)*y + f(x)*y' = 0

Equate the coefficients :
μ'(x) + f(x) = 0 * μ(x)
f'(x) = -x * μ(x)

Differentiate both sides of the first equation with respect to : x
μ''(x) + f'(x)

Substitute -x * μ(x) for f'(x)
μ''(x) - x*μ(x) = 0

```

Figure 3.33: The computation of the adjoint of the given second order linear differential equation, Airy's equation (*SymIntegration/Examples/Differential Equations/Test SymIntegration Second Order Linear Homogeneous Equation Adjoint Equation/main.cpp*).

reductionoforder(const Symbolic &a(t), const Symbolic &b(t), const Symbolic &c(t), const Symbolic &y₁, const Symbolic &t)

to compute the second solution of a second order homogeneous differential equation with this form

$$a(t)y'' + b(t)y' + c(t)y = 0$$

with known $y_1(t)$ (the first solution) and a, b, c are functions of t , which can also be constants as well.

secondorderlineardiffeq_nonhomogeneousequationssolution(const Symbolic &a, const Symbolic &b, const Symbolic &c, const Symbolic &g, const Symbolic &y, const Symbolic &t)

to compute the general solution (particular solution and the complementary solution) of a second order nonhomogeneous differential equation with this form

$$ay'' + by' + cy = g$$

with a, b , and c as constants, and g , a function of t , a one term function (not a linear combination of functions).

secondorderlineardiffeq_nonhomogeneousequationssolution(const Symbolic &a, const Symbolic &b, const Symbolic &c, const SymbolicMatrix &g, const Symbolic &y, const Symbolic &t)

to compute the general solution (particular solution and the complementary solution) of a second order nonhomogeneous differential equation with this form

$$ay'' + by' + cy = g[1] + g[2] + \dots g[n]$$

with a, b , and c as constants, and g is a symbolic matrix with 1 column (symbolic vector to be exact) that consists of the term of $g(t)$, e.g. if we have $g(t) = 3e^{2t} + 2\sin t - 8e^t \cos 2t$, then $g[1] = 3e^{2t}$, $g[2] = 2\sin t$, $g[3] = -8e^t \cos 2t$.

II. Second Order Linear Equations

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Nonhomogeneous Second Order Linear Differential
Equations SymbolicMatrix rhs ]# ./main
W:
[ 3*e^(2*t) ]
[ 2*sin(t) ]
[-8*e^t*cos(2*t)]

Solution for y'' - 3y' - 4y = 3e^{2t} + 2 sin(t) - 8e^t cos(2t)

*****

For rhs = 3*e^(2*t)

The current solution is
Y(t) = -0.5*e^(2*t)
*****

For rhs = 2*sin(t)

The current solution is
Y(t) = -0.294118*sin(t)+0.176471*cos(t)
*****

For rhs = -8*e^t*cos(2*t)
Attempted to cast -8*cos(2*t) to double failed.
Attempted to cast -8*e^t to double failed.

The current solution is
Y(t) = 0.153846*e^t*sin(2*t)+0.769231*e^t*cos(2*t)
*****

The particular solution is
Y(t) = -0.5*e^(2*t)-0.294118*sin(t)+0.176471*cos(t)+0.153846*e^t*sin(2*t)+0.769231*e^t*cos(2*t)
The general solution for the homogeneous equation is:
c1*e^(4*t)+c2*e^(-t)
The general solution for the nonhomogeneous equation is:
c1*e^(4*t)+c2*e^(-t)+0.153846*e^t*sin(2*t)+0.769231*e^t*cos(2*t)
Time taken by function: 645443 microseconds

```

Figure 3.34: The computation of the particular solution for the second order nonhomogeneous differential equation (*SymIntegration/Examples/Differential Equations/Test SymIntegration Nonhomogeneous Second Order Linear Differential Equations SymbolicMatrix rhs/main.cpp*).

* The **attempted to cast $-8\cos 2t$ to double** and **attempted to cast $-8e^t$ to double** are warnings because the source code matching with the pattern of $g(t) = \exp(a * t)$ and $g(t) = \cos(a * t)$ based on the ordering, before it reaches the right pattern of $g(t) = a * \exp(c * t) * \cos(b * t)$, with a, b, c as constants. The `secondorderlineardiffeq_nonhomogeneousequationssolution()` function can handle these $g(t)$ functions:

$$\begin{aligned}
 g(t) &= e^{bt} \\
 g(t) &= ae^{bt} \\
 g(t) &= a \sin(bt) \\
 g(t) &= a \cos(bt) \\
 g(t) &= ae^{bt} \sin(ct) \\
 g(t) &= ae^{bt} \cos(ct)
 \end{aligned}$$

and the linear combinations of the $g(t)$ functions above.

`secondorderlineardiffeq_nonhomogeneousequationssolution(const Symbolic &a, const Symbolic &b, const Symbolic &c, Polynomial<double> &P(t))`

to compute the general solution (particular solution and the complementary solution) of a second order nonhomogeneous differential equation with this form

$$ay'' + by' + cy = P(t)$$

with a, b , and c as constants, with $P(t)$ is a polynomial. In order to solve and find the particular solution with polynomial on the right hand side, we initially assume that $Y(t)$ is a polynomial of the same degree as the nonhomogeneous term, $P(t)$, that is if $P(t) = t^2 - 1$, then $Y(t) = At^2 + Bt + C$. The function currently can handle polynomial of degree n , with n as positive integer.

```

Solution for y'' - 3y' - 4y = 4t^2 - 1
p(t) = 4t^2-1
The particular solution is
Y(t) = -t^(2)+1.5*t-1.375
The general solution for the homogeneous equation is:
c1*e^(4*t)+c2*e^(-t)
The general solution for the nonhomogeneous equation is:
c1*e^(4*t)+c2*e^(-t)-t^(2)+1.5*t-1.375
Solution for y'' - 3y' - 4y = 5t^3 + 2t^2 - 1
p(t) = 5t^3+2t^2-1
The particular solution is
Y(t) = -1.25*t^(3)+2.3125*t^(2)-5.34375*t+5.41406
The general solution for the homogeneous equation is:
c1*e^(4*t)+c2*e^(-t)
The general solution for the nonhomogeneous equation is:
c1*e^(4*t)+c2*e^(-t)-1.25*t^(3)+2.3125*t^(2)-5.34375*t+5.41406
Solution for y'' - 3y' - 4y = 5t - 10
p(t) = 5t-10
The particular solution is
Y(t) = -1.25*t+3.4375
The general solution for the homogeneous equation is:
c1*e^(4*t)+c2*e^(-t)
The general solution for the nonhomogeneous equation is:
c1*e^(4*t)+c2*e^(-t)-1.25*t+3.4375
Solution for y'' - 3y' - 4y = 50
p(t) = 50
The particular solution is
Y(t) = -12.5
The general solution for the homogeneous equation is:
c1*e^(4*t)+c2*e^(-t)
The general solution for the nonhomogeneous equation is:
c1*e^(4*t)+c2*e^(-t)-12.5
Time taken by function: 226665 microseconds

```

Figure 3.35: The computation of the particular solution for the second order nonhomogeneous differential equation with polynomial as the nonhomogeneous term (*SymIntegration/Examples/Differential Equations/Test SymIntegration Nonhomogeneous Second Order Linear Differential Equations Polynomial rhs/main.cpp*).

The code used to handle polynomial of degree n uses Gaussian elimination to obtain the coefficient A_0, A_1, A_n for $Y(t)$, the function `solve_nhsystem_resultonly0` handles this.

```

...
...

void secondorderlineardiffeq_nonhomogeneousequationssolution(const Symbolic &
    lhs_a, const Symbolic &lhs_b, const Symbolic &lhs_c, Polynomial<double> &
    rhs_function)
{
    Symbolic A("A"), t("t");
    Symbolic lhs_final, Yt_final;

```

```

Polynomial<double> p = rhs_function;
cout << "p(t) = " << p << endl;
int n = rhs_function.terms.front().second; // degree of the polynomial
if (n >= 0) // this is working for Polynomial of degree n, n =
    positive integer
{
    Symbolic Yt;
    for (int i = 0; i <= n; ++i)
    {
        Yt += A[i]*(t^(n-i)); // A[i] is symbolic A_{i} so we don't
            need to declare infinite number of Symbolic A1("A1"),... ,
            An("An")
    }
    Symbolic dy = df(Yt,t);
    Symbolic ddy = df(dy,t);

    lhs_final = ddy*lhs_a + dy* lhs_b + Yt*lhs_c;

    vector<vector<double>> mat_A(n+1, vector<double>(n+1));
    vector<vector<double>> vec_b(n+1, vector<double>(1));

    for (int i = 0; i <= n; ++i)
    {
        Symbolic coeff_ti = lhs_final.coeff(t,n-i);
        for (int j = 0; j <= n; ++j)
        {
            mat_A[i][j] = coeff_ti.coeff(A[j],1);
        }
    }

    int n1 = n;
    for (int i = 0 ; i < n+1 ; ++i)
    {
        if(p.terms.front().second == n1) // p.terms.front().second =
            get the degree of the polynomial
        {
            vec_b[i][0] = p.terms.front().first; // p.terms.front().
                first = get the coefficient
            p.terms.pop_front(); // remove value from front of list
        }
        else if(p.terms.front().second != n1)
        {
            vec_b[i][0] = 0;
        }
        n1 = n1-1;
    }

    vector<double> c_solution;

```

```

        solve_nhsystem_resultonly(mat_A,vec_b,c_solution);

        for (int i = 0 ; i < n+1 ; ++i)
        {
            Yt_final += c_solution[i]*(t^(n-i));
        }
        cout << "\nThe particular solution is\nY(t) = " << Yt_final << endl;
    }
}

```

Code 40: *src/dsolve.cpp*

secondorderlineardiffeq_nonhomogeneousequationsivpsolution(const Symbolic &a, const Symbolic &b, const Symbolic &c, const Symbolic &g, double y_0 , double y'_0 , const Symbolic &y, const Symbolic &t)

to compute the initial value problem general solution (particular solution and the complementary solution) of a second order nonhomogeneous differential equation with this form

$$ay'' + by' + cy = g$$

and the initial conditions

$$y(0) = y_0, \quad y'(0) = y'_0$$

with a, b , and c as constants, and g , a function of t , a one term function (not a linear combination of functions).

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Initial Value Problem for Nonhomogeneous Second Order Linear |
rced Vibrations ]# ./main

Solution for u'' + u' + 1.25u = 3 cos(t)

The particular solution is
Y(t) = 2.82353*sin(t)+0.705882*cos(t)

The general solution for the homogeneous equation is:
y(t) = e^(-0.5*t)*c1*cos(t)+e^(-0.5*t)*c2*sin(t)

The solution for the initial value problem is:
1.29412*e^(-0.5*t)*cos(t)+0.823529*e^(-0.5*t)*sin(t)+2.82353*sin(t)+0.705882*cos(t)

Time taken by function: 168117 microseconds

```

Figure 3.36: The computation of the initial value problem general solution for the second order nonhomogeneous differential equation (*SymIntegration/Examples/Differential Equations/Test SymIntegration Initial Value Problem for Nonhomogeneous Second Order Linear Differential Equations Forced Vibrations/main.cpp*).

secondorderlineardiffeq_springmasssystem(double a , double b , double c , const Symbolic &y, const Symbolic &t, double y_0 , double y'_0)

The computation of initial value problem solution for the second order linear differential equation for spring-mass system / mechanical vibration with damping, small damping or no damping. The output also returns δ (the phase), T_d (the quasi period), and t (the time which the mass first passes through its equilibrium position).

secondorderlineardiffeq_RLCserieselectriccircuit(double R , double L , double C , double q_0 , double q'_0)

to compute the charge $q(t)$ on the capacitor at any time given the resistor R , inductor L , and the capacitor C for the series circuit. We have q_0 to represent the initial charge on the

capacitor, and $q'_0 = I_0$ represents the initial current in the circuit.

secondorderlineardiffeq_nonhomogeneousequationsivpforcedvibrationsolution(const Symbolic &a, const Symbolic &b, const Symbolic &c, const Symbolic &g(t) double y₀, double y'₀, const Symbolic &y, const Symbolic &t)
to compute the initial value problem general solution (particular solution and the complementary solution) of a second order nonhomogeneous differential equation with this form

$$ay'' + by' + cy = g(t)$$

and the initial conditions

$$y(0) = y_0, \quad y'(0) = y'_0$$

with a, b , and c as constants, and g , a function of t , a one term function (not a linear combination of functions). The output also contains some forced vibration parameters : $F_0, \omega, \omega_0, \omega_{\max}, \Gamma, R, R_{\max}, \Delta, \delta$. If you want to plot it, the forcing function is represented by $g(t)$, and the solution is represented by the general solution of the initial value problem ($u(t)$) that can be expressed a sum of the complementary solution and the particular solution.

$$u(t) = u_c(t) + u_p(t)$$




Figure 3.37: A forced vibration with damping; solution of $u'' + 0.125u' + u = 3 \cos t, u(0) = 2, u'(0) = 0$ (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Forced Vibrations IVP Second Order ODE Solution and Forcing Function/main.cpp).

III. HIGHER ORDER LINEAR EQUATIONS

[SI*] General Theory of n th Order Linear Equations

An n th order linear differential equation is an equation of the form

$$P_0(t) \frac{d^n y}{dt^n} + P_1(t) \frac{d^{n-1} y}{dt^{n-1}} + \cdots + P_{n-1}(t) \frac{dy}{dt} + P_n(t) y = G(t) \quad (3.205)$$

We assume that the functions $P_0, \dots, P_n$ and G are continuous real-valued functions on some interval $I : \alpha < t < \beta$, and that P_0 is nowhere zero in this interval. Then, dividing Eq. (3.205) by $P_0(t)$, we obtain

$$L[y] = \frac{d^n y}{dt^n} + p_1(t) \frac{d^{n-1} y}{dt^{n-1}} + \dots + p_{n-1}(t) \frac{dy}{dt} + p_n(t)y = g(t) \quad (3.206)$$

The linear differential operator L of order n defined by Eq. (3.206) is similar to the second order operator. The mathematical theory associated with Eq. (3.206) is completely analogous to that for the second order linear equation; for this reason we simply state the results for the n th order problem. The proofs of most of the results are also similar to those for the second order equation and are usually left as exercises.

Since Eq. (3.206) involves the n th derivative of y with respect to t , it will, so to speak, require n integrations to solve Eq. (3.206). Each of these integrations introduces an arbitrary constant. Hence we can expect that, to obtain a unique solution, it is necessary to specify n initial conditions

$$y(t_0) = y_0, \quad y'(t_0) = y'_0, \quad \dots, y^{(n-1)}(t_0) = y_0^{(n-1)} \quad (3.207)$$

where t_0 may be any point in the interval I and $y_0, \dots, y_0^{(n-1)}$ is any set of prescribed real constants.

Theorem 3.9: Existence and Uniqueness

If the functions $p_1, p_2, \dots, p_n$, and g are continuous on the open interval I then there exists exactly one solution $y = \phi(t)$ of the differential equation (3.206) that also satisfies the initial conditions (3.207). This solution exists throughout the interval I .

The Homogeneous Equation

As in the corresponding second order problem, we first discuss the homogeneous equation

$$L[y] = y^{(n)} + p_1(t)y^{(n-1)} + \dots + p_{n-1}(t)y' + p_n(t)y = 0 \quad (3.208)$$

If the functions $y_1, y_2, \dots, y_n$ are solutions of Eq. (3.208), then it follows by direct computation that the linear combination

$$y = c_1 y_1(t) + c_2 y_2(t) + \dots + c_n y_n(t) \quad (3.209)$$

where $c_1, c_2, \dots, c_n$ are arbitrary constants, is also a solution of Eq. (3.208). For any choice of the point t_0 in I , and for any choice of $y_0, y'_0, \dots, y_0^{(n-1)}$, we must be able to determine $c_1, \dots, c_n$ so that the equations

$$\begin{aligned} c_1 y_1(t_0) + \dots + c_n y_n(t_0) &= y_0 \\ c_1 y'_1(t_0) + \dots + c_n y'_n(t_0) &= y'_0 \\ &\vdots \\ c_1 y_1^{(n-1)}(t_0) + \dots + c_n y_n^{(n-1)}(t_0) &= y_0^{(n-1)} \end{aligned} \quad (3.210)$$

are satisfied. Equations (3.210) can be solved uniquely for the constants $c_1, \dots, c_n$, provided that the determinant of coefficients is not zero. On the other hand, if the determinant of coefficients is zero, then it is always possible to choose values of $y_0, y'_0, \dots, y_0^{(n-1)}$ so that Eqs.

(3.210) do not have a solution. Hence a necessary and sufficient condition for the existence of a solution of Eqs. (3.210) for arbitrary values of $y_0, y'_0, \dots, y_0^{(n-1)}$ is that the Wronskian

$$W(y_1, \dots, y_n) = \begin{vmatrix} y_1 & y_2 & \cdots & y_n \\ y'_1 & y'_2 & \cdots & y'_n \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)} & y_2^{(n-1)} & \cdots & y_n^{(n-1)} \end{vmatrix} \quad (3.211)$$

is not zero at $t = t_0$. Since t_0 can be any point in the interval I , it is necessary and sufficient that $W(y_1, y_2, \dots, y_n)$ be nonzero at every point in the interval. Just as for the second order linear equation, it can be shown that if $y_1, y_2, \dots, y_n$ are solutions of Eq. (3.208), then $W(y_1, y_2, \dots, y_n)$ either is zero for every t in the interval I or else is never zero.

Theorem 3.10: Wronskian and Linear Combination

If the functions $p_1, p_2, \dots, p_n$ are continuous on the open interval I , if the functions $y_1, y_2, \dots, y_n$ are solutions of Eq. (3.208), and if $W(y_1, y_2, \dots, y_n) \neq 0$ for at least one point in I , then every solution of Eq. (3.208) can be expressed as a linear combination of the solutions $y_1, y_2, \dots, y_n$.

A set of solutions $y_1, y_2, \dots, y_n$ of Eq. (3.208) whose Wronskian is nonzero is referred to as a fundamental set of solutions. The existence of a fundamental set of solutions can be demonstrated precisely the same way as for the second order linear equation. Since all solutions of Eq. (3.208) are of the form (3.209), we use the term general solution to refer to an arbitrary linear combination of any fundamental set of solutions of Eq. (3.208).

Linear Dependence and Independence

The functions $f_1, f_2, \dots, f_n$ are said to be linearly dependent on an interval I if there exists a set of constants $k_1, k_2, \dots, k_n$, not all zero, such that

$$k_1 f_1(t) + k_2 f_2(t) + \cdots + k_n f_n(t) = 0 \quad (3.212)$$

for all t in I . The functions $f_1, \dots, f_n$ are said to be linearly independent on I if they are not linearly dependent there.

Theorem 3.11: Fundamental Set of Solutions on I

If $y_1(t), \dots, y_n(t)$ is a fundamental set of solutions of Eq. (3.208)

$$L[y] = y^{(n)} + p_1(t)y^{(n-1)} + \cdots + p_{n-1}(t)y' + p_n(t)y = 0$$

on an interval I , then $y_1(t), \dots, y_n(t)$ are linearly independent on I . Conversely, if $y_1(t), \dots, y_n(t)$ are linearly independent solutions of Eq. (3.208) on I , then they form a fundamental set of solutions on I .

The Nonhomogeneous Equation

Now consider the nonhomogeneous equation (3.206)

$$L[y] = y^{(n)} + p_1(t)y^{(n-1)} + \cdots + p_{n-1}(t)y' + p_n(t)y = g(t)$$

If Y_1 and Y_2 are any two solutions of Eq. (3.206), then it follows immediately from the linearity of the operator L that

$$L[Y_1 - Y_2](t) = L[Y_1](t) - L[Y_2](t) = g(t) - g(t) = 0$$

Hence the difference of any two solutions of the nonhomogeneous equation (3.206) is a solution of the homogeneous equation (3.208). Since any solution of the homogeneous equation can be expressed as a linear combination of a fundamental set of solutions $y_1, \dots, y_n$, it follows that any solution of Eq. (3.206) can be written as

$$y = c_1 y_1(t) + c_2 y_2(t) + \dots + c_n y_n(t) + Y(t) \quad (3.213)$$

where Y is some particular solution of the nonhomogeneous equation (3.206). The linear combination (3.213) is called the general solution of the nonhomogeneous equation (3.206). Thus the primary problem is to determine a fundamental set of solutions $y_1, \dots, y_n$ of the homogeneous equation (3.208). If the coefficients are constants, this is a fairly simple problem. If the coefficients are not constants, it is usually necessary to use numerical methods or series methods. The method of reduction of order also applies to n th order linear equations, but seldom useful for equations of higher than second order.

[SI*] **Homogeneous Equations with Constant Coefficients**

Consider the n th order linear homogeneous differential equation

$$L[y] = a_0 y^{(n)} + a_1 y^{(n-1)} + \dots + a_{n-1} y' + a_n y = 0 \quad (3.214)$$

where $a_0, a_1, \dots, a_n$ are real constants. From our knowledge of second order linear equations with constant coefficients, it is natural to anticipate that $y = e^{rt}$ is a solution of Eq. (3.214) for suitable values of r . Indeed,

$$L[e^{rt}] = e^{rt}(a_0 r^n + a_1 r^{n-1} + \dots + a_{n-1} r + a_n) = e^{rt} Z(r) \quad (3.215)$$

for all r , where

$$Z(r) = a_0 r^n + a_1 r^{n-1} + \dots + a_{n-1} r + a_n \quad (3.216)$$

For those values of r for which $Z(r) = 0$, it follows that $L[e^{rt}] = 0$ and $y = e^{rt}$ is a solution of Eq. (3.214). The polynomial $Z(r)$ is called the characteristic polynomial, and the equation $Z(r) = 0$ is the characteristic equation of the differential equation (3.214). A polynomial of degree n has n zeros, say, $r_1, r_2, \dots, r_n$, some of which may be equal; hence we can write the characteristic polynomial in the form

$$Z(r) = a_0(r - r_1)(r - r_2) \dots (r - r_n) \quad (3.217)$$

Real and Unequal Roots

If the roots of the characteristic equation are real and no two are equal, then we have n distinct solutions $e^{r_1 t}, e^{r_2 t}, \dots, e^{r_n t}$ of Eq. (3.214). If these functions are linearly independent, then the general solution of Eq. (3.214) is

$$y = c_1 e^{r_1 t} + c_2 e^{r_2 t} + \dots + c_n e^{r_n t} \quad (3.218)$$

One way to establish the linear independence of $e^{r_1 t}, e^{r_2 t}, \dots, e^{r_n t}$ is to evaluate their Wronskian determinant.

[SI*] **Example:**

Find the general solution of

$$y^{(4)} + y''' - 7y'' - y' + 6y = 0 \quad (3.219)$$

Also find the solution that satisfies the initial conditions

$$y(0) = 1, \quad y'(0) = 0, \quad y''(0) = -2, \quad y'''(0) = -1 \quad (3.220)$$

and plot its graph.

Solution:

Assuming that $y = e^{rt}$, we must determine r by solving the polynomial equation

$$r^4 + r^3 - 7r^2 - r + 6 = 0 \quad (3.221)$$

The roots of this equation are $r_1 = 1, r_2 = -1, r_3 = 2, r_4 = -3$. Therefore the general solution of Eq. (3.219) is

$$y = c_1 e^t + c_2 e^{-t} + c_3 e^{2t} + c_4 e^{-3t} \quad (3.222)$$

The initial conditions (3.220) require that $c_1, \dots, c_4$ satisfy the four equations

$$\begin{aligned} c_1 + c_2 + c_3 + c_4 &= 1 \\ c_1 - c_2 + 2c_3 - 3c_4 &= 0 \\ c_1 + c_2 + 4c_3 + 9c_4 &= -2 \\ c_1 - c_2 + 8c_3 - 27c_4 &= -1 \end{aligned} \quad (3.223)$$

By solving this system of four linear algebraic equations, we find that

$$c_1 = \frac{11}{8}, \quad c_2 = \frac{5}{12}, \quad c_3 = -\frac{2}{3}, \quad c_4 = -\frac{1}{8}$$

Therefore the solution of the initial value problem is

$$y = \frac{11}{8} e^t + \frac{5}{12} e^{-t} - \frac{2}{3} e^{2t} - \frac{1}{8} e^{-3t} \quad (3.224)$$

```

Polynomial : r^(4)+r^(3)-7*r^(2)-r+6
P:
(1,0)
(1,0)
(-7,0)
(-1,0)
(6,0)

P':
(4,0)
(3,0)
(-14,0)
(-1,0)

Initial guess:
(1.1,0)
(2,0)
(3,0)
(4,0)

Roots found at iteration: 6
*****

End of iteration

z_{i} final:
(1,0)
(2,0)
(-1,0)
(-3,0)

Unique vector:
(1,0)
(2,0)
(-1,0)
(-3,0)

Duplicate vector:

The general solution is:
y(t) = e^t*c[0]+e^(2*t)*c[1]+e^(-t)*c[2]+e^(-3*t)*c[3]

The initial value problem solution is:
y(t) = 1.375*e^t-0.666667*e^(2*t)+0.416667*e^(-t)-0.125*e^(-3*t)

Time taken by function: 109664 microseconds

```

Figure 3.38: The computation of the initial value problem general solution for the fourth order homogeneous differential equation with real and unequal roots (*SymIntegration/Examples/Differential Equations/Test SymIntegration IVP Solution for Higher Order Linear Differential Equations/main.cpp*).




Figure 3.39: The plot of the solution of the initial value problem from the differential equation (3.219) with the initial conditions from Eq. (3.220) (*Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D IVP Solution for Higher Order Homogeneous Linear Differential Equation/main.cpp*).

Complex Roots

If the characteristic equation has complex roots, they must occur in conjugate pairs, $\lambda \pm i\mu$, since the coefficients $a_0, \dots, a_n$ are real numbers. Provided that none of the roots is repeated,

the general solution of Eq. (3.214) is still of the form (3.218). However, just as for the second order equation, we can replace the complex-valued solution $e^{(\lambda+i\mu)t}$ and $e^{(\lambda-i\mu)t}$ by the real-valued solutions

$$e^{\lambda t} \cos(\mu t), \quad e^{\lambda t} \sin(\mu t) \quad (3.225)$$

obtained as the real and imaginary parts of $e^{(\lambda+i\mu)t}$. Thus, even though some of the roots of the characteristic equation are complex, it is still possible to express the general solution of Eq. (3.214) as a linear combination of real-valued solutions.

[SI*] **Example:**

Find the general solution of

$$y^{(4)} - y = 0 \quad (3.226)$$

Also find the solution that satisfies the initial conditions

$$y(0) = \frac{7}{2}, \quad y'(0) = -4, \quad y''(0) = \frac{5}{2}, \quad y'''(0) = -2 \quad (3.227)$$

and draw its graph.

Solution:

Substituting e^{rt} for y , we find that the characteristic equation is

$$r^4 - 1 = (r^2 - 1)(r^2 + 1) = 0$$

Therefore the roots are $r = 1, -1, i, -i$, and the general solution of Eq. (3.226) is

$$y = c_1 e^t + c_2 e^{-t} + c_3 \cos t + c_4 \sin t$$

If we impose the initial conditions (3.227), we find that

$$c_1 = 0, \quad c_2 = 3, \quad c_3 = \frac{1}{2}, \quad c_4 = -1$$

thus the solution of the given initial value problem is

$$y = 3e^{-t} + \frac{1}{2} \cos t - \sin t \quad (3.228)$$

Observe that the initial conditions (3.227) cause the coefficient c_1 of the exponentially growing term in the general solution to be zero. Therefore this term is absent in the solution (3.228), which describes an exponential decay to a steady oscillation. However, if the initial conditions are changed slightly, then c_1 is likely to be nonzero, and the nature of the solution changes enormously. For example, if the first three initial conditions remain the same, but the value of $y'''(0)$ is changed from -2 to $-\frac{15}{8}$, then the solution of the initial value problem becomes

$$y = \frac{1}{32} e^t + \frac{95}{32} e^{-t} + \frac{1}{2} \cos t - \frac{17}{16} \sin t \quad (3.229)$$

The coefficients in Eq. (3.229) differ only slightly from those in Eq. (3.228), but the exponentially growing term, even with the relatively small coefficient of $\frac{1}{32}$, completely dominates the solution by the time t is larger than about 4 or 5.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration IVP Solution for Higher Order Linear Differential Equations
]# ./main
Polynomial : r^(4)-1
P:
(1,0)
(0,0)
(0,0)
(0,0)
(-1,0)

P':
(4,0)
(0,0)
(0,0)
(0,0)

Initial guess:
(1,0)
(2,0)
(3,0)
(4,0)

Roots found at iteration: 31
*****

End of iteration

z (i) final:
(1,0)
(-1,0)
(0,1)
(0,-1)

The general solution is:
y(t) = e^(-t)*c[0]+c[1]*cos(t)+c[2]*sin(t)+e^t*c[3]

The initial value problem solution is:
y(t) = 3*e^(-t)+0.5*cos(t)-sin(t)

Time taken by function: 159680 microseconds

```

Figure 3.40: *The computation of the initial value problem general solution for the fourth order homogeneous differential equation with complex roots (SymIntegration/Examples/Differential Equations/Test SymIntegration IVP Solution for Higher Order Linear Differential Equations/main.cpp).*

```

P:
(1,0)
(0,0)
(0,0)
(0,0)
(-1,0)

P':
(4,0)
(0,0)
(0,0)
(0,0)
(0,0)

Initial guess:
(1,1,0)
(2,0)
(3,0)
(4,0)

Roots found at iteration: 31
*****

End of iteration

z {i} final:
(1,0)
(0,1)
(0,-1)
(-1,0)

Unique vector:
(1,0)
(0,1)
(0,-1)
(-1,0)

Duplicate vector:

The general solution is:
y(t) = e^t*c[0]+c[1]*cos(t)+c[2]*sin(t)+e^(-t)*c[3]

The initial value problem solution is:
y(t) = 0.03125*e^t+0.5*cos(t)-1.0625*sin(t)+2.96875*e^(-t)

Time taken by function: 168841 microseconds

```

Figure 3.41: The computation of the initial value problem general solution for the fourth order homogeneous differential equation with complex roots that will have the solution of Eq. (3.229) (*SymIntegration/Examples/Differential Equations/Test SymIntegration IVP Solution for Higher Order Linear Differential Equations/main.cpp*).




Figure 3.42: The plots of the solution of the initial value problem of Eqs. (3.228) and (3.229) (*Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D IVP Solution for Higher Order Homogeneous Linear Differential Equation with Complex Roots/main.cpp*).

Repeated Roots

If the roots of the characteristic equation are not distinct—that is, if some roots are repeated—then the solution (3.218) is clearly not the general solution of Eq. (3.214). Recall that if r_1 is a repeated root for the second order linear equation $a_0 y'' + a_1 y' + a_2 y = 0$, then two linearly independent solutions are $e^{r_1 t}$ and $t e^{r_1 t}$. For an equation of order n , if a root of $Z(r) = 0$, say $r = r_1$, has multiplicity s (where $s \leq n$), then

$$e^{r_1 t}, \quad t e^{r_1 t}, \quad t^2 e^{r_1 t}, \quad \dots, \quad t^{s-1} e^{r_1 t} \quad (3.230)$$

are corresponding solutions of Eq. (1).

If a complex root $\lambda \pm i\mu$ is repeated s times, the complex conjugate $\lambda - i\mu$ is also repeated s times. Corresponding to these $2s$ complex-valued solutions, we can find $2s$ real-valued solutions by noting that the real and imaginary parts of $e^{(\lambda+i\mu)t}, t e^{(\lambda+i\mu)t}, \dots, t^{s-1} e^{(\lambda+i\mu)t}$ are also linearly independent solutions:

$$\begin{aligned} e^{\lambda t} \cos(\mu t), \quad e^{\lambda t} \sin(\mu t), t e^{\lambda t} \cos(\mu t), \quad t e^{\lambda t} \sin(\mu t), \\ \dots t^{s-1} e^{\lambda t} \cos(\mu t), \quad t^{s-1} e^{\lambda t} \sin(\mu t), \end{aligned}$$

Hence the general solution of Eq. (3.214) can always be expressed as a linear combination of n real-valued solutions.

[SI*] Example:

Find the general solution of

$$y^{(4)} + 2y'' + y = 0 \quad (3.231)$$

The characteristic equation is

$$r^4 + 2r^2 + 1 = (r^2 + 1)(r^2 + 1) = 0$$

The roots are $r = i, i, -i, -i$, and the general solution of Eq. (3.231) is

$$c_1 \cos t + c_2 \sin t + c_3 t \cos t + c_4 t \sin t$$

```

P:
(1,0)
(0,0)
(2,0)
(0,0)
(1,0)

P':
(4,0)
(0,0)
(4,0)
(0,0)

Initial guess:
(1.1,0)
(2,0)
(3,0)
(4,0)

Roots found at iteration: 47
*****

End of iteration

z {i} final:
(0,-1)
(0,1)
(0,1)
(0,-1)

```

Figure 3.43: The computation of the initial value problem general solution for the fourth order homogeneous differential equation with repeated roots (*SymIntegration/Examples/Differential Equations/Test SymIntegration IVP Solution for Higher Order Linear Differential Equations/main.cpp*).


```

Roots found at iteration: 47
*****
End of iteration

z_{i} final:
(0,-1)
(0,1)
(0,1)
(0,-1)

Unique vector:

Duplicate vector:
(0,-1)
(0,1)
(0,1)
(0,-1)

Deleted Duplicate vector:
(0,-1)
(0,1)

Final Duplicate vector:
(0,1)

The general solution is:
y(t) = c[0]*cos(t)+c[1]*sin(t)+c[2]*cos(t)*t+c[3]*sin(t)*t

The initial value problem solution is:
y(t) = cos(t)-0.5*sin(t)+0.5*cos(t)*t-0.5*sin(t)*t

Time taken by function: 150408 microseconds

```

Figure 3.44: The computation of the initial value problem general solution for the fourth order homogeneous differential equation with repeated roots (*SymIntegration/Examples/Differential Equations/Test SymIntegration IVP Solution for Higher Order Linear Differential Equations/main.cpp*).

[SI*] Example:

Consider the spring-mass system, consisting of two unit masses suspended from springs with spring constants 3 and 2, respectively. Assume that there is no damping in the system.




Figure 3.45: A two-spring, two-mass system.

- (a) Show that the displacements u_1 and u_2 of the masses from their respective equilibrium

positions satisfy the equations

$$u_1'' + 5u_1 = 2u_2, \quad u_2'' + 2u_2 = 2u_1 \quad (3.232)$$

- (b) Solve the first of Eqs. (3.232) for u_2 and substitute into the second equation, thereby obtaining the following fourth order equation for u_1 :

$$u_1^{(4)} + 7u_1'' + 6u_1 = 0 \quad (3.233)$$

Find the general solution of Eq. (3.233)

- (c) Suppose that the initial conditions are

$$u_1(0) = 1, \quad u_1'(0) = 0, \quad u_2(0) = 2, \quad u_2'(0) = 0 \quad (3.234)$$

Use the first of Eqs. (3.232) and the initial conditions (3.234) to obtain values for $u_1''(0)$ and $u_1'''(0)$. Then show that the solution of Eq. (3.233) that satisfies the four initial conditions on u_1 is $u_1(t) = \cos t$. Show that the corresponding solution $u_2(t) = 2 \cos t$.

- (d) Now suppose that the initial conditions are

$$u_1(0) = -2, \quad u_1'(0) = 0, \quad u_2(0) = 1, \quad u_2'(0) = 0 \quad (3.235)$$

Proceed as in part (c) to show that the corresponding solutions are $u_1(t) = -2 \cos \sqrt{6}t$ and $u_2(t) = \cos \sqrt{6}t$.

- (e) Observe that the solutions obtained in parts (c) and (d) describe two distinct modes of vibration. In the first, the frequency of the motion is 1, and the two masses move in phase, both moving up or down together; the second mass moves twice as far as the first. The second motion has frequency $\sqrt{6}$, and the masses move out of phase with each other, one moving down while the other is moving up, and vice versa. In this mode the first mass moves twice as far as the second. For other initial conditions, not proportional to either of Eqs. (3.234) or (3.235), the motion of the masses is a combination of these two modes.

Solution:

- (a) Start by drawing free-body diagram for the two masses. If m_1 is displaced a distance u_1 and m_2 is displaced a distance u_2 , then we have the following two diagrams.




Figure 3.46: A two-spring, two-mass system.

Apply Newton's second law twice—once for m_1 and once for m_2 .

$$\begin{aligned} k_2(u_2 - u_1) - k_1 u_1 &= m_1 a_1 \\ -k_2(u_2 - u_1) &= m_2 a_2 \end{aligned}$$

Plug in the numbers for k_1, k_2, m_1 , and m_2 and use the fact that acceleration is the second derivative of position.

$$\begin{aligned} 2(u_2 - u_1) - 3u_1 &= u_1'' \\ -2(u_2 - u_1) &= u_2'' \end{aligned}$$

Therefore, the equations of motion are

$$u_1'' + 5u_1 = 2u_2 \tag{3.236}$$

$$u_2'' + 2u_2 = 2u_1 \tag{3.237}$$

(b) Divide both sides of equation (3.236) by 2

$$\frac{1}{2}u_1'' + \frac{5}{2}u_1 = u_2$$

and plug this formula for u_2 into equation (3.237).

$$\begin{aligned} \left(\frac{1}{2}u_1'' + \frac{5}{2}u_1\right)'' + 2\left(\frac{1}{2}u_1'' + \frac{5}{2}u_1\right) &= 2u_1 \\ \frac{1}{2}u_1^{(4)} + \frac{5}{2}u_1'' + 5u_1 &= 2u_1 \end{aligned}$$

$$u_1^{(4)} + 7u_1'' + 6u_1 = 0 \tag{3.238}$$

This is a homogeneous ODE with constant coefficients, so the solution is of the form $u_1 = e^{rt}$.

$$\begin{aligned}u_1 &= e^{rt} \\u_1' &= re^{rt} \\u_1'' &= r^2 e^{rt} \\u_1''' &= r^3 e^{rt} \\u_1^{(4)} &= r^4 e^{rt}\end{aligned}$$

Substitute these expressions into the ODE

$$r^4 e^{rt} + 7(r^2 e^{rt}) + 6(e^{rt}) = 0$$

Divide both sides by e^{rt} .

$$\begin{aligned}r^4 + 7r^2 + 6 &= 0 \\(r^2 + 6)(r^2 + 1) &= 0 \\r &= \{-\sqrt{6}i, \sqrt{6}i, -i, i\}\end{aligned}$$

Four solutions to equation (3.238) are then $u_1 = e^{-\sqrt{6}it}$, $u_1 = e^{\sqrt{6}it}$ and $u_1 = e^{-it}$ and $u_1 = e^{it}$. By the principle of superposition, the general solution for u_1 is a linear combination of these four.

$$\begin{aligned}u_1(t) &= C_1 e^{-\sqrt{6}it} + C_2 e^{\sqrt{6}it} + C_3 e^{-it} + C_4 e^{it} \\&= C_1(\cos \sqrt{6}t - i \sin \sqrt{6}t) + C_2(\cos \sqrt{6}t + i \sin \sqrt{6}t) + C_3(\cos t - i \sin t) + C_4(\cos t + i \sin t) \\&= (C_1 + C_2) \cos \sqrt{6}t + (-iC_1 + iC_2) \sin \sqrt{6}t + (C_3 + C_4) \cos t + (-iC_3 + iC_4) \sin t \\&= C_5 \cos \sqrt{6}t + C_6 \sin \sqrt{6}t + C_7 \cos t + C_8 \sin t\end{aligned}$$

(c) Suppose that the initial conditions are

$$\begin{aligned}u_1(0) &= 1, & u_1'(0) &= 0 \\u_2(0) &= 2, & u_2'(0) &= 0\end{aligned}$$

From equation (3.236), $u_1''(0)$ and $u_1'''(0)$ can be determined.

$$u_1'' = 2u_2 - 5u_1 \rightarrow \begin{cases} u_1''(0) = 2u_2(0) - 5u_1(0) = -1 \\ u_1'''(0) = 2u_2'(0) - 5u_1'(0) = 0 \end{cases}$$

Differentiate the general solution three times.

$$\begin{aligned}u_1(t) &= C_5 \cos \sqrt{6}t + C_6 \sin \sqrt{6}t + C_7 \cos t + C_8 \sin t \\u_1'(t) &= -C_5 \sqrt{6} \sin \sqrt{6}t + C_6 \sqrt{6} \cos \sqrt{6}t - C_7 \sin t + C_8 \cos t \\u_1''(t) &= -6C_5 \cos \sqrt{6}t - 6C_6 \sin \sqrt{6}t - C_7 \cos t - C_8 \sin t \\u_1'''(t) &= 6\sqrt{6}C_5 \sin \sqrt{6}t - 6\sqrt{6}C_6 \cos \sqrt{6}t + C_7 \sin t - C_8 \cos t\end{aligned}$$

Apply the initial conditions now to determine C_5, C_6, C_7 , and C_8 .

$$\begin{aligned} u_1(0) &= C_5 + C_7 = 1 \\ u_1'(0) &= C_6\sqrt{6} + C_8 = 0 \\ u_1''(0) &= -6C_5 - C_7 = -1 \\ u_1'''(0) &= -6\sqrt{6}C_6 - C_8 = 0 \end{aligned}$$

Solving this system of equations yields $C_5 = 0, C_6 = 0, C_7 = 1$, and $C_8 = 0$. Therefore,

$$u_1(t) = \cos t$$

Now that $u_1(t)$ is known, plug it back into equation (3.237) to find $u_2(t)$.

$$u_2'' + 2u_2 = 2 \cos t$$

This is a linear nonhomogeneous ODE, so its general solution can be expressed as a sum of the complementary solution and the particular solution.

$$u_2(t) = u_{2c}(t) + u_{2p}(t)$$

The complementary solution satisfies the associated homogeneous equation.

$$u_{2c}'' + 2u_{2c} = 0 \tag{3.239}$$

This is a homogeneous ODE with constant coefficients, so the solution is of the form $u_{2c} = e^{st}$.

$$u_{2c} = e^{st} \rightarrow u_{2c}' = se^{st} \rightarrow u_{2c}'' = s^2 e^{st}$$

Substitute these expressions into the ODE.

$$s^2 e^{st} + 2(e^{st}) = 0$$

To compute s , divide both sides by e^{st}

$$\begin{aligned} s^2 + 2 &= 0 \\ s &= \{-i\sqrt{2}, i\sqrt{2}\} \end{aligned}$$

As a result, two solutions to equation (3.239) are $u_{2c} = e^{-i\sqrt{2}t}$ and $u_{2c} = e^{i\sqrt{2}t}$. By the principle of superposition, the general solution for u_{2c} is a linear combination of these two.

$$\begin{aligned} u_{2c}(t) &= C_9 e^{-i\sqrt{2}t} + C_{10} e^{i\sqrt{2}t} \\ &= C_9 (\cos \sqrt{2}t - i \sin \sqrt{2}t) + C_{10} (\cos \sqrt{2}t + i \sin \sqrt{2}t) \\ &= (C_9 + C_{10}) \cos \sqrt{2}t + (-iC_9 + iC_{10}) \sin \sqrt{2}t \\ &= C_{11} \cos \sqrt{2}t + C_{12} \sin \sqrt{2}t \end{aligned}$$

On the other hand, the particular solution satisfies

$$u_{2p}'' + 2u_{2p} = 2 \cos t$$

Since the nonhomogeneous term is cosine and there are only even derivatives on the left, we use the trial solution $u_{2p} = A \cos t$. Plug this into the ODE to determine A .

$$\begin{aligned}(A \cos t)'' + 2(A \cos t) &= 2 \cos t \\ (-A \cos t) + 2A \cos t &= 2 \cos t \\ A \cos t &= 2 \cos t\end{aligned}$$

Matching the coefficients, we see that $A = 2$, which means $u_{2p}(t) = 2 \cos t$. The general solution for $u_2(t)$ is then

$$u_2(t) = C_{11} \cos \sqrt{2}t + C_{12} \sin \sqrt{2}t + 2 \cos t$$

Differentiate it with respect to t .

$$u_2'(t) = -C_{11}\sqrt{2}\sin \sqrt{2}t + C_{12}\sqrt{2}\cos \sqrt{2}t - 2 \sin t$$

Apply the initial conditions to determine C_{11} and C_{12} .

$$\begin{aligned}u_2(0) &= C_{11} + 2 = 2 \\ u_2'(0) &= C_{12}\sqrt{2} = 0\end{aligned}$$

Solving this system of equations yields $C_{11} = 0$ and $C_{12} = 0$. Therefore,

$$u_2(t) = 2 \cos t$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Spring Two Mass System Higher Order Linear Differential Equations ]# ./
main

Two-spring, two-mass system solution k {1} = 3, k {2} = 2, m {1} = 1, m {2} = 1,
with u_{1}(0) = 1, u_{1}'(0) = 0, u_{2}(0) = 2, u_{2}'(0) = 0

The equation of motion :
df(u1[t],t,t)-2*u2+5*u1 = 0
df(u2[t],t,t)+2*u2-2*u1 = 0

u2 == 0.5*df(u1[t],t,t)+2.5*u1

The homogeneous ODE with constant coefficients:
0.5*df(u1[t],t,t,t,t)+3.5*df(u1[t],t,t)+3*u1[t] = 0

*****
*****          For u_{1}(t)          *****
*****

P:
(1,0)
(0,0)
(7,0)
(0,0)
(6,0)

P':
(4,0)
(0,0)
(14,0)
(0,0)

Initial conditions (u_{1} and u_{2}):
(1,0)
(0,0)
(2,0)
(0,0)

Initial conditions (u_{1}):
(1,0)
(0,0)
(-1,0)
(0,0)

```

Figure 3.47: The computation of the initial value problem general solution for the fourth order homogeneous differential equation that represent the two-spring, two-mass system for part (c) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Two Spring Two Mass System Higher Order Linear Differential Equations/main.cpp*).

```

*****
*****          For u_{1}(t)          *****
*****

P:
(1,0)
(0,0)
(7,0)
(0,0)
(6,0)

P':
(4,0)
(0,0)
(14,0)
(0,0)

Initial conditions (u_{1} and u_{2}):
(1,0)
(0,0)
(2,0)
(0,0)

Initial conditions (u_{1}):
(1,0)
(0,0)
(-1,0)
(0,0)

Initial guess for the roots (generated randomly):
(5.59957,0)
(2.82398,0)
(6.61124,0)
(3.93247,0)

Roots found at iteration: 26
*****

End of iteration

z {i} final:
(0,1)
(0,-1)
(0,2.44949)
(0,-2.44949)

The general solution is:
u_{1}(t) = c[0]*cos(t)+c[1]*sin(t)+c[2]*cos(2.44949*t)+c[3]*sin(2.44949*t)

The initial value problem solution is:
u_{1}(t) = cos(t)

The equation of motion 2 with u_{1}(t):
df(u2[t],t,t)+2*u2[t]-2*cos(t) = 0

```

Figure 3.48: The computation of the initial value problem general solution for the fourth order homogeneous differential equation that represent the two-spring, two-mass system for part (c) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Two Spring Two Mass System Higher Order Linear Differential Equations/main.cpp*).

```

*****
*****          For u_{2}(t)          *****
*****

P:
(1,0)
(0,0)
(2,0)

P':
(2,0)
(0,0)

Initial conditions (u_{2}):
(2,0)
(0,0)

Initial guess for the roots (generated randomly):
(5.19157,0)
(5.97163,0)

Roots found at iteration: 33
*****

End of iteration

z_{i} final:
(0,-1.41421)
(0,1.41421)

The general solution is:
u_{2}(t) = c[0]*cos(1.41421*t)+c[1]*sin(1.41421*t)+2*cos(t)

The initial value problem solution is:
u_{2}(t) = 2*cos(t)

Time taken by function: 200112 microseconds

```

Figure 3.49: The computation of the initial value problem general solution for the fourth order homogeneous differential equation that represent the two-spring, two-mass system for part (c) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Two Spring Two Mass System Higher Order Linear Differential Equations/main.cpp*).

(d) Suppose secondly that the initial conditions are

$$\begin{aligned} u_1(0) &= -2, & u_1'(0) &= 0 \\ u_2(0) &= 1, & u_2'(0) &= 0 \end{aligned}$$

From equation (3.236), $u_1''(0)$ and $u_1'''(0)$ can be determined.

$$u_1'' + 5u_1 = 2u_2 \quad \begin{cases} u_1''(0) = 2u_2(0) - 5u_1(0) = 12 \\ u_1'''(0) = 2u_2'(0) - 5u_1'(0) = 0 \end{cases}$$

Differentiate the general solution three times

$$\begin{aligned} u_1'(t) &= -C_5\sqrt{6}\sin\sqrt{6}t + C_6\sqrt{6}\cos\sqrt{6}t - C_7\sin t + C_8\cos t \\ u_1''(t) &= -6C_5\cos\sqrt{6}t - 6C_6\sin\sqrt{6}t - C_7\sin t - C_8\cos t \\ u_1'''(t) &= 6\sqrt{6}C_5\sin\sqrt{6}t - 6\sqrt{6}C_6\cos\sqrt{6}t + C_7\sin t - C_8\cos t \end{aligned}$$

Apply the initial conditions now to determine C_5, C_6, C_7 , and C_8 .

$$\begin{aligned} u_1(0) &= C_5 + C_7 = -2 \\ u_1'(0) &= C_6\sqrt{6} + C_8 = 0 \\ u_1''(0) &= -6C_5 - C_7 = 12 \\ u_1'''(0) &= -6\sqrt{6}C_6 - C_8 = 0 \end{aligned}$$

Solving this system of equations yields $C_5 = -2, C_6 = 0, C_7 = 0$, and $C_8 = 0$. Therefore,

$$u_1(t) = -2\cos\sqrt{6}t$$

Now that $u_1(t)$ is known, plug it back into equation (2) to find $u_2(t)$.

$$u_2'' + 2u_2 = -4 \cos \sqrt{6}t$$

This is a linear nonhomogeneous ODE, so its general solution can be expressed as a sum of the complementary solution and the particular solution.

$$u_2(t) = u_{2c}(t) + u_{2p}(t)$$

The complementary solution satisfies the associated homogeneous equation.

$$u_{2c}'' + 2u_{2c} = 0 \tag{3.240}$$

This is a homogeneous ODE with constant coefficients, so the solution is of the form $u_{2c} = e^{st}$.

$$u_{2c} = e^{st} \rightarrow u_{2c}' = se^{st} \rightarrow u_{2c}'' = s^2 e^{st}$$

Substitute these expressions into the ODE.

$$s^2 e^{st} + 2(e^{st}) = 0$$

To compute s , divide both sides by e^{st}

$$\begin{aligned} s^2 + 2 &= 0 \\ s &= \{-i\sqrt{2}, i\sqrt{2}\} \end{aligned}$$

As a result, two solutions to equation (3.240) are $u_{2c} = e^{-i\sqrt{2}t}$ and $u_{2c} = e^{i\sqrt{2}t}$. By the principle of superposition, the general solution for u_{2c} is a linear combination of these two.

$$\begin{aligned} u_{2c}(t) &= C_9 e^{-i\sqrt{2}t} + C_{10} e^{i\sqrt{2}t} \\ &= C_9 (\cos \sqrt{2}t - i \sin \sqrt{2}t) + C_{10} (\cos \sqrt{2}t + i \sin \sqrt{2}t) \\ &= (C_9 + C_{10}) \cos \sqrt{2}t + (-iC_9 + iC_{10}) \sin \sqrt{2}t \\ &= C_{11} \cos \sqrt{2}t + C_{12} \sin \sqrt{2}t \end{aligned}$$

On the other hand, the particular solution satisfies

$$u_{2p}'' + 2u_{2p} = -4 \cos \sqrt{6}t$$

Since the nonhomogeneous term is cosine and there are only even derivatives on the left, we use the trial solution $u_{2p} = A \cos \sqrt{6}t$. Plug this into the ODE to determine A .

$$\begin{aligned} (A \cos \sqrt{6}t)'' + 2(A \cos \sqrt{6}t) &= -4 \cos \sqrt{6}t \\ (-6A \cos \sqrt{6}t) + 2A \cos \sqrt{6}t &= -4 \cos \sqrt{6}t \\ -4A \cos \sqrt{6}t &= -4 \cos \sqrt{6}t \end{aligned}$$

Matching the coefficients, we see that $A = 1$, which means $u_{2p}(t) = \cos \sqrt{6}t$. The general solution for $u_2(t)$ is then

$$u_2(t) = C_{11} \cos \sqrt{2}t + C_{12} \sin \sqrt{2}t + \cos \sqrt{6}t$$

Differentiate it with respect to t .

$$u_2'(t) = -C_{11}\sqrt{2}\sin\sqrt{2}t + C_{12}\sqrt{2}\cos\sqrt{2}t - \sqrt{6}\sin\sqrt{6}t$$

Apply the initial conditions to determine C_{11} and C_{12} .

$$u_2(0) = C_{11} + 1 = 1$$

$$u_2'(0) = C_{12}\sqrt{2} = 0$$

Solving this system of equations yields $C_{11} = 0$ and $C_{12} = 0$. Therefore,

$$u_2(t) = \cos\sqrt{6}t$$

```
*****
*****                               For u_{1}(t)                               *****
*****

P:
(1,0)
(0,0)
(7,0)
(0,0)
(6,0)

P':
(4,0)
(0,0)
(14,0)
(0,0)

Initial conditions (u_{1} and u_{2}):
(-2,0)
(0,0)
(1,0)
(0,0)

Initial conditions (u_{1}):
(-2,0)
(0,0)
(12,0)
(0,0)

Initial guess for the roots (generated randomly):
(4.55549,0)
(3.71947,0)
(4.50372,0)
(4.74454,0)

Roots found at iteration: 29
```

Figure 3.50: The computation of the initial value problem general solution for the fourth order homogeneous differential equation that represent the two-spring, two-mass system for part (d) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Two Spring Two Mass System Higher Order Linear Differential Equations/main.cpp*).

```

*****
End of iteration

z_{i} final:
(0,2.44949)
(0,-2.44949)
(0,-1)
(0,1)

The general solution is:
u_{1}(t) = c[0]*cos(t)+c[1]*sin(t)+c[2]*cos(2.44949*t)+c[3]*sin(2.44949*t)

The initial value problem solution is:
u_{1}(t) = -2*cos(2.44949*t)

The equation of motion 2 with u_{1}(t):
df(u2[t],t,t)+2*u2[t]+4*cos(2.44949*t) = 0

*****
*****          For u_{2}(t)          *****
*****

P:
(1,0)
(0,0)
(2,0)

P':
(2,0)
(0,0)

Initial conditions (u_{2}):
(1,0)
(0,0)

Initial guess for the roots (generated randomly):
(5.0858,0)
(5.39588,0)

Roots found at iteration: 33

```

Figure 3.51: The computation of the initial value problem general solution for the fourth order homogeneous differential equation that represent the two-spring, two-mass system for part (d) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Two Spring Two Mass System Higher Order Linear Differential Equations/main.cpp*).

```

*****
*****          For u_{2}(t)          *****
*****
P:
(1,0)
(0,0)
(2,0)

P':
(2,0)
(0,0)

Initial conditions (u_{2}):
(1,0)
(0,0)

Initial guess for the roots (generated randomly):
(5.0858,0)
(5.39588,0)

Roots found at iteration: 33
*****

End of iteration

z_{i} final:
(0,1.41421)
(0,-1.41421)

The general solution is:
u_{2}(t) = c[0]*cos(1.41421*t)+c[1]*sin(1.41421*t)+1*cos(2.44949*t)

The initial value problem solution is:
u_{2}(t) = 1*cos(2.44949*t)

Time taken by function: 207694 microseconds

```

Figure 3.52: The computation of the initial value problem general solution for the fourth order homogeneous differential equation that represent the two-spring, two-mass system for part (d) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Two Spring Two Mass System Higher Order Linear Differential Equations/main.cpp*).

[SI*] The Method of Undetermined Coefficients

A particular solution Y of the nonhomogeneous n th order linear equation with constant coefficients

$$L[y] = a_0 y^{(n)} + a_1 y^{(n-1)} + \cdots + a_{n-1} y' + a_n y = g(t) \quad (3.241)$$

can be obtained by the method of undetermined coefficients, provided that $g(t)$ is of an appropriate form. Although the method of undetermined coefficients is not as general as the method of variation of parameters, it is usually much easier to use when it is applicable. Just as for the second order linear equation, when the constant coefficient linear differential operator L is applied to a polynomial $A_0 t^m + A_1 t^{m-1} + \cdots + A_m$, an exponential function $e^{\alpha t}$, a sine function $\sin \beta t$, or a cosine function $\cos \beta t$, the result is a polynomial, an exponential function, or a linear combination of sine and cosine functions, respectively.

The main difference in using this method for higher order equations stems from the fact that roots of the characteristic polynomial equation may have multiplicity greater than 2.

[SI*] Example:

Find the general solution of

$$y''' - 3y'' + 3y' - y = 4e^t \quad (3.242)$$

Solution:

The characteristic polynomial for the homogeneous equation corresponding to Eq. (3.242) is

$$r^3 - 3r^2 + 3r - 1 = (r - 1)^3$$

so the general solution of the homogeneous equation is

$$y_c(t) = c_1 e^t + c_2 t e^t + c_3 t^2 e^t \quad (3.243)$$

To find a particular solution $Y(t)$ of Eq. (3.242), we start by assuming that $Y(t) = Ae^t$. However, since e^t , te^t , and t^2e^t are all solutions of the homogeneous equation, we must multiply this initial choice by t^3 . Thus our final assumption is that $Y(t) = At^3e^t$, where A is an undetermined coefficient. To find the correct value for A , we differentiate $Y(t)$ three times, substitute for y and its derivatives in Eq. (3.242), and collect terms in the resulting equation. In this way we obtain

$$6Ae^t = 4e^t$$

Thus $A = \frac{2}{3}$ and the particular solution is

$$Y(t) = \frac{2}{3}t^3e^t \quad (3.244)$$

The general solution of Eq. (3.242) is the sum of $y_c(t)$ from Eq. (3.243) and $Y(t)$ from Eq. (3.244), that is

$$y = c_1e^t + c_2te^t + c_3t^2e^t + \frac{2}{3}t^3e^t$$

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Nonhomogeneous Higher Order Linear Differential
Equations General Solution with Method of Undetermined Coefficients ]# ./main

General solution for y''' - 3y'' + 3y' - y = 4 exp(t)

P:
(1,0)
(-3,0)
(3,0)
(-1,0)

P':
(3,0)
(-6,0)
(3,0)

Initial guess for the roots (generated randomly):
(3.34839,0)
(3.12548,0)
(3.94203,0)

Roots found at iteration: 25
*****

End of iteration

z {i} final:
(1,0)
(1,0)
(1,0)

Deleted Duplicate vector:
(1,0)

Final Duplicate vector:
(1,0)

The general solution of the homogeneous equation is:
y(t) = e^t*c[0]+e^t*c[1]*t+e^t*c[2]*t^2(2)

The general solution of the nonhomogeneous equation is:
Y(t) = 0.666667*e^t*t^3(3)

The general solution of the differential equation is:
Y(t) = e^t*c[0]+e^t*c[1]*t+e^t*c[2]*t^2+0.666667*e^t*t^3(3)

Time taken by function: 380497 microseconds
```

Figure 3.53: The computation of the general solution for the Eq. (3.242) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Nonhomogeneous Higher Order Linear Differential Equations General Solution with Method of Undetermined Coefficients/main.cpp*).

[SI*] **Example:**

Find the particular solution of the equation

$$y^{(4)} + 2y'' + y = 3 \sin t - 5 \cos t \quad (3.245)$$

Solution:

The general solution of the homogeneous equation is

$$y_c(t) = c_1 \cos t + c_2 \sin t + c_3 t \cos t + c_4 t \sin t$$

corresponding to the roots $r = i, i, -i$, and $-i$ of the characteristic equation. Our initial assumption for a particular solution is $Y(t) = A \sin t + B \cos t$, but we must multiply this choice by t^2 to make it different from all solutions of the homogeneous equation. Thus our final assumption is

$$Y(t) = At^2 \sin t + Bt^2 \cos t$$

Next, we differentiate $Y(t)$ four times, substitute into the differential equation (3.245), and collect terms, obtaining finally

$$-8A \sin t - 8B \cos t = 3 \sin t - 5 \cos t$$

Thus $A = -\frac{3}{8}$, $B = \frac{5}{8}$, and the particular solution of Eq. (3.245) is

$$Y(t) = -\frac{3}{8}t^2 \sin t + \frac{5}{8}t^2 \cos t \quad (3.246)$$

```

End of iteration

z {i} final:
(0,1)
(0,-1)
(0,1)
(0,-1)

Deleted Duplicate vector:
(0,-1)
(0,1)

Final root vector:
(0,1)
n_occurrence = 2

The general solution of the homogeneous equation is:
y(t) = c[0]*cos(t)+c[1]*sin(t)+c[2]*cos(t)*t+c[3]*sin(t)*t
*****

For rhs = 3*sin(t)
Y(t) particular = -8*A*cos(t)-8*A*sin(t)
Y(t) assumed = cos(t)*t^(2)+sin(t)*t^(2)

The general solution of this nonhomogeneous equation is:
Y_{i}(t) = -0.375*t^(2)*sin(t)
*****

For rhs = -5*cos(t)
Y(t) particular = -8*A*cos(t)-8*A*sin(t)
Y(t) assumed = cos(t)*t^(2)+sin(t)*t^(2)

The general solution of this nonhomogeneous equation is:
Y_{i}(t) = 0.625*t^(2)*cos(t)

The general solution of the differential equation is:
Y(t) = c[0]*cos(t)+c[1]*sin(t)+c[2]*cos(t)*t+c[3]*sin(t)*t-0.375*t^(2)*sin(t)+0.625*t^(2)*cos(t)

Time taken by function: 788510 microseconds

```

Figure 3.54: The computation of the general solution for the Eq. (3.245) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Nonhomogeneous Higher Order Linear Differential Equations General Solution with Method of Undetermined Coefficients/main.cpp*).

[SI*] The method of undetermined coefficients can be used whenever it is possible to guess the correct form for $Y(t)$. However, this is usually impossible for differential equations not having constant coefficients, or for nonhomogeneous terms other than the type described previously.

[SI*] **The Method of Variation of Parameters**

The method of variation of parameters for determining a particular solution of the nonhomogeneous n th order linear differential equation

$$L[y] = y^{(n)} = y^{(n)} + p_1(t)y^{(n-1)} + \cdots + p_{n-1}(t)y' + p_n(t)y = g(t) \quad (3.247)$$

is a direct extension of the method for the second order differential equation. As before, to use the method of variation of parameters, it is first necessary to solve the corresponding homogeneous differential equation. In general, this may be difficult unless the coefficients are constants. However, the method of variation of parameters is still more general than the method of undetermined coefficients in that it leads to an expression for the particular solution for any continuous function g , whereas the method of undetermined coefficients is restricted in practice to a limited class of functions g .

Suppose then that we know a fundamental set of solutions $y_1, y_2, \dots, y_n$ of the homogeneous equation. Then the general solution of the homogeneous equation is

$$y_c(t) = c_1y_1(t) + c_2y_2(t) + \cdots + c_ny_n(t) \quad (3.248)$$

The method of variation of parameters for determining a particular solution of Eq. (3.247) rests on the possibility of determining n functions $u_1, u_2, \dots, u_n$, such that $Y(t)$ is of the form

$$Y(t) = u_1(t)y_1(t) + u_2(t)y_2(t) + \cdots + u_n(t)y_n(t) \quad (3.249)$$

Since we have n functions to determine, we will have to specify n conditions. One of these is clearly that Y satisfy Eq. (3.247). The other $n - 1$ conditions are chosen so as to make the calculations as simple as possible. Since we hardly expect a simplification in determining Y if we must solve high order differential equations for $u_1, \dots, u_n$ it is natural to impose conditions to suppress the terms that lead to higher derivatives of $u_1, \dots, u_n$. From Eq. (3.249) we obtain

$$Y' = (u_1y_1' + u_2y_2' + \cdots + u_ny_n') + (u_1'y_1 + u_2'y_2 + \cdots + u_n'y_n) \quad (3.250)$$

where we have omitted the independent variable t on which each function in Eq. (3.250) depends. Thus the first condition that we impose is that

$$u_1'y_1 + u_2'y_2 + \cdots + u_n'y_n = 0 \quad (3.251)$$

Continuing this process in a similar manner through $n - 1$ derivatives of Y gives

$$Y^{(m)} = u_1y_1^{(m)} + u_2y_2^{(m)} + \cdots + u_ny_n^{(m)}, \quad m = 0, 1, 2, \dots, n - 1 \quad (3.252)$$

and the following $n - 1$ conditions on the functions $u_1, \dots, u_n$:

$$u_1'y_1^{(m-1)} + u_2'y_2^{(m-1)} + \cdots + u_n'y_n^{(m-1)}, \quad m = 1, 2, \dots, n - 1 \quad (3.253)$$

The n th derivative of Y is

$$Y^{(n)} = (u_1y_1^{(n)} + \cdots + u_ny_n^{(n)}) + (u_1'y_1^{(n-1)} + \cdots + u_n'y_n^{(n-1)}) \quad (3.254)$$

Finally, we impose the condition that Y must be a solution of Eq. (3.247). On substituting for the derivatives of Y from Eqs. (3.252) and (3.254), collecting terms, and making use of the fact that $L[y_i] = 0$ $i = 1, 2, \dots, n$, we obtain

$$u_1'y_1^{(n-1)} + u_2'y_2^{(n-1)} + \cdots + u_n'y_n^{(n-1)} = g \quad (3.255)$$

Equation (3.255), coupled with the $n - 1$ equations (3.253), gives n simultaneous linear nonhomogeneous algebraic equations for $u'_1, u'_2, \dots, u'_n$:

$$\begin{aligned} y_1 u'_1 + y_2 u'_2 + \cdots + y_n u'_n &= 0 \\ y'_1 u'_1 + y'_2 u'_2 + \cdots + y'_n u'_n &= 0 \\ y''_1 u'_1 + y''_2 u'_2 + \cdots + y''_n u'_n &= 0 \\ &\vdots \\ y_1^{(n-1)} u'_1 + y_2^{(n-1)} u'_2 + \cdots + y_n^{(n-1)} u'_n &= 0 \end{aligned} \quad (3.256)$$

The system (3.256) is a linear algebraic system for the unknown quantities $u'_1, \dots, u'_n$. By solving this system and then integrating the resulting expressions, you can obtain the coefficients $u_1, \dots, u_n$. A sufficient condition for the existence of a solution of the system of equations (3.256) is that the determinant of coefficients is nonzero for each value of t . However, the determinant of coefficients is precisely $W(y_1, y_2, \dots, y_n)$, and it is nowhere zero since $y_1, \dots, y_n$ is a fundamental set of solutions of the homogeneous equation. Hence it is possible to determine $u'_1, \dots, u'_n$. Using Cramer's rule, we can write the solution of the system of equations (3.256) in the form

$$u'_m(t) = \frac{g(t)W_m(t)}{W(t)}, \quad m = 1, 2, \dots, n \quad (3.257)$$

Here $W(t) = W(y_1, y_2, \dots, y_n)(t)$, and W_m is the determinant obtained from W by replacing the m th column by the column $(0, 0, \dots, 0, 1)$. With this notation a particular solution of Eq. (3.247) is given by

$$Y(t) = \sum_{m=1}^n y_m(t) \int_{t_0}^t \frac{g(s)W_m(s)}{W(s)} ds \quad (3.258)$$

where t_0 is arbitrary. Although the procedure is straightforward, the algebraic computations involved in determining $Y(t)$ from Eq. (3.258) become more and more complicated as n increases. In some cases the calculations may be simplified to some extent by using Abel's identity

$$W(t) = W(y_1, \dots, y_n)(t) = ce^{[-\int p_1(t) dt]}$$

The constant c can be determined by evaluating W at some convenient point.

[SI*] **Example:**

Given that $y_1(t) = e^t$, $y_2(t) = te^t$, and $y_3(t) = e^{-t}$ are solutions of the homogeneous equation corresponding to

$$y''' - y'' - y' + y = \sin(t) \quad (3.259)$$

determine a particular solution of Eq. (3.259) in terms of an integral.

Solution:

We use Eq. (3.258). First, we have

$$W(t) = W(e^t, te^t, e^{-t})(t) = \begin{vmatrix} e^t & te^t & e^{-t} \\ e^t & (t+1)e^t & -e^{-t} \\ e^t & (t+2)e^t & e^{-t} \end{vmatrix} = 4e^t$$

Next, we can use minors associated with the i th column for each $W_i(t)$

$$W_1(t) = \begin{vmatrix} 0 & te^t & e^{-t} \\ 0 & (t+1)e^t & -e^{-t} \\ 1 & (t+2)e^t & e^{-t} \end{vmatrix} = -2t - 1$$

$$W_2(t) = \begin{vmatrix} e^t & 0 & e^{-t} \\ e^t & 0 & -e^{-t} \\ e^t & 1 & e^{-t} \end{vmatrix} = 2$$

$$W_3(t) = \begin{vmatrix} e^t & te^t & 0 \\ e^t & (t+1)e^t & 0 \\ e^t & (t+2)e^t & 1 \end{vmatrix} = e^{2t}$$

Substituting these results in Eq. (3.258), we have

$$\begin{aligned} Y(t) &= e^t \int_{t_0}^t \frac{\sin(s)(-1-2s)}{4e^s} ds + te^t \int_{t_0}^t \frac{\sin(s)(2)}{4e^s} ds + e^{-t} \int_{t_0}^t \frac{\sin(s)(e^{2s})}{4e^s} ds \\ &= \frac{1}{4} \int_{t_0}^t \{e^{t-s}[-1+2(t-s)] + e^{-(t-s)}\} \sin(s) ds \end{aligned}$$

```
General solution for y''' - y'' - y' + y = sin(t)

P:
(1,0)
(-1,0)
(-1,0)
(1,0)

P':
(3,0)
(-2,0)
(-1,0)

Initial guess for the roots (generated randomly):
(5.492,0)
(4.59702,0)
(4.358,0)

Roots found at iteration: 22
*****

End of iteration

z_{i} final:
(1,0)
(-1,0)
(1,0)

Sorted Duplicate vector:
(1,0)
(1,0)

Final root vector:
(-1,0)
(1,0)

The general solution of the homogeneous equation is:
y(t) = e^(-t)*c[0]+e^t*c[1]+e^t*c[2]*t
```

Figure 3.55: The computation of the particular solution for the Eq. (3.259) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Nonhomogeneous Higher Order Linear Differential Equations General Solution with Variation of Parameters/main.cpp*).

```

The general solution of the homogeneous equation is:
y(t) = e^(-t)*c[0]+e^t*c[1]+e^t*c[2]*t

W(t):
[ e^(-t)      e^t      e^t*t ]
[ -e^(-t)     e^t      e^t*t+e^t ]
[  e^(-t)     e^t      e^t*t+2*e^t ]

det (W(t)) = 4*e^t

W_{i}(t):
[ 0      e^t      e^t*t ]
[ 0      e^t      e^t*t+e^t ]
[ 1      e^t      e^t*t+2*e^t ]

det (W_{i}(t)):
e^(2*t)

W_{i}(t):
[ e^(-t)      0      e^t*t ]
[ -e^(-t)     0      e^t*t+e^t ]
[  e^(-t)     1      e^t*t+2*e^t ]

det (W_{i}(t)):
-2*t-1

W_{i}(t):
[ e^(-t)  e^t  0 ]
[ -e^(-t) e^t  0 ]
[  e^(-t) e^t  1 ]

det (W_{i}(t)):
2

The particular solution of the differential equation is:
Y(t) = e^(-t)*(int(0.25*e^t*sin(t),t))+e^t*(int(-0.5*t*sin(t)*e^(-t),t))+e^t*(int(-0.25*sin(t)*e^(-t),t))+e^t*t*(int(0.5*sin(t)*e^(-t),t))

Time taken by function: 4659706 microseconds

```

Figure 3.56: The computation of the particular solution for the Eq. (3.259) (*SymIntegration/Examples/Differential Equations/Test SymIntegration Nonhomogeneous Higher Order Linear Differential Equations General Solution with Variation of Parameters/main.cpp*).

i. Functions in SymIntegration Related to Higher Order Linear Equations

[SI*] We will list the basic functions that can be used for solving higher order linear differential equations:

AberthEhrlich(const vector<complex<double>> & $\vec{P}$, const vector<complex<double>> & x_0 , int N)

to compute the root of higher order polynomial P with Aberth-Ehrlich formula, P is a vector of size n with complex entry, the configuration is like this:

$$\vec{P} = \begin{bmatrix} (a_1, b_1) \\ (a_2, b_2) \\ \vdots \\ (a_n, b_n) \end{bmatrix}$$

$$\text{polynomial}(\vec{P}) = (a_1, b_1)r^{n-1} + (a_2, b_2)r^{n-2} + \dots + (a_n, b_n)$$

so if we want to compute a polynomial of order 6, then $\vec{P}$ has to be a size of 7. N is the number of maximum iteration for the Abert-Ehrlich method, and x_0 is the initial guess / approximation vector with complex entry and has to be a size of $n - 1$, the entry of x_0 is preferable to be distinct and in decimal (e.g. 1.23 or 2.55), not round number to avoid error due to rounding.

This function is written in **src/numericalmethod.cpp** since it is a numerical method, but very useful in Differential Equations field and application thus we write it here.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Aberth-Ehrlich Method to Compute Roots of Polynomial ]# ./main
Polynomial : r^(4)+r^(3)-7*r^(2)-r+6
P:
(1,0)
(1,0)
(-7,0)
(-1,0)
(6,0)

P':
(4,0)
(3,0)
(-14,0)
(-1,0)

Initial guess:
(1,0)
(2,0)
(3,0)
(4,0)

iteration: 0

z {i} new:
(1,0)
(2,0)
(2.29412,0)
(8.10931,0)

iteration: 1

z {i} new:
(1,0)
(2,0)
(0.789054,0)
(-7.71719,0)

iteration: 2

z {i} new:
(1,0)
(2,0)
(-0.628758,0)
(-3.16734,-1.77748e-15)

iteration: 3

z {i} new:
(1,0)
(2,1.49372e-47)
(-0.996208,-4.54674e-17)
(-3.00002,-5.32888e-16)

```

Figure 3.57: The root-finding computation with Aberth-Ehrlich method (*SymIntegration/Examples/Numerical Methods/Test SymIntegration Aberth-Ehrlich Method to Compute Roots of Polynomial/main.cpp*).

```

iteration: 4
z {i} new:
(1,0)
(2,1.47984e-48)
(-1,7.65354e-17)
(-3,-1.65491e-16)

iteration: 5
z {i} new:
(1,0)
(2,-4.2281e-49)
(-1,-4.59293e-17)
(-3,2.01903e-16)

iteration: 6
z {i} new:
(1,0)
(2,-1.66768e-49)
(-1,7.65354e-17)
(-3,-1.65491e-16)

Roots found at iteration: 6
*****

End of iteration
z {i} final:
(1,0)
(2,0)
(-1,0)
(-3,0)

Time taken by function: 9088 microseconds

```

Figure 3.58: The root-finding computation with Aberth-Ehrlich method (*SymIntegration/Examples/Numerical Methods/Test SymIntegration Aberth-Ehrlich Method to Compute Roots of Polynomial-main.cpp*).

higherorderlineardiffeq_homogeneousequationsivpsolution(const vector<complex<double>> & $\vec{P}$, const vector<complex<double>> & $\vec{ic}$, const vector<complex<double>> & $\vec{x}_0$, int N)
 this function computes the general solution and the initial value solution for higher order homogeneous linear differential equation (the order of the differential equation is n th order)

$$(a_1, b_1)y^{(n)} + (a_2, b_2)y^{(n-1)} + \dots + (a_n, b_n)y' + (a_{n+1}, b_{n+1})y = 0$$

the input $\vec{P}$ is the coefficients vector of the characteristic equation representing the differential equation

$$\vec{P} = \begin{bmatrix} (a_1, b_1) \\ (a_2, b_2) \\ \vdots \\ (a_n, b_n) \\ (a_{n+1}, b_{n+1}) \end{bmatrix}$$

it is a vector with size of $n + 1$, as we can see the characteristic equation from $\vec{P}$ is

$$\text{polynomial}(\vec{P}) = (a_1, b_1)r^{n-1} + (a_2, b_2)r^{n-2} + \dots + (a_n, b_n)r + (a_{n+1}, b_{n+1})$$

usually we will deal with the condition where $b_1, b_2, \dots, b_n, b_{n+1} = 0$ (the coefficients are real number). The input has to be in complex form because there is a possibility to obtain complex roots.

Then we will have another input $\vec{ic}$ that represents the initial condition with this configura-

tion:

$$\begin{aligned} y(0) &= ic_1 \\ y'(0) &= ic_2 \\ &\vdots \\ y^{(n-1)}(0) &= ic_n \end{aligned}$$

thus

$$\vec{ic} = \begin{bmatrix} ic_1 \\ ic_2 \\ \vdots \\ ic_n \end{bmatrix}$$

it is a vector with size of n , and the input of $\vec{ic}$ is also in complex form. The last two parameters are N , the number of maximum iteration for the Abert-Ehrlich method in order to compute the roots of the characteristic equation, and $\vec{x}_0$ is the initial guess / approximation vector with complex entry, it has to be a size of n , the entry of $\vec{x}_0$ is preferable to be distinct and in decimal (e.g. 1.23 or 2.55), not round number to avoid error due to rounding. The last two parameters can be omitted and automated in the main source code, with Mersenne Twister engine, we can generate a good $\vec{x}_0$ that can approximate the roots.

higherorderlineardiffeq_homogeneousequationsivpsolution(const vector<complex<double>> & $\vec{P}$, const vector<complex<double>> & $\vec{ic}$)

this function produces the the same output with the previous function but with less input, this function computes the general solution and the initial value solution for higher order homogeneous linear differential equation (the order of the differential equation is n th order)

$$(a_1, b_1)y^{(n)} + (a_2, b_2)y^{(n-1)} + \cdots + (a_n, b_n)y' + (a_{n+1}, b_{n+1})y = 0$$

the input $\vec{P}$ is the coefficients vector of the characteristic equation representing the differential equation

$$\vec{P} = \begin{bmatrix} (a_1, b_1) \\ (a_2, b_2) \\ \vdots \\ (a_n, b_n) \\ (a_{n+1}, b_{n+1}) \end{bmatrix}$$

it is a vector with size of $n + 1$, as we can see the characteristic equation from $\vec{P}$ is

$$\text{polynomial}(\vec{P}) = (a_1, b_1)r^{n-1} + (a_2, b_2)r^{n-2} + \cdots + (a_n, b_n)r + (a_{n+1}, b_{n+1})$$

usually we will deal with the condition where $b_1, b_2, \dots, b_n, b_{n+1} = 0$ (the coefficients are real number). The input has to be in complex form because there is a possibility to obtain complex roots.

Then we will have another input $\vec{ic}$ that represents the initial condition with this configura-

tion:

$$\begin{aligned} y(0) &= ic_1 \\ y'(0) &= ic_2 \\ &\vdots \\ y^{(n-1)}(0) &= ic_n \end{aligned}$$

thus

$$\vec{ic} = \begin{bmatrix} ic_1 \\ ic_2 \\ \vdots \\ ic_n \end{bmatrix}$$

it is a vector with size of n , and the input of $\vec{ic}$ is also in complex form. This function omitted $\vec{x}_0$ (we generate it with Mersenne Twister that is normally distributed with $\mu = 5$ and $\sigma = 1.2$), and it also set $N = 100$ in the source code **src/dsolve.cpp**.

```
P:
(1,0)
(-3,0)
(2,0)
(-6,0)
(1,0)
(-3,0)

P':
(5,0)
(-12,0)
(6,0)
(-12,0)
(1,0)

Initial guess:
(1,1,0)
(2,0)
(3,0)
(4,0)
(5,1,0)

*****

End of iteration

z {i} final:
(0,1)
(0,-1)
(3,0)
(0,-1)
(0,1)

Unique vector:
(3,0)

Duplicate vector:
(0,1)
(0,-1)
(0,-1)
(0,1)
```

Figure 3.59: The computation for the initial value problem of $y^{(5)} - 3y^{(4)} + 2y^{(3)} - 6y'' + y' - 3$ with initial condition $y(0) = 1, y'(0) = 0, y''(0) = -2, y^{(3)}(0) = -1, y^{(4)}(0) = 3$ (**SymIntegration/Examples/Numerical Methods/Test SymIntegration IVP Solution for Higher Order Linear Differential Equations/main.cpp**).

```

*****
End of iteration

z {i} final:
(0,1)
(0,-1)
(3,0)
(0,-1)
(0,1)

Unique vector:
(3,0)

Duplicate vector:
(0,1)
(0,-1)
(0,-1)
(0,1)

Sorted Duplicate vector:
(0,-1)
(0,-1)
(0,1)
(0,1)

Deleted Duplicate vector:
(0,-1)
(0,1)

Final Duplicate vector:
(3,0)
(0,1)

The general solution is:
y(t) = e^(3*t)*c[0]+c[1]*cos(t)+c[2]*sin(t)+c[3]*cos(t)*t+c[4]*sin(t)*t
n_occurrence = 2

The initial value problem solution is:
y(t) = 1*cos(t)-0.5*sin(t)+0.5*cos(t)*t-0.5*sin(t)*t

Time taken by function: 351427 microseconds

```

Figure 3.60: The computation for the initial value problem of $y^{(5)} - 3y^{(4)} + 2y^{(3)} - 6y'' + y' - 3$ with initial condition $y(0) = 1, y'(0) = 0, y''(0) = -2, y^{(3)}(0) = -1, y^{(4)}(0) = 3$ (*SymIntegration/Examples/Numerical Methods/Test SymIntegration IVP Solution for Higher Order Linear Differential Equations/main.cpp*).

higherorderlineardiffeq_vectorize(const Symbolic &F(y,t), const Symbolic &y, const Symbolic &t, int n)

this function takes the coefficient of the ordinary differential equation with order of n , with this form:

$$F(y,t) = a_0 y^{(n)} + a_1 y^{(n-1)} + \cdots + a_{n-1} y' + a_n y = 0$$

with y as the function of t , $y(t)$.

It will return this complex vector (as the final result):

$$P = \begin{bmatrix} (a_0, 0) \\ (a_1, 0) \\ \vdots \\ (a_{n-1}, 0) \\ (a_n, 0) \end{bmatrix}$$

```

The ODE with constant coefficients:
1/2*df(u1[t],t,t,t,t)+7/2*df(u1[t],t,t)+3*u1[t] = 0

The vector coefficient of homogeneous ODE with constant coefficients:
(1,0)
(0,0)
(7,0)
(0,0)
(6,0)

Time taken by function: 96071 microseconds

```

Figure 3.61: The vectorization of an ODE (*SymIntegration/Examples/Numerical Methods/Test SymIntegration Vectorize ODE/main.cpp*).

higherorderlineardiffeq_twospringtwomasssystem(double k_1 , double k_2 , double m_1 , double m_2 , Symbolic & t , const vector<complex<double>> & $\vec{ic}$)

this function computes the equation of motion for two-spring, two-mass system, and the initial value problem solution with given input of m_1 as the mass for the first box / object, m_2 as the mass for the second box / object, k_1 is the spring constant between the ceiling / wall with the first box, and k_2 is the spring constant between the first and second box. Lastly, t is the independent variable that represents time. The solution will be given as $u_1(t)$ and $u_2(t)$.

Then we will have another input $\vec{ic}$ that represents the initial condition with this configuration:

$$\begin{aligned} u_1(0) &= ic_1 \\ u_1'(0) &= ic_2 \\ u_2(0) &= ic_3 \\ u_2'(0) &= ic_4 \end{aligned}$$

thus

$$\vec{ic} = \begin{bmatrix} ic_1 \\ ic_2 \\ ic_3 \\ ic_4 \end{bmatrix}$$

it is a vector with size of $n = 4$, it is the order of the differential equation for the system after we solve the equations of motion, the input of $\vec{ic}$ is in complex form.

higherorderlineardiffeq_nonhomogeneousequationsgeneralsolution(const vector<complex<double>> & $\vec{P}$, const Symbolic & $g(t)$, const Symbolic & t)

this function computes the general solution of a nonhomogeneous higher order linear differential equation with this form

$$a_0 y^{(n)}(t) + a_1 y^{(n-1)}(t) + \cdots + a_{n-1} y'(t) + a_n y(t) = g(t)$$

with

$$\vec{P} = \begin{bmatrix} (a_1, b_1) \\ (a_2, b_2) \\ \vdots \\ (a_n, b_n) \\ (a_{n+1}, b_{n+1}) \end{bmatrix}$$

as a vector with size of $n + 1$ that represents the differential equation of order n , $b_1, b_2, \dots, b_{n+1}$ will have value of zero.

The function $g(t)$ is the right hand side function, it is a single function only, not a linear combination of functions.

Currently, it can handle several types of $g(t)$ as listed below:

$$\begin{aligned} g(t) &= ae^t \\ g(t) &= a \sin(t) \\ g(t) &= a \cos(t) \end{aligned}$$

with a as constant.

The code that handle or recognize certain $g(t)$ function can be made with **if** conditional, here is a snippet of what is already written

```
void higherorderlineardiffeq_nonhomogeneousequationsgeneralsolution(const
    vector<complex<double>> &P, const Symbolic &rhs_function, const Symbolic
    &t)
{
    ...
    ...

    Symbolic A_solve;
    Symbolic F = rhs_function;
    cout << "\nY(t) particular = "<< Yt_particular_total << endl;
    cout << "\nY(t) assumed = "<< Yt_assumed << endl;

    // Case 1 : g(t) = a*exp(t)
    if (df(F,t,0) == df(F,t,1) && df(F,t,1) == df(F,t,2) && df(F,t,0) ==
        df(F,t,2)) // meaning that the rhs_function is in the form of a*
        exp(t)
    {
        //double a = df(F,t,0)/exp(t);
        Symbolic lhs_nh = Yt_particular_total.coeff(exp(t),1);
        A_solve = solve(lhs_nh*exp(t)- rhs_function, A).front().rhs;
        Yt_final = A_solve*Yt_assumed ;
    }
    // Case 2 : g(t) = a*sin(t) or g(t) = a*cos(t)
    if (df(F,t,0) == -df(F,t,2) && df(F,t,0) != df(F,t,1) && df(F,t,0) ==
        df(F,t,4)) // meaning that the rhs_function is in the form of a*
        sin(t) or a* cos(t)
    {
        if (df(F/sin(t),t) == 0) // meaning that the rhs_function is in the
            form of a*sin(t)
        {
            double a = F/sin(t);
            Symbolic lhs_nh = Yt_particular_total.coeff(A*sin(t),1);
            A_solve = a/lhs_nh;
            Yt_final = A_solve*Yt_assumed.coeff(sin(t),1)*sin(t) ;
        }
        if (df(F/cos(t),t) == 0) // meaning that the rhs_function is in the
            form of a*cos(t)
        {
            double a = F/cos(t);
            Symbolic lhs_nh = Yt_particular_total.coeff(A*cos(t),1);
            A_solve = a/lhs_nh;
            Yt_final = A_solve*Yt_assumed.coeff(cos(t),1)*cos(t) ;
        }
    }

    ...
    ...
}
```

```
}
```

Code 41: *src/dsolve.cpp*

higherorderlineardiffeq_nonhomogeneousequationsgeneralsolution(const vector<complex<double>> & $\vec{P}$, const SymbolicMatrix & A , const Symbolic & t)

this function computes the general solution of a nonhomogeneous higher order linear differential equation with this form

$$a_0 y^{(n)}(t) + a_1 y^{(n-1)}(t) + \cdots + a_{n-1} y'(t) + a_n y(t) = A$$

with

$$\vec{P} = \begin{bmatrix} (a_1, b_1) \\ (a_2, b_2) \\ \vdots \\ (a_n, b_n) \\ (a_{n+1}, b_{n+1}) \end{bmatrix}$$

as a vector with size of $n + 1$ that represents the differential equation of order n , $b_1, b_2, \dots, b_{n+1}$ will have value of zero.

The matrix A acts as the right hand side, a linear combination of functions that is represented as a matrix with 1 column, each row represent unique function, for example:

$$A = \begin{bmatrix} 3 \sin t \\ -5 \cos t \end{bmatrix} = 3 \sin t - 5 \cos t$$

higherorderlineardiffeq_nonhomogeneousequations_variationofparameters(const vector<complex<double>> & $\vec{P}$, const Symbolic & $g(t)$, const Symbolic & t)

this function use the method of variation of parameters to compute the particular solution of a nonhomogeneous higher order linear differential equation with this form

$$a_0 y^{(n)}(t) + a_1 y^{(n-1)}(t) + \cdots + a_{n-1} y'(t) + a_n y(t) = g(t)$$

with

$$\vec{P} = \begin{bmatrix} (a_1, b_1) \\ (a_2, b_2) \\ \vdots \\ (a_n, b_n) \\ (a_{n+1}, b_{n+1}) \end{bmatrix}$$

as a vector with size of $n + 1$ that represents the differential equation of order n , $b_1, b_2, \dots, b_{n+1}$ will have value of zero.

IV. SERIES SOLUTIONS OF SECOND ORDER LINEAR EQUATIONS

[SI*] **Power Series**

Finding the general solution of a linear differential equation depends on determining a fundamental set of solutions of the homogeneous equation. We will discuss the use of power series to construct fundamental sets of solutions of second order linear differential equations whose coefficients are functions of the independent variable.

1. A power series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ is said to converge at a point x if

$$\lim_{m \rightarrow \infty} \sum_{n=0}^m a_n(x - x_0)^n$$

exists for that x . The series certainly converges for $x = x_0$; it may converge for all x , or it may converge for some values of x and not for others.

2. The series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ is said to converge absolutely at a point x if the series

$$\sum_{n=0}^{\infty} |a_n(x - x_0)|^n = \sum_{n=0}^{\infty} |a_n| |x - x_0|^n$$

converges. It can be shown that if the series converges absolutely, then the series also converges; however, the converse is not necessarily true.

3. One of the most useful tests for the absolute convergence of a power series is the ratio test. If $a_n \neq 0$, and if, for a fixed value of x ,

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}(x - x_0)^{n+1}}{a_n(x - x_0)^n} \right| = |x - x_0| \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = |x - x_0|L$$

then the power series converges absolutely at that value of x if $|x - x_0|L < 1$ and diverges if $|x - x_0|L > 1$. If $|x - x_0|L = 1$, the test is inconclusive.

4. If the power series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ converges at $x = x_1$, it converges absolutely for $|x - x_0| < |x_1 - x_0|$; and if it diverges at $x = x_1$, it diverges for $|x - x_0| > |x_1 - x_0|$.
5. There is a nonnegative number ρ , called the radius of convergence, such that $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ converges absolutely for $|x - x_0| < \rho$ and diverges for $|x - x_0| > \rho$. For a series that converges only at x_0 , we define ρ to be zero; for a series that converges for all x ; we say that ρ is infinite. If $\rho > 0$, then the interval $|x - x_0| < \rho$ is called the interval of convergence. The series may either converge or diverge when $|x - x_0| = \rho$.
6. Suppose that $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ and $\sum_{n=0}^{\infty} b_n(x - x_0)^n$ converge to $f(x)$ and $g(x)$, respectively, for $|x - x_0| < \rho, \rho > 0$.

The series can be added or subtracted termwise, and

$$f(x) \pm g(x) = \sum_{n=0}^{\infty} (a_n \pm b_n)(x - x_0)^n$$

the resulting series converges at least for $|x - x_0| < \rho$.

7. Suppose that $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ and $\sum_{n=0}^{\infty} b_n(x - x_0)^n$ converge to $f(x)$ and $g(x)$, respectively, for $|x - x_0| < \rho, \rho > 0$.

The series can be formally multiplied, and

$$f(x)g(x) = \left[\sum_{n=0}^{\infty} a_n(x - x_0)^n \right] \left[\sum_{n=0}^{\infty} b_n(x - x_0)^n \right] = \sum_{n=0}^{\infty} c_n(x - x_0)^n$$

where $c_n = a_0b_n + a_1b_{n-1} + \cdots + a_nb_0$. The resulting series converges at least for $|x - x_0| < \rho$.

Further, if $g(x) \neq 0$, the series can be formally divided, and

$$\frac{f(x)}{g(x)} = \sum_{n=0}^{\infty} d_n(x - x_0)^n$$

In most cases the coefficients d_n can be most easily obtained by equating coefficients in the equivalent relation

$$\begin{aligned} \sum_{n=0}^{\infty} a_n(x - x_0)^n &= \left[\sum_{n=0}^{\infty} d_n(x - x_0)^n \right] \left[\sum_{n=0}^{\infty} b_n(x - x_0)^n \right] \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n d_k b_{n-k} \right) (x - x_0)^n \end{aligned}$$

In the case of division the radius of convergence of the resulting power series may be less than ρ .

8. The function f is continuous and has derivatives of all orders for $|x - x_0| < \rho$. Further, $f', f'', \dots$ can be computed by differentiating the series termwise; that is,

$$\begin{aligned} f'(x) &= a_1 + 2a_2(x - x_0) + \cdots + na_n(x - x_0)^{n-1} + \cdots \\ &= \sum_{n=1}^{\infty} na_n(x - x_0)^{n-1} \\ f''(x) &= 2a_2 + 6a_3(x - x_0) + \cdots + n(n-1)a_n(x - x_0)^{n-2} + \cdots \\ &= \sum_{n=2}^{\infty} n(n-1)a_n(x - x_0)^{n-2} \end{aligned}$$

and so forth, and each of the series converges absolutely for $x - x_0 < \rho$.

9. The value of a_n is given by

$$a_n = \frac{f^{(n)}(x_0)}{n!}$$

The series is called the Taylor' series for the function f about $x = x_0$.

10. If $\sum_{n=0}^{\infty} a_n(x - x_0)^n = \sum_{n=0}^{\infty} b_n(x - x_0)^n$ for each x in some open interval with center x_0 , then $a_n = b_n$ for $n = 0, 1, 2, 3, \dots$. In particular, if $\sum_{n=0}^{\infty} a_n(x - x_0)^n = 0$ for each such x , then $a_0 = a_1 = \cdots = a_n = \cdots = 0$.

[SI*] A function f that has a Taylor series expansion about $x = x_0$

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

with a radius of convergence $\rho > 0$, is said to be analytic at $x = x_0$. All of the familiar functions of calculus are analytic except perhaps at certain easily recognized points. For example, $\sin x$ and e^x are analytic everywhere except at $x = 0$, and $\tan x$ is analytic except at odd multiples of $\frac{\pi}{2}$.

In many respects the natural context for the use of power series is the complex plane.

[SI*] **Example:**

Determine the radius of convergence of the given power series

(a)

$$\sum_{n=0}^{\infty} (x-3)^n$$

(b)

$$\sum_{n=0}^{\infty} \frac{n}{2^n} x^n$$

(c)

$$\sum_{n=0}^{\infty} \frac{x^{2n}}{n!}$$

(d)

$$\sum_{n=0}^{\infty} 2^n x^n$$

(e)

$$\sum_{n=1}^{\infty} \frac{(2x+1)^n}{n^2}$$

(f)

$$\sum_{n=1}^{\infty} \frac{(x-x_0)^n}{n^2}$$

(g)

$$\sum_{n=1}^{\infty} \frac{(-1)^n n^2 (x+2)^n}{4^n}$$

(h)

$$\sum_{n=1}^{\infty} \frac{n! x^n}{n^n}$$

Solution:

(a) Apply the ratio test to determine the condition in which the provided series converges.

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{(x-3)^{n+1}}{(x-3)^n} \right| \\ &= \lim_{n \rightarrow \infty} |x-3| \\ &= |x-3| \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } |x-3| < 1 \\ \text{unknown if } |x-3| = 1 \\ \text{divergent if } |x-3| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $-1 < x-3 < 1$, or $2 < x < 4$, we see that the center of convergence is at $x = 3$ and the radius of convergence is 1.

(b) Apply the ratio test to determine the condition in which the provided series converges.

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{n+1}{2^{n+1}} x^{n+1}}{\frac{n}{2^n} x^n} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{1}{2} \frac{n+1}{n} x \right| \\
 &= \lim_{n \rightarrow \infty} \frac{1}{2} \frac{n+1}{n} |x| \\
 &= \frac{1}{2} \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right) |x| \\
 &= \frac{1}{2} |x|
 \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } \frac{1}{2}|x| < 1 \\ \text{unknown if } \frac{1}{2}|x| = 1 \\ \text{divergent if } \frac{1}{2}|x| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $|x| < 2$, or $-2 < x < 2$, we see that the center of convergence is at $x = 0$ and the radius of convergence is 2.

(c) Apply the ratio test to determine the condition in which the provided series converges.

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{x^{2(n+1)}}{(n+1)!}}{\frac{x^{2n}}{n!}} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{n!}{(n+1)!} \frac{x^{2n+2}}{x^{2n}} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{1}{n+1} x^2 \right| \\
 &= \lim_{n \rightarrow \infty} \frac{1}{n+1} x^2 \\
 &= 0x^2 \\
 &= 0
 \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } 0x^2 < 1 \\ \text{unknown if } 0x^2 = 1 \\ \text{divergent if } 0x^2 > 1 \end{cases}$$

From the condition of convergence, which can also be written as $|x| < \sqrt{\frac{1}{0}} = \infty$, or $-\infty < x < \infty$, we see that the center of convergence is at $x = 0$ and the radius of convergence is ∞ .

(d) Apply the ratio test to determine the condition in which the provided series converges.

$$\begin{aligned}\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{2^{n+1}x^{n+1}}{2^n x^n} \right| \\ &= \lim_{n \rightarrow \infty} |2x| \\ &= \lim_{n \rightarrow \infty} 2|x| \\ &= 2|x|\end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } 2|x| < 1 \\ \text{unknown if } 2|x| = 1 \\ \text{divergent if } 2|x| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $|x| < \frac{1}{2}$, or $-\frac{1}{2} < x < \frac{1}{2}$, we see that the center of convergence is at $x = 0$ and the radius of convergence is $\frac{1}{2}$.

(e) Apply the ratio test to determine the condition in which the provided series converges.

$$\begin{aligned}\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{(2x+1)^{n+1}}{(n+1)^2}}{\frac{(2x+1)^n}{n^2}} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{n^2}{(n+1)^2} (2x+1) \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{n^2}{n^2 \left(1 + \frac{1}{n}\right)^2} (2x+1) \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{1}{\left(1 + \frac{1}{n}\right)^2} (2x+1) \right| \\ &= \lim_{n \rightarrow \infty} \frac{1}{\left(1 + \frac{1}{n}\right)^2} |2x+1| \\ &= |2x+1|\end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } |2x+1| < 1 \\ \text{unknown if } |2x+1| = 1 \\ \text{divergent if } |2x+1| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $-1 < 2x+1 < 1$, or $-1 < x < 0$, we see that the center of convergence is at $x = -\frac{1}{2}$ and the radius of convergence is $\frac{1}{2}$.

(f) Apply the ratio test to determine the condition in which the provided series converges.

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{(x-x_0)^{n+1}}{(n+1)^2}}{\frac{(x-x_0)^n}{n^2}} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{n^2}{(n+1)^2} (x-x_0) \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{n^2}{n^2 \left(1 + \frac{1}{n}\right)^2} (x-x_0) \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{1}{\left(1 + \frac{1}{n}\right)^2} (x-x_0) \right| \\
 &= \lim_{n \rightarrow \infty} \frac{1}{\left(1 + \frac{1}{n}\right)^2} |x-x_0| \\
 &= |x-x_0|
 \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } |x-x_0| < 1 \\ \text{unknown if } |x-x_0| = 1 \\ \text{divergent if } |x-x_0| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $-1 < x-x_0 < 1$, or $-1+x_0 < x < 1+x_0$, we see that the center of convergence is at $x = x_0$ and the radius of convergence is 1.

(g) Apply the ratio test to determine the condition in which the provided series converges.

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+1}(n+1)^2(x+2)^{n+1}}{4^{n+1}}}{\frac{(-1)^n n^2 (x+2)^n}{4^n}} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1}}{(-1)^n} \frac{(n+1)^2}{n^2} \frac{4^n}{4^{n+1}} \frac{(x+2)^{n+1}}{(x+2)^n} \right| \\
 &= \lim_{n \rightarrow \infty} \left| (-1) \frac{n^2 \left(1 + \frac{1}{n}\right)^2}{n^2} \frac{1}{4} (x+2) \right| \\
 &= \lim_{n \rightarrow \infty} \left| (-1) \left(1 + \frac{1}{n}\right)^2 \frac{1}{4} (x+2) \right| \\
 &= \lim_{n \rightarrow \infty} \frac{1}{4} \left(1 + \frac{1}{n}\right)^2 |x+2| \\
 &= \frac{1}{4} |x+2|
 \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } \frac{1}{4}|x+2| < 1 \\ \text{unknown if } \frac{1}{4}|x+2| = 1 \\ \text{divergent if } \frac{1}{4}|x+2| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $|x+2| < 4$, or $-6 < x < 2$, we see that the center of convergence is at $x = -2$ and the radius of convergence is 4.

(h) Apply the ratio test to determine the condition in which the provided series converges.

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{(n+1)!x^{n+1}}{(n+1)^{n+1}}}{\frac{n!x^n}{n^n}} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{(n+1)!}{n!} \cdot \frac{n^n}{(n+1)^{n+1}} \cdot \frac{x^{n+1}}{x^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| (n+1) \frac{n^n}{n^{n+1} \left(1 + \frac{1}{n}\right)^{n+1}} x \right| \\ &= \lim_{n \rightarrow \infty} \left| (n+1) \frac{1}{n \left(1 + \frac{1}{n}\right)^{n+1}} x \right| \\ &= \lim_{n \rightarrow \infty} \frac{n+1}{n} \frac{1}{\left(1 + \frac{1}{n}\right)^{n+1}} |x| \\ &= \left(\lim_{n \rightarrow \infty} \frac{n+1}{n} \right) \left[\lim_{n \rightarrow \infty} \frac{1}{\left(1 + \frac{1}{n}\right)^{n+1}} \right] |x| \\ &= \left(\lim_{n \rightarrow \infty} \frac{1 + \frac{1}{n}}{1} \right) \left(\frac{1}{e} \right) |x| \\ &= (1) \left(\frac{1}{e} \right) |x| \\ &= \frac{1}{e} |x| \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } \frac{1}{e}|x| < 1 \\ \text{unknown if } \frac{1}{e}|x| = 1 \\ \text{divergent if } \frac{1}{e}|x| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $|x| < e$, or $-e < x < e$, we see that the center of convergence is at $x = 0$ and the radius of convergence is e .

[SI*] **Example:**

Determine the Taylor series about the point x_0 for the given function. Also determine the radius of convergence of the series.

- (a) $\sin x, x_0 = 0$
- (b) $e^x, x_0 = 0$
- (c) $x, x_0 = 1$
- (d) $x^2, x_0 = -1$
- (e) $\ln(x), x_0 = 1$
- (f) $\frac{1}{1-x}, x_0 = 0$
- (g) $\frac{1}{1+x}, x_0 = 0$
- (h) $\frac{1}{1-x}, x_0 = 2$

Solution:

- (a) The Taylor series expansion for a function $f(x)$ about the point $x = x_0$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

Since $x_0 = 0$, this formula reduces to

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

The aim then is to determine the n th derivative of $\sin x$ evaluated at $x = 0$. Start taking derivatives and try to observe a pattern.

$$\begin{aligned} n = 0 : f^{(0)}(x) &= \sin x & \rightarrow & f^{(0)}(0) = 0 \\ n = 1 : f^{(1)}(x) &= \cos x & \rightarrow & f^{(1)}(0) = 1 \\ n = 2 : f^{(2)}(x) &= -\sin x & \rightarrow & f^{(2)}(0) = 0 \\ n = 3 : f^{(3)}(x) &= -\cos x & \rightarrow & f^{(3)}(0) = -1 \\ n = 4 : f^{(4)}(x) &= \sin x & \rightarrow & f^{(4)}(0) = 0 \\ n = 5 : f^{(5)}(x) &= \cos x & \rightarrow & f^{(5)}(0) = 1 \\ & \vdots & & \end{aligned}$$

Since $f^{(0)}(0) = 0$, the sum can be started from $n = 1$.

$$f(x) = \sin x = \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

All the even valued of n result in $f^{(n)}(0)$ being zero, so sum over the odd integers only by making the substitution $n = 2k + 1$.

$$\sin x = \sum_{k=0}^{\infty} \frac{f^{(2k+1)}(0)}{(2k+1)!} x^{2k+1}$$

Based on the results above, $f^{(2k+1)}(0) = (-1)^k$. Therefore,

$$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2k+1}$$

Apply the ratio test to determine the condition in which the series converges.

$$\begin{aligned}
 \lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| &= \lim_{k \rightarrow \infty} \left| \frac{\frac{(-1)^{k+1}}{(2(k+1)+1)!} x^{2(k+1)+1}}{\frac{(-1)^k}{(2k+1)!} x^{2k+1}} \right| \\
 &= \lim_{k \rightarrow \infty} \left| \frac{(-1)^{k+1} (2k+1)! x^{2k+3}}{(-1)^k (2k+3)! x^{2k+1}} \right| \\
 &= \lim_{k \rightarrow \infty} \left| -\frac{1}{(2k+3)(2k+2)} x^2 \right| \\
 &= \lim_{k \rightarrow \infty} \frac{1}{(2k+3)(2k+2)} x^2 \\
 &= 0x^2
 \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } 0x^2 < 1 \\ \text{unknown if } 0x^2 = 1 \\ \text{divergent if } 0x^2 > 1 \end{cases}$$

From the condition of convergence, which can also be written as $|x| < \sqrt{\frac{1}{0}} = \infty$, or $-\infty < x < \infty$, we see that the center of convergence is at $x = 0$ and the radius of convergence is ∞ .

(b) The Taylor series expansion for a function $f(x)$ about the point $x = x_0$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

Since $x_0 = 0$, this formula reduces to

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

The aim then is to determine the n th derivative of e^x evaluated at $x = 0$. Start taking derivatives and try to observe a pattern.

$$\begin{aligned}
 n = 0: \quad f^{(0)}(x) &= e^x & \rightarrow & \quad f^{(0)}(0) = 1 \\
 n = 1: \quad f^{(1)}(x) &= e^x & \rightarrow & \quad f^{(1)}(0) = 1 \\
 n = 2: \quad f^{(2)}(x) &= e^x & \rightarrow & \quad f^{(2)}(0) = 1 \\
 n = 3: \quad f^{(3)}(x) &= e^x & \rightarrow & \quad f^{(3)}(0) = 1 \\
 n = 4: \quad f^{(4)}(x) &= e^x & \rightarrow & \quad f^{(4)}(0) = 1 \\
 n = 5: \quad f^{(5)}(x) &= e^x & \rightarrow & \quad f^{(5)}(0) = 1 \\
 &\vdots & &
 \end{aligned}$$

Notice that $f^{(n)}(0) = 1$ for all n . Therefore,

$$f(x) = e^x = \sum_{n=0}^{\infty} \frac{1}{n!} x^n$$

Apply the ratio test to determine the condition in which the series converges.

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{1}{(n+1)!} x^{n+1}}{\frac{1}{n!} x^n} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{n!}{(n+1)!} \frac{x^{n+1}}{x^n} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{1}{n+1} x \right| \\
 &= \lim_{n \rightarrow \infty} \frac{1}{n+1} |x| \\
 &= 0|x|
 \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } 0 < |x| < 1 \\ \text{unknown if } |x| = 1 \\ \text{divergent if } |x| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $|x| < \frac{1}{0} = \infty$, or $-\infty < x < \infty$, we see that the center of convergence is at $x = 0$ and the radius of convergence is ∞ .

(c) The Taylor series expansion for a function $f(x)$ about the point $x = x_0$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

Since $x_0 = 1$, this formula becomes

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(1)}{n!} (x - 1)^n$$

The aim then is to determine the n th derivative of x evaluated at $x = 1$. Start taking derivatives and try to observe a pattern.

$$\begin{aligned}
 n = 0 : \quad f^{(0)}(x) &= x & \rightarrow & \quad f^{(0)}(1) = 1 \\
 n = 1 : \quad f^{(1)}(x) &= 1 & \rightarrow & \quad f^{(1)}(1) = 1 \\
 n = 2 : \quad f^{(2)}(x) &= 0 & \rightarrow & \quad f^{(2)}(1) = 0 \\
 n = 3 : \quad f^{(3)}(x) &= 0 & \rightarrow & \quad f^{(3)}(1) = 0 \\
 n = 4 : \quad f^{(4)}(x) &= 0 & \rightarrow & \quad f^{(4)}(1) = 0 \\
 n = 5 : \quad f^{(5)}(x) &= 0 & \rightarrow & \quad f^{(5)}(1) = 0 \\
 &\vdots & &
 \end{aligned}$$

Notice that $f^{(n)}(1) = 0$ for $n > 1$. Therefore,

$$\begin{aligned} f(x) &= x = \frac{f^{(0)}(1)}{0!}(x-1)^0 + \frac{f^{(1)}(1)}{1!}(x-1)^1 + \sum_{n=2}^{\infty} \frac{f^{(n)}(1)}{n!}(x-1)^n \\ &= (x-1)^0 + (x-1)^1 + \sum_{n=2}^{\infty} \frac{0}{n!}(x-1)^n \\ &= 1 + (x-1) \end{aligned}$$

Since this series is finite, it is convergent for any value of x . That is, the radius of convergence is ∞ .

(d) The Taylor series expansion for a function $f(x)$ about the point $x = x_0$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!}(x-x_0)^n$$

Since $x_0 = -1$, this formula becomes

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(-1)}{n!}(x+1)^n$$

The aim then is to determine the n th derivative of x^2 evaluated at $x = -1$. Start taking derivatives and try to observe a pattern.

$$\begin{aligned} n=0: \quad f^{(0)}(x) &= x^2 \quad \rightarrow \quad f^{(0)}(-1) = 1 \\ n=1: \quad f^{(1)}(x) &= 2x \quad \rightarrow \quad f^{(1)}(-1) = -2 \\ n=2: \quad f^{(2)}(x) &= 2 \quad \rightarrow \quad f^{(2)}(-1) = 2 \\ n=3: \quad f^{(3)}(x) &= 0 \quad \rightarrow \quad f^{(3)}(-1) = 0 \\ n=4: \quad f^{(4)}(x) &= 0 \quad \rightarrow \quad f^{(4)}(-1) = 0 \\ n=5: \quad f^{(5)}(x) &= 0 \quad \rightarrow \quad f^{(5)}(-1) = 0 \\ &\vdots \end{aligned}$$

Notice that $f^{(n)}(-1) = 0$ for $n > 2$. Therefore,

$$\begin{aligned} f(x) &= x = \frac{f^{(0)}(-1)}{0!}(x+1)^0 + \frac{f^{(1)}(-1)}{1!}(x+1)^1 + \frac{f^{(2)}(-1)}{2!}(x+1)^2 + \sum_{n=3}^{\infty} \frac{f^{(n)}(-1)}{n!}(x+1)^n \\ &= (x+1)^0 - 2(x+1)^1 + (x+1)^2 + \sum_{n=3}^{\infty} \frac{0}{n!}(x+1)^n \\ &= 1 - 2(x+1) + (x+1)^2 \end{aligned}$$

Since this series is finite, it is convergent for any value of x . That is, the radius of convergence is ∞ .

(e) The Taylor series expansion for a function $f(x)$ about the point $x = x_0$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!}(x-x_0)^n$$

Since $x_0 = 1$, this formula becomes

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(1)}{n!} (x-1)^n$$

The aim then is to determine the n th derivative of $\ln x$ evaluated at $x = 1$. Start taking derivatives and try to observe a pattern.

$$\begin{aligned} n=0: & f^{(0)}(x) = \ln x \rightarrow f^{(0)}(1) = 0 \\ n=1: & f^{(1)}(x) = x^{-1} \rightarrow f^{(1)}(1) = 1 \\ n=2: & f^{(2)}(x) = -x^{-2} \rightarrow f^{(2)}(1) = -1 \\ n=3: & f^{(3)}(x) = 2x^{-3} \rightarrow f^{(3)}(1) = 2 \\ n=4: & f^{(4)}(x) = -6x^{-4} \rightarrow f^{(4)}(1) = -6 \\ n=5: & f^{(5)}(x) = 24x^{-5} \rightarrow f^{(5)}(1) = 24 \\ & \vdots \end{aligned}$$

Since $f^{(0)}(1) = 0$, the series can start at $n = 1$.

$$f(x) = \ln x = \sum_{n=1}^{\infty} \frac{f^{(n)}(1)}{n!} (x-1)^n$$

Notice that $f^{(n)}(1) = (-1)^{n-1}(n-1)!$ for $n > 0$.

$$\begin{aligned} \ln x &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}(n-1)!}{n!} (x-1)^n \\ &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} (x-1)^n \end{aligned}$$

Apply the ratio test to determine the condition in which the series converges.

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^n}{n+1} (x-1)^{n+1}}{\frac{(-1)^{n-1}}{n} (x-1)^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| (-1) \frac{n}{n+1} (x-1) \right| \\ &= \lim_{n \rightarrow \infty} \frac{n}{n+1} |x-1| \\ &= \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n}} |x-1| \\ &= |x-1| \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } |x-1| < 1 \\ \text{unknown if } |x-1| = 1 \\ \text{divergent if } |x-1| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $-1 < x - 1 < 1$, $0 < x < 2$, we see that the center of convergence is at $x = 1$ and the radius of convergence is 1.

(f) The Taylor series expansion for a function $f(x)$ about the point $x = x_0$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

Since $x_0 = 0$, this formula becomes

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

The aim then is to determine the n th derivative of $\frac{1}{1-x}$ evaluated at $x = 0$. Start taking derivatives and try to observe a pattern.

$$\begin{aligned} n = 0: & \quad f^{(0)}(x) = (1-x)^{-1} \quad \rightarrow \quad f^{(0)}(0) = 1 \\ n = 1: & \quad f^{(1)}(x) = (1-x)^{-2} \quad \rightarrow \quad f^{(1)}(0) = 1 \\ n = 2: & \quad f^{(2)}(x) = 2(1-x)^{-3} \quad \rightarrow \quad f^{(2)}(0) = 2 \\ n = 3: & \quad f^{(3)}(x) = 6(1-x)^{-4} \quad \rightarrow \quad f^{(3)}(0) = 6 \\ n = 4: & \quad f^{(4)}(x) = 24(1-x)^{-5} \quad \rightarrow \quad f^{(4)}(0) = 24 \\ n = 5: & \quad f^{(5)}(x) = 120(1-x)^{-6} \quad \rightarrow \quad f^{(5)}(0) = 120 \\ & \quad \vdots \end{aligned}$$

Notice that $f^{(n)}(0) = n!$.

$$\begin{aligned} f(x) &= \frac{1}{1-x} = \sum_{n=0}^{\infty} \frac{n!}{n!} x^n \\ &= \sum_{n=0}^{\infty} x^n \end{aligned}$$

Apply the ratio test to determine the condition in which the series converges.

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{x^n} \right| \\ &= \lim_{n \rightarrow \infty} |x| \\ &= |x| \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } |x| < 1 \\ \text{unknown if } |x| = 1 \\ \text{divergent if } |x| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $-1 < x < 1$, we see that the center of convergence is at $x = 0$ and the radius of convergence is 1.

(g) The Taylor series expansion for a function $f(x)$ about the point $x = x_0$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

Since $x_0 = 0$, this formula becomes

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

The aim then is to determine the n th derivative of $\frac{1}{1+x}$ evaluated at $x = 0$. Start taking derivatives and try to observe a pattern.

$$\begin{aligned} n = 0: & f^{(0)}(x) = (1+x)^{-1} \rightarrow f^{(0)}(0) = 1 \\ n = 1: & f^{(1)}(x) = -(1+x)^{-2} \rightarrow f^{(1)}(0) = -1 \\ n = 2: & f^{(2)}(x) = 2(1+x)^{-3} \rightarrow f^{(2)}(0) = 2 \\ n = 3: & f^{(3)}(x) = -6(1+x)^{-4} \rightarrow f^{(3)}(0) = -6 \\ n = 4: & f^{(4)}(x) = 24(1+x)^{-5} \rightarrow f^{(4)}(0) = 24 \\ n = 5: & f^{(5)}(x) = -120(1+x)^{-6} \rightarrow f^{(5)}(0) = -120 \\ & \vdots \end{aligned}$$

Notice that $f^{(n)}(0) = (-1)^n n!$.

$$\begin{aligned} f(x) &= \frac{1}{1+x} = \sum_{n=0}^{\infty} \frac{(-1)^n n!}{n!} x^n \\ &= \sum_{n=0}^{\infty} (-1)^n x^n \end{aligned}$$

Apply the ratio test to determine the condition in which the series converges.

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} x^{n+1}}{(-1)^n x^n} \right| \\ &= \lim_{n \rightarrow \infty} |(-1)x| \\ &= \lim_{n \rightarrow \infty} |x| \\ &= |x| \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } |x| < 1 \\ \text{unknown if } |x| = 1 \\ \text{divergent if } |x| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $-1 < x < 1$, we see that the center of convergence is at $x = 0$ and the radius of convergence is 1.

(h) The Taylor series expansion for a function $f(x)$ about the point $x = x_0$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

Since $x_0 = 2$, this formula becomes

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(2)}{n!} (x - 2)^n$$

The aim then is to determine the n th derivative of $\frac{1}{1-x}$ evaluated at $x = 2$. Start taking derivatives and try to observe a pattern.

$$\begin{aligned} n = 0 : \quad f^{(0)}(x) &= (1-x)^{-1} \quad \rightarrow \quad f^{(0)}(2) = -1 \\ n = 1 : \quad f^{(1)}(x) &= (1-x)^{-2} \quad \rightarrow \quad f^{(1)}(2) = 1 \\ n = 2 : \quad f^{(2)}(x) &= 2(1-x)^{-3} \quad \rightarrow \quad f^{(2)}(2) = -2 \\ n = 3 : \quad f^{(3)}(x) &= 6(1-x)^{-4} \quad \rightarrow \quad f^{(3)}(2) = 6 \\ n = 4 : \quad f^{(4)}(x) &= 24(1-x)^{-5} \quad \rightarrow \quad f^{(4)}(2) = -24 \\ n = 5 : \quad f^{(5)}(x) &= 120(1-x)^{-6} \quad \rightarrow \quad f^{(5)}(2) = 120 \\ &\vdots \end{aligned}$$

Notice that $f^{(n)}(2) = (-1)^{n+1}n!$.

$$\begin{aligned} f(x) &= \frac{1}{1-x} = \sum_{n=0}^{\infty} \frac{(-1)^{n+1}n!}{n!} (x-2)^n \\ &= \sum_{n=0}^{\infty} (-1)^{n+1} (x-2)^n \end{aligned}$$

Apply the ratio test to determine the condition in which the series converges.

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+2}(x-2)^{n+1}}{(-1)^{n+1}(x-2)^n} \right| \\ &= \lim_{n \rightarrow \infty} |(-1)(x-2)| \\ &= \lim_{n \rightarrow \infty} |x-2| \\ &= |x-2| \end{aligned}$$

According to this test, the series is

$$\begin{cases} \text{convergent if } |x-2| < 1 \\ \text{unknown if } |x-2| = 1 \\ \text{divergent if } |x-2| > 1 \end{cases}$$

From the condition of convergence, which can also be written as $-1 < x-2 < 1$, or $1 < x < 3$, we see that the center of convergence is at $x = 2$ and the radius of convergence is 1.

```

Time taken by function: 2445077 microseconds
root@ ~/SourceCodes/CPP/C++ Create Library/Test Taylor Series ]# ./main
Taylor series for sin(x) , x_{0} = 0
x-1/6*x^(3)+1/120*x^(5)-1/5040*x^(7)+1/362880*x^(9)
Taylor series for exp(x) , x_{0} = 0
x+1/2*x^(2)+1/6*x^(3)+1/24*x^(4)+1/120*x^(5)+1/720*x^(6)+1/5040*x^(7)+1/40320*x^(8)+1/362880*x^(9)+1
Taylor series for x, x_{0} = 1
x
Taylor series for x^(2), x_{0} = -1
x^(2)
Taylor series for ln(x), x_{0} = 1
9*x-18*x^(2)+28*x^(3)-31.5*x^(4)+25.2*x^(5)-14*x^(6)+5.14286*x^(7)-1.125*x^(8)+1/9*x^(9)-2.82897
Taylor series for 1/(1-x), x_{0} = 0
x+x^(2)+x^(3)+x^(4)+x^(5)+x^(6)+x^(7)+x^(8)+x^(9)+1
Taylor series for 1/(1+x), x_{0} = 0
-x+x^(2)-x^(3)+x^(4)-x^(5)+x^(6)-x^(7)+x^(8)-x^(9)+1
Taylor series for 1/(1-x), x_{0} = 2
4097*x-7423*x^(2)+7937*x^(3)-5503*x^(4)+2561*x^(5)-799*x^(6)+161*x^(7)-19*x^(8)+x^(9)-1023
Time taken by function: 2445077 microseconds
    
```

Figure 3.62: The computation for the Taylor series expansion for function $f(x)$ in the example (a) to (h) (*SymIntegration/Examples/Differential Equations/Test Taylor Series/main.cpp*).

[SI*] Series Solutions Near an Ordinary Point

We now consider methods of solving second order linear equations when the coefficients are functions of the independent variable. In this section we will denote the independent variable by x . It is sufficient to consider the homogeneous equation

$$P(x) \frac{d^2 y}{dx^2} + Q(x) \frac{dy}{dx} + R(x)y = 0 \quad (3.260)$$

since the procedure for the corresponding nonhomogeneous equation is similar. Many problems in mathematical physics lead to equations of the form (3.260) having polynomial coefficients; examples include the Bessel equation

$$x^2 y'' + xy' + (x^2 - \nu^2)y = 0$$

where ν is a constant, and the Legendre equation

$$(1 - x^2)y'' - 2xy' + \alpha(\alpha + 1)y = 0$$

where α is a constant. For this reason, as well as to simplify the algebraic computations, we primarily consider the case in which the functions P , Q , and R are polynomials. However, as we will see, the method of solution is also applicable when P , Q , and R are general analytic functions.

For the present, then, suppose that P , Q , and R are polynomials and that they have no common factors. Suppose also that we wish to solve Eq. (3.260) in the neighborhood of a point x_0 . The solution of Eq. (3.260) in an interval containing x_0 is closely associated with the behavior of P in that interval.

A point x_0 such that $P(x_0) \neq 0$ is called an ordinary point. Since P is continuous, it follows that there is an interval about x_0 in which $P(x)$ is never zero. In that interval we can divide Eq. (3.260) by $P(x)$ to obtain

$$y'' + p(x)y' + q(x)y = 0 \quad (3.261)$$

where $p(x) = \frac{Q(x)}{P(x)}$ and $q(x) = \frac{R(x)}{P(x)}$ are continuous functions. Hence, according to the existence and uniqueness Theorem, there exists in that interval a unique solution of Eq. (3.260) that also satisfies the initial conditions $y(x_0) = y_0, y'(x_0) = y'_0$ for arbitrary values of y_0 and y'_0 . In this section we discuss the solution of Eq. (3.260) in the neighborhood of an ordinary point.

On the other hand, if $P(x_0) = 0$, then x_0 is called a singular point of Eq. (3.260). In this case at least one of $Q(x_0)$ and $R(x_0)$ is not zero. Consequently, at least one of the coefficients p and q in Eq. (3.261) becomes unbounded as $x \rightarrow x_0$, and therefore the uniqueness Theorem does not apply in this case.

We now take up the problem of solving Eq. (3.260) in the neighborhood of an ordinary point x_0 . We look for solutions of the form

$$y = a_0 + a_1(x - x_0) + \cdots + a_n(x - x_0)^n + \cdots = \sum_{n=0}^{\infty} a_n(x - x_0)^n \quad (3.262)$$

and assume that the series converges in the interval $|x - x_0| < \rho$ for some $\rho > 0$. While at first sight it may appear unattractive to seek a solution in the form of a power series, this is actually a convenient and useful form for a solution. Within their intervals of convergence, power series behave very much like polynomials and are easy to manipulate both analytically and numerically. Indeed, even if we can obtain a solution in terms of elementary functions, such as exponential and trigonometric functions, we are likely to need a power series or some equivalent expression if we want to evaluate them numerically or to plot their graphs. The most practical way to determine the coefficients a_n is to substitute the series (3.262) and its derivatives for y, y' , and y'' in Eq. (3.260).

[SI*] **Example:**

Find a series solution of the equation

$$y'' + y = 0, \quad -\infty < x < \infty \quad (3.263)$$

Solution:

As we know, a fundamental set of solutions of this equation are $\sin x$ and $\cos x$, so series methods are not needed to solve this equation. However, this example illustrates the use of power series in a relatively simple case. For Eq. (3.263), $P(x) = 1, Q(x) = 0$, and $R(x) = 1$; hence every point is an ordinary point.

We look for a solution in the form of a power series about $x_0 = 0$

$$y = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n + \cdots = \sum_{n=0}^{\infty} a_nx^n \quad (3.264)$$

and assume that the series converges in some interval $|x| < \rho$. Differentiating Eq. (3.264) term by term yields

$$y' = a_1 + 2a_2x + \cdots + na_nx^{n-1} + \cdots = \sum_{n=1}^{\infty} na_nx^{n-1} \quad (3.265)$$

$$y'' = 2a_2 + 6a_3x + \cdots + n(n-1)a_nx^{n-2} + \cdots = \sum_{n=2}^{\infty} n(n-1)a_nx^{n-2} \quad (3.266)$$

Substituting the series (3.264) and (3.266) for y and y'' in Eq. (3.263) gives

$$\sum_{n=2}^{\infty} n(n-1)a_nx^{n-2} + \sum_{n=0}^{\infty} a_nx^n = 0$$

To combine the two series, we need to rewrite at least one of them so that both series display the same generic term. Thus, in the first sum, we shift the index of summation by replacing n by $n + 2$ and starting the sum at 0 rather than at 2. We obtain

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

or

$$\sum_{n=0}^{\infty} [(n+2)(n+1)a_{n+2} + a_n] x^n = 0$$

For this equation to be satisfied for all x , the coefficient of each power of x must be zero; hence we conclude that

$$(n+2)(n+1)a_{n+2} + a_n = 0, \quad n = 0, 1, 2, 3, \dots \quad (3.267)$$

Equation (3.267) is referred to as a recurrence relation. The successive coefficients can be evaluated one by one by writing the recurrence relation first for $n = 0$, then for $n = 1$, and so forth. In this example Eq. (3.267) relates each coefficient to the second one before it. Thus the even-numbered coefficients ($a_0, a_2, a_4, \dots$) and the odd-numbered ones ($a_1, a_3, a_5, \dots$) are determined separately. For the even-numbered coefficients we have

$$\begin{aligned} a_2 &= -\frac{a_0}{2 \cdot 1} = -\frac{a_0}{2!} \\ a_4 &= -\frac{a_2}{4 \cdot 3} = \frac{a_0}{4!} \\ a_6 &= -\frac{a_4}{6 \cdot 5} = -\frac{a_0}{6!} \end{aligned}$$

These results suggest that in general, if $n = 2k$, then

$$a_n = a_{2k} = \frac{(-1)^k}{(2k)!} a_0, \quad k = 1, 2, 3, \dots \quad (3.268)$$

We can prove Eq. (3.268) by mathematical induction. First, observe that it is true for $k = 1$. Next, assume that it is true for an arbitrary value of k and consider the case $k + 1$. We have

$$a_{2k+2} = -\frac{a_{2k}}{(2k+2)(2k+1)} = -\frac{(-1)^k}{(2k+2)(2k+1)(2k)!} a_0 = \frac{(-1)^{k+1}}{(2k+2)!} a_0$$

Hence Eq. (3.268) is also true for $k + 1$, and consequently it is true for all positive integers k . Similarly, for the odd-numbered coefficients

$$\begin{aligned} a_3 &= -\frac{a_1}{3 \cdot 2} = -\frac{a_1}{3!} \\ a_5 &= -\frac{a_3}{5 \cdot 4} = \frac{a_1}{5!} \\ a_7 &= -\frac{a_5}{7 \cdot 6} = -\frac{a_1}{7!} \end{aligned}$$

and in general, if $n = 2k + 1$, then

$$a_n = a_{2k+1} = \frac{(-1)^k}{(2k+1)!} a_1, \quad k = 1, 2, 3, \dots \quad (3.269)$$

Substituting these coefficients into Eq. (3.264)

$$\begin{aligned}
 y &= a_0 + a_1x + a_2x^2 + \cdots + a_nx^n + \cdots \\
 &= a_0 + a_1x - \frac{a_0}{2!}x^2 - \frac{a_1}{3!}x^3 + \frac{a_0}{4!}x^4 + \frac{a_1}{5!}x^5 \\
 &\quad + \cdots + \frac{(-1)^na_0}{(2n)!}x^{2n} + \frac{(-1)^na_1}{(2n+1)!}x^{2n+1} + \cdots \\
 &= a_0 \left[1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots + \frac{(-1)^n}{(2n)!}x^{2n} + \cdots \right] \\
 &\quad + a_1 \left[x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots + \frac{(-1)^n}{(2n+1)!}x^{2n+1} + \cdots \right] \\
 &= a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!}x^{2n} + a_1 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!}x^{2n+1}
 \end{aligned} \tag{3.270}$$

Now that we have formally obtained two series solutions of Eq. (3.263), we can test them for convergence. Using the ratio test, we can show that each of the series in Eq. (3.270) converges for all x , and this justifies retroactively all the steps used in obtaining the solutions. Indeed, we recognize that the first series in Eq. (3.270) is exactly the Taylor series for $\cos x$ about $x = 0$ and that the second is the Taylor series for $\sin x$ about $x = 0$. Thus, as expected, we obtain the solution $y = a_0 \cos x + a_1 \sin x$.

Notice that no conditions are imposed on a_0 and a_1 ; hence they are arbitrary. From Eqs. (3.264) and (3.265) we see that y and y' evaluated at $x = 0$ are a_0 and a_1 , respectively. Since the initial conditions $y(0)$ and $y'(0)$ can be chosen arbitrarily, it follows that a_0 and a_1 should be arbitrary until specific initial conditions are stated.




Figure 3.63: The polynomial approximations to $\cos x$. The value of n is the degree of the approximating polynomial (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Polynomial Approximation to Cosine/main.cpp).




Figure 3.64: The polynomial approximations to $\sin x$. The value of n is the degree of the approximating polynomial (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Polynomial Approximation to Cosine/main.cpp).

The graphs show how the partial sums of the series in Eq. (3.270) approximate $\cos x$ and $\sin x$. As the number of terms increases, the interval over which the approximation is satisfactory become longer, and for each x in this interval the accuracy of the approximation improves. However, you should always remember that a truncated power series provides only a local approximation of the solution in a neighborhood of the initial point $x = 0$; it cannot adequately represent the solution for large $|x|$.

[SI*] If we had not known that $\sin x$ and $\cos x$ form a fundamental set of solutions of Eq. (3.263) and had simply solved Eq. (3.263) using series methods, we would still have obtained the solution (3.270). In recognition of the fact that the differential equation (3.263) often occurs in applications, we might decide to give the two solutions of Eq. (3.270) special names

$$C(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n}, \quad S(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} \quad (3.271)$$

Then we might ask what properties these functions have. For instance, can we be sure that $C(x)$ and $S(x)$ form a fundamental set of solutions? It follows at once from the series expansions that $C(0) = 1$ and $S(0) = 0$. By differentiating the series for $C(x)$ and $S(x)$ term by term, we find that

$$S'(x) = C(x), \quad C'(x) = -S(x) \quad (3.272)$$

Thus, at $x = 0$ we have $S'(0) = 1$ and $C'(0) = 0$. Consequently, the Wronskian of C and S at $x = 0$ is

$$W(C, S)(0) = \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1 \quad (3.273)$$

so these functions do indeed form a fundamental set of solutions. By substituting $-x$ for x in each of Eqs. (3.271), we obtain $C(-x) = C(x)$, and $S(-x) = -S(x)$. Moreover, by calculating with the infinite series, we can show that the functions $C(x)$ and $S(x)$ have all the usual analytical and algebraic properties of the cosine and sine functions, respectively.

The function $\sin x$ can be defined as the unique solution of the initial value problem $y'' + y = 0, y(0) = 0, y'(0) = 1$; similarly, $\cos x$ can be defined as the unique solution of the initial value problem $y'' + y = 0, y(0) = 1, y'(0) = 0$. Many other functions that are important in mathematical physics are also defined as solutions of certain initial value problems. For most of these functions there is no simpler or more elementary way to approach them.

[SI*] **Example:**

Find a series solution in powers of x of Airy's equation

$$y'' - xy = 0, \quad -\infty < x < \infty \quad (3.274)$$

Solution:

For this equation $P(x) = 1, Q(x) = 0$, and $R(x) = -x$; hence every point is an ordinary point. We assume that

$$y = \sum_{n=0}^{\infty} a_n x^n \quad (3.275)$$

and that the series converges in some interval $|x| < \rho$. The series for y'' is given by Eq. (3.266), we can rewrite it as

$$y'' = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n \quad (3.276)$$

Substituting the series (3.275) and (3.276) for y and y'' in Eq. (3.274), we obtain

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n - x \sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} a_n x^{n+1} \quad (3.277)$$

Next, we shift the index of summation in the series on the right side of Eq. (3.277) by replacing n by $n-1$ and starting the summation at 1 rather than zero. Thus we have

$$2 \cdot 1a_2 + \sum_{n=1}^{\infty} (n+2)(n+1)a_{n+2}x^n = \sum_{n=1}^{\infty} a_{n-1}x^n$$

Again, for this equation to be satisfied for all x in some interval, the coefficients of like powers of x must be equal; hence $a_2 = 0$, and we obtain the recurrence relation

$$(n+2)(n+1)a_{n+2} = a_{n-1}, \quad \text{for } n = 1, 2, 3, \dots \quad (3.278)$$

Since a_{n+2} is given in terms of a_{n-1} , the a 's are determined in steps of three. Thus a_0 determines a_3 , which in turn determines $a_6, \dots$; a_1 determines a_4 , which in turn determines $a_7, \dots$; and a_2 determines a_5 , which in turn determines $a_8, \dots$. Since $a_2 = 0$, we immediately conclude that $a_5 = a_8 = a_{11} = \dots = 0$.

For the sequence $a_0, a_3, a_6, a_9, \dots$ we set $n = 1, 4, 7, 10, \dots$ in the recurrence relation:

$$\begin{aligned} a_3 &= \frac{a_0}{2 \cdot 3} \\ a_6 &= \frac{a_3}{5 \cdot 6} = \frac{a_0}{2 \cdot 3 \cdot 5 \cdot 6} \\ a_9 &= \frac{a_6}{8 \cdot 9} = \frac{a_0}{2 \cdot 3 \cdot 5 \cdot 6 \cdot 8 \cdot 9} \end{aligned}$$

These results suggest the general formula

$$a_{3n} = \frac{a_0}{2 \cdot 3 \cdot 5 \cdot 6 \cdots (3n-1)(3n)}, \quad n \geq 4$$

For the sequence $a_1, a_4, a_7, a_{10}, \dots$ we set $n = 2, 5, 8, 11, \dots$ in the recurrence relation:

$$\begin{aligned} a_4 &= \frac{a_1}{3 \cdot 4} \\ a_7 &= \frac{a_4}{6 \cdot 7} = \frac{a_0}{3 \cdot 4 \cdot 6 \cdot 7} \\ a_{10} &= \frac{a_7}{9 \cdot 10} = \frac{a_0}{3 \cdot 4 \cdot 6 \cdot 7 \cdot 9 \cdot 10} \end{aligned}$$

In general, we have

$$a_{3n+1} = \frac{a_1}{3 \cdot 4 \cdot 6 \cdot 7 \cdots (3n)(3n+1)}, \quad n \geq 4$$

Thus the general solution of Airy's equation is

$$\begin{aligned} y = a_0 &\left[1 + \frac{x^3}{2 \cdot 3} + \frac{x^6}{2 \cdot 3 \cdot 5 \cdot 6} + \cdots + \frac{x^{3n}}{2 \cdot 3 \cdots (3n-1)(3n)} + \cdots \right] \\ &+ a_1 \left[x + \frac{x^4}{3 \cdot 4} + \frac{x^7}{3 \cdot 4 \cdot 6 \cdot 7} + \cdots + \frac{x^{3n+1}}{3 \cdot 4 \cdots (3n)(3n+1)} + \cdots \right] \end{aligned} \quad (3.279)$$

Having obtained these two series solutions, we can now investigate their convergence. Because of the rapid growth of the denominators of the terms in the series (3.279), we might expect these series to have a large radius of convergence. Indeed, it is easy to use the ratio test to show that both these series converge for all x .

Assuming for the moment that the series do converge for all x , let y_1 and y_2 denote the functions defined by the expressions in the first and second sets of brackets, respectively, in Eq. (3.279). Then, by choosing first $a_0 = 1, a_1 = 0$ and then $a_0 = 0, a_1 = 1$, it follows that y_1 and y_2 are individually solutions of Eq. (3.274). Notice that y_1 satisfies the initial conditions $y_1(0) = 1, y_1'(0) = 0$ and that y_2 satisfies the initial conditions $y_2(0) = 0, y_2'(0) = 1$. Thus $W(y_1, y_2)(0) = 1 \neq 0$, and consequently y_1 and y_2 are a fundamental set of solutions. Hence the general solution of Airy's equation is

$$y = a_0 y_1(x) + a_1 y_2(x), \quad -\infty < x < \infty$$




Figure 3.65: The polynomial approximations to the solution $y_1(x)$ of Airy's equation. The value of n is the degree of the approximating polynomial (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Solution of Airy's Equation Ai(x)/main.cpp).




Figure 3.66: The polynomial approximations to the solution $y_2(x)$ of Airy's equation. The value of n is the degree of the approximating polynomial (Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Solution of Airy's Equation Bi(x)/main.cpp).

The partial sums provide provide local approximations to the solutions in a neighborhood of the origin. Although the quality of the approximation improves as the number of terms increases, no polynomial can adequately represent y_1 and y_2 for large $|x|$. A practical way to estimate the interval in which a given partial sum is reasonably accurate is to compare the graphs of that partial sum and the next one, obtained by including one more term. As soon

as the graphs begin to separate noticeably, we can be confident that the original partial sum is no longer accurate.

Observe that both y_1 and y_2 are monotone for $x > 0$ and oscillatory for $x < 0$. The oscillations are not uniform but, rather, decay in amplitude and increase in frequency as the distance from the origin increases.

The solutions y_1 and y_2 of Airy's equation are not elementary functions that you have already encountered in calculus. However, because of their importance in some physical applications, these functions have been extensively studied, and their properties are well known to applied mathematicians and scientists.

[SI*] Example:

Find a series solution in powers of $x - 1$ of Airy's equation.

Solution:

The point $x = 1$ is an ordinary point of Eq. (3.274), and thus we look for a solution of the form

$$y = \sum_{n=0}^{\infty} a_n (x - 1)^n$$

where we assume that the series converges in some interval $|x - 1| < \rho$. Then

$$y' = \sum_{n=1}^{\infty} n a_n (x - 1)^{n-1} = \sum_{n=0}^{\infty} (n + 1) a_{n+1} (x - 1)^n$$

and

$$y'' = \sum_{n=2}^{\infty} n(n - 1) a_n (x - 1)^{n-2} = \sum_{n=0}^{\infty} (n + 2)(n + 1) a_{n+2} (x - 1)^n$$

Substituting for y and y'' in Eq. (3.274), we obtain

$$\begin{aligned} y'' - xy &= 0 \\ \sum_{n=0}^{\infty} (n + 2)(n + 1) a_{n+2} (x - 1)^n - x \sum_{n=0}^{\infty} a_n (x - 1)^n &= 0 \\ \sum_{n=0}^{\infty} (n + 2)(n + 1) a_{n+2} (x - 1)^n &= x \sum_{n=0}^{\infty} a_n (x - 1)^n \end{aligned} \tag{3.280}$$

Now to equate the coefficients of like powers of $(x - 1)$, we must express x , the coefficient of y in Eq. (3.274), in powers of $x - 1$; that is, we write $x = 1 + (x - 1)$. Note that this is precisely the Taylor series for x about $x = 1$. Then Eq. () takes the form

$$\begin{aligned} \sum_{n=0}^{\infty} (n + 2)(n + 1) a_{n+2} (x - 1)^n &= [1 + (x - 1)] \sum_{n=0}^{\infty} a_n (x - 1)^n \\ &= \sum_{n=0}^{\infty} a_n (x - 1)^n + \sum_{n=0}^{\infty} a_n (x - 1)^{n+1} \end{aligned}$$

Shifting the index of summation in the second series on the right gives

$$\sum_{n=0}^{\infty} (n + 2)(n + 1) a_{n+2} (x - 1)^n = \sum_{n=0}^{\infty} a_n (x - 1)^n + \sum_{n=1}^{\infty} a_{n-1} (x - 1)^n$$

Equating coefficients of like powers of $x - 1$, we obtain

$$\begin{aligned} 2a_2 &= a_0 \\ (3 \cdot 2)a_3 &= a_1 + a_0 \\ (4 \cdot 3)a_4 &= a_2 + a_1 \\ (5 \cdot 4)a_5 &= a_3 + a_2 \\ &\vdots \end{aligned}$$

The general recurrence relation is

$$(n+2)(n+1)a_{n+2} = a_n + a_{n-1}, \quad \text{for } n \geq 1 \quad (3.281)$$

Solving for the first few coefficients a_n in terms of a_0 and a_1 , we find that

$$\begin{aligned} a_2 &= \frac{a_0}{2} \\ a_3 &= \frac{a_1}{6} + \frac{a_0}{6} \\ a_4 &= \frac{a_2}{12} + \frac{a_1}{12} = \frac{a_0}{24} + \frac{a_1}{12} \\ a_5 &= \frac{a_3}{20} + \frac{a_2}{20} = \frac{a_0}{30} + \frac{a_1}{120} \end{aligned}$$

Hence

$$\begin{aligned} y &= a_0 \left[1 + \frac{(x-1)^2}{2} + \frac{(x-1)^3}{6} + \frac{(x-1)^4}{24} + \frac{(x-1)^5}{30} + \dots \right] \\ &\quad + a_1 \left[(x-1) + \frac{(x-1)^3}{6} + \frac{(x-1)^4}{12} + \frac{(x-1)^5}{120} + \dots \right] \end{aligned} \quad (3.282)$$

In general, when the recurrence relation has more than two terms, as in Eq. (3.281), the determination of a formula for a_n in terms a_0 and a_1 will be fairly complicated, if not impossible. In this example such a formula is not readily apparent. Lacking such a formula, we cannot test the two series in Eq. (3.282) for convergence by direct methods such as the ratio test. However, even without knowing the formula for a_n , it is possible to establish that the two series in Eq. (3.282) converge for all x . Further, they define functions y_3 and y_4 , that are a fundamental set of solutions of the Airy equation (3.274). Thus

$$y = a_0 y_3(x) + a_1 y_4(x)$$

is the general solution of Airy's equation for $-\infty < x < \infty$.

[SI*] If we look for a solution of Eq. (3.260) of the form $y = \sum_{n=0}^{\infty} a_n (x - x_0)^n$, then the coefficients $P(x)$, $Q(x)$, and $R(x)$ in Eq. (3.260) must also be expressed in powers of $x - x_0$. Alternatively, we can make the change of variable $x - x_0 = t$, obtaining a new differential equation for y as a function of t , and then look for solutions of this new equation of the form $\sum_{n=0}^{\infty} a_n t^n$. When we have finished the calculations, we replace t by $x - x_0$.

When we have found the two sets of solutions, for example of Airy's equation. The functions y_1 and y_2 defined by the series in Eq. (3.279) are a fundamental set of solutions of Eq. (3.274) for all x , and this is also true for the functions y_3 and y_4 defined by the series in Eq.

(3.282). According to the general theory of second order linear equations, each of the first two functions can be expressed as a linear combination of the latter two functions and vice versa.

It is not particularly important if we are unable to determine the general coefficient a_n in terms of a_0 and a_1 . What is essential is that we can determine as many coefficients as we want. Thus we can find as many terms in the two series solutions as we want, even if we cannot determine the general term.

[SI*] **Example:**

$$1. \ y'' - y = 0, \ x_0 = 0$$

$$2. \ y'' - xy' - y = 0, \ x_0 = 0$$

[SI*] We have considered the problem of finding solutions of

$$P(x)y'' + Q(x)y' + R(x)y = 0 \tag{3.283}$$

where P, Q , and R are polynomials in the neighborhood of an ordinary point x_0 . Assuming that Eq. (3.283) does have a solution $y = \phi(x)$ and that ϕ has a Taylor series

$$y = \phi(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n \tag{3.284}$$

which converges for $|x - x_0| < \rho$, where $\rho > 0$, we found that the a_n can be determined by directly substituting the series (3.284) for y in Eq. (3.283).

[SI*] **Euler Equations; Regular Singular Points**

[SI*] **Series Solution Near a Regular Singular Point**

[SI*] **Bessel's Equation**

i. **Compute Power Series Solution for Homogeneous Second Order Linear Differential Equations with Constant Coefficients**

Consider the homogeneous second order linear differential equation

$$y'' + 3y' + 2y = 0$$

with initial conditions

$$\begin{aligned} y(0) &= 1 \\ y'(0) &= 2 \end{aligned}$$

We will compute the power series solution around an ordinary point $x_0 = 0$.

We have created a function previously in SymIntegration to compute this type differential equation: `secondorderlineardiffeq_ivpsolution(1,3,2,y,t,0,1,2)`.

The exact solution is

$$y(x) = 4e^{-x} - 3e^{-2x}$$

Now, we want to compute the power series solution for this differential equation to see whether the series solutions converge to the exact solution or not as the terms goes to ∞ .

To find a power series solution for a second order linear differential equation $P(x)y'' + Q(x)y' + R(x)y = 0$ near an ordinary (regular) point $x_0 = 0$, assume a standard power series

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

We will have

$$y(x) = \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = a_1 + 2a_2 x + 3a_3 x^2 + 4a_4 x^3 + \dots$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = 2a_2 + 6a_3 x + 12a_4 x^2 + \dots$$

How to compute the recurrence relation:

1. **Plug in series:** Substitute y, y' , and y'' into the differential equation.
2. **Multiply and distribute:** Multiply the coefficients like x or x^2 into the summation terms, adjusting indices so every power of x matches (e.g., changing x^{n-1} or x^n to a common power like x^k or x^{n-2}).
3. **Combine sums:** Merge all summations into a single expression of the format $\sum c_n x^n = 0$.
4. **Set coefficients to zero:** Equate the coefficient of each power of x to 0 to generate the algebraic equation.
5. **Isolate the highest index:** Solve the equation for the term with the highest subscript (like a_{n+2} or a_n) in terms of lower subscripts. This equation is the recurrence relation.

6. **Iterate coefficients:** Compute sequential terms $(a_2, a_3, a_4, \dots)$ as multiples of the free parameters a_0 and a_1 to write the final general solution $y(x) = a_0 y_1(x) + a_1 y_2(x)$

For the differential equation mentioned above, we substitute $y = \sum_{n=0}^{\infty} a_n x^n$ then differentiate it twice and substitute them into the differential equation to find the recurrence relation. Knowing that we only deal with constant coefficients now, the recurrence formula is easy to obtain.

Always remember that a single universal or "master" formula does not exist for the power series coefficients of every second-order differential equation. Instead, the power series method provides a systematic algorithmic procedure to construct the general solution for linear differential equations with analytic coefficients.

To compute the power series solution of a homogeneous linear differential equation with constant coefficients, we can use the recurrence relation derived from matching coefficients. For an m -th order linear homogeneous differential equation with constant coefficients b_k :

$$\sum_{k=0}^m b_k y^{(k)}(x) = 0$$

Assuming a power series solution $y(x) = \sum_{n=0}^{\infty} a_n x^n$, substituting the derivatives yields the recurrence relation for the coefficients a_n :

$$a_{n+m} = -\frac{1}{b_m \prod_{j=1}^m (n+j)} \sum_{k=0}^{m-1} b_k a_{n+k} \prod_{j=1}^k (n+j)$$

To solve a homogeneous second order linear differential equation with constant coefficients using a power series method in C++, we can represent the series solution as a polynomial class where the coefficients are calculated recurrence-wise.

We have created a class in SymIntegration:

class HigherOrderODE_Homogeneous_PowerSeriesSolver

This class is defined in **dsolve.h** and the computation code for the functions of the class is in **dsolve.cpp**.

```
...

#ifndef HIGHERORDERODE_HOMOGENEOUS_POWERSERIESSOLVER_H
#define HIGHERORDERODE_HOMOGENEOUS_POWERSERIESSOLVER_H

class HigherOrderODE_Homogeneous_PowerSeriesSolver {
private:
    vector<double> ode_coefficients; // ODE coefficients [b_0, b_1, ..., b_m]
    vector<double> initial_values; // Initial conditions [y(0), y'(0), ..., y^(m-1)(0)]
    int order; // Order of the differential equation (m)
    vector<double> coefficients;
```

```

// Helper function to calculate Pochhammer symbol / falling factorial
// factors
double get_factorial_multiplier(int n, int k) const
{
    double multiplier = 1.0;
    for (int j = 1; j <= k; ++j)
    {
        multiplier *= (n + j);
    }
    return multiplier;
}
public:
// Constructor initializes the terms (series degree+1) and base boundary
// conditions
HigherOrderODE_Homogeneous_PowerSeriesSolver(const vector<double>&
    ode_coeffs, const vector<double>& init_conditions) ;

// Computes coefficients up to order N using the recurrence relation:
void computeSeries(int terms) ;
vector<double> coefficientsvector(int terms) ;
double evaluateAt(double x, int terms) ;
void printSeries() const ;
};

#endif

...

```

Code 42: *include/symintegral/dsolve.h*

The **dsolve.cpp** that handles the function to compute the power series solution and approximation at x can be seen below

```

HigherOrderODE_Homogeneous_PowerSeriesSolver::
HigherOrderODE_Homogeneous_PowerSeriesSolver(const vector<double>& ode_coeffs
, const vector<double>& init_conditions)
{
    ode_coefficients = ode_coeffs;
    initial_values = init_conditions;
    order = ode_coeffs.size() - 1;
}

void HigherOrderODE_Homogeneous_PowerSeriesSolver::computeSeries(int terms)
{
    if (terms < 2)
    {
        return;
    }
}

```



```
coefficients.resize(terms,0.0);

// Step 1: Assign initial conditions to the first m series terms
// Recall that  $y^{(k)}(0) = k! * \text{coefficients}_k$ , so  $\text{coefficients}_k = y^{(k)}(0) / k!$ 
double factorial = 1.0;
for (int k = 0; k <= order - 1 && k < terms; ++k)
{
    if (k > 0)
    {
        factorial *= k;
    }
    coefficients[k] = initial_values[k] / factorial;
}

// Step 2: Use recurrence relation to compute subsequent coefficients
double b_m = ode_coefficients[order];
if (b_m == 0.0)
{
    std::cerr << "Error: Leading coefficient b_m cannot be zero." <<
        endl;
}

for (int n = 0; n <= terms - order - 1; ++n)
{
    double sum_terms = 0.0;
    for (int k = 0; k < order; ++k)
    {
        double mult = get_factorial_multiplier(n, k);
        sum_terms += ode_coefficients[k] * coefficients[n + k] * mult
            ;
    }

    double divisor = b_m * get_factorial_multiplier(n, order);
    coefficients[n + order] = -sum_terms / divisor;
}
...
}

vector<double> HigherOrderODE_Homogeneous_PowerSeriesSolver::coefficientsvector(
    int terms)
{
    if (terms < 2)
    {
        return {};
    }
}
```

```
...

    return coefficients;
}

// Evaluates the power series at a specific value of x
double HigherOrderODE_Homogeneous_PowerSeriesSolver::evaluateAt(double x, int
    terms)
{
    double current_x_power = 1.0;
    double sum = 0.0;

    for (int i = 0; i <= terms; ++i)
    {
        sum += coefficients[i] * current_x_power;
        current_x_power *= x;
    }
    return sum;
}

void HigherOrderODE_Homogeneous_PowerSeriesSolver::printSeries() const
{
    cout << "\nSeries solution: \n" << endl;
    cout << "y(x) = ";
    bool first = true;
    for (size_t i = 0; i < coefficients.size(); ++i)
    {
        if (coefficients[i] == 0.0)
        {
            continue;
        }
        if (!first && coefficients[i] > 0)
        {
            cout << " + ";
        }
        ...
    }
    cout << " + ... \n";
}
```

Code 43: *src/dsolve.cpp*

```

IVP Solve for  $y'' + 3y' + 2y = 0$  with  $y(0) = 1, y'(0) = 2$ 

The general solution is:
 $c1*e^{(-t)}+c2*e^{(-2*t)}$ 

The solution for the initial value problem is:
 $4*e^{(-t)}-3*e^{(-2*t)}$ 

 $y' = -4*e^{(-t)}+6*e^{(-2*t)}$ 

Critical value  $t_{\{m\}} = 0.405465$ 

Maximum value  $y_{\{m\}} = 1.33333$ 

```

Figure 3.67: The exact solution of $y'' + 3y' + 2y = 0$ computed with `secondorderlineardiff_eq_ivpsolution(1,3,2,y,t,0,1,2)` (`SymIntegration/Examples/Differential Equations/Test SymIntegration IVPSolve Second Order Linear Equations/main.cpp`).

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second
Order Linear Differential Equation with Constant Coefficients ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second
Order Linear Differential Equation with Constant Coefficients ]# ./main

( 2 ) y + ( 3 ) y^(1) + ( 1 ) y^(2) = 0

Initial conditions:
y(0) = 1
y^(1) (0) = 2

Series solution:

y(x) = 1 + 2x - 4x^2 + 3.33333x^3 - 1.83333x^4 + 0.766667x^5 - 0.261111x^6 + 0.0753968x^7 - 0.0189484x^8 + 0.00422178x^9 - 0.000845459x^10 + 0.0
0015382x^11 - 2.5645e-05x^12 + 3.94603e-06x^13 - 5.63764e-07x^14 + 7.51716e-08x^15 + ...

Time taken by function: 472 microseconds

```

Figure 3.68: The power series solution of $y'' + 3y' + 2y = 0$ computed with `class HigherOrderODE_Homogeneous_PowerSeriesSolver`, $n = 16$ (highest degree of the solution is at $n - 1$) (`SymIntegration/Examples/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential Equation with Constant Coefficients/main.cpp`).




Figure 3.69: The plot of the exact solution and the series solution with various terms for the differential equation $y'' + 3y' + 2y = 0$, we combine SymIntegration library with Hamzstplot, thus SymIntegration is used to compute the series solution and then plot it with Hamzstplot (**Hamzstplot/Examples with Makefile/Differential Equations/Plot 2D Comparing Series Solutions with Exact Solution for Second Order Homogeneous Linear Differential Equation/main.cpp**).

ii. Compute Power Series Solution for Homogeneous Second Order Linear Differential Equations with Variable Coefficients Near an Ordinary Point x_0

To solve a homogeneous second order linear differential equation with variable coefficients around a point x_0 , we must first classify the point x_0 . If x_0 is an ordinary point ($P(x_0) \neq 0$), we use a standard power series expansion. If x_0 is a regular singular point, use the Frobenius method.

The general homogeneous second order linear differential equation is structured as

$$P(x)y'' + Q(x)y' + R(x)y = 0$$

with the initial condition:

$$\begin{aligned} y(0) &= y_0 \\ y'(0) &= dy_0 \end{aligned}$$

1. For a homogeneous second order linear differential equation of the form:

$$P(x)y'' + Q(x)y' + R(x)y = 0$$

where $P(x)$, $Q(x)$, and $R(x)$ are polynomials or analytic functions, we find a series solution depending on the nature of the point x_0 where we expand the series.

2. Ordinary or Singular Points

- **Ordinary Point** ($P(x_0) \neq 0$): We can assume a direct Taylor power series centered at x_0 :

$$y(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n$$

Plugging this into the differential equation yields a recurrence relation that defines all a_n based on two arbitrary constants (a_0 and a_1), producing two linearly independent solutions $y_1(x)$ and $y_2(x)$. The general solution is $y(x) = c_1 y_1(x) + c_2 y_2(x)$.

- **Regular Singular Point** ($P(x_0) = 0$, but limits for $\frac{(x-x_0)Q}{P}$ and $\frac{(x-x_0)^2 R}{P}$ are finite): We use the Frobenius method, assuming a solution multiplied by a fractional or real power r :

$$y(x) = (x - x_0)^r \sum_{n=0}^{\infty} a_n (x - x_0)^n$$

An indicial equation determines the value of r , leading to the series expansion.

Since there is no universal formula for the recurrence relation, the specific recurrence relation for the coefficients a_n changes completely depending on the functions $P(x)$, $Q(x)$, and $R(x)$. The clean principle of superposition and straightforward power series substitution generally fails or becomes non-algorithmic for second order nonlinear equations.

Below is the full mathematical mechanism to compute the power series solution of a homogeneous second order linear differential equation around a point x_0 :

1. Point Classification

First, write the differential equation in standard form:

$$y'' + p(x)y' + q(x)y = 0$$

where $p(x) = \frac{Q(x)}{P(x)}$ and $q(x) = \frac{R(x)}{P(x)}$. We will compute the power series solution about the given point x_0 , if $x_0 \neq 0$ then the center will be shifted by x_0 .

- **Power Series Method (Ordinary Point)**

If $p(x)$ and $q(x)$ are analytic at x_0 ($P(x_0) \neq 0$), we assume the solution as a standard Taylor series:

$$y(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n$$

- **Frobenius Method (Regular Singular Point)**

If $p(x)$ or $q(x)$ diverges at x_0 ($P(x_0) = 0$), but $\lim_{x \rightarrow x_0} (x - x_0)p(x)$ and $\lim_{x \rightarrow x_0} (x - x_0)^2 q(x)$ finite (remain analytic), x_0 is a regular singular point. We assume a Frobenius series:

$$y(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^{n+r}, \quad a_0 \neq 0$$

2. Compute Derivatives

After we have classified the point x_0 and obtain the $y(x)$, we now differentiate the assumed series term-by-term to find expressions for y' and y'' .

- **Power Series Method (Ordinary Point)**

$$y(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n (x - x_0)^{n-1}$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n (x - x_0)^{n-2}$$

Substituting these equations above into the differential equation yields a recurrence relation determining a_n for $n \geq 2$ based on arbitrary constants a_0 and a_1 .

- **Frobenius Method (Regular Singular Point)**

$$y(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^{n+r}, \quad a_0 \neq 0$$

$$y'(x) = \sum_{n=0}^{\infty} (n+r) a_n (x - x_0)^{n+r-1}$$

$$y''(x) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n (x - x_0)^{n+r-2}$$

Plugging this into the differential equation produces the indicial equation (a quadratic equation in r). Solving it gives roots r_1 and r_2 , which dictate the structure of the recurrence relation for a_n .

3. Shifting the Index of Summation, Substitute and Align Powers

Substitute $y(x)$, $y'(x)$, and $y''(x)$ back into the original differential equation. To combine the terms into a single summation, we must perform two adjustments:

- **Shift Indices:** Change the dummy variables (e.g., let $k = n - 2$) so that every term features the exact same power of x (typically x^k or $(x - x_0)^k$).

- **Align Starting Limits:** Expand the lower terms of the series explicitly until all remaining summations start at the same index value.

Frobenius Method (Regular Singular Point)

After we substitute y, y', y'' into the differential equation multiplied by x^2 to clear denominators:

$$x^2 y'' + x \cdot p(x) y' + q(x) y = 0$$

Expand $p(x) = \sum_{k=0}^{\infty} p_k x^k$ and $q(x) = \sum_{k=0}^{\infty} q_k x^k$ as standard Taylor series.

Combine Cauchy products of series so every summation term matches a unified power exponent of x^{n+r} . If a summation index starts at $n = 0$ but evaluates with an offset x^{n+r-2} , re-index or shift the summation counter $n \rightarrow n + 2$ (adjusting the summation bounds and general coefficient subscripts simultaneously) to factor out a common x^{n+r} baseline.

4. Solve the Indicial and Recurrence Equations

Factor out the common power of x^k . For the differential equation to equal zero for all x , the coefficient of every power of x must independently equal zero.

- **Power Series Method (Ordinary Point)**

$$\sum_{k=0}^{\infty} f(a_k, a_{k+1}, a_{k+2}) x^k = 0$$

Because this equation must equal zero for all values of x , the identity principle states that the coefficients of each power of x must independently equal zero. Set the inner expression to zero to obtain the recurrence relation:

$$f(a_k, a_{k+1}, a_{k+2}) = 0$$

Solve the recurrence relation for higher coefficients, to express the higher-order coefficients $(a_2, a_3, a_4, \dots)$ in terms of the arbitrary starting constraints a_0 and a_1

- **Frobenius Method (Regular Singular Point)**

Setting the lowest power coefficient (x^r) to zero yields a quadratic equation in terms of r / the indicial equation:

$$f(r) = 0$$

$$r(r-1) + p_0 r + q_0 = 0$$

Solve this to find two roots, r_1 and r_2 (where $\text{Re}(r_1) \geq \text{Re}(r_2)$).

Equate the general coefficient of x^{n+r} to zero for $n \geq 1$, producing a linear relation linking a_n to preceding coefficients $a_0, a_1, \dots, a_{n-1}$:

$$a_n(n+r)(n+r-1) + \sum_{k=0}^n [p_{n-k}(k+r) + q_{n-k}] a_k = 0$$

(a) **Case 1: $r_1 - r_2$ is not an integer**

When we get the roots that do not differ by an integer, both solutions are pure power series:

$$y_1(x) = (x - x_0)^{r_1} \sum_{n=0}^{\infty} a_n (x - x_0)^n, \quad (a_0 \neq 0)$$

$$y_2(x) = (x - x_0)^{r_2} \sum_{n=0}^{\infty} b_n (x - x_0)^n, \quad (b_0 \neq 0)$$

(b) **Case 2: $r_1 = r_2$**

When the roots are identical, the second linearly independent solution introduces a logarithmic term:

$$y_1(x) = (x - x_0)^{r_1} \sum_{n=0}^{\infty} a_n x^n, \quad (a_0 \neq 0)$$

$$y_2(x) = y_1(x - x_0) \ln(x - x_0) + (x - x_0)^{r_1} \sum_{n=1}^{\infty} b_n (x - x_0)^n$$

(c) **Case 3: $r_1 - r_2$ is a positive integer**

When the roots differ by a positive integer ($r_1 - r_2 \in \mathbb{N}, \mathbb{N} \in \mathbb{Z}^+$), the second solution may contain a logarithmic term depending on whether a coefficient log-jam occurs:

$$y_1(x) = (x - x_0)^{r_1} \sum_{n=0}^{\infty} a_n (x - x_0)^n, \quad (a_0 \neq 0)$$

$$y_2(x) = C y_1(x) \ln(x - x_0) + (x - x_0)^{r_2} \sum_{n=0}^{\infty} b_n (x - x_0)^n, \quad (b_0 \neq 0)$$

where C can be 0.

$r_1 - r_2$	First solution ($y_1(x)$)	Second solution ($y_2(x)$)
Not an integer ($r_1 - r_2 \notin \mathbb{Z}$)	$\sum_{n=0}^{\infty} a_n x^{n+r_1}$	$\sum_{n=0}^{\infty} b_n x^{n+r_2}$
Equal roots ($r_1 = r_2$)	$\sum_{n=0}^{\infty} a_n x^{n+r_1}$	$y_1(x) \ln(x) + \sum_{n=1}^{\infty} b_n x^{n+r_1}$
differ by a positive integer ($r_1 - r_2 \in \mathbb{N}$)	$\sum_{n=0}^{\infty} a_n x^{n+r_1}$	$C y_1(x) \ln(x) + \sum_{n=0}^{\infty} b_n x^{n+r_2}$

Table 3.3: Depending on the difference $\Delta = r_1 - r_2$, the fundamental solutions $y_1(x)$ and $y_2(x)$ take different forms.

5. Compute the Final General Solution

$$y(x) = a_0 y_1(x) + a_1 y_2(x)$$

where $y_1(x)$ and $y_2(x)$ are two linearly independent fundamental solutions.

We would like to demonstrate a class in SymIntegration that we have created:

class SecondOrderODE_Homogeneous_PowerSeriesSolver

this **class** is able to compute the power series solution of a homogeneous second order linear differential equation with variable coefficients near an ordinary point (not singular), we name linear differential equation as ODE (Ordinary Differential Equation).

We will use this **class** to compute the solution for the famous Airy differential equation

$$y'' - xy = 0$$

We have computed the solution near $x_0 = 0$ first (no shifting), then near a regular point $x_0 = 1$ in the examples above under the **Series Solutions Near an Ordinary Point** part. In this section we will use SymIntegration to compute the series solutions.

A power series solution for a homogeneous second order linear differential equation with variable coefficients, formulated as $P(x)y'' + Q(x)y' + R(x)y = 0$, can be computed systematically by shifting the variable to center the equations around x_0 , and then evaluating Taylor expansion derivatives recursively.

The general solution of Airy's equation near $x_0 = 0$ is

$$y = a_0 \left[1 + \frac{x^3}{2 \cdot 3} + \frac{x^6}{2 \cdot 3 \cdot 5 \cdot 6} + \cdots + \frac{x^{3n}}{2 \cdot 3 \cdots (3n-1)(3n)} + \cdots \right] \\ + a_1 \left[x + \frac{x^4}{3 \cdot 4} + \frac{x^7}{3 \cdot 4 \cdot 6 \cdot 7} + \cdots + \frac{x^{3n+1}}{3 \cdot 4 \cdots (3n)(3n+1)} + \cdots \right]$$

it is the same equation as Eq. (3.279), that is composed by the function $Ai(x)$ and $Bi(x)$, two linearly independent solutions to the Airy differential equation.

$$Ai(x) = 1 + \frac{x^3}{2 \cdot 3} + \frac{x^6}{2 \cdot 3 \cdot 5 \cdot 6} + \cdots + \frac{x^{3n}}{2 \cdot 3 \cdots (3n-1)(3n)} + \cdots \\ Bi(x) = x + \frac{x^4}{3 \cdot 4} + \frac{x^7}{3 \cdot 4 \cdot 6 \cdot 7} + \cdots + \frac{x^{3n+1}}{3 \cdot 4 \cdots (3n)(3n+1)} + \cdots$$

the general solution of Airy's equation near $x_0 = 1$ is

$$y = a_0 \left[1 + \frac{(x-1)^2}{2} + \frac{(x-1)^3}{6} + \frac{(x-1)^4}{24} + \frac{(x-1)^5}{30} + \cdots \right] \\ + a_1 \left[(x-1) + \frac{(x-1)^3}{6} + \frac{(x-1)^4}{12} + \frac{(x-1)^5}{120} + \cdots \right]$$

it is the same equation as Eq. (3.282) with

$$Ai(x) = 1 + \frac{(x-1)^2}{2} + \frac{(x-1)^3}{6} + \frac{(x-1)^4}{24} + \frac{(x-1)^5}{30} + \cdots \\ Bi(x) = (x-1) + \frac{(x-1)^3}{6} + \frac{(x-1)^4}{12} + \frac{(x-1)^5}{120} + \cdots$$

In SymIntegration if we want to compute the power series solution for $Ai(x)$, then we will set the initial condition as

$$y(0) = 1 \\ y'(0) = 0$$

if we want to compute the power series solution for $Bi(x)$, then we will set the initial condition as

$$y(0) = 0 \\ y'(0) = 1$$

Below is the **main.cpp** source code to compute the $Ai(x)$ solution of the Airy differential equation $y'' - xy = 0$ near $x_0 = 0$.

```
int main(void)
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    Polynomialcoeff P({1.0, 0.0, 0.0});
    Polynomialcoeff Q({0.0, 0.0, 0.0});
    Polynomialcoeff R({0.0, -1.0, 0.0});

    double x0 = 0.0;

    double y0 = 1.0;
    double dy0 = 0.0;
    int num_terms = 14;

    double test_x = 0.45;

    SecondOrderODE_Homogeneous_PowerSeriesSolver solver(P, Q, R, x0, y0, dy0);
    solver.computeSeries(num_terms);

    //cout << std::fixed << std::setprecision(10);
    //solver.printCoefficients();
    solver.printSolution();

    double approximation = solver.evaluateAt(test_x, 10);
    cout << "\nx = " << test_x << std::endl;
    cout << "Power Series Approximation at x : " << std::fixed << std::
        setprecision(6) << approximation << endl;

    ...

    return 0;
}
```

Code 44: *SymIntegration/Examples/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential Equation with Variable Coefficients/main.cpp*

IV. Series Solutions of Second Order Linear Equations

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear D
ifferential Equation with Variable Coefficients ]# ./main
( 1 ) y'' + ( 0 ) y' + ( - 1x ) y
Initial conditions: y(0) = 1 , y'(0) = 0

x_{0} = 0
Computed Power Series Coefficients:
c_{0} = 1
c_{1} = 0
c_{2} = -0
c_{3} = 0.166667
c_{4} = -0
c_{5} = -0
c_{6} = 0.00555556
c_{7} = -0
c_{8} = -0
c_{9} = 7.71605e-05
c_{10} = -0
c_{11} = -0
c_{12} = 5.84549e-07
c_{13} = -0

Series solution:

y(x) = 1 + 0.166667*(x - 0)^3 + 0.00555556*(x - 0)^6 + 7.71605e-05*(x - 0)^9 + 5.84549e-07*(x - 0)^12 + ...

x = 0.45
Power Series Approximation at x : 1.015234
Time taken by function: 851 microseconds

```

Figure 3.70: The power series solution of Airy differential equation $y'' - xy = 0$ for $Ai(x)$ with initial condition $y(0) = 1$, $y'(0) = 0$ computed with class `SecondOrderODE_Homogeneous_PowerSeriesSolver`, $n = 14$ (highest degree of the solution is at $n - 1$) near a regular point $x_0 = 0$ (`SymIntegration/Examples/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential Equation with Variable Coefficients/main.cpp`).

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear D
ifferential Equation with Variable Coefficients ]# ./main
( 1 ) y'' + ( 0 ) y' + ( - 1x ) y
Initial conditions: y(0) = 0 , y'(0) = 1

x_{0} = 0
Computed Power Series Coefficients:
c_{0} = 0
c_{1} = 1
c_{2} = -0
c_{3} = -0
c_{4} = 0.0833333
c_{5} = -0
c_{6} = -0
c_{7} = 0.00198413
c_{8} = -0
c_{9} = -0
c_{10} = 2.20459e-05
c_{11} = -0
c_{12} = -0
c_{13} = 1.4132e-07

Series solution:

y(x) = 1*(x - 0) + 0.0833333*(x - 0)^4 + 0.00198413*(x - 0)^7 + 2.20459e-05*(x - 0)^10 + 1.4132e-07*(x - 0)^13 + ...

x = 0.45
Power Series Approximation at x : 0.453425
Time taken by function: 861 microseconds

```

Figure 3.71: The power series solution of Airy differential equation $y'' - xy = 0$ for $Bi(x)$ with initial condition $y(0) = 0$, $y'(0) = 1$ computed with class `SecondOrderODE_Homogeneous_PowerSeriesSolver`, $n = 14$ (highest degree of the solution is at $n - 1$) near a regular point $x_0 = 0$ (`SymIntegration/Examples/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential Equation with Variable Coefficients/main.cpp`).

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential
Equation with Variable Coefficients ]# ./main
( 1 ) y'' + ( 0 ) y' + ( - 1x ) y
Initial conditions: y(0) = 1 , y'(0) = 0

x_{0} = 1
Computed Power Series Coefficients:
c_{0} = 1
c_{1} = 0
c_{2} = 0.5
c_{3} = 0.166667
c_{4} = 0.0416667
c_{5} = 0.0333333
c_{6} = 0.00694444
c_{7} = 0.00178571
c_{8} = 0.000719246
c_{9} = 0.000121252
c_{10} = 2.78329e-05
c_{11} = 7.64089e-06
c_{12} = 1.12943e-06
c_{13} = 2.27396e-07

Series solution:
y(x) = 1 + 0.5*(x - 1)^2 + 0.166667*(x - 1)^3 + 0.0416667*(x - 1)^4 + 0.0333333*(x - 1)^5 + 0.00694444*(x - 1)^6 + 0.00178571*(x - 1)^7 + 0.000719246*(x - 1)^8 + 0.000121
252*(x - 1)^9 + 2.78329e-05*(x - 1)^10 + 7.64089e-06*(x - 1)^11 + 1.12943e-06*(x - 1)^12 + 2.27396e-07*(x - 1)^13 + ...

x = 0.45
Power Series Approximation at x : 1.118827
Time taken by function: 877 microseconds

```

Figure 3.72: The power series solution of Airy differential equation $y'' - xy = 0$ for $Ai(x)$ with initial condition $y(0) = 1$, $y'(0) = 0$ computed with class `SecondOrderODE_Homogeneous_PowerSeriesSolver`, $n = 14$ (highest degree of the solution is at $n - 1$) near a regular point $x_0 = 1$ (`SymIntegration/Examples/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential Equation with Variable Coefficients/main.cpp`).

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential
Equation with Variable Coefficients ]# ./main
( 1 ) y'' + ( 0 ) y' + ( - 1x ) y
Initial conditions: y(0) = 0 , y'(0) = 1

x_{0} = 1
Computed Power Series Coefficients:
c_{0} = 0
c_{1} = 1
c_{2} = 0
c_{3} = 0.166667
c_{4} = 0.0833333
c_{5} = 0.00833333
c_{6} = 0.00833333
c_{7} = 0.00218254
c_{8} = 0.000297619
c_{9} = 0.000146054
c_{10} = 2.75573e-05
c_{11} = 4.03339e-06
c_{12} = 1.31524e-06
c_{13} = 2.02505e-07

Series solution:
y(x) = 1*(x - 1) + 0.166667*(x - 1)^3 + 0.0833333*(x - 1)^4 + 0.00833333*(x - 1)^5 + 0.00833333*(x - 1)^6 + 0.00218254*(x - 1)^7 + 0.000297619*(x - 1)^8 + 0.000146054*(x
- 1)^9 + 2.75573e-05*(x - 1)^10 + 4.03339e-06*(x - 1)^11 + 1.31524e-06*(x - 1)^12 + 2.02505e-07*(x - 1)^13 + ...

x = 0.45
Power Series Approximation at x : 0.468836
Time taken by function: 875 microseconds

```

Figure 3.73: The power series solution of Airy differential equation $y'' - xy = 0$ for $Bi(x)$ with initial condition $y(0) = 0$, $y'(0) = 1$ computed with class `SecondOrderODE_Homogeneous_PowerSeriesSolver`, $n = 14$ (highest degree of the solution is at $n - 1$) near a regular point $x_0 = 1$ (`SymIntegration/Examples/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential Equation with Variable Coefficients/main.cpp`).

Remark

Homogeneous second order linear ordinary differential equations with variable coefficients appear as $y'' + p(x)y' + q(x)y = 0$. To compute the solutions we can use advanced techniques like power series method, Frobenius method, or reduction of order because constant-root methods fail. They model advanced systems where parameters change with location or time, such as quantum mechanics, heat flow in non-uniform shapes, planetary orbits with changing gravity, non-uniform vibrating strings, electrical circuits with time-varying components and wave propagation. Some

major physical and engineering applications:

- **Vibrating Circular Membranes:** Modeled using Bessel's differential equation to find the harmonic frequencies and nodal lines of a struck drumhead. Bessel's equation arises when studying vibrations of circular membranes, acoustics in tapered horns, or heat flow in cylindrical coordinates.
- **Heat Conduction in Cylinders:** Solves steady-state and transient temperature distributions inside cylindrical pipes or solid metal rods.
- **Quantum Mechanics:** The Schrödinger equation reduces to this form for potentials like the quantum harmonic oscillator or hydrogen atom, yielding Legendre polynomials for angular wave functions. The behavior of a particle in a varying potential energy field reduces to a variable-coefficient ODE such as Airy's or Hermite's differential equations.
- **Electromagnetic Waveguides:** Determines transverse electric and magnetic field distributions inside coaxial cables and cylindrical wave guides.
- **Acoustics and Optics:** Explains diffraction patterns, sound waves traveling through a medium with changing density or temperature (like the atmosphere or ocean layers), planetary motion perturbations, and sound waves in tapered horns.
- **Astronomy and Aerospace:** Satellite motion and orbital mechanics under central forces with changing radial distances or atmospheric drag gradients. Legendre's differential equation models potential fields in spherical coordinates, gravity of planets, and electrostatics.
- **Non-Uniform Mechanical Systems:** Vibrations of a hanging chain or a variable cross-section beam where mass and tension change with position.
- **Electrical Engineering:** RLC circuits where resistance, inductance, or capacitance change over time (e.g., signal modulators, or time-varying filters). Transmission lines with position-dependent resistance, capacitance, or inductance use these models to compute voltage and current loss over varying distances.
- **Advanced Engineering:** Non-uniform transmission lines and tapered beams use these equations to track variable voltage drop, current dispersion, or structural load limits.

Common types of homogeneous second order linear ordinary differential equations with variable coefficients :

- **Cauchy-Euler Equation:** Used in scaling-invariant mechanical and aerospace stress problems, elasticity and radial fluid flow, it is defined by

$$x^2 y'' + axy' + by = 0$$

- **Bessel Equation:** Applied to heat conduction in cylinders and circular wave propagation, it is defined by

$$x^2 y'' + xy' + (x^2 - \nu^2)y = 0$$

- **Legendre Equation:** Found in spherical harmonic problems in electrostatics and quantum wave functions. Legendre's equation used in physics problems involving spherical coordinates (like gravitational and electrostatic potentials), it is defined by

$$(1 - x^2)y'' - 2xy' + n(n + 1)y = 0$$

iii. Compute Power Series Solution for Homogeneous Second Order Linear Differential Equations with Variable Coefficients Near a Singular Point x_0

iv. Maclaurin Series to Handle Non-Elementary Integral

[SI*] A Maclaurin series is an infinite polynomial expansion of a function centered specifically at zero, functioning as a special case of the Taylor series. It allows us to approximate complex functions like exponentials and trigonometric ratios using basic powers of x . The general formula for Maclaurin series for function $f(x)$ is defined as

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = f(0) + f'(0)x + \frac{f''(0)}{2!} + \frac{f'''(0)}{3!}x^3 + \dots$$

In the **Series Solutions Near an Ordinary Point** section we encounter Airy's differential equation which has series solutions, for a second order differential equation

$$\frac{d^2y}{dx^2} - xy = 0$$

the general solution can be expressed as a linear combination of two independent Airy functions, $Ai(x)$ and $Bi(x)$:

$$y(x) = c_1 Ai(x) + c_2 Bi(x)$$

with

$$Ai(x) = \frac{1}{\pi} \int_0^{\infty} \cos\left(\frac{t^3}{3} + xt\right) dt$$

and

$$Bi(x) = \frac{1}{\pi} \int_0^{\infty} \left[\sin\left(\frac{t^3}{3} + xt\right) + e^{-\frac{t^3}{3} + xt} \right] dt$$

The integral $\int \sin\left(\frac{t^3}{3} + xt\right) dt$ and $\int \cos\left(\frac{t^3}{3} + xt\right) dt$ cannot be evaluated in terms of elementary functions, thus we need to expand it by using Maclaurin series.

1. Definite Integral Form (Airy Function)

The boundary conditions are from 0 to ∞ , the integral is directly related to the Airy function of the first kind:

$$Ai(x) = \frac{1}{\pi} \int_0^{\infty} \cos\left(\frac{t^3}{3} + xt\right) dt$$

By multiplying both sides by π , we get:

$$\pi Ai(x) = \int_0^{\infty} \cos\left(\frac{t^3}{3} + xt\right) dt$$

2. Indefinite Integral via Series Expansion

To approximate or express the indefinite integral, you can use a Taylor series expansion centered at $t = 0$ (the Maclaurin series).

Expand the Cosine Term

Recall the standard Maclaurin series for $\cos(\theta)$:

$$\cos(\theta) = \sum_{n=0}^{\infty} \frac{(-1)^n \theta^{2n}}{(2n)!} = 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \dots$$

Substitute the Argument

Substitute $\theta = \left(\frac{t^3}{3} + xt\right)$:

$$\cos\left(\frac{t^3}{3} + xt\right) = 1 - \frac{1}{2}\left(\frac{t^3}{3} + xt\right)^2 + \frac{1}{24}\left(\frac{t^3}{3} + xt\right)^4 - \dots$$

Expanding the first few terms:

$$\cos\left(\frac{t^3}{3} + xt\right) = 1 - \frac{1}{2}\left(\frac{t^6}{9} + \frac{2xt^4}{3} + x^2t^2\right) + \dots$$

$$\cos\left(\frac{t^3}{3} + xt\right) = 1 - \frac{x^2}{2}t^2 - \frac{x}{3}t^4 - \frac{1}{18}t^6 + \dots$$

Integrate Term-by-Term

Integrating each power of t with respect to t :

$$\begin{aligned}\int \cos\left(\frac{t^3}{3} + xt\right) dt &= \int \left(1 - \frac{x^2}{2}t^2 - \frac{x}{3}t^4 - \frac{1}{18}t^6 + \dots\right) dt \\ &= t - \frac{x^2}{6}t^3 - \frac{x}{15}t^5 - \frac{1}{126}t^7 + \dots + C\end{aligned}$$

For the $\int \sin\left(\frac{t^3}{3} + xt\right) dt$:

1. Expand the Sine Term

Recall the Maclaurin series for the sine term:

$$\sin(\theta) = \sum_{n=0}^{\infty} \frac{(-1)^n \theta^{2n+1}}{(2n+1)!}$$

Substitute $\theta = \frac{t^3}{3} + xt$

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} \left(\frac{t^3}{3} + xt\right)^{2n+1}$$

2. Decompose with Binomial Theorem

$$\begin{aligned}\left(\frac{t^3}{3} + xt\right)^{2n+1} &= \sum_{k=0}^{2n+1} \binom{2n+1}{k} \left(\frac{t^3}{3}\right)^{2n+1-k} (xt)^k \\ \left(\frac{t^3}{3} + xt\right)^{2n+1} &= \sum_{k=0}^{2n+1} \binom{2n+1}{k} \frac{x^k}{3^{2n+1-k}} t^{6n+3-2k}\end{aligned}$$

3. Integrate

integrate it with respect to variable t :

$$\begin{aligned}\int \sin\left(\frac{t^3}{3} + xt\right) dt &= \sum_{n=0}^{\infty} \sum_{k=0}^{2n+1} \frac{(-1)^n}{(2n+1)!} \binom{2n+1}{k} \frac{x^k}{3^{2n+1-k}} \int t^{6n+3-2k} dt \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{2n+1} \frac{(-1)^n \binom{2n+1}{k} x^k t^{6n+4-2k}}{(2n+1)! \cdot 3^{2n+1-k} (6n+4-2k)} + C\end{aligned}$$

Thus the linearly independent series components:

$$\begin{aligned} y_1(x) &= 1 + \sum_{n=1}^{\infty} \frac{x^{3n}}{2 \cdot 3 \cdot 5 \cdot 6 \dots (3n-1)(3n)}, \quad (\text{Associated with } Ai(x)) \\ &= 1 + \sum_{n=1}^{\infty} \frac{x^{3n}}{3^n n! [2 \cdot 5 \dots (3n-1)]} \end{aligned}$$

$$\begin{aligned} y_2(x) &= x + \sum_{n=1}^{\infty} \frac{x^{3n+1}}{3 \cdot 4 \cdot 6 \cdot 7 \dots (3n)(3n+1)}, \quad (\text{Associated with } Bi(x)) \\ &= x + \sum_{n=1}^{\infty} \frac{x^{3n+1}}{3^n n! [1 \cdot 4 \dots (3n+1)]} \end{aligned}$$

Theorem 3.12: The Binomial Theorem

The Binomial theorem provides a fast way to expand any two-term expression raised to a power, the coefficients for each term is computed with Pascal' triangle or combination rule.

The expansion of $(x + y)^n$ with integer $n \geq 0$:

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$$

with

- n as the power or exponent, n is an integer.
- x and y as the two terms inside the parentheses.
- $\binom{n}{k}$ as the Binomial coefficient (n choose k).
- Exponents of x start at n and decrease by one each step $(n, n-1, \dots, 0)$.
- Exponents of y start at 0 and increase by one each step $(0, 1, \dots, n)$.

Definition 3.1: Bernoulli Numbers

Bernoulli numbers (B_n) form a sequence of rational numbers used in number theory, calculus, and expansions of trigonometric functions. They are most easily defined via an Exponential Generating Function or computed step-by-step using a recursive formula.

The Bernoulli numbers are the coefficients in the Taylor series expansion of the function $\frac{x}{e^x - 1}$

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}$$

To compute B_n recursively starting with $B_0 = 1$, use the identity for $n \geq 1$:

$$\sum_{k=0}^n \binom{n+1}{k} B_k = 0$$

which rearranges to the explicit iterative formula:

$$B_n = -\frac{1}{n+1} \sum_{k=0}^{n-1} \binom{n+1}{k} B_k$$

The first 7 terms of Bernoulli numbers:

$$B_0 = 1$$

$$B_1 = -0.5$$

$$B_2 = 0.166667$$

$$B_3 = 0$$

$$B_4 = -0.0333333$$

$$B_5 = 0$$

$$B_6 = 0.0238095$$

The Bernoulli numbers have appeared in many important results, including the series expansions of trigonometric and hyperbolic trigonometric functions, the Euler-Maclaurin summation formula, the evaluation of the Riemann zeta function, and Fermat's Last Theorem.

[SI*] We will list the Maclaurin series for several functions:

$$\begin{aligned}
 \cos(x) &= \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots, \quad -\infty < x < \infty \\
 \sin(x) &= \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots, \quad -\infty < x < \infty \\
 \tan(x) &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1} 2^{2n} (2^{2n-1} - 1) B_{2n}}{(2n)!} x^{2n-1} = x + \frac{x^3}{3} + \frac{2x^5}{15} + \frac{16x^7}{315} + \dots, \quad |x| < \frac{\pi}{2} \\
 \csc(x) &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1} 2(2^{2n-1} - 1) B_{2n}}{(2n)!} x^{2n-1} = \frac{1}{x} + \frac{x}{6} + \frac{7x^3}{360} + \frac{31x^5}{15120} + \dots, \quad 0 < |x| < \pi \\
 \sec(x) &= \sum_{n=0}^{\infty} \frac{E_{2n}}{(2n)!} x^{2n} = 1 + \frac{x^2}{2} + \frac{5x^4}{24} + \frac{61x^6}{720} + \dots, \quad |x| < \frac{\pi}{2} \\
 \cot(x) &= \sum_{n=0}^{\infty} \frac{(-1)^n 2^{2n} B_{2n}}{(2n)!} x^{2n-1} = \frac{1}{x} - \frac{x}{3} + \frac{x^3}{45} - \frac{2x^5}{945} - \dots, \quad 0 < |x| < \pi \\
 \sinh(x) &= \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!} = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \dots, \quad x \in \mathbb{C} \text{ or } x \in \mathbb{R} \\
 \cosh(x) &= \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \dots, \quad x \in \mathbb{C} \text{ or } x \in \mathbb{R} \\
 \tanh(x) &= \sum_{n=1}^{\infty} \frac{2^{2n} (2^{2n} - 1) B_{2n}}{(2n)!} x^{2n-1} = x - \frac{x^3}{3} + \frac{2}{15} x^5 - \frac{17}{315} x^7 + \dots, \quad |x| < \frac{\pi}{2} \\
 e^x &= \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \\
 \ln(1+x) &= \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots, \quad -1 < x \leq 1
 \end{aligned}$$

B_{2n} is Bernoulli numbers and E_n is Euler numbers, the function $\ln(x)$ has no Maclaurin series because $\ln(x)$ is undefined at $x = 0$. Since $\sinh(x) = \frac{e^x - e^{-x}}{2}$ and $\cosh(x) = \frac{e^x + e^{-x}}{2}$, we can substitute the known Maclaurin series for e^x and e^{-x} and add or subtract them term by term.

If we want to compute the Maclaurin series for $f(x) = \ln(x+2)$, we will need to use substitution and using the logarithm arithmetic rule:

$$\ln\left(2\left(1 + \frac{x}{2}\right)\right) = \ln(2) + \ln\left(1 + \frac{x}{2}\right)$$

By substituting $u = \frac{x}{2}$, we will have

$$\ln(1+u) = u - \frac{u^2}{2} + \frac{u^3}{3} - \frac{u^4}{4} + \dots$$

thus

$$\ln\left(1 + \frac{x}{2}\right) = \frac{x}{2} - \frac{x^2}{8} + \frac{x^3}{24} - \frac{x^4}{64} + \dots$$

Finally, we will obtain the Maclaurin series for $\ln(x+2)$

$$\begin{aligned}\ln(x+2) &= \ln\left(2\left(1+\frac{x}{2}\right)\right) \\ &= \ln(2) + \ln\left(1+\frac{x}{2}\right) \\ &= \ln(2) + \frac{x}{2} - \frac{x^2}{8} + \frac{x^3}{24} - \frac{x^4}{64} + \dots \\ &= \ln(2) + \sum_{n=1}^{\infty} \frac{(-1)^{n-1}x^n}{n2^n}\end{aligned}$$

v. Frobenius Method

[SI*] In mathematics, the method of Frobenius, named after Ferdinand Georg Frobenius, is a way to find an infinite series solution for a linear second-order ordinary differential equation of the form

$$x^2y'' + p(x)xy' + q(x)y = 0 \quad (3.285)$$

with $y' = \frac{dy}{dx}$ and $y'' = \frac{d^2y}{dx^2}$.

In the vicinity of the regular singular point $x = 0$. One can divide by x^2 to obtain a differential equation of the form

$$y'' + \frac{p(x)}{x}y' + \frac{q(x)}{x^2}y = 0$$

which will not be solvable with regular power series methods if either $\frac{p(x)}{x}$ or $\frac{q(x)}{x^2}$ is not analytic at $x = 0$. The Frobenius method enables one to create a power series solution to such a differential equation, provided that $p(x)$ and $q(x)$ are themselves analytic at 0 or, being analytic elsewhere, both their limits at 0 exist (and are finite).

[SI*] Frobenius' contribution was not so much in all the possible forms of the series solutions involved. These forms had all been established earlier, by Lazarus Fuchs. The indicial polynomial and its role had also been established by Fuchs.

A first contribution by Frobenius to the theory was to show that—as regards a first, linearly independent solution, which then has the form of an analytical power series multiplied by an arbitrary power r of the independent variable—the coefficients of the generalized power series obey a recurrence relation, so that they can always be straightforwardly calculated.

A second contribution by Frobenius was to show that, in cases in which the roots of the indicial equation differ by an integer, the general form of the second linearly independent solution can be obtained by a procedure which is based on differentiation with respect to the parameter r .

A large part of Frobenius' 1873 publication was devoted to proofs of convergence of all the series involved in the solutions, as well as establishing the radii of convergence of these series.

[SI*] The method of Frobenius is to seek a power series solution of the form

$$y(x) = x^r \sum_{k=0}^{\infty} A_k x^k, \quad A_0 \neq 0$$

Differentiating:

$$y'(x) = \sum_{k=0}^{\infty} (k+r) A_k x^{k+r-1}$$

$$y''(x) = \sum_{k=0}^{\infty} (k+r-1)(k+r) A_k x^{k+r-2}$$

Substituting the above differentiation into the original ODE Eq. (3.283):

$$\begin{aligned} & x^2 \sum_{k=0}^{\infty} (k+r-1)(k+r) A_k x^{k+r-2} + xp(x) \sum_{k=0}^{\infty} (k+r) A_k x^{k+r-1} + q(x) \sum_{k=0}^{\infty} A_k x^{k+r} \\ &= \sum_{k=0}^{\infty} (k+r-1)(k+r) A_k x^{k+r} + p(x) \sum_{k=0}^{\infty} (k+r) A_k x^{k+r} + q(x) \sum_{k=0}^{\infty} A_k x^{k+r} \\ &= \left[(k+r-1)(k+r) A_k x^{k+r} + p(x)(k+r) A_k x^{k+r} + q(x) A_k x^{k+r} \right] \\ &= \sum_{k=0}^{\infty} \left[(k+r-1)(k+r) A_k x^{k+r} + p(x)(k+r) A_k x^{k+r} + q(x) A_k x^{k+r} \right] \\ &= \sum_{k=0}^{\infty} [(k+r-1)(k+r) + p(x)(k+r) + q(x)] A_k x^{k+r} \\ &= [r(r-1) + p(x)r + q(x)] A_0 x^r \\ &+ \sum_{k=1}^{\infty} [(k+r-1)(k+r) + p(x)(k+r) + q(x)] A_k x^{k+r} = 0 \end{aligned}$$

vi. The Airy Differential Equation

[SI*] In mathematics, the Airy function $Ai(x)$ is a special function named after the British astronomer George Biddell Airy. The function $Ai(x)$ and the related function $Bi(x)$ are linearly independent solutions to the differential equation

$$\frac{d^2 y}{dx^2} - xy = 0$$

known as the Airy equation or the Stokes equation. This is a second-order linear ordinary differential equation with a non-constant coefficient (x).

Because the solution of the linear differential equation

$$\frac{d^2 y}{dx^2} - ky = 0$$

is oscillatory for $k < 0$ and exponential for $k > 0$, the Airy functions are oscillatory for $x < 0$ and exponential for $x > 0$. In fact, the Airy equation is the simplest second-order linear differential equation with a turning point (a point where the character of the solutions changes from oscillatory to exponential).

[SI*] The general solution to the Airy differential equation $y'' - xy = 0$ is expressed as a linear combination of two independent Airy functions, $Ai(x)$ and $Bi(x)$:

$$y(x) = c_1 Ai(x) + c_2 Bi(x)$$

where c_1 and c_2 are real constants, $Ai(x)$ is the Airy function of the first kind, which decays exponentially as $x \rightarrow \infty$ and oscillates as $x \rightarrow -\infty$. $Bi(x)$ is the Airy function of the second kind, which grows exponentially as $x \rightarrow \infty$ and oscillates as $x \rightarrow -\infty$. Both $Ai(x)$ and $Bi(x)$ share the same amplitude and frequency of oscillation when $x \rightarrow -\infty$, but they differ in phase by $\frac{\pi}{2}$.

For real values of x , the Airy function of the first kind can be defined by the improper Riemann integral

$$Ai(x) = \frac{1}{\pi} \int_0^{\infty} \cos\left(\frac{t^3}{3} + xt\right) dt$$

which converges by Dirichlet's test. Up to scalar multiplication, $Ai(x)$ is the solution subject to the condition $y \rightarrow 0$ as $x \rightarrow \infty$. The standard choice for the other solution is the Airy function of the second kind, denoted by $Bi(x)$. It is defined as the solution with the same amplitude of oscillation as $Ai(x)$ as $x \rightarrow \infty$ which differs in phase by $\frac{\pi}{2}$:

$$Bi(x) = \frac{1}{\pi} \int_0^{\infty} \left[\sin\left(\frac{t^3}{3} + xt\right) + e^{-\frac{t^3}{3} + xt} \right] dt$$

Power Series Solutions

Because the coefficient is variable, the equation is solved using a power series

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

The recurrence relation:

$$a_{n+2} = \frac{a_{n-1}}{(n+1)(n+2)}, \quad \text{for } n \geq 1$$

Thus the linearly independent series components:

$$\begin{aligned} y_1(x) &= 1 + \sum_{n=1}^{\infty} \frac{x^{3n}}{2 \cdot 3 \cdot 5 \cdot 6 \dots (3n-1)(3n)}, \quad (\text{Associated with } Ai(x)) \\ &= 1 + \sum_{n=1}^{\infty} \frac{x^{3n}}{3^n n! [2 \cdot 5 \dots (3n-1)]} \\ y_2(x) &= x + \sum_{n=1}^{\infty} \frac{x^{3n+1}}{3 \cdot 4 \cdot 6 \cdot 7 \dots (3n)(3n+1)}, \quad (\text{Associated with } Bi(x)) \\ &= x + \sum_{n=1}^{\infty} \frac{x^{3n+1}}{3^n n! [1 \cdot 4 \dots (3n+1)]} \end{aligned}$$

For negative x ($x < 0$) solutions are oscillatory, behaving similarly to sine and cosine functions with changing amplitude. For positive x ($x > 0$) $Ai(x)$ decays exponentially toward zero, while $Bi(x)$ grows exponentially.

[SI*] The Airy differential equation ($y'' - xy = 0$) is applied widely in physics, quantum mechanics, optics, and wave propagation to model phenomena near turning points, such as light diffraction, supernumerary rainbows, and electrons in triangular potential wells. Its solutions, called Airy functions, describe fields that change from oscillating to exponential behavior, it is able to describe physical phenomena where wave behavior transitions between oscillatory and exponential decay.

Application of the Airy differential equation:

1. Quantum Mechanics

- **Triangular Wells:** Solves the Schrödinger equation for particles in a linear constant force field or confined by a triangular potential well. Describes electron wave functions in triangular potential wells.
- **Semiconductor Heterojunctions:** Explains the behavior and trapping of electrons near interfaces under uniform electric fields.
- **Uniform Approximations:** Approximates wave behavior via the WKB method near classical turning points where potential equals energy.
- **WKB Approximation:** Provides uniform semiclassical approximations near classical turning points where potentials change linearly.

2. Optics and Wave Propagation

- **Optical Caustics:** Models light intensity patterns near bright edges and caustics, such as the formation of rainbows.
- **Electromagnetic Diffraction:** Describes light and radio wave diffraction patterns near shadow boundaries.

3. Acoustics and Electromagnetism

- **Underwater Sound:** Computes transparent boundary conditions for parabolic wave equations simulating sound in the ocean.
- **Radio Waves:** Analyzes tropospheric radar propagation and radio wave signals curving through the atmosphere.
- **Atmospheric and Underwater Acoustics:** Used in parabolic equation simulations for tracking sound or radar wave propagation through the troposphere and oceans.

vii. The Hermite Differential Equation

[SI*] In mathematics, the Hermite polynomials are a classical orthogonal polynomial sequence. The polynomials arise in:

- Signal processing as Hermitian wavelets for wavelet transform analysis.
- Probability, such as the Edgeworth series, as well as in connection with Brownian motion.
- Combinatorics, as an example of an Appell sequence, obeying the umbral calculus.
- Numerical analysis as Gaussian quadrature.
- Physics, where they give rise to the eigenstates of the quantum harmonic oscillator; and they also occur in some cases of the heat equation (when the term xu_x is present).
- Systems theory in connection with nonlinear operations on Gaussian noise.
- Random matrix theory in Gaussian ensembles.

Hermite polynomials were defined by Pierre-Simon Laplace in 1810, though in scarcely recognizable form, and studied in detail by Pafnuty Chebyshev in 1859. Chebyshev's work was overlooked, and they were named later after Charles Hermite, who wrote on the polynomials in 1864, describing them as new. They were not new, although Hermite was the first to define the multidimensional polynomials.

Like the other classical orthogonal polynomials, the Hermite polynomials can be defined from several different starting points. Noting from the outset that there are two different standardizations in common use, one convenient method is as follows:

- The "**probabilist's Hermite polynomials**" are given by

$$He_n(x) = (-1)^n e^{\frac{x^2}{2}} \frac{d^n}{dx^n} e^{-\frac{x^2}{2}}$$

- The "**physicist's Hermite polynomials**" are given by

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2}$$

These equations have the form of a Rodrigues' formula and can also be written as

$$He_n(x) = \left(x - \frac{d}{dx} \right)^n \cdot 1$$

$$H_n(x) = \left(2x - \frac{d}{dx} \right)^n \cdot 1$$

The two definitions are not exactly identical; each is rescaling of the other:

$$He_n(x) = 2^{-\frac{n}{2}} H_n \left(\frac{x}{\sqrt{2}} \right)$$

$$H_n(x) = 2^{\frac{n}{2}} He_n \left(\sqrt{2}x \right)$$

The polynomials He_n are sometimes denoted by H_n , especially in probability theory, because

$$\frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

is the probability density function for the normal distribution with expected value 0 and standard deviation 1. The probabilist's Hermite polynomials are also called the monic Hermite polynomials, because they are monic.

- The first eleven probabilist's Hermite polynomials are:

$$\begin{aligned} He_0(x) &= 1 \\ He_1(x) &= x \\ He_2(x) &= x^2 - 1 \\ He_3(x) &= x^3 - 3x \\ He_4(x) &= x^4 - 6x^2 + 3 \\ He_5(x) &= x^5 - 10x^3 + 15x \\ He_6(x) &= x^6 - 15x^4 + 45x^2 - 15 \\ He_7(x) &= x^7 - 21x^5 + 105x^3 - 105x \\ He_8(x) &= x^8 - 28x^6 + 210x^4 - 420x^2 + 105 \\ He_9(x) &= x^9 - 36x^7 + 378x^5 - 1260x^3 + 945x \\ He_{10}(x) &= x^{10} - 45x^8 + 630x^6 - 3150x^4 + 4725x^2 - 945 \end{aligned}$$

- The first eleven physicist's Hermite polynomials are:

$$H_0(x) = 1$$

$$H_1(x) = 2x$$

$$H_2(x) = 4x^2 - 2$$

$$H_3(x) = 8x^3 - 12x$$

$$H_4(x) = 16x^4 - 48x^2 + 12$$

$$H_5(x) = 32x^5 - 160x^3 + 120x$$

$$H_6(x) = 64x^6 - 480x^4 + 720x^2 - 120$$

$$H_7(x) = 128x^7 - 1344x^5 + 3360x^3 - 1680x$$

$$H_8(x) = 256x^8 - 3584x^6 + 13440x^4 - 13440x^2 + 1680$$

$$H_9(x) = 512x^9 - 9216x^7 + 48384x^5 - 80640x^3 + 30240x$$

$$H_{10}(x) = 1024x^{10} - 23040x^8 + 161280x^6 - 403200x^4 + 302400x^2 - 30240$$

	Physicist's	Probabilist's
Symbol	H_n	He_n
Head coefficient	2^n	1
Differential operator	$(-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2}$	$(-1)^n e^{\frac{x^2}{2}} \frac{d^n}{dx^n} e^{-\frac{x^2}{2}}$
Orthogonal to	e^{-x^2}	$e^{-\frac{1}{2}x^2}$
Inner product	$\int H_m(x) H_n(x) \frac{e^{-x^2}}{\sqrt{\pi}} dx = 2^n n! \delta_{mn}$	$\int He_m(x) He_n(x) \frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx = n! \delta_{nm}$
Generating function	$e^{xt-t^2} = \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!}$	$e^{xt-\frac{1}{2}t^2} = \sum_{n=0}^{\infty} He_n(x) \frac{t^n}{n!}$
Rodrigues' formula	$\left(2x - \frac{d}{dx}\right)^n \cdot 1$	$\left(x - \frac{d}{dx}\right)^n \cdot 1$
Recurrence relation	$H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x)$	$He_{n+1}(x) = xHe_n(x) - nHe_{n-1}(x)$

Table 3.4: The differences between probabilist's Hermite polynomials and physicist's Hermite polynomials.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hermite Polynomials ]# make
g++ -c -o main.o main.cpp
g++ -o main -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hermite Polynomials ]# ./main
Physicist's Hermite polynomials:
H_{0}(x) = 1
H_{1}(x) = 2*x
H_{2}(x) = 4*x^2-2
H_{3}(x) = 8*x^3-12*x
H_{4}(x) = 16*x^4-48*x^2+12
H_{5}(x) = 32*x^5-160*x^3+120*x
H_{6}(x) = 64*x^6-480*x^4+720*x^2-120
H_{7}(x) = 128*x^7-1344*x^5+3360*x^3-1680*x
H_{8}(x) = 256*x^8-3584*x^6+13440*x^4-13440*x^2+1680
H_{9}(x) = 512*x^9-9216*x^7+48384*x^5-80640*x^3+30240*x
H_{10}(x) = 1024*x^10-23040*x^8+161280*x^6-403200*x^4+302400*x^2-30240

Probabilist's Hermite polynomials:
He_{0}(x) = 1
He_{1}(x) = x
He_{2}(x) = x^2-1
He_{3}(x) = x^3-3*x
He_{4}(x) = x^4-6*x^2+3
He_{5}(x) = x^5-10*x^3+15*x
He_{6}(x) = x^6-15*x^4+45*x^2-15
He_{7}(x) = x^7-21*x^5+105*x^3-105*x
He_{8}(x) = x^8-28*x^6+210*x^4-420*x^2+105
He_{9}(x) = x^9-36*x^7+378*x^5-1260*x^3+945*x
He_{10}(x) = x^10-45*x^8+630*x^6-3150*x^4+4725*x^2-945

Time taken by function: 196326 microseconds

```

Figure 3.74: The first eleven physicist's and probabilist's Hermite polynomials (SymIntegration/Examples/Differential Equations/Test SymIntegration Hermite Polynomials/main.cpp).




Figure 3.75: The probabilist's Hermite polynomials (Hamzstplot/Examples with Makefile/Differential Equation-s/Plot 2D Probabilist's Hermite Polynomials/main.cpp).




Figure 3.76: The physicist's Hermite polynomials (*Hamzstplot/Examples with Makefile/Differential Equations/-Plot 2D Physicist's Hermite Polynomials/main.cpp*).

[SI*] The probabilist's Hermite polynomials are solutions of the Sturm-Liouville differential equation

$$\left(e^{-\frac{1}{2}x^2}y'\right)' + \lambda e^{-\frac{1}{2}x^2}y = 0$$

where λ is a constant. Imposing the boundary condition that y should be polynomially bounded at infinity, the equation has solutions only if λ is a non-negative integer, and the solution is uniquely given by

$$y(x) = C_1 He_\lambda(x)$$

where C_1 is a constant.

Rewriting the differential equation as an eigenvalue problem

$$L[y] = y'' - xy' = -\lambda y$$

the Hermite polynomials $He_\lambda(x)$ may be understood as eigenfunctions of the differential operator $L[y]$. This eigenvalue problem is called the Hermite equation, although the term is also used for the closely related equation

$$y'' - xy' = -\lambda y$$

whose solution is uniquely given in terms of physicist's Hermite polynomials in the form

$$y(x) = C_1 H_\lambda(x)$$

where C_1 denotes a constant, after imposing the boundary condition that y should be polynomially bounded at infinity.

The general solutions to the above second-order differential equations are in fact linear combinations of both Hermite polynomials and confluent hypergeometric functions of the first kind. For example, for the physicist's Hermite equation

$$y'' - xy' + \lambda y = 0, \quad -\infty < x < \infty$$

the general solution takes the form

$$y(x) = C_1 H_\lambda(x) + C_2 h_\lambda(x)$$

where C_1 and C_2 are constants, $H_\lambda(x)$ are physicist's Hermite polynomials (of the first kind), and $h_\lambda(x)$ are physicist Hermite functions (of the second kind). The latter functions are compactly represented as

$$h_\lambda(x) = {}_1F_1\left(-\frac{\lambda}{2}; \frac{1}{2}; x^2\right)$$

where ${}_1F_1(a; b; z)$ are Confluent hypergeometric functions of the first kind. The conventional Hermite polynomials may also be expressed in terms of confluent hypergeometric functions. With more general boundary conditions, the Hermite polynomials can be generalized to obtain more general analytic functions for complex-valued λ . An explicit formula of Hermite polynomials in terms of contour integrals is also possible.

[SI*] **Recurrence Relation**

The sequence of probabilist's Hermite polynomials also satisfies the recurrence relation

$$He_{n+1}(x) = xHe_n(x) - He'_n(x)$$

Individual coefficients are related by the following recursion formula:

$$a_{n+1,k} = \begin{cases} -(k+1)a_{n,k+1}, & k = 0 \\ a_{n,k-1} - (k+1)a_{n,k+1}, & k > 0 \end{cases}$$

with $a_{0,0} = 1, a_{1,0} = 0, a_{1,1} = 1$.

For the physicist's Hermite polynomials, assuming

$$H_n(x) = \sum_{k=0}^n a_{n,k} x^k$$

we have

$$H_{n+1}(x) = 2xH_n(x) - H'_n(x)$$

Individual coefficients are related by the following recursion formula:

$$a_{n+1,k} = \begin{cases} -a_{n,k+1}, & k = 0 \\ 2a_{n,k-1}, & k > 0 \end{cases}$$

with $a_{0,0} = 1, a_{1,0} = 0, a_{1,1} = 2$.

The Hermite polynomials constitute an Appell sequence, i.e., they are a polynomial sequence satisfying the identity

$$\begin{aligned} He'_n(x) &= nHe_{n-1}(x) \\ H'_n(x) &= 2nH_{n-1}(x) \end{aligned}$$

An integral recurrence that is deduced is as follows:

$$\begin{aligned} He_{n+1}(x) &= (n+1) \int_0^x He_n(t) dt - He'_n(0) \\ H_{n+1}(x) &= 2(n+1) \int_0^x H_n(t) dt - H'_n(0) \end{aligned}$$

Equivalently, by Taylor-expanding,

$$\begin{aligned} He_n(x+y) &= \sum_{k=0}^n \binom{n}{k} x^{n-k} He_k(y) = 2^{-\frac{n}{2}} \sum_{k=0}^n \binom{n}{k} He_{n-k}(x\sqrt{2}) He_k(y\sqrt{2}) \\ H_n(x+y) &= \sum_{k=0}^n \binom{n}{k} H_k(x) (2y)^{n-k} = 2^{-\frac{n}{2}} \sum_{k=0}^n \binom{n}{k} H_{n-k}(x\sqrt{2}) H_k(y\sqrt{2}) \end{aligned}$$

These umbral identities are self-evident and included in the differential operator representation

$$\begin{aligned} He_n(x) &= e^{-\frac{D^2}{2}} x^n \\ H_n(x) &= 2^n e^{-\frac{D^2}{4}} x^n \end{aligned}$$

In consequence, for the m -th derivatives the following relations hold:

$$\begin{aligned} He_n^{(m)}(x) &= \frac{n!}{(n-m)!} He_{n-m}(x) = m! \binom{n}{m} He_{n-m}(x) \\ H_n^{(m)}(x) &= 2^m \frac{n!}{(n-m)!} H_{n-m}(x) = 2^m m! \binom{n}{m} H_{n-m}(x) \end{aligned}$$

It follows that the Hermite polynomials also satisfy the recurrence relation.

$$\begin{aligned} He_{n+1}(x) &= x He_n(x) - n He_{n-1}(x) \\ H_{n+1}(x) &= 2x H_n(x) - 2n H_{n-1}(x) \end{aligned}$$

These last relations, together with the initial polynomials $H_0(x)$ and $H_1(x)$, can be used in practice to compute the polynomials quickly.

Turán's inequalities are

$$H_n(x)^2 - H_{n-1}(x) H_{n+1}(x) = (n-1)! \sum_{i=0}^{n-1} \frac{2^{n-1}}{i!} H_i(x)^2 > 0$$

Moreover, the following multiplication theorem holds:

$$\begin{aligned} H_n(\gamma x) &= \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \gamma^{n-2i} (\gamma^2 - 1)^i \binom{n}{2i} \frac{(2i)!}{i!} H_{n-2i}(x) \\ He_n(\gamma x) &= \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \gamma^{n-2i} (\gamma^2 - 1)^i \binom{n}{2i} \frac{(2i)!}{i!} 2^{-i} He_{n-2i}(x) \end{aligned}$$

[SI*] Explicit Expression

The physicist's Hermite polynomials can be written explicitly as

$$H_n(x) = \begin{cases} n! \sum_{l=0}^{\frac{n}{2}} \frac{(-1)^{\frac{n}{2}-l}}{(2l)! (\frac{n}{2}-l)!} (2x)^{2l}, & \text{for even } n, \\ n! \sum_{l=0}^{\frac{n-1}{2}} \frac{(-1)^{\frac{n-1}{2}-l}}{(2l+1)! (\frac{n-1}{2}-l)!} (2x)^{2l+1}, & \text{for odd } n, \end{cases}$$

These two equations may be combined into one using the floor function:

$$H_n(x) = n! \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^m}{m! (n-2m)!} (2x)^{n-2m}$$

The probabilist's Hermite polynomials He have similar formulas, which may be obtained from these by replacing the power of $2x$ with the corresponding power of $\sqrt{2}x$ and multiplying the entire sum by $2^{-\frac{n}{2}}$:

$$He_n(x) = n! \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^m}{m!(n-2m)!} \frac{x^{n-2m}}{2^m}$$

Inverse Explicit Expression

The inverse of the above explicit expressions, that is, those for monomials in terms of probabilist's Hermite polynomials He are

$$x^n = n! \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} \frac{1}{2^m m!(n-2m)!} He_{n-2m}(x)$$

The corresponding expressions for the physicist's Hermite polynomials H follow directly by properly scaling this:

$$x^n = \frac{n!}{2^n} \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} \frac{1}{m!(n-2m)!} H_{n-2m}(x)$$

[SI*] Generating Function

The Hermite polynomials are given by the exponential generating function

$$\begin{aligned} e^{xt - \frac{1}{2}t^2} &= \sum_{n=0}^{\infty} He_n(x) \frac{t^n}{n!} \\ e^{2xt - t^2} &= \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!} \end{aligned}$$

This equality is valid for all complex values of x and t , and can be obtained by writing the Taylor expansion at x of the entire function $z \rightarrow e^{-z^2}$ (in the physicist's case). One can also derive the (physicist's) generating function by using the Cauchy's integral formula to write the Hermite polynomials as

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2} = (-1)^n e^{x^2} \frac{n!}{2\pi i} \oint_{\gamma} \frac{e^{-z^2}}{(z-x)^{n+1}} dz$$

Using this in the sum

$$\sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!}$$

one can evaluate the remaining integral using the calculus of residues and arrive at the desired generating function.

A slight generalization states

$$e^{2xt - t^2} H_k(x - t) = \sum_{n=0}^{\infty} \frac{H_{n+k}(x) t^n}{n!}$$

viii. Classical Orthogonal Polynomials

[SI*] Orthogonal polynomials are infinite sequences of polynomials where any two distinct polynomials in the set have an inner product of zero, defined with respect to a specific interval

and weight function. They behave like perpendicular vectors and serve as foundational tools in approximation theory, quantum physics, and numerical integration.

Core properties:

- **Inner Product:** Defined via an integral over an interval $[a, b]$ with a non-negative weight function $w(x)$, this is the orthogonality condition:

$$\int_a^b p_n(x)p_m(x)w(x) dx = 0, \quad \text{for } n \neq m$$

- **Degree Sequence:** Each $p_n(x)$ represents a distinct polynomial of exact degree n .
- **Real Roots:** The roots of an orthogonal polynomial of degree n are real, distinct, and located strictly inside the orthogonality interval.
- **Recurrence Relations:** Any system of orthogonal polynomials satisfies a simple three-term recurrence relation, allowing fast generation of higher degrees.

Any standard basis $\{1, x, x^2, \dots\}$ can be turned into orthogonal polynomials using the Gram-Schmidt process.

Classical families:

- **Legendre Polynomials:** Orthogonal on $[-1, 1]$ with a weight function of 1; used in potential theory and spherical harmonics.
- **Chebyshev Polynomials:** Orthogonal on $[-1, 1]$ with weight $(1 - x^2)^{-\frac{1}{2}}$; crucial for minimax polynomial approximation and numerical interpolation.
- **Hermite Polynomials:** Orthogonal on $(-\infty, \infty)$ with weight e^{-x^2} ; heavily applied in quantum harmonic oscillators.
- **Laguerre Polynomials:** Orthogonal on $[0, \infty)$ with weight e^{-x} ; used in quantum mechanics for hydrogen wavefunctions.

[SI*] Classical orthogonal polynomials-such as Legendre, Chebyshev, Hermite and Laguerre polynomials-are widely used in mathematical physics (in particular, the theory of random matrices), numerical analysis, and engineering to solve differential equations, approximate complex functions efficiently, and improve numerical integration accuracy / compute integrals.

Core mathematical tasks:

- **Differential Equations:** Serve as exact eigenfunction solutions to classical Sturm-Liouville differential systems.
- **Spectral Methods:** Act as global basis functions to convert differential equations into algebraic equations for fast computer simulations.
- **Numerical Integration (Quadrature):** Form the foundation of Gaussian quadrature rules for precise numerical integration of smooth functions. Gaussian quadrature rules use roots of these polynomials to compute definite integrals with high precision using few points.

[SI*] Major function families and their applications:

- **Legendre Polynomials:** Applied in potential theory and problems with spherical symmetry (like gravitational and electrostatic potentials).
- **Chebyshev Polynomials:** Used in numerical analysis for polynomial interpolation, minimizing interpolation errors (minimax approximation), and efficient numerical quadrature.

- **Hermite Polynomials:** Applied in quantum mechanics to solve the quantum harmonic oscillator wave equation, and in probability for Hermite transform and random processes (probability models with Gaussian distributions).
- **Laguerre Polynomials:** Used in quantum chemistry (solving Schrödinger equation for hydrogen-like atoms), as well as in chemical physics and control theory involving systems with exponential decay over semi-infinite.
- **Jacobi Polynomials:** Applied in boundary value problems and specialized spectral methods where endpoint behavior needs precise tuning.

[SI*] In mathematics, the classical orthogonal polynomials are the most widely used orthogonal polynomials: the Hermite polynomials, Laguerre polynomials, Jacobi polynomials (including as a special case the Gegenbauer polynomials, Chebyshev polynomials, and Legendre polynomials).

Classical orthogonal polynomials appeared in the early 19th century in the works of Adrien-Marie Legendre, who introduced the Legendre polynomials. In the late 19th century, the study of continued fractions to solve the moment problem by P.L. Chebyshev and then A.A. Markov and T.J. Stieltjes led to the general notion of orthogonal polynomials.

For given polynomials $Q, L : \mathbb{R} \rightarrow \mathbb{R}$ and $\forall n \in \mathbb{N}_0$ the classical orthogonal polynomials $f_n : \mathbb{R} \rightarrow \mathbb{R}$ are characterized by being solutions of the differential equation

$$Q(x)f_n'' + L(x)f_n' + \lambda_n f_n = 0$$

with to be determined constants $\lambda_n \in \mathbb{R}$.

[SI*] In general, the orthogonal polynomials P_n with respect to a weight $W : \mathbb{R} \rightarrow \mathbb{R}^+$ satisfy

$$\deg P_n = n, n = 0, 1, 2, \dots$$

$$\int P_m(x)P_n(x)W(x) dx = 0, \quad m \neq n$$

The relations above define P_n up to multiplication by a number. Various normalisations are used to fix the constant, e.g.

$$\int P_n(x)^2 W(x) dx = 1$$

The classical orthogonal polynomials correspond to the following three families of weights:

$$(\text{Jacobi}) \quad W(x) = \begin{cases} (1-x)^\alpha(1+x)^\beta, & -1 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

$$(\text{Physicist's Hermite}) \quad W(x) = e^{-x^2}$$

$$(\text{Laguerre}) \quad W(x) = \begin{cases} x^\alpha e^{-x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

[SI*] **Jacobi Polynomials**

Jacobi polynomials are a standard class of orthogonal polynomials denoted as $P_n^{(\alpha, \beta)}(x)$ that are defined on the interval $[-1, 1]$ with respect to the weight function $(1-x)^\alpha(1+x)^\beta$.

The Jacobi polynomials are defined by

$$P_n^{(\alpha, \beta)}(x) = \frac{\Gamma(\alpha + n + 1)}{n! \Gamma(\alpha + \beta + n + 1)} \sum_{m=0}^n \binom{n}{m} \frac{\Gamma(\alpha + \beta + n + m + 1)}{\Gamma(\alpha + m + 1)} \left(\frac{x-1}{2} \right)^m$$

The general formula to compute the first n Jacobi polynomials are given by

$$P_n^{(\alpha, \beta)}(x) = \sum_{k=0}^n \binom{n+\alpha}{n-k} \binom{n+k+\alpha+\beta}{k} \left(\frac{x-1}{2}\right)^k$$

[SI*] **Hermite Polynomials**

The Hermite polynomials are defined by

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2} = e^{\frac{x^2}{2}} \left(x - \frac{d}{dx}\right)^n e^{-\frac{x^2}{2}}$$

They satisfy the orthogonality condition

$$\int_{-\infty}^{\infty} H_n(x) H_m(x) e^{-x^2} dx = \sqrt{\pi} 2^n n! \delta_{mn}$$

and the differential equation

$$y'' - 2xy' + 2ny = 0$$

[SI*] **Laguerre Polynomials**

The generalised Laguerre polynomials are defined by

$$L_n^{(\alpha)}(x) = \frac{x^{-\alpha} e^x}{n!} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha})$$

(the classical Laguerre polynomials correspond to $\alpha = 0$).

They satisfy the orthogonality relation

$$\int_0^{\infty} x^{\alpha} e^{-x} L_n^{(\alpha)}(x) L_m^{(\alpha)}(x) dx = \frac{\Gamma(n+\alpha+1)}{n!} \delta_{n,m}$$

and the differential equation

$$xy'' + (\alpha + 1 - x)y' + ny = 0$$

[SI*] **Differential Equation**

The classical orthogonal polynomials arise from a differential equation of the form

$$Q(x)y'' + L(x)y' + \lambda y = 0 \quad (3.286)$$

where Q is a given quadratic (at most) polynomial, and L is a given linear polynomial. The function y , and the constant λ , are to be found.

This is a Sturm-Liouville type of equation. Such equations generally have singularities in their solution functions y except for particular values of λ . They can be thought of as eigenvector/eigenvalue problems: Letting D be the differential operator,

$$D(y) = Qy'' + Ly'$$

and changing the sign of λ , the problem is to find the eigenvectors (eigenfunctions) y , and the corresponding eigenvalues λ , such that y does not have singularities and $D(y) = \lambda y$.

The solutions of this differential equation have singularities unless λ takes on specific values. There is a series of numbers $\lambda_0, \lambda_1, \lambda_2, \dots$ that led to a series of polynomial solutions $P_0, P_1, P_2, \dots$ if one of the following sets of conditions are met:

1. Q is actually quadratic, L is linear, Q has two distinct real roots, the root of L lies strictly between the roots of Q and the leading terms of Q and L have the same sign.
2. Q is not actually quadratic, but is linear, L is linear, the roots of Q and L are different, and the leading terms of Q and L have the same sign if the root of L is less than the root of Q , or vice versa.
3. Q is just a nonzero constant, L is linear, and the leading term of L has the opposite sign of Q .

These three cases lead to the Jacobi-like, Laguerre-like, and Hermite-like polynomials, respectively.

In each of these three cases, we have the following:

- The solutions are a series of polynomials $P_0, P_1, P_2, \dots$, each P_n having degree n , and corresponding to a number λ_n .
- The interval of orthogonality is bounded by whatever roots Q has.
- The root of L is inside the interval of orthogonality.
- Letting $R(x) = e^{\int \frac{L(x)}{Q(x)} dx}$, the polynomials are orthogonal under the weight function $W(x) = \frac{R(x)}{Q(x)}$.
- $W(x)$ has no zeros or infinities inside the interval, though it may have zeros or infinities at the end points.
- $W(x)$ gives a finite inner product to any polynomials.
- $W(x)$ can be made to be greater than 0 in the interval (Negate the entire differential equation if necessary so that $Q(x) > 0$ inside the interval).

Because of the constant of integration, the quantity $R(x)$ is determined only up to an arbitrary positive multiplicative constant. It will be used only in homogeneous differential equations (where this doesn't matter) and in the definition of the weight function (which can also be indeterminate).

[SI*] **Rodrigues' Formula**

Under the assumptions of the preceding section, $P_n(x)$ is proportional to $\frac{1}{W(x)} \frac{d^n}{dx^n} (W(x)[Q(x)]^n)$. This is known as Rodrigues's formula, after Olinde Rodrigues. It is often written as

$$P_n(x) = \frac{1}{e_n W(x)} \frac{d^n}{dx^n} (W(x)[Q(x)]^n) \quad (3.287)$$

where the numbers e_n depend on the standardization.

[SI*] **The Numbers λ_n**

Under the assumptions of the preceding section, we have

$$\lambda_n = -n \left(\frac{n-1}{2} Q'' + L' \right) \quad (3.288)$$

Since Q is quadratic and L is linear, Q'' and L' are constants, so these are just numbers.

[SI*] **Second Form for the Differential Equation**

Let

$$R(x) = e^{\int \frac{L(x)}{Q(x)} dx}$$

Then

$$(Ry')' = Ry'' + R'y' = Ry'' + \frac{RL}{Q}y'$$

Now multiply the differential equation

$$Qy'' + Ly' + \lambda y = 0$$

by $\frac{R}{Q}$, getting

$$\begin{aligned} Ry'' + \frac{RL}{Q}y' + \frac{R\lambda}{Q}y &= 0 \\ (Ry')' + \frac{R\lambda}{Q}y &= 0 \end{aligned}$$

This is the standard Sturm-Liouville form for the equation.

[SI*] **Third Form for the Differential Equation**

Let

$$S(x) = \sqrt{R(x)} = e^{\int \frac{L(x)}{2Q(x)} dx}$$

Then

$$S' = \frac{SL}{2Q}$$

Now multiply the differential equation

$$Qy'' + Ly' + \lambda y = 0$$

by $\frac{S}{Q}$, thus

$$\begin{aligned} Sy'' + \frac{SL}{Q}y' + \frac{S\lambda}{Q}y &= 0 \\ Sy'' + 2S'y' + \frac{S\lambda}{Q}y &= 0 \end{aligned}$$

since $(Sy)'' = Sy'' + 2S'y' + S''y$, so

$$(Sy)'' + \left(\frac{S\lambda}{Q} - S'' \right) y = 0$$

or, letting $u = Sy$,

$$u'' + \left(\frac{\lambda}{Q} - \frac{S''}{S} \right) u = 0$$

[SI*] **Formulas Involving Derivatives**

Under the assumptions of the preceding section, let $P_n^{(r)}$ denote the r -th derivative of P_n . $P_n^{(r)}$ is a polynomial of degree $n - r$. Then we have the following:

- **Orthogonality:** For fixed r , the polynomial sequence $P_r^{(r)}, P_{r+1}^{(r)}, P_{r+2}^{(r)}, \dots$ are orthogonal, weighted by WQ^r .
- **Generalized Rodrigues' formula:** $P_n^{(r)}$ is proportional to

$$\frac{1}{W(x)[Q(x)]^r} \frac{d^{n-r}}{dx^{n-r}} (W(x)[Q(x)]^n)$$

- **Differential equation:** $P_n^{(r)}$ is a solution of $Qy'' + (rQ' + L)y' + [\lambda_n - \lambda_r]y = 0$, where λ_r is the same function as λ_n , that is,

$$\lambda_r = -r \left(\frac{r-1}{2} Q'' + L' \right)$$

- **Differential equation, second form:** $P_n^{(r)}$ is a solution of $(RQ^r y')' + [\lambda_n - \lambda_r]RQ^{r-1}y = 0$.

There are also some mixed recurrences. In each of these, the numbers a, b , and c depend on n and r , and are unrelated in the various formulas.

- $P_n^{(r)} = aP_{n+1}^{(r+1)} + bP_n^{(r+1)} + cP_{n-1}^{(r+1)}$
- $P_n^{(r)} = (ax + b)P_n^{(r+1)} + cP_{n-1}^{(r+1)}$
- $QP_n^{(r+1)} = (ax + b)P_n^{(r)} + cP_{n-1}^{(r)}$

There are an enormous number of other formulas involving orthogonal polynomials in various ways. Here is a tiny sample of them, relating to the Chebyshev, associated Laguerre, and Hermite polynomials:

- $2T_m(x)T_n(x) = T_{m+n}(x) + T_{m-n}(x)$
- $H_{2n}(x) = (-4)^n n! L_n^{-\frac{1}{2}}(x^2)$
- $H_{2n+1}(x) = 2(-4)^n n! x L_n^{\frac{1}{2}}(x^2)$

[SI*] Orthogonality

The differential equation for a particular λ may be written (omitting explicit dependence on x)

$$Q\ddot{y}_n + L\dot{y}_n + \lambda_n y_n = 0$$

multiplying by $\left(\frac{R}{Q}\right) y_m$ yields

$$Ry_m \ddot{y}_n + \frac{R}{Q} L y_m \dot{y}_n + \frac{R}{Q} \lambda_n y_m y_n = 0$$

and reversing the subscripts yields

$$Ry_n \ddot{y}_m + \frac{R}{Q} L y_n \dot{y}_m + \frac{R}{Q} \lambda_m y_n y_m = 0$$

subtracting and integrating:

$$\int_a^b \left[R(y_m \ddot{y}_n - y_n \ddot{y}_m) + \frac{R}{Q} L(y_m \dot{y}_n - y_n \dot{y}_m) \right] dx + (\lambda_n - \lambda_m) \int_a^b \frac{R}{Q} y_m y_n dx = 0$$

it can be seen that

$$\frac{d}{dx} [R(y_m \dot{y}_n - y_n \dot{y}_m)] = R(y_m \ddot{y}_n - y_n \ddot{y}_m) + R \frac{L}{Q} (y_m \dot{y}_n - y_n \dot{y}_m)$$

so that

$$[R(y_m \dot{y}_n - y_n \dot{y}_m)]_a^b + (\lambda_n - \lambda_m) \int_a^b \frac{R}{Q} y_m y_n dx = 0$$

If the polynomials y are such that the term on the left is zero, and $\lambda_m \neq \lambda_n$ for $m \neq n$, then the orthogonality relationship will hold:

$$\int_a^b \frac{R}{Q} y_m y_n dx = 0$$

for $m \neq n$.

[SI*] Derivation from Differential Equation

All of the polynomial sequences arising from the differential equation above are equivalent, under scaling and/or shifting of the domain, and standardizing of the polynomials, to more restricted classes. Those restricted classes are exactly "classical orthogonal polynomials."

- Every Jacobi-like polynomial sequence can have its domain shifted and/or scaled so that its interval of orthogonality is $[-1, 1]$, and has $Q = 1 - x^2$. They can then be standardized into the Jacobi polynomials $P_n^{(\alpha, \beta)}$. There are several important subclasses of these: Gegenbauer, Legendre, and two types of Chebyshev.
- Every Laguerre-like polynomial sequence can have its domain shifted, scaled, and/or reflected so that its interval of orthogonality is $[0, \infty)$, and has $Q = x$. They can then be standardized into the Associated Laguerre polynomials $L_n^{(\alpha)}$. The plain Laguerre polynomials L_n are a subclass of these.
- Every Hermite-like polynomial sequence can have its domain shifted and/or scaled so that its interval of orthogonality is $(-\infty, \infty)$, and has $Q = 1$ and $L(0) = 0$. They can then be standardized into the Hermite polynomials H_n .

Because all polynomial sequences arising from a different equation in the manner described above are trivially equivalent to the classical polynomials, the actual classical polynomials are always used.

1. Jacobi Polynomials

Jacobi polynomials are a standard class of orthogonal polynomials denoted as $P_n^{(\alpha, \beta)}(x)$ that are defined on the interval $[-1, 1]$ with respect to the weight function $(1 - x)^\alpha(1 + x)^\beta$. They generalize other important polynomials and appear widely in differential equations and physical sciences (the Gegenbauer polynomials, and thus also the Legendre, Zernike and Chebyshev polynomials are special cases of the Jacobi polynomials). The Jacobi polynomials were introduced by Carl Gustav Jacob Jacobi.

The variety of characterizations of the Jacobi polynomials:

- The general formula to compute the first n Jacobi polynomials are given by

$$P_n^{(\alpha, \beta)}(x) = \sum_{k=0}^n \binom{n + \alpha}{n - k} \binom{n + k + \alpha + \beta}{k} \left(\frac{x - 1}{2}\right)^k$$

- For $\alpha, \beta > -1$ (this puts the root of L inside the interval of orthogonality) the Jacobi polynomials are given by the Rodrigues' formula

$$P_n^{(\alpha, \beta)}(x) = \frac{(-1)^n}{2^n n!} (1 - x)^{-\alpha} (1 + x)^{-\beta} \frac{d^n}{dx^n} \{ (1 - x)^\alpha (1 + x)^\beta (1 - x^2)^n \}$$

If $\alpha = \beta = 0$, then it reduces to the Legendre polynomials:

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$$

- They are normalised (standardized) by

$$P_n^{(\alpha, \beta)}(1) = \binom{n + \alpha}{n}$$

and satisfy the orthogonality condition

$$\int_{-1}^1 (1-x)^\alpha (1+x)^\beta P_m^{(\alpha,\beta)}(x) P_n^{(\alpha,\beta)}(x) dx = \frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)n!} \delta_{nm}$$

with $\alpha, \beta > -1$. As defined, they do not have unit norm with respect to the weight. This can be corrected by dividing by the square root of the right hand side of the equation above, when $n = m$.

Although it does not yield an orthonormal basis, an alternative normalization is sometimes preferred due to its simplicity:

$$P_n^{(\alpha,\beta)}(1) = \binom{n+\alpha}{n}$$

- The polynomials have the symmetry relation

$$P_n^{(\alpha,\beta)}(-x) = (-1)^n P_n^{(\beta,\alpha)}(x)$$

thus the other terminal value is

$$P_n^{(\alpha,\beta)}(-1) = (-1)^n \binom{n+\beta}{n}$$

- The k th derivative of the explicit expression leads to

$$\frac{d^k}{dx^k} P_n^{(\alpha,\beta)}(x) = \frac{\Gamma(\alpha+\beta+n+1+k)}{2^k \Gamma(\alpha+\beta+n+1)} P_{n-k}^{(\alpha+k,\beta+k)}(x)$$

- The generating function of the Jacobi polynomials is given by

$$\sum_{n=0}^{\infty} P_n^{(\alpha,\beta)}(x) t^n = 2^{(\alpha+\beta)} R^{-1} (1-t+R)^{-\alpha} (1+t+R)^{-\beta}$$

where

$$R = R(x, t) = (1 - 2xt + t^2)^{\frac{1}{2}}$$

and the branch of square root is chosen so that $R(x, 0) = 1$.

- The Jacobi polynomials reduce to other classical polynomials.
Ultraspherical

$$C_n^{(\lambda)}(x) = \frac{(2\lambda)_n}{\left(\lambda + \frac{1}{2}\right)_n} P_n^{(\lambda-\frac{1}{2}, \lambda-\frac{1}{2})}(x)$$

$$P_n^{(\alpha,\alpha)}(x) = \frac{(\alpha+1)_n}{(2\alpha+1)_n} C_n^{(\alpha+\frac{1}{2})}(x)$$

Legendre

$$P_n(x) = C_n^{(\frac{1}{2})}(x) = P_n^{(0,0)}(x)$$

Chebyshev

$$T_n(x) = \frac{P_n^{(-\frac{1}{2}, -\frac{1}{2})}(x)}{P_n^{(-\frac{1}{2}, -\frac{1}{2})}(1)}$$

$$U_n(x) = C_n^{(1)}(x) = (n+1) \frac{P_n^{(\frac{1}{2}, \frac{1}{2})}(x)}{P_n^{(\frac{1}{2}, \frac{1}{2})}(1)}$$

$$V_n(x) = \frac{P_n^{(-\frac{1}{2}, \frac{1}{2})}(x)}{P_n^{(-\frac{1}{2}, \frac{1}{2})}(1)}$$

$$W_n(x) = (2n+1) \frac{P_n^{(\frac{1}{2}, -\frac{1}{2})}(x)}{P_n^{(\frac{1}{2}, -\frac{1}{2})}(1)}$$

$$T_n^*(x) = T_n(2x-1)$$

$$U_n^*(x) = U_n(2x-1)$$

Laguerre

$$\lim_{\beta \rightarrow \infty} P_n^{(\alpha, \beta)} \left(1 - \left(\frac{2x}{\beta} \right) \right) = L_n^{(\alpha)}(x)$$

$$\lim_{\alpha \rightarrow \infty} P_n^{(\alpha, \beta)} \left(\left(\frac{2x}{\alpha} \right) - 1 \right) = (-1)^n L_n^{(\beta)}(x)$$

Hermit

$$\lim_{\alpha \rightarrow \infty} \alpha^{-\frac{1}{2}n} P_n^{(\alpha, \alpha)} \left(\alpha^{-\frac{1}{2}} x \right) = \frac{H_n(x)}{2^n n!} L_n^{(\beta)}(x)$$

- The Jacobi polynomials appear as the eigenfunctions of the Markov process on $[-1, +1]$

$$\mathbb{L} = (1-x^2) \frac{\partial^2}{\partial x^2} + (px-q) \frac{\partial}{\partial x}$$

defined up to the time it hits the boundary. For $p = -(\beta + \alpha + 2)$, $q = \beta - \alpha$, we have

$$\mathbb{L} P_n^{(\alpha, \beta)} = -n(n + \alpha + \beta + 1) P_n^{(\alpha, \beta)}$$

Thus this process is named the Jacobi process.

- The Jacobi polynomials are solutions to the singular Sturm-Liouville eigenvalue problem

$$(1-x^2)y'' + (\beta - \alpha - (\alpha + \beta + 2)x)y' + n(n + \alpha + \beta + 1)y = 0$$

- The Jacobi polynomials can also be expressed as a finite terminating Gauss hypergeometric series ${}_2F_1 \left(-n, n + \alpha + \beta + 1; \alpha + 1; \frac{1-x}{2} \right)$

$$P_n^{(\alpha, \beta)}(x) = \frac{(\alpha + 1)_n}{n!} {}_2F_1 \left(-n, n + \alpha + \beta + 1; \alpha + 1; \frac{1-x}{2} \right)$$

where $(\alpha + 1)_n$ is Pochhammer's symbol (for the rising factorial). In this case, the series for the hypergeometric function is finite, therefore one obtains the following equivalent expression:

$$P_n^{(\alpha, \beta)}(x) = \frac{\Gamma(\alpha + n + 1)}{n! \Gamma(\alpha + \beta + n + 1)} \sum_{m=0}^n \binom{n}{m} \frac{\Gamma(\alpha + \beta + n + m + 1)}{\Gamma(\alpha + m + 1)} \left(\frac{x-1}{2} \right)^m$$

• Special cases

$$P_0^{(\alpha, \beta)}(x) = 1$$

$$P_1^{(\alpha, \beta)}(x) = (\alpha + 1)(\alpha + \beta + 2) \frac{x-1}{2}$$

$$P_2^{(\alpha, \beta)}(x) = \frac{(\alpha + 1)(\alpha + 2)}{2} + (\alpha + 2)(\alpha + \beta + 3) \frac{x-1}{2} + \frac{(\alpha + \beta + 3)(\alpha + \beta + 4)}{2} \left(\frac{x-1}{2} \right)^2$$

⋮

$$P_n^{(\alpha, \beta)}(x) = \frac{\Gamma(1 + 2n + \alpha + \beta)}{\Gamma(1 + n) \Gamma(1 + n + \alpha + \beta)} \left(\frac{x}{2} \right)^n + \text{lower-degree terms}$$

Thus, the leading coefficient is

$$\frac{\Gamma(1 + 2n + \alpha + \beta)}{2^n n! \Gamma(1 + n + \alpha + \beta)}$$

Special Cases

The Jacobi polynomials with $\alpha = \beta$ are called the Gegenbauer polynomials (with parameter $\gamma = \alpha + \frac{1}{2}$).

For $\alpha = \beta = 0$, these are called the Legendre polynomials (for which the interval of orthogonality is $[-1, 1]$ and the weight function is simply 1):

$$P_0(x) = 1$$

$$P_1(x) = x$$

$$P_2(x) = \frac{3x^2 - 1}{2}$$

$$P_3(x) = \frac{5x^3 - 3x}{2}$$

For $\alpha = \beta = -\frac{1}{2}$, one obtains the Chebyshev polynomials of the first, and for $\alpha = \beta = \frac{1}{2}$, one obtains the Chebyshev polynomials of the second kind.

The Jacobi-like polynomials, once they have had their domain shifted and scaled so that the interval of orthogonality is $[-1, 1]$, still have two parameters to be determined. They are α and β in the Jacobi polynomials, written $P_n^{(\alpha, \beta)}$. We have $Q(x) = 1 - x^2$ and $L(x) = \beta - \alpha - (\alpha + \beta + 2)x$. Both α and β are required to be greater than -1 (This puts the root of L inside the interval of orthogonality).

When α and β are not equal, these polynomials are not symmetrical about $x = 0$.

The Jacobi polynomial $P_n^{(\alpha, \beta)}$ is, up to scaling, the unique polynomial solution of the Sturm-Liouville problem:

$$(1 - x^2)y'' + (\beta - \alpha - [\alpha + \beta + 2]x)y' + \lambda y = 0, \quad \text{with } \lambda = n(n + 1 + \alpha + \beta)$$

The other solution involves the logarithm function. Bochner's theorem states that the Jacobi polynomials are uniquely characterized as polynomial solutions to Sturm-Liouville problems with polynomial coefficients.

2. Gegenbauer Polynomials

In mathematics, Gegenbauer polynomials $C_n^{(\alpha)}$ are orthogonal polynomials on the interval $[-1, 1]$ with respect to the weight function $(1 - x^2)^{\alpha - \frac{1}{2}}$. They are named after Leopold Gegenbauer. When one sets parameters α and β in the Jacobi polynomials equal to each other, one obtains the Gegenbauer or ultraspherical polynomials. They are written $C_n^{(\alpha)}$, and defined as

$$C_n^{(\alpha)}(x) = \frac{\Gamma(2\alpha + n)\Gamma\left(\alpha + \frac{1}{2}\right)}{\Gamma(2\alpha)\Gamma\left(\alpha + n + \frac{1}{2}\right)} P_n^{(\alpha - \frac{1}{2}, \alpha - \frac{1}{2})}(x)$$

We have $Q(x) = 1 - x^2$ and $L(x) = -(2\alpha + 1)x$. The parameter α is required to be greater than $-\frac{1}{2}$. The accepted standardization sets $C_n^{(0)}(1) = \frac{2}{n}$.

The parameter α is closely related to the derivatives of $C_n^{(\alpha)}$:

$$C_n^{(\alpha+1)}(x) = \frac{1}{2\alpha} \frac{d}{dx} C_{n+1}^{(\alpha)}(x)$$

or, more generally:

$$C_n^{(\alpha+m)}(x) = \frac{\Gamma(\alpha)}{2^m \Gamma(\alpha + m)} C_{n+m}^{(\alpha)[m]}(x)$$

All the other classical Jacobi-like polynomials (Legendre, etc) are special cases of the Gegenbauer polynomials, obtained by choosing a value of α and choosing a standardization.

The variety of characterizations of the Gegenbauer polynomials:

- The polynomials can be defined in terms of their generating function:

$$\frac{1}{(1 - 2xt + t^2)^\alpha} = \sum_{n=0}^{\infty} C_n^{(\alpha)}(x) t^n, \quad 0 \leq |x| < 1, |t| \leq 1, \alpha > 0$$

- The recurrence relation for Gegenbauer polynomials is

$$\begin{aligned} C_0^{(\alpha)}(x) &= 1 \\ C_1^{(\alpha)}(x) &= 2\alpha x \\ C_{n+1}^{(\alpha)}(x) &= \frac{1}{n+1} \left[2x(n+\alpha)C_n^{(\alpha)}(x) - (n+2\alpha-1)C_{n-1}^{(\alpha)}(x) \right] \end{aligned}$$

with $n = 1, 2, 3, \dots$, they are orthogonal polynomials depending on a parameter α .

- Gegenbauer polynomials are particular solutions of the Gegenbauer differential equation:

$$(1 - x^2)y'' - (2\alpha + 1)xy' + n(n + 2\alpha)y = 0$$

when $\alpha = \frac{1}{2}$, the equation reduces to the Legendre equation, and the Gegenbauer polynomials reduce to the Legendre polynomials. When $\alpha = 1$, the equation reduces to the Chebyshev differential equation, and the Gegenbauer polynomials reduce to the Chebyshev polynomials of the second kind.

- They are given as Gaussian hypergeometric series in certain cases where the series is in fact finite:

$$C_n^{(\alpha)}(z) = \frac{(2\alpha)_n}{n!} {}_2F_1 \left(-n, 2\alpha + n; \alpha + \frac{1}{2}; \frac{1-z}{2} \right)$$

Here $(2\alpha)_n$ is the rising factorial. Explicitly,

$$C_n^{(\alpha)}(z) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \frac{\Gamma(n - k + \alpha)}{\Gamma(\alpha)k!(n - 2k)!} (2z)^{n-2k}$$

From this it is also easy to obtain the value at unit argument:

$$C_n^{(\alpha)}(1) = \frac{\Gamma(2\alpha + n)}{\Gamma(2\alpha)n!}$$

- They are special cases of the Jacobi polynomials:

$$C_n^{(\alpha)}(x) = \frac{(2\alpha)_n}{\left(\alpha + \frac{1}{2}\right)_n} P_n^{\left(\alpha - \frac{1}{2}, \alpha - \frac{1}{2}\right)}(x)$$

in which $(\theta)_n$ represents the rising factorial of θ .

One therefore also has the Rodrigues formula

$$C_n^{(\alpha)}(x) = \frac{(-1)^n}{2^n n!} \frac{\Gamma\left(\alpha + \frac{1}{2}\right) \Gamma(n + 2\alpha)}{\Gamma(2\alpha) \Gamma\left(\alpha + n + \frac{1}{2}\right)} (1 - x^2)^{-\alpha + \frac{1}{2}} \frac{d^n}{dx^n} \left[(1 - x^2)^{n + \alpha - \frac{1}{2}} \right]$$

- An alternative normalization sets $C_n^{(\alpha)}(1) = 1$. Assuming this alternative normalization, the derivatives of Gegenbauer are expressed in terms of Gegenbauer:

$$\begin{aligned} \frac{d^q}{dx^q} C_{q+2j+1}^{(\alpha)}(x) &= \frac{2^q (q + 2j + 1)}{(q - 1)! \Gamma(q + 2j + 2\alpha + 1)} \sum_{i=0}^j \frac{(2i + \alpha + 1) \Gamma(2i + 2\alpha + 1)}{(2i + 1)! (j - i)!} \\ &\times \frac{\Gamma(q + j + i + \alpha + 1)}{\Gamma(j + i + \alpha + 2)} (q + j - i - 1)! C_{2i+1}^{(\alpha)}(x) \end{aligned}$$

Orthogonality and Normalization

For a fixed $\alpha > -\frac{1}{2}$, the polynomials are orthogonal on $[-1, 1]$ with respect to the weighting function

$$w(z) = (1 - z^2)^{\alpha - \frac{1}{2}}$$

To wit, for $n \neq m$,

$$\int_{-1}^1 C_n^{(\alpha)}(x) C_m^{(\alpha)}(x) (1-x^2)^{\alpha-\frac{1}{2}} dx = 0$$

They are normalized by

$$\int_{-1}^1 [C_n^{(\alpha)}(x)]^2 (1-x^2)^{\alpha-\frac{1}{2}} dx = \frac{\pi 2^{1-2\alpha} \Gamma(n+2\alpha)}{n!(n+\alpha)[\Gamma(\alpha)]^2}$$

Applications

The Gegenbauer polynomials appear naturally as extensions of Legendre polynomials in the context of potential theory and harmonic analysis. The Newtonian potential in $\mathbb{R}^n$ has the expansion, valid with $\alpha = \frac{n-2}{2}$,

$$\frac{1}{|x-y|^{n-2}} = \sum_{k=0}^{\infty} \frac{|x|}{|y|^{k+n-2}} C_k^{(\alpha)}\left(\frac{x \cdot y}{|x||y|}\right)$$

When $n = 3$, this gives the Legendre polynomial expansion of the gravitational potential. Similar expressions are available for the expansion of the Poisson kernel in a ball.

It follows that the quantities $C_k^{(\frac{n-2}{2})}(x \cdot y)$ are spherical harmonics, when regarded as a function of x only. They are, in fact, exactly the zonal spherical harmonics, up to a normalizing constant.

Gegenbauer polynomials also appear in the theory of positive-definite functions.

The Askey-Gasper inequality reads

$$\sum_{j=0}^n \frac{C_j^{(\alpha)}(x)}{\binom{2\alpha+j-1}{j}} \geq 0, \quad x \geq -1, \alpha \geq \frac{1}{4}$$

In spectral methods for solving differential equations, if a function is expanded in the basis of Chebyshev polynomials and its derivative is represented in a Gegenbauer/ultraspherical basis, then the derivative operator becomes a diagonal matrix, leading to fast banded matrix methods for large problems.

3. Legendre Polynomials

The differential equation

$$(1-x^2)y'' - 2xy' + \lambda y = 0, \quad \text{with } \lambda = n(n+1)$$

is Legendre's equation.

The second form of the differential equation is:

$$\frac{d}{dx}[(1-x^2)y'] + \lambda y = 0$$

The recurrence relation is

$$(n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x)$$

A mixed recurrence is

$$P_{n+1}^{[r+1]}(x) = P_{n+1}^{[r+1]}(x) + (2n+1)P_n^{[r]}(x)$$

Rodrigues' formula is

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} ([x^2 - 1]^n)$$

Associated Legendre polynomials

The associated Legendre polynomials, denoted by $P_l^{(m)}(x)$ where l and m are integers with $0 \leq m \leq l$, are defined as

$$P_l^{(m)}(x) = (-1)^m (1-x^2)^{\frac{m}{2}} P_l^{[m]}(x)$$

The m in parentheses (to avoid confusion with an exponent) is a parameter. The m in brackets denotes the m -th derivative of the Legendre polynomial.

These "polynomials" are misnamed—they are not polynomials when m is odd.

They have a recurrence relation:

$$(l+1-m)P_{l+1}^{(m)}(x) = (2l+1)xP_l^{(m)}(x) - (l+m)P_{l-1}^{(m)}(x)$$

For fixed m , the sequence $P_m^{(m)}, P_{m+1}^{(m)}, P_{m+2}^{(m)}, \dots$ are orthogonal over $[-1, 1]$, with weight 1.

For given m , $P_l^{(m)}(x)$ are the solutions of

$$(1-x^2)y'' - 2xy' + \left[\lambda - \frac{m^2}{1-x^2} \right] y = 0, \quad \text{with } \lambda = l(l+1)$$

4. Chebyshev Polynomials

The Chebyshev polynomials are two sequences of orthogonal polynomials related to cosine and sine functions, notated as $T_n(x)$ and $U_n(x)$. They can be defined in several equivalent ways, one of which starts with trigonometric functions:

- The Chebyshev polynomials of the first kind T_n are defined by

$$T_n(\cos \theta) = \cos(n\theta), \quad n = 0, 1, 2, 3, \dots$$

The recurrence relation is

$$\begin{aligned} T_0(x) &= 1 \\ T_1(x) &= x \\ T_{n+1}(x) &= 2xT_n(x) - T_{n-1}(x) \end{aligned}$$

- The Chebyshev polynomials of the second kind U_n are defined by

$$U_n(\cos \theta) = \frac{\sin((n+1)\theta)}{\sin \theta}, \quad n = 0, 1, 2, 3, \dots$$

or

$$U_n = \frac{1}{n+1} T'_{n+1}$$

The recurrence relation is

$$\begin{aligned} U_0(x) &= 1 \\ U_1(x) &= 2x \\ U_{n+1}(x) &= 2xU_n(x) - U_{n-1}(x) \end{aligned}$$

That these expressions define polynomials in $\cos \theta$ is not obvious at first sight but can be shown using de Moivre's formula.

The Chebyshev polynomials T_n are polynomials with the largest possible leading coefficient whose absolute value on the interval $[-1, 1]$ is bounded by 1, they are also the "extremal" polynomials for many other properties.

In 1952, Cornelius Lanczos showed that the Chebyshev polynomials are important in approximation theory for the solution of linear systems; the roots of $T_n(x)$, which are also called Chebyshev nodes, are used as matching points for optimizing polynomial interpolation. The resulting interpolation polynomial minimizes the problem of Runge's phenomenon and provides an approximation that is close to the best polynomial approximation to a continuous function under the maximum norm, also called the "minimax" criterion. This approximation leads directly to the method of Clenshaw-Curtis quadrature.

These polynomials were named after Pafnuty Chebyshev. The letter T is used because of the alternative transliterations of the name *Chebyshev* as *Tchebycheff*, *Tchebyshev* (French) or *Tschebyschow* (German).

Trigonometric and Complex Definition

An equivalent way to state the Chebyshev trigonometric definition is via exponentiation of a complex number: given a complex number $z = a + bi$ with absolute value of one,

$$z^n = T_n(a) + ibU_{n-1}(a)$$

That $\cos(nx)$ is an n -th degree polynomial in $\cos(x)$ and can be seen by observing that $\cos(nx)$ is the real part of one side of de Moivre's formula:

$$\cos n\theta + i \sin n\theta = (\cos \theta + i \sin \theta)^n$$

The real part of the other side is a polynomial in $\cos x$ and $\sin x$, in which all powers of $\sin x$ are even and thus replaceable through the identity $\cos^2 x + \sin^2 x = 1$. By the same reasoning, $\sin nx$ is the imaginary part of the polynomial, in which all powers of $\sin x$ are odd and thus, if one factor of $\sin x$ is factored out, the remaining factors can be replaced to create a polynomial of degree $n - 1$ in $\cos x$.

For x outside the interval $[-1, 1]$, the above definition implies

$$T_n(x) = \begin{cases} \cos(n \arccos x), & \text{if } |x| \leq 1 \\ \cosh(n \operatorname{arccosh} x), & \text{if } x \geq 1 \\ (-1)^n \cosh(n \operatorname{arccosh} (-x)), & \text{if } x \leq -1 \end{cases}$$

Generating Function

The ordinary generating function for T_n is

$$\sum_{n=0}^{\infty} T_n(x) t^n = \frac{1 - tx}{1 - 2tx + t^2}$$

There are several other generating functions for the Chebyshev polynomials; the exponential generating function is

$$\begin{aligned} \sum_{n=0}^{\infty} T_n(x) \frac{t^n}{n!} &= \frac{1}{2} \left(e^{t(x - \sqrt{x^2 - 1})} + e^{t(x + \sqrt{x^2 - 1})} \right) \\ &= e^{tx} \cosh(t\sqrt{x^2 - 1}) \end{aligned}$$

The generating function relevant for 2-dimensional potential theory and multipole expansion is

$$\sum_{n=0}^{\infty} T_n(x) \frac{t^n}{n!} = \ln \left(\frac{1}{\sqrt{1 - 2tx + t^2}} \right)$$

The ordinary generating function for U_n is

$$\sum_{n=0}^{\infty} U_n(x) t^n = \frac{1}{1 - 2tx + t^2}$$

and the exponential generating function is

$$\sum_{n=0}^{\infty} U_n(x) \frac{t^n}{n!} = e^{tx} \left(\cosh(t\sqrt{x^2 - 1}) + \frac{x}{\sqrt{x^2 - 1}} \sinh(t\sqrt{x^2 - 1}) \right)$$

The differential equation

$$(1 - x^2)y'' - xy' + \lambda y = 0, \quad \text{with } \lambda = n^2$$

is Chebyshev's equation.

The first six Chebyshev polynomials of the first kind $T_n(x)$ are

$$\begin{aligned} T_0(x) &= 1 \\ T_1(x) &= x \\ T_2(x) &= 2x^2 - 1 \\ T_3(x) &= 4x^3 - 3x \\ T_4(x) &= 8x^4 - 8x^2 + 1 \\ T_5(x) &= 16x^5 - 20x^3 + 5x \end{aligned}$$

The first six Chebyshev polynomials of the second kind $U_n(x)$ are

$$\begin{aligned} U_0(x) &= 1 \\ U_1(x) &= 2x \\ U_2(x) &= 4x^2 - 1 \\ U_3(x) &= 8x^3 - 4x \\ U_4(x) &= 16x^4 - 12x^2 + 1 \\ U_5(x) &= 32x^5 - 32x^3 + 6x \end{aligned}$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Chebyshev Polynomials ]# ./main
Chebyshev polynomials of the first kind:
T_{0}(x) = 1
T_{1}(x) = x
T_{2}(x) = 2*x^(2)-1
T_{3}(x) = 4*x^(3)-3*x
T_{4}(x) = 8*x^(4)-8*x^(2)+1
T_{5}(x) = 16*x^(5)-20*x^(3)+5*x
T_{6}(x) = 32*x^(6)-48*x^(4)+18*x^(2)-1
T_{7}(x) = 64*x^(7)-112*x^(5)+56*x^(3)-7*x
T_{8}(x) = 128*x^(8)-256*x^(6)+160*x^(4)-32*x^(2)+1
T_{9}(x) = 256*x^(9)-576*x^(7)+432*x^(5)-120*x^(3)+9*x
T_{10}(x) = 512*x^(10)-1280*x^(8)+1120*x^(6)-400*x^(4)+50*x^(2)-1

Chebyshev polynomials of the second kind:
U_{0}(x) = 1
U_{1}(x) = 2*x
U_{2}(x) = 4*x^(2)-1
U_{3}(x) = 8*x^(3)-4*x
U_{4}(x) = 16*x^(4)-12*x^(2)+1
U_{5}(x) = 32*x^(5)-32*x^(3)+6*x
U_{6}(x) = 64*x^(6)-80*x^(4)+24*x^(2)-1
U_{7}(x) = 128*x^(7)-192*x^(5)+80*x^(3)-8*x
U_{8}(x) = 256*x^(8)-448*x^(6)+240*x^(4)-40*x^(2)+1
U_{9}(x) = 512*x^(9)-1024*x^(7)+672*x^(5)-160*x^(3)+10*x
U_{10}(x) = 1024*x^(10)-2304*x^(8)+1792*x^(6)-560*x^(4)+60*x^(2)-1

Time taken by function: 174549 microseconds

```

Figure 3.77: The first eleven Chebyshev polynomials of the first kind and the second kind (*SymIntegration/Examples/Differential Equations/Test SymIntegration Jacobi Polynomials/main.cpp*).

Rodrigues' formula is

$$T_n(x) = \frac{\Gamma\left(\frac{1}{2}\right) \sqrt{1-x^2}}{(-2)^n \Gamma\left(n + \frac{1}{2}\right)} \frac{d^n}{dx^n} \left([1-x^2]^{n-\frac{1}{2}} \right)$$

These polynomials have the property that, in the interval of orthogonality,

$$T_n(x) = \cos(n \arccos(x))$$

This means that all their local minima and maxima have values of -1 and $+1$, that is, the polynomials are "level." Because of this, expansion of functions in terms of Chebyshev polynomials is sometimes used for polynomial approximations in computer math libraries.

Some authors use versions of these polynomials that have been shifted so that the interval of orthogonality is $[0, 1]$ or $[-2, 2]$.

5. Laguerre Polynomials

The most general Laguerre-like polynomials, after the domain has been shifted and scaled, are the Associated Laguerre polynomials (also called generalized Laguerre polynomials), denoted by $L_n^{(\alpha)}$. There is a parameter α , which can be any real number strictly greater than -1 . The parameter is put in parentheses to avoid confusion with an exponent. The plain Laguerre polynomials are simply the $\alpha = 0$ version of these:

$$L_n(x) = L_n^{(0)}(x)$$

In mathematics, the Laguerre polynomials, named after Edmond Laguerre (1834-1886), are nontrivial solutions of Laguerre's differential equation:

$$xy'' + (1-x)y' + ny = 0, \quad y = L(x)$$

which is a second-order linear differential equation. This equation has nonsingular solutions only if n is a non-negative integer.

The Laguerre polynomials denoted by $L_n(x)$, and defined as solutions to Laguerre's differential equation can be explicitly computed using the Rodrigues's formula:

$$L_n(x) = \frac{e^x}{n!} \frac{d^n}{dx^n} (x^n e^{-x})$$

Sometimes the name Laguerre polynomials is used for solutions of

$$xy'' + (\alpha + 1 - x)y' + \lambda y = 0, \quad \text{with } \lambda = n$$

where $\lambda = n$ is still a non-negative integer. Then they are also named generalized Laguerre polynomials, or rarely Sonine polynomials.

The second form of the differential equation is

$$(x^{\alpha+1} e^{-x} y')' + \lambda x^{\alpha} e^{-x} y = 0$$

More generally, a Laguerre function is a solution when n is not necessarily a non-negative integer.

The variety of characterizations of the Laguerre polynomials:

- The Laguerre polynomials are also used for Gauss-Laguerre quadrature to numerically compute integrals of the form

$$\int_0^{\infty} f(x) e^{-x} dx$$

- These polynomials, usually denoted by $L_0, L_1, \dots$, are a polynomial sequence which can be defined by the Rodrigues' formula,

$$L_n(x) = \frac{e^x}{n!} \frac{d^n}{dx^n} (e^{-x} x^n) = \frac{1}{n!} \left(\frac{d}{dx} - 1 \right)^n x^n$$

or with $\alpha \neq 0$

$$L_n^{(\alpha)}(x) = \frac{x^{-\alpha} e^x}{n!} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}) = \frac{x^{-\alpha}}{n!} \left(\frac{d}{dx} - 1 \right)^n x^{n+\alpha}$$

The parameter α is closely related to the derivatives of $L_n^{(\alpha)}$:

$$L_n^{(\alpha+1)}(x) = -\frac{d}{dx} L_{n+1}^{(\alpha)}(x)$$

or, more generally:

$$L_n^{(\alpha+m)}(x) = (-1)^m L_{n+m}^{(\alpha)}(x)$$

The generating function for them with $\alpha \neq 0$ is

$$\sum_{n=0}^{\infty} t^n L_n^{(\alpha)}(x) = \frac{1}{(1-t)^{\alpha+1}} e^{-\frac{tx}{1-t}}$$

The generating function for them with $\alpha = 0$ is

$$\sum_{n=0}^{\infty} t^n L_n(x) = \frac{1}{1-t} e^{-\frac{tx}{1-t}}$$

- They are orthogonal polynomials on the interval $[0, \infty)$ with respect to an inner product and the weighting function e^{-x} :

$$\langle f, g \rangle = \int_0^{\infty} f(x)g(x)e^{-x} dx$$

or

$$\int_0^{\infty} e^{-x} L_m(x) L_n(x) dx = \delta_{mn}$$

The rook polynomials in combinatorics are more or less the same as Laguerre polynomials, up to elementary changes of variables.

- The recurrence relation for the standard Laguerre polynomials without α ($\alpha = 0$):

$$L_0(x) = 1$$

$$L_1(x) = 1 - x$$

$$L_2(x) = \frac{1}{2}(x^2 - 4x + 2)$$

$$L_3(x) = \frac{1}{6}(-x^3 + 9x^2 - 18x + 6)$$

$$L_4(x) = \frac{1}{24}(x^4 - 16x^3 + 72x^2 - 96x + 24)$$

$$L_5(x) = \frac{1}{120}(-x^5 + 25x^4 - 200x^3 + 600x^2 - 600x + 120)$$

$\vdots$

$$L_{k+1}(x) = \frac{(2k+1-x)L_k(x) - kL_{k-1}(x)}{k+1}, \quad k \geq 1$$

The generalized Laguerre polynomials with α ($\alpha \neq 0$) are given by:

$$L_0^{(\alpha)}(x) = 1$$

$$L_1^{(\alpha)}(x) = 1 - x + \alpha$$

$$L_2^{(\alpha)}(x) = \frac{1}{2}x^2 - (\alpha+2)x + \frac{1}{2}(\alpha+1)(\alpha+2)$$

$$L_3^{(\alpha)}(x) = -\frac{1}{6}x^3 + \frac{1}{2}(\alpha+3)x^2 - \frac{1}{2}(\alpha+2)(\alpha+3)x + \frac{1}{6}(\alpha+1)(\alpha+2)(\alpha+3)$$

$\vdots$

$$L_{k+1}^{(\alpha)}(x) = \frac{(2k+1+\alpha-x)L_k^{(\alpha)}(x) - (k+\alpha)L_{k-1}^{(\alpha)}(x)}{k+1}, \quad k \geq 1$$

The coefficient of the leading term is $\frac{(-1)^n}{n!}$, the constant term, which the value at 0, is

$$L_n^{(\alpha)}(0) = \binom{n+\alpha}{n} = \frac{\Gamma(n+\alpha+1)}{n!\Gamma(\alpha+1)}$$

The discriminant is

$$\text{Disc}\left(L_n^{(\alpha)}\right) = \prod_{j=1}^n j^{j-2n+2} (j+\alpha)^{j-1}$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Laguerre Polynomials ]# ./main
Laguerre polynomials with  $\alpha = 0$  :
L_{0}(x) = 1
L_{1}(x) = -x+1
L_{2}(x) = 1/2*x^(2)-2*x+1
L_{3}(x) = -1/6*x^(3)+1.5*x^(2)-3*x+1
L_{4}(x) = 1/24*x^(4)-0.666667*x^(3)+3*x^(2)-4*x+1
L_{5}(x) = -1/120*x^(5)+0.288333*x^(4)-1.66667*x^(3)+5*x^(2)-5*x+1
L_{6}(x) = 1/720*x^(6)-0.05*x^(5)+0.625*x^(4)-3.33333*x^(3)+7.5*x^(2)-6*x+1
L_{7}(x) = -1/5040*x^(7)+0.0897222*x^(6)-0.175*x^(5)+1.45833*x^(4)-5.83333*x^(3)+10.5*x^(2)-7*x+1
L_{8}(x) = 1/40320*x^(8)-0.0015873*x^(7)+0.0388889*x^(6)-0.466667*x^(5)+2.91667*x^(4)-9.33333*x^(3)+14*x^(2)-8*x+1
L_{9}(x) = -1/362880*x^(9)+0.000223214*x^(8)-0.00714286*x^(7)+0.116667*x^(6)-1.05*x^(5)+5.25*x^(4)-14*x^(3)+18*x^(2)-9*x+1
L_{10}(x) = 1/3628800*x^(10)-2.75573e-05*x^(9)+0.0011607*x^(8)-0.0238095*x^(7)+0.291667*x^(6)-2.1*x^(5)+8.75*x^(4)-20*x^(3)+22.5*x^(2)-10*x+1
Laguerre polynomials with  $\alpha = 1$  :
L_{0}(x) = 1
L_{1}(x) = -x+2
L_{2}(x) = 1/2*x^(2)-3*x+3
L_{3}(x) = -1/6*x^(3)+2*x^(2)-6*x+4
L_{4}(x) = 1/24*x^(4)-0.833333*x^(3)+5*x^(2)-10*x+5
L_{5}(x) = -1/120*x^(5)+0.25*x^(4)-2.5*x^(3)+10*x^(2)-15*x+6
L_{6}(x) = 1/720*x^(6)-0.0583333*x^(5)+0.875*x^(4)-5.83333*x^(3)+17.5*x^(2)-21*x+7
L_{7}(x) = -1/5040*x^(7)+0.0111111*x^(6)-0.233333*x^(5)+2.33333*x^(4)-11.6667*x^(3)+28*x^(2)-28*x+8
L_{8}(x) = 1/40320*x^(8)-0.00178571*x^(7)+0.05*x^(6)-0.7*x^(5)+5.25*x^(4)-21*x^(3)+42*x^(2)-36*x+9
L_{9}(x) = -1/362880*x^(9)+0.000248016*x^(8)-0.00892857*x^(7)+0.166667*x^(6)-1.75*x^(5)+10.5*x^(4)-35*x^(3)+60*x^(2)-45*x+10
L_{10}(x) = 1/3628800*x^(10)-3.03131e-05*x^(9)+0.00136409*x^(8)-0.0327381*x^(7)+0.458333*x^(6)-3.85*x^(5)+19.25*x^(4)-55*x^(3)+82.5*x^(2)-55*x+11
Time taken by function: 977588 microseconds
    
```

Figure 3.78: The first eleven Laguerre polynomials with $\alpha = 0$ and $\alpha = 1$ (SymIntegration/Examples/Differential Equations/Test SymIntegration Laguerre Polynomials/main.cpp).

Furthermore

$$xL'_n(x) = nL_n(x) - nL_{n-1}(x)$$

- In solution of some boundary value problems, the characteristic values can be useful:

$$L_k(0) = 1, \quad L'_k(0) = -k$$

The closed form is

$$L_n(x) = \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{k!} x^k$$

- Polynomials of negative index can be expressed using the ones with positive index

$$L_{-n}(x) = e^x L_{n-1}(-x)$$

- Laguerre functions are defined by confluent hypergeometric functions and Kummer's transformation as

$$L_n^{(\alpha)}(x) := \binom{n+\alpha}{n} M(-n, \alpha+1, x) = \frac{(\alpha+1)_n}{n!} {}_1F_1(-n; \alpha+1; x)$$

where $(a)_n$ is the Pochhammer symbol (the rising factorial), $\binom{n+\alpha}{n}$ is a generalized binomial coefficient. When n is an integer the function reduces to a polynomial of degree n . It has the alternative expression

$$L_n^{(\alpha)}(x) = \frac{(-1)^n}{n!} U(-n, \alpha+1, x)$$

in terms of Kummer's function of the second kind.

- The closed form for these generalized Laguerre polynomials of degree n is

$$L_n^{(\alpha)}(x) = \sum_{i=0}^n (-1)^i \binom{n+\alpha}{n-i} \frac{x^i}{i!}$$

derived by applying Leibniz's theorem for differentiation of a product to Rodrigues' formula.

- Laguerre polynomials have a differential operator representation, much like the closely related Hermite polynomials. Namely, let $D = \frac{d}{dx}$ and consider the differential operator $M = xD^2 + (\alpha + 1)D$. Then

$$e^{-tM}x^n = (-1)^n t^n n! L_n^{(\alpha)}\left(\frac{x}{t}\right)$$

- Laguerre's equation can be manipulated into a form that is more useful in applications:

$$u = x^{\frac{\alpha-1}{2}} e^{-\frac{x}{2}} L_n^{(\alpha)}(x)$$

which is a solution of

$$u'' + \frac{2}{x}u' + \left[\frac{\lambda}{x} - \frac{1}{4} - \frac{\alpha^2 - 1}{4x^2}\right]u = 0, \quad \text{with } \lambda = n + \frac{\alpha + 1}{2}$$

This can be further manipulated. When $l = \frac{\alpha-1}{2}$ is an integer, and $n \geq l + 1$:

$$u = x^l e^{-\frac{x}{2}} L_{n-l-1}^{(2l+1)}(x)$$

is a solution of

$$u'' + \frac{2}{x}u' + \left[\frac{\lambda}{x} - \frac{1}{4} - \frac{l(l+1)}{x^2}\right]u = 0, \quad \text{with } \lambda = n$$

The solution is often expressed in terms of derivatives instead of associated Laguerre polynomials:

$$u = x^l e^{-\frac{x}{2}} L_{n+l}^{[2l+1]}(x)$$

- Given the generating function specified above, the polynomials may be expressed in terms of a contour integral

$$L_n^{(\alpha)}(x) = \frac{1}{2\pi i} \oint_C \frac{e^{-\frac{xt}{1-t}}}{(1-t)^{\alpha+1} t^{n+1}} dt$$

where the contour circles the origin once in a counterclockwise direction without enclosing the essential singularity at 1.

- Differentiating the power series representation of a generalized Laguerre polynomial k times leads to

$$\frac{d^k}{dx^k} L_n^{(\alpha)}(x) = \begin{cases} (-1)^k L_{n-k}^{(\alpha+k)}(x), & \text{if } k \leq n \\ 0, & \text{otherwise} \end{cases}$$

This points to a special case ($\alpha = 0$) of the formula above: for integer $\alpha = k$ the generalized polynomial may be written

$$L_n^{(k)}(x) = (-1)^k \frac{d^k L_{n+k}(x)}{dx^k}$$

The shift by k sometimes causing confusion with the usual parenthesis notation for a derivative.

Moreover, the following equation holds:

$$\frac{1}{k!} \frac{d^k}{dx^k} x^\alpha L_n^{(\alpha)}(x) = \binom{n+\alpha}{k} x^{\alpha-k} L_n^{(\alpha-k)}(x)$$

which generalizes with Cauchy's formula to

$$L_n^{(\alpha')}(x) = (\alpha' - \alpha) \binom{\alpha' + n}{\alpha' - \alpha} \int_0^x \frac{t^\alpha (x-t)^{\alpha'-\alpha-1}}{x^{\alpha'}} L_n^{(\alpha)}(t) dt$$

The derivative with respect to the second variable α has the form,

$$\frac{d}{d\alpha} L_n^{(\alpha)}(x) = \sum_{i=0}^{n-1} \frac{L_i^{(\alpha)}(x)}{n-i}$$

The generalized Laguerre polynomials obey the differential equation

$$xL_n^{(\alpha)''}(x) + (\alpha + 1 - x)L_n^{(\alpha)'}(x) - nL_n^{(\alpha)}(x) = 0$$

which may be compared with the equation obeyed by the k -th derivative of the ordinary Laguerre polynomial,

$$xL_n^{[k]''}(x) + (k + 1 - x)L_n^{[k]'}(x) + (n - k)L_n^{[k]}(x) = 0$$

where $L_n^{[k]}(x) \equiv \frac{d^k L_n(x)}{dx^k}$ for this equation only.

In Sturm-Liouville form, the differential equation is

$$-(x^{\alpha+1}e^{-x} \cdot L_n^{(\alpha)}(x)')' = n \cdot x^\alpha e^{-x} \cdot L_n^{(\alpha)}(x)$$

which shows that $L_n^{(\alpha)}$ is an eigenvector for the eigenvalue n .

- The Laguerre polynomials arise in quantum mechanics, in the radial part of the solution of the Schrödinger equation for a one-electron atom. They also describe the static Wigner functions of oscillator systems in quantum mechanics in phase space. They further enter in the quantum mechanics of the Morse potential and of the 3D isotropic harmonic oscillator.

Physicists often use a definition for the Laguerre polynomials that is larger, by a factor of $(n!)$, than the definition used here.

- The generalized Laguerre polynomials are used to describe the quantum wavefunction for hydrogen atom orbitals. The convention used expresses the generalized Laguerre polynomials as

$$L_n^{(\alpha)}(x) = \frac{\Gamma(\alpha + n + 1)}{\Gamma(\alpha + 1)n!} {}_1F_1(-n; \alpha + 1; x)$$

where ${}_1F_1(a; b; x)$ is the confluent hypergeometric function. In the physics literature, the generalized Laguerre polynomials are instead defined as

$$\bar{L}_n^{(\alpha)}(x) = \frac{[\Gamma(\alpha + n + 1)]^2}{\Gamma(\alpha + 1)n!} {}_1F_1(-n; \alpha + 1; x)$$

The physics version is related to the standard version by

$$\bar{L}_n^{(\alpha)}(x) = (n + \alpha)! L_n^{(\alpha)}(x)$$

There is yet another, albeit less frequently used, convention in the physics literature

$$\tilde{L}_n^{(\alpha)}(x) = (-1)^\alpha \bar{L}_{n-\alpha}^{(\alpha)}$$

6. Hermite Polynomials

The differential equation

$$y'' - 2xy' + \lambda y = 0, \quad \text{with } \lambda = 2n$$

is Hermite's equation.

The second form of the differential equation is

$$(e^{-x^2} y')' + e^{-x^2} \lambda y = 0$$

The third form is

$$(e^{-\frac{x^2}{2}} y')'' + (\lambda + 1 - x^2)(e^{-\frac{x^2}{2}} y) = 0$$

The recurrence relation is

$$H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x)$$

this is the physicist's Hermite polynomial.

Rodrigues' formula is

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} (e^{-x^2})$$

One can define the associated Hermite functions

$$\psi_n(x) = (h_n)^{-\frac{1}{2}} e^{-\frac{x^2}{2}} H_n(x)$$

Because the multiplier is proportional to the square root of the weight function, these functions are orthogonal over $(-\infty, \infty)$ with no weight function.

The third form of the differential equation above, for the associated Hermite functions, is

$$\psi'' + (\lambda + 1 - x^2)\psi = 0$$

The associated Hermite functions arise in many areas of mathematics and physics. In quantum mechanics, they are the solutions of Schrödinger's equation for the harmonic oscillator. They are also eigenfunctions (with eigenvalue $(-i^n)$) of the continuous Fourier transform.

Many authors, particularly probabilists, use an alternate definition of the Hermite polynomials, with a weight functions of $e^{-\frac{x^2}{2}}$ instead of e^{-x^2} . If the notation He is used for these Hermite polynomials, and H for those above, then these may be characterized by

$$He_n(x) = 2^{-\frac{n}{2}} H_n\left(\frac{x}{\sqrt{2}}\right)$$

ix. Functions in SymIntegration Related to Series Solutions of Second Order Linear Equations

[SI*] We will list the basic functions that can be used related to series solutions of second order linear differential equations:

class HigherOrderODE_Homogeneous_PowerSeriesSolver

this class encapsulates the mathematical array elements and generation rules to compute the series solution for homogeneous higher order linear differential equations with constant coefficients

$$b_m y^{(m)} + b_{m-1} y^{(m-1)} + \dots + b_0 y = 0$$

with initial conditions

$$\begin{aligned} y(0) &= ic_0 \\ y'(0) &= ic_1 \\ &\vdots \\ y^{(m-1)}(0) &= ic_{m-1} \end{aligned}$$

around an ordinary point $x_0 = 0$, with $b_i \in \mathbb{R}, i = 0, 1, 2, \dots, m$ and $ic_j \in \mathbb{R}, j = 0, 1, 2, \dots, m - 1$.

Inside the **private:** we have defined **vector<double> ode_coefficients** to store the ordinary differential equation coefficients ($b_0, b_1, \dots, b_m$) the entries represents the coefficients starting from y to $y^{(m)}$, **vector<double> initial_values** to store the initial conditions value ($ic_0, ic_1, \dots, ic_{m-1}$), and **vector<double> coefficients** to dynamically stores the c_n (the coefficients of the power series solution).

The functions related to this class are:

- **computeSeries(int n)**
a **void** function to compute the power series solution with n as the terms that we specify for the series solution. This is the core algorithmic function, if you want to change the differential equation type, we only need to modify this specific recurrence formula.
- **coefficientsvector(int n)**
a **vector<double>** function to save the coefficients of the power series solution with n as the terms that we specify for the series solution.
- **evaluateAt(double x0, int n)**
a **double** function to compute the approximation of the power series solution at x_0 ($y(x_0)$), the number of terms is n and the highest degree of the power series will be $n - 1$.
- **printSeries() const**
a **void** function to display the series on the command line interface (terminal / xterm) up until the n -th terms that we have specified in the **computeSeries(int n)** function, we need to call this function (**computeSeries(int n)**) first, before we are able to display the series. This function formats output nicely by handling sign changes and omitting zero-value coefficients.

This class uses **vector<double>** to store computed elements sequentially, serving as the look-back pipeline for the relation step. The recurrence loop is using **coefficients[n+order]**, it is isolated directly by shifting index bounds inside a standard **for** loop.

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second
Order Linear Differential Equation with Constant Coefficients ]# ./main

( 1 ) y + ( 0 ) y^(1) + ( 1 ) y^(2) = 0

Initial conditions:
y(0) = 1
y^(1) (0) = 0

Series solution:

y(x) = 1 - 0.5x^2 + 0.0416667x^4 - 0.00138889x^6 + 2.48016e-05x^8 - 2.75573e-07x^10 + ...

x = 0.45

Power Series Approximation at x : 0.980447

Time taken by function: 496 microseconds

```

Figure 3.79: The power series solution of $y'' + y = 0$ computed with class **HigherOrderODE_Homogeneous_PowerSeriesSolver**, $n = 11$ (highest degree of the solution is at $n - 1$) and then compute the approximation of the series solution at $x_0 = 0.45$ (**SymIntegration/Examples/Differential Equations/Test SymIntegration Series Solutions for Homogeneous Second Order Linear Differential Equation with Constant Coefficients/main.cpp**).

class Polynomialcoeff

An elegant way to implement a power series solution for an ordinary differential equation (ODE) in C++ is to create a polynomial class. This class uses a vector to store the coefficient of a polynomial, suppose if we declare and create a polynomial like this

Polynomialcoeff P({1.0, 0.0, 2.0, -3.0});

then it will create a polynomial of name **P** with this configuration:

$$P(x) = 1 + 2x^2 - 3x^3$$

this class is very useful in order to compute series solution for second order homogeneous ODE with variable coefficients. This class is written in **/include/symintegral/dsolve.h** only.

The functions related to this class are:

- **shift_around(double x_0) const**
a **Polynomialcoeff** function to create a shifted polynomial around the point x_0 , it will return a new polynomial $p(x_{new})$ where $x_{new} = x - x_0$. This implementation applies a fast synthetic division algorithm (nested loops inside **shift_around**) to dynamically construct Taylor expansions for $P(x)$, $Q(x)$, and $R(x)$ centered at x_0 .
- **shiftToCenter(double x_0) const**
a **Polynomialcoeff** function to create a shifted polynomial around the point x_0 , it is an alternative besides **shift_around(double x_0)**.
- **shift(double x_0) const**
a **Polynomialcoeff** function to create a shifted polynomial around the point x_0 , it is an alternative besides **shift_around(double x_0)**.
- **trim()**
a **void** function that remove the trailing zeros to maintain accurate degree.
- **get_coeff(size_t m)**
a **double** function to retrieve the coefficient of the polynomial at degree m , given that the highest order of the polynomial is equal or greater than m .
- **sum_coeff() const**
a **double** function to retrieve the sum of the coefficients of the polynomial, we can call it in the source code with: **P.sum_coeff()**, given that we have declared a **Polynomialcoeff P** with its entries.
- **maxDegree() const**
an **int** function to return the maximum degree of the polynomial, we can call it in the

source code with: **P.maxDegree()**, given that we have declared a **Polynomialcoeff P** with its entries.

- **evaluateAt(double x) const**

a **double** function to evaluate the polynomial at a specific x value using Horner's method.

The **Polynomialcoeff Class** encapsulates variable coefficient arrays. It uses a secure access function **get_coeff()** to smoothly manage degree requests beyond the polynomial's boundary without crashing or forcing unnecessary manual padding. Thus, it is very useful when it is used in **class SecondOrderODE_Homogeneous_PowerSeriesSolver**.

class SecondOrderODE_Homogeneous_PowerSeriesSolver

this class is an elegant way to solve a homogeneous second order linear differential equation with variable coefficients

$$P(x)y'' + Q(x)y' + R(x)y = 0$$

by using power series this class encapsulates the mathematical array elements and generation rules to compute the series solution around an ordinary point $x_0 \in \mathbb{R}$, x_0 can be 0 or other real number. It validates that the layout center point (x_0) is an ordinary / regular point ($P(x_0) \neq 0$) before processing. This class will compute the power series solution of a homogeneous second order linear differential equation with polynomial coefficients around an ordinary point x_0 , the solution can be written as a Taylor series centered at x_0

$$y(x) = \sum_{n=0}^{\infty} c_n (x - x_0)^n$$

By shifting the coordinate system such that $x_{new} = x - x_0$, we can rewrite the coefficient polynomials $P(x)$, $Q(x)$, and $R(x)$ in terms of $(x - x_0)$ using Taylor expansion. This simplifies the computation to expanding around $x_{new,0} = 0$.

Inside the **private:** we have defined **vector<double> coefficients** to dynamically stores the c_n (the coefficients of the power series solution) the use of **std::vector** allow flexible calculation depths without memory leaks, then we also define **Polynomialcoeff P, Q, R** to represents $P(x)$, $Q(x)$, and $R(x)$ respectively.

This **class** encapsulates the differential equation coefficients that use standard recurrence mechanisms derived by shifting indices of the power series. For an equation $P(x)y'' + Q(x)y' + R(x) = 0$, the algorithm isolates the highest index element c_{n+2} dynamically based on previous computed indices $c_0, c_1, c_2, \dots, c_{n+1}$.

```
...

#ifndef SECONDORDERODE_HOMOGENEOUS_POWERSERIESSOLVER_H
#define SECONDORDERODE_HOMOGENEOUS_POWERSERIESSOLVER_H
class SecondOrderODE_Homogeneous_PowerSeriesSolver {
private:
    Polynomialcoeff P; // Coefficient for y''
    Polynomialcoeff Q; // Coefficient for y'
    Polynomialcoeff R; // Coefficient for y
    double x0;
    double y0; // for y(0)
```



```
double dy0; // for y'(0)
vector<double> coefficients; // Power series coefficients [c_0, c_1,
    c_2, ...]

public:
    SecondOrderODE_Homogeneous_PowerSeriesSolver(
        const Polynomialcoeff& P_input,
        const Polynomialcoeff& Q_input,
        const Polynomialcoeff& R_input,
        double x0_input,
        double y0_input,
        double dy0_input) ;

    void computeSeries(int terms) ;
    vector<double> coefficientsvector(int terms) ;
    double evaluateAt(double x, int terms) ;
    void printCoefficients() const ;
    void printSolution() const ;

};

#endif

...
```

Code 45: *include/symintegral/dsolve.h*

The functions related to this class are:

- **computeSeries(int n)**
a **void** function to compute the power series solution with n as the terms that we specify for the series solution. It dynamically shifts indices and unrolls the algorithmic Cauchy product loop configurations to automatically decouple the targeted c_{n+2} parameter from past historical weights.
- **coefficientsvector(int n)**
a **vector<double>** function to save the coefficients of the power series solution with n as the terms that we specify for the series solution.
- **evaluateAt(double x_0 , int n)**
a **double** function to compute the approximation of the power series solution at x_0 ($y(x_0)$), the number of terms is n and the highest degree of the power series will be $n - 1$. This function applies Horner's-like incremental power aggregation to minimize floating-point multiplication errors. The Horner-adjacent evaluation approach avoids using the expensive **std::pow** operations inside loops by incrementally multiplying the current x value. When we employ Horner's evaluation schema, it can safely aggregate terms without experiencing float overflows or scaling precision decay at localized evaluations.
- **printCoefficients() const**
a **void** function to display the coefficients of the solution ($y(x)$) for the differential equation, we need to call this function (**computeSeries(int n)**) first, before we are able to display the coefficients.

- **printSolution() const**

a **void** function to display the solution ($y(x)$) on the command line interface (terminal / xterm) up until the n -th terms that we have specified in the **computeSeries(int n)** function, we need to call this function (**computeSeries(int n)**) first, before we are able to display the series. This function formats output nicely by handling sign changes and omitting zero-value coefficients.

Recurrence Tracking

By substituting $y(x_{new}) = \sum_{n=0}^{\infty} c_n x_{new}^n$ into the differential equation, the total coefficient multiplying x_{new}^m aggregates contributions from the j -th derivative coefficients. **Isolating the Next Coefficient**

At step m , the coefficient of x_{new}^m uniquely relies on known entries up to c_{m+1} , alongside exactly one unknown entry: $p_0(m+2)(m+1)c_{m+2}$. Isolating c_{m+2} yields:

$$c_{m+2} = \frac{-1}{P(x_0)(m+2)(m+1)} \left[\sum_{j=1}^m p_j(m-j+2)(m-j+1)c_{m-j+2} + q_j(m-j+1)c_{m-j+1} + r(j)c_{m-j} \right]$$

The algorithm safely loops up to n terms, resolving one higher order coefficient with every incremental iteration step.

bernoullinumbers(int n)

to compute the n -th term of Bernoulli numbers.

probabilists_Hermitepolynomials(int n , const Symbolic & x)

to compute the first n probabilist's Hermite polynomials.

physicists_Hermitepolynomials(int n , const Symbolic & x)

to compute the first n physicist's Hermite polynomials.

Jacobipolynomials(double α , double β , int n , const Symbolic & x)

to compute the first n Jacobi polynomials.

IV. Series Solutions of Second Order Linear Equations

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Jacobian Polynomials ]# ./main
Jacobian polynomials:
 $\alpha = 2, \beta = 1$ 
P_{0}(x) = 1
P_{1}(x) = 2.5*x+0.5
P_{2}(x) = 1.5*x+5.25*x^2-0.75
P_{3}(x) = -3.5*x+3.5*x^2+10.5*x^3-0.5
P_{4}(x) = -2.5*x-11.25*x^2+7.5*x^3+20.625*x^4+0.625
P_{5}(x) = 4.21875*x-8.4375*x^2-30.9375*x^3+15.4688*x^4+40.2188*x^5+0.46875
P_{6}(x) = 3.28125*x+18.0469*x^2-24.0625*x^3-78.2031*x^4+31.2812*x^5+78.2031*x^6-0.546875
P_{7}(x) = -4.8125*x+14.4375*x^2+62.5625*x^3-62.5625*x^4-187.688*x^5+62.5625*x^6+151.938*x^7-0.4375
P_{8}(x) = -3.9375*x-25.5938*x^2+51.1875*x^3+191.953*x^4-153.562*x^5-435.094*x^6+124.312*x^7+295.242*x^8+0.492188
P_{9}(x) = 5.33203*x-21.3281*x^2-106.641*x^3+159.961*x^4+543.867*x^5-362.578*x^6-984.141*x^7+246.035*x^8+574.082*x^9+0.410156
P_{10}(x) = 4.51172*x+33.8379*x^2-90.2344*x^3-383.496*x^4+460.195*x^5+1457.29*x^6-832.734*x^7-2185.93*x^8+485.762*x^9+1117.25*x^10-0.451172

Legendre polynomials:
 $\alpha = 0, \beta = 0$ 
P_{0}(x) = 1
P_{1}(x) = x
P_{2}(x) = 1.5*x^2-0.5
P_{3}(x) = -1.5*x+2.5*x^3
P_{4}(x) = -3.75*x^2+4.375*x^4+0.375
P_{5}(x) = 1.875*x-8.75*x^3+7.875*x^5
P_{6}(x) = 6.5625*x^2-19.6875*x^4+14.4375*x^6-0.3125
P_{7}(x) = -2.1875*x+19.6875*x^3-43.3125*x^5+26.8125*x^7
P_{8}(x) = -9.84375*x^2+54.1406*x^4-93.8438*x^6+59.2734*x^8+0.273438
P_{9}(x) = 2.46094*x-36.0938*x^3+140.766*x^5-281.094*x^7+94.9609*x^9
P_{10}(x) = -13.5352*x^2-117.305*x^4+351.914*x^6-427.324*x^8+180.426*x^10-0.246094

Gegenbauer polynomials:
 $\alpha = 1, \beta = 1$ 
P_{0}(x) = 1
P_{1}(x) = 2*x
P_{2}(x) = 3.75*x^2-0.75
P_{3}(x) = -3*x+3*x^3
P_{4}(x) = -8.75*x^2+13.125*x^4+0.625
P_{5}(x) = 3.75*x-22.5*x^3+24.75*x^5
P_{6}(x) = 14.7656*x^2-54.1406*x^4+46.9219*x^6-0.546875
P_{7}(x) = -4.375*x+48.125*x^3-125.125*x^5+89.375*x^7
P_{8}(x) = -21.6562*x^2+140.766*x^4-281.531*x^6+170.93*x^8+0.492188
P_{9}(x) = 4.92188*x-85.3125*x^3+383.906*x^5-621.562*x^7+328.047*x^9
P_{10}(x) = 29.3262*x^2-293.262*x^4+997.09*x^6-1353.19*x^8+631.49*x^10-0.451172

Time taken by function: 8256343 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Jacobian Polynomials ]#

```

Figure 3.80: The first eleven Jacobi polynomials, Legendre polynomials, and Gegenbauer polynomials (SymIntegration/Examples/Differential Equations/Test SymIntegration Jacobian Polynomials/main.cpp).

firstkind_Chebyshevpolynomials(int n , const Symbolic & x)
to compute the first n Chebyshev polynomials of the first kind.

secondkind_Chebyshevpolynomials(int n , const Symbolic & x)
to compute the first n Chebyshev polynomials of the second kind.

Gegenbauerpolynomials(double α , int n , const Symbolic & x)
to compute the first n Gegenbauer polynomials with parameter α .

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Differential Equations/Test SymIntegration Gegenbauer Polynomials ]# ./main
Gegenbauer polynomials with  $\alpha = 1$  :
C_{0}(x) = 1
C_{1}(x) = 2*x
C_{2}(x) = 4*x^2-1
C_{3}(x) = 8*x^3-4*x
C_{4}(x) = 16*x^4-12*x^2+1
C_{5}(x) = 32*x^5-32*x^3+6*x
C_{6}(x) = 64*x^6-80*x^4+24*x^2-1
C_{7}(x) = 128*x^7-192*x^5+80*x^3-8*x
C_{8}(x) = 256*x^8-448*x^6+240*x^4-40*x^2+1
C_{9}(x) = 512*x^9-1024*x^7+672*x^5-160*x^3+10*x
C_{10}(x) = 1024*x^10-2304*x^8+1792*x^6-560*x^4+60*x^2-1

Time taken by function: 139430 microseconds

```

Figure 3.81: The first eleven Gegenbauer polynomials with $\alpha = 1$ (SymIntegration/Examples/Differential Equations/Test SymIntegration Gegenbauer Polynomials/main.cpp).

Laguerrepolynomials(double α , int n , const Symbolic & x)
to compute the first n generalized Laguerre polynomials with parameter α .

V. THE LAPLACE TRANSFORM

[SI*] **Solution of Initial Value Problems**

[SI*]

VI. SYSTEMS OF FIRST ORDER LINEAR EQUATIONS

[SI*] **Basic Theory**

[SI*]

VII. NUMERICAL METHODS

[SI*] **The Euler or Tangent Line Method**

[SI*] **The Runge-Kutta Method**

VIII. NONLINEAR DIFFERENTIAL EQUATIONS

[S^{*}] **The Phase Plane**

IX. PARTIAL DIFFERENTIAL EQUATIONS

[SI*] **The Phase Plane**

X. BOUNDARY VALUE PROBLEMS AND STURM-LIOUVILLE THEORY

[SI*] **The Occurrence of Two-Point Boundary Value Problems**

[SI*] **Nonhomogeneous Boundary Value Problems**

Chapter 4

Probability and Statistics

"Education is the best provision for old age." - Aristotle

At a glance, people might be thinking that probability is only numerical computation, the probability that a head will show up on a coin that is tossed upward is around 0.5 on a fair coin. After more thorough reading, statistics and probability are actually more symbolic than numeric. We just substitute the random variable with a number that we want to test / to know the probability a certain event will occur is how much. That is the main reason, why we put statistics and probability computation in SymIntegration.

On August 23rd, 2025 we have created a new source code **statistics.cpp** that will handle the probability and statistics computation, like computing probability mass function of a certain discrete distribution, the moment generating function and to compute the mean and variance. It could come in handy one day.

I. PROCESSING AND SUMMARIZING DATA WITH STATISTICS

[SI*] Statistical methods are used to analyze data from an industrial process (manufacturing, energy sources, drug testing). Statistical methods are involved with gathering of information or **scientific data**, which then will be processed to extract the knowledge from the data and improve the **quality** of the process.

i. Descriptive Statistics Computation

Suppose we have a univariate data of a Mathematics examination result for 25 students, compute its' mean, median, mode, standard deviation, and variance.

Solution:

To compute the mean we will use this formula:

$$\bar{x} = \frac{\sum x}{n} \quad (4.1)$$

Mode is the value that occur frequently, and Median is the middle value after the data is sorted, if the size of the data is even, then the median is the average of the two middle data.

Standard deviation can be computed with this formula:

$$S_x = \sqrt{\frac{\sum x^2}{n} - (\bar{x})^2} \quad (4.2)$$

Standard deviation is the square root of variance.

In SymIntegration there is a function that can be used to compute the standard descriptive statistics, it is:

double descriptivestatistics(vector<double>)

the input is a vector with double as the data type, since it can also represents integer.

We have saved the grade data in a textfile named **vectorx.txt**, we can just load / call it from the **main.cpp** file.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Load Vector from textfile and Compute Descriptive Statistics ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Load Vector from textfile and Compute Descriptive Statistics ]# ./main

Statistics Descriptive
Mode : 6
Median : 6
Mean : 6.2
Quantile 1/4: 5
Quantile 3/4: 8
Variance : 6.41667
Standard deviation : 2.53311

Time taken by function: 833 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Load Vector from textfile and Compute Descriptive Statistics ]#
```

Figure 4.1: The computation of descriptive statistics with SymIntegration took 833 microseconds (*SymIntegration/Examples/Statistics/Test SymIntegration Load Vector from textfile and Compute Descriptive Statistics/main.cpp*).

II. RANDOM VARIABLES AND PROBABILITY DISTRIBUTIONS

Definition 4.1: Random Variable

A random variable is a function that associates a real number with each element in the sample space.

We shall use a capital letter, say X , to denote a random variable and its corresponding small letter, x in this case, for one of its values.

Definition 4.2: Discrete Sample Space

If a sample space contains a finite number of possibilities or an unending sequence with as many elements as there are whole numbers, it is called a discrete sample space.

Definition 4.3: Continuous Sample Space

If a sample space contains an infinite number of possibilities equal to the number of points on a line segment, it is called a continuous sample space.

i. Discrete Probability Distributions

Definition 4.4: Probability Mass Function

If X is discrete, the probability mass function of X is defined as

$$f(x) = P(X = x), \quad \forall x \in \mathbb{R} \quad (4.3)$$

the probability is concentrated at only discrete values in the range of X , and is zero for all other values. f must also obey the basic rules of probability. That is, f must be non-negative

$$f(x) \geq 0$$

and the sum of all probabilities should add to 1

$$\sum_x f(x) = 1$$

Definition 4.5: Discrete Cumulative Distribution Function

The cumulative distribution function $F(x)$ of a discrete random variable X with probability distribution $f(x)$ is

$$F(x) = P(X \leq x) = \sum_{t \leq x} f(t), \quad \text{for } -\infty < x < \infty \quad (4.4)$$

ii. Continuous Probability Distributions

Definition 4.6: Probability Density Function

If X is continuous, its range is the entire set of real numbers $\mathbb{R}$. The probability of any specific value x is only one out of the infinitely many possible values in the range of X , which means that

$$P(X = x) = 0$$

for all $x \in \mathbb{R}$. The probability mass is spread so thinly over the range of values, that it can be measured only over intervals $[a, b] \subset \mathbb{R}$, rather than at specific points.

the probability density function of X that takes on values in any interval $[a, b] \subset \mathbb{R}$ is defined as

$$P(X \in [a, b]) = \int_a^b f(x) dx \quad (4.5)$$

the density function f must satisfy the basic laws of probability

$$f(x) \geq 0, \quad \forall x \in \mathbb{R}$$

and

$$\int_{-\infty}^{\infty} f(x) dx = 1$$

Definition 4.7: Continuous Cumulative Distribution Function

The function $f(x)$ is a probability density function (pdf) for the continuous random variable X , defined over the set of real numbers, if

$$f(x) \geq 0, \quad \forall x \in \mathbb{R} \quad (4.6)$$

$$\int_{-\infty}^{\infty} f(x) dx = 1 \quad (4.7)$$

$$P(a < X < b) = \int_a^b f(x) dx \quad (4.8)$$

Definition 4.8: Cumulative Distribution Function

For any random variable X , whether discrete or continuous, we can define the cumulative distribution function (cdf) as

$$F : \mathbb{R} \rightarrow [0, 1]$$

that gives the probability of observing a value at most some given value x

$$F(x) = P(X \leq x), \quad \forall -\infty < x < \infty \quad (4.9)$$

when X is discrete, F is given as

$$F(x) = P(X \leq x) = \sum_{u \leq x} f(u) \quad (4.10)$$

and when X is continuous, F is given as

$$F(x) = P(X \leq x) = \int_{-\infty}^x f(u) du \quad (4.11)$$

iii. Joint Probability Distributions

Definition 4.9: Joint Probability Distribution

The function $f(x, y)$ is a joint probability distribution or probability mass function of the discrete random variables X and Y if

$$f(x, y) \geq 0, \quad \forall (x, y) \quad (4.12)$$

$$\sum_x \sum_y f(x, y) = 1 \quad (4.13)$$

$$P(X = x, Y = y) = f(x, y) \quad (4.14)$$

For any region A in the xy plane,

$$P[(X, Y) \in A] = \sum_A f(x, y) \quad (4.15)$$

Definition 4.10: Joint Density Function

The function $f(x, y)$ is a joint density function of the continuous random variables X and Y if

$$f(x, y) \geq 0, \quad \forall (x, y) \quad (4.16)$$

$f(x, y)$ is a surface lying above the xy plane, and $P[(X, Y) \in A]$, where A is any region in the xy plane. The joint density function is equal to the volume of the right cylinder bounded by the base A and the surface.

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = 1 \quad (4.17)$$

$$P[(X, Y) \in A] = \int \int_A f(x, y) \, dx \, dy \quad (4.18)$$

for any region A in the xy plane.

Definition 4.11: Marginal Distributions

Given the joint probability distribution $f(x, y)$ of the discrete random variables X and Y , the probability distribution $g(x)$ of X alone is obtained by summing $f(x, y)$ over the values of Y . Similarly, the probability distribution $h(y)$ of Y alone is obtained by summing $f(x, y)$ over the values of X .

We define $g(x)$ and $h(y)$ to be the marginal distributions of X and Y , respectively. When X and Y are continuous random variables, summations are replaced by integrals.

The marginal distributions of X alone and Y alone are

$$\begin{aligned} g(x) &= \sum_y f(x, y) \\ h(y) &= \sum_x f(x, y) \end{aligned} \quad (4.19)$$

for the discrete case, and

$$\begin{aligned} g(x) &= \int_{-\infty}^{\infty} f(x, y) \, dy \\ h(y) &= \int_{-\infty}^{\infty} f(x, y) \, dx \end{aligned} \quad (4.20)$$

for the continuous case.

Definition 4.12: Conditional Distribution

Let X and Y be two random variables, discrete or continuous. The conditional distribution of the random variable Y given that $X = x$ is

$$f(y|x) = \frac{f(x,y)}{g(x)} \quad (4.21)$$

provided $g(x) > 0$.

Similarly, the conditional distribution of X given that $Y = y$ is

$$f(x|y) = \frac{f(x,y)}{h(y)} \quad (4.22)$$

provided $h(y) > 0$.

If we wish to find the probability that the discrete random variable X falls between a and b when it is known that the discrete variable $Y = y$, we evaluate

$$P(a < X < b | Y = y) = \sum_{a < x < b} f(x|y) \quad (4.23)$$

where the summation extends over all values of X between a and b . When X and Y are continuous, we evaluate

$$P(a < X < b | Y = y) = \int_a^b f(x|y) dx \quad (4.24)$$

Definition 4.13: Statistically Independent

Let X and Y be two random variables, discrete or continuous, with joint probability distribution $f(x,y)$ and marginal distributions $g(x)$ and $h(y)$, respectively. The random variables X and Y are said to be statistically independent if and only if

$$f(x,y) = g(x)h(y) \quad (4.25)$$

for all (x,y) within their range.

Definition 4.14: Mutually Statistically Independent

Let $X_1, X_2, \dots, X_n$ be n random variables, discrete or continuous, with joint probability distribution $f(x_1, x_2, \dots, x_n)$ and marginal distribution $f_1(x_1), f_2(x_2), \dots, f_n(x_n)$, respectively. The random variables $X_1, X_2, \dots, X_n$ are said to be mutually statistically independent if and only if

$$f(x_1, x_2, \dots, x_n) = f_1(x_1)f_2(x_2) \dots f_n(x_n) \quad (4.26)$$

for all $(x_1, x_2, \dots, x_n)$ within their range.

Definition 4.15: Mean

Let X be a random variable with probability distribution $f(x)$. The mean, or expected value, of X is

$$\mu = E(X) = \sum_x x f(x) \quad (4.27)$$

if X is discrete, and

$$\mu = E(X) = \int_{-\infty}^{\infty} x f(x) dx \quad (4.28)$$

if X is continuous.

Theorem 4.1: Expected Value of a Random Variable $g(X)$

Let X be a random variable with probability distribution $f(x)$. The expected value of the random variable $g(X)$ is

$$\mu_{g(X)} = E[g(X)] = \sum_x g(x) f(x) \quad (4.29)$$

if X is discrete, and

$$\mu_{g(X)} = E[g(X)] = \int_{-\infty}^{\infty} g(x) f(x) dx \quad (4.30)$$

if X is continuous.

Definition 4.16: Mean for Two Random Variables

Let X and Y be random variables with joint probability distribution $f(x, y)$. The mean, or expected value, of the random variable $g(X, Y)$ is

$$\mu_{g(X,Y)} = E[g(X, Y)] = \sum_x \sum_y g(x, y) f(x, y) \quad (4.31)$$

if X and Y are discrete, and

$$\mu_{g(X,Y)} = E[g(X, Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f(x, y) dx dy \quad (4.32)$$

if X and Y are continuous.

Definition 4.17: Variance

Let X be a random variable with probability distribution $f(x)$ and mean μ . The variance of X is

$$\sigma^2 = E[(X - \mu)^2] = \sum_x (x - \mu)^2 f(x) \quad (4.33)$$

if X is discrete, and

$$\sigma^2 = E[(X - \mu)^2] = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx \quad (4.34)$$

if X is continuous.

The positive square root of the variance, σ , is called the standard deviation of X . The quantity $x - \mu$ is called the deviation of an observation from its mean.

Theorem 4.2: Variance

The variance of a random variable X is

$$\sigma^2 = E(X^2) - \mu^2 \quad (4.35)$$

Theorem 4.3: Variance of Random Variable $g(X)$

Let X be a random variable with probability distribution $f(x)$. The variance of the random variable $g(X)$ is

$$\sigma_{g(X)}^2 = E\{[g(X) - \mu_{g(X)}]^2\} = \sum_x [g(x) - \mu_{g(X)}]^2 f(x) \quad (4.36)$$

if X is discrete, and

$$\sigma_{g(X)}^2 = E\{[g(X) - \mu_{g(X)}]^2\} = \int_{-\infty}^{\infty} [g(x) - \mu_{g(X)}]^2 f(x) dx \quad (4.37)$$

Definition 4.18: Covariance

Let X and Y be random variables with joint probability distribution $f(x, y)$. The covariance of X and Y is

$$\sigma_{XY} = E[(X - \mu_X)(Y - \mu_Y)] = \sum_x \sum_y (x - \mu_X)(y - \mu_Y) f(x, y) \quad (4.38)$$

if X and Y are discrete, and

$$\sigma_{XY} = E[(X - \mu_X)(Y - \mu_Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x - \mu_X)(y - \mu_Y) f(x, y) dx dy \quad (4.39)$$

if X and Y are continuous.

The covariance between two random variables is a measure of the nature of the association between the two.

If large values of X often result in large values of Y or small values of X result in small values of Y , positive $X - \mu_X$ will often result in positive $Y - \mu_Y$ and negative $X - \mu_X$ will often result in negative $Y - \mu_Y$.

Thus, the product $(X - \mu_X)(Y - \mu_Y)$ will tend to be positive. On the other hand, if large X values often result in small Y values, the product $(X - \mu_X)(Y - \mu_Y)$ will tend to be negative.

When X and Y are statistically independent, it can be shown that the covariance is zero. The converse, however, is not generally true. Two variables may have zero covariance and still not be statistically independent. Note that the covariance only describes the linear relationship between two random variables. Therefore, if a covariance between X and Y is zero, X and Y may have a nonlinear relationship, which means that they are not necessarily independent.

Theorem 4.4: Covariance

The covariance of two random variables X and Y with means μ_X and μ_Y , respectively, is given by

$$\sigma_{XY} = E(XY) - \mu_X \mu_Y \quad (4.40)$$

Definition 4.19: Correlation

Let X and Y be random variables with covariance σ_{XY} and standard deviations σ_X and σ_Y , respectively. The correlation coefficient of X and Y is

$$\rho_{XY} = \frac{\sigma_{XY}}{\sigma_X \sigma_Y} \quad (4.41)$$

it should be clear that ρ_{XY} is free of the units of X and Y . The correlation coefficient satisfies the inequality $-1 \leq \rho_{XY} \leq 1$. It assumes a value of zero when $\sigma_{XY} = 0$.

iv. Covariance and Vectorized Moments

Given two random variables, X and Y , with respective means, μ_X and μ_Y , the covariance is defined by

$$\text{Cov}(X, Y) = E[(X - \mu_X)(Y - \mu_Y)] = E[XY] - \mu_X\mu_Y$$

The second formula follows by expansion. Notice also that

$$\text{Cov}(X, X) = \text{Var}(X)$$

the covariance is a common measure of the relationship between the two random variables, where if $\text{Cov}(X, Y) = 0$, we say the random variables are uncorrelated. Furthermore, if $\text{Cov}(X, Y) \neq 0$, the sign of the covariance gives an indication of the direction of the relationship.

Another important concept is the correlation coefficient,

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \quad (4.42)$$

It is a normalized form of the covariance with $-1 \leq \rho_{XY} \leq 1$. Values nearing ± 1 indicate a very strong linear relationship between X and Y , whereas values near or at 0 indicate a lack of a linear relationship.

Note that if X and Y are independent random variables, then $\text{Cov}(X, Y) = 0$ and hence $\rho_{XY} = 0$. However, the opposite case does not always hold, since, in general $\rho_{XY} = 0$ does not imply independence. Nevertheless for jointly normal random variables it does.

Consider now a random vector $\mathbf{X} = (X_1, \dots, X_n)$, taken as a column vector. It can be described by moments in an analogous manner to a scalar random variable. A key quantity is the mean vector,

$$\mu_X := [E[X_1], E[X_2], \dots, E[X_n]]^T$$

Furthermore, the covariance matrix is the matrix defined by the expectation (taken element-wise) of the (outer product) random matrix given by $(X - \mu_X)(X - \mu_X)^T$, and is expressed as

$$\Sigma_X = \text{Cov}(X) = E[(X - \mu_X)(X - \mu_X)^T] \quad (4.43)$$

As can be verified, the i, j -th element of Σ_X is $\text{Cov}(X_i, X_j)$ and hence the diagonal elements are the variances.

The covariance matrix is a fundamental concept in statistics and data analysis, providing insight into the relationships between multiple variables. It captures the extent to which two variables change together and is essential for the various applications such as the Principal Component Analysis (PCA), financial analysis and multivariate statistics. By understanding and interpreting the covariance matrix, we can make more informed decisions and uncover deeper patterns in data.

v. Linear Combinations and Transformations

We now consider linear transformations applied to random vectors. For any collection of random variables,

$$E[X_1 + \dots + x_n] = E[X_1] + \dots + E[X_n]$$

For uncorrelated random variables,

$$\text{Var}(X_1 + \cdots + X_n) = \text{Var}(X_1) + \cdots + \text{Var}(X_n)$$

More generally, if we allow the random variables to be correlated, then

$$\text{Var}(X_1 + \cdots + X_n) = \text{Var}(X_1) + \cdots + \text{Var}(X_n) + 2 \sum_{i < j} \text{Cov}(X_i, X_j) \quad (4.44)$$

Note that the right-hand side of Eq. (4.51) is the sum of the elements of the matrix $\text{Cov}(X_1, \dots, X_n)$. this is a special case of a more general affine transformation, where we take a random vector $\mathbf{X} = (x_1, \dots, x_n)$ with covariance matrix Σ_X , and an $m \times n$ matrix A and m vector $\mathbf{b}$. We then set

$$\mathbf{Y} = A\mathbf{X} + \mathbf{b} \quad (4.45)$$

In this case, the new random vector $\mathbf{Y}$ exhibits mean and covariance

$$\begin{aligned} E[\mathbf{Y}] &= AE[\mathbf{X}] + \mathbf{b} \\ \text{Cov}(\mathbf{Y}) &= A\Sigma_X A^T \end{aligned} \quad (4.46)$$

Now to retrieve Eq. (4.51), we set $1 \times n$ matrix $A = [1, \dots, 1]$ and observe that $A\Sigma_X A^T$ is a sum of all of the elements of Σ_X .

vi. The Cholesky Decomposition and Generating Random Vectors

If you wish to create an n -dimensional random vector $\mathbf{Y}$ with some specified mean vector μ_Y and covariance matrix Σ_Y . That is, μ_Y and Σ_Y are known.

The formulas in Eqs. (4.53) yield a potential recipe for such a task if we are given a random vector $\mathbf{X}$ with zero-mean and identity-covariance matrix ($\Sigma_X = I$). FOr example, in the context of Monte Carlo random variable generation, creating such a random vector $\mathbf{X}$ is trivial-just generate a sequence of n i.i.d. normal $(0, 1)$ random variables.

Now, apply the affine transformation Eq. (4.52) on $\mathbf{X}$ with $\mathbf{b} = \mu_Y$ and a matrix A that satisfies

$$\Sigma_Y = AA^T \quad (4.47)$$

Now Eq. (4.53) guarantees that $\mathbf{Y}$ has the desired μ_Y and Σ_Y .

vii. Generate Random Number in SymIntegration

Generating random number is the essence or basic need in statistical inference, it is very useful as well if we want to simulate physics phenomena or forecasting weather with random initial condition.

In SymIntegration we have two functions to generate random number, the first one is:

randomnumberint(int a, int b, int n)

with a and b as the range so the number generated will be in $[a, b]$, and n is the amount of random number that we want to generate. This function will generate integer random number only.

The second function is:

randomnumberreal(double a, double b, int n)

with a and b as the range so the number generated will be in $[a, b]$, and n is the amount of random number that we want to generate. This function will generate real random number, in C++ we use **double** data type to represent the real number here.

The functions are written in **src/statistics.cpp** with the header that corresponds to this is in **/include/symintegral/statistics.h**. We are using uniform distribution to generate the random number to ensure that the probability of each number in the range to be picked is the same, equal opportunity for all.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Generate Random Number from Function ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Generate Random Number from Function ]# ./main
Random integer:
19
129
123
133
93
58
173
82
88
28
173
Random double:
1.06899
0.683921
4.21156
3.94917
2.27386
0.971918
4.0864
3.09706
2.44087
3.7999
Time taken by function: 759 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Generate Random Number from Function ]# []
```

Figure 4.2: The random number generation with SymIntegration took 759 microseconds (*SymIntegration/Examples/Statistics/Test SymIntegration Generate Random Number from Function/main.cpp*).

We are using the standard C++ code **std::random_device rd** as a uniform random bit generator that is intended to produce non-deterministic random numbers, typically by accessing a hardware entropy source or a system-provided source of randomness. Its purpose is to provide a high-quality, truly random seed for other pseudo-random number generators. the **rd** is an object of type **std::random_device**.

Then, we also use **std::mt19937 generate** as the Mersenne Twister engine, a widely used and well-regarded pseudo-random number generator (PRNG). It produces a long sequence of pseudo-random numbers based on an initial seed. The **generate** is an object of type **std::mt19937**,

the **rd()** part calls the **operator()** of the **std::random_device** object, which returns a single random value. This value is then used as the seed to initialize the **std::mt19937** engine **generate**.

std::random_device provides a non-deterministic seed, ensuring that each run of the program will likely produce a different sequence of random numbers. This seed then initializes **std::mt19937**, which is a high-quality and efficient pseudo-random number generator suitable for most non-cryptographic applications. Subsequent calls to **generate()** or distributions associate with **generate** will produce the pseudo-random numbers.

viii. Create Covariance Matrix in SymIntegration

The general form of a covariance matrix is as follow:

$$\Sigma_X = \begin{bmatrix} \sigma_{11} & \sigma_{12} & \dots & \sigma_{1n} \\ \sigma_{21} & \sigma_{22} & \dots & \sigma_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{n1} & \sigma_{n2} & \dots & \sigma_{nn} \end{bmatrix} = \begin{bmatrix} \text{Cov}(X_1, X_1) & \text{Cov}(X_1, X_2) & \dots & \text{Cov}(X_1, X_n) \\ \text{Cov}(X_2, X_1) & \text{Cov}(X_2, X_2) & \dots & \text{Cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(X_n, X_1) & \text{Cov}(X_n, X_2) & \dots & \text{Cov}(X_n, X_n) \end{bmatrix}$$

we can tell that covariance matrix is a symmetric matrix with the property of $a_{ij} = a_{ji}$. The formula to compute σ_{ij} can be seen at Eq. (4.47).

For sample covariance the formula is as follow:

$$\sigma_{xy} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{n - 1}$$

Now suppose we have a mid-term exam results by Sine, Sweden, and Bludut in Linear Algebra and Differential Equations. Make a covariance matrix.

Student	Linear Algebra	Differential Equations
Sine	80	70
Sweden	63	20
Bludut	100	50

Table 4.1: The mid-term exam results.

With SymIntegration we can create the covariance matrix with this function:
covariancematrix(vector<vector<double>>)

The output will be a **SymbolicMatrix** or **Matrix<Symbolic>**, a class that is originally made from SymbolicC++. The goodness of this SymbolicMatrix is that it can also contain symbolic input.

The result will be show that

$$\begin{aligned} \text{Cov}(X, X) &= \sigma_{11} = 343 \\ \text{Cov}(X, Y) &= \sigma_{12} = 260 \\ \text{Cov}(Y, Y) &= \sigma_{22} = 633.33 \end{aligned}$$

the covariance for X (the Linear Algebra mid-term score) and Y (the Differential Equations mid-term score) is 260, as this is a positive number, it means that when X increases (or decreases) Y also increases (or decreases).

While the diagonal elements are 343 and 633.33 indicate the variance in data sets X and Y respectively. X shows the lowest variance, while Y shows the highest variance.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Create Covariance Matrix from Textfile ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Create Covariance Matrix from Textfile ]# ./main
Data:
      80      70
      63      20
     100      50

The covariance matrix:
[ 343  260 ]
[ 260 633.333]

Time taken by function: 1265 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Create Covariance Matrix from Textfile ]#
```

Figure 4.3: The covariance matrix for this example case. (*SymIntegration/Examples/Statistics/Test SymIntegration Create Covariance Matrix from Textfile/main.cpp*).

ix. Create Cholesky Decomposition in SymIntegration

Perform a Cholesky decomposition on

$$\Sigma_X = \begin{bmatrix} 4 & 12 & -16 \\ 12 & 37 & -43 \\ -16 & -43 & 98 \end{bmatrix}$$

We will use this formula

$$A_{ii} = \sqrt{\Sigma_{X_{ii}} - \sum_{k=0}^{i-1} (A_{ik})^2}$$

The result is

$$\begin{bmatrix} 4 & 12 & -16 \\ 12 & 37 & -43 \\ -16 & -43 & 98 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 6 & 1 & 0 \\ -8 & 5 & 3 \end{bmatrix} \begin{bmatrix} 2 & 6 & -8 \\ 0 & 1 & 5 \\ 0 & 0 & 3 \end{bmatrix}$$

In SymIntegration the Cholesky decomposition can be performed with this function:
choleskyDecomposition(vector<vector<double>> matrix)

The function is **void**, the goodness of using **void** is we don't need to have **return ..** at the end of the function, and we can call the result / the output after we call the function.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Cholesky Decomposition ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Cholesky Decomposition ]# ./main
Data:
      4      12     -16
      12     37     -43
     -16    -43     98

A =
      2      0      0
      6      1      0
     -8      5      3

A^{T} =
      2      6     -8
      0      1      5
      0      0      3

Time taken by function: 889 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Cholesky Decomposition ]#
```

Figure 4.4: The Cholesky decomposition for a square matrix of size 3, we load the matrix from textfile *matrix.txt*.
 (SymIntegration/Examples/Statistics/Test SymIntegration Cholesky Decomposition/main.cpp).

III. DISCRETE PROBABILITY DISTRIBUTIONS

[SI*] We will start with the Bernoulli process, then we will cover the definition of other discrete distributions.

Definition 4.20: Bernoulli Process

An experiment that consists of repeated trials, each with two possible outcomes that may be labeled success or failure is called a Bernoulli process.

The Bernoulli process must possess the following properties:

1. The experiment consists of repeated trials.
2. Each trial results in an outcome that may be classified as a success or a failure.
3. The probability of success, denoted by p , remains constant from trial to trial.
4. The repeated trials are independent.

Definition 4.21: Binomial Distribution

The number X of successes in n Bernoulli trials is called a binomial random variable. The probability distribution of this discrete random variable is called the binomial distribution, and its values will be denoted by $b(x; n, p)$.

A Bernoulli trial can result in a success with probability p and a failure with probability $q = 1 - p$. Then the probability distribution of the binomial random variable X , the number of successes in n independent trials, is

$$b(x; n, p) = \binom{n}{x} p^x (q^{n-x}), \quad x = 0, 1, 2, \dots, n \quad (4.48)$$

Theorem 4.5: Mean and Variance of the Binomial Distribution

The mean and variance of the binomial distribution $b(x; n, p)$ are

$$\mu = np \quad (4.49)$$

$$\sigma^2 = npq \quad (4.50)$$

Definition 4.22: Negative Binomial Distribution

The number X of trials required to produce k successes in a negative binomial experiment is called a negative binomial random variable, and its probability distribution is called the negative binomial distribution.

If repeated independent trials can result in a success with probability p and a failure with probability $q = 1 - p$, then the probability distribution of the random variable, X , the number of the trial on which the k -th success occurs, is

$$b^*(x; k, p) = \binom{x-1}{k-1} p^k q^{x-k}, \quad x = k, k+1, k+2, \dots \quad (4.51)$$

Definition 4.23: Geometric Distribution

If repeated independent trials can result in a success with probability p and a failure with probability $q = 1 - p$, then the probability distribution of the random variable X , the number of the trial on which the first success occurs, is

$$g(x; p) = pq^{x-1}, \quad x = 1, 2, 3, \dots \quad (4.52)$$

Theorem 4.6: The Mean and Variance of the Geometric Distribution

The mean and variance of a random variable following the geometric distribution are

$$\mu = \frac{1}{p} \quad (4.53)$$

$$\sigma^2 = \frac{1-p}{p^2} \quad (4.54)$$

Definition 4.24: Poisson Process

Experiments yielding numerical values of a random variable X , the number of outcomes occurring during a given time interval or in a specified region, are called Poisson experiments. A Poisson experiment can generate observations for the random variable X representing the number of telephone calls received per hour by an office, the number of days school is closed due to snow during the winter, or the number of games postponed due to rain during a baseball season.

Properties of the Poisson process:

1. The number of outcomes occurring in one time interval or specified region of space is independent of the number that occur in any other disjoint time interval or region. In this sense we say that the Poisson process has no memory.
2. The probability that a single outcome will occur during a very short time interval or in a small region is proportional to the length of the time interval or the size of the region and does not depend on the number of outcomes occurring outside this time interval or region.
3. The probability that more than one outcome will occur in such a short time interval or fall in such a small region is negligible.

The number X of outcomes occurring during a Poisson experiment is called a Poisson random variable, and its probability distribution is called the Poisson distribution.

If you want to learn more on non-homogeneous Poisson process, it allows for a varying rate of events over time or space, and also the average rate (λ) can become a function of time or space.

Definition 4.25: Poisson Distribution

The probability distribution of the Poisson random variable X , representing the number of outcomes occurring in a given time interval or specified region denoted by t , is

$$p(x; \Delta t) = \frac{e^{-\lambda t} (\lambda t)^x}{x!}, \quad x = 0, 1, 2, \dots \quad (4.55)$$

where λ is the average number of outcomes per unit time, distance, area, or volume and $e = 2.71828$.

Use cases of Poisson distribution:

1. Traffic flow: Modeling the number of cars passing through a toll booth in a given time period.
2. Call Centers: Predicting the number of incoming calls during specific hours.
3. Insurance Claims: Estimating the number of insurance claims within a certain timeframe.
4. Web Server Requests: Analyzing the number of requests a server receives in a fixed time interval.
5. Epidemiology: Studying the occurrence of diseases or rare events in a population.

Practical applications of Poisson distribution:

1. Network Security: Analyzing the number of security breaches or attacks on a network within a specific timeframe.
2. Inventory Management: Estimating the number of items sold in a store during a particular hour.
3. Quality Control: Assessing the number of defects in a manufacturing process.
4. Biology and Genetics: Studying the distribution of mutations in a DNA sequence.
5. Finance: Predicting the number of defaults in a loan portfolio.

Theorem 4.7: Mean and Variance of the Poisson Distribution

Both the mean and the variance of the Poisson distribution $p(x; \lambda t)$ are

$$\mu = \sigma^2 = \lambda t \quad (4.56)$$

where t is the specific "time", "distance", "area", or "volume" of interest.

Theorem 4.8: Approximation of Binomial Distribution by a Poisson Distribution

Let X be a binomial random variable with probability distribution $b(x; n, p)$. When $n \rightarrow \infty$, $p \rightarrow 0$, and $np \xrightarrow{n \rightarrow \infty} \mu$ remains constant,

$$b(x; n, p) \xrightarrow{n \rightarrow \infty} p(x; \mu) \quad (4.57)$$

i. Functions in SymIntegration Related to Discrete Distribution

[SI*] To compute the probability mass function, cumulative density function, mean, variance, and moment generating function for the discrete distributions above we can use these functions in SymIntegration:

binomialpmf(x,n,p)

binomialcdf(x,n,p)

binomialmean(x,n,p)

binomialvar(x,n,p)

binomialmgf(x,n,p)

negativebinomialpmf(x,k,p)

negativebinomialmean(x,k,p)

negativebinomialvar(x,k,p)

negativebinomialmgf(x,k,p)

geometricpmf(x,p)

geometricmean(x,p)

geometricvar(x,p)

geometricmgf(x,p)

poissonpmf(x,λt)

poissoncdf(x,λt)

poissonmean(x,λt)

poissonvar(x,λt)

poissonmgf(x,λts)


```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Discrete Distributions ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Discrete Distributions ]# ./main

binomialpmf(2;4,0.75) = 0.210938
binomialcdf(8;15,0.4) = 0.904953
binomialmean(8;15,0.4) = 6
binomialvar(8;15,0.4) = 3.6
binomialmgf(8;15,0.4) = 0.4*e^t+0.6
negativebinomialpmf(6;4,0.55) = 0.1853
negativebinomialmean(6;4,0.55) = 7.27273
negativebinomialvar(6;4,0.55) = 5.95041
negativebinomialmgf(6;4,0.55) = 0.0915063*(-0.45*e^t+1)^(-4)
geometricpmf(5;0.01) = 0.00960596
geometricmean(5;0.01) = 100
geometricvar(5;0.01) = 9900
geometricmgf(5;0.01) = 0.01*(-0.99*e^t+1)^(-1)
poissonpmf(6;4) = 0.104196
poissoncdf(6;4) = 0.889326
poissonmean(6;4) = 4
poissonvar(6;4) = 4
poissonmgf(6;4) = e^(4*e^t-4)

Time taken by function: 56539 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Discrete Distributions ]# 

```

Figure 4.5: The functions in *SymIntegration* to compute the probability mass function, cumulative distribution function, mean, variance, and moment generating function for several discrete probability distributions (*SymIntegration/Examples/Statistics/Test SymIntegration Discrete Distributions/main.cpp*).

[SI*] To generate random numbers that have certain discrete distributions we can use these functions in *SymIntegration*:

vrandn_bernoulli(p, n)

vrandn_binomial(p, n)

with n as the number of random number we want to generate, the result will be in **std::vector<double>**.

[SI*] To generate n unique integer random numbers that have uniform distribution with range $[a, b]$ we can use this function in *SymIntegration*:

vrandn_uniqueinteger(a, b, n)

IV. CONTINUOUS PROBABILITY DISTRIBUTIONS

[SI*] We will show the definitions and theorems of continuous distributions, starting from uniform distribution.

i. Uniform Distribution

Definition 4.26: Uniform Distribution

The density function of the continuous uniform random variable X on the interval $[A, B]$ is

$$f(x; A, B) = \begin{cases} \frac{1}{B-A}, & A \leq x \leq B, \\ 0, & \text{elsewhere} \end{cases} \quad (4.58)$$

the density function forms a rectangle with base $B - A$ and constant height $\frac{1}{B-A}$. The uniform distribution is often called the rectangular distribution. This is the simplest continuous distributions in all of statistics.

Theorem 4.9: The Mean and Variance

The mean and variance of the uniform distribution are

$$\mu = \frac{A + B}{2} \quad (4.59)$$

$$\sigma^2 = \frac{(B - A)^2}{12} \quad (4.60)$$

ii. Normal Distribution

Definition 4.27: Normal Distribution

This is the most important continuous probability distribution in the entire field of statistics.

A continuous random variable X having the bell-shaped distribution is called a normal random variable. The density of the normal random variable X , with mean μ and variance σ^2 , is

$$n(x; \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2\sigma^2}(x-\mu)^2}, \quad -\infty < x < \infty \quad (4.61)$$

where $\pi = 3.14159$ and $e = 2.71828$.

The properties of the normal curve:

1. The mode, which is the point on the horizontal axis where the curve is a maximum, occurs at $x = \mu$.
2. The curve is symmetric about a vertical axis through the mean μ .
3. The curve has its points of inflection at $x = \mu \pm \sigma$; it is concave downward if $\mu - \sigma < X < \mu + \sigma$ and is concave upward otherwise.
4. The normal curve approaches the horizontal axis asymptotically as we proceed in either direction away from the mean.
5. The total area under the curve and above the horizontal axis is equal to 1.

The normal distribution finds enormous application as a limiting distribution. Under certain conditions, the normal distribution provides a good continuous approximation to the binomial and hypergeometric distributions.

Theorem 4.10: The Mean and Variance

The mean and variance of $n(x; \mu, \sigma)$ are μ and σ^2 . Hence, the standard deviation is σ .

The curve of any continuous probability distribution or density function is constructed so that the area under the curve bounded by two ordinates $x = x_1$ and $x = x_2$ equals the probability that the random variable X assumes a value between $x = x_1$ and $x = x_2$. Thus, for the normal curve we will have

$$P(x_1 < X < x_2) = \int_{x_1}^{x_2} n(x; \mu, \sigma) dx = \frac{1}{\sqrt{2\pi}\sigma} \int_{x_1}^{x_2} e^{-\frac{1}{2\sigma^2}(x-\mu)^2} dx \quad (4.62)$$

Definition 4.28: Standard Normal Distribution

We are able to transform all the observations of any normal random variable X into a new set of observations of a normal random variable Z with mean 0 and variance 1. This can be done by means of the transformation

$$Z = \frac{X - \mu}{\sigma} \quad (4.63)$$

Whenever X assumes a value x , the corresponding value of Z is given by

$$z = \frac{x - \mu}{\sigma}$$

Therefore, if X falls between the values $x = x_1$ and $x = x_2$, the random variable Z will fall between the corresponding values

$$z_1 = \frac{x_1 - \mu}{\sigma}$$

and

$$z_2 = \frac{x_2 - \mu}{\sigma}$$

Consequently, we may write

$$\begin{aligned} P(x_1 < X < x_2) &= \frac{1}{\sqrt{2\pi}\sigma} \int_{x_1}^{x_2} e^{-\frac{1}{2\sigma^2}(x-\mu)^2} dx \\ &= \frac{1}{\sqrt{2\pi}(1)} \int_{x_1}^{x_2} e^{-\frac{1}{2(1)^2}(z)^2} dz \\ &= \int_{x_1}^{x_2} n(z; 0, 1) dz \\ &= P(z_1 < Z < z_2) \end{aligned} \quad (4.64)$$

where Z is seen to be a normal random variable with mean 0 and variance 1.

The distribution of a normal random variable with mean 0 and variance 1 is called a standard normal distribution.

Theorem 4.11: Normal Approximation to the Binomial

If X is a binomial random variable with mean $\mu = np$ and variance $\sigma^2 = npq$, then the limiting form of the distribution of

$$Z = \frac{X - np}{\sqrt{npq}}$$

as $n \rightarrow \infty$, is the standard normal distribution $n(z; 0, 1)$.

Definition 4.29: Normal Approximation to the Binomial Distribution

Let X be a binomial random variable with parameters n and p . For large n , X has approximately a normal distribution with $\mu = np$ and $\sigma^2 = npq = np(1 - p)$ and

$$\begin{aligned} P(X \leq x) &= \sum_{k=0}^x b(k; n, p) \\ &= P\left(Z \leq \frac{x + 0.5 - np}{\sqrt{npq}}\right) \end{aligned}$$

as the area under normal curve to the left of $x + 0.5$, and the approximation will be good if np and $np(1 - p)$ are greater than or equal to 5.

[SI*] Computing the cdf of a Normal Distribution

When computing for the probability density function of a normal distribution we can use the Eq (4.53) directly, but the problem is when we have to compute the cumulative distribution function of a normal distribution, it will take an advanced level of integration that has no known indefinite integral, take a look at Eq. (4.53)

$$P(X = x) = n(x; \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2\sigma^2}(x-\mu)^2}, \quad -\infty < x < \infty$$

When we want to compute the cdf we will integrate the pdf just like this

$$\begin{aligned} P(X \leq x) &= \int_{-\infty}^x \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2\sigma^2}(x-\mu)^2} dx \\ &= \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^x e^{-\frac{1}{2\sigma^2}(x-\mu)^2} dx \\ &= \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^x e^{-\frac{1}{2\sigma^2}(x^2 - 2\mu x + \mu^2)} dx \end{aligned}$$

The computation for the integral above is not easy analytically since there is no elementary indefinite integral for

$$\int e^{-x^2} dx$$

Now, we would like to introduce to the Gaussian integral, also known as Euler-Poisson integral, it is the integral of the Gaussian function $f(x) = e^{-x^2}$ over the entire real line. The integral is

$$\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi} \quad (4.65)$$

Carl Friedrich Gauss published the precise integral in 1809. The integral has a wide range of applications, e.g. with a slight change of variables it is used to compute the normalizing constant of the normal distribution. In physics this type of integral appears frequently in quantum mechanics.

The same integral with finite limits is closely related to both the error function and the cumulative distribution function of the normal distribution. So we can approximate the cdf of normal distribution with error function.

The Gaussian integral can be solved analytically through the methods of multivariable calculus. Thus if we have an arbitrary Gaussian function the definite integral is

$$\int_{-\infty}^{\infty} e^{-a(x+b)^2} dx = \sqrt{\frac{\pi}{a}} \quad (4.66)$$

In short, the cdf for a normal distribution can be related to error function, so the cdf formula is

$$\phi(x) = P(X \leq x) = \frac{1}{2} \operatorname{erf} \left(-\frac{x}{\sqrt{2}} \right) \quad (4.67)$$

for $\mu = 0, \sigma = 1$.

$$\phi(x) = P(X \leq x) = \frac{1}{2} \operatorname{erf} \left(-\frac{\frac{x-\mu}{\sigma}}{\sqrt{2}} \right) \quad (4.68)$$

for $\mu \neq 0, \sigma \neq 1$. The error function itself can be computed with Gamma function or Hypergeometric function

$$\operatorname{erf}(x) = 1 - \frac{\Gamma \left(\frac{1}{2}, x^2 \right)}{\sqrt{\pi}} \quad (4.69)$$

$$\operatorname{erf}(x) = \frac{2x}{\sqrt{\pi}} {}_2F_1 \left(-\frac{1}{2}, \frac{3}{2}; -x^2 \right) \quad (4.70)$$

Luckily, the standard library for C++ `<cmath>` already has the function to compute the error function `erfc(x)` that we can use and embed in SymIntegration. Undergraduate students usually only taught to read the cdf for normal distribution from statistics table, the derivative to compute the cdf itself is rarely taught at undergraduate level. This is the background, or the engine behind all that values at the statistics table. How integral plays an important part in statistics, it really shows that statistics actually more symbolic than numeric.

iii. Gamma Distribution

Definition 4.30: Gamma Function

The Gamma distribution derives its name from the well-known gamma function. The Gamma function is defined by

$$\Gamma(\alpha) = \int_0^{\infty} x^{\alpha-1} e^{-x} dx, \quad \text{for } \alpha > 0 \quad (4.71)$$

Simple properties of the gamma function:

- (a) $\Gamma(n) = (n-1)(n-2) \dots (1)\Gamma(1)$, for a positive integer n .
- (b) $\Gamma(n) = (n-1)!$ for a positive integer n .
- (c) $\Gamma(1) = 1$.
- (d) $\Gamma(\frac{1}{2}) = \sqrt{\pi}$

Definition 4.31: Gamma Distribution

The continuous random variable X has a gamma distribution, with parameters α and β , if its density function is given by

$$f(x; \alpha, \beta) = \begin{cases} \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\beta}}, & x > 0 \\ 0, & \text{elsewhere} \end{cases} \quad (4.72)$$

where $\alpha > 0$ and $\beta > 0$. Often we call α as shape and β as scale.

The exponential distribution is a special case of the gamma distribution. The exponential and gamma distributions play an important role in both queuing theory and reliability problems.

The special gamma distribution for which $\alpha = 1$ is called the exponential distribution.

The Lower Incomplete Gamma Function $\gamma(s, x)$

In order to compute the cdf of several continuous distributions, e.g. chi-squared and Gamma, without using external library like **Boost** we need to know this function: the lower incomplete Gamma function.

The lower incomplete Gamma function is defined by the integral:

$$\gamma(s, x) = \int_0^x t^{s-1} e^{-t} dt$$

the definite integral above is quite hard to compute, so we approximate it with the Taylor expansion, the expansion of the exponential function and term by term integration give the following expansion:

$$\gamma(s, x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{n+s}}{n!(n+s)}$$

We can use the series expansion above and create the C++ code to compute the cdf of Gamma distribution or chi-squared distribution. The lower incomplete Gamma function will come in handy for other statistics related computation.

The relations for the Gamma functions:

$$\gamma(s, x) + \Gamma(s, x) = \Gamma(s)$$

with $\Gamma(s, x)$ as the upper incomplete Gamma function.

The Gamma Distribution' cdf with Lower Incomplete Gamma Function $\gamma(s, x)$

The Gamma cdf formula is an integral that represents the probability of a Gamma-distributed

random variable being less than or equal to a specific value, it is defined as

$$\begin{aligned}
 P(X \leq x) &= F(x; \alpha, \beta) \\
 &= \frac{1}{\Gamma(\alpha)} \int_0^x t^{\alpha-1} e^{-\beta t} dt \\
 &= \frac{\gamma\left(\alpha, \frac{x}{\beta}\right)}{\Gamma(\alpha)}
 \end{aligned}$$

With α as the shape parameter and β as the scale parameter. Be careful, some textbooks use rate parameter: $\lambda = \frac{1}{\beta}$ instead of using β . Computer can only use this formula due to the complex integral, the cdf is calculated using numerical methods.

iv. Exponential Distribution

Definition 4.32: Exponential Distribution

The continuous random variable X has an exponential distribution, with parameter β , if its density function is given by

$$f(x; \beta) = \begin{cases} \frac{1}{\beta} e^{-\frac{x}{\beta}}, & x > 0 \\ 0, & \text{elsewhere} \end{cases} \quad (4.73)$$

where $\beta > 0$. Sometimes we also write exponential distribution density function this way

$$f(x; \beta) = \begin{cases} \lambda e^{-\lambda x}, & x > 0 \\ 0, & \text{elsewhere} \end{cases} \quad (4.74)$$

The exponential distribution is more appropriate when the memoryless property is justified. If the failure of the component is a result of gradual or slow wear, then the exponential does not apply and either the gamma or the Weibull distribution may be more appropriate.

Theorem 4.12: The Mean and Variance

The mean and variance of the gamma distribution are

$$\begin{aligned}
 \mu &= \alpha\beta \\
 \sigma^2 &= \alpha\beta^2
 \end{aligned} \quad (4.75)$$

The mean and variance of the exponential distribution are

$$\begin{aligned}
 \mu &= \beta \\
 \sigma^2 &= \beta^2
 \end{aligned} \quad (4.76)$$

v. Beta Distribution

Definition 4.33: Beta Function

A beta function is defined by

$$B(\alpha, \beta) = \int_0^1 x^{\alpha-1} (1-x)^{\beta-1} dx = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, \quad \text{for } \alpha, \beta > 0 \quad (4.77)$$

where $\Gamma(\alpha)$ is the gamma function.

Definition 4.34: Beta Distribution

The continuous random variable X has a beta distribution with parameters $\alpha > 0$ and $\beta > 0$ if its density function is given by

$$f(x) = \begin{cases} \frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1}, & 0 < x < 1 \\ 0, & \text{elsewhere} \end{cases} \quad (4.78)$$

Note that the uniform distribution on $(0, 1)$ is a beta distribution with parameters $\alpha = 1$ and $\beta = 1$. The major reason why we use Beta distribution is because it is defined in the range of $[0, 1]$ so a Beta distribution is a very natural distribution to use when we are talking about probabilities and we want to specify about a prior knowledge of the probabilities of something accruing.

Applications:

- 1. Survival Analysis**

In fields like medicine and engineering, the Beta distribution is used to analyze survival times or failure rates, aiding in understanding product reliability or treatment effectiveness. By modeling these distributions, researchers gain insights into how long products last or how treatments perform over time. This statistical tool helps assess durability and reliability, crucial for making informed decisions in various industries.

- 2. Time Management**

In project management, the Beta distribution helps estimate task durations and overall project completion times. It's like a tool that assists in planning and scheduling. By using this distribution, project managers can better understand the range of possible timeframes for completing tasks or the entire project. This aids in making more informed decisions and managing time effectively.

- 3. Risk Management**

Risk management is essential in finance and insurance. It's about figuring out the chances of different events happening and how they could affect things. By looking at probabilities, it helps make smart choices to reduce risks.

- 4. Genetics**

The beta distribution is a part of the population genetics that not only allows scientists to keep tabs on genes at the population level but as well makes them understand how the genes mutate within the whole group. Allele frequencies, or the different version of genes, must be borne in mind while studying pedigrees and evolution. Utilizing the Beta distribution data, the researchers are able to track the changes in the allele distribution on behalf of the intergenerational population. Such data is informative as far as evolutionary patterns and the nature of ecosystems is known.

- 5. Quality Control**

In manufacturing industries, the Beta distribution serves as a crucial tool for characterizing the variability in product quality. By employing this distribution, businesses gain valuable insights into the distribution of defects or deviations from desired specifications within their production processes. This understanding enables effective control measures to be implemented, ensuring products consistently meet or exceed quality standards. By leveraging the Beta distribution, manufacturers can optimize processes, minimize waste, and enhance overall product reliability, fostering customer satisfaction and trust.

The shape of the distribution is determined entirely by the parameters α and β :

- $\alpha = 1, \beta = 1$: Uniform distribution on $(0, 1)$
- $\alpha > 1, \beta > 1$: Unimodal distribution with a peak within the interval $(0, 1)$. As α and β increase, the peak becomes sharper and more centered.
- $\alpha < 1, \beta < 1$: U-shaped distribution with peaks at the endpoints of the interval.
- $\alpha > 1, \beta < 1$: Right-skewed distribution
- $\alpha < 1, \beta > 1$: Left-skewed distribution

The Beta (α, β) distribution converges to the normal distribution as $\alpha, \beta \rightarrow \infty$. This convergence can be seen by reducing the variance in the graph of the beta distribution. The probability mass then becomes more concentrated around the mean, and the distribution becomes more bell-shaped.

Definition 4.35: Moment Generating Function for Beta Distribution

The moment generating function (mgf) for beta distribution is defined by

$$M_X(\alpha; \beta; t) = 1 + \sum_{i=1}^{\infty} \left(\prod_{j=0}^{i-1} \frac{\alpha + j}{\alpha + \beta + j} \right) \frac{t^i}{i!} \quad (4.79)$$

this function is a series representation of a beta distribution that can be used to compute moments like mean and variance by computing the derivative of the mgf.

The mgf of the beta distribution can be related to the Kummer's confluent hypergeometric function (of the first kind). It is a power series solution to Kummer's differential equation (or confluent hypergeometric equation), an important equation in physics, chemistry, and engineering. It is used to solve problems involving normal and slow rotationally symmetric TE modes in azimuthally magnetized circular ferrite waveguides.

Thus, this is the step by step to obtain the moment generating function for beta distribution

$$\begin{aligned} M_X(\alpha; \beta; t) &= E[e^{tX}] \\ &= \int_0^1 e^{tx} f(x; \alpha, \beta) dx \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 e^{tx} x^{\alpha-1} (1-x)^{\beta-1} dx \\ &= \frac{1}{B(\alpha, \beta)} \int_0^1 e^{tx} x^{\alpha-1} (1-x)^{\beta-1} dx \\ &= \sum_{n=0}^{\infty} \frac{\alpha^{(n)}}{(\alpha + \beta)^n} \frac{t^n}{n!} \\ &= 1 + \sum_{k=1}^{\infty} \left(\prod_{r=0}^{k-1} \frac{\alpha + r}{\alpha + \beta + r} \right) \frac{t^k}{k!} \\ &= {}_1F_1(\alpha; \alpha + \beta; t) \end{aligned}$$

where

$$x^{(n)} = x(x+1)(x+2) \dots (x+n-1)$$

is the rising factorial, also known as the "Pochhammer symbol."

We want to put it simply like this, the mgf of the beta distribution is:

$$M_X(\alpha; \beta; t) = {}_1F_1(\alpha; \alpha + \beta; t)$$

where

$${}_1F_1(\alpha; \beta; t) = \sum_{n=0}^{\infty} \frac{\alpha^{(n)} t^n}{(\beta)^n n!} \quad (4.80)$$

and

$${}_1F_1(\alpha; \alpha + \beta; t) = \sum_{n=0}^{\infty} \frac{\alpha^{(n)} t^n}{(\alpha + \beta)^n n!} \quad (4.81)$$

The good news is the **GNU scientific library** and **Boost** are able to numerically compute the hypergeometric function ${}_1F_1$ directly for a small t ($|t| < 1$), but not for the symbolic computation. So we add this to SymIntegration, thus SymIntegration is able to compute hypergeometric function ${}_1F_1$ numerically and symbolically, with limitation of course, the speed is slower than GSL and Boost since symbolic function will has more cost overall, knowing we can do more after that, to integrate it symbolically or derivate it.

It is handled in **src/specialfunctions.cpp** along with the include in **include/symintegral/specialfunctions.h**

```
Symbolic hypergeometric_1F1(double a, double b, const Symbolic &s, int
    max_iterations) {
    Symbolic sum = 1;

    for (int i = 1; i <= max_iterations; ++i)
    {
        sum += (rising_pochhammer(a, i) * (s^(i))) / (rising_pochhammer
            (b, i) * factorial(i));
    }
    return sum;
}

double hypergeometric_1F1(double a, double b, double s, int max_iterations) {
    Symbolic sum = 1;

    for (int i = 1; i <= max_iterations; ++i)
    {
        sum += (rising_pochhammer(a, i) * (s^(Symbolic(i)))) / (
            rising_pochhammer(b, i) * factorial(i));
    }
    return sum;
}
```

Code 46: *src/specialfunctions.cpp*

We put the **max_iterations** as 5, since it gives the same numeric answer as Boost and GSL.

```
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test/Test Hypergeometric 1F1 GSL vs Boost ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lgsl -lcblas
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test/Test Hypergeometric 1F1 GSL vs Boost ]# ./main
GSL: Hypergeometric 1F1 (2.0, 3.0, 0.5): 1.405115e+00
Boost: MGF of Beta(2, 3) at t=0.5 is: 1.22758

Time taken by function: 273 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test/Test Hypergeometric 1F1 GSL vs Boost ]#
```

Figure 4.6: The computation of the hypergeometric function and the mgf of beta distribution ($\alpha = 2, \beta = 3, t = 0.5$) with GSL and Boost has a very fast speed. (*SymIntegration/Examples/Calculus/Test Hypergeometric 1F1 GSL vs Boost/main.cpp*).

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta ]# ./main

for n = 5

Hypergeometric_1F1(2, 3, z) = 0.666667*z+0.25*z^2+0.0666667*z^3+0.0138889*z^4+0.00238095*z^5+1

Hypergeometric_1F1(2, 3, 0.5) = 1.40511

mgf Beta (2, 3, 0.5) = 1.22758

Time taken by function: 24444 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta ]#
```

Figure 4.7: The computation of the complete hypergeometric function and the mgf of beta distribution ($\alpha = 2, \beta = 3, t = 0.5$) with SymIntegration, it is slower than GSL and Boost but able to use symbolic of "z". (*SymIntegration/Examples/Calculus/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta/main.cpp*).

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta ]# make clean
rm -f main.o main
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta ]# ./main

for n = 5

Hypergeometric_1F1(2, 3, z) = 0.666667*z+0.25*z^2+0.0666667*z^3+0.0138889*z^4+0.00238095*z^5+1

mgf Beta (2, 3, 0.5) = 1.22758

Time taken by function: 19414 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta ]#
```

Figure 4.8: The computation of the symbolic hypergeometric function and the mgf of beta distribution ($\alpha = 2, \beta = 3, t = 0.5$) with SymIntegration. (*SymIntegration/Examples/Calculus/Test SymIntegration Compute Hypergeometric Function 1F1 and mgf Beta/main.cpp*).

Definition 4.36: Survival Function

The survival function, also known as the complementary cumulative distribution function (ccdf), is the probability that X takes a value greater than x , or

$$P(X > x)$$

it is defined as

$$S(x; \alpha, \beta) = 1 - F(x; \alpha, \beta) = 1 - \frac{B(x; \alpha, \beta)}{B(\alpha, \beta)} \quad (4.82)$$

In R, the survival function can be calculated using the **pbeta** function with the **lower.tail = FALSE** argument.

Real life applications of Beta distribution:

- 1. Crop Yield Analysis**

Farmers and researchers use the Beta distribution to understand how crop yields vary across different areas of land. This helps them decide where to plant crops, how to allocate resources, and manage risks.

- 2. Sport Analytics**

In sports, the Beta distribution helps analyze player performance metrics like shooting accuracy or goal-scoring rates. Coaches and analysts use this information to make strategic decisions.

- 3. Environmental Risk Assessment**

Environmental scientists use the Beta distribution to model uncertainty in environmental factors like pollutant concentrations. This helps assess potential risks and guide policy decisions.

- 4. Clinical Trials**

In medical research, the Beta distribution models treatment outcomes or response rates in clinical trials. This helps researchers design trials, determine sample sizes, and accurately analyze results.

- 5. Weather Forecasting**

Meteorologists use the Beta distribution to model the distribution of weather phenomena such as rainfall or wind speeds. This helps predict weather patterns and assess the likelihood of extreme events.

Theorem 4.13: Mean and Variance

The mean and variance of a beta distribution with parameters α and β are

$$\mu = \frac{\alpha}{\alpha + \beta} \quad (4.83)$$

$$\sigma^2 = \frac{\alpha\beta}{(\alpha + \beta)^2(\alpha + \beta + 1)} \quad (4.84)$$

The first two theoretical moments for Beta distribution are the mean (μ) and variance (σ^2). When we are doing observation or experiment we obtain samples and we can compute the sample mean and variance with

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n}$$

$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}$$

Now by equating $\bar{x} = \mu$ and $s^2 = \sigma^2$ we will obtain two equations with two unknowns (α, β). Solving these equations simultaneously gives the estimates for α and β . This is called the method of moments, the drawback is that it can be inefficient and sensitive to outliers. Other methods, such as maximum likelihood estimation, often provide more accurate estimates.

The Beta Distribution' cdf with Regularized Incomplete Beta Function $I_x(\alpha, \beta)$

The beta cdf can be expressed using the regularized incomplete beta function, which is defined as the incomplete beta function ($B(x; \alpha, \beta)$) divided by the complete beta function ($B(\alpha, \beta)$). This can be written as:

$$F_X(x) = \frac{B(x; \alpha, \beta)}{B(\alpha, \beta)} = I_x(\alpha, \beta)$$

where $I_x(\alpha, \beta)$ is the regularized incomplete beta function.

The incomplete beta function is defined as an integral:

$$B(x; \alpha, \beta) = \int_0^x u^{\alpha-1} (1-u)^{\beta-1} du$$

And the complete beta function can be expressed using the Gamma function:

$$B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta)}$$

The incomplete beta function and its regularized form are crucial in statistics, particularly in calculating probabilities related to beta distribution and in determining cumulative distribution functions (cdf) for distributions like the binomial, F -distribution, and Student's t -distribution.

Characteristics of the cdf and sf:

1. Both cdf and sf are increasing functions on the interval $(0, 1)$.
2. The cdf approaches 0 as x approaches 0 and approaches 1 as x approaches 1.
3. The cdf approaches 1 as x approaches 0 and approaches 0 as x approaches 1.
4. The sum of the cdf and sf at any point x is equal to 1.
5. The shapes of the cdf and sf depend on the values of α and β . For example, if $\alpha > 1$ and $\beta > 1$, the cdf is the S-shaped and the sf is concave down.

vi. Chi-Squared Distribution

Definition 4.37: Chi-Squared Distribution

The chi-squared distribution with ν degrees of freedom is the distribution of a sum of the squares of k independent standard normal random variables.

The chi-squared distribution χ_ν^2 is a special case of the gamma distribution and the univariate Wishart distribution. Specifically if $X \sim \chi_\nu^2$ then $X \sim \text{Gamma}(\alpha = \frac{\nu}{2}, \beta = 2)$ (where α is the shape parameter and β is the scale parameter of the gamma distribution) and $X \sim W_1(1, k)$.

The probability density function of the chi-squared distribution is

$$f(x; \nu) = \frac{1}{\Gamma(\nu/2)2^{\nu/2}} x^{\nu/2-1} e^{-x/2}, \quad x > 0 \quad (4.85)$$

The cumulative distribution function is

$$F(x; \nu) = \frac{\gamma(\frac{\nu}{2}, \frac{x}{2})}{\Gamma(\frac{\nu}{2})} = P\left(\frac{\nu}{2}, \frac{x}{2}\right) \quad (4.86)$$

where $\gamma(s, t)$ is the lower incomplete gamma function and $P(s, t)$ is the regularized gamma function.

The chi-squared distribution is one of the most widely used probability distributions in inferential statistics, notably in hypothesis testing and in construction of confidence intervals.

The lower incomplete gamma function is given by

$$\begin{aligned} \gamma(a, x) &= \int_0^x t^{a-1} e^{-t} dt \\ &= a^{-1} x^a e^{-x} {}_1F_1(1; 1+a; x) \\ &= a^{-1} x^a {}_1F_1(a; 1+a; -x) \end{aligned}$$

where ${}_1F_1(a; b; x)$ is the confluent hypergeometric function of the first kind.

By definition, the lower and upper incomplete gamma functions satisfy

$$\Gamma(a, x) + \gamma(a, x) = \Gamma(a) \quad (4.87)$$

Theorem 4.14: The Mean and Variance

The mean and variance of the chi-squared distribution are

$$\mu = \nu \quad (4.88)$$

$$\sigma^2 = 2\nu \quad (4.89)$$

The Chi-Squared Distribution' cdf with Lower Incomplete Gamma Function $\gamma(s, x)$

The chi-squared cdf is expressed using the incomplete Gamma function as

$$\begin{aligned} P(X \leq x) &= F(x; \nu) \\ &= \frac{1}{\Gamma(\frac{\nu}{2})} \gamma\left(\frac{\nu}{2}, \frac{x}{2}\right) \end{aligned}$$

with ν as the degrees of freedom, Γ is the Gamma function, and γ is the lower incomplete Gamma function.

We can also write the cdf in a series expansion for computation:

$$F(x; \nu) = \left(\frac{x}{2}\right)^{\left(\frac{\nu}{2}\right)} e^{-\frac{x}{2}} \sum_{m=0}^{\infty} \frac{\left(\frac{x}{2}\right)^m}{\Gamma\left(\frac{\nu}{2} + m + 1\right)}$$

vii. t -Distribution

In many experimental scenarios, knowledge of σ is certainly no more reasonable than knowledge of the population mean μ . Often, in fact, an estimate of σ must be supplied by the same sample information that produced the sample average $\bar{x}$. As a result, a natural statistic to consider to deal with inferences on μ is

$$T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \quad (4.90)$$

since S is the sample analog to σ . If the sample size is small, the values of S^2 fluctuate considerably from sample to sample and the distribution T deviates appreciably from that of a standard normal distribution.

If the sample size is large enough, say $n \geq 30$, the distribution of T does not differ considerably from the standard normal. However, for $n < 30$, it is useful to deal with the exact distribution of T . In developing the sampling distribution of T , we shall assume that our random sample was selected from a normal population. We can then write

$$T = \frac{(\bar{X} - \mu)/(\sigma/\sqrt{n})}{\sqrt{S^2/\sigma^2}} = \frac{Z}{\sqrt{V/(n-1)}} \quad (4.91)$$

where

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

has the standard normal distribution and

$$V = \frac{(n-1)S^2}{\sigma^2} \quad (4.92)$$

has a chi-squared distribution with $\nu = n - 1$ degrees of freedom. In sampling from normal populations, we can show that $\bar{X}$ and S^2 are independent, and consequently so are Z and V .

Theorem 4.15: t -Distribution

Let Z be a standard normal random variable and V a chi-squared random variable with ν degrees of freedom. If Z and V are independent, then the distribution of the random variable T , where

$$T = \frac{Z}{\sqrt{\frac{V}{\nu}}} \quad (4.93)$$

with $\nu = n - 1$, is given by the density function

$$h(t) = \frac{\Gamma\left[\frac{\nu+1}{2}\right]}{\Gamma\left(\frac{\nu}{2}\right)\sqrt{\pi\nu}} \left(1 + \frac{t^2}{\nu}\right)^{-\frac{\nu+1}{2}}, \quad -\infty < t < \infty \quad (4.94)$$

This is known as the t -distribution with ν degrees of freedom.

The t -distribution represents the structure that occurs if all of the values of $\frac{\bar{x}-\mu}{s/\sqrt{n}}$ are formed, where $\bar{x}$ and s are taken from samples of size n from a $n(x; \mu, \sigma)$ distribution.

The Student's t -Distribution' cdf with Regularized Incomplete Beta Function $I_x(\alpha, \beta)$

To compute the cdf of t -distribution we will use this formula:

$$F_\nu(t) = \frac{1}{2} I_x\left(\frac{\nu}{2}, \frac{1}{2}\right)$$

for $t \leq 0$.

$$F_\nu(t) = 1 - \frac{1}{2} I_x\left(\frac{\nu}{2}, \frac{1}{2}\right)$$

for $t \geq 0$, where

$$x = \frac{\nu}{\nu + t^2}$$

and $\nu = n - 1$ is the degrees of freedom.

viii. F -Distribution

In probability theory and statistics, the F -distribution or F -ratio, also known as Snedecor's F distribution or the Fisher Snedecor distribution, is a continuous probability distribution that arises frequently as the null distribution of a test statistic, most notably in the analysis of variance (ANOVA) and other F -tests.

As you can see, the F -distribution finds enormous application in comparing sample variances. Applications of the F -distribution are found in problems involving two or more samples.

The statistic F is defined to be the ratio of two independent chi-squared random variables, each divided by its number of degrees of freedom. Hence, we can write

$$F = \frac{U/\nu_1}{V/\nu_2} \quad (4.95)$$

where U and V are independent random variables having chi-squared distributions with ν_1 and ν_2 degrees of freedom.

Theorem 4.16: F-Distribution

Let U and V be two independent random variables having chi-squared distributions with ν_1 and ν_2 degrees of freedom, respectively. Then the distribution of the random variable $F = \frac{U/\nu_1}{V/\nu_2}$ is given by the density function

$$h(f) = \begin{cases} \frac{\Gamma\left(\frac{\nu_1 + \nu_2}{2}\right) \left(\frac{\nu_1}{\nu_2}\right)^{\frac{\nu_1}{2}}}{\Gamma\left(\frac{\nu_1}{2}\right) \Gamma\left(\frac{\nu_2}{2}\right)} \frac{f^{\frac{\nu_1}{2} - 1}}{\left(1 + \frac{\nu_1 f}{\nu_2}\right)^{\frac{\nu_1 + \nu_2}{2}}}, & f > 0 \\ 0, & f \leq 0 \end{cases} \quad (4.96)$$

This is known as the F -distribution with ν_1 and ν_2 degrees of freedom. The curve of the F -distribution depends not only on the two parameters ν_1 and ν_2 but also on the order in which we state them.

The statistics table provides the f -value that you are looking for given the various combinations of the degrees of freedom ν_1 and ν_2 . Hence the f -value with 6 and 10 degrees of freedom ν_1 and ν_2 , leaving an area of 0.05 to the right, is

$$f_{0.05} = 3.22$$

Theorem 4.17: The F-Distribution with Degrees of Freedom

Writing $f_\alpha(\nu_1, \nu_2)$ for f_α with ν_1 and ν_2 degrees of freedom, we obtain

$$f_{1-\alpha}(\nu_1, \nu_2) = \frac{1}{f_\alpha(\nu_2, \nu_1)} \quad (4.97)$$

The F-Distribution' pdf with Beta Function

The pdf of the F -distribution for a variable x with numerator degrees of freedom ν_1 and denominator degrees of freedom ν_2 is defined as:

$$f(x; \nu_1, \nu_2) = \frac{1}{B\left(\frac{\nu_1}{2}, \frac{\nu_2}{2}\right)} \left(\frac{\nu_1}{\nu_2}\right)^{\frac{\nu_1}{2}} x^{\frac{\nu_1}{2} - 1} \left(1 + \frac{\nu_1}{\nu_2} x\right)^{-\frac{\nu_1 + \nu_2}{2}}$$

where $B(a, b)$ is the beta function, which can be defined in terms of the gamma function:

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$$

the gamma function $\Gamma()$ can be computed easily and available in standard C++ library as **tgamma()**.

The pdf of F -distribution is always greater than or equal to zero for all valid x . The integral of the pdf over its entire domain $(0, \infty)$ is equal to 1.

The F -Distribution' cdf with Incomplete Beta Function

The regularized incomplete beta function, denoted $I_x(a, b)$, is defined as:

$$I_x(a, b) = \frac{B_x(a, b)}{B(a, b)} = \frac{\int_0^x t^{a-1}(1-t)^{b-1} dt}{\int_0^1 t^{a-1}(1-t)^{b-1} dt}$$

where $B_x(a, b)$ is the incomplete beta function, and $B(a, b)$ is the complete beta function.

The cdf of an F -distribution with ν_1 and ν_2 degrees of freedom is given by the following formula:

$$F(x; \nu_1, \nu_2) = P(X \leq x) = I_z\left(\frac{\nu_1}{2}, \frac{\nu_2}{2}\right)$$

where the variable z is defined as

$$z = \frac{\nu_1 x}{\nu_1 x + \nu_2}$$

Using the identity $I_z(a, b) = 1 - I_{1-z}\left(\frac{\nu_2}{2}, \frac{\nu_1}{2}\right)$

$$F(x; \nu_1, \nu_2) = 1 - I_{1-z}\left(\frac{\nu_2}{2}, \frac{\nu_1}{2}\right)$$

where $1 - z$ is:

$$1 - z = 1 - \frac{\nu_1 x}{\nu_1 x + \nu_2} = \frac{\nu_2}{\nu_1 x + \nu_2}$$

The F -Distribution Connection to the F -Statistic

The F -distribution is often derived as the ratio of two chi-squared distributions, which is the basis for the F -test. The F -statistic is calculated as

$$F = \frac{\text{Variance}_{\text{between}}}{\text{Variance}_{\text{within}}}$$

or

$$F = \frac{\text{Mean Square}_{\text{between}}}{\text{Mean Square}_{\text{within}}}$$

In the context of ANOVA, this can be written as:

$$F = \frac{SS_{\text{between}}/df_{\text{between}}}{SS_{\text{within}}/df_{\text{within}}}$$

The F -Distribution with Two Sample Variances

Suppose that random samples of size n_1 and n_2 are selected from two normal populations with variances σ_1^2 and σ_2^2 , respectively. We know that

$$\chi_1^2 = \frac{(n_1 - 1)S_1^2}{\sigma_1^2}$$

$$\chi_2^2 = \frac{(n_2 - 1)S_2^2}{\sigma_2^2}$$

are random variables having chi-squared distributions with $\nu_1 = n_1 - 1$ and $\nu_2 = n_2 - 1$ degrees of freedom. Furthermore, since the samples are selected at random, we are dealing with independent random variables. Furthermore, since the samples are selected at random, we are dealing with independent random variables.

Theorem 4.18: F-Distribution with Two Sample Variances

If S_1^2 and S_2^2 are the variances of independent random samples of size n_1 and n_2 taken from normal populations with variances σ_1^2 and σ_2^2 , respectively, then

$$F = \frac{S_1^2/\sigma_1^2}{S_2^2/\sigma_2^2} = \frac{\sigma_2^2 S_1^2}{\sigma_1^2 S_2^2} \quad (4.98)$$

has an F -distribution with $\nu_1 = n_1 - 1$ and $\nu_2 = n_2 - 1$ degrees of freedom.

The F -distribution is used in two-sample situations to draw inferences about the population variances. However, the F -distribution can also be applied to many other types of problems involving sample variances. In fact, the F -distribution is called the variance ratio distribution. The F -distribution plays an important role in the analysis of variance.

Three things one must bear in mind, regarding these fundamental sampling distributions:

1. One cannot use Central Limit Theorem unless σ is known. When σ is not known, it should be replaced by s , the sample standard deviation, in order to use the Central Limit Theorem.
2. The T statistic is not a result of the Central Limit Theorem and $x_1, x_2, \dots, x_n$ must come from a $n(x; \mu, \sigma)$ distribution in order for $\frac{\bar{x} - \mu}{s/\sqrt{n}}$ to be a t -distribution; s is, of course, merely an estimate of σ .
3. While the notion of degrees of freedom is new at this point, the concept should be very intuitive, since it is reasonable that the nature of the distribution of S and also t should depend on the amount of information in the sample $x_1, x_2, \dots, x_n$.

ix. Weibull Distribution and Hazard Rates

For a random variable T , representing the lifetime of an individual or a component, an interesting quantity is the instantaneous chance of failure at any time, given that the component has been operating without failure up to time x . This can be expressed as

$$h(x) = \lim_{\delta \rightarrow 0} \frac{1}{\delta} P(T \in [x, x + \delta) | T > x)$$

Alternatively, by using the conditional probability and noticing that the pdf $f(x)$ satisfies $f(x)\delta \approx P(x \leq T < x + \delta)$ for small δ , we can express the above as

$$h(x) = \frac{f(x)}{1 - F(x)} \quad (4.99)$$

Here the function $h(\cdot)$ is called the hazard rate, and it is a common method of viewing the distribution for lifetime random variables T . In fact, we can reconstruct the cdf $F(x)$ by

$$1 - F(x) = e^{-\int_0^x h(t) dt} \quad (4.100)$$

Hence, every continuous non-negative random variable can be described uniquely by its hazard rate. The Weibull distribution is naturally defined through the hazard rate by considering hazard rate functions that have a specific simple form. It is a distribution with

$$h(x) = \lambda x^{\alpha-1} \quad (4.101)$$

where λ is positive and α takes on any real value. Notice that the parameter α gives the Weibull distribution different modes of behavior. If $\alpha = 1$ then the hazard rate is constant, in which case the Weibull distribution is actually an exponential distribution with rate λ . If $\alpha > 1$, then the hazard rate increases over time. This depicts a situation of "aging components", i.e. the longer a component has lived, the higher the instantaneous chance of failure. This is sometimes called Increasing Failure Rate (IFR). Conversely, $\alpha < 1$ depicts a situation where the longer a component has lasted, the lower the chance of it failing (as it perhaps the case with totalitarian political regimes). This is sometimes called Decreasing Failure Rate (DFR).

Based on Eq. (4.108) and using Eq. (4.107), we obtain the cdf and pdf

$$\begin{aligned} F(x) &= 1 - e^{-\frac{\lambda}{\alpha} x^\alpha} \\ f(x) &= \lambda x^{\alpha-1} e^{-\frac{\lambda}{\alpha} x^\alpha} \end{aligned} \quad (4.102)$$

where the bijection from λ to θ is

$$\begin{aligned} \lambda &= \alpha \theta^{-\alpha} \\ \theta &= \left(\frac{\alpha}{\lambda} \right)^{\frac{1}{\alpha}} \end{aligned} \quad (4.103)$$

In this case, θ is called the scale parameter and α is the shape parameter.

x. Cauchy Distribution

Also known as the Loretz distribution, this is a distribution without mean.

Definition 4.38: Cauchy Distribution

At first glance, a plot of the pdf looks very similar to the normal distribution. However, it is fundamentally different as its mean and standard deviation are undefined. The pdf of the Cauchy distribution is given by

$$f(x) = \frac{1}{\pi \gamma \left(1 + \left(\frac{x - x_0}{\gamma} \right)^2 \right)} \quad (4.104)$$

where x_0 is the location parameter at which the peak is observed and γ is the scale parameter.

The physical example of Cauchy distributed random variable is like this: consider a drone hovering stationary in the sky at unit height. A pivoting laser is attached to its undercarriage, which pivots back and forth as it shoots pulses at the ground. At any point the laser fires, it makes an angle θ from the vertical $(-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2})$.

Since the laser fires at a high frequency as it is pivoting, we can assume that the angle θ is distributed uniformly on $[-\frac{\pi}{2}, \frac{\pi}{2}]$. For each shot from the laser, a point can be measured, X , horizontally on the ground from the point above which the drone is hovering. We can now

consider the horizontal measurement as a new random variable, X . Hence the cdf is

$$\begin{aligned} F_X(x) &= P(\tan(\theta) \leq x) = P(\theta \leq \text{atan}(x)) \\ &= F_\theta(\text{atan}(x)) \\ &= \begin{cases} 0, & \text{atan}(x) \leq -\frac{\pi}{2} \\ \frac{1}{\pi}\text{atan}(x), & \text{atan}(x) \in (-\frac{\pi}{2}, \frac{\pi}{2}) \\ 1, & \text{atan}(x) \geq \frac{\pi}{2} \end{cases} \end{aligned} \quad (4.105)$$

Now since it always holds that $\text{atan}(x) \in (-\frac{\pi}{2}, \frac{\pi}{2})$ we can obtain the density by taking the derivative of $\frac{1}{\pi}\text{atan}(x)$ which evaluates to

$$f(x) = \frac{1}{\pi(1+x^2)}$$

This is a special case of the more complicated density Eq. (4.111), with x_0 and $\gamma = 1$. Importantly, the expectation integral,

$$\int_{-\infty}^{\infty} xf(x) dx$$

is not defined since each of the one-sided improper integrals does not converge.

xi. Bivariate Normal

One of the most ubiquitous families of multivariate distributions is the multivariate normal distribution. Similarly to the fact that a scalar (univariate) normal distribution is parametrized by the mean μ and the variance σ^2 , a multivariate normal distribution is parametrized by the mean vector μ_X and the covariance matrix Σ_X .

Definition 4.39: Standard Multivariate Normal Distribution

For the standard multivariate normal distribution having $\mu_X = \mathbf{0}$ mean and $\Sigma_X = I$, the pdf for the random vector $\mathbf{X} = (X_1, \dots, X_n)$ is

$$f(\mathbf{x}) = (2\pi)^{-\frac{n}{2}} e^{-\frac{1}{2}\mathbf{x}^T \mathbf{x}} \quad (4.106)$$

Now, in general, using an affine transformation, it can be shown that for arbitrary μ_X and Σ_X (positive definite)

$$f(\mathbf{x}) = |\Sigma_X|^{-\frac{1}{2}} (2\pi)^{-\frac{n}{2}} e^{-\frac{1}{2}(\mathbf{x}-\mu_X)^T \Sigma_X^{-1}(\mathbf{x}-\mu_X)} \quad (4.107)$$

where $|\Sigma_X|$ is the determinant of Σ_X .

Definition 4.40: Bivariate Normal Distribution

From Eq. (4.114) In the case of $n = 2$, this becomes the bivariate normal distribution with a density represented as

$$f_{XY}(x, y; \sigma_X, \sigma_Y, \mu_X, \mu_Y, \rho) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \times \exp \left[-\frac{1}{2(1-\rho^2)} \left(\frac{(x-\mu_X)^2}{\sigma_X^2} - \frac{2\rho(x-\mu_X)(y-\mu_Y)}{\sigma_X\sigma_Y} + \frac{(y-\mu_Y)^2}{\sigma_Y^2} \right) \right] \quad (4.108)$$

Here the elements of the mean and covariance matrix are

$$\mu_X = \begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}$$

$$\Sigma_Y = \begin{bmatrix} \sigma_X^2 & \sigma_X\sigma_Y\rho \\ \sigma_X\sigma_Y\rho & \sigma_Y^2 \end{bmatrix}$$

Note that $\rho \in (-1, 1)$ is the correlation coefficient.

xii. Functions in SymIntegration Related to Continuous Distribution

[SI*] To compute the probability density function, cumulative density function, mean, variance, and moment generating function for the continuous distributions above we can use these functions in SymIntegration:

uniformpdf(x,a,b)
uniformcdf(x,a,b)
uniformmgf(x,a,b)
uniformmean(x,a,b)
uniformvar(x,a,b)

normalpdf(x,μ,σ)
normalcdf(x,μ,σ)
normalmgf(x,μ,σ)
normalmean(x,μ,σ)
normalvar(x,μ,σ)

gammapdf(x,α, β)
gammacdf(x,α, β)
gammamgf(x,α, β)
gammamean(x,α, β)
gammavar(x,α, β)

exponentialpdf(x,λ)
exponentialcdf(x,λ)

exponentialmgf(x, λ)
exponentialmean(x, λ)
exponentialvar(x, λ)

betapdf(x, α, β)
betacdf(x, α, β)
betamgf(x, α, β)
betamean(x, α, β)
betavar(x, α, β)

```
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Continuous Distributions ]# ./main

uniformpdf(1;1,5) = 0.25
uniformcdf(3;1,5) = 0.5
uniformmgf(x;1,5) = 0.25*e^(5*t)*t^(-1)-0.25*e^t*t^(-1)
uniformmean(x;1,5) = 3
uniformvar(x;1,5) = 1.33333
normalpdf(362;300,50) = 0.00369875
normalcdf(362;300,50) = 0.892512
normalmgf(362;300,50) = e^(300*t+1250*t^(2))
normalmean(362;300,50) = 300
normalvar(362;300,50) = 2500
gammampdf(1,2,0.2) = 0.168449
gammacdf(1,2,0.2) = 0.959572
gammamgf(1,2,0.2) = (-0.2*t+1)^(-2)
gammamean(1,2,0.2) = 0.4
gammavar(1,2,0.2) = 0.08
exponentialpdf(8,0.2) = 0.0403793
exponentialcdf(8,0.2) = 0.798103
exponentialmgf(8,0.2) = (-5*t+1)^(-1)
exponentialmean(8,0.2) = 5
exponentialvar(8,0.2) = 25
betapdf(0.8,3,2) = 1.536
betacdf(0.8,3,2) = 0.8192
betamgf(0.8,3,2) = 0.6*t+0.2*t^(2)+0.047619*t^(3)+0.00892857*t^(4)+0.00138889*t^(5)+1
betamean(0.8,3,2) = 0.6
betavar(0.8,3,2) = 0.04
Time taken by function: 321002 microseconds
```

Figure 4.9: The functions in SymIntegration to compute the probability density function, cumulative distribution function, mean, variance, and moment generating function for several continuous probability distributions (SymIntegration/Examples/Statistics/Test SymIntegration Continuous Distributions/main.cpp).

[SI*] To generate random numbers that have certain continuous distributions we can use these

functions in SymIntegration:

```

vrandn_normal( $\mu$ ,  $\sigma$ ,  $n$ )
vrandn_exponential( $\lambda$ ,  $n$ )
vrandn_chisquared( $\nu$ ,  $n$ )
vrandn_fdist( $\nu_1$ ,  $\nu_2$ ,  $n$ )
vrandn_tdist( $\nu$ ,  $n$ )
vrandn_erlang( $k$ ,  $\lambda$ ,  $n$ )
vrandn_gamma( $\alpha$ ,  $\beta$ ,  $n$ )
vrandn_beta( $\alpha$ ,  $\beta$ ,  $n$ )

```

with n as the number of random number we want to generate, the result will be in `std::vector<double>`.

[SI*] To compute the quantile of certain distributions we can use these functions in SymIntegration:

```
zquantile( $p$ )
```

(p is the desired probability, we are using Wichura's algorithm, a robust algorithm for accurate computation of z quantile, especially for extreme quantiles. This numerical approximation algorithm is very useful to compute the inverse cdf / quantile value of the normal distribution. Given a probability p , the inverse cdf returns the value x as the z quantile such that probability of a random variable being less than or equal to x is p , $P(X \leq x) = p$. We take it from John Burkardt's CPP consortium codes, named ASA 241, then blend it into SymIntegration.)

```
tquantile( $x_0$ ,  $\nu$ ,  $p$ )
```

(x_0 is the initial value for the approximation, you can input any value you want, ν is the degrees of freedom, p is the desired probability, we are using iterative method / Newton-Raphson method to find the root of $F(x) - p = 0$, where $F(x)$ is the cdf of the Student's t -distribution)

svec to replace `vector<string>`

We create a typedef defined in `include/symintegral/symintegrationc++.h` to represents `vector<string>`, so we don't need to write `vector<string>` anymore, we can use **svec** instead. This is very useful for data analysis.

Some explanations:

1. We used to call **Boost** to compute the pdf and cdf of some continuous distributions, but now we are releasing dependencies on **Boost** to make the program runs faster. For example, we used to use this code to compute the pdf of Beta distribution:

```

#include <boost/math/distributions/beta.hpp> //Faster computing of
      beta cdf and pdf

...

Symbolic betapdf(double x, double alpha, double beta)
{
    boost::math::beta_distribution<> my_beta(alpha,beta);
    double pdf_value = boost::math::pdf(my_beta,x);
}

```

```
}  
  
...
```

Code 47: *src/statistics.cpp*

Now we are using this code to compute the pdf of Beta distribution:

```
...  
  
double betapdf(double x, double alpha, double beta)  
{  
    if (x < 0.0 || x > 1.0 || alpha <= 0.0 || beta <= 0.0)  
    {  
        // Handle invalid input: PDF is 0 outside [0,1] or parameters are  
        // invalid  
        return 0.0;  
    }  
    else  
    {  
        // Calculate log of the Beta function using log-Gamma functions  
        double log_beta_function = std::lgamma(alpha) + std::lgamma(beta)  
            - std::lgamma(alpha + beta);  
  
        // Calculate log of the numerator  
        double log_numerator = (alpha - 1.0) * std::log(x) + (beta -  
            1.0) * std::log(1.0 - x);  
  
        // Calculate the log of the PDF  
        double log_pdf = log_numerator - log_beta_function;  
  
        // Return the exponentiated value  
        return std::exp(log_pdf);  
    }  
}  
  
...
```

Code 48: *src/statistics.cpp*

we can compute the pdf of Beta distribution by using the **lgamma** function from the **<cmath>** header to compute the logarithm of the Gamma function, which is more numerically stable for large values.

2. To generate random number of certain distribution, we make use of **std** library that already have a lot of continuous and discrete distributions covered.

First thing needed is to create / obtain a non-deterministic seed for the random number engine. Two good options currently are:

std::random_device

(create a non-deterministic seed if available and entropy > 0)

or

std::chrono::system_clock::now().time_since_epoch().count()

(provides a more robust seed than a fixed value)

Then we will use **std::mt19937** (Mersenne Twister engine) to generate the random number.

Here is an example of a function in **src/statistics.cpp** to generate a random number that has normal distribution:

```
...

std::vector<double> vrandn_normal(double mu, double sigma, int n)
{
    std::chrono::system_clock::now().time_since_epoch().count()

    std::default_random_engine generator(
        std::chrono::system_clock::now().time_since_epoch().count());

    std::vector<double> vec;
    std::normal_distribution<double> distribution(mu, sigma);
    for(int i=1; i<n; i++)
    {
        vec.push_back(static_cast<double>(distribution(generator)))
    }
    return vec;
}

...
```

Code 49: *src/statistics.cpp*

3. The C++ Standard Library does not directly offer a **std::beta_distribution**. However, a Beta distribution can be simulated using two independently generated Gamma distributions.

The relationship is as follows:

If X is a random variable from a Gamma distribution with shape parameter α and scale parameter 1 ($\text{Gamma}(\alpha, 1)$), and Y is a random variable from a Gamma distribution with shape parameter β and scale parameter 1 ($\text{Gamma}(\beta, 1)$), then the random variable

$$Z = \frac{X}{X + Y}$$

follows a Beta distribution with shape parameters α and β (denoted $\text{Beta}(\alpha, \beta)$).

xiii. Generate Gamma Distributed Random Number in SymIntegration and Plot the Histogram with Hamzstplot

We will demonstrate how to generate random number with Gamma distribution then plot the data as histogram with Hamzstplot.

We will create a C++ source file **main.cpp**, and a Makefile (or compile it manually).

```
#include <iostream>
#include "symintegrationc++.h"
#include <bits/stdc++.h>
#include <cmath>

#include <chrono>

using namespace std::chrono;
using namespace std;
using namespace SymbolicConstant;

int main(void)
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    std::vector<double> x = vrandn_gamma(2,1.5,50);
    for(int i=0;i<50;i++)
    {
        cout<< x[i] << endl;
    }

    // Get ending timepoint
    auto stop = high_resolution_clock::now();
    auto duration = duration_cast<microseconds>(stop - start);

    cout << "\nTime taken by function: " << duration.count() << " microseconds"
        << endl;

    return 0;
}
```

Code 50: *main.cpp*

```
CFLAGS = -ggdb
DEFINES = -DDEBUGGA
INCLUDES =
LIBS = -lstdc++ -lsymintegration
MAIN = main.o
```

```
CC=g++

.cc.o:
$(CC) -c $(CFLAGS) $(DEFINES) $(INCLUDES) $<

all:: main

gnuplot_i.o:
main.o: main.cpp

main: $(MAIN)
$(CC) -o $@ $(CFLAGS) $(MAIN) $(LIBS)

clean:
rm -f $(MAIN) main
```

Code 51: *Makefile*

From the current working directory containing the **main.cpp** and the makefile, type:

```
make
./main
```

to save the output you can also type this:

```
./main > gamma.dat
```

If you are still learning about Makefile, or feel more confident with manual compiling, then instead of **make** you can type this:

```
g++ -o main main.cpp -lsymintegration
./main
```

```
g++ -o main -gddb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Generate Random Number from Gamma Distribution ]# ./main
1.49028
4.10471
1.345
0.465866
1.34767
2.26934
1.07946
4.17375
1.48454
3.77183
2.87478
0.318233
3.1834
1.90781
3.69633
1.51896
4.32441
2.56319
6.54034
2.51703
3.25444
4.40127
1.68972
0.734017
3.02267
6.22162
4.24151
1.61816
5.48238
3.12266
1.6428
5.69689
1.54791
3.17954
5.72097
5.6629
4.99465
2.66944
6.83603
2.46991
1.13762
2.35658
0.503396
3.15512
5.12802
4.61204
4.52307
5.98776
4.05892
0
Time taken by function: 1008 microseconds
```

Figure 4.10: The random number generation that has Gamma distribution with parameters $\alpha = 2, \beta = 1.5$, we generate 50 random numbers. (SymIntegration/Examples/Statistics/Test SymIntegration Generate Random Number from Gamma Distribution/main.cpp).




Figure 4.11: The random number generation that has been saved in `gamma.dat`. (*SymIntegration/Examples/Statistics/Test SymIntegration Generate Random Number from Gamma Distribution/main.cpp*).

Now, if we want to plot the histogram, we will need to add **Hamzstplot** library when compiling and add few more lines of code.

We will create again a C++ source file **main.cpp**, and a Makefile (or compile it manually).

```
#include <cmath>
#include <hamzstplot/hamzstplot.h>
#include <random>
#include "symintegrationc++.h"

using namespace std;

int main() {
    using namespace hamzstplot;

    std::vector<double> x =vrandn_gamma(2, 1.5,1000);

    auto h = hist(x);
    std::cout << "Histogram with " << h->num_bins() << " bins" << std::endl;

    show();

    return 0;
}
```

```
}
```

Code 52: *main.cpp*

```
CFLAGS = -ggdb
DEFINES = -DDEBUGGA
INCLUDES =
LIBS = -lstdc++ -lhamzstplot -lsymintegration
MAIN = main.o
CC=g++

.cc.o:
$(CC) -c $(CFLAGS) $(DEFINES) $(INCLUDES) $<

all:: main

gnuplot_i.o:
main.o: main.cpp

main: $(MAIN)
$(CC) -o $@ $(CFLAGS) $(MAIN) $(LIBS)

clean:
rm -f $(MAIN) main
```

Code 53: *Makefile*




Figure 4.12: The histogram of 1000 random numbers that has been generated with Gamma distribution of parameters $\alpha = 2, \beta = 1.5$. (*Hamzstplot/Examples with Makefile/Statistics/Plot 2D Histogram of Random Number/main.cpp*).

V. STATISTICAL INFERENCE

[SI*] The action of statistical inference involves using mathematical techniques to make conclusions about unknown population parameters based on collected data. The field of statistical inference employs a variety of stochastic models to analyze and put forward efficient methods for carrying out such analyses.

Analysis and methods of statistical inference can be categorized as either frequentist (also known as classical) or Bayesian. The former is based on the assumption that population parameters of some underlying distribution, or probability law, exist and are fixed, but are yet unknown. The process of statistical inference then deals with making conclusions about these parameters based on sampled data. In the latter Bayesian case, it is only assumed that there is a prior distribution of the parameters. In this case, the key process deals with analyzing a posterior distribution (of the parameters)-an outcome of the inference process.

In general, a statistical inference process involves data, a model, and analysis. The data is assumed to be comprised of random samples from the model. The goal of the analysis is then to make informed statements about population parameters of the model based on the data. Such statements typically take one of the following forms:

1. Point Estimation

Determination of a single value (or vector of values) representing a best estimate of the parameter / parameters. In this case, the notion of "best" can be defined in different ways.

2. Confidence Intervals

Determination of a range of values where the parameter lies. Under the model and the statistical process used, it is guaranteed that the parameter lies within this range with a pre-specified probability.

3. Hypothesis Tests

The process of determining if the parameter lies in a given region, in the complement of that region, or fails to take on a specific value. Such tests often represent a scientific hypothesis in a very natural way

[SI*] A Random Sample

When carrying out (frequentist) statistical inference, we assume there is some underlying distribution $F(x, \theta)$ from which we are sampling, where θ is the scalar or vector-valued unknown parameter we wish to know. Importantly, we assume that each observation is statistically independent and identically distributed as the rest. That is, from a probabilistic perspective, the observations are taken as independent and identically distributed (i.i.d.) random variables. In mathematical statistics language, this is called a random sample. We denote the random variables of the observations by $X_1, \dots, X_n$ and their respective values $x_1, \dots, x_n$.

Typically, we compute statistics from the random sample. For example, two common standard statistics include the sample mean and sample variance. However, we can model

these statistics as random variables

$$\begin{aligned}\bar{X} &= \frac{1}{n} \sum_{i=1}^n X_i \\ S^2 &= \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2\end{aligned}\tag{4.109}$$

Note that for S^2 , the denominator is $n - 1$ to make S^2 an unbiased estimator.

In general, the phrase statistic implies a quantity calculated based on the sample. When working with data, the sample mean and sample variance are nothing but numbers computed from our sample observations. However, in the statistical inference paradigm, we associate random variables to these values, since they themselves are functions of the random sample.

To illustrate the fact that $\bar{X}$ and S^2 are random variables, assume we have sampled data from certain distribution (e.g. exponential). If we collect $n = 10$ observations, then the sample mean and sample variance are random variables. The point to see is that $\bar{X}$ and S^2 are themselves random variable with underlying distributions.

[SI*] Statistical inference consists of some methods by which one makes inferences or generalizations about a population. The trend today is to distinguish between the classical method of estimating a population parameter, whereby inferences are based strictly on information obtained from a random sample selected from the population, and the Bayesian method, which utilizes prior subjective knowledge about the probability distribution of the unknown parameters in conjunction with the information provided by the sample data.

On this book, the sections on confidence interval before the Bayesian statistics section are the so-called classical methods to estimate unknown population parameters such as the mean, the proportion, and the variance by computing statistics from random samples and applying the theory of sampling distributions.

i. Sampling from a Normal Population

[SI*] It is often assumed that the distribution $F(x; \theta)$ is a normal distribution, and hence $\theta = (\mu, \sigma^2)$. This assumption is called the normality assumption, and is sometimes justified due to the central limit theorem. Under the normality assumption, the distribution of the random variables $\bar{X}$ and S^2 as well as transformations of them are well known. The following three distributional relationships play a key role:

$$\begin{aligned}\bar{X} &\text{Normal} \left(\mu, \frac{\sigma^2}{n} \right) \\ \frac{(n-1)S^2}{\sigma^2} &\chi_{n-1}^2 \\ T &:= \frac{\bar{X} - \mu}{S/\sqrt{n}} t_{n-1}\end{aligned}\tag{4.110}$$

Here, " " denotes as "distributed as," and implies that the statistics on the left hand side of the " " symbols are distributed according to the distribution on the right hand side. The notation

χ_{n-1}^2 and t_{n-1} denotes a chi-squared and t -distribution, respectively, each with $n - 1$ degrees of freedom. The chi-squared distribution is a gamma distribution with parameters $\alpha = \frac{n}{2}$ and $\beta = 2$, some textbooks, e.g. [19], use another parameter thus state that $\lambda = \frac{1}{2}$ because $\lambda = \frac{1}{\beta}$, for Gamma distribution in this book we use parameters α and β .

ii. The Central Limit Theorem

[SI*] The Central Limit Theorem (CLT) is one of the most fundamental results of probability and statistics. It has several versions and many generalizations, they all have one thing in common: summation of a large number of random quantities, each with finite variance, yield a sum that is approximately normally distributed. This is the main reason that the normal distribution is ubiquitous in nature and present throughout the universe.

Theorem 4.19: Central Limit Theorem

If $\bar{X}$ is the mean of a random sample of size n taken from a population with mean μ and finite variance σ^2 , then the limiting form of the distribution of

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

as $n \rightarrow \infty$, is the standard normal distribution $n(z; 0, 1)$.

iii. Point Estimation

[SI*] A point estimate of some population parameter θ is a single value $\hat{\theta}$ of a statistic $\hat{\Theta}$. For example, the value $\bar{x}$ of the statistic $\bar{X}$, computed from a sample of size n , is a point estimate of the population parameter μ .

Definition 4.41: Unbiased Estimator

A statistic $\hat{\Theta}$ is said to be an unbiased estimator of the parameter θ if

$$\mu_{\hat{\Theta}} = E(\hat{\Theta}) = \theta$$

Definition 4.42: Most Efficient Estimator

If we consider all possible unbiased estimators of some parameter θ , the one with the smallest variance is called the most efficient estimator of θ .

iv. Confidence Interval as a Concept

[SI*] Even the most efficient unbiased estimator is unlikely to estimate the population parameter exactly. It is true that estimation accuracy increases with large samples, but there is still no reason we should expect a point estimate to be exactly equal to the population parameter it is supposed to estimate. There are many situations in which it is preferable to determine an interval within which we would expect to find the value of the parameter. Such an interval

is called an interval estimate.

An interval estimate of a population parameter θ is an interval of the form $\hat{\theta}_L < \theta < \hat{\theta}_U$, where $\hat{\theta}_L$ and $\hat{\theta}_U$ depend on the value of the statistic $\hat{\Theta}$ for a particular sample and also on the sampling distribution of $\hat{\Theta}$.

[SI*] Since different samples will generally yield different values of $\hat{\Theta}$ and, therefore different values of $\hat{\theta}_L$ and $\hat{\theta}_U$, these endpoints of the interval are values of corresponding random variables $\hat{\Theta}_L$ and $\hat{\Theta}_U$. From the sampling distribution of $\hat{\Theta}$ we shall be able to determine $\hat{\Theta}_L$ and $\hat{\Theta}_U$ such that $P(\hat{\Theta}_L < \theta < \hat{\Theta}_U)$ is equal to any positive fractional value we care to specify. If, for instance, we find $\hat{\Theta}_L$ and $\hat{\Theta}_U$ such that

$$P(\hat{\Theta}_L < \theta < \hat{\Theta}_U) = 1 - \alpha$$

for $0 < \alpha < 1$, then we have a probability of $1 - \alpha$ of selecting a random sample that will produce an interval containing θ . The interval $\hat{\theta}_L < \theta < \hat{\theta}_U$, computed from the selected sample, is called a $100(1 - \alpha)\%$ confidence interval, the fraction $1 - \alpha$ is called the confidence coefficient or the degree of confidence, and the endpoints, $\hat{\theta}_L$ and $\hat{\theta}_U$, are called the lower and upper confidence limits. Thus, when $\alpha = 0.05$, we have a 95% confidence interval, and when $\alpha = 0.01$ we obtain a wider 99% confidence interval. The wider the confidence interval is, the more confident we can be that the interval contains the unknown parameter.

Definition 4.43: Confidence Interval on μ, σ^2 Known

If $\bar{x}$ is the mean of a random sample of size n from a population with known variance σ^2 , a $100(1 - \alpha)\%$ confidence interval for μ is given by

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

where $z_{\alpha/2}$ is the z -value leaving an area of $\alpha/2$ to the right.

Theorem 4.20: Error in Confidence Interval

If $\bar{x}$ is used as an estimate of μ , we can be $100(1 - \alpha)\%$ confident that the error will not exceed $z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$.

Theorem 4.21: Sample Size for Confidence Interval

If $\bar{x}$ is used as an estimate of μ , we can be $100(1 - \alpha)\%$ confident that the error will not exceed a specified amount e when the sample size is

$$n = \left(\frac{z_{\alpha/2} \sigma}{e} \right)^2$$

Definition 4.44: One-Sided Confidence Bounds on μ, σ^2 Known

If $\bar{X}$ is the mean of a random sample of size n from a population with variance σ^2 , the one-sided $100(1 - \alpha)\%$ confidence bounds for μ are given by

upper-one sided bound: $\bar{x} + z_\alpha \frac{\sigma}{\sqrt{n}}$.

lower-one sided bound: $\bar{x} - z_\alpha \frac{\sigma}{\sqrt{n}}$.

[SI*] If we have a random sample from a normal distribution, then the random variable

$$T = \frac{\bar{X} - \mu}{S/\sqrt{n}}$$

has a Student' t -distribution with $n - 1$ degrees of freedom. Here S is the sample standard deviation. In this situation, with σ unknown, T can be used to construct a confidence interval on μ .

Definition 4.45: Confidence Interval on μ, σ^2 Unknown

If $\bar{x}$ and s are the mean and standard deviation of a random sample from a normal population with unknown variance σ^2 , a $100(1 - \alpha)\%$ confidence interval for μ is given by

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} < \mu < \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

where $t_{\alpha/2}$ is the t -value with $\nu = n - 1$ degrees of freedom, leaving an area of $\alpha/2$ to the right.

[SI*] We should emphasize that for σ known we exploited the Central Limit Theorem, whereas for σ unknown we made use of the sampling distribution of the random variable T .

[SI*] Often statisticians recommend that even when normality cannot be assumed, σ is unknown, and $n \geq 30$, s can replace σ and the confidence interval

$$\bar{x} \pm z_{\alpha/2} \frac{s}{\sqrt{n}}$$

may be used. This is often referred to as a large-sample confidence interval.

[SI*] A common way to calculate a confidence interval is by using the following formula:

Confidence interval = Point estimate $\pm$ Margin of Error

The sample mean is often used as the point estimate, and the required margin on error depends on the desired confidence level and the variability of the data, calculated as a critical value (e.g., a z -value or t -value) multiplied by the standard error of the mean.

[SI*] The confidence interval provides a range of plausible values for a population parameter. The confidence level indicates the reliability of the method used to create the interval. For example, a 95% confidence interval for the average height of a certain plant species suggests that we are 95% confident that the true average height of the species falls between 10 and 15.

[SI*] Sometimes, other than the population mean, the experimenter may also be interested in predicting the possible value of a future observation. For instance, in quality control, the experimenter may need to use the observed data to predict a new observation.

Definition 4.46: Prediction Interval of a Future Observation, σ^2 Known

For a normal distribution of a measurements with unknown mean μ and known variance σ^2 , a $100(1 - \alpha)\%$ prediction interval of a future observation X_0 is

$$\bar{x} - z_{\alpha/2}\sigma\sqrt{1 + \frac{1}{n}} < X_0 < \bar{x} + z_{\alpha/2}\sigma\sqrt{1 + \frac{1}{n}}$$

where $z_{\alpha/2}$ is the z -value leaving an area of $\alpha/2$ to the right.

Definition 4.47: Prediction Interval of a Future Observation, σ^2 Unknown

For a normal distribution of a measurements with unknown mean μ and unknown variance σ^2 , a $100(1 - \alpha)\%$ prediction interval of a future observation X_0 is

$$\bar{x} - t_{\alpha/2}s\sqrt{1 + \frac{1}{n}} < X_0 < \bar{x} + t_{\alpha/2}s\sqrt{1 + \frac{1}{n}}$$

where $t_{\alpha/2}$ is the t -value with $\nu = n - 1$ degrees of freedom, leaving an area of $\alpha/2$ to the right.

[SI*] If we have two populations with means μ_1 and μ_2 and variance σ_1^2 and σ_2^2 , respectively, a point estimator of the difference between μ_1 and μ_2 is given by statistic $\bar{X}_1 - \bar{X}_2$.

Definition 4.48: Confidence Interval for $\mu_1 - \mu_2$, σ_1^2 and σ_2^2 Known

If $\bar{x}_1$ and $\bar{x}_2$ are means of independent random samples of sizes n_1 and n_2 from populations with known variances σ_1^2 and σ_2^2 , respectively, a $100(1 - \alpha)\%$ confidence interval for $\mu_1 - \mu_2$ is given by

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2}\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} < \mu_1 - \mu_2 < (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2}\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

where $z_{\alpha/2}$ is the z -value leaving an area of $\alpha/2$ to the right.

[SI*] Consider the case where σ_1^2 and σ_2^2 are unknown. If $\sigma_1^2 = \sigma_2^2 = \sigma^2$, we obtain a standard normal variable of the form

$$Z = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\sigma^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

The two random variables

$$\frac{(n_1 - 1)S_1^2}{\sigma^2} \quad \text{and} \quad \frac{(n_2 - 1)S_2^2}{\sigma^2}$$

have chi-squared distributions with $n_1 - 1$ and $n_2 - 1$ degrees of freedom, respectively. Furthermore, they are independent chi-squared variables, since the random samples were selected independently. Consequently their sum

$$V = \frac{(n_1 - 1)S_1^2}{\sigma^2} + \frac{(n_2 - 1)S_2^2}{\sigma^2} = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{\sigma^2}$$

has a chi-squared distribution with $\nu = n_1 + n_2 - 2$ degrees of freedom.

Since the preceding expressions for Z and V can be shown to be independent, it follows that the statistic

$$T = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\sigma^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}} \cdot \frac{1}{\frac{(n_1-1)S_1^2 + (n_2-1)S_2^2}{\sigma^2(n_1+n_2-2)}}$$

has the t -distribution with $\nu = n_1 + n_2 - 2$ degrees of freedom.

Definition 4.49: Pooled Estimate of Variance

A point estimate of the unknown common variance σ^2 can be obtained by pooling the sample variances. Denoting the pooled estimator by S_p^2 , we have the following

$$S_p^2 = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2}$$

Substituting S_p^2 in the T statistic, we obtain the less cumbersome form

$$T = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{S_p \sqrt{\left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

Using the T statistic, we have

$$P(-t_{\alpha/2} < T < t_{\alpha/2}) = 1 - \alpha$$

where $t_{\alpha/2}$ is the t -value with $n_1 + n_2 - 2$ degrees of freedom, above which we find an area of $\alpha/2$. Substituting for T in the inequality, we write

$$P \left(-t_{\alpha/2} < \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{S_p \sqrt{\left(\frac{1}{n_1} + \frac{1}{n_2} \right)}} < t_{\alpha/2} \right) = 1 - \alpha$$

After the usual mathematical manipulations, the difference of the sample means $\bar{x}_1 - \bar{x}_2$ and the pooled variance are computed and then the following $100(1 - \alpha)\%$ confidence interval for $\mu_1 - \mu_2$ is obtained.

The value of s_p^2 is easily seen to be a weighted average of the two sample variances s_1^2 and s_2^2 , where the weights are the degrees of freedom.

Definition 4.50: Confidence Interval for $\mu_1 - \mu_2$, $\sigma_1^2 = \sigma_2^2$ but Both Unknown

If $\bar{x}_1$ and $\bar{x}_2$ are means of independent random samples of sizes n_1 and n_2 from approximately normal populations with unknown but equal variances, a $100(1 - \alpha)\%$ confidence interval for $\mu_1 - \mu_2$ is given by

$$(\bar{x}_1 - \bar{x}_2) - t_{\alpha/2} s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} < \mu_1 - \mu_2 < (\bar{x}_1 - \bar{x}_2) + t_{\alpha/2} s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$$

where s_p is the pooled estimate of the population standard deviation and $t_{\alpha/2}$ is the t -value with $\nu = n_1 + n_2 - 2$ degrees of freedom leaving an area of $\alpha/2$ to the right.

Definition 4.51: Confidence Interval for $\mu_1 - \mu_2$, $\sigma_1^2 \neq \sigma_2^2$ and Both Unknown

If $\bar{x}_1$ and s_1^2 and $\bar{x}_2$ and s_2^2 are the means and variance of independent random samples of sizes n_1 and n_2 , respectively, from approximately normal populations with unknown and unequal variances, a $100(1 - \alpha)\%$ confidence interval for $\mu_1 - \mu_2$ is given by

$$(\bar{x}_1 - \bar{x}_2) - t_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} < \mu_1 - \mu_2 < (\bar{x}_1 - \bar{x}_2) + t_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

where $t_{\alpha/2}$ is the t -value with

$$\nu_{\text{raw}} = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\frac{\left(\frac{s_1^2}{n_1} \right)^2}{n_1 - 1} + \frac{\left(\frac{s_2^2}{n_2} \right)^2}{n_2 - 1}}$$

$\nu = \lfloor \nu_{\text{raw}} \rfloor$ is the degrees of freedom leaving an area of $\alpha/2$ to the right.

[SI*] Paired Observations

We shall consider estimation procedures for the difference of two means when the samples are not independent and the variances of the two populations are not necessarily equal.

The conditions of the two populations are not assigned randomly to experimental units. Rather, each homogeneous experimental unit receives both population conditions; as a result, each experimental unit has a pair of observations, one for each observation. For example, if we run a test on a new diet using 15 individuals, the weights before and after going on the diet form the information for our two samples. The two populations are "before" and "after," and the experimental unit is the individual. Obviously, the observations in a pair have something in common. To determine if the diet is effective, we consider the differences $d_1, d_2, \dots, d_n$ in the paired observations. These differences are the values of a random sample $D_1, D_2, \dots, D_n$ from a population of differences that we shall assume to be normally distributed with mean $\mu_D = \mu_1 - \mu_2$ and variance σ_D^2 . We estimate σ_D^2 by s_d^2 , the variance of the differences that constitute our sample. the point estimator of μ_D is given by $\bar{D}$.

The reader may visualize the i -th pair difference as

$$D_i = X_{1i} - X_{2i}$$

Since the two observations are taken on the same experimental unit, they are not independent and, in fact,

$$\text{Var}(D_i) = \text{Var}(X_{1i} - X_{2i}) = \sigma_1^2 + \sigma_2^2 - 2\text{Cov}(X_{1i}, X_{2i})$$

One certainly expects that if the unit is homogeneous, the covariance is positive. As a result, the gain in quality of the confidence interval over that obtained without pairing will be greatest when there is homogeneity within units and large differences as one goes from unit to unit.

[SI*] Although pairing should indeed reduce variance and hence reduce the standard error of the point estimate, the degrees of freedom are reduced by reducing the problem to a one-sample problem. As a result, the $t_{\alpha/2}$ point attached to the standard error is adjusted accordingly. Thus, pairing may be counterproductive.

[SI*] A $100(1 - \alpha)\%$ confidence interval for μ_D can be established by writing

$$P(-t_{\alpha/2} < T < t_{\alpha/2}) = 1 - \alpha$$

where $T = \frac{\bar{D} - \mu_D}{s_d / \sqrt{n}}$ and $t_{\alpha/2}$, as before, is a value of the t -distribution with $n - 1$ degrees of freedom.

Definition 4.52: Confidence interval for $\mu_D = \mu_1 - \mu_2$ for Paired Observations

If $\bar{d}$ and s_d are the mean and standard deviation, respectively, of the normally distributed differences of n random pairs of measurements, a $100(1 - \alpha)\%$ confidence interval for $\mu_D = \mu_1 - \mu_2$ is

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} < \mu_D < \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

where $t_{\alpha/2}$ is the t -value with $\nu = n - 1$ degrees of freedom, leaving an area of $\alpha/2$ to the right.

[SI*] Single Sample: Estimating a Proportion

A point estimator of the proportion p in a binomial experiment is given by the statistic $\hat{P} = \frac{X}{n}$, where X represents the number of successes in n trials. Therefore, the sample proportion $\hat{p} = \frac{x}{n}$ will be used as the point estimate of the parameter p .

If the unknown proportion p is not expected to be too close to 0 or 1, we can establish a confidence interval for p by considering the sampling distribution of $\hat{P}$. Designating a failure in each binomial trial by the value 0 and a success by the value 1, the number of successes, can be interpreted as the sum of n values consisting only of 0 and 1s, and $\hat{p}$ is just the sample mean of these n values. Hence, by the Central Limit Theorem, for n sufficiently large, $\hat{P}$ is approximately normally distributed with mean

$$\mu_{\hat{P}} = E(\hat{P}) = E\left(\frac{X}{n}\right) = \frac{np}{n} = p$$

and variance

$$\sigma_{\hat{p}}^2 = \sigma_{\frac{X}{n}} = \frac{\sigma_X^2}{n^2} = \frac{npq}{n^2} = \frac{pq}{n}$$

Therefore, we can assert that

$$P(-z_{\alpha/2} < Z < z_{\alpha/2}) = 1 - \alpha$$

with

$$Z = \frac{\hat{p} - p}{\sqrt{\frac{pq}{n}}}$$

and $z_{\alpha/2}$ is the z-value leaving an area of $\alpha/2$ to the right. Substituting for Z , we write

$$P\left(-z_{\alpha/2} < \frac{\hat{p} - p}{\sqrt{\frac{pq}{n}}} < z_{\alpha/2}\right) = 1 - \alpha$$

When n is large, very little error is introduced by substituting the point estimate $\hat{p} = \frac{x}{n}$ for the p under the radical sign. Then we can write

$$P\left(\hat{p} - z_{\alpha/2}\sqrt{\frac{pq}{n}} < p < \hat{p} + z_{\alpha/2}\sqrt{\frac{pq}{n}}\right) \approx 1 - \alpha$$

On the other hand, by solving for p in the quadratic equality above,

$$-z_{\alpha/2} < \frac{\hat{p} - p}{\sqrt{\frac{pq}{n}}} < z_{\alpha/2}$$

we obtain another form of the confidence interval for p with limits

$$\frac{\hat{p} + \frac{z_{\alpha/2}^2}{2n}}{1 + \frac{z_{\alpha/2}^2}{n}} \pm \frac{z_{\alpha/2}}{1 + \frac{z_{\alpha/2}^2}{n}} \sqrt{\frac{\hat{p}\hat{q}}{n} + \frac{z_{\alpha/2}^2}{4n^2}}$$

Definition 4.53: Large Sample Confidence Intervals for p

If $\hat{p}$ is the proportion of successes in a random sample of size n and $\hat{q} = 1 - \hat{p}$, an approximate $100(1 - \alpha)\%$ confidence interval, for the binomial parameter p is given by (method 1)

$$\hat{p} - z_{\alpha/2}\sqrt{\frac{pq}{n}} < p < \hat{p} + z_{\alpha/2}\sqrt{\frac{pq}{n}}$$

or by (method 2)

$$\frac{\hat{p} + \frac{z_{\alpha/2}^2}{2n}}{1 + \frac{z_{\alpha/2}^2}{n}} - \frac{z_{\alpha/2}}{1 + \frac{z_{\alpha/2}^2}{n}} \sqrt{\frac{\hat{p}\hat{q}}{n} + \frac{z_{\alpha/2}^2}{4n^2}} < p < \frac{\hat{p} + \frac{z_{\alpha/2}^2}{2n}}{1 + \frac{z_{\alpha/2}^2}{n}} + \frac{z_{\alpha/2}}{1 + \frac{z_{\alpha/2}^2}{n}} \sqrt{\frac{\hat{p}\hat{q}}{n} + \frac{z_{\alpha/2}^2}{4n^2}}$$

where $z_{\alpha/2}$ is the z-value leaving an area of $\alpha/2$ to the right.

When n is small and the unknown proportion p is believed to be close to 0 or to 1, the confidence-interval procedure established here is unreliable and, therefore, should not be

used. To be on the safe side, one should require both $n\hat{p}$ and $n\hat{q}$ to be greater than or equal to 5.

Theorem 4.22: Error in estimating p by $\hat{p}$

If $\hat{p}$ is used as an estimate of p , we can be $100(1 - \alpha)\%$ confident that the error will not exceed $z_{\alpha/2} \sqrt{\frac{\hat{p}\hat{q}}{n}}$.

Theorem 4.23: Choice of Sample Size

If $\hat{p}$ is used as an estimate of p , we can be $100(1 - \alpha)\%$ confident that the error will be less than a specified amount e when the sample size is approximately

$$n = \frac{z_{\alpha/2}^2 \hat{p}\hat{q}}{e^2}$$

Theorem 4.24: Determine Sample Size Without $\hat{p}$

If $\hat{p}$ is used as an estimate of p , we can be at least $100(1 - \alpha)\%$ confident that the error will not exceed a specified amount e when the sample size is

$$n = \frac{z_{\alpha/2}^2}{4e^2}$$

[SI*] **Two Samples: Estimating the Difference Between Two Proportions**

Consider the problem where we wish to estimate the difference between two binomial parameters p_1 and p_2 . First, we select independent random samples of sizes n_1 and n_2 from the two binomial populations with means $n_1 p_1$ and $n_2 p_2$ and variances $n_1 p_1 q_1$ and $n_2 p_2 q_2$, respectively; then we determine the numbers x_1 and x_2 of people in each sample with the target criteria (e.g. defective, lung cancer, etc) and form the proportions

$$\begin{aligned}\hat{p}_1 &= \frac{x_1}{n} \\ \hat{p}_2 &= \frac{x_2}{n}\end{aligned}$$

A point estimator of the difference between the two proportions, $p_1 - p_2$, is given by the statistic $\hat{P}_1 - \hat{P}_2$. Therefore, the difference of the sample proportions, $\hat{p}_1 - \hat{p}_2$, will be used as the point estimate of $p_1 - p_2$.

We know that $\hat{P}_1$ and $\hat{P}_2$ are each approximately normally distributed, with means p_1 and p_2 and variances $\frac{p_1 q_1}{n_1}$ and $\frac{p_2 q_2}{n_2}$, respectively. Choosing independent samples from the two populations ensures that the variables $\hat{P}_1$ and $\hat{P}_2$ will be independent, and then by the reproductive property of the normal distribution, we conclude that $\hat{P}_1 - \hat{P}_2$ is approximately normally distributed with mean

$$\mu_{\hat{P}_1 - \hat{P}_2} = p_1 - p_2$$

and variance

$$\sigma_{\hat{P}_1 - \hat{P}_2}^2 = \frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}$$

Therefore, we can assert that

$$P(-z_{\alpha/2} < Z < z_{\alpha/2}) = 1 - \alpha$$

where

$$Z = \frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}}}$$

and $z_{\alpha/2}$ is the value above which we find an area of $\alpha/2$ to the right under the standard normal curve. Substituting for Z , we write

$$P\left(-z_{\alpha/2} < \frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}}} < z_{\alpha/2}\right)$$

After performing the usual mathematical manipulations, we replace p_1, p_2, q_1 , and q_2 under the radical sign by their estimates $\hat{p}_1 = \frac{x_1}{n_1}, \hat{p}_2 = \frac{x_2}{n_2}, \hat{q}_1 = 1 - \hat{p}_1$, and $\hat{q}_2 = 1 - \hat{p}_2$, provided that $n_1 \hat{p}_1, n_1 \hat{q}_1, n_2 \hat{p}_2$, and $n_2 \hat{q}_2$ are all greater than or equal to 5.

Definition 4.54: Large Sample Confidence Interval for $p_1 - p_2$

If $\hat{p}_1$ and $\hat{p}_2$ are the proportions of successes in random samples of sizes n_1 and n_2 , respectively, $\hat{q}_1 = 1 - \hat{p}_1$, and $\hat{q}_2 = 1 - \hat{p}_2$, an approximate $100(1 - \alpha)\%$ confidence interval for the difference of two binomial parameters, $p_1 - p_2$, is given by

$$(\hat{p}_1 - \hat{p}_2) - z_{\alpha/2} \sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}} < p_1 - p_2 < (\hat{p}_1 - \hat{p}_2) + z_{\alpha/2} \sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}}$$

where $z_{\alpha/2}$ is the z -value leaving an area of $\alpha/2$ to the right.

[SI*] Single Sample: Estimating the Variance

If a sample of size n is drawn from a normal population with variance σ^2 and the sample variance s^2 is computed, we obtain a value of the statistic S^2 . This computed sample variance is used as a point estimate of σ^2 . Hence, the statistic S^2 is called an estimator of σ^2 .

An interval estimate of σ^2 can be established by using this statistic

$$X^2 = \frac{(n-1)S^2}{\sigma^2}$$

The statistic X^2 has a chi-squared distribution with $n - 1$ degrees of freedom when samples are chosen from a normal population. We may write

$$P(\chi_{1-\frac{\alpha}{2}}^2 < X^2 < \chi_{\alpha/2}^2) = 1 - \alpha$$

where $\chi_{1-\frac{\alpha}{2}}^2$ and $\chi_{\alpha/2}^2$ are values of the chi-square distribution with $n - 1$ degrees of freedom, leaving areas of $1 - \frac{\alpha}{2}$ and $\frac{\alpha}{2}$, respectively, to the right. Substituting for X^2 , we write

$$P\left[\chi_{1-\frac{\alpha}{2}}^2 < \frac{(n-1)S^2}{\sigma^2} < \chi_{\alpha/2}^2\right] = 1 - \alpha$$

Dividing each term in the inequality by $(n-1)S^2$ and then inverting each term, we obtain

$$P \left[\frac{(n-1)S^2}{\chi_{\alpha/2}^2} < \sigma^2 < \frac{(n-1)S^2}{\chi_{1-\frac{\alpha}{2}}^2} \right] = 1 - \alpha$$

Definition 4.55: Confidence Interval for σ^2

If s^2 is the variance of a random sample of size n from a normal population, a $100(1-\alpha)\%$ confidence interval for σ^2 is

$$\frac{(n-1)S^2}{\chi_{\alpha/2}^2} < \sigma^2 < \frac{(n-1)S^2}{\chi_{1-\frac{\alpha}{2}}^2}$$

where $\chi_{1-\frac{\alpha}{2}}^2$ and $\chi_{\alpha/2}^2$ are values of the chi-square distribution with $n-1$ degrees of freedom, leaving areas of $1-\frac{\alpha}{2}$ and $\frac{\alpha}{2}$, respectively, to the right.

[SI*] Two Samples: Estimating the Ratio of Two Variances

A point estimate of the ratio of two population variances $\frac{\sigma_1^2}{\sigma_2^2}$ is given by the ratio $\frac{s_1^2}{s_2^2}$ of the sample variances. Hence, the statistic $\frac{s_1^2}{s_2^2}$ is called an estimator of $\frac{\sigma_1^2}{\sigma_2^2}$.

If σ_1^2 and σ_2^2 are the variances of normal populations, we can establish an interval estimate of by using the statistic

$$F = \frac{\sigma_2^2 S_1^2}{\sigma_1^2 S_2^2}$$

The random variable F has an F -distribution with $\nu_1 = n_1 - 1$ and $\nu_2 = n_2 - 1$ degrees of freedom. Therefore, we may write

$$P(f_{1-\frac{\alpha}{2}}(\nu_1, \nu_2) < F < f_{\frac{\alpha}{2}}(\nu_1, \nu_2)) = 1 - \alpha$$

where $f_{1-\frac{\alpha}{2}}(\nu_1, \nu_2)$ and $f_{\frac{\alpha}{2}}(\nu_1, \nu_2)$ are the values of the F -distribution with ν_1 and ν_2 degrees of freedom, leaving areas of $1-\frac{\alpha}{2}$ and $\frac{\alpha}{2}$, respectively, to the right.

Substituting for F , we write

$$P \left(f_{1-\frac{\alpha}{2}}(\nu_1, \nu_2) < \frac{\sigma_2^2 S_1^2}{\sigma_1^2 S_2^2} < f_{\frac{\alpha}{2}}(\nu_1, \nu_2) \right) = 1 - \alpha$$

Multiplying each term in the inequality by $\frac{S_2^2}{S_1^2}$ and then inverting each term, we obtain

$$P \left(\frac{S_1^2}{S_2^2} \frac{1}{f_{\frac{\alpha}{2}}(\nu_1, \nu_2)} < \frac{\sigma_1^2}{\sigma_2^2} < \frac{S_1^2}{S_2^2} \frac{1}{f_{1-\frac{\alpha}{2}}(\nu_1, \nu_2)} \right) = 1 - \alpha$$

the results enable us to replace the quantity $f_{1-\frac{\alpha}{2}}(\nu_1, \nu_2)$ by $\frac{1}{f_{\frac{\alpha}{2}}(\nu_2, \nu_1)}$. Therefore,

$$P \left(\frac{S_1^2}{S_2^2} \frac{1}{f_{\frac{\alpha}{2}}(\nu_1, \nu_2)} < \frac{\sigma_1^2}{\sigma_2^2} < \frac{S_1^2}{S_2^2} f_{\frac{\alpha}{2}}(\nu_2, \nu_1) \right) = 1 - \alpha$$

For any two independent random samples of sizes n_1 and n_2 selected from two normal populations, the ratio of the sample variances $\frac{s_1^2}{s_2^2}$ is computed, then the confidence interval can be obtained.

Definition 4.56: The Confidence Interval for $\frac{\sigma_1^2}{\sigma_2^2}$

If s_1^2 and s_2^2 are the variances of independent samples of sizes n_1 and n_2 , respectively, from normal populations, then a $100(1 - \alpha)\%$ confidence interval for $\frac{\sigma_1^2}{\sigma_2^2}$ is

$$\frac{s_1^2}{s_2^2} \frac{1}{f_{\frac{\alpha}{2}}(v_1, v_2)} < \frac{\sigma_1^2}{\sigma_2^2} < \frac{s_1^2}{s_2^2} f_{\frac{\alpha}{2}}(v_2, v_1)$$

where $f_{\frac{\alpha}{2}}(v_1, v_2)$ is an f -value with $v_1 = n_1 - 1$ and $v_2 = n_2 - 1$ degrees of freedom, leaving an area of $\frac{\alpha}{2}$ to the right, and $f_{\frac{\alpha}{2}}(v_2, v_1)$ is a similar f -value with $v_2 = n_2 - 1$ and $v_1 = n_1 - 1$ degrees of freedom.

v. Functions in SymIntegration Related to Confidence Interval

[SI*] To compute several computations related to confidence interval, we can use these functions in SymIntegration:

confidenceinterval_onesampletwosides(vector<double> data, double σ , double α)

this function computes a two sides $100(1 - \alpha)\%$ confidence interval for μ with a given σ (the population standard deviation) that is known. The formula comes from the two sides probability statement that makes use of the Central Limit Theorem:

$$P\left(-z_{\alpha/2} < \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} < z_{\alpha/2}\right) = 1 - \alpha$$

confidenceinterval_onesampleonesided(vector<double> data, double σ , double α)

this function computes a one-sided $100(1 - \alpha)\%$ confidence interval for μ with a given σ (the population standard deviation) that is known. The formula comes from the one-sided probability statement that makes use of the Central Limit Theorem:

$$P\left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} < z_{\alpha/2}\right) = 1 - \alpha$$

for the upper bound, and

$$P\left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} > -z_{\alpha/2}\right) = 1 - \alpha$$

for the lower bound.

confidenceinterval_onesampletwosides(vector<double> data, double α)

this function computes a two sides $100(1 - \alpha)\%$ confidence interval for μ with unknown σ (the population standard deviation) so we use T statistics that has a Student's t -distribution to approximate σ with s (sample standard deviation).

The formula comes from the two sides probability statement that makes use of the Central Limit Theorem:

$$P\left(-t_{\alpha/2} < \frac{\bar{X} - \mu}{S/\sqrt{n}} < t_{\alpha/2}\right) = 1 - \alpha$$

confidenceinterval_onesampleonesided(vector<double> data, double α)

this function computes a one-sided $100(1 - \alpha)\%$ confidence interval for μ with unknown σ (the population standard deviation) so we use T statistics that has a Student's t -distribution to approximate σ with s (sample standard deviation).

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for One Sample ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for One Sample ]# ./main

*****

Confidence interval one sample two sides
Mean data = 2.6,  $\sigma$  = 0.3, n = 36

The 99% confidence interval is :

2.47121 <  $\mu$ 1 < 2.72879

*****

*****

Confidence interval one sample one-sided confidence bounds
Mean data = 6.2,  $\sigma$  = 2, n = 25

The 95% confidence interval is :

Upper one-sided bound: 6.85794
Lower one-sided bound: 5.54206

*****

*****

Confidence interval one sample two sides
Mean data = 10,  $\sigma$  = 0.282843, n = 7

The 95% confidence interval is :

9.73841 <  $\mu$ 1 < 10.2616

*****

*****

Confidence interval one sample one-sided confidence bounds
Mean data = 10,  $\sigma$  = 0.282843, n = 7

The 95% confidence interval is :

Upper one-sided bound: 10.2077
Lower one-sided bound: 9.79227

*****

Time taken by function: 1319 microseconds
```

Figure 4.13: The computation of one sample confidence interval for two sides and one-sided confidence bounds with σ known and σ unknown (*SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval for One Sample/main.cpp*).

predictioninterval(vector<double> data, double σ , double α)

this function computes a $100(1 - \alpha)\%$ prediction interval of a future observation with known σ .

predictioninterval(vector<double> data, double α)

this function computes a $100(1 - \alpha)\%$ prediction interval of a future observation with unknown σ .

```
-----
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Prediction Interval for One Sample ]# ./main
*****

Prediction interval for one sample data,  $\sigma$  known
Mean data = 257300,  $\sigma$  = 25000, n = 50

The 95% prediction interval is :

207813 < X0 < 306787

*****

*****

Prediction interval for one sample data,  $\sigma$  unknown
Mean data = 96.24, s = 0.878714, n = 30

The 95% prediction interval is :

94.4131 < X0 < 98.0669

*****

Time taken by function: 973 microseconds
-----
```

Figure 4.14: *The computation of one sample prediction interval with σ known and σ unknown (SymIntegration/Examples/Statistics/Test SymIntegration Prediction Interval for One Sample/main.cpp).*

Sweden Sexy, Walnut PUPULU, Sine Bam Bam, Belang CUPULU, Bludut Hasta La Meo
The functions for two samples will be described with the example in the next subsection.

vi. Compute Confidence Interval for $\mu_1 - \mu_2$, σ_1^2 and σ_2^2 Known with SymIntegration

A study was conducted in which two types of engines, A and B , were compared. Gas mileage, in miles per gallon, was measured. Fifty experiments were conducted using engine type A and 75 experiments were done with engine type B . The gasoline used and other conditions were held constant. The average gas mileage was 36 miles per gallon for engine A and 42 miles per gallon for engine B . Find a 96% confidence interval on $\mu_B - \mu_A$, where μ_A and μ_B are population mean gas mileages for engines A and B , respectively. Assume that the population standard deviations are 6 and 8 for engines A and B , respectively.

Solution:

The point estimate of $\mu_B - \mu_A$ is $\bar{x}_B - \bar{x}_A = 42 - 36 = 6$. Using $\alpha = 0.04$, we find

$$z_{0.02} = 2.05$$

Hence, with substitution in the formula, the 96% confidence interval is

$$6 - 2.05\sqrt{\frac{64}{75} + \frac{36}{50}} < \mu_B - \mu_A < 6 + 2.05\sqrt{\frac{64}{75} + \frac{36}{50}}$$

or simply $3.43 < \mu_B - \mu_A < 8.57$.

In SymIntegration to compute the confidence interval for $\mu_1 - \mu_2$, σ_1^2 and σ_2^2 known we will use this function:

confidenceinterval_knownsigma(vector<double> data1, vector<double> data2, double σ_1 , double σ_2 , double α)

The input is vector **data1** (corresponding to X_1 , a random variable that represents the gas mileage for engine A), vector **data2** (corresponding to X_2 , a random variable that represents the gas mileage for engine B), the known population standard deviation σ_1 (corresponding to **data1**) and σ_2 (corresponding to **data2**), and the level of significance α to create the $100(1 - \alpha)\%$ confidence interval for $\mu_1 - \mu_2$.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Variances Known ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Variances Known ]# ./main

*****

Confidence interval with population variances known
Mean data 1 = 36, n1 = 50
Mean data 2 = 42, n2 = 75

The 96% confidence interval is :

3.42393 <  $\mu_1 - \mu_2$  < 8.57607

*****

Time taken by function: 830 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Variances Known ]#
```

Figure 4.15: The computation of confidence interval for $\mu_1 - \mu_2$, σ_1^2 and σ_2^2 known with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval Variances Known/main.cpp).

vii. Compute Confidence Interval for $\mu_1 - \mu_2$, $\sigma_1^2 = \sigma_2^2$ but Both Unknown with SymIntegration

The article "Macroinvertebrate Community Structure as an Indicator of Acid Mine Pollution," published in the *Journal of Environmental Pollution*, reports on an investigation undertaken in Cane Creek, Alabama, to determine the relationship between selected physiochemical parameters and different measures of macroinvertebrate community structure. One facet of the investigation was an evaluation of the effectiveness of a numerical species diversity index to indicate aquatic degradation due to acid mine drainage. Conceptually, a high index of macroinvertebrate species diversity should indicate an unstressed aquatic system, while a low diversity index should indicate a stressed aquatic system.

Two independent sampling stations were chosen for this study, one located downstream from the acid mine discharge point and the other located upstream. For 12 monthly samples collected at the downstream station, the species diversity index had a mean value $\bar{x}_1 = 3.11$ and a standard deviation $s_1 = 0.771$, while 10 monthly samples collected at the upstream station had a mean index value $\bar{x}_2 = 2.04$ and a standard deviation $s_2 = 0.448$. Find a 90% confidence interval for the difference between the population means for the two locations, assuming that the populations are approximately normally distributed with equal variances.

Solution:

Let μ_1 and μ_2 represent the population means, respectively, for the species diversity indices at the downstream and upstream stations. We wish to find a 90% confidence interval for $\mu_1 - \mu_2$. Our point estimate of $\mu_1 - \mu_2$ is

$$\bar{x}_1 - \bar{x}_2 = 3.11 - 2.04 = 1.07$$

The pooled estimate, s_p^2 , of the common variance, σ^2 , is

$$\begin{aligned} s_p^2 &= \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2} \\ &= \frac{(11)(0.771^2) + (9)(0.448^2)}{12 + 10 - 2} \\ &= 0.417 \end{aligned}$$

Taking the square root, we obtain $s_p = 0.646$. Using $\alpha = 0.1$, we obtain

$$t_{0.05} = 1.725$$

for $\nu = n_1 + n_2 - 2 = 20$ degrees of freedom. Therefore, the 90% confidence interval for $\mu_1 - \mu_2$ is

$$1.07 - (1.725)(0.646)\sqrt{\frac{1}{12} + \frac{1}{10}} < \mu_1 - \mu_2 < 1.07 + (1.725)(0.646)\sqrt{\frac{1}{12} + \frac{1}{10}}$$

which simplifies to $0.593 < \mu_1 - \mu_2 < 1.547$.

In SymIntegration to compute the confidence interval for $\mu_1 - \mu_2$, with $\sigma_1^2 = \sigma_2^2$ but both population variances are unknown we will use this function:

`confidenceinterval_sameunknownsigma(vector<double> data1, vector<double> data2, double  $\alpha$ )`

The input is vector **data1** (corresponding to X_1 , a random variable that represents index of macroinvertebrate species diversity at the downstream location), vector **data2** (corresponding to X_2 , a random variable that represents index of macroinvertebrate species diversity at the upstream location), and the level of significance α to create the $100(1 - \alpha)\%$ confidence interval for $\mu_1 - \mu_2$.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Variances Same but Unknown ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Variances Same but Unknown ]# ./main

*****

Confidence interval with same population variances but both unknown
Mean data 1 = 3.11, n1 = 12
Mean data 2 = 2.04, n2 = 10

The 90% confidence interval is :

0.592974 <  $\mu_1 - \mu_2$  < 1.54703

*****

Time taken by function: 753 microseconds

```

Figure 4.16: The computation of confidence interval for $\mu_1 - \mu_2$, $\sigma_1^2 = \sigma_2^2$ but both are unknown with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval Variances Same but Unknown/main.cpp).

viii. Compute Confidence Interval for $\mu_1 - \mu_2$, $\sigma_1^2 \neq \sigma_2^2$ and Both Unknown with SymIntegration

A study was conducted by the Department of Zoology at the Virginia Tech to estimate the difference in the amounts of the chemical orthophosphorus measured at two different stations on the James River. Orthophosphorus was measured in milligrams per liter. Fifteen samples were collected from station 1, and 12 samples were collected from station 2. The 15 samples from station 1 had an average orthophosphorus content of 3.84 milligrams per liter and a standard deviation of 3.07 milligrams per liter, while the 12 samples from station 2 had an average content of 1.49 milligrams per liter and a standard deviation of 0.80 milligram per liter. Find a 95% confidence interval for the difference in the true average orthophosphorus contents at these two stations, assuming that the observations came from normal populations with different variances.

Solution:

For station 1, we have $\bar{x}_1 = 3.84$, $s_1 = 3.07$, and $n_1 = 15$. For station 2, $\bar{x}_2 = 1.49$, $s_2 = 0.80$, and $n_2 = 12$. We wish to find a 95% confidence interval for $\mu_1 - \mu_2$. Since the population variances are assumed to be unequal, we can only find an approximate 95% confidence interval based on the t -distribution with ν degrees of freedom, where

$$\nu = \frac{\left(\frac{3.07^2}{15} + \frac{0.80^2}{12}\right)^2}{\left(\frac{\left(\frac{3.07^2}{15}\right)^2}{14}\right) + \left(\frac{\left(\frac{0.80^2}{12}\right)^2}{11}\right)} = 16.3 \approx 16$$

Our point estimate of $\mu_1 - \mu_2$ is

$$\bar{x}_1 - \bar{x}_2 = 3.84 - 1.49 = 2.35$$

Using $\alpha = 0.05$, we find that

$$t_{0.025} = 2.120$$

for $\nu = 16$ degrees of freedom. Therefore, the 95% confidence interval for $\mu_1 - \mu_2$ is

$$2.35 - 2.120\sqrt{\frac{3.07^2}{15} + \frac{0.80^2}{12}} < \mu_1 - \mu_2 < 2.35 + 2.120\sqrt{\frac{3.07^2}{15} + \frac{0.80^2}{12}}$$

which simplifies to $0.60 < \mu_1 - \mu_2 < 4.10$. Hence, we are confident that the interval from 0.60 to 4.10 milligrams per liter contains the difference of the true average orthophosphorus contents for these two locations.

In SymIntegration to compute the confidence interval for $\mu_1 - \mu_2$, with $\sigma_1^2 \neq \sigma_2^2$ and both population variances are unknown we will use this function:

confidenceinterval_unequalunknownsigma(vector<double> data1, vector<double> data2, double α)

The input is vector **data1** (corresponding to X_1 , a random variable that represents the amount of the chemical orthophosphorus measured at station 1), vector **data2** (corresponding to X_2 , a random variable that represents the amount of the chemical orthophosphorus measured at station 2), and the level of significance α to create the $100(1 - \alpha)\%$ confidence interval for $\mu_1 - \mu_2$.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Variances Unequal and Unknown ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Variances Unequal and Unknown ]# ./main

*****

Confidence interval with unequal and unknown population variances
Mean data 1 = 3.84, s1 = 3.07, n1 = 15
Mean data 2 = 1.49, s2 = 0.8, n2 = 12

The 95% confidence interval is :

0.599749 <  $\mu_1 - \mu_2$  < 4.10025

*****

Time taken by function: 781 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Variances Unequal and Unknown ]# 

```

Figure 4.17: The computation of confidence interval for $\mu_1 - \mu_2$, $\sigma_1^2 \neq \sigma_2^2$ and both are unknown with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval Variances Unequal and Unknown/main.cpp).

ix. Compute Confidence Interval for $\mu_D = \mu_1 - \mu_2$, Paired Observations with SymIntegration

Fortune magazine (March 1997) reported the total returns to investors for the 10 years prior to 1996 and also for 1996 for 431 companies. The total returns for 10 of the companies are listed below. Find a 95% confidence interval for the mean change in percent return to investors.

Company	Total Returns 1986 - 1996	Total Returns 1996
Coca-Cola	29.8%	43.3%
Mirage Resorts	27.9%	25.4%
Merck	22.1%	24.0%
Microsoft	44.5%	88.3%
Johnson & Johnson	22.2%	18.1%
Intel	43.8%	131.2%
Pfizer	21.7%	34.0%
Procter & Gamble	21.9%	32.1%
Berkshire Hathaway	28.3%	6.2%
S&P 500	11.8%	20.3%

Table 4.2: Total return to investors for 10 years prior to 1996 and also for 1996 for 10 companies.

Solution:

We wish to find 95% confidence interval for $\mu_1 - \mu_2$. Since the observations are paired, $\mu_1 - \mu_2 = \mu_D$. The point estimate of μ_D is $\bar{d} = 14.89$. The standard deviation, s_d , of the sample differences is

$$s_d = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (d_i - \bar{d})^2} = 30.4868$$

Using $\alpha = 0.05$, we find that $t_{0.025} = 2.262$ for $\nu = n - 1 = 9$ degrees of freedom. Therefore the 95% confidence interval is

$$14.89 - (2.262) \frac{30.4868}{\sqrt{10}} < \mu_D < 14.89 + (2.262) \frac{30.4868}{\sqrt{10}}$$

which yields $-6.92 < \mu_D < 36.70$.

In SymIntegration to compute the confidence interval for $\mu_D = \mu_1 - \mu_2$ paired observations we can use this function:

confidenceinterval_paired(vector<double> data1, vector<double> data2, double α)

The input is vector **data1** (corresponding to X_1 , a random variable that represents the "after" state), vector **data2** (corresponding to X_2 , a random variable that represents the "before" state), and the level of significance α to create the $100(1 - \alpha)\%$ confidence interval for $\mu_D = \mu_1 - \mu_2$.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Paired Observations ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Paired Observations ]# ./main

*****

Confidence interval for paired observations
Mean d = 14.89, sd = 30.4868, n = 10

The 95% confidence interval is :

-6.91892 <  $\mu_D$  < 36.6989

*****

Time taken by function: 742 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval Paired Observations ]# 

```

Figure 4.18: *The computation of confidence interval for $\mu_D = \mu_1 - \mu_2$ paired observations with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval Paired Observations/main.cpp).*

x. Compute Confidence Interval for Proportion p with SymIntegration

In a random sample of $n = 500$ families owning television sets in the city of Ontario, Canada, it is found that $x = 340$ subscribe to HBO, and the rest to BAM, a channel for dogs and by dogs all around the world. Find a 95% confidence interval for the actual proportion of families with television sets in this city that subscribe to HBO.

Solution:

The point estimate of p is

$$\hat{p} = \frac{340}{500} = 0.68$$

We find that

$$z_{0.025} = 1.96$$

Therefore, using method 1 from Definition (4.53), the 95% confidence interval for p is

$$0.68 - 1.96\sqrt{\frac{(0.68)(0.32)}{500}} < p < 0.68 + 1.96\sqrt{\frac{(0.68)(0.32)}{500}}$$

which simplifies to $0.6391 < p < 0.7209$.

In SymIntegration to compute the confidence interval for proportion p we can use this function:
confidenceinterval_proportion(vector<string> data, const string &inputString, double α)

The input is a string vector **data**, **inputString** as the string that we want to compute the confidence interval for the proportion p , and the level of significance α to create the $100(1 - \alpha)\%$ confidence interval for p .

In SymIntegration we are using method 2 from Definition (4.53) to compute the confidence interval for p , the exact results are only until 2 digit decimals for this example between method 1 and method 2. Note that although method 2 yields more accurate results, it is more complicated to calculate (not a problem if we are using computer), and the gain in accuracy that it provides diminishes when the sample size is large enough. Hence, method 1 is commonly used in practice, or college examination.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Proportion ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Proportion ]# ./main

*****

Confidence interval for proportion
Data size = 500
Amount of HBO in the data = 340
Proportion for HBO, p = 0.68, q = 0.32

The 95% confidence interval is :

0.637873 < p < 0.719382

*****

Time taken by function: 1128 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Proportion ]# 

```

Figure 4.19: *The computation of confidence interval for proportion p (the proportion of families that subscribe to HBO in Ontario, Canada) with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval for Proportion/main.cpp).*

xi. Compute Confidence Interval for the Difference Between Two Proportions $p_1 - p_2$ with SymIntegration

A certain change in a process for manufacturing component parts is being considered. Samples are taken under both the existing and the new process so as to determine if the new process results in an improvement. If 75 of 1500 items from the existing process are found to be defective and 80 of 2000 items from the new process are found to be defective, find a 90% confidence interval for the true difference in the proportion of defectives between the existing and the new process.

Solution:

Let p_1 and p_2 be the true proportions of defectives for the existing and new processes, respectively. Hence

$$\hat{p}_1 = \frac{75}{1500} = 0.05$$
$$\hat{p}_2 = \frac{80}{2000} = 0.04$$

and

$$z_{0.05} = 1.645$$

Therefore, substituting into the formula we will obtain

$$(0.05 - 0.04) - 1.645 \sqrt{\frac{(0.05)(0.95)}{1500} + \frac{(0.04)(0.96)}{2000}} < p_1 - p_2 < (0.05 - 0.04) + 1.645 \sqrt{\frac{(0.05)(0.95)}{1500} + \frac{(0.04)(0.96)}{2000}}$$

we find the 90% confidence interval to be $-0.0017 < p_1 - p_2 < 0.0217$. Since the interval contains the value 0, there is no reason to believe that the new process produces a significant decrease in the proportion of defectives over the existing method.

In SymIntegration to compute the confidence interval for the difference between two proportions $p_1 - p_2$ we can use this function:

confidenceinterval_differencebetweentwoproportions(vector<string> data1, vector<string> data2, const string &inputString, double α)

The input is a string vector **data1**, string vector **data2**, **inputString** as the string (e.g. Defective, Summa Cum laude, Lung Cancer, etc) that we want to compute the confidence interval for the proportion p , and the level of significance α to create the $100(1 - \alpha)\%$ confidence interval for $p_1 - p_2$.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Difference Between Proportions ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Difference Between Proportions ]# ./main
*****

Confidence interval for difference between two proportions
Data 1 size = 1500, Data 2 size = 2000
Amount of DEFECTIVE in the data 1 = 75
Amount of DEFECTIVE in the data 2 = 80
Proportion for DEFECTIVE, p1 = 0.05, q1 = 0.95
Proportion for DEFECTIVE, p2 = 0.04, q2 = 0.96

The 90% confidence interval is :

-0.00173124 < p1 - p2 < 0.0217312

*****

Time taken by function: 4008 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Difference Between Proportions ]# 

```

Figure 4.20: The computation of confidence interval for the difference between two proportions of the existing process and the new process in manufacturing component parts (*SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval for Difference Between Proportions/main.cpp*).

xii. Compute Confidence Interval for Variance σ^2 with SymIntegration

The following are the weights, in decagrams, of 10 packages of grass seed distributed by a certain company: 46.4, 46.1, 45.8, 47.0, 46.1, 45.9, 45.8, 46.9, 45.2, and 46.0. Find a 95% confidence interval for the variance of the weights of all such packages of grass seed distributed by this company, assuming a normal population.

Solution:

First we find

$$\begin{aligned}s^2 &= \frac{n \sum_{i=1}^n x_i^2 - (\sum_{i=1}^n x_i)^2}{n(n-1)} \\&= \frac{(10)(21,273.12) - (461.2)^2}{(10)(9)} \\&= 0.286\end{aligned}$$

To obtain a 95% confidence interval, we choose $\alpha = 0.05$. Then, using inverse cdf formula for chi-squared with $\nu = 9$ degrees of freedom, we find

$$\begin{aligned}\chi_{0.025}^2 &= 19.023 \\ \chi_{0.975}^2 &= 2.700\end{aligned}$$

The notation of $\chi_{0.025}^2$ means that after the $\chi_{0.025}^2$ value there is 0.025 area to the right, and an area of 0.975 to the left.

Therefore, the 95% confidence interval for σ^2 is

$$\frac{(9)(0.286)}{19.023} < \sigma^2 < \frac{(9)(0.286)}{2.700}$$

or simply $0.135 < \sigma^2 < 0.953$.

In SymIntegration to compute the confidence interval for variance σ^2 we can use this function:
confidenceinterval_variance(vector<doubles> data, double α)

The input is a double vector **data**, and the level of significance α to create the $100(1 - \alpha)\%$ confidence interval for σ^2 .

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Variance ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Variance ]# ./main

*****

Confidence interval for variance
Data size = 10
Mean sample = 46.12, variance sample = 0.286222

The 95% confidence interval is :

0.135411 <  $\sigma^2$  < 0.953856

*****

Time taken by function: 10225 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Variance ]# 

```

Figure 4.21: The computation of confidence interval for variance σ^2 of the weights of grass seed distributed by certain company with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval for Variance/main.cpp).

xiii. Compute Confidence Interval for Ratio of Two Variances $\frac{\sigma_1^2}{\sigma_2^2}$ with SymIntegration

A confidence interval for the difference in the mean orthophosphorus contents, measured in milligrams per liter, at two stations on the James River was constructed in the previous computation example by assuming the normal population variance to be unequal. Justify this assumption by constructing 98% for $\frac{\sigma_1^2}{\sigma_2^2}$ and for $\frac{\sigma_1}{\sigma_2}$, where σ_1^2 and σ_2^2 are the variances of the populations of orthophosphorus contents at station 1 and station 2, respectively.

Solution:

We will use new dummy data as **station1.txt** and **station2.txt** with $n_1 = 15$, $n_2 = 12$, $s_1 = 3.07479$ and $s_2 = 0.805983$.

For a 98% confidence interval, $\alpha = 0.02$, and we will obtain

$$\begin{aligned} f_{0.01}(14, 11) &\approx 4.30 \\ f_{0.01}(11, 14) &\approx 3.87 \end{aligned}$$

Therefore, the 98% confidence interval for $\frac{\sigma_1^2}{\sigma_2^2}$ is

$$\left(\frac{3.07479^2}{0.805983^2} \right) \left(\frac{1}{4.30} \right) < \frac{\sigma_1^2}{\sigma_2^2} < \left(\frac{3.07479^2}{0.805983^2} \right) (3.87) \quad (3.87)$$

which simplifies to $3.38995 < \frac{\sigma_1^2}{\sigma_2^2} < 56.991$. Taking square roots of the confidence limits, we find that a 98% confidence interval for $\frac{\sigma_1}{\sigma_2}$ is

$$1.84118 < \frac{\sigma_1}{\sigma_2} < 7.49912$$

Since this interval does not allow for the possibility of $\frac{\sigma_1}{\sigma_2}$ being equal to 1, we were correct in assuming that $\sigma_1 \neq \sigma_2$ or $\sigma_1^2 \neq \sigma_2^2$.

In SymIntegration to compute the confidence interval for the ratio of two variances $\frac{\sigma_1^2}{\sigma_2^2}$ we can use this function:

confidenceinterval_ratiotwovariances(vector<doubles> data1, vector<doubles> data2, double α)

The input is the first data as a double vector **data1**, the second data as a double vector **data2**, and the level of significance α to create the $100(1 - \alpha)\%$ confidence interval for σ^2 .

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Ratio of Two Variances ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Ratio of Two Variances ]# ./main

*****

Confidence interval for the ratio of two variances
Mean data 1 = 4.68074, s1 = 3.07479, n1 = 15
Mean data 2 = 1.14591, s2 = 0.805983, n2 = 12

The 98% confidence interval is :

3.38995 <  $\sigma_1^2 / \sigma_2^2$  < 56.2368
1.84118 <  $\sigma_1 / \sigma_2$  < 7.49912

*****

Time taken by function: 938 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Confidence Interval for Ratio of Two Variances ]# 

```

Figure 4.22: The computation of confidence interval for ratio of two variances $\frac{\sigma_1^2}{\sigma_2^2}$ and the ration of two standard deviations $\frac{\sigma_1}{\sigma_2}$ of the populations of orthophosphorus contents in two different stations with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Confidence Interval for Ratio of Two Variances/main.cpp).

xiv. Maximum Likelihood Estimation

[SI*] Maximum likelihood estimation (MLE) is one of the most important approaches to estimation in all of statistical inference. It is a statistical method for estimating the parameters of a probability distribution by finding the parameter values that make the observed data most probable.

To do that, it maximizes the likelihood function, which represents the probability of the observed data given the parameters. The maximum likelihood estimate is the parameter value that results in this highest probability.

[SI*] Some key concepts in MLE:

1. **Likelihood function**

This function is essentially the joint probability density function (pdf) of the observed data, viewed as a function of the distribution's parameters.

2. **Maximization**

The goal is to find the parameter value(s) that maximize the likelihood function. This is often done by taking the natural logarithm of the likelihood function, which simplifies calculations because products become sums.

3. **Log-likelihood**

The logarithm is taken because it is often easier to differentiate. Maximizing the log-likelihood is equivalent to maximizing the likelihood itself.

4. **Estimator**

The process involves taking the derivative of the log-likelihood function, setting it to zero, and solving for the parameter(s) to find the maximum.

[SI*] The applications of MLE:

1. **Statistics and machine learning**

MLE is a powerful and widely used technique in both statistics and machine learning for optimizing models.

2. **Regression**

For example, the least-squares method used in linear regression produces the same result as MLE under certain assumptions.

3. **Model fitting**

It helps find the best-fitting distribution to a set of data by finding the parameters that make that data most likely to have occurred.

Definition 4.57: Maximum Likelihood Estimator

Given independent observations $x_1, x_2, \dots, x_n$ from a probability density function (continuous case) or probability mass function (discrete case) $f(x; \theta)$, the maximum likelihood estimator $\hat{\theta}$ is that which maximizes the likelihood function

$$L(x_1, x_2, \dots, x_n; \theta) = f(x; \theta) = f(x_1, \theta)f(x_2, \theta) \dots f(x_n, \theta)$$

[SI*] **Example:**

Consider a Poisson distribution with probability mass function

$$f(x|\mu) = \frac{e^{-\mu}\mu^x}{x!}, \quad x = 0, 1, 2, \dots$$

Suppose that a random sample $x_1, x_2, \dots, x_n$ is taken from the distribution. What is the maximum likelihood estimate of μ ?

Solution:

The likelihood function is

$$L(x_1, x_2, \dots, x_n; \mu) = \prod_{i=1}^n f(x_i | \mu) = \frac{e^{-n\mu} \mu^{\sum_{i=1}^n x_i}}{\prod_{i=1}^n x_i!}$$

Now consider

$$\begin{aligned} \ln L(x_1, x_2, \dots, x_n; \mu) &= -n\mu + \sum_{i=1}^n x_i \ln \mu - \ln \prod_{i=1}^n x_i! \\ \frac{\partial \ln L(x_1, x_2, \dots, x_n; \mu)}{\partial \mu} &= -n + \sum_{i=1}^n \frac{x_i}{\mu} \end{aligned}$$

Solving for $\hat{\mu}$, the maximum likelihood estimator, involves setting the derivative to zero and solving for the parameter. Thus,

$$\hat{\mu} = \sum_{i=1}^n \frac{x_i}{n} = \bar{x}$$

the second derivative of the log-likelihood is negative, which implies that the solution above indeed is a maximum. Since μ is the mean of the Poisson distribution, the sample average would certainly seem like a reasonable estimator.

[SI*] **Example:**

Consider a random sample $x_1, x_2, \dots, x_n$ from a normal distribution $N(\mu, \sigma)$. Find the maximum likelihood estimators for μ and σ^2 .

Solution:

The likelihood function for the normal distribution is

$$L(x_1, x_2, \dots, x_n; \mu, \sigma^2) = \frac{1}{(2\pi)^{\frac{n}{2}} (\sigma^2)^{\frac{n}{2}}} e^{\left[-\frac{1}{2} \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma}\right)^2\right]}$$

taking logarithm gives us

$$\ln L(x_1, x_2, \dots, x_n; \mu, \sigma^2) = -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln \sigma^2 - \frac{1}{2} \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma}\right)^2$$

Hence,

$$\begin{aligned} \frac{\partial \ln L}{\partial \mu} &= \sum_{i=1}^n \left(\frac{x_i - \mu}{\sigma^2}\right) \\ \frac{\partial \ln L}{\partial \sigma^2} &= -\frac{n}{2\sigma^2} + \frac{1}{2(\sigma^2)^2} \sum_{i=1}^n (x_i - \mu)^2 \end{aligned}$$

Setting both derivatives equal to 0, we obtain

$$\sum_{i=1}^n x_i - n\mu = 0$$

and

$$n\sigma^2 = \sum_{i=1}^n (x_i - \mu)^2$$

Thus, the maximum likelihood estimator of μ is given by

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n x_i = \bar{x}$$

On the other hand, the maximum likelihood estimator of σ^2 is

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$$

Checking the second-order partial derivative matrix confirms that the solution results in a maximum of the likelihood function. It is interesting to note the distinction between the maximum likelihood estimator of σ^2 and the unbiased estimator S^2 . The numerators are identical, of course, and the denominator is the degrees of freedom $n - 1$ for the unbiased estimator and n for the maximum likelihood estimator. Maximum likelihood estimators do not necessarily enjoy the property of unbiasedness. However, they do have very important asymptotic properties.

[SI*] The method of maximum likelihood cannot be applied without knowledge of the underlying distribution.

[SI*] **MLE for Binomial Distribution**

Suppose that there are n trials $x_1, x_2, \dots, x_n$ from a Bernoulli process with parameter p , the probability of a success. That is, the probability of r successes is given by

$$\binom{n}{r} p^r (1-p)^{n-r}$$

work out the maximum likelihood estimator for the parameter p .

Solution:

The likelihood function is

$$\begin{aligned} L(x_1, x_2, \dots, x_n; p) &= \prod_{i=1}^n f(x_i; p) \\ &= \prod_{i=1}^n p^{x_i} (1-p)^{1-x_i} \\ &= p^{n\bar{x}} (1-p)^{n(1-\bar{x})} \end{aligned}$$

Hence,

$$\ln L = n [\bar{x} \ln(p) + (1 - \bar{x}) \ln(1 - p)]$$

taking derivative with respect to p and setting the derivative to zero, we obtain

$$\frac{\partial \ln(L)}{\partial p} = n \left(\frac{\bar{x}}{p} - \frac{1 - \bar{x}}{1 - p} \right) = 0$$

which yields

$$\frac{\bar{x}}{p} - \frac{1 - \bar{x}}{1 - p} = 0$$

Therefore

$$\hat{p} = \bar{x}$$

[SI*] **MLE for Negative Binomial Distribution**

Consider the observation X from the negative binomial distribution

$$b^*(x; k, p) = \binom{x-1}{k-1} p^k (1-p)^{x-k}, \quad x = k, k+1, k+2, \dots$$

Find the maximum likelihood estimator for p , assuming k is known.

Solution:

From the density function $b^*(x; p) = \binom{x-1}{k-1} p^k (1-p)^{x-k}$, we obtain

$$\ln L = \ln \binom{x-1}{k-1} + k \ln p + (n-k) \ln(1-p)$$

Setting

$$\frac{\partial \ln L}{\partial p} = \frac{k}{p} - \frac{n-k}{1-p} = 0$$

we obtain

$$\hat{p} = \frac{k}{n}$$

[SI*] **MLE for Uniform Distribution**

Consider a random sample of $x_1, \dots, x_n$ from a uniform distribution $U(0, \theta)$ with unknown parameter θ , where $\theta > 0$. Determine the maximum likelihood estimator of θ .

Solution:

The likelihood function of θ is

$$L(x_1, \dots, x_n; \theta) = \frac{1}{\theta^n}, \text{ for } \max\{x_1, \dots, x_n\} < \theta < \infty$$

Since $\frac{1}{\theta^n}$ is a decreasing function of θ , the maximum only achieves at the lower bound, which is $\max\{x_1, \dots, x_n\}$. This is called the largest order statistic of the sample.

[SI*] **MLE for Lognormal Distribution**

Consider the lognormal distribution with the density function

$$f(x; \mu, \sigma) = \begin{cases} \frac{1}{\sqrt{2\pi}\sigma x} e^{-\frac{1}{2\sigma^2} [\ln(x)-\mu]^2}, & x \geq 0 \\ 0, & x < 0 \end{cases}$$

Suppose we have a random sample $x_1, x_2, \dots, x_n$ from a lognormal distribution. Write out the likelihood function then develop the maximum likelihood estimators of μ and σ^2 .

Solution:

The likelihood function is

$$\begin{aligned} L(x_1, x_2, \dots, x_n; \mu, \sigma) &= \prod_{i=1}^n f(x_i; \mu, \sigma) \\ &= \prod_{i=1}^n \left[\frac{1}{\sqrt{2\pi\sigma} x_i} e^{-\frac{[\ln(x_i) - \mu]^2}{2\sigma^2}} \right] \\ &= \frac{1}{(2\pi)^{\frac{n}{2}} \sigma^n \prod_{i=1}^n x_i} e^{-\frac{1}{2\sigma^2} \sum_{i=1}^n [\ln(x_i) - \mu]^2} \end{aligned}$$

It is easy to obtain

$$\ln(L) = -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln(\sigma^2) - \sum_{i=1}^n \ln(x_i) - \frac{1}{2\sigma^2} \sum_{i=1}^n [\ln(x_i) - \mu]^2$$

So, setting

$$\frac{\partial \ln L}{\partial \mu} = \frac{1}{\sigma^2} \sum_{i=1}^n [\ln(x_i) - \mu] = 0$$

we obtain

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n \ln(x_i)$$

and setting

$$\frac{\partial \ln L}{\partial \sigma^2} = -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^n [\ln(x_i) - \mu]^2 = 0$$

we get

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n [\ln(x_i) - \hat{\mu}]^2$$

[SI*] MLE for Gamma Distribution

Consider a random sample of $x_1, x_2, \dots, x_n$ coming from the gamma distribution with density function

$$f(x; \alpha, \beta) = \begin{cases} \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\beta}}, & x > 0 \\ 0, & \text{elsewhere} \end{cases}$$

for $\alpha, \beta > 0$.

Determine the maximum likelihood estimation for parameters α and β .

Solution:

The likelihood function is

$$\begin{aligned} L(x_1, \dots, x_n; \alpha, \beta) &= \frac{1}{\beta^{\alpha n} \Gamma(\alpha)^n} \prod_{i=1}^n \left(x_i^{\alpha-1} e^{-\frac{x_i}{\beta}} \right) \\ &= \frac{1}{\beta^{\alpha n} \Gamma(\alpha)^n} \left(\prod_{i=1}^n x_i \right)^{\alpha-1} e^{-\sum_{i=1}^n \left(\frac{x_i}{\beta} \right)} \end{aligned}$$

The log-likelihood can be expressed as

$$\begin{aligned} \ln L &= -n\alpha \ln(\beta) - n \ln(\Gamma(\alpha)) + (\alpha - 1) \sum_{i=1}^n \ln(x_i) - \frac{1}{\beta} \sum_{i=1}^n x_i \\ &= C(\underline{x}) - \alpha n \ln \beta - \frac{n\bar{x}}{\beta} \end{aligned}$$

where $C(\underline{x})$ is a function of $\underline{x}$. Taking logarithm of $\ln L$ and setting it to be zero, we obtain the maximum likelihood estimation for β to be

$$\hat{\beta} = \frac{\bar{x}}{\alpha}$$

[SI*] **MLE for Weibull Distribution**

Consider a random sample of $x_1, x_2, \dots, x_n$ observations from a Weibull distribution with parameters α and β and density function

$$f(x) = \begin{cases} \alpha \beta x^{\beta-1} e^{-\alpha x^\beta}, & x > 0 \\ 0, & \text{elsewhere} \end{cases}$$

for $\alpha, \beta > 0$.

Determine the maximum likelihood estimation for parameters α and β .

Solution:

The likelihood function is

$$\begin{aligned} L(x_1, \dots, x_n; \alpha, \beta) &= \prod_{i=1}^n f(x_i; \alpha, \beta) = (\alpha \beta)^n \prod_{i=1}^n x_i^{\beta-1} e^{-\alpha x_i^\beta} \\ &= (\alpha \beta)^n e^{-\alpha \sum_{i=1}^n x_i^\beta} \left(\prod_{i=1}^n x_i \right)^{\beta-1} \end{aligned}$$

So, the log-likelihood can be expressed as

$$\ln L = n[\ln(\alpha) + \ln(\beta)] - \alpha \sum_{i=1}^n x_i^\beta + (\beta - 1) \sum_{i=1}^n \ln(x_i)$$

To solve for the maximum likelihood estimate, we need to solve the following two equations

$$\begin{aligned} \frac{\partial \ln L}{\partial \alpha} &= 0 \\ \frac{\partial \ln L}{\partial \beta} &= 0 \end{aligned}$$

xv. Compute Maximum Likelihood Estimation for Continuous Distribution

It is known that a sample consisting of the values 12, 11.2, 13.5, 12.3, 13.8, and 11.9 comes from a population with the density function

$$f(x; \theta) = \begin{cases} \frac{\theta}{x^{\theta+1}}, & x > 1 \\ 0, & \text{elsewhere} \end{cases}$$

where $\theta > 0$. Find the maximum likelihood estimate of θ .

Solution:

The likelihood function of n observations from this population can be written as

$$L(x_1, x_2, \dots, x_n; \theta) = \prod_{i=1}^n \frac{\theta}{x_i^{\theta+1}} = \frac{\theta^n}{(\prod_{i=1}^n x_i)^{\theta+1}}$$

which implies that

$$\ln L(x_1, x_2, \dots, x_n; \theta) = n \ln(\theta) - (\theta + 1) \sum_{i=1}^n \ln(x_i)$$

Setting $0 = \frac{\partial \ln L}{\partial \theta} = \frac{n}{\theta} - \sum_{i=1}^n \ln(x_i)$ results in

$$\begin{aligned} \hat{\theta} &= \frac{n}{\sum_{i=1}^n \ln(x_i)} \\ &= \frac{6}{\ln(12) + \ln(11.2) + \ln(13.5) + \ln(12.3) + \ln(13.8) + \ln(11.9)} \\ &= 0.3970 \end{aligned}$$

Since the second derivative of L is $-\frac{n}{\theta^2}$ which is always negative, the likelihood function does achieve its maximum value at $\hat{\theta}$.

In `SymIntegration`, we can compute the mle for any distribution given that we provide the pdf or the pmf, it is assumed that the parameters for the distribution is at least one and at maximum two. Normal distribution has two parameters: μ and σ^2 , Gamma distribution has two parameters: α and β , exponential distribution has one parameter $\lambda = \frac{1}{\beta}$.

The function that is used in `SymIntegration` to compute the maximum likelihood estimation is: **mle(const Symbolic &f(x; μ, σ^2), const Symbolic &x, const Symbolic & μ , const Symbolic & σ^2 , int n)**

The first input is the Symbolic function of the pdf or pmf, we are using a normal distribution as the example above, then the random variable x , and the first parameter μ , then the second parameter σ^2 , and last the number of samples.

For this case, we can put any symbolic for the second parameter (e.g. s, γ, ρ , etc) because the pdf does not have the second parameter, thus it won't really matter with the end result what symbolic second parameter that we put into the function **mle**.

If we want a numeric answer then we will make use of **Equations rules** and then implement the **.subst_all(rules)** into the **mle** result.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Maximum Likelihood Estimation ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Maximum Likelihood Estimation ]# ./main
f(x;θ) = θ*x^(-θ-1)
estimated θ = 6*(ln(x[1])+ln(x[2])+ln(x[3])+ln(x[4])+ln(x[5])+ln(x[6]))^(-1)
estimated θ = 0.396975

Time taken by function: 42617 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Maximum Likelihood Estimation ]# 

```

Figure 4.23: The computation of maximum likelihood estimation for parameter θ with the density function $f(x; \theta) = \frac{\theta}{x^{\theta+1}}, x > 1$ (*SymIntegration/Examples/Statistics/Test SymIntegration Maximum Likelihood Estimation/main.cpp*).

The source code that handle this maximum likelihood estimation is located in **src/statistics-mle.cpp** and **include/statistics-mle.h**

xvi. Bayesian Statistics

[S*]

VI. HYPOTHESIS TESTING

i. One Sample Hypothesis Test for Comparing Means

[SI*] Hypothesis testing is a statistical process used to make a decision about a population based on sample data. It involves creating a null hypothesis (a statement of no effect or no difference) and an alternative hypothesis, then using sample evidence and probability to test the null hypothesis. The goal is to determine if there is enough evidence to reject the null hypothesis in favor of the alternative.

[SI*] How Hypothesis Testing Works

1. **State the hypotheses:** Formulate a null hypothesis (H_0) and an alternative hypothesis (H_1).
 - Null Hypothesis (H_0): A specific statement of no significant difference between a hypothesized value and the value estimated from a sample (e.g., a new drug has no effect).
 - Alternative Hypothesis (H_1): A statement that contradicts the null hypothesis (e.g., the new drug does have an effect).
2. **Formulate an analysis plan:** Choose a statistical test and define a significance level (α). The choice of test depends on the nature of the data and the question being asked.

The test statistic measures how much the sample data deviates from what we did expect if the null hypothesis were true. Different tests use different statistics:

- (a) Z-test: Used when population variance is known and sample size is large.
 - (b) Student's t -test: Used when the sample size is small or population variance unknown.
 - (c) Chi-Squared test: Used for categorical data to compare observed vs expected counts.
3. **Analyze the sample data:** Collect a sample data and perform the chosen statistical test. This involves calculating a test statistic and a P -value.

4. Interpret the results:

(a) Using P -value:

Compare the P -value to the significance level (α) to decide whether to reject or fail to reject the null hypothesis.

- If the P -value is less than the significance level (α), you reject the null hypothesis.
- If the P -value is greater than or equal to the significance level, you fail to reject the null hypothesis.

The P -value approach has been adopted extensively by users of applied statistics. The approach is designed to give the user an alternative to a mere "reject" or "do not reject" conclusion. The P -value computation also gives the user important information when the z value falls well into the ordinary critical region.

(b) Using Critical Value:

We compute the test statistic, for example if we know that the sample size is more than 30 or we know the population variance then we use z statistic (otherwise, we use t statistic), thus we use this formula:

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$

For the case of hypothesis testing with

$$H_0 : \mu = \mu_0$$

$$H_1 : \mu < \mu_0$$

If the test statistic $z < a$ then reject H_0 , if $z \geq a$ then we do not reject H_0 . With a as the critical value

$$a = \mu_0 - z_\alpha \frac{\sigma}{\sqrt{n}}$$

For the case of hypothesis testing with

$$H_0 : \mu = \mu_0$$

$$H_1 : \mu > \mu_0$$

If the test statistic $z > b$ then reject H_0 , if $z \leq b$ then we do not reject H_0 . With b as the critical value

$$b = \mu_0 + z_\alpha \frac{\sigma}{\sqrt{n}}$$

For the case of hypothesis testing with

$$H_0 : \mu = \mu_0$$

$$H_1 : \mu \neq \mu_0$$

If the test statistic $z < a$ or $z > b$ then reject H_0 , if $z \geq a$ then we do not reject H_0 . With a and b as the critical value

$$a = \mu_0 - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

$$b = \mu_0 + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

[SI*] Examples that are using hypothesis testing:

- Testing a new drug by comparing a group that receives the drug to a control group to see if there is a statistically significant difference in outcomes.
- Assessing if a change in a production process has resulted in a statistical difference in product yield.

[SI*] Types of hypothesis testing:

1. Two-tailed

$$H_0 : \mu = c$$

$$H_1 : \mu \neq c$$

If the hypothesis test is two-tailed, with variance known and large sample size then the P -value can be computed with:

$$P = P(|Z| > z_{\frac{\alpha}{2}}) = 2P(Z < -z_{\frac{\alpha}{2}})$$

with

$$z_{\frac{\alpha}{2}} = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$

2. One-tailed (Left-tailed)

$$H_0 : \mu \geq c$$

$$H_1 : \mu < c$$

If the hypothesis test is one-tailed (left-tailed), with variance known and large sample size then the P -value can be computed with:

$$P = P(Z < z_{\frac{\alpha}{2}})$$

with

$$z_{\frac{\alpha}{2}} = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$

3. One-tailed (Right-tailed)

$$H_0 : \mu \leq c$$

$$H_1 : \mu > c$$

If the hypothesis test is one-tailed (right-tailed), with variance known and large sample size then the P -value can be computed with:

$$P = P(Z > z_{\frac{\alpha}{2}})$$

with

$$z_{\frac{\alpha}{2}} = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$

[SI*] We will explore hypothesis testing through a few specific practical hypothesis tests. Recall the general hypothesis test formulation where we partition the parameter space Θ as follows:

$$H_0 : \theta \in \Theta_0, \quad H_1 : \theta \in \Theta_1$$

One of the most common cases for a single population is to consider θ as μ , the population mean, in which case $\Theta = \mathbb{R}$. Often, we wish to test if the population mean is equal to some value, μ_0 . This allows us to construct a two-sided hypothesis test as follows:

$$H_0 : \mu = \mu_0, \quad H_1 : \mu \neq \mu_0 \quad (4.111)$$

However, one could instead chose to construct a one-sided hypothesis test, as

$$H_0 : \mu \leq \mu_0, \quad H_1 : \mu > \mu_0 \quad (4.112)$$

or alternatively, in the opposite direction

$$H_0 : \mu \geq \mu_0, \quad H_1 : \mu < \mu_0 \quad (4.113)$$

the choice of setting up which one out of these three hypothesis tests, depend on the context of the problem.

Once the hypothesis is established, the general approach involves calculating the test statistic, along with the corresponding p -value, and then finally making some statement about the null hypothesis based on some chosen level of significance.

[SI*] Possible situations for testing a statistical hypothesis

	H_0 is true	H_0 is false
Do not reject H_0	Correct decision	Type II error
Reject H_0	Type I error	Correct decision

Table 4.3

[SI*] Statistical Hypothesis with Discrete Data

A certain type of cold vaccine is known to be only 25% effective after a period of 2 years. To determine if a new and somewhat more expensive vaccine is superior in providing protection against the same virus for a longer period of time, suppose that 20 people are chosen at random and inoculated. If more than 8 of those receiving the new vaccine surpass the 2-year period without contracting the virus, the new vaccine will be considered superior to the one presently in use. We are essentially testing the null hypothesis that the new vaccine is equally effective after a period of 2 years as the one now commonly used. The alternative hypothesis is that the new vaccine is in fact superior. The hypothesis testing is usually written as follows

$$H_0 : p = 0.25$$

$$H_1 : p > 0.25$$

The test statistic on which we base our decision is X , the number of individuals in our test group who receive protection from the new vaccine for a period of at least 2 years. The possible values of X , from 0 to 20, are divided into two groups: those numbers less than or equal to 8 and those greater than 8. All possible scores greater than 8 constitute the critical region. The last number that we observe in passing into the critical region is called the critical value.

Therefore, if $x > 8$, we reject H_0 in favor of the alternative hypothesis H_1 . If $x \leq 8$, we fail to reject H_0 .

1. The Probability of Type I Error

The probability of committing a type I error, also called the level of significance, is denoted by the Greek letter α .

In our illustration, a type I error will occur when more than 8 individuals inoculated with the new vaccine surpass the 2-year period without contracting the virus and researchers conclude that the new vaccine is better when it is actually equivalent to the one in use. Hence, if X is the number of individuals who remain free of the virus for at least 2 years,

$$\begin{aligned}
 \alpha &= P(\text{type I error}) \\
 &= P(X > 8, p = 0.25) \\
 &= \sum_{x=9}^{20} b(x; 20, 0.25) \\
 &= 1 - \sum_{x=0}^8 b(x; 20, 0.25) \\
 &= 1 - 0.9591 \\
 &= 0.0409
 \end{aligned}$$

We say that the null hypothesis, $p = 0.25$, is being tested at the $\alpha = 0.0409$ level of

significance. Sometimes the level of significance is called the size of the test. A critical region of size 0.0409 is very small, and therefore it is unlikely that a type I error will be committed. Consequently, it would be most unusual for more than 8 individuals to remain immune to a virus for a 2-year period using a new vaccine that is essentially equivalent to the one now on the market.

2. The Probability of Type II Error

The probability of committing a type II error, denoted by β , is impossible to compute unless we have a specific alternative hypothesis. If we test the null hypothesis that $p = 0.25$ against the alternative hypothesis that $p = 0.5$, then we are able to compute the probability of not rejecting H_0 when it is false.

We simply find the probability of obtaining 8 or fewer in the group to surpass the 2-year period when $p = 0.5$. In this case,

$$\begin{aligned}\beta &= P(\text{type II error}) \\ &= P(X \leq 8, p = 0.5) \\ &= \sum_{x=0}^8 b(x; 20, 0.5) \\ &= 0.2517\end{aligned}$$

This is a rather high probability, indicating a test procedure in which it is quite likely that we shall reject the new vaccine when, in fact, it is superior to what is now in use. Ideally, we like to use a test procedure for which the type I and type II error probabilities are both small.

The probability of committing both types of error can be reduced by increasing the sample size.

[SI*] Statistical Hypothesis with Continuous Data

Consider the null hypothesis that the average weight of male students in a certain college is 68 kilograms against the alternative hypothesis that is unequal to 68. That is, we wish to test

$$\begin{aligned}H_0 : \mu &= 68 \\ H_1 : \mu &\neq 68\end{aligned}$$

The alternative hypothesis allows for the possibility that $\mu < 68$ or $\mu > 68$.

Reject H_0 ($\mu \neq 68$)	Do not reject H_0 ($\mu = 68$)	Reject H_0 ($\mu \neq 68$)
--------------------------------	------------------------------------	--------------------------------

Table 4.4: Critical region (in light cyan)

Assume the standard deviation of the population of weights to be $\sigma = 3.6$. For large samples, we may substitute s for σ if no other estimate of σ is available. Our decision statistic, based on a random sample of size $n = 36$, will be $\bar{X}$, the most efficient estimator of μ . From the Central Limit Theorem, we know that the sampling distribution of $\bar{X}$ is approximately normal with standard deviation

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}} = \frac{3.6}{6} = 0.6$$

1. The Probability of Type I Error

The probability of committing a type I error, or the level of significance of our test, is equal to the sum of the areas that have been shaded in each tail of the distribution.




Figure 4.24: Critical region for testing $\mu = 68$ versus $\mu \neq 68$.

Therefore

$$\alpha = P(\bar{X} < 67, \mu = 68) + P(\bar{X} > 69, \mu = 68)$$

The z-values corresponding to $\bar{x}_1 = 67$ and $\bar{x}_2 = 69$ when H_0 is true are

$$z_1 = \frac{67 - 68}{0.6} = -1.67$$

$$z_2 = \frac{69 - 68}{0.6} = 1.67$$

Therefore,

$$\begin{aligned}\alpha &= P(Z < -1.67) + P(Z > 1.67) \\ &= 2P(Z < -1.67) \\ &= 0.0950\end{aligned}$$

The 9.5% of all samples of size 36 would lead us to reject $\mu = 68$ kilograms when, in fact, it is true. To reduce α , we have a choice of increasing the sample size or widening the fail-to-reject region.

Suppose that we increase the sample size to $n = 64$. Then $\sigma_{\bar{X}} = \frac{3.6}{8} = 0.45$. Now

$$z_1 = \frac{67 - 68}{0.45} = -2.22$$

$$z_2 = \frac{69 - 68}{0.45} = 2.22$$

Hence,

$$\begin{aligned}\alpha &= P(Z < -2.22) + P(Z > 2.22) \\ &= 2P(Z < -2.22) \\ &= 0.0264\end{aligned}$$

The reduction in α is not sufficient by itself to guarantee a good testing procedure. We must also evaluate β for various alternative hypotheses.

2. The Probability of Type II Error

If it is important to reject H_0 when the true mean is some value $\mu \geq 70$ or $\mu \leq 66$, then the probability of committing a type II error should be computed and examined for the alternatives $\mu = 66$ and $\mu = 70$. Because of symmetry, it is only necessary to consider the probability of not rejecting the null hypothesis that $\mu = 68$ when the alternative $\mu = 70$ is true. A type II error will result when the sample mean $\bar{x}$ falls between 67 and 69 when H_1 is true. Therefore, we find that

$$\beta = P(67 \leq \bar{X} \leq 69, \mu = 70)$$




Figure 4.25: Probability of type II error for testing $\mu = 68$ versus $\mu = 70$.

The z-values corresponding to $\bar{x}_1 = 67$ and $\bar{x}_2 = 69$ when H_1 is true are

$$\begin{aligned} z_1 &= \frac{67 - 70}{0.45} = -6.67 \\ z_2 &= \frac{69 - 70}{0.45} = -2.22 \end{aligned}$$

Therefore

$$\begin{aligned} \beta &= P(-6.67 < Z < -2.22) \\ &= P(Z < -2.22) - P(Z < -6.67) \\ &= 0.0132 - 0.0000 \\ &= 0.0132 \end{aligned}$$

If the true value of μ is the alternative $\mu = 66$, the value of β will again be 0.0132. For all possible values of $\mu < 66$ or $\mu > 70$, the value of β will be even smaller when $n = 64$, and consequently there would be little chance of not rejecting H_0 when it is false.

Definition 4.58: Important Properties of a Test of Hypothesis

1. The type I error and type II error are related. A decrease in the probability of one generally results in an increase in the probability of the other.
2. The size of the critical region, and therefore the probability of committing a type I error, can always be reduced by adjusting the critical value(s).
3. An increase in the sample size n will reduce α and β simultaneously.
4. If the null hypothesis is false, β is a maximum when the true value of a parameter approaches the hypothesized value. The greater the distance between the true value and the hypothesized value, the smaller β will be.

Definition 4.59: Power of a Test

The power of a test is the probability of rejecting H_0 given that a specific alternative is true.

The power of a test can be computed as $1 - \beta$. Often different types of tests are compared by contrasting power properties. To produce a desirable power, say, greater than 0.8, one must either increase α or increase the sample size.

Definition 4.60: Test Procedure for a Single Mean (Variance Known)

For large sample size $n > 30$ and known population variance we will use Z statistics

$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} > z_{\alpha/2}$$

or

$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} < -z_{\alpha/2}$$

If $-z_{\alpha/2} < z < z_{\alpha/2}$, do not reject H_0 . Rejection of H_0 , of course, implies acceptance of the alternative hypothesis $\mu \neq \mu_0$. With this definition of the critical region, it should be clear that there will be probability α of rejecting H_0 (falling into the critical region) when, indeed, $\mu = \mu_0$.

[SI*] Although it is easier to understand the critical region written in terms of z , we can write the same critical region in terms of the computed average $\bar{x}$. The following can be written as an identical decision procedure:

Reject H_0 if $\bar{x} < a$ or $\bar{x} > b$, where

$$a = \mu_0 - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

$$b = \mu_0 + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$




Figure 4.26: Critical region for the alternative hypothesis $\mu \neq \mu_0$.

[SI*] Tests on one-sided hypotheses on the mean involve the same statistic described in the two-sided case. The difference, of course, is that the critical region is only in one tail of the standard normal distribution. For example, suppose that we seek to test

$$H_0 : \mu = \mu_0$$

$$H_1 : \mu > \mu_0$$

The signal that favors H_1 comes from large values of z . Thus, rejection of H_0 results when the computed $z > z_\alpha$. Obviously, if the alternative is $H_1 : \mu < \mu_0$, the critical region is entirely in the lower tail and thus rejection results from $z < -z_\alpha$. Although in a one-sided testing case the null hypothesis can be written as $H_0 : \mu \leq \mu_0$ or $H_0 : \mu \geq \mu_0$, it is usually written as $H_0 : \mu = \mu_0$.




Figure 4.27: Critical region for the alternative hypothesis $\mu > \mu_0$.

Definition 4.61: The t -Statistic for a Test on a Single Mean

For two-sided hypothesis with unknown population variance

$$H_0 : \mu = \mu_0$$

$$H_1 : \mu \neq \mu_0$$

We reject H_0 at significance level α when computed t -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

exceeds $t_{\frac{\alpha}{2}, n-1}$ or is less than $-t_{\frac{\alpha}{2}, n-1}$.

The sample variance formula is:

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

with

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

GlanzFreya' Guide 4.1: On Type I Error for Hypothesis Testing

Type I error has the same analogy when we make a verdict that an innocent person is guilty.

1. Significance Level (α)

The significance level is a predefined maximum probability of committing type I error we're willing to accept.

The lower the significance level is, the lower the risk of committing a type I error.

With the significance level, we can find the critical value to reject the null hypothesis in a hypothesis test. The critical value for right-tailed hypothesis test:

$$\text{Critical Value} = \mu + Z \left(\frac{\sigma}{\sqrt{n}} \right)$$

The critical value for left-tailed hypothesis test:

$$\text{Critical Value} = \mu - Z \left(\frac{\sigma}{\sqrt{n}} \right)$$

For two-tailed hypothesis test, we use both critical values above so the rejection area will be at $\bar{x} > \mu + Z \left(\frac{\sigma}{\sqrt{n}} \right)$ and at $\bar{x} < \mu - Z \left(\frac{\sigma}{\sqrt{n}} \right)$.

with μ as the population mean, σ as the population standard deviation, if σ is unknown, we can estimate it using sample standard deviation, s , n as the sample size, Z as the Z statistics associated with a given α . If σ is unknown or the sample size is less than 30, we would use T statistics to produce a more realizable result.

2. Sample Size (n)

When we increase the sample size the probability of type I error would decrease. This means that the precision of the test statistic improves.

3. Data Variability

If the data variability decreases, (i.e., the population standard deviation becomes smaller), we would expect to have a smaller probability of committing a type I error.

GlanzFreyer' Guide 4.2: On Type II Error for Hypothesis Testing

Type II error has the same analogy when we make a verdict that a guilty person is innocent.

1. Significance Level (α)

A decrease in the significant level causes an increase in the probability of the type II error or a decrease in the power.

2. Sample Size (n)

When we increase the sample size the probability of type II error would decrease.

3. Data Variability

Across different significance levels, if the data variability decreases, (i.e., the population standard deviation becomes smaller), the probability of committing a type II error would decrease.

4. Effect Size

The effect size is the magnitude of the difference between the null hypothesis and the alternative hypothesis.

If the effect size increases, it is easier to detect a true effect, and the probability of a type II error decreases.




Figure 4.28: Probability of type I and type II errors.

[SI*] The hypothesis-testing approach to statistical inference is very closely related to the confidence interval approach. Confidence interval estimation involves computation of bounds within which it is "reasonable" for the parameter in question to lie. For the case of a single population mean μ with σ^2 known, the structure of both hypothesis testing and confidence interval

estimation is based on the random variable

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

It turns out that the testing of $H_0 : \mu = \mu_0$ against $H_1 : \mu \neq \mu_0$ at a significance level α is equivalent to computing a $100(1 - \alpha)\%$ confidence interval on μ and rejecting H_0 if μ_0 is outside the confidence interval. If μ_0 is inside the confidence interval, the hypothesis is not rejected. Recall that with an observed value $\bar{x}$, failure to reject H_0 at significance level α implies that

$$-z_{\alpha/2} \leq \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} \leq z_{\alpha/2}$$

which is equivalent to

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu_0 \leq \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

ii. Functions in SymIntegration Related to Hypothesis Testing

[SI*] To compute several computations related to confidence interval, we can use these functions in SymIntegration:

hypothesistest(vector<double> data, double μ , double σ , double α , double effect_size)

this function computes the two-tailed hypothesis test, if $n < 30$ it will automatically use T statistics and compute the sample standard deviation instead of using the given σ , otherwise it will use Z statistics and use the given σ .

hypothesistest_righttailed(vector<double> data, double μ , double σ , double α , double effect_size)

this function computes the right-tailed hypothesis test.

hypothesistest_lefttailed(vector<double> data, double μ , double σ , double α , double effect_size)

this function computes the left-tailed hypothesis test.

hypothesistest_knownvariances_twotailed(vector<double> data1, vector<double> data2, double d_0 , double σ_1 , double σ_2 , double α , double δ)

this function computes the two-tailed hypothesis test for two means with known variances, thus we will use Z statistics. The parameter δ is the difference for the specific alternative, we can choose any small number here, for example for the right-tailed $H_1 : \mu > \mu_0$ the specific alternative will be $\mu = \mu_0 + \delta$, for the left-tailed $H_1 : \mu < \mu_0$ the specific alternative will be $\mu = \mu_0 - \delta$. The parameter δ here is used to compute the power of the test and β (type II error).

hypothesistest_knownvariances_righttailed(vector<double> data1, vector<double> data2, double d_0 , double σ_1 , double σ_2 , double α , double δ)

hypothesistest_knownvariances_lefttailed(vector<double> data1, vector<double> data2, double d_0 , double σ_1 , double σ_2 , double α , double δ)

hypothesistest_equalunknownvariances_twotailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

this function computes the two-tailed hypothesis test for two means with equal but unknown variances, thus we will use T statistics with $\nu = n_1 + n_2 - 2$ degrees of freedom.

hypothesistest_equalunknownvariances_righttailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

hypothesistest_equalunknownvariances_lefttailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

hypothesistest_unequalunknownvariances_twotailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

this function computes the two-tailed hypothesis test for two means with unequal and unknown variances, thus we will use T statistics with $\nu = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\frac{\left(\frac{s_1^2}{n_1}\right)^2}{n_1-1} + \frac{\left(\frac{s_2^2}{n_2}\right)^2}{n_2-1}}$ degrees of freedom.

hypothesistest_unequalunknownvariances_righttailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

hypothesistest_unequalunknownvariances_lefttailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

hypothesistest_paired_twotailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

this function computes the two-tailed hypothesis test for paired observations using T statistics $t = \frac{\bar{D}-d_0}{s_D/\sqrt{n}}$ with $\nu = n - 1$ degrees of freedom.

hypothesistest_paired_righttailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

hypothesistest_paired_lefttailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

iii. Compute Right-Tailed Hypothesis Testing with SymIntegration

A random sample of 100 deaths in the United States during the past year showed an average life span of 71.8 years. Assuming a population standard deviation of 8.9 years, does this seem to indicate that the mean life span today is greater than 70 years? Use a 0.05 level of significance.

Solution:

We construct the one-sided(right-tailed) hypothesis test as

$$H_0 : \mu = 70 \text{ years}$$

$$H_1 : \mu > 70 \text{ years}$$

with $\alpha = 0.05$. The critical region can be computed by finding the inverse of the cdf of Z / inverse of normal cdf with $\mu = 0, \sigma = 1$ (standard normal distribution), also known as the normal quantile function or the probit function.

To compute the inverse cdf of Z we use Wichura's algorithm, this algorithm calculates the percentage point z_p of the standard normal distribution corresponding to a given lower tail area p , in this case $p = \alpha$.

$$P(Z > z_\alpha) = 0.05$$

With Wichura's algorithm we will obtain:

$$z_\alpha = 1.645$$

thus the critical region is:

$$z > 1.645$$

where

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$

We are using the data from textfile **Vector.txt** that has 100 data of recorded deaths with mean

$$\bar{x} = 71.8842$$

the standard deviation from the sample is

$$s = 6.63$$

while from the problem it is known that the population standard deviation is

$$\sigma = 8.9$$

because we are using Z statistics so we will use σ instead of s , and hence

$$z_{\text{computed}} = \frac{71.8 - 70}{8.9 / \sqrt{100}} = 2.02$$

Decision: Reject H_0 (because $z_{\text{computed}} > z_\alpha$) and conclude that the mean life span today is greater than 70 years.

If we want to use P -value method, then we need to compute


$$P = P(Z > z_{\text{computed}}) = P(Z > 2.02) = 0.0216$$

As a result, we reach the same conclusion with P -value method, since the P -value $< \alpha$ then we have to reject H_0 .

In SymIntegration we have some functions to compute the hypothesis test, for this case we will use this function:

hypothesistest_righttailed(vector<double>, double μ_0 , double σ , double α , double δ)

The input is a vector for the data, then the μ_0 that will be used for the H_0 , σ is the population standard deviation, if the sample size is less than 30 then the function will use s (sample standard deviation) with t statistics instead, α is the level of significance / type I error, δ is the effect size that is defined with $\delta = |\bar{x} - \mu|$.



Line	Value
1	60.90
2	70.91
3	65.96
4	66.89
5	71.00
6	61.10
7	61.15
8	68.03
9	69.77
10	71.07
11	61.07
12	65.94
13	71.10
14	64.10
15	62.10
16	60.91
17	70.87
18	80.78
19	60.82
20	65.95
21	64
22	72
23	72
24	72
25	75
26	76
27	77
28	77
29	79
30	80
31	81
32	82
33	80
34	80
35	80
36	80
37	80
38	80
39	88
40	82
41	80
42	80
43	80
44	80

Figure 4.29: The data that is used for this example is saved as **Vectory.txt**.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Test ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Test ]# ./main

Sample size : 100

Mode : 72
Median : 72
Mean : 71.8842
Quantile 1/4: 70
Quantile 3/4: 75.25
Variance : 44.0116
Standard deviation : 6.63412

*****

Hypothesis testing right-tailed

H0:  $\mu = 70$ 
H1:  $\mu > 70$ 

The probability of a type I error (alpha): 0.05
The probability of a type II error (beta): 0.352857
Power: 0.647143

Critical value: 71.4639
Critical region :  $z > 1.64485$ 
Computed z : 2.02247

*Reject the null hypothesis when the sample average is greater than 71.4639

*Reject the null hypothesis when the computed z is greater than 1.64485

P-value: 0.0215638

*Reject null hypothesis if P-value  $\leq 0.05$  , we fail to reject the null hypothesis if P-value  $> 0.05$ 

*****

Time taken by function: 1750 microseconds

```

Figure 4.30: *The computation of one-tailed hypothesis testing (right-tailed) with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Right-Tailed Hypothesis Testing/main.cpp).*

iv. Compute Left-Tailed Hypothesis Testing with SymIntegration

The SineBam Electric Institute has published figures on the number of kilowatt hours used annually by various home appliances. It is claimed that a vacuum cleaner uses an average of 46 kilowatt hours per year. If a random sample of 12 homes included in a planned study indicates that vacuum cleaners use an average of 42 kilowatt hours per year with a standar deviation of 11.9 kilowatt hours, does this suggest at the 0.05 level of significance that vacuum cleaners use, on average, less than 46 kilowatt hours annually? Assume the population of kilowatt hours to be normal.

Solution:

We construct the one-sided(left-tailed) hypothesis test as

$$H_0 : \mu = 46 \text{ kilowatt hours}$$

$$H_1 : \mu < 46 \text{ kilowatt hours}$$

with $\alpha = 0.05$. The critical region can be computed by finding the inverse of the cdf of t .

We use Newton-Raphson method to compute the value of t_{α} that leaves an area of α to the left.

$$P(T \leq t_{\alpha}) = 0.05$$

$$t_{\alpha} = -1.796$$

thus the critical region is:

$$t < -1.796$$

where

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

We are using the data from textfile **Vector.txt** that has 12 dummy data of kilowatt hours mean

$$\bar{x} = 46$$

the standard deviation from the sample is

$$s = 0$$

because the dummy data are all 46, while from the problem it is known that the sample standard deviation is

$$s = 11.9$$

so we will use $s = 11.9$, and hence

$$t_{\text{computed}} = \frac{42 - 46}{11.9/\sqrt{12}} = -1.16$$

Decision: Do not reject H_0 (because $t_{\text{computed}} > t_{\alpha}$) and conclude that the average number of kilowatt hours used annually by home vacuum cleaners is not significantly less than 46.

If we want to use P -value method, then we need to compute

$$P = P(T < t_{\text{computed}}) = P(T < -1.16) \approx 0.135$$

As a result, we reach the same conclusion with P -value method, since the P -value $> \alpha$ then we fail to reject H_0 .

In `SymIntegration` we have some functions to compute the hypothesis test, for this case we will use this function:

`hypothesistest_lefttailed(vector<double>, double  $\mu_0$ , double  $\sigma$ , double  $\alpha$ , double  $\delta$ )`

The input is a vector for the data, then the μ_0 that will be used for the H_0 , σ is the population standard deviation, if the sample size is less than 30 then the function will read σ as s (sample standard deviation) and we will use t statistics instead, α is the level of significance / type I error, δ is the effect size that is defined with $\delta = |\bar{x} - \mu|$.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Left-Tailed Hypothesis Testing ]# ./main
Sample size : 12
Mode : 46
Median : 46
Mean : 46
Quantile 1/4: 46
Quantile 3/4: 46
Variance : 0
Standard deviation : 0

*****

Hypothesis testing left-tailed

H0:  $\mu = 46$ 
H1:  $\mu < 46$ 

The probability of a type I error (alpha): 0.05
The probability of a type II error (beta): 0.729687
Power: 0.270313

Critical value: 39.8307
Critical region :  $t < -1.79588$ 
Computed t : -1.1644

*Reject the null hypothesis when the sample average is less than 39.8307

*Reject the null hypothesis when the computed t is less than -1.79588

P-value: 0.134446

*Reject null hypothesis if P-value  $\leq 0.05$  , we fail to reject the null hypothesis if P-value  $> 0.05$ 

*****

Time taken by function: 1215 microseconds
```

Figure 4.31: The computation of one-tailed hypothesis testing (left-tailed) with `SymIntegration` (`SymIntegration/Examples/Statistics/Test SymIntegration Left-Tailed Hypothesis Testing/main.cpp`).

v. Compute Two-Tailed Hypothesis Testing with SymIntegration

A manufacturer of sports equipment has developed a new synthetic fishing line that the company claims has a mean breaking strength of 8 kilograms with a standard deviation of 0.5 kilogram. Test the hypothesis that $\mu = 8$ kilograms against the alternative that $\mu \neq 8$ kilograms if a random sample of 50 lines is tested and found to have a mean breaking strength of 7.8 kilograms. Use a 0.01 level of significance.

Solution:

We construct the two-sided(two-tailed) hypothesis test as

$$H_0 : \mu = 8 \text{ kilograms}$$

$$H_1 : \mu \neq 8 \text{ kilograms}$$

with $\alpha = 0.01$.

The critical region can be computed by finding the inverse of the cdf of Z / inverse of normal cdf with $\mu = 0, \sigma = 1$ (standard normal distribution), also known as the normal quantile function or the probit function.

For this two-tailed case, we will have

$$P(Z < -z_{\alpha/2}) = 0.005$$

$$P(Z > z_{\alpha/2}) = 0.005$$

With Wichura's algorithm we will obtain:

$$z_{\alpha/2} = 2.575$$

due to the symmetry of standard normal distribution, the critical regions are:

$$z < -2.575 \quad \text{and} \quad z > 2.575$$

where

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$

We are using the data from textfile **Vector.txt** that has 50 data of recorded deaths with mean

$$\bar{x} = 7.8$$

the standard deviation from the sample is

$$s = 1.03$$

while from the problem it is known that the population standard deviation is

$$\sigma = 0.5$$

because we are using Z statistics so we will use σ instead of s , and hence

$$z_{\text{computed}} = \frac{7.8 - 8}{0.5 / \sqrt{50}} = -2.823843$$

Decision: Reject H_0 (because $z_{\text{computed}} < z_{\alpha}$) and conclude that the mean breaking strength of the synthetic fishing line is not 8 kilograms, but is in fact, less than 8 kilograms based on the sample data.

Since the test in this example is two-tailed, the desired P -value is twice the area of the shaded region to the left of $z = -2.83$. Therefore, we have

$$P = P(|Z| > 2.83) = 2P(Z < -2.83) = 0.0046$$

In SymIntegration we have some functions to compute the hypothesis test, for this case of two-tailed hypothesis testing we will use this function:

hypothesistest(vector<double>, double μ_0 , double σ , double α , double δ)

The input is a vector for the data, then the μ_0 that will be used for the H_0 , σ is the population standard deviation, if the sample size is less than 30 then the function will use s (sample standard deviation) with t statistics instead, α is the level of significance / type I error, δ is the effect size that is defined with $\delta = |\bar{x} - \mu|$.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two-Tailed Hypothesis Testing ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two-Tailed Hypothesis Testing ]# ./main

Sample size : 50

Mode : 7
Median : 8
Mean : 7.8
Quantile 1/4: 7
Quantile 3/4: 8
Variance : 1.06122
Standard deviation : 1.03016

*****

Hypothesis testing two-tailed

H0:  $\mu = 8$ 
H1:  $\mu \neq 8$ 

The probability of a type I error (alpha): 0.01
The probability of a type II error (beta): 0.199421
Power: 0.800579

Critical value (upper bound): 8.18214
Critical value (lower bound): 7.81786
Critical region :  $z < -2.57583$  and  $z > 2.57583$ 
Computed  $z$  : -2.82843

*Reject the null hypothesis when the sample average is greater than 8.18214 or less than 7.81786
*Reject the null hypothesis when the computed  $z$  is greater than 2.57583 or smaller than -2.57583

P-value: 0.00467773

*Reject null hypothesis if P-value  $\leq 0.01$  , we fail to reject the null hypothesis if P-value  $> 0.01$ 

*****

Time taken by function: 1459 microseconds
```

Figure 4.32: The computation of two-tailed hypothesis testing with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Two-Tailed Hypothesis Testing/main.cpp).

vi. Two Samples Hypothesis Tests for Comparing Means

[SI*] Tests concerning two means represent a set of very important analytical tools for the scientist or engineer. Two independent random samples of sizes n_1 and n_2 , respectively, are drawn from two populations with means μ_1 and μ_2 and variances σ_1^2 and σ_2^2 . We know that the random variable

$$Z = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

has a standard normal distribution. Here we assuming that n_1 and n_2 are sufficiently large that the Central Limit Theorem applies. Of course, if the two populations are normal, the statistic above has a standard normal distribution even for small n_1 and n_2 . Obviously, if we can assume that $\sigma_1 = \sigma_2 = \sigma$, the statistic above reduces to

$$Z = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sigma \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

The two statistics above serve as a basis for the development of the test procedures involving two means. The equivalence between tests and confidence intervals, along with the technical detail involving tests on one mean, allow a simple transition to tests on two means.

[SI*] The two-sided hypothesis on two means can be written generally as

$$H_0 : \mu_1 - \mu_2 = d_0$$

Obviously, the alternative can be two sided or one sided. Again, the distribution used is the distribution of the test statistic under H_0 . Values $\bar{x}_1$ and $\bar{x}_2$ are computed and, for σ_1 and σ_2 known, the test statistic is given by

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - d_0}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

with a two-tailed critical region in the case of a two-sided alternative. That is, reject H_0 in favor of $H_1 : \mu_1 - \mu_2 \neq d_0$ if $z > z_{\alpha/2}$ or $z < -z_{\alpha/2}$. One-tailed critical regions are used in the case of the one-sided alternatives.

[SI*] Unknown But Equal Variances

The more prevalent situations involving tests on two means are those in which variances are unknown. If the scientist involved is willing to assume that both distributions are normal and that $\sigma_1 = \sigma_2 = \sigma$, the pooled t -test (two-sample t -test) may be used.

Definition 4.62: Two Sample Pooled t -Test

For the two-sided hypothesis

$$H_0 : \mu_1 = \mu_2$$

$$H_1 : \mu_1 \neq \mu_2$$

we reject H_0 at significance level α when the computed t -statistic

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - d_0}{s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

where

$$s_p^2 = \frac{s_1^2(n_1 - 1) + s_2^2(n_2 - 1)}{n_1 + n_2 - 2}$$

exceeds $t_{\frac{\alpha}{2}, n_1 + n_2 - 2}$ or is less than $-t_{\frac{\alpha}{2}, n_1 + n_2 - 2}$.

[SI*] Unknown but Unequal Variances

There are situations where the analyst is not able to assume that $\sigma_1 = \sigma_2$. Recall that, if the populations are normal, the statistic

$$T' = \frac{(\bar{X}_1 - \bar{X}_2) - d_0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

has an approximate t -distribution with approximate degrees of freedom

$$\nu = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\frac{\left(\frac{s_1^2}{n_1}\right)^2}{n_1 - 1} + \frac{\left(\frac{s_2^2}{n_2}\right)^2}{n_2 - 1}}$$

As a result, the test procedure is to not reject H_0 when

$$-t_{\frac{\alpha}{2}, \nu} < t' < t_{\frac{\alpha}{2}, \nu}$$

with ν given as above. Again, as in the case of the pooled t -test, one-sided alternatives suggest one-sided critical regions.

[SI*] Paired Observations

A study of the two-sample t -test or confidence interval on the difference between means should suggest the need for experimental design. It was suggested that the conditions of the two populations (often referred to as the two treatments) should be assigned randomly to the experimental units. This is done to avoid biased results due to systematic differences between experimental units. In other words, in hypothesis-testing jargon, it is important that any significant difference found between means be due to the different conditions of the populations and not due to the experimental units in the study.

Testing of two means can be accomplished when data are in the form of paired observations. In this pairing structure, the conditions of the two populations (treatments) are assigned

randomly within homogeneous units. Computation of the confidence interval for $\mu_1 - \mu_2$ in the situation with paired observations is based on the random variable

$$T = \frac{\bar{D} - \mu_D}{S_d / \sqrt{n}}$$

where $\bar{D}$ and S_d are random variables representing the sample mean and standard deviation of the differences of the observations in the experimental units. As in the case of the pooled t -test, the assumption is that the observations from each population are normal. This two-sample problem is essentially reduced to a one-sample problem by using the computed differences $d_1, d_2, \dots, d_n$. Thus, the hypothesis reduces to

$$H_0 : \mu_D = d_0$$

The computed test statistic is then given by

$$t = \frac{\bar{d} - d_0}{s_d / \sqrt{n}}$$

Critical regions are constructed using the t -distribution with $n - 1$ degrees of freedom.

[SI*] **Choice of Sample Size for Testing Means**

In most practical circumstances, the experiment should be planned, with a choice of sample size made prior to the data-taking process if possible. the sample size is usually determined to achieve good power for a fixed α and fixed specific alternative. This fixed alternative may be in the form of $\mu - \mu_0$ in the case of a hypothesis involving a single mean or $\mu_1 - \mu_2$ in the case of a problem involving two means.

Suppose that we wish to test the hypothesis

$$H_0 : \mu = \mu_0$$

$$H_1 : \mu > \mu_0$$

with a significance level α , when the variance σ^2 is known. For a specific alternative, say $\mu = \mu_0 + \delta$, the power of the test is

$$1 - \beta = P(\bar{X} > a \text{ when } \mu = \mu_0 + \delta)$$

Therefore,

$$\begin{aligned} \beta &= P(\bar{X} < a \text{ when } \mu = \mu_0 + \delta) \\ &= P\left(\frac{\bar{X} - (\mu_0 + \delta)}{\sigma / \sqrt{n}} < \frac{a - (\mu_0 + \delta)}{\sigma / \sqrt{n}} \text{ when } \mu = \mu_0 + \delta\right) \end{aligned}$$

Under the alternative hypothesis $\mu = \mu_0 + \delta$, the statistic

$$\frac{\bar{X} - (\mu_0 + \delta)}{\sigma / \sqrt{n}}$$

is the standard normal variable Z . So

$$\beta = P\left(Z < \frac{a - \mu_0}{\sigma / \sqrt{n}} - \frac{\delta}{\sigma / \sqrt{n}}\right) = P\left(Z < z_\alpha - \frac{\delta}{\sigma / \sqrt{n}}\right)$$

from which we conclude that

$$-z_\beta = z_\alpha - \frac{\delta\sqrt{n}}{\sigma}$$

and hence the choice of sample size

$$n = \frac{(z_\alpha + z_\beta)^2 \sigma^2}{\delta^2}$$

a result that is also true when the alternative hypothesis is $H_1 : \mu < \mu_0$.

In the case of a two-tailed test, we obtain the power $1 - \beta$ for a specified alternative when

$$n \approx \frac{(z_{\alpha/2} + z_\beta)^2 \sigma^2}{\delta^2}$$

vii. Compute Right-Tailed Hypothesis Testing on Two Means with Unknown But Equal Variances with SymIntegration

An experiment was performed to compare the abrasive wear of two different laminated materials. Twelve pieces of material 1 were tested by exposing each piece to a machine measuring wear. Ten pieces of material 2 were similarly tested. In each case, the depth of wear was observed. The samples of material 1 gave an average (code) wear of 85 units with a sample standard deviation of 4, while the samples of material 2 gave an average of 81 with a sample standard deviation of 5. Can we conclude at the 0.05 level of significance that the abrasive wear of material 1 exceeds that of material 2 by more than 2 units? Assume the populations to be approximately normal with equal variances.

Solution:

Let μ_1 and μ_2 represent the population means of the abrasive wear for material 1 and material 2, respectively.

We construct the one-sided (right-tailed) hypothesis test as

$$H_0 : \mu_1 - \mu_2 = 2$$

$$H_1 : \mu_1 - \mu_2 > 2$$

with $\alpha = 0.05$. The critical region can be computed by finding the inverse of the cdf of t .

$$P(T > t_\alpha) = 0.05$$

thus, we will obtain

$$t_\alpha = 1.725$$

where

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - d_0}{s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

with $\nu = 20$ degrees of freedom.

We are using data from textfiles **Vectorx.txt** and **Vectorx.txt**, the computations are:

$$\bar{x}_1 = 85, \quad s_1 = 4, \quad n_1 = 12$$

$$\bar{x}_2 = 81, \quad s_2 = 5, \quad n_2 = 10$$

Hence

$$s_p = \sqrt{\frac{(11)(16) + (9)(25)}{12 + 10 - 2}} = 4.478$$

$$t_{\text{computed}} = \frac{(85 - 81) - 2}{4.478 \sqrt{\frac{1}{12} + \frac{1}{10}}} = 1.04$$

$$P = P(T > 1.04) \approx 0.16$$

Decision: Do not reject H_0 . We are unable to conclude that the abrasive wear of material 1 exceeds that of material 2 by more than 2 units.

In SymIntegration we have some functions to compute the hypothesis test for tests on two means, for this case we will use this function:

hypothesistest_equalunknownvariances_righttailed(vector<double> data1, vector<double> data2, double d_0 , double α)

The textfile **Vectorx.txt** will be **data1** (the corresponding sample mean is $\bar{x}_1$ and sample standard deviation is s_1), the textfile **Vectorx.txt** will be **data2** (the corresponding sample mean is $\bar{x}_2$ and sample standard deviation is s_2), d_0 is the difference we want to test in the null hypothesis $H_0 : \mu_1 - \mu_2 = d_0$, and α is the level of significance.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Samples Right-Tailed Hypothesis Testing ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Samples Right-Tailed Hypothesis Testing ]# ./main

Sample size : 12

Mode : 88
Median : 88
Mean : 85.0417
Quantile 1/4: 80
Quantile 3/4: 88
Variance : 16.1117
Standard deviation : 4.01394
Sample size : 10

Mode : 83
Median : 83
Mean : 81.07
Quantile 1/4: 78
Quantile 3/4: 85.025
Variance : 25.3379
Standard deviation : 5.03368

*****

Hypothesis testing right-tailed

H0:  $\mu_1 - \mu_2 = 2$ 
H1:  $\mu_1 - \mu_2 > 2$ 

The probability of a type I error (alpha): 0.05
The probability of a type II error (beta): 0.95
Power: 0.05

Critical value: 3.73604
Critical region :  $t > 1.72472$ 
Computed t : 1.02295

*Reject the null hypothesis when the computed t is greater than 1.72472

P-value: 0.159273

*Reject null hypothesis if P-value <= 0.05 , we fail to reject the null hypothesis if P-value > 0.05

*****

Time taken by function: 1422 microseconds
```

Figure 4.33: The computation of right-tailed hypothesis testing for two samples tests on two means with same but unknown variances (SymIntegration/Examples/Statistics/Test SymIntegration Two Samples Right-Tailed Hypothesis Testing/main.cpp).

viii. Compute Two-Tailed Hypothesis Testing on Two Means with Known Variances with SymIntegration

A random sample of size $n_1 = 25$, taken from a normal population with a standard deviation $\sigma_1 = 5.2$, has a mean $\bar{x}_1 = 81$. A second random sample of size $n_2 = 36$, taken from a different normal population with a standard deviation $\sigma_2 = 3.4$, has a mean $\bar{x}_2 = 76$. Test the hypothesis that $\mu_1 = \mu_2$ against the alternative, $\mu_1 \neq \mu_2$. Quote a P -value in your conclusion.

Solution:

We construct the two-sided (two-tailed) hypothesis test as

$$H_0 : \mu_1 - \mu_2 = 0$$

$$H_1 : \mu_1 - \mu_2 \neq 0$$

with $\alpha = 0.05$, since the problem does not state α , we assume this α to be used to compute the critical region.

Meanwhile to only make a decision, we don't really need α , we only need to compute

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - d_0}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}} = \frac{(81 - 76) - 0}{\sqrt{\frac{5.2^2}{25} + \frac{3.4^2}{36}}} = 4.22$$

So, the P -value is

$$P(|Z| > 4.22) = 2P(Z < -4.22) = 0.00002428$$

Since P value is smaller than our chosen $\alpha = 0.05$ then we reject the null hypothesis $H_0 : \mu_1 - \mu_2 = 0$, even if we use smaller $\alpha = 0.01$ or $\alpha = 0.0001$ we will still reject the null hypothesis based on P -value method. We conclude that $\mu_1 > \mu_2$.

In SymIntegration we have some functions to compute the hypothesis test for tests on two means, for this case we will use this function:

hypothesistest_knownvariances_twotailed(vector<double> data1, vector<double> data2, double d_0 , double σ_1 , double σ_2 , double α , double δ)

The textfile **Vectorx.txt** will be **data1** (the corresponding sample mean is $\bar{x}_1$ and known population standard deviation is σ_1), the textfile **Vectorx.txt** will be **data2** (the corresponding sample mean is $\bar{x}_2$ and known population standard deviation is σ_2), d_0 is the difference we want to test in the null hypothesis $H_0 : \mu_1 - \mu_2 = d_0$, α is the level of significance, δ is the difference for the specific alternative, we can choose any small number here, for example for the right-tailed $H_1 : \mu > \mu_0$ the specific alternative will be $\mu = \mu_0 + \delta$, for the left-tailed $H_1 : \mu < \mu_0$ the specific alternative will be $\mu = \mu_0 - \delta$. The parameter δ here is used to compute the power of the test and β (type II error).


```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing ]# ./main

*****

Hypothesis testing two-tailed, known variances

H0:  $\mu_1 - \mu_2 = 0$ 
H1:  $\mu_1 - \mu_2 \neq 0$ 

Mean data 1 = 81,  $\sigma_1 = 5.2$ ,  $n_1 = 25$ 
Mean data 2 = 76,  $\sigma_2 = 3.4$ ,  $n_2 = 36$ 

The probability of a type I error (alpha): 0.05
The probability of a type II error (beta): 0.264582
Power: 0.735418

Critical value (upper bound): 1.5745
Critical value (lower bound): -1.5745
Critical region :  $z < -1.95996$  and  $z > 1.95996$ 
Computed z : 4.22169

*Reject the null hypothesis when the computed z is greater than 1.95996 or less than -1.95996

P-value: 2.42482e-05

*Reject null hypothesis if P-value  $\leq 0.05$  , we fail to reject the null hypothesis if P-value  $> 0.05$ 

*****

Time taken by function: 969 microseconds

```

Figure 4.34: The computation of two-tailed hypothesis testing for two samples tests on two means with known variances (*SymIntegration/Examples/Statistics/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing/main.cpp*).

ix. Compute Two-Tailed Hypothesis Testing on Two Means with Unequal and Unknown Variances with SymIntegration

In a study conducted at Virginia Tech, the plasma ascorbic acid levels of pregnant women were compared for smokers versus nonsmokers. Thirty-two women in the last three months of pregnancy, free of major health disorders and ranging in age from 15 to 32 years, were selected for the study. Prior to the collection 20 ml of blood, the participants were told to avoid breakfast, forgo their vitamin supplements, and avoid foods high in ascorbic acid content. From the blood samples, the following plasma ascorbic acid values were determined, in milligrams per 100 milliliters:

Is there sufficient evidence to conclude that there is a difference between plasma ascorbic acid levels of smokers and nonsmokers? Assume that the two sets of data came from normal populations with unequal variances. Use a P -value.

Nonsmokers		Smokers
0.97	1.16	0.48
0.72	0.86	0.71
1.00	0.85	0.98
0.81	0.58	0.68
0.62	0.57	1.18
1.32	0.64	1.36
1.24	0.98	0.78
0.99	1.09	1.64
0.90	0.92	
0.74	0.78	
0.88	1.24	
0.94	1.18	

Table 4.5: Plasma ascorbic acid values.

Solution:

We construct the two-sided (two-tailed) hypothesis test as

$$H_0 : \mu_1 - \mu_2 = 0$$

$$H_1 : \mu_1 - \mu_2 \neq 0$$

with $\alpha = 0.05$, since the problem does not state α , we assume this α to be used to compute the critical region.

The degrees of freedom is calculated as

$$\nu = \frac{\left(\frac{0.391478^2}{8} + \frac{0.214414^2}{24} \right)^2}{\frac{\left(\frac{0.391478^2}{8} \right)^2}{7} + \frac{\left(\frac{0.214414^2}{24} \right)^2}{23}} = 8.44$$

Hence, we use 8 degrees of freedom. Now, we compute the t_{computed}

$$t_{\text{computed}} = \frac{(\bar{x}_1 - \bar{x}_2) - d_0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}} = \frac{0.91583 - 0.97625}{\sqrt{\frac{0.391478^2}{8} + \frac{0.214414^2}{24}}} \approx -0.42$$

The P -value is

$$P(|T| > 0.42) = 2P(T < -0.42) = 0.688207$$

Decision: We fail to reject the null hypothesis, thus there is no difference between plasma ascorbic acid levels of smokers and nonsmokers.

In SymIntegration we have some functions to compute the hypothesis test for tests on two means, for this case we will use this function:

hypothesistest_unequalunknownvariances_twotailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

The textfile **Vectorx.txt** will be **data1** (the corresponding sample mean is $\bar{x}_1$ and the sample standard deviation is s_1), the textfile **Vectorx.txt** will be **data2** (the corresponding sample mean is $\bar{x}_2$ and the sample standard deviation is s_2), d_0 is the difference we want to test in the null hypothesis $H_0 : \mu_1 - \mu_2 = d_0$, α is the level of significance, δ is the difference for the specific alternative, we can choose any small number here, for example for the right-tailed $H_1 : \mu > \mu_0$ the specific alternative will be $\mu = \mu_0 + \delta$, for the left-tailed $H_1 : \mu < \mu_0$ the specific alternative will be $\mu = \mu_0 - \delta$. The parameter δ here is used to compute the power of the test and β (type II error).

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing Unequal Unknown Variances ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing Unequal Unknown Variances ]# ./main
*****
Hypothesis testing two-tailed, unequal and unknown variances

H0:  $\mu_1 - \mu_2 = 0$ 
H1:  $\mu_1 - \mu_2 \neq 0$ 

Mean data 1 = 0.915833, s1 = 0.214414, n1 = 24
Mean data 2 = 0.97625, s2 = 0.391478, n2 = 8

The probability of a type I error (alpha): 0.05
The probability of a type II error (beta): 0.00179547
Power: 0.998205

Critical value (upper bound): 0.210049
Critical value (lower bound): -0.210049
Critical region :  $t < -2.306$  and  $t > 2.306$ 
Computed t : -0.416197

*Reject the null hypothesis when the computed t is greater than 2.306 or less than -2.306

P-value: 0.688207

*Reject null hypothesis if P-value <= 0.05 , we fail to reject the null hypothesis if P-value > 0.05
*****
Time taken by function: 937 microseconds
```

Figure 4.35: The computation of two-tailed hypothesis testing for two samples tests on two means with unequal and unknown variances (SymIntegration/Examples/Statistics/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing Unequal Unknown Variances/main.cpp).

x. Compute Two-Tailed Hypothesis Testing on Paired Observations with SymIntegration

In a study conducted in the Forestry and Wildlife Department at Virginia Tech, J. A. Wesson examined the influence of the drug succinylcholine on the circulation levels of androgens in the blood. Blood samples were taken from wild, free-ranging deer immediately after they had received an intramuscular injection of succinylcholine administered using darts and a capture gun. A second blood sample was obtained from each deer 30 minutes after the first sample, after which the deer was released. The levels of androgens at time of capture and 30 minutes later, measured in nanograms per milliliter (ng/mL), for 15 deer.

Assuming that the populations of androgen levels at time of injection and 30 minutes later are normally distributed, test at the 0.05 level of significance whether the androgen concentrations are altered after 30 minutes.

Deer	Androgen At Time of Injection	Androgen 30 Minutes after Injection	d_i
1	2.76	7.02	4.26
2	5.18	3.10	-2.08
3	2.68	5.44	2.76
4	3.05	3.99	0.94
5	4.10	5.21	1.11
6	7.05	10.26	3.21
7	6.60	13.91	7.31
8	4.79	18.53	13.74
9	7.39	7.91	0.52
10	7.30	4.85	-2.45
11	11.78	11.10	-0.68
12	3.90	3.74	-0.16
13	26.00	94.03	68.03
14	67.48	94.03	26.55
15	17.04	41.70	24.66

Table 4.6: The level of androgens at time of capture and 30 minutes later, measured in nanograms per milliliter (ng/mL).

Solution:

Let μ_1 and μ_2 be the average androgen concentration at the time of injection and 30 minutes later, respectively.

We construct the two-sided (two-tailed) hypothesis test as

$$H_0 : \mu_D = \mu_2 - \mu_1 = 0$$

$$H_1 : \mu_D = \mu_2 - \mu_1 \neq 0$$

with $\alpha = 0.05$. The critical region can be computed by finding the inverse of the cdf of t .

$$P(T < t_{\alpha/2}) = 0.025$$

thus, we will obtain

$$t_{\alpha/2} = 2.145$$

so the critical region is when $t < -2.145$ or $t > 2.145$, where

$$t_{\text{computed}} = \frac{\bar{d} - d_0}{s_D / \sqrt{n}}$$

with $\nu = 14$ degrees of freedom.

Therefore,

$$t_{\text{computed}} = \frac{9.848 - 0}{18.474 / \sqrt{15}} = 2.06$$

Though the t -statistic is not significant at the 0.05 level, we obtain the P -value

$$P = P(|T| > 2.06) \approx 0.06$$

As a result, there is some evidence that there is a difference in mean circulating levels of androgen.

The assumption of no interaction would imply that the effect on androgen levels of the deer is roughly the same in the data for both treatments, i.e., at the time of injection of succinylcholine and 30 minutes following injection. This can be expressed with the two factors switching roles; for example, the difference in treatments is roughly the same across the units (i.e., the deer). There certainly are some deer/treatment combinations for which the no interaction assumption seems to hold, but there is hardly any strong evidence that the experimental units are homogeneous.

However, the nature of the interaction and the resulting increase in $\text{Var}(\bar{D})$ appear to be dominated by a substantial difference in the treatments. This is further demonstrated by the fact that 11 of the 15 deer exhibited positive signs for the computed d_i and the negative d_i (for deer 2, 10, 11, and 12) are small in magnitude compared to the 12 positive ones. Thus, it appears that the mean level of androgen is significantly higher 30 minutes following injection than at injection, and the conclusions may be stronger than $p = 0.06$ would suggest.

In SymIntegration we have some functions to compute the hypothesis test for tests on paired observations, for this case we will use this function:

hypothesistest_paired_twotailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

The textfile **Vectorx.txt** will be **data1** (the corresponding sample mean is $\bar{x}_1$), the textfile **Vectorx.txt** will be **data2** (the corresponding sample mean is $\bar{x}_2$), we make a new vector **vector<double> datad** in the function to compute the difference between **data1** and **data2**, thus $\text{datad}[i] = \text{data1}[i] - \text{data2}[i]$, for $i = 1, 2, \dots, n$, with n is the number of the paired observations, **datad**, **data1** and **data2** all have the same number of sample size: n . d_0 is the difference we want to test in the null hypothesis $H_0 : \mu_D = \mu_2 - \mu_1 = d_0$, α is the level of significance, δ is the difference for the specific alternative, we can choose any small number here, for example for the right-tailed $H_1 : \mu_D > d_0$ the specific alternative will be $\mu_D = d_0 + \delta$, for the left-tailed $H_1 : \mu_D < d_0$ the specific alternative

will be $\mu_D = d_0 - \delta$. The parameter δ here is used to compute the power of the test and β (type II error).

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing Paired Observations ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing Paired Observations ]# ./main

*****

Hypothesis testing two-tailed, paired observations

H0:  $\mu_d = \mu_1 - \mu_2 = 0$ 
H1:  $\mu_d = \mu_1 - \mu_2 \neq 0$ 

Mean data 1 = 21.6547, s1 = 30.9207, n1 = 15
Mean data 2 = 11.8067, s2 = 16.6364, n2 = 15
Mean d = 9.848, sd = 18.4736, nd = 15

The probability of a type I error (alpha): 0.05
The probability of a type II error (beta): 0.0734377
Power: 0.926562

Critical value (upper bound): 10.2304
Critical value (lower bound): -10.2304
Critical region :  $t < -2.14479$  and  $t > 2.14479$ 
Computed t : 2.06463

*Reject the null hypothesis when the computed t is greater than 2.14479 or less than -2.14479

P-value: 0.0579983

*Reject null hypothesis if P-value  $\leq 0.05$  , we fail to reject the null hypothesis if P-value  $> 0.05$ 

*****

Time taken by function: 1068 microseconds
```

Figure 4.36: The computation of two-tailed hypothesis testing for paired observations (*SymIntegration/Examples/Statistics/Test SymIntegration Two Samples Two-Tailed Hypothesis Testing Paired Observations/main.cpp*).

xi. Hypothesis Test on a Single Proportion

[SI*] We shall consider the problem of testing the hypothesis that the proportion of successes in a binomial experiment equals some specified value. That is, we are testing the null hypothesis H_0 that $p = p_0$, where p is the parameter of the binomial distribution. The alternative hypothesis may be one of the usual one-sided or two-sided alternatives:

$$p < p_0, p > p_0, \text{ or } p \neq p_0$$

The appropriate random variable on which we base our decision criterion is the binomial random variable X , although we could just as well use the statistic $\hat{p} = \frac{X}{n}$. Values of X that are far from the mean $\mu = np_0$ will lead to the rejection of the null hypothesis. Because X is a discrete binomial variable, it is unlikely that a critical region can be established whose size is exactly equal to a prespecified value α . For this reason it is preferable, in dealing with small samples, to base our decisions on P -values. TO test the hypothesis

$$H_0 : p = p_0$$

$$H_1 : p < p_0$$

we use the binomial distribution to compute the P -value

$$P = P(X \leq x \text{ when } p = p_0)$$

The value x is the number of successes in our sample of size n . If this P -value is less than or equal to α , our test is significant at the α level and we reject H_0 in favor of H_1 . Similarly, to test the hypothesis

$$H_0 : p = p_0$$

$$H_1 : p > p_0$$

at the α -level of significance, we compute

$$P = P(X \geq x \text{ when } p = p_0)$$

and reject H_0 in favor of H_1 if this P -value is less than or equal to α . Finally, to test the hypothesis

$$H_0 : p = p_0$$

$$H_1 : p \neq p_0$$

at the α -level of significance, we compute

$$P = 2P(X \leq x \text{ when } p = p_0) \quad \text{if } x < np_0$$

or

$$P = 2P(X \geq x \text{ when } p = p_0) \quad \text{if } x > np_0$$

and reject H_0 in favor of H_1 if the computed P -value is less than or equal to α .

[SI*] The binomial probabilities can be obtained from the actual binomial formula when n is small. For large n , approximation procedures are required. When the hypothesized value p_0 is very close to 0 or 1, the Poisson distribution, with parameter $\lambda = np_0$ may be used. However, the normal curve approximation, with parameters $\mu = np_0$ and $\sigma^2 = np_0q_0$, is usually preferred

for large n and is very accurate as long as p_0 is not extremely close to 0 or to 1. If we use the normal approximation, the z -value for testing $p = p_0$ is given by

$$z = \frac{x - np_0}{\sqrt{np_0q_0}} = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0q_0}{n}}}$$

which is a value of the standard normal variable Z . Hence, for a two-tailed test at the α -level of significance, the critical region is $z < -z_{\alpha/2}$ or $z > z_{\alpha/2}$. For the one-sided alternative $p < p_0$, the critical region is $z < -z_\alpha$, and for the alternative $p > p_0$, the critical region is $z > z_\alpha$.

xii. Compute Right-Tailed Hypothesis Testing on a Single Proportion with SymIntegration

A commonly prescribed drug for relieving nervous tension is believed to be only 60% effective. Experimental results with a new drug administered to a random sample of 100 adults who were suffering from nervous tension show that 70 received relief. Is this sufficient evidence to conclude that the new drug is superior the the one commonly prescribed? Use a 0.05 level of significance.

Solution:

We construct the one-sided(right-tailed) hypothesis test as

$$\begin{aligned}H_0 : p &= 0.6 \\H_1 : p &> 0.6\end{aligned}$$

with $\alpha = 0.05$. The critical region can be computed by finding the inverse of the cdf of Z

$$P(Z > z_\alpha) = 0.05$$

we will obtain $z_\alpha = 1.645$, thus the critical region is

$$z > 1.645$$

Now for the computation

$$z_{\text{computed}} = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0 q_0}{n}}} = \frac{0.7 - 0.6}{\sqrt{\frac{(0.6)(0.4)}{100}}} = 2.04$$

and the P -value

$$P = P(Z > 2.04) = 0.0206134$$

Decision: P -value is less than α . Reject H_0 and conclude that the new drug is superior.

In SymIntegration we have some functions to compute the hypothesis test for tests on a single proportion, for this case we will use this function:

hypothesistest_proportion_righttailed(vector<string> data, const string &inputString, double p_0 , double α)

The textfile **drug.txt** will be **data**, the **inputString** is the proportion for the string (e.g. RELIEF, Defective, CUPULU, PLUTUTUT, etc) that we want to compute in the **data**, the p_0 is the hypothesized value in the null hypothesis $H_0 : p = p_0$ and the alternative hypothesis $H_1 : p > p_0$, and α is the level of significance.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing for Proportion ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing for Proportion ]# ./main
*****

Right-tailed hypothesis testing on a single proportion

H0: p = 0.6
H1: p > 0.6
Data size = 100
Amount of RELIEF in the data = 70
Proportion for RELIEF, p = 0.7, q = 0.3
Critical region : z > 1.64485
Computed z : 2.04124

*****

*Reject the null hypothesis when the computed z is greater than 1.64485

P-value: 0.0206134

*Reject null hypothesis if P-value <= 0.05 , we fail to reject the null hypothesis if P-value > 0.05

*****

Time taken by function: 816 microseconds

```

Figure 4.37: The computation of two-tailed hypothesis testing on a single proportion (*SymIntegration/Examples/Statistics/Test SymIntegration Hypothesis Testing on a Single Proportion/main.cpp*).

xiii. Hypothesis Tests on Two Proportions

[SI*] Situations often arise where we wish to test the hypothesis that two proportions are equal. In general, we wish to test the null hypothesis that two proportions, or binomial parameters, are equal. That is, we are testing $p_1 = p_2$ against one of the alternatives $p_1 < p_2$, $p_1 > p_2$, or $p_1 \neq p_2$. Of course, this is equivalent to testing the null hypothesis that $p_1 - p_2 = 0$ against one of the alternatives $p_1 - p_2 < 0$, $p_1 - p_2 > 0$, or $p_1 - p_2 \neq 0$. The statistic on which we base our decision is the random variable $\hat{P}_1 - \hat{P}_2$. Independent samples of sizes n_1 and n_2 are selected at random from two binomial populations and the proportions of successes $\hat{P}_1$ and $\hat{P}_2$ for the two samples are computed.

[SI*] In our construction of confidence intervals for p_1 and p_2 we noted, for n_1 and n_2 sufficiently large, that the point estimator $\hat{P}_1$ minus $\hat{P}_2$ was approximately normally distributed with mean

$$\mu_{\hat{P}_1 - \hat{P}_2} = p_1 - p_2$$

and variance

$$\sigma_{\hat{P}_1 - \hat{P}_2}^2 = \frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}$$

Therefore, our critical region(s) can be established by using the standard normal variable

$$Z = \frac{(\hat{P}_1 - \hat{P}_2) - (p_1 - p_2)}{\sqrt{\frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}}}$$

When H_0 is true, we can substitute $p_1 = p_2 = p$ and $q_1 = q_2 = q$ (where p and q are the common values) in the preceding formula for Z to give the form

$$Z = \frac{\hat{P}_1 - \hat{P}_2}{\sqrt{pq \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

To compute a value of Z , however, we must estimate the parameters p and q that appear in the radical. Upon pooling the data from both samples, the pooled estimate of the proportion p is

$$\hat{p} = \frac{x_1 + x_2}{n_1 + n_2}$$

where x_1 and x_2 are the numbers of successes in each of the two samples. Substituting $\hat{p}$ for p and $\hat{q} = 1 - \hat{p}$ for q , the z -value for testing $p_1 = p_2$ is determined from the formula

$$z = \frac{\hat{P}_1 - \hat{P}_2}{\sqrt{\hat{p}\hat{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

The critical regions for the appropriate alternative hypotheses are set up as before, using critical points of the standard normal curve. Hence, for the alternative $p_1 \neq p_2$ at the α -level of significance, the critical region is $z < -z_{\alpha/2}$ or $z > z_{\alpha/2}$. For a test where the alternative is $p_1 < p_2$, the critical region is $z < -z_{\alpha}$, and when the alternative is $p_1 > p_2$, the critical region is $z > z_{\alpha}$.

xiv. Compute Right-Tailed Hypothesis Testing on Two Proportions with SymIntegration

A vote is to be taken among the residents of a town and the surrounding county to determine whether a proposed chemical plant should be constructed. The construction site is within the town limits, and for this reason many voters in the county believe that the proposal will pass because of the large proportion of the town voters who favor the construction. To determine if there is a significant difference in the proportions of town voters and county voters favoring the proposal, a poll is taken. If 120 of 200 town voters favor the proposal and 240 of 500 county residents favor it, would you agree that the proportion of town voters favoring the proposal is higher than the proportion of county voters? Use an $\alpha = 0.05$ level of significance.

Solution:

Let p_1 and p_2 be the true proportions of voters in the town and county, respectively, favoring the proposal.

We construct the right-tailed hypothesis test as

$$H_0 : p_1 = p_2$$

$$H_1 : p_1 > p_2$$

with $\alpha = 0.05$. The critical region can be computed by finding the inverse of the cdf of Z

$$P(Z > z_\alpha) = 0.05$$

thus, we will obtain

$$z_\alpha = 1.64485$$

so the critical region is when

$$z > 1.64485$$

Now for further computations

$$\hat{p}_1 = \frac{x_1}{n_1} = \frac{120}{200} = 0.60$$

$$\hat{p}_2 = \frac{x_2}{n_2} = \frac{240}{500} = 0.48$$

$$\hat{p} = \frac{x_1 + x_2}{n_1 + n_2} = \frac{120 + 240}{200 + 500} = 0.51$$

Therefore,

$$z_{\text{computed}} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}\hat{q}\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} = \frac{0.60 - 0.48}{\sqrt{(0.51)(0.49)\left(\frac{1}{200} + \frac{1}{500}\right)}} = 2.86972$$

The P -value is

$$P = P(Z > 2.86972) = 0.0020$$

Decision: Reject H_0 and agree that the proportion of town voters favoring the proposal is higher than the proportion of county voters.

In SymIntegration we have some functions to compute the hypothesis test for tests on two proportions, for this case we will use this function:

hypothesistest_twoproportions_righttailed(vector<string> data1, vector<string> data2, const string &inputString, double α)

The textfile **town.txt** will be **data1**, textfile **county.txt** will be **data2** the **inputString** is the proportion for the string (e.g. FAVOR, SWEDEN, CUPULU, PLUTUTUT, etc) that we want to compute in the **data1** and **data2**, and α is the level of significance.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing on Two Proportions ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing on Two Proportions ]# ./main

*****

Right-tailed hypothesis testing on two proportions

H0: p1 = p2
H1: p1 > p2
Data 1 size = 200, Data 2 size = 500
Amount of FAVOR in the data 1 = 120
Amount of FAVOR in the data 2 = 240
Proportion for FAVOR
p1 = 0.6, q1 = 0.4
p2 = 0.48, q2 = 0.52
pooled estimate of p = 0.514286, pooled estimate of q = 0.485714
Critical region : z > 1.64485
Computed z : 2.86972

*****

*Reject the null hypothesis when the computed z is greater than 1.64485

P-value: 0.00205418

*Reject null hypothesis if P-value <= 0.05 , we fail to reject the null hypothesis if P-value > 0.05

*****

Time taken by function: 1575 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing on Two Proportions ]#
```

Figure 4.38: *The computation of two-tailed hypothesis testing on two proportions (SymIntegration/Examples/Statistics/Test SymIntegration Hypothesis Testing on Two Proportions/main.cpp).*

xv. One- and Two Sample Tests Concerning Variances

[SI*] Applications of one- and two-sample tests on variances are certainly not difficult to motivate. Engineers and scientists are confronted with studies in which they are required to demonstrate that measurements involving products or processes adhere to specifications set by consumers. The specifications are often met if the process variance is sufficiently small. Attention is also focused on comparative experiments between methods or processes, where inherent reproducibility or variability must formally be compared. In addition to determine if the equal variance assumption is violated, a test comparing two variances is often applied prior to conducting a t -test on two means.

[SI*] Let us first consider the problem of testing the null hypothesis H_0 that the population variance σ^2 equals a specified value σ_0^2 against one of the usual alternatives $\sigma^2 < \sigma_0^2$, $\sigma^2 > \sigma_0^2$, or $\sigma^2 \neq \sigma_0^2$. The appropriate statistic on which to base our decision is the chi-squared statistic, which was used to construct a confidence interval for σ^2 . Therefore, if we assume that the distribution of the population being sampled is normal, the chi-squared value for testing $\sigma^2 = \sigma_0^2$ is given by

$$\chi^2 = \frac{(n-1)s^2}{\sigma_0^2}$$

where n is the sample size, s^2 is the sample variance, and σ_0^2 is the value of σ^2 given by the null hypothesis. If H_0 is true, χ^2 is a value of the chi-squared distribution with $\nu = n - 1$ degrees of freedom. Hence, for a two-tailed test at the α -level of significance, the critical region is $\chi^2 < \chi_{1-\frac{\alpha}{2}}^2$ or $\chi^2 > \chi_{\frac{\alpha}{2}}^2$. For the one-sided alternative $\sigma^2 < \sigma_0^2$, the critical region is $\chi^2 < \chi_{1-\alpha}^2$, and for the one-sided alternative $\sigma^2 > \sigma_0^2$, the critical region is

$$\chi^2 > \chi_{\alpha}^2$$

[SI*] **Robustness of χ^2 -Test to Assumption to Normality**

Various tests depend, at least theoretically, on the assumption of normality. In general, many procedures in applied statistics have theoretical underpinnings that depend on the normal distribution. These procedures vary in the degree of their dependency on the assumption of normality.

A procedure that is reasonably insensitive to the assumption is called a robust procedure (i.e., robust to normality). The χ^2 -test on a single variance is very nonrobust to normality (i.e., the practical success of the procedure depends on normality). As a result, the P -value computed may be appreciably different from the actual P -value if the population sampled is not normal. Indeed, it is quite feasible that a statistically significant P -value may not truly signal $H_1 : \sigma \neq \sigma_0$; rather, a significant value may be a result of the violation of the normality assumptions. Therefore, the analyst should approach the use of this particular χ^2 -test with caution.

[SI*] Now let us consider the problem of testing the equality of the variances σ_1^2 and σ_2^2 of two populations. That is, we shall test the null hypothesis H_0 that $\sigma_1^2 = \sigma_2^2$ against one of the usual alternatives

$$\begin{aligned}\sigma_1^2 &< \sigma_2^2 \\ \sigma_1^2 &> \sigma_2^2 \\ \sigma_1^2 &\neq \sigma_2^2\end{aligned}$$

For independent random samples of sizes n_1 and n_2 , respectively, from the two populations, the f -value for testing $\sigma_1^2 = \sigma_2^2$ is the ratio

$$f = \frac{s_1^2}{s_2^2}$$

where s_1^2 and s_2^2 are the variances computed from the two samples. If the two populations are approximately normally distributed and the null hypothesis is true, the ratio $f = \frac{s_1^2}{s_2^2}$ is a value of the F -distribution with $\nu = n_1 - 1$ and $\nu_2 = n_2 - 1$ degrees of freedom. Therefore, the critical regions of size α corresponding to the one-sided alternatives $\sigma_1^2 < \sigma_2^2$ and $\sigma_1^2 > \sigma_2^2$ are, respectively, $f < f_{1-\alpha}(\nu_1, \nu_2)$ and $f > f_{\alpha}(\nu_1, \nu_2)$. For the two-sided alternative $\sigma_1^2 \neq \sigma_2^2$, the critical region is $f < f_{1-\frac{\alpha}{2}}(\nu_1, \nu_2)$ or $f > f_{\frac{\alpha}{2}}(\nu_1, \nu_2)$.

xvi. Compute Right-Tailed Hypothesis Testing on One-Sample Variance with SymIntegration

A manufacturer of car batteries claims that the life of the company's batteries is approximately normally distributed with a standard deviation equal to 0.9 year. If a random sample of 10 of these batteries has a standard deviation of 1.2 years, do you think that $\sigma > 0.9$ year? Use a 0.05 level of significance.

Solution:

We construct the right-tailed hypothesis test as

$$\sigma^2 = 0.81$$

$$\sigma^2 > 0.81$$

with $\alpha = 0.05$. The critical region can be computed by finding the inverse of the cdf of chi-squared distribution

$$P(\chi^2 > \chi_\alpha^2) = 0.05$$

thus, we will obtain

$$\chi_\alpha^2 = 16.919$$

so the critical region is when

$$\chi^2 > 16.919$$

Now to compute χ_{computed}^2

$$\chi_{\text{computed}}^2 = \frac{(n-1)s^2}{\sigma_0^2} = \frac{(9)(1.44)}{0.81} = 16.0077$$

The P -value is

$$P(\chi_\alpha^2 > 16.0) = 0.06672$$

Decision: The χ^2 statistic is not significant at the 0.05 level. However, based on the P -value 0.06672, there is evidence that $\sigma > 0.9$.

In SymIntegration for one-sample right-tailed hypothesis test concerning variance we will use this function:

hypothesistest_onesamplevariance_righttailed(vector<double> data, double σ_0 , double α)

The textfile **battery.txt** will be **data**, $\sigma_0 = 0.9$, and $\alpha = 0.05$ is the level of significance.


```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing on One-Sample Variance ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing on One-Sample Variance ]# ./main
*****

Right-tailed hypothesis testing on one-sample variance

H0:  $\sigma^2 = 0.81$ 
H1:  $\sigma^2 > 0.81$ 
Mean data = 2.475,  $s^2 = 1.44069$ , n = 10
Critical region :  $\chi^2 > 16.919$ 
Computed  $\chi^2$  : 16.0077

*****

*Reject the null hypothesis when the computed  $\chi^2$  is greater than 16.919

P-value: 0.0667207

*Reject null hypothesis if P-value  $\leq 0.05$  , we fail to reject the null hypothesis if P-value  $> 0.05$ 

*****

Time taken by function: 1585 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing on One-Sample Variance ]# 

```

Figure 4.39: The computation for one-sample right-tailed hypothesis test concerning variance (*SymIntegration/Examples/Statistics/Test SymIntegration Hypothesis Testing on One-Sample Variance/main.cpp*).

xvii. Compute Two-Tailed Hypothesis Testing on Two-Samples Variances with SymIntegration

In testing for the difference in the abrasive wear of the two materials, we assumed that the two unknown population variances were equal. Were we justified in making this assumption? Use a 0.10 level of significance.

Solution:

We construct the right-tailed hypothesis test as

$$H_0 : \sigma_1^2 = \sigma_2^2$$

$$H_1 : \sigma_1^2 \neq \sigma_2^2$$

with $\alpha = 0.10$. The critical region can be computed by finding the inverse of the cdf of F distribution

$$P(F > f_{\alpha/2}(\nu_1, \nu_2)) = 0.05$$

$$P(F < f_{1-\frac{\alpha}{2}}(\nu_1, \nu_2)) = 0.05$$

thus we will obtain

$$f_{\alpha/2}(\nu_1, \nu_2) = f_{0.05}(11, 9) = 3.10249$$

$$f_{1-\frac{\alpha}{2}}(\nu_1, \nu_2) = f_{0.95}(11, 9) = 0.345277$$

Therefore, the null hypothesis is rejected when $f < 0.34$ or $f > 3.10$, where $f = \frac{s_1^2}{s_2^2}$ with $\nu_1 = 11$ and $\nu_2 = 9$ degrees of freedom.

The computations for F statistic is

$$f = \frac{s_1^2}{s_2^2} = \frac{16}{25} = 0.64$$

Decision: Do not reject H_0 . Conclude that there is insufficient evidence that the variances differ.

In SymIntegration for two-samples two-tailed hypothesis test concerning variances we will use this function:

hypothesistest_twosamplesvariances_twotailed(vector<double> data1, vector<double> data2, double α)

The textfile **Vectorx.txt** will be **data1**, the textfile **Vector.y.txt** will be **data2**, and $\alpha = 0.1$ is the level of significance.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing on Two-Samples Variances ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing on Two-Samples Variances ]# ./main
*****

Two-tailed hypothesis testing on two-samples variances

H0:  $\sigma_1^2 = \sigma_2^2$ 
H1:  $\sigma_1^2 \neq \sigma_2^2$ 
Mean data 1 = 85.0417,  $s_1^2 = 16.1117$ ,  $n_1 = 12$ 
Mean data 2 = 81.07,  $s_2^2 = 25.3379$ ,  $n_2 = 10$ 
Critical region :  $f > 3.10249$  and  $f < 0.345277$ 
Computed f : 0.635875

*****

*Reject the null hypothesis when the computed f is greater than 3.10249 or less than 0.345277

P-value: 0.472498

*Reject null hypothesis if P-value  $\leq 0.1$  , we fail to reject the null hypothesis if P-value  $> 0.1$ 

*****

Time taken by function: 935 microseconds

```

Figure 4.40: *The computation for two-samples two-tailed hypothesis test concerning variances (SymIntegration/Examples/Statistics/Test SymIntegration Hypothesis Testing on Two-Samples Variances/main.cpp).*

xviii. Goodness-of-Fit Test

[SI*] We have been concerned with the testing of statistical hypotheses about single population parameters such as μ , σ^2 , and p . Now we shall consider a test to determine if a population has a specified theoretical distribution.

xix. Test for Independence (Categorical Data)

[SI*] The chi-squared test can also be used to test the hypothesis of independence of two variables of classification.

A contingency table with r rows and c columns is referred to as an $r \times c$ table. The row and column totals are called marginal frequencies.

Definition 4.63: Test for Independence

Calculate

$$\chi^2 = \sum_i \frac{(o_i - e_i)^2}{e_i}$$

where the summation extends over all rc cells in the $r \times c$ contingency table.

If $\chi^2 > \chi^2_{\alpha}$ with $\nu = (r - 1)(c - 1)$ degrees of freedom, reject the null hypothesis of independence at the α -level of significance; otherwise, we fail to reject the null hypothesis.

[SI*] It is important that the statistic on which we base our decision has a distribution that is only approximated by the chi-squared distribution. The computed χ^2 -values depend on the cell frequencies and consequently are discrete. The continuous chi-squared distribution seems to approximate the discrete sampling distribution of χ^2 very well, provided that the number of degrees of freedom is greater than 1. In a 2×2 contingency table, where we have only 1 degree of freedom, a correction called Yates' correction for continuity is applied. The corrected formula then becomes

$$\chi^2(\text{corrected}) = \sum_i \frac{(|o_i - e_i| - 0.5)^2}{e_i}$$

If the expected cell frequencies are large, the corrected and uncorrected results are almost the same. When the expected frequencies are between 5 and 10, Yates' correction should be applied. For expected frequencies less than 5, the Fisher-Irwin exact test should be used. The Fisher-Irwin test may be avoided, however, by choosing a larger sample.

xx. Compute Test for Independence for Categorical Data with SymIntegration

Suppose that we wish to determine whether the opinions of the are independent of their levels of income. Members of a random sample of 1000 registered voters from the state of Illinois are classified as to whether they are in a low, medium, or high income bracket and whether or not the favor the tax reform.

Tax Reform	Income Level			Total
	Low	Medium	High	
For	182	213	203	598
Against	154	138	110	402
Total	336	351	313	1000

Table 4.7: A random sample of 1000 registered voters from the state of Illinois are classified based on their income level and whether or not they favor the tax reform.

Our decision to accept or reject the null hypothesis, H_0 , of independence between a voter's opinion concerning the tax reform and his or her level of income is based upon how good a fit we have between the observed frequencies in each of the 6 cells of the Table and the frequencies that we would expect for each cell under the assumption that H_0 is true. To find these expected frequencies, let us define the following events:

L : A person selected is in the low-income level.

M : A person selected is in the medium-income level.

H : A person selected is in the high-income level.

F : A person selected is for the tax reform.

A : A person selected is against the tax reform.

By using the marginal frequencies, we can list the following probability estimates

$$P(L) = \frac{336}{1000}, \quad P(M) = \frac{351}{1000}, \quad P(H) = \frac{313}{1000}$$

$$P(F) = \frac{598}{1000}, \quad P(A) = \frac{402}{1000}$$

The expected frequencies are obtained by multiplying each cell probability by the total number of observations. We round these frequencies to one decimal. Thus, the expected number of low-income voters in our sample who favor the tax reform is estimated to be

$$P(L \cap F) \times 1000 = P(L)P(F) \times 1000 = \frac{(336)(598)}{1000} = 200.9$$

when H_0 is true. The general rule for obtaining the expected frequency of any cell is given by the following formula;

$$\text{expected frequency} = \frac{(\text{column total}) \times (\text{row total})}{\text{grand total}}$$

We will find that

$$\begin{aligned}\chi^2 &= \frac{(182 - 200.9)^2}{200.9} + \frac{(213 - 209.9)^2}{209.9} + \frac{(203 - 187.2)^2}{187.2} \\ &\quad + \frac{(154 - 135.1)^2}{135.1} + \frac{(138 - 141.1)^2}{141.1} + \frac{(110 - 125.8)^2}{125.8} \\ &= 7.878212\end{aligned}$$

with $\alpha = 0.05$ we will have

$$\chi_{0.05}^2 = 5.991464$$

with $\nu = (r - 1)(c - 1) = (2 - 1)(3 - 1) = 2$. If $\chi^2 > \chi_{0.05}^2$ then reject the null hypothesis; otherwise, we fail to reject the null hypothesis.

Another way is to use P -value, which can be obtained by

$$P = P(\chi^2 > 7.878212) = 0.019466$$

In this case, the null hypothesis is rejected ($\chi^2 > \chi_{0.05}^2$ and P -value is less than $\alpha = 0.5$) and we conclude that a voter's opinion concerning the tax reform and his or her level of income are not independent.

In SymIntegration, to compute test for independence for categorical data we can use this function:

hypothesistest_categoricaldata(vector<vector<double>> data, double α)

The textfile **Voters.txt** will be **data** in matrix form, and $\alpha = 0.05$ is the level of significance.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hypothesis Testing for Independence with Categorical Data ]# ./main
sum of column 0 = 336
sum of column 1 = 351
sum of column 2 = 313
sum of row 0 = 598
sum of row 1 = 402

*****

Hypothesis testing on categorical data

H0: Independent
H1: Not independent
Number of rows = 2, number of columns = 3
Degrees of freedom = 2

Observed Frequencies:
      182.000000      213.000000      203.000000
      154.000000      138.000000      110.000000

Expected Frequencies:
      200.928000      209.898000      187.174000
      135.072000      141.102000      125.826000
Critical region :  $\chi^2 > 5.991464$ 
Computed  $\chi^2$  : 7.878212

*****

*Reject the null hypothesis when the computed  $\chi^2$  is greater than 5.991464

P-value: 0.019466

*Reject null hypothesis if P-value <= 0.050000 , we fail to reject the null hypothesis if P-value > 0.050000

*****

Time taken by function: 1748 microseconds
```

Figure 4.41: The computation for test for independence for categorical data (*SymIntegration/Examples/Statistic-s/Test SymIntegration Hypothesis Testing for Independence with Categorical Data/main.cpp*).

xxi. Compute Two-Sample Case Study using t - and F - Tests with the Box-and-Whisker and Quantile-Quantile Plot with SymIntegration and Hamzstplot

In a data analysis study conducted by personnel at the Statistics Consulting Center at Virginia Tech, two different materials, alloy A and alloy B , were compared in terms of breaking strength. Alloy B is more expensive, but it should certainly be adopted if it can be shown to be stronger than alloy A . The consistency of performance of the two alloys should also be taken into account. Random samples of beams made from each alloy were selected, and strength was measured in units of 0.001-inch deflection as a fixed force was applied at both ends of the beam. Twenty specimens were used for each of the two alloys you can see it in **alloyA.txt** and **alloyB.txt** for this case example working directory.

Solution:

It is important that the engineer compare the two alloys. Of concern is average strength and reproducibility. It is of interest to determine if there is a severe violation of the normality assumption required of both the t - and F -tests.

We will construct the hypothesis test :

$$H_0 : \mu_A = \mu_B$$

$$H_1 : \mu_A > \mu_B$$

we will use the t -test to confirm whether alloy A and alloy B has the same average strength or alloy B has greater average strength than alloy A .

The sample means and standar deviations are

$$\bar{x}_A = 83.55$$

$$s_A = 3.663$$

$$\bar{x}_B = 79.70$$

$$s_B = 3.097$$

With $\alpha = 0.05$. Now, we will compute

$$t_{\text{computed}} = \frac{(\bar{x}_A - \bar{x}_B) - d_0}{\sqrt{\frac{s_A^2}{n_A} + \frac{s_B^2}{n_B}}} = 3.58953$$

The P -value is

$$P = P(T > 3.58953) = 0.0004897$$

Afterwards, we will construct another hypothesis test for the F -test:

$$H_0 : \sigma_A^2 = \sigma_B^2$$

$$H_1 : \sigma_A^2 \neq \sigma_B^2$$

we will have

$$f_{\text{computed}} = \frac{s_A^2}{s_B^2} = 1.39929$$

The P -value can be computed with

$$P = 2P(F > 1.39929) = 2(1 - P(F < 1.39929)) = 0.4709$$

Decision: Rejects H_0 in favor of H_1 due to the P -value from t -test is smaller than α ($0.0004897 < 0.05$), it means that the strength of alloy A is greater than alloy B . Then for the second test, the F -test, we fail to reject H_0 as the P -value is greater than α ($0.4709 > 0.05$), thus it means that the variances of alloy A and alloy B are different.

In **SymIntegration**, to help for the t -test and F -test for this example case we can use these two functions:

hypothesistest_unequalunknownvariances_righttailed(vector<double> data1, vector<double> data2, double d_0 , double α , double δ)

hypothesistest_twosamplesvariances_twotailed(vector<double> data1, vector<double> data2, double α)

The textfile **alloyA.txt** will be **data1**, textfile **alloyB.txt** will be **data2**, $d_0 = 0$, $\delta = 1$ (to compute β and the power of the test), and $\alpha = 0.05$ is the level of significance.

To plot the box-and-whisker or the normal quantile-quantile plot we can use **Hamzstplot** and **SymIntegration**, we join the textfile into one **alloy.txt** that has the first 20 data that represents alloy A and the next 20 data represents data for alloy B , we also create a textfile **alloyname.txt** to name the box-and-whisker plot for each alloy.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Hamzstplot 1.51/Examples with Makefile/Statistics/Plot Box-and-Whisker Plots for Two Alloys ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lhamzstplot -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Hamzstplot 1.51/Examples with Makefile/Statistics/Plot Box-and-Whisker Plots for Two Alloys ]# ./main
*****

Right-tailed hypothesis testing, unequal and unknown variances

H0:  $\mu_1 - \mu_2 = 0$ 
H1:  $\mu_1 - \mu_2 > 0$ 

Mean data 1 = 83.55, s1 = 3.66312, n1 = 20
Mean data 2 = 79.7, s2 = 3.09669, n2 = 20

The probability of a type I error (alpha): 0.05
The probability of a type II error (beta): 0.227298
Power: 0.772702

Critical value: 1.27596
Critical region :  $t > 1.6883$ 
Computed t : 3.58953

*Reject the null hypothesis when the computed t is greater than 1.6883

P-value: 0.000489718

*Reject null hypothesis if P-value  $\leq 0.05$  , we fail to reject the null hypothesis if P-value  $> 0.05$ 

*****

Right-tailed hypothesis testing on two-samples variances

H0:  $\sigma_1^2 = \sigma_2^2$ 
H1:  $\sigma_1^2 > \sigma_2^2$ 
Mean data 1 = 83.55, s12 = 13.4184, n1 = 20
Mean data 2 = 79.7, s22 = 9.58947, n2 = 20
Critical region :  $f > 2.16825$ 
Computed f : 1.39929

*****

*Reject the null hypothesis when the computed f is greater than 2.16825

P-value: 0.235461

*Reject null hypothesis if P-value  $\leq 0.05$  , we fail to reject the null hypothesis if P-value  $> 0.05$ 

*****

```

Figure 4.42: The computation for both *t*- and *F*-test for the alloys data (*SymIntegration/Examples/Statistics/Test SymIntegration and Hamzstplot Proc t-Test and Plot Box-and-Whisker Plots for Two Alloys/main.cpp*).

```

*****
Two-tailed hypothesis testing on two-samples variances

H0:  $\sigma_1^2 = \sigma_2^2$ 
H1:  $\sigma_1^2 \neq \sigma_2^2$ 
Mean data 1 = 83.55,  $s_1^2 = 13.4184$ ,  $n_1 = 20$ 
Mean data 2 = 79.7,  $s_2^2 = 9.58947$ ,  $n_2 = 20$ 
Critical region :  $f > 2.52645$  and  $f < 0.395812$ 
Computed f : 1.39929

*****

*Reject the null hypothesis when the computed f is greater than 2.52645 or less than 0.395812

P-value: 0.470922

*Reject null hypothesis if P-value  $\leq 0.05$  , we fail to reject the null hypothesis if P-value  $> 0.05$ 

*****
Press ENTER to continue...

```

Figure 4.43: The computation for both *t*- and *F*-test for the alloys data (*SymIntegration/Examples/Statistics/Test SymIntegration and Hamzstplot Proc t-Test and Plot Box-and-Whisker Plots for Two Alloys/main.cpp*).




Figure 4.44: The box-and-whisker plot for the alloys data (*SymIntegration/Examples/Statistics/Test SymIntegration and Hamzstplot Proc t-Test and Plot Box-and-Whisker Plots for Two Alloys/main.cpp*).

For the quantile-quantile plot, we make use of this function in *SymIntegration*:
normalquantile(vector<double> data, c)

With **data** is the sample data, e.g. alloy A or alloy B, the data is saved in textfile, and **c** is the scale to limit the *x*-axis range.

Basically, the normal quantile-quantile plot needs to follow this process:

1. Sort the **vector<double>** for the data, from lowest to highest, e.g. **std::sort(alloyA.begin(), alloyA.end())**.

2. Call the function to create a **vector<double>** of the normal quantile, e.g. **normalquantile(alloyA, 0.3)**, 0.3 is the scale to shrink the x -axis range / limit.

3. We will then have this sequence of 2-tuple:

$$(x_i, y_i) = (\text{data}_i - \overline{\text{data}}, \text{data}_i)$$

with $\overline{\text{data}}$ is the mean sample for alloy A or alloy B . We can then use the scatter plot from this quantile-quantile that have been generated : (x_i, y_i) .




Figure 4.45: The normal quantile-quantile plot of data for alloy A (*SymIntegration/Examples/Statistics/Test SymIntegration and Hamzstplot Plot 2D Normal Quantile-Quantile Plot/main.cpp*).




Figure 4.46: The normal quantile-quantile plot of data for alloy A (*SymIntegration/Examples/Statistics/Test SymIntegration and Hamzstplot Plot 2D Normal Quantile-Quantile Plot/main.cpp*).

xxii. One-Way Analysis of Variance (ANOVA)

[SI*] In the estimation and hypothesis testing section, we were restricted in each case to considering no more than two population parameters. Such was the case, for example, in testing for the equality of two population means using independent samples from normal populations with common but unknown variance, where it was necessary to obtain a pooled estimate of σ^2 .

Suppose in an industrial experiment that an engineer is interested in how the mean absorption of moisture in concrete varies among 5 different concrete aggregates. The samples are exposed to moisture for 48 hours. It is decided that 6 samples are to be tested for each aggregate, requiring a total of 30 samples to be tested.

The model for this situation may be set up as follows. There are 6 observations taken from each of 5 populations with means $\mu_1, \mu_2, \dots, \mu_5$, respectively. We may wish to test

$$\begin{aligned} H_0 : \mu_1 = \mu_2 = \dots = \mu_5 \\ H_1 : \text{At least two of the means are not equal} \end{aligned}$$

In addition, we may be interested in making individual comparisons among these 5 population means.

In the analysis-of-variance procedure, it is assumed that whatever variation exists among the aggregate average is attributed to

1. Variation in absorption among observation within aggregate types
2. Variation among aggregate types, that is, due to differences in the chemical composition of the aggregates.

[SI*] Random samples of size n are selected from each of k populations. The k different populations are classified on the basis of a single criterion such as different treatments or groups. Today the term **treatment** is used generally to refer to the various classifications, whether they be different aggregates, different analysts, different fertilizers, or different regions of the country.

[SI*] **Assumption and Hypotheses in One-Way ANOVA**

It is assumed that the k populations are independent and normally distributed with means $\mu_1, \mu_2, \dots, \mu_k$ and common variance σ^2 . These assumptions are made more palatable by randomization. We wish to derive appropriate methods for testing the hypothesis

$$\begin{aligned} H_0 : \mu_1 = \mu_2 = \dots = \mu_k \\ H_1 : \text{At least two of the means are not equal} \end{aligned}$$

Let y_{ij} denote the j -th observation from the i -th treatment and arrange the data to become a table. Here, Y_i is the total of all observations in the sample from the i -th treatment, $\bar{y}_i$ is the mean of all observations in the sample from the i -th treatment, $Y_{..}$ is the total of all nk observations, and $\bar{y}_{..}$ is the mean of all nk observations.

[SI*] **Model for One-Way ANOVA**

Each observation may be written in the form

$$Y_{ij} = \mu_i + \epsilon_{ij}$$

where ϵ_{ij} measures the deviation of the j -th observation of the i -th sample from the corresponding treatment mean. The ϵ_{ij} -term represents random error and plays the same role as the error terms in the regression models. An alternative preferred form of this equation is obtained by substituting $\mu_i = \mu + \alpha_i$, subject to the constraint

$$\sum_{i=1}^k \alpha_i = 0$$

Hence, we may write

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}$$

where μ is just the grand mean of all the μ_i , that is,

$$\mu = \frac{1}{k} \sum_{i=1}^k \mu_i$$

and α_i is called the effect of the i -th treatment.

The null hypothesis that the k population means are equal against the alternative that at least two of the means are unequal may now be replaced by the equivalent hypothesis

$$H_0 : \alpha_1 = \alpha_2 = \cdots = \alpha_k$$

$$H_1 : \text{At least one of the } \alpha_i \text{ is not equal to zero}$$

[SI*] Resolution of Total Variability into Components

Our test will be based on a comparison of two independent estimates of the common population variance σ^2 . These estimates will be obtained by partitioning the total variability of our data, designated by the double summation

$$\sum_{i=1}^k \sum_{j=1}^n (y_{ij} - \bar{y}_{..})^2$$

into two components.

Theorem 4.25: Sum of Squares Identity

The formula is

$$\sum_{i=1}^k \sum_{j=1}^n (y_{ij} - \bar{y}_{..})^2 = n \sum_{i=1}^k (\bar{y}_{i.} - \bar{y}_{..})^2 + \sum_{i=1}^k \sum_{j=1}^n (y_{ij} - \bar{y}_{i.})^2$$

Definition 4.64: Three Important Measures of Variability

It will be convenient in what follows to identify the terms of the sum-of-squares identity by the following notation:

SST (Total sum of squares)

$$\sum_{i=1}^k \sum_{j=1}^{n_i} (y_{ij} - \bar{y}_{..})^2$$

SSA (Treatment sum of squares)

$$n \sum_{i=1}^k (\bar{y}_{i.} - \bar{y}_{..})^2$$

SSE (Error sum of squares)

$$\sum_{i=1}^k \sum_{j=1}^{n_i} (y_{ij} - \bar{y}_{i.})^2$$

The sum-of-squares identity can then be represented symbolically by the equation

$$SST = SSA + SSE$$

The identity above expresses how between-treatment and within-treatment variation add to the total sum of squares.

Theorem 4.26: The Expected Value of SSA

The formula is

$$E(SSA) = (k - 1)\sigma^2 + n \sum_{i=1}^k \alpha_i^2$$

Definition 4.65: Treatment Mean Square

If H_0 is true, an estimate of σ^2 , based on $k - 1$ degrees of freedom, is provided by this expression:

$$s_1^2 = \frac{SSA}{k - 1}$$

If H_0 is true and thus each α_i in $E(SSA)$ is equal to zero, we see that

$$E\left(\frac{SSA}{k - 1}\right) = \sigma^2$$

and s_1^2 is an unbiased estimate of σ^2 . However, if H_1 is true, we have

$$E\left(\frac{SSA}{k - 1}\right) = \sigma^2 + \frac{n}{k - 1} \sum_{i=1}^k \alpha_i^2$$

and s_1^2 estimates σ^2 plus an additional term, which measures variation due to the systematic effects.

Definition 4.66: Error Mean Square

A second and independent estimate of σ^2 , based on $k(n - 1)$ degrees of freedom, is this familiar formula:

$$s^2 = \frac{SSE}{k(n - 1)} \quad (4.114)$$

It is instructive to point out the importance of the expected values of the mean squares indicated above.

[SI*] Use of F -Test in ANOVA

The estimate s^2 is unbiased regardless of the truth or falsity of the null hypothesis. It is important to note that the sum-of-squares identity has partitioned not only the total variability of the data, but also the total number of degrees of freedom. That is,

$$nk - 1 = k - 1 + k(n - 1)$$

[SI*] F -Ratio for Testing Equality of Means

When H_0 is true, the ratio $f = \frac{s_1^2}{s^2}$ is a value of the random variable F having the F -distribution with $k - 1$ and $k(n - 1)$ degrees of freedom. Since s_1^2 overestimates σ^2 when H_0 is false, we have a one-tailed test with the critical region entirely in the right tail of the distribution.

The null hypothesis H_0 is rejected at the α -level of significance when

$$f > f_\alpha[k - 1, k(n - 1)]$$

Another approach, the P -value approach, suggests that the evidence in favor or against H_0 is

$$P = P\{f[k - 1, k(n - 1)] > f\}$$

When H_1 is true, the presence of the condition $E(s_1^2) > E(s^2)$ suggests that the F -ratio be used in the context of a one-sided upper-tailed test. That is, we would expect the numerator

s_1^2 to exceed the denominator.

The computations for an analysis-of-variance problem are usually summarized in tabular form.

[SI*] Assumptions for one-way ANOVA:

1. The dependent variable is approximately normally distributed within each group. This assumption is more critical for smaller sample sizes.
2. The samples are selected at random and should be independent of one another.
3. All groups have equal standard deviations.
4. Each data point should belong to one and only one group / treatment. There should be no overlap or sharing of data points between groups / treatments.

[SI*] There are two main types of ANOVA:

1. One-way ANOVA

This is the most basic form of ANOVA and is used when there is only one independent variable with more than two treatments or groups. It assesses whether there are any statistically significant differences among the means of the groups.

2. Two-way ANOVA

Extending the one-way ANOVA, two-way ANOVA involves two independent variables. It allows for examination of the main effects of each variable as well as the interaction between them. The interaction effect explores whether the effect of one variable on the dependent variable is different depending on the level of the other variable.

GlanzFreya' Guide 4.3: How to Perform One-Way ANOVA

One-way ANOVA is a type of hypothesis test where only one factor is considered. We use F -statistic to perform a one-way analysis of variance.

The steps are:

1. Define the null and alternative hypothesis

$$H_0 : \mu_1 = \mu_2 = \cdots = \mu_k$$

$$H_1 : \text{At least two of the means are not equal}$$

2. Compute the degrees of freedom between and within the treatments / groups.

$$df_{\text{between}} = k - 1$$

$$df_{\text{within}} = N - k$$

$$df_{\text{total}} = N - 1$$

with N is the number of samples in all treatments / groups combined, and k is the number of treatments / groups.

3. Compute the mean of all samples in each treatment / group, then compute the grand mean ($\mu_1, \dots, \mu_k, \mu_{\text{grand}}$).
4. Compute the sum of squares total and sum of squares within, then compute the sum of squares between (SSA, SST, SSE).
5. Compute the variance between and within samples

$$s_1^2 = \frac{SSA}{k - 1}$$

$$s^2 = \frac{SSE}{k(n - 1)}$$

6. Compute the F value with this formula:

$$F_{\text{value}} = \frac{s_1^2}{s^2}$$

7. Compute the P value with this formula:

$$P_{\text{value}} = 1 - P(X \leq F_{\text{value}}) = 1 - F(F_{\text{value}}; k - 1, N - k)$$

with $F(F_{\text{value}}; k - 1, N - k)$ is the cdf of F -distribution with numerator degree of freedom $k - 1$ and denominator degree of freedom $N - k$.

8. If $F_{\text{value}} < P_{\text{value}}$ then we fail to reject the null hypothesis.

If $F_{\text{value}} > P_{\text{value}}$ then we reject the null hypothesis.

The fundamental strategy of ANOVA is to systematically examine variability within treatments / groups being compared and also examine variability among the groups being compared.

xxiii. Compute One-Way ANOVA with SymIntegration

Suppose in an industrial experiment that an engineer is interested in how the mean absorption of moisture in concrete varies among 5 different concrete aggregates. The samples are exposed to moisture for 48 hours. It is decided that 6 samples are to be tested for each aggregate, requiring a total of 30 samples to be tested.

The model for this situation may be set up as follows. There are 6 observations taken from each of 5 populations with means $\mu_1, \mu_2, \dots, \mu_5$, respectively.

Aggregate:	1	2	3	4	5	
	551	595	639	417	563	
	457	580	615	449	631	
	450	508	511	517	522	
	731	583	573	438	613	
	499	633	648	415	656	
	632	517	677	555	679	
Total	3320	3416	3663	2791	3664	16,854
Mean	553.33	569.33	610.50	465.17	610.67	561.80

Table 4.8: The absorption of moisture in concrete aggregates.

Test the hypothesis $\mu_1 = \mu_2 = \mu_3 = \mu_4 = \mu_5$ at the 0.05 level of significance for the data on absorption of moisture by various types of cement aggregates.

Solution:

The hypotheses are

$$H_0 : \mu_1 = \mu_2 = \dots = \mu_5$$

$$H_1 : \text{At least two of the means are not equal}$$

$$\alpha = 0.05$$

Critical region :

$$f > 2.76$$

with $\nu_1 = 4$ and $\nu_2 = 25$ degrees of freedom. The sum-of-squares computations give

$$SST = 209,377$$

$$SSA = 85,356$$

$$SSE = 209,377 - 85,356 = 124,021$$

Decision: Reject H_0 and conclude that the aggregates do not have the same mean absorption. The P -value for $f = 4.30$ is 0.0088, which is smaller than $\alpha = 0.05$.

This example has equal sample size from aggregate 1 to aggregate 5, however in real life, during experimental work, one often loses some of the desired observations. Experimental animals may die, experimental material may be damaged, or human subjects may drop out of a study. For analysis with unequal sample size, we just need to adjust the formula for *SST*, *SSA* and *SSE*.

In *SymIntegration*, the function to compute the one-way analysis-of-variance / ANOVA is: **ANOVA(vector<vector<double>>)**

The input is a matrix that is defined as **vector<vector<double>>** from the standard C++ library. It is a **void** function, so it returns only the table and nothing else.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration ANOVA ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration ANOVA ]# ./main

Matrix A :
551.000000    595.000000    639.000000    417.000000    563.000000
457.000000    580.000000    615.000000    449.000000    631.000000
450.000000    508.000000    511.000000    517.000000    522.000000
731.000000    583.000000    573.000000    438.000000    613.000000
499.000000    633.000000    648.000000    415.000000    656.000000
632.000000    517.000000    677.000000    555.000000    679.000000

One-Way ANOVA:

Source      DF      SS      MS      F      P
Model        4    85356.466667    21339.116667    4.301536    0.008752
Error       25   124020.333333    4960.813333
Total       29   209376.800000

R-Square      Grand Mean      Root MSE
0.407669      561.800000      70.433041

Time taken by function: 1417 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration ANOVA ]#
```

Figure 4.47: The computation of one-way analysis-of-variance / ANOVA for the cement aggregates with *SymIntegration*, the execution time for the function is around 1417 microseconds without dependencies of using any external libraries besides *SymIntegration*, it is 1.76 times faster than using *Boost*. (*SymIntegration/Examples/Statistics/Test SymIntegration One-Way ANOVA/main.cpp*).

We would like to explain why we tried to use **Boost** as comparison. The library **Boost** is very robust and solid, it includes all pdf and cdf computation for almost all distributions that are known today. So we tried to use **Boost** to compute the cdf for *F*-distribution to compute the *P*-value. It resulted in 2494 microseconds for the execution time of the ANOVA function. It becomes slower because we are using two libraries **Boost** and **SymIntegration**. If the reader wants to compare only using **Boost** and create the ANOVA table, they can compare the execution time, **Boost** only, **SymIntegration** only, and **Boost + SymIntegration** (this one will be slower than the previous two options).

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration ANOVA ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration ANOVA ]# ./main

Matrix A :
  551.000000    595.000000    639.000000    417.000000    563.000000
  457.000000    580.000000    615.000000    449.000000    631.000000
  450.000000    508.000000    511.000000    517.000000    522.000000
  731.000000    583.000000    573.000000    438.000000    613.000000
  499.000000    633.000000    648.000000    415.000000    656.000000
  632.000000    517.000000    677.000000    555.000000    679.000000

One-Way ANOVA:

Source      DF      SS      MS      F      P
Model        4    85356.466667    21339.116667    4.301536    0.008752
Error       25   124020.333333    4960.813333
Total       29   209376.800000

R-Square      Grand Mean    Root MSE
0.407669      561.800000      70.433041

Time taken by function: 2494 microseconds

```

Figure 4.48: The computation of one-way analysis-of-variance / ANOVA for the cement aggregates with Boost and SymIntegration, the execution time for the function is around 2494 microseconds.

```

...

double Fcdf(double x, double r1, double r2)
{
    //boost::math::fisher_f_distribution<> fisher_f(r1,r2);
    //double cdf_value = boost::math::cdf(fisher_f,x);

    /*The cumulative distribution function (CDF) for Fisher's F distribution
    It has at least 11 decimal digit precision with boost::math::
       fisher_f_distribution
    */

    double z ;
    double cdf_value;

    z = (x*r1)/(r2 + r1*x);
    cdf_value = incbeta(z,r1/2.0, r2/2.0);

    return cdf_value;
}

...
...

void ANOVA(vector<vector<double>> matrix)
{
    vector<vector<double>> anovamatrix;

```

```
vector<double> total_column;
vector<double> mean_column;
vector<double> sst_column;
vector<double> ssa_column;
int C = matrix[0].size();
int N = 0; // computing total data

for(int i = 0 ; i < C ; ++i)
{
    double sum = 0;
    int R = matrix.size();
    for (int j = 0 ; j < R ; ++j)
    {
        sum += matrix[j][i]; // sum of the column

        if(matrix[j][i] !=0 )
        {
            N = N+1;
        }
        else if(matrix[j][i] == 0 )
        {
            N = N;
        }
    }

    total_column.push_back(sum);
    mean_column.push_back(sum/(R));
    //cout << "sum of column " << i << " = " << total_column[i] << endl;
    //cout << "mean sum of column " << i << " = " << mean_column[i] <<
    endl;
}

...
...

double SSE = SST-SSA;

int df_model = C-1;
int df_total = N-1;
int df_error = df_total - df_model;

double s1_square = SSA/(df_model);
double s1 = SSE/(df_error);
double computed_f = s1_square/s1;

int r1 = df_model;
int r2 = df_error;
```

```

double p_value = 1 - Fcdf(computed_f,r1,r2);
double r_square = 1 - SSE/SST;

cout << "\nSource" << setw(10) << "DF" << setw(23) << "SS" << setw(23) << "
    MS" << setw(23) << "F" << setw(23) << "P" << endl;
cout << "\nModel" << setw(10) << df_model << setw(23) << SSA << setw(23) <<
    s1_square << setw(23) << computed_f << setw(23) << p_value << endl;
cout << "\nError" << setw(10) << df_error << setw(23) << SSE << setw(23) <<
    s1 << setw(23) << endl;
cout << "\nTotal" << setw(10) << df_total << setw(23) << SST << endl;

cout << "\nR-Square" << setw(23) << "Grand Mean" << setw(23) << "Root MSE"
    << endl;
cout << r_square << setw(23) << y_bar << setw(23) << sqrt(s1) << endl;
}

```

Code 54: *src/statistics.cpp*

To compute the cdf of F -distribution is not an easy task. We need to use numerical integration of the pdf of F -distribution, most people will use a well-tested library like **Boost** that provides high accuracy and handles potential numerical issues. We use manual implementation made by **Lewis Van Winkle** with zlib license, he created a C file that compute incomplete Beta function, that is very important in order to compute the cdf of F -distribution and student's t -distribution by continued fraction using Lentz's algorithm. The article can be read here: <https://codeplea.com/incomplete-beta-function-c>

xxiv. Two-Way Analysis of Variance (ANOVA)

[SI*] Consider a situation where it is of interest to study the effects of two factors, A and B , on some response. For example, in a chemical experiment, we would like to vary simultaneously the reaction pressure and reaction time and study the effect of each on the yield. In a biological experiment, it is of interest to study the effects of drying time and temperature on the amount of solids (percent by weight) left in samples of yeast.

[SI*] Interaction and experimental error are separated in the two-factor experiment only if multiple observations are taken at the various treatment combinations. For maximum efficiency, there should be the same number n of observations at each combination. These should be true replications, not just repeated measurements.

[SI*] **Two-Factor Analysis of Variance**

To present general formulas for the analysis of variance of a two-factor experiment using repeated observations in a completely randomized design, we shall consider the case of n replications of the treatment combinations determined by a levels of factor A and b levels of factor B . The observations may be classified by means of a rectangular array where the rows represent the levels of factor A and the columns represent the levels of factor B . Each treatment combination defines a cell in our array. thus, we have ab cells, each cell containing n observations. Denoting the k -th observation taken at the i -th level of factor A and the j -th level of factor B by y_{ijk} . The observations in the (ij) -th cell constitute a random sample

A	B				Total	Mean
	1	2	...	b		
1	y_{111}	y_{121}	...	y_{1b1}	$Y_{1..}$	$\bar{y}_{1..}$
	y_{112}	y_{122}	...	y_{1b2}		
	$\vdots$	$\vdots$		$\vdots$		
	y_{11n}	y_{12n}	...	y_{1bn}		
2	y_{211}	y_{221}	...	y_{2b1}	$Y_{2..}$	$\bar{y}_{2..}$
	y_{212}	y_{222}	...	y_{2b2}		
	$\vdots$	$\vdots$		$\vdots$		
	y_{21n}	y_{22n}	...	y_{2bn}		
$\vdots$	$\vdots$	$\vdots$		$\vdots$	$\vdots$	$\vdots$
a	y_{a11}	y_{a21}	...	y_{ab1}	$Y_{a..}$	$\bar{y}_{a..}$
	y_{a12}	y_{a22}	...	y_{ab2}		
	$\vdots$	$\vdots$		$\vdots$		
	y_{a1n}	y_{a2n}	...	y_{abn}		
Total	$Y_{.1}$	$Y_{.2}$	...	$Y_{.b}$	$Y_{...}$	$\bar{y}_{...}$
Mean	$\bar{y}_{.1}$	$\bar{y}_{.2}$	...	$\bar{y}_{.b}$		

Table 4.9: two-Factor Experiment with n Replications.

of size n from a population that is assumed to be normally distributed with mean μ_{ij} and variance σ^2 . All ab populations are assumed to have the same variance σ^2 .

Let us define the following useful symbols:

Y_{ij} = sum of the observations in the (ij) -th cell,
 $Y_{i..}$ = sum of the observations for the i -th level of factor A ,
 $Y_{.j.}$ = sum of the observations for the j -th level of factor B ,
 $Y_{...}$ = sum of all abn observations,
 $\bar{y}_{ij}$ = mean of the observations in the (ij) th cell,
 $\bar{y}_{i..}$ = mean of the observations for the (i) th level of factor A ,
 $\bar{y}_{.j.}$ = mean of the observations for the (j) th level of factor B ,
 $\bar{y}_{...}$ = mean of all abn observations.

Unlike in the one-factor situation, here we are assuming that the populations, where n independent identically distributed observations are taken, are combinations of factors. Also we will assume throughout that an equal number (n) of observations are taken at each factor combination.

[SI*] **Model and Hypotheses for the Two-Factor Problem**

Each observation may be written in the form

$$y_{ijk} = \mu_{ij} + \epsilon_{ijk}$$

where ϵ_{ijk} measures the deviations of the observed y_{ijk} values in the (ij) -th cell from the population mean μ_{ij} . If we let $(\alpha\beta)_{ij}$ denote the interaction effect of the i -th level of factor A and the j -th level of factor B , α_i the effect of the i -th level of factor A , β_j the effect of the j -th level of factor B , and μ the overall mean, we can write

$$\mu_{ij} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij}$$

and then

$$y_{ijk} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + \epsilon_{ijk}$$

on which we impose the restrictions

$$\sum_{i=1}^a a_i = 0, \quad \sum_{j=1}^b \beta_j = 0, \quad \sum_{i=1}^a (\alpha\beta)_{ij} = 0, \quad \sum_{j=1}^b (\alpha\beta)_{ij} = 0$$

The three hypotheses to be tested are as follows:

1. The first hypothesis

$$H'_0 : \alpha_1 = \alpha_2 = \cdots = \alpha_a = 0$$

$$H'_1 : \text{At least one of } \alpha_i \text{ is not equal to zero}$$

2. The second hypothesis

$$H''_0 : \beta_1 = \beta_2 = \cdots = \beta_b = 0$$

$$H''_1 : \text{At least one of } \beta_j \text{ is not equal to zero}$$

3. The third hypothesis

$$H'''_0 : (\alpha\beta)_{11} = (\alpha\beta)_{12} = \cdots = (\alpha\beta)_{ab} = 0$$

$$H'''_1 : \text{At least one of } (\alpha\beta)_{ij} \text{ is not equal to zero}$$

Theorem 4.27: Sum-of-Squares Identity

The formula is

$$\begin{aligned} \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^n (y_{ijk} - \bar{y}_{...})^2 &= bn \sum_{i=1}^a (\bar{y}_{i..} - \bar{y}_{...})^2 + an \sum_{j=1}^b (\bar{y}_{.j.} - \bar{y}_{...})^2 \\ &\quad + n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...})^2 + \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^n (y_{ijk} - \bar{y}_{ij.})^2 \end{aligned}$$

Symbolically we write the sum-of-squares identity as

$$SST = SSA + SSB + SS(AB) + SSE$$

where SSA and SSB are called the sums of squares for the main effects A and B , respectively, $SS(AB)$ is called the interaction sum of squares for A and B , and SSE is the error sum of squares. The degrees of freedom are partitioned according to the identity

$$abn - 1 = (a - 1) + (b - 1) + (a - 1)(b - 1) + ab(n - 1)$$

[SI*] Formation of Mean Squares

If we divide each of the sums of squares on the right side of the sum-of-squares identity by its corresponding number of degrees of freedom, we obtain the four statistics

$$\begin{aligned} S_1^2 &= \frac{SSA}{a - 1} \\ S_2^2 &= \frac{SSB}{b - 1} \\ S_3^2 &= \frac{SS(AB)}{(a - 1)(b - 1)} \\ S^2 &= \frac{SSE}{ab(n - 1)} \end{aligned}$$

All of these variance estimates are independent estimates of σ^2 under the condition that there are no effects α_i , β_j , and, of course, $(\alpha\beta)_{ij}$. If we interpret the sums of squares as functions of the independent random variables $y_{111}, y_{112}, \dots, y_{abn}$, it is not difficult to verify that

$$\begin{aligned} E(S_1^2) &= E\left[\frac{SSA}{a - 1}\right] = \sigma^2 + \frac{nb}{a - 1} \sum_{i=1}^a \alpha_i^2 \\ E(S_2^2) &= E\left[\frac{SSB}{b - 1}\right] = \sigma^2 + \frac{na}{b - 1} \sum_{j=1}^b \beta_j^2 \\ E(S_3^2) &= E\left[\frac{SS(AB)}{(a - 1)(b - 1)}\right] = \sigma^2 + \frac{n}{(a - 1)(b - 1)} \sum_{i=1}^a \sum_{j=1}^b (\alpha\beta)_{ij}^2 \\ E(S^2) &= E\left[\frac{SSE}{ab(n - 1)}\right] = \sigma^2 \end{aligned}$$

from which we immediately observe that all four estimates of σ^2 are unbiased when H'_0 , H''_0 , and H'''_0 are true.

Definition 4.67: F-Test for Factor A

To test the hypothesis H'_0 , that the effects of factors A are all equal to zero, we compute the following ratio:

$$f_1 = \frac{s_1^2}{s^2}$$

which is a value of the random variable F_1 having the F -distribution with $a - 1$ and $ab(n - 1)$ degrees of freedom when H'_0 is true. The null hypothesis is rejected at the α -level of significance when $f_1 > f_\alpha[a - 1, ab(n - 1)]$.

Definition 4.68: F-Test for Factor B

To test the hypothesis H''_0 , that the effects of factors B are all equal to zero, we compute the following ratio:

$$f_2 = \frac{s_2^2}{s^2}$$

which is a value of the random variable F_2 having the F -distribution with $b - 1$ and $ab(n - 1)$ degrees of freedom when H''_0 is true. The null hypothesis is rejected at the α -level of significance when $f_2 > f_\alpha[b - 1, ab(n - 1)]$.

Definition 4.69: F-Test for Interaction

To test the hypothesis H'''_0 , that the interaction effects are all equal to zero, we compute the following ratio:

$$f_3 = \frac{s_3^2}{s^2}$$

which is a value of the random variable F_3 having the F -distribution with $(a - 1)(b - 1)$ and $ab(n - 1)$ degrees of freedom when H'''_0 is true. We conclude that, at the α -level of significance, interaction is present when $f_3 > f_\alpha[(a - 1)(b - 1), ab(n - 1)]$.

Source of Variation	Sum of Squares	Degrees of Freedom	Mean Square	Computed f
A	SSA	$a - 1$	$s_1^2 = \frac{SSA}{a-1}$	$f_1 = \frac{s_1^2}{s^2}$
B	SSB	$b - 1$	$s_2^2 = \frac{SSB}{b-1}$	$f_2 = \frac{s_2^2}{s^2}$
AB	$SS(AB)$	$(a - 1)(b - 1)$	$s_3^2 = \frac{SS(AB)}{(a-1)(b-1)}$	$f_3 = \frac{s_3^2}{s^2}$
Error	SSE	$ab(n - 1)$	$s^2 = \frac{SSE}{ab(n-1)}$	
Total	SST	$abn - 1$		

Table 4.10: Analysis of Variance for the Two-Factor Experiment with n Replications.

xxv. Compute Two-Way ANOVA with SymIntegration

In an experiment conducted to determine which of 3 different missile systems is preferable, the propellant burning rate for 24 static firings was measured. Four different propellant types were used. The experiment yielded duplicate observations of burning rates at each combination of the treatments.

Missile System	Propellant Type			
	b_1	b_2	b_3	b_4
a_1	34.0	30.1	29.8	29.0
	32.7	32.8	26.7	28.9
a_2	32.0	30.2	28.7	27.6
	33.2	29.8	28.1	27.8
a_3	28.4	27.3	29.7	28.8
	29.3	28.9	27.3	29.1

Table 4.11: Propellant Burning Rates.

Test the following hypotheses:

- (a) H'_0 : there is no difference in the mean propellant burning rates when different missile systems are used.
- (b) H''_0 : there is no difference in the mean propellant burning rates of the 4 propellant types.
- (c) H'''_0 : there is no interaction between the different missile systems and the different propellant types.

Solution:

- (a) The first hypothesis

$$H'_0 : \alpha_1 = \alpha_2 = \alpha_3 = 0$$

$$H'_1 : \text{At least one of } \alpha_i \text{ is not equal to zero}$$

- (b) The second hypothesis

$$H''_0 : \beta_1 = \beta_2 = \beta_3 = \beta_4 = 0$$

$$H''_1 : \text{At least one of } \beta_j \text{ is not equal to zero}$$

- (c) The third hypothesis

$$H'''_0 : (\alpha\beta)_{11} = (\alpha\beta)_{12} = \cdots = (\alpha\beta)_{34} = 0$$

$$H'''_1 : \text{At least one of } (\alpha\beta)_{ij} \text{ is not equal to zero}$$

With the sum-of-squares formula to do the computation, we will have this resulting table for the analysis of variance:

Source of Variation	Sum of Squares	Degrees of Freedom	Mean Square	Computed f
Missile system	14.52	2	7.26	5.84
Propellant type	40.08	3	13.36	10.75
Interaction	22.16	6	3.69	2.97
Error	14.91	12	1.24	
Total	91.68	23		

Table 4.12: Analysis of Variance for the Two-Factor Experiment Data.

We can compute the "model" (with 11 degrees of freedom), "error", "total" analysis of variance before we test the system, type, and system by type interaction.

In SymIntegration to compute two-way analysis of variance (ANOVA) we can use this function: **twoway_ANOVA(vector<vector<double>> matrix, int a, int b)**

The input is a matrix that is defined as **vector<vector<double>>** from the standard C++ library, then a is the number of levels of factor A (in this case $a = 3$), and b is the number of levels of factor B (in this case $b = 4$). It is a **void** function, so it returns only the table and nothing else.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two-Way ANOVA ]# ./main
*****
Two-way table of averages
      33.350000      31.450000      28.250000      28.950000      30.500000
      32.600000      30.000000      28.400000      27.700000      29.675000
      28.850000      28.100000      28.500000      28.950000      28.600000
-----
      31.600000      29.850000      28.383333      28.533333
*****

Hypotheses for the Two-Factor Problem
H0':  $\alpha_{\{1\}} = \alpha_{\{2\}} = \dots = \alpha_{\{a\}} = 0$ 
H1': At least one of the  $\alpha_{\{i\}}$  is not equal to zero

H0'':  $\beta_{\{1\}} = \beta_{\{2\}} = \dots = \beta_{\{b\}} = 0$ 
H1'': At least one of the  $\beta_{\{j\}}$  is not equal to zero

H0''':  $(\alpha\beta)_{\{11\}} = (\alpha\beta)_{\{12\}} = \dots = (\alpha\beta)_{\{ab\}} = 0$ 
H1''': At least one of the  $(\alpha\beta)_{\{ij\}}$  is not equal to zero
*****

Analysis of Variance
Source      DF      SS      MS      F      P
Model       11      76.768333      6.978939      5.616853      0.002986
Error       12      14.910000      1.242500
Total       23      91.678333

R-Square    0.837366      Grand Mean    29.591667      Root MSE    1.114675

Analysis of Variance for the Two-Factor Experiment with n replications
Source      DF      Type III SS      MS      F      P
A           2      14.523333      7.261667      5.844400      0.016898
B           3      40.081667      13.360556      10.752962      0.001020
A * B       6      22.163333      3.693889      2.972949      0.051168

Time taken by function: 1542 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Two-Way ANOVA ]#

```

Figure 4.49: The computation of two-way analysis-of-variance / ANOVA for the propellant rate data. (*SymIntegration/Examples/Statistics/Test SymIntegration Two-Way ANOVA/main.cpp*).

Learning from the two-way ANOVA table we can say that :

The F -test on the model ($P = 0.002986$) is testing the accumulation of the two main effects and the interaction.

- Reject H_0' and conclude that different missile systems result in different mean propellant burning rates. The P -value is approximately 0.016898.
- Reject H_0'' and conclude that the mean propellant burning rates are not the same for the four propellant types. The P -value is approximately 0.0010.
- Interaction is barely insignificant at the 0.05 level, but the P -value of approximately 0.05116 would indicate that interaction must be taken seriously.

At this point we should draw some type of interpretation of the interaction. It should be emphasized that statistical significance of a main effect merely implies that marginal means are significantly different. However, if we take a look at the two-way table of averages it is apparent that more important information exists in the body of the table, trends that are inconsistent with the trend depicted by marginal averages. The table of averages certainly suggests that the effect of propellant type depends on the system being used. For example, for system 3 the propellant-type effect does not appear to be important, although it does have a large effect if either system 1 or system 2 is used. This explains the "significant" interaction between these two factors.

xxvi. Three-Way Analysis of Variance (ANOVA)

[SI*] We consider an experiment with three factors, A, B , and C , at a, b , and c levels, respectively, in a completely randomized experimental design. Assume again that we have n observations for each of the abc treatment combinations.

Definition 4.70: Model for the Three-Factor Experiment

The model for the three-factor experiment is

$$y_{ijkl} = \mu + \alpha_i + \beta_j + \gamma_k + (\alpha\beta)_{ij} + (\alpha\gamma)_{ik} + (\beta\gamma)_{jk} + (\alpha\beta\gamma)_{ijk} + \epsilon_{ijkl}$$

with $i = 1, 2, \dots, a$; $j = 1, 2, \dots, b$; $k = 1, 2, \dots, c$; and $l = 1, 2, \dots, n$, where α_i, β_j , and γ_k are the main effects and $(\alpha\beta)_{ij}, (\alpha\gamma)_{ik}$, and $(\beta\gamma)_{jk}$ are the two-factor interaction effects that have the same interpretation as in the two-factor experiment.

The term $(\alpha\beta\gamma)_{ijk}$ is called the three-factor interaction effect, a term that represents a nonadditivity of the $(\alpha\beta)_{ij}$ over the different levels of the factor C . The sum of all main effects is zero and the sum over any subscript of the two- and three-factor interaction effects is zero. In many experimental situations, these higher-order interactions are insignificant and their mean squares reflect only random variation, but we shall outline the analysis in its most general form.

Definition 4.71: Sum of Squares for a Three-Factor Experiment

The general philosophy concerning the analysis is the same as the one- and two-factor experiments. The sum of squares is partitioned into eight terms, each representing a source of variation from which we obtain independent estimates of σ^2 when all the main effects and interaction effects are zero. If the effects of any given factor or interaction are not all zero, then the mean square will estimate the error variance plus a component due to the systematic effect in question.

$$SSA = bcn \sum_{i=1}^a (\bar{y}_{i...} - \bar{y}_{....})^2$$

$$SSB = acn \sum_{j=1}^b (\bar{y}_{.j..} - \bar{y}_{....})^2$$

$$SSC = abn \sum_{k=1}^c (\bar{y}_{..k.} - \bar{y}_{....})^2$$

$$SS(AB) = cn \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij..} - \bar{y}_{i...} - \bar{y}_{.j..} + \bar{y}_{....})^2$$

$$SS(AC) = bn \sum_{i=1}^a \sum_{k=1}^c (\bar{y}_{i.k.} - \bar{y}_{i...} - \bar{y}_{..k.} + \bar{y}_{....})^2$$

$$SS(BC) = an \sum_{j=1}^b \sum_{k=1}^c (\bar{y}_{.jk.} - \bar{y}_{.j..} - \bar{y}_{..k.} + \bar{y}_{....})^2$$

$$SS(ABC) = n \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^c (\bar{y}_{ijk.} - \bar{y}_{ij..} - \bar{y}_{i.k.} - \bar{y}_{.jk.} + \bar{y}_{i...} + \bar{y}_{.j..} + \bar{y}_{..k.} - \bar{y}_{....})^2$$

$$SST = \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^c \sum_{l=1}^n (y_{ijkl} - \bar{y}_{....})^2$$

$$SSE = \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^c \sum_{l=1}^n (y_{ijkl} - \bar{y}_{ijk.})^2$$

The averages in the formulas are defined as follows:

$\bar{y}_{....}$ = average of all $abcn$ observations.

$\bar{y}_{i...}$ = average of the observations for the i -th level of factor A .

$\bar{y}_{.j..}$ = average of the observations for the j -th level of factor B .

$\bar{y}_{..k.}$ = average of the observations for the k -th level of factor C .

$\bar{y}_{ij..}$ = average of the observations for the i -th level of factor A and the j -th level of factor B .

$\bar{y}_{i.k.}$ = average of the observations for the i -th level of factor A and the k -th level of factor C .

$\bar{y}_{.jk.}$ = average of the observations for the j -th level of factor B and the k -th level of factor C .

$\bar{y}_{ijk.}$ = average of the observations for the (ijk) -th treatment combination.

Source of Variation	Sum of Squares	Degrees of Freedom	Mean Square	Computed f
A	SSA	$a - 1$	$s_1^2 = \frac{SSA}{a-1}$	$f_1 = \frac{s_1^2}{s_2^2}$
B	SSB	$b - 1$	$s_2^2 = \frac{SSB}{b-1}$	$f_2 = \frac{s_2^2}{s_2^2}$
C	SSC	$c - 1$	$s_3^2 = \frac{SSC}{c-1}$	$f_3 = \frac{s_3^2}{s_2^2}$
AB	$SS(AB)$	$(a - 1)(b - 1)$	$s_4^2 = \frac{SS(AB)}{(a-1)(b-1)}$	$f_4 = \frac{s_4^2}{s_2^2}$
AC	$SS(AC)$	$(a - 1)(c - 1)$	$s_5^2 = \frac{SS(AC)}{(a-1)(c-1)}$	$f_5 = \frac{s_5^2}{s_2^2}$
BC	$SS(BC)$	$(b - 1)(c - 1)$	$s_6^2 = \frac{SS(BC)}{(b-1)(c-1)}$	$f_6 = \frac{s_6^2}{s_2^2}$
ABC	$SS(ABC)$	$(a - 1)(b - 1)(c - 1)$	$s_7^2 = \frac{SS(ABC)}{(a-1)(b-1)(c-1)}$	$f_7 = \frac{s_7^2}{s_2^2}$
Error	SSE	$abc(n - 1)$	$s^2 = \frac{SSE}{ab(n-1)}$	
Total	SST	$abcn - 1$		

Table 4.13: Analysis of Variance for the Three-Factor Experiment with n Replications.

xxvii. Compute Three-Way ANOVA with SymIntegration

In the production of a particular material, three variables are of interest: A , the operator effect (three operators); B , the catalyst used in the experiment (three catalysts); and C , the washing time of the product following the cooling process (15 minutes and 20 minutes). Three runs were made at each combination of factors. It was felt that all interactions among the factors should be studied. Perform an analysis of variance to test for significant effects.

Solution:

The three-way ANOVA table computation result:

The ANOVA table shows that none of the interactions show a significant effect at the $\alpha = 0.05$ level.

To understand how we do the hypothesis testing with the P -value, we should take a look at:

1. The P -value for A is 0.000126, if we have chosen $\alpha = 0.05$, it means that we reject the null hypothesis that is saying

$$H'_0 : \alpha_1 = \alpha_2 = \alpha_3 = 0$$

$$H'_1 : \text{At least one of } \alpha_i \text{ is not equal to zero}$$

It means that the operator / factor A is significant in production of a particular material.

		Washing Time, C					
		15 Minutes			20 Minutes		
		Catalyst, B			Catalyst, B		
Operator, A	1	2	3	1	2	3	
1	10.7	10.3	11.2	10.9	10.5	12.2	
	10.8	10.2	11.6	12.1	11.1	11.7	
	11.3	10.5	12.0	11.5	10.3	11.0	
2	11.4	10.2	10.7	9.8	12.6	10.8	
	11.8	10.9	10.5	11.3	7.5	10.2	
	11.5	10.5	10.2	10.9	9.9	11.5	
3	13.6	12.0	11.1	10.7	10.2	11.9	
	14.1	11.6	11.0	11.7	11.5	11.6	
	14.5	11.5	11.5	12.7	10.9	12.2	

Table 4.14: Data for production of a particular material with three factors.

2. The P -value for B is 0.0010, if we have chosen $\alpha = 0.05$, it means that we reject the null hypothesis that is saying

$$H_0'' : \beta_1 = \beta_2 = \beta_3 = 0$$

$$H_1'' : \text{At least one of } \beta_j \text{ is not equal to zero}$$

It means that the catalyst effect / factor B is significant in production of a particular material.

3. The P -value for C is 0.1686, if we have chosen $\alpha = 0.05$, it means that we do not reject the null hypothesis that is saying

$$H_0''' : \gamma_1 = \gamma_2 = 0$$

$$H_1''' : \text{At least one of } \gamma_k \text{ is not equal to zero}$$

It means that the washing time / factor C is not significant in production of a particular material.

However, the P -value for BC is 0.0610; thus, it should not be ignored. The operator and catalyst effects are significant, while the effect of washing time is not significant.

Suppose we want to see the impact of interaction BC , about dealing with the effect that the interaction between catalyst and washing time is having on the test on the washing time main effect (factor C).

The BC interaction is significant at approximately 0.06 level. Suppose, however, we observe a two-way table of means between BC (the two-way table is provided in the result from SymIntegration' three-way ANOVA function).

It is clear why washing time was found not to be significant. A non-thorough analyst may get the impression that washing time can be eliminated from any future study which yield is being

Source of Variation	Sum of Squares	Degrees of Freedom	Mean Square	Computed f	P -Value
A	13.98	2	6.99	11.64	0.000126
B	10.18	2	5.09	8.48	0.00096
C	1.185	1	1.185	1.97	0.168585
AB	4.77	4	1.19	1.987	0.117196
AC	2.91	2	1.46	2.43	0.1027
BC	3.63	2	1.82	3.03	0.060982
ABC	4.91	4	1.23	2.04	0.1089
Error	21.613	36	0.60		
Total	63.19	53			

Table 4.15: Analysis of Variance for the Three-Factor Experiment with $n = 3$ Replications.

measured. However, it is obvious how the effect of washing time changes from a negative effect for the first catalyst to what appears to be a positive effect for the third catalyst. If we merely focus on the data for catalyst 1, a simple comparison between the means at the two washing times will produce a simple t -statistic:

$$t = \frac{12.19 - 11.29}{\sqrt{0.6 \left(\frac{2}{9}\right)}} = 2.5$$

which is significant at a level less than 0.02. Thus, an important negative effect of washing time for catalyst 1 might very well be ignored if the analyst makes the incorrect broad interpretation of the insignificant F -ratio for washing time.

To analyze the three variables in the three-way ANOVA in order to create the source code to compute the ANOVA table, we should take a note that:

1. We start with the input as a square matrix with the number of rows, R , and the number of columns, C .
2. The factor A will be the main row, there will be R/a row for each factor $i = 1, 2, a$ that will represents factor A , in this case $a = 3$.
3. The factor B will be composed of b columns, since it is three-way then the total columns will be $b \times c$ columns each number of b columns corresponds to each factor C , in this case 1 column of factor C has 3 columns from factor B .
4. The total number of entries / data is $N = abc n$.

We will depends on a lot of **for** loop with linear algebra in the source code.

In SymIntegration to compute three-way analysis of variance (ANOVA) we can use this function:

threeway_ANOVA(vector<vector<double>> matrix, int a, int b, int c)

The input is a matrix that is defined as **vector<vector<double>>** from the standard C++ library, then a is the number of levels of factor A (in this case $a = 3$), b is the number of levels of factor

B (in this case $b = 3$), and c is the number of levels of factor C (in this case $c = 2$). It is a **void** function, so it returns only the table and nothing else.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Three-Way ANOVA ]# ./main
*****
Two-way table of averages, A (row) and B (column)
  11.216667      10.483333      11.616667      | 11.105556
  11.116667      10.266667      10.650000      | 10.677778
  12.883333      11.283333      11.550000      | 11.905556
-----
      11.738889      10.677778      11.272222
*****
Two-way table of averages, A (row) and C (column)
  10.955556      11.255556      | 11.105556
  10.855556      10.500000      | 10.677778
  12.322222      11.488889      | 11.905556
-----
      11.377778      11.081481
*****
Two-way table of averages, B (row) and C (column)
  12.188889      11.288889      | 11.738889
  10.855556      10.500000      | 10.677778
  11.088889      11.455556      | 11.272222
-----
      11.377778      11.081481
*****

```

Figure 4.50: The computation of three-way analysis-of-variance / ANOVA for the production of a particular material showing all the two-way tables from all possible combinations of 2 factors (*SymIntegration/Examples/Statistics/Test SymIntegration Three-Way ANOVA/main.cpp*).

```

Hypotheses for the Three-Factor Problem
H0^{(1)}:  $\alpha_{\{1\}} = \alpha_{\{2\}} = \dots = \alpha_{\{a\}} = 0$ 
H1^{(1)}: At least one of the  $\alpha_{\{i\}}$  is not equal to zero

H0^{(2)}:  $\beta_{\{1\}} = \beta_{\{2\}} = \dots = \beta_{\{b\}} = 0$ 
H1^{(2)}: At least one of the  $\beta_{\{j\}}$  is not equal to zero

H0^{(3)}:  $\gamma_{\{1\}} = \gamma_{\{2\}} = \dots = \gamma_{\{c\}} = 0$ 
H1^{(3)}: At least one of the  $\gamma_{\{k\}}$  is not equal to zero

H0^{(4)}:  $(\alpha\beta)_{\{11\}} = (\alpha\beta)_{\{12\}} = \dots = (\alpha\beta)_{\{ab\}} = 0$ 
H1^{(4)}: At least one of the  $(\alpha\beta)_{\{ij\}}$  is not equal to zero

H0^{(5)}:  $(\alpha\gamma)_{\{11\}} = (\alpha\gamma)_{\{12\}} = \dots = (\alpha\gamma)_{\{ac\}} = 0$ 
H1^{(5)}: At least one of the  $(\alpha\gamma)_{\{ik\}}$  is not equal to zero

H0^{(6)}:  $(\beta\gamma)_{\{11\}} = (\beta\gamma)_{\{12\}} = \dots = (\beta\gamma)_{\{bc\}} = 0$ 
H1^{(6)}: At least one of the  $(\beta\gamma)_{\{jk\}}$  is not equal to zero

H0^{(7)}:  $(\alpha\beta\gamma)_{\{111\}} = (\alpha\beta\gamma)_{\{112\}} = \dots = (\alpha\beta\gamma)_{\{abc\}} = 0$ 
H1^{(7)}: At least one of the  $(\alpha\beta\gamma)_{\{ijk\}}$  is not equal to zero

*****

Analysis of Variance

Source      DF      SS      MS      F      P
Model       17      41.579259  2.445839  4.073883  0.000194
Error       36      21.613333  0.600370
Total       53      63.192593

R-Square    Grand Mean    Root MSE
0.657977    11.229630     0.774836

Analysis of Variance for the Three-Factor Experiment with n replications

Source      DF      Type III SS      MS      F      P
A           2      13.982593      6.991296  11.644972  0.000126
B           2      10.182593      5.091296  8.480259  0.000960
C           1      1.185185      1.185185  1.974090  0.168585
A * B       4      4.774074      1.193519  1.987970  0.117196
A * C       2      2.913704      1.456852  2.426589  0.102655
B * C       2      3.633704      1.816852  3.026218  0.060982
A * B * C   4      4.907407      1.226852  2.043492  0.108896
Error       36      21.613333      0.600370
-----
Total       53      63.192593

Time taken by function: 2697 microseconds

```

Figure 4.51: The computation of three-way analysis-of-variance / ANOVA for the production of a particular material (*SymIntegration/Examples/Statistics/Test SymIntegration Three-Way ANOVA/main.cpp*).

VII. LINEAR REGRESSION AND CORRELATION

i. Pearson's Correlation

[SI*] In Darrell Huff's book **How to Lie with Statistics**, he refers to a study that claimed that non-smoking college students gained better grades than students who smoked. The study concluded that 'smoking leads to poor college grades.' Huff questioned the findings of the study, arguing that instead of smoking being the cause of poor grades, perhaps 'low grades depressed students and caused them to smoke,' or maybe 'students with lower grades were more sociable and therefore more likely to smoke.' Measuring the relationship between two sets of data is a useful tool to establish how one variable changes with another, but it is limited; it does not tell you why this relationship exists. the data in the study clearly showed a relationship between smoking and poor grades, but it does not explain why this relationship exists.

Data that consists of measurements of two variables taken from each individual in a sample is called bivariate data; the relationship between the two variables is referred to as the correlation.

For bivariate data we try to classify one of two variables as 'independent' and the other as 'dependent.' The independent variable is the one that can be controlled by the person conducting the experiment or study; it is hypothesised to 'cause' some kind of effect to the dependent variable. the dependent variable is the variable that is just observed without being controlled, and is supposed to show the 'effect.'

[SI*] The line of best fit on a scatter diagram is drawn to give the best representation of the correlation between the two variables. The line of best fit is the one that has an approximately even spread of the data points either side of it.

[SI*] **Pearson's product moment correlation coefficient**

Pearson product moment correlation coefficient (PMCC) is a number, usually denoted by r , which can take any value between -1 and $+1$. Its sign indicates the type of correlation, and its magnitude (size) indicates the strength of correlation.

1. $r = +1$ means that there is perfect positive correlation
2. $r = 0$ means that there is no correlation
3. $r = -1$ means that there is perfect negative correlation

If $-0.5 < r < 0.5$, it is difficult to draw a line of best fit that has any meaning, because the data is just too scattered.

The formula to compute the PMCC is

$$r = \frac{S_{xy}}{S_x S_y} \quad (4.115)$$

where S_{xy} is the covariance of x and y , S_x is the standard deviation of the x values, and S_y is the standard deviation of the y values.

The formula that act as the foundation blocks are:

$$\bar{x} = \frac{\sum x}{n} \quad (4.116)$$

$$\bar{y} = \frac{\sum y}{n} \quad (4.117)$$

$$S_x = \sqrt{\frac{\sum x^2}{n} - (\bar{x})^2} \quad (4.118)$$

$$S_y = \sqrt{\frac{\sum y^2}{n} - (\bar{y})^2} \quad (4.119)$$

$$S_{xy} = \frac{\sum xy}{n} - (\bar{x})(\bar{y}) \quad (4.120)$$

with n as the number of observations / data points.
[SI*] The formula for the regression line is

$$y - \bar{y} = \frac{S_{xy}}{(S_x)^2}(x - \bar{x}) \quad (4.121)$$

ii. Linear Regression

[SI*] Often, in practice, we need to solve problems involving sets of variables when it is known that there exists some inherent relationship among the variables. For example, in an industrial situation it may be known that the tar content in the outlet stream in a chemical process is related to the inlet temperature. It may be of interest to develop a method of prediction, that is, a procedure for estimating the tar content for various levels of the inlet temperature from experimental information. In this case the tar content is the dependent variable, or response, and the inlet temperature is the independent variable, or regressor. A reasonable form of a relationship between the response Y and the regressor x is the linear relationship

$$Y = \beta_0 + \beta_1 x$$

with β_0 as the intercept and β_1 as the slope. If the relationship is exact, then it is a deterministic relationship between two scientific variables and there is no random or probabilistic component to it. However, the relationship is not deterministic (i.e., a given x does not always give the same value for Y).

The concept of regression analysis deals with finding the best relationship between Y and x , quantifying the strength of that relationship, and using methods that allow for prediction of the response values given values of the regressor x .

[SI*] In most application of regression, the linear equation $Y = \beta_0 + \beta_1 x$ is an approximation that is a simplification of something unknown and much more complicated.

Thus, an analysis of the relationship between Y and x requires the statement of a statistical model. The model includes the set

$$\{(x_i, y_i); i = 1, 2, \dots, n\}$$

of data involving n pairs of (x, y) values. The value y_i depends on x_i , via a linear structure that also has the random component involved.

The basis for the use of a statistical model relates to how the random variable Y moves with x and the random component.

Definition 4.72: Simple Linear Regression Model

the response Y is related to the independent variable x through the equation

$$Y = \beta_0 + \beta_1 x + \epsilon$$

β_0 and β_1 are unknown intercept and slope parameters, respectively, and ϵ is a random variable that is assumed to be distributed with $E(\epsilon) = 0$ and $\text{Var}(\epsilon) = \sigma^2$. The quantity σ^2 is often called the error variance or residual variance.

Definition 4.73: The Fitted Regression Line

Suppose we denote the estimates b_0 for β_0 and b_1 for β_1 . Then the estimated or fitted regression line is given by

$$\hat{y} = b_0 + b_1x$$

where $\hat{y}$ is the predicted or fitted value. Obviously, the fitted line is the estimate of the true regression line.

Definition 4.74: Residual: Error in Fit

Given a set of regression data $\{(x_i, y_i); i = 1, 2, \dots, n\}$ and a fitted model, $\hat{y}_i = b_0 + b_1x$, the i th residual ϵ_i , is given by

$$\epsilon_i = y_i - \hat{y}_i, \quad i = 1, 2, \dots, n$$

Definition 4.75: Estimating the Regression Coefficients

In order to find b_0 and b_1 , the estimates of β_0 and β_1 , so that the sum of squares of the residuals is a minimum. The residual sum of squares is often called the sum of squares of the errors about the regression line and is denoted by SSE . This minimization procedure for estimating the parameters is called the method of least squares.

Given the sample $\{(x_i, y_i); i = 1, 2, \dots, n\}$, the least squares estimates b_0 and b_1 of the regression coefficients β_0 and β_1 are computed from the formulas

$$b_1 = \frac{n \sum_{i=1}^n x_i y_i - (\sum_{i=1}^n x_i)(\sum_{i=1}^n y_i)}{n \sum_{i=1}^n x_i^2 - (\sum_{i=1}^n x_i)^2} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

and

$$b_0 = \frac{\sum_{i=1}^n y_i - b_1 \sum_{i=1}^n x_i}{n}$$

Thus, the estimated regression line is given by

$$\hat{y} = b_0 + b_1x$$

[SI*] In addition to the assumptions that the error term in the model

$$Y_i = \beta_0 + \beta_1 x_i + \epsilon_i$$

is a random variable with mean 0 and constant variance σ^2 , suppose that we make the further assumption that $\epsilon_1, \epsilon_2, \dots, \epsilon_n$ are independent from run to run in the experiment. This provides a foundation for finding the means and variances for the estimators of β_0 and β_1 .

It is important to remember that our values of b_0 and b_1 , based on a given sample of n observations, are only estimates of true parameters β_0 and β_1 . If the experiment is repeated over and over again, each time using the same fixed values of x , the resulting estimates of β_0 and β_1 will most likely differ from experiment to experiment. These different estimates

may be viewed as values assumed by the random variables B_0 and B_1 , while b_0 and b_1 are specific realizations.

Since the values of x remain fixed, the values of B_0 and B_1 depend on the variations in the values of y or, more precisely, on the values of the random variables, $Y_1, Y_2, \dots, Y_n$. The distributional assumptions imply that the $Y_i, i = 1, 2, \dots, n$, are also independently distributed, with mean $\mu_{Y|x} = \beta_0 + \beta_1 x_i$ and equal variances σ^2 ; that is,

$$\sigma_{Y|x_i}^2 \quad \text{for } i = 1, 2, \dots, n$$

[SI*] **Mean and Variance of Estimators**

We show that the estimator B_1 is unbiased for β_1 and demonstrate the variances of both B_0 and B_1 . This will begin a series of developments that lead to hypothesis testing and confidence interval estimation on the intercept and slope.

Since the estimator

$$B_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \frac{\sum_{i=1}^n (x_i - \bar{x})Y_i}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

is of the form $\sum_{i=1}^n c_i Y_i$, where

$$c_i = \frac{x_i - \bar{x}}{\sum_{i=1}^n (x_i - \bar{x})^2}, \quad i = 1, 2, \dots, n$$

we may conclude that B_1 has a $n(\mu_{B_1}, \sigma_{B_1})$ distribution with

$$\begin{aligned} \mu_{B_1} &= \frac{\sum_{i=1}^n (x_i - \bar{x})(\beta_0 + \beta_1 x_i)}{\sum_{i=1}^n (x_i - \bar{x})^2} = \beta_1 \\ \sigma_{B_1}^2 &= \frac{\sum_{i=1}^n (x_i - \bar{x})^2 \sigma_{Y_i}^2}{[\sum_{i=1}^n (x_i - \bar{x})^2]^2} = \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \end{aligned}$$

it can also be shown that the random variable B_0 is normally distributed with mean

$$\mu_{B_0} = \beta_0$$

and variance

$$\sigma_{B_0}^2 = \frac{\sum_{i=1}^n x_i^2}{n \sum_{i=1}^n (x_i - \bar{x})^2} \sigma^2$$

From the foregoing results, it is apparent that the least squares estimators for β_0 and β_1 are both unbiased estimators.

[SI*] The parameter σ^2 , the model error variance, reflects random variation or experimental error variation around the regression line. It is advantageous to use the notation

$$\begin{aligned} S_{xx} &= \sum_{i=1}^n (x_i - \bar{x})^2 \\ S_{yy} &= \sum_{i=1}^n (y_i - \bar{y})^2 \\ S_{xy} &= \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) \end{aligned}$$

Now we may write the error sum of squares as follows:

$$\begin{aligned}
 SSE &= \sum_{i=1}^n (y_i - b_0 - b_1 x_i)^2 \\
 &= \sum_{i=1}^n [(y_i - \bar{y}) - b_1(x_i - \bar{x})]^2 \\
 &= \sum_{i=1}^n (y_i - \bar{y})^2 - 2b_1 \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) + b_1^2 \sum_{i=1}^n (x_i - \bar{x})^2 \\
 &= S_{yy} - 2b_1 S_{xy} + b_1^2 S_{xx} \\
 &= S_{yy} - b_1 S_{xy}
 \end{aligned}$$

with $b_1 = \frac{S_{xy}}{S_{xx}}$.

Theorem 4.28: Unbiased Estimate of Variance

An unbiased estimate of σ^2 is

$$s^2 = \frac{SSE}{n-2} = \sum_{i=1}^n \frac{(y_i - \hat{y})^2}{n-2} = \frac{S_{yy} - b_1 S_{xy}}{n-2}$$

The important parameter σ^2 estimated by s^2 is called a mean squared error, depicting a type of mean (division by $n-2$) of the squared residuals.

[SI*] We know that B_1 follows a normal distribution. It turns out that under the normality assumption, we are able to conclude that $\frac{(n-2)S^2}{\sigma^2}$ is a chi-squared variable with $n-2$ degrees of freedom, independent of the random variable B_1 .

Thus, the statistic

$$T = \frac{(B_1 - \beta_1)/(\sigma/\sqrt{S_{xx}})}{S/\sigma} = \frac{B_1 - \beta_1}{S/\sqrt{S_{xx}}}$$

has a t -distribution with $n-2$ degrees of freedom. The statistic T can be used to construct a $100(1-\alpha)\%$ confidence interval for the coefficient β_1

Definition 4.76: Confidence Interval for β_1

A $100(1-\alpha)\%$ confidence interval for the parameter β_1 in the regression line $\mu_{Y|x} = \beta_0 + \beta_1 x$ is

$$b_1 - t_{\frac{\alpha}{2}} \frac{s}{\sqrt{S_{xx}}} < \beta_1 < b_1 + t_{\frac{\alpha}{2}} \frac{s}{\sqrt{S_{xx}}}$$

where $t_{\frac{\alpha}{2}}$ is a value of the t -distribution with $n-2$ degrees of freedom that leaves an area of $\frac{\alpha}{2}$ to the right.

[SI*] Confidence intervals and hypothesis testing on the coefficient β_0 may be established from the fact that B_0 is also normally distributed. It is not difficult to show that

$$T = \frac{B_0 - \beta_0}{S \sqrt{\frac{\sum_{i=1}^n x_i^2}{n S_{xx}}}}$$

has a t -distribution with $n - 2$ degrees of freedom from which we may construct a $100(1 - \alpha)\%$ confidence interval for β_0

Definition 4.77: Confidence Interval for β_0

A $100(1 - \alpha)\%$ confidence interval for the parameter β_0 in the regression line $\mu_{Y|x} = \beta_0 + \beta_1 x$ is

$$b_0 - t_{\frac{\alpha}{2}} \frac{s}{\sqrt{nS_{xx}}} \sqrt{\sum_{i=1}^n x_i^2} < \beta_0 < b_0 + t_{\frac{\alpha}{2}} \frac{s}{\sqrt{nS_{xx}}} \sqrt{\sum_{i=1}^n x_i^2}$$

where $t_{\frac{\alpha}{2}}$ is a value of the t -distribution with $n - 2$ degrees of freedom that leaves an area of $\frac{\alpha}{2}$ to the right.

[SI*] Coefficient of determination, represented by R^2 , is a measure of the proportion of variability explained by the fitted model. The analysis-of-variance approach makes use of the error sum of squares $SSE = \sum_{i=1}^n (x_i - \hat{y}_i)^2$ and the total corrected sum of squares $SST = \sum_{i=1}^n (y_i - \bar{y})^2$. The SSE value is the variation due to error, or variation unexplained. Clearly, if $SSE = 0$, all variation is explained. The quantity that represents variation explained is $SST - SSE$. The R^2 is

$$R^2 = 1 - \frac{SSE}{SST}$$

Note that if the fit is perfect, all residuals are zero, and thus $R^2 = 1.0$. But if SSE is only slightly smaller than SST , $R^2 \approx 0$.

The R^2 criterion is dangerous to use for comparing competing models for the same data set. Adding additional terms to the model decreases SSE and thus increases R^2 (or at least does not decrease it). This implies that R^2 can be made artificially high by an unwise practice of overfitting (i.e., the inclusion of too many model terms).

[SI*] **Prediction**

There are several reasons for building a linear regression. One, of course, is to predict response values at one or more values of the independent variable.

The equation $\hat{y} = b_0 + b_1 x$ may be used to predict or estimate the mean response $\mu_{Y|x_0}$ at $x = x_0$, where x_0 is not necessarily one of the prechosen values, or it may be used to predict a single value y_0 of the variable Y_0 , when $x = x_0$. We would expect the error of prediction to be higher in the case of a single predicted value than in the case where a mean is predicted. This, then, will affect the width of our intervals for the values being predicted.

Suppose that the experimenter wishes to construct a confidence interval for $\mu_{Y|x_0}$. We shall use the point estimator $\hat{Y}_0 = B_0 + B_1 x_0$ to estimate $\mu_{Y|x_0} = \beta_0 + \beta_1 x_0$. It can be shown that the sampling distribution of $\hat{Y}_0$ is normal with mean

$$\mu_{Y|x_0} = E(\hat{Y}_0) = E(B_0 + B_1 x_0) = \beta_0 + \beta_1 x_0 = \mu_{Y|x_0}$$

and variance

$$\sigma_{\hat{Y}_0}^2 = \sigma_{B_0 + B_1 x_0}^2 = \sigma_{\bar{Y} + B_1(x_0 - \bar{x})}^2 = \sigma^2 \left[\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{S_{xx}} \right]$$

the latter following from the fact that $\text{Cov}(\bar{Y}, B_1) = 0$. Thus, a $100(1 - \alpha)\%$ confidence

interval on the mean response $\mu_{Y|x_0}$ can now be constructed from the statistic

$$T = \frac{\hat{Y}_0 - \mu_{Y|x_0}}{S \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{S_{xx}}}}$$

which has a t -distribution with $n - 2$ degrees of freedom.

Definition 4.78: Confidence Interval for $\mu_{Y|x_0}$

A $100(1 - \alpha)\%$ confidence interval for the mean response $\mu_{Y|x_0}$ is

$$\hat{y}_0 - t_{\frac{\alpha}{2}} s \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{S_{xx}}} < \mu_{Y|x_0} < \hat{y}_0 + t_{\frac{\alpha}{2}} s \sqrt{\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{S_{xx}}}$$

where $t_{\frac{\alpha}{2}}$ is a value of the t -distribution with $n - 2$ degrees of freedom that leaves an area of $\frac{\alpha}{2}$ to the right.

[SI*] Prediction Interval

Another type of interval that is often misinterpreted and confused with that $\mu_{Y|x_0}$ is the prediction interval for a future observed response.

To obtain a prediction interval for any single value y_0 of the variable Y_0 , it is necessary to estimate the variance of the differences between the ordinates $\hat{y}_0$, obtained from the computed regression lines in repeated sampling when $x = x_0$, and the corresponding true ordinate y_0 . We can think of the difference $\hat{y}_0 - y_0$ as a value of the random variable $\hat{Y}_0 - Y_0$, whose sampling distribution can be shown to be normal with mean

$$\mu_{\hat{Y}_0 - Y_0} = E(\hat{Y}_0 - Y_0) = E[B_0 + B_1 x_0 - (\beta_0 + \beta_1 x_0 + \epsilon_0)] = 0$$

and variance

$$\sigma_{\hat{Y}_0 - Y_0}^2 = \sigma_{B_0 + B_1 x_0 - \epsilon_0} = \sigma_{\hat{Y} + B_1(x_0 - \bar{x}) - \epsilon_0} = \sigma^2 \left[1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{S_{xx}} \right]$$

Thus, a $100(1 - \alpha)\%$ prediction interval for a single predicted value y_0 can be constructed from the statistic

$$T = \frac{\hat{Y}_0 - Y_0}{S \sqrt{1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{S_{xx}}}}$$

which has a t -distribution with $n - 2$ degrees of freedom.

Definition 4.79: Prediction Interval for y_0

A $100(1 - \alpha)\%$ prediction interval for a single response y_0 is given by

$$\hat{y}_0 - t_{\frac{\alpha}{2}} s \sqrt{1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{S_{xx}}} < y_0 < \hat{y}_0 + t_{\frac{\alpha}{2}} s \sqrt{1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{S_{xx}}}$$

where $t_{\frac{\alpha}{2}}$ is a value of the t -distribution with $n - 2$ degrees of freedom that leaves an area of $\frac{\alpha}{2}$ to the right..

iii. Simple Regression Line Plot and Pearson's Correlation Computation

Suppose we have a bivariate data as follow Compute the Pearson's correlation (PMCC), then plot

Distance, x (km)	4	8	5	10	6
Time, y (minutes)	15	35	12	40	24

Table 4.16: Distance travelled to school and travel time

the data on a scatter diagram and draw the line of best fit.

Solution:

We will have

$$\begin{aligned}\bar{x} &= \frac{\sum x}{n} = 6.6 \\ \bar{y} &= \frac{\sum y}{n} = 25.2 \\ S_x &= \sqrt{\frac{\sum x^2}{n} - (\bar{x})^2} = 2.154 \\ S_y &= \sqrt{\frac{\sum y^2}{n} - (\bar{y})^2} = 10.907 \\ S_{xy} &= \frac{\sum xy}{n} - (\bar{x})(\bar{y}) = 22.48\end{aligned}$$

Hence

$$r = \frac{S_{xy}}{S_x S_y} = 0.957$$

The value of r is close to $+1$, showing that there is strong positive correlation between the distance travelled to school and the time taken.

Now we will compute the regression line formula

$$\begin{aligned}y - \bar{y} &= \frac{S_{xy}}{(S_x)^2}(x - \bar{x}) \\ y - 25.2 &= \frac{22.48}{(2.154)^2}(x - 6.6) \\ y &= 4.85x - 6.78\end{aligned}$$

To plot the bivariate data on a scatter diagram and the regression line we can use Hamzstlab Mathematics, and the backend computation can be done with SymIntegration.

In SymIntegration the function to compute Pearson's correlation is:

rpearson(M,n)

with M as the matrix of size $n \times 2$, the first column of the matrix represents the independent variable, x , while the second column of the matrix represents the dependent variable, y .

The function to compute the regression line is:
regressionline(const SymbolicMatrix &M, int n)

with M as the matrix of size $n \times 2$, the result will be a symbolic function, $y(x)$.

```
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
./main ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Load Vector from textfile and Compute Regression Line ]#
W:
[ 4 15]
[ 8 35]
[ 5 12]
[10 40]
[ 6 24]

x bar: 6.6
y bar: 25.2
sum x: 33
sum y: 126
sum xy: 944
sum x^2: 241
sum y^2: 3770

Pearson's correlation coefficient, r :
0.956835

Regression line, y = 4.84483*x-6.77586

Time taken by function: 7136 microseconds
```

Figure 4.52: The computation of Pearson's correlation (PMCC), r , and the regression line equation with *SymIntegration* took 7136 microseconds. (*SymIntegration/Examples/Statistics/Test SymIntegration Load Vector from textfile and Compute Regression Line/main.cpp*).

iv. Compute Linear Regression with Confidence Interval and Prediction Interval with SymIntegration

One of the more challenging problems confronting the water pollution control field is presented by the tanning industry. Tannery wastes are chemically complex. They are characterized by high values of chemical oxygen demand, volatile solids, and other pollution measures. Consider the experimental data, which were obtained from 33 samples of chemically treated waste in a study conducted at Virginia Tech. Readings on x , the percent reduction in total solids, and y , the percent reduction in chemical oxygen demand, were recorded.

Solids Reduction, $x(\%)$	Oxygen Demand Reduction, $y(\%)$	Solids Reduction, $x(\%)$	Oxygen Demand Reduction, $y(\%)$
3	5	36	34
7	11	37	36
11	21	38	38
15	16	39	37
18	16	39	36
27	28	39	45
29	27	40	39
30	25	41	41
30	35	42	40
31	30	42	44
31	40	43	37
32	32	44	44
33	34	45	46
33	32	46	46
34	34	47	49
36	37	50	51
36	38		

Table 4.17: Measures of reduction in solids and oxygen demand.

Estimate the regression line for the pollution data of Table 4.10. Find a 95% confidence interval for β_0 and β_1 in the regression line $\mu_{Y|x} = \beta_0 + \beta_1 x$ based on the pollution data, construct 95% confidence limits for the mean response $\mu_{Y|x_i}$ and the 95% prediction interval for y_i for all x_i with $i = 1, 2, \dots, 33$.

Solution:

$$\begin{aligned}\sum_{i=1}^{n=33} x_i &= 1104 \\ \sum_{i=1}^{n=33} y_i &= 1124 \\ \sum_{i=1}^{n=33} x_i y_i &= 41,355 \\ \sum_{i=1}^{n=33} x_i^2 &= 41,086\end{aligned}$$

Therefore,

$$\begin{aligned}b_1 &= \frac{(33)(41,355) - (1104)(1124)}{(33)(41,086) - (1104)^2} = 0.903643 \\ b_0 &= \frac{1124 - (0.903643)(1104)}{33} = 3.829633\end{aligned}$$

Thus, the estimated regression line is given by

$$\hat{y} = 3.8296 + 0.9036x$$

Now, to compute the confidence interval we will need to compute the "S" sum formulas

$$\begin{aligned}S_{xx} &= 4152.18 \\ S_{yy} &= 3713.88 \\ S_{xy} &= 3752.09\end{aligned}$$

The 95% confidence interval for the parameter β_1 is

$$b_1 - t_{0.025} \frac{s}{\sqrt{S_{xx}}} < \beta_1 < b_1 + t_{0.025} \frac{s}{\sqrt{S_{xx}}}$$

Therefore,

$$0.801425 < \beta_1 < 1.00586$$

The 95% confidence interval for the parameter β_0 is

$$b_0 - t_{0.025} \frac{s}{\sqrt{nS_{xx}}} \sqrt{\sum_{i=1}^n x_i^2} < \beta_0 < b_0 + t_{0.025} \frac{s}{\sqrt{nS_{xx}}} \sqrt{\sum_{i=1}^n x_i^2}$$

Therefore,

$$0.222861 < \beta_0 < 7.43641$$

In SymIntegration, we have another function that can compute the estimated regression line, the $100(1 - \alpha)\%$ confidence interval for β_0 and β_1 in the regression line $\mu_{Y|x} = \beta_0 + \beta_1 x$, that will also compute the one-way ANOVA table for the regression line, and print the table of the original sample data and the fitted / regressed value and the $100(1 - \alpha)\%$ confidence limits for the mean response $\mu_{Y|x_i}$ and the $100(1 - \alpha)\%$ prediction interval for y_i for all x_i with $i = 1, 2, \dots, n$.

This function is:
regressionline(vector<vector<double>> matrix, double α)

With **matrix** as a $n \times 2$ matrix that can be saved as textfile (.txt), and α is the level of significance. While the other function at the previous example: **regressionline(const SymbolicMatrix &M, int n)** resulting in Symbolic output, this function is a **void** function that shows the output only on the computer screen.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Linear Regression with Confidence Interval and Prediction Interval
*****
Simple Linear Regression
Sxx = 4152.18 , Syy = 3713.88 , Sxy = 3752.09
SSE = 323.327, SSR = 3390.55, SST = 3713.88
R-Squared = 91.2941 %
Computed f = 325.08, P-value = 0

An unbiased estimate of  $\sigma^2$  is :
s2 = 10.4299
The estimated regression line is given by:
y = 3.82963 + 0.903643x
*****
Analysis of Variance

```

Source	DF	SS	MS	F	P
Regression	1	3390.55	3390.55	325.08	0
Residual Error	31	323.327	10.4299		
Total	32	3713.88			

```

R-Squared          S
91.2941 %          3.22954

If P value is less than the level of significance 0.05 then reject H0:  $\beta_1 = 0$ 

Rejection of H0 may suggest that the relationship is, indeed, linear.
The failure to reject H0:  $\beta_1 = 0$  suggests that there is no linear relationship between Y and x.
*****
A 95% confidence interval for the parameter  $\beta_1$  is :
0.801425 <  $\beta_1$  < 1.00586
A 95% confidence interval for the parameter  $\beta_0$  is :
0.222861 <  $\beta_0$  < 7.43641
```

Figure 4.53: The computation of the estimated regression line, the $100(1 - \alpha)\%$ confidence interval for β_0 and β_1 in the regression line $\mu_{Y|x} = \beta_0 + \beta_1 x$, the one-way ANOVA table. (*SymIntegration/Examples/Statistics/Test SymIntegration Linear Regression with Confidence Interval and Prediction Interval/main.cpp*).

Obs	y_data	Fit	SE Fit	95% CI	95% PI
1	5	6.54056	1.62659	(3.22311 , 9.85802)	(-0.834391 , 13.9155)
2	11	10.1551	1.44014	(7.21795 , 13.0923)	(2.94324 , 17.367)
3	21	13.7697	1.25801	(11.204 , 16.3354)	(6.70095 , 20.8385)
4	16	17.3843	1.08238	(15.1768 , 19.5918)	(10.4375 , 24.331)
5	16	20.0952	0.957084	(18.1432 , 22.0472)	(13.2254 , 26.965)
6	28	28.228	0.648619	(26.9051 , 29.5509)	(21.5098 , 34.9462)
7	27	30.0353	0.604898	(28.8016 , 31.269)	(23.3341 , 36.7365)
8	25	30.9389	0.588247	(29.7392 , 32.1387)	(24.2439 , 37.634)
9	35	30.9389	0.588247	(29.7392 , 32.1387)	(24.2439 , 37.634)
10	30	31.8426	0.575493	(30.6688 , 33.0163)	(25.1521 , 38.533)
11	40	31.8426	0.575493	(30.6688 , 33.0163)	(25.1521 , 38.533)
12	32	32.7462	0.566897	(31.59 , 33.9024)	(26.0588 , 39.4336)
13	34	33.6499	0.562652	(32.5023 , 34.7974)	(26.964 , 40.3358)
14	32	33.6499	0.562652	(32.5023 , 34.7974)	(26.964 , 40.3358)
15	34	34.5535	0.562855	(33.4056 , 35.7015)	(27.8675 , 41.2395)
16	37	36.3608	0.576484	(35.185 , 37.5365)	(29.67 , 43.0516)
17	38	36.3608	0.576484	(35.185 , 37.5365)	(29.67 , 43.0516)
18	34	36.3608	0.576484	(35.185 , 37.5365)	(29.67 , 43.0516)
19	36	37.2644	0.589604	(36.0619 , 38.4669)	(30.5689 , 43.96)
20	38	38.1681	0.606595	(36.9309 , 39.4052)	(31.4662 , 44.8699)
21	37	39.0717	0.62714	(37.7927 , 40.3508)	(32.362 , 45.7814)
22	36	39.0717	0.62714	(37.7927 , 40.3508)	(32.362 , 45.7814)
23	45	39.0717	0.62714	(37.7927 , 40.3508)	(32.362 , 45.7814)
24	39	39.9754	0.659904	(38.6478 , 41.3029)	(33.2562 , 46.6945)
25	41	40.879	0.677548	(39.4971 , 42.2609)	(34.1489 , 47.6091)
26	40	41.7826	0.706746	(40.3412 , 43.2241)	(35.0401 , 48.5252)
27	44	41.7826	0.706746	(40.3412 , 43.2241)	(35.0401 , 48.5252)
28	37	42.6863	0.738195	(41.1807 , 44.1919)	(35.9297 , 49.4429)
29	44	43.5899	0.771621	(42.0162 , 45.1637)	(36.8179 , 50.362)
30	46	44.4936	0.806777	(42.8481 , 46.139)	(37.7045 , 51.2827)
31	46	45.3972	0.843448	(43.677 , 47.1174)	(38.5896 , 52.2048)
32	49	46.3009	0.881443	(44.5031 , 48.0986)	(39.4733 , 53.1285)
33	51	49.0118	1.00185	(46.9685 , 51.0551)	(42.1155 , 55.9081)

Time taken by function: 3486 microseconds

Figure 4.54: The table that shows the fitted $\hat{y}_i$, SE fit, the confidence interval for the mean response $\mu_{Y|x_i}$ and prediction interval for y_i . (SymIntegration/Examples/Statistics/Test SymIntegration Linear Regression with Confidence Interval and Prediction Interval/main.cpp).

The term "SE Fit" is used in computing confidence intervals on mean response, the formula is:

$$\text{SE Fit}_i = t_{\frac{\alpha}{2}} s \sqrt{\frac{1}{n} + \frac{(x_i - \bar{x})^2}{S_{xx}}}$$

v. Multiple Linear Regression

[SI*] In most research problems where regression analysis is applied, more than one independent variable is needed in the regression model. The complexity of most scientific mechanisms is such that in order to be able to predict an important response, a multiple regression model is needed. When this model is linear in the coefficients, it is called a multiple linear regression model. For the case of k independent variables $x_1, x_2, \dots, x_k$, the mean $Y|x_1, x_2, \dots, x_k$ is given by the multiple linear regression model

$$\mu_{Y|x_1, x_2, \dots, x_k} = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k$$

and the estimated response is obtained from the sample regression equation

$$\hat{y} = b_0 + b_1 x_1 + \dots + b_k x_k$$

where each regression coefficient β_i is estimated by b_i from the sample data using the method of least squares.

Definition 4.80: Multiple Linear Regression Model

We obtain the least squares estimators of the parameters $\beta_0, \beta_1, \dots, \beta_k$ by fitting the multiple linear regression model

$$\mu_{Y|x_1, x_2, \dots, x_k} = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k$$

to the data points

$$\{(x_{1i}, x_{2i}, \dots, x_{ki}, y_i); \quad i = 1, 2, \dots, n \text{ and } n > k\}$$

where y_i is the observed response to the values $x_{1i}, x_{2i}, \dots, x_{ki}$ of the k independent variables $x_1, x_2, \dots, x_k$. Each observation $(x_{1i}, x_{2i}, \dots, x_{ki}, y_i)$ is assumed to satisfy the following equation.

$$y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_k x_{ki} + \epsilon_i$$

or

$$\begin{aligned} y_i &= \hat{y}_i + e_i \\ &= b_0 + b_1 x_{1i} + b_2 x_{2i} + \dots + b_k x_{ki} + e_i \end{aligned}$$

where ϵ_i and e_i are the random error and residual, respectively, associated with the response y_i and fitted value $\hat{y}_i$.

[SI*] As in the case of simple linear regression, it is assumed that the ϵ_i are independent and identically distributed with mean 0 and common variance σ^2 .

In using the concept of least squares to arrive at estimates $b_0, b_1, \dots, b_k$, we minimize the expression

$$\begin{aligned} SSE &= \sum_{i=1}^n e_i^2 \\ &= \sum_{i=1}^n (y_i - b_0 - b_1 x_{1i} - b_2 x_{2i} - \dots - b_k x_{ki})^2 \end{aligned}$$

Differentiating SSE in turn with respect to $b_0, b_1, \dots, b_k$ and equating to zero, we generate the set of $k + 1$ normal equations for multiple linear regression.

Multiple Linear Regression Parameter Estimation with Matrices

In fitting a multiple linear regression model, particularly when the number of variables exceeds two, a knowledge of matrix theory can facilitate the mathematical manipulations considerably.

Suppose that the experimenter has k independent variables $x_1, x_2, \dots, x_k$ and n observations $y_1, y_2, \dots, y_n$, each of which can be expressed by the equation

$$y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_k x_{ki} + \epsilon_i$$

This model essentially represents n equations describing how the response values are generated in the scientific process.

Definition 4.81: General Linear Model

Using matrix notation, we can write the following equation that represents the multiple linear regression equation:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

where

$$\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}, \quad \mathbf{X} = \begin{bmatrix} 1 & x_{11} & x_{21} & \dots & x_{k1} \\ 1 & x_{12} & x_{22} & \dots & x_{k2} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 1 & x_{1n} & x_{2n} & \dots & x_{kn} \end{bmatrix}$$

$$\boldsymbol{\beta} = \begin{bmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_k \end{bmatrix}, \quad \boldsymbol{\epsilon} = \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{bmatrix}$$

Then the least squares method for estimation of $\boldsymbol{\beta}$, involves finding $\mathbf{b}$ for which

$$SSE = (\mathbf{y} - \mathbf{X}\mathbf{b})^T (\mathbf{y} - \mathbf{X}\mathbf{b})$$

is minimized. This minimization process involves solving for $\mathbf{b}$ in the equation

$$\frac{\partial}{\partial \mathbf{b}} (SSE) = \mathbf{0}$$

The result reduces to the solution of $\mathbf{b}$ in

$$(\mathbf{X}^T \mathbf{X})\mathbf{b} = \mathbf{X}^T \mathbf{y}$$

Notice the nature of the $\mathbf{X}$ matrix. Apart from the initial element, the i th row represents the x -values that give rise to the response y_i . Writing

$$\mathbf{X}^T \mathbf{X} = \begin{bmatrix} n & \sum_{i=1}^n x_{1i} & \sum_{i=1}^n x_{2i} & \dots & \sum_{i=1}^n x_{ki} \\ \sum_{i=1}^n x_{1i} & \sum_{i=1}^n x_{1i}^2 & \sum_{i=1}^n x_{1i}x_{2i} & \dots & \sum_{i=1}^n x_{1i}x_{ki} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^n x_{ki} & \sum_{i=1}^n x_{ki}x_{1i} & \sum_{i=1}^n x_{ki}x_{2i} & \dots & \sum_{i=1}^n x_{ki}^2 \end{bmatrix}$$

and

$$\mathbf{X}^T \mathbf{y} = \begin{bmatrix} \sum_{i=1}^n y_i \\ \sum_{i=1}^n x_{1i} y_i \\ \vdots \\ \sum_{i=1}^n x_{ki} y_i \end{bmatrix}$$

If the matrix $\mathbf{X}^T \mathbf{X}$ is not singular, the formula for parameter estimation in multiple linear regression is

$$\mathbf{b} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$$

where $\mathbf{b}$ is the vector of estimated coefficients, $\mathbf{X}$ is the matrix of independent variables (with a column of ones for the intercept), $\mathbf{X}^T$ is the transpose of $\mathbf{X}$, and $\mathbf{y}$ is the vector of the dependent variable. This formula uses the method of least squares to find the coefficients that minimize the sum of squared errors, which is the vertical distance between the observed and predicted values.

Model:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

with:

$\mathbf{y}$: Vector of the dependent variable

$\mathbf{X}$: Matrix of independent variables (including a column of 1 for the intercept)

$\boldsymbol{\beta}$: Vector of population parameters to be estimated

$\boldsymbol{\epsilon}$: Vector of error terms

Estimation:

$$\mathbf{b} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$$

with:

$\mathbf{b}$: Vector of estimated coefficients (intercept and slopes)

$\mathbf{X}^T$: Transpose of the matrix $\mathbf{X}$.

$(\mathbf{X}^T \mathbf{X})^{-1}$: The inverse of the matrix product $\mathbf{X}^T \mathbf{X}$.

The formula finds the set of coefficients ($\mathbf{b}$) that produce a regression line (or plane/hyper-plane) that best fits the data. "Best fit" is defined by the least-squares criterion, meaning the sum of the squared differences between the actual y values and the predicted $\hat{y}$ values is minimized. The matrix algebra approach provides a direct way to calculate the coefficients without having to manually solve a system of linear equations for each parameter, which would be necessary for a large number of independent variables.

To evaluate the model we can use R^2 (R-squared) to measure the proportion of the variance in the dependent variable that is predictable from the independent variables. A modified version of R^2 , adjusted R-squared is a better measure when comparing models with different numbers of predictors. Another measure is the p -value, it is often used to determine the statistical significance of the independent variables, indicating whether their relationship with the dependent variable is likely due to chance.

[SI*] **Properties of the Least Squares Estimators**

The means and variances of the estimators $b_0, b_1, \dots, b_k$ are readily obtained under certain assumptions on the random errors $\epsilon_1, \epsilon_2, \dots, \epsilon_k$ that are identical to those made in the case of simple linear regression. When we assume these errors to be independent, each with mean 0

and variance σ^2 , it can be shown that $b_0, b_1, \dots, b_k$ are, respectively, unbiased estimators of the regression coefficients $\beta_0, \beta_1, \dots, \beta_k$.

In addition, the variances of the b 's are obtained through the elements of the inverse of the A matrix. Note that the off-diagonal elements of $A = X^T X$ represents sums of products of elements in the columns of X , while the diagonal elements of A represents sums of squares of elements in the columns of X .

The inverse matrix, A^{-1} , apart from the multiplier σ^2 , represents the variance-covariance matrix of the estimated regression coefficients.

That is, the elements of the matrix $A^{-1}\sigma^2$ display the variances of $b_0, b_1, \dots, b_k$ on the main diagonal and covariances on the off-diagonal. For example, in a $k = 2$ multiple linear regression problem, we might write

$$(X^T X)^{-1} = \begin{bmatrix} c_{00} & c_{01} & c_{02} \\ c_{10} & c_{11} & c_{12} \\ c_{20} & c_{21} & c_{22} \end{bmatrix}$$

with the elements below the main diagonal determined through the symmetry of the matrix. Then we can write

$$\begin{aligned} \sigma_{b_i}^2 &= c_{ii}\sigma^2, \quad i = 0, 1, 2 \\ \sigma_{b_i b_j} &= \text{Cov}(b_i, b_j) = c_{ij}\sigma^2, \quad i \neq j \end{aligned}$$

Theorem 4.29: Unbiased Estimate for Multiple Linear Regression

For the linear regression equation

$$y = X\beta + \epsilon$$

an unbiased estimate of σ^2 is given by the error or residual mean square

$$s^2 = \frac{SSE}{n - k - 1}$$

where

$$SSE = \sum_{i=1}^n e_i^2 = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

The error and regression sums of squares take on the same form and play the same role as in the simple linear regression case. In fact, the sum-of-squares identity

$$\sum_{i=1}^n (y_i - \bar{y})^2 = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2 + \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

continues to hold, and we retain our previous notation, namely

$$SST = SSR + SSE$$

with

$$SST = \sum_{i=1}^n (y_i - \bar{y})^2$$

and

$$SSR = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$$

There are k degrees of freedom associated with SSR , and, as always, SST has $n - 1$ degrees of freedom. Therefore, after subtraction, SSE has $n - k - 1$ degrees of freedom.

[SI*] **Analysis of Variance in Multiple Regression**

The partition of the total sum of squares into its components, the regression and error sums of squares, plays an important role. An analysis of variance can be conducted to shed light on the quality of the regression equation. A useful hypothesis that determines if a significant amount of variation is explained by the model is

$$H_0 : \beta_1 = \beta_2 = \beta_3 = \cdots = \beta_k = 0$$

The analysis of variance involves an F -test via a table. The test is an upper-tailed test.

Source	Sum of Squares	Degrees of Freedom	Mean Squares	F
Regression	SSR	k	$MSR = \frac{SSR}{k}$	$f = \frac{MSR}{MSE}$
Error	SSE	$n - (k + 1)$	$MSE = \frac{SSE}{n - (k + 1)}$	
Total	SST	$n - 1$		

Table 4.18: The analysis of variance table that uses F -test.

Rejection of H_0 implies that the regression equation differs from a constant. That is, at least one regressor variable is important.

Further utility of the mean square error (or residual mean square) lies in its use in hypothesis testing and confidence interval estimation. In addition, the mean square error plays an important role in situations where the scientist is searching for the best from a set of competing models. Many model-building criteria involve the statistic s^2 .

[SI*] **Inferences in Multiple Linear regression**

A knowledge of the distributions of the individual coefficient estimators enables the experimenter to construct confidence intervals for the coefficients and to test hypotheses about them.

Recall that the $b_j (j = 0, 1, 2, \dots, k)$ are normally distributed with mean β_j and variance $c_{jj}\sigma^2$. Thus, we can use the statistic

$$t = \frac{b_j - \beta_{j0}}{s\sqrt{c_{jj}}}$$

with $n - k - 1$ degrees of freedom to test hypotheses and construct confidence intervals on β_j . For example, if we wish to test

$$H_0 : \beta_j = \beta_{j0}$$

$$H_1 : \beta_j \neq \beta_{j0}$$

we compute the above t -statistic and do not reject H_0 if $-t_{\alpha/2} < t < t_{\alpha/2}$, where $t_{\alpha/2}$ has $n - k - 1$ degrees of freedom.

[SI*] Individual t -Tests for Variable Screening

The t -test most often used in multiple regression is the one that tests the importance of individual coefficients (i.e., $H_0 : \beta_j = 0$ against the alternative $H_1 : \beta_j \neq 0$). These tests often contribute to what is termed variable screening, where the analyst attempts to arrive at the most useful model (i.e., the choice of which regressors to use). It should be emphasized here that if a coefficient is found insignificant (i.e., the hypothesis $H_0 : \beta_j = 0$ is not rejected), the conclusion drawn is that the variable is insignificant (i.e., explains an insignificant amount of variation in y), in the presence of the other regressors in the model.

[SI*] Inferences on Mean Response and Prediction

One of the most useful inferences that can be made regarding the quality of the predicted response y_0 corresponding to the values $x_{10}, x_{20}, \dots, x_{k0}$ is the confidence interval on the mean response $\mu_{Y|x_{10}, x_{20}, \dots, x_{k0}}$. We are interested in constructing a confidence interval on the mean response for the set of conditions given by

$$\mathbf{x}_0^T = [1, x_{10}, x_{20}, \dots, x_{k0}]$$

We augment the conditions on the x 's by the number 1 in order to facilitate the matrix notation. Normality in the ϵ_i produces normality in the b_j and the mean and variance are still the same. So is the covariance between b_i and b_j , for $i \neq j$. Hence,

$$\hat{y} = b_0 + \sum_{j=1}^k b_j x_{j0}$$

is likewise normally distributed and is, in fact, an unbiased estimator for the mean response on which we are attempting to attach a confidence interval. The variance of $\hat{y}_0$, written in matrix notation simply as a function of σ^2 , $(\mathbf{X}^T \mathbf{X})^{-1}$, and the condition vector $\mathbf{x}_0^T$ is

$$\sigma_{\hat{y}_0}^2 = \sigma^2 \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0$$

If this expression is expanded for a given case, say $k = 2$, it is readily seen that it appropriately accounts for the variance of the b_j and the covariance of b_i and b_j , for $i \neq j$. After σ^2 is replaced by s^2 , the $100(1 - \alpha)\%$ confidence interval on $\mu_{Y|x_{10}, x_{20}, \dots, x_{k0}}$ can be constructed from the statistic

$$T = \frac{\hat{y}_0 - \mu_{Y|x_{10}, x_{20}, \dots, x_{k0}}}{s \sqrt{\mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0}}$$

which has a t -distribution with $n - k - 1$ degrees of freedom.

Definition 4.82: Confidence Interval for $\mu_{Y|x_{10}, x_{20}, \dots, x_{k0}}$

A $100(1 - \alpha)\%$ confidence interval for the mean response $\mu_{Y|x_{10}, x_{20}, \dots, x_{k0}}$ is

$$\hat{y}_0 - t_{\frac{\alpha}{2}} s \sqrt{\mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0} < \mu_{Y|x_{10}, x_{20}, \dots, x_{k0}} < \hat{y}_0 + t_{\frac{\alpha}{2}} s \sqrt{\mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0}$$

where $t_{\frac{\alpha}{2}}$ is a value of the t -distribution with $n - k - 1$ degrees of freedom.

The quantity $s \sqrt{\mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0}$ is often called the standard error of prediction.

[SI*] As in the case of simple linear regression, we need to make a clear distinction between the confidence interval on a mean response and the prediction interval on an observed response.

The latter provides a bound within which we can say with a preselected degree of certainty that a new observed response will fall.

A prediction interval for a single predicted response y_0 is once again established by considering the difference $\hat{y}_0 - y_0$. The sampling distribution can be shown to be normal with mean

$$\mu_{\hat{y}_0 - y_0} = 0$$

and variance

$$\sigma_{\hat{y}_0 - y_0}^2 = \sigma^2 [1 + \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0]$$

Thus, a $100(1 - \alpha)\%$ prediction interval for a single prediction value y_0 can be constructed from the statistic

$$T = \frac{\hat{y}_0 - y_0}{s \sqrt{1 + \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0}}$$

which has t -distribution with $n - k - 1$ degrees of freedom.

Definition 4.83: Prediction Interval for y_0

A $100(1 - \alpha)\%$ prediction interval for a single response y_0 is given by

$$\hat{y}_0 - t_{\frac{\alpha}{2}} s \sqrt{1 + \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0} < y_0 < \hat{y}_0 + t_{\frac{\alpha}{2}} s \sqrt{1 + \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0}$$

where $t_{\frac{\alpha}{2}}$ is a value of the t -distribution with $n - k - 1$ degrees of freedom.

[SI*] Choice of a Fitted Model through Hypothesis Testing

In many regression situations, individual coefficients are of importance to the experimenter. For example, in an economics application, $\beta_1, \beta_2, \dots$ might have some particular significance, and then confidence intervals and tests of hypotheses on these parameters would be of interest to the economist.

The experimenter using regression analysis is also interested in deletion of variables when the situation dictates that, in addition to arriving at a workable prediction equation, he or she must find the "best regression" involving only variables that are useful predictors.

One criterion that is commonly used to illustrate the adequacy of a fitted regression model is the coefficient of determination, R^2 .

Definition 4.84: R^2

The formula is

$$R^2 = \frac{SSR}{SST} = 1 - \frac{SSE}{SST}$$

The regression sum of squares can be used to give some indication concerning whether or not the model is an adequate explanation of the true situation. We can test the hypothesis H_0 that the regression is not significant by merely forming the ratio

$$f = \frac{SSR/k}{SSE/(n - k - 1)} = \frac{SSR/k}{s^2} = \frac{MSR}{MSE}$$

and rejecting H_0 at the α -level of significance when $f > f_\alpha(k, n - k - 1)$.

[SI*] **The Adjusted Coefficient of Determination R^2_{adj}**

Adjusted R^2 is a variation on R^2 that provides an adjustment for degrees of freedom. The coefficient of determination cannot decrease as terms are added to the model. In other words, R^2 does not decrease as the error degrees of freedom $n - k - 1$ are reduced, the latter result being produced by an increase in k , the number of model terms.

Definition 4.85: Adjusted R^2

The formula is

$$R^2_{\text{adj}} = 1 - \frac{SSE/(n - k - 1)}{SST/(n - 1)}$$

vi. Compute Multiple Linear Regression with SymIntegration

A study was done on a diesel-powered light-duty pickup truck to see if humidity, air temperature, and barometric pressure influence emission of nitrous oxide (in ppm). Emission measurements were taken at different times, with varying experimental conditions. The data are given in table below. The model is

$$\mu_{Y|x_1, x_2, x_3} = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3$$

or, equivalently,

$$y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \beta_3 x_{3i} + \epsilon_i, \quad i = 1, 2, \dots, 20$$

Fit this multiple linear regression model to the given data and then estimate the amount of nitrous oxide emitted for the conditions where humidity is 50%, temperature is 76⁰ F, and barometric pressure is 29.30.

Nitrous Oxide, y	Humidity, x_1	Temp., x_2	Pressure, x_3	Nitrous Oxide, y	Humidity x_1	Temp., x_2	Pressure, x_3
0.90	72.4	76.3	29.18	1.07	23.2	76.8	29.38
0.91	41.6	70.3	29.35	0.94	47.4	86.6	29.35
0.96	34.3	77.1	29.24	1.10	31.5	76.9	29.63
0.89	35.1	68.0	29.27	1.10	10.6	86.3	29.56
1.00	10.7	79.0	29.78	1.10	11.2	86.0	29.48
1.10	12.9	67.4	29.39	0.91	73.3	76.3	29.40
1.15	8.3	66.8	29.69	0.87	75.4	77.9	29.28
1.03	20.1	76.9	29.48	0.78	96.6	78.7	29.29
0.77	72.2	77.7	29.09	0.82	107.4	86.8	29.03
1.07	24.0	67.7	29.60	0.95	54.9	70.9	29.37

Table 4.19: Source: Charles T. Hare, "Light-Duty Diesel Emission Correction Factors for Ambient Conditions," EPA-600/2-77-116. U.S. Environmental Protection Agency.

Solution:

The solution of the set of estimating equations yields the unique estimates

$$b_0 = -3.507778$$

$$b_1 = -0.002625$$

$$b_2 = 0.000799$$

$$b_3 = 0.154155$$

Therefore, the regression equation is

$$\hat{y} = -3.507778 - 0.002625x_1 + 0.000799x_2 + 0.154155x_3$$

For 50% humidity, a temperature of 76⁰ F, and a barometric pressure of 29.30, the estimated amount of nitrous oxide emitted is

$$\begin{aligned} \hat{y} &= -3.507778 - 0.002625(50) + 0.000799(76) + 0.154155(29.30) \\ &= 0.9384 \end{aligned}$$

In SymIntegration, to compute the multiple regression model' coefficients we can use this function:

multipleregression(vector<vector<double>> &X, vector<vector<double>> &y)

with X as the matrix of independent variables, including a column if 1 as the first column for the intercept, and y is the vector of the dependent variable. The computation requires inverse operation, thus it is a bit expensive on the computational side since computing inverse of a matrix requires some works before, such as computing the determinant of the corresponding matrix and the adjugate matrix as well.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Multiple Linear
Regression ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Multiple Linear Regression ]# ./main

Matrix X :
    1.000000    72.400000    76.300000    29.180000
    1.000000    41.600000    70.300000    29.350000
    1.000000    34.300000    77.100000    29.240000
    1.000000    35.100000    68.000000    29.270000
    1.000000    10.700000    79.000000    29.780000
    1.000000    12.900000    67.400000    29.390000
    1.000000     8.300000    66.800000    29.690000
    1.000000    20.100000    76.900000    29.480000
    1.000000    72.200000    77.700000    29.090000
    1.000000    24.000000    67.700000    29.600000
    1.000000    23.200000    76.800000    29.380000
    1.000000    47.400000    86.600000    29.350000
    1.000000    31.500000    76.900000    29.630000
    1.000000    10.600000    86.300000    29.560000
    1.000000    11.200000    86.000000    29.480000
    1.000000    73.300000    76.300000    29.400000
    1.000000    75.400000    77.900000    29.280000
    1.000000    96.600000    78.700000    29.290000
    1.000000   107.400000    86.800000    29.030000
    1.000000    54.900000    70.900000    29.370000

Vector y :
    0.900000
    0.910000
    0.960000
    0.890000
    1.000000
    1.100000
    1.150000
    1.030000
    0.770000
    1.070000
    1.070000
    0.940000
    1.100000
    1.100000
    0.910000
    0.870000
    0.780000
    0.820000
    0.950000
```

Figure 4.55: The computation of the multiple regression model with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Multiple Linear Regression/main.cpp).

```

Multiple regression with matrix algebra:

b[0] = -3.507778
b[1] = -0.002625
b[2] = 0.000799
b[3] = 0.154155

(X^{T} X)^{-1} X^{T} y:
      -3.507778
      -0.002625
       0.000799
       0.154155

y          y estimated
0.900000   0.861376
0.910000   0.963638
0.960000   0.971276
0.890000   0.966530
1.000000   1.117988
1.100000   1.042824
1.150000   1.100667
1.030000   1.045388
0.770000   0.849145
1.070000   1.046299
1.070000   1.021756
0.940000   0.961436
1.100000   1.038587
1.100000   1.090168
1.100000   1.076021
0.910000   0.892927
0.870000   0.870194
0.780000   0.816725
0.820000   0.754766
0.950000   0.932288

For 50% humidity, 76^{0} F and 29.30 barometric pressure, the estimated multiple regression is:
0.938434

Time taken by function: 6299 microseconds

```

Figure 4.56: *The computation of the multiple regression model with SymIntegration (SymIntegration/Examples/Statistics/Test SymIntegration Multiple Linear Regression/main.cpp).*

vii. Compute Inferences in Multiple Linear Regression with SymIntegration

The percent survival rate of sperm in a certain type of animal semen, after storage, was measured at various combinations of concentrations of three materials used to increase chance of survival. The data are given in table below. Test the hypothesis that $\beta_2 = -2.5$ at the 0.05 level of significance against the alternative that $\beta_2 > -2.5$. Then construct a 95% confidence interval for the mean response and the prediction interval for y_0 .

y (% survival)	x_1 (weight %)	x_2 (weight %)	x_3 (weight %)
25.5	1.74	5.30	10.80
31.2	6.32	5.42	9.40
25.9	6.22	8.41	7.20
38.4	10.52	4.63	8.50
18.4	1.19	11.60	9.40
26.7	1.22	5.85	9.90
26.4	4.10	6.62	8.00
25.9	6.32	8.72	9.10
32.0	4.08	4.42	8.70
25.2	4.15	7.60	9.20
39.7	10.15	4.83	9.40
35.7	1.72	3.12	7.60
26.5	1.70	5.30	8.20

Table 4.20: Data for survival rate of sperm.

Solution:

$$H_0 : \beta_2 = -2.5$$

$$H_1 : \beta_2 > -2.5$$

Computations:

$$t = \frac{b_2 - \beta_{20}}{s\sqrt{c_{22}}} = \frac{-1.8618 + 2.5}{2.073\sqrt{0.0166}} = 2.390$$

$$P = P(T > 2.390) = 0.04$$

The signal that favors H_1 comes from large values of t . The P -value corresponding to $t = 2.390$ computed by the formula $P(T > 2.390)$ is coming from $H_1 : \beta_2 > -2.5$, since the sign is $>$ in H_1 thus we need to compute the P -value with $>$ sign.

Decision: Reject H_0 and conclude that $\beta_2 > -2.5$.

In SymIntegration to compute the estimation of the multiple linear regression model along with the ANOVA table and inferences in multiple linear regression we can use this function: **multipleregressionfull(vector<vector<double>> &X, vector<vector<double>> &y, double α)**

with X as the vector X as the matrix of independent variables as in the General Linear Model $y = X\beta + \epsilon$, y is the vector of the dependent variable, and α is the level of significance for computing inferences in multiple linear regression.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Multiple Linear Regression Animal Semen ]# ./main

Matrix X :
      1.000000      1.740000      5.300000     10.800000
      1.000000      6.320000      5.420000      9.400000
      1.000000      6.220000      8.410000      7.200000
      1.000000     10.520000      4.630000      8.500000
      1.000000      1.190000     11.600000      9.400000
      1.000000      1.220000      5.850000      9.900000
      1.000000      4.100000      6.620000      8.000000
      1.000000      6.320000      8.720000      9.100000
      1.000000      4.080000      4.420000      8.700000
      1.000000      4.150000      7.600000      9.200000
      1.000000     10.150000      4.830000      9.400000
      1.000000      1.720000      3.120000      7.600000
      1.000000      1.700000      5.300000      8.200000

Vector y :
      25.500000
      31.200000
      25.900000
      38.400000
      18.400000
      26.700000
      26.400000
      25.900000
      32.000000
      25.200000
      39.700000
      35.700000
      26.500000

Multiple regression with matrix algebra:

b[0] = 39.153886
b[1] = 1.016159
b[2] = -1.861393
b[3] = -0.343152

y = 39.153886 + 1.016159 x1 + -1.861393 x2 + -0.343152 x3

```

Figure 4.57: The computation of the multiple regression model with ANOVA and inferences on mean response and prediction (*SymIntegration/Examples/Statistics/Test SymIntegration Multiple Linear Regression Animal Semen/main.cpp*).

```

*****
Analysis of Variance

Source           DF          SS          MS          F          P-value
Regression       3          399.452877      133.150959      30.983038      0.000045
Residual Error   9          38.677892       4.297544
Total           12          438.130769

Rejection of H0 implies that the regression equation differs from a constant

R-Square         Grand Mean         Root MSE
0.911721         29.038462         2.073052

Variable         DF          Parameter Estimate      Standard Error      t value      P-value
Intercept        1          39.153886              5.887382            6.650475      0.000047
x1               1          1.016159              0.190911            5.322683      0.000240
x2               1          -1.861393             0.267338            -6.962706     0.000033
x3               1          -0.343152             0.617067            -0.556103     0.295848

Obs   y_data      Fit          SE Fit          95% CI          95% PI          Residual
1     25.500000    27.816643    3.201540    (24.615103 , 31.018182) (22.138446 , 33.494839) -2.316643
2     31.200000    32.404840    1.774690    (30.630150 , 34.179530) (27.390702 , 37.418978) -1.204840
3     25.900000    27.779020    3.073862    (24.705158 , 30.852882) (22.171820 , 33.386219) -1.879020
4     38.400000    38.691119    2.899519    (35.791600 , 41.590638) (33.177566 , 44.204672) -0.291119
5     18.400000    16.125418    3.571770    (12.553647 , 19.697188) (10.230539 , 22.020297)  2.274582
6     26.700000    26.356941    2.343384    (24.013558 , 28.700325) (21.114470 , 31.599412)  0.343059
7     26.400000    28.404945    1.831029    (26.573916 , 30.235973) (23.370590 , 33.439299) -2.004945
8     25.900000    26.442355    2.201496    (24.240859 , 28.643850) (21.261752 , 31.622957) -0.542355
9     32.000000    32.237628    1.770072    (30.467556 , 34.007699) (27.225123 , 37.250133) -0.237628
10    25.200000    26.178759    1.565245    (24.613514 , 27.744003) (21.234869 , 31.122649) -0.978759
11    39.700000    37.649205    2.956527    (34.692677 , 40.605732) (32.105460 , 43.192950)  2.050795
12    35.700000    32.985589    3.314084    (29.671504 , 36.299673) (27.243184 , 38.727993)  2.714411
13    26.500000    28.427540    2.226526    (26.201014 , 30.654067) (23.236252 , 33.618829) -1.927540

Time taken by function: 5086 microseconds

```

Figure 4.58: The computation of the multiple regression model with ANOVA and inferences on mean response and prediction (*SymIntegration/Examples/Statistics/Test SymIntegration Multiple Linear Regression Animal Semen/main.cpp*).

The f -test in the analysis of variance indicates that a significant amount of variability is explained. The R^2 value of 0.9117 implies that the model explains 91.17% of the variability in the response.

Note the parameter estimates, the standard errors, and the t -statistics shown in the output. The standard errors are computed from square roots of diagonal elements of $(X^T X)^{-1}s^2$, $s^2 = MSE = RMSE^2$.

In this illustration, the variable x_3 is insignificant in the presence of x_1 and x_2 based on the t -test and the corresponding P -value of 0.295848.

The formula to compute the t -value for the variable intercept, x_1, x_2, x_3 is based from the hypothesis test

$$H_0 : \beta_i = 0$$

$$H_1 : \beta_i > 0$$

we then use this formula:

$$t = \frac{b_i - \beta_{i0}}{s\sqrt{c_{ii}}}$$

with $\beta_{i0} = 0$, s is the RMSE (Root Mean Square Error), and c_{ii} is the diagonal entry of the matrix $(X^T X)^{-1}$.

The P -value itself can be computed with

$$P(T > |t|)$$

Take a look at the P -value at the analysis of variance (ANOVA) table, it means that we reject H_0 with

$$H_0 : \beta_1 = \beta_2 = \beta_3 = \cdots = \beta_k = 0$$

This should not be misinterpreted. Although it does indicate that the regression explained by the model is significant, this does not rule out the following possibilities:

1. The linear regression model for this set of x 's is not the only model that can be used to explain the data; indeed, there may be other models with transformations on the x 's that give a larger value of the F -statistic.
2. The model might have been more effective with the inclusion of other variables in addition to x_1, x_2 , and x_3 or perhaps with the deletion of one or more of the variables in the model, say x_3 , which has a $P = 0.2958$.

Regarding the pitfalls in the use of R^2 as a criterion for comparing competing models. These pitfalls are certainly relevant in multiple linear regression. In fact, in its employment in multiple regression, the dangers are even more pronounced since the temptation to overfit is so great. One should always keep in mind that $R^2 \approx 1.0$ can always be achieved at the expense of error degrees of freedom when an excess of model terms is employed. However, $R^2 = 1$, describing a model with a near perfect fit, does not always result in a model that predicts well.

Chapter 5

Operations Research Computation with SymIntegration

"Without mathematics, there's nothing you can do. Everything around you is mathematics. Everything around you is numbers." - Shakuntala Devi

Operations research is a branch of mathematics which is concerned with the application of scientific methods and techniques to decision making problems and with establishing the best or optimal solutions. Computers and mathematical algorithms have impacted our lives in many ways. In many engineering disciplines, it has become common to perform simulations for the rational design and the optimization of all kinds of devices and processes. Simulations can be much cheaper than performing many experiments and they provide theoretical and quantitative insights. Examples are airplanes, combustion engines, antennas, and the construction of bridges and other buildings.

The optimum seeking methods are also known as mathematical programming techniques and are generally studied as a part of operations research.

Methods of operations research:

- Mathematical programming techniques
 1. Calculus methods
 2. Calculus of variations
 3. Nonlinear programming
 4. Geometric programming
 5. Quadratic programming
 6. Linear programming
 7. Dynamic programming
 8. Integer programming
 9. Stochastic programming
 10. Separable programming
 11. Multiobjective programming

12. Network methods: CPM and PERT
13. Game theory
- Stochastic process techniques
 1. Statistical decision theory
 2. Markov processes
 3. Queueing theory
 4. Renewal theory
 5. Simulation methods
 6. Reliability theory
- Statistical methods
 1. Regression analysis
 2. Cluster analysis, pattern recognition
 3. Design of experiments
 4. Discriminate analysis (factor analysis)

I. LINEAR PROGRAMMING

[SI*] Linear programming is concerned with the optimization (minimization or maximization) of a linear function while satisfying a set of linear equality and/or inequality constraints or restrictions. The linear programming problem was first conceived by George B. Dantzig around 1947 while he was working as a Mathematical Advisor to the United States Air Force Comptroller on developing a mechanized planning tool for a time-staged deployment, training, and logistical supply program.

In 1949, George B. Dantzig published the "simplex method" for solving linear programs. Since that time a number of individuals have contributed to the field of linear programming in many different ways, including theoretical development, computational aspects, and exploration of new applications of the subject [4].

[SI*] The standard form of linear programming in matrix form:

Minimize

$$c^t x \tag{5.1}$$

subject to the constraints

$$Ax = b \tag{5.2}$$

and

$$x \geq 0 \tag{5.3}$$

where

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}, \quad c = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

It can be seen that there are m equations and n decision variables in a linear programming problem.

We may assume that $m < n$, for if $m > n$ there would be $m - n$ redundant equations which could be eliminated (it is the basic knowledge of linear algebra, read more here [26]). The case where $m = n$ is of no interest, for then there is either a unique solution x which satisfies Eqs. (5.2) and (5.3) (in which case there can be no optimization) or no solution, in which case the constraints are inconsistent.

the case $m < n$ corresponds to an underdetermined set of linear equations which, if they have one solution, have an infinite number of solutions. The problem of linear programming is to find one of these solutions satisfying Eqs.(5.2) and (8.3) and yielding the minimum of f .

[SI*] A linear programming problem may have

1. A unique and finite optimum solution.
2. An infinite number of optimal solutions.
3. An unbounded solution.
4. No solution.
5. A unique feasible point.

Assuming that the linear programming is properly formulated, the following general geometrical characteristics can be noted from the graphical solution.

1. The feasible region is a convex polygon.
2. The optimum value occurs at an extreme point or vertex of the feasible region.

[SI*] To represent an optimization problem as a linear program, several assumptions that are implicit in the linear programming formulation are needed, they are:

1. **Proportionality**

Given a variable x_j , its contribution to cost is $c_j x_j$ and its contribution to the i th constraint is $a_{ij} x_j$. This means that if x_j is doubled, so is its contribution to cost and to each of the constraints.

2. **Additivity**

This assumption guarantees that the total cost is the sum of the individual costs, and that the total contribution to the i th restriction is the sum of the individual contributions of the individual activities. In other words, there are no substitution or interaction effects among the activities.

3. **Divisibility**

This assumption ensures that the decision variables can be divided into any fractional levels so that noninteger values for the decision variables are permitted.

4. **Deterministic**

The coefficients c_j , a_{ij} , and b_i are all known deterministically. Any probabilistic or stochastic element inherent in demand, costs, prices, resource availabilities, usages, and so on are all assumed to be approximated by these coefficients through some deterministic equivalent.

It is important to recognize that if a linear programming problem is being used to model a given situation, then the aforementioned assumptions are implied to hold, at least over some anticipated operating range for the activities.

Despite the seemingly restrictive assumptions, linear programs are among the most widely used models today. They represent several systems quite satisfactorily, and they are capable of providing a large amount of information besides simply a solution. Moreover, they are also often used to solve certain types of nonlinear optimization problems via (successive) linear approximations and constitute an important tool in solution methods for linear discrete optimization problems having integer restricted variables.

[SI*] **Problem Manipulation**

Recall that a linear program is a problem of minimizing or maximizing a linear function in the presence of linear inequality and/or equality constraints. By simple manipulation the problem can be transformed from one form to another equivalent form.

1. Inequalities and Equations

An inequality can be easily transformed into an equation. To illustrate, consider the constraint given by

$$\sum_{j=1}^n a_{ij}x_j \geq b_i$$

this constraint can be put in an equation form by subtracting the nonnegative surplus or slack variable x_{n+i} (sometimes denoted by s_i) leading to

$$\sum_{j=1}^n a_{ij}x_j - x_{n+i} = b_i$$

and

$$x_{n+i} \geq 0$$

Similarly, the constraint

$$\sum_{j=1}^n a_{ij}x_j \leq b_i$$

is equivalent to

$$\sum_{j=1}^n a_{ij}x_j + x_{n+i} = b_i$$

and

$$x_{n+i} \geq 0$$

2. Standard Form of a Linear Programming Problem

The general linear programming problem can be stated in the following standard form

- In scalar form.

Minimize

$$f(x_1, x_2, \dots, x_n) = c_1x_1 + c_2x_2 + \dots + c_nx_n \quad (5.4)$$

subject to the constraints

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &= b_2 \\ &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n &= b_m \end{aligned} \quad (5.5)$$

and

$$\begin{aligned} x_1 &\geq 0 \\ x_2 &\geq 0 \\ &\vdots \\ x_n &\geq 0 \end{aligned} \tag{5.6}$$

- In compact form.

Minimize

$$f(x_1, x_2, \dots, x_n) = \sum_{j=1}^n c_j x_j \tag{5.7}$$

subject to the constraints

$$\sum_{j=1}^n a_{ij} x_j = b_i, \quad i = 1, 2, \dots, m \tag{5.8}$$

and

$$x_j \geq 0, \quad j = 1, 2, \dots, n \tag{5.9}$$

- In matrix form.

Minimize

$$\mathbf{c}^t \mathbf{x} \tag{5.10}$$

subject to the constraints

$$A\mathbf{x} = \mathbf{b} \tag{5.11}$$

and

$$\mathbf{x} \geq \mathbf{0} \tag{5.12}$$

where

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}, \quad \mathbf{c} = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

It can be seen that there are m equations and n decision variables in a linear programming problem, and the superscript T is used to indicate the transpose. In some linear programming books, e.g. [32], vector $\mathbf{x}$ is written with uppercase $\mathbf{X}$ which is usually confusing for those who are used to use lowercase for vectors only, and uppercase for matrix. In this book vectors will be denoted always with lowercase either with rightarrow ($\vec{x}$) or bolded ($\mathbf{x}$), while a matrix will be denoted with big uppercase without rightarrow or being bolded and will always be written first that it is a matrix.

[SI*] **Definitions and Theorems**

Some of the more powerful methods for solving linear programming problems take advantage of these characteristics.

1. Point in n -dimensional space. A point x in an n -dimensional space is characterized by an ordered set of n values of coordinates $(x_1, x_2, \dots, x_n)$. The coordinates of x are also called the components of x .
2. Line segment in n -dimensions (L). If the coordinates of two points A and B are given by $x_j^{(1)}$ and $x_j^{(2)}$ ($j = 1, 2, \dots, n$), the line segment (L) joining these points is the collection of points $x(\lambda)$ whose coordinates are given by

$$x_j = \lambda x_j^{(1)} + (1 - \lambda)x_j^{(2)}, \quad j = 1, 2, \dots, n$$

where $0 \leq \lambda \leq 1$.

Thus

$$L = \{x | x = \lambda x^{(1)} + (1 - \lambda)x^{(2)}\} \quad (5.13)$$

In one dimension, for example, it is easy to see that the definition is in accordance with our experience

$$x(\lambda) - x^{(1)} = \lambda(x^{(2)} - x^{(1)}), \quad 0 \leq \lambda \leq 1 \quad (5.14)$$

hence

$$x(\lambda) = (1 - \lambda)x^{(1)} + \lambda x^{(2)} \quad (5.15)$$

3. In n -dimensional space, the set of points coordinates satisfy a linear equation

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = \mathbf{a}^T \mathbf{x} = b \quad (5.16)$$

is called a hyperplane.

Thus, the hyperplane H is given by

$$H(A, b) = \{x | A^T x = b\} \quad (5.17)$$

A hyperplane has $(n - 1)$ dimensions in n -dimensional space. For example, in a 3-dimensional space, it is a plane and in 2-dimensional space, it is a line. The set of points whose coordinates satisfy a linear inequality like $a_1x_1 + \dots + a_nx_n \leq b$ is called a closed half space, closed due to the inclusion of equality sign in the above inequality.

A hyperplane partitions E^n into two closed half-spaces so that

$$H^+ = \{x | A^T x \geq b\} \quad (5.18)$$

and

$$H^- = \{x | A^T x \leq b\} \quad (5.19)$$

4. Convex set is a collection of points such that if $x^{(1)}$ and $x^{(2)}$ are any two points in the collection, the line segment joining them is also in the collection. If S denotes the convex set, it can be defined mathematically as follows:

$$\text{if } x^{(1)}, x^{(2)} \in S, \text{ then } x \in S$$

where

$$x = \alpha x^{(2)} + (1 - \alpha)x^{(1)}, \quad 0 \leq \alpha \leq 1$$

As an assumption, the set containing only one point is convex.

A convex polygon consists of a set of points having the property that the line segment joining any two points in the set is entirely in the convex set. In problems having more than two decision variables, the feasible region is called a convex polyhedron.

5. Convex polyhedron is a set of points common to one or more half-spaces. In particular, a convex polygon is the intersection of one or more half planes.
6. Vertex is a point in the convex set which does not lie on a line segment joining two other points of the set.
Thus, for example, every point on the circumference of a circle and each corner point of a polygon can be called a vertex or extreme point.
7. Feasible solution is any solution in linear programming problem that satisfies the constraints

$$Ax = b \quad (5.20)$$

and

$$x \geq 0 \quad (5.21)$$

8. Basic solution is a solution in which $(n - m)$ variables are set equal to zero.
The basic solution can be obtained by setting $(n - m)$ variables to zero and solving the constraint Eqs. (5.5) simultaneously.
9. Basis is the collection of variables that are not set equal to zero to obtain the basic solution.
10. Basic feasible solution is a basic solution which satisfies the nonnegativity conditions of Eqs. (5.6).
11. Nondegenerate basic feasible solution is a basic feasible solution which has got exactly m positive x_i .
12. Optimal solution is a feasible solution which optimizes the objective function.
13. Optimal basic solution is a basic feasible solution for which the objective function is optimal.

Theorem 5.1: Intersection of Convex Sets

The intersection of any number of convex sets is also convex.

Theorem 5.2: Feasible Region in Linear Programming

The feasible region of a linear programming problem is convex.

Theorem 5.3: Local Minimum in Linear Programming

Any local minimum solution is global for a linear programming problem.

Theorem 5.4: Basic Feasible Solution in Linear Programming

Every basic feasible solution is an extreme point of the convex set of feasible solutions.

Theorem 5.5: Closed and Bounded Convex Polyhedron

Let S be a closed, bounded convex polyhedron with x_{i^e} , $i = 1$ to p as the set of its extreme points. Then any vector $x \in S$ can be written as

$$\begin{aligned} x &= \sum_{i=1}^p \lambda_i x_{i^e} \\ \lambda_i &\geq 0 \\ \sum_{i=1}^p \lambda_i &= 1 \end{aligned} \tag{5.22}$$

Theorem 5.6: Closed Convex Polyhedron

Let S be a closed convex polyhedron. Then the minimum of a linear function over S is attained at an extreme point of S .

$$\begin{aligned} x &= \sum_{i=1}^p \lambda_i x_{i^e} \\ \lambda_i &\geq 0 \\ \sum_{i=1}^p \lambda_i &= 1 \end{aligned} \tag{5.23}$$

For the proof of each theorem above, you can read it here [32].

i. Simplex Method

Find the amount of each type of food to be purchased in order to meet exactly the daily requirements of a person at minimum cost. Assume that a person, on the average, requires 3000 calories and 100 grams of protein.

	Bread	Meat	Potatoes	Cabbage	Milk
Calories	2500	3000	600	100	600
Protein (grams)	80	150	20	10	40
Cost (USD/kg)	3	10	1	2	3

Table 5.1: The calorie values and the protein contents of five types of foods along with their costs.

Solution:

Let x_1, x_2, x_3, x_4 , and x_5 denote the amount of bread, meat, potatoes, cabbage, and milk to be purchased in kg per day. Then the objective function to be minimized is given by

$$f = 3x_1 + 10x_2 + x_3 + 2x_4 + 3x_5 \quad (5.24)$$

The constraints due to the calorie and protein requirements are respectively given by

$$\begin{aligned} 2500x_1 + 3000x_2 + 600x_3 + 100x_4 + 600x_5 &= 3000 \\ 80x_1 + 150x_2 + 20x_3 + 10x_4 + 40x_5 &= 100 \end{aligned} \quad (5.25)$$

The nonnegativity requirements of the decision variables are given by

$$x_i \geq 0, \quad i = 1, 2, 3, 4, 5 \quad (5.26)$$

Since $m = 2$ (the number of equality constraints) and $n = 5$ (the number of decision variables), a basic solution can be obtained by setting any of the $(n - m)$ variables equal to zero and solving the constraint equations. The number of basic solutions is

$$\frac{5!}{3!2!} = 10$$

Out of these ten basic solutions, the solution which has all the variables nonnegative and makes the objective function given by Eq. (5.4) assume a minimum value will be the optimum.

The simplex computation makes use of Gaussian elimination to compute the values of the basic variables and the objective function for all basic solutions.

This is the initial simplex tableau:

$$S = \begin{bmatrix} x_1 & x_2 & x_3 & x_4 & x_5 & f & C \\ 2500 & 3000 & 600 & 100 & 600 & 0 & 3000 \\ 80 & 150 & 20 & 10 & 40 & 0 & 100 \\ -3 & -10 & -1 & -2 & -3 & 1 & 0 \end{bmatrix} \quad (5.27)$$

Thus we make this matrix A that represents the constraints equations only:

$$A = \begin{bmatrix} 2500 & 3000 & 600 & 100 & 600 & 0 & 3000 \\ 80 & 150 & 20 & 10 & 40 & 0 & 100 \end{bmatrix} \quad (5.28)$$

The matrix A is the one that will be the input for the C++ codes, the objective function will be inputted as a vector, to create the matrix A we make two options, the first one is loading the matrix from textfile (**matrix.txt**) and the second one is to input the coefficients manually in the C++ source code (**main.cpp**).

To load a **std::vector** from a text file in C++, we need to include **<fstream>** for file operations, **<vector>** for using vector, **<string>** for reading lines, and **<iostream>** for input/output.

The loaded vector will then be saved as Symbolic matrix **B_mat** that will be used as the input to do the simplex computation.

To do the simplex computation we use this function in SymIntegration:
Symbolic simplexmethod(const SymbolicMatrix &A, vector<double> f, int C, int R, int n)

with A as the SymbolicMatrix that represent the constraint equations that we have defined in Eq. (5.8), f as the vector of objective function, C as the number of column for matrix A , R as the number of row for matrix A , and n is the number of decision variables.

```

xterm
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Simplex Method Function ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Simplex Method Function ]# ./main
Simplex Method :
A :
[2500 3000 600 100 600 0 3000]
[ 80 150 20 10 40 0 100]

*****Basic Variables*****
*****
0 1
*****

My wife is the most beautiful Goddess, she teaches me this !
A :
[2500 3000 0 0 0 0 3000]
[ 80 150 0 0 0 0 100]

A (in reduced row form) :
[2500 3000 0 0 0 0 3000]
[ 0 54 0 0 0 0 4 ]

A (in row reduced echelon form) :
[ 1 1.2 0 0 0 0 1.2 ]
[ 0 1 0 0 0 0 0.0740741]

Simplified final matrix :
[ 1 1.2 1.2 ]
[ 0 1 0.0740741]

Solution:
x_{1} = 1.11111
x_{2} = 0.0740741

Value of the objective function : 4.07407

*****Basic Variables*****
*****
0 2
*****

My wife is the most beautiful Goddess, she teaches me this !
A :
[2500 0 600 0 0 0 3000]
[ 80 0 20 0 0 0 100]

A (in reduced row form) :
[2500 0 600 0 0 0 3000]
[ 0 0 0.8 0 0 0 4 ]

```

Figure 5.1: The function in *SymIntegration* to compute the linear programming problem with simplex method (*SymIntegration/Examples/Operations Research/Test SymIntegration Simplex Method Function/main.cpp*).

```

x_{5} = -5.55112e-16
Value of the objective function : 5
*****Basic Variables*****
3 4
*****
My wife is the most beautiful Goddess, she teaches me this !
A :
[ 0  0  0  100 600  0 3000]
[ 0  0  0  10  40  0  100]
A (in reduced row form) :
[ 0  0  0  100 600  0 3000]
[ 0  0  0  10  40  0  100]
A (in row reduced echelon form) :
[ 0  0  0  0  0.0333333  0.2  0  1 ]
[ 0  0  0  0  0.1  0.4  0  1 ]
Simplified final matrix :
[0.0333333  0.2  1 ]
[ 0.1  0.4  1 ]
Solution:
x_{4} = -30
x_{5} = 10
Value of the objective function : -30
*****End of Simplex Method*****
Objective Function Value      Decision Variables
4.1                          0  1
5                            0  2
4.7                          0  3
4                            0  4
5                            1  2
-6.7                         1  3
5                            1  4
5                            2  3
5                            2  4
-30                          3  4
0
Time taken by function: 51473 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Simplex Method Function ]#

```

Figure 5.2: The function in SymIntegration to compute the linear programming problem with simplex method takes 51473 microseconds (SymIntegration/Examples/Operations Research/Test SymIntegration Simplex Method Function/main.cpp).

Due to the C++ coding, the meaning of decision variable 0 is for x_1 , 1 for x_2 and so on, from the result we know that the feasible and optimum solution occurs when we use basic variables 0 and 4, or, x_1 and x_5 , referring to bread and milk. With the values for the basic variables:

$$\begin{aligned}
 x_1 &= 1.15385 \\
 x_5 &= 0.192308 \\
 f &= 4.03846
 \end{aligned}$$

In real life, you cannot really buy 0.192308 kg of milk at that exact decimal value, so we usually round it to the nearest possible value that we can use in real life / in the market.

For comparison, we also try to use **Armadillo** to handle the Gaussian elimination part, but it becomes slower because we need to add another library when compiling.

I. Linear Programming

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Simplex Method Function ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -std=c++ -lsymintegration -larmadillo
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Simplex Method Function ]# ./main
Simplex Method :
A :
[2500 3000 600 100 600 0 3000]
[ 80 150 20 10 40 0 100]

*****Basic Variables*****
0 1
*****

My wife is the most beautiful Goddess, she teaches me this !
A :
[2500 3000 0 0 0 0 3000]
[ 80 150 0 0 0 0 100]

A (in reduced row form) :
[2500 3000 0 0 0 0 3000]
[ 0 54 0 0 0 0 4 ]

A (in row reduced echelon form) :
[ 1 1.2 0 0 0 0 1.2 ]
[ 0 1 0 0 0 0 0.0740741]

Simplified final matrix :
[ 1 1.2 1.2 ]
[ 0 1 0.0740741]

Solution:
x (1) = 1.11111
x (2) = 0.0740741

Value of the objective function : 4.07407
```

Figure 5.3: Comparing computing with simplex method with *SymIntegration* and *Armadillo* (*SymIntegration/Examples/Operations Research/Test SymIntegration Simplex Method Function/main.cpp*).

```
*****Basic Variables*****
3 4
*****

My wife is the most beautiful Goddess, she teaches me this !
A :
[ 0 0 0 100 600 0 3000]
[ 0 0 0 10 40 0 100]

A (in reduced row form) :
[ 0 0 0 100 600 0 3000]
[ 0 0 0 10 40 0 100]

A (in row reduced echelon form) :
[ 0 0 0 0.0333333 0.2 0 1 ]
[ 0 0 0 0.1 0.4 0 1 ]

Simplified final matrix :
[0.0333333 0.2 1 ]
[ 0.1 0.4 1 ]

Solution:
x (4) = -30
x (5) = 10

Value of the objective function : -30

*****End of Simplex Method*****

Objective Function Value      Decision Variables
4.1                          0 1
5                             0 2
4.7                          0 3
4                             0 4
5                             1 2
-6.7                         1 3
5                             1 4
5                             2 3
5                             2 4
-30                          3 4
0

Time taken by function: 81639 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Simplex Method Function ]#
```

Figure 5.4: The function in *SymIntegration* to compute the linear programming problem with simplex method and *Armadillo* for the Gaussian elimination part takes 82639 microseconds (*SymIntegration/Examples/Operations Research/Test SymIntegration Simplex Method Function/main.cpp*).

So we remove the dependencies toward **Armadillo** to make things easier and faster.

ii. Functions in SymIntegration Related to Linear Programming

[SI*] We will list the basic functions that can be used in linear programming with SymIntegration:

allCombinations(int n , int k , int $start_idx$, vector<int>& $current_comb$, vector<vector<int>>& all_combs)

this void function returns the all the possible index combination of k from $n + 1$ items, the $start_idx$ is set at 0, since C++ has the first index set to 0. You have to declare empty vector<int> $current_comb$ and empty vector<vector<int>> all_combs before run this void function.

contiguousCombinations(int n , int k , int $start_idx$)

this function returns the contiguous index from combination for choosing k elements from $n + 1$ elements, the $start_idx$ is set at 0, since C++ has the first index set to 0.

For example for all contiguous combinations of 4 out of 6 elements we will have

0	1	2	3
0	1	2	5
0	1	4	5
0	3	4	5
1	2	3	4
2	3	4	5

discontiguousCombinations(int n , int k , int $start_idx$)

this function returns the discontiguous index from combination for choosing k elements from $n + 1$ elements, the $start_idx$ is set at 0, since C++ has the first index set to 0.

For example for all discontiguous combinations of 4 out of 6 elements we will have

0	1	2	4
0	1	3	4
0	1	3	5
0	2	3	4
0	2	3	5
0	2	4	5
1	2	3	5
1	2	4	5
1	3	4	5

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Operations Research/Test SymIntegration Contiguous and Discontinuous Comb:
ion ]# ./main
All contiguous combinations of 4 from 6 items:
contiguous matrix:
    0          1          2          3
    0          1          2          5
    0          1          4          5
    0          3          4          5
    1          2          3          4
    2          3          4          5

All discontinuous combinations of 4 from 6 items:
discontinuous matrix:
    0          1          2          4
    0          1          3          4
    0          1          3          5
    0          2          3          4
    0          2          3          5
    0          2          4          5
    1          2          3          5
    1          2          4          5
    1          3          4          5

Time taken by function: 1285 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Operations Research/Test SymIntegration Contiguous and Discontinuous Comb:

```

Figure 5.5: Generating contiguous and discontinuous index combinations for choosing 4 elements from 6 elements (*SymIntegration/Examples/Operations Research/Test SymIntegration Contiguous and Discontinuous Combination/main.cpp*).

Symbolic simplexmethod(const SymbolicMatrix &A, vector<double> f, int C, int R, int n)
 this function will do simplex computation with A as the SymbolicMatrix that represent the constraint equations that we have defined in Eq. (5.8), f as the vector of objective function, C as the number of column for matrix A , R as the number of row for matrix A , and n is the number of decision variables.

II. NONLINEAR PROGRAMMING

[SI*]

Chapter 6

Numerical Linear Algebra

"All the girls in the world were divided into two classes: one class included all the girls in the world except her, and they had all the usual human feelings and were very ordinary girls; while the other class -herself alone- had no weaknesses and was superior to all humanity." - Anna Karenina (Leo Tolstoy)

WHY we create this chapter when we have created a book in 2023 [10] that contain a full summary of Elementary Linear Algebra? Because it is still using open source C++ library like **Eigen**, **Armadillo**, **GiNaC**, **SymbolicC++** that are created by someone else, we only learn to call the functions needed, but not deep enough. Thus starting from 2025 we are creating library for C++ that not only able to compute and solve problems in calculus, differential equations, linear programming, it grows and can also cover the field of linear algebra, so we are going to introduce a lot of features, functions in SymIntegration that can be used in Numerical Linear Algebra problems.

Instead of using open source library that already established, after branching out from **SymbolicC++**, SymIntegration can do a lot more, we learn more everyday, theoretically and practically, how to convert algorithms to a working C++ codes that can manage matrix multiplication, addition, LU decomposition, computing eigenvalues, and many more. If you want to read the full comprehensive summary read [10], in this book we will only cover the basic examples, and function in SymIntegration to solve Numerical Linear Algebra problems, there won't be theorems and explanations that are long here.

I. SOLUTIONS OF SYSTEMS OF LINEAR EQUATIONS

[SI*] Consider the nonhomogeneous linear system

$$\begin{array}{ccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \dots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \dots & + & a_{2n}x_n & = & b_2 \\ \vdots & & \vdots & & & & \vdots & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \dots & + & a_{mn}x_n & = & b_m \end{array} \quad (6.1)$$

it consists of m equations with n variables that need to be solved. We can also write it in matrix vector term like this:

$$Ax = b \quad (6.2)$$

with x is a vector of size n and b is a vector of size m . The entire solution set of $Ax = b$ can be obtained by translating the solution set of $Ax = 0$ by the vector x_0 .

When you have matrix of size $m \times n$, you just cannot apply Gaussian elimination directly to an $m \times n$ matrix problem. If you have more equations than unknowns ($m > n$), the your problem is overdetermined and you have no solution, which means you need to use something like the least squares method. If you have $Ax = b$, then instead of having $x = A^{-1}b$ (when $n = m$), then you have to do

$$x = (A^T A)^{-1} A^T b$$

In the case where you have less equations then unknowns ($m < n$), then your problem is underdetermined and you have an infinity of solutions, like the general solution of homogeneous linear system. In that case, you either pick one at random (e.g. setting some of the unknowns to an arbitrary value as parameter), or you need to use regularization, which means trying to add some extra constraints.

For the case when we have Eq. (6.1) with $m = n$ we will use Gaussian elimination, but when the size of the square matrix is very large, it is better and faster to use numerical method like *LU*-decomposition.

i. Functions in SymIntegration Related to Basic Matrix-Vector Operations

[SI*] We will list the basic functions that can be used in matrix, vector operations and loading textfile with SymIntegration:

loadMatrixFromFile(const string& filename)

this function will reads a textfile and return **vector<vector<double>>** as the output, e.g. **vector<vector<double>> matrixA loadMatrixFromFile("matrix.txt")** is a declaration of data type of **vector<vector<double>>** with name **matrixA** and the data is taken from **matrix.txt** that is in the current working directory.

loadIntMatrixFromFile(const string& filename)

this function will reads a textfile and return **vector<vector<int>>** as the output.

loadComplexMatrixFromFile(const string& filename)

this function will reads a textfile and return **vector<vector<complex<double>>>** as the output.

loadVectorFromFile(const string& filename)

this function will reads a textfile and return **vector<double>** as the output.

loadIntVectorFromFile(const string& filename)

this function will reads a textfile and return **vector<int>** as the output.

loadComplexVectorFromFile(const string& filename)

this function will reads a textfile and return **vector<complex<double>>** as the output.

print3DMatrix(vector<vector<vector<double>>>)

this function will display / print the 3D matrix that has the data type of **double**, **vector<vector<vector<double>>>**.

printMatrix(vector<vector<double>>)

this function will display / print the matrix that has the data type of **double**, **vector<vector<double>>**.

printIntMatrix(vector<vector<int>>)

this function will display / print the matrix that has the data type of **integer**, **vector<vector<int>>**.

printVector(vector<double>)

this function will display / print a vector that has the data type of **double**, **vector<double>**.

printIntVector(vector<int>)

this function will display / print a vector that has the data type of **integer**, **vector<int>**.

vector<double> flattenDoubleMatrix(vector<vector<double>> matrix)

this function will flatten a matrix with **double** data type entries into a **vector<double>**.

vector<double> Create3DMatrixfromVector(vector<double> vector), int d , int r , int c

this function will construct a 3D matrix with **double** (with size $d \times r \times c$) data type entries from a vector.

vector<vector<double>> matricizeDoubleVector(vector<double> vec, int R , int C)

this function will make a matrix of size $R \times C$ from a vector **vec** with size n , this condition has to be fulfilled: $n = R * C$.

save3DMatrixdouble(vector<vector<double>>, const string& filename)

this function save the 3D matrix / **vector<vector<vector<double>>>** of **double** data type with **filename** that we specify, e.g. "matrix.txt" or "matrix.csv."

save3DMatrixint(vector<vector<int>>, const string& filename)

this function save the 3D matrix / **vector<vector<vector<int>>>** of **integer** data type with **filename** that we specify, e.g. "matrix.txt" or "matrix.csv."

saveMatrixdouble(vector<vector<double>>, const string& filename)

this function save the matrix / **vector<vector<double>>** of **double** data type with **filename** that we specify, e.g. "matrix.txt" or "matrix.csv."

saveMatrixint(vector<vector<double>>, const string& filename)

this function save the matrix / **vector<vector<double>>** of **int** datatype with **filename** that we specify, e.g. "matrix.txt" or "matrix.csv."

saveVectordouble(vector<double>, const string& filename)

this function save the vector / **vector<double>** of **double** datatype with **filename** that we specify, e.g. "vec.txt" or "vec.csv."

saveVectorint(vector<int>, const string& filename)

this function save the vector / **vector<int>** of **integer** datatype with **filename** that we specify, e.g. "vec.txt" or "vec.csv."

saveVectorstring(vector<string>, const string& filename)

this function save the vector / **vector<string>** of **string** datatype with **filename** that we specify, e.g. "vec.txt" or "vec.csv."

addRow(vector<vector<double>> matrix, vector<double> vector_x, int k)

this function adds a row column (**vector_x**) into the k -throw in the matrix.

addColumn(vector<vector<double>> matrix, vector<double> vector_x, int k)

this function adds a vector column (**vector_x**) into the k -th column in the matrix.

deleteRow(vector<vector<double>>&, int)

this function deletes a certain row from a matrix.

deleteColumn(vector<vector<double>>&, int)

this function deletes a certain column from a matrix.

getRow(vector<vector<double>>, int)

this function extracts a certain row from a matrix.

getColumn(vector<vector<double>>, int)

this function extracts a certain column from a matrix.

The indices will be starting from 0 for the first row / first column, so if you want to take the first row you will type like this **getRow(matrixA,0)**, with **matrixA** is the matrix with **vector<vector<double>>** data type.

```
#include<bits/stdc++.h>
#include<iostream>
#include "symintegrationc++.h"
#include<vector>
#include <chrono>
#include <string>
#include <iomanip> // For std::setprecision
using namespace std::chrono;
using namespace std;

// Driver program
int main()
{
    vector<vector<double>> MatrixA = loadMatrixFromFile("matrix.txt");

    int R = MatrixA.size();
    int C = MatrixA[0].size();

    cout << "Column-1" << endl;
    for (int i = 0; i<R; i++)
    {
```



```

        // take the first column
        cout << getColumn(MatrixA,0)[i] << endl;
    }

    cout << "Row-2" << endl;
    for (int i = 0; i<C; i++)
    {
        // take the second row
        cout << getRow(MatrixA,1)[i] << endl;
    }

    cout << "Vector from row 1:"<<endl;
    vector<double> vectora = getRow(MatrixA,0) ;
    printVector(vectora);

    return 0;
}

```

Code 55: Test SymIntegration getRow and getColumn Example/main.cpp

multiply(vector<vector<double>>, vector<vector<double>>)

this function will perform multiplication of two matrices represented by vector<vector<double>>. For matrices multiplication, order matters, thus in general for non-similar matrices A and B , $AB \neq BA$.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrices Multiplication ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrices Multiplication ]# ./main
A:
    1.000000    2.000000
    3.000000    4.000000
    0.000000    1.000000

B:
    4.000000    3.000000
    2.000000    1.000000

A*B:
    8.000000    5.000000
   20.000000   13.000000
    2.000000    1.000000

Time taken by function: 977 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrices Multiplication ]#

```

Figure 6.1: The multiplication between two matrices, we load the matrices from textfiles **matrixA.txt** and **matrixB.txt** (SymIntegration/Examples/Linear Algebra/Test SymIntegration Matrices Multiplication/main.cpp).

add(vector<vector<double>>, vector<vector<double>>)

this function will perform addition of two matrices represented by vector<vector<double>>.

subtract(vector<vector<double>>, vector<vector<double>>)

this function will perform subtraction of two matrices represented by vector<vector<double>>.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrices Addition ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrices Addition ]# ./main
A:
    4.000000    3.000000
    2.000000    1.000000

B:
    1.000000    0.000000
    2.000000    3.000000

A + B:
    5.000000    3.000000
    4.000000    4.000000

Time taken by function: 1074 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrices Addition ]#

```

Figure 6.2: The addition between two matrices, we load the matrices from textfiles *matrixA.txt* and *matrixB.txt* (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Matrices Addition/main.cpp*).

transpose(const vector<vector<double>> &)

this function will perform transpose operation of a matrix represented by `vector<vector<double>>`.

dmat to replace `vector<vector<double>>`

We create a typedef defined in `include/symintegral/symintegrationnc++.h` to represents `vector<vector<double>>`, so we don't need to write `vector<vector<double>>` anymore, we can use **dmat** instead.

The compiler will process the typedef declaration when compiling each .cpp file that includes `include/symintegral/symintegrationnc++.h`, making the aliased types available for use within those files.

dvec to replace `vector<double>`

svec to replace `vector<string>`

smat to replace `vector<vector<string>>`

complexd to replace `complex<double>`

cmat to replace `vector<vector<complex<double>>>`

cvec to replace `vector<complex<double>>`

createVector(int n, double k)

this function will create a vector represented by `vector<double>` with size of n and populated with k as the entries.

createMatrix(int R, int C, double k)

this function will create a matrix represented by `vector<vector<double>>` with size of $R \times C$ and populated with k as the entries.

createIdentityMatrix(int n)

this function will create an identity matrix with size of n .

createMatrixFromColumnVectors(const vector<vector<double>>& columnVectors)

this function will create a matrix represented by `vector<vector<double>>` from column vectors that are composed by $\{v_1, v_2, \dots, v_c\}$ and all the column vectors are of the same size of r , thus the matrix will have the size of $r \times c$.

```
int main()
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    dvec veca = loadVectorFromFile("veca.txt");
    dvec vecb = loadVectorFromFile("vecb.txt");
    dvec b = loadVectorFromFile("b.txt");

    dmat columnVectors = {veca, vecb, b};
    dmat newmatrix = createMatrixFromColumnVectors(
        columnVectors);

    printMatrix(newmatrix);

    // Get ending timepoint
    auto stop = high_resolution_clock::now();
    auto duration = duration_cast<microseconds>(stop - start
    );

    cout << "\nTime taken by function: " << duration.count()
        << " microseconds" << endl;

    return 0;
}
```

Code 56: *Test SymIntegration Create Matrix from Column Vectors/main.cpp*

VandermondeMatrix(const vector<double>& vector_x, int n)

this function will return a Vandermonde matrix with type **vector<vector<double>>** from input of vector x (with size of m) and n as the number of column for the Vandermonde matrix.

The Vandermonde matrix:

$$A = \begin{bmatrix} 1 & x_1 & x_1^2 & \dots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \dots & x_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_m & x_m^2 & \dots & x_m^{n-1} \end{bmatrix}$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Vandermonde Matrix ]# ./main
Vector x:
1
2
3
4
5

Vandermonde Matrix(x,3):
      1.000000      1.000000      1.000000
      1.000000      2.000000      4.000000
      1.000000      3.000000      9.000000
      1.000000      4.000000     16.000000
      1.000000      5.000000     25.000000

Time taken by function: 788 microseconds

```

Figure 6.3: The function to create Vandermonde matrix $(x,3)$. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Vandermonde Matrix/main.cpp*).

multiplymatrixvector(vector<vector<double>> &A, vector<double> &x)

this function will multiply matrix A and vector x .

scalarmultiplication(vector<vector<double>> &A, double k)

to compute the scalar multiplication of a matrix with k .

ii. Solve Nonhomogeneous System of Linear Equation with Gaussian Elimination in SymIntegration

Suppose we have a nonhomogeneous linear system like this:

$$\begin{aligned} 0.0333x_1 + 0.2x_2 &= 1 \\ 0.1x_1 + 0.4x_2 &= 1 \end{aligned}$$

In the form of matrix and vector following the Eq. (6.2) we will have:

$$A = \begin{bmatrix} 0.0333 & 0.2 \\ 0.1 & 0.4 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

We are going to solve for

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

In SymIntegration to solve a nonhomogeneous linear system with Gaussian elimination and back substitution you can use:

solve_nhsystem(vector<vector<double>> &, vector<vector<double>> &, vector<double> &)
 it is a **void** function that returns nothing, and you can still call the solution after the function is called. The source code that handles this computation is located in **src/linearalgebra.cpp**.

Algorithm for **solve_nhsystem**:

1. Read input from textfiles of **matrix.txt** for matrix A and **vectorb.txt** for vector $\mathbf{b}$.
2. The function will perform a merge for the matrix A and vector $\mathbf{b}$ to become an augmented matrix.
3. Then the function will perform forward elimination, that will iterate through each row to create zeros below the main diagonal. Ensuring for each pivot element (diagonal element) it will be non-zero. If it is zero, the function will swap the current row with a row below that has a non-zero element in the same column (partial pivoting). If no such row exists, then it indicates that there is no unique solution. The rank of the matrix is not full. Then for each row below the current pivot row, it calculates a ratio and performs row operations to make the element below the pivot zero.
4. After forward elimination, the function will perform back substitution to obtain the solution. Once the matrix is in row echelon form (upper triangular), it solves for the last variable directly, then it substitutes the value of the last variable into the second-to-last equation to solve the second-to-last variable, and so on, working upwards till x_1 .

When we code with C++ we need to remember that there is some truncation error, and precision error, like floating-point inaccuracies when we comparing values to zero or performing divisions.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination Void Function for Square Mat
tion Void Function for Square Matrix Load from Textfiles ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination Void Function for Square Mat
rix Load from Textfiles ]# ./main
Augmented Matrix:
0.033300      0.200000      1.000000
0.100000      0.400000      1.000000

Augmented Matrix in reduced row form:
0.100000      0.400000      1.000000
0.000000      0.066800      0.667000

Solution:
x1 = 9.98503
x2 = -29.94012

Time taken by function: 1151 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination Void Function for Square Mat
rix Load from Textfiles ]# ./main
Augmented Matrix:
0.033300      0.200000      1.000000
0.100000      0.400000      1.000000

Augmented Matrix in reduced row form:
0.100000      0.400000      1.000000
0.000000      0.066800      0.667000

Solution:
x1 = 9.98503
x2 = -29.94012

Time taken by function: 1159 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination Void Function for Square Mat
rix Load from Textfiles ]# ./main
Augmented Matrix:
0.033300      0.200000      1.000000
0.100000      0.400000      1.000000

Augmented Matrix in reduced row form:
0.100000      0.400000      1.000000
0.000000      0.066800      0.667000

Solution:
x1 = 9.98503
x2 = -29.94012

Time taken by function: 978 microseconds

```

Figure 6.4: The Gaussian elimination with forward elimination and back substitution, we load the matrix from textfiles *matrix.txt* and *vectorb.txt*. We run the code 3 times to see the running time of the function from 978 microseconds to 1151 microseconds (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Gaussian Elimination Void Function for Square Matrix Load from Textfiles/main.cpp*).

We use **void** function for this Gaussian elimination because in C++ a void function is a function that does not return any value to the calling code. Instead of a specific data type (like int, double, string) as its return type, some functions in *SymIntegration* uses **void** for ease of use, it can be used for printing output to the console or a file, modifying global variables or variables passed by reference, performing calculations and storing results in external structures, and calling other functions to achieve a specific task.

We also compare it with Armadillo' function `solve(A,B,solve_opts::force_approx)`, turns out we can gain faster execution time, it is probably because we code it ourselves and can make it more efficient to return the solution of x . In Armadillo you need to print the solution yourself. In *SymIntegration* currently, we are printing out the augmented matrix, the reduced row form of the augmented matrix, and the solution when you execute / run the function `solve_nhsystem`.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimina
tion Void Function for Square Matrix Load from Textfiles/Gaussian Elimination fo
r Square Matrix with Armadillo ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -larmadillo
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination Void Function for Square Ma
trix Load from Textfiles/Gaussian Elimination for Square Matrix with Armadillo ]# ./main
Matrix A:
0.0333 0.2000
0.1000 0.4000

Vector B:
1.0000
1.0000

Solution:
-29.9401
9.9850

Time taken by function: 1551 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination Void Function for Square Ma
trix Load from Textfiles/Gaussian Elimination for Square Matrix with Armadillo ]# 

```

Figure 6.5: The comparison of solving a linear system with Armadillo, we load the matrix from textfiles *matrixA.txt* and *vectorB.txt*. (DianFreya Math Physics Simulator/Source Codes/C++/C+++ Gnuplot SymbolicC++/ch23-Numerical Linear Algebra/Gaussian Elimination for Square Matrix with Armadillo/main.cpp).

The code that is using Armadillo (so the reader can check it themselves):

```

#include <iostream>
#include <iomanip> // to declare the manipulator of setprecision()
#include <fstream>
#include <bits/stdc++.h> //for setw(6) at display() function
#include <vector>
#include <armadillo>
#include <chrono>

using namespace std::chrono;
using namespace std;
using namespace arma;

// Driver code
int main(int argc, char** argv)
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    mat A;
    A.load("matrixA.txt");
    mat B;
    B.load("vectorB.txt");
    mat X;
    X = solve(A,B,solve_opts::force_approx);

    cout <<"Matrix A:" << "\n" << A <<endl;
    cout <<"Vector B:" << "\n" << B <<endl;
    cout <<"Solution:" << "\n" << X <<endl;

    // Get ending timepoint

```

```
    auto stop = high_resolution_clock::now();
    auto duration = duration_cast<microseconds>(stop - start);

    cout << "\nTime taken by function: " << duration.count() << " microseconds"
          << endl;
}
```

Code 57: *main.cpp*

II. DETERMINANT

[SI*] Suppose we have a 2×2 matrix A

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

the matrix A is invertible if $ad - bc \neq 0$, and the expression $ad - bc$ is called the determinant.

$$\det(A) = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Thus, the inverse of A can be expressed in terms of the determinant as

$$A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \quad (6.3)$$

Now, we will find the formula that applicable to square matrices of all orders, not only of size 2.

Definition 6.1: Minor and Cofactor

If A is a square matrix, then the minor of entry a_{ij} is denoted by M_{ij} and is defined to be the determinant of the submatrix that remains after the i -th row and j -th column are deleted from A . The number

$$C_{ij} = (-1)^{i+j} M_{ij} \quad (6.4)$$

is called the cofactor of entry a_{ij} .

Theorem 6.1: Row or Column of Cofactors

If A is an $n \times n$ matrix, then regardless of which row or column of A is chosen, the number obtained by multiplying the entries in that row or column by the corresponding cofactors and adding the resulting products is always the same.

Definition 6.2: Cofactor Expansion

If A is an $n \times n$ matrix, then the number obtained by multiplying the entries in any row or column of A by the corresponding cofactors and adding the resulting products is called the determinant of A , and the sums themselves are called cofactor expansions of A . That is,

$$\det(A) = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj} \quad (6.5)$$

(cofactor expansion along the j -th column)

$$\det(A) = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in} \quad (6.6)$$

(cofactor expansion along the i -th row)

[SI*] To find the determinant of size n that is very large will need more computational cost, since we will peel the cofactor expansion along the i -th row or j -th column down to the matrix of size 2×2 . It is an ordered-recursive computation technique.

[SI*] The concept of a determinant is mathematically defined only for square matrices. Non-square matrices, also known as rectangular matrices, do not possess a determinant in the conventional sense.

The determinant of a square matrix represents the scaling factor of the volume (or area in 2D) when a linear transformation is applied. This geometric is inherently tied to dimensions being equal. The determinant is also crucial for calculating the inverse of a matrix, which is only defined for square, non-singular matrices.

[SI*] **Alternative Concepts for Non-Square Matrices**

While a determinant doesn't exist for non-square matrices, related concepts can be used in certain contexts:

- **Singular Values**

Non-square matrices have singular values, which are related to the scaling of principal axes in the transformation defined by the matrix. The product of singular values can be related to a "volume" in a generalized sense.

- **Gramian Determinant**

For a matrix A , the Gramian matrix $A^T A$ (or AA^T) is always square. Its determinant, known as the Gramian determinant, can be used in some applications involving non-square matrices.

i. Functions in SymIntegration Related to Determinant

[SI*] We will list the basic functions that can be used related to determinant and inverse:

determinant(const vector<vector<double>> &matrix)

this function computes the determinant of a square matrix.

inverse(vector<vector<double>> &matrix)

this function computes the inverse of a square matrix that is non-empty. The computation of an inverse of a matrix is a complex one, it needs adjugate, cofactor and determinant computation.

ii. Compute Determinant of a Square Matrix with SymIntegration

Suppose we have a matrix A

$$A = \begin{bmatrix} 3 & -2 & 7 \\ -2 & 4 & -3 \\ -1 & 9 & 4 \end{bmatrix}$$

to compute the determinant with SymIntegration we will use this function:

determinant(const vector<vector<double>> &matrix)

The full C++ source code can be seen below:

```
#include<bits/stdc++.h>
#include<iostream>
#include "symintegrationc++.h"
#include<vector>
#include <chrono>
#include <algorithm> // For std::next_permutation
#include <string>
using namespace std::chrono;
using namespace std;

// Driver program
int main()
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    //string filename = "matrix.txt";

    vector<vector<double>> doubleMatrix = loadMatrixFromFile("matrix.txt");

    cout << "\nMatrix A : " << endl;
    printMatrix(doubleMatrix);
    cout << "\nDeterminant of matrix A : " << determinant(doubleMatrix) << endl
        ;

    // Get ending timepoint
    auto stop = high_resolution_clock::now();
    auto duration = duration_cast<microseconds>(stop - start);

    cout << "\nTime taken by function: " << duration.count() << " microseconds"
        << endl;

    return 0;
}
```

Code 58: Test SymIntegration Compute Determinant/main.cpp

```
Time taken by function: 751 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Determinant ]# ./main

Matrix A :
      3.000000      -2.000000      7.000000
     -2.000000      4.000000     -3.000000
     -1.000000      9.000000      4.000000

Determinant of matrix A : 9.000000

Time taken by function: 751 microseconds
```

Figure 6.6: The computation of the determinant of matrix A (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Determinant/main.cpp*).

III. EUCLIDEAN VECTOR SPACES

[SI*] Engineers and physicists represent vectors in two dimensions or in three dimensions by arrows. The direction of the arrowhead specifies the direction of the vector and the length of the arrow specifies the magnitude. The tail of the arrow is called the initial point of the vector and the tip the terminal point.

We usually denote vectors with boldface such as $\mathbf{a}, \mathbf{b}, \mathbf{v}, \mathbf{w}$ or with an arrow above it $\vec{a}, \vec{b}, \vec{v}, \vec{w}$, and we denote scalars in lowercase letters like a, b, c, d . When we want to indicate that a vector has initial point A and terminal point B , we will write

$$\mathbf{v} = \vec{AB}$$

[SI*] We denote the length of a vector $\mathbf{v}$ by the symbol $\|\mathbf{v}\|$, which is read as norm of $\mathbf{v}$, or the magnitude of $\mathbf{v}$.

Definition 6.3: Norm

If $\mathbf{v} = (v_1, v_2, \dots, v_n)$ is a vector in $\mathbb{R}^n$, then the norm of $\mathbf{v}$ is denoted by $\|\mathbf{v}\|$, and is defined by the formula

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2} \quad (6.7)$$

[SI*] A vector of norm 1 is called a unit vector. Such vectors are useful for specifying a direction when length is not relevant to the problem at hand. If $\mathbf{v}$ is any nonzero vector in $\mathbb{R}^n$, then

$$\mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v} \quad (6.8)$$

defines a unit vector that is in the same direction as $\mathbf{v}$. The process of multiplying a nonzero vector by the reciprocal of its length to obtain a unit vector is called normalizing $\mathbf{v}$.

Definition 6.4: Distance in $\mathbb{R}^n$

If $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ are points in $\mathbb{R}^n$, then we denote the distance between $\mathbf{u}$ and $\mathbf{v}$ by $d(\mathbf{u}, \mathbf{v})$ and define it to be

$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\| = \sqrt{(u_1 - v_1)^2 + (u_2 - v_2)^2 + \dots + (u_n - v_n)^2} \quad (6.9)$$

Definition 6.5: Angle Between Two Vectors in $\mathbb{R}^2$ and $\mathbb{R}^3$

If $\mathbf{u}$ and $\mathbf{v}$ are nonzero vectors in $\mathbb{R}^2$ or $\mathbb{R}^3$, and if θ is the angle between $\mathbf{u}$ and $\mathbf{v}$, then the dot product of $\mathbf{u}$ and $\mathbf{v}$ is denoted by $\mathbf{u} \cdot \mathbf{v}$ and is defined as

$$\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta \quad (6.10)$$

If $\mathbf{u} = \mathbf{0}$ or $\mathbf{v} = \mathbf{0}$, then we define $\mathbf{u} \cdot \mathbf{v}$ to be 0.

Since $0 \leq \theta \leq \pi$, it follows from the properties of the cosine function that

- θ is acute if $\mathbf{u} \cdot \mathbf{v} > 0$.
- θ is obtuse if $\mathbf{u} \cdot \mathbf{v} < 0$.
- $\theta = \frac{\pi}{2}$ if $\mathbf{u} \cdot \mathbf{v} = 0$.

Definition 6.6: Dot Product in $\mathbb{R}^n$

If $\mathbf{u}$ and $\mathbf{v}$ are vectors in $\mathbb{R}^n$, then the dot product or the Euclidean inner product of $\mathbf{u}$ and $\mathbf{v}$ is denoted by $\mathbf{u} \cdot \mathbf{v}$ and is defined by

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + \cdots + u_nv_n \quad (6.11)$$

Definition 6.7: Orthogonal and Orthonormal

Two nonzero vectors $\mathbf{u}$ and $\mathbf{v}$ in $\mathbb{R}^n$ are said to be orthogonal (or perpendicular) if $\mathbf{u} \cdot \mathbf{v} = 0$.

The zero vector ($\mathbf{0}$) in $\mathbb{R}^n$ is orthogonal to every vector in $\mathbb{R}^n$.

A nonempty set of vectors in $\mathbb{R}^n$ is called an orthogonal set if all pairs of distinct vectors in the set are orthogonal.

An orthogonal set of unit vectors is called an orthonormal set.

[SI*] The vector equation

$$\mathbf{n} \cdot \vec{P_0P} = 0$$

is the representation of a line through the point $P_0(x_0, y_0)$ that has normal $\mathbf{n} = (a, b)$ or a plane through the point $P_0(x_0, y_0, z_0)$ that has normal $\mathbf{n} = (a, b, c)$.

For line, the vector $\vec{P_0P}$ can be expressed in terms of components as

$$\vec{P_0P} = (x - x_0, y - y_0)$$

For plane, the vector $\vec{P_0P}$ can be expressed in terms of components as

$$\vec{P_0P} = (x - x_0, y - y_0, z - z_0)$$

[SI*] The point-normal equation of the line:

$$a(x - x_0) + b(y - y_0) = 0 \quad (6.12)$$

Theorem 6.2: Equation for Line and Plane

If a and b are constants that are not both zero, then an equation of the form

$$ax + by + c = 0 \quad (6.13)$$

represents a line in $\mathbb{R}^2$ with normal $\mathbf{n} = (a, b)$.

If a, b , and c are constants that are not all zero, then an equation of the form

$$ax + by + cz + d = 0 \quad (6.14)$$

represents a plane in $\mathbb{R}^3$ with normal $\mathbf{n} = (a, b, c)$.

[SI*] There are other useful ways of specifying lines and planes.

A unique line in $\mathbb{R}^2$ or $\mathbb{R}^3$ is determined by a point $\mathbf{x}_0$ on the line and a nonzero vector $\mathbf{v}$ parallel to the line.

A unique plane in $\mathbb{R}^3$ is determined by a point $\mathbf{x}_0$ in the plane and two collinear vectors $\mathbf{v}_1$ and $\mathbf{v}_2$ parallel to the plane. The best way to visualize this is to translate the vectors so their initial points are at $\mathbf{x}_0$.

Theorem 6.3: Vector and Parametric Equations of Lines in $\mathbb{R}^2$ and $\mathbb{R}^3$

Let L be the line in $\mathbb{R}^2$ or $\mathbb{R}^3$ that contains the point $\mathbf{x}_0$ and is parallel to the nonzero vector $\mathbf{v}$. Then the equation of the line through $\mathbf{x}_0$ that is parallel to $\mathbf{v}$ is

$$\mathbf{x} = \mathbf{x}_0 + t\mathbf{v} \quad (6.15)$$

If $\mathbf{x}_0 = \mathbf{0}$, then the line passes through the origin and the equation has the form

$$\mathbf{x} = t\mathbf{v} \quad (6.16)$$

the variable t is called a parameter, and it varies from $-\infty$ to ∞ .

Theorem 6.4: Vector and Parametric Equations of Planes in $\mathbb{R}^3$

Let W be the plane in $\mathbb{R}^3$ that contains the point $\mathbf{x}_0$ and is parallel to the noncollinear vectors $\mathbf{v}_1$ and $\mathbf{v}_2$. Then an equation of the plane through $\mathbf{x}_0$ that is parallel to $\mathbf{v}_1$ and $\mathbf{v}_2$ is given by

$$\mathbf{x} = \mathbf{x}_0 + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 \quad (6.17)$$

If $\mathbf{x}_0 = \mathbf{0}$, then the plane passes through the origin and the equation has the form

$$\mathbf{x} = t_1\mathbf{v}_1 + t_2\mathbf{v}_2 \quad (6.18)$$

Definition 6.8: Parametric Equations of Lines in $\mathbb{R}^n$

If $\mathbf{x}_0$ and $\mathbf{v}$ are vectors in $\mathbb{R}^n$, and if $\mathbf{v}$ is nonzero, then the equation

$$\mathbf{x} = \mathbf{x}_0 + t\mathbf{v} \quad (6.19)$$

defines the line through $\mathbf{x}_0$ that is parallel to $\mathbf{v}$. In the special case where $\mathbf{x}_0 = \mathbf{0}$, the line is said to pass through the origin.

Definition 6.9: Parametric Equations of Planes in $\mathbb{R}^n$

If $\mathbf{x}_0$, $\mathbf{v}_1$ and $\mathbf{v}_2$ are vectors in $\mathbb{R}^n$, and if $\mathbf{v}_1$ and $\mathbf{v}_2$ are not collinear, then the equation

$$\mathbf{x} = \mathbf{x}_0 + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 \quad (6.20)$$

defines the plane through $\mathbf{x}_0$ that is parallel to $\mathbf{v}_1$ and $\mathbf{v}_2$. In the special case where $\mathbf{x}_0 = \mathbf{0}$, the plane is said to pass through the origin.

Definition 6.10: Cross Product

If $\mathbf{u} = (u_1, u_2, u_3)$ and $\mathbf{v} = (v_1, v_2, v_3)$ are vectors in 3-space, then the cross product $\mathbf{u} \times \mathbf{v}$ is the vector defined by

$$\mathbf{u} \times \mathbf{v} = (u_2v_3 - u_3v_2, u_3v_1 - u_1v_3, u_1v_2 - u_2v_1) \quad (6.21)$$

or, in determinant notation,

$$\mathbf{u} \times \mathbf{v} = \left(\begin{vmatrix} u_2 & u_3 \\ v_2 & v_3 \end{vmatrix}, -\begin{vmatrix} u_1 & u_3 \\ v_1 & v_3 \end{vmatrix}, \begin{vmatrix} u_1 & u_2 \\ v_1 & v_2 \end{vmatrix} \right) \quad (6.22)$$

Definition 6.11: Scalar Triple Product

If $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are vectors in 3-space, then

$$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix} \quad (6.23)$$

is called the scalar triple product of $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$, and $\mathbf{w} = (w_1, w_2, w_3)$.

It follows that

$$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \mathbf{w} \cdot (\mathbf{u} \times \mathbf{v}) = \mathbf{v} \cdot (\mathbf{w} \times \mathbf{u})$$

Theorem 6.5: Geometric Interpretation of Determinants

(a) The absolute value of the determinant

$$\det \begin{bmatrix} u_1 & u_2 \\ v_1 & v_2 \end{bmatrix}$$

is equal to the area of the parallelogram in 2-space determined by the vectors $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$.

(b) The absolute value of the determinant

$$\det \begin{bmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{bmatrix}$$

is equal to the volume of the parallelepiped in 3-dimension determined by the vectors $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$, and $\mathbf{w} = (w_1, w_2, w_3)$.

i. Functions in SymIntegration Related to Euclidean Vector Spaces

[SI*] We will list the basic functions that can be used related to Euclidean vector spaces:

distance(const vector<double> &vectorx, const vector<double> &vectory)

to compute the distance between two points (the point can be represented as n -tuple / array / **vector<double>**).

distance(const Symbolic &f, const Symbolic &x, const Symbolic &y, const Symbolic &z, const vector<double> &vectorx)

to compute the distance between a plane / a line that is defined with function $f(x, y, z)$ / $f(x, y)$ and a point that is represented as vector x .

dot(const vector<double> &vectorx, const vector<double> &vectory)

to compute the dot product / Euclidean inner product of two vectors.

angle(const vector<double> &vectorx, const vector<double> &vectory)

to compute the angle between two vectors.

norm(const vector<double> &vectorx)

to compute the norm of a vector.

add(vector<double> &vectorA, vector<double> &vectorB)

to compute the addition between two vectors.

subtract(vector<double> &vectorA, vector<double> &vectorB)

to compute subtraction between two vectors.

scalarmultiplication(vector<double> &vectorA, double k)

to compute the scalar multiplication of a vector with k .

orthogonalprojection(vector<double> vectorx, vector<double> vectory)

to compute the orthogonal projection of x on y .

vectorequation(const Symbolic &f, const Symbolic &x, const Symbolic &y, const Symbolic &z)

to compute the vector equation of the form $x = x_0 + tv_1 + sv_2$ that represents a plane / a line through x_0 and parallel to v_1 and/or v_2 (for a line we will only have one v) with given input of a function $f(x, y, z) = 0$.

vectorequationdecomp(const Symbolic &f, const Symbolic &x, const Symbolic &y, const Symbolic &z, vector<double> &vectorv1, vector<double> &vectorv2, vector<double> &vectorv3)

to compute all the vectors that compose the vector equation / parametric equations of the form $x = x_0 + tv_1 + sv_2$ that represents a plane / a line through x_0 and parallel to v_1 and/or v_2 (for a line we will only have one v) with given input of a function $f(x, y, z) = 0$. It is a **void** function, so we will need to declare the vector v_1, v_2 , and v_3 before calling the function, afterwards we can print the vectors.

crossproduct(vector<double> &vectoru, vector<double> &vectorv)

to compute the cross product of $u \times v$.

scalartripleproduct(vector<double> &vectoru, vector<double> &vectorv, vector<double> &vectorw)

to compute the scalar triple product of u, v , and w .

areaoftriangle(vector<double> &x, vector<double> &y, vector<double> &z)

to compute the area of triangle in $\mathbb{R}^n$, given 3 vertices that we input as **vector<double>** type, x, y, z . We use the formula with two known sides and the angle between these two sides

$$A = \frac{1}{2}ab \sin(c)$$

it is more efficient than using determinant formula to compute the area.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Area of triangle in n-dimension ]# ./main
Vertices, a :
-1
4
b :
3
1
c :
2
6
Area of triangle : 8.5
Time taken by function: 808 microseconds
```

Figure 6.7: The computation of the area of triangle given 3 vertices: a, b and c in $\mathbb{R}^2$ (SymIntegration/Examples/-Linear Algebra/Test SymIntegration Area of triangle in n-dimension/main.cpp).

ii. Compute Norm, Dot Product, Angle with SymIntegration

Suppose we have two vectors :

$$\mathbf{x} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\mathbf{y} = \begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix}$$

we can compute the norm, dot product, and the angle between $\mathbf{x}$ and $\mathbf{y}$ with SymIntegration' functions.

$$\begin{aligned} \|\mathbf{x}\| &= 1 \\ \|\mathbf{y}\| &= 2.82843 \\ \mathbf{x} \cdot \mathbf{y} &= 2 \\ \theta &= \cos^{-1} \left(\frac{\mathbf{x} \cdot \mathbf{y}}{\|\mathbf{x}\| \|\mathbf{y}\|} \right) = 0.70710678 \text{ rad} = 45^\circ \end{aligned}$$

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Euclidean Vector Spaces ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Euclidean Vector Spaces ]# ./main

Vector x :
0
0
1

Vector y :
0
2
2

norm(x) : 1
norm(y) : 2.82843
dot(x,y) : 2
angle(x,y) in degree: 45

Time taken by function: 689 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Euclidean Vector Spaces ]#
```

Figure 6.8: The computation of the norm, dot product, and angle of two vectors $\mathbf{x}$ and $\mathbf{y}$ (SymIntegration/Examples/-Linear Algebra/Test SymIntegration Euclidean Vector Spaces/main.cpp).

iii. Compute Orthogonal Projection on a Line with SymIntegration

Let $\mathbf{u} = (2, -1, 3)$ and $\mathbf{a} = (4, -1, 2)$. Find the vector component of $\mathbf{u}$ along $\mathbf{a}$ and the vector component of $\mathbf{u}$ orthogonal to $\mathbf{a}$.

Solution:

$$\begin{aligned}\mathbf{u} \cdot \mathbf{a} &= (2)(4) + (-1)(-1) + (3)(2) = 15 \\ \|\mathbf{a}\|^2 &= 4^2 + (-1)^2 + 2^2 = 21\end{aligned}$$

Then the vector component of $\mathbf{u}$ along $\mathbf{a}$ is

$$\begin{aligned}\text{proj}_{\mathbf{a}} \mathbf{u} &= \frac{\mathbf{u} \cdot \mathbf{a}}{\|\mathbf{a}\|^2} \mathbf{a} \\ &= \frac{15}{21} (4, -1, 2) \\ &= \left(\frac{20}{7}, -\frac{5}{7}, \frac{10}{7} \right)\end{aligned}$$

and the vector component of $\mathbf{u}$ orthogonal to $\mathbf{a}$ is

$$\begin{aligned}\mathbf{u} - \text{proj}_{\mathbf{a}} \mathbf{u} &= (2, -1, 3) - \left(\frac{20}{7}, -\frac{5}{7}, \frac{10}{7} \right) \\ &= \left(-\frac{6}{7}, -\frac{2}{7}, \frac{11}{7} \right)\end{aligned}$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Orthogonal Projection ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Orthogonal Projection ]# ./main

Vector u :
2
-1
3

Vector a :
4
-1
2

norm(a)^2 : 29.6985

dot(u,a) : 15

Orthogonal projection of u on a:
2.85714
-0.714286
1.42857

Vector component of u orthogonal to a:
-0.857143
-0.285714
1.57143

u + a :
6
-2
5

u - a :
-2
0
1

5*u :
10
-5
15

Time taken by function: 1066 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Orthogonal Projection ]# 

```

Figure 6.9: The computation of the orthofonal projection of vector u on a and some other vector operations. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Orthogonal Projection/main.cpp*).

iv. Compute Vector and Parametric Equations of a Plane in $\mathbb{R}^3$ with SymIntegration

Find vector and parametric equations of the plane $x - y + 2z = 5$.

Solution:

We will find the parametric equations first. We can do this by solving the equation for any one of the variables in terms of the other two and then using those two variables as parameters.

For example, solving for x in terms of y and z yields

$$x = 5 + y - 2z$$

and then using y and z as parameters t and s , respectively, yields the parametric equations

$$x = 5 + t - 2s$$

$$y = t$$

$$z = s$$

To obtain a vector equation of the plane we rewrite these parametric equations as

$$(x, y, z) = (5 + t - 2s, t, s)$$

or, equivalently, as

$$(x, y, z) = (5, 0, 0) + t(1, 1, 0) + s(-2, 0, 1)$$

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Vector and Parametric Equations ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Vector and Parametric Equations ]# ./main

Vector equation of the plane x - y + 2z - 5 = 0:
[t-2*s+5]
[ t ]
[ s ]

The parametric equations in vector terms:
(x,y,z) = v1 + v2*t + v2*s
v1 =
5
0
0

v2 =
1
1
0

v3 =
-2
0
1

Time taken by function: 29519 microseconds
```

Figure 6.10: The computation of vector equation of the plane in parametric equations term and then in vector terms. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Vector and Parametric Equations/main.cpp*).

IV. GENERAL VECTOR SPACES

[SI*] In Euclidean Vector Spaces we developed properties of vectors in $\mathbb{R}^2$ and $\mathbb{R}^3$, we noticed patterns in various formulas that enabled us to extend the notion of a vector to an n -tuple of real numbers. Although n -tuple took us outside the realm of our "visual experience," it gave us a valuable tool for understanding and studying systems of linear equations.

Definition 6.12: Vector Space Axioms

Let V be an arbitrary nonempty set of objects on which two operations are defined: addition, and multiplication by scalars.

- By addition we mean a rule for associating with each pair of objects u and v in V an object $u + v$, called the sum of u and v .
- By scalar multiplication we mean a rule for associating with each scalar k and each object u in V an object ku , called the scalar multiple of u by k .

If the following axioms are satisfied by all objects u, v, w in V and all scalars k and m , then we call V a vector space and we call the objects in V vectors

1. If u and v are objects in V , then $u + v$ is in V
2. $u + v = v + u$
3. $u + (v + w) = (u + v) + w$
4. There is an object 0 in V , called a zero vector for V , such that

$$0 + u = u + 0 = u$$

for all u in V

5. For each u in V , there is an object $-u$ in V , called a negative of u , such that

$$u + (-u) = (-u) + u = 0$$

6. If k is any scalar and u is any object in V , then ku is in V .
7. $k(u + v) = ku + kv$
8. $(k + m)u = ku + mu$
9. $k(mu) = (km)(u)$
10. $1u = u$

Definition 6.13: Subspace

A subset W of a vector space V is called a subspace of V if W is itself a vector space under the addition and scalar multiplication defined on V .

In general, to show that a nonempty set W with two operations is a vector space one must verify the ten vector space axioms.

Definition 6.14: Linear Combination

If w is a vector in a vector space V , then w is said to be a linear combination of the vectors $v_1, v_2, \dots, v_r$ in V if w can be expressed in the form

$$w = k_1 v_1 + k_2 v_2 + \dots + k_r v_r \quad (6.24)$$

where $k_1, k_2, \dots, k_r$ are scalars. These scalars are called the coefficients of the linear combination.

Theorem 6.6: Linear Combination and Subspace

If $S = \{w_1, w_2, \dots, w_r\}$ is a nonempty set of vectors in a vector space V , then:

1. The set W of all possible linear combinations of the vectors in S is a subspace of V .
2. The set W in part (a) is the "smallest" subspace of V that contains all of the vectors in S in the sense that any other subspace that contains those vectors contains W .

Definition 6.15: Span of a Subspace

The subspace of a vector space V that is formed from all possible linear combinations of the vectors in a nonempty set S is called the span of S , and we say that the vectors in S span that subspace. If $S = \{w_1, w_2, \dots, w_r\}$, then we denote the span of S by

$$\text{span}\{w_1, w_2, \dots, w_r\} \quad \text{or} \quad \text{span}(S)$$

[SI*] The standard unit vectors in $\mathbb{R}^n$ are

$$e_1 = (1, 0, 0, \dots, 0), \quad e_2 = (0, 1, 0, \dots, 0), \dots, \quad e_n = (0, 0, 0, \dots, 1)$$

these vectors span $\mathbb{R}^n$ since every vector $v = (v_1, v_2, \dots, v_n)$ in $\mathbb{R}^n$ can be expressed as

$$v = v_1 e_1 + v_2 e_2 + \dots + v_n e_n$$

Theorem 6.7: Solution of Homogeneous Linear System

The solution set of a homogeneous linear system $Ax = 0$ in n unknowns is a subspace of $\mathbb{R}^n$.

[SI*] The solution set of every homogeneous system of m equations in n unknowns is a subspace of $\mathbb{R}^n$. But, the solution set of a nonhomogeneous system of m equations in n unknowns is a

subspace of $\mathbb{R}^n$ is never true, since there are two possible scenarios: first, the system may not have any solution at all, and second, if there are solutions, then the solution set will not be closed under either addition or under scalar multiplication.

Theorem 6.8: Span and Linear Combination

If $S = \{v_1, v_2, \dots, v_r\}$ and $S' = \{w_1, w_2, \dots, w_k\}$ are nonempty sets of vectors in a vector space V , then

$$\text{span}\{v_1, v_2, \dots, v_r\} = \text{span}\{w_1, w_2, \dots, w_k\}$$

if and only if each vector in S is a linear combination of those in S' , and each vector in S' is a linear combination of those in S .

[SI*] In a rectangular xy -coordinate system every vector in the plane can be expressed in exactly one way as a linear combination of the standard unit vectors.

Definition 6.16: Linear Independence

If $S = \{v_1, v_2, \dots, v_r\}$ is a nonempty set of vectors in a vector space V , then the vector equation

$$k_1 v_1 + k_2 v_2 + \dots + k_r v_r = \mathbf{0}$$

has at least one solution, namely

$$k_1 = 0, \quad k_2 = 0, \quad \dots, \quad k_r = 0$$

we call this the trivial solution. If this is the only solution, then S is said to be a linearly independent set. If there are solutions in addition to the trivial solution, then S is said to be a linearly dependent set.

[SI*] To show that a linear system has a nontrivial solutions you can

1. Solve it directly with Gaussian Elimination, when you have this vector equation

$$k_1 v_1 + k_2 v_2 + \dots + k_r v_r = \mathbf{0}$$

then turns it into homogeneous system and make a coefficient matrix out of it.

2. Check the determinant of the coefficient matrix, if the determinant is zero then the coefficient matrix has infinitely many solutions / nontrivial solutions. If the determinant is not zero then the system has only the trivial solution, and the vectors are linearly independent.

Theorem 6.9: Interpretation of Linear Independence

A set S with two or more vectors is

- (a) Linearly dependent if and only if at least one of the vectors in S is expressible as a linear combination of the other vectors in S .
- (b) Linearly independent if and only if no vector in S is expressible as a linear combination of the other vector in S .

Theorem 6.10: Sets With One or Two Vectors

- (a) A finite set that contains $\mathbf{0}$ is linearly dependent.
- (b) A set with exactly one vector is exactly independent if and only if that vector is not $\mathbf{0}$.
- (c) A set with exactly two vectors is linearly independent if and only if neither vector is a scalar multiple of the other.

Theorem 6.11: Geometric Interpretation of Linear Independence

Let $S = \{v_1, v_2, \dots, v_r\}$ be a set of vectors in $\mathbb{R}^n$. If $r > n$, then S is linearly dependent.

[SI*] Sometimes linear dependence of functions can be deduced from known identities. But, there is no general method that can be used to determine whether a set of functions is linear independent or linear dependent.

Definition 6.17: Wronskian

If $f_1 = f_1(x)$, $f_2 = f_2(x)$, $\dots$, $f_n = f_n(x)$ are functions that are $n - 1$ times differentiable on the interval $(-\infty, \infty)$, then the determinant

$$\begin{vmatrix} f_1(x) & f_2(x) & \dots & f_n(x) \\ f_1'(x) & f_2'(x) & \dots & f_n'(x) \\ \vdots & \vdots & & \vdots \\ f_1^{(n-1)} & f_2^{(n-1)} & \dots & f_n^{(n-1)} \end{vmatrix}$$

is called the Wronskian of $f_1, f_2, \dots, f_n$.

Theorem 6.12: Wronskian and Linear Independence of Functions

If the functions $f_1, f_2, \dots, f_n$ have $n - 1$ continuous derivatives on the interval $(-\infty, \infty)$, and if the Wronskian of these functions is not identically zero on $(-\infty, \infty)$, then these functions form a linearly independent set of vectors in $C^{(n-1)}(-\infty, \infty)$.

[SI*] Rectangular coordinate systems are common, but they are not essential. There are nonrectangular coordinate systems in 2-space, 3-space, to the n -space with equal spacing, skew axes (the major axis is not orthogonal to each other) with equal spacing or unequal spacing. Our common rectangular coordinate system has perpendicular axes.

Definition 6.18: Basis for a Vector Space

If V is any vector space and $S = \{v_1, v_2, \dots, v_n\}$ is a finite set of vectors in V , then S is called a basis for V if the following two conditions hold:

- (a) S is linearly independent.
- (b) S spans V .

[SI*] To prove the vectors $v_1, v_2, \dots, v_n$ are linearly independent we must show that

$$c_1 v_1 + c_2 v_2 + \dots + c_n v_n = \mathbf{0} \quad (6.25)$$

has only the trivial solution.

To prove that the vectors $v_1, v_2, \dots, v_n$ span $\mathbb{R}^n$, we must show that every vector $\mathbf{b} = (b_1, b_2, \dots, b_n)$ in $\mathbb{R}^n$ can be expressed as

$$c_1 v_1 + c_2 v_2 + \dots + c_n v_n = \mathbf{b} \quad (6.26)$$

From the two equations above we will have two linear systems, the first is homogeneous (to prove for linear independence) and the second one is nonhomogeneous system (to prove that the vectors span), both systems will have the same coefficient matrix. The nonhomogeneous system is consistent for all values of $b_1, b_2, \dots, b_n$.

Theorem 6.13: Uniqueness of Basis Representation

If $S = \{v_1, v_2, \dots, v_n\}$ is a basis for a vector space V , then every vector v in V can be expressed in the form $v = c_1 v_1 + c_2 v_2 + \dots + c_n v_n$ in exactly one way.

Definition 6.19: Coordinate Vector

If $S = \{v_1, v_2, \dots, v_n\}$ is a basis for a vector space V (the order of the vectors in S remains fixed), and

$$v = c_1 v_1 + c_2 v_2 + \dots + c_n v_n$$

is the expression for a vector v in terms of the basis S , then the scalars $c_1, c_2, \dots, c_n$ are called the coordinates of v relative to the basis S . The vector $(c_1, c_2, \dots, c_n)$ in $\mathbb{R}^n$ constructed from these coordinates is called the coordinate vector of v relative to S ; it is denoted by

$$(v)_S = (c_1, c_2, \dots, c_n)$$

Theorem 6.14: Number of Vectors in a Basis

All bases for a finite-dimensional vector space have the same number of vectors.

Theorem 6.15: Number of Vectors in a Basis 2

Let V be a finite-dimensional vector space, and let $\{v_1, v_2, \dots, v_n\}$ be any basis

- (a) If a set has more than n vectors, then it is linearly dependent.
- (b) If a set has fewer than n vectors, then it does not span V .

Definition 6.20: Dimension

The dimension of a finite-dimensional vector space V is denoted by $\dim(V)$ and is defined to be the number of vectors in a basis for V . In addition, the zero vector space is defined to have dimension zero.

Engineers often use the term degrees of freedom as a synonym for dimension.

[SI*] Dimensions of some Vector Spaces:

$$\dim(\mathbb{R}^n) = n$$

$$\dim(P_n) = n + 1$$

$$\dim(M_{mn}) = m \times n$$

e.g. the standard basis for matrix 3×5 / M_{35} has 15 vectors.

Theorem 6.16: Plus/Minus Theorem

Let S be a nonempty set of vectors in a vector space V .

- (a) If S is a linearly independent set, and if v is a vector in V that is outside of $\text{span}(S)$, then the set $S \cup \{v\}$ that results by inserting v into S is still linearly independent.
- (b) If v is a vector in S that is expressible as a linear combination of other vectors in S , and if $S - \{v\}$ denotes the set obtained by removing v from S , then S and $S - \{v\}$ span the same space; that is,

$$\text{span}(S) = \text{span}(S - \{v\})$$

[SI*] In general, to show that a set of vectors $\{v_1, v_2, \dots, v_n\}$ is a basis for a vector space V , we must show that the vectors are linearly independent and span V . If we happen to know that V has dimension n , so that $\{v_1, v_2, \dots, v_n\}$ contains the right number of vectors for a basis, then it suffices to check either linear independence or spanning, the remaining condition will hold automatically.

Theorem 6.17: Basis and Dimension

Let V be an n -dimensional vector space, and let S be a set in V with exactly n vectors. Then S is a basis for V if and only if S spans V or S is linearly independent.

Theorem 6.18: Basis Rules

Let S be a finite set of vectors in a finite-dimensional vector space V .

- (a) If S spans V but is not a basis for V , then S can be reduced to a basis for V by removing appropriate vectors from S .
- (b) If S is linearly independent set that is not already a basis for V , then S can be enlarged to a basis for V by inserting appropriate vectors in S .

Theorem 6.19: Dimension of a Vector Space and Its Subspaces

If W is a subspace of a finite-dimensional vector space V , then:

- (a) W is finite-dimensional.
- (b) $\dim(W) \leq \dim(V)$.
- (c) $W = V$ if and only if $\dim(W) = \dim(V)$.

[SI*] A basis is the vector space generalization of a coordinate system, changing bases is akin to changing coordinate axes in $\mathbb{R}^2$ and $\mathbb{R}^3$.

[SI*] If $S = \{v_1, v_2, \dots, v_n\}$ is a basis for a finite-dimensional vector space V , and if

$$(v)_S = (c_1, c_2, \dots, c_n)$$

is the coordinate vector of v relative to S , then the mapping

$$v \rightarrow (v)_S \tag{6.27}$$

creates a connection (a one-to-one correspondence) between vectors in the general vector space V and vectors in the familiar vector space $\mathbb{R}^n$. We call $v \rightarrow (v)_S$ the coordinate map from V to $\mathbb{R}^n$.

[SI*] We will find it convenient to express coordinate vectors in the matrix form

$$[v]_S = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \tag{6.28}$$

[SI*] If we change the basis for a vector space V from an old basis $B = \{u_1, u_2, \dots, u_n\}$ to a new basis $B' = \{u'_1, u'_2, \dots, u'_n\}$, then for each vector v in V , the old coordinate vector $[v]_B$ is related to the new coordinate vector $[v]_{B'}$ by the equation

$$[v]_B = P_{B' \rightarrow B} [v]_{B'} \tag{6.29}$$

where the columns of P are the coordinate vectors of the new basis vectors relative to the old basis; that is, the column vectors of P are

$$[u'_1]_B, [u'_2]_B, \dots, [u'_n]_B \tag{6.30}$$

[SI*] The matrix P is called the transition matrix from B' to B . For emphasis, we will often denote it by

$$P_{B' \rightarrow B}$$

This matrix can be expressed in terms of its column vectors as

$$P_{B' \rightarrow B} = [[u'_1]_B \mid [u'_2]_B \mid \dots \mid [u'_n]_B] \tag{6.31}$$

A column vector $[u'_1]_B$ is a column vector u'_1 represented in old bases $B = \{u_1, u_2, \dots, u_n\}$ terms.

Similarly, the transition matrix from B to B' can be expressed in terms of its column vectors as

$$P_{B \rightarrow B'} = [[u_1]_{B'} \mid [u_2]_{B'} \mid \dots \mid [u_n]_{B'}] \tag{6.32}$$

A column vector $[u_1]_{B'}$ is a column vector u_1 represented in new bases $B' = \{u'_1, u'_2, \dots, u'_n\}$ terms.

[SI*] Consider this old bases in $\mathbb{R}^n$

$$B = \{u_1, u_2, \dots, u_n\}$$

and the new bases in $\mathbb{R}^n$

$$B' = \{u'_1, u'_2, \dots, u'_n\}$$

We can use transition matrix $P_{B \rightarrow B'}$ to obtain old bases from multiplying $P_{B \rightarrow B'}$ with new bases

$$\begin{aligned} P_{B \rightarrow B'}[\mathbf{u}'_1] &= [\mathbf{u}_1] \\ P_{B \rightarrow B'}[\mathbf{u}'_2] &= [\mathbf{u}_2] \\ &\vdots \\ P_{B \rightarrow B'}[\mathbf{u}'_n] &= [\mathbf{u}_n] \end{aligned}$$

in converse, we can use transition matrix $P_{B' \rightarrow B}$ to obtain new bases from multiplying $P_{B' \rightarrow B}$ with old bases

$$\begin{aligned} P_{B' \rightarrow B}[\mathbf{u}_1] &= [\mathbf{u}'_1] \\ P_{B' \rightarrow B}[\mathbf{u}_2] &= [\mathbf{u}'_2] \\ &\vdots \\ P_{B' \rightarrow B}[\mathbf{u}_n] &= [\mathbf{u}'_n] \end{aligned}$$

[SI*] The columns of the transition matrix from an old basis to a new basis are the coordinate vectors of the old basis relative to the new basis.

[SI*] Let $\mathbf{v}$ or $[\mathbf{v}]_B$ be any vector in V , the changing of bases will change the vector $\mathbf{v}$ as well. We know that vector $\mathbf{v}$ can be represented in old bases $B = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ as

$$\mathbf{v} = k_1\mathbf{u}_1 + k_2\mathbf{u}_2 + \dots + k_n\mathbf{u}_n$$

this vector $\mathbf{v}$ will change when we use new bases $B' = \{\mathbf{u}'_1, \mathbf{u}'_2, \dots, \mathbf{u}'_n\}$ into $\mathbf{v}'$ or $[\mathbf{v}]_{B'}$

$$\mathbf{v}' = k'_1\mathbf{u}'_1 + k'_2\mathbf{u}'_2 + \dots + k'_n\mathbf{u}'_n$$

we can write them in vector notation as

$$\mathbf{v} = [\mathbf{v}]_B = \begin{bmatrix} k_1 \\ k_2 \\ \vdots \\ k_n \end{bmatrix}, \quad \mathbf{v}' = [\mathbf{v}]_{B'} = \begin{bmatrix} k'_1 \\ k'_2 \\ \vdots \\ k'_n \end{bmatrix}$$

We are able to determine $[\mathbf{v}]_B$ and $[\mathbf{v}]_{B'}$ when we know the new and old bases by using the transition matrix

$$[\mathbf{v}]_B = P_{B' \rightarrow B}[\mathbf{v}]_{B'} \tag{6.33}$$

$$[\mathbf{v}]_{B'} = P_{B \rightarrow B'}[\mathbf{v}]_B \tag{6.34}$$

with

$$(P_{B' \rightarrow B})(P_{B \rightarrow B'}) = I_n \tag{6.35}$$

GlanzFreya' Guide 6.1: A procedure for computing $P_{B \rightarrow B'}$

1. Form the matrix

$$[B' \mid B]$$

with $B = \{u_1, u_2, \dots, u_n\}$ as old bases and $B' = \{u'_1, u'_2, \dots, u'_n\}$ as new bases. The bases will be treated as column vectors when constructing the matrix above.

2. Use elementary row operations, e.g. Gaussian elimination / Gauss-Jordan elimination to reduce the matrix to reduced row echelon form.

3. The resulting matrix will be

$$[I \mid P_{B \rightarrow B'}]$$

[SI*] Suppose now that B and B' are bases for a finite-dimensional vector space V . Since multiplication by $P_{B' \rightarrow B}$ maps coordinate vectors relative to the basis B' into coordinate vectors relative to a basis B , and $P_{B \rightarrow B'}$ maps coordinate vectors relative to B into coordinate vectors relative to B' , it follows that for every vector v in V we have

$$[v]_B = P_{B' \rightarrow B} [v]_{B'} \quad (6.36)$$

$$[v]_{B'} = P_{B \rightarrow B'} [v]_B \quad (6.37)$$

Theorem 6.20: Transition to Standard Basis for $\mathbb{R}^n$

Let $B' = \{u_1, u_2, \dots, u_n\}$ be any basis for the vector space $\mathbb{R}^n$ and let $S = \{e_1, e_2, \dots, e_n\}$ be the standard basis for $\mathbb{R}^n$. If the vectors in these bases are written in column form / column vector, then

$$P_{B' \rightarrow S} = [u_1 \mid u_2 \mid \dots \mid u_n] \quad (6.38)$$

[SI*] Changing basis allows you to convert a matrix from a complicated form to a simple form. It is often possible to represent a matrix in a basis where the only nonzero elements are on the diagonal, which is exceptionally simple. These diagonal elements will be the eigenvalues of the matrix. This is especially helpful in solving linear systems of differential equations. Often in physics, engineering, logistics, and probably lots of other places, you have a system of differential equations which all depend on each other. In order to solve the system directly, you would have to solve all equations at once, which is hard. We can use matrices to describe this system.

By changing basis, you may be able to make that matrix diagonal, which effectively separates the differential equations from each other, so you can solve just one at a time. This is comparatively easy. Real life situations that uses changing basis are

- (a) Quantum mechanics, solving the Schrödinger equation to describe the state of matter on the quantum level.
- (b) Electrical engineering, understanding the time-evolution of an electrical circuit.
- (c) Mechanical engineering, understanding the motion of a linear mechanical system, such as multiple spring-mass system.
- (d) Solid mechanics. Computing the principal stresses/strains in a material involves a change of basis. With the principal stresses, we can compute the maximum normal

stress and maximum shear stress the material experiences. And these maxima tell us whether the material will become damaged or not.

Changing basis allows you to convert a matrix from a complicated form to a simple form. Rotation is essentially a change of basis. If you ever employed a linear coordinate change, you changed bases. Let say you have a cube that is rotated arbitrarily and is not orthogonal the coordinate axes, and you're trying to determine if a point is inside of it. It is a bit tricky to determine directly. If you can change to a basis where the cube is orthogonal to the coordinate axes, determining if the point is inside, then it becomes trivial.

[SI*] Row space, column space, and null space are some important vector spaces that are associated with matrices. Understanding them will make us know the relationships between the solutions of a linear system and properties of its coefficient matrix.

Definition 6.21: Row and Column Vectors

For an $m \times n$ matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

the vectors

$$\begin{aligned} \mathbf{r}_1 &= [a_{11} \quad a_{12} \quad \dots \quad a_{1n}] \\ \mathbf{r}_2 &= [a_{21} \quad a_{22} \quad \dots \quad a_{2n}] \\ &\vdots \\ \mathbf{r}_m &= [a_{m1} \quad a_{m2} \quad \dots \quad a_{mn}] \end{aligned}$$

in $\mathbb{R}^n$ that are formed from the rows of A are called the row vectors of A , and the vectors

$$\mathbf{c}_1 = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix}, \quad \mathbf{c}_n = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

in $\mathbb{R}^m$ formed from the columns of A are called the column vectors of A .

Definition 6.22: Row Space and Column Space

If A is an $m \times n$ matrix, then the subspace of $\mathbb{R}^n$ spanned by the row vectors of A is called the row space of A .

The subspace of $\mathbb{R}^m$ spanned by the column vectors of A is called the column space of A .

The solution space of the homogeneous system of equations

$$Ax = 0$$

which is a subspace of $\mathbb{R}^n$, is called the null space of A .

[SI*] Suppose that

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

If $c_1, c_2, \dots, c_n$ denote the column vectors of A , then the product Ax can be expressed as a linear combination of these vectors with coefficients from x

$$Ax = x_1 c_1 + x_2 c_2 + \cdots + x_n c_n \quad (6.39)$$

Thus, a linear system, $Ax = b$, of m equations in n unknowns can be written as

$$x_1 c_1 + x_2 c_2 + \cdots + x_n c_n = b \quad (6.40)$$

from which we conclude that $Ax = b$ is consistent if and only if b is expressible as a linear combination of the column vectors of A .

Theorem 6.21: Column Space and Consistent Linear System

A system of linear equations $Ax = b$ is consistent if and only if b is in the column space of A .

Theorem 6.22: Null Space and Consistent Linear System

If x_0 is any solution of a consistent linear system $Ax = b$, and if $S = \{v_1, v_2, \dots, v_k\}$ is a basis for the null space of A , then every solution of $Ax = b$ can be expressed in the form

$$x = x_0 + c_1 v_1 + c_2 v_2 + \cdots + c_k v_k \quad (6.41)$$

Conversely, for all choices of scalars $c_1, c_2, \dots, c_k$, the vector x in this formula is a solution of $Ax = b$.

[SI*] The equation

$$x = x_0 + c_1 v_1 + c_2 v_2 + \cdots + c_k v_k$$

gives a formula for the general solution of $Ax = b$.

The vector $\mathbf{x}_0$ is called a particular solution of $A\mathbf{x} = \mathbf{b}$.

The remaining part of the formula $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_k\mathbf{v}_k$ is called the general solution of $A\mathbf{x} = \mathbf{0}$.

The general solution of a consistent linear system can be expressed as the sum of a particular solution of that system and the general solution of the corresponding homogeneous system.

Theorem 6.23: Elementary Row Operations for Null Space, Row Space, and Column Space

Elementary row operations do not change the null space of a matrix.

Elementary row operations do not change the row space of a matrix.

Elementary row operations change the column space of a matrix.

Theorem 6.24: Bases for the Row and Column Spaces

If a matrix R is in row echelon form, then the row vectors with the leading 1's (the nonzero row vectors) form a basis for the row space of R , and the column vectors with the leading 1's of the row vectors form a basis for the column space of R .

Theorem 6.25: Row Equivalent Matrices and Column Vectors

If A and B are row equivalent matrices, then:

- (a) A given set of column vectors of A is linearly independent if and only if the corresponding column vectors of B are linearly independent.
- (b) A given set of column vectors of A forms a basis for the column space of A if and only if the corresponding column vectors of B form a basis for the column space of B .

[SI*] Vector spaces are used extensively in electrical engineering, particularly in the analysis and design of electrical systems. Here are some examples of applications of vector spaces in electrical engineering:

- (a) Analysis of electric circuits: In electric circuits, currents and voltages can be represented as vectors in a vector space. By applying the principles of linear algebra to these vectors, it's possible to analyze the behavior of the circuit and determine the values of circuit variables such as voltage, current, and power.
- (b) Design of control systems: Control systems are used to regulate and stabilize the behavior of complex electrical systems. Vector spaces can be used to represent the state of a control system, and linear algebra can be used to design control algorithms that stabilize the system.
- (c) Signal processing: In signal processing, signals such as audio, video, and data can be represented as vectors in a vector space. By applying techniques from linear algebra, it's possible to analyze and process these signals, for example, by applying filters or compressing data.

- (d) Electromagnetics: Vector spaces are used to describe the behavior of electromagnetic fields, which are represented as vector fields. Maxwell's equations, which govern the behavior of electromagnetic fields, are expressed using vector calculus.

Theorem 6.26: Dimension of Row Space and Column Space

The row space and column space of a matrix A have the same dimension.

- [SI*] The number of nonzero rows of a reduced row echelon form matrix R is the same as the number of leading 1's in matrix R .

Definition 6.23: Rank and Nullity

The common dimension of the row space and column space of a matrix A is called the rank of A and is denoted by $\text{rank}(A)$.

The dimension of the null space of A is called the nullity of A and is denoted by $\text{nullity}(A)$.

- [SI*] Since the rank of a matrix A of size $m \times n$ is the common dimension of its row and column space, it follows that the rank is at most the smaller of m and n .

$$\text{rank}(A) \leq \min(m, n)$$

Theorem 6.27: Dimension Theorem for Matrices

If A is a matrix with n columns, then

$$\text{rank}(A) + \text{nullity}(A) = n$$

- [SI*] The number of leading variables is the same as the number of leading 1's in the reduced row echelon form of matrix A , which is the rank of A .

The number of free variables is the same as the number of parameters in the general solution of

$$Ax = 0$$

which is the nullity of A .

Theorem 6.28: Rank and Nullity

If A is an $m \times n$ matrix, then

- (a) $\text{rank}(A)$ = the number of leading variables in the general solution of $Ax = 0$.
- (b) $\text{nullity}(A)$ = the number of parameters in the general solution of $Ax = 0$.

- [SI*] In many applications the equations in a linear system correspond to physical constraints or conditions that must be satisfied. In general, the most desirable systems are those that have the same number of constraints as unknowns, since such systems often have a unique solution.

Unfortunately, it is not always possible to match the number of constraints and unknowns.

When linear systems have more constraints than unknowns like this

$$\begin{array}{rrrr} x_1 & + & x_2 & + & x_3 & = & b_1 \\ x_1 & + & 2x_2 & + & 5x_3 & = & b_2 \\ -x_1 & + & 6x_2 & + & 7x_3 & = & b_3 \\ -x_1 & - & 3x_2 & + & x_3 & = & b_4 \end{array}$$

it is called overdetermined systems.

When linear systems have fewer constraints than unknowns like this

$$\begin{array}{rrrr} x_1 & + & x_2 & + & x_3 & = & b_1 \\ x_1 & + & 2x_2 & + & 5x_3 & = & b_2 \end{array}$$

it is called underdetermined systems.

Theorem 6.29: Consistent Linear System

If $Ax = b$ is a consistent linear system of m equations in n unknowns, and if A has rank r , then the general solution of the system contains $n - r$ parameters.

Theorem 6.30: Overdetermined and Underdetermined Systems

Let A be an $m \times n$ matrix.

(a) **Overdetermined Case**

If $m > n$, then the linear system $Ax = b$ is inconsistent for at least one vector b in $\mathbb{R}^n$.

(b) **Underdetermined Case**

If $m < n$, then for each vector b in $\mathbb{R}^n$ the linear system $Ax = b$ is either inconsistent or has infinitely many solutions.

[SI*] There are six important vector spaces associated with a matrix A and its transpose A^T :

1. Row space of A
2. Column space of A
3. Null space of A
4. Row space of A^T
5. Column space of A^T
6. Null space of A^T

Transposing a matrix converts row vectors into column vectors and conversely, so except for a difference in notation, the row space of A^T is the same as the column space of A , and the column space of A^T is the same as the row space of A .

Out of the six spaces listed above, only the following four are distinct:

1. Row space of A

2. Column space of A
3. Null space of A
4. Null space of A^T

These are called the fundamental spaces of a matrix A .

Theorem 6.31: Rank for Matrix and Its Transpose

If A is any matrix, then

$$\text{rank}(A) = \text{rank}(A^T)$$

[SI*] If A is an $m \times n$ matrix, then

$$\text{rank}(A) + \text{nullity}(A) = m \quad (6.42)$$

It is possible to express the dimensions of all four fundamental spaces in terms of size and rank of A .

$$\begin{aligned} \dim[\text{row}(A)] &= r \\ \dim[\text{col}(A)] &= r \\ \dim[\text{null}(A)] &= n - r \\ \dim[\text{null}(A^T)] &= m - r \end{aligned} \quad (6.43)$$

The four formulas above provide an algebraic relationship between the size of a matrix and the dimension of its fundamental spaces.

[SI*] If A is an $m \times n$ matrix, then the null space of A consists of those vectors that are orthogonal to each of the row vectors of A .

Definition 6.24: Orthogonal Complement

If W is a subspace of $\mathbb{R}^n$, then the set of all vectors in $\mathbb{R}^n$ that are orthogonal to every vector in W is called the orthogonal complement of W and is denoted by the symbol $W^\perp$.

Theorem 6.32: Orthogonal Complement Basic

If W is a subspace of $\mathbb{R}^n$, then:

- (a) $W^\perp$ is a subspace of $\mathbb{R}^n$.
- (b) The only vector common to W and $W^\perp$ is $\mathbf{0}$.
- (c) The orthogonal complement of $W^\perp$ is W .

Theorem 6.33: Orthogonal Complement for Matrix

If A is an $m \times n$ matrix, then:

- (a) The null space of A and the row space of A are orthogonal complement in $\mathbb{R}^n$.
- (b) The null space of A^T and the column space of A are orthogonal complements in $\mathbb{R}^m$.

[SI*] Digital data are commonly stored in matrix form, and many techniques for improving transmission speed use the rank of a matrix. Rank plays a role to measure the "redundancy" in a matrix in the sense that if A is an $m \times n$ matrix of rank k , then $n - k$ of the column vectors and $m - k$ of the row vectors can be expressed in terms of k linearly independent column or row vectors. The essential idea in many data compression schemes is to approximate the original data set by a data set with smaller rank that conveys nearly the same information, then eliminate redundant vectors in the approximating set.

i. General Vector Spaces Application in Engineering

[SI*] Row and Column spaces

Row and column spaces are fundamental in engineering for understanding system constraints, reachability, and data relationship, appearing in robotics (workspace), structural analysis (load distribution), signal processing (data filtering), and data science (dimensionality reduction), representing the span of possible outputs (column space) and system constraints / input dimensions (row space) for linear systems defined by matrices.

Column Space (Image Space): The set of all possible output vectors ($\mathbf{b}$) from a transformation $A\mathbf{x} = \mathbf{b}$, meaning the "reach" or achievable states/positions.

Row Space: The space spanned by the rows of a matrix, revealing inherent constraints or relationships within the system's equations.

Rank: The dimension of both the row and column spaces, indicating the effective number of independent variables or dimensions.

Applications in engineering:

1. Robotics & Kinematics

Workspace determination: The column space defines the reachable workspace for a robot's end-effector, as the final position is a linear combination (span) of arm segment movements.

2. Structural Analysis & Loads

Load paths: The row space helps understand how applied forces (loads) are distributed and constrained within a structure, determining if a structure can support given forces.

3. Data Analysis & Machine Learning

Dimensionality reduction: Identifying the rank of a data matrix helps reduce dimensions (Principal Component Analysis), keeping the most significant components (in the column/row space).

4. Signal Processing

System modeling: Matrices represent filters or transformations. The column space shows what signals can be produced, while the row space indicates filtering properties.

5. System of Linear Equations (Civil/Electrical Engineering)

Solvability: For $A\mathbf{x} = \mathbf{b}$, the system has solutions if $\mathbf{b}$ is in the column space of A (i.e., $\mathbf{b}$ can be formed by the columns of A).

Constraints: The row space defines the constraints that must be met for a solution to exist.

In essence, the column space answers "what can I do or produce?", while the row space answers "what are the inherent rules or limitations?" within a linear engineering model.

[SI*] **Null space**

The solution space of $Ax = 0$ is the null space (or kernel), representing all input vectors x that a transformation A maps to the zero vector, crucial in engineering for finding equilibrium states, system redundancies, identifying dependencies (like in structural analysis or circuits), and understanding stability, showing which inputs cause "no output" or collapse, fundamentally linked to rank and nullity in signal processing, control systems, and computer graphics.

For a matrix A , the null space, or kernel, $\text{null}(A)$ is the set of all vectors x such that $Ax = 0$. The zero vector $x = 0$ is always a trivial solution. If non-zero solutions exist, they form a subspace (a line, a plane, etc., through the origin) with infinite solutions, indicating dependency or redundancy in the system.

Applications in engineering:

1. **Structural Engineering (Statics)**

Finding a forces in a truss (e.g. $Ax = 0$) where A relates forces to equilibrium, x are unknown forces, and 0 is zero net force. Non-zero solutions (null space vectors) represent forces that can exist without causing external loads, indicating redundant members or mechanisms where forces can balance in multiple ways (like a slack cable).

2. **Electrical Engineering (Circuit Analysis)**

Nodal analysis often leads to $Ax = 0$, where x are node voltages, A is the admittance matrix. A non-trivial null space means there are floating nodes or independent loops (Kirchoff's Current Law) where voltages can exist without external sources.

3. **Control Systems & Signal Processing**

Analyzing system stability or observability ($Ax = 0$). A non-trivial null space indicates unobservable states or internal dynamics (like oscillations) that aren't detectable from the system's outputs.

4. **Computer Graphics & Robotics**

Finding transformations (e.g., in 3D rotations, $Ax = 0$ can define axes). In kinematics, it can represent null-space motion, allowing for movement (like arm articulation) that doesn't change the end-effector's position, useful for obstacle avoidance.

5. **Data Analysis & Machine Learning (Dimensionality Reduction)**

Principal Component Analysis (PCA) uses eigenvectors, related to $Ax = \lambda x$ (eigenvalue problem), but the null space helps find dimensions where data has zero variance, revealing intrinsic data structure.

ii. **Matrix Transformations from $\mathbb{R}^n$ to $\mathbb{R}^m$**

[SI*] Functions of the form

$$w = F(x)$$

has independent variable x as a vector in $\mathbb{R}^n$ and the dependent variable w as a vector in $\mathbb{R}^m$. The special class of such functions are called matrix transformations. Such transformations are fundamental in the study of linear algebra and have important application in physics, computer graphics, engineering, social sciences, and various branches of mathematics.

Definition 6.25: Transformation

If V and W are vector spaces, and if f is a function with domain V and codomain W , then we say that f is a transformation from V to W or that f maps V to W , which we denote by writing

$$f : V \rightarrow W$$

In the special case where $V = W$, the transformation is also called an operator on V .

[SI*] Suppose that $f_1, f_2, \dots, f_m$ are real-valued functions of n variables

$$\begin{aligned} w_1 &= f_1(x_1, x_2, \dots, x_n) \\ w_2 &= f_2(x_1, x_2, \dots, x_n) \\ &\vdots \\ w_m &= f_m(x_1, x_2, \dots, x_n) \end{aligned} \tag{6.44}$$

These m equations assign a unique point $(w_1, w_2, \dots, w_m)$ in $\mathbb{R}^m$ to each point $(x_1, x_2, \dots, x_n)$ in $\mathbb{R}^n$ and thus define a transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$.

If we denote this transformation by T , then

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

and

$$T(x_1, x_2, \dots, x_n) = (w_1, w_2, \dots, w_m)$$

[SI*] In the special case where the equations are linear and can be expressed in the form

$$\begin{aligned} w_1 &= a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n \\ w_2 &= a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n \\ &\vdots \\ w_m &= a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n \end{aligned} \tag{6.45}$$

which we can write in matrix notation as

$$\begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_m \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \tag{6.46}$$

or more briefly as

$$w = Ax \tag{6.47}$$

Although we could view this as a linear system, we will view it instead as a transformation that maps the column vector x in $\mathbb{R}^n$ into the column vector w in $\mathbb{R}^m$.

We call this a matrix transformation (or matrix operator if $m = n$), and we denote it by

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

thus

$$w = T_A(x)$$

The matrix transformation T_A is called multiplication by A , and the matrix A is called the standard matrix for the transformation.

On occasion, we also use the notation

$$x \vec{T}_A w \quad (6.48)$$

which is read " T_A maps x into w ."

Theorem 6.34: Properties of Matrix Transformations

For every matrix A the matrix transformation $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ has the following properties for all vectors u and v in $\mathbb{R}^n$ and for every scalar k :

- (a) $T_A(0) = 0$
- (b) $T_A(ku) = kT_A(u)$ [Homogeneity property]
- (c) $T_A(u + v) = T_A(u) + T_A(v)$ [Additivity property]
- (d) $T_A(u - v) = T_A(u) - T_A(v)$

[SI*] A matrix transformation maps linear combinations of vectors in $\mathbb{R}^n$ into the corresponding linear combinations in $\mathbb{R}^m$

$$T_A(k_1u_1 + k_2u_2 + \cdots + k_ru_r) = k_1T_A(u_1) + k_2T_A(u_2) + \cdots + k_rT_A(u_r) \quad (6.49)$$

Depending on whether n -tuples and m -tuples are regarded as vectors or points, the geometric effect of a matrix transformation

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

is to map each vector (point) in $\mathbb{R}^n$ into a vector (point) in $\mathbb{R}^m$.

Theorem 6.35: Same Matrix Transformations

If $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $T_B : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are matrix transformations, and if

$$T_A(x) = T_B(x)$$

for every vector x in $\mathbb{R}^n$, then $A = B$.

[SI*] For any matrix A with size of $m \times n$ and standard basis vectors $e_1, e_2, \dots, e_n$ for $\mathbb{R}^n$

$$Ae_j, \quad j = 1, 2, \dots, n$$

will result as the j -th column of A since every entry of e_j is 0 except for the j -th, which is 1.

[SI*] If 0 is the $m \times n$ matrix, then

$$T_0(x) = 0x = 0$$

so multiplication by zero maps every vector in $\mathbb{R}^n$ into the zero vector in $\mathbb{R}^m$. We call T_0 the zero transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$.

[SI*] If I is the $n \times n$ identity matrix, then

$$T_I(x) = Ix = x$$

so multiplication by I maps every vector in $\mathbb{R}^n$ into itself. We call T_I the identity operator on $\mathbb{R}^n$.

[SI*] To find the standard matrix for a matrix transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$ by considering the effect of that transformation on the standard basis vectors for $\mathbb{R}^n$.

Suppose that A is unknown and that

$$e_1, e_2, \dots, e_n$$

are the standard basis vectors for $\mathbb{R}^n$. Suppose also that the images of these vectors under the transformation T_A are

$$T_A(e_1) = Ae_1, \quad T_A(e_2) = Ae_2, \dots, \quad T_A(e_n) = Ae_n$$

Ae_j is a linear combination of the columns of A in which the successive coefficients are the entries of e_j , so the product of Ae_j is just the j -th column of the matrix A . Thus,

$$A = [T_A(e_1) \mid T_A(e_2) \mid \dots \mid T_A(e_n)] \quad (6.50)$$

is the standard matrix for a matrix transformation.

[SI*] **Reflection Operators**

Some of the most basic matrix operators on $\mathbb{R}^2$ and $\mathbb{R}^3$ are those that map each point into its symmetric image about a fixed line or a fixed plane; these are called reflection operators.

[SI*] **Projection Operators**

Matrix operators on $\mathbb{R}^2$ and $\mathbb{R}^3$ that map each point into its orthogonal projection on a fixed line or plane are called projection operators, or more precisely, orthogonal projection operators.

[SI*] **Rotation Operators**

Matrix operators on $\mathbb{R}^2$ and $\mathbb{R}^3$ that move points along circular arcs called rotation operators.

[SI*] **Dilations and Contractions**

If k is a nonnegative scalar, then the operator

$$T(x) = kx$$

on $\mathbb{R}^2$ or $\mathbb{R}^3$ has the effect of increasing or decreasing the length of each vector by a factor of k .

1. If $0 \leq k \leq 1$ the operator is called a contraction with factor k .
2. If $k > 1$ it is called a dilation with factor k .
3. If $k = 1$, then T is the identity operator and can be regarded either as a contraction or a dilation.

[SI*] **Expansion and Compressions**

In a dilation or contraction of $\mathbb{R}^2$ or $\mathbb{R}^3$, all coordinates are multiplied by a factor k . If only one of the coordinates is multiplied by k , then the resulting operator is called an expansion or compression.

[SI*] **Shears**

A matrix operator of the form $T(x, y) = (x + ky, y)$ translates a point (x, y) in the xy -plane parallel to the x -axis by an amount ky that is proportional to the y -coordinate of the point.

[SI*] Now, when we see a square matrix of size 2×2 or 3×3 , if it is in reduced row echelon form and fit the standard matrix we can describe it as matrix operator. For example

$$A_1 = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$$

A_1 is a matrix corresponds to a shear in the x -direction with factor 3.

$$A_2 = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}$$

A_2 is a matrix corresponds to a shear in the x -direction with factor 2, and the y -direction with factor -2 .

$$A_3 = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}$$

A_3 is a matrix corresponds to an expansion in the x -direction with factor 3.

[SI*] Suppose that T_A is a matrix transformation from $\mathbb{R}^n$ to $\mathbb{R}^k$ and T_B is a matrix transformation from $\mathbb{R}^k$ to $\mathbb{R}^m$. If x is a vector in $\mathbb{R}^n$, then T_A maps this vector into a vector $T_A(x)$ in $\mathbb{R}^k$, and T_B maps that vector into the vector $T_B(T_A(x))$ in $\mathbb{R}^m$. This process creates a transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$ that we call the composition of T_B with T_A and denote by the symbol

$$T_B \circ T_A$$

which is read " T_B circle T_A ."

The transformation T_A is performed first

$$(T_B \circ T_A)(x) = T_B(T_A(x)) \quad (6.51)$$

this composition is itself a matrix transformation since

$$(T_B \circ T_A)(x) = T_B(T_A(x)) = B(T_A(x)) = B(Ax) = (BA)x$$

which shows that it is multiplication by BA

$$T_B \circ T_A = T_{BA} \quad (6.52)$$

[SI*] The standard matrix for a composition is the product of the standard matrices in the appropriate order.

[SI*] Composition not always commutative, when we let $T_A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the reflection about the line $y = x$, and let $T_B : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the orthogonal projection on the y -axis. $T_B \circ T_A$ and $T_A \circ T_B$ will have different effects on a vector x graphically.

$$T_B \circ T_A \neq T_A \circ T_B$$

For another case, let $T_A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the reflection about the y -axis, and let $T_B : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the reflection about the x -axis. In this case $T_B \circ T_A$ and $T_A \circ T_B$ are the same; both map every vector $x = (x, y)$ into its negative $-x = (-x, -y)$:

$$(T_A \circ T_B)(x, y) = T_A(x, -y) = (-x, -y)$$

$$(T_B \circ T_A)(x, y) = T_B(-x, y) = (-x, -y)$$

thus

$$T_B \circ T_A = T_A \circ T_B$$

the standard matrices for T_A and T_B also commute

$$[T_B \circ T_A] = [T_A \circ T_B] = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

The operator $T(\mathbf{x}) = -\mathbf{x}$ on $\mathbb{R}^2$ or $\mathbb{R}^3$ is called the reflection about the origin.

Definition 6.26: One-to-One Matrix Transformations

A matrix transformation $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is said to be one-to-one if T_A maps distinct vectors (points) in $\mathbb{R}^n$ into distinct vectors (points) in $\mathbb{R}^m$.

[SI*] Rotation operators on $\mathbb{R}^2$ are one-to-one since distinct vectors that are rotated through the same angle have distinct images.

The orthogonal projection of $\mathbb{R}^3$ on the xy plane is not one-to-one, because if we map a vertical line, the whole points in that vector / vertical line will have the same image, remember that the image of (x, y, z) for orthogonal projection on xy plane is $(x, y, 0)$.

Theorem 6.36: One-to-One Matrix Transformations

If A is an $n \times n$ matrix and $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is the corresponding matrix operator, then the following statements are equivalent.

- (a) A is invertible.
- (b) The range of T_A is in $\mathbb{R}^n$.
- (c) T_A is one-to-one.

[SI*] If $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a one-to-one matrix operator, and if $T_{A^{-1}} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is its inverse, then the standard matrices for these operators are related by the equation

$$T_{A^{-1}} = T_A^{-1} \quad (6.53)$$

[SI*] **Standard Matrix for T^{-1}**

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the operator that rotates each vector in $\mathbb{R}^2$ through the angle θ , so the standard matrix is

$$[T] = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

it is evident geometrically that to undo the effect of T , one must rotate each vector in $\mathbb{R}^2$ through the angle $-\theta$. But this is exactly what the operator T^{-1} does, since the standard matrix for T^{-1} is

$$[T^{-1}] = [T]^{-1} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} \cos(-\theta) & -\sin(-\theta) \\ \sin(-\theta) & \cos(-\theta) \end{bmatrix}$$

which is the standard matrix for a rotation through the angle $-\theta$.

[SI*] If $f_1, f_2, \dots, f_m$ are any functions of the n variables $x_1, x_2, \dots, x_n$, then the equations

$$\begin{aligned} w_1 &= f_1(x_1, x_2, \dots, x_n) \\ w_2 &= f_2(x_1, x_2, \dots, x_n) \\ &\vdots \\ w_m &= f_m(x_1, x_2, \dots, x_n) \end{aligned}$$

define a transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ that maps the vector $x = (x_1, x_2, \dots, x_n)$ into the vector $(w_1, w_2, \dots, w_m)$. But it is only the case where these equations are linear that T is a matrix transformation.

Theorem 6.37: Algebraic Properties of Transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation if and only if the following relationships hold for all vectors u and v in $\mathbb{R}^n$ and for every scalar k :

- (a) $T(u + v) = T(u) + T(v)$ (additivity property)
- (b) $T(ku) = kT(u)$ (homogeneity property)

Theorem 6.38: Linear Transformation and Matrix Transformation

Every linear transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$ is a matrix transformation, and conversely, every matrix transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$ is a linear transformation.

[SI*] The inverse of reflection in $\mathbb{R}^2$ and $\mathbb{R}^3$ is the reflection itself toward the same axis / plane. The inverse of dilation is contraction. The inverse of rotation through an angle θ is the rotation through an angle $-\theta$ about the same axis (in $\mathbb{R}^2$ or $\mathbb{R}^3$).

[SI*] By inspection

1. The orthogonal projection on the x -axis in $\mathbb{R}^2$ is not one-to-one because the second row is all 0's, so the determinant is 0.
2. The reflection about the y -axis in $\mathbb{R}^2$ is one-to-one because the determinant is -1 .
3. The reflection about the line $y = x$ in $\mathbb{R}^2$ is one-to-one because the determinant is -1 .
4. The contraction with factor $k > 0$ in $\mathbb{R}^2$ is one-to-one because the determinant is $k^2 \neq 0$.
5. The rotation about the positive z -axis in $\mathbb{R}^3$ is one-to-one because the determinant is 1.
6. The reflection about the xy -plane in $\mathbb{R}^3$ is one-to-one because the determinant is -1 .
7. The dilation with factor $k > 0$ in $\mathbb{R}^3$ is one-to-one because the determinant is $k^3 \neq 0$.

If today we can see the application of matrix transformation in photo editing with computer, in the future the automatic matrix transformation in $\mathbb{R}^3$ will be able to create a better and faster optimal engine, machine, computer' hardware, by resizing it to make it smaller but still with great proportion in a system as a whole, or to create a waste plant that convert trashes to electricity that need compact engine, smaller but able to do bigger job, that perhaps can take 10 million tons of trashes a day with a centralized machine as small as 24 inch CRT TV that able to control the gigantic medium to process the trashes.

iii. Dynamical Systems and Markov Chains

- [SI*] A dynamical system is a finite set of variables whose values change with time. The value of a variable at a point in time is called the state of the variable at that time, and the vector formed from these states is called the state of the dynamical system at that time.
- [SI*] In many dynamical systems the states of the variables are not known with certainty but can be expressed as probabilities; such dynamical systems are called stochastic processes. If an experiment or observation has n possible outcomes, then the probabilities of those outcomes must be nonnegative fractions whose sum is 1.

In a stochastic process with n possible states, the state vector at each time t has the form

$$\mathbf{x}(t) = \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_n(t) \end{bmatrix}$$

with $x_i(t)$ is the probability that the system is in state i with $i = 1, 2, \dots, n$. The entries in this vector must add up to 1 since they account for all n possibilities. In general, a vector with nonnegative entries that add up to 1 is called a probability vector.

- [SI*] A square matrix, each of whose columns is a probability vector, is called a stochastic matrix. Such matrices commonly occur in formulas that relate successive states of a stochastic process.

For example, the state vectors $\mathbf{x}(k+1)$ and $\mathbf{x}(k)$ are related by an equation of the form

$$\mathbf{x}(k+1) = P\mathbf{x}(k) \quad (6.54)$$

in which

$$\begin{bmatrix} P_A & (1 - P_B) \\ (1 - P_A) & P_B \end{bmatrix}$$

is a stochastic matrix with:

P_A is the probability that event A will occur again at time $k+1$ after it occurs at time k .
 $1 - P_A$ is the probability that event A will not occur at time $k+1$ (event B occurs at $k+1$) after event A occurs at time k .

P_B is the probability that event B will occur again at time $k+1$ after it occurs at time k .
 $1 - P_B$ is the probability that event B will not occur at time $k+1$ (event A occurs at $k+1$) after event B occurs at time k .

Definition 6.27: Markov Chain

A Markov chain is a dynamical system whose state vectors at a succession of time interval are probability vectors and for which the state vectors at successive time intervals are related by an equation of the form

$$\mathbf{x}(k+1) = P\mathbf{x}(k)$$

in which $P = [p_{ij}]$ is a stochastic matrix and p_{ij} is the probability that the system will be in state i at time $t = k+1$ if it is in state j at time $t = k$. The matrix P is called the transition matrix for the system.

[SI*] In a Markov chain with an initial state of $x(0)$, the successive state vectors are

$$x(1) = Px(0), \quad x(2) = Px(1), \quad x(3) = Px(2), \quad x(4) = Px(3), \dots$$

For brevity, it is common to denote $x(k) = x_k$, which allows us to write the successive state vectors more briefly as

$$x_1 = Px_0, \quad x_2 = Px_1, \quad x_3 = Px_2, \quad x_4 = Px_3, \dots \quad (6.55)$$

Alternatively, these state vectors can be expressed in terms of the initial state vector x_0 as

$$\begin{aligned} x_1 &= Px_0 \\ x_2 &= P(Px_0) = P^2x_0 \\ x_3 &= P(P^2x_0) = P^3x_0 \\ x_4 &= P(P^3x_0) = P^4x_0 \end{aligned}$$

from which it follows that

$$x_k = P^k x_0 \quad (6.56)$$

[SI*] The matrix

$$P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

is stochastic and hence can be regarded as the transition matrix for a Markov chain. A real life example for this transition matrix is the changing of day and night. It will always happens like that with probability 1, after day we will have night, and vice versa. A simple calculation shows that

$$P^2 = I$$

from which it follows that

$$I = P^2 = P^4 = P^6 = \dots$$

and

$$P = P^3 = P^5 = P^7 = \dots$$

Thus, the successive states in the Markov chain with initial vector x_0 are

$$x_0, Px_0, x_0, Px_0, x_0, \dots$$

which oscillate between x_0 and Px_0 . Thus, the Markov chain does not stabilize unless both components of x_0 are $\frac{1}{2}$.

Definition 6.28: Regular Markov Chain

A stochastic matrix P is said to be regular if P or some positive power of P has all positive entries, and a Markov chain whose transition matrix is regular is said to be a regular Markov chain.

Theorem 6.39: Long-term Behavior of Markov Chains: Steady-state Vector

If P is the transition matrix for a regular Markov chain, then:

- (a) There is a unique probability vector q such that $Pq = q$.
- (b) For any initial probability vector x_0 , the sequence of the state vectors

$$x_0, Px_0, \dots, P^k x_0$$

converges to q .

The vector q is called the steady-state vector of the Markov chain. It can be found by rewriting the equation as

$$(I - P)q = 0$$

and then solving this equation for q subject to the requirement that q be a probability vector.

[SI*] Example:

Suppose that at some initial point in time 100,000 people live in a certain city and 25,000 people live in a village on the mountain. The Regional Planning Commission determines that each year 10% of the city population moves to the village for retirement or choose a healthier and more meaningful lifestyle and 3% of the village population moves to the city for study or work purpose.

- (a) Assuming that the total population remains constant, make a table that shows the populations of the city and the village over a five-year period (round to the nearest integer).
- (b) Over the long term, how will the population be distributed between the city and the village?

Solution:

- (a) Let state 1 be living in the city, and state 2 be living in the village on a mountain. Then the transition matrix is

$$P = \begin{bmatrix} 0.9 & 0.03 \\ 0.1 & 0.97 \end{bmatrix}$$

the initial state vector is

$$x_0 = \begin{bmatrix} 100,000 \\ 25,000 \end{bmatrix}$$

The populations over five-year period will be

$$Px_0 = \begin{bmatrix} 90,750 \\ 34,250 \end{bmatrix}, \quad P^2x_0 = \begin{bmatrix} 82,702 \\ 42,298 \end{bmatrix}, \quad P^3x_0 = \begin{bmatrix} 75,701 \\ 49,299 \end{bmatrix}$$

$$P^4x_0 = \begin{bmatrix} 69,610 \\ 55,390 \end{bmatrix}, \quad P^5x_0 = \begin{bmatrix} 64,311 \\ 60,689 \end{bmatrix}$$

- (b) The system

$$(I - P)q = 0$$

Year	1	2	3	4	5
City	90,750	82,702	75,701	69,610	64,311
Village	34,250	42,298	49,299	55,390	60,689

Table 6.1: The populations of the city and the village over a five-year period.

$$\begin{bmatrix} 0.1 & -0.03 \\ -0.1 & 0.03 \end{bmatrix} \begin{bmatrix} q_1 \\ q_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The general solution is

$$q_1 = \frac{3}{10}s, \quad q_2 = s$$

since

$$q_1 + q_2 = 1$$

we will have

$$\begin{aligned} \frac{3}{10}s + s &= 1 \\ \frac{13}{10}s &= 1 \\ s &= \frac{10}{13} = 0.7692308 \end{aligned}$$

and

$$q_1 = \frac{3}{13} = 0.23076922$$

the steady-state vector is

$$\mathbf{q} = \begin{bmatrix} \frac{3}{13} \\ \frac{10}{13} \end{bmatrix} = \begin{bmatrix} 0.23076922 \\ 0.7692308 \end{bmatrix}$$

Over the long term, the population of 125,000 will be distributed with $\frac{3}{13}(125,000) = 28,846$ in the city and $\frac{10}{13}(125,000) = 96,154$ in the village.

[SI*] Example:

Physical traits are determined by the genes that an offspring receives from its parents. In the simplest case a trait in the offspring is determined by one pair of genes, one member of the pair inherited from the male parent and the other from the female parent. Typically, each gene in a pair can assume one of two forms, called alleles, denoted by A and a . This leads to three possible pairings

$$AA, \quad Aa, \quad aa$$

called genotypes (the pairs Aa and aA determine the same trait and hence are not distinguished from one another). It is shown in the study of heredity that if a parent of known genotype is crossed with a random parent of unknown genotype probabilities given in the following table, which can be viewed as a transition matrix for a Markov process: Thus, for example, the offspring of a parent of genotype AA that is crossed at random with a parent of unknown genotype will have a 50% chance of being AA , a 50% chance of being Aa , and no chance of being aa .

	AA	Aa	aa
AA	$\frac{1}{2}$	$\frac{1}{4}$	0
Aa	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
aa	0	$\frac{1}{4}$	$\frac{1}{2}$

Table 6.2: The row represents the genotype of offspring and the column represents the genotype of parent.

- (a) Show that the transition matrix is regular
 (b) Find the steady-state vector, and discuss its physical interpretation.

Solution:

- (a) Let state 1 be the genotype being AA , state 2 be the genotype being Aa , and state 3 be the genotype being aa . Then the transition matrix is

$$P = \begin{bmatrix} \frac{1}{2} & \frac{1}{4} & 0 \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ 0 & \frac{1}{4} & \frac{1}{2} \end{bmatrix}$$

the transition matrix is said to be regular if P or some positive power of P has all positive entries, thus

$$P^2 = \begin{bmatrix} \frac{3}{8} & \frac{1}{4} & \frac{1}{8} \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{8} & \frac{1}{4} & \frac{3}{8} \end{bmatrix}$$

since P^2 has positive entries, then the transition matrix P is regular.

- (b) Now to find the steady-state vector $\mathbf{q}$ we will use the formula

$$(I - P)\mathbf{q} = \mathbf{0}$$

then solving this equation for $\mathbf{q}$ we will have

$$\begin{bmatrix} \frac{1}{2} & -\frac{1}{4} & 0 \\ -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \\ 0 & -\frac{1}{4} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} q_1 \\ q_2 \\ q_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

the general solution of this system is

$$q_1 = s, \quad q_2 = 2s, \quad q_3 = s$$

for $\mathbf{q}$ to be a probability vector, we must have

$$\begin{aligned} q_1 + q_2 + q_3 &= 1 \\ s + 2s + s &= 1 \\ 4s &= 1 \\ s &= \frac{1}{4} \end{aligned}$$

the steady-state vector q is

$$q = \begin{bmatrix} q_1 \\ q_2 \\ q_3 \end{bmatrix} = \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ \frac{1}{4} \end{bmatrix}$$

over long-term, there is a 50% chance the offspring will have genotype Aa , there will be more offspring with genotype being Aa compared to genotype being AA or aa .

iv. Functions in SymIntegration Related to General Vector Spaces

[SI*] We will list the basic functions that can be used related to general vector spaces:

spanningtest(vector<vector<double>> &matrix)

to compute the determinant of a matrix (composed by column vectors), and if the determinant is not equal to 0 then it spans $\mathbb{R}^n$, with n as the dimension of the matrix.

linearindependencetest(vector<vector<double>> &matrix)

to determine whether the column vectors from the coefficient matrix are linearly independent or linearly dependent in $\mathbb{R}^n$, with n is the length / size of the column vectors.

basistest(vector<vector<double>> &matrix)

to determine whether the column vectors from the coefficient matrix are the basis in $\mathbb{R}^n$, with n is the length / size of the column vectors. Basis means that the column vectors span $\mathbb{R}^n$ and are linearly independent.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Test for Basis ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Test for Basis ]# ./main

Matrix:
      1.000000      4.000000      5.000000
      2.000000      9.000000      8.000000
      2.000000      9.000000      9.000000
     -1.000000     -4.000000     -5.000000

The column vectors span R^{4}

Augmented Matrix:
      1.000000      4.000000      5.000000      0.000000      0.000000
      2.000000      9.000000      8.000000      0.000000      0.000000
      2.000000      9.000000      9.000000      0.000000      0.000000
     -1.000000     -4.000000     -5.000000      0.000000      0.000000

No unique solution or infinite solutions exist.

The column vectors are the basis in R^{4}

Time taken by function: 1454 microseconds
```

Figure 6.11: The basis test for coefficient matrix A that is composed of column vectors v_1, v_2 and v_3 (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Test for Basis/main.cpp*).

coordinatevector(vector<vector<double>> &A, vector<vector<double>> &b)

to compute the coordinate vector of v relative to the basis $S = \{v_1, v_2, \dots, v_n\}$. Vector v will be the input for b and the basis S is consisting of column vectors that will be the input for matrix A .

basistransition_withcoordinatevector(vector<vector<double>> &oldbasis, vector<vector<double>> &newbasis, vector<double> &w)

to compute the transition matrix from B' to B , the transition matrix from B to B' , the coordinate vector $[w]_B$, and $[w]_{B'}$.

rowspacebasis(vector<vector<double>> &A)

to obtain the basis for the row space, with A as the matrix, it can be non square matrix as well.

columnspacebasis(vector<vector<double>> &A)

to obtain the basis for the column space, with A as the matrix, it can be non square matrix as well.

homogeneouslinearsystembasis(vector<vector<double>> &A)

to obtain the basis for the solution space of the linear system $Ax = 0$, with A as the matrix, it can be non square matrix as well.

normalizehomogeneouslinearsystembasis_matrix(vector<vector<double>> &A)

to obtain the normalize basis for the solution space of the linear system $Ax = 0$, with A as the matrix, it can be non square matrix as well. The output is a matrix with row vector as the basis, you can then transpose it if you want the basis as the column vector.

vector<double> reflection_yaxis(vector<double> &x)

to compute the reflection of vector x about y -axis in $\mathbb{R}^2$.

vector<double> reflection_xaxis(vector<double> &x)

to compute the reflection of vector x about x -axis in $\mathbb{R}^2$.

vector<double> reflection_linex(vector<double> &x)

to compute the reflection of vector x about the line $y = x$ in $\mathbb{R}^2$.

vector<double> orthogonalprojection_xaxis(vector<double> &x)

to compute the orthogonal projection of vector x on the x -axis in $\mathbb{R}^2$.

vector<double> orthogonalprojection_yaxis(vector<double> &x)

to compute the orthogonal projection of vector x on the y -axis in $\mathbb{R}^2$.

vector<double> rotation_ccw(vector<double> &x, double θ)

to compute the image of vector x under a counterclockwise rotation of θ radians in $\mathbb{R}^2$.

vector<double> contractiondilation(vector<double> &x, double k)

to compute the contraction (if $0 \leq k < 1$) or dilation (if $k > 1$) of vector x with factor k in $\mathbb{R}^n$.

vector<double> compressionexpansion_xdirection(vector<double> &x, double k)

to compute the compression (if $0 \leq k < 1$) or expansion (if $k > 1$) of vector x in the x -direction with factor k in $\mathbb{R}^2$.

vector<double> compressionexpansion_ydirection(vector<double> &x, double k)

to compute the compression (if $0 \leq k < 1$) or expansion (if $k > 1$) of vector x in the y -direction with factor k in $\mathbb{R}^2$.

vector<double> shear_xdirection(vector<double> &x, double k)

to compute the shear of vector x in the x -direction with factor k in $\mathbb{R}^2$.

vector<double> shear_ydirection(vector<double> &x, double k)

to compute the shear of vector x in the y -direction with factor k in $\mathbb{R}^2$.

vector<double> reflection_xyplane(vector<double> &x)

to compute the reflection of vector x about xy -plane in $\mathbb{R}^3$.

vector<double> reflection_xzplane(vector<double> &x)

to compute the reflection of vector x about xz -plane in $\mathbb{R}^3$.

vector<double> reflection_yzplane(vector<double> &x)

to compute the reflection of vector x about yz -plane in $\mathbb{R}^3$.

vector<double> orthogonalprojection_xyplane(vector<double> &x)

to compute the orthogonal projection of vector x on the xy -plane in $\mathbb{R}^3$.

vector<double> orthogonalprojection_xzplane(vector<double> &x)

to compute the orthogonal projection of vector x on the xz -plane in $\mathbb{R}^3$.

vector<double> orthogonalprojection_yzplane(vector<double> &x)

to compute the orthogonal projection of vector x on the yz -plane in $\mathbb{R}^3$.

vector<double> rotation3d_xaxis_ccw(vector<double> &x, double θ)

to compute the the image of vector x under a counterclockwise rotation about the x -axis through an angle θ radians in $\mathbb{R}^3$.

vector<double> rotation3d_yaxis_ccw(vector<double> &x, double θ)

to compute the the image of vector x under a counterclockwise rotation about the y -axis through an angle θ radians in $\mathbb{R}^3$.

vector<double> rotation3d_zaxis_ccw(vector<double> &x, double θ)

to compute the the image of vector x under a counterclockwise rotation about the z -axis through an angle θ radians in $\mathbb{R}^3$.

vector<double> compressionexpansion3d_xdirection(vector<double> &x, double k)

to compute the compression (if $0 \leq k < 1$) or expansion (if $k > 1$) of vector x in the x -direction with factor k in $\mathbb{R}^3$.

vector<double> compressionexpansion3d_ydirection(vector<double> &x, double k)

to compute the compression (if $0 \leq k < 1$) or expansion (if $k > 1$) of vector x in the y -direction with factor k in $\mathbb{R}^3$.

vector<double> compressionexpansion3d_zdirection(vector<double> &x, double k)

to compute the compression (if $0 \leq k < 1$) or expansion (if $k > 1$) of vector x in the z -direction with factor k in $\mathbb{R}^3$.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrix Transformation ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrix Transformation ]# ./main

Vector x:
5
1

Reflection about the y-axis:
-5
1

Reflection about the x-axis:
5
-1

Reflection about the line y=x:
1
5

Orthogonal projection on the x-axis:
5
0

Orthogonal projection on the y-axis:
0
1

Rotation through an angle  $\pi/6$  :
3.83013
3.36603

Vector x in 3d:
5
1
7

Reflection about the xy-plane:
5
1
-7

Reflection about the xz-plane:
5
-1
7

Reflection about the yz-plane:
-5
1
7

```

Figure 6.12: The computation for matrix transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Matrix Transformation/main.cpp*).

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrix Transformation ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Matrix Transformation ]# ./main

Vector x:
5
1

Reflection about the y-axis:
-5
1

Reflection about the x-axis:
5
-1

Reflection about the line y=x:
1
5

Orthogonal projection on the x-axis:
5
0

Orthogonal projection on the y-axis:
0
1

Rotation through an angle  $\pi/6$  :
3.83013
3.36603

Vector x in 3d:
5
1
7

Reflection about the xy-plane:
5
1
-7

Reflection about the xz-plane:
5
-1
7

Reflection about the yz-plane:
-5
1
7

```

Figure 6.13: The computation for matrix transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Matrix Transformation/main.cpp*).

MarkovChain(vector<vector<double>>& P, vector<double>& x0, int k)

to compute state vector x_k with P as the transition matrix, x_0 as the initial state vector, and k is to represents the time $t = k$.

v. Determine Linear Independence in $\mathbb{R}^n$ with SymIntegration

Determine whether the vectors

$$v_1 = (1, 2, 2, -1)$$

$$v_2 = (4, 9, 9, -4)$$

$$v_3 = (5, 8, 9, -5)$$

in $\mathbb{R}^4$ are linearly dependent or linearly independent.

Solution:

The linear independence or linear dependence of these vectors is determined by whether there exist nontrivial solutions of the vector equation

$$k_1 v_1 + k_2 v_2 + k_3 v_3 = \mathbf{0}$$

or, equivalently, of

$$k_1(1, 2, 2, -1) + k_2(4, 9, 9, -4) + k_3(5, 8, 9, -5) = (0, 0, 0, 0)$$

Equating corresponding components on the two sides yields the homogeneous linear system

$$k_1 + 4k_2 + 5k_3 = 0$$

$$2k_1 + 9k_2 + 8k_3 = 0$$

$$2k_1 + 9k_2 + 9k_3 = 0$$

$$-k_1 - 4k_2 - 5k_3 = 0$$

This system has only the trivial solution

$$k_1 = 0, \quad k_2 = 0, \quad k_3 = 0$$

from which you can conclude that v_1, v_2 , and v_3 are linearly independent.

```
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Linear Independence ]# ./main
Error: Could not open file b.txt

Matrix:
      1.000000      4.000000      5.000000
      2.000000      9.000000      8.000000
      2.000000      9.000000      9.000000
     -1.000000     -4.000000     -5.000000
No unique solution or infinite solutions exist.

The column vectors are linearly independent.

Time taken by function: 1051 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Linear Independence ]#
```

Figure 6.14: The computation to determine whether the column vectors v_1, v_2 , and v_3 are linearly independent or linearly dependent in $\mathbb{R}^4$. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Orthogonal Projection/main.cpp*).

vi. Compute Coordinate Vector Relative to Basis $S = \{v_1, v_2, v_3\}$ with SymIntegration

The vectors

$$v_1 = (1, 2, 1)$$

$$v_2 = (2, 9, 0)$$

$$v_3 = (3, 3, 4)$$

form a basis for $\mathbb{R}^3$. Find the coordinate vector of $v = (5, -1, 9)$ relative to the basis $S = \{v_1, v_2, v_3\}$.

Solution:

To find $(v)_S$ we must first express v as a linear combination of the vectors in S ; that is, we must find values of c_1, c_2 , and c_3 such that

$$v = c_1 v_1 + c_2 v_2 + c_3 v_3$$

or, in terms of components,

$$(5, -1, 9) = c_1(1, 2, 1) + c_2(2, 9, 0) + c_3(3, 3, 4)$$

Equating corresponding components gives

$$c_1 + 2c_2 + 3c_3 = 5$$

$$2c_1 + 9c_2 + 3c_3 = -1$$

$$c_1 + 4c_3 = 9$$

Solving this system we obtain $c_1 = 1, c_2 = -1, c_3 = 2$. Therefore,

$$(v)_S = (1, -1, 2)$$

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Coordinate Vectors Relative to Standard Bases ]# ./main

Matrix:
      1.000000      2.000000      3.000000
      2.000000      9.000000      3.000000
      1.000000      0.000000      4.000000

(v)_{S}:
1.000000
-1.000000
2.000000

Time taken by function: 1099 microseconds
```

Figure 6.15: The computation to find the coordinate vector of $v = (5, -1, 9)$ relative to the basis $S = \{v_1, v_2, v_3\}$. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Coordinate Vectors Relative to Standard Bases/main.cpp*).

vii. Compute Transition Matrix ($P_{B' \rightarrow B}$) from Old Basis (B') to New Basis (B) and the Coordinate Vectors $[w]_B$ and $[w]_{B'}$ with SymIntegration

Consider the bases $B = [u_1, u_2]$ and $B' = [u'_1, u'_2]$ for $\mathbb{R}^2$ where

$$u_1 = \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \quad u_2 = \begin{bmatrix} 4 \\ -1 \end{bmatrix}$$

$$u'_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad u'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- (a) Find the transition matrix from B' to B (from old basis to new basis)
- (b) Find the transition matrix from B to B' (from new basis to old basis)
- (c) Compute the coordinate vector $[w]_B$ where

$$w = \begin{bmatrix} 3 \\ -5 \end{bmatrix}$$

then compute $[w]_{B'}$

Solution:

(a)

$$\left[\begin{array}{cc|cc} 2 & 4 & 1 & -1 \\ 2 & -1 & 3 & -1 \end{array} \right]$$

reduces to

$$\left[\begin{array}{cc|cc} 1 & 0 & \frac{13}{10} & -\frac{1}{2} \\ 0 & 1 & -\frac{2}{5} & 0 \end{array} \right]$$

so

$$P_{B' \rightarrow B} = \begin{bmatrix} \frac{13}{10} & -\frac{1}{2} \\ -\frac{2}{5} & 0 \end{bmatrix}$$

(b)

$$\left[\begin{array}{cc|cc} 1 & -1 & 2 & 4 \\ 3 & -1 & 2 & -1 \end{array} \right]$$

reduces to

$$\left[\begin{array}{cc|cc} 1 & 0 & 0 & -\frac{5}{2} \\ 0 & 1 & -2 & -\frac{13}{2} \end{array} \right]$$

so

$$P_{B \rightarrow B'} = \begin{bmatrix} 0 & -\frac{5}{2} \\ -2 & -\frac{13}{2} \end{bmatrix}$$

(c)

$$\left[\begin{array}{cc|cc} 2 & 4 & 1 & 0 \\ 2 & -1 & 0 & 1 \end{array} \right]$$

reduces to

$$\left[\begin{array}{cc|cc} 1 & 0 & \frac{1}{10} & \frac{2}{5} \\ 0 & 1 & \frac{1}{5} & -\frac{1}{5} \end{array} \right]$$

so

$$P_{E \rightarrow B} = \begin{bmatrix} \frac{1}{10} & \frac{2}{5} \\ \frac{1}{5} & -\frac{1}{5} \end{bmatrix}$$

where E is the standard basis for $\mathbb{R}^2$. Hence

$$[w]_B = P_{E \rightarrow B}[w]_E \begin{bmatrix} \frac{1}{10} & \frac{2}{5} \\ \frac{1}{5} & -\frac{1}{5} \end{bmatrix} \begin{bmatrix} 3 \\ -5 \end{bmatrix} = \begin{bmatrix} -\frac{17}{10} \\ \frac{8}{5} \end{bmatrix}$$

and

$$[w]_{B'} = P_{B \rightarrow B'}[w]_B \begin{bmatrix} 0 & -\frac{5}{2} \\ -2 & -\frac{13}{2} \end{bmatrix} \begin{bmatrix} -\frac{17}{10} \\ \frac{8}{5} \end{bmatrix} = \begin{bmatrix} -4 \\ -7 \end{bmatrix}$$

If we are to compute $[w]_{B'}$ directly, then we will have

$$\left[\begin{array}{cc|cc} 1 & -1 & 1 & 0 \\ 3 & -1 & 0 & 1 \end{array} \right]$$

reduces to

$$\left[\begin{array}{cc|cc} 1 & 0 & -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 & -\frac{3}{2} & \frac{1}{2} \end{array} \right]$$

so

$$P_{E \rightarrow B'} = \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ -\frac{3}{2} & \frac{1}{2} \end{bmatrix}$$

thus

$$[w]_{B'} = P_{E \rightarrow B'}[w]_E \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ -\frac{3}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 3 \\ -5 \end{bmatrix} = \begin{bmatrix} -4 \\ -7 \end{bmatrix}$$

we obtain the same result, it is because $[w]_{B'} = P_{B \rightarrow B'}[w]_B = P_{E \rightarrow B'}[w]_E$.

In SymIntegration, we can compute the transition matrix from B' to B , the transition matrix from B to B' , the coordinate vector $[w]_B$, and $[w]_{B'}$ with this function:
basistransition_withcoordinatevector(vector<vector<double>> &oldbasis, vector<vector<double>> &newbasis, vector<double> &w))

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Transition from Old Basis to New Basis ]# ./main
*****
New basis (B):
      2.000000      4.000000
      2.000000     -1.000000

Old basis (B'):
      1.000000     -1.000000
      3.000000     -1.000000

Transition matrix from old basis to new basis:
      2.000000      4.000000      1.000000     -1.000000
      2.000000     -1.000000      3.000000     -1.000000

Transition matrix from elementary basis to new basis:
      2.000000      4.000000      1.000000      0.000000
      2.000000     -1.000000      0.000000      1.000000

Transition matrix from elementary basis to old basis:
      1.000000     -1.000000      1.000000      0.000000
      3.000000     -1.000000      0.000000      1.000000

[ I | transition from B' to B]:
      1.000000      0.000000      1.300000     -0.500000
      0.000000      1.000000     -0.400000     -0.000000

[ I | transition from E to B]:
      1.000000      0.000000      0.100000      0.400000
      0.000000      1.000000      0.200000     -0.200000

```

Figure 6.16: The computation to compute the transition matrix from B' to B , the transition matrix from B to B' , the coordinate vector $[w]_B$, and $[w]_{B'}$. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Transition from Old Basis to New Basis/main.cpp).

```

From matrix [ B | B' ],
P_{B' -> B}:
      1.300000      -0.500000
     -0.400000      -0.000000

From matrix [ B' | B ],
P_{B -> B'}:
      0.000000      -2.500000
     -2.000000     -6.500000

P_{E -> B}:
      0.100000      0.400000
      0.200000     -0.200000

P_{E -> B'}:
     -0.500000      0.500000
     -1.500000      0.500000

w:
 3.000000
-5.000000

w_{B}:
-1.700000
 1.600000

w_{B'}:
-4.000000
-7.000000

```

Figure 6.17: The computation to compute the transition matrix from B' to B , the transition matrix from B to B' , the coordinate vector $[w]_B$, and $[w]_{B'}$. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Transition from Old Basis to New Basis/main.cpp).

viii. Compute Basis for Row Space and a Column Space of a Matrix with SymIntegration

Find a basis for the row space and the column space of the matrix

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix}$$

Solution:

Since elementary row operations do not change the row space of a matrix, we can find a basis for the row space of A by finding a basis for the row space of any row echelon form of A . Reducing A to row echelon form, we obtain

$$R = \begin{bmatrix} 1 & -3 & 4.5 & -0.5 & 4 & 1 \\ 0 & 0 & 1 & 3 & -2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The nonzero row vectors of R form a basis for the row space of R and hence form a basis for the row space of A . These basis vectors are

$$r_1 = [1 \quad -3 \quad 4.5 \quad -0.5 \quad 4 \quad 1]$$

$$r_2 = [0 \quad 0 \quad 1 \quad 3 \quad -2 \quad -6]$$

$$r_3 = [0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 5]$$

The problem of finding a basis for the column space of a matrix A is complicated by the fact that an elementary row operation can alter its column space. However, the good news is that elementary row operations do not alter dependence relationships among the column vectors.

The row echelon form of A is R , keeping in mind that A and R can have different column spaces, we cannot find a basis for the column space of A directly from the column vectors of R .

However, if we can find a set of column vectors of R that forms a basis for the column space of R , then the corresponding column vectors of A will form a basis for the column space of A .

Since the first, third, and fifth columns of R contain the leading 1's of the row vectors, the vectors

$$c'_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad c'_2 = \begin{bmatrix} 4.5 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad c'_3 = \begin{bmatrix} -0.5 \\ -2 \\ 1 \\ 0 \end{bmatrix}$$

form a basis for the column space of R . Thus, the corresponding column vectors of A , which are

$$c_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \\ -1 \end{bmatrix}, \quad c_2 = \begin{bmatrix} 4 \\ 9 \\ 9 \\ -4 \end{bmatrix}, \quad c_3 = \begin{bmatrix} 5 \\ 8 \\ 9 \\ -5 \end{bmatrix}$$

form a basis for the column space of A .

In SymIntegration, to obtain the basis for the row space we can use this function:
rowspacebasis(vector<vector<double>> &A)

with A as the matrix, it can be non square matrix as well.

To obtain the basis for the column space we can use this function:
columnspacebasis(vector<vector<double>> &A)

with A as the matrix, it can be non square matrix as well.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Row Space Basis and Column Space Basis ]# ./main
*****
A:
      1.000000      -3.000000      4.000000      -2.000000      5.000000      4.000000
      2.000000      -6.000000      9.000000      -1.000000      8.000000      2.000000
      2.000000      -6.000000      9.000000      -1.000000      9.000000      7.000000
      -1.000000      3.000000      -4.000000      2.000000      -5.000000      -4.000000

R:
      1.000000      -3.000000      4.500000      -0.500000      4.000000      1.000000
      0.000000      0.000000      1.000000      3.000000      -2.000000      -6.000000
      0.000000      0.000000      0.000000      0.000000      1.000000      5.000000
      0.000000      0.000000      0.000000      0.000000      0.000000      0.000000

Basis for the row space of the matrix A:
      1.000000      -3.000000      4.500000      -0.500000      4.000000      1.000000
      0.000000      0.000000      1.000000      3.000000      -2.000000      -6.000000
      0.000000      0.000000      0.000000      0.000000      1.000000      5.000000
*****
*****
A:
      1.000000      -3.000000      4.000000      -2.000000      5.000000      4.000000
      2.000000      -6.000000      9.000000      -1.000000      8.000000      2.000000
      2.000000      -6.000000      9.000000      -1.000000      9.000000      7.000000
      -1.000000      3.000000      -4.000000      2.000000      -5.000000      -4.000000

R:
      1.000000      -3.000000      4.500000      -0.500000      4.000000      1.000000
      0.000000      0.000000      1.000000      3.000000      -2.000000      -6.000000
      0.000000      0.000000      0.000000      0.000000      1.000000      5.000000
      0.000000      0.000000      0.000000      0.000000      0.000000      0.000000

Basis for the column space of the matrix A:
      1.000000      4.000000      5.000000
      2.000000      9.000000      8.000000
      2.000000      9.000000      9.000000
      -1.000000      -4.000000      -5.000000
*****
Time taken by function: 2870 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Row Space Basis and Column Space Basis ]#
```

Figure 6.18: The computation to obtain the basis for the row space and column space of matrix A . (SymIntegration/Examples/Linear Algebra/Test SymIntegration Row Space Basis and Column Space Basis/main.cpp).

ix. Compute Basis for the Solution Space of the Linear System $Ax = \mathbf{0}$ with SymIntegration

Find the rank, nullity, and the basis for the solution space of the linear system $Ax = \mathbf{0}$ (homogeneous system), with matrix

$$A = \begin{bmatrix} -1 & 2 & 0 & 4 & 5 & -3 \\ 3 & -7 & 2 & 0 & 1 & 4 \\ 2 & -5 & 2 & 4 & 6 & 1 \\ 4 & -9 & 2 & -4 & -4 & 7 \end{bmatrix}$$

Solution:

The reduced row echelon form of A is

$$R = \begin{bmatrix} 1 & 0 & -4 & -28 & -37 & 13 \\ 0 & 1 & -2 & -12 & -16 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Since this matrix has two leading 1's, its row and column spaces are two dimensional and

$$\text{rank}(A) = 2$$

To find the nullity of A , we must find the dimension of the solution space of the linear system $Ax = \mathbf{0}$. This system can be solved by reducing its augmented matrix to reduced row echelon form. The resulting matrix will be R with an additional last column of zeros, and hence the corresponding system of equations will be

$$x_1 - 4x_3 - 28x_4 - 37x_5 + 13x_6 = 0$$

$$x_2 - 2x_3 - 12x_4 - 16x_5 + 5x_6 = 0$$

Solving these equations for the leading variables yields

$$x_1 = 4x_3 + 28x_4 + 37x_5 - 13x_6$$

$$x_2 = 2x_3 + 12x_4 + 16x_5 - 5x_6$$

from which we obtain the general solution

$$x_1 = 4r + 28s + 37t - 13u$$

$$x_2 = 2r + 12s + 16t - 5u$$

$$x_3 = r$$

$$x_4 = s$$

$$x_5 = t$$

$$x_6 = u$$

or in column vector form

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = r \begin{bmatrix} 4 \\ 2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} 28 \\ 12 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} 37 \\ 16 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + u \begin{bmatrix} -13 \\ -5 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

Because the four vectors form a basis for the solution space,

$$\text{nullity}(A) = 4$$

In `SymIntegration`, to obtain the basis for the solution space of the linear system $Ax = 0$ we can use this function:

homogeneouslinearsystembasis(vector<vector<double>> &A)

with A as the matrix, it can be non square matrix as well.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Homogeneous Linear System Basis for the Solution Space ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Homogeneous Linear System Basis for the Solution Space ]# ./main

*****

A:
      -1.000000      2.000000      0.000000      4.000000      5.000000     -3.000000
      3.000000     -7.000000      2.000000      0.000000      1.000000      4.000000
      2.000000     -5.000000      2.000000      4.000000      6.000000      1.000000
      4.000000     -9.000000      2.000000     -4.000000     -4.000000      7.000000

Rank(A) = 2
nullity(A) = 4
dim(A) = 6

R:
      1.000000      0.000000     -4.000000     -28.000000     -37.000000     13.000000
      0.000000      1.000000     -2.000000     -12.000000     -16.000000      5.000000
      0.000000      0.000000      0.000000      0.000000      0.000000      0.000000
      0.000000      0.000000      0.000000      0.000000      0.000000      0.000000

Basis for the solution space of Ax = 0 :
      4.000000     28.000000     37.000000     -13.000000
      2.000000     12.000000     16.000000      -5.000000
      1.000000      0.000000      0.000000      0.000000
      0.000000      1.000000      0.000000      0.000000
      0.000000      0.000000      1.000000      0.000000
      0.000000      0.000000      0.000000      1.000000

*****

Time taken by function: 1589 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Homogeneous Linear System Basis for the Solution Space ]#
```

Figure 6.19: The computation to obtain the basis for the solution space of the linear system $Ax = 0$. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Homogeneous Linear System Basis for the Solution Space/main.cpp*).

x. Compute State Vector $\mathbf{x}_k$ in a Markov Chain with SymIntegration

Suppose that Glanz is going to find a nice place to do workout, improve her physical, spiritual and mental power, exercise with Tai Chi, FoY, and Ashtanga Yoga. There are 3 places on the mountain she lives on that can accomodate her needs, they are VP (Valhalla Projection), PB (Puncak Bintang), and Kathmandu. Based on data from 2020 Glanz' migration pattern of moving to certain place can be modeled by a Markov chain with transition matrix

$$P = \begin{bmatrix} 0.97 & 0.51 & 0.6 \\ 0.02 & 0.33 & 0.3 \\ 0.01 & 0.16 & 0.1 \end{bmatrix}$$

That is

$p_{11} = 0.97$ = probability that Glanz will go to VP tomorrow when she already in VP today.

$p_{12} = 0.51$ = probability that Glanz will go to VP tomorrow when she already in PB today.

$p_{13} = 0.6$ = probability that Glanz will go to VP tomorrow when she already in Kathmandu today.

$p_{21} = 0.02$ = probability that Glanz will go to PB tomorrow when she already in VP today.

$p_{22} = 0.33$ = probability that Glanz will go to PB tomorrow when she already in PB today.

$p_{23} = 0.3$ = probability that Glanz will go to PB tomorrow when she already in Kathmandu today.

$p_{31} = 0.01$ = probability that Glanz will go to Kathmandu tomorrow when she already in VP today.

$p_{32} = 0.16$ = probability that Glanz will go to Kathmandu tomorrow when she already in PB today.

$p_{33} = 0.1$ = probability that Glanz will go to Kathmandu tomorrow when she already in Kathmandu today.

Assuming that t is in day and Glanz started all her new cycle of becoming Double-Star Hunter, Nature Golden Child and True Earth Rune bearer from Puncak Bintang at time $t = 0$ (in 2020). Track Glanz' probable locations over a 7-days and after 46-days period. Then determine the steady-state vector $\mathbf{q}$, if exist.

Solution:

Let $x_1(k)$, $x_2(k)$, and $x_3(k)$ be the probabilities that Glanz is in VP, PB, and Kathmandu, respectively, at time $t = k$, and let

$$\mathbf{x}(k) = \begin{bmatrix} x_1(k) \\ x_2(k) \\ x_3(k) \end{bmatrix}$$

be the state vector at that time. Since we know with certainty that Glanz is in PB at time $t = 0$, the initial state vector is

$$\mathbf{x}(0) = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

The state vector over a 7-days period and then after 46-days are

$$\mathbf{x}(1) = P\mathbf{x}(0) = \begin{bmatrix} 0.51 \\ 0.33 \\ 0.16 \end{bmatrix}, \quad \mathbf{x}(2) = P\mathbf{x}(1) = \begin{bmatrix} 0.759 \\ 0.1671 \\ 0.0739 \end{bmatrix}, \quad \mathbf{x}(3) = P\mathbf{x}(2) = \begin{bmatrix} 0.8658 \\ 0.0925 \\ 0.0417 \end{bmatrix}$$

$$\begin{aligned}
x(4) = Px(3) &= \begin{bmatrix} 0.912 \\ 0.0604 \\ 0.0276 \end{bmatrix}, & x(5) = Px(4) &= \begin{bmatrix} 0.9320 \\ 0.0464 \\ 0.0215 \end{bmatrix}, & x(6) = Px(5) &= \begin{bmatrix} 0.9407 \\ 0.0404 \\ 0.0189 \end{bmatrix} \\
x(7) = Px(6) &= \begin{bmatrix} 0.9444 \\ 0.0378 \\ 0.0178 \end{bmatrix}, \dots, x(46) = Px(45) &= \begin{bmatrix} 0.9473 \\ 0.0358 \\ 0.0169 \end{bmatrix}
\end{aligned}$$

Now we take a look again at the entries at matrix P

$$P = \begin{bmatrix} 0.97 & 0.51 & 0.6 \\ 0.02 & 0.33 & 0.3 \\ 0.01 & 0.16 & 0.1 \end{bmatrix}$$

Since the entries of P are positive, the Markov chain is regular and hence has a unique steady-state vector q . To find q we will solve the system $(I - P)q = 0$, which we can write as

$$\begin{aligned}
(I - P)q &= 0 \\
\begin{bmatrix} 0.03 & -0.51 & -0.6 \\ -0.02 & 0.67 & -0.3 \\ -0.01 & -0.16 & 0.9 \end{bmatrix} \begin{bmatrix} q_1 \\ q_2 \\ q_3 \end{bmatrix} &= \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}
\end{aligned}$$

One-way to determine q_1, q_2, q_3 is to reduce the matrix $(I - P)$ into reduced row echelon form then take parameter solution and then use substitution of the knowledge that $q_1 + q_2 + q_3 = 1$.

Since Gauss-Jordan elimination especially for fraction is not an easy task to be done, thus after we retrieve three equations from $(I - P)q$, all equations are homogeneous (equal to 0), then substitute this

$$q_3 = 1 - q_1 - q_2$$

so we have three equations with 2 unknowns. It is easy to be done now, we can just take two out of the three equations (random picking is not a problem) and then we can compute the value for q_1 and q_2 with simple elimination method from algebra, after that we can get the value for q_3 . This is the method of thinking of how we solve with C++ as well.

We will obtain

$$\begin{aligned}
q_1 &= 0.947260 \\
q_2 &= 0.035842 \\
q_3 &= 0.016897
\end{aligned}$$

the steady-state vector is

$$\begin{bmatrix} q_1 \\ q_2 \\ q_3 \end{bmatrix} = \begin{bmatrix} 0.947260 \\ 0.035842 \\ 0.016897 \end{bmatrix}$$

which is consistent with the results obtained above after 46-days period ($x(46)$), the 46-days state vector converge to the steady-state vector.

In SymIntegration to compute state vector x_k we can use this function:
MarkovChain(vector<vector<double>>& P, vector<double>& x0, int k)

with P as the transition matrix, x_0 as the initial state vector, and k is to represents the time $t = k$.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Steady-State Vector for Markov Chain ]# ./main
P =
      0.970000      0.510000      0.600000
      0.020000      0.330000      0.300000
      0.010000      0.160000      0.100000

x_{0} =
0.000000
1.000000
0.000000

State vector, k = 46 :
0.947261
0.035842
0.016897

Time taken by function: 1977 microseconds
```

Figure 6.20: The computation to obtain the state vector x_k at time $t = k = 46$ (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Steady-State Vector for Markov Chain/main.cpp*).

V. EIGENVALUES AND EIGENVECTORS

- [SI*] The word *eigen* is from the German word, meaning "characteristic". The underlying idea first appeared in the study of rotational motion but was later used to classify various kinds of surfaces and to describe solutions to certain differential equations. It has applications in various fields such as computer graphics, mechanical vibrations, heat flow, population dynamics, quantum mechanics, and economics.
- [SI*] Let A be a square matrix, $A \in \mathbb{R}^{n \times n}$. Many of the important properties of a matrix are summarized by its eigenvalues and eigenvectors [?].

Definition 6.29: Eigenvalue and Eigenvector

Let $\mathbb{C}$ denote the complex plane. $\lambda \in \mathbb{C}$ is an eigenvalue of A (an $n \times n$ matrix) if and only if there is some $x \in \mathbb{C}^n$, $x \neq 0$ (a nonzero vector x in $\mathbb{R}^n$ is called an eigenvector of A / or the matrix operator T_A), such that

$$Ax = \lambda x \quad (6.57)$$

for some scalar λ . The scalar λ is called an eigenvalue of A .

Definition 6.30: Spectrum and Spectral Radius

Let $\sigma(A)$ denote the set of eigenvalues of A ; we refer to it as the spectrum of A . If $Ax = \lambda x$ for a nonzero x , then x is an eigenvector corresponding to $\lambda \in \sigma(A)$. The spectral radius of A is

$$\rho(A) \equiv \max\{|\lambda| : \lambda \in \sigma(A)\} \quad (6.58)$$

If x is an eigenvector, then so is any scalar multiple of x ; hence eigenvectors are defined only up to a proportionality constant. Also $\lambda \in \sigma(A)$ if and only if

$$\det(A - \lambda I) = 0 \quad (6.59)$$

the characteristic equation of A , implying that $\lambda \in \sigma(A)$ if and only if $A - \lambda I$ is singular.

- [SI*] Eigenvalue problems have a very different character from the problems involving square or rectangular linear systems of equations. For a system of equations, the domain of A could be one space and the range could be a different one. For example, matrix A maps n -vectors of polynomial coefficients to m -vectors of sampled polynomial values. To ask about the eigenvalues of such an A would be meaningless.

Eigenvalue problems make sense only when the range and the domain spaces are the same. This reflects the fact that in applications, eigenvalues are generally used where a matrix is to be compounded iteratively, either explicitly as a power A^k or implicitly in a functional form such as e^{tA} .

Eigenvalues and eigenvectors are useful for two reasons, one algorithmic, the other physical. Algorithmically, eigenvalue analysis can simplify solutions of certain problems by reducing a coupled system to a collection of scalar problems. Physically, eigenvalue analysis can give insight into the behavior of evolving systems governed by linear equations. The most

familiar examples in this latter class are the study of resonance (e.g., of musical instruments when struck or plucked or bowed) and of stability (e.g., of fluid flows subjected to small perturbations). In such cases eigenvalues tend to be particularly useful for analyzing behavior for large times t .

[SI*] In general, the image of a vector x under multiplication by a square matrix A differs from x in both magnitude and direction.

However, in the special case where x is an eigenvector of A , multiplication by A leaves the direction unchanged.

For example, in $\mathbb{R}^2$ or $\mathbb{R}^3$ multiplication by A maps each eigenvector x of A along the same line through the origin as x .

Depending on the sign and magnitude of the eigenvalue λ corresponding to x , the operation

$$Ax = \lambda x$$

compresses or stretches x by a factor of λ , with a reversal of direction in the case where λ is negative.

[SI*] Note first that the equation

$$Ax = \lambda x$$

can be rewritten as

$$Ax = \lambda Ix$$

or equivalently, as

$$(\lambda I - A)x = 0$$

For λ to be an eigenvalue of A this equation must have a nonzero solution for x .

Theorem 6.40: Characteristic Equation

If A is an $n \times n$ matrix, then λ is an eigenvalue of A if and only if it satisfies the equation

$$\det(\lambda I - A) = 0 \tag{6.60}$$

This is called the characteristic equation of A .

[SI*] When the $\det(\lambda I - A)$ is expanded, the result is a polynomial $p(\lambda)$ of degree n that is called the characteristic polynomial of A .

Since a polynomial of degree n has at most n distinct roots, it follows that the equation

$$\lambda^n + c_1\lambda^{n-1} + \cdots + c_n = 0 \tag{6.61}$$

has at most n distinct solutions and consequently that an $n \times n$ matrix has at most n distinct eigenvalues. Some of these solutions may be complex numbers, and it is possible for a matrix to have complex eigenvalues, even if that matrix itself has real entries.

Even if a matrix is real, some of its eigenvalues may be complex. Physically, this is related to the phenomenon that real dynamical systems can have motions that oscillate as well as grow

or decay. Algorithmically, it means that even if the input to a matrix eigenvalue problem is real, the output may have to be complex.

Theorem 6.41: Eigenvalues for Triangular Matrix

If A is an $n \times n$ triangular matrix (upper triangular, lower triangular, or diagonal), then the eigenvalues of A are the entries on the main diagonal of A .

Theorem 6.42: Eigenvalues and Eigenvectors

If A is an $n \times n$ matrix, the following statements are equivalent.

- (a) λ is an eigenvalue of A .
- (b) The system of equations $(\lambda I - A)x = \mathbf{0}$ has nontrivial solutions.
- (c) There is nonzero vector x such that $Ax = \lambda x$.
- (d) λ is a solution of the characteristic equation $\det(\lambda I - A) = 0$

[SI*] Eigenvectors corresponding to an eigenvalue λ of a matrix A are the nonzero vectors that satisfy the equation

$$(\lambda I - A)x = \mathbf{0}$$

these eigenvectors are the nonzero vectors in the null space of the matrix $\lambda I - A$. We call this null space the eigenspace of A corresponding to λ .

The eigenspace of A corresponding to the eigenvalue λ is the solution space of the homogeneous system $(\lambda I - A)x = \mathbf{0}$.

[SI*] Once the eigenvalues and eigenvectors of a matrix A are found, it is a simple matter to find the eigenvalues and eigenvectors of any positive integer power of A .

If λ is an eigenvalue of A and x is a corresponding eigenvector, then

$$A^2x = A(Ax) = A(\lambda x) = \lambda(Ax) = \lambda(\lambda x) = \lambda^2x$$

which shows that λ^2 is an eigenvalue of A^2 and that x is a corresponding eigenvector.

Theorem 6.43: Eigenvalue for Powers of a Matrix

If k is a positive integer, λ is an eigenvalue of a matrix A , and x is a corresponding eigenvector, then λ^k is an eigenvalue of A^k and x is a corresponding eigenvector.

Theorem 6.44: Eigenvalue and Invertibility

A square matrix A is invertible if and only if $\lambda = 0$ is not an eigenvalue of A .

[SI*] It is often useful to decompose a matrix into the product of other matrices. A particularly useful decomposition is the Jordan decomposition.

Suppose that A is a $n \times n$ nonsingular square matrix and has n distinct eigenvalues. Then

$$A = NDN^{-1} \quad (6.62)$$

where D is a diagonal matrix with the distinct eigenvalues on the diagonal, and the columns of N are right eigenvectors of A .

$$D = \begin{bmatrix} \lambda_1 & 0 & 0 & \dots & 0 \\ 0 & \lambda_2 & 0 & \dots & 0 \\ 0 & 0 & \lambda_3 & \dots & 0 \\ \vdots & 0 & \dots & \ddots & 0 \\ 0 & 0 & 0 & \dots & \lambda_n \end{bmatrix} \quad (6.63)$$

Equivalently we can express

$$A = N^{-1}DN \quad (6.64)$$

where the diagonal elements of D are the left eigenvalues of A and the rows of N are left eigenvectors of A .

[SI*] **Defective Eigenvalues and Matrices**

If a matrix A has the characteristic polynomial $(z - 2)^3$, so there is a single eigenvalue $\lambda = 2$ of algebraic multiplicity 3. In the case of this, we can choose three independent eigenvectors, for example e_1, e_2 , and e_3 , so the geometric multiplicity is also 3.

An eigenvalue whose algebraic multiplicity exceeds its geometric multiplicity is a defective eigenvalue. A matrix that has one or more defective eigenvalues is a defective matrix. Any diagonal matrix is nondefective. For such a matrix, both the algebraic and the geometric multiplicities of an eigenvalue λ are equal to the number of its occurrences along the diagonal.

[SI*] **Determinant and Trace**

The trace of $A \in \mathbb{C}^{m \times m}$ is the sum of its diagonal elements:

$$\text{tr}(A) = \sum_{i=1}^m a_{ii}$$

both the trace and determinant are related simply to the eigenvalues.

Theorem 6.45: Determinant and Trace Relationship with Eigenvalues

The determinant and trace of a matrix A are equal to the product and the sum of the eigenvalues of A , counted with algebraic multiplicity

$$\det(A) = \prod_{i=1}^m \lambda_i \quad (6.65)$$

$$\text{tr}(A) = \sum_{i=1}^m \lambda_i \quad (6.66)$$

[SI*] **Unitary Diagonalization**

It sometimes happens that not only does an $m \times m$ matrix A have m linearly independent

eigenvectors, but these can be chosen to be orthogonal. In such a case, A is unitarily diagonalizable, that is, there exists a unitary matrix Q such that

$$A = Q\Lambda Q^* \quad (6.67)$$

This factorization is both an eigenvalue decomposition and a singular value decomposition, aside from the matter of the signs (possibly complex) of the entries of A .

Theorem 6.46: Hermitian Matrix and Eigenvalues

A hermitian matrix is unitarily diagonalizable, and its eigenvalues are real.

Theorem 6.47: Normal Matrix and Unitarily Diagonalizable

A matrix is unitarily diagonalizable, if and only if it is normal ($A^*A = AA^*$).

[SI*] **Unitary Triangularization (Schur Factorization)**

One final matrix factorization is actually one that is most useful in numerical analysis, because all matrices, including defective ones, can be factored in this way. A Schur factorization of a matrix A is a factorization

$$A = QTQ^* \quad (6.68)$$

where Q is unitary and T is upper-triangular. Note that since A and T are similar, the eigenvalues of A necessarily appear on the diagonal of T .

Theorem 6.48: Schur Factorization

Every square matrix A has a Schur factorization.

[SI*] **Eigenvalue-Revealing Factorizations**

1. A diagonalization $A = X\Lambda X^{-1}$ exists if and only if A is nondefective.
2. A unitary diagonalization $A = Q\Lambda Q^*$ exists if and only if A is normal.
3. A unitary triangularization (Schur factorization) $A = QTQ^*$ always exists.

To compute eigenvalues, we shall construct one of these factorizations. In general, this will be the Schur factorization, since this applies without restriction to all matrices. Moreover, since unitary transformations are involved, the algorithms that result tend to be numerically stable.

If A is normal, then the Schur form comes out diagonal, and in particular, if A is hermitian, then we can take advantage of this symmetry throughout the computation and reduce A to diagonal form with half as much work or less than is required for general A .

[SI*] **Eigenvalue Algorithms**

For computing eigenvalues and eigenvectors most of the available algorithms proceed in two phases: first, a preliminary reduction from full to structured form; then, an iterative process for the final convergence.

The method of computing eigenvalue by using rootfinding technique on characteristic polynomial is a bad strategy [28]. Polynomial rootfinding is an ill-conditioned problem in general, even when the underlying eigenvalue problem is well-conditioned. Polynomial rootfinding

is a mainstream topic in scientific computing—precisely because it is so rarely the best way to solve applied problems.

It is well known that no formula exists for expressing the roots of an arbitrary polynomial, given its coefficients. This impossibility result is one of the crowning achievements of a body of mathematical work carried out by Abel, Galois, and others in the nineteenth century. Abel proved in 1824 that no analogue of the quadratic formula can exist for polynomials of degree 5 or more. There could be no computer program that would produce the exact roots of an arbitrary polynomial in a finite number of steps. It follows that the same conclusion applies to the more general problem of computing eigenvalues of matrices.

An eigenvalue solver cannot be based on the same kind of techniques that we have used so far for solving linear systems. Methods like Householder reflections and Gaussian elimination would solve linear systems of equations exactly in a finite number of steps if they could be implemented in exact arithmetic. By contrast, any eigenvalue solver must be iterative.

The goal of an eigenvalue solver is to produce sequences of numbers that converge rapidly towards eigenvalues. In this respect eigenvalue computations are more representative of scientific computing than solutions of linear systems of equations.

The need to iterate may seem discouraging at first, but the algorithms available in the field converge extraordinary quickly. In most cases it is possible to compute sequences of numbers that double or triple the numbers of digits of accuracy at every step. Thus, although computing eigenvalues is an "unsolvable" problem in principle, in practice it differs from the solution of linear systems by only a small constant factor, typically closer to 1 than 10.

[SI*] Properties of Eigenvalues

1. Eigenvectors with distinct eigenvalues are linearly independent.
2. Singular matrices have zero eigenvalues.
3. If A is a square matrix, then $\lambda = 0$ is not an eigenvalue of A .
4. For a scalar multiple of a matrix, if A is a square matrix and λ is an eigenvalue of A then $k\lambda$ is an eigenvalue of kA .
5. For matrix powers, if A is a square matrix and λ is an eigenvalue of A and $n \geq 0$ is an integer, then λ^n is an eigenvalue of A^n .
6. For polynomial of matrix, if A is a square matrix, λ is an eigenvalue of A and $p(x)$ is a polynomial in variable x , then $p(\lambda)$ is the eigenvalue of matrix $p(A)$.
7. If A is a square matrix, λ is an eigenvalue of A , then λ^{-1} is an eigenvalue of A^{-1} .
8. If A is a square matrix, λ is an eigenvalue of A , then λ is an eigenvalue of A^T .

[SI*] Properties of Eigenvectors

1. Eigenvectors provide a concise representation of a matrix, as they capture the most important features of the matrix.
2. Eigenvectors can be used to solve linear systems of equations, which arise in many practical applications.
3. Eigenvectors can be used to diagonalize a matrix, which simplifies many computations, e.g. power and exponentials of the matrix.

4. Eigenvector computation can be computationally expensive, particularly for large matrices.
5. Eigenvectors may not exist for all matrices, or they may be complex numbers.
6. Eigenvectors may not be unique, as a matrix may have multiple eigenvectors corresponding to the same eigenvalue.

[SI*] **Applications of Eigenvalues and Eigenvectors**

1. **Stability Analysis**

In engineering, they help analyze the stability of systems.

2. **Quantum Mechanics**

In physics, they are used to study the states of quantum systems.

3. **Principal Component Analysis (PCA)**

In machine learning, they are used to reduce the dimensionality of data. The eigenvectors corresponding to the largest eigenvalues capture the most important features of the data, helping reduce complexity while retaining important patterns.

4. **Google's PageRank Algorithm**

Eigenvalues and eigenvectors play a key role in Google's PageRank algorithm, which determines the importance of web pages based on link structures. In PageRank, each page is represented as a node, and the links between pages form a matrix. By calculating the eigenvalues and eigenvectors of this matrix, the most important pages (with the highest eigenvector values) are identified. These pages are considered more relevant and are ranked higher in search results. The eigenvector corresponding to the eigenvalue of 1 determines the relative importance of each page in the network. This method allows Google to rank pages based on their connectivity, rather than just the number of incoming links.

5. **Markov Processes**

In a Markov process, eigenvalues are used to find the stationary distribution, which represents the long-term stable state of the system. The eigenvector corresponding to eigenvalue 1 gives the steady-state probabilities. Additionally, the second-largest eigenvalue determines the rate of convergence to equilibrium.

6. **Electrical Engineering**

In RLC circuits, the analysis of the currents in the loops can be solved using eigenvalues and eigenvectors. The system of differential equations representing the circuit can be transformed into a matrix form. The eigenvalues and eigenvectors of this matrix provide the general solution for the system, which describes how the current in each loop behaves over time. The eigenvectors represent the natural modes of the circuit, while the eigenvalues determine the rates of change. This method helps in understanding the long-term behavior of the RLC circuit.

7. **Mechanical Engineering**

Eigenvalues and eigenvectors enable us to "decompose" a linear process into smaller, more manageable tasks. When stress is applied to a "plastic" solid, for example, the deformation can be divided into "principal directions," or the directions where the deformation is greatest. The eigenvectors in the principal directions are the eigenvectors, and the associated eigenvalue is the percentage deformation in each principal direction. Also mechanical engineers utilize eigenvalues and eigenvectors to analyze dynamic systems such as rotating machinery and vehicle dynamics. Eigenvalues determine the

stability of rotational speeds or vehicle maneuvers, while eigenvectors represent the modes of motion.

8. System of Communication

Claude Shannon utilized eigenvalues to calculate the theoretical limit of how much information can be carried via a communication channel such as a telephone line or the air. The eigenvectors and eigenvalues of the communication channel (represented as a matrix) are calculated, and then the eigenvalues are waterfilled. The eigenvalues are then essentially the gains of the channel's fundamental modes, which are recorded by the eigenvectors.

9. Bridge Construction

The smallest magnitude eigenvalue of a system that models the bridge is the natural frequency of the bridge. Engineers use this knowledge to guarantee that their structure are stable.

10. Automobile Stereo System Design

Eigenvalue analysis is also employed in the design of car stereo systems, where it aids in the reproduction of car vibration caused by music.

11. Oil Exploration

Eigenvalue analysis is commonly used by oil firms to explore land for oil. Because oil, dirt, and other substances all produce linear systems with varying eigenvalues, eigenvalue analysis can help pinpoint where oil reserves lie. Oil companies set up probes all-around a site to pick up waves created by a massive truck vibrating the ground. The waves are modified when they move through the different substances in the earth. The oil corporations are directed to possible drilling sites based on the study of these waves.

GlanzFreya' Guide 6.2: Compute Basis for the Eigenspace Corresponding to an Eigenvalue

These are the methods / procedure to compute basis for the eigenspace corresponding to λ_i .

Suppose we have square matrix A with size of $n \times n$, and then we want to compute its basis for the eigenspace.

1. Compute the eigenvalues, with formula

$$\det(\lambda I - A) = 0$$

we will obtain characteristic equation that will result in polynomial of degree n , thus with the help of root finding formula (Bisection, Newton-Raphson) or root quadratic formula we will obtain at most n distinct eigenvalues.

2. After we obtain $\lambda_1, \lambda_2, \dots, \lambda_n$, each value of the eigenvalue if substituted into the equation

$$(\lambda_i I - A)x = 0$$

can be used to determine the basis for eigenspace corresponding to λ_i , with $i = 1, 2, \dots, n$ and x is an eigenvector of A corresponding to λ_i .

3. Compute from $i = 1, 2, \dots, n$ the matrix

$$(\lambda_i I - A)x = 0$$

Solve the system using Gaussian elimination to obtain reduced row echelon form for matrix

$$(\lambda_i I - A)$$

we will obtain the solution for x in parameter term, for example like this:

$$\begin{aligned} x_1 &= 2s \\ x_2 &= 3t - s \\ x_3 &= t \\ &\vdots \\ x_n &= s \end{aligned}$$

with s, t are example of parameter.

GlanzFreya' Guide 6.3: Compute Basis for the Eigenspace Corresponding to an Eigenvalue

4. The eigenvectors x of A corresponding to λ_i are the nonzero vector of the form

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} 2s \\ 3t - s \\ t \\ \vdots \\ s \end{bmatrix} = \begin{bmatrix} 2s \\ -s \\ 0 \\ \vdots \\ s \end{bmatrix} + \begin{bmatrix} 0 \\ 3t \\ t \\ \vdots \\ 0 \end{bmatrix} = s \begin{bmatrix} 2 \\ -1 \\ 0 \\ \vdots \\ 1 \end{bmatrix} + t \begin{bmatrix} 0 \\ 3 \\ 1 \\ \vdots \\ 0 \end{bmatrix}$$

thus we obtain that these eigenvectors below

$$\begin{bmatrix} 2 \\ -1 \\ 0 \\ \vdots \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 3 \\ 1 \\ \vdots \\ 0 \end{bmatrix}$$

are linearly independent and form a basis for the eigenspace corresponding to λ_i .

5. Do the computation till $i = n$.

GlanzFreya' Guide 6.4: The Power Method for Finding the Largest Eigenvalue and its Eigenvector

Power method is a numerical method to compute the largest eigenvalue along with its eigenvector of a general square matrix A , it does not have to be symmetric.

Remark: this Power method and the two following iterative methods are only working if the matrix has real eigenvalues.

Suppose we have a matrix A with size $n \times n$.

The algorithm for power method:

1. Start with a random initial vector v with size n .
2. Create a **for** loop with highest number of iterations that we can define ourselves ($N = 50, N = 100, N = 200$), in this loop we compute

$$\begin{aligned} w &= Av \\ ||w|| &= \sqrt{w \cdot w} \\ w_{\text{normalize}} &= \frac{w}{||w||} \\ v &= w \end{aligned}$$

3. After the iteration finishes, we will obtain the eigenvector v and we can compute the eigenvalue with this formula

$$\lambda = \frac{v^T Av}{v^T v}$$

this formula is called the Rayleigh quotient, it produces an approximate eigenvalue when given an estimated eigenvector.

We will proceed with another two methods to approximate all eigenvalues and eigenvectors, these three methods combine work amazingly well for all general square real matrix (not only work for symmetric matrix, e.g. Jacobi eigenvalue algorithm only works for real symmetric matrix).

GlanzFreya' Guide 6.5: The Deflation Method for Approximating all the Eigenvalues

The method of deflation proceeds after the power method by reducing the matrix A to an $(n-1) \times (n-1)$ matrix for finding the next largest eigenvalue (the second largest eigenvalue) of this matrix, till the smallest eigenvalue.

Suppose we rank the eigenvalues of matrix A like this

$$\lambda_1 > \lambda_2 > \dots > \lambda_n$$

with the corresponding eigenvector v_i for $\lambda_i, i = 1, 2, \dots, n$

1. We perform the power method to obtain v_1 and λ_1 , multiply v_1 by a permutation matrix P which interchanges the largest element and the first element

$$Pv_1 = w_1$$

2. We compute the elementary matrix R such that

$$R^{-1} = [w_1 \quad e_2 \quad e_3]$$

and R is the inverse of R^{-1} with

$$Rw_1 = e_1$$

3. Let

$$B = RPAP^{-1}R^{-1} = RPAPR^{-1}$$

because $P^{-1} = P$, then

$$\begin{aligned} Be_1 &= RPAPR^{-1}e_1 \\ &= RPAPw_1 \\ &= RPAv_1 \\ &= \lambda_1 RPv_1 \\ &= \lambda_1 e_1 \end{aligned}$$

thus e_1 is an eigenvector of B with eigenvalue λ_1 and B must be an upper triangular matrix with λ_1 as the first element on the leading diagonal.

$$B = \begin{bmatrix} \lambda_1 & \dots & x^1 \\ \vdots & \ddots & \vdots \\ 0 & \dots & x^n \end{bmatrix}$$

4. Delete the first column and the first row of matrix $B^{k-1} = RPAP^{-1}R^{-1}$ to give an $(n-1) \times (n-1)$ matrix B^k , then perform the power method on this matrix B^k to obtain the next eigenvalue, with k as the indicator for the iteration number ($k = 1, 2, \dots, n-1$).
5. Repeat till we have computed all the eigenvalues.

GlanzFreya' Guide 6.6: The Inverse Iteration with Shift for Computing all the Eigenvectors

The method of deflation is able to compute all the eigenvalues, now we want to compute all the eigenvectors correspond to each of the eigenvalue.

The inverse iteration method [22] is a natural generalization of the power iteration method. If A is an invertible matrix with real, nonzero eigenvalues $\{\lambda_1, \dots, \lambda_n\}$, then the eigenvalues of A^{-1} are $\{\frac{1}{\lambda_1}, \dots, \frac{1}{\lambda_n}\}$. Thus if

$$|\lambda_1| > |\lambda_2| > \dots > |\lambda_n|$$

then

$$|\frac{1}{\lambda_1}| < |\frac{1}{\lambda_2}| < \dots < |\frac{1}{\lambda_n}|$$

and so by applying the power method iteration on A^{-1} , we can obtain the eigenvector x_n and the eigenvalue λ_n (the smallest eigenvalue).

The advantage of inverse iteration is that it can be easily adapted to find any eigenvalue of the matrix A , instead of just the extreme ones. Observe that for any $\mu \in \mathbb{R}$, the matrix $B = A - \mu I$ has eigenvalues $\{\lambda_1 - \mu, \dots, \lambda_n - \mu\}$. In particular, by choosing μ to be close to an eigenvalue λ_j of A , we can ensure that $\lambda_j - \mu$ is the eigenvalue of B of the smallest magnitude. Then by applying inverse iteration on B , an approximation to x_j and λ_j can be obtained, with $j = 1, 2, \dots, n$.

This version of the algorithm is known as inverse iteration with shift

1. Pick some μ close to the desired eigenvalue, in this case since we already know all the eigenvalues (from the deflation method) we will write it like this

$$\mu_i = \lambda_i - 0.1$$

2. Generate a random vector of size n , we name it v^0 , normalize it so that $\|v^0\| = 1$
3. For $k = 1, 2, \dots, N$ (N is the maximum number of iterations)
Solve for w

$$(A - \mu I)w = v^{(k-1)}$$

then let

$$v^{(k)} = \frac{w}{\|w\|}$$

The inverse iteration with shift method works really well in cases where the eigenvalues are already known.

GlanzFreya' Guide 6.7: The Rayleigh Quotient Iteration

We have learned the method for obtaining an eigenvalue estimate from an eigenvector estimate (the Rayleigh quotient), and another method for obtaining an eigenvector estimate from an eigenvalue estimate (inverse iteration). The Rayleigh quotient iteration idea is to use continually improving eigenvalue estimates to increase the rate of convergence of inverse iteration at every step.

The algorithm:

1. We want to compute the eigenvalues of a matrix A with size $n \times n$.
2. Generate random vector with length of n .

$$\mathbf{v}^{(0)}$$

and normalize it so that $\|\mathbf{v}^{(0)}\| = 1$.

3. For $k = 1, 2, \dots, N$ (define N as the maximum number of iterations)
Solve

$$(A - \lambda^{(k-1)}I)\mathbf{w} = \mathbf{v}^{(k-1)}$$

for $\mathbf{w}$, we need to compute $(A - \lambda^{(k-1)}I)^{-1}$.

Set

$$\mathbf{v}^{(k)} = \frac{\mathbf{w}}{\|\mathbf{w}\|}$$

it normalizes $\mathbf{w}$.

The Rayleigh quotient is

$$\lambda^{(k)} = (\mathbf{v}^{(k)})^T A \mathbf{v}^{(k)}$$

The convergence of this algorithm is spectacular: each iteration triples the number of digits of accuracy.

Theorem 6.49: Rayleigh Quotient Iteration

Rayleigh quotient iteration converges to an eigenvalue/eigenvector pair for all except a set of measure zero of starting vectors $\mathbf{v}^{(0)}$. When it converges, the convergence is ultimately cubic in the sense that if λ_i is an eigenvalue of A and $\mathbf{v}^0$ is sufficiently close to the eigenvector $\mathbf{q}_i$, then

$$\|\mathbf{v}^{(k+1)} - (\pm \mathbf{q}_i)\| = \mathcal{O}(\|\mathbf{v}^{(k)} - (\pm \mathbf{q}_i)\|^3) \quad (6.69)$$

and

$$|\lambda^{(k+1)} - \lambda_i| = \mathcal{O}(|\lambda^{(k)} - \lambda_i|^3) \quad (6.70)$$

as $k \rightarrow \infty$. The $\pm$ signs are not necessarily the same on the two sides Eq. (6.62).

Definition 6.31: Compute Real Eigenvalues with QR Algorithm with Shifts

What makes the QR iteration fly is the introduction of shifts $A \rightarrow A - \mu I$ at each step. This idea leads to cubic convergence, thanks to its implicit connection with the Rayleigh quotient iteration, the downside is that it can only be used if the matrix has real eigenvalues only, if there is even one complex eigenvalue, then this method will result in matrix A that is getting bigger as the iteration goes by.

The Lecture 29 from Trefethen [28] is a great source of learning for this QR algorithm with shifts.

The shifted QR algorithm:

1. Define the number of iterations as N .
2. For $k = 1, 2, \dots, N$, set $A^{(k-1)} = A$ and compute $A^{(k-1)} - \mu^{(k)} I$, we choose $\mu^{(k)}$ as the a_{nn} , the entry of matrix $A^{(k-1)}$ at the last column and last row, the bottom right entry.

3. Compute the QR decomposition

$$A^{(k-1)} - \mu^{(k)} I = Q^{(k)} R^{(k)}$$

we use QR decomposition with Householder reflection instead of Gram-Schmidt.

4. Compute matrix A for the next iteration

$$A^{(k)} = R^{(k)} Q^{(k)} + \mu^{(k)} I$$

set $A = A^{(k)}$ and go to the next iteration till finish / converge.

We will obtain all the eigenvalues at the diagonal of matrix $A^{(N)}$, after that we can use inverse iteration with shift to compute the corresponding eigenvector for each eigenvalue. The source code can be seen at [src/linearalgebra.cpp](#) at the function `(vector<complex<double>>> QRAlgorithmwithShifts(vector<vector<complex<double>>>& A, int iterations) )`.

The funny thing why we make the entry and input all in complex term, even if QR algorithm with shifts is said to only work when the matrix has real eigenvalues only, is that one day with new advancement in science it will not become a waste, there is a possibility that this algorithm / iteration can be developed so it can work to compute complex eigenvalues as well.

[SI*] If a real matrix A has complex conjugate eigenvalues, a basic QR algorithm (even with simple real shifts) will fail to converge to a purely upper triangular matrix. However, it will converge to a real quasi-upper triangular matrix (also known as the real Schur form), where complex eigenvalues are represented by 2×2 blocks on the diagonal.

If the initial matrix A is real and you only use real shifts ($s_k \in \mathbb{R}$), all subsequent matrices

A_k in the iteration

$$\begin{aligned} A_k - s_k I &= Q_k R_k \\ A_{k+1} &= R_k Q_k + s_k I \end{aligned}$$

remain real.

A matrix with complex eigenvalues cannot be similar to a real upper triangular matrix. Therefore, the normal QR algorithm cannot "converge" to a diagonal representation of these eigenvalues. The algorithm might cycle, where the matrix A_k switches between different forms without the off-diagonal elements tending to zero.

To handle complex eigenvalues in real matrices, the QR algorithm is modified into Francis double step QR algorithm, this algorithm accelerate convergence for complex pairs, it uses a "double shift" (s_1, s_2) instead of one shift (s). This allows the algorithm to work with real arithmetic while effectively handling complex eigenvalues, avoiding complex numbers altogether. Double shift QR will successfully finds both real and complex eigenvalues by converging to a real schur form, real schur form is a quasi-upper triangular matrix. In this form, complex eigenvalues appear as a 2×2 block on the diagonal

$$A_2 \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

The eigenvalues of this 2×2 block are complex conjugate pairs, solvable via the quadratic formula (solve $|\lambda I - A_2|$)

$$\begin{aligned} \lambda_1 &= \frac{-b + \sqrt{b^2 - 4ac}}{2a} \\ \lambda_2 &= \frac{-b - \sqrt{b^2 - 4ac}}{2a} \end{aligned}$$

with

$$\begin{aligned} a &= 1 \\ b &= -a_{11} - a_{22} \\ c &= (a_{11}a_{22}) - (a_{12}a_{21}) \end{aligned}$$

[SI*] Francis Double Step QR Algorithm

The QR algorithm with double shift, often called the Francis double step / double-shift QR algorithm, is a robust numerical method to compute all eigenvalues (and the Schur form) of a real non-Hermitian (nonsymmetric) matrix. It is an advanced, fast method for finding eigenvalues of real matrices, by applying two shifts simultaneously in each QR iteration to accelerate convergence, particularly for complex conjugate eigenvalues, we can use the eigenvalues of the bottom-right 2×2 block as shifts (the Francis shift) to rapidly converge to the Schur form, allowing deflation and real-arithmetic computation.

Notes: In many textbooks and sources they are mentioning about using the eigenvalues of the bottom-right 2×2 block as shifts, this could be problematic if the eigenvalues are complex, then it will lead to complex arithmetic, thus from our experience and trial in coding for this, we choose to use the algorithm that is using s, t, x, y, z from Lecture 4 ETH Zurich.

How this algorithm works:

1. Hessenberg Form

First, the matrix is reduced to upper Hessenberg form (nearly triangular) for efficiency, and it keeps the eigenvalues of the original matrix invariant.

2. Shifts

Instead of one shift, two shifts (s_1, s_2) are chosen, either the eigenvalues of the trailing 2×2 submatrix of H (bottom-right 2×2 block) as shifts or if we don't want to use eigenvalues at the bottom-right 2×2 matrix, skip this and continue on to the **Initialize "Bulge"** phase.

3. Simultaneous QR Step

A single "double shift" QR step effectively performs two single shifts at once, but keeps the computation in real arithmetic, avoiding complex numbers for real matrices.

4. Initialize "Bulge"

Instead of calculating $QR = (H - s_1 I)(H - s_2 I)$, calculate the first column of this matrix, which only has 3 non-zero entries. Determine a Householder matrix P_0 that maps this column to a multiple of e_1 .

In order to compute the first householder reflector, P_0 . Given an upper Hessenberg matrix H , calculate the first column of $G = H^2 - sH + tI$, where

$$\begin{aligned} s &= h_{n-1,n-1} + h_{n,n} \\ t &= (h_{n-1,n-1} * h_{n,n}) - (h_{n-1,n} * h_{n,n-1}) \end{aligned}$$

The first three components of $Ge_1 = x$ are

$$\begin{aligned} x &= h_{1,1}^2 + h_{1,2}h_{2,1} - sh_{1,1} + t \\ y &= h_{2,1}(h_{1,1} + h_{2,2} - s) \\ z &= h_{2,1}h_{3,2} \end{aligned}$$

this will populate the first three entries for the target vector x .

Construct the Householder vector v . Compute a reflector P_0 that maps $x = [x, y, z]^T$ to a multiple of $e_1 = [1, 0, 0]^T$ ($P_0 x = \alpha e_1$).

Let

$$x = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} h_{1,1}^2 + h_{1,2}h_{2,1} - sh_{1,1} + t \\ h_{2,1}(h_{1,1} + h_{2,2} - s) \\ h_{2,1}h_{3,2} \end{bmatrix}$$

Calculate v

$$v = x + \text{sgn}(x_1) \|x\|_2 e_1$$

If the sign of the first component of vector x is negative then $\text{sgn}(x_1) = 1$, if the sign of the first component of vector x is positive then $\text{sgn}(x_1) = -1$. Normalize v

$$v = \frac{v}{\|v\|_2}$$

Define the first Householder reflector P_0

$$P_0 = I - 2vv^T$$

given that v is already normalized. If v is not normalized yet, then use this formula

$$P_0 = I - 2 \frac{vv^T}{||v||^2}$$

5. Bulge Chasing

When we apply P_0 to H

$$H \leftarrow P_0^T H P_0$$

This creates a "bulge" (a 3×3 submatrix) below the Hessenberg form, which is then "chased" down the matrix through a series of transformations (e.g. Householder transformations / rotations or Givens rotations) until it disappears.

We use a sequence of Householder reflectors $P_1, P_2, \dots, P_{n-2}$ to eliminate this "bulge," bringing the matrix to quasi-upper triangular form / real Schur form.

6. Deflation

When the subdiagonal elements become near zero, the matrix "deflates," splitting into smaller subproblems, revealing eigenvalues.

7. Iteration

Repeat until the matrix is in real Schur form (quasi-upper triangular)

It is significantly faster than the basic QR algorithm, it handles complex eigenvalues for real matrices efficiently using only real numbers (Francis double shift), and achieve rapid (often quadratic or better) convergence.

Instead of explicitly computing the QR-decomposition of $(H - s_1 I)(H - s_2 I)$, which is expensive, the implicit Q theorem is used to achieve the same result with far less computation. By utilizing the implicit approach and working with Hessenberg matrices, each step costs $O(n^2)$ flops, rather than $O(n^3)$.

There is also another method called Multi-shift QR algorithm, it generalizes the double shift to use multiple simultaneous shifts for even faster convergence, especially on modern parallel hardware.

Algorithm 3.5 The Francis' Double step QR algorithm

```

1: Let  $H_0 = H \in \mathbb{R}^{n \times n}$  be an upper Hessenberg matrix. This algorithm computes its
   real Schur form  $H = UTU$  using the Francis double step QR algorithm.  $T$  is a quasi
   upper triangular matrix.
2:  $p := n$ ; /*  $p$  indicates the 'actual' matrix size. */
3: while  $p > 2$  do
4:    $q := p - 1$ ;
5:    $s := H_{q,q} + H_{p,p}$ ;  $t := H_{q,q}H_{p,p} - H_{q,p}H_{p,q}$ ;
6:   /* compute first 3 elements of first column of  $M$  */
7:    $x := H_{1,1}^2 + H_{1,2}H_{2,1} - sH_{1,1} + t$ ;
8:    $y := H_{2,1}(H_{1,1} + H_{2,2} - s)$ ;
9:    $z := H_{2,1}H_{3,2}$ ;
10:  for  $k = 0$  to  $p - 3$  do
11:    Determine the Householder reflector  $P$  with  $P^T [x; y; z]^T = \alpha e_1$ ;
12:     $r := \max\{1, k\}$ ;
13:     $H_{k+1:k+3, r:n} := P^T H_{k+1:k+3, r:n}$ ;
14:     $r := \min\{k + 4, p\}$ ;
15:     $H_{1:r, k+1:k+3} := H_{1:r, k+1:k+3} P$ ;
16:     $x := H_{k+2, k+1}$ ;  $y := H_{k+3, k+1}$ ;
17:    if  $k < p - 3$  then
18:       $z := H_{k+4, k+1}$ ;
19:    end if
20:  end for
21:  Determine the Givens rotation  $P$  with  $P^T [x; y]^T = \alpha e_1$ ;
22:   $H_{q:p, p-2:n} := P^T H_{q:p, p-2:n}$ ;
23:   $H_{1:p, p-1:p} := H_{1:p, p-1:p} P$ ;
24:  /* check for convergence */
25:  if  $|H_{p,q}| < \varepsilon (|H_{q,q}| + |H_{p,p}|)$  then
26:     $H_{p,q} := 0$ ;  $p := p - 1$ ;  $q := p - 1$ ;
27:  else if  $|H_{p-1, q-1}| < \varepsilon (|H_{q-1, q-1}| + |H_{q,q}|)$  then
28:     $H_{p-1, q-1} := 0$ ;  $p := p - 2$ ;  $q := p - 1$ ;
29:  end if
30: end while

```

Figure 6.21: The Francis double-step QR algorithm from ETH Zürich' lecture 4, March 14th, 2018: the QR algorithm [3].

i. Diagonalization

[SI*] A basis for $\mathbb{R}^n$ that consists of eigenvectors of an $n \times n$ matrix A can be used to study geometric properties of A and to simplify various numerical computations.

[SI*] The matrix product

$$P^{-1}AP$$

is called a similarity transformation of the matrix A . Such products are important in the study of eigenvectors and eigenvalues.

Definition 6.32: Similarity

If A and B are square matrices, then we say that B is similar to A if there is an invertible matrix P such that

$$B = P^{-1}AP$$

[SI*] Similar matrices have many properties in common. For example, if $B = P^{-1}AP$, then it

follows that A and B have the same determinant, since

$$\begin{aligned}\det(B) &= \det(P^{-1}AP) \\ &= \det(P^{-1}) \det(A) \det(P) \\ &= \frac{1}{\det(P)} \det(A) \det(P) \\ &= \det(A)\end{aligned}$$

In general, any property that is shared by all similar matrices is called a similarity invariant. The list of similarity invariants:

- (a) A and $P^{-1}AP$ have the same determinant.
- (b) A is invertible if and only if $P^{-1}AP$ is invertible.
- (c) A and $P^{-1}AP$ have the same rank.
- (d) A and $P^{-1}AP$ have the same nullity.
- (e) A and $P^{-1}AP$ have the same trace.
- (f) A and $P^{-1}AP$ have the same characteristic polynomial.
- (g) A and $P^{-1}AP$ have the same eigenvalues.
- (h) If λ is an eigenvalue of A and hence of $P^{-1}AP$, then the eigenspace of A corresponding to λ and the eigenspace of $P^{-1}AP$ corresponding to λ have the same dimension.

Definition 6.33: Diagonalizable

A square matrix A is said to be diagonalizable if it is similar to some diagonal matrix; that is, if there exists an invertible matrix P such that

$$P^{-1}AP$$

is diagonal. In this case the matrix P is said to diagonalize A .

Theorem 6.50: Diagonalizable and Eigenvectors

If A is an $n \times n$ matrix, the following statements are equivalent.

- (a) A is diagonalizable.
- (b) A has n linearly independent eigenvectors.

[SI*] Matrix A cannot always be diagonalized.

After we obtain the eigenvalues for matrix A with size of $n \times n$, we compute these matrices

$$\begin{aligned}A_1 &= \lambda_1 I - A \\ A_2 &= \lambda_2 I - A \\ &\vdots \\ A_n &= \lambda_n I - A\end{aligned}$$

We know that the eigenvectors corresponding to an eigenvalue λ_i of a matrix A are the nonzero vectors that satisfy the equation

$$(\lambda_i I - A)x = \mathbf{0}$$

these eigenvectors are the nonzero vectors in the null space of the matrix $\lambda_i I - A$. With $i, j = 1, 2, \dots, n$ and $i \neq j$, thus $\lambda_i \neq \lambda_j$, for repeated root / repeated eigenvalue it will share the same set of eigenvectors.

But, instead of computing the eigenvectors for each matrix $A_1, A_2, \dots, A_n$ we can just compute the rank of each matrix above, then we can obtain the nullity (the dimension of the null space) and find the sum of all nullity.

$$n = \text{rank}(A) + \text{nullity}(A)$$

If

$$\sum_{i=1}^n \text{nullity}(A_i) \neq n$$

then there is no matrix P that diagonalizes A .

Theorem 6.51: Linearly Independent Eigenvectors

If $v_1, v_2, \dots, v_k$ are eigenvectors of a matrix A corresponding to distinct eigenvalues, then $\{v_1, v_2, \dots, v_k\}$ is a linearly independent set.

Theorem 6.52: Distinct Eigenvalues

If an $n \times n$ matrix A has n distinct eigenvalues, then A is diagonalizable.

[SI*] We can apply diagonalization to compute high powers of a square matrix A .

If A happens to be diagonalizable, then the computations can be simplified by diagonalizing A .

Suppose that A is a diagonalizable $n \times n$ matrix, that P diagonalizes A , and that

$$P^{-1}AP = \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix} = D$$

Squaring both sides of this equation yields

$$(P^{-1}AP)^2 = \begin{bmatrix} \lambda_1^2 & 0 & \dots & 0 \\ 0 & \lambda_2^2 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \lambda_n^2 \end{bmatrix} = D^2$$

We can rewrite the left side of this equation as

$$(P^{-1}AP)^2 = P^{-1}APP^{-1}AP = P^{-1}AIA P = P^{-1}A^2P$$

from which we obtain the relationship

$$P^{-1}A^2P = D^2$$

More generally, if k is a positive integer, then a similar computation will show that

$$P^{-1}A^kP = D^k = \begin{bmatrix} \lambda_1^k & 0 & \dots & 0 \\ 0 & \lambda_2^k & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \lambda_n^k \end{bmatrix}$$

which we can rewrite as

$$A^k = PD^kP^{-1} = P \begin{bmatrix} \lambda_1^k & 0 & \dots & 0 \\ 0 & \lambda_2^k & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \lambda_n^k \end{bmatrix} P^{-1} \quad (6.71)$$

Computing the right side of this formula involves only three matrix multiplication and the powers of the diagonal entries of D . For matrices of large size and high powers of λ , this involves substantially fewer operations than computing A^k directly.

Theorem 6.53: Eigenvalues of Powers of a Matrix

If λ is an eigenvalue of a square matrix A and $\mathbf{x}$ is a corresponding eigenvector, and if k is any positive integer, then λ^k is an eigenvalue of A^k and $\mathbf{x}$ is a corresponding eigenvector.

[SI*] If we have a matrix of size 3×3 with characteristic polynomial of

$$(\lambda - 1)(\lambda - 2)^2$$

thus, the eigenspace corresponding to $\lambda = 1$ is at most one-dimensional, and the eigenspace corresponding to $\lambda = 2$ is at most two-dimensional.

If λ_0 is an eigenvalue of an $n \times n$ matrix A , then the dimension of the eigenspace corresponding to λ_0 is called the geometric multiplicity of λ_0 , and the number of times that $\lambda - \lambda_0$ appears as a factor in the characteristic polynomial of A is called the algebraic multiplicity of λ_0 .

Theorem 6.54: Geometric and Algebraic Multiplicity

If A is a square matrix, then:

- (a) For every eigenvalue of A , the geometric multiplicity is less than or equal to the algebraic multiplicity.
- (b) A is diagonalizable if and only if the geometric multiplicity of every eigenvalue is equal to the algebraic multiplicity.

ii. Complex Vector Spaces

[SI*] The characteristic equation of any square matrix can have complex solutions, the notion of complex eigenvalues and eigenvectors arise naturally, even within the context of matrices with real entries.

[SI*] If $z = a + bi$ is a complex number, then:

- $\operatorname{Re}(z) = a$ and $\operatorname{Im}(z) = b$ are called the real part of z and the imaginary part of z , respectively.
- $|z| = \sqrt{a^2 + b^2}$ is called the modulus (or absolute value) of z .
- $\bar{z} = a - bi$ is called the complex conjugate of z .
- $z\bar{z} = a^2 + b^2 = |z|^2$
- The angle ϕ is called an argument of z .
- $\operatorname{Re}(z) = |z| \cos \phi$
- $\operatorname{Im}(z) = |z| \sin \phi$
- $e^{i\phi} = \cos \phi + i \sin \phi$
- $z = |z|(\cos \phi + i \sin \phi)$ is called the polar form of z .

[SI*] It is possible for the characteristic equation of a matrix A with real entries to have imaginary solutions, for example

$$A = \begin{bmatrix} -2 & -1 \\ 5 & 2 \end{bmatrix}$$

$$p(\lambda) = \det(\lambda I - A) = \lambda^2 + 1 = 0$$

the characteristic polynomial above has solutions

$$\lambda^2 + 1 = 0$$

$$\lambda = \pm i$$

[SI*] A vector space in which scalars are allowed to be complex numbers is called a complex vector space.

Definition 6.34: Complex Vector Space

If n is a positive integer, then a complex n -tuple is a sequence of n complex numbers $(v_1, v_2, \dots, v_n)$. The set of all complex n -tuples is called complex n -space and is denoted by $\mathbb{C}^n$. Scalars are complex numbers, and the operations of addition, subtraction, and scalar multiplication are performed componentwise.

[SI*] If $v_1, v_2, \dots, v_n$ are complex numbers, then we call $\mathbf{v} = (v_1, v_2, \dots, v_n)$ a vector in $\mathbb{C}^n$ and $v_1, v_2, \dots, v_n$ its components. Some examples of vectors in $\mathbb{C}^3$ are

$$\mathbf{u} = (1 + 8i, 5, -3i)$$

$$\mathbf{v} = (1, 0, 2i)$$

$$\mathbf{w} = (\pi i, 6 - \sqrt{10}i, \frac{1}{2}i)$$

Every vector

$$\mathbf{v} = (v_1, v_2, \dots, v_n) = (a_1 + b_1i, a_2 + b_2i, \dots, a_n + b_ni)$$

in $\mathbb{C}^n$ can be split into real and imaginary parts as

$$\mathbf{v} = (a_1, a_2, \dots, a_n) + i(b_1, b_2, \dots, b_n)$$

which can be denoted as

$$v = \operatorname{Re}(v) + i \operatorname{Im}(v)$$

where

$$\operatorname{Re}(v) = (a_1, a_2, \dots, a_n)$$

$$\operatorname{Im}(v) = (b_1, b_2, \dots, b_n)$$

The vector

$$\bar{v} = (\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n) = (a_1 - b_1i, a_2 - b_2i, \dots, a_n - b_ni)$$

is called the complex conjugate of v and can be expressed in terms of $\operatorname{Re}(v)$ and $\operatorname{Im}(v)$ as

$$\bar{v} = (a_1, a_2, \dots, a_n) - i(b_1, b_2, \dots, b_n) = \operatorname{Re}(v) - i\operatorname{Im}(v) \quad (6.72)$$

It follows that the vectors in $\mathbb{R}^n$ can be viewed as those vectors in $\mathbb{C}^n$ whose imaginary part is zero.

A vector v in $\mathbb{C}^n$ is in $\mathbb{R}^n$ if and only if

$$\bar{v} = v$$

[SI*] We call a matrix A a real matrix if its entries are required to be real numbers and a complex matrix if its entries are allowed to be complex numbers. The standard operations on real matrices carry over to complex matrices without change, and all of familiar properties of matrices continue to hold.

If A is a complex matrix, then $\operatorname{Re}(A)$ and $\operatorname{Im}(A)$ are the matrices formed from the real and imaginary parts of the entries of A , and the $\bar{A}$ is the matrix formed by taking the complex conjugate of each entry in A .

Theorem 6.55: Algebraic Properties of the Complex Conjugate for Vectors

If u and v are vectors in $\mathbb{C}^n$, and if k is a scalar, then:

- (a) $\overline{\bar{u}} = u$
- (b) $\overline{k u} = k \bar{u}$
- (c) $\overline{u + v} = \bar{u} + \bar{v}$
- (d) $\overline{u - v} = \bar{u} - \bar{v}$

Theorem 6.56: Algebraic Properties of the Complex Conjugate for Matrices

If A is an $m \times k$ complex matrix and B is a $k \times n$ complex matrix, then:

- (a) $\overline{\bar{A}} = A$
- (b) $\overline{(A^T)} = (\bar{A})^T$
- (c) $\overline{AB} = \bar{A}\bar{B}$

Definition 6.35: Complex Euclidean Inner Product

If $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ are vectors in $\mathbb{C}^n$, then the complex Euclidean inner product of $\mathbf{u}$ and $\mathbf{v}$ (also called complex dot product) is denoted by $\mathbf{u} \cdot \mathbf{v}$ and is defined as

$$\mathbf{u} \cdot \mathbf{u} = u_1 \overline{v_1} + u_2 \overline{v_2} + \dots + u_n \overline{v_n} + \quad (6.73)$$

We also define the Euclidean norm on $\mathbb{C}^n$ to be

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{|v_1|^2 + |v_2|^2 + \dots + |v_n|^2} \quad (6.74)$$

As in the real case, we call $\mathbf{v}$ a unit vector in $\mathbb{C}^n$ if $\|\mathbf{v}\| = 1$, and we say two vectors $\mathbf{u}$ and $\mathbf{v}$ are orthogonal if $\mathbf{u} \cdot \mathbf{v} = 0$.

[SI*] If $\mathbf{u}$ and $\mathbf{v}$ are column vectors in $\mathbb{R}^n$, then their dot product can be expressed as

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v} = \mathbf{v}^T \mathbf{u}$$

The analogous formulas in $\mathbb{C}^n$ are

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \overline{\mathbf{v}} = \overline{\mathbf{v}}^T \mathbf{u} \quad (6.75)$$

Theorem 6.57: Properties of Complex Euclidean Inner Product

If $\mathbf{u}, \mathbf{v}$, and $\mathbf{w}$ are vectors in $\mathbb{C}^n$, and if k is a scalar, then the complex Euclidean inner product has the following properties:

- (a) $\mathbf{u} \cdot \mathbf{v} = \overline{\mathbf{v} \cdot \mathbf{u}}$ (Antisymmetry property)
- (b) $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$ (Distributive property)
- (c) $k(\mathbf{u} \cdot \mathbf{v}) = (k\mathbf{u}) \cdot \mathbf{v}$ (Homogeneity property)
- (d) $\mathbf{u} \cdot k\mathbf{v} = \overline{k}(\mathbf{u} \cdot \mathbf{v})$ (Antihomogeneity property)
- (e) $\mathbf{v} \cdot \mathbf{v} \geq 0$ and $\mathbf{v} \cdot \mathbf{v} = 0$ if and only if $\mathbf{v} = \mathbf{0}$ (Positivity property)

[SI*] Except for the use of complex scalars, the notions of linear combination, linear independence, subspace, spanning, basis, and dimension carry over without change to $\mathbb{C}^n$.

[SI*] Eigenvalues and eigenvectors are defined for complex matrices exactly as for real matrices.

If A is an $n \times n$ matrix with complex entries, then the complex roots of the characteristic equation

$$\det(\lambda I - A) = 0$$

are called complex eigenvalues of A .

As in the real case, λ is a complex eigenvalue of A if and only if there exists a nonzero vector $\mathbf{x}$ in $\mathbb{C}^n$ such that

$$A\mathbf{x} = \lambda\mathbf{x}$$

Each such x is called a complex eigenvector of A corresponding to λ .

The complex eigenvectors of A corresponding to λ are the nonzero solutions of the linear system

$$(\lambda I - A)x = \mathbf{0}$$

and the set of all such solutions is a subspace of $\mathbb{C}^n$, called the eigenspace of A corresponding to λ .

Theorem 6.58: Conjugate Pairs in Eigenvectors and Eigenvalues

If λ is an eigenvalue of a real $n \times n$ matrix A , and if x is a corresponding eigenvector, then $\bar{\lambda}$ is also an eigenvalue of A , and $\bar{x}$ is a corresponding eigenvector.

Theorem 6.59: Determining Eigenvalues of a 2×2 Matrix

If A is a 2×2 matrix with real entries, then the characteristic equation of A is

$$\lambda^2 - \text{tr}(A)\lambda + \det(A) = 0 \quad (6.76)$$

and

- (a) A has two distinct real eigenvalues if $\text{tr}(A)^2 - 4\det(A) > 0$;
- (b) A has one repeated real eigenvalue if $\text{tr}(A)^2 - 4\det(A) = 0$;
- (c) A has two complex conjugate eigenvalues if $\text{tr}(A)^2 - 4\det(A) < 0$

Theorem 6.60: Symmetric Matrices Have Real Eigenvalues

If A is a real symmetric matrix, then A has real eigenvalues.

Theorem 6.61: Geometric Interpretation of Complex Eigenvalues

The eigenvalues of the real matrix

$$C = \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \quad (6.77)$$

are $\lambda = a \pm bi$. If a and b are not both zero, then this matrix can be factored as

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} = \begin{bmatrix} |\lambda| & 0 \\ 0 & |\lambda| \end{bmatrix} \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \quad (6.78)$$

where ϕ is the angle from the positive x -axis to the ray that joins the origin to the point (a, b) .

[SI*] Geometrically, a multiplication by a matrix of form

$$C = \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \quad (6.79)$$

can be viewed as a rotation through the angle ϕ followed by a scaling with factor $|\lambda|$. Since the characteristic equation of C is

$$(\lambda - a)^2 + b^2 = 0$$

from which it follows that the eigenvalues of C are

$$\lambda = a \pm bi$$

Assuming that a and b are not both zero, let ϕ be the angle from the positive x -axis to the ray that joins the origin to the point (a, b) . The angle ϕ is an argument of the eigenvalue $\lambda = a + bi$, therefore

$$\begin{aligned} a &= |\lambda| \cos \phi \\ b &= |\lambda| \sin \phi \end{aligned}$$

It follows from this that the matrix in (6.68) can be written as

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} = \begin{bmatrix} |\lambda| & 0 \\ 0 & |\lambda| \end{bmatrix} \begin{bmatrix} \frac{a}{|\lambda|} & -\frac{b}{|\lambda|} \\ \frac{b}{|\lambda|} & \frac{a}{|\lambda|} \end{bmatrix} = \begin{bmatrix} |\lambda| & 0 \\ 0 & |\lambda| \end{bmatrix} \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix}$$

Theorem 6.62: Similar Matrix to Real Matrix with Complex Eigenvalues

Let A be a real 2×2 matrix with complex eigenvalues $\lambda = a \pm bi$ (where $b \neq 0$). If x is an eigenvector of A corresponding to $\lambda = a - bi$, then the matrix

$$P = [\operatorname{Re}(x) \quad \operatorname{Im}(x)]$$

is invertible and

$$A = P \begin{bmatrix} a & -b \\ b & a \end{bmatrix} P^{-1}$$

[SI*] Let

$$A = PCP^{-1} = PSR_\theta P^{-1} = P \begin{bmatrix} |\lambda| & 0 \\ 0 & |\lambda| \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} P^{-1} \quad (6.80)$$

If we now view P as the transition matrix from the basis $B = \{\operatorname{Re}(x), \operatorname{Im}(x)\}$ to the standard basis then computing a product Ax_0 can be broken down into a three-step process:

1. Map x_0 from standard coordinates into B -coordinates by forming the product $P^{-1}x_0$.
2. Rotate and scale the vector $P^{-1}x_0$ by forming the product $SR_\theta P^{-1}x_0$.
3. Map the rotated and scaled vector back to standard coordinates to obtain $Ax_0 = PSR_\theta P^{-1}x_0$.

iii. Differential Equations

[SI*] Differential equations are equations involving functions and their derivatives. Linear algebra, eigenvalues and eigenvectors can be applied to solving systems of differential equations.

[SI*] A differential equation is an equation involving unknown functions and their derivatives. The order of a differential equation is the order of the highest derivative it contains. The simplest differential equations are the first-order equations of the form

$$y' = ay \quad (6.81)$$

where $y = f(x)$ is an unknown differentiable function to be determined, the basic definition of derivative of a function $y = f(x)$ with respect to x is

$$y' = \frac{dy}{dx}$$

where a is a constant. As with most differential equations, this equation has infinitely many solutions; they are the functions of the form

$$y = ce^{ax} \tag{6.82}$$

where c is an arbitrary constant, we call this function $y = ce^{ax}$ as the general solution of $y' = ay$. Below is the step to obtain this general solution

$$\begin{aligned} y' &= ay \\ \frac{dy}{dx} &= ay \\ \frac{dy}{y} &= adx \\ \int \frac{dy}{y} &= \int adx \\ \ln |y| &= ax + c \\ e^{\ln |y|} &= e^{ax+c} \\ y &= ce^{ax} \end{aligned}$$

Thus every function of this form is a solution of $y' = ay$, if we substitute y then

$$y' = cae^{ax} = ay$$

[SI*] Often a physical problem that leads to a differential equation imposes some conditions that enable us to isolate one particular solution from the general solution.

For example, if we have a differential equation

$$y' = 5y$$

the general solution is

$$y = ce^{5x}$$

now, there is another condition, that is called initial condition that can be represented with this equation

$$y(0) = 6$$

it means $y = 6$ when $x = 0$, then we substitute the initial condition into $y = ce^{5x}$ to obtain the constant c

$$\begin{aligned} y &= ce^{5x} \\ y(0) &= ce^{5(0)} \\ 6 &= ce^0 \\ c &= 6 \end{aligned}$$

so with this initial condition we are able to obtain a single line, or a single solution for certain condition, instead of infinitely many solutions, that is

$$y = 6e^{5x}$$

the equation above is the only solution for the differential equation $y' = 5y$ with initial condition $y(0) = 6$.

The problem of solving a differential equation subject to an initial condition is called an initial-value problem.

[SI*] We will be concerned with solving systems of differential equations of the form

$$\begin{aligned} y_1' &= a_{11}y_1 + a_{12}y_2 + \cdots + a_{1n}y_n \\ y_2' &= a_{21}y_1 + a_{22}y_2 + \cdots + a_{2n}y_n \\ &\vdots \\ y_n' &= a_{n1}y_1 + a_{n2}y_2 + \cdots + a_{nn}y_n \end{aligned} \quad (6.83)$$

where $y_1 = f_1(x)$, $y_2 = f_2(x)$, $\dots$, $y_n = f_n(x)$ are functions to be determined, and the a_{ij} are constants. In matrix notation, we can write it as

$$\begin{bmatrix} y_1' \\ y_2' \\ \vdots \\ y_n' \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$$

or, more briefly as

$$\mathbf{y}' = A\mathbf{y} \quad (6.84)$$

where the notation $\mathbf{y}'$ denotes the vector obtained by differentiating each component of $\mathbf{y}$.

[SI*] The basic idea for solving a system

$$\mathbf{y}' = A\mathbf{y}$$

whose coefficient matrix is not diagonal is to introduce a new unknown vector $\mathbf{u}$ that is related to the unknown vector $\mathbf{y}$ by an equation of the form $\mathbf{y} = P\mathbf{u}$ in which P is an invertible matrix that diagonalizes A . Of course, such a matrix may or may not exist, but if it does then we can rewrite the equation $\mathbf{y}' = A\mathbf{y}$ as

$$P\mathbf{u}' = A(P\mathbf{u})$$

[SI*] If A a square matrix of size $n \times n$ is diagonalizable and

$$\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$$

is a solution of the system $\mathbf{y}' = A\mathbf{y}$, then there will exists a matrix P such that

$$P = [p_1 \ p_2 \ \cdots \ p_n]$$

where $\mathbf{p}_i$ is an eigenvector corresponding to λ_i . Matrix P diagonalizes A , and

$$D = P^{-1}AP = \begin{bmatrix} \lambda_1 & 0 & 0 & \dots & 0 \\ 0 & \lambda_2 & 0 & \dots & 0 \\ 0 & 0 & \lambda_3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & \lambda_n \end{bmatrix}$$

Then we make the substitution

$$\mathbf{y} = P\mathbf{u}$$

$$\mathbf{y}' = P\mathbf{u}'$$

that will yield the diagonal system

$$\mathbf{u}' = D\mathbf{u}$$

or

$$\begin{aligned} u_1' &= \lambda_1 u_1 \\ u_2' &= \lambda_2 u_2 \\ &= \vdots \\ u_n' &= \lambda_n u_n \end{aligned}$$

the solution of this system is

$$\begin{aligned} u_1 &= c_1 e^{\lambda_1 x} \\ u_2 &= c_2 e^{\lambda_2 x} \\ &= \vdots \\ u_n &= c_n e^{\lambda_n x} \end{aligned}$$

the equation $\mathbf{y} = P\mathbf{u}$ will be

$$\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = [\mathbf{p}_1 \ \mathbf{p}_2 \ \dots \ \mathbf{p}_n] \begin{bmatrix} c_1 e^{\lambda_1 x} \\ c_2 e^{\lambda_2 x} \\ \vdots \\ c_n e^{\lambda_n x} \end{bmatrix} = \mathbf{p}_1 c_1 e^{\lambda_1 x} + \mathbf{p}_2 c_2 e^{\lambda_2 x} + \dots + \mathbf{p}_n c_n e^{\lambda_n x}$$

so the general solution of the system is a linear combination of $e^{\lambda_1 x}, e^{\lambda_2 x}, \dots, e^{\lambda_n x}$ where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of A .

A little note here, The vector column for matrix P that we sometimes wrote $\mathbf{p}_i$ is another notation for $\mathbf{x}_i$ the eigenvector of A corresponding to λ_i .

$$P = [\mathbf{p}_1 \ \mathbf{p}_2 \ \dots \ \mathbf{p}_n] = [\mathbf{x}_1 \ \mathbf{x}_2 \ \dots \ \mathbf{x}_n]$$

Theorem 6.63: Diagonalizable Matrix A and the General Solution

If the coefficient matrix A of the system $\mathbf{y}' = A\mathbf{y}$ is diagonalizable, then the general solution of the system can be expressed as

$$\mathbf{y} = c_1 e^{\lambda_1 x} \mathbf{x}_1 + c_2 e^{\lambda_2 x} \mathbf{x}_2 + \cdots + c_n e^{\lambda_n x} \mathbf{x}_n \quad (6.85)$$

where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of A , and $\mathbf{x}_i$ is an eigenvector of A corresponding to λ_i .

[SI*] Example:

Consider the system of differential equation $\mathbf{y}' = A\mathbf{y}$, where A is a 2×2 matrix. For what values of $a_{11}, a_{12}, a_{21}, a_{22}$ do the component solutions $y_1(t), y_2(t)$ tend to zero as $t \rightarrow \infty$? In particular, what must be true about the determinant and the trace of A for this to happen?

Solution:

From the theorem we know that

$$\mathbf{y} = c_1 e^{\lambda_1 t} \mathbf{t}_1 + c_2 e^{\lambda_2 t} \mathbf{t}_2 \quad (6.86)$$

where λ_1, λ_2 are the eigenvalues of A , and $\mathbf{t}_i$ is an eigenvector of A corresponding to λ_i . Now if we take a look at the eigenvector components it will be like this

$$\mathbf{t}_1 = \begin{bmatrix} t_{11} \\ t_{12} \end{bmatrix}, \quad \mathbf{t}_2 = \begin{bmatrix} t_{21} \\ t_{22} \end{bmatrix}$$

thus

$$\begin{aligned} y_1(t) &= c_1 e^{\lambda_1 t} t_{11} + c_2 e^{\lambda_2 t} t_{21} \\ y_2(t) &= c_1 e^{\lambda_1 t} t_{12} + c_2 e^{\lambda_2 t} t_{22} \end{aligned}$$

when $t \rightarrow \infty$, $y_1(t)$ and $y_2(t)$ will tend to zero if and only if

$$\lambda_1 < 0 \quad \text{and} \quad \lambda_2 < 0$$

Now, we will check the condition for the eigenvalues λ_1 and λ_2 with the formula

$$\det(\lambda I - A) = 0$$

$$\begin{vmatrix} \lambda - a_{11} & -a_{12} \\ -a_{21} & \lambda - a_{22} \end{vmatrix} = \lambda^2 - (a_{11} + a_{22})\lambda + (a_{11}a_{22} - a_{12}a_{21}) = 0$$

The eigenvalues formula for the characteristic equation above is

$$\lambda_{1,2} = \frac{(a_{11} + a_{22}) \pm \sqrt{(a_{11} + a_{22})^2 - 4(a_{11}a_{22} - a_{12}a_{21})}}{2(1)}$$

we know that

$$\begin{aligned} \text{tr}(A) &= a_{11} + a_{22} \\ \det(A) &= a_{11}a_{22} - a_{12}a_{21} \end{aligned}$$

we can substitute into the eigenvalues formula

$$\lambda_{1,2} = \frac{\text{tr}(A) \pm \sqrt{(\text{tr}(A))^2 - 4(1)(\det(A))}}{2(1)}$$

$$\lambda_{1,2} = \frac{\text{tr}(A) \pm \sqrt{(\text{tr}(A))^2 - 4(\det(A))}}{2}$$

in order to obtain negative eigenvalues then the conditions need to be fulfilled are

$$\begin{aligned} (\text{tr}(A))^2 - 4(\det(A)) &\geq 0 \\ \text{tr}(A) &< 0 \\ \text{tr}(A) + \sqrt{(\text{tr}(A))^2 - 4(\det(A))} &< 0 \\ \text{tr}(A) - \sqrt{(\text{tr}(A))^2 - 4(\det(A))} &< 0 \end{aligned}$$

if the 4 conditions above are met, then as $t \rightarrow \infty$, $y_1(t)$ and $y_2(t)$ will go to zero.

iv. Functions in SymIntegration Related to Eigenvalues and Eigenvectors

[SI*] We will list the basic functions that can be used related to eigenvalues and eigenvectors:

GaussJordanComplexMatrix(vector<vector<complex<double>>> &A)

to reduce complex matrix A into reduced row echelon form with the Gauss-Jordan elimination, the input is a complex matrix, so it can handle real matrix (complex matrix input with no imaginary part in the entries) and complex matrix (complex matrix input with imaginary part in the entries) as well.

PermutationMatrixMax(vector<double>& vectorx)

to returns permutation matrix P that will swap the first row of vector x with the row that contain the maximum / largest value. P initially is an identity matrix, then the first row of P will be swapped with the row of P that has the maximum value in vector x . The permutation matrix P multiply vector x will swap two elements in vector x thus making the element with largest value as the first element / first row for x .

penrose(vector<double>& vectorx, vector<double>& vectory)

to compute the matrix M such that $Mx = y$ with Moore-Penrose pseudoinverse method.

PowerMethod(vector<vector<double>>& A, int N)

to compute the largest eigenvalue and its corresponding eigenvector of a real square matrix A , N is the maximum number of iteration for the numerical method that will be used for the computation.

EigenvaluesEigenvectorsApproximation(vector<vector<double>>& A, int N)

to compute all the eigenvalues and eigenvectors of real square matrix A , N is the maximum number of iteration for the numerical method that will be used for the computation.

RayleighQuotientIteration(vector<vector<double>>& A, int N, double a, double b)

to compute an eigenvalue and the corresponding eigenvector of real square matrix with Rayleigh quotient iteration with A as the real square matrix, N is the number of iteration, a is

the lower bound for uniform distribution, b is the upper bound for the uniform distribution, we are using uniform distribution to generate random μ and σ as the input to generate the initial vector v^0 that is normally distributed with mean μ and standard deviation σ , this initial vector is a prerequisite for the Rayleigh quotient iteration. The key here is to choose a and b carefully so we can pinpoint the range where the eigenvalue could be located.

ComplexRayleighQuotientIteration(vector<vector<complex<double>>> &A, int N_{Rayleigh} , int N double a , double b)

to compute all the real or complex eigenvalues and the corresponding eigenvector of real square matrix or complex square matrix with A as the matrix that we want to compute the eigenvalues and the corresponding eigenvector, N_{Rayleigh} is the maximum number of iteration for the Rayleigh quotient iteration, N is the maximum number of the outermost iteration till we obtain all the correct eigenvalues, a is the lower bound for uniform distribution, b is the upper bound for the uniform distribution, we are using uniform distribution to generate random μ and σ as the input to generate the initial complex vector v^0 that is normally distributed with mean μ and standard deviation σ , this initial vector is a prerequisite for the Rayleigh quotient iteration. The key here is to choose a and b carefully so we can pinpoint the range where the eigenvalue could be located.

QRAlgorithmwithShifts(vector<vector<complex<double>>> &A, int N)

to compute the eigenvalues and its corresponding eigenvector, given that the matrix has only real eigenvalues with A as the matrix, and N as the number of iterations we want to use till the matrix converge.

diagonalization(vector<vector<double>>& A)

to compute the matrix P that diagonalizes matrix A and also to compute $P^{-1}AP$.

printComplexMatrix(const ComplexMatrix &A)

to print complex matrix A .

printComplexVector(const ComplexVector &x)

to print complex vector x .

vector<vector<double>> ComplextoRealMatrix(const ComplexMatrix &A)

to convert complex matrix A into real matrix, we only take the real argument of $A(\text{re}(A))$.

complex<double> complexdivision(complex<double> z , double k)

to divide a complex number z by a scalar k .

vector<complex<double>> complexvecrand_normal(double μ , double σ , int n)

to produce a vector of random complex number of length n that has normal distribution with mean μ and standard deviation σ .

vector<complex<double>> complexvecrand_uniform(double a , double b , int)

to produce a vector of random complex number of length n that has uniform distribution with range $[a, b]$.

complex<double> conjugate(complex<double> y)

to return the conjugate of a complex number $y = a + bi$.

double complexnorm (const ComplexVector &x)

to compute the norm of a complex vector x .

complex<double> complexdotproduct (const ComplexVector &x, const ComplexVector &y)

to compute the complex dot product of two complex vectors x and y , denoted by $x \cdot y$.

ComplexVector addComplexVectors(const ComplexVector &x, const ComplexVector &y)

to perform add operation on two complex vectors x and y .

ComplexVector subtractComplexVectors(const ComplexVector &x, const ComplexVector &y)

to perform subtraction operation on two complex vectors x and y .

ComplexVector scalarmultiplicationComplexVector(const ComplexVector &x, double k)

to perform a scalar multiplication on complex vector x with scalar k .

ComplexVector complexnumbermultiplicationComplexVector(const ComplexVector &x, complex<double> z)

to perform a complex number multiplication on complex vector x with $z = a + bi$.

getcolumnComplexMatrix(const ComplexMatrix& A, int i)

to obtain the i -th column of complex matrix A .

ComplexMatrix addComplexMatrices(const ComplexMatrix &A, const ComplexMatrix &B)

to perform add operation on two complex matrices A and B .

ComplexMatrix subtractComplexMatrices(const ComplexMatrix &A, const ComplexMatrix &B)

to perform subtraction operation on two complex matrices A and B .

ComplexMatrix multiplyComplexMatrices(const ComplexMatrix &A, const ComplexMatrix &B)

to perform multiplication operation on two complex matrices A and B .

ComplexMatrix scalarmultiplicationComplexMatrix(const ComplexMatrix &A, double k)

to perform a scalar multiplication on a complex matrix A with scalar k .

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Complex Vector Spaces ]# ./main
Matrix A:
      (1,2)      (3,-1)
      (0,-1)      (4,5)

Matrix B:
      (2,0)      (1,3)
      (-1,4)      (0,-2)

Vector x:
      (1,1)
      (0,1)
      (3,-1)

Vector y:
      (1,1)
      (2,0)
      (0,4)

x . y = (-2,-10)
y . x = (-2,10)

|| x || = 3.61
|| y || = 4.69

x + y :
      (2,2)
      (2,1)
      (3,3)

x - y :
      (0,0)
      (-2,1)
      (3,-5)

x * 3 :
      (3,3)
      (0,3)
      (9,-3)

A + B:
      (3,2)      (4,2)
      (-1,3)      (4,3)

A - B:
      (-1,2)      (2,-4)
      (1,-5)      (4,7)

A * B:
      (3,17)      (-7,-1)
      (-24,9)      (13,-9)

A * 3:
      (3,6)      (9,-3)
      (0,-3)      (12,15)

Time taken by function: 1382 microseconds

```

Figure 6.22: The computation related to complex number, complex vectors and matrices. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Complex Vector Spaces/main.cpp*).

ComplexMatrix complexnumbermultiplicationComplexMatrix(const ComplexMatrix &A, complex<double> z)

to perform a multiplication between a complex matrix A with a complex number $z = a + bi$.

ComplexVector multiplycomplexmatrixvector(const ComplexMatrix &A, const ComplexVector &x)

to perform multiplication operation between a complex matrix A and a complex vector x .

ComplexMatrix createSubMatrix(vector<vector<complex<double>>> A, int i_0 , int i_1 , int j_0 , int j_1)

to create a sub matrix from matrix A from row $i_0 + 1$ to row $i_1 + 1$ and column $j_0 + 1$ to

column $j_1 + 1$, always remember that index of matrix and vector in C++ starts from zero.

ComplexMatrix createIdentityComplexMatrix(int n)

to create a size n identity complex matrix.

ComplexMatrix TransposeComplexMatrix(ComplexMatrix &A)

to return the transpose of a complex matrix A .

ComplexMatrix ComplexMatrixInverse(ComplexMatrix &A)

to compute the inverse of a complex matrix complex matrix A .

IVPSolution_firstorderdiffeq(vector<vector<complex<double>>> &A, vector<complex<double>> &x, double a, double b)

to compute the solution satisfying the initial condition for the system of first-order differential equations with A as the matrix that represents the system of first-order differential equations, x as the vector that represents the initial value $y_i(0)$, a is the lower bound for uniform distribution, b is the upper bound for the uniform distribution, we are using uniform distribution to generate random μ and σ as the input to generate the initial complex vector v^0 that is normally distributed with mean μ and standard deviation σ , this initial vector is a prerequisite for the Rayleigh quotient iteration. The key here is to choose a and b carefully so we can pinpoint the range where the eigenvalue could be located.

FrancisDoubleStepQRReal(vector<vector<complex<double>>>& A, int N)

to compute the real and complex eigenvalues from real matrix with Francis double step QR algorithm using real arithmetic in all the iterations.

FrancisDoubleStepQR(vector<vector<complex<double>>>& A, int N)

to compute the real and complex eigenvalues from real (or complex) matrix with Francis double step QR algorithm using complex arithmetic in all the iterations.

v. Compute Moore-Penrose Pseudoinverse with SymIntegration

In eigenvalue and eigenvector computation, when we use power method we can only compute the largest eigenvalue, to compute other eigenvalues, then we need to deflate the matrix, that is why we need to know about this Moore-Penrose pseudoinverse.

Suppose you know vector v and the identity vector e_1 , and you want to compute matrix M such that

$$Mv = e_1$$

Solution:

One simple way is to use

$$M = e_1 \cdot v^+$$

where v^+ is the Moore-Penrose pseudoinverse of v (treated as a column vector).

For a non-zero column vector v :

$$v^+ = \frac{v^T}{v^T v}$$

where $v^T v$ is the dot product of v and itself.

So

$$M = \frac{e_1 v^T}{v^T v}$$

The matrix will map v to e_1 .

Now, if we want to use an example with vector

$$v = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix}$$

and

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

thus

$$M = \frac{e_1 v^T}{v^T v}$$

$$M = \frac{1}{v \cdot v} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \begin{bmatrix} 5 & 6 & 7 \end{bmatrix} = \frac{1}{110} \begin{bmatrix} 5 & 6 & 7 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0.045455 & 0.54545 & 0.063636 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$(v \cdot v = v^T v)$$

In SymIntegration to obtain the matrix M we can use this function:
penrose(vector<double>& vectorx, vector<double>& vectory)

vi. Compute Real Eigenvalues and Its Corresponding Eigenvectors with SymIntegration

Find the eigenvalues and eigenvectors of

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 4 & -17 & 8 \end{bmatrix}$$

Solution:

The characteristic polynomial of A is

$$\det(\lambda I - A) = \det \begin{bmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -1 \\ -4 & 17 & \lambda - 8 \end{bmatrix} = \lambda^3 - 8\lambda^2 + 17\lambda - 4$$

The eigenvalues of A must therefore satisfy the cubic equations

$$\lambda^3 - 8\lambda^2 + 17\lambda - 4 = 0$$

To solve this equation, we will begin by searching for integer solutions. This task can be simplified by exploiting the fact that all integer solutions (if there are any) of a polynomial equation with integer coefficients

$$\lambda^n + c_1\lambda^{n-1} + \cdots + c_n = 0$$

must be divisors of the constant term, c_n . Thus, the only possible integer solutions are the divisors of -4 , that is $\pm 1, \pm 2, \pm 4$. Successively substituting these values in the cubic equations / characteristic equation shows that $\lambda = 4$ is an integer solution. As a consequence $\lambda - 4$ must be a factor of the left side of the characteristic equation. Dividing $\lambda - 4$ into $\lambda^3 - 8\lambda^2 + 17\lambda - 4$ shows that the characteristic equation can be rewritten as

$$(\lambda - 4)(\lambda^2 - 4\lambda + 1) = 0$$

Thus, the remaining solutions satisfy the quadratic equation

$$\lambda^2 - 4\lambda + 1 = 0$$

which can be solved by the quadratic formula. Thus, the eigenvalues of A are

$$\lambda = 4$$

$$\lambda = 2 + \sqrt{3}$$

$$\lambda = 2 - \sqrt{3}$$

It could be harder if we have square matrix of size 4×4 or more, thus we would opt for computing eigenvalues and eigenvectors with numerical method.

In SymIntegration, this function can be used to compute all the eigenvalues and eigenvectors of real square matrix:

EigenvaluesEigenvectorsApproximation(vector<vector<double>>& **A**, int **N**)

with A as the matrix that we want to compute the eigenvalues and eigenvectors, N is the number of iterations for the numerical methods that we are going to deploy. The numerical methods are: the power method (as the initial / start to compute the largest eigenvalue and its corresponding eigenvector), the deflation method (to compute all the eigenvalues), and last is the inverse iteration with shift (to compute all the eigenvectors, given the eigenvalues are known).

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation with Power Method, Deflation Method and Inverse Iterati
on with Shift ]# ./main

A:
      0.000000      1.000000      0.000000
      0.000000      0.000000      1.000000
      4.000000     -17.000000      8.000000

Eigenvalue(1) = 4.008940

Corresponding Eigenvector:
-0.060523
-0.242091
-0.968364

Eigenvalue(2) = 3.721781

Corresponding Eigenvector:
0.069184
0.258199
0.963611

Eigenvalue(3) = 0.267949

Corresponding Eigenvector:
0.963611
0.258199
0.069184

Eigenvectors (in column vectors):
-0.060523      0.069184      0.963611
-0.242091      0.258199      0.258199
-0.968364      0.963611      0.069184

Time taken by function: 3375 microseconds
```

Figure 6.23: The computation to approximate all eigenvalues and eigenvectors of matrix A . (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Eigenvalues Approximation with Power Method, Deflation Method and Inverse Iteration with Shift/main.cpp*).

As a comparison we try another matrix, suppose we have a matrix A

$$A = \begin{bmatrix} -4 & 12 & -12 \\ 0 & 5 & -6 \\ -2 & -2 & 10 \end{bmatrix}$$

We try to approximate the eigenvalues and eigenvectors with **Eigen** and **SymIntegration** to compare the computation speed / computation cost. Both have the same results, only **Eigen** has faster time due to its only have header library. But the compiling time takes quite a long time on our laptop for **Eigen** compared to **SymIntegration**.

V. Eigenvalues and Eigenvectors

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation with Power Method, Deflation Method and Inverse Iterati
on with Shift ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation with Power Method, Deflation Method and Inverse Iterati
on with Shift ]# ./main

A:
      -4.000000      12.000000     -12.000000
      0.000000      5.000000      -6.000000
      -2.000000     -2.000000     10.000000

Eigenvalue(1) = 13.678780

Corresponding Eigenvector:
0.686585
0.413453
-0.598045

Eigenvalue(2) = -4.678780

Corresponding Eigenvector:
-0.985014
-0.090874
-0.146591

Eigenvalue(3) = 2.000000

Corresponding Eigenvector:
0.666667
0.666667
0.333333

Eigenvectors (in column vectors):
      0.686585      -0.985014      0.666667
      0.413453      -0.090874      0.666667
      -0.598045     -0.146591      0.333333

Time taken by function: 3433 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation with Power Method, Deflation Method and Inverse Iterati
on with Shift ]# 

```

Figure 6.24: The computation to approximate all eigenvalues and eigenvectors of matrix A with SymIntegration with SymIntegration. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Eigenvalues Approximation with Power Method, Deflation Method and Inverse Iteration with Shift/main.cpp).

```

g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation/Compute Eigenvalues and Eigenvectors with Eigen ]# ./ma
n
Matrix A:
-4 12 -12
 0 5 -6
-2 -2 10

Eigenvalues:
λ_1 = -4.67878 + 0i
λ_2 = 13.6788 + 0i
λ_3 = 2 + 0i

Eigenvectors:
x[λ_1] = (-0.985014 + 0i, -0.0908736 + 0i, -0.146591 + 0i )
x[λ_2] = (0.686585 + 0i, 0.413453 + 0i, -0.598045 + 0i )
x[λ_3] = (0.666667 + 0i, 0.666667 + 0i, 0.333333 + 0i )

Time taken by function: 3107 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation/Compute Eigenvalues and Eigenvectors with Eigen ]# 

```

Figure 6.25: The computation to approximate all eigenvalues and eigenvectors of matrix A with Eigen. (SymIntegration/Examples/Linear Algebra/Compute Eigenvalues and Eigenvectors with Eigen/main.cpp).

We also want to test Rayleigh quotient iteration to compute the eigenvalue and the corresponding eigenvector, this method is said to converge fast, but we need to run it multiple times to obtain

all the eigenvalues, with SymIntegration.

In SymIntegration, this function can be used to compute an eigenvalue and the corresponding eigenvector of real square matrix with Rayleigh quotient iteration:

RayleighQuotientIteration(vector<vector<double>>& A, int N, double a, double b)

with A as the real square matrix, N is the number of iteration, a is the lower bound for uniform distribution, b is the upper bound for the uniform distribution, we are using uniform distribution to generate random μ and σ as the input to generate the initial vector v^0 that is normally distributed with mean μ and standard deviation σ , this initial vector is a prerequisite for the Rayleigh quotient iteration. The key here is to choose a and b carefully so we can pinpoint the range where the eigenvalue could be located.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation with Rayleigh Quotient Iteration ]
# ./main

v initial vector_{normalize} =
0.372864
-0.386006
0.843784

 $\lambda^{\{0\}}$  = 12.0216

Eigenvalue = 4.13413

Corresponding Eigenvector:
-0.0561771
-0.233992
-0.970614

Time taken by function: 4678 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation with Rayleigh Quotient Iteration ]
# ./main

v initial vector_{normalize} =
0.692503
0.428555
0.580328

 $\lambda^{\{0\}}$  = 0.619297

Eigenvalue = 0.267949

Corresponding Eigenvector:
-0.963611
-0.258199
-0.0691842

Time taken by function: 4678 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Eigenvalues Approximation with Rayleigh Quotient Iteration ]
# ./main

v initial vector_{normalize} =
0.687873
0.657233
-0.308019

 $\lambda^{\{0\}}$  = 3.60263

Eigenvalue = 3.73205

Corresponding Eigenvector:
0.0691842
0.258199
0.963611

Time taken by function: 4679 microseconds
```

Figure 6.26: The computation to approximate an eigenvalue and its corresponding eigenvector of matrix A with Rayleigh quotient iteration (SymIntegration/Examples/Linear Algebra/Test SymIntegration Eigenvalues Approximation with Rayleigh Quotient Iteration/main.cpp).

vii. Compute Matrix P that Diagonalizes a Matrix A with SymIntegration

Find a matrix P that diagonalizes

$$A = \begin{bmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{bmatrix}$$

Solution:

The characteristic equation of A is

$$(\lambda - 1)(\lambda - 2)^2 = 0$$

and we found the following bases for the eigenspaces:

For $\lambda = 2$

$$p_1 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}, \quad p_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

For $\lambda = 1$

$$p_3 = \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$$

There are three basis vectors in total, so the matrix

$$P = \begin{bmatrix} -1 & 0 & -2 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$

diagonalizes A . As a check, you should verify that

$$P^{-1}AP = \begin{bmatrix} -1 & 0 & -2 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 21 & 0 & 3 \end{bmatrix} \begin{bmatrix} -1 & 0 & -2 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

In general, there is no preferred order for the columns of P . Since the i -th diagonal entry of $P^{-1}AP$ is an eigenvalue for the i -th column vector of P , changing the order of the columns of P just changes the order of the eigenvalues on the diagonal of $P^{-1}AP$.

In SymIntegration we can use this function to compute the matrix P that diagonalizes matrix A and also to compute $P^{-1}AP$:

diagonalization(vector<vector<double>>& A)

with A as the matrix that we want to diagonalize, we are using the power method, deflation method, and inverse iteration with shift to obtain the matrix P (matrix with eigenvectors as the columns), thus the result here for P is the matrix with normalize eigenvectors, the end result is still the same, we will have the eigenvalues on the diagonal of $P^{-1}AP$.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Diagonalization ]# ./main

A:
      0.000000      0.000000     -2.000000
      1.000000      2.000000      1.000000
      1.000000      0.000000      3.000000

Eigenvalue(1) = 2.000000
Eigenvalue(2) = 2.000000
Eigenvalue(3) = 1.000000

P:
     -0.592140     -0.551064      0.816497
      0.546572      0.626624     -0.408248
      0.592140      0.551064     -0.408248

P^{-1}:
      1.081702     -7.888875     10.052280
      0.652340      8.476911     -7.172231
      2.449490     -0.000000      2.449490

P^{-1}AP:
      2.000000      0.000000      0.000000
      0.000000      2.000000     -0.000000
     -0.000000     -0.000000      1.000000

Time taken by function: 4514 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Diagonalization ]# 

```

Figure 6.27: The computation to obtain matrix P that diagonalizes matrix A , then compute P^{-1} and $P^{-1}AP$. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Diagonalization/main.cpp*).

viii. Compute Complex Eigenvalues and Its Corresponding Eigenvector using Rayleigh Quotient Iteration with SymIntegration

Find the eigenvalues and bases for the eigenspaces of

$$A = \begin{bmatrix} -2 & -1 \\ 5 & 2 \end{bmatrix}$$

Solution:

The characteristic polynomial of A is

$$\begin{vmatrix} \lambda + 2 & 1 \\ -5 & \lambda - 2 \end{vmatrix} = \lambda^2 + 1 = (\lambda - i)(\lambda + i)$$

so the eigenvalues of A are $\lambda = i$ and $\lambda = -i$. Note that these eigenvalues are complex conjugates. To find the eigenvectors we must solve the system

$$\begin{bmatrix} \lambda + 2 & 1 \\ -5 & \lambda - 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

substitute with $\lambda = i$, the system becomes

$$\begin{bmatrix} i + 2 & 1 \\ -5 & i - 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

We could solve this system by reducing the augmented matrix

$$\begin{bmatrix} i + 2 & 1 & 0 \\ -5 & i - 2 & 0 \end{bmatrix}$$

to reduced row echelon form by Gauss-Jordan elimination. Observe that the reduced row echelon form must have a row of zeros in order to have a basis for the corresponding $\lambda = i$, thus each row of the matrix above must be a scalar multiple of the other.

We can just set one of the row to zero and then do the parameterization. Now we set the first row to zero, so we will have

$$\begin{bmatrix} 0 & 0 & 0 \\ -5 & i - 2 & 0 \end{bmatrix}$$

We divide the second row with $-\frac{1}{5}$ then interchange the rows

$$\begin{bmatrix} 1 & \frac{2}{5} - \frac{i}{5} & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

then do the parameterization to get the general solution

$$\begin{aligned} x_1 &= \left(-\frac{2}{5} + \frac{i}{5} \right) t \\ x_2 &= t \end{aligned}$$

you can check by substituting the values of x_1 and x_2 into the first row

$$\begin{bmatrix} i + 2 & 1 & 0 \\ -5 & i - 2 & 0 \end{bmatrix}$$

The equation will hold for

$$(i + 2)x_1 + x_2 = 0$$

you can verify this by hand.

So, the eigenspace corresponding to $\lambda = i$ is one-dimensional and consists of all complex scalar multiples of the basis vector

$$\mathbf{x} = \begin{bmatrix} -\frac{2}{5} + \frac{i}{5} \\ 1 \end{bmatrix}$$

As a check, let us confirm that

$$A\mathbf{x} = i\mathbf{x}$$

$$A\mathbf{x} = \begin{bmatrix} -2 & -1 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} -\frac{2}{5} + \frac{i}{5} \\ 1 \end{bmatrix} = \begin{bmatrix} -2\left(-\frac{2}{5} + \frac{i}{5}\right) - 1 \\ 5\left(-\frac{2}{5} + \frac{i}{5}\right) + 2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{5} - \frac{2i}{5} \\ i \end{bmatrix} = i\mathbf{x}$$

We could find the basis for the eigenspace corresponding to $\lambda = -i$ in a similar way, but to save time and be more efficient we have a theorem that implies

$$\bar{\mathbf{x}} = \begin{bmatrix} -\frac{2}{5} - \frac{i}{5} \\ 1 \end{bmatrix}$$

must be a basis for this eigenspace. The following computations confirm that $\bar{\mathbf{x}}$ is an eigenvector of A corresponding to $\lambda = -i$:

$$A\bar{\mathbf{x}} = \begin{bmatrix} -2 & -1 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} -\frac{2}{5} - \frac{i}{5} \\ 1 \end{bmatrix} = \begin{bmatrix} -2\left(-\frac{2}{5} - \frac{i}{5}\right) - 1 \\ 5\left(-\frac{2}{5} - \frac{i}{5}\right) + 2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{5} + \frac{2i}{5} \\ -i \end{bmatrix} = -i\bar{\mathbf{x}}$$

In SymIntegration, to compute the approximation of a complex eigenvalue we can use Rayleigh quotient iteration that has been modified so it can work with complex number. The function that can be used is:

ComplexRayleighQuotientIteration(vector<vector<complex<double>>> &A, int N_{Rayleigh} , int N double a , double b)

with A as the matrix that we want to compute the eigenvalues and the corresponding eigenvector, N_{Rayleigh} is the maximum number of iteration for the Rayleigh quotient iteration, N is the maximum number of the outermost iteration till we obtain all the correct eigenvalues, a is the lower bound for uniform distribution, b is the upper bound for the uniform distribution, we are using uniform distribution to generate random μ and σ as the input to generate the initial complex vector $\mathbf{v}^0$ that is normally distributed with mean μ and standard deviation σ , this initial vector is a prerequisite for the Rayleigh quotient iteration. The key here is to choose a and b carefully so we can pinpoint the range where the eigenvalue could be located.


```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Complex Eigenvalues with Rayleigh Quotient ]# ./main
Matrix A :
      (-2,0)      (-1,0)
      (5,0)       (2,0)

v initial vector_{normalize} =
(0.195,0.535)
(0.752,0.331)

λ^{0} = (2,2.03)

Eigenvalue = (0,1)

Corresponding Eigenvector:
(0.113,-0.392)
(-0.619,0.671)

Time taken by function: 5184 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Complex Eigenvalues with Rayleigh Quotient ]# ./main
Matrix A :
      (-2,0)      (-1,0)
      (5,0)       (2,0)

v initial vector_{normalize} =
(0.621,-0.0174)
(0.71,0.333)

λ^{0} = (2.2,-1.31)

Eigenvalue = (0,-1)

Corresponding Eigenvector:
(0.359,0.193)
(-0.912,-0.0275)

Time taken by function: 5538 microseconds

```

Figure 6.28: The computation to obtain complex eigenvalue of matrix A one by one with Rayleigh quotient iteration (the earlier / beta version of the code). (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Complex Eigenvalues with Rayleigh Quotient/main.cpp*).

We compile it then run it twice and it produces different eigenvalues

$$\begin{aligned}\lambda_1 &= (0,1) = i \\ \lambda_2 &= (0,-1) = -i\end{aligned}$$

which are corrects, and the computed corresponding eigenvector is already normalized.

Rayleigh quotient has a similar atmosphere like bisection method in root finding, in term of choosing a and b when we call the function, the different here is that eigenvalue problem is scientific computing, it is far more complex and deeper than root finding problem.

After some trial and error, to make it more efficient we modify and update the source code so it will produce a vector filled with the correct eigenvalues and a matrix filled with the eigenvectors, that is showing the eigenvector as row vector (depends on the future usage, the matrix can be transposed so it will show the eigenvector as column vector).

```

*****
Iter: 2

v initial vector_{normalize} =
(0.66,0.275)
(0.378,0.588)

 $\lambda^{\{0\}} = (1.6, -1.71)$ 

Eigenvalue = (0,-1)

Corresponding Eigenvector:
(-0.407,-0.0268)
(0.842,-0.354)

Ax:
(-0.0268,0.407)
(-0.354,-0.842)

 $\lambda x$ :
(-0.0268,0.407)
(-0.354,-0.842)

Eigenvalue accepted= (0,-1)

Eigenvalues :
(0,1)
(0,-1)

Eigenvectors (represented by row):
(-0.322,-0.251) (0.392,0.824)
(-0.407,-0.0268) (0.842,-0.354)

Time taken by function: 13710 microseconds

```

Figure 6.29: *The computation to obtain all the complex eigenvalues of matrix A along with its corresponding eigenvector, it stops at iteration 2 (the latest version of the code made on January 1st, 2026). (SymIntegration/Examples/Linear Algebra/Test SymIntegration Complex Eigenvalues with Rayleigh Quotient/main.cpp).*

If we run the compiled code again, we will probably end up with different eigenvectors, yet the eigenvalues are still the same. This proves the eigenvectors' property that says: *eigenvectors may not be unique, as a matrix may have multiple eigenvectors corresponding to the same eigenvalue.*

```

*****
Iter: 4
v initial vector_{normalize} =
(0.358,0.744)
(0.16,-0.541)

 $\lambda^{\{0\}}$  = (-2.11,1.88)

Eigenvalue = (0,1)

Corresponding Eigenvector:
(-0.356,-0.2)
(0.512,0.756)

Ax:
(0.2,-0.356)
(-0.756,0.512)

 $\lambda x$ :
(0.2,-0.356)
(-0.756,0.512)

Eigenvalue accepted= (0,1)

Eigenvalues :
(0,-1)
(0,1)

Eigenvectors (represented by row):
(-0.0169,0.408) (-0.374,-0.833)
(-0.356,-0.2) (0.512,0.756)

Time taken by function: 22872 microseconds

```

Figure 6.30: *The computation to obtain all the complex eigenvalues of matrix A along with its corresponding eigenvector, this time it stops at iteration 4 (the latest version of the code made on January 1st, 2026). ([SymIntegration/Examples/Linear Algebra/Test SymIntegration Complex Eigenvalues with Rayleigh Quotient/main.cpp](#)).*

We modify the source code since we saw the defect when the earlier version of complex Rayleigh quotient iteration resulting in an eigenvalue of $\lambda = (1, -1) = 1 - i$, which is wrong when we check that $Ax \neq \lambda x$.

```

Time taken by function: 5011 microseconds
root@ ~:/SourceCodes/CPP/C++ Create Library/Test SymIntegration Complex Eigenvalues with Rayleigh Quotient
main

Matrix A :
      (-2,0)      (-1,0)
      (5,0)       (2,0)

v initial vector_{normalize} =
(0.419,0.504)
(0.477,0.585)

 $\lambda^{\{0\}} = (2.26, -0.0304)$ 

Eigenvalue = (1,-1)

Corresponding Eigenvector:
(-0.282,0.12)
(0.657,-0.689)

Ax:
(-0.0923,0.449)
(-0.0974,-0.778)

 $\lambda x$ :
(-0.162,0.402)
(-0.0325,-1.35)

Time taken by function: 5011 microseconds

```

Figure 6.31: The computation to obtain an eigenvalue of matrix A ended up with the wrong eigenvalue since $Ax \neq \lambda x$ (the earlier / beta version of the code). (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Complex Eigenvalues with Rayleigh Quotient/main.cpp*).

If you want to use the earlier / beta version of the code you just need to delete the outermost **for** loop and adjust a bit the **ComplexRayleighQuotientIteration()** function in **src/linearalgebra.cpp**.

The most robust and generally used method for dense, non-symmetric matrices with complex eigenvalues is the QR algorithm with implicit shifts. For finding a specific eigenpair when a good initial guess is available, inverse iteration and Rayleigh quotient iteration are highly effective.

Rayleigh quotient iteration often exhibits cubic convergence for Hermitian matrices and rapid convergence for general matrices, it is a powerful extension of inverse iteration that uses the Rayleigh quotient as the shift μ .

ix. Compute Real Eigenvalues and Its Corresponding Eigenvector using QR Algorithm with Shifts with SymIntegration

Compute the eigenvalues and its corresponding eigenvector for

$$A = \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}$$

Solution:

At iteration $k = 1$, set $A^{(0)} = A$, and we will choose $\mu^1 = a_{22} = 3$, thus

$$\mu^{(1)}I = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$$

$$A - \mu^{(1)}I = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

the QR decomposition of $A - \mu^{(1)}I$ is

$$Q^{(1)} = \begin{bmatrix} -0.447 & -0.894 \\ -0.894 & 0.447 \end{bmatrix}, \quad R^{(1)} = \begin{bmatrix} -2.24 & -0.447 \\ 0 & -0.894 \end{bmatrix}$$

$$R^{(1)}Q^{(1)} = \begin{bmatrix} 1.4 & 1.8 \\ 0.8 & -0.4 \end{bmatrix}$$

$$R^{(1)}Q^{(1)} + \mu^{(1)}I = \begin{bmatrix} 4.4 & 1.8 \\ 0.8 & 2.6 \end{bmatrix}$$

we will set $A^{(1)} = R^{(1)}Q^{(1)} + \mu^{(1)}I$ and proceed to the next iteration $k = 2$, repeat the iteration till $A^{(k)} = R^{(k)}Q^{(k)} + \mu^{(k)}I$ become a matrix with zero below the diagonal, the diagonal will be the eigenvalues of the matrix A .

We use 6 iterations and we obtain

$$A^{(6)} = R^{(6)}Q^{(6)} + \mu^{(6)}I = \begin{bmatrix} 5 & -1 \\ 0 & 2 \end{bmatrix}$$

hence the eigenvalues of matrix A are

$$\lambda_1 = 5$$

$$\lambda_2 = 2$$

the corresponding eigenvector for $\lambda_1 = 5$

$$x_1 = \begin{bmatrix} 0.707 \\ 0.707 \end{bmatrix}$$

and the corresponding eigenvector for $\lambda_2 = 2$

$$x_2 = \begin{bmatrix} 0.447 \\ -0.894 \end{bmatrix}$$

In SymIntegration, to compute the eigenvalues and its corresponding eigenvector, given that the matrix has only real eigenvalues we can use this function:

QRAlgorithmwithShifts(vector<vector<complex<double>>>& A, int N)

with A as the matrix, and N as the number of iterations we want to use till the matrix converge.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Real Eigenvalues and Eigenvectors with QR Algorithm with Shifts ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Real Eigenvalues and Eigenvectors with QR Algorithm with Shifts ]# ./main

Iteration: 1
μ = (3,0)

A:
    (4,0)    (1,0)
    (2,0)    (3,0)

μI =
    (3,0)    (0,0)
    (0,0)    (3,0)

A - μI =
    (1,0)    (1,0)
    (2,0)    (0,0)

Q:
    (-0.447,0)  (-0.894,0)
    (-0.894,0)  (0.447,0)

R:
    (-2.24,0)  (-0.447,0)
    (0,0)     (-0.894,0)

RQ =
    (1.4,0)    (1.8,0)
    (0.8,0)    (-0.4,0)

A_{new} =
    (4.4,0)    (1.8,0)
    (0.8,0)    (2.6,0)

Iteration: 2
μ = (2.6,0)

A:
    (4.4,0)    (1.8,0)
    (0.8,0)    (2.6,0)

μI =
    (2.6,0)    (0,0)
    (0,0)     (2.6,0)

```

Figure 6.32: The computation to obtain all eigenvalues of matrix A and the corresponding eigenvector for each eigenvalue. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real Eigenvalues and Eigenvectors with QR Algorithm with Shifts/main.cpp*).

```

Iteration: 2
μ = (2.6,0)
A:
  (4.4,0)   (1.8,0)
  (0.8,0)   (2.6,0)
μI =
  (2.6,0)   (0,0)
  (0,0)     (2.6,0)
A - μI =
  (1.8,0)   (1.8,0)
  (0.8,0)   (0,0)
Q:
  (-0.914,0) (-0.406,0)
  (-0.406,0) (0.914,0)
R:
  (-1.97,0)  (-1.64,0)
  (0,0)      (-0.731,0)
RQ =
  (2.47,0)   (-0.703,0)
  (0.297,0)  (-0.668,0)
A_{new} =
  (5.07,0)   (-0.703,0)
  (0.297,0)  (1.93,0)
Iteration: 3
μ = (1.93,0)
A:
  (5.07,0)   (-0.703,0)
  (0.297,0)  (1.93,0)
μI =
  (1.93,0)   (0,0)
  (0,0)      (1.93,0)
A - μI =
  (3.14,0)   (-0.703,0)
  (0.297,0)  (0,0)
Q:
  (-0.996,0) (-0.0943,0)
  (-0.0943,0) (0.996,0)

```

Figure 6.33: The computation to obtain all eigenvalues of matrix A and the corresponding eigenvector for each eigenvalue. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real Eigenvalues and Eigenvectors with QR Algorithm with Shifts/main.cpp*).

```

Iteration: 5
μ = (2,0)

A:
      (5,0)      (-1,0)
(4.3e-06,0)      (2,0)

μI =
      (2,0)      (0,0)
      (0,0)      (2,0)

A - μI =
      (3,0)      (-1,0)
(4.3e-06,0)      (0,0)

Q:
      (-1,0) (-1.43e-06,0)
(-1.43e-06,0)      (1,0)

R:
      (-3,0)      (1,0)
      (0,0) (1.43e-06,0)

RQ =
      (3,0)      (1,0)
(-2.05e-12,0) (1.43e-06,0)

A_{new} =
      (5,0)      (1,0)
(-2.05e-12,0)      (2,0)

Iteration: 6
μ = (2,0)

A:
      (5,0)      (1,0)
(-2.05e-12,0)      (2,0)

μI =
      (2,0)      (0,0)
      (0,0)      (2,0)

A - μI =
      (3,0)      (1,0)
(-2.05e-12,0)      (0,0)

Q:
      (-1,0) (6.83e-13,0)
(6.83e-13,0)      (1,0)

```

Figure 6.34: The computation to obtain all eigenvalues of matrix A and the corresponding eigenvector for each eigenvalue. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real Eigenvalues and Eigenvectors with QR Algorithm with Shifts/main.cpp*).


```

A:
      (5,0)      (1,0)
(-2.05e-12,0)   (2,0)

μI =
      (2,0)      (0,0)
      (0,0)      (2,0)

A - μI =
      (3,0)      (1,0)
(-2.05e-12,0)   (0,0)

Q:
      (-1,0) (6.83e-13,0)
(6.83e-13,0) (1,0)

R:
      (-3,0)      (-1,0)
      (0,0)      (0,0)

RQ =
      (3,0)      (-1,0)
      (0,0)      (0,0)

A_{new} =
      (5,0)      (-1,0)
      (0,0)      (2,0)

*****

End of iteration

*****

Eigenvalue(1) = (5,0)

Corresponding Eigenvector:
(0.707,0)
(0.707,0)

Eigenvalue(2) = (2,0)

Corresponding Eigenvector:
(0.447,0)
(-0.894,0)

Eigenvectors (in column vectors):
      (0.707,0)   (0.447,0)
      (0.707,0)   (-0.894,0)

Eigenvalues:
(5,0)
(2,0)

Time taken by function: 6656 microseconds

```

Figure 6.35: The computation to obtain all eigenvalues of matrix A and the corresponding eigenvector for each eigenvalue. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real Eigenvalues and Eigenvectors with QR Algorithm with Shifts/main.cpp*).

x. Compute Real and Complex Eigenvalues from Real Matrix using Francis Double Step QR Algorithm with SymIntegration

Compute the eigenvalues of the real matrix

$$A = \begin{bmatrix} 7 & 3 & 4 & -11 & -9 & -2 \\ -6 & 4 & -5 & 7 & 1 & 12 \\ -1 & -9 & 2 & 2 & 9 & 1 \\ -8 & 0 & -1 & 5 & 0 & 8 \\ -4 & 3 & -5 & 7 & 2 & 10 \\ 6 & 1 & 4 & -11 & -7 & -1 \end{bmatrix}$$

Solution:

We will compute the Hessenberg matrix of A

$$H = \begin{bmatrix} 7 & 7.2761 & 5.8120 & -0.1397 & 9.0152 & 7.9363 \\ -12.3693 & 4.1307 & 18.9685 & -1.2071 & 10.6833 & 2.4160 \\ 0 & -7.1603 & 2.4478 & -0.5656 & -4.1814 & -3.2510 \\ 0 & 0 & -8.5988 & 2.9151 & -3.4169 & 5.7230 \\ 0 & 0 & 0 & 1.0464 & -2.8351 & -10.9792 \\ 0 & 0 & 0 & 0 & 1.4143 & 5.3415 \end{bmatrix}$$

Next, we will compute the first householder reflector, P_0 . Given an upper Hessenberg matrix H , calculate the first column of $G = H^2 - sH + tI$, where

$$s = h_{n-1,n-1} + h_{n,n}$$

$$t = (h_{n-1,n-1} * h_{n,n}) - (h_{n-1,n} * h_{n,n-1})$$

The first three components of $Ge_1 = x$ are

$$x = h_{1,1}^2 + h_{1,2}h_{2,1} - sh_{1,1} + t$$

$$y = h_{2,1}(h_{1,1} + h_{2,2} - s)$$

$$z = h_{2,1}h_{3,2}$$

this will populate the first three entries for the target vector x .

Construct the Householder vector v . Compute a reflector P_0 that maps $x = [x, y, z]^T$ to a multiple of $e_1 = [1, 0, 0]^T$ ($P_0x = \alpha e_1$).

Let

$$x = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} h_{1,1}^2 + h_{1,2}h_{2,1} - sh_{1,1} + t \\ h_{2,1}(h_{1,1} + h_{2,2} - s) \\ h_{2,1}h_{3,2} \end{bmatrix}$$

Calculate v

$$v = x + \text{sgn}(x_1) \|x\|_2 e_1$$

If the sign of the first component of vector x is negative then $\text{sgn}(x_1) = 1$, if the sign of the first component of vector x is positive then $\text{sgn}(x_1) = -1$. Normalize v

$$v = \frac{v}{\|v\|_2}$$

Define the first Householder reflector P_0

$$P_0 = I - 2vv^T$$

given that v is already normalized. If v is not normalized yet, then use this formula

$$P_0 = I - 2 \frac{vv^T}{||v||^2}$$

Apply P_0 to the top-left $\times n$ submatrix of H . After P_0 has been applied, H is transformed into

$$H = P_0^T H P_0$$

After computing P_0 to act on the first column, the subsequent reflectors $P_1, P_2, \dots, P_{n-2}$ are calculated to restore the Hessenberg form (chase the "bulge" created by P_0 down the matrix).

The "bulge" exists in the subdiagonal elements that are now non-zero. The second Householder reflector P_1 is designed to eliminate these new non-zeros.

Identify the "vector" to nullify. Look at the first column of the current matrix H below the subdiagonal. Because of the initial transformation, the elements $h_{3,1}$ and $h_{4,1}$ are now non-zero. Let x be a small vector composed of these elements:

$$x = \begin{bmatrix} h_{2,1} \\ h_{3,1} \\ h_{4,1} \end{bmatrix}$$

In many implementations, the bulge starts at $h_{3,1}, h_{4,1}$, but the reflector P_1 typically operates on a 3×3 submatrix starting from the first non-zero element of the bulge, $h_{2,1}$.

Then define the Householder reflector P_1 formulation. Compute a reflector that maps x to a multiple of e_1 ($P_1 x = \alpha e_1$)

Let

$$x = \begin{bmatrix} h_{2,1} \\ h_{3,1} \\ h_{4,1} \end{bmatrix}$$

Calculate v

$$v = x + \text{sgn}(x_1) ||x||_2 e_1$$

If the sign of the first component of vector x is negative then $\text{sgn}(x_1) = 1$, if the sign of the first component of vector x is positive then $\text{sgn}(x_1) = -1$. Normalize v

$$v = \frac{v}{||v||_2}$$

Define the first Householder reflector P_1

$$P_1 = I - 2vv^T$$

given that v is already normalized. If v is not normalized yet, then use this formula

$$P_1 = I - 2 \frac{vv^T}{||v||^2}$$

Basically, after successfully computing P_0 and updating the matrix H creating the first bulge, iterate for $k = 1$ to $k = n - 2$. Identify the "bulge" (non-zero elements below the subdiagonal) created by the previous step. Compute a new Householder reflector P_k to eliminate these elements in the current column. Apply P_k to H and update the matrix H .

$$H \leftarrow P_k^T H P_k$$

Repeat this iteration to compute $P_0, P_1, \dots, P_{n-2}$ (to push the bulge off the bottom right of the matrix) for N amount of times, till H converges to the quasi-upper triangular matrix (real Schur form).

The matrix P_k is orthogonal ($P_k = P_k^T$).

In SymIntegration to compute the real and complex eigenvalues from real matrix, we can use this function:

FrancisDoubleStepQRReal(vector<vector<complex<double>>>& A, int N)

with A as the matrix in complex term, even if the entries are all real, no imaginary parts. N is the number of iterations that we want to test.

Another function in SymIntegration that can be used to compute Francis double step QR algorithm with complex arithmetic is:

FrancisDoubleStepQR(vector<vector<complex<double>>>& A, int N)

In this case, we try to put $N = 8$ and it converges to the eigenvalues already. We can also put a stop iteration condition if all the eigenvalues already converges then the iteration will stop, here is an example for the stop criteria

$$|\lambda_i^k| - |\lambda_i^{k-1}| < 10^{-6}$$

for $i = 1, 2, \dots, n$ and k is the last/ current iteration.

```
*****
End of 8 iterations
*****
H updated :
      5.000012      -5.999927      7.326973      6.485041      -8.955880      20.790016
      5.999985      5.000037      4.726162      2.538411      -14.009292      2.792811
      0.000000      -0.000002      0.904127      5.837251      11.286555      1.661152
      0.000000      0.000000      -0.687738      1.079478      -3.042816      10.446256
      0.000000      -0.000000      0.000000      0.011822      4.023762      -4.918176
      0.000000      -0.000000      -0.000000      0.000000      -0.000000      2.961084

Eigenvalues:
(5.000024, -5.999956)
(5.000024, 5.999956)
(0.991803, -2.001703)
(0.991803, 2.001703)
(4.023762, -0.000000)
(2.961083, 0.000000)

Time taken by function: 466193 microseconds
```

Figure 6.36: The computation to obtain all complex and real eigenvalues of matrix A with real entries, no imaginary parts with $N = 8$. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm/main.cpp).

V. Eigenvalues and Eigenvectors

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -std=c++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm ]# ./main

A =
(7,0) (3,0) (4,0) (-11,0) (-9,0) (-2,0)
(6,0) (4,0) (-5,0) (7,0) (1,0) (12,0)
(-1,0) (-9,0) (2,0) (2,0) (9,0) (1,0)
(-8,0) (0,0) (-1,0) (5,0) (0,0) (8,0)
(-4,0) (3,0) (-5,0) (7,0) (2,0) (10,0)
(6,0) (1,0) (4,0) (-11,0) (-7,0) (-1,0)

H_0 =
(7,0) (7.27607,0) (5.81205,0) (-0.139701,0) (9.0152,0) (7.93634,0)
(12.3693,0) (4.13072,0) (18.9685,0) (-1.20707,0) (10.6833,0) (2.41595,0)
(0,0) (-7.16034,0) (2.44776,0) (-0.565594,0) (-4.1814,0) (-3.25096,0)
(0,0) (0,0) (-8.59877,0) (2.9151,0) (-3.41686,0) (5.72297,0)
(0,0) (0,0) (0,0) (1.04644,0) (-2.8351,0) (-10.9792,0)
(0,0) (0,0) (0,0) (1.41429,0) (5.34152,0)

H_0 =
7.000000 7.276069 5.812049 -0.139701 9.015201 7.936343
12.369317 4.130719 18.968509 -1.207073 10.683309 2.415951
0.000000 -7.160342 2.447765 -0.565594 -4.181396 -3.250955
0.000000 0.000000 -8.598771 2.915100 -3.416858 5.722969
0.000000 0.000000 0.000000 1.046436 -2.835101 -10.979178
0.000000 0.000000 0.000000 0.000000 1.414293 5.341517

```

Figure 6.37: The computation to obtain all complex and real eigenvalues of matrix A with real entries, no imaginary parts. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm/main.cpp).

```

Iteration: 1

H =
7.000000 7.276069 5.812049 -0.139701 9.015201 7.936343
12.369317 4.130719 18.968509 -1.207073 10.683309 2.415951
0.000000 -7.160342 2.447765 -0.565594 -4.181396 -3.250955
0.000000 0.000000 -8.598771 2.915100 -3.416858 5.722969
0.000000 0.000000 0.000000 1.046436 -2.835101 -10.979178
0.000000 0.000000 0.000000 0.000000 1.414293 5.341517

S = 2.506416
t = 0.384040
x = 121.839129
y = 106.676737
z = -88.568536

x =
121.839129
106.676737
-88.568536

w =
-184.578127
-0.000000
-0.000000

u =
-62.738999
106.676737
-88.568536

uT =
-62.738999 106.676737 -88.568536

uTu =
23160.493693

uuT =
3936.181938 -6692.791623 5556.701269
-6692.791623 11379.926135 -9448.202418
5556.701269 -9448.202418 7844.385621

||u|| = 0.000043

```

Figure 6.38: The computation to obtain all complex and real eigenvalues of matrix A with real entries, no imaginary parts. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm/main.cpp).

```

P_{0}:
    0.660095      0.577949      -0.479843      0.000000      0.000000      0.000000
    0.577949      0.017298      0.815890      0.000000      0.000000      0.000000
    -0.479843      0.815890      0.322606      0.000000      0.000000      0.000000
    0.000000      0.000000      0.000000      1.000000      0.000000      0.000000
    0.000000      0.000000      0.000000      0.000000      1.000000      0.000000
    0.000000      0.000000      0.000000      0.000000      0.000000      1.000000

B_{0}:
    7.372561      18.102312      7.417642      -0.518446      14.131712      8.194987
    -0.820541      7.072521      -1.487345      -0.563083      1.983574      1.976173
    -3.427433      14.845075      -0.866598      -1.100268      3.041574      -2.885829
    4.126061      -7.015647      -2.774018      2.915100      -3.416858      5.722969
    0.000000      0.000000      0.000000      1.046436      -2.835101      -10.979178
    0.000000      0.000000      0.000000      0.000000      1.414293      5.341517

x_{1}:
-0.820541
-3.427433
4.126061

P_{1} reflector:
    1.000000      0.000000      0.000000      0.000000      0.000000      0.000000
    0.000000      0.151215      0.631631      -0.760379      0.000000      0.000000
    0.000000      0.631631      0.529966      0.565843      0.000000      0.000000
    0.000000      -0.760379      0.565843      0.318819      0.000000      0.000000
    0.000000      0.000000      0.000000      0.000000      1.000000      0.000000
    0.000000      0.000000      0.000000      0.000000      0.000000      1.000000

H^{1}:
    7.372561      7.816770      15.071721      -9.732688      14.131712      8.194987
    -5.426321      5.509396      8.980460      -12.198117      4.819205      -5.875578
    -0.000000      -1.150449      4.112520      -7.813504      0.931413      2.957123
    0.000000      -0.594085      0.782781      -0.500893      -0.876574      -1.310975
    0.000000      -0.795688      0.592119      0.333624      -2.835101      -10.979178
    0.000000      0.000000      0.000000      0.000000      1.414293      5.341517

```

Figure 6.39: The computation to obtain all complex and real eigenvalues of matrix A with real entries, no imaginary parts. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm/main.cpp).

```

H^{1}:
    7.372561      7.816770      15.071721      -9.732688      14.131712      8.194987
    -5.426321      5.509396      8.980460      -12.198117      4.819205      -5.875578
    -0.000000      -1.150449      4.112520      -7.813504      0.931413      2.957123
    0.000000      -0.594085      0.782781      -0.500893      -0.876574      -1.310975
    0.000000      -0.795688      0.592119      0.333624      -2.835101      -10.979178
    0.000000      0.000000      0.000000      0.000000      1.414293      5.341517

x_{2}:
-1.150449
-0.594085
-0.795688

P_{2} reflector:
    1.000000      0.000000      0.000000      0.000000      0.000000      0.000000
    0.000000      1.000000      0.000000      0.000000      0.000000      0.000000
    0.000000      0.000000      0.757007      0.390914      0.523571      0.000000
    0.000000      0.000000      0.390914      0.371118      -0.842293      0.000000
    0.000000      0.000000      0.523571      -0.842293      -0.128125      0.000000
    0.000000      0.000000      0.000000      0.000000      0.000000      1.000000

H^{2}:
    7.372561      7.816770      15.003708      -9.623271      14.278258      8.194987
    -5.426321      5.509396      4.553051      -5.075539      14.358833      -5.875578
    -0.000000      -1.519733      -0.084844      0.199860      7.096121      -4.022294
    -0.000000      0.000000      0.953422      -2.803829      3.387732      9.917138
    -0.000000      0.000000      0.454532      -2.161796      3.665199      4.059200
    0.000000      0.000000      0.740483      -1.191249      -0.181207      5.341517

x_{3}:
0.953422
0.454532
0.740483

```

Figure 6.40: The computation to obtain all complex and real eigenvalues of matrix A with real entries, no imaginary parts. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm/main.cpp).

```

P_{3} reflector:
    1.000000    0.000000    0.000000    0.000000    0.000000    0.000000
    0.000000    1.000000    0.000000    0.000000    0.000000    0.000000
    0.000000    0.000000    1.000000    0.000000    0.000000    0.000000
    0.000000    0.000000    0.000000    0.739125    0.352369    0.574047
    0.000000    0.000000    0.000000    0.352369    0.524048    -0.775377
    0.000000    0.000000    0.000000    0.574047    -0.775377    -0.263174

H^{3}:
    7.372561    7.816770    15.003708    2.622721    -2.262645    -18.751949
   -5.426321    5.509396    4.553051    -2.064713    10.292050    -12.500810
   -0.000000   -1.519733   -0.004844    0.339186    6.907930    -4.328878
   -0.000000    0.000000    1.289934    5.489574    -8.475220    -7.994189
   -0.000000    0.000000    0.000000    1.111672    0.136337    -3.600585
    0.000000   -0.000000    0.000000    0.635934    -1.194952    0.576976

x_{4}:
1.111672
0.635934

P_{4} reflector:
    1.000000    0.000000    0.000000    0.000000    0.000000    0.000000
    0.000000    1.000000    0.000000    0.000000    0.000000    0.000000
    0.000000    0.000000    1.000000    0.000000    0.000000    0.000000
    0.000000    0.000000    0.000000    1.000000    0.000000    0.000000
    0.000000    0.000000    0.000000    0.000000    0.868910    0.496546
    0.000000    0.000000    0.000000    0.000000    0.496546    -0.868910

H^{4}:
    7.372561    7.816770    15.003708    2.622721    -11.275212    15.153374
   -5.426321    5.509396    4.553051    -2.064713    2.726372    15.961311
   -0.000000   -1.519733   -0.004844    0.339186    3.846664    7.187618
   -0.000000    0.000000    1.289934    5.489574    -11.326063    2.730697
   -0.000000    0.000000    0.000000    1.280713    -1.821932    2.228287
   -0.000000    0.000000   -0.000000   -0.000000   -0.177346    2.535245

```

Figure 6.41: The computation to obtain all complex and real eigenvalues of matrix A with real entries, no imaginary parts. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm/main.cpp*).

```

H^{4}:
    7.372561    7.816770    15.003708    2.622721    -11.275212    15.153374
   -5.426321    5.509396    4.553051    -2.064713    2.726372    15.961311
   -0.000000   -1.519733   -0.004844    0.339186    3.846664    7.187618
   -0.000000    0.000000    1.289934    5.489574    -11.326063    2.730697
   -0.000000    0.000000    0.000000    1.280713    -1.821932    2.228287
   -0.000000    0.000000   -0.000000   -0.000000   -0.177346    2.535245

x_{5}:
-1.821932
-0.177346

P_{5} last:
    0.004704    -0.096882
   -0.096882    1.995296

P_{5} reflector:
    1.000000    0.000000    0.000000    0.000000    0.000000    0.000000
    0.000000    1.000000    0.000000    0.000000    0.000000    0.000000
    0.000000    0.000000    1.000000    0.000000    0.000000    0.000000
    0.000000    0.000000    0.000000    1.000000    0.000000    0.000000
    0.000000    0.000000    0.000000    0.000000    1.000000    0.000000
    0.000000    0.000000    0.000000    0.000000    0.000000    0.995296

H^{5}:
    7.372561    7.816770    15.003708    2.622721    -11.275212    15.082091
   -5.426321    5.509396    4.553051    -2.064713    2.726372    15.886227
   -0.000000   -1.519733   -0.004844    0.339186    3.846664    7.153807
   -0.000000    0.000000    1.289934    5.489574    -11.326063    2.717852
   -0.000000    0.000000    0.000000    1.280713    -1.821932    2.217805
   -0.000000    0.000000   -0.000000   -0.000000   -0.176512    2.511449

```

Figure 6.42: The computation to obtain all complex and real eigenvalues of matrix A with real entries, no imaginary parts. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm/main.cpp*).

```

H^(5):
      7.372561      7.816770      15.003708      2.622721      -11.275212      15.082091
     -5.426321      5.509396      4.553051     -2.064713      2.726372      15.886227
      0.000000     -1.519733     -0.084844      0.339186      3.846664      7.153807
      0.000000      0.000000      1.289934      5.489574     -11.326063      2.717852
      0.000000      0.000000      0.000000      1.280713     -1.821932      2.217805
      0.000000      0.000000     -0.000000     -0.000000     -0.176512      2.511449

Iteration: 2

H =
      7.372561      7.816770      15.003708      2.622721      -11.275212      15.082091
     -5.426321      5.509396      4.553051     -2.064713      2.726372      15.886227
      0.000000     -1.519733     -0.084844      0.339186      3.846664      7.153807
      0.000000      0.000000      1.289934      5.489574     -11.326063      2.717852
      0.000000      0.000000      0.000000      1.280713     -1.821932      2.217805
      0.000000      0.000000     -0.000000     -0.000000     -0.176512      2.511449

s = 0.689517
t = -4.184220
x = 2.670627
y = -66.160096
z = 8.246560

X =
2.670627
-66.160096
8.246560

W =
-66.725530
-0.000000
-0.000000

U =
-64.054903
-66.160096
8.246560

u =
-64.054903
-66.160096
8.246560

```

Figure 6.43: The computation to obtain all complex and real eigenvalues of matrix A with real entries, no imaginary parts. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Compute Real and Complex Eigenvalues with Francis Double Step QR Algorithm/main.cpp*).

xi. Compute Inverse of a Complex Matrix with Gauss-Jordan Elimination with SymIntegration

Find the inverse of

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 3 \\ 1 & 0 & 8 \end{bmatrix}$$

Solution:

We want to reduce A to the identity matrix by row operations and simultaneously apply these operations to I to produce A^{-1} . To accomplish this we will adjoin the identity matrix to the right side of A , thereby producing a partitioned matrix of the form

$$[A|I]$$

Then we will apply row operations to this matrix until the left side is reduced to I ; these operations will convert the right side to A^{-1} , so the final matrix will have the form.

$$[I|A^{-1}]$$

After the computations we will obtain

$$A^{-1} = \begin{bmatrix} -40 & 16 & 9 \\ 13 & -5 & -3 \\ 5 & -2 & -1 \end{bmatrix}$$

If a matrix A is not invertible, then it will be impossible to reduce A to I_n by elementary row operations. This will be signaled by a row of zeros appearing on the left side of the partition at some stage of the inversion algorithm.

Reduced row echelon form (rref) is an advanced row echelon form (ref) where pivots are all 1, and there are zeros both above and below each pivot, forming an identity matrix on the coefficient side.

The algorithm steps are as follow:

1. Form an augmented matrix

$$[A|I]$$

2. Perform forward elimination to make matrix A into row echelon form.

- (a) Find the leftmost non-zero column (pivot column).
- (b) Swap rows if needed to get a non-zero entry (pivot) at the top of this column.
- (c) Multiply the pivot row by a scalar to make the pivot equal to 1 (leading one).
- (d) Add multiples of this pivot row to rows below it to create zeros in the rest of the pivot column.
- (e) Repeat for the submatrix below the first pivot.

3. Perform backward elimination to make the matrix into reduced row echelon form.

- (a) Starting from the last pivot (bottom right), use that pivot row to create zeros above it in its column.
 - (b) Work upwards and leftwards, clearing entries above each subsequent pivot.
4. Once in reduced row echelon form (rref), given the matrix is invertible, the matrix will look like

$$[I|A^{-1}]$$

it reads as (identity matrix | inverse matrix). Besides finding the inverse of a matrix this can also be used for solving systems of linear equations.

In SymIntegration to reduce complex matrix A into reduced row echelon form with the Gauss-Jordan elimination we can use this function:

GaussJordanComplexMatrix(vector<vector<complex<double>>> &A)

with A as the matrix that we want to compute the inverse, in the source code we are showing with **cout** the process to obtain the inverse, you can comment the process so you can get the final result, A^{-1} , the process can be kept / still shown for learning purpose.

In this case, the matrix A is real, but we can input it in complex matrix term with no imaginary part, so the entries for imaginary parts will be all zeros.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gauss Jordan Method Finding Inverse for Complex Matrix ]# ./main
A:
      (1,0)      (2,0)      (3,0)
      (2,0)      (5,0)      (3,0)
      (1,0)      (0,0)      (8,0)
Swap row
A_{rref} :
      (2,0)      (5,0)      (3,0)
      (1,0)      (2,0)      (3,0)
      (1,0)      (0,0)      (8,0)
A^{-1} :
      (0,0)      (1,0)      (0,0)
      (1,0)      (0,0)      (0,0)
      (0,0)      (0,0)      (1,0)
Eliminate elements below the pivot, i = 0, k = 1
A_{rref} :
      (2,0)      (5,0)      (3,0)
      (0,0)      (-0.5,0)      (1.5,0)
      (1,0)      (0,0)      (8,0)
A^{-1} :
      (0,0)      (1,0)      (0,0)
      (1,0)      (-0.5,0)      (0,0)
      (0,0)      (0,0)      (1,0)
Eliminate elements below the pivot, i = 0, k = 2
A_{rref} :
      (2,0)      (5,0)      (3,0)
      (0,0)      (-0.5,0)      (1.5,0)
      (0,0)      (-2.5,0)      (6.5,0)
A^{-1} :
      (0,0)      (1,0)      (0,0)
      (1,0)      (-0.5,0)      (0,0)
      (0,0)      (-0.5,0)      (1,0)
Swap row
A_{rref} :
      (2,0)      (5,0)      (3,0)
      (0,0)      (-2.5,0)      (6.5,0)
      (0,0)      (-0.5,0)      (1.5,0)
A^{-1} :
      (0,0)      (1,0)      (0,0)
      (0,0)      (-0.5,0)      (1,0)
      (1,0)      (-0.5,0)      (0,0)
```

Figure 6.44: The computation to obtain matrix A^{-1} with Gauss-Jordan elimination for matrix A with no imaginary part. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Gauss Jordan Method Finding Inverse for Complex Matrix/main.cpp).

```

Eliminate elements below the pivot, i = 1, k = 2
A_{rrref} :
      (2,0)      (5,0)      (3,0)
      (0,0)      (-2.5,0)    (6.5,0)
      (0,0)      (0,0)      (0.2,0)
A^{-1} :
      (0,0)      (1,0)      (0,0)
      (0,0)      (-0.5,0)    (1,0)
      (1,0)      (-0.4,0)    (-0.2,0)
Swap row
A_{rrref} :
      (2,0)      (5,0)      (3,0)
      (0,0)      (-2.5,0)    (6.5,0)
      (0,0)      (0,0)      (0.2,0)
A^{-1} :
      (0,0)      (1,0)      (0,0)
      (0,0)      (-0.5,0)    (1,0)
      (1,0)      (-0.4,0)    (-0.2,0)
After forward elimination:
A_{rrref} :
      (2,0)      (5,0)      (3,0)
      (0,0)      (-2.5,0)    (6.5,0)
      (0,0)      (0,0)      (0.2,0)
A^{-1} :
      (0,0)      (1,0)      (0,0)
      (0,0)      (-0.5,0)    (1,0)
      (1,0)      (-0.4,0)    (-0.2,0)
After leading 1:
A_{rrref} :
      (1,0)      (2.5,0)      (1.5,0)
      (-0,-0)    (1,-0)      (-2.6,-0)
      (0,0)      (0,0)      (1,0)
A^{-1} :
      (0,0)      (0.5,0)      (0,0)
      (-0,-0)    (0.2,-0)      (-0.4,-0)
      (5,0)      (-2,0)      (-1,0)
Eliminate elements above the pivot, i = 1, k = 1
A_{rrref} :
      (1,0)      (2.5,0)      (1.5,0)
      (-0,0)      (1,0)      (0,0)
      (0,0)      (0,0)      (1,0)
A^{-1} :
      (0,0)      (0.5,0)      (0,0)
      (13,0)     (-5,-0)      (-3,-0)
      (5,0)      (-2,0)      (-1,0)

```

Figure 6.45: The computation to obtain matrix A^{-1} with Gauss-Jordan elimination for matrix A with no imaginary part. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Gauss Jordan Method Finding Inverse for Complex Matrix/main.cpp).

```

      (0,0)      (-2,0)      (-1,0)
Eliminate elements above the pivot, i = 0, k = 1
A_{rref} :
      (1,0)      (0,0)      (1.5,0)
      (-0,0)     (1,0)      (0,0)
      (0,0)      (0,0)      (1,0)
A^{-1} :
      (-32.5,0)   (13,0)      (7.5,0)
      (13,0)      (-5,-0)     (-3,-0)
      (5,0)       (-2,0)      (-1,0)
Eliminate elements above the pivot, i = 0, k = 2
A_{rref} :
      (1,0)      (0,0)      (0,0)
      (-0,0)     (1,0)      (0,0)
      (0,0)      (0,0)      (1,0)
A^{-1} :
      (-40,0)     (16,0)      (9,0)
      (13,0)      (-5,-0)     (-3,-0)
      (5,0)       (-2,0)      (-1,0)
After backward elimination:
A_{rref} :
      (1,0)      (0,0)      (0,0)
      (-0,0)     (1,0)      (0,0)
      (0,0)      (0,0)      (1,0)
A^{-1} :
      (-40,0)     (16,0)      (9,0)
      (13,0)      (-5,-0)     (-3,-0)
      (5,0)       (-2,0)      (-1,0)
AA^{-1} :
      (1,0)      (0,0)      (0,0)
      (0,0)      (1,0)      (0,0)
      (0,0)      (0,0)      (1,0)
Time taken by function: 4610 microseconds

```

Figure 6.46: The computation to obtain matrix A^{-1} with Gauss-Jordan elimination for matrix A with no imaginary part. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Gauss Jordan Method Finding Inverse for Complex Matrix/main.cpp).

xii. Compute Initial Value Problem Solution of First-Order Linear Systems of Differential Equations with SymIntegration

(a) Solve the system

$$\begin{aligned}y_1' &= y_1 + y_2 \\ y_2' &= 4y_1 - 2y_2\end{aligned}$$

(b) Find the solution that satisfies the initial conditions $y_1(0) = 1, y_2(0) = 6$.

Solution:

(a) The coefficient matrix for the system is

$$\begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix}$$

Matrix A will be diagonalized by any matrix P whose columns are linearly independent eigenvectors of A . Since

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 1 & -1 \\ -4 & \lambda + 2 \end{vmatrix} = \lambda^2 + \lambda - 6 = (\lambda + 3)(\lambda - 2)$$

the eigenvalues of A are

$$\begin{aligned}\lambda_1 &= 2 \\ \lambda_2 &= -3\end{aligned}$$

By definition,

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

is an eigenvector of A corresponding to λ if and only if $\mathbf{x}$ is a nontrivial solution of

$$\begin{bmatrix} \lambda - 1 & -1 \\ -4 & \lambda + 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

If $\lambda = 2$, this system becomes

$$\begin{bmatrix} 1 & -1 \\ -4 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Solving this system yields

$$\begin{aligned}x_1 - x_2 &= 0 \\ x_1 &= x_2 \\ x_2 &= t\end{aligned}$$

so

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Thus,

$$\mathbf{p}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

is a basis for the eigenspace corresponding to $\lambda = 2$.

Similarly, you can show that

$$\mathbf{p}_2 = \begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix}$$

is a basis for the eigenspace corresponding to $\lambda = -3$. Thus,

$$P = [\mathbf{p}_1 \ \mathbf{p}_2] = \begin{bmatrix} 1 & -\frac{1}{4} \\ 1 & 1 \end{bmatrix}$$

diagonalizes A , and

$$D = P^{-1}AP = \begin{bmatrix} 2 & 0 \\ 0 & -3 \end{bmatrix}$$

The substitution

$$\begin{aligned} \mathbf{y} &= P\mathbf{u} \\ \mathbf{y}' &= P\mathbf{u}' \end{aligned}$$

yields the "diagonal system"

$$\mathbf{u}' = D\mathbf{u} = \begin{bmatrix} 2 & 0 \\ 0 & -3 \end{bmatrix} \mathbf{u}$$

or

$$\begin{aligned} u_1' &= 2u_1 \\ u_2' &= -3u_2 \end{aligned}$$

The solution of this system is

$$\begin{aligned} u_1 &= c_1 e^{2x} \\ u_2 &= c_2 e^{-3x} \end{aligned}$$

or

$$\mathbf{u} = \begin{bmatrix} c_1 e^{2x} \\ c_2 e^{-3x} \end{bmatrix}$$

now we go back to the equation $\mathbf{y} = P\mathbf{u}$ and substitute $\mathbf{u}$ that we just compute above

$$\mathbf{y} = P\mathbf{u} = \begin{bmatrix} 1 & -\frac{1}{4} \\ 1 & 1 \end{bmatrix} \begin{bmatrix} c_1 e^{2x} \\ c_2 e^{-3x} \end{bmatrix} = \begin{bmatrix} c_1 e^{2x} - \frac{1}{4} c_2 e^{-3x} \\ c_1 e^{2x} + c_2 e^{-3x} \end{bmatrix}$$

or

$$\begin{aligned} y_1 &= c_1 e^{2x} - \frac{1}{4} c_2 e^{-3x} \\ y_2 &= c_1 e^{2x} + c_2 e^{-3x} \end{aligned}$$

(b) If we substitute the initial conditions $y_1(0) = 1, y_2(0) = 6$ into

$$\begin{aligned} y_1 &= c_1 e^{2x} - \frac{1}{4} c_2 e^{-3x} \\ y_2 &= c_1 e^{2x} + c_2 e^{-3x} \end{aligned}$$

we will obtain

$$\begin{aligned} c_1 - \frac{1}{4}c_2 &= 1 \\ c_1 + c_2 &= 6 \end{aligned}$$

Solving this system, we obtain

$$\begin{aligned} c_1 &= 2 \\ c_2 &= 4 \end{aligned}$$

so the solution satisfying the initial condition is

$$\begin{aligned} y_1 &= 2e^{2x} - \frac{1}{4}4e^{-3x} \\ y_2 &= 2e^{2x} + 4e^{-3x} \end{aligned}$$

In SymIntegration, to compute the solution satisfying the initial condition for the system of first-order differential equations is

IVPSolution_firstorderdiffeq(vector<vector<complex<double>>> &A, vector<complex<double>> &x, double a, double b)

with A as the matrix that represents the system of first-order differential equations, x as the vector that represents the initial value $y_i(0)$, a is the lower bound for uniform distribution, b is the upper bound for the uniform distribution, we are using uniform distribution to generate random μ and σ as the input to generate the initial complex vector v^0 that is normally distributed with mean μ and standard deviation σ , this initial vector is a prerequisite for the Rayleigh quotient iteration. The key here is to choose a and b carefully so we can pinpoint the range where the eigenvalue could be located.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Solve Initial Value P
problem of First Order Differential Equations ]# ./main

Matrix A :
      (1,0)
      (4,0)      (-2,0)

Eigenvalues :
      (2,0)
      (-3,0)

Eigenvectors (represented bycolumn):
      (0.707,0)      (-0.243,0)
      (0.707,0)      (0.97,0)

Augmented Matrix:
      (0.707,0)      (-0.243,0)      (1,0)
      (0.707,0)      (0.97,0)      (6,0)

After backward elimination:
      (1,0)      (0,0)      (2.83,0)
      (0,0)      (1,0)      (4.12,0)

The solution satisfying the initial condition is :
[ 2*e^(2*x)-e^(-3*x) ]
[ 2*e^(2*x)+4*e^(-3*x) ]

Time taken by function: 24883 microseconds
```

Figure 6.47: The computation to obtain the initial value problem solution of system of first-order differential equations. (SymIntegration/Examples/Linear Algebra/Test SymIntegration Solve Initial Value Problem of First Order Differential Equations/main.cpp).

VI. INNER PRODUCT SPACES

[SI*] We will use the most important properties of the dot product on $\mathbb{R}^n$ as axioms, which, if satisfied by the vectors in a vector space V , will enable us to extend the notions of length, distance, angle, and perpendicularity to general vector spaces.

Definition 6.36: Inner Product

An inner product on a real vector space V is a function that associates a real number $\langle u, v \rangle$ with each pair of vectors in V in such a way that the following axioms are satisfied for all vectors u, v , and w in V and all scalars k .

1. $\langle u, v \rangle = \langle v, u \rangle$ [Symmetry axiom]
2. $\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$ [Additivity axiom]
3. $\langle ku, v \rangle = k \langle u, v \rangle$ [Homogeneity axiom]
4. $\langle v, v \rangle \geq 0$ and $\langle v, v \rangle = 0$ if and only if $v = 0$ [Positivity axiom]

A real vector space with an inner product is called a real inner product space. This definition applies only to real vector spaces (vector spaces with real scalars).

[SI*] The inner product space axioms will be satisfied automatically if we define the inner product of two vectors u and v in $\mathbb{R}^n$ to be

$$\langle u, v \rangle = u \cdot v = u_1v_1 + u_2v_2 + \cdots + u_nv_n$$

This inner product is commonly called the Euclidean inner product (or the standard inner product) on $\mathbb{R}^n$ to distinguish it from other possible inner products that might be defined on $\mathbb{R}^n$. We call $\mathbb{R}^n$ with the Euclidean inner product Euclidean n -space.

[SI*] Inner products can be used to define notions of norm and distance in a general inner product space just as we did with dot products in $\mathbb{R}^n$.

Definition 6.37: Norm and Distance

If V is a real inner product space, then the norm (or length) of a vector v in V is denoted by $\|v\|$ and is defined by

$$\|v\| = \sqrt{\langle v, v \rangle} = \sqrt{v \cdot v}$$

and the distance between two vectors is denoted by $d(u, v)$ and is defined by

$$d(u, v) = \|u - v\| = \sqrt{\langle u - v, u - v \rangle} = \sqrt{(u - v) \cdot (u - v)}$$

A vector of norm 1 is called a unit vector.

Theorem 6.64: Properties of Real Inner Product Spaces

If $\mathbf{u}$ and $\mathbf{v}$ are vectors in a real inner product space V , and if k is a scalar, then:

- (a) $\|\mathbf{v}\| \geq 0$ with equality if and only if $\mathbf{v} = \mathbf{0}$.
- (b) $\|k\mathbf{v}\| = |k|\|\mathbf{v}\|$.
- (c) $d(\mathbf{u}, \mathbf{v}) = d(\mathbf{v}, \mathbf{u})$.
- (d) $d(\mathbf{u}, \mathbf{v}) \geq 0$ with equality if and only if $\mathbf{u} = \mathbf{v}$.

[SI*] There are various applications when we modify Euclidean inner product by weighting each term differently. If

$$w_1, w_2, \dots, w_n$$

are positive real numbers, which we will call weights, and if $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ are vectors in $\mathbb{R}^n$, then it can be shown that the formula

$$\langle \mathbf{u}, \mathbf{v} \rangle = w_1 u_1 v_1 + w_2 u_2 v_2 + \dots + w_n u_n v_n \quad (6.87)$$

defines an inner product on $\mathbb{R}^n$ that we call the weighted Euclidean inner product with weights $w_1, w_2, \dots, w_n$.

[SI*] Suppose that some physical experiment has n possible numerical outcomes

$$x_1, x_2, \dots, x_n$$

and that a series of m repetitions of the experiment yields these values with various frequencies. Specifically, suppose that x_1 occurs f_1 times, x_2 occurs f_2 times, and so forth. Since there are a total of m repetitions of the experiment, it follows that

$$f_1 + f_2 + \dots + f_n = m$$

Thus, the arithmetic average of the observed numerical values (denoted by $\bar{x}$) is

$$\bar{x} = \frac{f_1 x_1 + f_2 x_2 + \dots + f_n x_n}{f_1 + f_2 + \dots + f_n} = \frac{1}{m} (f_1 x_1 + f_2 x_2 + \dots + f_n x_n) \quad (6.88)$$

if we let

$$\begin{aligned} \mathbf{f} &= (f_1, f_2, \dots, f_n) \\ \mathbf{x} &= (x_1, x_2, \dots, x_n) \\ w_1 &= w_2 = \dots = w_n = 1/m \end{aligned}$$

then we can express $\bar{x}$ as the weighted Euclidean inner product

$$\bar{x} = \langle \mathbf{f}, \mathbf{x} \rangle = w_1 f_1 x_1 + w_2 f_2 x_2 + \dots + w_n f_n x_n$$

[SI*] Norm and distance depend on the inner product being used. If the inner product is changed, then the norms and distances between vectors also change.

[SI*] If V is an inner product space, then the set of points in V that satisfy

$$\|\mathbf{u}\| = 1$$

is called the unit sphere or sometimes the unit circle in V .

Case 1:

Let the Euclidean inner product be

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_1 + u_2 v_2$$

If $\mathbf{u} = (x, y)$, then

$$\|\mathbf{u}\| = \langle \mathbf{u}, \mathbf{u} \rangle^{1/2} = \sqrt{x^2 + y^2}$$

so the equation of the unit circle is

$$\begin{aligned} \sqrt{x^2 + y^2} &= 1 \\ x^2 + y^2 &= 1 \end{aligned}$$

the graph of this equation is a circle with radius of 1 centered at the origin.

Case 2:

Let the weighted Euclidean inner product be

$$\langle \mathbf{u}, \mathbf{v} \rangle = \frac{1}{9} u_1 v_1 + \frac{1}{4} u_2 v_2$$

If $\mathbf{u} = (x, y)$, then

$$\|\mathbf{u}\| = \langle \mathbf{u}, \mathbf{u} \rangle^{1/2} = \sqrt{\frac{1}{9} x^2 + \frac{1}{4} y^2}$$

so the equation of the unit circle is

$$\begin{aligned} \sqrt{\frac{1}{9} x^2 + \frac{1}{4} y^2} &= 1 \\ \frac{x^2}{9} + \frac{y^2}{4} &= 1 \end{aligned}$$

the graph of this equation is an ellipse centered at the origin.

What we see geometrically here is two different shapes, the first is a circle, the second is an ellipse. Since both share the same norm of $\|\mathbf{u}\| = 1$, the Euclidean inner product will produce the equation for a circle, while the weighted Euclidean inner product will produce the equation for an ellipse.

[SI*] The Euclidean inner product and the weighted Euclidean inner products are special cases of a general class of inner products on $\mathbb{R}^n$ called matrix inner products.

To define this class of inner products, let $\mathbf{u}$ and $\mathbf{v}$ be vectors in $\mathbb{R}^n$ that are expressed in column form, and let A be an invertible $n \times n$ matrix. It can be shown that if $\mathbf{u} \cdot \mathbf{v}$ is the Euclidean inner product on $\mathbb{R}^n$, then the formula

$$\langle \mathbf{u}, \mathbf{v} \rangle = A\mathbf{u} \cdot A\mathbf{v} \tag{6.89}$$

also defines an inner product; it is called the inner product on $\mathbb{R}^n$ generated by A .

If $\mathbf{u}$ and $\mathbf{v}$ are in column form, then $\mathbf{u} \cdot \mathbf{v}$ can be written as $\mathbf{v}^T \mathbf{u}$ from which it follows that

$$\langle \mathbf{u}, \mathbf{v} \rangle = (\mathbf{A}\mathbf{v})^T \mathbf{A}\mathbf{u}$$

or, equivalently as

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{v}^T \mathbf{A}^T \mathbf{A} \mathbf{u} \quad (6.90)$$

[SI*] The standard Euclidean inner product on $\mathbb{R}^n$ is generated by the $n \times n$ identity matrix, since setting $\mathbf{A} = \mathbf{I}$ yields

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{I} \mathbf{u} \cdot \mathbf{I} \mathbf{v} = \mathbf{u} \cdot \mathbf{v}$$

and the weighted Euclidean inner product

$$\langle \mathbf{u}, \mathbf{v} \rangle = w_1 u_1 v_1 + w_2 u_2 v_2 + \cdots + w_n u_n v_n$$

is generated by the matrix

$$\mathbf{A} = \begin{bmatrix} \sqrt{w_1} & 0 & 0 & \cdots & 0 \\ 0 & \sqrt{w_2} & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & \sqrt{w_n} \end{bmatrix}$$

matrix $\mathbf{A}$ is a diagonal matrix with entries of $\sqrt{w_i}$ at the row- i -th, column- i -th, and zeros elsewhere.

We can observe that $\mathbf{A}^T \mathbf{A}$ will become $n \times n$ diagonal matrix whose diagonal entries are the weights $w_1, w_2, \dots, w_n$.

[SI*] The weighted Euclidean inner product $\langle \mathbf{u}, \mathbf{v} \rangle = 3u_1 v_1 + 2u_2 v_2$ is the inner product on $\mathbb{R}^2$ generated by matrix

$$\mathbf{A} = \begin{bmatrix} \sqrt{3} & 0 \\ 0 & \sqrt{2} \end{bmatrix}$$

[SI*] If $\mathbf{U}$ and $\mathbf{V}$ are $n \times n$ matrices, then the formula

$$\langle \mathbf{U}, \mathbf{V} \rangle = \text{tr}(\mathbf{U}^T \mathbf{V}) \quad (6.91)$$

defines an inner product on the vector space M_{nn} . This can be proved by confirming that the four inner product space axioms are satisfied.

For the 2×2 matrices we will have

$$\mathbf{U} = \begin{bmatrix} u_1 & u_2 \\ u_3 & u_4 \end{bmatrix}, \quad \text{and} \quad \mathbf{V} = \begin{bmatrix} v_1 & v_2 \\ v_3 & v_4 \end{bmatrix}$$

This yields

$$\langle \mathbf{U}, \mathbf{V} \rangle = \text{tr}(\mathbf{U}^T \mathbf{V}) = u_1 v_1 + u_2 v_2 + u_3 v_3 + u_4 v_4$$

which is, just the dot product of the corresponding entries in the two matrices. The norm of a matrix $\mathbf{U}$ relative to this inner product is

$$\|\mathbf{U}\| = \langle \mathbf{U}, \mathbf{U} \rangle^{1/2} = \sqrt{u_1^2 + u_2^2 + u_3^2 + u_4^2}$$

[SI*] Recall from calculus that the arc length of a curve $y = f(x)$ over an interval $[a, b]$ is given by the formula

$$L = \int_a^b \sqrt{1 + [f'(x)]^2} dx$$

the arc length is not the same with $\|f\|$, which is the length of f when f is viewed as a vector in $C[a, b]$.

Theorem 6.65: Algebraic Properties of Inner Products

If u, v , and w are vectors in a real inner product space V , and if k is a scalar, then:

- (a) $\langle 0, v \rangle = \langle v, 0 \rangle = 0$
- (b) $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$
- (c) $\langle p, v - w \rangle = \langle u, v \rangle - \langle u, w \rangle$
- (d) $\langle u - v, w \rangle = \langle u, w \rangle - \langle v, w \rangle$
- (e) $k \langle u, v \rangle = \langle u, kv \rangle$

[SI*] The angle θ between two vectors u and v in $\mathbb{R}^n$ is

$$\theta = \cos^{-1} \left(\frac{u \cdot v}{\|u\| \|v\|} \right) \quad (6.92)$$

it followed from the Cauchy-Schwarz inequality that

$$-1 \leq \frac{u \cdot v}{\|u\| \|v\|} \leq 1 \quad (6.93)$$

as required for the inverse cosine to be defined.

Theorem 6.66: Cauchy-Schwarz inequality

If u and v are vectors in a real inner product space V , then

$$|\langle u, v \rangle| \leq \|u\| \|v\| \quad (6.94)$$

[SI*] The following two alternative forms of the Cauchy-Schwarz inequality are useful to know:

$$\langle u, v \rangle^2 \leq \langle u, u \rangle \langle v, v \rangle \quad (6.95)$$

$$\langle u, v \rangle^2 \leq \|u\|^2 \|v\|^2 \quad (6.96)$$

[SI*] From the Cauchy-Schwarz inequality

$$-1 \leq \frac{u \cdot v}{\|u\| \|v\|} \leq 1$$

we will have

$$-1 \leq \frac{\langle u, v \rangle}{\|u\| \|v\|} \leq 1 \quad (6.97)$$

there is a unique angle θ in radian measure for which

$$\cos \theta = \frac{\langle u, v \rangle}{\|u\| \|v\|}, \quad 0 \leq \theta \leq \pi \quad (6.98)$$

this enables us to define the angle θ between $\mathbf{u}$ and $\mathbf{v}$ to be

$$\theta = \cos^{-1} \left(\frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\| \|\mathbf{v}\|} \right) \quad (6.99)$$

Not to be confused with the angle between two vectors in Euclidean vector spaces that is definitely use Euclidean inner product ($\mathbf{u} \cdot \mathbf{v}$), the numerator here is the inner product which can be Euclidean inner product, weighted Euclidean inner product, or inner products generated by matrices.

Theorem 6.67: Properties of Length and Distance in General Inner Product Spaces

If $\mathbf{u}, \mathbf{v}$, and $\mathbf{w}$ are vectors in a real inner product space V , and if k is any scalar, then

- (a) $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$ (Triangle inequality for vectors)
- (b) $d(\mathbf{u}, \mathbf{v}) \leq d(\mathbf{u}, \mathbf{w}) + d(\mathbf{w}, \mathbf{v})$ (Triangle inequality for distances)

[SI*] There is only an occasional need to compute angles in vector spaces other than $\mathbb{R}^2$ and $\mathbb{R}^3$.

A problem of more interest in general vector spaces is ascertaining whether the angle between vectors is $\frac{\pi}{2}$. If $\mathbf{u}$ and $\mathbf{v}$ are nonzero vectors, then the angle between them is $\theta = \frac{\pi}{2}$ if and only if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

Definition 6.38: Orthogonality

Two vectors $\mathbf{u}$ and $\mathbf{v}$ in an inner product space are called orthogonal if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

[SI*] **Example:**

The vectors $\mathbf{u} = (1, 1)$ and $\mathbf{v} = (1, -1)$ are orthogonal with respect to the Euclidean inner product on $\mathbb{R}^2$, since

$$\mathbf{u} \cdot \mathbf{v} = (1)(1) + (1)(-1) = 0$$

However, they are not orthogonal with respect to the weighted Euclidean inner product $\langle \mathbf{u}, \mathbf{v} \rangle = 3u_1v_1 + 5u_2v_2$, since

$$\langle \mathbf{u}, \mathbf{v} \rangle = 3(1)(1) + 5(1)(-1) = -2 \neq 0$$

Theorem 6.68: Generalized Theorem of Pythagoras

If $\mathbf{u}$ and $\mathbf{v}$ are orthogonal vectors in an inner product space, then

$$\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2$$

Definition 6.39: Orthogonal Complements

If W is a subspace of an inner product space V , then the set of all vectors in V that are orthogonal to every vector in W is called the orthogonal complement of W and is denoted by the symbol $W^\perp$.

Theorem 6.69: Orthogonality in Subspace of Inner Product Space

If W is a subspace of an inner product space V , then:

- (a) $W^\perp$ is a subspace of V .
- (b) $W \cap W^\perp = \{0\}$

[SI*] The set $W^\perp$ contains at least the zero vector (zero vector is perpendicular toward all vectors in $\mathbb{R}^n$), since

$$\langle 0, w \rangle = 0$$

for every vector w in W . Thus, we need to show that $W^\perp$ is closed under addition and scalar multiplication. To do this, suppose that u and v are vectors in $W^\perp$, so that for every vector w in W we have $\langle u, w \rangle = 0$ and $\langle v, w \rangle = 0$. It follows from the additivity and homogeneity axioms of inner products that

$$\begin{aligned}\langle u + v, w \rangle &= \langle u, w \rangle + \langle v, w \rangle = 0 + 0 = 0 \\ \langle ku, w \rangle &= k \langle u, w \rangle = k(0) = 0\end{aligned}$$

since 0 is in W and $W^\perp$, thus it proves that $u + v$ and ku are in $W^\perp$.

If there exists a vector z in both W and $W^\perp$, then z is orthogonal to itself; that is,

$$\langle z, z \rangle = 0$$

It follows from the positivity axiom for inner products that

$$z = 0$$

Theorem 6.70: Orthogonal Complement of $W^\perp$

If W is a subspace of a finite-dimensional inner product space V , then the orthogonal complement of $W^\perp$ is W ; that is,

$$(W^\perp)^\perp = W$$

i. Gram-Schmidt Process

[SI*] In many problems involving vector spaces, we are free to choose any basis for the vector space that seems appropriate. In inner product spaces, the solution of a problem is often greatly simplified by choosing a basis in which the vectors are orthogonal to one another.

Definition 6.40: Orthogonal and Orthonormal

A set of two or more vectors in a real inner product space is said to be orthogonal if all pairs of distinct vectors in the set are orthogonal. An orthogonal set in which each vector has norm 1 is said to be orthonormal.

[SI*] If v is a nonzero vector in an inner product space, then with $k = \frac{1}{\|v\|}$

$$\left\| \frac{1}{\|v\|} v \right\| = \left| \frac{1}{\|v\|} \right| \|v\| = \frac{1}{\|v\|} \|v\| = 1$$

from which we see that multiplying a nonzero vector by the reciprocal of its norm produces a vector of norm 1. This process is called normalizing v .

Any orthogonal set of nonzero vectors can be converted to an orthonormal set by normalizing each of its vectors.

[SI*] The set $S = \{v_1, v_2, v_3\}$ is orthonormal if

$$\begin{aligned} \langle v_1, v_2 \rangle &= \langle v_1, v_3 \rangle = \langle v_2, v_3 \rangle = 0 \\ \|v_1\| &= \|v_2\| = \|v_3\| = 1 \end{aligned}$$

[SI*] In $\mathbb{R}^2$ any two nonzero perpendicular vectors are linearly independent because neither is a scalar multiple of the other. In $\mathbb{R}^3$ any three nonzero mutually perpendicular vectors are linearly independent because no one lies in the plane of the other two.

Theorem 6.71: Orthogonal and Linear Independent

If $S = \{v_1, v_2, \dots, v_n\}$ is an orthogonal set of nonzero vectors in an inner product space, then S is linearly independent.

[SI*] In an inner product space, a basis consisting of orthonormal vectors is called an orthonormal basis, and a basis consisting of orthogonal vectors is called an orthogonal basis. An example of an orthonormal basis is the standard basis for $\mathbb{R}^n$ with the Euclidean inner product.

[SI*] If we want to express a vector u as a linear combination of basis vectors

$$S = \{v_1, v_2, \dots, v_n\}$$

we need to convert the vector equation

$$u = c_1 v_1 + c_2 v_2 + \dots + c_n v_n$$

to a linear system and solve for the coefficients $c_1, c_2, \dots, c_n$.

Theorem 6.72: Coordinates Relative to Orthonormal Bases

(a) If $S = \{v_1, v_2, \dots, v_n\}$ is an orthogonal basis for an inner product space V , and if u is any vector in V , then

$$u = \frac{\langle u, v_1 \rangle}{\|v_1\|^2} v_1 + \frac{\langle u, v_2 \rangle}{\|v_2\|^2} v_2 + \dots + \frac{\langle u, v_n \rangle}{\|v_n\|^2} v_n \quad (6.100)$$

(b) If $S = \{v_1, v_2, \dots, v_n\}$ is an orthonormal basis for an inner product space V , and if u is any vector in V , then

$$u = \langle u, v_1 \rangle v_1 + \langle u, v_2 \rangle v_2 + \dots + \langle u, v_n \rangle v_n \quad (6.101)$$

[SI*] The coordinate vector of a vector u in V relative to an orthogonal basis $S = \{v_1, v_2, \dots, v_n\}$ is

$$(u)_S = \left(\frac{\langle u, v_1 \rangle}{\|v_1\|^2}, \frac{\langle u, v_2 \rangle}{\|v_2\|^2}, \dots, \frac{\langle u, v_n \rangle}{\|v_n\|^2} \right) \quad (6.102)$$

and relative to an orthonormal basis $S = \{v_1, v_2, \dots, v_n\}$ is

$$u = \langle u, v_1 \rangle v_1 + \langle u, v_2 \rangle v_2 + \dots + \langle u, v_n \rangle v_n \quad (6.103)$$

Theorem 6.73: Projection Theorem

If W is a finite-dimensional subspace of an inner product space V , then every vector u in V can be expressed in exactly one way as

$$u = w_1 + w_2 \quad (6.104)$$

where w_1 is in W and w_2 is in $W^\perp$.

[SI*] The vector w_1 and w_2 are commonly denoted by

$$\begin{aligned} w_1 &= \text{proj}_W u \\ w_2 &= \text{proj}_{W^\perp} u \end{aligned} \quad (6.105)$$

They are called the orthogonal projection of u on W and the orthogonal projection of u on $W^\perp$, respectively.

The vector w_2 is also called the component of u orthogonal to W . The formula for u in term of orthogonal projection is

$$u = \text{proj}_W u + \text{proj}_{W^\perp} u \quad (6.106)$$

or it can also be written as

$$u = \text{proj}_W u + (u - \text{proj}_W u) \quad (6.107)$$

Theorem 6.74: Formulas for Calculating Orthogonal Projections

Let W be a finite-dimensional subspace of an inner product space V .

(a) If $\{v_1, v_2, \dots, v_r\}$ is an orthogonal basis for W , and u is any vector in V , then

$$\text{proj}_W u = \frac{\langle u, v_1 \rangle}{\|v_1\|^2} v_1 + \frac{\langle u, v_2 \rangle}{\|v_2\|^2} v_2 + \dots + \frac{\langle u, v_r \rangle}{\|v_r\|^2} v_r \quad (6.108)$$

(b) If $\{v_1, v_2, \dots, v_r\}$ is an orthonormal basis for W , and u is any vector in V , then

$$\text{proj}_W u = \langle u, v_1 \rangle v_1 + \langle u, v_2 \rangle v_2 + \dots + \langle u, v_r \rangle v_r \quad (6.109)$$

Theorem 6.75: Orthonormal Basis in Inner Product Space

Every nonzero finite-dimensional inner product space has an orthonormal basis.

[SI*] Let W be any nonzero finite-dimensional subspace of an inner product space, and suppose that $\{u_1, u_2, \dots, u_r\}$ is any basis for W . The following steps will produce an orthogonal basis $\{v_1, v_2, \dots, v_r\}$ for W .

1. Let $v_1 = u_1$
2. We can obtain a vector v_2 that is orthogonal to v_1 by computing the component u_2 that is orthogonal to the space W_1 spanned by v_1 .

$$v_2 = u_2 - \text{proj}_{W_1} u_2 = u_2 - \frac{\langle u_2, v_1 \rangle}{\|v_1\|^2} v_1$$

3. To construct a vector v_3 that is orthogonal to both v_1 and v_2 , we compute the component of u_3 orthogonal to the space W_2 spanned by v_1 and v_2 .

$$v_3 = u_3 - \text{proj}_{W_2} u_3 = u_3 - \frac{\langle u_3, v_1 \rangle}{\|v_1\|^2} v_1 - \frac{\langle u_3, v_2 \rangle}{\|v_2\|^2} v_2$$

the linear independence of $\{u_1, u_2, \dots, u_r\}$ ensures that $v_3 \neq 0$. Continuing in this way we will produce an orthogonal set of vectors after r steps. Since orthogonal sets are linearly independent, this set will be an orthogonal basis for the r -dimensional space W .

Definition 6.41: The Gram-Schmidt Process

To convert a basis $\{u_1, u_2, \dots, u_r\}$ into an orthogonal basis $\{v_1, v_2, \dots, v_r\}$, perform the following computations:

$$\begin{aligned} v_1 &= u_1 \\ v_2 &= u_2 - \frac{\langle u_2, v_1 \rangle}{\|v_1\|^2} v_1 \\ v_3 &= u_3 - \frac{\langle u_3, v_1 \rangle}{\|v_1\|^2} v_1 - \frac{\langle u_3, v_2 \rangle}{\|v_2\|^2} v_2 \\ v_4 &= u_4 - \frac{\langle u_4, v_1 \rangle}{\|v_1\|^2} v_1 - \frac{\langle u_4, v_2 \rangle}{\|v_2\|^2} v_2 - \frac{\langle u_4, v_3 \rangle}{\|v_3\|^2} v_3 \\ &\vdots \\ v_r &= u_r - \frac{\langle u_r, v_1 \rangle}{\|v_1\|^2} v_1 - \frac{\langle u_r, v_2 \rangle}{\|v_2\|^2} v_2 - \dots - \frac{\langle u_r, v_{r-1} \rangle}{\|v_{r-1}\|^2} v_{r-1} \end{aligned} \tag{6.110}$$

To convert the orthogonal basis into an orthonormal basis $\{q_1, q_2, \dots, q_r\}$, normalize the orthogonal basis vectors.

Theorem 6.76: Extending Orthonormal Sets to Orthonormal Bases

If W is a finite-dimensional inner product space, then

- (a) Every orthogonal set of nonzero vectors in W can be enlarged to an orthogonal basis for W .
- (b) Every orthonormal set in W can be enlarged to an orthonormal basis for W .

ii. **QR-Decomposition**

- [SI*] The QR-decomposition / factorization is the thread that connects most of the algorithms of numerical linear algebra [28], including methods for least squares, eigenvalue, and singular value problems, as well as iterative methods for all of these and also for systems of equations.
- [SI*] QR-decomposition is very important in the field of numerical algorithms, including those for computing eigenvalues of large matrices.
- [SI*] If A is an $m \times n$ matrix with linearly independent column vectors, and if Q is the matrix that results by applying Gram-Schmidt process to the column vectors of A , what relationship, if any exist between A and Q ?

To solve the problem above, suppose that the column vectors of A are $u_1, u_2, \dots, u_n$ and the orthonormal column vectors of Q are $q_1, q_2, \dots, q_n$. Thus, A and Q can be written in partitioned form as

$$A = [u_1 \mid u_2 \mid \dots \mid u_n]$$

and

$$Q = [q_1 \mid q_2 \mid \dots \mid q_n]$$

it follows that $u_1, u_2, \dots, u_n$ are expressible in terms of the vectors $q_1, q_2, \dots, q_n$ as

$$\begin{aligned} u_1 &= \langle u_1, q_1 \rangle q_1 + \langle u_1, q_2 \rangle q_2 + \dots + \langle u_1, q_n \rangle q_n \\ u_2 &= \langle u_2, q_1 \rangle q_1 + \langle u_2, q_2 \rangle q_2 + \dots + \langle u_2, q_n \rangle q_n \\ &\vdots \\ u_n &= \langle u_n, q_1 \rangle q_1 + \langle u_n, q_2 \rangle q_2 + \dots + \langle u_n, q_n \rangle q_n \end{aligned}$$

Recall that the j -th column vector of a matrix product is a linear combination of the column vectors of the first factor with coefficients coming from the j -th column of the second factor, it follows that these relationships can be expressed in matrix form as

$$[u_1 \mid u_2 \mid \dots \mid u_n] = [q_1 \mid q_2 \mid \dots \mid q_n] \begin{bmatrix} \langle u_1, q_1 \rangle & \langle u_2, q_1 \rangle & \dots & \langle u_n, q_1 \rangle \\ \langle u_1, q_2 \rangle & \langle u_2, q_2 \rangle & \dots & \langle u_n, q_2 \rangle \\ \vdots & \vdots & \dots & \vdots \\ \langle u_1, q_n \rangle & \langle u_2, q_n \rangle & \dots & \langle u_n, q_n \rangle \end{bmatrix}$$

or more briefly as

$$A = QR \tag{6.111}$$

where R is the second factor in the product. This equation tells that it is a factorization of A into the product of a matrix Q with orthonormal column vectors and an invertible upper triangular matrix R . However, it is a property of the Gram-Schmidt process that for $j \geq 2$, the vector q_j is orthogonal to $u_1, u_2, \dots, u_{j-1}$. Thus, all entries below the main diagonal of R are zero, and R has the form

$$\begin{bmatrix} \langle u_1, q_1 \rangle & \langle u_2, q_1 \rangle & \dots & \langle u_n, q_1 \rangle \\ 0 & \langle u_2, q_2 \rangle & \dots & \langle u_n, q_2 \rangle \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & \langle u_n, q_n \rangle \end{bmatrix} \tag{6.112}$$

R is invertible and the diagonal entries are nonzero.

Theorem 6.77: QR-Decomposition

If A is an $m \times n$ matrix with linearly independent column vectors, then A can be factored as

$$A = QR$$

where Q is an $m \times n$ matrix with orthonormal column vectors, and R is an $n \times n$ invertible upper triangular matrix.

[SI*] Some Characteristics of QR-decomposition:

1. QR -decomposition always exists, but may not be unique.
2. If the determinant of matrix A is not 0 and all diagonal entries of $R > 0$, then this decomposition is unique.
3. There are two types of QR -decomposition: full and reduced.

QR -decomposition is also applicable for $m \times n$ matrices, where $m \geq n$. In that case, Q will be a $m \times m$ matrix and R will be an $m \times n$ matrix.

iii. Functions in SymIntegration Related to Inner Product Spaces

[SI*] We will list the basic functions that can be used related to inner product spaces:

gramschmidt(vector<vector<double>> &A)

to compute the orthonormal basis from basis that become the column vector of matrix A .

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gram-Schmidt ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gram-Schmidt ]# ./main

A:
      1.000000      0.000000      0.000000
      1.000000      1.000000      0.000000
      1.000000      1.000000      1.000000

Orthogonal basis:
      1.000000      -0.666667      0.000000
      1.000000      0.333333      -0.500000
      1.000000      0.333333      0.500000

Orthonormal basis:
      0.577350      -0.816497      0.000000
      0.577350      0.408248      -0.707107
      0.577350      0.408248      0.707107

Time taken by function: 3110 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gram-Schmidt ]#
```

Figure 6.48: The computation for Gram-Schmidt orthogonalization method. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Gram-Schmidt/main.cpp*).

QRDecomposition(vector<vector<double>> &A, vector<vector<double>> &Q, vector<vector<double>> &R)

to compute the QR -decomposition of matrix A .

QRDecompositionComplex(ComplexMatrix &, ComplexMatrix &, ComplexMatrix &)

to compute the QR -decomposition of a complex matrix A .

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration QR Decomposition Complex Eigenvalues Manual ]# ./main

Matrix A :
(1,0)      (0,0)      (0,0)
(1,0)      (1,0)      (0,0)
(1,0)      (1,0)      (1,0)

Matrix Q :
(0.577,0)  (-0.816,0) (-2.36e-16,0)
(0.577,0)  (0.408,0)  (-0.707,0)
(0.577,0)  (0.408,0)  (0.707,0)

Matrix R :
(1.73,0)   (1.15,0)   (0.577,0)
(0,0)      (0.816,0)   (0.408,0)
(0,0)      (0,0)      (0.707,0)

Time taken by function: 1128 microseconds

```

Figure 6.49: The QR -decomposition of a complex matrix A . (*SymIntegration/Examples/Linear Algebra/Test SymIntegration QR Decomposition Complex Matrix/main.cpp*).

QRDecompositionHouseholderReflections(vector<vector<complex<double>>>& A)
to compute the QR -decomposition of a complex matrix A with Householder reflections.

iv. Compute QR-Decomposition Using Gram-Schmidt with SymIntegration

Find the QR-decomposition of

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$$

Solution:

The column vectors of A are

$$u_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \quad u_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \quad u_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Applying the Gram-Schmidt process with normalization to these column vectors yields the orthonormal vectors

$$q_1 = \begin{bmatrix} \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{bmatrix}, \quad q_2 = \begin{bmatrix} -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix}, \quad q_3 = \begin{bmatrix} 0 \\ -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

Thus

$$R = \begin{bmatrix} \langle u_1, q_1 \rangle & \langle u_2, q_1 \rangle & \langle u_3, q_1 \rangle \\ 0 & \langle u_2, q_2 \rangle & \langle u_3, q_2 \rangle \\ 0 & 0 & \langle u_3, q_3 \rangle \end{bmatrix} = \begin{bmatrix} \frac{3}{\sqrt{3}} & \frac{2}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ 0 & \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ 0 & 0 & \frac{1}{\sqrt{2}} \end{bmatrix}$$

It follows that the QR-decomposition of A is

$$\begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{3}} & -\frac{2}{\sqrt{6}} & 0 \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \frac{3}{\sqrt{3}} & \frac{2}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ 0 & \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ 0 & 0 & \frac{1}{\sqrt{2}} \end{bmatrix}$$

To compute the QR-decomposition in SymIntegration we use this function:

QRDecomposition(vector<vector<double>> &A, vector<vector<double>> &Q, vector<vector<double>> &R)

With A as the input matrix that we want to decompose, matrices Q and R will be the decomposition result but we need to declare them first as square matrices with same size as A before we call the function.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration QR Decomposition ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration QR Decomposition ]# ./main

A:
      1.000000      0.000000      0.000000
      1.000000      1.000000      0.000000
      1.000000      1.000000      1.000000

Q:
      0.577350     -0.816497      0.000000
      0.577350      0.408248     -0.707107
      0.577350      0.408248      0.707107

R:
      1.732051      1.154701      0.577350
      0.000000      0.816497      0.408248
      0.000000      0.000000      0.707107

Time taken by function: 1341 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration QR Decomposition ]# 

```

Figure 6.50: The computation to obtain the QR-decomposition of matrix A with Gram-Schmidt orthogonalization method. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration QR Decomposition/main.cpp*).

Implementing QR-decomposition in this SymIntegration' function is using the Gram-Schmidt orthogonalization, another method is by using Householder transformations. Gram-Schmidt is simpler to implement but less numerically stable. Libraries like LAPACK or Eigen are using Householder reflections for their stability. As a comparison, in another section we will talk about SymIntegration' function to compute QR-decomposition with Householder reflections / transformations.

v. Compute QR-Decomposition Using Householder Reflections with SymIntegration

Find the QR-decomposition with Householder reflections of

$$A = \begin{bmatrix} -1 & -1 & 1 \\ 1 & 3 & 3 \\ -1 & -1 & 5 \end{bmatrix}$$

Solution:

Suppose we define matrix A with size $m \times n$.

The number of iterations depends on the number of row and the number of column, if $m = n$ then we will have $m - 1$ iterations, if $m < n$ then we will have m iterations.

First, at iteration $j = 1$ we will have

$$x = \begin{bmatrix} -1 \\ 1 \\ -1 \end{bmatrix}$$

(we define vector x as the first column of matrix A with criteria, if $j > 1$ then $x(1 : j - 1) = 0$, e.g. at iteration $j = 2$ we will take the second column of matrix A for vector x and we will make the first entry of vector x to be zero, at iteration $j = 3$ we will take the second column of matrix A for vector x and we will make the first and second entry of vector x to be zero).

then

$$w = \pm \|x\| e_1 \begin{bmatrix} \sqrt{3} \\ 0 \\ 0 \end{bmatrix}$$

Construct the householder vector v

$$v = x - w \begin{bmatrix} -2.7321 \\ 1 \\ -1 \end{bmatrix}$$

Construct the Householder reflector H_j

$$H_1 = I - 2 \frac{vv^T}{v^T v} = \begin{bmatrix} -0.5774 & 0.5774 & -0.5774 \\ 0.5774 & 0.7887 & 0.211 \\ -0.5774 & -0.211 & 0.7887 \end{bmatrix}$$

we will obtain

$$R_1 = H_1 A = \begin{bmatrix} 1.73 & 2.89 & -1.73 \\ 0 & 1.58 & 4 \\ 0 & 0.423 & 4 \end{bmatrix}$$

and

$$Q_1 = H_1 = \begin{bmatrix} -0.5774 & 0.5774 & -0.5774 \\ 0.5774 & 0.7887 & 0.211 \\ -0.5774 & -0.211 & 0.7887 \end{bmatrix}$$

we will set $A = R$ for the next iteration.

Continuing at iteration $j = 2$

$$x = \begin{bmatrix} 0 \\ 1.58 \\ 0.423 \end{bmatrix}$$

$$w = \pm \|x\| e_2 \begin{bmatrix} 0 \\ -1.63 \\ 0 \end{bmatrix}$$

Construct the householder vector v

$$v = x - w \begin{bmatrix} 0 \\ 3.21 \\ 0.423 \end{bmatrix}$$

Construct the Householder reflector H_j

$$H_2 = I - 2 \frac{vv^T}{v^T v} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -0.966 & -0.259 \\ 0 & -0.259 & 0.966 \end{bmatrix}$$

we will obtain

$$R_2 = H_2 R_1 = \begin{bmatrix} 1.73 & 2.89 & -1.73 \\ 0 & -1.63 & -4.9 \\ 0 & 0 & 2.83 \end{bmatrix}$$

and

$$Q_2 = H_1 H_2 = \begin{bmatrix} -0.5774 & -0.408 & -0.707 \\ 0.5774 & -0.816 & 0 \\ -0.5774 & -0.408 & 0.707 \end{bmatrix}$$

At the end of iteration we will obtain the QR-decomposition:

$$Q = H_1 H_2$$

and

$$R = H_2 H_1 A$$

This method uses a sequence of reflections to zero out the entries below the diagonal of the matrix A . Each reflection is a transformation that flips the matrix across a hyperplane. The sequence of reflections is stored in the orthogonal matrix Q , and the result of these reflections is the upper triangular matrix R . This method is generally more numerically stable than the Gram-Schmidt process.

Implementing Householder transformations is significantly more complex than Gram-Schmidt but yields better results for real-world numerical problems.

In SymIntegration, we can compute the QR-decomposition with Householder reflections with this function:

QRDecompositionHouseholderReflections(vector<vector<complex<double>>>& A)

with A as the matrix that we want to decompose.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration QR Algorithm with Householder Reflections ]# ./main

Matrix A :
      (-1,0)      (-1,0)      (1,0)
      (1,0)       (3,0)       (3,0)
      (-1,0)      (-1,0)      (5,0)

Iteration : 1

x:
(-1,0)
(1,0)
(-1,0)

w:
(1.73,0)
(0,0)
(0,0)

v:
(-2.73,0)
(1,0)
(-1,0)

R:
      (1.73,0)      (2.89,0)      (-1.73,0)
      (0,0)       (1.58,0)       (4,0)
      (0,0)       (0.423,0)      (4,0)

Householder Matrix / Q:
      (-0.577,0)      (0.577,0)      (-0.577,0)
      (0.577,0)      (0.789,0)      (0.211,0)
      (-0.577,0)      (0.211,0)      (0.789,0)
```

Figure 6.51: The computation to obtain the QR-decomposition of matrix A with Householder reflections method. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration QR Decomposition with Householder Reflections/main.cpp*).

```

Iteration : 2

x:
(0,0)
(1.58,0)
(0.423,0)

w:
(0,0)
(-1.63,0)
(0,0)

v:
(0,0)
(3.21,0)
(0.423,0)

R:
(1.73,0) (2.89,0) (-1.73,0)
(0,0) (-1.63,0) (-4.9,0)
(0,0) (0,0) (2.83,0)

Householder Matrix / Q:
(-0.577,0) (-0.408,0) (-0.707,0)
(0.577,0) (-0.816,0) (0,0)
(-0.577,0) (-0.408,0) (0.707,0)

*****

End of iteration

*****

Q:
(-0.577,0) (-0.408,0) (-0.707,0)
(0.577,0) (-0.816,0) (0,0)
(-0.577,0) (-0.408,0) (0.707,0)

R:
(1.73,0) (2.89,0) (-1.73,0)
(0,0) (-1.63,0) (-4.9,0)
(0,0) (0,0) (2.83,0)

Q * R:
(-1,0) (-1,0) (1,0)
(1,0) (3,0) (3,0)
(-1,0) (-1,0) (5,0)

Time taken by function: 2877 microseconds

```

Figure 6.52: The computation to obtain the QR-decomposition of matrix A with Householder reflections method. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration QR Decomposition with Householder Reflections/main.cpp*).

At the heart of the Householder method is an idea originally proposed by Alston Householder in 1958. This is an ingenious way of designing the unitary matrices Q_k so that $Q_n \dots Q_2 Q_1 A = R$ is an upper-triangular.

The matrix Q_k is chosen to introduce zeros below the diagonal in the k -th column while preserving all the zeros previously introduced [28].

VII. DIAGONALIZATION AND QUADRATIC FORMS

[SI*] We have found conditions that guaranteed the diagonalizability of an $n \times n$ matrix, now we will check what class or classes of matrices might actually satisfy those conditions.

Definition 6.42: Orthogonal Matrix

A square matrix A is said to be orthogonal if its transpose is the same as its inverse, that is, if

$$A^{-1} = A^T$$

or, equivalently, if

$$AA^T = A^T A = I$$

[SI*] The standard matrix for the counterclockwise rotation of $\mathbb{R}^2$ through an angle θ is

$$A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

This matrix is orthogonal for all choices of θ since

$$A^T A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Theorem 6.78: Equivalent Statement for Orthogonal Matrix

The following are equivalent for an $n \times n$ matrix A .

- (a) A is orthogonal.
- (b) The row vectors of A form an orthonormal set in $\mathbb{R}^n$ with the Euclidean inner product.
- (c) The column vectors of A form an orthonormal set in $\mathbb{R}^n$ with the Euclidean inner product.

[SI*] If the row vectors of A are $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_n$ then AA^T can be expressed as

$$AA^T = \begin{bmatrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_n \end{bmatrix} \begin{bmatrix} \mathbf{r}_1 & \mathbf{r}_2 & \dots & \mathbf{r}_n \end{bmatrix} = \begin{bmatrix} \mathbf{r}_1 \cdot \mathbf{r}_1 & \mathbf{r}_1 \cdot \mathbf{r}_2 & \dots & \mathbf{r}_1 \cdot \mathbf{r}_n \\ \mathbf{r}_2 \cdot \mathbf{r}_1 & \mathbf{r}_2 \cdot \mathbf{r}_2 & \dots & \mathbf{r}_2 \cdot \mathbf{r}_n \\ \vdots & \vdots & & \vdots \\ \mathbf{r}_n \cdot \mathbf{r}_1 & \mathbf{r}_n \cdot \mathbf{r}_2 & \dots & \mathbf{r}_n \cdot \mathbf{r}_n \end{bmatrix}$$

it follows that $AA^T = I$ if and only if

$$\mathbf{r}_1 \cdot \mathbf{r}_1 = \mathbf{r}_2 \cdot \mathbf{r}_2 = \dots = \mathbf{r}_n \cdot \mathbf{r}_n = 1$$

and

$$\mathbf{r}_i \cdot \mathbf{r}_j = 0, \quad i \neq j$$

which are true if and only if $\{\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_n\}$ is an orthonormal set in $\mathbb{R}^n$.

Theorem 6.79: Fundamental Properties of Orthogonal Matrices

- (a) The inverse of an orthogonal matrix is orthogonal.
- (b) A product of orthogonal matrices is orthogonal.
- (c) If A is orthogonal, then $\det(A) = 1$ or $\det(A) = -1$.

Theorem 6.80: Orthogonal Matrices as Linear Operators

If A is an $n \times n$ matrix, then the following are equivalent

- (a) A is orthogonal.
- (b) $\|Ax\| = \|x\|$ for all x in $\mathbb{R}^n$.
- (c) $Ax \cdot Ay = x \cdot y$ for all x and y in $\mathbb{R}^n$.

[SI*] If A is orthogonal matrix and $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is multiplication by A , then we will call T_A orthogonal operator on $\mathbb{R}^n$.

The orthogonal operators on $\mathbb{R}^n$ are precisely those operators that leave the lengths of all vectors unchanged.

[SI*] Orthonormal bases for inner product spaces are convenient because many familiar formulas hold for such bases.

Theorem 6.81: Change of Orthonormal Basis

If S is an orthonormal basis for an n -dimensional inner product space V , and if

$$(u)_S = (u_1, u_2, \dots, u_n)$$

$$(v)_S = (v_1, v_2, \dots, v_n)$$

then:

- (a) $\|u\| = \sqrt{u_1^2 + u_2^2 + \dots + u_n^2}$
- (b) $d(u, v) = \sqrt{(u_1 - v_1)^2 + (u_2 - v_2)^2 + \dots + (u_n - v_n)^2}$
- (c) $\langle u, v \rangle = u_1 v_1 + u_2 v_2 + \dots + u_n v_n$

[SI*] Transitions between orthonormal bases for an inner product space are of special importance in geometry and various applications.

Theorem 6.82: Transition Matrix for Orthonormal Basis

Let V be a finite-dimensional inner product space. If P is the transition matrix from one orthonormal basis for V to another orthonormal basis for V , then P is an orthogonal matrix.

[SI*] **Example:**

What conditions must a and b satisfy for the matrix

$$\begin{bmatrix} a+b & b-a \\ a-b & b+a \end{bmatrix}$$

to be orthogonal?

Solution:

We will compute $A^T A$ then determine a and b from there

$$A^T A = \begin{bmatrix} 2(a^2 + b^2) & 0 \\ 0 & 2(a^2 + b^2) \end{bmatrix}$$

To be orthogonal the condition of $A^T A = I$ has to be fulfilled, thus

$$2(a^2 + b^2) = 1$$

$$a^2 + b^2 = \frac{1}{2}$$

so a and b must satisfy $a^2 + b^2 = \frac{1}{2}$.

[SI*] **Example:**

Find a, b , and c for which the matrix

$$\begin{bmatrix} a & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ b & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ c & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}$$

is orthogonal. Are the values of a, b , and c are unique?

Solution:

The row vectors of an orthogonal matrix are an orthonormal set. Thus

$$\|r_1\|^2 = 1$$

$$a^2 + \left(\frac{1}{\sqrt{2}}\right)^2 + \left(-\frac{1}{\sqrt{2}}\right)^2 = 1$$

$$a^2 + 1 = 1$$

$$a = 0$$

$$\|r_2\|^2 = 1$$

$$b^2 + \left(\frac{1}{\sqrt{6}}\right)^2 + \left(\frac{1}{\sqrt{6}}\right)^2 = 1$$

$$b^2 + \frac{2}{6} = 1$$

$$b = \pm \frac{2}{\sqrt{6}}$$

$$\|r_3\|^2 = 1$$

$$c^2 + \left(\frac{1}{\sqrt{3}}\right)^2 + \left(\frac{1}{\sqrt{3}}\right)^2 = 1$$

$$c^2 + \frac{2}{3} = 1$$

$$c = \pm \frac{1}{\sqrt{3}}$$

The column vectors of an orthogonal matrix are an orthonormal set, from which it is clear that b and c must have opposite signs. Thus the only possibilities are

$$a = 0, \quad b = \frac{2}{\sqrt{6}}, \quad c = -\frac{1}{\sqrt{3}}$$

or

$$a = 0, \quad b = -\frac{2}{\sqrt{6}}, \quad c = \frac{1}{\sqrt{3}}$$

- [SI*] Multiplication by a 3×3 orthogonal matrix A is a rotation about some axis if $\det(A) = 1$ and is a rotation about some axis followed by a reflection about some coordinate plane if $\det(A) = -1$.
- [SI*] A linear operator on $\mathbb{R}^2$ is called rigid if it does not change the lengths of vectors, and it is called angle preserving if it does not change the angle between nonzero vectors.

1. Linear operators that are rigid: rotations about the origin, reflections about any line through the origin.
2. Linear operators that are angle preserving: All rigid operators on $\mathbb{R}^2$ are angle preserving. Dilations and contractions are angle preserving operators that are not rigid.

Definition 6.43: Orthogonally Similar

If A and B are square matrices, then we say that A and B are orthogonally similar if there is an orthogonal matrix P such that $P^T A P = B$.

- [SI*] If A is orthogonally similar to some diagonal matrix, say

$$P^T A P = D$$

then we say that A is orthogonally diagonalizable and that P orthogonally diagonalizes A .

Theorem 6.83: Conditions for Orthogonal Diagonalizability

If A is an $n \times n$ matrix, then the following are equivalent

- (a) A is orthogonally diagonalizable.
- (b) A has an orthonormal set of n eigenvectors.
- (c) A is symmetric.

Theorem 6.84: Properties of Symmetric Matrices

If A is a symmetric matrix, then:

- (a) The eigenvalues of A are all real numbers.
- (b) Eigenvectors from different eigenspaces are orthogonal.

- [SI*] Procedure for orthogonally diagonalizing a symmetric matrix.

1. Find a basis for each eigenspace of A .

2. Apply the Gram-Schmidt process to each of these bases to obtain an orthonormal basis for each eigenspace.
3. Form the matrix P whose columns are the vectors constructed in step 2. This matrix will orthogonally diagonalize A , and the eigenvalues on the diagonal of $D = P^T A P$ will be in the same order as their corresponding eigenvectors in P .

Eigenvectors from different eigenspaces are orthogonal, and applying Gram-Schmidt process ensures that the eigenvectors within the same eigenspace are orthonormal. It follows that the entire set of eigenvectors obtained by this procedure will be orthonormal.

[SI*] If A is a symmetric matrix that is orthogonally diagonalized by

$$P = [\mathbf{u}_1 \ \mathbf{u}_2 \ \dots \ \mathbf{u}_n]$$

and if $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of A corresponding to the unit eigenvectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$, then we know that $D = P^T A P$, where D is a diagonal matrix with the eigenvalues in the diagonal positions. It follows from this that the matrix A can be expressed as

$$A = P D P^T = [\mathbf{u}_1 \ \mathbf{u}_2 \ \dots \ \mathbf{u}_n] \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix} \begin{bmatrix} \mathbf{u}_1^T \\ \mathbf{u}_2^T \\ \vdots \\ \mathbf{u}_n^T \end{bmatrix} = [\lambda_1 \mathbf{u}_1 \ \lambda_2 \mathbf{u}_2 \ \dots \ \lambda_n \mathbf{u}_n] \begin{bmatrix} \mathbf{u}_1^T \\ \mathbf{u}_2^T \\ \vdots \\ \mathbf{u}_n^T \end{bmatrix}$$

Multiplying out, we obtain the formula

$$A = \lambda_1 \mathbf{u}_1 \mathbf{u}_1^T + \lambda_2 \mathbf{u}_2 \mathbf{u}_2^T + \dots + \lambda_n \mathbf{u}_n \mathbf{u}_n^T \quad (6.113)$$

which is called a spectral decomposition of A .

Note that in each term of the spectral decomposition of A has the form $\lambda \mathbf{u} \mathbf{u}^T$, where $\mathbf{u}$ is a unit eigenvector of A in column form, and λ is an eigenvalue of A corresponding to $\mathbf{u}$. Since $\mathbf{u}$ has size $n \times 1$, it follows that the product $\mathbf{u} \mathbf{u}^T$ has size $n \times n$.

It can be proved that $\mathbf{u} \mathbf{u}^T$ is the standard matrix for the orthogonal projection of $\mathbb{R}^n$ on the subspace spanned by the vector $\mathbf{u}$.

The spectral decomposition of A tells that the image of a vector $\mathbf{x}$ under multiplication by a symmetric matrix A can be obtained by projecting $\mathbf{x}$ orthogonally on the lines (one-dimensional subspaces) determined by the eigenvectors of A , then scaling those projections by the eigenvalues, and then adding the scaled projections.

The terminology spectral decomposition is derived from the fact that the set of all eigenvalues of a matrix A is sometimes called the spectrum of A . The terminology eigenvalue decomposition is due to a paper entitled "A Singularly Valuable Decomposition: The SVD of a Matrix," written by Professor Dan Kalman.

[SI*] If A is an $n \times n$ matrix that is not orthogonally diagonalizable, it may still be possible to achieve considerable simplification in the form of $P^T A P$ by choosing the orthogonal matrix P appropriately.

Theorem 6.85: Schur's Theorem

If A is an $n \times n$ matrix with real entries and real eigenvalues, then there is an orthogonal matrix P such that $P^T A P$ is an upper triangular matrix of the form

$$P^T A P = \begin{bmatrix} \lambda_1 & X & X & \dots & X \\ 0 & \lambda_2 & X & \dots & X \\ 0 & 0 & \lambda_3 & \dots & X \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \lambda_n \end{bmatrix} \quad (6.114)$$

in which $\lambda_1, \lambda_2, \dots, \lambda_n$ are eigenvalues of the matrix A repeated according to multiplicity.

[SI*] It is common to denote the upper triangular matrix in Schur's Theorem by S in which case that the equation from Schur's Theorem can be rewritten as

$$A = P S P^T \quad (6.115)$$

which is called a Schur decomposition of A .

Theorem 6.86: Hessenberg's Theorem

If A is an $n \times n$ matrix, then there is an orthogonal matrix P such that $P^T A P$

$$P^T A P = \begin{bmatrix} X & X & \dots & X & X & X \\ X & X & \dots & X & X & X \\ 0 & X & \ddots & X & X & X \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & X & X & X \\ 0 & 0 & \dots & 0 & X & X \end{bmatrix} \quad (6.116)$$

or in other words every square matrix with real entries is orthogonally similar to a matrix in which each entry below the first subdiagonal is zero. Such a matrix is said to be in upper Hessenberg form.

* The first subdiagonal is $\{a_{i+1,i}\}$ with $i = 1, 2, \dots, n$.

[SI*] It is common to denote the upper Hessenberg matrix by H , in which case that equation can be rewritten as

$$A = P H P^T \quad (6.117)$$

which is called an upper Hessenberg decomposition of A .

In many numerical algorithms the initial matrix is first converted to upper Hessenberg form to reduce the amount of computation in subsequent parts of the algorithm.

[SI*] **Example:**

Let A be a diagonalizable matrix with the property that eigenvectors from distinct eigenvalues are orthogonal. Must A be symmetric?

Solution:

Suppose A is a matrix with size of $n \times n$, if we want to diagonalize A , then we will have to:

1. Compute the eigenvectors of A :

$$\{\lambda_1, \lambda_2, \dots, \lambda_n\}$$

2. Find the basis for each eigenspace of A by solving

$$(\lambda_i I - A)x = \mathbf{0}$$

with $i = 1, 2, \dots, n$. The basis $\{v_1, v_2, \dots, v_n\}$ will be called eigenvectors of A .

3. Apply the Gram-Schmidt process to each of the bases / the eigenvectors we obtain from step (2) to convert $\{v_1, v_2, \dots, v_n\}$ into an orthogonal set $\{u_1, u_2, \dots, u_n\}$.

We can then normalize $\{u_1, u_2, \dots, u_n\}$ to become an orthonormal set.

4. The matrix P that diagonalizes A consist of the orthonormal set $\{u_1, u_2, \dots, u_n\}$ as the column vectors.

$$P = [u_1 \mid u_2 \mid \dots \mid u_n]$$

The Gram-Schmidt process will ensure that columns of P corresponding to the same eigenvalue are an orthonormal set. Since eigenvectors from distinct eigenvalues are orthogonal, this means that P will be an orthogonal matrix. Then since A is orthogonally diagonalizable, it must be symmetric. From Theorem that states if A is an $n \times n$ matrix, whenever A is orthogonally diagonalizable with n orthonormal set of eigenvectors then A is symmetric as well.

GlanzFreya' Guide 6.8: Hessenberg Decomposition

If A is an $n \times n$ matrix that is not orthogonally diagonalizable, we can use Hessenberg decomposition to decompose every square matrix with real entries into a matrix in which each entry below the first subdiagonal is zero.

$$A = PHP^T$$

and

$$H = P^T AP$$

matrices H and A are orthogonally similar.

The algorithm to obtain matrix H :

1. Set variable:
 $row = 1$;
2. Iterate from first column to the second-to-last column, $j = 0, 1, \dots, n - 2$.

Extract a sub-column from matrix A and assign it as vector x , starting from entry at row $j + 1$ down to the bottom. Vector x will have the length of $j - row$.

3. Create vector w from vector x ,

$$w = \begin{bmatrix} \pm \|x\| \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

Vector w is a vector with the first entry is the norm of vector x with sign either plus or minus, and the rest entries are zero, the sign is depending on the first entry of vector x , if the first entry of vector x is negative then the sign will be positive at vector w , if the first entry of vector x is positive then the sign will be negative at vector w .

4. Compute vector v
5. Construct a Householder reflector

$$\hat{H}_{j+1} = I_{n-row} - 2 \frac{vv^T}{v^T v}$$

thus

$$H_{j+1} = \begin{bmatrix} 1 & 0 \\ 0 & \hat{H}_{j+1} \end{bmatrix}$$

$row = row + 1$, $H_{j+1}AH_{j+1} \mapsto A$, we will use this $H_{j+1}AH_{j+1}$ as A at the next iteration, repeat till the last iteration.

After finish we will have

$$P = H_1 H_2 \dots H_n$$

$$P^T = H_n H_{n-1} \dots H_1$$

i. Quadratic Forms

[SI*] Expressions of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n$$

occurred in our study of linear equations and linear systems. If $a_1, a_2, \dots, a_n$ are treated as fixed constants, then this expression is a real-valued function of the n variables $x_1, x_2, \dots, x_n$ and is called a linear form on $\mathbb{R}^n$.

All variables in a linear form occur to the first power and there are no products of variables.

[SI*] Quadratic forms on $\mathbb{R}^n$ are functions of the form

$$a_1x_1^2 + a_2x_2^2 + \cdots + a_nx_n^2 + \sum a_kx_ix_j$$

The notation of $\sum a_kx_ix_j$ represents all possible terms of $a_kx_ix_j$ in which $x_i \neq x_j$.

The terms of the form $a_kx_ix_j$ are called cross product terms. It is common to combine the cross product terms involving x_ix_j with those involving x_jx_i to avoid duplication.

[SI*] A general quadratic form on $\mathbb{R}^2$ would typically be expressed as

$$a_1x_1^2 + a_2x_2^2 + 2a_3x_1x_2 \quad (6.118)$$

in matrix form it can be expressed as

$$\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} a_1 & a_3 \\ a_3 & a_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{x}^T A \mathbf{x}$$

and a general quadratic form on $\mathbb{R}^3$ can be expressed as

$$a_1x_1^2 + a_2x_2^2 + a_3x_3^2 + 2a_4x_1x_2 + 2a_5x_1x_3 + 2a_6x_2x_3 \quad (6.119)$$

or, in matrix form

$$\begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix} \begin{bmatrix} a_1 & a_4 & a_5 \\ a_4 & a_2 & a_6 \\ a_5 & a_6 & a_3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \mathbf{x}^T A \mathbf{x}$$

The matrix A in these formulas is symmetric, that its diagonal entries are the coefficients of the squared terms, and its off-diagonal entries are half the coefficients of the cross product terms.

[SI*] In general, if A is a symmetric $n \times n$ matrix and $\mathbf{x}$ is an $n \times 1$ column vector of variables, then we call the function

$$Q_A(\mathbf{x}) = \mathbf{x}^T A \mathbf{x} \quad (6.120)$$

the quadratic form associated with A . It can be expressed in dot product notation as

$$\mathbf{x}^T A \mathbf{x} = \mathbf{x} \cdot A \mathbf{x} = A \mathbf{x} \cdot \mathbf{x} \quad (6.121)$$

In the case where A is a diagonal matrix, the quadratic form $\mathbf{x}^T A \mathbf{x}$ has no cross product terms; for example, if A has diagonal entries $\lambda_1, \lambda_2, \dots, \lambda_n$, then

$$\begin{aligned} \mathbf{x}^T A \mathbf{x} &= \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \\ &= \lambda_1x_1^2 + \lambda_2x_2^2 + \cdots + \lambda_nx_n^2 \end{aligned}$$

[SI*] There are three kinds of problems that occur in applications of quadratic forms:

1. If $\mathbf{x}^T A \mathbf{x}$ is a quadratic form on $\mathbb{R}^2$ or $\mathbb{R}^3$, what kind of curve or surface is represented by the equation $\mathbf{x}^T A \mathbf{x} = k$?
2. If $\mathbf{x}^T A \mathbf{x}$ is a quadratic form on $\mathbb{R}^n$, what conditions must A satisfy for $\mathbf{x}^T A \mathbf{x}$ to have positive values for $\mathbf{x} \neq \mathbf{0}$?
3. If $\mathbf{x}^T A \mathbf{x}$ is a quadratic form on $\mathbb{R}^n$, what are its maximum and minimum values if $\mathbf{x}$ is constrained to satisfy $\|\mathbf{x}\| = 1$?

[SI*] Many of the techniques for solving the problems above are based on simplifying the quadratic form $\mathbf{x}^T A \mathbf{x}$ by making a substitution

$$\mathbf{x} = P\mathbf{y} \quad (6.122)$$

that expresses the variables $x_1, x_2, \dots, x_n$ in terms of new variables $y_1, y_2, \dots, y_n$. If P is invertible, then we call () a change of variable, and if P is orthogonal, then we call () an orthogonal change of variable.

If we change the variable $\mathbf{x} = P\mathbf{y}$ in the quadratic form $\mathbf{x}^T A \mathbf{x}$, then we obtain

$$\mathbf{x}^T A \mathbf{x} = (P\mathbf{y})^T A (P\mathbf{y}) = \mathbf{y}^T P^T A P \mathbf{y} = \mathbf{y}^T (P^T A P) \mathbf{y} \quad (6.123)$$

Since the matrix $B = P^T A P$ is symmetric, the effect of the change of variable is to produce a new quadratic form $\mathbf{y}^T B \mathbf{y}$ in the variables $y_1, y_2, \dots, y_n$.

In particular, if we choose P to orthogonally diagonalize A , then the new quadratic form will be $\mathbf{y}^T D \mathbf{y}$, where D is a diagonal matrix with the eigenvalues of A on the main diagonal; that is,

$$\begin{aligned} \mathbf{x}^T A \mathbf{x} &= \mathbf{y}^T D \mathbf{y} = \begin{bmatrix} y_1 & y_2 & \cdots & y_n \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \\ \mathbf{x}^T A \mathbf{x} &= \mathbf{y}^T D \mathbf{y} = \lambda_1 y_1^2 + \lambda_2 y_2^2 + \cdots + \lambda_n y_n^2 \end{aligned}$$

Theorem 6.87: The Principal Axes Theorem

If A is a symmetric $n \times n$ matrix, then there is an orthogonal change of variable that transforms the quadratic form $\mathbf{x}^T A \mathbf{x}$ into a quadratic form $\mathbf{y}^T D \mathbf{y}$ with no cross product terms. Specifically, if P orthogonally diagonalizes A , then making the change of variable $\mathbf{x} = P\mathbf{y}$ in the quadratic form $\mathbf{x}^T A \mathbf{x}$ yields the quadratic form

$$\mathbf{x}^T A \mathbf{x} = \mathbf{y}^T D \mathbf{y} = \lambda_1 y_1^2 + \lambda_2 y_2^2 + \cdots + \lambda_n y_n^2 \quad (6.124)$$

in which $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of A corresponding to the eigenvectors that form the successive columns of P .

[SI*] If A is a symmetric $n \times n$ matrix, then the quadratic form $\mathbf{x}^T A \mathbf{x}$ is a real-valued function whose range is the set of all possible values for $\mathbf{x}^T A \mathbf{x}$ as $\mathbf{x}$ varies over $\mathbb{R}^n$.

It can be shown that an orthogonal change of variable $\mathbf{x} = P\mathbf{y}$ does not alter the range of a quadratic form; that is, the set of all values for $\mathbf{x}^T A \mathbf{x}$ as $\mathbf{x}$ varies over $\mathbb{R}^n$ is the same as the set of all values for $\mathbf{y}^T (P^T A P) \mathbf{y}$ as $\mathbf{y}$ varies over $\mathbb{R}^n$.

[SI*] Recall that a conic section or conic is a curve that results by cutting a double-napped cone with a plane. The most important conic sections are ellipses, hyperbolas, and parabolas, which result when the cutting plane does not pass through the vertex.

Circles are special cases of ellipses that results when the cutting plane is perpendicular to the axis of symmetry of the cone. If the cutting plane passes through the vertex, then the resulting intersection is called a degenerate conic. The possibilities are a point, a pair of intersecting lines or a single line.

[SI*] Quadratic forms in $\mathbb{R}^2$ arise naturally in the study of conic sections. For example it is shown in analytic geometry that an equation of the form

$$ax^2 + 2bxy + cy^2 + dx + ey + f = 0 \quad (6.125)$$

in which a, b , and c are not all zero, represents a conic section. If $d = e = 0$ in (6.125) then there are no linear terms, so the equation becomes

$$ax^2 + 2bxy + cy^2 + f = 0 \quad (6.126)$$

and is said to represent central conic. These include circles, ellipses, and hyperbolas, but not parabolas. Furthermore, if $b = 0$, then there is no cross product term, and the equation

$$ax^2 + cy^2 + f = 0 \quad (6.127)$$

is said to represent a central conic in standard position.

If we take the constant f in Equations (6.126) and (6.127) to the right side and let $k = -f$, then we can rewrite these equations in matrix form as

$$\begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} a & b \\ b & c \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = k \quad (6.128)$$

this one corresponds to Equation (6.126) in which there is a cross product term $2bxy$.

$$\begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & c \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = k \quad (6.129)$$

This one corresponds to Equation (6.127) in which there is no cross product term. Geometrically, the existence of a cross product term signals that the graph of the quadratic form is rotated about the origin.

The three-dimensional analogs of the Equation (6.128) is

$$\begin{bmatrix} x & y & z \end{bmatrix} \begin{bmatrix} a & d & e \\ d & b & f \\ e & f & c \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = k \quad (6.130)$$

The three-dimensional analogs of the Equation (6.129) is

$$\begin{bmatrix} x & y & z \end{bmatrix} \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = k \quad (6.131)$$

If a, b , and c are not all zero, then the graphs of these equations in $\mathbb{R}^3$ are called central quadrics in standard position.

[SI*] An equation of the form

$$ax^2 + 2bxy + cy^2 + f = 0 \quad (6.132)$$

represents a central conic. If $b = 0$, then the conic is in standard position, and if $b \neq 0$, it is rotated. It is an easy matter to identify central conics in standard position by matching the equation with one of the standard forms:

Ellipse

$$\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1 \quad (6.133)$$

with $\alpha > 0, \beta > 0$.

Hyperbola type 1

$$\frac{x^2}{\alpha^2} - \frac{y^2}{\beta^2} = 1 \quad (6.134)$$

with $\alpha > 0, \beta > 0$.

Hyperbola type 2

$$\frac{y^2}{\beta^2} - \frac{x^2}{\alpha^2} = 1 \quad (6.135)$$

with $\alpha > 0, \beta > 0$.

[SI*] The equation

$$9x^2 + 16y^2 - 144 = 0$$

can be rewritten as

$$\frac{x^2}{16} + \frac{y^2}{9} = 1$$

which is the ellipse with $\alpha = 4, \beta = 3$.

[SI*] If a central conic is rotated out of standard position, then it can be identified by first rotating the coordinate axes to put it in standard position and then matching the resulting equation with one of standard forms.

To find a rotation that eliminates the cross product term in the equation

$$ax^2 + 2bxy + cy^2 = k \quad (6.136)$$

it will be convenient to express the equation in the matrix form

$$\mathbf{x}^T A \mathbf{x} = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} a & b \\ b & c \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = k \quad (6.137)$$

and look for a change of variable

$$\mathbf{x} = P \mathbf{x}'$$

that diagonalizes A and for which $\det(P) = 1$. The transition matrix

$$P = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \quad (6.138)$$

has the effect of rotating the xy -axes of a rectangular coordinate system through an angle θ , our problem reduces to finding θ that diagonalizes A , thereby eliminating the cross product

term in Equation (6.136). If we make this change of variable, then in the $x'y'$ -coordinate system. Equation (6.137) will become

$$\begin{aligned} \mathbf{x}^T A \mathbf{x} &= (P\mathbf{x}')^T A P\mathbf{x}' = \mathbf{x}'^T P^T A P\mathbf{x}' = \mathbf{x}'^T D\mathbf{x}' \\ \mathbf{x}'^T D\mathbf{x}' &= \begin{bmatrix} x' & y' \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} x' \\ y' \end{bmatrix} = k \end{aligned} \quad (6.139)$$

where λ_1 and λ_2 are the eigenvalues of A . The conic can now be identified by writing Equation (6.139) in the form

$$\lambda_1 x'^2 + \lambda_2 y'^2 = k \quad (6.140)$$

and performing the necessary algebra to match it with one of the standard forms.

For example, if λ_1, λ_2 , and k are positive, then Equation (6.140) represents an ellipse with an axis of length $2\sqrt{\frac{k}{\lambda_1}}$ in the x' -direction and $2\sqrt{\frac{k}{\lambda_2}}$ in the y' -direction.

The first column vector of P , which is a unit eigenvector corresponding to λ_1 , is along the positive x' -axis; and the second column vector of P , which is a unit eigenvector corresponding to λ_2 , is a unit vector along the y' -axis. These are called the principal axes of the ellipse.




Figure 6.53: The principal axes of the ellipse, the conic above is of form $\lambda_1 x'^2 + \lambda_2 y'^2 = k$, if λ_1, λ_2 , and k are positive, then it is an ellipse with $\alpha = \sqrt{\frac{k}{\lambda_1}}$ and $\beta = \sqrt{\frac{k}{\lambda_2}}$.

[SI*] If we have a conic with equation

$$ax^2 + 2bxy + cy^2 = k$$

with $b \neq 0$, the cross product term from the equation above ($2bxy$) can be eliminated by a rotation through an angle θ that satisfies

$$\cot 2\theta = \frac{a - c}{2b} \quad (6.141)$$

Definition 6.44: Terminology in Quadratic Form

A quadratic form $x^T Ax$ is said to be

- **Positive definite** if $x^T Ax > 0$ for $x \neq 0$.
- **Negative definite** if $x^T Ax < 0$ for $x \neq 0$.
- **Indefinite** if $x^T Ax$ has both positive and negative values.

Theorem 6.88: Terminology for Known Symmetric Matrix and Eigenvalues

If A is a symmetric matrix, then:

- $x^T Ax$ is positive definite if and only if all eigenvalues of A are positive.
- $x^T Ax$ is negative definite if and only if all eigenvalues of A are negative.
- $x^T Ax$ is indefinite if and only if A has at least one positive eigenvalue and at least one negative eigenvalue.

[SI*] A quadratic form for which $x^T Ax \geq 0$ if $x \neq 0$ is called positive semidefinite, and one for which $x^T Ax \leq 0$ if $x \neq 0$ is called negative semidefinite.

$x^T Ax$ is positive semidefinite if and only if all eigenvalues of A are nonnegative and is negative semidefinite if and only if all eigenvalues of A are nonpositive.

[SI*] **Example:**

It is not usually possible to tell from the signs of the entries in a symmetric matrix A whether that matrix is positive definite, negative definite, or indefinite. For example, the entries of the matrix

$$A = \begin{bmatrix} 3 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{bmatrix}$$

are nonnegative, but the matrix is indefinite since its eigenvalues are

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 4 \\ \lambda_3 &= -2 \end{aligned}$$

If we write the quadratic form, it will be

$$x^T Ax = \begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix} \begin{bmatrix} 3 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 3x_1^2 + 2x_1x_2 + 2x_1x_3 + 4x_2x_3$$

now, if we use different input for x

$$x^T Ax = 4$$

for $x_1 = 0, x_2 = 1$, and $x_3 = 1$.

$$x^T Ax = -4$$

for $x_1 = 0, x_2 = 1$, and $x_3 = -1$. Thus the quadratic form is indefinite.

[SI*] If $\mathbf{x}^T B \mathbf{x} = k$ is the equation of a conic, and if $k \neq 0$, then we can divide through by k and rewrite the equation in the form

$$\mathbf{x}^T A \mathbf{x} = 1 \quad (6.142)$$

where $A = (1/k)B$. If we now rotate the coordinate axes to eliminate the cross product term (if any) in this equation, then the equation of the conic in the new coordinate system will be of the form

$$\lambda_1 x'^2 + \lambda_2 y'^2 = 1 \quad (6.143)$$

in which λ_1 and λ_2 are the eigenvalues of A . The particular type of conic represented by this equation will depend on the signs of the eigenvalues λ_1 and λ_2

- $\mathbf{x}^T A \mathbf{x} = 1$ represents an ellipse if $\lambda_1 > 0$ and $\lambda_2 > 0$.
- $\mathbf{x}^T A \mathbf{x} = 1$ has no graph if $\lambda_1 < 0$ and $\lambda_2 < 0$.
- $\mathbf{x}^T A \mathbf{x} = 1$ represents a hyperbola if λ_1 and λ_2 have opposite signs.

In the case of the ellipse, Equation (6.143) can be rewritten as

$$\frac{x'^2}{\left(\frac{1}{\sqrt{\lambda_1}}\right)^2} + \frac{y'^2}{\left(\frac{1}{\sqrt{\lambda_2}}\right)^2} = 1 \quad (6.144)$$

so the axes of the ellipse have lengths $2/\sqrt{\lambda_1}$ and $2/\sqrt{\lambda_2}$.

Theorem 6.89: Classifying Conic Sections

If A is a symmetric 2×2 matrix, then:

- (a) $\mathbf{x}^T A \mathbf{x} = 1$ represents an ellipse if A is positive definite.
- (b) $\mathbf{x}^T A \mathbf{x} = 1$ has no graph if A is negative definite.
- (c) $\mathbf{x}^T A \mathbf{x} = 1$ represents a hyperbola if A is indefinite.

[SI*] Positive definite matrices are the most important symmetric matrices in applications, so it will be useful to learn a little more about them. We already know that a symmetric matrix is positive definite if and only if its eigenvalues are all positive.

[SI*] We define the k -th principal submatrix of an $n \times n$ matrix A to be the $k \times k$ submatrix consisting of the first k rows and columns of A .

Theorem 6.90: Determinant Test for Symmetric Matrix

A symmetric matrix A is positive definite if and only if the determinant of every principal submatrix is positive.

ii. Hermitian, Unitary, and Normal Matrices

[SI*] The transpose operation is less important for complex matrices than for real matrices.

Definition 6.45: Conjugate Transpose

If A is a complex matrix, then the conjugate transpose of A , denoted by A^* , is defined by

$$A^* = \overline{A}^T \quad (6.145)$$

[SI*] The order in which the transpose and conjugation operations are performed in computing $A^* = \overline{A}^T$ does not matter

$$(\overline{A^T}) = (\overline{A})^T$$

[SI*] In the case where A has real entries we have

$$A^* = (\overline{A})^T = A^T$$

so A^* is the same as A^T for real matrices.

[SI*] **Example:**

Find the conjugate transpose A^* of the matrix

$$A = \begin{bmatrix} 2+i & -3i & 0 \\ 3 & 1-5i & i \end{bmatrix}$$

Solution:

We have

$$\overline{A} = \begin{bmatrix} 2-i & 3i & 0 \\ 3 & 1+5i & -i \end{bmatrix}$$

and hence

$$A^* = \overline{A}^T = \begin{bmatrix} 2-i & 3 \\ 3i & 1+5i \\ 0 & -i \end{bmatrix}$$

Theorem 6.91: Algebraic Properties of the Conjugate Transpose Operation

If k is a complex scalar, and if A, B , and C are complex matrices whose sizes are such that the stated operations can be performed, then:

- (a) $(A^*)^* = A$
- (b) $(A + B)^* = A^* + B^*$
- (c) $(A - B)^* = A^* - B^*$
- (d) $(kA)^* = \overline{k}A^*$
- (e) $(AB)^* = B^*A^*$

[SI*] **Remark**

Note that the relationship

$$u \cdot v = \overline{v}^T u$$

can be expressed in terms of the conjugate transpose as

$$u \cdot v = v^* u \quad (6.146)$$

Definition 6.46: Hermitian and Unitary Matrices

A square complex matrix A is said to be unitary if

$$A^{-1} = A^* \quad (6.147)$$

and is said to be Hermitian if

$$A^* = A \quad (6.148)$$

[SI*] If A is a real matrix, then

$$A^* = A^T$$

in which case, the unitary matrix formula for real matrix becomes

$$A^{-1} = A^T$$

and the Hermitian matrix formula for real matrix becomes

$$A^T = A$$

The unitary matrices are complex generalizations of the real orthogonal matrices and Hermitian matrices are complex generalizations of the real symmetric matrices.

[SI*] Hermitian matrices are easy to recognize because their diagonal entries are real, and the entries that are symmetrically positioned across the main diagonal are complex conjugates. For example,

$$A = \begin{bmatrix} 2 & 3i & 2+6i \\ -3i & -6 & 5-7i \\ 2-6i & 5+7i & 1 \end{bmatrix}$$

is Hermitian.

Theorem 6.92: Eigenvalues for Hermitian Matrix

The eigenvalues of a Hermitian matrix are real numbers.

Theorem 6.93: Eigenvectors for Hermitian Matrix

If A is a Hermitian matrix, then eigenvectors from different eigenspaces are orthogonal.

Theorem 6.94: Equivalent Statements for Unitary Matrix

If A is an $n \times n$ matrix with complex entries, then the following are equivalent.

- (a) A is unitary.
- (b) $\|Ax\| = \|x\|$ for all x in $\mathbb{C}^n$.
- (c) $Ax \cdot Ay = x \cdot y$ for all x and y in $\mathbb{C}^n$.
- (d) The column vectors of A form an orthonormal set in $\mathbb{C}^n$ with respect to the complex Euclidean inner product.
- (e) The row vectors of A form an orthonormal set in $\mathbb{C}^n$ with respect to the complex Euclidean inner product.

[SI*] Unitary matrices are the complex analogs of the real orthogonal matrices.

Definition 6.47: Unitary Diagonalizability

A square complex matrix is said to be unitarily diagonalizable if there is a unitary matrix P such that $P^*AP = D$ is a complex diagonal matrix. Any such matrix P is said to unitarily diagonalize A .

Theorem 6.95: Unitary Diagonalizability for Hermitian Matrix

Every $n \times n$ Hermitian matrix A has an orthonormal set of n eigenvectors and is unitarily diagonalized by any $n \times n$ matrix P whose column vectors form an orthonormal set of eigenvectors of A .

[SI*] The procedure for unitarily diagonalizing Hermitian matrix A is exactly the same as that for orthogonally diagonalizing a symmetric matrix:

1. Find a basis for each eigenspace of A .
2. Apply the Gram-Schmidt process to each of these bases to obtain orthonormal bases for the eigenspaces.
3. Form the matrix P whose column vectors are the basis vectors obtained in Step 2. This will be a unitary matrix and will unitarily diagonalize A .

[SI*] We defined a square matrix with real entries to be skew-symmetric if

$$A^T = -A$$

A skew-symmetric matrix must have zeros on the main diagonal, and each entry off the main diagonal must be the negative of its mirror image about the main diagonal. Here is an example.

$$A = \begin{bmatrix} 0 & 3 & -5 \\ -3 & 0 & 6 \\ 5 & -6 & 0 \end{bmatrix}$$

[SI*] The complex analogs of the skew-symmetric matrices are the matrices for which

$$A^* = -A$$

Such matrices are said to be skew-Hermitian.

A skew-Hermitian matrix A has the property

$$A^* = \overline{A}^T = -A$$

it must be that A has zeros or pure imaginary numbers on the main diagonal, and that the complex conjugate of each entry off the main diagonal is the negative of the real number of its mirror image about the main diagonal. Here is an example.

$$A = \begin{bmatrix} i & 2-i & 6 \\ -2-i & 3i & 4i \\ -6 & 4i & 0 \end{bmatrix}$$

[SI*] Hermitian matrices enjoy many, but not all, of the properties of real symmetric matrices. For example, we know that real symmetric matrices are orthogonally diagonalizable and Hermitian matrices are unitarily diagonalizable. However, whereas the real symmetric matrices are the only orthogonally diagonalizable matrices, the Hermitian matrices do not constitute the entire class of unitarily diagonalizable complex matrices; that is, there exist unitarily diagonalizable matrices that are not Hermitian.

Specifically, it can be proved that a square complex matrix A is unitarily diagonalizable if and only if

$$AA^* = A^*A \tag{6.149}$$

Matrices with this property are said to be normal. Normal matrices include the Hermitian, skew-Hermitian, and unitary matrices in the complex case and the symmetric, skew-symmetric, and orthogonal matrices in the real case. The nonzero skew-symmetric matrices are particularly interesting because they are examples of real matrices that are not orthogonally diagonalizable but are unitarily diagonalizable.

[SI*] We have seen that Hermitian matrices have real eigenvalues.

The eigenvalues of skew-Hermitian matrix are either zero or purely imaginary (have real part of zero) and that the eigenvalues of unitary matrices have modulus 1, $|\lambda| = 1$.




Figure 6.54

iii. Functions in SymIntegration Related to Diagonalization and Quadratic Forms

[SI*] We will list the basic functions that can be used related to diagonalization and quadratic forms:

quadraticmultiplication(vector<vector<double>> &matrixA, vector<double> &vectorx)
to compute the quadratic form of $x^T Ax$.

complex<double> complexquadraticmultiplication(ComplexMatrix &A, ComplexVector &x)

to compute the complex quadratic multiplication ($\bar{x}^T Ax$) of a square complex matrix A with size $n \times n$ and vector x with size n .

Remember that in real space $\mathbb{R}^n$ the quadratic form is

$$x^T Ax = x \cdot Ax = Ax \cdot x$$

the analogous quadratic form in complex space $\mathbb{C}^n$ is

$$x \cdot Ax = Ax \cdot x = \bar{x}^T Ax$$

SpectralDecomposition(vector<vector<complex<double>>> &A, double a, double b)

to compute the spectral decomposition of matrix A , with a as the lower bound for uniform distribution, b as the upper bound for the uniform distribution, we are using uniform distribution to generate random μ and σ as the input to generate the initial complex vector v^0 that is normally distributed with mean μ and standard deviation σ , this initial vector is a prerequisite for the Rayleigh quotient iteration. The key here is to choose a and b carefully so we can pinpoint the range where the eigenvalue could be located.

iv. Compute Spectral Decomposition with SymIntegration

Compute the spectral decomposition of the matrix

$$A = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

Solution:

The matrix A has eigenvalues $\lambda_1 = -3$ and $\lambda_2 = 2$ with corresponding eigenvectors that already normalized (orthonormal)

$$u_1 = \frac{x_1}{||x_1||} = \begin{bmatrix} \frac{1}{\sqrt{5}} \\ -\frac{2}{\sqrt{5}} \end{bmatrix}, \quad u_2 = \frac{x_2}{||x_2||} = \begin{bmatrix} \frac{2}{\sqrt{5}} \\ \frac{1}{\sqrt{5}} \end{bmatrix}$$

so a spectral decomposition of A is

$$A = \lambda_1 u_1 u_1^T + \lambda_2 u_2 u_2^T$$

$$A = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} = (-3) \begin{bmatrix} \frac{1}{5} & -\frac{2}{5} \\ -\frac{2}{5} & \frac{4}{5} \end{bmatrix} + (2) \begin{bmatrix} \frac{4}{5} & \frac{2}{5} \\ \frac{2}{5} & \frac{1}{5} \end{bmatrix}$$

where the 2×2 matrices on the right side are the standard matrices for the orthogonal projections onto the eigenspaces corresponding to $\lambda_1 = -3$ and $\lambda_2 = 2$, respectively.

In SymIntegration, to compute spectral decomposition of a matrix we can use this function: **SpectralDecomposition(vector<vector<complex<double>>> &A, double a, double b)** with A as the matrix, a as the lower bound for uniform distribution, b as the upper bound for the uniform distribution, we are using uniform distribution to generate random μ and σ as the input to generate the initial complex vector v^0 that is normally distributed with mean μ and standard deviation σ , this initial vector is a prerequisite for the Rayleigh quotient iteration. The key here is to choose a and b carefully so we can pinpoint the range where the eigenvalue could be located.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Spectral Decomposition ]# ./main
Matrix A :
      (1,0)      (2,0)
      (2,0)      (-2,0)

Eigenvalues :
(-3,0)
(2,0)

Eigenvectors (represented by column):
(-0.447,0)  (0.894,0)
(0.894,0)   (0.447,0)

Spectral decomposition for  $\lambda_{\{1\}} = (-3,0) *$ 
(0.2,0)      (-0.4,0)
(-0.4,0)      (0.8,0)

Spectral decomposition for  $\lambda_{\{2\}} = (2,0) *$ 
(0.8,0)      (0.4,0)
(0.4,0)      (0.2,0)

Time taken by function: 10308 microseconds
```

Figure 6.55: The spectral decomposition of matrix A . (SymIntegration/Examples/Linear Algebra/Test SymIntegration Spectral Decomposition/main.cpp).

v. Compute Hessenberg Decomposition with SymIntegration

Compute the Hessenberg decomposition of the matrix

$$A = \begin{bmatrix} 1 & 0 & 2 & 3 \\ -1 & 0 & 5 & 2 \\ 2 & -2 & 0 & 0 \\ 2 & -1 & 2 & 0 \end{bmatrix}$$

Solution:

To compute the upper Hessenberg form H ($H_{ij} = 0$ for $i > j + 1$) for Hessenberg decomposition with Householder reflections, you iteratively apply similarity transformations using Householder reflectors to zero out entries below the subdiagonal, preserving eigenvalues. You construct these reflectors (H_k) by reflecting specific vectors (e.g. the first column's tail) towards a desired form, forming

$$A_k = H_k A_{k-1} H_k$$

until the matrix becomes upper Hessenberg (all entries below subdiagonal are zero), with the final

$$H = H_1 H_2 \dots H_{n-2}$$

and

$$H = P^T A P$$

First, at iteration $j = 1$ we will have

$$x = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}$$

(we define vector x as the part of column j below the diagonal $A(j+1 : n, j)$).

For the vector w , the choice of the sign for the first entry is the opposite of the first element of vector x .

$$w = \begin{bmatrix} \pm ||x|| \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix}$$

Construct the householder vector v

$$v = w - x = \begin{bmatrix} 4 \\ -2 \\ -2 \end{bmatrix}$$

Construct the Householder reflector $\hat{H}_j$

$$\hat{H}_1 = I - 2 \frac{vv^T}{v^T v} = \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \\ \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \end{bmatrix}$$

Create $n \times n$ Householder matrix H_j by embedding $\hat{H}_j$ into the lower-right $(n - j) \times (n - j)$ block, with ones on the diagonal and zeros elsewhere in the top-left $j \times j$ block.

$$H_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ 0 & \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \\ 0 & \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \end{bmatrix}$$

Accumulate

$$P = PH_j$$

starts with $P = I$.

Thus

$$H_1 A H_1 = \begin{bmatrix} 1 & \frac{10}{3} & \frac{1}{3} & \frac{4}{3} \\ 3 & 0 & -\frac{4}{3} & -\frac{5}{3} \\ 0 & 3 & \frac{2}{3} & -\frac{2}{3} \\ 0 & 4 & \frac{8}{3} & -\frac{2}{3} \end{bmatrix}$$

Do the transformation at the end of each iteration:

$$A_{new} = H_j A_{old} H_j$$

Repeat the iteration till $j = n - 2$, after $n - 2$ steps the matrix A will be in upper Hessenberg form, and P will hold the accumulated orthogonal transformations.

HessenbergDecomposition(vector<vector<complex<double>>> &A)

with A as the square matrix that we want to compute the upper Hessenberg decomposition of A .

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Hessenberg Decomposition ]# ./main

Matrix A :
(1,0)      (0,0)      (2,0)      (3,0)
(-1,0)     (0,0)     (5,0)     (2,0)
(2,0)     (-2,0)     (0,0)     (0,0)
(2,0)     (-1,0)     (2,0)     (0,0)

X:
(-1,0)
(2,0)
(2,0)

W:
(3,0)
(0,0)
(0,0)

V:
(4,0)
(-2,0)
(-2,0)

vTv:
(24,0)

vvT:
(16,0)      (-8,0)      (-8,0)
(-8,0)      (4,0)      (4,0)
(-8,0)      (4,0)      (4,0)

P:
(1.33,0)    (-0.667,0)    (-0.667,0)
(-0.667,0)  (0.333,0)    (0.333,0)
(-0.667,0)  (0.333,0)    (0.333,0)

I - 2P:
(-0.333,0)  (0.667,0)    (0.667,0)
(0.667,0)   (0.667,0)   (-0.333,0)
(0.667,0)   (-0.333,0)  (0.667,0)

H_{i}:
(1,0)      (0,0)      (0,0)      (0,0)
(0,0)     (-0.333,0)  (0.667,0)  (0.667,0)
(0,0)     (0.667,0)  (0.667,0)  (-0.333,0)
(0,0)     (0.667,0)  (-0.333,0) (0.667,0)

```

Figure 6.56: The Hessenberg decomposition of matrix A. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Hessenberg Decomposition/main.cpp*).

```

H_{i}:
(1,0)      (0,0)      (0,0)      (0,0)
(0,0)      (-0.333,0)  (0.667,0)  (0.667,0)
(0,0)      (0.667,0)  (0.667,0)  (-0.333,0)
(0,0)      (0.667,0)  (-0.333,0)  (0.667,0)

H_{i} A H_{i}:
(1,0)      (3.33,0)    (0.333,0)  (1.33,0)
(3,0)      (0,0)      (-1.33,0)  (-1.67,0)
(0,0)      (3,0)     (0.667,0)  (-0.667,0)
(0,0)      (4,0)     (2.67,0)  (-0.667,0)

Householder Matrix:
(1,0)      (0,0)      (0,0)      (0,0)
(0,0)      (-0.333,0)  (0.667,0)  (0.667,0)
(0,0)      (0.667,0)  (0.667,0)  (-0.333,0)
(0,0)      (0.667,0)  (-0.333,0)  (0.667,0)

x:
(3,0)
(4,0)

w:
(-5,0)
(0,0)

v:
(-8,0)
(-4,0)

vTv:
(80,0)

vTvT:
(64,0)      (32,0)
(32,0)      (16,0)

P:
(1.6,0)      (0.8,0)
(0.8,0)      (0.4,0)

I - 2P:
(-0.6,0)      (-0.8,0)
(-0.8,0)      (0.6,0)

H_{i}:
(1,0)      (0,0)      (0,0)      (0,0)
(0,0)      (1,0)      (0,0)      (0,0)
(0,0)      (0,0)      (-0.6,0)  (-0.8,0)
(0,0)      (0,0)      (-0.8,0)  (0.6,0)

```

Figure 6.57: The Hessenberg decomposition of matrix A. (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Hessenberg Decomposition/main.cpp*).

```

H_{i}:
(1,0)      (0,0)      (0,0)      (0,0)
(0,0)      (1,0)      (0,0)      (0,0)
(0,0)      (0,0)      (-0.6,0)   (-0.8,0)
(0,0)      (0,0)      (-0.8,0)   (0.6,0)

H_{i} A H_{i}:
(1,0)      (3.33,0)    (-1.27,0)   (0.533,0)
(3,0)      (0,0)      (2.13,0)    (0.0667,0)
(0,0)      (-5,0)     (0.773,0)   (2.59,0)
(0,0)      (0,0)     (-0.747,0)  (-0.773,0)

Householder Matrix:
(1,0)      (0,0)      (0,0)      (0,0)
(0,0)      (-0.333,0)  (-0.933,0)  (-0.133,0)
(0,0)      (0.667,0)  (-0.133,0)  (-0.733,0)
(0,0)      (0.667,0)  (-0.333,0)  (0.667,0)

*****

End of iteration

*****

P^{T} A P:
(1,0)      (3.33,0)    (-1.27,0)   (0.533,0)
(3,0)      (0,0)      (2.13,0)    (0.0667,0)
(0,0)      (-5,0)     (0.773,0)   (2.59,0)
(0,0)      (0,0)     (-0.747,0)  (-0.773,0)

Time taken by function: 5124 microseconds

```

Figure 6.58: The Hessenberg decomposition of matrix A . (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Hessenberg Decomposition/main.cpp*).

VIII. LINEAR TRANSFORMATIONS

[SI*] A matrix transformation $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is defined as a mapping of the form

$$T_A(\mathbf{x}) = A\mathbf{x}$$

in which A is an $m \times n$ matrix.

Definition 6.48: Linear Transformation

If $T : V \rightarrow W$ is a function from a vector space V to a vector space W , then T is called a linear transformation from V to W if the following two properties hold for all vectors $\mathbf{u}$ and $\mathbf{v}$ in V and for all scalars k :

- (a) $T(k\mathbf{u}) = kT(\mathbf{u})$ (Homogeneity property)
- (b) $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ (Additivity property)

In the special case where $V = W$, the linear transformation T is called a linear operator on the vector space V .

[SI*] The homogeneity and additivity properties of a linear transformation $T : V \rightarrow W$ can be used in combination to show that if $\mathbf{v}_1$ and $\mathbf{v}_2$ are vectors in V and k_1 and k_2 are any scalars, then

$$T(k_1\mathbf{v}_1 + k_2\mathbf{v}_2) = k_1T(\mathbf{v}_1) + k_2T(\mathbf{v}_2)$$

More generally, if $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r$ are vectors in V and $k_1, k_2, \dots, k_r$ are any scalars, then

$$T(k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + \dots + k_r\mathbf{v}_r) = k_1T(\mathbf{v}_1) + k_2T(\mathbf{v}_2) + \dots + k_rT(\mathbf{v}_r) \quad (6.150)$$

Theorem 6.96: Linear Transformation

If $T : V \rightarrow W$ is a linear transformation, then:

- (a) $T(\mathbf{0}) = \mathbf{0}$
- (b) $T(\mathbf{u} - \mathbf{v}) = T(\mathbf{u}) - T(\mathbf{v})$ for all $\mathbf{u}$ and $\mathbf{v}$ in V

[SI*] Matrix Transformations

We have based the definition of a general linear transformation on the homogeneity and additivity properties of matrix transformations, it follows that a matrix transformation $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is also a linear transformation in this more general sense with $V = \mathbb{R}^n$ and $W = \mathbb{R}^m$.

[SI*] The Zero Transformation

Let V and W be any two vector spaces. The mapping $T : V \rightarrow W$ such that $T(\mathbf{v}) = \mathbf{0}$ for every $\mathbf{v}$ in V is a linear transformation called the zero transformation. To see that T is linear, observe that

$$\begin{aligned} T(\mathbf{u} + \mathbf{v}) &= \mathbf{0} \\ T(\mathbf{u}) &= \mathbf{0} \\ T(\mathbf{v}) &= \mathbf{0} \\ T(k\mathbf{u}) &= \mathbf{0} \end{aligned}$$

Therefore,

$$\begin{aligned}T(\mathbf{u} + \mathbf{v}) &= T(\mathbf{u}) + T(\mathbf{v}) \\T(k\mathbf{u}) &= kT(\mathbf{u})\end{aligned}$$

[SI*] **The Identity Operator**

Let V be any vector space. The mapping $I : V \rightarrow V$ defined by $I(\mathbf{v}) = \mathbf{v}$ is called the identity operator on V . I is linear.

[SI*] **Dilation and Contraction Operators**

If V is a vector space and k is any scalar, then the mapping $T : V \rightarrow V$ given by $T(\mathbf{x}) = k\mathbf{x}$ is a linear operator on V , for if c is any scalar and if $\mathbf{u}$ and $\mathbf{v}$ are any vectors in V , then

$$\begin{aligned}T(c\mathbf{u}) &= k(c\mathbf{u}) = c(k\mathbf{u}) = cT(\mathbf{u}) \\T(\mathbf{u} + \mathbf{v}) &= k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v} = T(\mathbf{u}) + T(\mathbf{v})\end{aligned}$$

If $0 < k < 1$, then T is called the contraction of V with factor k , and if $k > 1$, it is called the dilation of V with factor k .

[SI*] **A Linear Transformation from P_n to P_{n+1}**

Let $\mathbf{p} = p(x) = c_0 + c_1x + \cdots + c_nx^n$ be a polynomial in P_n , and define the transformation $T : P_n \rightarrow P_{n+1}$ by

$$T(\mathbf{p}) = T(p(x)) = xp(x) = c_0x + c_1x^2 + \cdots + c_nx^{n+1}$$

This transformation is linear because for any scalar k and any polynomials $\mathbf{p}_1$ and $\mathbf{p}_2$ in P_n we have

$$T(k\mathbf{p}) = T(kp(x)) = x(kp(x)) = k(xp(x)) = kT(\mathbf{p})$$

and

$$\begin{aligned}T(\mathbf{p}_1 + \mathbf{p}_2) &= T(p_1(x) + p_2(x)) \\&= x(p_1(x) + p_2(x)) \\&= xp_1(x) + xp_2(x) \\&= T(\mathbf{p}_1) + T(\mathbf{p}_2)\end{aligned}$$

[SI*] **A Linear Transformation Using an Inner Product**

Let V be an inner product space, let $\mathbf{v}_0$ be any fixed vector in V , and let $T : V \rightarrow R$ be the transformation

$$T(\mathbf{x}) = \langle \mathbf{x}, \mathbf{v}_0 \rangle$$

that maps a vector $\mathbf{x}$ into its inner product with $\mathbf{v}_0$. This transformation is linear, for if k is any scalar, and if $\mathbf{u}$ and $\mathbf{v}$ are any vectors in V , then it follows from properties of inner products that

$$\begin{aligned}T(k\mathbf{u}) &= \langle k\mathbf{u}, \mathbf{v}_0 \rangle = k \langle \mathbf{u}, \mathbf{v}_0 \rangle = kT(\mathbf{u}) \\T(\mathbf{u} + \mathbf{v}) &= \langle \mathbf{u} + \mathbf{v}, \mathbf{v}_0 \rangle = \langle \mathbf{u}, \mathbf{v}_0 \rangle + \langle \mathbf{v}, \mathbf{v}_0 \rangle = T(\mathbf{u}) + T(\mathbf{v})\end{aligned}$$

[SI*] **Example: Transformations on Matrix Spaces**

Let M_{nn} be the vector space of $n \times n$ matrices. In each part determine whether the transformation is linear.

(a) $T_1(A) = A^T$

(b) $T_2(A) = \det(A)$

Solution:

(a) It follows that

$$\begin{aligned} T_1(kA) &= (kA)^T = kA^T = kT_1A \\ T_1(A+B) &= (A+B)^T = A^T + B^T = T_1(A) + T_1(B) \end{aligned}$$

so T_1 is linear.

(b) It follows that

$$T_2(kA) = \det(kA) = k^n \det(A) = k^n T_2(A)$$

Thus, T_2 is not homogeneous and hence not linear if $n > 1$. Note that additivity also fails because we showed that $\det(A+B)$ and $\det(A) + \det(B)$ are not generally equal.

[SI*] **Translation Is Not Linear**

A Theorem states that a linear transformation maps $\mathbf{0}$ to $\mathbf{0}$. This property is useful for identifying transformations that are not linear. For example, if $\mathbf{x}_0$ is a fixed nonzero vector in $\mathbb{R}^2$, then the transformation

$$T(\mathbf{x}) = \mathbf{x} + \mathbf{x}_0$$

has the geometric effect of translating each point $\mathbf{x}$ in a direction parallel to $\mathbf{x}_0$ through a distance of $\|\mathbf{x}_0\|$. This cannot be a linear transformation since $T(\mathbf{0}) = \mathbf{x}_0$, so T does not map $\mathbf{0}$ to $\mathbf{0}$.




Figure 6.59: $T(\mathbf{x}) = \mathbf{x} + \mathbf{x}_0$ translates each point $\mathbf{x}$ along a line parallel to $\mathbf{x}_0$ through a distance $\|\mathbf{x}_0\|$.

[SI*] **The Evaluation Transformation**

Let V be a subspace of $F(-\infty, \infty)$, let

$$x_1, x_2, \dots, x_n$$

be distinct real numbers, and let $T : V \rightarrow \mathbb{R}^n$ be the transformation

$$T(f) = (f(x_1), f(x_2), \dots, f(x_n)) \quad (6.151)$$

that associates with f the n -tuple of function values at $x_1, x_2, \dots, x_n$. We call this the evaluation transformation on V at $x_1, x_2, \dots, x_n$. Thus, for example, if

$$x_1 = -1, \quad x_2 = 2, \quad x_3 = 4$$

and if $f(x) = x^2 - 1$, then

$$T(f) = (f(x_1), f(x_2), f(x_3)) = (0, 3, 15)$$

The evaluation transformation is linear, for if k is any scalar, and if f and g are any functions in V , then

$$\begin{aligned} T(kf) &= ((kf)(x_1), (kf)(x_2), \dots, (kf)(x_n)) \\ &= (kf(x_1), kf(x_2), \dots, kf(x_n)) \\ &= k(f(x_1), f(x_2), \dots, f(x_n)) \\ &= kT(f) \end{aligned}$$

and

$$\begin{aligned} T(f+g) &= ((f+g)(x_1), (f+g)(x_2), \dots, (f+g)(x_n)) \\ &= (f(x_1) + g(x_1), f(x_2) + g(x_2), \dots, f(x_n) + g(x_n)) \\ &= (f(x_1), f(x_2), \dots, f(x_n)) + (g(x_1), g(x_2), \dots, g(x_n)) \\ &= T(f) + T(g) \end{aligned}$$

[SI*] If $\mathbb{R}^n \rightarrow \mathbb{R}^n$ is a matrix transformation, say multiplication by A , and if $e_1, e_2, \dots, e_n$ are the standard basis vectors for $\mathbb{R}^n$, then A can be expressed as

$$A = [T(e_1) \mid T(e_2) \mid \dots \mid T(e_n)]$$

It follows from this that the image of any vector $v = (c_1, c_2, \dots, c_n)$ in $\mathbb{R}^n$ under multiplication by A can be expressed as

$$T(v) = c_1 T(e_1) + c_2 T(e_2) + \dots + c_n T(e_n)$$

This formula tells us that for a matrix transformation the image of any vector is expressible as a linear combination of the images of the standard basis vectors.

Theorem 6.97: Linear Transformations from Images of Basis Vectors

Let $T : V \rightarrow W$ be a linear transformation, where V is finite dimensional. If $S = \{v_1, v_2, \dots, v_n\}$ is a basis for V , then the image of any vector v in V can be expressed as

$$T(v) = c_1 T(v_1) + c_2 T(v_2) + \dots + c_n T(v_n) \quad (6.152)$$

where $c_1, c_2, \dots, c_n$ are the coefficients required to express v as a linear combination of the vectors in S .

[SI*] **A Linear Transformation from $C^1(-\infty, \infty)$ to $F(-\infty, \infty)$**

Let $V = C^1(-\infty, \infty)$ be the vector space of functions with continuous first derivatives on $(-\infty, \infty)$, and let $W = F(-\infty, \infty)$ be the vector space of all real-valued functions defined

on $(-\infty, \infty)$. Let $D : V \rightarrow W$ be the transformation that maps a function $f = f(x)$ into its derivative, that is

$$D(f) = f'(x)$$

From the properties of differentiation, we have

$$D(f + g) = D(kf) = kD(f)$$

and

$$D(f + g) = D(f) + D(g)$$

Thus, D is a linear transformation.

[SI*] **An Integral Transformation**

Let $V = C(-\infty, \infty)$ be the vector space of continuous functions on the interval $(-\infty, \infty)$, let $W = C^1(-\infty, \infty)$ be the vector space of functions with continuous first derivatives on $(-\infty, \infty)$, and let $J : V \rightarrow W$ be the transformation that maps a function f in V into

$$J(f) = \int_0^x f(t) dt$$

For example, if $f(x) = x^2$, then

$$J(f) = \int_0^x t^2 dt = \frac{t^3}{3} \Big|_0^x = \frac{x^3}{3}$$

The transformation $J : V \rightarrow W$ is linear, for if k is any constant, and if f and g are any functions in V , then properties of the integral imply that

$$\begin{aligned} J(kf) &= \int_0^x kf(t) dt \\ &= k \int_0^x f(t) dt \\ &= kJ(f) \end{aligned}$$

$$\begin{aligned} J(f + g) &= \int_0^x (f(t) + g(t)) dt \\ &= \int_0^x f(t) dt + \int_0^x g(t) dt \\ &= J(f) + J(g) \end{aligned}$$

[SI*] If A is an $m \times n$ matrix, then the null space of A consists of all vectors x in $\mathbb{R}^n$ such that $Ax = \mathbf{0}$, and the column space of A consists of all vectors b in $\mathbb{R}^m$ for which there is at least one vector x in $\mathbb{R}^n$ such that $Ax = b$.

From the viewpoint of matrix transformations, the null space of A consists of all vectors in $\mathbb{R}^n$ that multiplication by A maps into $\mathbf{0}$, and the column space of A consists of all vectors in $\mathbb{R}^m$ that are images of at least one vector in $\mathbb{R}^n$ under multiplication by A .

Definition 6.49: Kernel and Range

If $T : V \rightarrow W$ is a linear transformation, then the set of vectors in V that maps T into $\mathbf{0}$ is called the kernel of T , and is denoted by $\ker(T)$.

The set of all vectors in W that are images under T of at least one vector in V is called the range of T and is denoted by $R(T)$.

[SI*] Kernel and Range of a Matrix Transformation

If $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is multiplication by the $m \times n$ matrix A , then, the kernel of T_A is the null space of A , and the range of T_A is the column space of A .

[SI*] Kernel and Range of the Zero Transformation

Let $T : V \rightarrow W$ be the zero transformation. Since T maps every vector in V into $\mathbf{0}$, it follows that $\ker(T) = V$. Moreover, since $\mathbf{0}$ is the only image under T of vectors in V , it follows that $R(T) = \{\mathbf{0}\}$.

[SI*] Kernel and Range of the Identity Transformation

Let $I : V \rightarrow V$ be the identity operator. Since $I(v) = v$ for all vectors in V , every vector in V is the image of some vector (itself); thus $R(I) = V$. Since the only vector that I maps into $\mathbf{0}$ is $\mathbf{0}$, it follows that $\ker(I) = \{\mathbf{0}\}$.

[SI*] Kernel and Range of an Orthogonal Projection

The points that T maps into $\mathbf{0} = (0, 0, 0)$ are precisely those on the z -axis, so $\ker(T)$ is the set of points of the form $(0, 0, z)$.

T maps the points in $\mathbb{R}^3$ to the xy -plane, where each point in that plane is the image of each point on the vertical line above it. Thus $R(T)$ is the set of points in the form $(x, y, 0)$.

[SI*] Kernel and Range of a Rotation

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear operator that rotates each vector in the xy -plane through the angle θ , it follows that $R(T) = \mathbb{R}^2$. Moreover, the only vector that rotates into $\mathbf{0}$ is $\mathbf{0}$, so $\ker(T) = \{\mathbf{0}\}$.

[SI*] Kernel of a Differentiation Transformation

Let $V = C^1(-\infty, \infty)$ be the vector space of functions with continuous first derivatives on $(-\infty, \infty)$, let $W = F(-\infty, \infty)$ be the vector space of all real-valued functions defined on $(-\infty, \infty)$, and let $D : V \rightarrow W$ be the differentiation transformation $D(f) = f'(x)$. The kernel of D is the set of functions in V with derivative zero. From calculus, this is the set of constant functions on $(-\infty, \infty)$.

[SI*] Most of the time, $\ker(T)$ and $R(T)$ turned out to be subspaces. They were either zero subspace or the entire vector space. In another case, the kernel was a line through the origin, and the range was a plane through the origin, both of which are subspaces of $\mathbb{R}^3$.

Theorem 6.98: Kernel and Range of a Linear Transformation

If $T : V \rightarrow W$ is a linear transformation, then:

- (a) The kernel of T is a subspace of V .
- (b) The range of T is a subspace of W .

[SI*] To show that $\ker(T)$ is a subspace, we must show that it contains at least one vector and is closed under addition and scalar multiplication.

[SI*] To show that $R(T)$ is a subspace, we must show that it contains at least one vector and is closed under addition and scalar multiplication.

[SI*] Application to Differential Equations

Differential equations of the form

$$y'' + \omega^2 y = 0 \quad (6.153)$$

with ω as a positive constant. The equation above arise in the study of vibrations. The set of all solutions of this equation on the interval $(-\infty, \infty)$ is the kernel of the linear

transformation $D : C^2(-\infty, \infty) \rightarrow C(-\infty, \infty)$, given by

$$D(y) = y'' + \omega^2 y$$

It is proved in standard textbooks on differential equations that the kernel is a two-dimensional subspace of $C^2(-\infty, \infty)$, so that if we can find two linearly independent solutions of the differential equation $D(y) = 0$, then all other solutions can be expressed as linear combinations of those two. We will find the general solution of

$$y'' + \omega^2 y = 0$$

the characteristic equation is $r^2 + \omega^2 = 0$ with the roots

$$r = \pm i\omega$$

the roots are conjugate complex numbers, hence the general solution is

$$y = c_1 e^{r_1 x} + c_2 e^{r_2 x}$$

and

$$\begin{aligned} y_1(x) &= e^{i\omega x} \\ y_2(x) &= e^{-i\omega x} \end{aligned}$$

the functions $y_1(x)$ and $y_2(x)$ are solutions of the differential equation. Based on the theorem, if y_1 and y_2 are solutions, then any linear combinations of y_1 and y_2 are also a solution.

$$\begin{aligned} y_1(x) + y_2(x) &= e^{i\omega x} + e^{-i\omega x} \\ &= (\cos \omega x + i \sin \omega x) + (\cos \omega x - i \sin \omega x) \\ &= 2 \cos \omega x \end{aligned} \tag{6.154}$$

and

$$\begin{aligned} y_1(x) - y_2(x) &= e^{i\omega x} - e^{-i\omega x} \\ &= (\cos \omega x + i \sin \omega x) - (\cos \omega x - i \sin \omega x) \\ &= 2i \sin \omega x \end{aligned} \tag{6.155}$$

neglecting the constant multipliers 2 and $2i$, respectively, we have obtained a pair of real-valued solutions

$$\begin{aligned} y_1(x) &= \cos \omega x \\ y_2(x) &= \sin \omega x \end{aligned}$$

they are solutions of the differential equations $D(y) = 0$. These functions are linearly independent since neither is a scalar multiple of the other, and thus

$$y = c_1 \cos \omega x + c_2 \sin \omega x \tag{6.156}$$

is a "general solution" of $D(y) = 0$ in the sense that every choice of c_1 and c_2 produces a solution, and every solution is of this form.

Definition 6.50: Rank and Nullity of Linear Transformations

Let $T : V \rightarrow W$ be a linear transformation. If the range of T is finite-dimensional, then its dimension is called the rank of T ; and if the kernel of T is finite-dimensional, then its dimension is called the nullity of T . The rank of T is denoted by $\text{rank}(T)$ and the nullity of T by $\text{nullity}(T)$.

Theorem 6.99: Dimension Theorem for Linear Transformations

If $T : V \rightarrow W$ is a linear transformation from an n -dimensional vector space V to a vector space W , then

$$\text{rank}(T) + \text{nullity}(T) = n \quad (6.157)$$

[SI*] In the special case where A is an $m \times n$ matrix and $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is multiplication by A , the kernel of T_A is the null space of A , and the range of T_A is the column space of A . Thus, it follows that

$$\text{rank}(T_A) + \text{nullity}(T_A) = n$$

[SI*] **Example:**

Let v_1, v_2 , and v_3 be vectors in a vector space V , and let $T : V \rightarrow \mathbb{R}^3$ be a linear transformation for which

$$T(v_1) = (2, -1, 4)$$

$$T(v_2) = (-3, 2, 1)$$

$$T(v_3) = (0, 5, 1)$$

Find $T(3v_1 - 2v_2 + v_3)$.

Solution:

The computation is quite simple

$$\begin{aligned} T(3v_1 - 2v_2 + v_3) &= 3T(v_1) - 2T(v_2) + T(v_3) \\ &= 3(2, -1, 4) - 2(-3, 2, 1) + (0, 5, 1) \\ &= (6, -3, 12) - (-6, 4, 2) + (0, 5, 1) \\ &= (12, -2, 11) \end{aligned}$$

[SI*] **Example:**

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear operator given by the formula

$$T(x, y) = (x - 3y, -2x + 6y)$$

(a) Which of the following vectors are in $R(T)$: $(1, -2), (3, 1), (-2, 4)$?

(b) Which of the following vectors are in $\ker(T)$: $(1, -3), (3, 1), (-6, -2)$?

Solution:

(a) We can use this method

$$\begin{aligned} x - 3y &= b_1 \\ -2x + 6y &= b_2 \end{aligned}$$

with (b_1, b_2) is the vector we want to test whether it is in $R(T)$ or not.

$$\begin{aligned}x - 3y &= 1 \\ -2x + 6y &= -2\end{aligned}$$

the equations are linear combination of each other, thus $(1, -2)$ is in $R(T)$.

$$\begin{aligned}x - 3y &= 3 \\ -2x + 6y &= 1\end{aligned}$$

the equations are not linear combination of each other, thus $(3, 1)$ is not in $R(T)$.

$$\begin{aligned}x - 3y &= -2 \\ -2x + 6y &= 4\end{aligned}$$

the equations are linear combination of each other, thus $(-2, 4)$ is in $R(T)$.

(b) We will compute one by one

$$T(1, -3) = (1 - 3(-3), -2(1) + 6(-3)) = (10, -20)$$

$(1, -3)$ is not in $\ker(T)$.

$$T(3, 1) = (3 - 3(1), -2(3) + 6(1)) = (0, 0)$$

$(3, 1)$ is in $\ker(T)$

$$T(-6, -2) = (-6 - 3(-2), -2(-6) + 6(-2)) = (0, 0)$$

$(-6, -2)$ is in $\ker(T)$

[SI*] **Example:**

Let $T : P_2 \rightarrow P_3$ be the linear transformation defined by

$$T(p(x)) = xp(x)$$

(a) Which of the following are in $\ker(T)$: $x^2, 0, 1 + x$?

(b) Which of the following are in $R(T)$: $x + x^2, 1 + x, 3 - x^2$?

Solution:

(a) We will compute one by one

$$T(x^2) = x(x^2) = x^3$$

x^2 is not in $\ker(T)$.

$$T(0) = x(0) = 0$$

0 is in $\ker(T)$.

$$T(1 + x) = x(1 + x) = x + x^2$$

$1 + x$ is not in $\ker(T)$. The only vector that will map $T(p(x))$ into 0 is 0 .

(b) Since $x + x^2 = x(1 + x)$, $x + x^2$ is in $R(T)$.

Neither $1 + x$ nor $3 - x^2$ can be expressed as $x p(x)$ with $p(x)$ in P_2 , so neither are in $R(T)$.

[SI*] Example:

Find a basis for the kernel of

(a) the linear operator $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by the formula

$$T(x, y) = (x - 3y, -2x + 6y)$$

(b) the linear transformation $T : P_2 \rightarrow P_3$ defined by

$$T(p(x)) = xp(x)$$

Solution:

(a) The kernel of $T(x, y)$ is the set of all vectors $(3x, x)$ or the line $y = 3x$.

The vector $(3, 1)$ is a basis for this space. If you input the basis or multiplication of the basis in the $T(x, y) = (x - 3y, -2x + 6y)$ we will obtain $\mathbf{0} = (0, 0)$.

(b) Since we need to obtain $T(x) = 0$, then

$$T(0) = x(0) = 0$$

the basis for the kernel is $\{0, 0, 0\}$ from standard basis of $P_2 : \{1, x, x^2\}$.

[SI*] Example:

Find a basis for the range of

(a) the linear operator $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by the formula

$$T(x, y) = (x - 3y, -2x + 6y)$$

(b) the linear transformation $T : P_2 \rightarrow P_3$ defined by

$$T(p(x)) = xp(x)$$

(c) the matrix operator $T_A : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ is multiplication by the 3×4 matrix A

$$A = \begin{bmatrix} 4 & 1 & -2 & -3 \\ 2 & 1 & 1 & -4 \\ 6 & 0 & -9 & 9 \end{bmatrix}$$

Solution:

(a) Since $x - 3y$ and $-2x + 6y$ can assume any real value, $R(T)$ is the set of all vectors $(x, -2x)$ or the line $y = -2x$. The vector $(1, -2)$ is a basis for this space.

(b) $R(T)$ is the set of all polynomials in P^3 with constant term 0. Thus the basis for the range / $R(T)$ is $\{x, x^2, x^3\}$.

- (c) A basis for $R(T)$ is a basis for the column space of A .

First we will transform A into its row echelon form

$$A_{ref} = \begin{bmatrix} 1 & \frac{1}{2} & \frac{1}{2} & -2 \\ 0 & 1 & 4 & -5 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

The columns of A corresponding to the columns of A_{ref} containing leading 1s are

$$\begin{bmatrix} 4 \\ 2 \\ 6 \end{bmatrix}, \quad \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} -3 \\ -4 \\ 9 \end{bmatrix}$$

Thus, $(4, 2, 6)$, $(1, 1, 0)$, and $(-3, -4, 9)$ form a basis for $R(T)$.

[SI*] **Example:**

Let T be multiplication by the matrix A .

$$\begin{bmatrix} 1 & -1 & 3 \\ 5 & 6 & -4 \\ 7 & 4 & 2 \end{bmatrix}$$

Find

- (a) a basis for the range of T .
- (b) a basis for the kernel of T .
- (c) the rank and nullity of T .
- (d) the rank and nullity of A .

Solution:

- (a) A row echelon form of A is

$$A_{ref} = \begin{bmatrix} 1 & -1 & 3 \\ 0 & 1 & \frac{19}{11} \\ 0 & 0 & 0 \end{bmatrix}$$

thus columns 1 and 2 of A is

$$\begin{bmatrix} 1 \\ 5 \\ 7 \end{bmatrix}, \quad \begin{bmatrix} -1 \\ 6 \\ 4 \end{bmatrix}$$

form a basis for the column space of A , which is the range of T .

- (b) The reduced row echelon form of A is

$$A_{rref} = \begin{bmatrix} 1 & 0 & \frac{14}{11} \\ 0 & 1 & -\frac{19}{11} \\ 0 & 0 & 0 \end{bmatrix}$$

so a general solution of $Ax = \mathbf{0}$ is

$$\begin{aligned} x_1 &= -\frac{14}{11}s \\ x_2 &= \frac{19}{11}s \\ x_3 &= s \end{aligned}$$

so

$$\begin{bmatrix} -14 \\ 19 \\ 11 \end{bmatrix}$$

is a basis for the null space of A , which is $\ker(T)$.

(c) $R(T)$ is two-dimensional, so $\text{rank}(T) = 2$.

$\ker(T)$ is one-dimensional, so $\text{nullity}(T) = 1$.

(d) The column space of A is two-dimensional, so $\text{rank}(A) = 2$. The null space of A is one-dimensional, so $\text{nullity}(A) = 1$.

[SI*] Example:

Describe the kernel and range of

- (a) the orthogonal projection on the xz -plane.
- (b) the orthogonal projection on the yz -plane.
- (c) the orthogonal projection on the plane defined by the equation $y = x$.

Solution:

- (a) The kernel is the y -axis, the range is the entire xy -plane.
- (b) The kernel is the x -axis, the range is the entire yz -plane.
- (c) The kernel is the line through the origin perpendicular to the plane $y = x$, the range is the entire plane $y = x$.

[SI*] Example:

In each part, use the given information to find the nullity of the linear transformation T .

- (a) $T : \mathbb{R}^5 \rightarrow \mathbb{R}^7$ has rank 3.
- (b) $T : P_4 \rightarrow P_3$ has rank 1.
- (c) The range of $T : \mathbb{R}^6 \rightarrow \mathbb{R}^3$ is $\mathbb{R}^3$.
- (d) $T : M_{22} \rightarrow M_{22}$ has rank 3.

Solution:

- (a) $\text{nullity}(T) = 5 - \text{rank}(T) = 2$
- (b) $\dim(P_4) = 5$, so $\text{nullity}(T) = 5 - \text{rank}(T) = 4$
- (c) Since $R(T) = \mathbb{R}^3$, T has rank 3.

$$\text{nullity}(T) = 6 - \text{rank}(T) = 3$$

- (d) $\text{nullity}(T) = 4 - \text{rank}(T) = 1$

[SI*] Example:

Let A be a 5×7 matrix with rank 4.

- (a) What is the dimension of the solution space of $Ax = \mathbf{0}$?
- (b) Is $Ax = \mathbf{b}$ consistent for all vectors $\mathbf{b}$ in $\mathbb{R}^5$? Explain.

Solution:

- (a) Remember that the dimension of matrix $m \times n$ is n (the number of column), thus

$$\dim(A) = \text{nullity}(A) + \text{rank}(A)$$

$\text{nullity}(A) = 7 - \text{rank}(A) = 3$, so the solution space of $Ax = \mathbf{0}$ has dimension 3.

- (b) No, since $\text{rank}(A) = 4$ while $\mathbb{R}^5$ has 5 dimension. To be consistent it has to be in $\mathbb{R}^4$.

[SI*] Example:

Let $T : V \rightarrow \mathbb{R}^3$ be a linear transformation from any vector space to $\mathbb{R}^3$. Give a geometric description of $R(T)$.

Solution:

$R(T)$ must be a subspace of $\mathbb{R}^3$, thus the possibilities are a line through the origin, a plane through the origin, the origin only, or all of $\mathbb{R}^3$. It is the definition of subspaces of $\mathbb{R}^3$ from the General Vector Spaces chapter.

[SI*] Example:

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be multiplication by

$$A = \begin{bmatrix} 1 & 3 & 4 \\ 3 & 4 & 7 \\ -2 & 2 & 0 \end{bmatrix}$$

- (a) Show that the kernel of T is a line through the origin, and find parametric equations for it.
- (b) Show that the range of T is a plane through the origin, and find an equation for it.

Solution:

- (a) A reduced row echelon form of matrix A is

$$A_{rref} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

A general solution of $Ax = \mathbf{0}$ is

$$x_1 - x_2 = 0$$

$$x_2 + x_3 = 0$$

$$x_1 = -t$$

$$x_2 = -t$$

$$x_3 = t$$

so

$$\begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix}$$

is a basis for the null space of A , the kernel of T . Vector $(-1, -1, 1)$ is a line through the origin, the parametric equation is

$$(x, y, z) = (-t - t, t) = (-1, -1, 1)t$$

with variable t as the parameter.

- (b) From the reduced row echelon form of matrix A , the columns of A corresponds to the columns of A_{rref} containing leading 1s are

$$\begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 2 \end{bmatrix}$$

thus, $(1, 3, -2)$ and $(3, 4, 2)$ form a basis for $R(T)$ (both vectors span $\mathbb{R}^3$ and not a linear combination to the other). The geometric description of this range of T is a plane through the origin with parametric equations

$$(x, y, z) = (s, 3s, -2s) + (3t, 4t, 2t) = (1, 3, -2)s + (3, 4, 2)t$$

with s, t as the parameters variables.

[SI*] **Example:**

- (a) Show that if a_1, a_2, b_1 , and b_2 are any scalars, then the formula

$$F(x, y) = (a_1x + b_1y, a_2x + b_2y)$$

defines a linear operator on $\mathbb{R}^2$.

- (b) Does the formula $F(x, y) = (a_1x^2 + b_1y^2, a_2x^2 + b_2y^2)$ define a linear operator on $\mathbb{R}^2$? Explain.

Solution:

- (a) We will need to show that $F(x, y)$ follows two properties which are homogeneity and additivity. First, for the homogeneity property

$$\begin{aligned} F(kx, ky) &= (a_1kx + b_1ky, a_2kx + b_2ky) \\ &= k(a_1x + b_1y, a_2x + b_2y) \\ &= kF(x, y) \end{aligned}$$

thus, $F(x, y)$ follows the homogeneity property. Now for the next property,

$$\begin{aligned} F(x + v, y + w) &= (a_1(x + v) + b_1(y + w), a_2(x + v) + b_2(y + w)) \\ &= (a_1x + b_1y, a_2x + b_2y) + (a_1v + b_1w, a_2v + b_2w) \\ &= F(x, y) + F(v, w) \end{aligned}$$

it proven that $F(x, y)$ follows the additivity property. Since $F(x, y)$ follows homogeneity and additivity property then it is a linear operator on $\mathbb{R}^2$.

- (b) No, because it does not follow the homogeneity property

$$\begin{aligned} F(kx, ky) &= (a_1k^2x^2 + b_1k^2y^2, a_2k^2x^2 + b_2k^2y^2) \\ &= k^2(a_1x^2 + b_1y^2, a_2x^2 + b_2y^2) \\ &= k^2F(x, y) \\ &\neq kF(x, y) \end{aligned}$$

[SI*] Example:

Let $\{v_1, v_2, \dots, v_n\}$ be a basis for a vector space V , and let $T : V \rightarrow W$ be a linear transformation. Show that if

$$T(v_1) = T(v_2) = T(v_n) = \mathbf{0}$$

then T is the zero transformation.

Solution:

The zero transformation will have these properties

$$T(v_i + v_j) = 0$$

$$T(v_i) = 0$$

$$T(v_j) = 0$$

$$T(kv_i) = 0$$

for $i, j = 1, 2, \dots, n$, with $i \neq j$. Since $T : V \rightarrow W$ is a linear transformation then it will follow the homogeneity and additivity properties

$$\begin{aligned} T(v_i + v_j) &= T(v_i) + T(v_j) \\ &= 0 + 0 \\ &= 0 \end{aligned}$$

$$\begin{aligned} T(kv_i) &= kT(v_i) \\ &= k(0) \\ &= 0 \end{aligned}$$

thus, T is the zero transformation.

[SI*] Example:

Let $\{v_1, v_2, \dots, v_n\}$ be a basis for a vector space V , and let $T : V \rightarrow V$ be a linear operator. Show that if

$$T(v_1) = v_1, \quad T(v_2) = v_2, \dots, T(v_n) = v_n$$

then T is the identity transformation on V .

Solution:

Let $w = c_1v_1 + c_2v_2 + \dots + c_nv_n$ be any vector in V . Then

$$\begin{aligned} T(w) &= c_1T(v_1) + c_2T(v_2) + \dots + c_nT(v_n) \\ &= c_1v_1 + c_2v_2 + \dots + c_nv_n \\ &= w \end{aligned}$$

Since w was an arbitrary vector in V , T must be the identity operator.

[SI*] Example:

Prove: If $\{v_1, v_2, \dots, v_n\}$ is a basis for V and $w_1, w_2, \dots, w_n$ are vectors in W , not necessarily distinct, then there exists a linear transformation $T : V \rightarrow W$ such that

$$T(v_1) = w_1, \quad T(v_2) = w_2, \dots, T(v_n) = w_n$$

Solution:

If $T : V \rightarrow W$ is a linear transformation, then it has to obey these two properties

$$\begin{aligned} T(\mathbf{0}) &= \mathbf{0} \\ T(v_i - v_j) &= T(v_i) - T(v_j) \end{aligned}$$

for all $i, j = 1, 2, \dots, n$ and $i \neq j$. Now we know that since $\{v_1, v_2, \dots, v_n\}$ is a basis for V , then

$$v_i - v_j \neq 0$$

thus

$$\begin{aligned} T(v_i - v_j) &= T(v_i) - T(v_j) \\ &= w_i - w_j \end{aligned}$$

if we make an assumption that $w_1, w_2, \dots, w_n$ are vectors in W not necessarily distinct, then we will encounter this (assuming $w_i = w_j$)

$$\begin{aligned} T(v_i - v_j) &= w_i - w_j \\ &= 0 \end{aligned}$$

the consequence is $v_i - v_j$ will be in kernel of T too besides 0 . Now, let x be any vector in V with $x = c_1 v_1 + c_2 v_2 + \dots + c_n v_n$, $c_1, c_2, \dots, c_n$ are scalars then

$$\begin{aligned} T(x) &= T(c_1 v_1 + c_2 v_2 + \dots + c_n v_n) \\ &= c_1 T(v_1) + c_2 T(v_2) + \dots + c_n T(v_n) \\ &= c_1 w_1 + c_2 w_2 + \dots + c_n w_n \end{aligned}$$

$T(x) \in W$, and for any scalars k

$$\begin{aligned} T(kx) &= T(c_1 k v_1 + c_2 k v_2 + \dots + c_n k v_n) \\ &= k c_1 T(v_1) + k c_2 T(v_2) + \dots + k c_n T(v_n) \\ &= k(c_1 w_1 + c_2 w_2 + \dots + c_n w_n) \end{aligned}$$

T follows the homogeneity and additivity properties, thus the linear transformation T is proven to exist.

[SI*] **Example:**

Let $D : P_3 \rightarrow P_2$ be the differentiation transformation $D(p) = p'(x)$. What is the kernel of D ?

Solution:

If $p'(x) = 0$, then $p(x)$ is a constant, so $\ker(D)$ consists of all constant polynomials.

[SI*] **Example:**

Let V be the vector space of real-valued functions with continuous derivatives of all orders on the interval $(-\infty, \infty)$, and let $W = F(-\infty, \infty)$ be the vector space of real-valued functions defined on $(-\infty, \infty)$.

(a) Find a linear transformation $T : V \rightarrow W$ whose kernel is P_3 .

(b) Find a linear transformation $T : V \rightarrow W$ whose kernel is P_n .

Solution:

(a) The fourth derivative of any polynomial of degree 3 or less is 0, so

$$T(f(x)) = f^{(4)}(x)$$

has $\ker(T) = P_3$.

- (b) The $n + 1$ -th derivative of any polynomial of degree n or less is 0, so

$$T(f(x)) = f^{(n+1)}(x)$$

has $\ker(T) = P_n$.

[SI*] The vector space $\mathbb{R}^n$ is the "mother" of all real n -dimensional vector spaces in the sense that any such space might differ from $\mathbb{R}^n$ in the notation used to represent vectors, but not in its algebraic structure.

Definition 6.51: One-to-One

If $T : V \rightarrow W$ is a linear transformation from a vector space V to a vector space W , then T is said to be one-to-one if T maps distinct vectors in V into distinct vectors in W .

Definition 6.52: Onto

If $T : V \rightarrow W$ is a linear transformation from a vector space V to a vector space W , then T is said to be onto (or onto W) if every vector in W is the image of at least one vector in V .

Theorem 6.100: Kernel and One-to-One

If $T : V \rightarrow W$ is a linear transformation, then the following statements are equivalent.

- (a) T is one-to-one.
- (b) $\ker(T) = \{\mathbf{0}\}$

Theorem 6.101: Linear Operator in a Finite-Dimensional Vector Space

If V is a finite-dimensional vector space, and if $T : V \rightarrow V$ is a linear operator, then the following statements are equivalent.

- (a) T is one-to-one.
- (b) $\ker(T) = \{\mathbf{0}\}$.
- (c) T is onto [i.e., $R(T) = V$]

[SI*] Dilations and Contractions Are One-to-One and Onto

If V is a finite-dimensional vector space and c is any nonzero scalar, then the linear operator $T : V \rightarrow V$ defined by $T(v) = cv$ is one-to-one and onto.

We know that the operator T is onto (and hence one-to-one) for if v is any vector in V then that vector is the image of the vector $(1/c)v$.

[SI*] Matrix Operators

If $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is the matrix operator $T_A(x) = Ax$, then it follows from theorem in the Eigenvalues and Eigenvectors section that

- (a) The range of T_A is $\mathbb{R}^n$.
- (b) T_A is one-to-one.

so T_A is one-to-one and onto if and only if A is invertible.

[SI*] **Shifting Operators**

Let $V = \mathbb{R}^\infty$ be the sequence and consider the linear "shifting operators" on V defined by

$$\begin{aligned} T_1(u_1, u_2, \dots, u_n, \dots) &= (0, u_1, u_2, \dots, u_n, \dots) \\ T_2(u_1, u_2, \dots, u_n, \dots) &= (u_2, u_3, \dots, u_n, \dots) \end{aligned}$$

T_1 is one-to-one because distinct dequences in $\mathbb{R}^\infty$ obviously have distinct images. This operator is not onto because no vector in $\mathbb{R}^\infty$ maps into the sequence $(1, 0, 0, \dots, 0, \dots)$, for example.

T_2 is not one-to-one because the vectors $(1, 0, 0, \dots, 0, \dots)$ and $(2, 0, 0, \dots, 0, \dots)$ both map into $(0, 0, 0, \dots, 0, \dots)$. This operator is onto because every possible sequence of real numbers can be obtained with an appropriate choice of the numbers $u_2, u_3, \dots, u_n, \dots$.

[SI*] **Basic Transformations That Are One-to-One and Onto**

The linear transformation $T_1 : P_3 \rightarrow \mathbb{R}^4$ and $T_2 : M_{22} \rightarrow \mathbb{R}^4$ defined by

$$\begin{aligned} T_1(a + bx + cx^2 + dx^3) &= (a, b, c, d) \\ T_2\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) &= (a, b, c, d) \end{aligned}$$

are both one-to-one and onto. Their kernels contain only the zero vector.

[SI*] **A One-to-One Linear Transformation**

Let $T : P_n \rightarrow P_{n+1}$ be the linear transformation

$$T(\mathbf{p}) = T(p(x)) = xp(x)$$

If

$$\begin{aligned} \mathbf{p} &= p(x) = c_0 + c_1x + \dots + c_nx^n \\ \mathbf{q} &= q(x) = d_0 + d_1x + \dots + d_nx^n \end{aligned}$$

are distinct polynomials, then they differ in at least one coefficient. Thus,

$$\begin{aligned} T(\mathbf{p}) &= c_0x + c_1x^2 + \dots + c_nx^{n+1} \\ T(\mathbf{q}) &= d_0x + d_1x^2 + \dots + d_nx^{n+1} \end{aligned}$$

also differ in at least one coefficient. It follows that T is one-to-one since it maps distinct polynomials $\mathbf{p}$ and $\mathbf{q}$ into distinct polynomials $T(\mathbf{p})$ and $T(\mathbf{q})$.

[SI*] **A Transformation That Is Not One-to-One**

Let

$$D : C^1(-\infty, \infty) \rightarrow F(-\infty, \infty)$$

be the differentiation transformation. This linear transformation is not one-to-one because it maps functions that differ by constant into the same function. For example,

$$D(x^2) = D(x^2 + 1) = 2x$$

Remember that $F(-\infty, \infty)$ represents the set of functions with derivatives of all orders on $(-\infty, \infty)$, while $C^1(-\infty, \infty)$ represents functions with the first derivatives on $(-\infty, \infty)$ that is continuous, and $C(-\infty, \infty)$ is the set of continuous functions on $(-\infty, \infty)$.

[SI*] Let $T : V \rightarrow W$ be a linear transformation, where V and W are finite-dimensional. Then

1. If $\dim(W) < \dim(V)$, then T cannot be one-to-one.
2. If $\dim(V) < \dim(W)$, then T cannot be onto.

These observations tell us, that any linear transformation from $\mathbb{R}^3$ to $\mathbb{R}^2$ must map some distinct points of $\mathbb{R}^3$ into the same point in $\mathbb{R}^2$, and it also tells us that there is no linear transformation that maps $\mathbb{R}^2$ onto all of $\mathbb{R}^3$.

Definition 6.53: Isomorphism

If a linear transformation $T : V \rightarrow W$ is both one-to-one and onto, then T is said to be an isomorphism, and the vector spaces V and W are said to be isomorphic.

[SI*] Isomorphic vector spaces have the same "algebraic form," even though they may consist of different kinds of objects.

The isomorphism

$$a_0 + a_1x + a_2x^2 \vec{T}(a_0, a_1, a_2)$$

matches up vector operations in P_2 and $\mathbb{R}^3$.

Theorem 6.102: Isomorphism for n -Dimensional Vector Space

Every real n -dimensional vector space is isomorphic to $\mathbb{R}^n$.

[SI*] Let V be a real n -dimensional vector space. To prove that V is isomorphic to $\mathbb{R}^n$ we must find a linear transformation $T : V \rightarrow \mathbb{R}^n$ that is one-to-one and onto. For this purpose, let

$$v_1, v_2, \dots, v_n$$

be any basis for V , let

$$u = k_1v_1 + k_2v_2 + \dots + k_nv_n \quad (6.158)$$

be the representation of a vector u in V as a linear combination of the basis vectors, and define the transformation $T : V \rightarrow \mathbb{R}^n$ by

$$T(u) = (k_1, k_2, \dots, k_n) \quad (6.159)$$

We will show that T is an isomorphism (linear, one-to-one, and onto). To prove the linearity, let u and v be vectors in V , let c be a scalar, and let

$$\begin{aligned} u &= k_1v_1 + k_2v_2 + \dots + k_nv_n \\ v &= d_1v_1 + d_2v_2 + \dots + d_nv_n \end{aligned} \quad (6.160)$$

be the representations of u and v as linear combinations of the basis vectors. Then it follows that

$$\begin{aligned} T(cu) &= T(ck_1v_1 + ck_2v_2 + \dots + ck_nv_n) \\ &= (ck_1, ck_2, \dots, ck_n) \\ &= c(k_1, k_2, \dots, k_n) \\ &= cT(u) \end{aligned}$$

and it follows that

$$\begin{aligned}
 T(\mathbf{u} + \mathbf{v}) &= T((k_1 + d_1)\mathbf{v}_1 + (k_2 + d_2)\mathbf{v}_2 + \cdots + (k_n + d_n)\mathbf{v}_n) \\
 &= (k_1 + d_1, k_2 + d_2, \dots, k_n + d_n) \\
 &= (k_1, k_2, \dots, k_n) + (d_1, d_2, \dots, d_n) \\
 &= T(\mathbf{u}) + T(\mathbf{v})
 \end{aligned}$$

which shows that T is linear. To show that T is one-to-one, we must show that if $\mathbf{u}$ and $\mathbf{v}$ are distinct vectors in V , then so are their images in $\mathbb{R}^n$. But if $\mathbf{u} \neq \mathbf{v}$, and if the representations of these vectors in terms of the basis vectors, then we must have $k_i \neq d_i$ for at least one i . Thus,

$$T(\mathbf{u}) = (k_1, k_2, \dots, k_n) \neq (d_1, d_2, \dots, d_n) = T(\mathbf{v})$$

which shows that $\mathbf{u}$ and $\mathbf{v}$ have distinct images under T . Finally, the transformation T is onto, for if

$$\mathbf{w} = (k_1, k_2, \dots, k_n)$$

is any vector in $\mathbb{R}^n$, then it follows that $\mathbf{w}$ is the image under T of the vector

$$\mathbf{u} = k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + \cdots + k_n\mathbf{v}_n$$

Remark

Note that the isomorphism T of the foregoing proof is the coordinate map

$$\mathbf{u} \vec{T}(k_1, k_2, \dots, k_n) = (\mathbf{u})_S$$

that maps $\mathbf{u}$ into its coordinate vector with respect to the basis $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$. Since there are generally many possible bases for a given vector space V , there are generally many possible isomorphisms between V and $\mathbb{R}^n$, one for each different basis.

[SI*] The Natural Isomorphism from P_{n-1} to $\mathbb{R}^n$

The mapping

$$a_0 + a_1x + \cdots + a_{n-1}x^{n-1} \xrightarrow{T} (a_0, a_1, \dots, a_{n-1})$$

from P_{n-1} to $\mathbb{R}^n$ is one-to-one, onto, and linear. This is called the natural isomorphism from P_{n-1} to $\mathbb{R}^n$ because, as the following computations show, it maps the natural basis $\{1, x, x^2, \dots, x^{n-1}\}$ for P_{n-1} into the standard basis for $\mathbb{R}^n$:

$$\begin{array}{rclcl}
 1 & = & 1 + 0x + 0x^2 + \cdots + 0x^{n-1} & \xrightarrow{T} & (1, 0, 0, \dots, 0) \\
 x & = & 0 + x + 0x^2 + \cdots + 0x^{n-1} & \xrightarrow{T} & (0, 1, 0, \dots, 0) \\
 \vdots & & \vdots & & \vdots \\
 x^{n-1} & = & 0 + 0x + 0x^2 + \cdots + 0x^{n-2} + x^{n-1} & \xrightarrow{T} & (0, 0, 0, \dots, 0, 1)
 \end{array}$$

[SI*] The Natural Isomorphism from M_{22} to $\mathbb{R}^4$

The matrices

$$E_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad E_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad E_4 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

form a basis for the vector space M_{22} of 2×2 matrices. An isomorphism $T : M_{22} \rightarrow \mathbb{R}^4$ can be constructed by first writing a matrix A in M_{22} in terms of the basis vectors as

$$A = \begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix} = a_1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + a_2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + a_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + a_4 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

and then defining T as

$$T(A) = (a_1, a_2, a_3, a_4)$$

Thus, for example,

$$\begin{bmatrix} 1 & -3 \\ 4 & 6 \end{bmatrix} \xrightarrow{T} (1, -3, 4, 6)$$

More generally, this idea can be used to show that the vector space M_{mn} of $m \times n$ matrices with real entries is isomorphic to $\mathbb{R}^{mn}$.

[SI*] **Differentiation by Matrix Multiplication**

Consider the differentiation transformation $D : P_3 \rightarrow P_2$ on the vector space of polynomials of degree three or less. If we map P_3 and P_2 into $\mathbb{R}^4$ and $\mathbb{R}^3$, respectively, by the natural isomorphisms, then the transformation D produces a corresponding matrix transformation from $\mathbb{R}^4$ to $\mathbb{R}^3$. Specifically, the derivative transformation

$$a_0 + a_1x + a_2x^2 + a_3x^3 \xrightarrow{T} a_1 + 2a_2x + 3a_3x^2$$

produces the matrix transformation

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} a_1 \\ 2a_2 \\ 3a_3 \end{bmatrix}$$

Thus, for example, the derivative

$$\frac{d}{dx}(2 + x + 4x^2 - x^3) = 1 + 8x - 3x^2$$

can be calculated as the matrix product

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 4 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 8 \\ -3 \end{bmatrix}$$

This idea is useful for constructing numerical algorithms to perform derivative calculations.

[SI*] In the case where V is a real n -dimensional inner product space, both V and $\mathbb{R}^n$ have, in addition to their algebraic structure, a geometric structure arising from their respective inner products. Thus, it is reasonable to inquire if there exists an isomorphism from V to $\mathbb{R}^n$ that preserves the geometric structure as well as the algebraic structure. For example, we would want orthogonal vectors in V to have orthogonal counterparts in $\mathbb{R}^n$, and we would want orthonormal sets in V to correspond to orthonormal sets in $\mathbb{R}^n$.

In order for an isomorphism to preserve geometric structure, it obviously has to preserve inner products, since notions of length, angle, and orthogonality are all based on the inner product. Thus, if V and W are inner product spaces, then we call an isomorphism $T : V \rightarrow W$ an inner product space isomorphism if

$$\langle T(u), T(v) \rangle = \langle u, v \rangle$$

It can be proved that if V is any real n -dimensional inner product space and $\mathbb{R}^n$ has the Euclidean inner product (the dot product), then there exists an inner product space

isomorphism from V to $\mathbb{R}^n$. Under such an isomorphism, the inner product space V has the same algebraic and geometric structure as $\mathbb{R}^n$. In this sense, every n -dimensional inner product space is a "carbon copy" of $\mathbb{R}^n$ with the Euclidean inner product that differs only in the notation used to represent vectors.

[SI*] **An Inner Product Space Isomorphism**

Let $\mathbb{R}^n$ be the vector space of real n -tuples in comma-delimited form, let M_n be the vector space of real $n \times 1$ matrices, let $\mathbb{R}^n$ have the Euclidean inner product $\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u} \cdot \mathbf{v}$, and let M_n have the inner product $\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u}^T \mathbf{v}$ in which $\mathbf{u}$ and $\mathbf{v}$ are expressed in column form. The mapping $T : \mathbb{R}^n \rightarrow M_n$ defined by

$$(v_1, v_2, \dots, v_n) \xrightarrow{T} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$$

is an inner product space isomorphism, so the distinction between the inner product space $\mathbb{R}^n$ and the inner product space M_n is essentially notational, a fact that we have used many times in this text.

i. **Functions in SymIntegration Related to Linear Transformations**

[SI*] We will list the basic functions that can be used related to linear transformations:

IX. NUMERICAL METHODS

[SI*] From the first section on this chapter of Numerical Linear Algebra, we have learned about Gaussian elimination and backward substitution to obtain the solution for systems of linear equations, they are fine for small-scale problems, but Gaussian elimination and Gauss-Jordan elimination are not suitable for large-scale problems in which compute roundoff error, memory usage, and speed are concerns. For large-scale and special case we can use numerical methods to solve systems of linear equations, these are the available methods currently:

1. **LU Decomposition**

A matrix factorization technique into lower triangular L and upper triangular U , it is used to solve systems of linear equations. It transforms a complex matrix problem into two simpler problems that can be solved sequentially using forward and backward substitution.

2. **QR Decomposition**

more stable than LU decomposition, but slower to converge.

3. **Cholesky Decomposition**

It is used if the system is symmetric.

4. **Singular Value Decomposition (SVD)**

If the system is not square (the number of equations is not the same as the number of variables).

Decomposition has better efficiency, for a given matrix A , the costly decomposition step is performed only once. This is faster for repeated solving of $A\mathbf{x} = \mathbf{b}$ with different $\mathbf{b}$ vectors compared to reinverting the matrix each time.

i. Functions in SymIntegration Related to Numerical Methods

[SI*] We will list the basic functions that can be used related to numerical methods:

LUsolve_nhsystem(vector<vector<double>> &A, vector<vector<double>> &b, vector<double> &x)

to find the solution of a linear system $Ax = b$ with LU -decomposition.

LUDecomposition(vector<vector<double>> &A, vector<vector<double>> &L, vector<vector<double>> &U)

to compute the LU -decomposition of matrix A , it is a void function so you have to call the matrix L and U later on in the source code, remember also this function doesn't compute the solution of the linear system.

SVD(vector<vector<double>> &A)

to compute singular value decomposition (SVD) with A as the matrix that we want to apply the SVD, SVD relies heavily on eigenvalues and eigenvectors computation for all the U, Σ, V^T matrices.

ii. LU -Decomposition

The credit for popularizing the matrix formulation of the LU -decomposition is often given to the British mathematician Alan Turing for his work on the subject in 1948. Turing, one of the great geniuses of the twentieth century is the founder of the field of artificial intelligence. Sadly, Turing, a homosexual, was tried and convicted of "gross indecency" in 1952, in violation of the then-existing British statutes. Depressed, he committed suicide at age 41 by eating an apple laced with cyanide. Lately before passed away in the 21st century, Queen Elizabeth II has done public apology for what has happened for Alan Turing, a world war II hero, without him Nazi cannot be stopped, this shows how close minded people in the past are, they read less, comprehend less, but judge more, just like the time when church burned Jeanne D'arc, or burn Galileo Galilei paper works, then apology after hundreded of years and made Jeanne D'arc a saint. It is human nature to judge what they don't know, out of fear, people are different just like the world full of color, so by learning science, computer and engineering, hopefully the result will make people have better understanding, compassion, and have higher quality of life.

Definition 6.54: LU -Decomposition

A factorization of a square matrix A as $A = LU$, where L is lower triangular and U is upper triangular is called an LU -decomposition (or LU -factorization) of A .

Not every square matrix has an LU -decomposition.

To systemically solve for entries in L and U from matrix A , the formula for matrix U :

$$\forall j \begin{cases} i = 0 \rightarrow U_{ij} = A_{ij} \\ i > 0 \rightarrow U_{ij} = A_{ij} - \sum_{k=0}^{i-1} L_{ik}U_{kj} \end{cases} \quad (6.161)$$

the formula for matrix L :

$$\forall i \begin{cases} j = 0 \rightarrow L_{ij} = \frac{A_{ij}}{U_{jj}} \\ j > 0 \rightarrow L_{ij} = \frac{A_{ij} - \sum_{k=0}^{j-1} L_{ik}U_{kj}}{U_{jj}} \end{cases} \quad (6.162)$$

In LU -decomposition we are using Doolittle algorithm, it is an efficient method for LU -decomposition, enabling various applications in numerical analysis and linear algebra. Applications on Doolittle algorithm:

1. Solving Linear Equations

LU -decomposition simplifies solving $Ax = b$ by solving $Lz = b$ then $Ux = y$.

2. Matrix Inversion

Decompose A into LU , then invert L and U .

3. Determinant Calculation

The determinant of A is the product of the diagonal elements of U . The formula is

$$\det(A) = \det(L) \times \det(U)$$

since $\det(L) = 1$, thus we only need to compute $\det(U)$.

With variations like partial pivoting, it creates numerical stability, to made it more robust against numerical errors that can occur during the process.

Applications of LU -decomposition:

1. Structural Engineering

It is used to analyze forces and stresses in bridges and buildings, ensuring efficient and safe designs then model the structure's behavior. It can be used with modal flexibility matrices to identify and locate damage in a structure by analyzing changes in its' structural properties.

2. Finite Element Analysis (FEA)

FEA relies heavily on matrix-based methods. LU -decomposition is used to solve the resulting stiffness matrices, which represent the relationship between forces and displacements in a structure.

3. Heat Transfer

It solves the systems of equations that result from discretizing partial differential equations used to model heat flow. It is used to solve for temperature distribution in objects when modeled using finite difference or finite element methods.

4. Fluid Dynamics

It is applied in computational fluid dynamics (CFD) to solve the pressure and velocity fields in fluid flow, which is often modeled by a system of linear equations.

5. Geotechnical Engineering

It can be applied in solving problems related to soil mechanics and foundation design that involve large systems of equations.

6. Computer Graphics

It helps in transforming 3D objects, like rotating and scaling models, for smoother rendering.

7. Robotics

It assists in solving kinematic equations, enabling real-time movement adjustments for robots.

8. Weather Prediction

It speeds up the solving of climate models and weather simulations for accurate forecasting.

9. Electrical Engineering

It is used in circuit analysis to solve systems of equations for designing and optimizing electrical circuits. *LU*-decomposition is a core component of the numerical iterative algorithms used to solve the power flow problem in power systems. It is used in methods like the Method of Moments for analyzing electromagnetic radiation and scattering, especially when a matrix needs to be decomposed repeatedly for different configurations, saving computation time. It can also be applied to matrix equations that arise in antenna design, particularly when optimizing parameters on a fixed geometry.

10. Modeling Complex Systems

Chemical processes often involve multiple interacting components, leading to a large system of linear equations that describe the process. *LU*-decomposition provides a structured way to solve these systems.

11. Steady-State and Transient Analysis

It is used in analyzing systems at steady-state and in transient simulations where the system evolves over time.

12. Chemical Reaction Networks

It is used for calculating concentrations of reactants and products in a system with multiple parallel or consecutive reactions.

13. Economics and Finance

It helps solve economic models for resource allocation and market predictions efficiently.

*Remember that *LU*-decompositions are not unique.

iii. Solve Nonhomogeneous System of Linear Equation with *LU*-Decomposition in SymIntegration

Solve the linear system

$$\begin{bmatrix} 1 & 1 & 1 \\ 4 & 3 & -1 \\ 3 & 5 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 4 \end{bmatrix}$$

Solution:

We can rewrite this system as

$$\begin{bmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 3 & -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 0 & -1 & -5 \\ 0 & 0 & -10 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 4 \end{bmatrix}$$

Let us define z_1, z_2 , and z_3 by the equation $Ux = z$

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & -1 & -5 \\ 0 & 0 & -10 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix}$$

then we can write the equation $Lz = b$

$$\begin{bmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 3 & -2 & 1 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 4 \end{bmatrix}$$

the equation $Lz = b$ can be written equivalently as

$$\begin{array}{rcl} z_1 & & = 1 \\ 4z_1 & + z_2 & = 6 \\ 3z_1 & - 2z_2 & + z_3 = 4 \end{array}$$

now by using forward substitution we will obtain

$$z_1 = 1, z_2 = 2, z_3 = 5$$

We go back to the equation $Ux = z$ and substitute z into the equation, which yields the linear system

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & -1 & -5 \\ 0 & 0 & -10 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix}$$

or equivalently,

$$\begin{array}{rcl} x_1 & + x_2 & + x_3 = 1 \\ & -x_2 & - 5x_3 = 2 \\ & & -/10x_3 = 5 \end{array}$$

now by using back substitution we will obtain

$$x_1 = 1, x_2 = 0.5, x_3 = -0.5$$

In SymIntegration, to make the matrices L and U with *LU*-decomposition we can use this function:

LUDecomposition(vector<vector<double>> &A, vector<vector<double>> &L, vector<vector<double>> &U)

the main source code for this function is located in **src/numericalmethod.cpp**.

```
#include<bits/stdc++.h>
#include<iostream>
#include "symintegrationc++.h"
#include<vector>
#include <chrono>
#include <algorithm> // For std::next_permutation
#include <string>
using namespace std::chrono;
using namespace std;

// Driver program
int main()
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    //string filename = "matrix.txt";

    dmat doubleMatrix = loadMatrixFromFile("matrix.txt");
    dmat vecb = loadMatrixFromFile("vectorb.txt");
    int n = doubleMatrix.size();
    int R = doubleMatrix.size();
    int C = doubleMatrix[0].size();

    // Declare the matrix L and U with the size.
    dmat L(n, vector<double>(n));
    dmat U(n, vector<double>(n));

    printMatrix(doubleMatrix);

    LUDecomposition(doubleMatrix, L, U);

    cout << "\nMatrix A : " << endl;
    printMatrix(doubleMatrix);
    cout << "\nMatrix L : " << endl;
    printMatrix(L);
    cout << "\nMatrix U : " << endl;
    printMatrix(U);

    // Get ending timepoint
    auto stop = high_resolution_clock::now();
    auto duration = duration_cast<microseconds>(stop - start);

    cout << "\nTime taken by function: " << duration.count() << " microseconds"
        << endl;
```

```

    return 0;
}

```

Code 59: Test SymIntegration LU Decomposition/main.cpp

```

root { ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration LU Decomposition }# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root { ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration LU Decomposition }# ./main
Data:
      1      1      1
      4      3     -1
      3      5      3

Matrix A :
      1.000000      1.000000      1.000000
      4.000000      3.000000     -1.000000
      3.000000      5.000000      3.000000

Matrix L :
      1.000000      0.000000      0.000000
      4.000000      1.000000      0.000000
      3.000000     -2.000000      1.000000

Matrix U :
      1.000000      1.000000      1.000000
      0.000000     -1.000000     -5.000000
      0.000000      0.000000     -10.000000

Time taken by function: 1485 microseconds

```

Figure 6.60: The LU-decomposition of matrix A (SymIntegration/Examples/Linear Algebra/Test SymIntegration LU Decomposition/main.cpp).

To find the solution with LU-decomposition, we use this function:
LUsolve_nhsystem(vector<vector<double>> &A, vector<vector<double>> &b, vector<double> &x)
 the main source code for this function is located in **src/linearalgebra.cpp**.

```

#include<bits/stdc++.h>
#include<iostream>
#include "symintegrationc++.h"
#include<vector>
#include <chrono>
#include <algorithm> // For std::next_permutation
#include <string>

using namespace std::chrono;
using namespace std;

// Driver program
int main()
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    dmat doubleMatrix = loadMatrixFromFile("matrix.txt");
    dmat vecb = loadMatrixFromFile("vectorb.txt");
    int n = doubleMatrix.size();

    cout << "Data:" << endl;
    printMatrix(doubleMatrix);

```



```

    dvec x;
    LUSolve_nhsystem(doubleMatrix, vecb, x);
    cout << "\nx : " << endl;
    printVector(x);

    // Get ending timepoint
    auto stop = high_resolution_clock::now();
    auto duration = duration_cast<microseconds>(stop - start);

    cout << "\nTime taken by function: " << duration.count() << " microseconds"
        << endl;

    return 0;
}

```

Code 60: Test SymIntegration LU Solve/main.cpp

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration LU Solve ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration LU Solve ]# ./main
Data:
      1.000000      1.000000      1.000000
      4.000000      3.000000     -1.000000
      3.000000      5.000000      3.000000

Solution for Lz = b:
z1 = 1.00000
z2 = 2.00000
z3 = 5.00000

Solution for Ux = z:
x1 = 1.00000
x2 = 0.50000
x3 = -0.50000

x :
1.000000
0.500000
-0.500000

Time taken by function: 1142 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration LU Solve ]#

```

Figure 6.61: Computing the solution for $Ax = b$ with LU-decomposition (SymIntegration/Examples/Linear Algebra/Test SymIntegration LU Solve/main.cpp).

We commented some **cout** commands and only show the solution vector x , it decreases the function time making it faster to show the final result.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration LU Solve ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration LU Solve ]# ./main
Data:
      1.000000      1.000000      1.000000
      4.000000      3.000000     -1.000000
      3.000000      5.000000      3.000000

x :
1.000000
0.500000
-0.500000

Time taken by function: 982 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration LU Solve ]#

```

Figure 6.62: Computing the solution for $Ax = b$ with LU-decomposition and only show the solution vector x (*SymIntegration/Examples/Linear Algebra/Test SymIntegration LU Solve/main.cpp*).

We also compare it with Gaussian elimination function, to see which algorithm could compute the solution vector x faster, *LU*-decomposition is faster than Gaussian elimination.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination ]# ./main
Data:
      1.000000      1.000000      1.000000
      4.000000      3.000000     -1.000000
      3.000000      5.000000      3.000000

Augmented Matrix:
      1.000000      1.000000      1.000000      1.000000
      4.000000      3.000000     -1.000000      0.000000
      3.000000      5.000000      3.000000      4.000000

Augmented Matrix in reduced row form:
      4.000000      3.000000     -1.000000      6.000000
      0.000000      2.750000      3.750000     -0.500000
      0.000000      0.000000      0.909091     -0.454545

Solution:
x1 = 1.00000
x2 = 0.50000
x3 = -0.50000

x :
1.000000
0.500000
-0.500000

Time taken by function: 1293 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gaussian Elimination ]#

```

Figure 6.63: Computing the solution for $Ax = b$ with Gaussian elimination (*SymIntegration/Examples/Linear Algebra/Test SymIntegration Gaussian Elimination Function/main.cpp*).

iv. Cholesky Decomposition

Cholesky decomposition or Cholesky factorization is a decomposition of a Hermitian, positive-definite matrix into the product of a lower triangular matrix and its conjugate transpose. The Cholesky decomposition is roughly twice as efficient as the LU decomposition for solving systems of linear equations. The Cholesky decomposition of a Hermitian positive-definite matrix Σ_X is a decomposition of the form

$$\Sigma_X = AA^T$$

where A is a real lower triangular matrix with positive diagonal entries, and A^T denotes the conjugate transpose of A . Every Hermitian positive-definite matrix (and thus also every real-valued symmetric positive-definite matrix) has a unique Cholesky decomposition.

$$\begin{bmatrix} \Sigma_{X_{11}} & \Sigma_{X_{12}} & \Sigma_{X_{13}} \\ \Sigma_{X_{21}} & \Sigma_{X_{22}} & \Sigma_{X_{23}} \\ \Sigma_{X_{31}} & \Sigma_{X_{32}} & \Sigma_{X_{33}} \end{bmatrix} = \begin{bmatrix} A_{11} & 0 & 0 \\ A_{21} & A_{22} & 0 \\ A_{31} & A_{32} & A_{33} \end{bmatrix} \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ 0 & A_{22} & A_{32} \\ 0 & 0 & A_{33} \end{bmatrix}$$

Every symmetric, positive definite matrix Σ_X can be decomposed into a product of a unique lower triangular matrix L and its transpose.

Definition 6.55: Cholesky Decomposition Algorithm

The following formulas are obtained by solving above lower triangular matrix and its transpose. These are the basis of Cholesky decomposition algorithm:

$$A_{ii} = \sqrt{\Sigma_{X_{ii}} - \sum_{k=0}^{i-1} (A_{ik})^2}$$

v. Singular Value Decomposition

[SI*] This is an extension of the diagonalization theory for $n \times n$ symmetric matrices to general $m \times n$ matrices.

The results have applications to compression, storage, and transmission of digitized information and form the basis for many of the best computational algorithms that are currently available for solving linear systems.

[SI*] Every symmetric matrix A can be expressed as

$$A = PDP^T \tag{6.163}$$

where P is an $n \times n$ orthogonal matrix of eigenvectors of A , and D is the diagonal matrix whose diagonal entries are the eigenvalues corresponding to the column vectors of P . We will call (6.163) an eigenvalue decomposition of A .

[SI*] If an $n \times n$ matrix A is not symmetric, then it does not have an eigenvalue decomposition, but it does have a Hessenberg decomposition

$$A = PHP^T$$

in which P is an orthogonal matrix and H is in upper Hessenberg form. Moreover, if A has real eigenvalues, then it has a Schur decomposition

$$A = PSP^T$$

in which P is an orthogonal matrix and S is upper triangular.

[SI*] The eigenvalue, Hessenberg, and Schur decompositions are important in numerical algorithms not only because the matrices D , H , and S have simpler forms than A , but also because the orthogonal matrices that appear in these factorizations do not magnify roundoff error. To see why this is so, suppose $\hat{x}$ is a column vector whose entries are known exactly and that

$$x = \hat{x} + e$$

is the vector that results when roundoff error is present in the entries of $\hat{x}$. If P is an orthogonal matrix, then the length-preserving property of orthogonal transformations implies that

$$\|Px - P\hat{x}\| = \|x - \hat{x}\| = \|e\|$$

which tells us that the error in approximating $P\hat{x}$ by Px has the same magnitude as the error in approximating $\hat{x}$ by x .

[SI*] There are two main paths that one might follow in looking for other kinds of decompositions of a general square matrix A : One might look for decompositions of the form

$$A = PJP^{-1}$$

in which P is invertible but not necessarily orthogonal, or one might look for decompositions of the form

$$A = U\Sigma V^T$$

in which U and V are orthogonal but not necessarily the same. The first path leads to decompositions in which J is either diagonal or a certain kind of block diagonal matrix, called a Jordan canonical form in honor of the French mathematician Camille Jordan. Jordan canonical forms are important theoretically and in certain applications, but they are of lesser importance numerically because of the roundoff difficulties that result from the lack of orthogonality in P .

Theorem 6.103: Matrices A and $A^T A$

If A is an $m \times n$ matrix, then:

- (a) A and $A^T A$ have the same null space.
- (b) A and $A^T A$ have the same row space.
- (c) A^T and $A^T A$ have the same column space.
- (d) A and $A^T A$ have the same rank.

Definition 6.56: Singular Value of A

If A is an $m \times n$ matrix, and if $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of $A^T A$, then the numbers

$$\sigma_1 = \sqrt{\lambda_1}, \quad \sigma_2 = \sqrt{\lambda_2}, \dots \quad \sigma_n = \sqrt{\lambda_n}$$

are called the singular values of A .

Example:

Find the singular values of the matrix

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$$

Solution:

The first step is to find the eigenvalues of the matrix

$$A^T A = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$$

The characteristic polynomial of $A^T A$ is

$$\lambda^2 - 4\lambda + 3 = (\lambda - 3)(\lambda - 1)$$

so the eigenvalues of $A^T A$ are

$$\lambda_1 = 3$$

$$\lambda_2 = 1$$

and the singular values of A in order of decreasing size are

$$\sigma_1 = \sqrt{\lambda_1} = \sqrt{3}$$

$$\sigma_2 = \sqrt{\lambda_2} = 1$$

[SI*] We will find it useful to extend the notion of a "main diagonal" to matrices that are not square. We define the main diagonal of an $m \times n$ matrix to be the line of entries that starts at the upper left corner and extends diagonally as far as it can go. We will refer to the entries on the main diagonal as the diagonal entries.

[SI*] Factorization of a general $m \times n$ matrix A is called singular value decomposition (abbreviated SVD).

Theorem 6.104: Singular Value Decomposition

If A is an $m \times n$ matrix, then A can be expressed in the form

$$A = U \Sigma V^T$$

where U and V are orthogonal matrices and Σ is an $m \times n$ matrix whose diagonal entries are the singular values of A and whose other entries are zero.

Theorem 6.105: Singular Value Decomposition (Expanded Form)

If A is an $m \times n$ matrix of rank k , then A can be factored as

$$A = U \Sigma V^T = \begin{bmatrix} \mathbf{u}_1 & \mathbf{u}_2 & \dots & \mathbf{u}_k & | & \mathbf{u}_{k+1} & \dots & \mathbf{u}_m \end{bmatrix} \begin{bmatrix} \sigma_1 & 0 & \dots & 0 & | \\ 0 & \sigma_2 & \dots & 0 & | \\ \vdots & \vdots & \ddots & \vdots & | \\ 0 & 0 & \dots & \sigma_k & | \end{bmatrix}$$

in which U, Σ , and V have sizes $m \times m$, $m \times n$, and $n \times n$, respectively, and in which

- (a) $V = [\mathbf{v}_1 \ \mathbf{v}_2 \ \dots \ \mathbf{v}_n]$ orthogonally diagonalizes $A^T A$.
- (b) The nonzero diagonal entries of Σ are $\sigma_1 = \sqrt{\lambda_1}$, $\sigma_2 = \sqrt{\lambda_2}$, $\dots$, $\sigma_k = \sqrt{\lambda_k}$, where $\lambda_1, \lambda_2, \dots, \lambda_k$ are the nonzero eigenvalues of $A^T A$ corresponding to the column vectors of V .
- (c) The column vector of V are ordered so that $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_k > 0$.
- (d) $\mathbf{u}_i = \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|} = \frac{1}{\sigma_i} A\mathbf{v}_i \quad (i = 1, 2, \dots, k)$
- (e) $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$ is an orthonormal basis for $\text{col}(A)$.
- (f) $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \mathbf{u}_{k+1}, \dots, \mathbf{u}_m\}$ is an extension of $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$ to an orthonormal basis for $\mathbb{R}^m$.

vi. Compute Singular Value Decomposition with SymIntegration

Find a singular value decomposition of the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$$

Solution:

We will have

$$A^T A = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$$

The characteristic polynomial of $A^T A$ is

$$\lambda^2 - 4\lambda + 3 = (\lambda - 3)(\lambda - 1)$$

so the eigenvalues of $A^T A$ are

$$\lambda_1 = 3$$

$$\lambda_2 = 1$$

and the corresponding singular values of A in order of decreasing size are

$$\sigma_1 = \sqrt{\lambda_1} = \sqrt{3}$$

$$\sigma_2 = \sqrt{\lambda_2} = 1$$

the eigenvectors corresponding to λ_1 and λ_2 are

$$v_1 = \begin{bmatrix} \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{bmatrix}, \quad v_2 = \begin{bmatrix} \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{bmatrix}$$

we know that $V = [v_1 | v_2]$ orthogonally diagonalizes $A^T A$. Thus, the vectors

$$u_1 = \frac{1}{\sigma_1} A v_1 = \frac{\sqrt{3}}{3} \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{6}}{3} \\ \frac{\sqrt{6}}{6} \\ \frac{\sqrt{6}}{6} \end{bmatrix}$$

$$u_2 = \frac{1}{\sigma_1} A v_2 = (1) \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{bmatrix} = \begin{bmatrix} 0 \\ -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{bmatrix}$$

are two of the three column vectors of U . Note that u_1 and u_2 are orthonormal, as expected.

We could extend the set $\{u_1, u_2\}$ to an orthonormal basis for $\mathbb{R}^3$. However, the computations will be easier if we first remove the messy radicals by multiplying u_1 and u_2 by appropriate scalars. Thus, we will look for a unit vector u_3 that is orthogonal to

$$\sqrt{6}u_1 = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}, \quad \sqrt{2}u_2 = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

to satisfy these two orthogonality conditions, the vector $\mathbf{u}_3$ must be a solution of the homogeneous linear system

$$\begin{bmatrix} 2 & 1 & 1 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

the general solution of this system is

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$$

Normalizing the vector on the right yields

$$\mathbf{u}_3 = \begin{bmatrix} -\frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{bmatrix}$$

Thus, the singular value decomposition of A is

$$A = U \Sigma V^T$$

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{6}}{3} & 0 & -\frac{1}{\sqrt{3}} \\ \frac{\sqrt{6}}{6} & -\frac{\sqrt{2}}{2} & \frac{1}{\sqrt{3}} \\ \frac{\sqrt{6}}{6} & \frac{\sqrt{2}}{2} & \frac{1}{\sqrt{3}} \end{bmatrix} \begin{bmatrix} \sqrt{3} & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \end{bmatrix}$$

In SymIntegration, to compute singular value decomposition (SVD) we can use this function:
SVD(vector<vector<double>>& A)

with A as the matrix that we want to apply the SVD, SVD relies heavily on eigenvalues and eigenvectors computation for all the U, Σ, V^T matrices.


```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration SVD ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration SVD ]# ./main

A:
      1.000000      1.000000
      0.000000      1.000000
      1.000000      0.000000

A^{T}A:
      2.000000      1.000000
      1.000000      2.000000

U:
      0.816497      0.000000     -0.577350
      0.408248      0.707107      0.577350
      0.408248     -0.707107      0.577350

Σ:
      1.732051      0.000000
      0.000000      1.000000
      0.000000      0.000000

V^{T}:
      0.707107      0.707107
     -0.707107      0.707107

UEV^{T}:
      1.000000      1.000000
     -0.000000      1.000000
      1.000000      0.000000

Time taken by function: 3312 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration SVD ]#

```

Figure 6.64: The computation to obtain the singular value decomposition of matrix A . (*SymIntegration/Examples/-Linear Algebra/Test SymIntegration SVD/main.cpp*).

Chapter 7

Numerical Analysis

Je n'avais pas peur, je suis né pour faire ça - Jeanne d'Arc

Numerical analysis is the study of algorithms that use numerical approximation (as opposed to symbolic manipulations) for the problems of mathematical analysis (as distinguished from discrete mathematics). It is the study of numerical methods that attempt at finding approximate solutions of problems rather than the exact ones. Numerical analysis finds application in all fields of engineering and the physical sciences, and in the 21st century also the life and social sciences, medicine, business and even the arts.

Modern numerical analysis begins by categorizing equations as belonging to different types and then devises methods of solution for generic problems of each type.

I. SOLUTIONS OF EQUATIONS IN ONE VARIABLE (ROOT FINDING)

[SI*] **Root Finding Methods**

In this section, we consider one of the most basic problems of numerical approximation, the root-finding problem. This process involves finding a root, or solution of an equation of the form

$$f(x) = 0$$

for a given function f .

Definition 7.1: Bisection Method

The Bisection Method is based on the Intermediate Value Theorem. Suppose f is a continuous function defined on the interval $[a, b]$ with $f(a)$ and $f(b)$ of opposite sign. By the Intermediate Value Theorem, there exists a number p in (a, b) with

$$f(p) = 0$$

We assume for simplicity that the root in the interval $[a, b]$ is unique. This method calls for a repeated halving of subintervals of $[a, b]$.

- The bisection method is slow to converge, N may become quite large before $|p - p_N|$ is sufficiently small, and a good intermediate approximation can be inadvertently discarded.
- The bisection method has the important property that it always converges to a solution.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Bisection Method with Function ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Bisection Method with Function ]# ./main
f = -cos(theta)-2*theta+2*sin(theta)+1
iteration  a          b          p          f(p)
1          1          2          1.5        -0.0757472
2          1          1.5        1.25        0.0826469
3          1.25        1.5        1.375       0.0172384
4          1.375       1.5        1.4375      -0.0256436
5          1.375       1.4375     1.40625     -0.00331926
6          1.375       1.40625    1.39062     0.00717788
7          1.39062     1.40625    1.39844     0.00198421
8          1.39844     1.40625    1.40234     -0.000653763
9          1.39844     1.40234    1.40039     0.000668658
10         1.40039     1.40234    1.40137     8.30682e-06
11         1.40137     1.40234    1.40186     -0.000322513
12         1.40137     1.40186    1.40161     -0.00015705
13         1.40137     1.40161    1.40149     -7.43579e-05
Procedure completed successfully
solution = 1.40149
Time taken by function: 30471 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Bisection Method with Function ]#

```

Figure 7.1: The computation for bisection method for $f(\theta_0) = 2\theta_0 - 2\sin \theta_0 + \cos \theta_0 - 1$ with function (*SymIntegration/Examples/Numerical Methods/Test SymIntegration Bisection Method with Function for Brachistochrone/main.cpp*).

The computation is using *SymIntegration*, with this function: **bisectionmethod**(f, x, a, b, N) that can be called to compute the the solution of a function with one variable, this method doesn't need to compute the derivative so it can be used to almost all functions.

As a comparison we try to measure the time if we apply the bisection method directly in the main source code, turns out it is faster than using the function.

```

g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Bisection Method ]# ./main
f = -cos(theta)-2*theta+2*sin(theta)+1
a = 1
b = 2
p = 1.5
f(p) = -0.0757472
f(b) = -0.765258
f(a) = 0.14264
f(a)*f(p) = -0.0108046
iteration  a          b          p          f(p)
1          1          2          1.5        -0.0757472
2          1          1.5        1.25        0.0826469
3          1.25        1.5        1.375       0.0172384
4          1.375        1.5        1.4375      -0.0256436
5          1.375        1.4375     1.40625     -0.00331926
6          1.375        1.40625    1.39062     0.00717788
7          1.39062     1.40625    1.39844     0.00198421
8          1.39844     1.40625    1.40234     -0.000653763
9          1.39844     1.40234    1.40039     0.000668658
10         1.40039     1.40234    1.40137     8.30682e-06
11         1.40137     1.40234    1.40186     -0.000322513
12         1.40137     1.40186    1.40161     -0.00015705
13         1.40137     1.40161    1.40149     -7.43579e-05
Procedure completed successfully

Time taken by function: 28906 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Bisection Method ]#

```

Figure 7.2: The computation for bisection method for $f(\theta_0) = 2\theta_0 - 2\sin \theta_0 + \cos \theta_0 - 1$ without function (SymIntegration/Examples/Numerical Methods/Test SymIntegration Bisection Method Manual for Brachistochrone/main.cpp).

Definition 7.2: Newton-Raphson Method

The Newton-Raphson Method is one of the most powerful and well known numerical methods for solving a root-finding problem.

Suppose that $f \in \mathbb{C}^2[a, b]$. Let $\bar{x} \in [a, b]$ be an approximate to p such that $f'(\bar{x}) \neq 0$ and $|p - \bar{x}|$ is "small." Consider the first Taylor polynomial for $f(x)$ expanded about $(\bar{x})$,

$$f(x) = f(\bar{x}) + (x - \bar{x})f'(\bar{x}) + \frac{(x - \bar{x})^2}{2}f''(\xi(x))$$

where $\xi(x)$ lies between x and $\bar{x}$. Since $f(p) = 0$, this equation with $x = p$ gives

$$0 = f(\bar{x}) + (p - \bar{x})f'(\bar{x}) + \frac{(p - \bar{x})^2}{2}f''(\xi(p))$$

Newton's method is derived by assuming that since $|p - \bar{x}|$ is small, the term involving $(p - \bar{x})^2$ is much smaller, so

$$0 \approx f(\bar{x}) + (p - \bar{x})f'(\bar{x})$$

Solving for p gives

$$p \approx \bar{x} - \frac{f(\bar{x})}{f'(\bar{x})}$$

Thus, the Newton-Raphson method, which starts with an initial approximation p_0 and generate the sequence $\{p_n\}_{n=0}^{\infty}$, is

$$p_n = p_{n-1} - \frac{f(p_{n-1})}{f'(p_{n-1})}, \quad \text{for } n \geq 1 \quad (7.1)$$

Starting with the initial approximation p_0 , the approximation p_1 is the x -intercept of the tangent line to the graph of f at $(p_0, f(p_0))$. The approximation p_2 is the x -intercept of the tangent line to the graph of f at $(p_1, f(p_1))$ and so on.

- Newton-Raphson method is a functional iteration technique of the form

$$p_n = g(p_{n-1})$$

for which

$$g(p_{n-1}) = p_{n-1} - \frac{f(p_{n-1})}{f'(p_{n-1})}, \quad \text{for } n \geq 1 \quad (7.2)$$

this is the functional iteration technique that was used to give the rapid convergence.

- It is clear that Newton-Raphson method cannot be continued if $f'(p_{n-1}) = 0$ for some n . This method is the most effective when f' is bounded away from zero near p .

The computation is using SymIntegration, with this function: **newtonmethod**(f, x, x_0, N) that can be called to compute the the solution of a function with one variable, given that the derivative is continuous and exist.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Newton's Method with function ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Newton's Method with function ]# ./main
Newton method :
f(x) = cos(x)-x
f'(x) = -sin(x)-1

      n      p_{n}
      0      0.78539816339742
      1      0.73953613351524
      2      0.73908517810601
      3      0.73908513321516

The procedure was successful.
solution =
0.73908513321516
Time taken by function: 5645 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Newton's Method with function ]#

```

Figure 7.3: The computation for root finding Newton-Raphson method for $f(x) = \cos(x) - x$. with function (SymIntegration/Examples/Numerical Methods/Test SymIntegration Newton's Method with function/main.cpp).

As a comparison we try to implement Newton-Raphson manually, in the main source code, it turns out calling the function resulting in faster speed.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Newton's Method Manual ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Newton's Method Manual ]# ./main
f(x) = cos(x)-x
f(x)[x=pi/4] = -0.0782914

f'(x) = -sin(x)-1
f'(x)[x=pi/4] = -1.70711

      n      p_{n}      p_{n} in Degree
      0      0.78539816339742      44.999999999999
      1      0.73953613351524      42.372299247846
      2      0.73908517810601      42.346461406149
      3      0.73908513321516      42.346458834093

The procedure was successful.
Time taken by function: 6779 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Newton's Method Manual ]#

```

Figure 7.4: The manual computation for root finding Newton-Raphson method for $f(x) = \cos(x) - x$. (SymIntegration/Examples/Numerical Methods/Test SymIntegration Newton's Method Manual/main.cpp).

[SI*] Root-finding algorithms for higher order polynomial are Durand-Kerner and Abert-Ehrlich, they are iterative numerical algorithms used to simultaneously find all roots (complex or real) of a polynomial, avoiding the need for deflation techniques. Both require initial complex approximations and offer high efficiency, with Aberth generally converging faster due to its cubic convergence compared to DK' quadratic rate.

Both are efficient at finding all roots of high-degree polynomials simultaneously. But can be sensitive to initial guess (bad guesses can fail to converge) and can be slow for repeated roots.

Definition 7.3: Durand-Kerner Method

The Durand-Kerner method, also known as the Weierstrass method, this algorithm updates all root approximations simultaneously in each iteration.

Durand-Kerner algorithm steps:

1. Initialize

Given a polynomial $P(z)$ of degree n , choose n distinct initial approximations for the roots: $z_1^{(0)}, z_2^{(0)}, \dots, z_n^{(0)}$ (typically chosen as complex numbers on a circle).

2. Iteration

For each root $i \in \{1, \dots, n\}$, calculate the update formula:

$$z_i^{(k+1)} = z_i^{(k)} - \frac{P(z_i^{(k)})}{\prod_{j=1, j \neq i}^n (z_i^{(k)} - z_j^{(k)})}$$

3. Terminate

Stop when the updates are smaller than the desired tolerance (converges quadratically).

Definition 7.4: Aberth-Ehrlich Method

The Aberth method improves upon Durand-Kerner, it uses a more complex Newton-like formula incorporating derivative $P'(z)$ and a corrective shift to achieve higher efficiency.

Aberth-Ehrlich algorithm steps:

1. Initialize

Choose n distinct initial approximations for the roots: $z_1^{(0)}, z_2^{(0)}, \dots, z_n^{(0)}$.

2. Iteration

For each root $i \in \{1, \dots, n\}$, calculate the Newton step and the shift component.
Newton step:

$$N(z_i) = \frac{P(z_i)}{P'(z_i)}$$

Shift:

$$S(z_i) = \sum_{j=1, j \neq i}^n \frac{1}{z_i - z_j}$$

Update:

$$z_i^{(new)} = z_i^{(old)} - \frac{N(z_i)}{1 - N(z_i)S(z_i)}$$

3. Terminate

Stop when the updates are within tolerance (converges cubically).

i. Functions in SymIntegration Related to Root Finding

[SI*] We will list the basic functions that can be used for root finding:

AberthEhrlich(const vector<complex<double>> & $\vec{P}$, const vector<complex<double>> & x_0 , int N)

to compute the root of higher order polynomial P with Aberth-Ehrlich formula, P is a vector of size n with complex entry, the configuration is like this:

$$\vec{P} = \begin{bmatrix} (a_1, b_1) \\ (a_2, b_2) \\ \vdots \\ (a_n, b_n) \end{bmatrix}$$

$$\text{polynomial}(\vec{P}) = (a_1, b_1)r^{n-1} + (a_2, b_2)r^{n-2} + \dots + (a_n, b_n)$$

so if we want to compute a polynomial of order 6, then $\vec{P}$ has to be a size of 7. N is the number of maximum iteration for the Aberth-Ehrlich method, and x_0 is the initial guess / approximation vector with complex entry and has to be a size of $n - 1$, the entry of x_0 is preferable to be distinct and in decimal (e.g. 1.23 or 2.55), not round number to avoid error due to rounding.

This function is written in **src/numericalmethod.cpp**.

II. INTERPOLATION AND POLYNOMIAL APPROXIMATION

[SI*] **Interpolation and the Lagrange Polynomial**

Taylor polynomials are not appropriate for interpolation, because the approximations are needed only at points close to x_0 . For ordinary computational purposes it is more efficient to use methods that include information at various points. To generalize the concept of linear interpolation, consider construction of a polynomial of degree at most n that passes through $n + 1$ points

$$(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$$

to approximate that $n + 1$ points, we will use the well-established Lagrange Interpolating Polynomial.

Theorem 7.1: n th Lagrange Interpolating Polynomial

If $x_0, x_1, \dots, x_n$ are $n + 1$ distinct numbers and f is a function whose values are given at these numbers, then a unique polynomial $P(x)$ of degree at most n exists with

$$f(x_k) = P(x_k)$$

for each $k = 0, 1, \dots, n$. This polynomial is given by

$$P(x) = f(x_0)L_{n,0}(x) + \dots + f(x_n)L_{n,n}(x) = \sum_{k=0}^n f(x_k)L_{n,k}(x) \quad (7.3)$$

where, for each $k = 0, 1, \dots, n$,

$$\begin{aligned} L_{n,k}(x) &= \frac{(x - x_0)(x - x_1) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0)(x_k - x_1) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)} \\ &= \prod_{i=0, i \neq k}^n \frac{(x - x_i)}{(x_k - x_i)} \end{aligned} \quad (7.4)$$

Theorem 7.2: Approximating a Function by an Interpolating Polynomial

Suppose $x_0, x_1, \dots, x_n$ are distinct numbers in the interval $[a, b]$ and $f \in C^{n+1}[a, b]$. Then, for each x in $[a, b]$, a number $\xi(x)$ in (a, b) exists with

$$f(x) = P(x) + \frac{f^{(n+1)}(\xi(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n) \quad (7.5)$$

where $P(x)$ is the interpolating polynomial given in Eq. (7.3).

[SI*] Example:

The example is taken from example 1, chapter 3.1 of [7].

Use the numbers (or nodes) $x_0 = 2, x_1 = 2.5$, and $x_2 = 4$ to find the second interpolating polynomial for $f(x) = \frac{1}{x}$.

Solution:

We need to determine the coefficient polynomials $L_0(x)$, $L_1(x)$, and $L_2(x)$. In nested form they are

$$\begin{aligned} L_0(x) &= \frac{(x - 2.5)(x - 4)}{(2 - 2.5)(2 - 4)} = (x - 6.5)x + 10 \\ L_1(x) &= \frac{(x - 2)(x - 4)}{(2.5 - 2)(2 - 4)} = \frac{(-4x + 24)x - 32}{3} \\ L_2(x) &= \frac{(x - 2)(x - 2.5)}{(4 - 2)(4 - 2.5)} = \frac{(x - 4.5)x + 5}{3} \end{aligned}$$

Since

$$\begin{aligned} f(x_0) &= f(2) = 0.5 \\ f(x_1) &= f(2.5) = 0.4 \\ f(x_2) &= f(4) = 0.25 \end{aligned}$$

we will have

$$\begin{aligned}
 P(x) &= \sum_{k=0}^2 f(x_k) L_k(x) \\
 &= 0.5((x - 6.5)x + 10) + 0.4 \left(\frac{(-4x + 24)x - 32}{3} \right) + 0.25 \left(\frac{(x - 4.5)x + 5}{3} \right) \\
 &= (0.05x - 0.425)x + 1.15
 \end{aligned}$$

We can now check with $x = 3$

$$\begin{aligned}
 f(3) &= \frac{1}{3} \approx 0.33333 \\
 P(3) &= 0.325
 \end{aligned}$$

it is quite a good approximation.

The C++ code is using SymIntegration and Armadillo to help us compute the polynomial Lagrange $P(x)$, we can input the initial nodes $x_0, x_1, \dots, x_N$ from textfile **vectorX.txt**, and then the function $f(x)$ can be inputted in the main source code symbolically.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Lagrange Interpolating Polynomial with Armadillo ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -larmadillo -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Lagrange Interpolating Polynomial with Armadillo ]# ./main
Vector x:
 2.0000
 2.5000
 4.0000

f(x) = x^(-1)

Vector f:
 0.5000
 0.4000
 0.2500

L :
 [ x^(2)-6.5*x+10 ]
 [ -1.33333*x^(2)+8*x-10.6667 ]
 [ 0.33333*x^(2)-1.5*x+1.66667 ]

P(x) = 0.05*x^(2)-0.425*x+1.15
P(3) = 0.325
f(3) = 1/3

Time taken by function: 39599 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Lagrange Interpolating Polynomial with Armadillo ]#

```

Figure 7.5: The computation to approximate $f(x) = \frac{1}{x}$ with nodes $x_0 = 2, x_1 = 2.5$, and $x_2 = 4$ using Lagrange interpolating polynomial with SymIntegration and Armadillo (*SymIntegration/Examples/Numerical Methods/Test SymIntegration Lagrange Interpolating Polynomial with Armadillo/main.cpp*).

Another C++ code that we create is using SymIntegration only, after doing work on linear algebra we add more functions so it can have better matrix and vector operations. The result is amazing, the speed almost twice faster than when we are using Armadillo.

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Numerical Methods/Test SymIntegration Lagrange Interpolating Polynomial ]# ./main
Vector x:
2
2.5
4

f(x) = x^(-1)

Vector f:
0.5
0.4
0.25

L :
[      x^(2)-6.5*x+10      ]
[ -1.33333*x^(2)+8*x-10.6667 ]
[ 0.333333*x^(2)-1.5*x+1.66667 ]

P(x) = 0.05*x^(2)-0.425*x+1.15
P(3) = 0.325
f(3) = 1/3

Time taken by function: 28588 microseconds

```

Figure 7.6: The computation to approximate $f(x) = \frac{1}{x}$ with nodes $x_0 = 2$, $x_1 = 2.5$, and $x_2 = 4$ using Lagrange interpolating polynomial with SymIntegration only (SymIntegration/Examples/Numerical Methods/Test SymIntegration Lagrange Interpolating Polynomial/main.cpp).

III. NUMERICAL METHODS FOR OPTIMIZATION

[SI*] Minimizing a single-variable function

[SI*] The numerical methods for minimizing a single-variable function can be broadly categorized into interval reduction methods and descent methods.

Interval reduction methods, like Golden Section search, iteratively narrow down an interval known to contain the minimum, while descent methods, such as gradient descent, use derivative information to iteratively move towards the minimum.

[SI*] The interval reduction methods:

1. Golden Section Search

This method is a bracketing method that works by iteratively narrowing down the interval containing the minimum. It uses the golden ratio to determine two points within the interval, evaluates the function at these points, and then discards a portion of the interval based on the function values. This process is repeated until the interval is sufficiently small.

2. Fibonacci Search

Similar to Golden Section Search, Fibonacci Search uses the Fibonacci sequence to determine the points at which to evaluate the function. It offers optimal performance for a fixed number of function evaluations.

[SI*] The descent methods:

1. Gradient Descent

This method uses the gradient (derivative) of the function to determine the direction of steepest descent and iteratively moves towards the minimum. At each step, the algorithm moves in the opposite direction of the gradient.

2. Newton's Method

This method uses both the first and second derivatives (gradient and Hessian) to find the minimum. It offers faster convergence than gradient descent but requires the calculation of the Hessian.

Many algorithms for geometry optimization are based on some variant of the Newton-Raphson (NR) scheme. In order to find the minima, the first derivative $f'(x) = 0$ (must be zero) and the second derivative $f''(x) > 0$ (must be positive). To find the maxima, the first derivative $f'(x) = 0$ (must be zero) and the second derivative $f''(x) < 0$ (must be negative).

[SI*] The other methods:

1. **Brent's Method**

A hybrid method that combines the bracketing approach of Golden Section Search with the local quadratic approximation of Newton's method, offering a good balance of robustness and efficiency.

2. **Downhill Simplex Method (Nelder-Mead)**

A direct search method that does not require derivative information. It works by iteratively moving and transforming a simplex (a geometric shape with $n + 1$ vertices in n -dimensional space) in the search space to find the minimum.

[SI*] The best method for minimizing a single-variable function depends on the specific characteristics of the function and the desired level of accuracy. For well-behaved functions, gradient-based methods like Newton's method can offer fast convergence. For functions whose derivatives are difficult or impossible to compute, direct search methods like Brent's method or the downhill simplex method may be more suitable.

[SI*] **Gradient Descent**

Gradient Descent (GD) is an iterative first-order optimisation algorithm, used to find a local maximum/minimum of a given function. This method is commonly used in machine learning (ML) and deep learning (DL) to minimize a cost/loss function (e.g. in a linear regression). Due to its importance and ease of implementation, this algorithm is usually taught at the beginning of almost all machine learning courses. It is also widely used in areas like control engineering (robotics, chemical), computer games, mechanical engineering.

- This method was proposed long before the era of modern computers by Augustin-Louis Cauchy in 1847.
- Gradient descent algorithm does not work for all functions. There are two specific requirements. A function has to be:

1. Differentiable

Differentiable means a function has a derivative for each point in its domain. Typical non-differentiable functions have a step or a cusp or a discontinuity, such as $f(x) = \frac{1}{x}$, $f(x) = \sqrt{|x|}$

2. Convex

For a univariate function, this means that the line segment connecting two function's points lays on or above its curve (it does not cross it). If it does, then it means that it has a local minimum which is not a global one.

In mathematics notation, for two points x_1 and x_2 laying on the function's curve this condition is expressed as:

$$f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2) \quad (7.6)$$

where λ denotes a point's location on a section line and its value has to be between 0 (left point) and 1 (right point), e.g. $\lambda = 0.5$ means a location in the middle.

Another way to check mathematically if a univariate function is convex is to calculate the second derivative and check if its value is always bigger than 0.

$$\frac{d^2 f(x)}{dx^2} > 0$$

- A gradient of an n -dimensional function $f(x)$ at a given point p is defined as

$$\nabla f(p) = \begin{bmatrix} \frac{\partial f}{\partial x_1}(p) \\ \frac{\partial f}{\partial x_2}(p) \\ \vdots \\ \frac{\partial f}{\partial x_n}(p) \end{bmatrix} \quad (7.7)$$

The upside-down triangle is a notation called nabla, or you can also call it del operator.

- **Gradient Descent Algorithm**

Gradient Descent Algorithm iteratively calculates the next point using gradient at the current position, scales it (by a learning rate) and subtracts obtained value from the current position (makes a step). It subtracts the value because we want to minimize the function (to maximize it would be adding). This process can be written as:

$$p_{n+1} = p_n - \gamma \nabla f(p_n) \quad (7.8)$$

There's an important parameter γ (read: gamma) which scales the gradient and thus controls the step size. In machine learning, gamma is called the learning rate and have a strong influence on performance.

The smaller the learning rate, the longer the GD converges, or may reach maximum iteration before reaching the optimum point.

If the learning rate is too big the algorithm may not converge to the optimal point (jump around) or even diverge completely.

The Gradient Descent method's steps are:

1. Choose a starting point (initialisation).
2. Calculate gradient at this point.
3. Make a scaled step in the opposite direction to the gradient (objective: minimize).
4. Repeat steps number 2 and 3 until one of the criteria is met:
 - Maximum number of iterations reached.
 - Step size is smaller than the tolerance (due to scaling or a small gradient)

- **Example:**

Find the global minimum of a function that is given by

$$f(x) = x^2 - x + 3$$

Solution:

We will compute the first and second derivative of $f(x)$.

$$\begin{aligned} \frac{df(x)}{dx} &= 2x - 1 \\ \frac{d^2 f(x)}{dx^2} &= 2 \end{aligned}$$

Because the second derivative is always bigger than 0, $f(x)$ is strictly convex.

Now we examine the first derivative and equate it to zero.

$$\begin{aligned}\frac{df(x)}{dx} &= 0 \\ 2x - 1 &= 0 \\ x &= 0.5\end{aligned}$$

$x = 0.5$ is the critical point.

Now, we will check with the gradient descent method, first we will choose the initial value and the parameter γ (the learning rate):

$$x_{old} = 2, \quad \gamma = 0.1$$

then we will have

$$\begin{aligned}x_{new} &= x_{old} - \gamma \nabla f(x_{old}) \\ x_{new} &= 2 - 0.1(2(2) - 1) = 1.7 \quad (i = 1) \\ x_{new} &= 1.7 - 0.1(2(1.7) - 1) = 1.46 \quad (i = 2) \\ x_{new} &= 1.46 - 0.1(2(1.46) - 1) = 1.268 \quad (i = 3) \\ &\vdots \\ x_{new} &= 0.500027 - 0.1(2(0.500027) - 1) = 0.500021 \quad (i = 50)\end{aligned}$$

It is proven then with the gradient descent method that the critical point is at $x = 0.5$. The computation can be done with **SymIntegration**, as long as the function $f(x)$ is differentiable and the derivative can be computed then we can use gradient descent method. Polynomials without the combination of trigonometry and transcendentals functions can be differentiated easily, even the fraction polynomial.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gradient Descent
]# make
g++ -c -o main.o main.cpp
g++ -o main -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gradient Descent
]# ./main
F(x) = x^(2)-x+3
i = 1
xnew = 1.7
f(xnew) = 4.19
i = 2
xnew = 1.46
f(xnew) = 3.6716
i = 3
xnew = 1.268
f(xnew) = 3.33982
i = 4
xnew = 1.1144
f(xnew) = 3.12749
i = 5
xnew = 0.99152
f(xnew) = 2.99159
i = 6
xnew = 0.893216
f(xnew) = 2.90462
i = 7
xnew = 0.814573
f(xnew) = 2.84896
```

Figure 7.7: The computation for minimization problem for function $f(x) = x^2 - x + 3$ with gradient descent method. (**SymIntegration/Examples/Operations Research/Test SymIntegration Gradient Descent/main.cpp**).

```

i = 45
xnew = 0.500065
f(xnew) = 2.75
i = 46
xnew = 0.500052
f(xnew) = 2.75
i = 47
xnew = 0.500042
f(xnew) = 2.75
i = 48
xnew = 0.500033
f(xnew) = 2.75
i = 49
xnew = 0.500027
f(xnew) = 2.75
i = 50
xnew = 0.500021
f(xnew) = 2.75
The gradient descent converges.
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gradient Descent ]# 

```

Figure 7.8: The gradient descent method took 50 iteration to converges with $\gamma = 0.1$. (*SymIntegration/Examples/Operations Research/Test SymIntegration Gradient Descent/main.cpp*).

- **Example:**

Find the global minimum of a function that is given by

$$f(x, y) = \frac{1}{2}x^2 + 2x + y^2 + y + 3$$

Solution:

In the case of a multivariate function, we need to compute the vector of derivatives in each main direction (along variable axes). Because we are interested only in a slope along one axis and we don't care about others, these derivatives are called partial derivatives.

We will have

$$\begin{aligned}
 z = f(x, y) &= \frac{1}{2}x^2 + 2x + y^2 + y + 3 \\
 \frac{\partial f}{\partial x} &= x + 2 \\
 \frac{\partial f}{\partial y} &= 2y + 1
 \end{aligned}$$

then the nabla is

$$\nabla f = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{bmatrix} = \begin{bmatrix} x + 2 \\ 2y + 1 \end{bmatrix}$$

at initialization we will input the initial value and the parameter γ

$$\begin{aligned}
 x_{old} &= -5 \\
 y_{old} &= 21 \\
 \gamma &= 0.1
 \end{aligned}$$

thus the gradient descent method

$$\begin{aligned}
 x_{new} &= x_{old} - \gamma \nabla f(x_{old}) \\
 x_{new} &= -5 - 0.1((-5) + 2) = -4.7 \quad (i = 1) \\
 x_{new} &= -4.7 - 0.1((-4.7) + 2) = -4.43 \quad (i = 2) \\
 x_{new} &= -4.43 - 0.1((-4.43) + 2) = -4.187 \quad (i = 3) \\
 &\vdots \\
 x_{new} &= -2.01718 - 0.1((-2.01718) + 2) = -2.01546 \quad (i = 50) \\
 y_{new} &= y_{old} - \gamma \nabla f(y_{old}) \\
 y_{new} &= 21 - 0.1(2(21) + 1) = 16.7 \quad (i = 1) \\
 y_{new} &= 16.7 - 0.1(2(16.7) + 1) = 13.26 \quad (i = 2) \\
 y_{new} &= 13.26 - 0.1(2(13.26) + 1) = 10.508 \quad (i = 3) \\
 &\vdots \\
 y_{new} &= -0.499616 - 0.1(2(-0.499616) + 1) = -0.499693 \quad (i = 50)
 \end{aligned}$$

and the value for function $f(x, y)$ at every iteration

$$\begin{aligned}
 f(x_{new}, y_{new}) &= 300.235 \quad (i = 1) \\
 f(x_{new}, y_{new}) &= 193.04 \quad (i = 2) \\
 f(x_{new}, y_{new}) &= 124.318 \quad (i = 3) \\
 &\vdots \\
 f(x_{new}, y_{new}) &= 0.75012 \quad (i = 50)
 \end{aligned}$$

so the function $f(x, y) = \frac{1}{2}x^2 + 2x + y^2 + y + 3$ reaches the minimum of 0.75012 at $(-2.01546, -0.499693)$.

```

xterm
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gradient Descent 2 Variables ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gradient Descent 2 Variables ]# ./main
F(x, y) = 0.5*x^(2)+2*x+y^(2)+y+3
i = 1
xnew = -4.7
ynew = 16.7
F(xnew, ynew) = 300.235
i = 2
xnew = -4.43
ynew = 13.26
F(xnew, ynew) = 193.04
i = 3
xnew = -4.187
ynew = 10.508
F(xnew, ynew) = 124.318
i = 4
xnew = -3.9683
ynew = 8.3064
F(xnew, ynew) = 80.2398
i = 5
xnew = -3.77147
ynew = 6.54512
F(xnew, ynew) = 51.9528
i = 6
xnew = -3.59432
ynew = 5.1361
F(xnew, ynew) = 33.7865
i = 7
xnew = -3.43489
ynew = 4.00888
F(xnew, ynew) = 22.1094

```

Figure 7.9: The computation for minimization problem for function $f(x, y) = \frac{1}{2}x^2 + 2x + y^2 + y + 3$ with gradient descent method. (SymIntegration/Examples/Operations Research/Test SymIntegration Gradient Descent 2 Variables/main.cpp).

```

i = 45
xnew = -2.02618
ynew = -0.499064
F(xnew,ynew) = 0.750344
i = 46
xnew = -2.02357
ynew = -0.499251
F(xnew,ynew) = 0.750278
i = 47
xnew = -2.02121
ynew = -0.499401
F(xnew,ynew) = 0.750225
i = 48
xnew = -2.01909
ynew = -0.499521
F(xnew,ynew) = 0.750182
i = 49
xnew = -2.01718
ynew = -0.499616
F(xnew,ynew) = 0.750148
i = 50
xnew = -2.01546
ynew = -0.499693
F(xnew,ynew) = 0.75012
Maximum number of iteration reached.
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Gradient Descent 2 Variables ]# 

```

Figure 7.10: The gradient descent method took 50 iteration to converges with $\gamma = 0.1$. (*SymIntegration/Examples/Operations Research/Test SymIntegration Gradient Descent 2 Variables/main.cpp*).

[SI*] Downhill Simplex

The downhill simplex, introduced by Nelder and Mead, also known as the Nelder-Mead' method [20], it is a method for solving multidimensional, unconstrained, numerical optimization problems. This type of problem is one of the most common in engineering and science, and this method is one of the simplest and fairly effective to solve the problem.

[SI*] We define a minimization problem as

$$\arg \min f(x) \quad (7.9)$$

$$x \in \mathbb{R}^N, f: \mathbb{R}^N \mapsto \mathbb{R}^1$$

[SI*] Method and simplex transforms

A simplex is a collection of connected points in $\mathbb{R}^N$. We call it a k -simplex where k is the degree that indicates how many points it contains. A k -simplex contains $k + 1$ points, so a triangle which has 3 points is a 2-simplex, a tetrahedron which has 4 points is a 3-simplex and so on.

When optimizing a function with N parameters we will use a simplex with $N + 1$ points. Meaning the degree k of the simplex is the same as the dimension of the parameter space. So for a problem with 2 parameters, $N = 2$, we use a 2-simplex, a triangle, that has three points. A 1 parameter problem would use a line segment in 1D and a 3 parameter problem would use a tetrahedron in 3D.

Each point on the simplex, denoted by $x_i, 0 \leq i \leq N$, corresponds to a complete set of parameters and therefore to a value of the objective function $f_i = f(x_i)$.

The core of the method is to keep the points sorted so that $f(x_0)$ is the current lowest value, i.e. the best point, and x_N is the worst. We then calculate the center of mass of all points except x_N :

$$c = \frac{1}{N} \sum_{i=0}^{N-1} x_i \quad (7.10)$$

Afterwards we attempt to replace the point x_N with one along the line formed by $\overrightarrow{x_N c}$ according to a set of transformations called reflection, expansion, and contraction. A

shrink transform that moves all points closer to the current best can also be performed. We associate each transform with a parameter, α for reflection, β for contraction, γ for expansion and δ for shrink, that should satisfy the following constraints:

- $\alpha > 0$
- $0 < \beta < 1$
- $\gamma > \alpha, \gamma > 1$
- $0 < \delta < 1$

[SI*] **Reflection**

The following illustrations are a 2D problem with a triangle simplex (can be used for function with two independent variable such as $f(x, y)$). The point x_0 with blue color is the current best, the point x_1 with green color is the second best, and the point x_2 with light red color is the third best / the one being transformed except for when the simplex performs a shrink and all but the best are changed. The blue lines indicate the current simplex and the light red lines are the new one.

For every iteration the reflected point x_r is the one tried first

$$x_r = c + \alpha(c - x_2) \quad (7.11)$$

If this point is better than the second best but not better than the current best such that

$$f(x_0) \leq f(x_r) < f(x_1)$$

we replace x_2 with x_r , otherwise we do either expansion, contraction or shrink.




Figure 7.11: x_2 is mirrored in c and creates a reflected point x_r .

[SI*] **Expansion**

If x_r is better than the current best,

$$f(x_r) < f(x_0)$$

we continue to explore and compute the expansion point x_e

$$x_e = c + \gamma(x_r - c) \quad (7.12)$$




Figure 7.12: The expansion point x_e is computed by moving further away from c .

Then one of two approaches can be used:

- Greedy minimization
in which the reflected and expanded points are compared and the better of the two is picked such that if

$$f(x_e) < f(x_r) < f(x_0)$$

we accept x_e ($x_2 = x_e$) and terminate the iteration.

Otherwise, if

$$f(x_r) \leq f(x_e)$$

then x_r is accepted ($x_2 = x_r$).

- Greedy expansion
we will accept x_e if

$$f(x_e) < f(x_1)$$

and

$$f(x_r) < f(x_0)$$

and skip comparing $f(x_e)$ to $f(x_r)$. This will keep the simplex as large as possible which can have some advantages for non-smooth functions.

[SI*] Contraction

If x_r is worse than the current second worst point, in 2D case it is x_1 , which means

$$f(x_r) \geq f(x_1)$$

we compute a contraction point using x_2 and x_r .

It will be

- Outside
if

$$f(x_1) \leq f(x_r) < f(x_2)$$

We compute the contraction point as

$$x_c^{out} = c + \beta(x_r - c) \tag{7.13}$$

and if

$$f(x_c^{out}) \leq f(x_r)$$

then x_c^{out} is accepted ($x_2 = x_c^{out}$). Otherwise, a shrink of the simplex is performed.




Figure 7.13: The contraction point x_c^{out} is computed outside the old simplex.

- Inside
if

$$f(x_r) \geq f(x_2)$$

We compute the contraction point as

$$x_c^{in} = c + \beta(x_2 - c) \quad (7.14)$$

and if

$$f(x_c^{in}) \leq f(x_2)$$

then x_c^{in} is accepted ($x_2 = x_c^{in}$). Otherwise, a shrink of the simplex is performed.




Figure 7.14: The contraction point x_c^{in} is computed inside the old simplex.

[SI*] Shrink

the last resort that will be performed if none of the criteria so far have been met is to shrink the entire simplex by moving all points closer to the current best points.

$$s_{i-1} = x_0 + \delta(x_i - x_0), \quad i = \{1, 2, \dots, N\} \quad (7.15)$$




Figure 7.15: *Shrink the simplex by moving every point closer to the current best.*

[SI*] Initialization and termination

When initializing the simplex, usually one point x_0 is to be given by the user (as the initial value / input), then the other points are created by placing them at certain distance μ from the initial point along the coordinate axes in the parameter space $\mathbb{R}^N$ of the objective function.

$$x_i = x_0 + \mu \hat{e}_i, \quad i = \{1, 2, \dots, N\} \quad (7.16)$$

The algorithm terminates in one of three different ways. Either by having the simplex become sufficiently small, such that the active search space is thought to not have any considerable variation in function value. Another criteria is for all the function values of the points of the simplex to have a similar enough value. And finally the algorithm must be set to run for a maximum number of iterations or function evaluations.




Figure 7.16: The flowchart for downhill simplex algorithm that is used to make the C++ code for computing the solution.

[SI*] The downhill simplex method has some pros and cons. Its major strength is that it does not require the gradient / derivative computation of f . the gradient of the objective function is often not known as an analytical expression and can therefore be very expensive to evaluate. If so it has to be done using some form of finite difference scheme. This computational cost increases with the dimensionality of the problem and is also subject to potential numerical stability problems if some care is not taken. Since downhill simplex does not require to compute the gradient of f , this makes this method applicable to non-smooth functions where the gradient may not exist on the whole domain.

This method has a memory footprint on the order of about $(N + 2) \times N$ since we always

have to store $N + 2$ sets of N parameters. Since we don't have to compute any gradient, then this method converges slowly compared to gradient based methods, generally because it requires more function evaluations.

[SI*] **Example:**

Find the value x and y that minimizes

$$f(\mathbf{x}) = f(x, y) = e^{-\frac{x^2+y^2}{2s^2}} - e^{-\frac{x^2+y^2}{2t^2}} + \frac{1}{10}x^2 + \frac{1}{10}y^2, \quad s = 0.75, t = 1 \quad (7.17)$$

Solution:

First, we choose the initial point:

$$\mathbf{x}_0 = \begin{bmatrix} 0.1 \\ 0.5 \end{bmatrix}$$

The other two points can be generated from $\mathbf{x}_0$ by placing a certain distance $\mu = 0.05$ that we choose, thus

$$\mathbf{x}_1 = \mathbf{x}_0 + \mu \hat{\mathbf{e}}_1 = \begin{bmatrix} 0.1 \\ 0.5 \end{bmatrix} + 0.05 \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0.15 \\ 0.5 \end{bmatrix}$$

$$\mathbf{x}_2 = \mathbf{x}_0 + \mu \hat{\mathbf{e}}_2 = \begin{bmatrix} 0.1 \\ 0.5 \end{bmatrix} + 0.05 \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0.1 \\ 0.55 \end{bmatrix}$$

We will have

$$f(\mathbf{x}_0) = -0.058442$$

$$f(\mathbf{x}_1) = -0.0604927$$

$$f(\mathbf{x}_2) = -0.066630$$

(It is unsorted, but we try to compute with unsorted vector first) From here we can compute $\mathbf{c}$

$$\mathbf{c} = \frac{1}{2} \sum_{i=0}^2 \mathbf{x}_i = \begin{bmatrix} 0.125 \\ 0.5 \end{bmatrix}$$

Now, we need to input the parameters as

$$\alpha = 1, \quad \beta = 0.5, \quad \gamma = 2, \quad \delta = 0.5$$

Then we will perform the reflection operation first,

$$\mathbf{x}_r = \mathbf{c} + \alpha(\mathbf{c} - \mathbf{x}_2) = \begin{bmatrix} 0.15 \\ 0.45 \end{bmatrix}$$

From here we will compare the value of $f(\mathbf{x}_0), f(\mathbf{x}_1), f(\mathbf{x}_2)$ and $f(\mathbf{x}_r)$.

If $f(\mathbf{x}_0) < f(\mathbf{x}_r) < f(\mathbf{x}_1)$ then

$$\mathbf{x}_2 = \mathbf{x}_r$$

and we go to iteration 2.

If $f(\mathbf{x}_r) < f(\mathbf{x}_0)$, then go to the expansion step.

If $f(x_r) \geq f(x_1)$, then go to the contraction step.

In this case we have

$$\begin{aligned} f(x_0) &= -0.058442 \\ f(x_1) &= -0.0604927 \\ f(x_2) &= -0.0666302 \\ f(x_r) &= -0.0523666 \end{aligned}$$

$f(x_r) \geq f(x_1)$ so we go and compute the contraction point, since $f(x_r) \geq f(x_2)$ we compute the inside contraction point

$$x_c^{in} = c + \beta(x_0 - c) = \begin{bmatrix} 0.1125 \\ 0.5 \end{bmatrix}$$

then we will have

$$\begin{aligned} f(x_0) &= -0.058442 \\ f(x_1) &= -0.0604927 \\ f(x_2) &= -0.0666302 \\ f(x_c) &= -0.0588848 \end{aligned}$$

since $f(x_c) > f(x_2)$ then we perform shrink that will replace x_1 with s_0 and x_2 with s_1

$$s_0 = x_0 + \delta(x_1 - x_0) = \begin{bmatrix} 0.1250 \\ 0.5250 \end{bmatrix}$$

$$s_1 = x_0 + \delta(x_2 - x_0) = \begin{bmatrix} 0.1375 \\ 0.5625 \end{bmatrix}$$

So we have reached the end of the first iteration, which is using the shrink and we get these values at the end of this iteration:

$$\begin{aligned} f(x_0) &= -0.058442 \\ f(x_1) &= -0.0604927 \\ f(x_2) &= -0.0666302 \end{aligned}$$

Continue till the last number of iteration. We use 6 iterations and we use C++ to help us in computing, thus we obtain the best vector x_0 that minimizes the function $f(x, y)$ is

$$x_0 = \begin{bmatrix} 0.1916 \\ 0.5 \end{bmatrix}$$

with $f(x_0) = -0.063016$.

Since it is only using 6 iterations, thus the real optimal solution can be different if we add more iterations, we also sort the vectors x_0, x_1, x_2 at the end of iteration 1, if it is moved before iteration 1 then we will get faster convergence. We try the code with sorted vector before iteration 1 and obtain $f(x_0) = -0.086404$ at the end of iteration 6.

We create the computation with **SymIntegration** and **Armadillo** and another one is with **SymIntegration** and **Eigen** for comparison, both produce the same answer but the one with **Eigen** has faster compiling time, we use 6 iterations with unsorted initial vectors then we sort the initial vectors and use 37 iterations, and then for the plotting in 3D we use Hamzstlab Mathematics with backend **ImGui** and **ImPlot3D** so we can change the value of s and t and the surface plot will change interactively as well.

```

Test Downhill Simplex with Armadillo
View Go Help
/root/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Armadillo/
you are using the root account. You may harm your system.
ctop
main main.cpp main.o Makefile
System
xterm
root [ ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Armadillo ]# time make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -larmadillo -lsymintegration

real    0m26.847s
user    0m25.979s
sys      0m0.868s
root [ ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Armadillo ]# ./main
f(x,y) = e^(-0.888889*x^2-0.888889*y^2)-e^(-0.5*x^2-0.5*y^2)+0.1*x^2+0.1*y^2

Total iteration:      6
Reflection:          0
Expansion:           3
Outside contraction:  0
Inside contraction:  1
Shrink:              2

x_{0}:
  0.1961
  0.5000

x_{1}:
  0.1719
  0.5000

x_{2}:
  0.2301
  0.5000

f(x,y) from x_{0}= -0.0630159990145
f(x,y) from x_{1}= -0.0616189114776
f(x,y) from x_{2}= -0.0610391545891
root [ ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Armadillo ]#

```

Figure 7.17: The computation of downhill simplex for $f(x,y) = e^{-\frac{x^2+y^2}{2s^2}} - e^{-\frac{x^2+y^2}{2t^2}} + \frac{1}{10}x^2 + \frac{1}{10}y^2$, $s = 0.75$, $t = 1$ with **SymIntegration** and **Armadillo** (**SymIntegration/Examples/Operations Research/Test Downhill Simplex with Armadillo/main.cpp**).

```

root@root: /root/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Eigen/
root account. You may harm your system.

main main.cpp main.o Makefile

root@root: ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Eigen ]# time make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration

real    0m19.034s
user    0m18.285s
sys      0m0.748s
root@root: ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Eigen ]# ./main
f(x,y) = e^(-0.888889*x^(2)-0.888889*y^(2))-e^(-0.5*x^(2)-0.5*y^(2))+0.1*x^(2)+0.1*y^(2)

Total iteration:      6
Reflection:           0
Expansion:             3
Outside contraction:  0
Inside contraction:   1
Shrink:                2

x_{0}:
0.196094
0.5
x_{1}:
0.171875
0.5
x_{2}:
0.230078
0.5

f(x,y) from x_{0}= -0.0630159990145
f(x,y) from x_{1}= -0.0616189114776
f(x,y) from x_{2}= -0.0610391545891
root@root: ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Eigen ]#

```

Figure 7.18: The computation of downhill simplex for $f(x,y) = e^{-\frac{x^2+y^2}{2s^2}} - e^{-\frac{x^2+y^2}{2t^2}} + \frac{1}{10}x^2 + \frac{1}{10}y^2$, $s = 0.75$, $t = 1$ with SymIntegration and Eigen (SymIntegration/Examples/Operations Research/Test Downhill Simplex with Eigen/main.cpp).

```

g++ -o main -gdb main.o -lstdc++ -larmadillo -lsymintegration

real    0m29.108s
user    0m28.112s
sys      0m0.893s
root [ ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Armadillo ]# ./main
Vector x_{0}:
  0.1000
  0.5000

Vector x_{1}:
  0.1500
  0.5000

Vector x_{2}:
  0.1000
  0.5000

After sorting,
x_{0}:
  0.1000
  0.5000

x_{1}:
  0.1500
  0.5000

x_{2}:
  0.1000
  0.5000

f(x,y) = e^(-0.888889*x^(2)-0.888889*y^(2))-e^(-0.5*x^(2)-0.5*y^(2))+0.1*x^(2)+0.1*y^(2)

Total iteration:    37
Reflection:         5
Expansion:          14
Outside contraction: 0
Inside contraction: 18
Shrink:             0

x_{0}:
  0.0973
  0.8914

x_{1}:
  0.0852
  0.8891

x_{2}:
  0.0629
  0.8977

f(x,y) from x_{0}= -0.0992255745363
f(x,y) from x_{1}= -0.0992180473087
f(x,y) from x_{2}= -0.0992086382243
root [ ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Armadillo ]# 

```

Figure 7.19: The computation of downhill simplex for $f(x,y) = e^{-\frac{x^2+y^2}{2s^2}} - e^{-\frac{x^2+y^2}{2t^2}} + \frac{1}{10}x^2 + \frac{1}{10}y^2$, $s = 0.75$, $t = 1$ with 37 iterations (SymIntegration/Examples/Operations Research/Test Downhill Simplex with Armadillo/main.cpp).

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Eigen ]#
time make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration

real    0m19.921s
user    0m19.108s
sys     0m0.760s
root [ ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Eigen ]# ./main
Vector x_{0}:
0.1
0.5
Vector x_{1}:
0.15
0.5
Vector x_{2}:
0.1
0.55
After sorting,
x_{0}:
0.1
0.55
x_{1}:
0.15
0.5
x_{2}:
0.1
0.5
f(x,y) = e^{(-0.888889*x^2-0.888889*y^2)}-e^{(-0.5*x^2-0.5*y^2)}+0.1*x^2+0.1*y^2

Total iteration:      37
Reflection:           5
Expansion:            14
Outside contraction:  0
Inside contraction:   18
Shrink:               0

x_{0}:
0.0972656
0.091406
x_{1}:
0.0051562
0.089063
x_{2}:
0.0628906
0.097656

f(x,y) from x_{0}= -0.0992255745363
f(x,y) from x_{1}= -0.0992180473087
f(x,y) from x_{2}= -0.0992086382243
root [ ~/SourceCodes/CPP/C++ Create Library/Test Downhill Simplex with Eigen ]#

```

Figure 7.20: The computation of downhill simplex for $f(x,y) = e^{-\frac{x^2+y^2}{2s^2}} - e^{-\frac{x^2+y^2}{2t^2}} + \frac{1}{10}x^2 + \frac{1}{10}y^2$, $s = 0.75$, $t = 1$ with 37 iterations (*SymIntegration/Examples/Operations Research/Test Downhill Simplex with Eigen/main.cpp*).

* the C++ codes are already updated so the initial vectors will be sorted before the **for** looping.

[SI*] Global Optimization

Global optimization is used for finding all global minima or maxima of a given function on a given domain disregarding any local optima. Global optimization is a large field with many applications, and many deterministic and stochastic optimization methods have been developed.

Deterministic methods include:

1. Branch-and-bound methods
2. Cutting-plane methods
3. Inner-and-outer approximation
4. Interval methods

Stochastic methods, on the other hand, are methods whose results depend on random numbers drawn while running the algorithm; the function to be optimized is still deterministic. Stochastic methods include:

1. Direct Monte Carlo sampling
2. Stochastic tunneling
3. Parallel tempering

Heuristic methods are strategies for searching the domain in an intelligent manner.

They include:

1. Differential evolution
2. Evolutionary computation (including, e.g., evolution strategies, genetic algorithms, genetic programming)
3. Graduated optimization
4. Simulated annealing
5. Swarm based optimization (including, e.g., particle-swarm optimization and ant colony optimization)
6. Taboo search

Other methods are Bayesian optimization and memetic algorithms.

IV. NUMERICAL DIFFERENTIATION

[SI*] The derivative of the function f at x_0 is

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} \quad (7.18)$$

This formula gives an obvious way to generate an approximation to $f'(x)$; simply compute

$$\frac{f(x_0 + h) - f(x_0)}{h}$$

for small values of h . Although this may be obvious, it is not very successful, due to roundoff error.

To approximate $f'(x_0)$, suppose first that $x_0 \in (a, b)$ where $f \in C^2[a, b]$, and that $x_1 = x_0 + h$ for some $h \neq 0$ that is sufficiently small to ensure that $x_1 \in [a, b]$. We construct the first Lagrange polynomial $P_{0,1}(x)$ for f determined by x_0 and x_1 , with its error term:

$$\begin{aligned} f(x) &= P_{0,1}(x) + \frac{(x - x_0)(x - x_1)}{2!} f''(\xi(x)) \\ &= \frac{f(x_0)(x - x_0 - h)}{-h} + \frac{f(x_0 + h)(x - x_0)}{h} + \frac{(x - x_0)(x - x_0 - h)}{2} f''(\xi(x)) \end{aligned}$$

for some $\xi(x)$ in $[a, b]$. Differentiating gives

$$\begin{aligned} f'(x) &= \frac{f(x_0 + h) - f(x_0)}{h} + D_x \left[\frac{(x - x_0)(x - x_0 - h)}{2} f''(\xi(x)) \right] \\ &= \frac{f(x_0 + h) - f(x_0)}{h} + \frac{2(x - x_0) - h}{2} f''(\xi(x)) + \frac{(x - x_0)(x - x_0 - h)}{2} D_x(f''(\xi(x))) \end{aligned}$$

so

$$f'(x) \approx \frac{f(x_0 + h) - f(x_0)}{h}$$

One difficulty with this formula is that we have no information about $D_x f''(\xi(x))$, so the truncation error cannot be estimated. When x is x_0 , however, the coefficient of $D_x f''(\xi(x))$ is 0, and the formula simplifies to

$$f'(x_0) = \frac{f(x_0 + h) - f(x_0)}{h} - \frac{h}{2} f''(\xi) \quad (7.19)$$

for small values of h , the difference quotient $\frac{f(x_0+h)-f(x_0)}{h}$ can be used to approximate $f'(x_0)$ with an error bounded by $M\frac{|h|}{2}$, where M is a bound on $|f''(\xi)|$ for $x \in [a, b]$. This formula is known as the forward-difference formula if $h > 0$ and the backward-difference formula if $h < 0$.

[SI*] To obtain general derivative approximation formulas, suppose that $\{x_0, x_1, \dots, x_n\}$ are $(n+1)$ distinct numbers in some interval I and that $f \in C^{n+1}(I)$. From Theorem 11.2 we will have,

$$f(x) = \sum_{k=0}^n f(x_k) L_k(x) + \frac{(x-x_0) \dots (x-x_n)}{(n+1)!} f^{(n+1)}(\xi(x))$$

for some $\xi(x)$ in I , where $L_k(x)$ denotes the k th Lagrange coefficient polynomial for f at $x_0, x_1, \dots, x_n$. Differentiating this expression gives

$$\begin{aligned} f'(x) &= \sum_{k=0}^n f(x_k) L'_k(x) + D_x \left[\frac{(x-x_0) \dots (x-x_n)}{(n+1)!} \right] f^{(n+1)}(\xi(x)) \\ &\quad + \frac{(x-x_0) \dots (x-x_n)}{(n+1)!} D_x [f^{(n+1)}(\xi(x))] \end{aligned}$$

Again, we have a problem estimating the truncation error unless x is one of the numbers x_j . In this case, the term multiplying $D_x [f^{(n+1)}(\xi(x))]$ is 0, and the formula becomes

$$f'(x_j) = \sum_{k=0}^n f(x_k) L'_k(x_j) + \frac{f^{(n+1)}(\xi(x_j))}{(n+1)!} \prod_{k=0, k \neq j}^n (x_j - x_k) \quad (7.20)$$

which is called an $(n+1)$ -point formula to approximate $f'(x_j)$.

In general, using more evaluation points in Eq. (7.20) produces greater accuracy, although the number of functional evaluations and growth of roundoff error discourages this somewhat. The most common formulas are those involving three and five evaluation points.

We first derive some useful three-point formulas and consider aspects of their errors. Since

$$L_0(x) = \frac{(x-x_1)(x-x_2)}{(x_0-x_1)(x_0-x_2)}$$

we have

$$L'_0(x) = \frac{2x - x_1 - x_2}{(x_0 - x_1)(x_0 - x_2)}$$

Similarly,

$$L'_1(x) = \frac{2x - x_0 - x_2}{(x_1 - x_0)(x_1 - x_2)}$$

$$L'_2(x) = \frac{2x - x_0 - x_1}{(x_2 - x_0)(x_2 - x_1)}$$

Hence, from Eq. (7.20),

$$\begin{aligned} f'(x_j) &= f(x_0) \left[\frac{2x_j - x_1 - x_2}{(x_0 - x_1)(x_0 - x_2)} \right] + f(x_1) \left[\frac{2x_j - x_0 - x_2}{(x_1 - x_0)(x_1 - x_2)} \right] \\ &\quad + f(x_2) \left[\frac{2x_j - x_0 - x_1}{(x_2 - x_0)(x_2 - x_1)} \right] + \frac{1}{6} f^{(3)}(\xi_j) \prod_{k=0, k \neq j}^2 (x_j - x_k) \end{aligned} \quad (7.21)$$

for each $j = 0, 1, 2$, where the notation ξ_j indicates that this point depends on x_j .

The three formulas from Eq. (7.21) become especially useful if the nodes are equally spaced, that is, when

$$\begin{aligned}x_1 &= x_0 + h \\x_2 &= x_0 + 2h\end{aligned}$$

for some $h \neq 0$.

Using Eq. (7.21) with $x_j = x_0$, $x_1 = x_0 + h$, and $x_2 = x_0 + 2h$ gives

$$f'(x_0) = \frac{1}{h} \left[-\frac{3}{2}f(x_0) + 2f(x_1) - \frac{1}{2}f(x_2) \right] + \frac{h^2}{3}f^{(3)}(\xi_0)$$

Doing the same for $x_j = x_1$ gives

$$f'(x_1) = \frac{1}{h} \left[-\frac{1}{2}f(x_0) + \frac{1}{2}f(x_2) \right] - \frac{h^2}{6}f^{(3)}(\xi_1)$$

and for $x_j = x_2$,

$$f'(x_2) = \frac{1}{h} \left[\frac{1}{2}f(x_0) - 2f(x_1) + \frac{3}{2}f(x_2) \right] + \frac{h^2}{3}f^{(3)}(\xi_2)$$

Since $x_1 = x_0 + h$ and $x_2 = x_0 + 2h$, these formulas can also be expressed as

$$\begin{aligned}f'(x_0) &= \frac{1}{h} \left[-\frac{3}{2}f(x_0) + 2f(x_0 + h) - \frac{1}{2}f(x_0 + 2h) \right] + \frac{h^2}{3}f^{(3)}(\xi_0) \\f'(x_0 + h) &= \frac{1}{h} \left[-\frac{1}{2}f(x_0) + \frac{1}{2}f(x_0 + 2h) \right] - \frac{h^2}{6}f^{(3)}(\xi_1) \\f'(x_0 + 2h) &= \frac{1}{h} \left[\frac{1}{2}f(x_0) - 2f(x_0 + h) + \frac{3}{2}f(x_0 + 2h) \right] + \frac{h^2}{3}f^{(3)}(\xi_2)\end{aligned}$$

As a matter of convenience, the variable substitution x_0 for $x_0 + h$ is used in the middle equation to change this formula to an approximation for $f'(x_0)$. A similar change, x_0 for $x_0 + 2h$, is used in the last equation. This gives three formulas for approximating $f'(x_0)$:

$$\begin{aligned}f'(x_0) &= \frac{1}{2h} [-3f(x_0) + 4f(x_0 + h) - f(x_0 + 2h)] + \frac{h^2}{3}f^{(3)}(\xi_0) \\f'(x_0) &= \frac{1}{2h} [-f(x_0 - h) + f(x_0 + h)] - \frac{h^2}{6}f^{(3)}(\xi_1) \\f'(x_0) &= \frac{1}{2h} [f(x_0 - 2h) - 4f(x_0 - h) + 3f(x_0)] + \frac{h^2}{3}f^{(3)}(\xi_2)\end{aligned}$$

Finally, note that since the last of these equations can be obtained from the first by simply replacing h with $-h$, there are actually only two formulas:

$$f'(x_0) = \frac{1}{2h} [-3f(x_0) + 4f(x_0 + h) - f(x_0 + 2h)] + \frac{h^2}{3}f^{(3)}(\xi_0) \quad (7.22)$$

where ξ_0 lies between x_0 and $x_0 + 2h$, and

$$f'(x_0) = \frac{1}{2h} [f(x_0 + h) - f(x_0 - h)] - \frac{h^2}{6} f^{(3)}(\xi_1) \quad (7.23)$$

where ξ_1 lies between $(x_0 - h)$ and $(x_0 + h)$.

Although the error in both Eqs. (7.22) and (7.23) are $O(h^2)$, the error in Eq. (7.23) are approximately half the error in Eq. (7.22). This is because Eq. (7.23) uses data on both sides of x_0 and Eq. (7.22) uses data on only one side. Note also that f needs to be evaluated at only two points in Eq. (7.23), whereas in Eq. (7.22) three evaluations are needed.

Similarly, the five-point formulas that involve evaluating the function at two more points is

$$f'(x_0) = \frac{1}{12h} [f(x_0 - 2h) - 8f(x_0 - h) + 8f(x_0 + h) - f(x_0 + 2h)] + \frac{h^4}{30} f^{(5)}(\xi) \quad (7.24)$$

where ξ lies between $x_0 - 2h$ and $x_0 + 2h$.

The other five-point formula is useful for end-point approximations, particularly with regard to the clamped cubic spline interpolation. It is

$$\begin{aligned} f'(x_0) = & \frac{1}{12h} [-25f(x_0) + 48f(x_0 + h) - 36f(x_0 + 2h) \\ & + 16f(x_0 + 3h) - 3f(x_0 + 4h)] + \frac{h^4}{5} f^{(5)}(\xi) \end{aligned} \quad (7.25)$$

where ξ lies between x_0 and $x_0 + 4h$. Left-endpoint approximations are found using this formula with $h > 0$ and right-endpoint approximations with $h < 0$.

[SI*] When we use three-point formula that use both sides of point, the optimal choice for h theoretically can be found when we find the minimum of function $e(h)$ (the roundoff errors) and solve it for h [7]

$$e(h) = \frac{\epsilon}{h} + \frac{h^2}{6} M \quad (7.26)$$

where

$$M = \max_{x \in [x_0 - a, x_0 + a]} |f'''(x)| \quad (7.27)$$

with a is a small number such as 0.1. With this formula we can determine the optimal choice of h only for certain cases. In practice, we cannot compute an optimal h to use in approximating the derivative, since we have no knowledge of the third derivative of the function or it will cost more.

GlanzFreyer' Guide 7.1: Approximating Second Order Derivative

Methods can also be derived to find approximations to higher derivatives of a function using only tabulated values of the function at various points. The derivation is algebraically tedious, however, so only a representative procedure will be presented. Expand a function f in a third Taylor polynomial about a point x_0 and evaluate at $x_0 + h$ and $x_0 - h$. Then

$$f(x_0 + h) = f(x_0) + f'(x_0)h + \frac{1}{2}f''(x_0)h^2 + \frac{1}{6}f'''(x_0)h^3 + \frac{1}{24}f^{(4)}(\xi_1)h^4$$

and

$$f(x_0 - h) = f(x_0) - f'(x_0)h + \frac{1}{2}f''(x_0)h^2 - \frac{1}{6}f'''(x_0)h^3 + \frac{1}{24}f^{(4)}(\xi_{-1})h^4$$

where $x_0 - h < \xi_{-1} < x_0 < \xi_1 < x_0 + h$.

If we add these equations, the term involving $f'(x_0)$ cancels and we obtain

$$f(x_0 + h) + f(x_0 - h) = 2f(x_0) + f''(x_0)h^2 + \frac{1}{24}[f^{(4)}(\xi_1) + f^{(4)}(\xi_{-1})]h^4$$

Solving this equation for $f''(x_0)$ gives

$$f''(x_0) = \frac{1}{h^2}[f(x_0 - h) - 2f(x_0) + f(x_0 + h)] - \frac{h^2}{24}[f^{(4)}(\xi_1) + f^{(4)}(\xi_{-1})] \quad (7.28)$$

Suppose $f^{(4)}$ is continuous on $[x_0 - h, x_0 + h]$. Since $\frac{1}{2}[f^{(4)}(\xi_1) + f^{(4)}(\xi_{-1})]$ is between $f^{(4)}(\xi_1)$ and $f^{(4)}(\xi_{-1})$, the Intermediate Value Theorem implies that a number ξ exists between ξ_1 and ξ_{-1} , and hence in $(x_0 - h, x_0 + h)$, with

$$f^{(4)}(\xi) = \frac{1}{2}[f^{(4)}(\xi_1) + f^{(4)}(\xi_{-1})]$$

This permits us to rewrite the approximation equation Eq. (7.28) as

$$f''(x_0) = \frac{1}{h^2}[f(x_0 - h) - 2f(x_0) + f(x_0 + h)] - \frac{h^2}{12}f^{(4)}(\xi) \quad (7.29)$$

for some ξ , where $x_0 - h < \xi < x_0 + h$.

[SI*] Example:

Let $f(x) = \ln(x)$ and $x_0 = 1.8$, use forward-difference formula to approximate $f'(1.8)$ with $h = 0.1, 0.01, 0.001$.

Then, let $f_2(x) = xe^x$ and $x_0 = 2.0$, use three-point formulas and five-point formulas to approximate $f'(2.0)$ with $h = 0.1$ and $h = -0.1$.

Solution:

Since $f'(x) = \frac{1}{x}$, the exact value of $f'(1.8)$ is 0.55555.

For f_2 , since $f_2'(x) = (x+1)e^x$, the exact value of $f_2'(x)$ is 22.167168.

In SymIntegration we can approximate the numerical differentiation problem with forward-difference formula:

numericaldifferentiation(const Symbolic &f, const Symbolic &x, double x_0 , double h)

where f is the function of x , x_0 is the value that we want to approximate, h is the step size.

For three-point formulas, the functions in SymIntegration are:

numericaldifferentiation3pointoneside(const Symbolic &f, const Symbolic &x, double x_0 , double h)

numericaldifferentiation3pointbothsides(const Symbolic &f, const Symbolic &x, double x_0 , double h)

For five-point formulas, the functions in SymIntegration are:

numericaldifferentiation5pointoneside(const Symbolic &f, const Symbolic &x, double x_0 , double h)

numericaldifferentiation5pointbothsides(const Symbolic &f, const Symbolic &x, double x_0 , double h)

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Numerical Differentiation ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Numerical Differentiation ]# ./main
For f(x)=ln(x)
f'(1.8) with h=0.001 : 0.5554012917
f'(1.8) with h=0.01 : 0.5540180376
f'(1.8) with h=0.1 : 0.5406722127

For f(x)=x*exp(x)
Three Point Formulas one side:
f'(2.0) with h=0.1 : 22.03230487
f'(2.0) with h=0.1 : 22.05452134
Three Point Formulas both sides:
f'(2.0) with h=0.1 : 22.22878688
f'(2.0) with h=0.2 : 22.41416066
Five Point Formulas one side:
f'(2.0) with h=0.1 : 22.16699562
Five Point Formulas both sides:
f'(2.0) with h=0.1 : 22.16591457

Time taken by function: 71978 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Numerical Differentiation ]#
```

Figure 7.21: The computation of numerical differentiation for $f(x) = \ln(x)$ with forward-difference formula and for $f(x) = xe^x$ with three-point and five-point formulas (SymIntegration/Examples/Numerical Methods/Test SymIntegration Numerical Differentiation/main.cpp).

[SI*] Richardson's Extrapolation

Richardson's extrapolation is used to generate high-accuracy results while using low-order formulas. Extrapolation can be applied whenever it is known that an approximation technique has an error term with a predictable form, one that depends on a parameter, usually the step size h .

Suppose that for each number $h \neq 0$ we have a formula $N(h)$ that approximates an unknown value M and that the truncation error involved with the approximation has the form

$$M - N(h) = K_1h + K_2h^2 + K_3h^3 + \dots,$$

for some collection of unknown constants $K_1, K_2, K_3, \dots$

Since the truncation error is $O(h)$, we would expect, for example, that

$$\begin{aligned} M - N(0.1) &\approx 0.1K_1 \\ M - N(0.01) &\approx 0.01K_1 \end{aligned}$$

and, in general, $M - N(h) \approx K_1 h$, unless there was a large variation in magnitude among the constants $K_1, K_2, K_3, \dots$

The object of extrapolation is to find an easy way to combine the rather inaccurate $O(h)$ approximations in an appropriate way to produce formulas with a higher-order truncation error. Suppose, for example, we could combine the $N(h)$ formulas so as to produce an $O(h^2)$ approximation formula, $\hat{N}(h)$, for M with

$$M - \hat{N}(h) = \hat{K}_2 h^2 + \hat{K}_3 h^3 + \dots,$$

for some, again unknown, collection of constants $\hat{K}_1, \hat{K}_2, \dots$. Then we would have

$$\begin{aligned} M - \hat{N}(0.1) &\approx 0.01\hat{K}_2 \\ M - \hat{N}(0.01) &\approx 0.0001\hat{K}_2 \end{aligned}$$

and so on. If the constants K_1 and $\hat{K}_2$ are roughly of the same magnitude, then the $\hat{N}(h)$ approximations would be much better than the corresponding $N(h)$ approximations. The extrapolation continues by combining the $\hat{N}(h)$ approximations in a manner that produces formulas with $O(h^3)$ truncation error, and so on.

To see specifically how we can generate these higher-order formulas, let us consider the formula for approximating M of the form

$$M = N(h) + K_1 h + K_2 h^2 + K_3 h^3 + \dots \quad (7.30)$$

Since the formula is assumed to hold for all positive h , consider the result when we replace the parameter h by half its value. Then we have the formula

$$M = N\left(\frac{h}{2}\right) + K_1 \frac{h}{2} + K_2 \frac{h^2}{4} + K_3 \frac{h^3}{8} + \dots$$

Subtracting Eq. (7.30) from twice this equation eliminates the term involving K_1 and gives

$$M = \left[N\left(\frac{h}{2}\right) + \left(N\left(\frac{h}{2}\right) - N(h) \right) \right] + K_2 \left(\frac{h^2}{2} - h^2 \right) + K_3 \left(\frac{h^3}{4} - h^3 \right) + \dots$$

to facilitate this discussion, we define $N_1(h) \equiv N(h)$ and

$$N_2(h) = N_1\left(\frac{h}{2}\right) + \left[N_1\left(\frac{h}{2}\right) - N_1(h) \right]$$

Then we have the $O(h^2)$ approximation formula for M :

$$M = N_2(h) - \frac{K_2}{2} h^2 - \frac{3K_3}{4} h^3 - \dots \quad (7.31)$$

If we now replace h by $\frac{h}{2}$ in this formula, we have

$$M = N_2\left(\frac{h}{2}\right) - \frac{K_2}{8} h^2 - \frac{3K_3}{32} h^3 - \dots \quad (7.32)$$

this can be combined with Eq. (7.31) to eliminate the h^2 term. Specifically, subtracting Eq. (7.31) from 4 times Eq. (7.32) gives

$$3M = 4N_2\left(\frac{h}{2}\right) - N_2(h) + \frac{3K_3}{8}h^3 + \dots$$

and dividing by 3 gives an $O(h^3)$ formula for approximating M :

$$M = \left[N_2\left(\frac{h}{2}\right) + \frac{N_2\left(\frac{h}{2}\right) - N_2(h)}{3} \right] + \frac{K_3}{8}h^3 + \dots$$

By defining

$$N_3(h) \equiv N_2\left(\frac{h}{2}\right) + \frac{N_2\left(\frac{h}{2}\right) - N_2(h)}{3}$$

we have the $O(h^3)$ formula:

$$M = N_3(h) + \frac{K_3}{8}h^3 + \dots$$

The process is continued by constructing an $O(h^4)$ approximation

$$N_4(h) = N_3\left(\frac{h}{2}\right) + \frac{N_3\left(\frac{h}{2}\right) - N_3(h)}{7}$$

an $O(h^5)$ approximation

$$N_5(h) = N_4\left(\frac{h}{2}\right) + \frac{N_4\left(\frac{h}{2}\right) - N_4(h)}{15}$$

and so on. In general, if M can be written in the form

$$M = N(h) + \sum_{j=1}^{m-1} K_j h^j + O(h^m) \quad (7.33)$$

then for each $j = 2, 3, \dots, m$, we have an $O(h^j)$ approximation of the form

$$N_j(h) = N_{j-1}\left(\frac{h}{2}\right) + \frac{N_{j-1}\left(\frac{h}{2}\right) - N_{j-1}(h)}{2^{j-1} - 1} \quad (7.34)$$

These approximations are generated by rows in the order indicated by the numbered entries. This is done to take best advantage of the highest-order formulas. The extrapolation can be applied whenever the truncation error for a formula has the form

$$\sum_{j=1}^{m-1} K_j h^{\alpha_j} + O(h^{\alpha_m})$$

for a collection of constants K_j and when $\alpha_1 < \alpha_2 < \alpha_3 < \dots < \alpha_m$.

$O(h)$	$O(h^2)$	$O(h^3)$	$O(h^4)$
1 : $N_1(h) \equiv N(h)$			
2 : $N_1\left(\frac{h}{2}\right) \equiv N\left(\frac{h}{2}\right)$	3 : $N_2(h)$		
4 : $N_1\left(\frac{h}{4}\right) \equiv N\left(\frac{h}{4}\right)$	5 : $N_2\left(\frac{h}{2}\right)$	6 : $N_3(h)$	
7 : $N_1\left(\frac{h}{8}\right) \equiv N\left(\frac{h}{8}\right)$	8 : $N_2\left(\frac{h}{4}\right)$	9 : $N_3\left(\frac{h}{4}\right)$	10 : $N_4(h)$

Table 7.1: The extrapolation table

[SI*] The centered difference formula in Eq. (7.23) to approximate $f'(x_0)$ can be expressed with an error formula

$$f'(x_0) = \frac{1}{2h}[f(x_0+h) - f(x_0-h)] - \frac{h^2}{6}f'''(x_0) - \frac{h^4}{120}f^{(5)}(x_0) - \dots$$

Since this error formula contains only even powers of h , extrapolation is more effective than as outlined in the opening discussin. In this case, we have the $O(h^2)$ approximation

$$f'(x_0) = N_1(h) - \frac{h^2}{6}f'''(x_0) - \frac{h^4}{120}f^{(5)}(x_0) - \dots \quad (7.35)$$

where

$$N_1(h) \equiv N(h) = \frac{1}{2h}[f(x_0+h) - f(x_0-h)]$$

Replacing h by $\frac{h}{2}$ in this formula gives the approximation

$$f'(x_0) = N_1\left(\frac{h}{2}\right) - \frac{h^2}{24}f'''(x_0) - \frac{h^4}{1920}f^{(5)}(x_0) - \dots$$

Subtracting Eq. (7.35) from 4 times this equation eliminates the $O(h^2)$ term that involves $f'''(x_0)$ and gives

$$3f'(x_0) = 4N_1\left(\frac{h}{2}\right) - N_1(h) + \frac{h^4}{160}f^{(5)}(x_0) + \dots$$

Dividing by 3 provides an $O(h^4)$ formula

$$f'(x_0) = N_2(h) + \frac{h^4}{480}f^{(5)}(x_0) + \dots$$

where

$$N_2(h) = N_1\left(\frac{h}{2}\right) + \frac{N_1\left(\frac{h}{2}\right) - N_1(h)}{3}$$

Continuing this procedure gives, for each $j = 2, 3, \dots$, and $O(h^{2j})$ approximation

$$N_j(h) = N_{j-1}\left(\frac{h}{2}\right) + \frac{N_{j-1}\left(\frac{h}{2}\right) - N_{j-1}(h)}{4^{j-1} - 1} \quad (7.36)$$

Notice that the denominator of the quotient is $4^{j-1} - 1$ instead of $2^{j-1} - 1$ because we are now eliminating powers of h^2 instead of powers of h . Since $\left(\frac{h}{2}\right)^2 = \frac{h^2}{4}$, the multipliers used to eliminate the powers of h^2 are powers of 4 instead of 2.

[SI*] **Example:**

Suppose that $x_0 = 2.0$, $h = 0.2$, and $f(x) = xe^x$, create the extrapolation table with Richardson's extrapolation.

Solution:

$$N_1(0.2) = N(0.2) = \frac{1}{0.4}[f(2.2) - 1.8] = 22.414160$$

$$N_1(0.1) = N(0.1) = 22.228786$$

$$N_1(0.05) = N(0.05) = 22.182564$$

The exact value of $f'(x) = xe^x + e^x$ at $x_0 = 2.0$ to six decimal places is 22.167168. We see that $N_1(0.05)$ has only one decimal place of accuracy, while $N_3(0.2)$ are exact up to 6 decimal places.

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Richardson's Extrapolation ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Richardson's Extrapolation ]# ./main
For f(x) = x *exp(x), x0 = 2, h = 0.2, order or error = 0(h^4), level =3
Richardson's Extrapolation table:

      22.41416066      0      0
      22.22878688      22.16699562      0
      22.18256486      22.16715752      22.16716831

      888

Time taken by function: 7573 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Richardson's Extrapolation ]#
```

Figure 7.22: The computation of Richardson's extrapolation for $x_0 = 2.0$, $h = 0.2$, and $f(x) = xe^x$ (*SymIntegration/Examples/Numerical Methods/Test SymIntegration Richardson's Extrapolation/main.cpp*).

We use the Richardson's extrapolation formula of this (with centered difference formula to approximate $f'(x_0)$):

$$N_j(h) = N_{j-1}\left(\frac{h}{2}\right) + \frac{N_{j-1}\left(\frac{h}{2}\right) - N_{j-1}(h)}{m^{j-1} - 1} \quad (7.37)$$

with m as the order of error as in $O(h^m)$.

In SymIntegration we can use this function to obtain the extrapolation table that is using the Eq. (7.37):

richardsonextrapolation(const Symbolic &f, const Symbolic &x, double x_0 , double h , int l , int m)

where f is the function of x , x_0 is the value that we want to approximate, h is the step size, l is the level / size of the square matrix, and m is the order of error.

V. NUMERICAL INTEGRATION

[SI*] The need often arises for evaluating the definite integral of a function that has no explicit antiderivative or whose antiderivative is not easy to obtain. The basic method involved in

approximating $\int_a^b f(x) dx$ is called numerical quadrature. It uses a sum

$$\sum_{i=0}^n a_i f(x_i)$$

to approximate $\int_a^b f(x) dx$.

The methods of quadrature are based on the interpolation polynomials. We first select a set of distinct nodes $\{x_0, x_1, \dots, x_n\}$ from the interval $[a, b]$. Then we integrate the Lagrange interpolating polynomial

$$P_n(x) = \sum_{i=0}^n f(x_i) L_i(x)$$

and its truncation error term over $[a, b]$ to obtain

$$\begin{aligned} \int_a^b f(x) dx &= \int_a^b \sum_{i=0}^n f(x_i) L_i(x) dx + \int_a^b \prod_{i=0}^n (x - x_i) \frac{f^{(n+1)}(\xi(x))}{(n+1)!} dx \\ &= \sum_{i=0}^n a_i f(x_i) + \frac{1}{(n+1)!} \int_a^b \prod_{i=0}^n (x - x_i) f^{(n+1)}(\xi(x)) dx \end{aligned}$$

where $\xi(x)$ is in $[a, b]$ for each x and

$$a_i = \int_a^b L_i(x) dx$$

for each $i = 0, 1, 2, \dots, n$.

The quadrature formula is, therefore,

$$\int_a^b f(x) dx \approx \sum_{i=0}^n a_i f(x_i)$$

with error given by

$$E(f) = \frac{1}{(n+1)!} \int_a^b \prod_{i=0}^n (x - x_i) f^{(n+1)}(\xi(x)) dx$$

[SI*] To derive the Trapezoidal rule for approximating $\int_a^b f(x) dx$. let $x_0 = a$, $x_1 = b$, $h = b - a$ and use the linear Lagrange polynomial:

$$P_1(x) = \frac{(x - x_1)}{(x_0 - x_1)} f(x_0) + \frac{(x - x_0)}{(x_1 - x_0)} f(x_1)$$

Then,

$$\begin{aligned} \int_a^b f(x) dx &= \int_{x_0}^{x_1} \left[\frac{(x - x_1)}{(x_0 - x_1)} f(x_0) + \frac{(x - x_0)}{(x_1 - x_0)} f(x_1) \right] dx \\ &\quad + \frac{1}{2} \int_{x_0}^{x_1} f''(\xi(x)) (x - x_0)(x - x_1) dx \end{aligned} \tag{7.38}$$

Since $(x - x_0)(x - x_1)$ does not change sign on $[x_0, x_1]$, the Weighted Mean Value Theorem for Integrals can be applied to the error term to give, for some ξ in (x_0, x_1) .

$$\begin{aligned} \int_{x_0}^{x_1} f''(\xi(x))(x - x_0)(x - x_1) dx &= f''(\xi(x)) \int_{x_0}^{x_1} (x - x_0)(x - x_1) dx \\ &= f''(\xi(x)) \left[\frac{x^3}{3} - \frac{(x_1 + x_0)}{2}x^2 + x_0x_1x \right]_{x_0}^{x_1} \\ &= -\frac{h^3}{6}f''(\xi) \end{aligned}$$

Consequently, Eq. (7.38) implies that

$$\begin{aligned} \int_a^b f(x) dx &= \left[\frac{(x - x_1)^2}{2(x_0 - x_1)}f(x_0) + \frac{(x - x_0)^2}{2(x_1 - x_0)}f(x_1) \right] - \frac{h^3}{12}f''(\xi) \\ &= \frac{(x_1 - x_0)}{2}[f(x_0) + f(x_1)] - \frac{h^3}{12}f''(\xi) \end{aligned}$$

Since $h = x_1 - x_0$, we have the following rule:

$$\int_a^b f(x) dx = \frac{h}{2}[f(x_0) + f(x_1)] - \frac{h^3}{12}f''(\xi) \quad (7.39)$$

this is called the Trapezoidal rule because when f is a function with positive values, $\int_a^b f(x) dx$ is approximated by the area in a trapezoid.

Since the error term for the Trapezoidal rule involves f'' , the rule gives the exact result when applied to any function whose second derivative is identically zero, that is, any polynomial of degree one or less.

[SI*] Simpson's rule results from integrating over $[a, b]$ the second Lagrange polynomial with nodes $x_0 = a$, $x_2 = b$, and $x_1 = a + h$, where $h = \frac{b-a}{2}$.

Therefore,

$$\begin{aligned} \int_a^b f(x) dx &= \int_{x_0}^{x_2} \left[\frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)}f(x_0) + \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)}f(x_1) \right. \\ &\quad \left. + \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}f(x_2) \right] dx \\ &\quad + \int_{x_0}^{x_2} \frac{(x - x_0)(x - x_1)(x - x_2)}{6}f^{(3)}(\xi(x)) dx \end{aligned}$$

Deriving Simpson's rule in this manner, however, provides only an $O(h^4)$ error term involving $f^{(3)}$. By approaching this problem in another way, a higher-order term involving $f^{(4)}$ can be derived.

To illustrate this alternative formula, suppose that f is expanded in the third Taylor polynomial about x_1 . Then for each x in $[x_0, x_2]$, a number $\xi(x)$ in (x_0, x_2) exists with

$$f(x) = f(x_1) + f'(x_1)(x - x_1) + \frac{f''(x_1)}{2}(x - x_1)^2 + \frac{f'''(x_1)}{6}(x - x_1)^3 + \frac{f^{(4)}(\xi(x))}{24}(x - x_1)^4$$

and

$$\begin{aligned} \int_{x_0}^{x_2} f(x) dx = & [f(x_1)(x - x_0) + \frac{f'(x_1)}{2}(x - x_1)^2 + \frac{f''(x_1)}{6}(x - x_1)^3 \\ & + \frac{f'''(x_1)}{24}(x - x_1)^4]_{x_0}^{x_2} + \frac{1}{24} \int_{x_0}^{x_2} f^{(4)}(\xi(x))(x - x_1)^4 dx \end{aligned} \quad (7.40)$$

since $(x - x_1)^4$ is never negative on $[x_0, x_2]$, the Weighted Mean Value Theorem for Integrals implies that

$$\frac{1}{24} \int_{x_0}^{x_2} f^{(4)}(\xi(x))(x - x_1)^4 dx = \frac{f^{(4)}(\xi_1)}{24} \int_{x_0}^{x_2} (x - x_1)^4 dx = \frac{f^{(4)}(\xi_1)}{120} (x - x_1)^5 \Big|_{x_0}^{x_2}$$

for some number ξ_1 in (x_0, x_2) .

However, $h = x_2 - x_1 = x_1 - x_0$, so

$$(x_2 - x_1)^2 - (x_0 - x_1)^2 = (x_2 - x_1)^4 - (x_0 - x_1)^4 = 0$$

whereas

$$(x_2 - x_1)^3 - (x_0 - x_1)^3 = 2h^3$$

$$(x_2 - x_1)^5 - (x_0 - x_1)^5 = 2h^5$$

Consequently, Eq. (7.40) can be rewritten as

$$\int_{x_0}^{x_2} f(x) dx = 2hf(x_1) + \frac{h^3}{3}f''(x_1) + \frac{f^{(4)}(\xi_1)}{60}h^5$$

If we now replace $f'''(x_1)$ by the approximation given in Eq. (7.29), we have

$$\begin{aligned} \int_{x_0}^{x_2} f(x) dx = & 2hf(x_1) + \frac{h^3}{3} \left[\frac{1}{h^2} [f(x_0) - 2f(x_1) + f(x_2)] - \frac{h^2}{12} f^{(4)}(\xi_2) \right] + \frac{f^{(4)}(\xi_1)}{60} h^5 \\ = & \frac{h}{3} [f(x_0) + 4f(x_1) + f(x_2)] - \frac{h^5}{12} \left[\frac{1}{3} f^{(4)}(\xi_2) - \frac{1}{5} f^{(4)}(\xi_1) \right] \end{aligned}$$

it can be shown by alternative methods that the values ξ_1 and ξ_2 in this expression can be replaced by a common value ξ in (x_0, x_2) . This gives Simpson's rule:

$$\int_{x_0}^{x_2} f(x) dx = \frac{h}{3} [f(x_0) + 4f(x_1) + f(x_2)] - \frac{h^5}{90} f^{(4)}(\xi) \quad (7.41)$$

Since the error term involves the fourth derivative of f , Simpson's rule gives exact results when applied to any polynomial of degree three or less.

[SI*] **Example:**

Compute the definite integral for $x^2, x^4, \frac{1}{x+1}, \sqrt{1+x^2}, \sin x, e^x$ using the Trapezoidal rule and Simpson's rule on the interval $[0, 2]$.

Solution:

VI. NUMERICAL SOLUTIONS TO BOUNDARY-VALUE PROBLEMS FOR ORDINARY
DIFFERENTIAL EQUATIONS

[SI*]

VII. NUMERICAL SOLUTIONS TO PARTIAL DIFFERENTIAL EQUATIONS

[SI*]

Chapter 8

Molecular Biology

"I'm the most beautiful dog in the world bih, I'm too good to be true bih." - Sine Bam Bam

This section is made to honor the science field of biology that had helped a lot of people, e.g. doctor who helps cure diseases thus make people live longer and be more productive. By learning biology we will know about certain plants or herbs that are able to cure deadly disease, for example Drontal is a medicine that costs USD 3 per tablet to cure worms for dogs, if you have no such USD 3 and you love your furry friend, then you have to know that mixing turmeric (blended turmeric) in your dog foods will eliminate the worms (ringworms, roundworms, and all kind of worms in your dog's intestine). How do we know that if we don't have search engine that can tell us turmeric is able to do such thing? By learning about the molecular / structure of tumeric and simulate or interact it with all kind of parasites, worms that can be simulated and computed with computer, not only we will be smarter but we will also find better and cheaper cure for all kinds of disease.

I. DNA TRANSCRIPTION AND TRANSLATION

[SI*] A DNA (deoxyribonucleic acid) molecule is the carrier of genetic information, structured as a double helix. It is composed of two strands, each with a backbone of alternating sugar and phosphate groups. The two strands are connected by four chemical bases: adenine (A), thymine (T), cytosine (C), and guanine (G), which form rungs on the "twisted ladder."

[SI*] Structures and components of DNA:

1. **Double helix**

DNA has the shape of a twisted ladder, formed by two strands that coil around each other.

2. **Backbone**

The sides of the ladder are made of a repeating sugar-phosphate backbone.

3. **Bases**

The "rungs" are made of pairs of four nitrogenous bases:

- Adenine (A) always pairs with Thymine (T).
- Cytosine (C) always pairs with Guanine (G).

4. **Nucleotides**

Each strand is a polymer of nucleotides, which are the basic units of DNA. Each

nucleotide consists of a sugar (deoxyribose), a phosphate group, and one of the four bases.

[SI*] Functions of DNA:

- **Genetic Information**

The specific order of sequence of the bases along the DNA backbone encodes the instructions for building and maintaining an organism.

- **Heredity**

This sequence of bases allows genetic information to be passed down from one generation to the next.

- **Protein Synthesis**

DNA's instructions are used to create proteins through a two-step process of transcription and translation.

1. **Transcription**

Enzymes "read" the DNA and create a copy in the form of a messenger RNA (mRNA) molecule.

2. **Translation**

The mRNA is then used as a template to link amino acids in the correct order to form a specific protein.

[SI*] DNA transcription and translation are the two core processes of protein synthesis that convert genetic information into functional proteins. Transcription is the first step, where a segment of DNA is copied into a messenger RNA (mRNA) molecule. The second step, translation, occurs on ribosomes, where the mRNA sequence is read and used to assemble a specific chain of amino acids, when then folds into a protein.

[SI*] Transcription is a process of creating an RNA copy from a DNA template. The purpose of transcription is to make a mobile copy of a gene that can be used to build a protein. Transcription happens in the nucleus in eukaryotes; in prokaryotes, it takes place in the cytoplasm.

The enzyme RNA polymerase reads the DNA and builds the new RNA strand. The end product is a messenger RNA (mRNA) molecule.

[SI*] Translation is the process of synthesizing a protein using the mRNA template. The purpose of translation is to convert the genetic code from mRNA into a sequence of amino acids that form a protein. Translation happens on ribosomes, which are located in the cytoplasm (or on the endoplasmic reticulum in eukaryotes).

Ribosomes is the cellular machinery where translation takes place. They are made of ribosomal RNA (rRNA) and proteins and hold the mRNA and tRNA in the correct positions. Transfer RNA (tRNA) acts as an adapter molecule, with one end carrying a specific amino acid and the other having an anticodon that matches a complementary codon on the mRNA.

Ribosomes read the mRNA sequence in three-nucleotide units called codons. Each codon specifies a particular amino acid, which is delivered to the ribosome by a transfer RNA (tRNA) molecule. The ribosome then links these amino acids together, forming a chain that grows until a stop codon is reached, at which point the completed polypeptide is released.

Key players:

1. **Ribosomes**

The molecular machies that read the mRNA

2. **Transfer RNA (tRNA)**

Molecules that bring the correct amino acid to the ribosome for each codon on the mRNA.

3. **Codons**

Three-nucleotide sequences on mRNA that specify which amino acid should be added next.

4. **Anticodons**

The three-nucleotide sequence on a tRNA that matches a specific mRNA codon.

The end product of translation is a chain of amino acids that folds into a functional protein.

The mRNA Genetic Code:

- **Codons**

Each codon is a sequence of three bases (Adenine, (A), Guanine (G), Cytosine (C), and Uracil (U), which replaces Thymine found in DNA). There are 64 possible codon combinations (permutation of 3 different letters from A,G,C,U $4 \times 4 \times 4$).

- **Reading Frame**

The sequence is always read in the 5' to 3' direction, starting from a specific start codon and ending at a stop codon.

- **Start Codon**

The universally recognized start codon is AUG, which codes for the amino acid Methionine (Met) and establishes the reading frame for the entire sequence.

- **Stop Codons**

There are three stop codons that signal the termination of translation: UAA, UAG, UGA. They do not code for any amino acid.

- **Degeneracy**

Most amino acids are coded by more than one codon, which is known as the degeneracy of the genetic code.

The process of translation:

1. **Initiation**

The ribosome assembles around the mRNA molecule. The first tRNA, carrying the amino acid methionine, binds to the start codon (AUG).

2. **Elongation**

The ribosome moves along the mRNA strand one codon at a time. A tRNA with an anticodon complementary to the next codon enters the ribosome. The amino acid on this new tRNA is added to the growing polypeptide chain via a peptide bond. The ribosome then translocates to the next codon to continue the process.

3. **Termination**

The ribosome reaches a stop codon (UAA, UAG, or UGA). A release factor binds to the stop codon instead of a tRNA. This triggers the release of the completed polypeptide chain from the ribosome. The ribosomal complex and the mRNA then dissociate.

[SI*] Together, transcription and translation are the two main steps of the "central dogma" of molecular biology: DNA → RNA → Protein.

The genetic information is first transcribed from the permanent DNA "blueprint" into a temporary mRNA "message." This message is then translated into the "machinery" of the cell, which are proteins.




Figure 8.1: DNA is transcribed into mRNA, mRNA is translated into amino acids. Each codon corresponds to one amino acid. Amino acids form polypeptide chains. Polypeptide chains fold into proteins.

		Second letter				
		U	C	A	G	
First letter	U	UUU } Phe UUC } UUA } Leu UUG }	UCU } UCC } Ser UCA } UCG }	UAU } Tyr UAC } UAA Stop UAG Stop	UGU } Cys UGC } UGA Stop UGG Trp	U C A G
	C	CUU } CUC } Leu CUA } CUG }	CCU } CCC } Pro CCA } CCG }	CAU } His CAC } CAA } Gln CAG }	CGU } CGC } Arg CGA } CGG }	U C A G
	A	AUU } AUC } Ile AUA } AUG Met	ACU } ACC } Thr ACA } ACG }	AAU } Asn AAC } AAA } Lys AAG }	AGU } Ser AGC } AGA } Arg AGG }	U C A G
	G	GUU } GUC } Val GUA } GUG }	GCU } GCC } Ala GCA } GCG }	GAU } Asp GAC } GAA } Glu GAG }	GGU } GGC } Gly GGA } GGG }	U C A G

Figure 8.2: The 64 possible codon combinations for mRNA.

[SI*] The 20 standard amino acids are the building blocks for proteins. these are categorized as either essential (must be obtained from the diet) or non-essential (can be synthesized by the body).

1. Alanine (Ala, A)
2. Arginine (Arg, R)
3. Asparagine (Asn, N)

4. Aspartic Acid (Asp, D)
5. Cysteine (Cys, C)
6. Glutamic Acid (Glu, E)
7. Glutamine (Gln, Q)
8. Glycine (Gly, G)
9. Histidine (His, H)
10. Isoleucine (Ile, I)
11. Leucine (Leu, L)
12. Lysine (Lys, K)
13. Methionine (Met, M)
14. Phenylalanine (Phe, F)
15. Proline (Pro, P)
16. Serine (Ser, S)
17. Threonine (Thr, T)
18. Tryptophan (Trp, W)
19. Tyrosine (Tyr, Y)
20. Valine (Val, V)

i. Transcription and translation from DNA to mRNA to Polypeptide with SymIntegration

Suppose we have a 12 DNA code that we want to translate into polypeptide, it is :

$$[A \ C \ G \ T \ C \ A \ G \ C \ A \ G \ C \ T]$$

DNA is a double helix, so the pair will be

$$\begin{bmatrix} A & C & G & T & C & A & G & C & A & G & C & T \\ T & G & C & A & G & T & C & G & T & C & G & A \end{bmatrix}$$

Then the mRNA that carries the genetic code from the DNA to the ribosome

$$\begin{bmatrix} A & C & G & T & C & A & G & C & A & G & C & T \\ T & G & C & A & G & T & C & G & T & C & G & A \\ A & C & G & U & C & A & G & C & A & G & C & U \end{bmatrix}$$

Then we convert RNA into a polypeptide, ribosomes reading the mRNA sequence in three-nucleotide units called codons, thus

$$\begin{bmatrix} ACG & UCA & GCA & GCU \\ Thr & Ser & Ala & Ala \end{bmatrix}$$

In SymIntegration we can use this function to perform the transcription and translation:
polypeptide(vector<string> DNA)

DNA is a data type of **vector<string>**, we load a textfile **DNA.txt** that is populated with the first half pair of DNA, it is treated as a vector in linear algebra, then after we make it into double helix DNA (becoming two strands) it become a 12×2 matrix, 12 is the number of the sequence of the nitrogenous bases (Adenine, Thymine, Guanine, Cytosine).

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DNA to mRNA to Polypeptide ]# make
g++ -c -o main.o main.cpp
g++ -o main -ggdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration DNA to mRNA to Polypeptide ]# ./main
DNA double helix:
A C G T C A G C A G C T
T G C A G T C G T C G A
A C G U C A G C A G C U

Polypeptide:
ACG UCA GCA GCU
Thr Ser Ala Ala

Time taken by function: 883 microseconds
```

Figure 8.3: The computation to obtain polypeptide from first half pair of DNA (*SymIntegration/Examples/Molecular Biology/Test SymIntegration DNA to mRNA to Polypeptide/main.cpp*).

The coding till we obtain the polypeptide takes place in **src/biology-dna.cpp** and **include/symintegral/biology-dna.h**. We are open for any kind of suggestions and improvements, it is only an example to demonstrate how transcription and translation from DNA to polypeptide works, we don't really take into account the real life situation where we have to start from the start codon (AUG) and stop at stop codon (UAA, UAG, UGA).

Chapter 9

Evolutionary Algorithms

"Walnut is here. Krem is here. We are kampung dogs from Pogor." - Walnut and Krem

Evolution algorithms are computer science-oriented techniques which imitate nature according to Darwin's principle of "survival of the fittest." A population of solutions is treated with genetic operations, namely selection, crossover and mutation in order to produce improved generations. This evolution approach can be applied in fluid dynamics, electrical engineering, engineering design, electromagnetic, scheduling, pattern recognition and signal processing.

I. GENETIC ALGORITHMS

[SI*] Genetic algorithms are search algorithms on the mechanics of natural selection and natural genetics. They combine survival of the fittest among string structures with a structured yet randomized information exchange to form a search algorithm with some of the innovative flair of human search. In every generation, a new set of artificial creatures (strings) is created using bits and pieces of the fittest of the old; an occasional new part is tried for good measure. While randomized, genetic algorithms are no simple random walk. They efficiently exploit historical information to speculate on new search points with expected improved performance.

[SI*] Genetic algorithms can be used for optimization. Optimization aims for efficient allocation of scarce resources. You find examples of optimization in any scientific or engineering discipline. For example, the minimum principle of physics produces many kinds of optimization problems. Agriculturists want to improve their cattle so that the cows are more healthy and produce better milk. Engineers are interested in making better paper machines, or in optimizing the fuel consumption of engines. As another type of example, consider designing optimal schedules for school classes or aircraft crews, or minimizing the lengths of wires in an electrical device [18].

The bottleneck of optimization is finding an algorithm which solves a given class of problems. There are no methods which can solve all optimization problems. A good solution now is better than a perfect solution in the future, any method which can be adapted to a variety of problems is of great interest and of great practical use.

[SI*] Usually, we do not have a ready-made optimization problem waiting for a solution. Rather, we have a phenomenon we are interested in. The phenomenon may be a physical process, a

financial situation or an abstract mathematical construct.

The first step is to explore the phenomenon using mathematical tools. After we have a model, we may proceed to simulate the phenomenon. We often must simplify the model to make simulations. Furthermore, we may have to design new computational methods to solve the problem in a reasonable time. When we have found a solution, we have to compare the results with the actual phenomenon to verify our model and solution.

When we have an optimization problem, we have to find a solution method. Here we have a wide variety of choices to make. First, we have to find out what kind of problem we have: Are the functions smooth, non-smooth or even non-continuous? Do we have discrete or continuous variables? Is the problem convex or not? Do we have many local maxima or is there just one maximum? Can we find a good initial guess for the solution?

In many real-world cases it is not possible to formulate a standard problem. Also, there may be so many local maxima that local optimization methods are useless. Optimization is hard, but finding the global maximum is much harder.

In these cases it is worthwhile to try genetic algorithms (GA). However, tailoring a GA to the problem may be far from nontrivial. It may be reasonable to look at other methods also: simulated annealing, tabu search and other heuristics, or local optimization with several different initial guesses.

[SI*] Evolution is a collective phenomenon of adapting to the environment and passing genes to next generations. If a combination of genes gives a reasonable chance of survival and of finding a mate, this set of genes may survive to the next generation, even if the combination is non-optimal.

Evolution, unfortunately, is not optimization. Consider the human eye, where the light sensors are "backwards." Or consider the appendix, or the way we are eating and breathing through our mouths. These nonoptimal solutions are due to the genes which our distant ancestors have.

Genetic algorithms (GA) are based on principles from genetics and evolution. GA can be used to solve optimization problems, although the greatest promise may be in machine learning and adaptive systems.

The power of mathematics is in technology transfer: we have models and methods which can describe many different phenomena and solve a wide variety of problems. GAs are one example of mathematical technology transfer: by simulating evolution we are actually able to solve optimization problems from a variety of sources.

Instead of a single sample from the solution space. GAs maintain a population of vectors. New solution vectors are generated using selection, recombination and mutation. The new vectors are evaluated for their fitness, and the old population is replaced with a new one. This is repeated until the algorithm converges or we run out of time or patience.

Often a good enough solution will do. If the search space of all possible solution is big enough, we may be satisfied with a solution which is better than a random guess.

Genetic algorithms are adaptable to different situations. If you can find a good coding for the problem, GAs can be quite efficient and reliable. However, there are cases where other methods are more appropriate. For example, if the problem is smooth and derivatives are available, combining a gradient-based local optimization method with a global strategy will be hard to beat.

[SI*] One of the problems of GAs is the optimal coding of a particular optimization task. There are many choices to make: are we using binary or real coding, what are the crossover, mutation, and selection operators, and how the values of the object function are mapped into function values.

The traditional genetic algorithms don't process real numbers. Instead, a binary representation is used. In practice, using binary coding for real variables is wasteful. Often the performance of binary coded genetic algorithms is not satisfactory either.

i. Functions in SymIntegration Related to Genetic Algorithms

[SI*] We will list the basic functions that can be used related to genetic algorithms:

GA_maximizationproblem_functionwithtwovariables_N_generations(const Symbolic &F, const Symbolic &x₁, const Symbolic &x₂, int n_population, double a_{j1}, double b_{j1}, double a_{j2}, double b_{j2}, int N)

this function compute the unconstrained optimization problem for function of two variables with this specifications

$$\max f(x_1, x_2) \text{ or } \min f(x_1, x_2)$$

with $F = f(x_1, x_2)$

$$a_{j1} \leq x_1 \leq b_{j1}$$

$$a_{j2} \leq x_2 \leq b_{j2}$$

n_population is the number of the chromosome population, *N* is the number of generations / iterations we want to compute.

ii. Optimization Problem with Simple Genetic Algorithms

Consider this example of unconstrained optimization problem :

$$\max f(x_1, x_2) = 21.5 + x_1 \sin(4\pi x_1) + x_2 \sin(20\pi x_2)$$

with

$$-3.0 \leq x_1 \leq 12.1$$

$$4.1 \leq x_2 \leq 5.8$$

We want to use genetic algorithms to find the solution.

Representation

First, we need to encode decision variables into binary strings. The length of the string depends on the required precision. For example, the domain of variable x_j is $[a_j, b_j]$ and the required precision is five places after the decimal point. The precision requirement implies that the range of domain of each variable should be divided into at least $(b_j - a_j) \times 10^5$ size ranges. The required bits (denoted with m_j) for a variable is calculated as follows:

$$\begin{aligned} 2^{m_j-1} &\leq (b_j - a_j) \times 10^5 \leq 2^{m_j} - 1 \\ (m_j - 1) \ln(2) &\leq \ln((b_j - a_j) \times 10^5) \leq m_j \ln(2) - \ln(1) \\ (m_j - 1) &\leq \frac{\ln((b_j - a_j) \times 10^5)}{\ln(2)} \leq m_j \end{aligned}$$

The mapping from a binary string to a real number for variable x_j is straightforward and completed as follows:

$$x_j = a_j + \text{decimal}(\text{substring}_j) \times \frac{b_j - a_j}{2^{m_j} - 1} \quad (9.1)$$

where $\text{decimal}(\text{substring}_j)$ represents the decimal value of substring_j for decision variable x_j .

Suppose that the precision is set as five places after the decimal point. The required bits for variables x_1 and x_2 is calculated as follows:

$$\begin{aligned} (12.1 - (-3.0)) \times 100,000 &= 1,510,000 \\ 2^{20} &< 1,510,000 \leq 2^{21}, \quad m_1 = 21 \\ (5.8 - (4.1)) \times 100,000 &= 170,000 \\ 2^{17} &< 1,510,000 \leq 2^{18}, \quad m_2 = 18 \\ m &= m_1 + m_2 = 21 + 18 = 39 \end{aligned}$$

The total length of a chromosome is 39 bits which can be represented as follows:

$$\begin{array}{cc} 000000101000100110100 & 101101011100101011 \\ \text{21bits} & \text{18bits} \end{array}$$

The corresponding values for variables x_1 and x_2 are given below: In standard C++ template you are able to convert binary number that is represented as **string** data type into decimal value with **stoi(...)** function.

	Binary Number	Decimal Number
x_1	000000101000100110100	20788
x_2	101101011100101011	186155

Table 9.1: The variable x_1 and x_2 with its corresponding binary and decimal number.

Initial Population

Initial population is randomly generated as follows:

$$\begin{aligned}
v_1 &= [110110101011100010001010110100101001001] \\
v_2 &= [100010010000101110010000001111010111010] \\
v_3 &= [101001010001001100101111001101000010100] \\
v_4 &= [011100011101000010110010101001100010110] \\
v_5 &= [011110011111101101100111001010010011100] \\
v_6 &= [111011101110011000100100101000100000010] \\
v_7 &= [001001101101010010001110010010100001000] \\
v_8 &= [000110110011000110010000111101101010010] \\
v_9 &= [011101111111010010011111100001010101100] \\
v_{10} &= [101001011000011101001010000110010010111]
\end{aligned}$$

The corresponding values are

$$\begin{aligned}
v_1 &= [x_1, x_2] = [9.90112, 4.69979] \\
v_2 &= [x_1, x_2] = [5.08353, 4.15101] \\
v_3 &= [x_1, x_2] = [6.73684, 5.6308] \\
v_4 &= [x_1, x_2] = [3.71332, 4.66294] \\
v_5 &= [x_1, x_2] = [4.19503, 5.62172] \\
v_6 &= [x_1, x_2] = [11.0913, 5.08449] \\
v_7 &= [x_1, x_2] = [-0.709624, 5.43648] \\
v_8 &= [x_1, x_2] = [-1.396, 4.30473] \\
v_9 &= [x_1, x_2] = [4.0755, 5.69819] \\
v_{10} &= [x_1, x_2] = [6.7636, 4.5459]
\end{aligned}$$

Evaluation

The process of evaluating the fitness of a chromosome consists of the following three steps:

1. Step 1

Convert the chromosome's genotype to its phenotype. Here, this means converting binary string into relative real values $x^k = (x_1^k, x_2^k), k = 1, 2, \dots, pop_size$.

2. Step 2

Evaluate the objective function $f(x^k)$.

3. Step 3

Convert the value of objective function into fitness. For the maximization problem, the fitness is simply equal to the value of objective function $eval(v_k) = f(x^k), k = 1, 2, \dots, pop_size$.

An evaluation function plays the role of the environment, and it rates chromosomes in terms of their fitness.

The fitness function values of above chromosomes are as follows:

$$\begin{aligned}
 eval(v_1) &= f(9.90112, 4.69979) = 10.9558 \\
 eval(v_2) &= f(5.08353, 4.15101) = 29.9485 \\
 eval(v_3) &= f(6.73684, 5.6308) = 20.0021 \\
 eval(v_4) &= f(3.71332, 4.66294) = 19.2558 \\
 eval(v_5) &= f(4.19503, 5.62172) = 20.713 \\
 eval(v_6) &= f(11.0913, 5.08449) = 36.2712 \\
 eval(v_7) &= f(-0.709624, 5.43648) = 26.9615 \\
 eval(v_8) &= f(-1.396, 4.30473) = 16.0388 \\
 eval(v_9) &= f(4.0755, 5.69819) = 29.1546 \\
 eval(v_{10}) &= f(6.7636, 4.5459) = 21.2002
 \end{aligned}$$

It is clear that chromosome v_6 is the strongest one and that chromosome v_1 is the weakest one.

Selection

In most practices, a roulette wheel approach is adopted as the selection procedure; it belongs to the fitness-proportional selection and can select a new population with respect to the probability distribution based on fitness values. The roulette wheel can be constructed as follows:

1. Calculate the fitness value $eval(v_k)$ for each chromosome v_k :

$$eval(v_k) = f(x^k), \quad k = 1, 2, \dots, pop_size$$

2. Calculate the total fitness for the population:

$$F = \sum_{k=1}^{pop_size} eval(v_k)$$

3. Calculate selection probability p_k for each chromosome v_k :

$$p_k = \frac{eval(v_k)}{F}, \quad k = 1, 2, \dots, pop_size$$

4. Calculate cumulative probability q_k for each chromosome v_k :

$$q_k = \sum_{j=1}^k p_j, \quad k = 1, 2, \dots, pop_size$$

The selection process begins by spinning the roulette wheel pop_size times: each time, a single chromosome is selected for a new population in the following way:

1. Step 1

Generate a random number r (pop_size times) from the range $[0, 1]$.

2. Step 2

If $r \leq q_1$, then select the first chromosome v_1 ; otherwise, select the k th chromosome v_k ($2 \leq k \leq pop_size$) such that $q_{k-1} < r \leq q_k$.

The total fitness F of the population is

$$F = \sum_{k=1}^{10} eval(v_k) = 230.501$$

The probability of a selection p_k for each chromosome $v_k (k = 1, 2, \dots, 10)$ is as follows:

$$\begin{array}{lll} p_1 = 0.0475303 & p_2 = 0.129928 & p_3 = 0.0867765 \\ p_4 = 0.0835386 & p_5 = 0.0898606 & p_6 = 0.157358 \\ p_7 = 0.116969 & p_8 = 0.0695822 & p_9 = 0.126483 \\ p_{10} = 0.919742 & & \end{array}$$

The cumulative probabilities q_k for each chromosome $v_k (k = 1, 2, \dots, 10)$ is as follows:

$$\begin{array}{lll} q_1 = 0.0475303 & q_2 = 0.177458 & q_3 = 0.264234 \\ q_4 = 0.347773 & q_5 = 0.437633 & q_6 = 0.594991 \\ q_7 = 0.71196 & q_8 = 0.781543 & q_9 = 0.908026 \\ q_{10} = 1 & & \end{array}$$

Now we are ready to spin the roulette wheel 10 times, and each time we select a single chromosome for a new population. Let us assume that a random sequence of 10 numbers from the range $[0, 1]$ is as follows:

$$\begin{array}{lll} 0.438817 & 0.585455 & 0.35496 \\ 0.303427 & 0.323658 & 0.915456 \\ 0.661517 & 0.417606 & 0.586972 \\ 0.947615 & & \end{array}$$

The first number $r_1 = 0.438817$ is greater than q_5 and smaller than q_6 , meaning that the chromosome v_6 is selected for the new population: the second number $r_2 = 0.585455$ is greater than q_5 and smaller than q_6 , meaning that the chromosome v_6 is again selected for the new population; and so on. Finally, the new population consists of the following chromosomes:

$$\begin{array}{ll} v'_1 & = [111011101110011000100100101000100000010] (v_6) \\ v'_2 & = [111011101110011000100100101000100000010] (v_6) \\ v'_3 & = [011110011111101101100111001010010011100] (v_5) \\ v'_4 & = [011100011101000010110010101001100010110] (v_4) \\ v'_5 & = [011100011101000010110010101001100010110] (v_4) \\ v'_6 & = [101001011000011101001010000110010010111] (v_{10}) \\ v'_7 & = [001001101101010010001110010010100001000] (v_7) \\ v'_8 & = [011110011111101101100111001010010011100] (v_5) \\ v'_9 & = [111011101110011000100100101000100000010] (v_6) \\ v'_{10} & = [101001011000011101001010000110010010111] (v_{10}) \end{array}$$

Crossover

Crossover used here is one-cut-point method, which randomly selects one cut-point and exchanges the right parts of two parents to generate offspring. Consider two chromosomes as follows, and the cut point is randomly selected after the 21st gene:

$$\begin{array}{ll} v_1 & = [111011101110011000100|1001010001000000010] \\ v_2 & = [101001011000011101001|010000110010010111] \end{array}$$

The resulting offspring by exchanging the right parts of their parents would be as follows:

$$\begin{array}{ll} v'_1 & = [111011101110011000100|010000110010010111] \\ v'_2 & = [101001011000011101001|1001010001000000010] \end{array}$$

The probability of crossover is set as $p_c = 0.25$, so we expect that, on average, 25% of chromosomes undergo crossover.

Now, we will generate a random number r (pop_size times) from the range $[0, 1]$, assume that the sequence of random numbers is

```
0.941361  0.522253  0.729774
0.41996   0.453728  0.601589
0.67799   0.764918  0.363573
0.0676645
```

This means that the chromosome v_{10} is the only chromosome selected for crossover, since we don't have the couple here, in this computation example, we will proceed to the next step -> mutation.

Otherwise, if we have at least 2 chromosomes that were chosen for crossover. We will then generate a random integer number pos from the range $[1, 38]$ (because 39 is the total length of a chromosome) as cutting point or in other words, the position of the crossover point. Assume that the generated number pos equals 1, then the two chromosomes are cut after the first bit, and offspring are generated by exchanging the right parts of them.

In the C++ code in SymIntegration, we assume that if the chosen chromosomes are odd number and more than 2, then the last one will not undergo any crossover, the chromosomes that undergone the crossover are the first even number only. In other words, we assume the mating is loyal only to another one, and not mating with another one.

Mutation

Mutation alters one or more genes with a probability equal to the mutation rate. Assume that the 23rd gene of the chromosome v_1 is selected for a mutation. Since the gene is 1, it would be flipped into 0. Thus the chromosome after mutation would be

$$\begin{aligned} v_1 &= [111011101110011000100|010000110010010111] \\ &\quad \downarrow \\ v'_1 &= [111011101110011000100|000000110010010111] \end{aligned}$$

The probability of mutation is set as $p_m = 0.01$, so we expect that, on average, 1% of total bit of population would undergo mutation. There are $m \times pop_size = 39 \times 10 = 390$ bits in the whole population; we expect 3.9 mutations per generation. Every bit has an equal chance to be mutated. Thus we need to generate a sequence of random numbers $r_k (k = 1, 2, \dots, 390)$ from the range $[0, 1]$, or we generate a pair sequence of random numbers of $r_{row}, row = 1, 2, \dots, m$ and $r_{column}, column = 1, 2, \dots, pop_size$ for the choice of bit that will undergo mutation.

After the mutation we will obtain the corresponding decimal values of variables $[x_1, x_2]$ and the new fitness are as follows:

$f(10.6194, 5.08449)$	= 36.7526
$f(11.0914, 5.08449)$	= 36.2746
$f(4.19503, 5.62172)$	= 20.713
$f(3.71332, 4.66294)$	= 19.2558
$f(3.71378, 4.66294)$	= 19.2367
$f(6.7636, 4.5459)$	= 21.2002
$f(-0.709624, 5.43648)$	= 26.9615
$f(4.19503, 5.62172)$	= 20.713
$f(11.0913, 5.08449)$	= 36.2712
$f(6.70461, 4.5459)$	= 25.9702

Now we just completed one iteration of genetic algorithm. The test run is terminated after 5 generations. We have obtained the best chromosome in the 5th generation as follows:

$$\mathbf{v}^* = (1111111111111111100101111100010111101011)$$

$$eval(\mathbf{v}^*) = f(12.0998, 5.70358) = 37.7174$$

Compared to the result of [11] that will have $eval(\mathbf{v}^*) = f(11.631407, 5.724824) = 38.818208$ at the 419th generation, we don't use a lot of generations, instead we only use 5 and re-initialize it again to see in the 5th generation the highest $eval(\mathbf{v}^*)$ will reach which value.

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration Genetic Algorithms for Mutivariable Functions Optimization
# ./main

Maximize this function: x1*sin(12.5664*x1)+x2*sin(20*x2)+21.5

-3 <= x_{1} <= 12.1
4.1 <= x_{2} <= 5.8

m_{1} = 21
m_{2} = 18

Initial population for x_{1}:
110110101011100010001
100010010000101110010
101001010001001100101
011100011101000010110
011110011111101101100
111011101110011000100
001001101101010010001
000110110011000110010
011101111111010010011
101001011000011101001

Corresponding Decimal for x_{1}:
1.79176e+06
1.12267e+06
1.35229e+06
932374
999276
1.95706e+06
318097
222770
982675
1.35601e+06

Corresponding value for x_{1}:
9.90112
5.08353
6.73684
3.71332
4.19503
11.0913
-0.709624
-1.396
4.0755
6.7636

```

Figure 9.1: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Mutivariable Functions Optimization/main.cpp).

```

Initial population for x_{2}:
010110100101001001
000001111010111010
111001101000010100
010101001100010110
111001010010011100
100101000100000010
110010010100001000
000111101101010010
111100001010101100
010000110010010111

Corresponding Decimal for x_{2}:
92489
7866
236052
86806
234652
151810
206088
31570
246444
68759

Corresponding value for x_{2}:
4.69979
4.15101
5.6308
4.66294
5.62172
5.08449
5.43648
4.30473
5.69819
4.5459

Initial population for x_{1} and x_{2}:
11011010101100010001010110100101001001
100010010000101110010000001111010111010
101001010001001100101111001101000010100
0111000111010000101100101001100010110
01111001111101101100111001010010011100
111011101110011000100100101000100000010
001001101101010010001110010010100001000
0001101100110001100100001111011010010
0111011111101001001111110000101010100
10100101100001110100101000011001001011

```

Figure 9.2: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Multivariable Functions Optimization/main.cpp).

```

*****
Procedure: Evaluation
Generation: 1
*****

F(x_{1}, x_{2}):
10.9558
29.9485
20.0021
19.2558
20.713
36.2712
26.9615
16.0388
29.1546
21.2002

```

Figure 9.3: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Multivariable Functions Optimization/main.cpp).

```

*****
Procedure: Selection
Generation: 1
*****

Total fitness for the population = 230.501

Selection probability, p_{k}:
0.0475303
0.129928
0.0867765
0.0835386
0.0898606
0.157358
0.116969
0.0695822
0.126483
0.0919742

Cumulative probability, q_{k}:
0.0475303
0.177458
0.264234
0.347773
0.437633
0.594991
0.71196
0.781543
0.908026
1

Roulette Wheel, random sequence of 10 numbers from the range [0,1]:
0.438817
0.585455
0.35496
0.303427
0.323658
0.915456
0.661517
0.417606
0.586972
0.947615

```

Figure 9.4: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (*SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Mutivariable Functions Optimization/main.cpp*).

```

New population :
111011101110011000100100101000100000010
111011101110011000100100101000100000010
01111001111101101100111001010010011100
011100011101000010110010101001100010110
011100011101000010110010101001100010110
101001011000011101001010000110010010111
0010011011010010001110010010100001000
01111001111101101100111001010010011100
111011101110011000100101000100000010
101001011000011101001010000110010010111

```

Figure 9.5: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (*SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Mutivariable Functions Optimization/main.cpp*).

```
*****
Procedure: Crossover
Generation: 1
*****

Random sequence of 10 numbers from the range [0,1]:
0.941361
0.522253
0.729774
0.41996
0.453728
0.601589
0.67799
0.764918
0.363573
0.0676645

Assume that the probability of crossover is 0.25

Crossover population :
10100101100001110100101000011001001011

Bits per chromosome = 39
Crossover position = 17
Crossover population final :
Crossover population index:
9

Population after crossover:
111011101110011000100100101000100000010
111011101110011000100100101000100000010
01111001111101101100111001010010011100
011100011101000010110010101001100010110
011100011101000010110010101001100010110
101001011000011101001010000110010010111
001001101101010010001110010010100001000
01111001111101101100111001010010011100
111011101110011000100100101000100000010
101001011000011101001010000110010010111
```

Figure 9.6: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (*SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Multivariable Functions Optimization/main.cpp*).

```

*****
Procedure: Mutation
Generation: 1
*****

Population size = 398

Assume that the probability of mutation is : 0.01

We expect 4 mutations per generation.
Every bit has an equal chance to be mutated.

Generating a sequence of random numbers for the row and column...

Mutated row:
1
2
5
10

Mutated column:
5
18
15
8

Population before mutation:
11011101110011000100100101000100000010
11011101110011000100100101000100000010
011110011111101101100111001010010011100
0111000111010000101100101001100010110
011000111010000101100101001100010110
1010010110000111010010100001100100111
0010011011010010001110010010100001000
01111001111101101100111001010010011100
11011101110011000100100101000100000010
1010010110000111010010100001100100111

Population after mutation:
11001101110011000100100101000100000010
11011101110011001100100101000100000010
01111001111110110111001010010011100
0111000111010000101100101001100010110
01110001110100101100101001100010110
1010010110000111010010100001100100111
0010011011010010001110010010100001000
011110011111101100111001010010011100
110111011100110001001010001000000010
1010010010000111010010100001100100111

```

Figure 9.7: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Mutivariable Functions Optimization/main.cpp).

```

*****
Procedure: Evaluating New Generation
Generation: 1
*****

New binary x1:
111001101110011000100
11011101110011001100
011110011111101101100
011100011101000010110
011100011101001010110
101001011000011101001
0010011011010010001
011110011111101100
11011101110011000100
101001001000011101001

New binary x2:
1001010001000000010
1001010001000000010
11001010010011100
010101001100010110
010101001100010110
010000110010010111
110010010100001000
111001010010011100
1001010001000000010
010000110010010111

```

Figure 9.8: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Mutivariable Functions Optimization/main.cpp).


```
New x1 :
10.6194
11.0914
4.19503
3.71332
3.71378
6.7636
-0.709624
4.19503
11.0913
6.70461

New x2:
5.08449
5.08449
5.62172
4.66294
4.66294
4.5459
5.43648
5.62172
5.08449
4.5459

New F(x_{1}, x_{2}) evaluation:
36.7526
36.2746
20.713
19.2558
19.2367
21.2002
26.9615
20.713
36.2712
25.9702
```

Figure 9.9: The computation of the simple genetic algorithms for the unconstrained optimization problem (maximization) (*SymIntegration/Examples/Genetic Algorithms/Test SymIntegration Genetic Algorithms for Mutivariable Functions Optimization/main.cpp*).

Chapter 10

Machine Learning

"Krem is not being naughty, Krem is only being Krem-self." - Krem

Machine learning is a subfield of artificial intelligence (AI) that allows machines to learn from data without being explicitly programmed.

Artificial neural networks are a subset or technique within machine learning. ANNs were first conceived decades ago and have been an important part of machine learning and artificial intelligence ever since. The basic idea behind neural networks is to combine linear combinations and nonlinear functions into layers such that they can approximate arbitrary functions. Neural networks have been used to great effect for example in image classification and recognition once the computational power and suitable algorithms to train large neural networks had become available [15].

Neural networks can be used for pattern recognition, e.g. digit recognition, letters recognition, image recognition. The problem of searching patterns in data is a fundamental one and has a long and successful history. For instance, the extensive astronomical observations of Tycho Brahe in the 16th century allowed Johannes Kepler to discover the empirical laws of planetary motion, which in turn provided a springboard for the development of classical mechanics. Similarly, the discovery of regularities in atomic spectra played a key role in the development and verification of quantum physics in the early twentieth century. The field of pattern recognition is concerned with the automatic discovery of regularities in data through the use of computer algorithms and with the use of these regularities to take actions such as classifying the data into different categories.

The use of artificial neural networks as a tool to help computers recognize patterns and simulate human intelligence had been introduced in the 1980s, by the early 2000s, Yann LeCun, Geoffrey Hinton, and Yoshua Bengio (recipients of the 2018 ACM A.M. Turing Award) were among a small group who remained committed to this approach. Deep learning methods (e.g., CNN) have been responsible for astonishing breakthroughs in computer vision, speech recognition, natural language processing, and robotics.

Artificial intelligence is now one of the fastest-growing areas in all of science and one of the most talked-about topics in society. The growth of and interest in AI is due to the recent advances in deep learning. These technologies are used by billions of people. Anyone who has a smartphone in their pocket can tangibly experience advances in natural language processing and computer

vision. In addition to the products we use every day, new advances in deep learning have given scientists powerful new tools-in areas ranging from medicine, to astronomy, to materials science.

I. INTRODUCTION

[SI*] The idea underlying an artificial neural network is to define a function taking a vector as the input and yielding a vector as the output. The analogy of artificial neural networks obviously stems from the neurons and axons in the nervous system and the brain, although the dependencies or connections implemented by axons are in general much more complicated (and still mostly unknown) than the ones between adjacent layers indicated.




Figure 10.1: The schematic diagram of an artificial neural network. Each neuron influences each neuron in the next layer.

It may be unsurprising that large enough neural networks with a certain number and structure of the hidden layers have been used very successfully in image-classification and image-recognition tasks.

In engineering, our aim is not to understand the brain per se, but to build better computer systems. The brain is an information processing device that has some incredible abilities and surpasses current engineering products in many domains, e.g., vision, speech recognition, and learning. These applications have evident economic utility if implemented on machines. If we can understand how the brain performs these functions, we can define solutions to these tasks as formal algorithms and implement them on computers.

The human brain is quite different from a computer. Whereas a computer generally has one processor, the brain is composed of a very large (10^{11}) number of processing units, namely, neurons, operating in parallel. The processing units are believed to be much simpler and slower than a processor in a computer. You can just compile a code and it will be able to

give you the initial value problem of a fourth order linear differential equation solution in few seconds, while on hands it will take probably around 10 minutes or more for human to compute the solution.

What makes the brain different, and is believed to provide its computational power, is the large connectivity. Neurons in the brain have connections, called synapses, to around 10^4 other neurons, all operating in parallel. In a computer, the processor is active and the memory is separate and passive, but it is believed that in the brain, both the processing and memory are distributed together over the network; processing is done by the neurons, and the memory is in the synapses between the neurons.

The difference between natural and artificial neural network might be the same as the classic example of the natural and artificial flying machines. An eagle flaps its wings; while a commercial airplane does not flap its wings but uses jet engines. The eagle and the airplane are two hardware implementations built for different purposes, satisfying different constraints. But both implement the same theory, which is aerodynamics.

[SI*] **Activation Functions**

Activation functions are critical in neural networks because they introduce non-linearity, allowing the model to learn complex patterns in data. Without them, a neural network would be equivalent to a linear regression model, no matter how many layers it had.

The activation function is applied to the weighted sum of inputs before producing the final output of a neuron. Activation functions decide whether a neuron should be activated based on the weighted sum of inputs and a bias term. They also make backpropagation possible by providing gradients for weight updates.

Suppose we have input vector $x = \{x_1, x_2, \dots, x_n\}$ with its corresponding weights $w = \{w_1, w_2, \dots, w_n\}$, thus the inputs will go to certain node and we will have to compute the sum formula first

$$\sum_{i=1}^n x_i w_i + b$$

with b as the bias term. Then we will put this sum formula into a certain activation function (sigmoid, tanh, ReLU, etc) that will be denoted as $f(\cdot)$ thus

$$f\left(\sum_{i=1}^n x_i w_i + b\right)$$

The activation function $f(\cdot)$ will be the output from a node that has the input vector x and its corresponding weight vector w .

Now, we will explain some activation functions that can be chosen depend on the case or application of the artificial neural networks:

ReLU (Rectified Linear Unit)

This activation function is a widely used activation function in deep learning that outputs the input directly if it is positive and returns zero otherwise. Its simplicity and efficiency make it a default choice in many neural network architectures, helping models learn complex patterns while reducing issues like the vanishing gradient problem (a common issue with sigmoid and tanh activation functions). It allows positive values to pass unchanged and sets

negative values to zero.

The output range: $(0, \infty)$.

It is defined as follow:

$$\text{ReLU}(x) = \max(0, x) \quad (10.1)$$

it returns x if x is greater than 0, if x is less than or equal to 0, the function returns 0.

ReLU's ability to output zero for negative inputs introduces sparsity in the network, meaning only a fraction of neurons activate at any given time. This can lead to more efficient and faster computation.

ReLU is computationally efficient as it involves only a thresholding operation. This simplicity makes it easy to implement and compute, which is important when training deep neural networks with millions of parameters. Although it seems like a piecewise linear function, ReLU is still a non-linear function.

Sigmoid

This activation function compresses input values into a range between 0 and 1, making it suitable for binary classification. However, sigmoid suffers from vanishing gradients, especially with very high or very low inputs.

The output range: $(0, 1)$.

It is defined as follow:

$$S(x) = \frac{1}{1 + e^{-x}} \quad (10.2)$$

The derivative of sigmoid function is known as the density of the logistic distribution

$$\frac{d}{dx} S(x) = f(x)(1 - f(x)) \quad (10.3)$$

Softmax

The softmax activation function converts raw output scores (logits) from a neural network into a probability distribution, where each value is between 0 and 1, and all values sum to exactly 1. Primarily used in the final layer of multi-class classification models, it makes output interpretable by highlighting the most likely class. It is a $\mathbb{R}^n \rightarrow \mathbb{R}^n$ function, it takes a vector of n size as the input and produce a vector of n size as the output.

The output range: $(0, 1)$.

For a vector of raw outputs (logits) $x = \{x_1, x_2, \dots, x_n\}$, the softmax function is defined as follows:

$$s_i = \sigma(x)_i = \frac{e^{x_i - \max(x)}}{\sum_{j=1}^n e^{x_j - \max(x)}} \quad (10.4)$$

the exponentiation makes all inputs positive and accentuates differences, then dividing by $\sum e^{x_j - \max(x)}$ ensures the sum of all output is 1 (normalization process). A direct implementation of e^{x_i} can easily cause an overflow error if x_i is large, to prevent this we often use $e^{x_i - \max(x)}$.

This will shift the inputs to a range close to zero, assuming the inputs themselves are not too far from each other. Crucially, it shifts them all to be negative (except the maximal x_j which turns into a zero). Negatives with large exponents "saturate" to zero rather than infinity, so we have a better chance of avoiding NaNs result.

The resulting probabilities indicates the confidence in each class i . This activation function is typically used at the end of Convolutional Neural Networks or Deep Neural Networks

for classifying inputs into one of n mutually exclusive categories, it also works best with Categorical Cross-Entropy Loss to calculate the difference between predicted and actual probability distributions.

Categorical Cross-Entropy loss measures the dissimilarity between the probability distributions predicted by a machine learning model and the true probability distributions of the target classes. It is the standard loss function for multi-class classification.

When using softmax as the output activation combined with Categorical Cross-Entropy loss, the derivative simplifies into a remarkably elegant formula:

$$\delta_i = \hat{y}_i - y_i$$

where $\hat{y}_i$ is the predicted probability from the softmax function for class i , y_i is the actual target for class i , which is typically 1 for correct class and 0 for incorrect class (one-hot encoded). To understand why this happens, we must apply the chain rule. The delta for the output layer is the derivative of the loss (L) with respect to the pre-activation input (x_i).

1. Categorical Cross-Entropy Loss

For a multi-class classification task, the loss function is defined as:

$$L = - \sum_{j=1}^n y_j \log(\hat{y}_j)$$

with n as the total number of classes, y_j is the actual output (usually a one-hot encoded vector), and $\hat{y}_j$ is the predicted probability for class j .

Because the true labels y_j are one-hot encoded, only the term corresponding to the true class ($y_{true} = 1$) remains, simplifying the formula to

$$L = - \log(\hat{y}_{true})$$

The multi-class classification is a classification problem where each sample belongs to only one class (mutually exclusive), it is different than multi-label classification where each sample may belong to multiple classes (or to no class).

2. The Softmax Derivative (Jacobian Matrix)

Because a change in a single pre-activation x_j affects all probabilities $\hat{y}_i$ (due to the normalization division). Since softmax has multiple inputs, with respect to which input element the partial derivative is computed the derivative of softmax requires a Jacobian matrix:

$$\frac{\partial \hat{y}_i}{\partial x_j} = \hat{y}_i(\delta_{ij} - \hat{y}_j) = \begin{cases} \hat{y}_i(1 - \hat{y}_i), & \text{if } i = j \\ -\hat{y}_i\hat{y}_j, & \text{if } i \neq j \end{cases}$$

where δ_{ij} is the Kronecker delta (equal to 1 if $i = j$, and 0 if $i \neq j$), thus

$$J_{softmax} = \begin{bmatrix} \frac{\partial s_1}{\partial x_1} & \frac{\partial s_1}{\partial x_2} & \cdots & \frac{\partial s_1}{\partial x_n} \\ \frac{\partial s_2}{\partial x_1} & \frac{\partial s_2}{\partial x_2} & \cdots & \frac{\partial s_2}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial s_n}{\partial x_1} & \frac{\partial s_n}{\partial x_2} & \cdots & \frac{\partial s_n}{\partial x_n} \end{bmatrix}$$

where n is the number of classification. When talking about the derivative of the softmax function, we actually talk about its Jacobian matrix, which is the matrix of all

first-order derivatives.

For vector functions like softmax it is imprecise to talk about a "gradient"; the Jacobian is the fully general derivative of a vector function.

3. Combining the Chain Rule

To get the final delta, you take the derivative of the loss with respect to the pre-activation x_i :

$$\delta_i = \frac{\partial L}{\partial x_i} = \sum_j \frac{\partial L}{\partial \hat{y}_j} \frac{\partial \hat{y}_j}{\partial x_i}$$

By substituting the derivatives and evaluating summation for one-hot encoded targets, the complex derivatives collapses into the simple difference between prediction and actual output:

$$\delta_i = \hat{y}_i - y_i$$

where $\hat{y}_i$ is the predicted probability from the softmax function for class i , y_i is the actual target for class i , which is typically 1 for correct class and 0 for incorrect class (one-hot encoded).

We note that the formula of $\delta_i = \hat{y}_i - y_i$ does not give a satisfactory result when we compute for **Feed-Forward Neural Network for Iris Dataset Classification with 1 Hidden Layer** subsection under the feed-forward neural network section, thus we use this formula instead:

$$\delta_j^L = f'(s_j^L) (y_j - \hat{y}_j)$$

with $f'(s_j^L)$ as the Jacobian matrix and the $y_j - \hat{y}_j$ as the error for class j , L is the last layer, the output layer. This formula gives amazing accuracy between 97% to 99%.

Tanh (Hyperbolic Tangent)

This activation function is often used in hidden layers of neural networks.

The output range: $(-1, 1)$.

It is defined as follow:

$$\tanh(x) = \frac{\sinh(x)}{\cosh(x)} = \frac{e^x - e^{-x}}{e^x + e^{-x}} \quad (10.5)$$

The derivative of *Tanh* is defined as

$$\frac{d}{dx} \left(\frac{e^x - e^{-x}}{e^x + e^{-x}} \right) = 1 - \tanh(x)^2 \quad (10.6)$$

Leaky ReLU

This activation function addresses the dying ReLU problem by allowing a small gradient for negative inputs, defined as

$$f(x) = \begin{cases} x, & \text{if } x > 0 \\ \alpha x, & \text{if } x \leq 0 \end{cases} \quad (10.7)$$

where α is a small positive constant (e.g., 0.01). This retains computational simplicity while ensuring neurons remain active. It is a useful alternative to ReLU when dying neurons are a concern.

In Parametric ReLU, it extends ReLU by making the slope α a learnable parameter during training. This flexibility allows the network to adaptively adjust the negative slope for

improved performance. It is often used in deeper architectures where a more adaptable activation is beneficial.

Exponential Linear Unit (ELU)

This activation function introduces smooth transitions for negative values. It mitigates the vanishing gradient problem and improves gradient flow, particularly in deep networks. However, it has slightly higher computational costs and is used where faster convergence is desired. It is defined as

$$f(x) = \begin{cases} x, & \text{if } x > 0 \\ \alpha(e^x - 1), & \text{if } x \leq 0 \end{cases} \quad (10.8)$$

where $\alpha > 0$.

Swish

This activation function is a smooth, non-monotonic activation function that often outperforms ReLU in deep architectures like EfficientNet. It enhances gradient flow but comes with a slightly higher computation cost. Swish is commonly used in modern CNNs and transformer-based models. It is defined as

$$f(x) = x \cdot \sigma(x) = x \cdot \frac{1}{1 + e^{-x}} \quad (10.9)$$

Function	Range	Gradient	Advantages	Disadvantages	Use Cases
Sigmoid	$(0, 1)$	Small for extremes	Smooth, probabilistic output	Vanishing gradients	Output layer for binary tasks
Tanh	$(-1, 1)$	Small for extremes	Zero centered	Vanishing gradients	RNNs
ReLU	$[0, \infty)$	Constant (1 for $x > 0$)	Sparse, efficient	Dying ReLU problems	CNNs, general DNNs
Leaky ReLU	$(-\infty, \infty)$	Constant	Mitigates dying ReLU problem	Requires tuning of α	Variants of CNNs, DNNs

Table 10.1: Comparison of activation functions.

[SI*] Backpropagation

Backward propagation of errors (backpropagation) is a key algorithm used to train neural networks by minimizing the difference between predicted and actual outputs. It works by propagating errors backward through the network, using the chain rule of calculus to compute gradients and then iteratively updating the weights and biases. Combined with optimization techniques like gradient descent, backpropagation enables the model to reduce loss across epochs and effectively learn complex patterns from data.

Mathematically, "training" means minimizing an error function, the sum of the squares of the errors in the outputs. Backpropagation corresponds to using a steepest-descent method for minimizing the error function. While useful for training networks, backpropagation has the

disadvantages that convergence is slow compared to Conjugate Gradient.

Minimization of a function of many variables is a mature field; there are numerous algorithms and more numerous variations of them. All are iterative, generating a sequence of sets of weights.

Backpropagation plays a critical role in how neural networks improve over time.

1. **Efficient Weight Update**

It computes the gradient of the loss function with respect to each weight using the chain rule making it possible to update weights efficiently.

2. **Scalability**

The backpropagation algorithm scales well to networks with multiple layers and complex architectures making deep learning feasible.

3. **Automated Learning**

With backpropagation the learning process becomes automated and the model can adjust itself to optimize its performance.

The working of backpropagation:

1. **Forward Pass Work**

In forward pass the input data is fed into the input layer. These inputs combined with their respective weights are passed to hidden layers. For example in a network with two hidden layers (h_1 and h_2) the output from h_1 serves as the input to h_2 . Before applying an activation function, a bias is added to the weighted inputs.

Each hidden layer computes the weighted sum of the inputs ($\sum_{i=1}^n x_i w_i$) then applies an activation function like ReLU (Rectified Linear Unit) to obtain the output. The output is passed to the next layer till reaches the output layer where an activation function such as softmax converts the weighted outputs into probabilities for classification.

2. **Error Calculation**

The network's prediction is compared against the actual target value using a loss function (e.g., Mean Squared Error) to determine the "cost" or error.

3. **Backward Pass (Backpropagation)**

The algorithm calculates the gradient of the loss function with respect to every weight in the network. It uses the chain rule to propagate the error from the output layer back to the input layer.

In the backward pass the error (the difference between the predicted and actual output) is propagated back through the network to adjust the weights and biases. One common method for error calculation is the Mean Squared Error given by

$$MSE = (y_i - \hat{y}_i)^2$$

with y_i as the actual output, $\hat{y}_i$ is the predicted output, $i = 1, 2, \dots, n$ and n is the number of classification of the outputs / possible outcome for the output.

Once the error is calculated the network adjust weights using gradients which are computed with the chain rule. These gradients indicate how much each weight and bias should be adjusted to minimize the error in the next iteration. The backward pass continues layer by layer ensuring that the network learns and improves its performance. The activation function through its derivative plays a crucial role in computing these gradients during backpropagation.

4. Weight Update

An optimization algorithm, usually a gradient descent, adjusts the weights by adding or subtracting (depends on the loss function that is used) a portion of the gradient (scaled by a learning rate) to minimize the error

[SI*] Gradient in Backpropagation

In a neural network, the gradient is the partial derivative of the loss function (error) with respect to each individual weight and bias using the multivariable chain rule. We use the gradient descent algorithm to find the best parameters that minimize the error. Gradient represents the direction and magnitude of the change needed to reduce the model's prediction error.

The algorithm iteratively moves, step by step, in the direction of steepest descent, toward the minimum point, the step size is the learning rate (η). During propagation, the network calculates how much each weight contributed to the total error.

- **Direction**

The sign of the gradient tells you whether to increase or decrease the weight to reduce loss.

- **Steepness**

The magnitude (absolute value) of the gradient indicates how sensitive the error is to that specific weight.

After the gradient is calculated, weights are updated using an optimization algorithm, most commonly Gradient Descent. The standard weight update formula for layer L using the gradient descent rule is

$$w_{new}^{(L)} = w_{old}^{(L)} - (\eta \times \frac{\partial C}{\partial w^{(L)}})$$

with η as the learning rate, it is a positive scalar, we can use fraction of real number like 0.01, remember that learning rate controls the step size. Gradient is denoted by $\frac{\partial C}{\partial w}$, C is the cost function, it is calculated using the chain rule to propagate error, we start from the output layer back to the input layer.

The gradient for a weight connecting neuron i to neuron j is typically computed as

$$\text{Weight Gradient : } \frac{\partial C}{\partial w_{ij}^{(L)}} = \delta_j^{(L)} \times y_i^{(L-1)}$$

and for the bias gradient at layer L

$$\text{Bias Gradient : } \frac{\partial C}{\partial b_j^{(L)}} = \delta_j^{(L)}$$

$$z_j^{(L)} = \sum_i w_{ij}^{(L)} y_i^{(L-1)} + b_j^{(L)}$$

$$y_j^{(L)} = \sigma(z_j^{(L)})$$

We compute the δ from the output layer with this formula

$$\delta_j^{(L)} = \frac{\partial C}{\partial y_j^{(L)}} \oplus \sigma'(z_j^{(L)})$$

$\oplus$ represents the hadamard product, $\delta_j^{(L)}$ is the error term that represents the error of the receiving neuron (neuron j at layer L), which includes the derivative of its activation function, $\sigma'(z_j^{(L)})$. $y_i^{(L-1)}$ is the post-activation output value of the sending neuron (neuron i at layer $L - 1$), while $y_j^{(L)}$ is the post-activation output value of the receiving neuron (neuron j at layer L).

For any previous hidden layer ($l < L$), we calculate the gradients by propagating the error backwards from the subsequent layer ($l + 1$), thus we compute δ for the hidden layer/s with this formula

$$\delta_i^{(l)} = \frac{\partial C}{\partial y_i^{(l)}} \oplus \sigma'(z_i^{(l)}) = \sum_j \left(w_{ij}^{(l+1)} \delta_j^{(l+1)} \right) \oplus \sigma'(z_i^{(l)})$$

and the formula to compute the weight and bias gradient connecting neuron k to neuron i to neuron j for hidden layer/s to input layer

$$\text{Weight Gradient : } \frac{\partial C}{\partial w_{ki}^{(l)}} = \delta_i^{(l)} \times y_k^{(l-1)}$$

$$\text{Bias Gradient : } \frac{\partial C}{\partial b_i^{(l)}} = \delta_i^{(l)}$$

with $\delta_i^{(l)}$ as the error term that represents the error of the neuron i at layer l , which includes the derivative of its activation function, $\sigma'(z_i^{(l)})$. $y_k^{(l-1)}$ is the post-activation output value of the sending neuron (neuron k at layer $l - 1$), while $y_i^{(l)}$ is the post-activation output value of the receiving neuron (neuron i at layer l). If l represents the first hidden layer then $l - 1$ is the input layer and $y_k^{(l-1)}$ is the k -th input variable.

In a neural network, the "delta" (δ) represents the error gradient with respect to the pre-activation (logit) of a layer.

If gradients become too small, the network suffers from the vanishing gradient problem, where early layers stop learning because their weights are no longer updated.

The delta output layer of a feed-forward neural network uses the post-activation neuron value to calculate the error. We start from the output layer to compute delta.

Calculate Post-Activation Output (y_j)

First we calculate the weighted sum (pre-activation, z_j) and then apply the activation function σ :

$$z_j = \sum_i (w_{ij} \cdot x_i) + b_j$$

$$y_j = \sigma(z_j)$$

Calculate the Error (e_j)

After forward pass, determine the error of the final output using a loss function, the first one is the mean squared error that uses the error formula as the difference between the actual post-activation output y_j and the actual output t_j :

$$e_j = t_j - y_j$$

thus

$$L = (t_j - y_j)^2$$

The second one is Categorical Cross-Entropy, if the output layer uses softmax as the activation function (e.g., multi-class problem) then the loss function / error is

$$L = - \sum_{j=1}^C t_j \log(y_j)$$

with C as the total number of classes, t_j is the true value (either 1 or 0 for one-hot encoded labels), y_j is the predicted probability for j -th class. For this loss function because the true values are 0 for incorrect classes, the formula simplifies to the negative natural logarithm of the predicted probability for the correct class, so if the prediction (y_j value) is high then the loss is low, vice versa.

Formula to Compute the Output Delta (δ_j)

The delta at the output layer is the first delta we compute in backpropagation, and it is computed by multiplying the error by the derivative of the activation function with respect to the pre-activation sum z_j :

$$\delta_j^{Output\ Layer} = (t_j - y_j) \cdot \sigma'(z_j)$$

The pre-activation (z_j) is needed for calculating the derivative of the activation function $\sigma'(z_j)$, which adjusts the gradient to account for the specific shape of the activation curve. The post-activation output (y_j) is essential for calculating the actual error signal because it represents the network's final prediction.

Afterwards, the error term for a hidden unit is calculated by backpropagating the error from the next layer. This process relies on the chain rule to determine how much a hidden unit contributed to the total error at the output layer.

Formula to Compute the Hidden layer' Delta (δ_j)

For a hidden unit j in layer L (assuming L is not the output layer), the neuron-specific delta δ_j^L for hidden neuron j relies on the deltas of all the neurons in the next layer that it connects to. It is defined mathematically as the derivative of the activation function multiplied by the weighted sum of the subsequent layer's deltas, and it is defined by the following formula:

$$\delta_j^L = \sigma'(z_j^L) \sum_k \left(w_{kj}^{L+1} \cdot \delta_k^{L+1} \right) \quad (10.10)$$

with z_j^L as the pre-activation of the hidden neuron j at layer L , δ_k^{L+1} as the error term of unit k in the next layer ($L + 1$), w_{kj}^{L+1} as the weight connecting the hidden unit j (current layer) to unit k (next layer), $\sigma'(z_j^L)$ as the derivative of the activation function evaluated at the pre-activation, z_j^L , and $\sum_k \left(w_{kj}^{L+1} \cdot \delta_k^{L+1} \right)$ is a summation over all neurons k in the subsequent layer that receive input from neuron j , we can also say that this is the backpropagated error signal from the next layer. Remember that the pre-activation, z_j , is the weighted sum from the previous layer plus the bias

$$z_j^L = \sum_i (w_{ij}^{L-1} \cdot x_i^{L-1}) + b_j^L$$

x_i^{L-1} is the neurons of the previous layer.

The post-activation value of the hidden neuron is used as an input to the weights of the next layer. Therefore, it dictates how much influence neuron j has on the network's final error. While the pre-activation and derivative $\sigma'(z_j)$ evaluates how sensitive the hidden neuron's output is to changes in its pre-activation. Multiplying by this derivative ensures the gradient descent updates only adjust weights in a way that respects the non-linear boundaries of the activation function.

[SI*] **Train Data and Test Data**

Training data is used to teach a model, while test data is used to evaluate its performance on unseen information, ensuring it generalizes than just memorizing. Training data is generally larger (e.g., 80%), used for fitting model parameters, whereas test data (e.g., 20%) remains isolated until final assessment.

Training data teaches the model patterns, weights, or features, it happens first. Test data checks the model's accuracy on new data, preventing overfitting, it is the final step to determine model reliability.

Basically the whole dataset will be divided into training data with large portion and into testing data with smaller portion (80/20 or 70/30).

[SI*] **Epoch in a Neural Network**

An epoch in a neural network represents one complete, single pass of the entire training dataset through the algorithm-both forward and backward. It is a hyperparameter determining how many times the model updates its internal weights using all training samples. Multiple epochs are usually required to minimize error and improve learning.

The different from batch, is that a batch is a subset of the dataset used to update weights, an iteration is a one update step using a single batch. The relationship formula is

$$1 \text{ Epoch} = \frac{\text{Total Dataset Size}}{\text{Batch Size}} \text{ iterations}$$

An iteration here refers to a single step of training where the model processes one batch of data and updates its parameters (weights and biases). Each iteration includes two main parts:

1. **Forward Propagation**

The input data in the batch is passed through the network to generate predictions, and the loss (error) is calculated.

2. **Backward Propagation**

The gradients are calculated, and the model's parameters are updated to reduce the loss.

We usually use multiple epochs because a single epoch is rarely sufficient to capture complex patterns, so the model re-reads the data, often updating weights after every batch to refine its understanding. Too many epochs can cause the model to learn the noise in the training data rather than general patterns, leading to high training accuracy but low validation accuracy (overfitting). The optimal number of epochs is the balance between overfitting and underfitting. An example is if you have 60,000 images of handwriting for digits 0-9 and a batch size of 500, it takes 120 iterations $\left(\frac{60,000}{500}\right)$ to complete one epoch.

[SI*] Confusion Matrix

A confusion matrix for a neural network is a table used to evaluate classification model performance by comparing predicted labels against true labels. It highlights True Positives (TP), True Negatives (TN), False Positives (FP), and False Negatives (FN), enabling identification of misclassifications, particularly for binary or multi-class classification problems.

The diagonal elements of the confusion matrix represent correct predictions (actual=predicted) while off-diagonal elements indicate errors.

The formula to compute the accuracy of the neural network is

$$\frac{TP + TN}{TP + TN + FP + FN}$$

The formula to compute the precision of the neural network is

$$\frac{TP}{TP + FP}$$

The formula to compute the recall/sensitivity of the neural network is

$$\frac{TP}{TP + FN}$$

The formula to compute the F1-score we need to compute the harmonic mean of precision and recall.

Table (10.2) is an example of a model analyzing 100 Kohanku Koi fishes:

- Actual Aeromonas (the number of fishes that is infected with Aeromonas Hydrophilia): 50
- Actual Not Aeromonas (the number of fishes that is not infected with Aeromonas Hydrophilia): 50

	Predicted: Aeromonas	Predicted: Not Aeromonas
Actual: Aeromonas	40(TP)	10(FN)
Actual: Not Aeromonas	5(FP)	45(TN)

Table 10.2: Binary confusion matrix example.

The confusion matrix is essential for unbalanced datasets (e.g., detecting a disease where 99% of fishes are healthy), as it reveals if the model is ignoring the minor class, which high accuracy alone might hide.

[SI*] Conjugate Gradient in Backpropagation

Previously we are talking about steepest descent / gradient descent method for finding the optimal weights in neural network, there is another method called Conjugate Gradient and it is known to train the neural networks faster than the steepest descent method. The Conjugate Gradient is a line-search optimization technique that improves upon steepest descent by generating conjugate search directions, reducing the "zigzagging" behavior and ensuring faster convergence. In the context of training a neural network using the Conjugate Gradient (CG) method, the algorithm minimizes the error (cost function) by treating it as

a local quadratic approximation. It solves for the optimal weights using the linear system $Ax = b$.

1. Vector x (The Weights)

This is the vector of all trainable parameters (weights and biases) in the neural network, flattened into a single one-dimensional vector, typically denoted as w or θ in literature. The goal is to find the optimal x that minimizes the error function $E(x)$.

2. Matrix A (The Hessian)

The Hessian matrix (H or $\nabla^2 E$) of the neural network's cost function, which contains all partial derivatives of the second order (second-order curvature). It describes the shape of the error surface, determining how the gradient changes. For conjugate gradient to work ideally, A must be symmetric and positive-definite (locally true near a minimum). In neural network training, the full Hessian is rarely explicitly computed due to its massive size ($N \times N$, where N is the number of parameters). Instead, the algorithm uses products of the Hessian with a vector, such as $Ap \approx \nabla^2 \cdot p$.

3. Vector b (The Gradient-related term)

A vector related to the gradient of the error function, frequently derived from the gradient $\nabla E(x)$.

If using a pure linear system $Ax = b$ for a quadratic error, b is a constant vector. In nonlinear optimization (neural network training), this represents components of the gradient, and the residual $r = b - Ax$ is replaced by the negative gradient of the error function ($-\nabla E(x)$).

Since neural network error surfaces are not strictly quadratic, non-linear conjugate gradient methods are used instead of solving the strict linear system.

Instead of calculating A (the Hessian), these algorithms compute the new search direction d_k as a linear combination of the current negative gradient and the previous search direction.

$$d_k = -\nabla E(k) + \beta_k d_{k-1}$$

where β_k is a scalar parameter, from now on we will denote $\nabla E(k)$ as g_k

$$\beta_k^{PRP} = \frac{g_k^T (g_k - g_{k-1})}{\|g_{k-1}\|^2}, \quad (\text{Polak-Ribiere})$$

if $\beta_k^{PRP} < 0$, it is often set to 0 to restart the algorithm, which helps avoid slow convergence. The Polak-Ribiere is very useful for unconstrained optimization and neural network training. It is often more efficient than Fletcher-Reeves method, because it has automatic restart that avoid jamming while optimizing.

$$\beta_k^{FR} = \frac{g_k^T g_k}{\|g_{k-1}\|^2}, \quad (\text{Fletcher-Reeves})$$

$$\beta_k^{DY} = \frac{g_k^T g_k}{d_{k-1}^T (g_k - g_{k-1})}, \quad (\text{Dai-Yuan})$$

$$\beta_k^{CD} = -\frac{g_k^T g_k}{d_{k-1}^T g_{k-1}}, \quad (\text{Conjugate Descent})$$

The Conjugate Gradient loop / algorithm:

1. Initialize weights and biases randomly.
2. Calculate the gradient $\frac{\partial E}{\partial w}$ for all weights and bias, concatenated into a single vector $\mathbf{g}$. The formula is

$$\mathbf{g}_k = \nabla f(\mathbf{w}_k)$$

it is calculated using backpropagation on the feed-forward neural network, where f is the loss function.

3. Initialize $\mathbf{d}_0 = -\mathbf{g}_0$, the first step is steepest descent.
4. Iterate:

- Find optimal η_k / the learning rate with strong Wolfe line search. The line search method determines the optimal step size (η_k) along the search direction ($\mathbf{d}_k$) to minimize loss function $f(\mathbf{w}_k + \eta_k \mathbf{d}_k)$ and it is crucial for ensuring stability. It is determined by minimizing $f(\mathbf{w}_k + \eta_k \mathbf{d}_k)$.

$$f(\mathbf{w}_k + \eta_k \mathbf{d}_k) = \min_{\eta \geq 0} f(\mathbf{w}_k + \eta \mathbf{d}_k)$$

it means that the orthogonality condition

$$\mathbf{g}_k^T \mathbf{d}_{k-1} = 0$$

is satisfied, where $\mathbf{g}_k$ represents the gradient of the objective function f at the value $\mathbf{w}_k$. Since it is difficult in practice to compute η_k , the inexact line search is introduced to compute approximate values for η_k , thus we will use strong Wolfe line search. The strong Wolfe conditions requires η_k to satisfy the following two conditions [21]:

$$\begin{aligned} f(\mathbf{w}_k + \eta_k \mathbf{d}_k) &\leq f(\mathbf{w}_k) + \delta \eta_k \mathbf{g}_k^T \mathbf{d}_k \quad (\text{sufficient decrease}) \\ |g(\mathbf{w}_k + \eta_k \mathbf{d}_k)^T \mathbf{d}_k| &\leq \sigma |\mathbf{g}_k^T \mathbf{d}_k| \quad (\text{strong curvature condition}) \end{aligned}$$

where $0 < \delta < \sigma < 1$, and $\mathbf{d}_k$ is the search direction, which is given by

$$\mathbf{d}_k = \begin{cases} -\mathbf{g}_k, & \text{if } k = 0 \\ -\mathbf{g}_k + \beta_k \mathbf{d}_{k-1}, & \text{if } k \geq 1 \end{cases}$$

The factor β_k determines how the CG methods differ, it is calculated by the algorithm to make $\mathbf{d}_k$ and $\mathbf{d}_{k-1}$ conjugate, or orthogonal. The β_k can be an indication if there is numerical instability or a poor search direction / small steps (the new gradient $\mathbf{g}_k$ almost similar to the old gradient $\mathbf{g}_{k-1}$) occurs while iterating, if β_k produces large values then there might be something wrong and we have to change our β_k formula since it will cause the training to fail or stall. Periodically resetting β_k to zero ensures the algorithm does not stay trapped in a poor, large-gradient region, this is called reverting to the steepest descent direction ($\mathbf{d}_{k=-\mathbf{g}_k}$).

- Update weights

$$\mathbf{w}_{k+1} = \mathbf{w}_k + \eta_k \mathbf{d}_k$$

- Calculate new gradient $\mathbf{g}_{k+1}$.
- Compute update coefficient β_k (e.g. with Polak-Ribiere or Fletcher-Reeves).
- Calculate new direction

$$\mathbf{d}_{k+1} = -\mathbf{g}_{k+1} + \beta_k \mathbf{d}_k, \quad k = 0, 1, \dots$$

5. Run until maximum iterations / last epoch or target error is reached.

Note: Moller uses a (temporary) small value for η_k , does a function evaluation in order to approximate the second derivative in the search direction, then uses the approximate second derivative to select a final η_k ; this uses two function calls. (If the new weights result in an increased error, they are rejected and some parameters are adjusted until the new weights reduce the error. In this case, a successful iteration takes more than two function calls. [13]).

Instead of backtracking like gradient descent, Conjugate Gradient creates "conjugate" directions that do not "undo" previous minimization steps, resulting in faster convergence. And unlike Newton's method, Conjugate Gradient does not require storing the Hessian matrix, making it efficient for large-scale feed-forward neural networks.

When the conjugate gradient method in a neural network stalls, it is typically due to a stuck search direction, a suboptimal line search, or a vanishing gradient problem. Unlike gradient descent, conjugate gradient needs precise line searches, is highly sensitive to the line search algorithm used to calculate the learning rate / step size η_k . If the loss is flat, it stops converging.

When a feed-forward neural network trained with a conjugate gradient algorithm consistently predicts only one class (for multi-class problem), it usually means the network is trapped in a local minimum, the learning rate is mismatched (check the line search algorithm, you should use the Wolfe conditions, if the learning rate is too large the CG method will diverge or get stuck, if it is too small the CG algorithm will make negligible progress and might just predict the majority class), or the architecture is incorrectly configured for multi-class classification.

To fix the local minimum trap, we need to make sure that the output layer is using softmax as the activation function and the loss function being used is categorical cross-entropy rather than binary cross-entropy. Don't forget for multi-class problem we need to use one-hot encoding for the actual output. Avoid using sigmoid activation function, which causes vanishing gradients, use ReLU activation function instead in the hidden layers.

Most common causes and solutions for conjugate gradient stagnation:

- **Invalid or Inaccurate Line Search**

Conjugate gradient relies on finding an optimal step size (η_k) to move along the conjugate direction (d_k). If the line search method (e.g. backtracking, secant) fails to find a good η_k , the algorithm stalls. If the step size is too large, the "conjugate" property of the search directions is lost, and the algorithm may oscillate or diverge just like standard stochastic gradient descent.

To fix this, use a more robust line search method that satisfies the Wolfe conditions to ensure sufficient decrease.

- **Stuck in Saddle Points**

The optimizer may have reached a saddle point where the gradient is zero or near-zero, but it is not a local minimum.

Normalize the inputs for weights and bias to each layer to have a mean of 0 and variance of 1. This keeps gradients in a healthy range and speeds up convergence.

- **Vanishing Gradients**

Deep networks can cause gradients to become negligible, resulting in tiny or no weight updates.

If the network is very deep, utilize residuals (skip connections) to mitigate vanishing gradients or avoid using sigmoid or tanh activation function and used LeakyReLU or ReLU instead.

- **Loss of Conjugacy**

Over many iterations, or due to non-convexity, the directions might lose their conjugacy. Resetting the conjugate gradient direction to the negative gradient (steepest descent) periodically can resolve this.

Reset the search direction to the steepest descent direction every N iterations (N is typically the number of parameters, the weights and bias).

- **Oscillating Loss**

An oscillating loss during epoch training with a conjugate gradient optimizer and ReLU activation function is usually caused by line search failures, non-convex gradient discontinuities (dying ReLUs), or an excessive learning rate.

Conjugate gradient heavily relies on exact or inexact line searches to determine the optimal step size. Because ReLU introduces sharp, non-convex boundaries, the function is piecewise linear. Standard line searches often fail to bracket the minimum, causing erratic jumps.

Ensure the line search algorithm satisfies the strong Wolfe conditions. If already does, tighten the tolerance parameters (e.g., lower the σ parameter in the curvature condition) to prevent the CG method from overshooting the optimal step in the parameter space.

The standard ReLU derivative is discontinuous. When a large batch of neurons "dies" (outputs 0), the gradient drops abruptly, causing the CG direction vectors to break orthogonality. The optimizer then "bounces" back and forth across these kinks. Use leaky ReLU or GELU instead to solve this. They provide a continuous, non-zero gradient across negative inputs, smoothing out the loss landscape and stabilizing the conjugate search directions.

If the initial learning rate or step bound is too large, CG will step outside the valid quadratic approximation of the loss surface. It then tries to correct itself backward on the next step, resulting in a zig-zagging or oscillating epoch loss.

Manually decrease the learning rate or the maximum step size. Test a value 10 times to 100 times smaller than the current setup to see if the oscillation smooths out.

- **Improper Initialization**

Poorly initialized weights and bias may land the model in a flat region of the loss landscape immediately.

Use Xavier/Glorot if you use tanh or sigmoid activation function or He initialization if you use ReLU activation function to ensure gradients are healthy.

The conjugate gradient that we are talking here is the nonlinear conjugate gradient that use inexact line search (strong Wolfe conditions), it is best for large-scale neural network training where exact search is too slow. Another type of conjugate gradient is scaled conjugate gradient with Levenberg-Marquardt approach to compute the step size, it is highly efficient for neural network training without manual tuning.

We use conjugate gradient when we have small to medium-size networks where rapid convergence is needed and the loss function is well-behaved (not extremely noisy). If we have very large datasets where calculating exact gradient at every step is too costly, then we

use steepest descent.

Applications of conjugate gradient:

1. Solving large sparse linear systems in computational physics and engineering.
2. Image reconstruction for medical imaging (CT scan, MRI).
3. Neural network training.
4. Structural analysis with finite element methods.
5. Computational fluid dynamics.

Moreover, their favorable properties, such as global convergence and low memory requirements, have qualified them to solve problems in diverse areas of science, such as data estimation, image restoration, and signal processing.

II. DATAFRAME IN C++ FOR NEURAL NETWORKS

[SI*] Neural networks have become a vast field with many applications in supervised machine learning. Supervised learning means that the input cases and the output cases are known and that their functional relationship is to be learned while the data may be noisy.

As the first stepping stone, we will need to be able to load the data, in this book we will use a simple `.csv` file that separates the column with comma (",") and the row with line break. The data in the `.csv` file is what we call dataframe. The popular programming language like Python, JULIA and R have packages to handle dataframe easily, with C++ we are not only gaining more speed, but also more power and flexibility that are future-proof for the future Artificial Intelligence, Machine Learning, and beyond this.

i. Loading Dataframe with Iris Dataset

Suppose we have a CSV-formatted data, with name of the file as "iris.csv" that classify iris flowers, this is the iris dataset. It contains 150 samples of iris flowers, with 4 features each

- Sepal length (in cm)
- Sepal width (in cm)
- Petal length (in cm)
- Petal width (in cm)

Each sample is labeled with one of three species

- Iris Setosa
- Iris Versicolor
- Iris Virginica

```
5.1,3.5,1.4,.2,"Setosa"  
4.9,3,1.4,.2,"Setosa"  
4.7,3.2,1.3,.2,"Setosa"  
4.6,3.1,1.5,.2,"Setosa"  
5,3.6,1.4,.2,"Setosa"  
5.4,3.9,1.7,.4,"Setosa"  
4.6,3.4,1.4,.3,"Setosa"  
5,3.4,1.5,.2,"Setosa"  
4.4,2.9,1.4,.2,"Setosa"  
4.9,3.1,1.5,.1,"Setosa"  
5.4,3.7,1.5,.2,"Setosa"  
4.8,3.4,1.6,.2,"Setosa"  
4.8,3,1.4,.1,"Setosa"  
4.3,3,1.1,.1,"Setosa"  
5.8,4,1.2,.2,"Setosa"  
5.7,4.4,1.5,.4,"Setosa"  
5.4,3.9,1.3,.4,"Setosa"  
5.1,3.5,1.4,.3,"Setosa"  
5.7,3.8,1.7,.3,"Setosa"  
5.1,3.8,1.5,.3,"Setosa"  
5.4,3.4,1.7,.2,"Setosa"
```

```

5.1,3.7,1.5,.4,"Setosa"
4.6,3.6,1,.2,"Setosa"
5.1,3.3,1.7,.5,"Setosa"
4.8,3.4,1.9,.2,"Setosa"
5,3,1.6,.2,"Setosa"
...
...
6.7,3.1,5.6,2.4,"Virginica"
6.9,3.1,5.1,2.3,"Virginica"
5.8,2.7,5.1,1.9,"Virginica"
6.8,3.2,5.9,2.3,"Virginica"
6.7,3.3,5.7,2.5,"Virginica"
6.7,3,5.2,2.3,"Virginica"
6.3,2.5,5,1.9,"Virginica"
6.5,3,5.2,2,"Virginica"
6.2,3.4,5.4,2.3,"Virginica"
5.9,3,5.1,1.8,"Virginica"

```

Code 61: *iris.csv*

From the data "iris.csv" above we can categorize the variables into 5 different variables / 5 columns of a matrix. We will have the variables in column vector-term like this:

$$\vec{x}_1 \quad \vec{x}_2 \quad \vec{x}_3 \quad \vec{x}_4 \quad \vec{x}_5$$

and

```

x1 = sepal_length
x2 = sepal_width
x3 = petal_length
x4 = petal_width
x5 = species

```

The variables x_1, x_2, x_3 , and x_4 are defined as **double** data type in C++, while the variable x_5 is defined as **string** data type.

The code below shows how to load a CSV with variables of these:

```

#include <iostream>
#include "symintegrationc++.h"
#include <bits/stdc++.h>
#include <cmath>
#include <chrono>

using namespace std::chrono;
using namespace std;
using namespace SymbolicConstant;

#include <iostream>
#include <fstream>

```

```
#include <vector>
#include <string>
#include <sstream>

struct IrisRecord
{
    double sepal_length, sepal_width, petal_length, petal_width;
    std::string species;
};

std::vector<IrisRecord> loadIris(const std::string& filename)
{
    std::vector<IrisRecord> data;
    std::ifstream file(filename);
    std::string line;

    while (std::getline(file, line))
    {
        if (line.empty()) continue;
        std::stringstream ss(line);
        std::string val;
        IrisRecord record;

        std::getline(ss, val, ','); record.sepal_length = std::stod(
            val);
        std::getline(ss, val, ','); record.sepal_width = std::stod(
            val);
        std::getline(ss, val, ','); record.petal_length = std::stod(
            val);
        std::getline(ss, val, ','); record.petal_width = std::stod(
            val);
        std::getline(ss, record.species, ',');

        data.push_back(record);
    }
    return data;
}

void showIrisVector(const std::vector<IrisRecord>& data)
{
    cout << "Species \t | Sepal length " << setw(10) << "| Sepal width "
        << setw(10) << "| Petal length " << setw(10) << "| Petal
        width " << endl;
    for (const auto& record : data)
    {
        cout << record.species
            << " \t | \t " << record.sepal_length << " \t | \t " <<
            record.sepal_width
```

```

        << " \t | \t " << record.petal_length << " \t | \t" << record
            .petal_width
        << endl;
    }
}

// Driver code
int main(int argc, char** argv)
{
    // Get starting timepoint
    auto start = high_resolution_clock::now();

    std::vector<IrisRecord> data = loadIris("iris.csv");

    int n = data.size();
    showIrisVector(data);
    cout << "\nNumber of data: " << n << endl;

    // Get ending timepoint
    auto stop = high_resolution_clock::now();
    auto duration = duration_cast<microseconds>(stop - start);

    cout << "\nTime taken by function: " << duration.count() << "
        microseconds" << endl;

    return 0;
}

```

Code 62: *Neural Networks/Test Iris DataFrame/main.cpp*

The basic needs when coding this is we know that each column represents which variable (and each variable is assigned to a certain suitable data type available in C++), we don't want to interchange the variable of petal length with sepal width, so we need to be sure which column represents which variable.

If we are able to load the data that contain not only data types of double, string, char, datetime, then we can then process it further with machine learning algorithms to recognize a pattern, or gain knowledge of the data.

- To "show" or print the contents of a `std::vector<IrisRecord>`, use a range-based **for** loop, which is the most concise method in modern C++.
- The most common method uses `std::ifstream` and `std::getline` to read the CSV-formatted data into vectors or arrays.

II. DataFrame in C++ for Neural Networks

```
root [ ~/SourceCodes/CPP/C++ Create Library/Test Iris DataFrame ]# ./main
Species      | Sepal length | Sepal width | Petal length | Petal width
"Setosa"      | 5.1          | 3.5         | 1.4         | 0.2
"Setosa"      | 4.9          | 3           | 1.4         | 0.2
"Setosa"      | 4.7          | 3.2         | 1.3         | 0.2
"Setosa"      | 4.6          | 3.1         | 1.5         | 0.2
"Setosa"      | 5           | 3.6         | 1.4         | 0.2
"Setosa"      | 5.4          | 3.9         | 1.7         | 0.4
"Setosa"      | 4.6          | 3.4         | 1.4         | 0.3
"Setosa"      | 5           | 3.4         | 1.5         | 0.2
"Setosa"      | 4.4          | 2.9         | 1.4         | 0.2
"Setosa"      | 4.9          | 3.1         | 1.5         | 0.1
"Setosa"      | 5.4          | 3.7         | 1.5         | 0.2
"Setosa"      | 4.8          | 3.4         | 1.6         | 0.2
"Setosa"      | 4.8          | 3           | 1.4         | 0.1
"Setosa"      | 4.3          | 3           | 1.1         | 0.1
"Setosa"      | 5.8          | 4           | 1.2         | 0.2
"Setosa"      | 5.7          | 4.4         | 1.5         | 0.4
"Setosa"      | 5.4          | 3.9         | 1.3         | 0.4
"Setosa"      | 5.1          | 3.5         | 1.4         | 0.3
"Setosa"      | 5.7          | 3.8         | 1.7         | 0.3
"Setosa"      | 5.1          | 3.8         | 1.5         | 0.3
"Setosa"      | 5.4          | 3.4         | 1.7         | 0.2
"Setosa"      | 5.1          | 3.7         | 1.5         | 0.4
"Setosa"      | 4.6          | 3.6         | 1           | 0.2
"Setosa"      | 5.1          | 3.3         | 1.7         | 0.5
"Setosa"      | 4.8          | 3.4         | 1.9         | 0.2
"Setosa"      | 5           | 3           | 1.6         | 0.2
"Setosa"      | 5           | 3.4         | 1.6         | 0.4
"Setosa"      | 5.2          | 3.5         | 1.5         | 0.2
"Setosa"      | 5.2          | 3.4         | 1.4         | 0.2
"Setosa"      | 4.7          | 3.2         | 1.6         | 0.2
"Setosa"      | 4.8          | 3.1         | 1.6         | 0.2
"Setosa"      | 5.4          | 3.4         | 1.5         | 0.4
"Setosa"      | 5.2          | 4.1         | 1.5         | 0.1
```

Figure 10.2: The loading of "iris.csv" data and show it as a dataframe (SymIntegration/Examples/Neural Networks/Test Iris DataFrame/main.cpp).

```
"Virginica"    | 6           | 2.2         | 5           | 1.5
"Virginica"    | 6.9         | 3.2         | 5.7         | 2.3
"Virginica"    | 5.6         | 2.8         | 4.9         | 2
"Virginica"    | 7.7         | 2.8         | 6.7         | 2
"Virginica"    | 6.3         | 2.7         | 4.9         | 1.8
"Virginica"    | 6.7         | 3.3         | 5.7         | 2.1
"Virginica"    | 7.2         | 3.2         | 6           | 1.8
"Virginica"    | 6.2         | 2.8         | 4.8         | 1.8
"Virginica"    | 6.1         | 3           | 4.9         | 1.8
"Virginica"    | 6.4         | 2.8         | 5.6         | 2.1
"Virginica"    | 7.2         | 3           | 5.8         | 1.6
"Virginica"    | 7.4         | 2.8         | 6.1         | 1.9
"Virginica"    | 7.9         | 3.8         | 6.4         | 2
"Virginica"    | 6.4         | 2.8         | 5.6         | 2.2
"Virginica"    | 6.3         | 2.8         | 5.1         | 1.5
"Virginica"    | 6.1         | 2.6         | 5.6         | 1.4
"Virginica"    | 7.7         | 3           | 6.1         | 2.3
"Virginica"    | 6.3         | 3.4         | 5.6         | 2.4
"Virginica"    | 6.4         | 3.1         | 5.5         | 1.8
"Virginica"    | 6           | 3           | 4.8         | 1.8
"Virginica"    | 6.9         | 3.1         | 5.4         | 2.1
"Virginica"    | 6.7         | 3.1         | 5.6         | 2.4
"Virginica"    | 6.9         | 3.1         | 5.1         | 2.3
"Virginica"    | 5.8         | 2.7         | 5.1         | 1.9
"Virginica"    | 6.8         | 3.2         | 5.9         | 2.3
"Virginica"    | 6.7         | 3.3         | 5.7         | 2.5
"Virginica"    | 6.7         | 3           | 5.2         | 2.3
"Virginica"    | 6.3         | 2.5         | 5           | 1.9
"Virginica"    | 6.5         | 3           | 5.2         | 2
"Virginica"    | 6.2         | 3.4         | 5.4         | 2.3
"Virginica"    | 5.9         | 3           | 5.1         | 1.8

Number of data: 150
Time taken by function: 6610 microseconds
```

Figure 10.3: The loading of "iris.csv" data and show it as a dataframe (SymIntegration/Examples/Neural Networks/Test Iris DataFrame/main.cpp).

III. SUPERVISED LEARNING: FEED-FORWARD NEURAL NETWORK (MULTILAYER PERCEPTRON)

[SI*] This is the first category of machine learning, the supervised learning which uses labeled data to train models that can predict outcomes for new data (e.g., classification or regression). A feed-forward neural network (FNN) is the simplest type of artificial neural networks where information moves in only one direction (forward flow) from input nodes, through hidden nodes (if any), to output nodes. It has no loops or cycles, the data never moves backward. It is used for pattern recognition, classification, and regression tasks like credit scoring or speech recognition. Other notable examples of the FNN usage are image recognition, financial forecasting, data transformation.

The feed-forward neural network has a structured layered design where data flows sequentially through each layer.

1. Input Layer

The input layer consists of neurons that receive the input data. Each neuron in the input layer represents a feature of the input data.

2. Hidden Layers

One or more hidden layers are placed between the input and output layers. These layers are responsible for learning the complex patterns in the data. Each neuron in a hidden layer applies a weighted sum of inputs followed by a non-linear activation function.

3. Output Layer

The output layer provides the final output of the network. The number of neurons in this layer corresponds to the number of classes in a classification problem or the number of outputs in a regression problem.

For classification problem, we use Softmax as the activation function from the output layer to determine the classification result.

In FNN, the neurons are connected by the corresponding weights, and non-linear activation functions (like Sigmoid or ReLU) are used in hidden layers. The weights are updated / adjusted to reduce prediction error using techniques like gradient descent, often through backpropagation. The process for training a feed-forward neural network:

1. Forward Propagation

During forward propagation the input data passes through the network and the output is calculated to generate a prediction. For classification, the output layer typically produces a set of scores or probabilities that represent the likelihood of the input belonging to specific classes.

The core mechanism of forward propagation:

- **Weighted Sum (z)**

We define the weighted sum with the notation of z and the formula is

$$z = W^L a^{(L-1)} + b^{(L)}$$

with z as the weighted sum that is already added with bias b at layer L , we define z, a, b as vectors and W as a matrix of size $m \times n$ that represents the weights from the n nodes of the current layer to the m nodes of the next layer, the size of the matrix and vectors have to match because they will become the next layer's weighted sum vector that will be activated with activation function.

- **Activation(a)**

The activation function is denoted with this formula

$$a^{(L)} = \sigma(z^{(L)})$$

where σ is a non-linear activation function like sigmoid, tanh, ReLU, or softmax.

The final layer's configuration depends on the type of classification:

- **Binary Classification**

Uses a single neuron with a sigmoid activation. An output of greater than 0.5 typically indicates the positive class.

- **Multi-class Classification**

Uses multiple neurons (one per class) with a softmax activation. Softmax converts raw scores into a probability distribution that sums to 1.

- **Result Interpretation**

The predicted class is usually the index of the neuron with the highest probability score.

2. **Loss Calculation**

The loss (or error) is calculated using a loss function such as Mean Squared Error (MSE) for regression tasks or Cross-Entropy Loss for classification tasks.

3. **Backpropagation**

In backpropagation the error is propagated back through the network to update the weights. The gradient of the loss function with respect to each weight is calculated and the weights are adjusted using gradient descent.

Gradient descent is an optimization algorithm used to minimize the loss function by iteratively updating the weights in the direction of the negative gradient. Common variants of gradient descent include:

1. **Batch Gradient Descent**

Updates weights after computing the gradient over the entire dataset.

2. **Stochastic Gradient Descent**

Updates weights for each training example individually.

3. **Mini-batch Gradient Descent**

It updates weights after computing the gradient over a small batch of training examples.

Evaluating the performance of the feed-forward neural network trained model involves several metrics:

1. **Accuracy**

The proportion of correctly classified instances out of the total instances.

2. **Precision**

The ratio of true positive predictions to the total predicted positives.

3. **Recall**

The ratio of true positive predictions to the actual positives.

4. **F1 Score**

The harmonic mean of precision and recall, providing a balance between the two.

5. **Confusion Matrix**

A table used to describe the performance of a classification model, showing the true positives, true negatives, false positives and false negatives.

Improving Convergence

Gradient descent has various advantages. It is simple, local; namely, the change in a weight uses only the values of the presynaptic and postsynaptic units and the error (suitably backpropagated). When on-line training is used, it does not need to store the training set and can adapt as the task to be learned changes. But gradient descent converges slowly, there are other methods that has faster convergence, e.g. conjugate gradient and second-order methods to the training of multilayer perceptrons.

Overtraining

A feed-forward neural network (multilayer perceptron) with d inputs, H neurons in hidden layer (hidden units), and K outputs has $H(d + 1)$ weights in the first layer (the $+1$ is for bias) and $K(H + 1)$ weights in the second layer. Both the space and time complexity of an MLP is $O(H \cdot (K + d))$. When e denotes the number of training epochs, training time complexity is $O(e \cdot H \cdot (K + d))$.

In an application, d and K are predefined and H is the parameter that we play with to tune the complexity of the model. An overcomplex model memorizes the noise in the training set and does not generalize to the validation set. For example, in the case of polynomial regression we notices that in the presence of noise or small samples, increasing the polynomial order leads to worse generalization. Similarly in an MLP, when the number of hidden units is large, the generalization accuracy deteriorates, and the bias/variance dilemma also holds for the MLP, as it does for any statistical estimator [2].

Structuring the Network

In some applications we know that the input has a local structure. For example, in computer vision we know that nearby pixels are correlated and there are local features like edges and corners; any object, for example, a handwritten digit, may be defined as a combination of such primitives. Similarly, in speech recognition, locality is in time and inputs close in time can be grouped as speech primitives. By combining these primitives, longer utterances, for example, speech phonemes, may be defined. In such a case when designing the MLP, hidden units are not connected to all input units because not all inputs are correlated. Instead, we define hidden units that define a window over the input space and are connected to only a small local subset of the inputs. This decreases the number of connections and therefore the number of free parameters.

We can repeat this in successive layers where each layer is connected to a small number of local units below and checks for a more complicated feature by combining the features below in a larger part of the input space until we get to the output layer.

For example, the input may be pixels. By looking at pixels, the first hidden layer units may learn to check for edges of various orientations. Then by combining edges, the second hidden layer units can learn to check for combinations of edges-for example, arcs, corners, line ends-and then combining them in upper layers, the units can look for semi-circles, rectangles, or in the case of a face recognition application, eyes, mouth, and so forth. This is the example of a hierarchical cone where features get more complex, abstract, and fewer in number as we go up the network until we get to classes. Such an architecture is called a convolutional neural network where the work of each hidden unit is considered to be a convolution of its input with its weight vector.

[SI*] The feed-forward neural network is the most successful model for searching patterns in data [5]. This model comprises multiple layers of logistic regression models rather than multiple perceptrons. For many applications, the resulting model can be significantly more compact, and hence faster to evaluate, than a support vector machine having the same generalization performance. the price to be paid for this compactness, is that the likelihood

function, which forms the basis for network training, is no longer a convex function of the model parameters.

[SI*] The linear model that will be the basic for the feed-forward neural network has this form

$$y(\mathbf{x}, \mathbf{w}) = f\left(\sum_{j=1}^M w_j \phi_j(\mathbf{x})\right) \quad (10.11)$$

where $f(\cdot)$ is a nonlinear activation function in the case of classification and is the identity in the case of regression. The difference between a linear model is that we will make the basic functions $\phi_j(\mathbf{x})$ depend on parameters and then to allow these parameters to be adjusted, along with the coefficients $\{w_j\}$, during training.

This leads to the basic neural network model, which can be described a series of functional transformations. First we construct M linear combinations of the input variables $x_1, \dots, x_D$ in the form

$$a_j = \sum_{i=1}^D w_{ji}^{(1)} x_i + w_{j0}^{(1)} \quad (10.12)$$

where $j = 1, 2, \dots, M$, and the superscript (1) indicates that the corresponding parameters are in the first 'layer' of the network. With $w_{ji}^{(1)}$ as weights and the parameters $w_{j0}^{(1)}$ as biases. The quantities a_j are known as activations. Each of them is then transformed using a differentiable, nonlinear activation function $h(\cdot)$ to give

$$z_j = h(a_j) \quad (10.13)$$

These quantities correspond to the outputs of the basis functions in (10.11) that, in the context of neural networks, are called hidden units. The nonlinear functions $h(\cdot)$ are generally chosen to be sigmoidal functions such as the logistic sigmoid or the tanh function. Following (10.11), these values are again linearly combined to give output unit activations

$$a_k = \sum_{j=1}^M w_{kj}^{(2)} z_j + w_{k0}^{(2)} \quad (10.14)$$

where $k = 1, 2, \dots, K$, and K is the total number of outputs. This transformation corresponds to the second layer of the network, and gain the $w_{k0}^{(2)}$ are bias parameters. Finally, the output unit activations are transformed using an appropriate activation function to give a set of network outputs y_k . The choice of activation function is determined by the nature of data and the assumed distribution of target variables. Thus for standard regression problems, the activation function is the identity so that $y_k = a_k$. Similarly, for multiple binary classification problems, each output unit activation is transformed using a logistic sigmoid function so that

$$y_k = \sigma(a_k) \quad (10.15)$$

where

$$\sigma(a) = \frac{1}{1 + e^{-a}} \quad (10.16)$$

For multiclass problem, a softmax activation function is used.

We can combine these various stages to give the overall network function that, for sigmoidal output unit activation, takes the form

$$y_k(\mathbf{x}, \mathbf{w}) = \sigma \left(\sum_{j=1}^M w_{kj}^{(2)} h \left(\sum_{i=1}^D w_{ji}^{(1)} x_i + w_{j0}^{(1)} \right) + w_{k0}^{(2)} \right) \quad (10.17)$$

where the set of all weight and bias parameters have been grouped together into a vector $\mathbf{w}$. Thus the neural network model is simply a nonlinear function from a set of input variables $\{x_i\}$ to a set of output variables $\{y_k\}$ controlled by a vector $\mathbf{w}$ of adjustable parameters. The process of evaluating (10.17) can then be interpreted as a forward propagation or information through the network.

The bias parameters in (10.12) can be absorbed into the set of weight parameters by defining an additional input variable x_0 whose value is clamped at $x_0 = 1$, so that (10.12) takes the form

$$a_j = \sum_{i=0}^D w_{ji}^{(1)} x_i \quad (10.18)$$

We can similarly absorb the second-layer biases into the second-layer weights, so that the overall network function becomes

$$y_k(\mathbf{x}, \mathbf{w}) = \sigma \left(\sum_{j=0}^M w_{kj}^{(2)} h \left(\sum_{i=0}^D w_{ji}^{(1)} x_i \right) \right) \quad (10.19)$$

The neural network model comprises two stages of processing, each of which resembles the perceptron model, and for this reason the feed-forward neural network is also known as the multilayer perceptron (MLP). A key difference compared to the perceptron, however, is that the neural network uses continuous sigmoidal nonlinearities in the hidden units, whereas the perceptron uses step-function nonlinearities. This means that the neural network function is differentiable with respect to the network parameters, and this property will play a central role in network training.

If the activation functions of all the hidden units in a network are taken to be linear, then for any such network we can always find an equivalent network without hidden units. This follows from the fact that the composition of successive linear transformations is itself a linear transformation. However, if the number of hidden units is smaller than either the number of input or output units, then the transformations that the network can generate are not the most general possible linear transformations from inputs to outputs because information is lost in the dimensionality reduction at the hidden units.

[SI*] Network Training

We have viewed neural networks as a general class of parametric nonlinear functions from a vector $\mathbf{x}$ of input variables to a vector $\mathbf{y}$ of output variables. A simple approach to the problem of determining the network parameters is to make an analogy with the discussion of polynomial curve fitting. Given a training set comprising a set of input vectors $\{\mathbf{x}_n\}$, where $n = 1, 2, \dots, N$, together with a corresponding set of target vectors $\{\mathbf{t}_n\}$, we minimize the error function

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \|\mathbf{y}(\mathbf{x}_n, \mathbf{w}) - \mathbf{t}_n\|^2 \quad (10.20)$$

However, we can provide a much more general view of network training by first giving a probabilistic interpretation to the network outputs.

Consider a single target variable t that can take any real value. We assume that t has Gaussian distribution with an x -dependent mean, which is given by the output of the neural network, so that

$$p(t|x, w) = N(t|y(x, w), \beta^{-1}) \quad (10.21)$$

where β is the precision (inverse variance) of the Gaussian noise. For the conditional distribution given by (10.21), it is sufficient to take the output unit activation function to be the identity, because such a network can approximate any continuous function from x to y . Given a data set of N independent, identically distributed observations $\mathbf{X} = \{x_1, x_2, \dots, x_N\}$, along with corresponding target values $\mathbf{t} = \{t_1, t_2, \dots, t_N\}$, we can construct the corresponding likelihood function

$$p(\mathbf{t}|\mathbf{X}, w, \beta) = \prod_{n=1}^N p(t_n|x_n, w, \beta)$$

Taking the negative logarithm, we obtain the error function

$$\frac{\beta}{2} \sum_{n=1}^N \{y(x_n, w) - t_n\}^2 - \frac{N}{2} \ln \beta + \frac{N}{2} \ln(2\pi) \quad (10.22)$$

which can be used to learn the parameters w and β . Here we consider the maximum likelihood approach of neural networks, it is usual to consider the minimization of an error function rather than the maximization of the (log) likelihood.

Now, consider first the determination of w . Maximizing the likelihood function is equivalent to minimizing the sum-of-squares error function given by

$$E(w) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, w) - t_n\}^2 \quad (10.23)$$

where we have discarded additive and multiplicative constants. The value of w found by minimizing $E(w)$ will be denoted by w_{ML} because it corresponds to the maximum likelihood solution. In practice, the nonlinearity of the network function $y(x_n, w)$ causes the error $E(w)$ to be nonconvex, and so in practice local maxima of the likelihood may be found, corresponding to local minima of the error function.

Having found w_{ML} , the value of β can be found by minimizing the negative log likelihood to give

$$\frac{1}{\beta_{ML}} = \frac{1}{N} \sum_{n=1}^N \{y(x_n, w_{ML}) - t_n\}^2 \quad (10.24)$$

Note that this can be evaluated once the iterative optimization required to find w_{ML} is completed. If we have multiple target variables, and we assume that they are independent conditional on x and w with shared noise precision β , then the conditional distribution of the target values is given by

$$p(\mathbf{t}|\mathbf{x}, w) = N(\mathbf{t}|\mathbf{y}(\mathbf{x}, w), \beta^{-1}\mathbf{I}) \quad (10.25)$$

Following the same argument as for a single target variable, we see that the maximum likelihood weights are determined by minimizing the sum-of-squares error function (10.20). The noise precision is then given by

$$\frac{1}{\beta_{ML}} = \frac{1}{NK} \sum_{n=1}^N \|\mathbf{y}(x_n, w_{ML}) - \mathbf{t}_n\|^2 \quad (10.26)$$

where K is the number of target variables. The assumption of independence can be dropped at the expense of a slightly more complex optimization problem.

There is a natural pairing of the error function (given by the negative log likelihood) and the output unit activation function. In the regression case, we can view the network as having an output activation function that is the identity, so that $y_k = a_k$. The corresponding sum-of-squares error function has the property

$$\frac{\partial E}{\partial a_k} = y_k - t_k \quad (10.27)$$

which we shall make use when discussing error backpropagation.

Now consider the case of binary classification in which we have a single target variable t such that $t = 1$ denotes class C_1 and $t = 0$ denotes class C_2 . We consider a network having a single output whose activation function is a logistic sigmoid

$$y = \sigma(a) = \frac{1}{1 + e^{-a}} \quad (10.28)$$

so that $0 \leq y(x, w) \leq 1$. We can interpret $y(x, w)$ as the conditional probability $p(C_1|x)$, with $p(C_2|x)$ given by $1 - y(x, w)$. The conditional distribution of targets given inputs is then a Bernoulli distribution of the form

$$p(t|x, w) = y(x, w)^t \{1 - y(x, w)\}^{1-t} \quad (10.29)$$

If we consider a training set of independent observations, then the error function, which is given by the negative log likelihood, is then a cross-entropy error function of the form

$$E(w) = - \sum_{n=1}^N \{t_n \ln(y_n) + (1 - t_n) \ln(1 - y_n)\} \quad (10.30)$$

where y_n denotes $y(x_n, w)$. Note that there is no analogue of the noise precision β because the target values are assumed to be correctly labelled. However, the model is easily extended to allow for labelling errors. Simard *et al.* (2003) found that using the cross-entropy error function instead of the sum-of-squares for a classification problem leads to faster training as well as improved generalization.

If we have K separate binary classifications to perform, then we can use a network having K outputs each of which has a logistic sigmoid activation function. Associated with each output is a binary class label $t_k \in \{0, 1\}$, where $k = 1, 2, \dots, K$. If we assume that the class labels are independent, given the input vector, then the conditional distribution of the targets is

$$p(\mathbf{t}|\mathbf{x}, w) = \prod_{k=1}^K y_k(\mathbf{x}, w)^{t_k} [1 - y_k(\mathbf{x}, w)]^{1-t_k} \quad (10.31)$$

Taking the negative logarithm of the corresponding likelihood function then gives the following error function

$$E(w) = - \sum_{n=1}^N \sum_{k=1}^K \{t_{nk} \ln(y_{nk}) + (1 - t_{nk}) \ln(1 - y_{nk})\} \quad (10.32)$$

where y_{nk} denotes $y_k(x_n, w)$. Again, the derivative of the error function with respect to the activation for a particular output unit takes the form (10.27) just as in the regression case.

It is interesting to contrast the neural network solution to this problem with the corresponding approach based on a linear classification model. Suppose that we are using a standard two-layer network. We see that the weight parameters in the first layer of the network are shared between the various outputs, whereas in the linear model each classification problem is solved independently. The first layer of the network can be viewed as performing a nonlinear feature extraction, and the sharing of features between the different outputs can save on computation and can also lead to improved generalization.

Finally, we consider the standard multiclass classification problem in which each input is assigned to one of K mutually exclusive classes. The binary target variables $t_k \in \{0, 1\}$ have a 1-of- K coding scheme indicating the class, and the network outputs are interpreted as $y_k(\mathbf{x}, \mathbf{w}) = p(t_k = 1 | \mathbf{x})$, leading to the following error function

$$E(\mathbf{w}) = - \sum_{n=1}^N \sum_{k=1}^K t_{kn} \ln(y_k(\mathbf{x}_n, \mathbf{w})) \quad (10.33)$$

We see that the output unit activation function, which corresponds to the canonical link, is given by the softmax function

$$y_k(\mathbf{x}, \mathbf{w}) = \frac{e^{a_k(\mathbf{x}, \mathbf{w})}}{\sum_j e^{a_j(\mathbf{x}, \mathbf{w})}} \quad (10.34)$$

which satisfies $0 \leq y_k \leq 1$ and $\sum_k y_k = 1$. Note that the $y_k(\mathbf{x}, \mathbf{w})$ are unchanged if a constant is added to all of the $a_k(\mathbf{x}, \mathbf{w})$, causing the error function to be constant for some directions in weight space. This degeneracy is removed if an appropriate regularization term is added to the error function.

Once again, the derivative of the error function with respect to the activation for a particular output unit takes the familiar form (10.27).

In summary, there is a natural choice of both output unit activation function and matching error function, according to the type of problem being solved. For regression we use linear outputs and a sum-of-squares error, for (multiple independent) binary classifications we use logistic sigmoid outputs and a cross-entropy error function, and for multiclass classification we use softmax outputs with the corresponding multiclass cross-entropy error function. For classification problems involving two classes, we can use a single logistic sigmoid output, or alternatively we can use a network with two outputs having a softmax output activation function.

Parameter Optimization

We turn next to the task of finding a weight vector $\mathbf{w}$, which minimizes the chosen function $E(\mathbf{w})$. First note that if we make a small step in weight space from $\mathbf{w}$ to $\mathbf{w} + \delta \mathbf{w}$ then the change in the error function is $\delta E \approx \delta \mathbf{w}^T \nabla E(\mathbf{w})$ points in the direction of greatest rate of increase of the error function. Because the error $E(\mathbf{w})$ is a smooth continuous function of $\mathbf{w}$, its smallest value will occur at a point in weight space such that the gradient of the error function vanishes, so that

$$\nabla E(\mathbf{w}) = 0 \quad (10.35)$$

as otherwise we could make a small step in the direction of $-\nabla E(\mathbf{w})$ and thereby further reduce the error. Points at which the gradient vanishes are called stationary points, and may be further classified into minima, maxima, and saddle points.

Our goal is to find a vector $\mathbf{w}$ such that $E(\mathbf{w})$ takes its smallest value. However, the error function typically has a highly nonlinear dependence on the weights and bias parameters,

and so there will be many points in weight space at which the gradient vanishes (or is numerically very small). Indeed, we see that for any point $\mathbf{w}$ that is a local minimum, there will be other points in weight space that are equivalent minima. For instance, in a two-layer network with M hidden neurons/units, each point in weight space is a member of a family of $M!2^M$ equivalent points.

Furthermore, there will typically be multiple inequivalent stationary points and in particular multiple inequivalent minima. A minimum that corresponds to the smallest value of the error function for any weight vector is said to be a global minimum. Any other minima corresponding to higher values of the error function are said to be local minima. For a successful application of neural networks, it may not be necessary to find the global minimum (and in general it will not be known whether the global minimum has been found) but it may be necessary to compare several local minima in order to find a sufficiently good solution.

Because there is clearly no hope of finding an analytical solution to the equation $\nabla E(\mathbf{w}) = 0$ we resort to iterative numerical procedures. The optimization of continuous nonlinear functions is a widely studied problem and there exists an extensive literature on how to solve it efficiently. Most techniques involve choosing some initial value $\mathbf{w}^{(0)}$ for the weight vector and then moving through weight space in a succession of steps of the form

$$\mathbf{w}^{(\tau+1)} = \mathbf{w}^{(\tau)} + \Delta \mathbf{w}^{(\tau)} \quad (10.36)$$

where τ labels the iteration step. Different algorithms involve different choices for the weight vector update $\Delta \mathbf{w}^{(\tau)}$. Many algorithms make use of gradient information and therefore require that, after each update, the value of $\nabla E(\mathbf{w})$ is evaluated at the new weight vector $\mathbf{w}^{(\tau+1)}$. In order to understand the importance of gradient information, it is useful to consider a local approximation to the error function based on a Taylor expansion.

Local Quadratic Approximation

Insight into the optimization problem, and into the various techniques for solving it, can be obtained by considering a local quadratic approximation to the error function.

Consider the Taylor expansion of $E(\mathbf{w})$ around some point $\hat{\mathbf{w}}$ in weight space

$$E(\mathbf{w}) \cong E(\hat{\mathbf{w}}) + (\mathbf{w} - \hat{\mathbf{w}})^T \mathbf{b} + \frac{1}{2}(\mathbf{w} - \hat{\mathbf{w}})^T \mathbf{H}(\mathbf{w} - \hat{\mathbf{w}}) \quad (10.37)$$

where cubic and higher terms have been omitted. Here $\mathbf{b}$ is defined to be the gradient of E evaluated at $\hat{\mathbf{w}}$

$$\mathbf{b} \equiv \nabla E|_{\mathbf{w}=\hat{\mathbf{w}}} \quad (10.38)$$

and the Hessian matrix $\mathbf{H} = \nabla \nabla E$ has elements

$$(\mathbf{H})_{ij} \equiv \frac{\partial^2 E}{\partial w_i \partial w_j} \Big|_{\mathbf{w}=\hat{\mathbf{w}}} \quad (10.39)$$

From Eq. (10.37), the corresponding local approximation to the gradient is given by

$$\nabla E \cong \mathbf{b} + \mathbf{H}(\mathbf{w} - \hat{\mathbf{w}}) \quad (10.40)$$

For points $\mathbf{w}$ that are sufficiently close to $\hat{\mathbf{w}}$, these expressions will give reasonable approximations for the error and its gradient.

Consider the particular case of a local quadratic approximation around a point $\mathbf{w}^*$ that is a

minimum of the error function. In this case there is no linear term, because $\nabla E = 0$ at $\mathbf{w}^*$, and Eq. (10.37) becomes

$$E(\mathbf{w}) = E(\mathbf{w}^*) + \frac{1}{2}(\mathbf{w} - \mathbf{w}^*)^T \mathbf{H}(\mathbf{w} - \mathbf{w}^*) \quad (10.41)$$

where the Hessian $\mathbf{H}$ is evaluated at $\mathbf{w}^*$. In order to interpret this geometrically, consider the eigenvalue equation for the Hessian matrix

$$\mathbf{H}\mathbf{u}_i = \lambda_i \mathbf{u}_i \quad (10.42)$$

where the eigenvectors $\mathbf{u}_i$ form a complete orthonormal set so that

$$\mathbf{u}_i^T \mathbf{u}_j = \delta_{ij} \quad (10.43)$$

We now expand $(\mathbf{w} - \mathbf{w}^*)$ as a linear combination of the eigenvectors in the form

$$\mathbf{w} - \mathbf{w}^* = \sum_i \alpha_i \mathbf{u}_i \quad (10.44)$$

This can be regarded as a transformation of the coordinate system in which the origin is translated to the point $\mathbf{w}^*$, and the axes are rotated to align with the eigenvectors (through the orthogonal matrix whose columns are the $\mathbf{u}_i$). Substituting Eq. (10.44) into Eq. (10.41), and using Eqs. (10.42) and (10.43), allows the error function to be written in the form

$$E(\mathbf{w}) = E(\mathbf{w}^*) + \frac{1}{2} \sum_i \lambda_i \alpha_i^2 \quad (10.45)$$

A matrix $\mathbf{H}$ is said to be positive definite if, and only if,

$$\mathbf{v}^T \mathbf{H} \mathbf{v} > 0, \quad \text{for all } \mathbf{v} \quad (10.46)$$

Because the eigenvectors $\{\mathbf{u}_i\}$ form a complete set, an arbitrary vector $\mathbf{v}$ can be written in the form

$$\mathbf{v} = \sum_i c_i \mathbf{u}_i \quad (10.47)$$

From Eqs (10.42) and (10.43), we then have

$$\mathbf{v}^T \mathbf{H} \mathbf{v} = \sum_i c_i^2 \lambda_i \quad (10.48)$$

and so $\mathbf{H}$ will be positive definite, if, and only if, all of its eigenvalues are positive. In the new coordinate system, whose basis vectors are given by the eigenvectors $\{\mathbf{u}_i\}$, the contours of constant E are ellipses centred on the origin. For a one-dimensional weight space, a stationary point $\mathbf{w}^*$ will be a minimum if

$$\left. \frac{\partial^2 E}{\partial w^2} \right|_{\mathbf{w}^*} > 0 \quad (10.49)$$

The corresponding result in D -dimensions is that the Hessian matrix, evaluated at $\mathbf{w}^*$, should be positive definite.

Use of Gradient Information

It is possible to evaluate the gradient of an error function efficiently by means of the backpropagation procedure. The use of this gradient information can lead to significant improvements

in the speed with which the minima of the error function can be located.

In the quadratic approximation to the error function, given in Eq. (10.37), the error surface is specified by the quantities $\mathbf{b}$ and $\mathbf{H}$, which contain a total of $W \frac{W+3}{2}$ independent elements (because the matrix $\mathbf{H}$ is symmetric), where W is the dimensionality of $\mathbf{w}$ (i.e., the total number of adaptive parameters in the network). The location of the minimum of this quadratic approximation therefore depends on $O(W^2)$ parameters, and we should not expect to be able to locate the minimum until we have gathered $O(W^2)$ independent pieces of information. If we do not make use of gradient information, we would expect to have to perform $O(W^2)$ function evaluations, each of which would require $O(W)$ steps. Thus, the computational effort needed to find the minimum using such an approach would be $O(W^3)$. Now compare this with an algorithm that makes use of the gradient information. Because each evaluation of ∇E brings W items of information, we might hope to find the minimum of the function in $O(W)$ gradient evaluations. As we shall see, by using error backpropagation, each such evaluation takes only $O(W)$ steps and so the minimum can now be found in $O(W^2)$ steps. For this reason, the use of gradient information forms the basis of practical algorithms for training neural networks.

Gradient Descent Optimization

The simplest approach to using gradient information is to choose the weight update in Eq. (10.36) to comprise a small step in the direction of the negative gradient, so that

$$\mathbf{w}^{(\tau+1)} = \mathbf{w}^{(\tau)} - \eta \nabla E(\mathbf{w}^{(\tau)}) \quad (10.50)$$

where the parameter $\eta > 0$ is known as the learning rate. After each such update, the gradient is re-evaluated for the new weight vector and the process repeated. Note that the error function is defined with respect to a training set, and so each step requires that the entire training set be processed in order to evaluate ∇E . Techniques that use the whole data set at once are called batch methods. At each step the weight vector is moved in the direction of the greatest decrease of the error function, and so this approach is known as gradient descent or steepest descent. Although such an approach might intuitively seem reasonable, in fact it turns out to be a poor algorithm.

For batch optimization, there are more efficient methods, such as conjugate gradients and quasi-Newton methods, which are much more robust and much faster than simple gradient descent [21]. Unlike gradient descent, these algorithms have the property that the error function always decreases at each iteration unless the weight vector has arrived at a local or global minimum.

In order to find a sufficiently good minimum, it may be necessary to run a gradient-based algorithm multiple times, each time using a differently randomly chosen starting point, and comparing the resulting performance on an independent validation set.

There is, however, an on-line version of gradient descent that has proved useful in practice for training neural networks on large data sets. Error functions based on maximum likelihood for a set of independent observations comprise a sum of terms, one for each data

$$E(\mathbf{w}) = \sum_{n=1}^N E_n(\mathbf{w}) \quad (10.51)$$

On-line gradient descent, also known as sequential gradient descent or stochastic gradient descent, makes an update to the weight vector based on one data point at a time, so that

$$\mathbf{w}^{(\tau+1)} = \mathbf{w}^{(\tau)} - \eta \nabla E_n(\mathbf{w}^{(\tau)}) \quad (10.52)$$

This update is repeated by cycling through the data either in sequence or by selecting points at random with replacement. There are of course intermediate scenarios in which the updates are based on batches of data points.

One advantage of on-line methods compared to batch methods is that the former handle redundancy in the data much more efficiently. To see, this consider an extreme example in which we take a data set and double its size by duplicating every data point. Note that this simply multiplies the error function by a factor of 2 and so is equivalent to using the original error function. Batch methods will require double the computational effort to evaluate the batch error function gradient, whereas on-line methods will be unaffected. Another property of on-line gradient descent is the possibility of escaping from local minima, since a stationary point with respect to the error function for the whole data set will generally not be a stationary point for each data point individually.

[SI*] **Error Backpropagation**

The multilayer perceptron architecture is sometimes called a backpropagation network. The term backpropagation is also used to describe the training of a multilayer perceptron using gradient descent applied to a sum-of-squares error function. Most training algorithms involve an iterative procedure for minimization of an error function, with adjustments to the weights being made in a sequence of steps. At each such step, we can distinguish between two distinct stages. In the first stage, the derivatives of the error function with respect to the weights must be evaluated. The important contribution of the backpropagation technique is in providing a computationally efficient method for evaluating such derivatives. Because it is at this stage that errors are propagated backwards through the network. In the second stage, the derivatives are then used to compute the adjustments to be made to the weights. The simplest such technique involves gradient descent. Thus, the first stage, namely the propagation of errors backwards through the network in order to evaluate derivatives, can be applied to many other kinds of network and not just the multilayer perceptron. It can also be applied to error functions other than just the simple sum-of-squares, and to the evaluation of other derivatives such as the Jacobian and Hessian matrices.

Evaluation of Error-Function Derivatives

We now derive the backpropagation algorithm for a general network having arbitrary feed-forward topology, arbitrary differentiable nonlinear activation functions, and a broad class of error function.

Many error functions of practical interest, for instance those defined by maximum likelihood for a set of i.i.d. data, comprise a sum of terms, one for each data point in the training set, so that

$$E(\mathbf{w}) = \sum_{n=1}^N E_n(\mathbf{w}) \quad (10.53)$$

Here we shall consider the problem of evaluating $\nabla E_n(\mathbf{w})$ for one such term in the error function. This may be used directly for sequential optimization, or the results can be accumulated over the training set in the case of batch methods.

Consider first a simple linear model in which the outputs y_k are linear combinations of the input variables x_i , so that

$$y_k = \sum_i w_{ki} x_i \quad (10.54)$$

together with an error function that, for a particular input pattern n , takes the form

$$E_n = \frac{1}{2} \sum_k (y_{nk} - t_{nk})^2 \quad (10.55)$$

where $y_{nk} = y_k(\mathbf{x}_n, \mathbf{w})$. The gradient of this error function with respect to a weight w_{ji} is given by

$$\frac{\partial E_n}{\partial w_{ji}} = (y_{nj} - t_{nj})x_{ni} \quad (10.56)$$

which can be interpreted as a 'local' computation involving the product of an 'error signal' $y_{nj} - t_{nj}$ associated with the output end of the link w_{ji} and the variable x_{ni} associated with the input end of the link.

In a general feed-forward network, each unit computes a weighted sum of its inputs of the form

$$a_j = \sum_i w_{ji} z_i \quad (10.57)$$

where z_i is the activation of a unit, or input, that sends a connection to unit j , and w_{ji} is the weight associated with that connection. We saw that biases can be included in this sum by introducing an extra unit, or input, with activation fixed at +1. We therefore do not need to deal with biases explicitly. The sum in (10.57) is transformed by a nonlinear activation function $h(\cdot)$ to give the activation z_j of unit j in the form

$$z_j = h(a_j) \quad (10.58)$$

Note that one or more variables z_i in the sum in (10.57) could be an input, and similarly, the unit j in (10.58) could be an output.

For each pattern in the training set, we shall suppose that we have supplied the corresponding input vector to the network and calculated the activations of all of the hidden and output units in the network by successive application of (10.57) and (10.58). This process is often called forward propagation because it can be regarded as a forward flow of information through the network.

Now consider the evaluation of the derivative of E_n with respect to a weight w_{ji} . The outputs of the various units will depend on the particular input pattern n . However, in order to keep the notation uncluttered, we shall omit the subscript n from the network variables. First we note that E_n depends on the weight w_{ji} only via the summed input a_j to unit j . We can therefore apply the chain rule for partial derivatives to give

$$\frac{\partial E_n}{\partial w_{ji}} = \frac{\partial E_n}{\partial a_j} \frac{\partial a_j}{\partial w_{ji}} \quad (10.59)$$

We now introduce a useful notation

$$\delta_j = \frac{\partial E_n}{\partial a_j} \quad (10.60)$$

where the δ 's are often referred to as errors for reasons. Using (10.57), we can write

$$\frac{\partial a_j}{\partial w_{ji}} = z_i \quad (10.61)$$

Substituting (10.60) and (10.61) into (10.59), we then obtain

$$\frac{\partial E_n}{\partial w_{ji}} = \delta_j z_i \quad (10.62)$$

Equation (10.62) tells us that the required derivative is obtained simply by multiplying the value of δ for the unit at the output end of the weight by the value of z for the unit at the input end of the weight (where $z = 1$ in the case of a bias). Note that this takes the same form as for the simple linear model considered at the start of this section. Thus, in order to evaluate the derivatives, we need only to calculate the value of δ_j for each hidden and output unit in the network, and then apply (10.62).

As we have seen already, for the output units, we have

$$\delta_k = y_k - t_k \quad (10.63)$$

provided we are using the canonical link as the output-unit activation function. To evaluate the δ 's for hidden units, we again make use of the chain rule for partial derivatives,

$$\delta_j = \frac{\partial E_n}{\partial a_j} = \sum_k \frac{\partial E_n}{\partial a_k} \frac{\partial a_k}{\partial a_j} \quad (10.64)$$

where the sum runs over all units k to which unit j sends connections. Note that the labelled k could include other hidden units and/or output units. In writing down (10.64), we are making use of the fact that variations in a_j give rise to variations in the error function only through variations in the variables a_k . If we now substitute the definition of δ given by (10.60) into (10.64), and make use of (10.57) and (10.58), we obtain the following backpropagation formula

$$\delta_j = h'(a_j) \sum_k w_{kj} \delta_k \quad (10.65)$$

which tells us that the value of δ for a particular hidden unit can be obtained by propagating the δ 's backwards from units higher up in the network. Note that the summation in (10.65) is taken over the first index on w_{kj} (corresponding to backward propagation of information through the network), whereas in the forward propagation

$$z_k = h \left(\sum_j w_{kj} z_j \right)$$

where the sum runs over all units that send connections to unit k (and a bias parameter is included in the summation). It is taken over the second index. Because we already know the values of the δ 's for the output units, it follows that recursively applying (10.65) we can evaluate the δ 's for all of the hidden units in a feed-forward network, regardless of its topology.

For batch methods, the derivative of the total error E can then be obtained by repeating the above steps for each pattern in the training set and then summing over all patterns:

$$\frac{\partial E}{\partial w_{ji}} = \sum_n \frac{\partial E_n}{\partial w_{ji}} \quad (10.66)$$

In the above derivation we have implicitly assumed that each hidden or output unit in the network has the same activation function $h(\cdot)$.

i. Backpropagation for Feed-Forward Neural Network

Consider a feed-forward neural network with 2 inputs, and one hidden layer with 2 neurons in the hidden layer, and 2 outputs. Assume that the neurons use sigmoid activation function for the forward and backward pass and there is no bias.




Figure 10.4: The schematic diagram of this example feed-forward neural network.

We will compute and simulate the backpropagation for this feed-forward neural network (FFN).

Let the input be

$$\mathbf{x} = \begin{bmatrix} 0.35 \\ 0.7 \end{bmatrix}$$

The initial weights matrix from the input to the hidden layer

$$\mathbf{W}_0 = \begin{bmatrix} 0.2 & 0.2 \\ 0.3 & 0.3 \end{bmatrix}$$

The initial weights matrix from the hidden layer to the output

$$\mathbf{W}_1 = \begin{bmatrix} 0.3 & 0.9 \\ 0.1 & 0.5 \end{bmatrix}$$

and the output

$$\mathbf{y} = \begin{bmatrix} 0.5 \\ 0.37 \end{bmatrix}$$

The initial calculation will compute the weighted sum at each node with the sum formula

$$a_j = f \left(\sum_{i=1}^2 x_i w_{ij} + b_j \right)$$

x_i represents the value of the i -th input, a_j represents the weighted sum of all the inputs and weights at each node, w_{ij} represents the weights between the i -th input and the j -th neuron, there is no bias, so $b_j = 0$.

After applying the activation function to a_j , we get the output of the neuron

$$o_j = \text{activation function}(a_j) = f \left(\sum_{i=1}^2 x_i w_{ij} + b_j \right)$$

The activation function is using the sigmoid function that returns a value between 0 and 1, introducing non-linearity into the model, thus

$$o_j = \frac{1}{1 + e^{-a_j}}$$

At h_1 node we will have

$$\begin{aligned} a_1 &= (w_{011}x_1) + (w_{012}x_2) \\ &= (0.2 * 0.35) + (0.2 * 0.7) \\ &= 0.21 \end{aligned}$$

Once we calculated the a_1 value, we can now proceed to find the $f(a_1)$ value (activation function computation)

$$\begin{aligned} f(a_1) &= \frac{1}{1 + e^{-0.21}} \\ &= 0.5523 \end{aligned}$$

Similarly, for the the second node in the hidden layer h_2 we will have

$$\begin{aligned} a_2 &= (w_{021}x_1) + (w_{022}x_2) \\ &= (0.3 * 0.35) + (0.3 * 0.7) \\ &= 0.315 \end{aligned}$$

$$\begin{aligned} f(a_2) &= \frac{1}{1 + e^{-0.315}} \\ &= 0.5781 \end{aligned}$$

The predicted output will be

$$\begin{aligned} o_1 &= (w_{111}h_1) + (w_{112}h_2) \\ &= (0.3 * 0.35) + (0.9 * 0.5781) \\ &= 0.68598 \end{aligned}$$

$$\begin{aligned} f(o_1) &= \frac{1}{1 + e^{-0.68598}} \\ &= 0.66507 \end{aligned}$$

and

$$\begin{aligned} o_2 &= (w_{121}h_1) + (w_{122}h_2) \\ &= (0.3 * 0.35) + (0.9 * 0.5781) \\ &= 0.34428 \end{aligned}$$

$$\begin{aligned} f(o_2) &= \frac{1}{1 + e^{-0.34428}} \\ &= 0.58523 \end{aligned}$$

thus we will have

$$\hat{y} = \begin{bmatrix} 0.66507 \\ 0.58523 \end{bmatrix}$$

The error calculation can be obtained by computing with this formula

$$\epsilon = \mathbf{y} - \hat{\mathbf{y}}$$

$$\epsilon = \begin{bmatrix} -0.165074 \\ -0.215231 \end{bmatrix}$$

The norm is

$$||\epsilon|| = 0.271244$$

the target is to minimize the norm of ϵ , to make it near zero. From here, we will do backpropagating process because the weights W_0 and W_1 that we have used are not sufficient to predict the output.

Backpropagation

First we will calculate the gradients with this formula to compute the change in each weight

$$\Delta w_{ij} = \eta \times \delta_j^{(L)} \times O_j^{(L-1)}$$

with η as the learning rate (we set it to 0.51), $\delta_j^{(L)}$ as the error term of the j -th neuron in the next layer L , and $O_j^{(L-1)}$ is the output value of the current layer / the sending neuron (layer $L - 1$). For this example, we will assume that layer L denotes the output layer.

To compute the error term, we can start from the first node in the last layer, the output layer

$$\begin{aligned} \delta y_1 &= f'(O_1^{(L)}) \left(\delta_k^{(L)} \right) \\ &= \hat{y}_1(1 - \hat{y}_1)(y_1 - \hat{y}_1) \\ &= 0.66507(1 - 0.66507)(-0.165074) \\ &= -0.03677 \end{aligned}$$

the second nodes in the output layer

$$\begin{aligned} \delta y_2 &= f'(O_1^{(L)}) \left(\delta_k^{(L)} \right) \\ &= \hat{y}_2(1 - \hat{y}_2)(y_2 - \hat{y}_2) \\ &= 0.585231(1 - 0.585231)(-0.215231) \\ &= -0.052244 \end{aligned}$$

now to the first node in the hidden layer

$$\begin{aligned} \delta h_1 &= f'(O_1^{(L-1)}) \sum_{k=1}^2 \left(w_{k,1} \cdot \delta_k^L \right) \\ &= 0.5523(1 - 0.5523)((0.3 \times -0.03677) + (0.1 \times -0.052244)) \\ &= -0.004019 \end{aligned}$$

the second node in the hidden layer

$$\begin{aligned} \delta h_2 &= f'(O_2^{(L-1)}) \sum_{k=1}^2 \left(w_{k,2} \cdot \delta_k^L \right) \\ &= 0.578105(1 - 0.578105)((0.9 \times -0.03677) + (0.5 \times -0.052244)) \\ &= -0.014443 \end{aligned}$$

After we have computed all the δ we will have the δ vectors

$$\delta_{output} = \begin{bmatrix} -0.03677 \\ -0.052244 \end{bmatrix}$$

$$\delta_{hidden} = \begin{bmatrix} -0.004019 \\ -0.014443 \end{bmatrix}$$

Now, we can do weights updates computation, for the weights from the hidden layer to the output layer

$$\begin{aligned} \Delta w_{11} &= \eta \times (\delta y_1) \times O_1^{(L-1)} \\ &= 0.51 \times -0.03677 \times 0.552308 \\ &= -0.010357 \end{aligned}$$

$$\begin{aligned} \Delta w_{12} &= \eta \times (\delta y_1) \times O_2^{(L-1)} \\ &= 0.51 \times -0.03677 \times 0.578105 \\ &= -0.010841 \end{aligned}$$

$$\begin{aligned} \Delta w_{21} &= \eta \times (\delta y_2) \times O_1^{(L-1)} \\ &= 0.51 \times -0.052244 \times 0.552308 \\ &= -0.0147159 \end{aligned}$$

$$\begin{aligned} \Delta w_{22} &= \eta \times (\delta y_2) \times O_2^{(L-1)} \\ &= 0.51 \times -0.052244 \times 0.578105 \\ &= -0.015403 \end{aligned}$$

for the weights from the input layer to the hidden layer

$$\begin{aligned} \Delta w_{011} &= \eta \times (\delta h_1) \times O_1^{L-2} \\ &= 0.51 \times -0.004019 \times 0.35 \\ &= -0.00071739 \end{aligned}$$

$$\begin{aligned} \Delta w_{012} &= \eta \times (\delta h_1) \times O_2^{L-2} \\ &= 0.51 \times -0.004019 \times 0.7 \\ &= -0.001434783 \end{aligned}$$

$$\begin{aligned} \Delta w_{021} &= \eta \times (\delta h_2) \times O_1^{(L-2)} \\ &= 0.51 \times -0.01443 \times 0.35 \\ &= -0.002575755 \end{aligned}$$

$$\begin{aligned} \Delta w_{022} &= \eta \times (\delta h_2) \times O_2^{(L-2)} \\ &= 0.51 \times -0.01443 \times 0.7 \\ &= -0.00515151 \end{aligned}$$

Thus, the new weights matrix for the iteration 1 is

$$W_0^{(1)} = \begin{bmatrix} 0.199283 & 0.198565 \\ 0.297422 & 0.294844 \end{bmatrix}$$

and the new hidden weights matrix is

$$W1^{(1)} = \begin{bmatrix} 0.289643 & 0.889159 \\ 0.085284 & 0.484597 \end{bmatrix}$$

We will use the weights matrix $W0^{(1)}$ and the hidden weights matrix $W1^{(1)}$ for the forward propagation to obtain new prediction for $\mathbf{y}$, after iteration 1, we will have

$$\hat{\mathbf{y}} = \begin{bmatrix} 0.662159 \\ 0.580954 \end{bmatrix}$$

The error calculation

$$\epsilon = \mathbf{y} - \hat{\mathbf{y}}$$

$$\epsilon = \begin{bmatrix} -0.162159 \\ -0.210954 \end{bmatrix}$$

The norm is

$$||\epsilon|| = 0.266077$$

Since the predicted vector $\hat{\mathbf{y}}$ is still not converge to $\mathbf{y}$ (the target output vector), the process of calculating the error (ϵ) and backpropagating continues until the desired output is reached and we will obtain the final weights matrix and the final hidden weights matrix that can be used for testing.

This process demonstrates how backpropagation iteratively updates weights by minimizing errors until the network accurately predicts the output.

For the C++ code, we use a lot of matrix and vector multiplication and we have to remember the rule of the multiplication for them, thus when we create the weights matrix, we transpose the weights matrix that represents $W0$ with the entries $w0_{ij}$ that represents the weights between the i -th input and the j -th neuron. That is why the initial weights matrix is like this

$$W0 = \begin{bmatrix} 0.2 & 0.2 \\ 0.3 & 0.3 \end{bmatrix}$$

instead of

$$W0 = \begin{bmatrix} 0.2 & 0.3 \\ 0.2 & 0.3 \end{bmatrix}$$

You will comprehend it when you remember the result of $W0\mathbf{x}$ corresponds to this neural network example, see Figure (10.4).

```

root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Neural Networks/Test Backpropagation FNN ]# ./main
Input:
0.35
0.7

Initial Weights :
      0.200000      0.200000
      0.300000      0.300000

Initial Hidden Layer Weights :
      0.300000      0.900000
      0.100000      0.500000

Biases:
0.000000
0.000000

Actual Output:
0.500000
0.370000

Feed-forward Neural Network with 1 hidden layer:

Hidden layer:
0.552308
0.578105

Predicted Output:
0.665074
0.585231

Error vector:
-0.165074
-0.215231

||Error|| = 0.271244

```

Figure 10.5: The computation of backpropagation process for the feed-forward neural network with *FNN_1hiddenlayer()* function (*SymIntegration/Examples/Neural Networks/Test Backpropagation FNN/main.cpp*).

```

*****
Backpropagation iteration
*****

Learning rate: 0.510000

Iteration = 1

Delta output vector:
-0.036770
-0.052244

Delta hidden layer vector:
-0.004019
-0.014443

New weights matrix:
      0.199283      0.198565
      0.297422      0.294844

New hiddenweights matrix:
      0.289643      0.889159
      0.085284      0.484597

Hidden layer:
0.551997
0.577004

Predicted Output:
0.662159
0.580954

Error vector:
-0.162159
-0.210954

||Error|| = 0.266077

```

Figure 10.6: The computation of backpropagation process for the feed-forward neural network with *FNN_1hiddenlayer()* function (*SymIntegration/Examples/Neural Networks/Test Backpropagation FNN/main.cpp*).

```
Iteration = 5

Delta output vector:
-0.034246
-0.048222

Delta hidden layer vector:
-0.002712
-0.012377

New weights matrix:
      0.197011      0.194022
      0.288050      0.276100

New hiddenweights matrix:
      0.250058      0.847861
      0.029354      0.426246

Hidden layer:
0.551014
0.572996

Predicted Output:
0.651039
0.564738

Error vector:
-0.151039
-0.194738

||Error|| = 0.246445
```

Figure 10.7: *The computation of backpropagation process for the feed-forward neural network with `FNN_1hiddenlayer()` function (`SymIntegration/Examples/Neural Networks/Test Backpropagation FNN/main.cpp`).*

ii. Feed-Forward Neural Network for Iris Dataset Classification with 1 Hidden Layer

Suppose we have data that are classified into n categories and each category is described by m variables. We want to create a neural network that can learn about the pattern of the data, so if we input a random number for the m variables we will know which category it belongs to based on the data that we have.

For this purpose, we will use the Iris dataset, it is a famous data of Iris flower species first introduced by the statistician Ronald Fisher in 1936, the Iris Setosa, Iris Versicolor, and Iris Virginica, there are 150 data that are classified into the three Iris species that we have mentioned, each iris has their corresponding features, represented by variables sepal length, sepal width, petal length, and petal width.

1. Data Preparation

We have a CSV file of the iris dataset, each line contains the measurements of one iris species.

2. Training

Training is where our network learns from its mistakes and gradually improves with backpropagation.

We feed some data into the neural network, letting it make predictions based on that data. We choose to feed 80% training data, which is 120 data, and we will have to shuffle the index when we code it before we train it, otherwise if we don't shuffle the index before we train the data, the computer will only learn about the first 120 data in the CSV file and it will never learn anything to be better.

3. Testing / Evaluation

After training we will obtain 120 bias for hidden layer, bias for output layer, weights from input to hiddenlayer and weights from hiddenlayer to output matrices that represent the model, we can use 20% of testing data, the leftover after we shuffle the index take the first 80% data for training the last 20% will be used to check which model can predict with highest accuracy.

Of course we can do that 20% testing, but in this book instead of using 20% we will test it on all the original data, all the 150 data with the best weights model we have obtained. Of course C++ can compute faster than Python and JULIA, it is a powerful programming language for implementing machine learning algorithms, so we have nothing to worry about, just run the code after long time of writing the code, we can learn more by doing this. Food for thoughts is like this, what if the last 20% of data all belong to Iris Setosa species only? and the model able to predict Iris Setosa 100% correctly without a doubt? Then it will probably make a false prediction in the end if we want the model to predict a flower with measurement of Iris Vesicolor.

Predict is where our network makes the prediction about the flower' species based on our input variables or testing data.

In this example we will create a Feed-forward neural network (multilayer peceptron) with 1 hidden layer to learn about the pattern of the iris dataset, it will have the architecture and configuration like this:

- **Input Layer:** 4 Neurons (representing sepal length, sepal width, petal length, petal width).
- **Hidden Layer(s):** 1 hidden layer is sufficient with 5-10 neurons in the hidden layer.
Activation function in the hidden layer: Sigmoid, or tanh, it is strong and stable. Another alternative is ReLU because of the fast training and avoiding vanishing gradient issues.
- **Output Layer:** 3 Neurons (representing Setosa, Versicolor, and Virginica).
Activation function in the output layer: Softmax, it is required for multi-class classification to output probabilities that sum to 1.
- **Loss Function:** Categorical Cross-Entropy.
- **Optimizer:** SGD (Stochastic Gradient Descent). Stochastic Gradient Descent for feed-forward neural networks updates model weights iteratively by calculating the gradient of the loss function based on a single sample or a small mini-batch, rather than the entire dataset. This approach allows for faster iteration and lower memory usage, which is ideal for large datasets, though it introduces high variance (noise) in parameter updates, which can help avoid local minima.

Small data size (150 data) doesn't need a lot of hidden layers, 2 is more than enough for iris dataset, but we only use 1 here.

Initialization

We initialize weights from input layer to hidden layer, it will become a matrix $W^{(0)}$ with size of 4×5 , then we initialize weights from hidden layer to the output layer, it is a matrix $W^{(1)}$ with size of 5×3 , then we initialize bias in the hidden layer, it is a vector b_0 of size 5, then we initialize bias in the output layer, a vector b_1 of size 3.

The matrix $W^{(0)}$ means that it represents the weights connecting the input neurons to the neurons in the hidden layer, we number it as 0, while matrix $W^{(1)}$ means that it represents the weights connecting the neurons in the hidden layer to the neurons in the output layer, we number it as 1, the superscript doesn't represent the updated value, it represents the layer location for the weights.

By the meaning of initialize, we generate random number (with double data type) with Mersenne Twister engine with range $[-1, 1]$.

The input for the training and testing function is:

1. Training matrix T that has the size of 120×4 , the order of the data is different with the original CSV data, because we have shuffled it first before it becomes an input to the function.
2. Vector t for the classification of the training matrix with corresponding order.
3. Testing matrix A that has the size of 150×4 , the order of the data is the same as the original CSV data, it is to test whether the model obtained in the training phase able to predict all the 150 data accurately.
4. Vector y that has the classification of the matrix A , it has the size of 150, the order of the data is the same as the original CSV data.

The first **for** loop is from 1 to 120, training phase, we check vector t , the first entry, if it is Setosa then we will create a new vector for this iteration

$$t_i = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

if it is Versicolor then we will create a new vector for this iteration

$$t_i = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

if it is Virginica then we will create a new vector for this iteration

$$t_i = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

This process of creating an identity vector / binary vector based on each classification / type is called one-hot encoding. The forward propagation phase will have a result of predicted probability by using Softmax activation function at the output layer, consider this output example $[0.2 \ 0.5 \ 0.3]$, it shows that the result leans toward Versicolor, the neural network predicts that based on the inputs the probability is 20% Setosa, 50% Versicolor, 30% Virginica.

Forward Propagation

In this phase, data flows from input to output, calculating the prediction based on the current weights.

A network with 4 input neurons, 1 hidden layer of 5 neurons, and an output layer of 3 neurons has:

Weights: $(4 \times 5) + (5 \times 3) = 35$

Biases: $5 + 3 = 8$

Total Parameters: 43

the hidden layer plays as the intermediate layer where transformations occur, to learn the complex pattern, while learning rate η is the hyperparameter that regulates the speed of weight updates.

The first general formula for the forward propagation from the input layer to the hidden layer is

$$h = f(W^{(0)}x + b)$$

with x as the input vector that represents the sepal length, sepal width, petal length, and petal width (4×1 vector), W is the weight matrix at the between the input layer and the hidden layer, b is the bias vector in the hidden layer, and f is the activation function (tanh), you can also try Sigmoid or ReLU for educational purpose.

The second general formula for the forward propagation from the hidden layer to the output layer is

$$y = f(W^{(1)}h + b)$$

with h as the input vector that represents the neurons in the hidden layer (5×1 vector), W is the weight matrix at the between the hidden layer and the output layer, b is the bias vector in the output layer, and f is the activation function (Softmax).

Loss Function(L)

The network computes the difference between predicted output $\hat{y}$ and the actual output y , often

using Mean Squared Error (MSE) or Cross-Entropy loss.

The error calculation can be computed with this formula:

$$e = y - \hat{y}$$

or if the output layer is using softmax then we will use cross-entropy loss that is defined as:

$$L = - \sum y \log(\hat{y})$$

Backpropagation

The error is sent backward through the network, using the chain rule to compute the gradient of the loss function with respect to each weight.

The backpropagation is the practice of fine-tuning the weights of a neural network based on the error rate obtained in the previous epoch. Proper tuning of the weights ensures lower error rates, making the model reliable by increasing its generalisation.

Weight Update

Weights are updated iteratively to reduce the loss / minimize the error (the error usually using Cross-Entropy Loss), by using an optimizer Stochastic Gradient Descent (SGD) formula, we will have the weight update formula

$$\begin{aligned} w_{ij}^{(L)} &= w_{ij}^{(L)} + \left(\eta \times \frac{\partial C}{\partial w^L} \right) \\ &= w_{ij}^{(L)} + \left(\delta_j^{(L)} \times O_i^{(L-1)} \right) \\ \delta_j^{(L)} &= f'(O_j^{(L)}) e_j \end{aligned}$$

with $\delta_j^{(L)}$ as the error term that represents the error of the receiving neuron (layer L), which includes the derivative of its activation function. O_i is the output value of the sending neuron (layer $L - 1$), e_j is the difference between the actual output of neuron j with the predicted output for neuron j ($y_j - \hat{y}_j$).

Gradient is denoted by $\frac{\partial C}{\partial w}$, C is the cost function, it is calculated using the chain rule to propagate error from the output layer back to the input.

The iterative process of feeding data and refining weights continues until the model's loss is minimized, resulting in optimized weights.

Bias Update

We update the bias after we compute the δ_j with this formula

$$b_j = b_j + \eta \cdot \delta_j$$

Classification Process

We do the normalization, scaling the input data, often using min-max scaling to a $[0, 1]$ range (with Sigmoid activation function) or $[-1, 1]$ range (with Tanh activation function), after the input signals pass through the optimal weighted connections, summed, and activated in the forward propagation phase we will reach the output layer, the final layer uses a Softmax activation function to produce a probability distribution across the three iris' species. The species with the highest probability is chosen.

III. Supervised Learning: Feed-Forward Neural Network (Multilayer Perceptron)

In the end, this feed-forward neural network can learn to classify objects or flowers with impressive accuracy, if it is trained well, we choose the optimal number of hidden layer, the activation function in each layer, and the optimal learning rate.

```
Epoch loss:6.315158
Epoch:994
Epoch loss:7.147247
Epoch:995
Epoch loss:6.726554
Epoch:996
Epoch loss:6.972265
Epoch:997
Epoch loss:6.941392
Epoch:998
Epoch loss:7.031226
Epoch:999
Epoch loss:6.588238
Epoch:1000
Epoch loss:4.846478

Model W0 matrix (from input to hidden layer):
      0.393350      3.921721      6.454273      1.608165      12.071175
      -2.648544     11.946345      6.058942      0.180194      9.000676
      2.442088     -11.245547     -10.595181     -2.257611     -18.769742
      0.583343      -6.368492     -6.489850     -1.133309     -10.643930

Model W1 matrix (from hidden layer to output):
      -3.357494      4.587249      5.600502
      3.581425      1.134718     -4.240889
      2.204437      1.369549     -1.116935
      2.784585     -0.994479     -3.142601
      1.424090      2.531305     -1.759481

Testing Feed-forward Neural Network with 1 hidden layer for Iris dataset:

Time taken by function: 18947334 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Neural Networks/Test FNN with Iris DataFrame 1 Hidden layer with Steepest Descent ]#
```

Figure 10.8: The training process computes the optimal $W^{(0)}$ and $W^{(1)}$ using 120 training data (80% of the original data), the updated weights with minimal loss from the backpropagation process with `FNN_1hiddenlayer_irisdataset_training` function, the epoch loss is zig-zagging due to the steepest descent formula, the loss cannot decrease continuously (`SymIntegration/Examples/Neural Network-s/Test FNN with Iris DataFrame 1 Hidden layer with Steepest Descent/main.cpp`).

```

*****
                                Next data
*****

Testing sample:30

Input layer:
5.900000
3.000000
5.100000
1.800000

Output layer:
0.000000
0.000000
1.000000

Predicted Output:
0.000004
0.020944
0.979052

Error vector:
-0.000004
-0.020944
0.020948

||Error|| = 0.029623

Average Loss over a batch of 30 samples : 0.194586

Correct prediction : 28
Wrong prediction : 2
Correct matrix:
      Total Correct      Setosa      Versicolor      Virginica      Model
      28.000000      9.000000      11.000000      8.000000      1.000000

Time taken by function: 24928 microseconds

```

Figure 10.9: The testing process with the optimal $W^{(0)}$ and $W^{(1)}$ for 30 testing sample (20% of the original data) with `FNN_1hiddenlayer_irisdataset_testing` function (`SymIntegration/Examples/Neural Network-s/Test FNN with Iris DataFrame 1 Hidden layer with Steepest Descent/main.cpp`).

```

*****
                                Next data
*****

Testing sample:150

Input layer:
5.900000
3.000000
5.100000
1.800000

Output layer:
0.000000
0.000000
1.000000

Predicted Output:
0.000004
0.020944
0.979052

Error vector:
-0.000004
-0.020944
0.020948

||Error|| = 0.029623

Average Loss over a batch of 150 samples : 0.161075

Correct prediction : 142
Wrong prediction : 8
Correct matrix:
      Total Correct      Setosa      Versicolor      Virginica      Model
      142.000000      50.000000      49.000000      43.000000      1.000000

Time taken by function: 1181561 microseconds

```

Figure 10.10: The testing process with the optimal $W^{(0)}$ and $W^{(1)}$ for 150 testing sample (100% of the original data) with `FNN_1hiddenlayer_irisdataset_testing` function (`SymIntegration/Examples/Neural Networks/Test FNN with Iris DataFrame 1 Hidden layer with Steepest Descent/main.cpp`).

The `FNN_1hiddenlayer_irisdataset_training` and `FNN_1hiddenlayer_irisdataset_testing` functions are an educative way to learn about neural network forward propagation, backpropagation

and the weights update algorithm. When we have finish till the last epoch, we know why we obtain the final weight that has the least error, you can see in the `src/artificialneuralnetworks.cpp` source code and you can uncomment or add something to print out the weights matrix, the output vector for the logit or output vector after the Softmax activation function.

It takes longer to code the training and testing in `vector<double>` and the `vector<vector<double>>` terms as a function for training and testing the neural network.

We want to compare it with the code that is using `class`, instead of function. This takes less codes and simpler, the structure is also better with `class`.

We have found a really good C++ codes with `class` from:
<https://github.com/HimanshuMohanty-Git24/Neural-Network-CPP.git>
it is made by Himanshu Mohanty with MIT License, and we use it then modify it, so the reader can compare the differences between the original code that we use, and the modified one that we then put into SymIntegration.

We will talk about the theory first: in C++, a `class` is a user-defined data type that serves as a blueprint or template for creating objects. It encapsulates data (attributes) and functions (methods) into a single unit, which is the core principle of Object-Oriented Programming.

A class is defined using the `class` keyword followed by the class name. The body is enclosed in curly braces and must end with a semicolon.

Key components:

- **Attributes**
Variables declared inside a class to hold data.
- **Methods**
Functions that belong to the class and define its behavior.
- **Access Specifiers**
public: Members are accessible from outside the class.
private: Members can only be accessed withing the class itself (default for class).
protected: Members are accessible in inherited classes.
- **Constructors & Destructors**
Special methods used to initialize objects when they are created and clean up resources when they are destroyed.

An object is an instance of a class. You create one by specifying the class name followed by the object name. The difference between class and struct is that class members are private by default while struct members are public by default and structs are used for simple data structure. You can use classes for complex objects with methods and inheritance, if you have tried to play computer games, or complex simulator, or physics engine, they are all using class so you can see the realistic details, effects, how far a person will fall back after hit by a shotgun at a certain distance, the lighting from sunlight that creates shadow of an object, almost like real life.

To create a C++ class across multiple files, you split it into a header file (`.h`) for the class blueprint and a source file (`.cpp`) for the logic. This modular approach speeds up compilation and keeps your project organized.

The header file contains the class declaration, which includes member variables like arrays and function prototypes.

The source file provides the actual logic for the functions declared in the header. You must include the include file `.h` to connect the definition to the declaration, in this book we compile all include files inside `symintegrationc++.h`, so we only need to put this one include file on top of the main source code. Use the class name and `::` before each function name to tell the compiler these functions belong to that specific class.

When compiling, you must include both the `main.cpp` and the `class.cpp` files (e.g., `g++ main.cpp class.cpp -o main`). We have created Makefile for this feed-forward neural network for iris dataset with `class`, so you only need to type `make` to compile it without the need to write the long `g++` compiling formula.

Now, we have learned the basic theory for `class`. Let see the code for the feed-forward neural network for iris dataset with `class`, first in the `include/artificialneuralnetworks.h` we will have

```
...
#ifndef FNN_NOHIDDENLAYER_IRIS_H
#define FNN_NOHIDDENLAYER_IRIS_H

class FNN_NoHiddenLayer_Iris {
private:
    double weights[4][3];
    double bias[3]; // Fixed-size array member
    double learning_rate;

public:
    FNN_NoHiddenLayer_Iris(); // Constructor

    vector<double> predict(const vector<double> &inputs);

    void train(vector<vector<double>> &X, vector<int> &y, int epochs);

    void save_model(const string &filename) ; // Void function declaration
    void load_model(const string& filename);
};

#endif
...
```

Code 63: `include/artificialneuralnetworks.h`

We include guards to prevent the file from being included multiple times, which causes errors.

Now for the `src/artificialneuralnetworks.cpp` source file

```
...
```

```

FNN_NoHiddenLayer_Iris::FNN_NoHiddenLayer_Iris()
{
    for (int i = 0; i < 4; ++i)
    {
        for (int j = 0; j < 3; ++j)
        {
            weights[i][j] = random_double(-0.5, 0.5);
        }
    }
    for (int j = 0; j < 3; ++j)
    {
        bias[j] = random_double(-0.5, 0.5);
    }
    learning_rate = 0.01;
}

double FNN_NoHiddenLayer_Iris::sigmoid(double x)
{
    return 1.0 / (1.0 + std::exp(-x));
}

double FNN_NoHiddenLayer_Iris::sigmoid_derivative(double x)
{
    return x * (1.0 - x);
}

vector<double> FNN_NoHiddenLayer_Iris::predict(const vector<double> &inputs)
{
    vector<double> outputs(3, 0.0);
    for (int j = 0; j < 3; ++j)
    {
        for (int i = 0; i < 4; ++i)
        {
            outputs[j] += inputs[i] * weights[i][j];
        }
        outputs[j] += bias[j];
        outputs[j] = SigmoidActivationfunction(outputs[j]);
    }
    return outputs;
}

...

```

Code 64: *src/artificialneuralnetworks.cpp*

The name of the class is **FNN_NoHiddenLayer_Iris** that initializes array for the weights and bias with fix size, and as we can see in the source file we have created functions of **predict**, **train**, **load_model**, **save_model** under this class.

After we have created the class, now we want to create the **main.cpp**, but in this case we will create two and name it as **train.cpp** and **predict.cpp**, the **train.cpp** has to be run first, and it will be used to obtain the best weights with least error to predict the iris species, while the **predict.cpp** will be used as a prediction cli (command line interface) where we can input the four variables for the irises (the sepal length, sepal width, petal length, and petal width) then the best model obtained from **train.cpp** will predict whether the inputted values belong to either iris Setosa, iris Versicolor or iris Virginica.

```
g++ -c -o predict.o predict.cpp
g++ -o predict -gddb predict.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Neural Networks/Test FNN Iris Dataset Predictor using Class with Steepest Descent ]# ./train
```

===== Iris Classification with Neural Network =====

Loading dataset...

DEBUG: Dataset loaded. Size: 150 samples, 4 features

DEBUG: Number of labels: 150

DEBUG: Label names:

DEBUG: - Iris-setosa

DEBUG: - Iris-versicolor

DEBUG: - Iris-virginica

DEBUG: Train set size: 120

DEBUG: Test set size: 150

Initializing Neural Network...

Starting Training Process

Training Progress:

[=====] 100% Epoch 1000/1000 - Loss: 0.2084

Training completed in 1.0358 seconds.

Evaluating the model on the test set:

[=====] 100%

Accuracy: 96.67%

Confusion Matrix:

Predicted >	Iris-setosa	Iris-versicolor	Iris-virginica
Iris-setosa	50	0	0
Iris-versicolor	0	47	3
Iris-virginica	0	2	48

Figure 10.11: The training process for the iris dataset without hidden layer and the activation function at the output layer is sigmoid with *FNN_NoHiddenLayer_Iris* class (*SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor using Class with Steepest Descent/train.cpp*).


```

Confusion Matrix:
      Predicted >
      Iris-setosa  Iris-versicolor  Iris-virginica
Iris-setosa      50                0                0
Iris-versicolor  0                47                3
Iris-virginica   0                2               48

Precision, Recall, and F1-score:

Iris-setosa:
Precision: 1.00
Recall:    1.00
F1-score:  1.00

Iris-versicolor:
Precision: 0.96
Recall:    0.94
F1-score:  0.95

Iris-virginica:
Precision: 0.94
Recall:    0.96
F1-score:  0.95

The model weights:
      0.187405      0.959035      -2.161684
      1.850920     -2.421174     -2.644875
      -2.292653     0.714200     3.410055
      -1.058507     -2.371125     3.601225

The model bias:
0.378021
1.410220
-1.612257

```

Figure 10.12: The evaluation of the model obtained from the training process for the iris dataset without hidden layer and the activation function at the output layer is sigmoid with `FNN_NoHiddenLayer_Iris` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor using Class with Steepest Descent/train.cpp`).

```

=====Iris Species Predictor=====

Enter four input features (or type 'exit' to quit):
1. Sepal length (cm): 4.5
2. Sepal width (cm): 5.1
3. Petal length (cm): 5.3
4. Petal width (cm): 4.6

Predicting...
[=====] 100%

Prediction Results:
-----
Predicted Species: Iris-virginica

Probabilities:
0.17%      Iris-setosa
0.00%      Iris-versicolor
99.99%     Iris-virginica

Enter four input features (or type 'exit' to quit):
1. Sepal length (cm): 

```

Figure 10.13: The prediction of which iris species belong to based on the given input of sepal length, sepal width, petal length, and petal width that user will give in the cli / terminal / xterm with `FNN_NoHiddenLayer_Iris` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor using Class with Steepest Descent/predict.cpp`).

We create another class as comparison, `FNN_1HiddenLayer_Iris`, we set the hidden layer to have 5 neurons (it can be changed by user for experimentation purpose) and the activation function at the hidden layer is sigmoid, then the activation function at the output layer is using softmax. The input neurons stay the same, 4 neurons, and the output neurons still 3 neurons / 3 classifications.

```

[=====] 99% Epoch 990/1000 - Loss:
: 0.0393

Training completed in 3.1363 seconds.

Evaluating the model on the test set:
[=====] 100%

Accuracy: 99.33%

Confusion Matrix:

    Predicted >      Iris-setosa  Iris-versicolor  Iris-virginica
Iris-setosa         50             0              0
Iris-versicolor      0             49             1
Iris-virginica       0             0             50

Precision, Recall, and F1-score:

Iris-setosa:
Precision: 1.00
Recall: 1.00
F1-score: 1.00

Iris-versicolor:
Precision: 1.00
Recall: 0.98
F1-score: 0.99

Iris-virginica:
Precision: 0.98
Recall: 1.00
F1-score: 0.99

```

Figure 10.14: The training process for the iris dataset with 1 hidden layer consisting of 5 neurons with sigmoid activation function at the hidden layer and the activation function at the output layer is softmax with `FNN_1HiddenLayer_Iris` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor 1 Hidden Layer using Class with Steepest Descent/train.cpp`).

```

The model weights:
    2.873544      1.181453      -1.193326
7.713539      -0.435190      0.408036
4.772574      9.565981      0.315779
-1.376258      -4.499668      -0.710437
-7.864477      -4.905031      -0.013974
-6.072253      -0.013974      -1.957176
-7.206076      -4.734083

The model hiddenweights:
    1.779509      1.400948      -0.717926
-0.608578      2.371403      4.947981
7.136587      -3.220675      -2.177980
2.291308      3.554297      -3.039231
2.814101      2.133686      -2.283374

The model bias in the hidden layer:
4.320484
0.342381
0.773372
4.789821
4.492461

The model bias in the output layer:
-0.669905
2.417736
4.461967

```

Figure 10.15: The training process for the iris dataset with 1 hidden layer consisting of 5 neurons with sigmoid activation function at the hidden layer and the activation function at the output layer is softmax with `FNN_1HiddenLayer_Iris` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor 1 Hidden Layer using Class with Steepest Descent/train.cpp`).

```

Accuracy: 98.67%
Confusion Matrix:
      Predicted >      Iris-setosa  Iris-versicolor  Iris-virginica
Iris-setosa          50             0              0
Iris-versicolor       0             48              2
Iris-virginica        0             0              50

Precision, Recall, and F1-score:

Iris-setosa:
Precision: 1.00
Recall:    1.00
F1-score:  1.00

Iris-versicolor:
Precision: 1.00
Recall:    0.96
F1-score:  0.98

Iris-virginica:
Precision: 0.96
Recall:    1.00
F1-score:  0.98

The model weights:
      0.613851      1.612227      -1.453502      5.644846      -1.401964
      0.938255      7.728365      4.515118      1.287156      9.714031
      -1.532171     -7.595027     -1.284246     -7.360822     -5.407066
      -1.808880     -3.781532     -1.591036     -6.282293     -2.025311

The model hiddenweights:
      5.011217      -2.101302      -1.622205
      1.588085      2.883951      -2.936433
      4.346169      -0.099619      -1.798401
      1.526841      3.492190      -3.579990
      2.273748      1.714440      -2.228758

The model bias in the hidden layer:
0.430315
8.906203
1.564328
6.755378
6.497567

The model bias in the output layer:
-2.732761
2.047286
6.477377

```

Figure 10.16: The training process for the iris dataset with 1 hidden layer consisting of 5 neurons with sigmoid activation function at the hidden layer and the activation function at the output layer is softmax with `FNN_1HiddenLayer_Iris` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor 1 Hidden Layer using Class with Steepest Descent/train.cpp`).

We have used standard gradient descent for training the neural network, now we want to compare if we are using a Conjugate Gradient (CG) algorithm for a feed-forward neural network. For a small dataset like Iris, it makes the neural network to converge faster than standard gradient descent. The approach involves representing the network weights as a vector, using backpropagation to compute the gradient, and using a line search to update weights along a conjugate direction, reducing MSE (Mean Squared Error).

```

Starting Training Process
Training Progress:
[=====] 100% Epoch 200/200 - Loss: 0.1882
Training completed in 16.8021 seconds.
Evaluating the model on the test set:
[=====] 100%
Accuracy: 98.67%
Confusion Matrix:
    Predicted >      Iris-setosa  Iris-versicolor  Iris-virginica
Iris-setosa         50             0              0
Iris-versicolor      0            48              2
Iris-virginica       0             0             50
Precision, Recall, and F1-score:
Iris-setosa:
Precision: 1.00
Recall: 1.00
F1-score: 1.00
Iris-versicolor:
Precision: 1.00
Recall: 0.96
F1-score: 0.98
Iris-virginica:
Precision: 0.96
Recall: 1.00
F1-score: 0.98
The model weights:
    -1.062172      0.314579      -0.529218      -1.157496      -0.764458
     0.164349      -0.401029      0.405046      -0.871413      -0.656090
    -0.529516      -0.368303      -0.055132      1.942797      -0.536916
     0.861717      -0.085202      -0.250408      2.666763      -0.354715
The model hiddenweights:
    -0.762174      -0.906450      0.233996
    -0.583184      -0.151056      0.851803
     0.150078      0.674797      0.686892
    -2.535796      0.608768      1.583251
     1.014366      0.083956      -0.385530

```

Figure 10.17: The training process for the iris dataset with 1 hidden layer consisting of 5 neurons with ReLU activation function at the hidden layer and the activation function at the output layer is softmax using conjugate gradient method with `FNN_1HiddenLayer_Iris_ConjugateGradient` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor 1 Hidden Layer using Class with Conjugate Gradient/train.cpp`).

III. Supervised Learning: Feed-Forward Neural Network (Multilayer Perceptron)

```
Predicted > Iris-setosa Iris-versicolor Iris-virginica
Iris-setosa      50         0         0
Iris-versicolor   0        48         2
Iris-virginica    0         0        50

Precision, Recall, and F1-score:

Iris-setosa:
Precision: 1.00
Recall: 1.00
F1-score: 1.00

Iris-versicolor:
Precision: 1.00
Recall: 0.96
F1-score: 0.98

Iris-virginica:
Precision: 0.96
Recall: 1.00
F1-score: 0.98

The model weights:
-1.062172      0.314579      -0.529218      -1.157496      -0.764458
 0.164349      -0.401029      0.405046      -0.871413      -0.656090
-0.529516      -0.368303      -0.055132      1.942797      -0.536916
 0.061717      -0.085202      -0.250408      2.666763      -0.354715

The model hiddenweights:
-0.762174      -0.906450      0.233996
-0.583184      -0.151056      0.851803
 0.150078      0.674797      0.686892
-2.535796      0.608768      1.583251
 1.014366      0.083956      -0.385530

The model bias in the hidden layer:
-0.234676
-0.074763
0.776133
0.430983
-0.022482

The model bias in the output layer:
2.603302
0.581319
-3.850410

Model trained and saved successfully!
```

Figure 10.18: The training process for the iris dataset with 1 hidden layer consisting of 5 neurons with ReLU activation function at the hidden layer and the activation function at the output layer is softmax using conjugate gradient method with `FNN_1HiddenLayer_Iris_ConjugateGradient` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor 1 Hidden Layer using Class with Conjugate Gradient/train.cpp`).

The conjugate gradient method for training neural network on this iris dataset able to achieve 98.67% of accuracy with loss of 0.1882 within 200 epochs for training of 80% of data (120 samples) and the testing is 150 samples, 100% of data. It is sometimes said that very high accuracy is overfitting and 80% of accuracy is the best because it is not too high nor too low, for some cases perhaps it is overfitting, but it depends on the case or problem that we are working on.

```

Initializing Neural Network...
Starting Training Process
Training Progress:
[=====] 100% Epoch 200/200 - Loss: 0.0957

Training completed in 16.1711 seconds.

Evaluating the model on the test set:
[=====] 100%

Accuracy: 98.00%

Confusion Matrix:
Predicted >      Iris-setosa  Iris-versicolor  Iris-virginica
Iris-setosa      50           0             0
Iris-versicolor  0           49            1
Iris-virginica   0           2            48

Precision, Recall, and F1-score:

Iris-setosa:
Precision: 1.00
Recall:    1.00
F1-score:  1.00

Iris-versicolor:
Precision: 0.96
Recall:    0.98
F1-score:  0.97

Iris-virginica:
Precision: 0.98
Recall:    0.96
F1-score:  0.97

The model weights:
-2.246316      0.032038      -0.108140      -0.698284      0.110260
-1.998872     -1.095444      1.460190      0.234610     -0.311328
 3.769575     -0.062567     -1.270523     -0.873604     -0.323579
 2.649634     -0.119740      0.108883     -1.249114      0.210217

The model hiddenweights:
-1.363625     -0.001630      1.513059
 0.440795     -1.212376      1.172248
 1.322513     -1.098335     -0.673634
-0.451245     1.459457     -0.940929
-0.701461     -1.498091     -0.666894

```

Figure 10.19: The training process for the iris dataset with 1 hidden layer consisting of 5 neurons with ReLU activation function at the hidden layer and the activation function at the output layer is softmax using conjugate gradient method with `FNN_1HiddenLayer_Iris_ConjugateGradient` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor 1 Hidden Layer using Class with Conjugate Gradient/train.cpp`).

```

Confusion Matrix:
      Predicted >      Iris-setosa  Iris-versicolor  Iris-virginica
Iris-setosa           50             0             0
Iris-versicolor        0             49             1
Iris-virginica         0             2             48

Precision, Recall, and F1-score:

Iris-setosa:
Precision: 1.00
Recall: 1.00
F1-score: 1.00

Iris-versicolor:
Precision: 0.96
Recall: 0.98
F1-score: 0.97

Iris-virginica:
Precision: 0.98
Recall: 0.96
F1-score: 0.97

The model weights:
      -2.246316      0.032038      -0.108140      -0.698284      0.110260
      -1.998872      -1.095444      1.460190      0.234610      -0.311328
      3.769575      -0.062567      -1.270523      -0.873604      -0.323579
      2.649634      -0.119740      0.108883      -1.249114      0.210217

The model hiddenweights:
      -1.363625      -0.001630      1.513059
      0.440795      -1.212376      1.172248
      1.322513      -1.098335      -0.673634
      -0.451245      1.459457      -0.940929
      -0.701461      -1.498091      -0.666894

The model bias in the hidden layer:
-0.653320
0.002011
1.277995
-1.167769
0.554871

The model bias in the output layer:
-0.650087
2.571865
-1.626273

Model trained and saved successfully!

```

Figure 10.20: The training process for the iris dataset with 1 hidden layer consisting of 5 neurons with ReLU activation function at the hidden layer and the activation function at the output layer is softmax using conjugate gradient method with `FNN_1HiddenLayer_Iris_ConjugateGradient` class (`SymIntegration/Examples/Neural Networks/Test FNN Iris Dataset Predictor 1 Hidden Layer using Class with Conjugate Gradient/train.cpp`).

In this C++ class code we are using a beta that we create on our own, since Polak-Ribiere, Fletcher Reeves or even OFR [31], they cannot converge and only lead to flat loss or oscillating at high loss or can only predict one class / stuck at local minimum even after a lot of re-training.

The beta formula that we are using is:

$$\beta_k^G = \begin{cases} \frac{\|g_k\|}{\|g_{k-1}\|}, & \text{if } \beta_{k-1} \geq \frac{\|g_k\|}{\|g_{k-1}\|} \\ 0.01, & \text{if } \beta_{k-1} < \frac{\|g_k\|}{\|g_{k-1}\|} \end{cases}$$

The 0.01 is the smaller decimal value of the input variables, when we see the sepal length, sepal width, petal length and petal width, they are all have 1 decimal number. This beta perhaps only work for iris dataset and need more improvements to be generalized for other dataset. This beta also sometimes stuck or has flat loss, but if you call the code again for another training (call `.train` again) then it will converge and has an amazing small loss ± 0.1 after 200 epochs. The code can be seen in `src/artificialneuralnetworks.cpp`

...

```
FNN_1HiddenLayer_Iris_ConjugateGradient::FNN_1HiddenLayer_Iris_ConjugateGradient()
```

```

{
    int n_classification = 3;
    int n_hiddenlayer = 5;
    int n_input = 4;
    for (int i = 0; i < n_input; ++i)
    {
        for (int j = 0; j < n_hiddenlayer; ++j)
        {
            weights[i][j] = He_initialization(n_input);
        }
    }
    for (int j = 0; j < n_hiddenlayer; ++j)
    {
        for (int k = 0; k < n_classification; ++k)
        {
            hiddenweights[j][k] = He_initialization(n_input);
        }
    }
    for (int j = 0; j < n_hiddenlayer; ++j)
    {
        biashidden[j] = He_initialization(n_input);
    }
    for (int k = 0; k < n_classification; ++k)
    {
        biasoutput[k] = He_initialization(n_input);
    }
    learning_rate = 0.51;
}

vector<double> FNN_1HiddenLayer_Iris_ConjugateGradient::predict(const vector<
double> &inputs)
{
    int n_classification = 3;
    int n_hiddenlayer = 5;
    int n_input = 4;
    vector<double> hiddenlayer(n_hiddenlayer, 0.0);
    vector<double> outputs(n_classification, 0.0);
    for (int j = 0; j < n_hiddenlayer; ++j)
    {
        for (int i = 0; i < n_input; ++i)
        {
            hiddenlayer[j] += inputs[i] * weights[i][j];
        }
        hiddenlayer[j] += biashidden[j];
        hiddenlayer[j] = ReLUActivationfunction(hiddenlayer[j]);
        //hiddenlayer[j] = TanhActivationfunction(hiddenlayer[j]);
    }
    for (int k = 0; k < n_classification; ++k)

```



```
{
    for (int j = 0; j < n_hiddenlayer; ++j)
    {
        outputs[k] += hiddenlayer[j] * hiddenweights[j][k];
    }
    outputs[k] += biasoutput[k];
}
vector<double>predicted_output_softmax =
    SoftMax_vectorresult_activationfunction(outputs);
return predicted_output_softmax;
}

void FNN_1HiddenLayer_Iris_ConjugateGradient::train(vector<vector<double>> &X,
    vector<int> &y, int epochs)
{
    ...
    ...
}
```

Code 65: *src/artificialneuralnetworks.cpp*

The class that handles the training and testing for the iris dataset specifically with 1 hidden layer, we initialize the weights and bias with He initialization because we are using ReLU as the activation function on the hidden layer.

iii. Functions in SymIntegration Related to Feed-Forward Neural Network

[SI*] We will list the basic functions that can be used for computing feed-forward neural networks:

FNN_1hiddenlayer(vector<double>& x , vector<vector<double>>& W , vector<vector<double>>& W_{hidden} , vector<double>& b , vector<double>& y)

this function computes and updates the weights parameter from input to output layer, with only 1 hidden layer that has m nodes, we need the input of the vector x as the vector of input layer with size of n , vector b as the bias with size of m , vector y as the vector of output layer, the matrix W as the $m \times n$ matrix that represents the weights from the input layer to the hidden layer, the matrix W_{hidden} as the $k \times m$ matrix from the hidden layer to the output layer. The activation function is using sigmoid, the parameter η (learning rate) is set to 0.51, and the iteration is set to 5, and these parameters can be modified in **src/artificialneuralnetwork.cpp**.

FNN_1hiddenlayer(vector<double>& x , vector<double>& y , int n_{hidden})

this function computes the same thing as the previous function, but we only need to give the input of x the input vector, and y the output vector, and n_{hidden} as the number of nodes in the hidden layer. The weight matrix W and the hiddenweight matrix W_{hidden} will be generated with Mersenne Twister engine with uniform distribution from 0.1 to 1. The activation function is using sigmoid, the parameter η (learning rate) is set to 0.51, and the iteration is set to 5.

```
Feed-forward Neural Network with 1 hidden layer type 2:
Weights from input to hidden layer:
    0.253954    0.490407
    0.606950    0.590103

Weights from hidden layer to output layer:
    0.219816    0.417713
    0.720708    0.139265

Hidden layer:
0.606391
0.651469

Predicted Output:
0.599990
0.628960

Error vector:
-0.099990
-0.258960

||Error|| = 0.277594

*****
                        Backpropagation iteration
*****

Learning rate: 0.510000

Iteration = 1

Delta output vector:
-0.023998
-0.060433

Delta hidden layer vector:
-0.011655
-0.004187

New weights matrix:
    0.251873    0.486246
    0.606203    0.588608

New hiddenweights matrix:
    0.212395    0.409740
    0.702018    0.119186

Hidden layer:
0.605522
0.651172
```

Figure 10.21: The computation of backpropagation process for the feed-forward neural network with **FNN_1hiddenlayer()** function (**SymIntegration/Examples/Neural Networks/Test Backpropagation FNN/main.cpp**).

```

Iteration = 5

Delta output vector:
-0.021779
-0.055120

Delta hidden layer vector:
-0.009554
-0.002689

New weights matrix:
           0.244511      0.471521
           0.603901      0.584005

New hiddenweights matrix:
           0.184541      0.379744
           0.631648      0.043407

Hidden layer:
0.602440
0.650257

Predicted Output:
0.588582
0.600790

Error vector:
-0.088582
-0.230790

||Error|| = 0.247206

Time taken by function: 6397 microseconds

```

Figure 10.22: The computation of backpropagation process for the feed-forward neural network with `FNN_1hiddenlayer()` function (*SymIntegration/Examples/Neural Networks/Test Backpropagation FNN/main.cpp*).

FNN_nohiddenlayer_irisdataset_training(vector<vector<double>>& X, vector<string>& y)

this function trains the neural network for iris dataset, with no hidden layer, so the input matrix X is a matrix of size $n \times 4$, 4 is the input variables (sepal length, sepal width, petal length, petal width) and n is the number of data that will be trained, normally 80% of total data, while vector y is a vector of string of the classification for the input variables, the classification consists of Iris Setosa, Iris Versicolor, and Iris Virginica. The activation function at the output layer is set to sigmoid, and can be changed by the user in `src/artificialneuralnetworks.cpp`.

FNN_nohiddenlayer_irisdataset_testing(vector<vector<double>>& X, vector<string>& y)
this function tests the optimal weight obtained from the function:

FNN_nohiddenlayer_irisdataset_training(vector<vector<double>>& X, vector<string>& y)

for iris dataset, with no hidden layer, so the input matrix X is a matrix of size $m \times 4$, 4 is the input variables (sepal length, sepal width, petal length, petal width) and m is the number of data that will be tested, normally 20% of total data, it is the data that is not trained that will be tested, while vector y is a vector of string of the classification for the input variables, the classification consists of Iris Setosa, Iris Versicolor, and Iris Virginica. The activation function at the output layer is set to sigmoid, and can be changed by the user in `src/artificialneuralnetworks.cpp`.

FNN_1hiddenlayer_irisdataset_training(vector<vector<double>>& X, vector<string>& y)

this function trains the neural network for iris dataset, with 1 hidden layer and 5 number of neurons in the hidden layer by default, so the input matrix X is a matrix of size $n \times 4$, 4 is the input variables (sepal length, sepal width, petal length, petal width) and n is the number of data that will be trained, normally 80% of total data, while vector y is a vector of string of the classification for the input variables, the classification consists of Iris Setosa, Iris Versicolor, and Iris Virginica. The activation function at the output layer is softmax,

the activation function at the hidden layer is sigmoid, and can be changed by the user in `src/artificialneuralnetworks.cpp`.

FNN_1hiddenlayer_irisdataset_testing(vector<vector<double>>& X, vector<string>& y)

this function tests the optimal weight obtained from the function:

FNN_1hiddenlayer_irisdataset_training(vector<vector<double>>& X, vector<string>& y)

for iris dataset, with 1 hidden layer and 5 number of neurons in the hidden layer by default, so the input matrix X is a matrix of size $m \times 4$, 4 is the input variables (sepal length, sepal width, petal length, petal width) and m is the number of data that will be tested, normally 20% of total data, it is the data that is not trained that will be tested, while vector y is a vector of string of the classification for the input variables, the classification consists of Iris Setosa, Iris Versicolor, and Iris Virginica. The activation function at the output layer is softmax, the activation function at the hidden layer is sigmoid, and can be changed by the user in `src/artificialneuralnetworks.cpp`.

IV. SUPERVISED LEARNING: SUPPORT VECTOR MACHINES (SVM)

[SI*] Support vector machines is a classification method based on maximum margin linear discriminants, that is, the goal is to find the optimal hyperplane (a line in 2D or a flat plane in 3D) that maximizes the gap or margin between the classes. Further, we can use the kernel trick to find the optimal nonlinear decision boundary between classes, which corresponds to a hyperplane in some high-dimensional "nonlinear" space.

V. UNSUPERVISED LEARNING: PRINCIPAL COMPONENT ANALYSIS (PCA)

[SI*] Unsupervised learning is the second type of machine learning which finds patterns in unlabeled data, grouping similar items (clustering) or reducing data dimensions. Principal Component Analysis reduces the complexity (dimensionality) of the data by combining highly correlated features into fewer, uncorrelated "principal components" that capture the most variance.

VI. HYBRID LEARNING: KERNEL METHODS

[SI*] A mathematical technique that takes non-linearly separable data and implicitly projects it into a higher-dimensional space where a flat line or plane can separate them. Crucially, it does this without the heavy computations of explicitly mapping the data.

[SI*] It is important to find a suitable data representation that facilitates data analysis. For example, for complex data such as text, sequences, images, and so on, we must typically extract or construct a set of attributes or features, so that we can represent the data instances as multivariate vectors.

That is, given a data instance vector x (e.g., a sequence), we need to find a mapping ϕ , so that $\phi(x)$ is the feature representation of x . Even when the input data is a numeric data matrix, if we wish to discover nonlinear relationships among the attributes, then a nonlinear mapping ϕ may be used, so that $\phi(x)$ represents a vector in the corresponding high-dimensional space comprising nonlinear attributes.

We use the term input space to refer to the data space for the input data x and feature space

to refer to the space of mapped vectors $\phi(x)$. Thus, given a set of data objects or instances x_i , and given a mapping function ϕ , we can transform them into feature vectors $\phi(x_i)$, which then allows us to analyze complex data instances via numeric analysis methods.

[SI*] **Sequence-based Features Example**

Consider a dataset of DNA sequences over the alphabet $\Sigma = \{A, C, G, T\}$. One simple feature space is to represent each sequence in terms of the probability distribution over symbols in Σ . That is, given a sequence x with length $|x| = m$, the mapping into feature space is given as

$$\phi(x) = \{P(A), P(C), P(G), P(T)\}$$

where $P(s) = \frac{n_s}{m}$ is the probability of observing symbol $s \in \Sigma$, and n_s is the number of times s appears in sequence x . Here the input space is the set of sequence Σ^* , and the feature space is $\mathbb{R}^4$. For example, if $x = ACAGCAGTA$, with $m = |x| = 9$, since A occurs four times, C and G occur twice, and T occurs once, we have

$$\phi(x) = \left(\frac{4}{9}, \frac{2}{9}, \frac{2}{9}, \frac{1}{9} \right) = (0.44, 0.22, 0.22, 0.11)$$

The mapping ϕ now allows one to compute the statistics over the data sample to make inferences about the population. For example, we may compute the mean symbol composition. We can also define the distance between any two sequences, for example

$$\begin{aligned} \delta(x, y) &= ||\phi(x) - \phi(y)|| \\ &= \sqrt{(0.44 - 0.4)^2 + (0.22 - 0.2)^2 + (0.22 - 0.4)^2 + (0.11 - 0)^2} \\ &= 0.22 \end{aligned}$$

We can compute larger feature spaces by considering, for example, the probability distribution over all substrings or words of size up to k over the alphabet Σ , and so on.

[SI*] **Nonlinear Features Example**

Consider the mapping ϕ that takes as input a vector $x = (x_1, x_2) \in \mathbb{R}^2$ and maps it to a "quadratic" feature space via the nonlinear mapping

$$\phi(x) = (x_1^2, x_2^2, \sqrt{2}x_1x_2)$$

with $\phi \in \mathbb{R}^3$. For example, the point $x = (5.9, 3)$ is mapped to the vector

$$\phi(x) = (5.9^2, 3^2, \sqrt{2}(5.9)(3)) = (34.81, 9, 25.03)$$

The main benefit of this transformation is that we may apply well-known linear analysis methods in the feature space. However, because the features are nonlinear combinations of the original attributes, this allows us to extract / mine the nonlinear patterns and relationships. For someone aware enough and have read Linear Algebra book this is actually a linear transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$.

[SI*] Whereas mapping into feature space allows one to analyze the data via algebraic and probabilistic modeling, the resulting feature space is usually very high-dimensional; it may even be infinite dimensional. Thus, transforming all the input points into feature space can be very expensive.

Kernel methods avoid explicitly transforming each points x in the input space into the mapped point $\phi(x)$ in the feature space. Instead, the input objects are represented via their $n \times n$ pairwise similarity values. The similarity function, called a kernel, is chosen so that it

represents a dot product in some high-dimensional feature space, yet it can be computed without directly constructing $\phi(x)$.

Let $\mathbb{I}$ denote the input space, which can comprise any arbitrary set of objects, and let $d = \{x_i\} \in \mathbb{I}, i = 1, \dots, n$ be a dataset comprising n objects in the input space. We can represent the pairwise similarity values between points in d via the $n \times n$ kernel matrix, defined as

$$K = \begin{bmatrix} K(x_1, x_1) & K(x_1, x_2) & \dots & K(x_1, x_n) \\ K(x_2, x_1) & K(x_2, x_2) & \dots & K(x_2, x_n) \\ \vdots & \vdots & \ddots & \vdots \\ K(x_n, x_1) & K(x_n, x_2) & \dots & K(x_n, x_n) \end{bmatrix}$$

where $K : \mathbb{I} \times \mathbb{I} \rightarrow \mathbb{R}$ is a kernel function on any two points in input space. However, we require that K corresponds to a dot product in some feature space. That is, for any $x_i, x_j \in \mathbb{I}$, the kernel function should satisfy the condition

$$K(x_i, x_j) = \phi(x_i)^T \phi(x_j) \quad (10.67)$$

where $\phi : \mathbb{I} \rightarrow \mathbb{F}$ is a mapping from the input space $\mathbb{I}$ to the feature space $\mathbb{F}$. Intuitively, this means that we should be able to compute the value of the dot product using the original input representation x , without having recourse to the mapping $\phi(x)$. Obviously, not just any arbitrary function can be used as a kernel; a valid kernel function must satisfy certain conditions so that Eq. (10.67) remains valid.

It is important to remark that the transpose operator for the dot product applies only when $\mathbb{F}$ is an abstract vector space with an inner product, the kernel is written as $K(x_i, x_j) = \langle \phi(x_i), \phi(x_j) \rangle$. However, for convenience we use the transpose operator, when $\mathbb{F}$ is an inner product space it should be understood that

$$\phi(x_i)^T \phi(x_j) = \langle \phi(x_i), \phi(x_j) \rangle$$

[SI*] Linear Kernel Example

Consider the identity mapping, $\phi(x) \rightarrow x$. This naturally leads to the linear kernel, which is simply the dot product between two input vectors, and thus satisfies Eq. (10.67):

$$\phi(x)^T \phi(y) = x^T y = K(x, y)$$

For example, consider the six points from the Iris dataset:

$$x_1 = \begin{bmatrix} 5.1 \\ 3.5 \\ 1.4 \\ 0.2 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 4.9 \\ 3.0 \\ 1.4 \\ 0.2 \end{bmatrix}, \quad x_3 = \begin{bmatrix} 4.7 \\ 3.2 \\ 1.3 \\ 0.2 \end{bmatrix}, \quad x_4 = \begin{bmatrix} 4.6 \\ 3.1 \\ 1.5 \\ 0.2 \end{bmatrix}, \quad x_5 = \begin{bmatrix} 5.0 \\ 3.6 \\ 1.4 \\ 0.2 \end{bmatrix}, \quad x_6 = \begin{bmatrix} 5.4 \\ 3.9 \\ 1.7 \\ 0.4 \end{bmatrix}$$

The formula to compute the entry for the i -th row and j -th column in the kernel matrix is

$$K(x_i, x_j) = x_i^T x_j$$

The kernel matrix for the linear kernel is

$$K = \begin{bmatrix} 40.26 & 37.49 & 37.03 & 36.45 & 40.10 & 43.65 \\ 37.49 & 35.01 & 34.49 & 33.98 & 37.30 & 40.62 \\ 37.03 & 34.49 & 34.06 & 33.53 & 36.88 & 40.15 \\ 36.45 & 33.98 & 33.53 & 33.06 & 36.30 & 39.56 \\ 40.10 & 37.30 & 36.88 & 36.30 & 39.96 & 43.50 \\ 43.65 & 40.62 & 40.15 & 39.56 & 43.50 & 47.42 \end{bmatrix}$$

VII. REINFORCEMENT LEARNING

- [SI*] Reinforcement learning is the third categories in machine learning, the algorithms learns to make decisions through trial and error in an interactive environment, aiming to maximize some notion of cumulative reward.

VIII. DEEP LEARNING: CONVOLUTIONAL NEURAL NETWORK

- [SI*] The convolutional neural network (CNN) is a type of artificial neural network that is effective for processing spatial relationships in complex images. CNN scans the image (or sequence of pixels) to learn spatial patterns, stroke thickness, curves, and angles. It is perfect when you are building an isolated character or digit classifier (e.g. recognizing single digits 0 – 9 or single alphabet letters).

The convolutional neural networks (CNNs) are mainly designed for image data they use convolution operations to reduce the number of parameters needed, it is one of two categories in deep learning (the other one is RNN / recurrent neural networks), deep learning is a subset of machine learning that involves neural networks with multiple layers. By mimicking the human visual system, CNNs automatically learn to detect spatial patterns-such as edges, shapes, and textures-without requiring manual feature extraction.

CNNs are specifically designed to process data with a grid-like topology, such as images, represented as grids of pixels. CNNs serve as a cornerstone technology in the field of computer vision and posses robust capabilities for processing multimedia data. Their applications beside digit recognition ranging from facial recognition (for security and surveillance), object classification to medical image analysis (assisting professionals by detecting anomalies in X-rays, MRIs, and CT scans).

- [SI*] **Key Components of CNN**

A standard CNN architecture consists of three main types of layers:

- **Convolutional Layer**

The core building block. It applies learnable filters (kernels) that slide over the input data to calculate dot products, generating "feature maps" that highlighting specific patterns like lines or curves..

- **Pooling Layer**

Applied after convolutional layers to reduce the spatial dimensions of the data (down-sampling). This minimizes computational load, limits memory usage, and helps prevent the model from overfitting.

- **Fully Connected (Dense) Layer**

Flattens the multi-dimensional feature maps into a 1D vector and connects every neuron to every neuron in the preceding and succeeding layers. It performs high-level reasoning and outputs the final classification scores or probabilities.

Between these layers, activation functions (most commonly ReLU) are used to introduce non-linearity into the network. This allows the model to learn complex patterns and relationships rather than just simple linear combinations.




Figure 10.23: The illustration of convolutional neural network pipeline for image classification.

[SI*] CNN: How Computers See Images

When we have an image saved in a computer, the computer sees the image as a matrix of numbers $[0, 255]$, it can be represented as 3D matrix of size $1080 \times 1920 \times 3$, 1080 represents the width of the image, 1920 represents the height of the image and 3 is the depth to represents the RGB color for each pixel.

RGB can be represented as vector of size 3, with domain $[0, 255] \in \mathbb{N}$, for example:

$[255, 0, 0] \rightarrow \text{Red}$
 $[0, 255, 0] \rightarrow \text{Green}$
 $[0, 0, 255] \rightarrow \text{Blue}$
 $[0, 0, 0] \rightarrow \text{Black}$
 $[255, 255, 255] \rightarrow \text{White}$
 $[0, 255, 255] \rightarrow \text{Cyan}$

If we are having a matrix with depth of 1, e.g. a picture of $1080 \times 1920 \times 1$, then we are dealing with a grayscale image, it means each pixel can take a value of $[0, 255]$, with 0 represents black, and 255 represents white, the image will not be colorful as the 3D matrix, because it is grayscale, it is easier and faster for CNN computation with grayscale matrix.




Figure 10.24: An example of a grayscale image.

[SI*] How Convolutional Neural Network Works

Suppose we have an image of size $6 \times 6 \times 1$, a yellowscale image with the yellowscale representation of the image like this:

$$I = \begin{bmatrix} 0 & 0 & 0 & 200 & 100 & 0 \\ 0 & 200 & 100 & 0 & 0 & 0 \\ 0 & 200 & 100 & 100 & 0 & 0 \\ 0 & 0 & 0 & 100 & 200 & 0 \\ 0 & 0 & 200 & 100 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

to extract a feature of that image we will use a kernel, M_{kernel} , a square matrix of size k , with $k < w$ and $k < h$, w is the width of the input image, and h is the height of the input image.

Thus we will take submatrix of $k \times k$ from the input image starting from the top left moving to the right and then down till finishes at the bottom right and each time we perform the operation with the kernel matrix M_{kernel} by using hadamard product (represented by $\oplus$), after the hadamard product operation we add all the number in the resulting matrix to obtain a single number that will be the input for the output map.

Suppose we will use the kernel of vertical edges of size 3:

$$M_{kernel} = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 0 & -1 \\ 1 & 0 & -1 \end{bmatrix}$$

We will start to show the example from the top left of matrix I , we will have

$$I_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 200 & 100 \\ 0 & 200 & 100 \end{bmatrix}$$

multiply it with the kernel we will obtain

$$I_1 \oplus M_{kernel} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 200 & 100 \\ 0 & 200 & 100 \end{bmatrix} \oplus \begin{bmatrix} 1 & 0 & -1 \\ 1 & 0 & -1 \\ 1 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -100 \\ 0 & 0 & -100 \end{bmatrix}$$

we add up all the entries in the resulting matrix thus we will obtain

$$o_{11} = \sum_{ij} I_1 \oplus M_{kernel} = -200$$

then we will move to the right 1 pixel and working on the next submatrix

$$I_2 = \begin{bmatrix} 0 & 0 & 200 \\ 200 & 100 & 0 \\ 200 & 100 & 100 \end{bmatrix}$$

multiply it with the kernel we will obtain

$$I_2 \oplus M_{kernel} = \begin{bmatrix} 0 & 0 & 200 \\ 200 & 100 & 0 \\ 200 & 100 & 100 \end{bmatrix} \oplus \begin{bmatrix} 1 & 0 & -1 \\ 1 & 0 & -1 \\ 1 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & -200 \\ 200 & 0 & 0 \\ 200 & 0 & -100 \end{bmatrix}$$

we add up all the entries in the resulting matrix thus we will obtain

$$o_{12} = \sum_{ij} I_2 \oplus M_{kernel} = 100$$

we iterate until the last bottom right submatrix of the input matrix, then we will have the output map of size $o_w \times o_h$ with

$$\begin{aligned} o_w &= w - k + 1 \\ o_h &= h - k + 1 \end{aligned}$$

thus we will obtain the output map like this:

$$O = \begin{bmatrix} o_{11} & o_{12} & \dots & o_{1,w-k+1} \\ o_{21} & o_{22} & \dots & o_{2,w-k+1} \\ \vdots & \vdots & \ddots & \vdots \\ o_{h-k+1,1} & o_{h-k+1,2} & \dots & o_{h-k+1,w-k+1} \end{bmatrix} = \begin{bmatrix} -200 & 100 & 100 & 300 \\ -200 & 200 & 0 & 200 \\ -300 & -100 & 100 & 300 \\ -200 & -200 & 0 & 200 \end{bmatrix}$$

```

root [ ~/SourceCodes/CPP/C++ Create Library/Test SymIntegration CNN Compute Output Map from Convolution Operation ]# ./main

Matrix A :
    0.000000    0.000000    0.000000    200.000000    100.000000    0.000000
    0.000000    200.000000    100.000000    0.000000    0.000000    0.000000
    0.000000    200.000000    100.000000    100.000000    0.000000    0.000000
    0.000000    0.000000    0.000000    100.000000    200.000000    0.000000
    0.000000    0.000000    200.000000    100.000000    0.000000    0.000000
    0.000000    0.000000    0.000000    0.000000    0.000000    0.000000

Kernel :
    1.000000    0.000000   -1.000000
    1.000000    0.000000   -1.000000
    1.000000    0.000000   -1.000000

i: 0
Submatrix at j: 0
    0.000000    0.000000    0.000000
    0.000000    200.000000    100.000000
    0.000000    200.000000    100.000000

Hadamard product:
    0.000000    0.000000   -0.000000
    0.000000    0.000000  -100.000000
    0.000000    0.000000  -100.000000

Submatrix at j: 1
    0.000000    0.000000    200.000000
    200.000000    100.000000    0.000000
    200.000000    100.000000    100.000000

Hadamard product:
    0.000000    0.000000  -200.000000
    200.000000    0.000000   -0.000000
    200.000000    0.000000  -100.000000

Submatrix at j: 2
    0.000000    200.000000    100.000000
    100.000000    0.000000    0.000000
    100.000000    100.000000    0.000000

Hadamard product:
    0.000000    0.000000  -100.000000
    100.000000    0.000000   -0.000000
    100.000000    0.000000   -0.000000

```

Figure 10.25: The computation of an output map from an input image that is represented by matrix A and the kernel to detect vertical edges with `CNN_2DConvolutionOperation()` function (`SymIntegration/Examples/Neural Networks/Test CNN Compute Output Map from Convolution Operation/main.cpp`).

```

i: 3
Submatrix at j: 0
0.000000      0.000000      0.000000
0.000000      0.000000      200.000000
0.000000      0.000000      0.000000

Hadamard product:
0.000000      0.000000      -0.000000
0.000000      0.000000      -200.000000
0.000000      0.000000      -0.000000

Submatrix at j: 1
0.000000      0.000000      100.000000
0.000000      200.000000      100.000000
0.000000      0.000000      0.000000

Hadamard product:
0.000000      0.000000      -100.000000
0.000000      0.000000      -100.000000
0.000000      0.000000      -0.000000

Submatrix at j: 2
0.000000      100.000000      200.000000
200.000000      100.000000      0.000000
0.000000      0.000000      0.000000

Hadamard product:
0.000000      0.000000      -200.000000
200.000000      0.000000      -0.000000
0.000000      0.000000      -0.000000

Submatrix at j: 3
100.000000      200.000000      0.000000
100.000000      0.000000      0.000000
0.000000      0.000000      0.000000

Hadamard product:
100.000000      0.000000      -0.000000
100.000000      0.000000      -0.000000
0.000000      0.000000      -0.000000

Output map:
-200.000000      100.000000      100.000000      300.000000
-200.000000      200.000000      0.000000      200.000000
-300.000000      -100.000000      100.000000      300.000000
-200.000000      -200.000000      0.000000      200.000000

Time taken by function: 5537 microseconds

```

Figure 10.26: The computation of an output map from an input image that is represented by matrix A and the kernel to detect vertical edges with `CNN_2DConvolutionOperation()` function (`SymIntegration/Examples/Neural Networks/Test CNN Compute Output Map from Convolution Operation/main.cpp`).

To detect many different features of an image we usually use multiple kernels, such as vertical edges kernel, horizontal edges kernel, sharpening kernel. When we use multiple kernels, we multiply each kernel matrix by the corresponding input image matrix from top left to bottom right. Each kernel matrix will produce an output map, thus with n number of kernels we will obtain $o_w \times o_h \times n$ output maps.

When we use a full color RGB image, we will have a 3D matrix multiplication, and the kernel will also be a 3D matrix because real images have depth. The resulting output map will still be a 2D matrix.




Figure 10.27: After the last convolution layer, we will flatten the 3D matrix into a vector then process it again till the output layer that produces the probability for the image classification.




Figure 10.28: The spatial dimensions gradually get smaller while the depth / the number of channels grows. The early layers work with large spatial maps and few channels, while the deeper layers work with small spatial maps and many channels.

The flattened vector after the last convolution layer is a distillation of the image's most important information or feature, compressed into a format that is useful for making predictions.

[SI*] **Kernels in Convolution Operation**

Convolution operation is performed by sliding a kernel (a filter) matrix over the input matrix, performing a Hadamard (element wise, represented by $\oplus$) product of the kernel with the underlying part of input matrix, summing up the result and recording it into the output matrix. When kernels slide over images, they apply operations like blurring, sharpening, and edge detection, depends on the kernel that is used. Each kernel type has a unique function, altering the image in specific ways.

The formula for Hadamard product, if we multiply 2 matrices A and B with same size the result is matrix C with the same size

$$c_{ij} = a_{ij} * b_{ij}$$

it is not the same as the usual matrix multiplication from linear algebra, so be careful.

The convolution operation purpose is to extract certain 'features' from an input image by using a 'kernel' that will generate an output feature map.

We will identify different type of kernel that can be used in standard image processing (e.g., Sobel X, Sobel Y, Laplacian, etc), but for the convolution neural network a convolution layer

uses filters / kernels that are essentially small grids of numbers e.g., (3×3) or (5×5) that slide across the image. These numbers are the weights. The network in CNN automatically learns the best values for these weights through backpropagation to detect things like edges, shapes, and textures.

The core features of convolutional layer weights:

- **Parameter Sharing:** Unlike standard fully connected layers where every pixel has its own unique weight, a convolutional layer reuses the exact same filter (weights) across the entire image.
- **Translation Invariance:** Because the same weights are shared, the network can detect an object regardless of where it appears in the image.
- **Fewer Parameters:** This sharing drastically reduces the total number of weights the network needs to learn, preventing overfitting.

Remember the fact that images are stored as matrices of pixel values in computer. Grayscale images are simple 2D matrices while colored images are a collection of 3 such 2D matrices used for storing each color value R, G, and B.

Convolution can be done on multiple dimensions including 1D, 2D, and 3D. 1D convolutions usually are done on linear time series. A popular example of 2D convolution is image processing as well as the filters that we use. 3D convolution has its use case in video event recognition and medical imaging.

In a CNNs, a 'feature' is a specific pattern or characteristic extracted from an image using convolutional filters, essentially representing a key aspect of the image like edges, lines, textures, or shapes, which are identified by the network during the process of feature extraction that is carried out by kernels.

Types of Kernels

Kernels can be of many types. In fact, any combination of values in a matrix can act as a kernel, but a good rule of thumb is to make sure the values add up to one to avoid distorting the overall values.

Basic convolution kernels are foundational in image processing, designed to perform simple yet essential operations on images. Each type of kernel has a specific structure and function, enabling it to highlight or modify particular aspects of the input data.

1. Identity Kernel

The simplest type of convolutional kernel. It just leave the input image as it is.

$$M_{kernel} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

2. Sharpen Kernel

It enhances the details and edges of an image. It does so by increasing the difference of pixel intensities, making the image appear sharper and clearer.

$$M_{kernel} = \begin{bmatrix} 0 & -1 & 0 \\ -1 & 5 & -1 \\ 0 & -1 & 0 \end{bmatrix}$$

3. Laplacian Kernel

It is a second-order derivative operator used for edge detection. It highlights regions of rapid intensity change and is particularly useful for finding edges in noisy images.

$$M_{kernel} = \begin{bmatrix} 0 & -1 & 0 \\ -1 & 4 & -1 \\ 0 & -1 & 0 \end{bmatrix}$$

4. Box-blur Kernel

It makes the image blurrier. It does so by averaging out the values of neighboring pixels. It smoothens out the image and reduces noise as well.

$$M_{kernel} = \frac{1}{9} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

5. Gaussian-blur Kernel

It makes the image blurrier with a weighted Gaussian distribution. This kernel assigns higher weights to the central pixels and progressively lower weights to the surrounding pixels, resulting in a more natural blur. This creating a smooth, blurred effect that minimizes the high-frequency noise in the image. The matrix can have different values based on different σ values (standard deviation).

$$M_{kernel} = \frac{1}{16} \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 1 \end{bmatrix}$$

6. Sobel Kernels

They are used for detecting edges in images by emphasizing the gradients. They calculate the first derivative of the image intensity in both horizontal and vertical directions, making them effective for edge detection.

Sobel X Kernel (Horizontal)

This kernel detects horizontal edges by emphasizing vertical gradients.

$$M_{kernel} = \begin{bmatrix} -1 & 0 & 1 \\ -2 & 0 & 2 \\ -1 & 0 & 1 \end{bmatrix}$$

Sobel Y Kernel (Vertical)

This kernel detects vertical edges by emphasizing horizontal gradients.

$$M_{kernel} = \begin{bmatrix} -1 & -2 & -1 \\ 0 & 0 & 0 \\ 1 & 2 & 1 \end{bmatrix}$$

7. Prewitt Kernels

Similar to Sobel kernels, they are used for detecting edges in images by emphasizing the gradients. These kernels emphasize regions with high intensity changes, which correspond to the edges. They compute the gradients in the horizontal and vertical directions, providing a simple way to detect edges in images.

Prewitt X Kernel (Horizontal)

This kernel emphasizes horizontal changes in intensity.

$$M_{kernel} = \begin{bmatrix} -1 & 0 & 1 \\ -1 & 0 & 1 \\ -1 & 0 & 1 \end{bmatrix}$$

Prewitt Y Kernel (Vertical)

This kernel emphasizes vertical changes in intensity.

$$M_{kernel} = \begin{bmatrix} -1 & -1 & -1 \\ 0 & 0 & 0 \\ 1 & 1 & 1 \end{bmatrix}$$

8. Specialized Convolution Kernels

These kernels are designed for specific tasks and applications in image processing and computer vision. These kernels go beyond basic and advanced operations to provide more focused and sophisticated feature extraction, texture analysis, and edge detection capabilities.

Gabor Kernels

Gabor kernels are used for texture analysis and feature extraction. They are particularly effective in edge detection and texture discrimination. Gabor kernels combine a Gaussian function and a sinusoidal wave, making them capable of capturing both spatial and frequency information.

The Gabor kernel is defined by the following equation:

$$G(x, y; \lambda, \theta, \phi, \sigma, \gamma) = e^{-\frac{x'^2 + \gamma^2 y'^2}{2\sigma^2}} \cos\left(2\pi \frac{x'}{\lambda} + \phi\right)$$

Where: $x' = x \cos(\theta) + y \sin(\theta)$

$y' = -x \sin(\theta) + y \cos(\theta)$

λ is the wavelength of the sinusoidal factor.

θ is the orientation of the normal to the parallel stripes of a Gabor function.

ϕ is the phase offset.

σ is the standard deviation of the Gaussian envelope.

γ is the spatial aspect ratio.

Scharr Kernels

Scharr kernels are similar to Sobel and Prewitt kernels but offer better rotational symmetry and more accurate edge detection. They are used for detecting edges with higher precision, making them suitable for applications requiring detailed edge information.

Scharr Kernel (Horizontal)

This kernel emphasizes horizontal changes in intensity with greater accuracy.

$$M_{kernel} = \begin{bmatrix} -3 & 0 & 3 \\ -10 & 0 & 10 \\ -3 & 0 & 3 \end{bmatrix}$$

Scharr Kernel (Vertical)

This kernel emphasizes vertical changes in intensity with greater accuracy.

$$M_{kernel} = \begin{bmatrix} -3 & -10 & -3 \\ 0 & 0 & 0 \\ 3 & 10 & 3 \end{bmatrix}$$

9. Kirsch Compass Kernel

This kernel is a non-linear edge detector that finds the maximum edge strength in a few predetermined directions. The operator takes a single kernel mask and rotates it in 45 degree increments through all 8 compass directions: N, NE, E, SE, S, SW, W, and NW.

Kirsch Kernel (North)

$$M_{kernel} = \begin{bmatrix} 5 & 5 & 5 \\ -3 & 0 & -3 \\ -3 & -3 & -3 \end{bmatrix}$$

Kirsch Kernel (East)

$$M_{kernel} = \begin{bmatrix} -3 & -3 & 5 \\ -3 & 0 & 5 \\ -3 & -3 & 5 \end{bmatrix}$$

Kirsch Maximum Response (All Directions)

This kernel also accomplishes total edge detection by using the maximum of all kernel directions.

$$M_{kernel} = \max \left(\begin{bmatrix} 5 & 5 & 5 \\ -3 & 0 & -3 \\ -3 & -3 & -3 \end{bmatrix}, \begin{bmatrix} -3 & 5 & 5 \\ -3 & 0 & 5 \\ -3 & -3 & -3 \end{bmatrix}, \begin{bmatrix} -3 & -3 & 5 \\ -3 & 0 & 5 \\ -3 & -3 & 5 \end{bmatrix}, \begin{bmatrix} -3 & -3 & -3 \\ -3 & 0 & 5 \\ -3 & 5 & 5 \end{bmatrix}, \right. \\ \left. \begin{bmatrix} -3 & -3 & -3 \\ -3 & 0 & -3 \\ 5 & 5 & 5 \end{bmatrix}, \begin{bmatrix} -3 & -3 & -3 \\ 5 & 0 & -3 \\ 5 & 5 & -3 \end{bmatrix}, \begin{bmatrix} 5 & -3 & -3 \\ 5 & 0 & -3 \\ 5 & -3 & -3 \end{bmatrix}, \begin{bmatrix} 5 & 5 & -3 \\ 5 & 0 & -3 \\ -3 & -3 & -3 \end{bmatrix} \right)$$

Applications of convolution kernels:

1. Image processing
Enhancing images, detecting edges, and removing noise.
2. Computer vision
Object detection, facial recognition, and scene understanding.
3. Medical imaging
Analyzing medical images for disease detection and diagnosis.
4. Robotics
Enabling robots to perceive and understand their environment.

In machine learning, during training the values inside the kernels get trained which act as feature detection for the images, in order to obtain more information from the images. During the training of CNNs, the values (weights) inside the kernels are adjusted through the process of backpropagation and optimization (such as gradient descent). These kernels are now trained feature detectors for images and can perform tasks like categorization and identification.

[SI*] Stride

The distance the kernel window moves at each step across the image is called the stride. We need to specify the stride in both dimensions of the image: one for moving left to right and one for moving top to bottom.

Whenever the stride in either direction is greater than 1, the sliding window will skip over some of the pixels in the input at each step. But, because we want high-quality features to use for classification, it is almost always set to one. Stride values are only increased when we wish to reduce the resolution of the image which occurs during the pooling stage of convolution.

[SI*] Padding

When performing the sliding window computation, we don't know what to do with the boundaries of the input. Staying entirely inside the input image means the kernel will never sit squarely over these boundary pixels like it does for every other pixel in the input. This could lead to a problem as all pixels aren't being treated the same.

We can leave it alone and force the kernel to stay within the image. But the image loses pixels: precisely $\frac{n-1}{2}$ for every odd n sized kernel. This isn't ideal for large kernels and small images.

This is why we use padding. Here, we also have options. We can pad the input boundaries with 0's just enough to make the output the original size. This can; however, dilute the influence of the border pixels. This method is called same padding. Another method used is duplicate padding where instead of zeros, the borders are padded with values duplicating the respective border values. This helps preserve features near the edges of the image.

[SI*] Convolutional Networks

Each input image comprises a set of pixel intensity values, and the desired output is a posterior probability distribution over the ten digit classes. We know that the identity of the digit is invariant under translations and scaling as well as (small) rotations. Furthermore, the network must also exhibit invariance to more subtle transformations such as elastic deformations.

Key property of images: nearby pixels are more strongly correlated than more distant pixels. Many of the modern approaches to computer vision exploit this property by extracting local features that depend only on small subregions of the image. Information from such features can then be merged in later stages of processing in order to detect higher-order features and ultimately to yield information about the image as whole. Also, local features that are useful in one region of the image are likely to be useful in other regions of the image, for instance if the object of interest is translated.

These notions are incorporated into convolutional neural networks through three mechanisms:

- Local receptive fields
- Weight sharing
- Subsampling

In the convolutional layer the units are organized into planes, each of which is called a feature map. Units in a feature map each take inputs only from a small subregion of the image, and all of the units in a feature map are constrained to share the same weight values. For instance, a feature map might consist of 100 units arranged in a 10×10 grid, with each

unit taking inputs from a 5×5 pixel patch of the image. The whole feature map therefore has 25 adjustable weight parameters plus one adjustable bias parameter. Input values from a patch are linearly combined using the weights and the bias, and the result transformed by a sigmoidal nonlinearity. If we think of the units as feature detectors, then all of the units in a feature map detect the same pattern but at different locations in the input image. Due to the weight sharing, the evaluation of the activations of these units is equivalent to a convolution of the image pixel intensities with a 'kernel' comprising the weight parameters. If the input image is shifted, the activations of the feature map will be shifted by the same amount but will otherwise be unchanged. This provides the basis for the (approximate) invariance of the network outputs to translations and distortions of the input image. Because we will typically need to detect multiple features in order to build an effective model, there will generally be multiple feature maps in the convolutional layer, each having its own set of weight and bias parameters.

The outputs of the convolutional units form the inputs to the subsampling layer of the network. For each feature map in the convolutional layer, there is a plane of units in the subsampling layer and each unit takes inputs from a small receptive field in the corresponding feature map of the convolutional layer. These units perform subsampling. For instance, each subsampling unit might take inputs from a 2×2 unit region in the corresponding feature map and would compute the average of those inputs, multiplied by an adaptive weight with the addition of an adaptive bias parameter, and then transformed using a sigmoidal nonlinear activation function. The receptive fields are chosen to be continuous and nonoverlapping so that there are half the number of rows and columns in the subsampling layer compared with the convolutional layer. In this way, the response of a unit in the subsampling layer will be relatively insensitive to small shifts of the image in the corresponding regions of the input space.

In a practical architecture, there may be several pairs of convolutional and subsampling layers. At each stage there is a larger degree of invariance to input transformations compared to the previous layer. There may be several feature maps in a given convolutional layer for each plane of units in the previous subsampling layer, so that the gradual reduction in spatial resolution is then compensated by an increasing number of features. The final layer of the network would typically be a fully connected, fully adaptive layer, with a softmax output nonlinearity in the case of multiclass classification.

The whole network can be trained by error minimization using backpropagation to evaluate the gradient of the error function. This involves a slight modification of the usual backpropagation algorithm to ensure that the shared-weight constraints are satisfied. Due to the use of local receptive fields, the number of weights in the network is smaller than if the network were fully connected. Furthermore, the number of independent parameters to be learned from the data is much smaller still, due to the substantial numbers of constraints on the weights.

[SI*] **Operational Principles of CNNs**

Within the architecture of convolutional neural networks, image data is initially input into CNN through the input layer. Following this, the network employs a sequence of layers, including the convolutional layer, batch normalization layer, ReLU layer, and pooling layer, to systematically extract and learn features from the input images [14].

After features are extracted, the final layers (often fully connected layers) classify the image based on the detected features. Once the network produces an output, a loss function measures the difference between the predicted output and the actual labels. Common loss functions include cross-entropy for classification tasks and mean squared error (MSE) or

mean absolute error (MAE) for regression tasks.

Then, perform backpropagation and optimization. During this process, training begins with the computation of gradients, utilizing the chain rule to ascertain the gradient of the loss with respect to the network's weights. This is followed by a weight adjustment phase, where optimization algorithms, such as stochastic gradient descent (SGD) or its advanced counterparts like Adam and RMSprop, refine the weights to minimize the loss. This intricate dance of forward (from input to loss determination) and backward (involving backpropagation and optimization) passes is iteratively executed over several epochs, continuing until the network reaches an optimal or satisfactory performance threshold.

CNNs are trained using backpropagation with gradient descent optimization. The goal is to adjust the weights to minimize the difference between the predicted output and the actual labels, typically represented by a loss function. Regularization techniques, such as dropout and weight decay, are often employed to prevent overfitting.

[SI*] **Input Layer**

The input layer in a convolutional neural network (CNN) serves as the initial stage where raw data is introduced into the model for subsequent processing. Typically, the input layer accepts multidimensional data, such as images represented by width, height, and channel dimensions. This layer plays a critical role in defining the structure of the data pipeline, as it must accommodate the specific dimensions and format of the dataset under consideration. For instance, an input layer for grayscale images would require a single channel, whereas for RGB images, it would require three channels corresponding to red, green, and blue. The input layer neither performs computation nor alters the data but serves as a conduit to prepare and structure the input for downstream convolutional layers, ensuring compatibility and facilitating efficient feature extraction in subsequent stages.

[SI*] **Convolutional Layer**

This is the core building block of a CNN. The layer's primary function is to detect local patterns using learned kernels (or filters). When processing an image, these kernels slide (or convolve) around the input data (such as an image) to produce a feature map, effectively transforming the input data into a form that makes it easier for the network to understand. The result of this convolution operation is a set of feature maps that give the spatial arrangement for the detected feature at different locations in the input.

The core concepts of convolutional layer:

- **Kernels/Filters**

Kernels are small, trainable matrices that slide across the input data to extract features such as edges, textures, or more abstract patterns. Each kernel is applied to the input data via convolution, producing a feature map as output. For example, a 3×3 kernel slides over an image, performing element-wise multiplication and summing the results.

- **Stride**

Stride determines how much the kernel shifts at each step during the convolution operation. Larger strides reduce the size of the output feature map but may lose fine-grained details. Typical strides: $S = 1$ (no skip) or $S = 2$ (downsampling, faster computation but less accurate).

- **Padding**

Padding is a technique used in convolutional operations to add extra rows and columns around the edges of the input tensor, enabling the preservation or modification of spatial dimensions after the operation. Common types of padding are

1. Valid Padding (no padding): In this approach, no additional rows or columns are added, resulting in an output tensor smaller than the input tensor due to the

reduction caused by the kernel size.

2. Same Padding: Here, padding is added such that the spatial dimensions of the output tensor match those of the input tensor. This is particularly useful for maintaining consistency in the feature map size across layers.
3. Full Padding: This type extends the input tensor by padding to ensure that every element of the input is fully convolved, leading to an output tensor larger than the input tensor.

Padding choices significantly impact the receptive field, computational efficiency, and the overall architecture of convolutional neural networks.

Formula for output size with stride S and padding P :

$$\text{Output size} = \frac{N + 2P - F}{S} + 1 \quad (10.68)$$

with N as the input size and F as the filter/kernel size.

The convolutional operation, a fundamental process in deep learning, involves applying a kernel K to an input tensor X to produce an output tensor Y . Mathematically, this is expressed as

$$Y[i \cdot j] = \sum_{m=0}^{M-1} \sum_{n=0}^{N-1} X[i + m, j + n] \cdot K[m, n] \quad (10.69)$$

where the kernel slides over the input tensor, performing element-wise multiplication and summing the results.

Additionally, convolutional layers benefit from weight sharing, wherein a single set of kernel parameters is applied across all spatial locations of the input tensor. This parameter-sharing mechanism significantly reduces the number of trainable parameters, thereby improving the model's computational efficiency and mitigating the risk of overfitting.

Feature maps are the outputs generated by applying a kernel to an input tensor, effectively highlighting regions where the kernel's learned pattern is most prominent. These maps serve as representations of specific features within the input data, facilitating the detection of patterns such as edges, textures, or more complex structures.

Each value in a feature map corresponds to a receptive field, which represents the specific region of the input image that influences that value. As the network depth increases, the receptive fields expand, allowing deeper layers to capture high-level, abstract features by aggregating information from larger regions of the input. This hierarchical representation is essential for learning complex patterns and achieving strong performance in tasks such as object recognition and semantic segmentation.

Convolutions are versatile operations in deep learning, tailored to different types of data and tasks.

1D convolutions are used for sequential data such as time series, signals, or text, with applications in tasks like speech recognition.

2D convolutions, the most common type, are applied to image data, processing 2D grayscale or RGB images to extract spatial features. For processing 3D data, such as volumetric scans (e.g., CT or MRI) or video frames.

3D convolutions introduce depth as a third dimension, capturing spatial-temporal information.

Dilated (atrous) convolutions expand the receptive field by introducing gaps between kernel elements, enabling models to capture broader contextual information without increasing computational parameters, which is particularly useful in semantic segmentation.

Lastly, depthwise separable convolutions decompose standard convolution into two steps: depthwise convolution, which applies a separate kernel to each input channel, and pointwise convolution, which uses a 1×1 kernel to combine channel outputs. This approach significantly reduces computational cost while maintaining accuracy, making it a key component in efficient architectures like MobileNet.

[SI*] **Pooling Layer**

Pooling layers are an essential component of convolutional neural networks (CNNs) used to reduce the spatial dimensions of feature maps.

Pooling layers serve several critical purposes in convolutional neural networks (CNNs). First, they facilitate dimensionality reduction by reducing the size of feature maps, thereby decreasing computational costs and memory requirements in subsequent layers. This ensures the network remains efficient, even with large input images. Second, pooling enhances translation invariance by summarizing features within a region, making the model robust to small shifts or distortions in the input. Lastly, pooling contributes to preventing overfitting by reducing spatial dimensions, acting as a form of implicit regularization that improves generalization.

Let $x^L \in \mathbb{R}^{H^L \times W^L \times D^L}$ be the input to the L -th layer, which is now a pooling layer. The pooling operation requires no parameter (i.e. w^i is null; hence, parameter learning is not needed for this layer). The spatial extent of the pooling ($H \times W$) is specified in the design of the CNN structure. Assume that H divides H^L and W divides W^L and the stride equals the pooling spatial extent, the output of pooling will be an order three tensor of size $H^{L+1} \times W^{L+1} \times D^{L+1}$, with

$$H^{L+1} = \frac{H^L}{H}, \quad W^{L+1} = \frac{W^L}{W}, \quad D^{L+1} = D^L \quad (10.70)$$

This formula ensures precise control over the spatial dimensions of the feature maps after pooling, which is essential for designing and understanding CNN architectures.

Pooling layers often use stride and padding to control the operation's behavior and output dimensions. The stride specifies the step size of the pooling window as it moves across the feature map. A stride of 2, for example, reduces the spatial dimensions by approximately half, enabling efficient dimensionality reduction while summarizing features.

Padding adds extra rows and columns around the edges of the feature map, similar to its role in convolution. This ensures that the pooling operation includes the boundary regions of the feature map, preserving spatial alignment and avoiding loss of information at the edges, especially in cases where maintaining the feature map's structure is crucial.

Pooling operations in convolutional neural networks (CNNs) come in various forms, each designed to summarize feature maps in different ways while reducing dimensions,

1. **Max Pooling:** This method retains the maximum value from a pooling region, effectively highlighting the most prominent feature. It is defined as

$$Y[i, j] = \max(X[i : i + F, j : j + F]) \quad (10.71)$$

where F is the window size.

Max pooling takes the maximum value from each $F \times F$ window of the input and assigns it to the output position (i, j) .

2. **Average Pooling:** In contrast, this method computes the average of values within the pooling region, capturing the general trend of features. It is defined as

$$Y[i, j] = \frac{1}{F^2} \sum_{k=0}^{F-1} \sum_{l=0}^{F-1} X[i+k, j+l] \quad (10.72)$$

This formula defines average pooling in a neural network. Here, X is the input feature map, and $Y[i, j]$ is the output at position (i, j) . A square window of size $F \times F$ slides over the input, and the values within the window are averaged. The result is assigned to the corresponding position in the output Y .

3. **Global Pooling:** It reduces each feature map to a single value, either by taking the maximum (global max pooling) or the average (global average pooling) over all spatial dimensions. This approach is commonly used in architectures like ResNet to replace fully connected layers, simplifying the model while maintaining performance.
4. **Adaptive Pooling:** It dynamically adjusts the output size to a fixed dimension, regardless of input size, making it particularly useful for handling variable-sized inputs.

An example of max pooling with filters: 2×2 and stride : 2

$$\begin{bmatrix} 1 & 1 & 2 & 5 \\ 5 & 6 & 7 & 8 \\ 3 & 2 & 1 & 0 \\ 1 & 2 & 3 & 4 \end{bmatrix} \xrightarrow{\text{Max Pooling}} \begin{bmatrix} 6 & 8 \\ 3 & 4 \end{bmatrix}$$

Advanced pooling methods extend the capabilities of standard pooling to address specific challenges in computer vision tasks:

- **Region of Interest (ROI) Pooling:** This technique is commonly used in object detection models like Faster-R-CNN to dynamically pool features from regions corresponding to detected objects, enabling precise localization and classification.
- **Feature Pyramid Pooling:** Employed in networks such as PSPNet, this method extracts multi-scale features by pooling at different scales, enhancing the model's ability to recognize objects of varying sizes and improving performance in tasks like semantic segmentation.
- **Fractional Pooling:** Designed to handle non-integer downsampling ratios, fractional pooling interpolates between neighboring values, allowing finer control over output dimensions and adapting to diverse input sizes more flexibly.

These techniques enhance pooling's adaptability, improving the performance of CNNs in complex tasks like object detection, segmentation, and multi-scale feature extraction.

[SI*] **Normalization Layer**

Normalization layers are essential for stabilizing and accelerating the training of deep neural networks. Normalization in neural networks involves adjusting the distribution of data, such as activations or inputs, to improve training and performance. It ensures stabilized learning by maintaining consistent distributions across layers, allowing gradient to flow predictably and reducing the risk of issues like exploding or vanishing gradients. Additionally, normalization facilitates faster convergence by minimizing internal covariate shift, which leads to

smoother and more efficient optimization. Finally, it enhances generalization, enabling the model to perform better on unseen data by encouraging robust and consistent learning of features.

1. **Batch Normalization (BN):** Batch normalization normalizes activations within a mini-batch by adjusting their distribution to have a mean of zero and a variance of one. Specifically, it computes the batch mean

$$\mu = \frac{1}{m} \sum_{i=1}^m x_i \quad (10.73)$$

and variance

$$\sigma^2 = \frac{1}{m} \sum_{i=1}^m (x_i - \mu)^2 \quad (10.74)$$

where m is the mini-batch size. Each activation is then normalized as

$$\hat{x}_i = \frac{x_i - \mu}{\sqrt{\sigma^2 + \epsilon}} \quad (10.75)$$

with a small constant ϵ added for numerical stability. Finally, learnable parameters γ (scale) and β (shift) are applied to allow the network to recover useful representations:

$$y_i = \gamma \hat{x}_i + \beta \quad (10.76)$$

Batch normalization reduces internal covariate shift, accelerates convergence, and provides implicit regularization. However, it may perform poorly with very small batch sizes and introduces computational overhead during training.

2. **Layer Normalization (LN):** Layer normalization normalizes activations across all features of a single input instance, making it independent of batch size. For an input with H features, the mean and variance are computed as

$$\mu = \frac{1}{H} \sum_{j=1}^H x_j \quad (10.77)$$

and

$$\sigma^2 = \frac{1}{H} \sum_{j=1}^H (x_j - \mu)^2 \quad (10.78)$$

Each feature is normalized using

$$\hat{x}_j = \frac{x_j - \mu}{\sqrt{\sigma^2 + \epsilon}} \quad (10.79)$$

followed by scaling and shifting with learnable parameters $y_j = \gamma \hat{x}_j + \beta$. This method works well for tasks like NLP and reinforcement learning, where batch dependencies are less relevant. Layer normalization is particularly useful in transformer architectures and recurrent neural networks, offering stable learning regardless of batch size.

3. **Instance Normalization (IN):** Instance normalization focuses on normalizing the spatial dimensions of each channel independently, making it effective for style transfer and

image generation tasks. For a feature map of height H and width W , the mean and variance for each channel c are calculated as

$$\mu_c = \frac{1}{HW} \sum_{i=1}^H \sum_{j=1}^W x_{i,j,c} \quad (10.80)$$

and

$$\sigma_c^2 = \frac{1}{HW} \sum_{i=1}^H \sum_{j=1}^W (x_{i,j,c} - \mu_c)^2 \quad (10.81)$$

The spatial dimensions are then normalized using

$$\hat{x}_{i,j,c} = \frac{x_{i,j,c} - \mu_c}{\sqrt{\sigma_c^2 + \epsilon}} \quad (10.82)$$

followed by scaling and shifting:

$$y_{i,j,c} = \gamma \hat{x}_{i,j,c} + \beta \quad (10.83)$$

Instance normalization normalizes each feature map independently, removing instance-specific variations while preserving structural features. This makes it especially effective for tasks involving stylized outputs.

In addressing the inherent limitations of conventional normalization strategies, advanced normalization techniques and their alternatives have emerged, presenting novel approaches to significantly enhance neural network performance. Batch renormalization stands as a notable extension of batch normalization (BN), specifically engineered to mitigate the instability encountered with smaller batch sizes. This method introduces adaptive scaling and shifting parameters, which serve to stabilize training by ensuring consistent statistical properties across a wider range of batch sizes, thereby improving the reliability and generalization of models trained under such conditions.

Furthermore, switchable normalization represents a dynamic approach wherein the network learns to optimally select from a repertoire of normalization techniques, including BN, layer normalization (LN), and instance normalization (IN), on a per-layer basis during training. This adaptive mechanism allows for a tailored normalization strategy, optimizing performance across diverse tasks and architectural configurations. Concurrently, a burgeoning area of research explores the feasibility of normalization-free networks, which entirely circumvent the need for explicit normalization layers. A prominent example is FixUp initialization, which employs specialized parameter initialization schemes and concomitant training adjustments. This methodology enables the effective training of deep neural networks without recourse to normalization, consequently reducing computational overhead and simplifying the overall model architecture.

[SI*] **Activation Layer**

Activation layers are a critical component of neural networks, introducing nonlinearity to the model, enabling it to learn complex patterns, and transforming raw inputs into useful features.

Activation functions also perform transformation, converting raw weighted sums into meaningful outputs for subsequent layers to process. Additionally, they influence gradient flow

during backpropagation, playing a vital role in convergence and stability by ensuring gradients are neither too large nor too small. These functions make activation layers essential for effective learning and performance.

We have describes the types of activation functions in the feed-forward neural network section, at the beginning of this chapter, thus we won't repeat again, readers can read that section if they are not well informed about activation functions.

Choosing an activation function depends on the specific requirements of the task and architecture. ReLU is the most commonly used activation function due to its simplicity, efficiency, and ability to mitigate the vanishing gradient problem, making it a default choice for many networks like CNNs and DNNs. Variants of ReLU, such as Leaky ReLU and Parametric ReLU, address issues like the dying ReLU" problem. For outputs requiring bounded ranges, functions like Tanh or sigmoid are more appropriate. In modern architectures, Swish has shown improved performance in certain scenarios, especially when its higher computational cost is acceptable. Ultimately, selecting the best activation function often relies on experimentation, as empirical results can vary based on the dataset and application.

[SI*] **Fully Connected Layer**

Fully connected (FC) layers are fundamental components of neural networks, especially in traditional architectures. They connect every input neuron to every output neuron, enabling the network to learn global patterns and make predictions.

A fully connected layer is a dense neural layer where every neuron is connected to all neurons in the previous layer, playing a critical role in aggregating and utilizing learned features. FC layers combine spatially distributed features extracted by convolutional layers into a compact representation, enabling high-level decision-making tasks such as classification or regression. They act as a transition point by flattening feature maps into a 1D vector before further processing. FC layers provide a global representation by integrating information from all preceding neurons, making them essential for tasks requiring comprehensive feature aggregation. However, they are computationally expensive due to the large number of parameters, proportional to the product of the input size and the number of neurons, which increases the risk of overfitting. Additionally, FC layers are vital for dimensionality transformation, reshaping data to suit downstream tasks effectively.

The output of a fully connected layer is computed as

$$y = f(Wx + b) \quad (10.84)$$

where W is the weight matrix of size $[\text{output_neurons} \times \text{input_neurons}]$, x is the input vector, b is the bias vector, and f is the activation function.

For an input vector x of size N and a fully connected layer with M neurons, the layer's functionality is defined by its weights and biases, output computation, and parameter count. The weight matrix W has dimensions $M \times N$ representing the connections between each input and neuron, while the bias vector b has size M , adding a trainable offset to each neuron. The output is computed in two steps: first, a linear transformation is applied, $z = Wx + b$, which aggregates the input features weighted by the learned parameters. Then, the activation function f is applied to the result, yielding the final output $y = f(z)$.

The total number of parameters in the FC layer, which includes $N \times M$ weights and M biases, is calculated as $\text{parameters} = (N \times M) + M$. For example, with an input vector of size 128 and an FC layer comprising 64 neurons, the parameter count would be $(128 \times 64) + 64 = 8256$. This high parameter count highlights the computation expense and potential overfitting risk associated with fully connected layers, making them critical yet resource-intensive component of neural networks.

Fully connected layers play a crucial role in aggregating features and making predictions. Despite their simplicity, they are computationally intensive and prone to overfitting. Modern architectures often replace or minimize their use with techniques like global pooling, attention mechanisms, or sparse connections.

[SI*] **Classification Layer**

In the end, a CNN often has a classification layer to make a prediction. If, for instance, the task is to classify an image among ten categories, this layer would output ten values, each one representing the probability of the image belonging to that class. In CNNs, fully connected layers serve as a crucial bridge between the feature extraction process and high-level decision-making tasks such as classification or regression. The transition to FC layers involves flattening the multidimensional feature maps produced by convolutional layers into a one-dimensional vector. For example, a $7 \times 7 \times 64$ feature map is reshaped into a 3136-dimensional vector, enabling the network to process spatially features, preparing them for classification or other downstream tasks.

FC layers are frequently employed as the final layers in CNNs, particularly in image classification tasks, where they act as output layers. In these scenarios, the final FC layer typically consists of C neurons, where C represents the number of target classes. A softmax activation function is then applied to this layer to produce class probabilities, ensuring the outputs are interpretable and suitable for classification. By integrating spatially learned features from convolutional layers into a global representation, FC layers enable CNNs to make accurate predictions, highlighting their essential role in combining feature extraction and task-specific decision-making.

[SI*] **Typical Model Structure of CNNs**

The fundamental principle underpinning convolutional neural networks lies in their utilization of shared local receptive fields and weights. This core concept enables efficient feature extraction from spatially correlated data. The seminal work in this field, LeNet, introduced by LeCun et al. in 1998, is widely recognized as the archetype of CNN architectures.

Over the past decade, convolutional neural networks have undergone significant evolution, driven by the need for improved performance, scalability, and efficiency in computer vision tasks. Each advancement in CNN architecture has introduced novel design principles, addressing critical challenges such as vanishing gradients, computational cost, and overfitting. Starting with AlexNet which demonstrated the power of deep learning on large-scale image datasets, modern architectures like ResNet, DenseNet, and EfficientNet have pushed the boundaries of depth, parameter efficiency, and computational optimization. This section explores the innovations and design philosophies of key CNN architectures, highlighting their contributions to the field and their practical applications in solving complex visual tasks.

1. **LeNet (1998)**

The original LeNet model featured a straightforward architecture comprising five

convolutional layers directly linked to corresponding pooling layers. Its initial application focused on the recognition of handwritten and printed characters, demonstrating remarkable performance for its time. It is worth noting that contemporary implementations of LeNet often incorporate modifications from the original design to suit a broader range of applications.

2. AlexNet (2012)

AlexNet marked a milestone in deep learning by winning the 2012 ImageNet competition, demonstrating the potential of deep CNNs for image classification. It introduced ReLU activation to accelerate training, dropout to reduce overfitting, and leveraged GPUs for efficient computation. The architecture consists of five convolutional layers and three fully connected layers, using large filters such as 11×11 and 5×5 . Despite its breakthrough, AlexNet's large parameter count (60 M) and computational requirements were significant drawbacks.

3. VGG (2014)

The VGG network demonstrated that deeper networks with smaller filters (3×3) could significantly improve performance. VGG-16 and VGG-19 are the most common configurations, comprising 13-16 convolutional layers followed by three fully connected layers. Max pooling is used for spatial downsampling. While VGG is simple and effective, its parameter count is exceedingly high (e.g., 138M for VGG-16), making it resource-intensive in terms of computation and memory.

4. Inception (GoogleLeNet, 2014)

Inception networks introduced inception modules, which apply convolutions of varying kernel sizes in parallel to capture multi-scale features. The architecture reduces computation through 1×1 convolutions for dimensionality reduction. Auxiliary classifiers improve gradient flow in deeper networks. Despite its efficiency and strong performance, the complexity of its design poses implementation challenges.

5. ResNet (2015)

ResNet introduced residual connections (skip connections) to address the vanishing gradient problem, enabling the training of extremely deep networks, such as ResNet-50 and ResNet-152. Residual connections allow gradients to flow directly through layers by learning residual mappings ($F(x) + x$), significantly improving training dynamics and performance. However, the increased depth of ResNet comes at the cost of high computational complexity.

6. MobileNet (2017)

MobileNet was designed for mobile and embedded systems, introducing depth-wise separable convolutions to reduce computation. Depthwise convolution applies a single filter per input channel, followed by pointwise convolution to combine channels. This lightweight design significantly reduces model size and complexity, albeit with a trade-off in accuracy for highly constrained model. Variants like MobileNetV2 and V3 further enhance performance and efficiency.

7. ShuffleNet (2017)

ShuffleNet targeted efficient computation on mobile devices by introducing grouped convolutions and channel shuffle mechanisms. Grouped convolutions divide channels into subsets for parallel computation, while channel shuffle ensures information exchange across groups. This design achieves a balance between accuracy and computational efficiency, though its implementation is more complex compared to simpler architectures.

8. GhostNet (2020)

GhostNet optimized lightweight CNNs by introducing ghost modules, which generate additional feature maps through inexpensive linear transformations instead of full convolutions. This reduces computational cost and parameter count while maintaining competitive accuracy. GhostNet is highly effective for resource-constrained devices, though extreme compression may lead to slight performance degradation.

[SI*] Applications of CNNs

Convolutional neural networks have revolutionized numerous fields by offering unparalleled performance in processing visual and spatial data. Their ability to learn hierarchical features and extract intricate patterns has enabled groundbreaking applications across industries.

Image classification is a fundamental task in computer vision where convolutional neural networks assign labels to images based on learned features. By extracting hierarchical patterns from raw image data, CNNs enable accurate categorization and identification across a wide range of applications. In medical imaging, CNNs classify X-rays, MRIs, and CT scans to assist in diagnosing diseases such as pneumonia or detecting tumors. For autonomous vehicles, they identify road signs, pedestrians, and objects to support safe navigation. In retail and commerce, CNNs categorize products based on visual attributes, enhancing search and recommendation systems.

Object detection is a key task in computer vision that combines classification and localization to identify objects in an image and draw bounding boxes around them. This task extends beyond merely recognizing what is in an image by also determining where the objects are located. It has diverse applications across various domains. In surveillance, object detection helps identify suspicious activities or unauthorized access in security footage. For autonomous vehicles, it is essential for real-time identification of vehicles, pedestrians, and obstacles to ensure safe navigation. In agriculture, object detection aids in identifying pests and diseases or estimating crop yield, improving efficiency and productivity. A significant milestone in object detection is the development of YOLO (You Only Look Once), a single-stage detection pipeline that revolutionized real-time object detection.

By combining detection and classification into a single step, YOLO dramatically reduced inference time while maintaining high accuracy, making it suitable for applications requiring low-latency detection. Its efficiency and reliability have made YOLO a widely adopted framework in real-world scenarios demanding real-time performance.

Face recognition systems are designed to identify or verify individuals by analyzing facial features from images or video streams. These systems have become integral in various domains. In security, face recognition is used for access control and surveillance, ensuring only authorized individuals can enter restricted areas. In consumer electronics, facial unlocking features enhance convenience and security in smartphones and laptops. In retail, face recognition enables personalized customer experiences by analyzing facial data to understand preferences and behaviors.

A notable advancement in this field is FaceNet, which employs deep embeddings to map faces into a high-dimensional feature space. This approach allows for highly accurate face verification and clustering of identities, enabling robust authentication systems and efficient video analytics. FaceNet's deployment in applications like secure authentication and surveillance highlights its effectiveness in real-world face recognition tasks.

Autonomous vehicles rely extensively on convolutional neural network (CNNs) for real-time scene understanding, integrating multiple vision tasks to ensure safe and efficient operation. Core tasks include object detection using models like YOLO for identifying vehicles, pedestrians, and obstacles in real-time, semantic segmentation with architectures like DeepLab to classify road scenes into meaningful regions and monocular depth estimation using CNNs to predict depth information from single-camera inputs. These capabilities enable autonomous systems to interpret complex environments dynamically. A prominent example is Tesla's Autopilot, which uses deep CNNs for multi-camera vision processing. This system powers advanced driver assistance features such as lane keeping, obstacle detection, and adaptive cruise control, forming the foundation for achieving full vehicle autonomy. By combining these techniques, autonomous vehicles demonstrate the potential of CNNs in transforming transportation. pg 42

i. Convolution Operation on an Image with OpenGL and STB Image

We will perform a convolution operation by sliding a kernel matrix over the input matrix (the input image). It is very handy to use OpenGL for this kind of task, since it is the fastest around, even if we need a steeper learning curve and need to code more it is worth the time and sweat.

A lot of people out there are coding with OpenCV for image manipulation, loading, or converting an image into greyscale, OpenCV requires at least 280 MB storage, and if we can do such image manipulation with smaller but faster library and then develop it ourselves from that, it is better that way.

So we are using STB image for the image loading part and showing the image on a new GUI window is thanks to OpenGL + vertex shaders + fragment shaders and the interactivity for performing the convolution operation with keyboard pressed events is thanks to GLFW.

The whole size for the source code is 17 KB, no OpenCV or any heavy dependencies, but we need OpenGL and GLFW which will be very useful in the future if we want to create physics simulation like Bullet or Project Chrono, or deep learning project like video recognition.

This code depends on shaders a lot to make the convolution operation works, shaders are little programs that rest on the GPU. These programs are run for each specific section of the graphics pipeline. In a basic sense, shaders are nothing more than programs transforming inputs to outputs. Shaders are also very isolated programs in that they're not allowed to communicate with each other; the only communication they have is via their inputs and outputs.

Shaders are written in the C-like language GLSL, GLSL is tailored for use with graphics and contains useful features specifically targeted at vector and matrix manipulation.

Shaders always begin with a version declaration, e.g. `#version 330` (meaning it uses OpenGL version 3.3), followed by a list of input and output variables, uniforms and its main function. Each shader's entry point is at its main function where we process any input variables and output the results in its output variables. If you want to comprehend about shaders more, you can read the comprehensive book by Joey de Vries [29].

Now we are going to talk about loading the image with an image-loading library, in this example we are using STB Image, it supports several popular formats like `.png` and `.jpg`. `stb_image.h` is a very popular single header image loading library by Sean Barrett that is able to load most popular file formats and is easy to integrate in your project. You just need to define it (`#define STB_IMAGE_IMPLEMENTATION`) then include the header (`#include "stb_image.h"`). By defining `STB_IMAGE_IMPLEMENTATION` the preprocessor modifies the header file such that it only contains the relevant definition source code, effectively turning the header file into a `.cpp` file.

In the computer world, OpenGL is considered as an API (Application Programming Interface) that provides us with a large set of functions that we can use to manipulate graphics and images. However, OpenGL by itself is not an API, but merely a specification, it specifies exactly what the result/output of each function should be and how it should perform. To perform kernel convolution on an image we cannot use OpenGL only, we need another libraries such as GLFW (it handles keyboard or mouse input events and gives us the bare necessities required for rendering

images/graphics to the screen), GLAD (this library is the manager library to handle the location of most OpenGL functions that we need, because most of its functions' location is not known at compile-time, thus this library retrieve the location of the functions that we need and store them in function pointers for later use).

Some definitions that are related to OpenGL:

- **Shader:** a small program that runs on the graphics card. Several stages of the graphics pipeline can use user-made shaders to replace existing functionality.
- **Vertex:** a collection of data that represent a single point.
- **Vertex Buffer Object (VBO):** a buffer object that allocates memory on the GPU and stores all the vertex data there for the graphics card to se.
- **Vertex Array Object (VAO):** stores buffer and vertex attribute state information.
- **Element Buffer Object (EBO):** a buffer object that stores indices on the GPU for indexed drawing.
- **Uniform:** a special type of GLSL variable that is global (each shader in a shader program can access this uniform variable) and only has to be set once.
- **Texture:** a special type of image used in shaders and usually wrapped around objects, giving the illusion an object is extremely detailed.
- **Texture Wrapping:** defines the mode that specifies how OpenGL should sample textures when texture coordinates are outside the range: (0,1).
- **Texture Filtering:** defines the mode that specifies how OpenGL should sample the texture when there are several texels (texture pixels) to choose from. This usually occurs when a texture is magnified.

A kernel (convolution matrix) is a small matrix-like array of values centered on the current pixel that multiplies surrounding pixel values by its kernel values and adds them all together to form a single value.

Kernels are an extremely useful tool for post-processing since they're quite easy to use and experiment with.

We will perform some convolution operations on this picture:



Figure 10.29: *This is a female Gerberian Shepsky dog who is about 1 month old, the one and only.*

We will perform the laplacian operation with this kernel / convolution matrix:

$$M_{kernel} = \begin{bmatrix} -1 & -1 & -1 \\ -1 & 8 & -1 \\ -1 & -1 & -1 \end{bmatrix}$$




Figure 10.30: The laplacian kernel operation on an image (*SymIntegration/Examples/Neural Networks/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/main.cpp*).

Next, we will perform the Sobel X operation with this kernel / convolution matrix:

$$M_{kernel} = \begin{bmatrix} -1 & 0 & 1 \\ -2 & 0 & 2 \\ -1 & 0 & 1 \end{bmatrix}$$




Figure 10.31: The Sobel X kernel operation on an image (*SymIntegration/Examples/Neural Networks/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/main.cpp*).

Next, we will perform the Sobel Y operation with this kernel / convolution matrix:

$$M_{kernel} = \begin{bmatrix} -1 & -2 & -1 \\ 0 & 0 & 0 \\ 1 & 2 & 1 \end{bmatrix}$$



Figure 10.32: The Sobel Y kernel operation on an image (*SymIntegration/Examples/Neural Networks/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/main.cpp*).

Next, we will perform the Schar horizontal operation with this kernel / convolution matrix:

$$M_{kernel} = \begin{bmatrix} -3 & 0 & 3 \\ -10 & 0 & 10 \\ -3 & 0 & 3 \end{bmatrix}$$




Figure 10.33: *The Schar horizontal operation on an image (SymIntegration/Examples/Neural Networks/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/main.cpp).*

Next, we will perform the Schar vertical operation with this kernel / convolution matrix:

$$M_{kernel} = \begin{bmatrix} -3 & -10 & -3 \\ 0 & 0 & 0 \\ 3 & 10 & 3 \end{bmatrix}$$



Figure 10.34: *The Schar vertical operation on an image (SymIntegration/Examples/Neural Networks/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/main.cpp).*

Next, we will perform the Kirsch west operation with this kernel / convolution matrix:

$$M_{kernel} = \begin{bmatrix} 5 & -3 & -3 \\ 5 & 0 & -3 \\ 5 & -3 & -3 \end{bmatrix}$$




Figure 10.35: *The Kisch west operation on an image (SymIntegration/Examples/Neural Networks/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/main.cpp).*

The C++ source code is modified from chapter 7 of the Learn OpenGL book [29], if we do not want to change the logic in the C++ code then we only need to play around in the **fragment.fs** (the fragment shader), this file handles the color output for the rendered image depends on the kernel / convolution matrix that we want to apply on the image or even just some nice RGB color interpolation or weighted grayscale. We don't need to recompile just call again **./main** and the image will be transformed / convoluted.

```
...

#version 330 core
out vec4 FragColor;

in vec3 ourColor;
in vec2 TexCoord;

// texture samplers
uniform sampler2D texture1;
uniform sampler2D texture2;
const float offset = 1.0 / 300.0;

void main()
{
    // linearly interpolate between both textures (10% texture1, 90% texture2)
    FragColor = mix(texture(texture1, TexCoord), texture(texture2, TexCoord),
        0.9);

    /* // Grayscale
    FragColor = texture(texture2, TexCoord);
    float average = (FragColor.r + FragColor.g + FragColor.b) / 3.0 ;
    FragColor = vec4(average, average, average, 1.0);
    */

    // /*
    // Weighted grayscale
    FragColor = texture(texture2, TexCoord);
    float average = 0.2126*FragColor.r + 0.7152*FragColor.g + 0.0722*FragColor.
        b ;
```

```

FragColor = vec4(average, average, average, 1.0);
// */
...

```

Code 66: *Neural Network/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/fragment.fs*

We have uncommented the weighted grayscale, so the image will be grayscaled like below when we do so, don't forget to keep all other convolution operations in comment.



Figure 10.36: *The weighted grayscale of an image (SymIntegration/Examples/Neural Networks/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/main.cpp).*

In the current updated code, we create a new logic so it is more efficient and easier to see the convoluted image while the GLFW GUI window still active.

In this new logic, we use **void processInput(GLFWwindow *window, Shader* activeShader)** instead of **void processInput(GLFWwindow *window)**, so the input variables become 2, so we can interchange the shader in use easily while the GLFW window is still open. This **void** function evaluates key presses in the main render loop to compile/bind the desired shader program on demand. Before starting the render loop, we initialize the base shader with this:

```
Shader *activeShader = new Shader("vertex.vs", "fragment.fs");
```

We are not using memory assignment (&m); instead we dynamically allocate it with **new** so it can be cleanly deleted and re-instantiated.

Then in the render loop we use the dynamically managed shader with **activeShader->use();**.




Figure 10.37: By pressing keyboard keys from 1 to 8, we can alternate and see the convoluted image (*SymIntegration/Examples/Neural Networks/Test Kernel Convolution Operation on an Image with OpenGL GLFW GLAD and STB Image/main.cpp*).

This is the prelude / introduction before we get into the real CNNs, even if in convolutional neural networks feature-extracting kernels are not predefined, so we will not be using any of the Sobel X, or Sobel Y or Laplacian kernel. In CNNs we will have to perform a lot of matrix computation with tons of randomly initialized kernels / weights (when training from scratch) to detect the pattern in an image, and we will still need to do the backpropagation to adjusting the weights and biases. The network learns the optimal weight matrices / kernel matrices during the training process to detect complex features like eyes in a face or wheels on a car.

In the image example above we can even create an image classification CNN to detect whether it is a certain race of a dog, determine the age of the dog, or even the gender of the dog.

Implementing a convolutional neural network (CNN) convolution operation in OpenGL involves utilizing the GPU's parallel processing capabilities. We typically execute convolutions by loading input feature maps and kernels as textures, and using an OpenGL Shading Language (GLSL) **fragment shader** attached to a Frame Buffer Object (FBO) to perform a sliding-window dot product.

OpenGL is designed for image rendering, it natively handles 2D textures, vector arithmetic, and linear interpolation-which map directly onto CNN components:

- **Input Feature Map:** Bound as a GLSL **sampler2D** containing the input values.
- **Filter/Kernel:** Passed as shader uniforms, often structured as **vec4** arrays or a 1D/2D texture depending on size.
- **Dot Product:** Calculated per pixel inside the fragment shader, utilizing functions like **dot()** or nested **for** loops.

The fragment shader is responsible for reading the neighboring pixels of the current coordinate, multiplying them by the kernel weights, and outputting the convolved value.

ii. Extract RGBA Matrices from an Image and then Create an Image from the RGBA Matrices with SymIntegration, OpenGL and STB Image

Remember that a computer sees an image as a matrix of numbers $[0, 255]$? it can be represented as a 3D matrix with 3 channels for RGB image, or with 4 channels for RGBA image, often the matrix A will have the same entries of 255.

We will use an example of Sine Bam Bam picture again, she is the best, one and only Gerberian Shepsky dog who is also a Mini Pincher mix, a very rare breed, 1 in a billion.

First we will add a few lines of code before the render loop **while (!glfwWindowShouldClose(window))**{}

```
...

vector<vector<int>> matrixR(height, vector<int>(width, 0.0));
vector<vector<int>> matrixG(height, vector<int>(width, 0.0));
vector<vector<int>> matrixB(height, vector<int>(width, 0.0));
vector<vector<int>> matrixA(height, vector<int>(width, 0.0));
// Save the RGBA matrix into 4 matrices
for (int y = 0; y < height; y++)
{
    for (int x = 0; x < width; x++)
    {
        // 1. Calculate the starting memory address of the pixel
        unsigned char* pixel_ptr = data + (y * width + x) * nrChannels;

        // 2. Extract components based on available channels (e.g., RGB or
        //    RGBA)
        // Cast to int to print numerical values rather than ASCII
        // characters
        int r = pixel_ptr[0];
        int g = (nrChannels >= 2) ? pixel_ptr[1] : 0;
        int b = (nrChannels >= 3) ? pixel_ptr[2] : 0;
        int a = (nrChannels >= 4) ? pixel_ptr[3] : 255;
        matrixR[y][x] = r;
        matrixG[y][x] = g;
        matrixB[y][x] = b;
        matrixA[y][x] = a;
    }
}

saveMatrixint(matrixR, "matrixR.txt");
saveMatrixint(matrixG, "matrixG.txt");
saveMatrixint(matrixB, "matrixB.txt");
saveMatrixint(matrixA, "matrixA.txt");
stbi_image_free(data);

...
```


Code 67: *Neural Network/Test Create RGBA Matrices with SymIntegration OpenGL GLFW GLAD and STB Image/main.cpp*

those lines of codes will extract the RGBA matrices into the current working directory.




Figure 10.38: When the program starts it will save separately each of the RGBA matrices from the second texture that we load (*/sources/Hamzstlab/LearnOpenGL/textures/6.png*) as *.txt* file in the current working directory (*SymIntegration/Examples/Neural Networks/Test Create RGBA Matrices with SymIntegration OpenGL GLFW GLAD and STB Image/main.cpp*).

After this we can copy the *matrixR.txt*, *matrixG.txt*, *matrixB.txt*, *matrixA.txt* into the *SymIntegration/Examples/Neural Networks/Test Create Image from RGBA Matrices with STB Image/* folder, if there are previous RGBA matrices then they will be overwritten.




Figure 10.39: Before we compile the program this is the current working directory (*SymIntegration/Examples/Neural Networks/Test Create Image from RGBA Matrices with STB Image/*).

open xterm/terminal in the current working directory and compile then run the program:
make
./main




Figure 10.40: After running the program we obtain the image, same as the image from */sources/Hamzst-lab/LearnOpenGL/textures/6.png* with 522 KB of size, this one has higher size of 721 KB, better resolution, sharper perhaps. (*SymIntegration/Examples/Neural Networks/Test Create Image from RGBA Matrices with STB Image/main.cpp*).

```

#include <iostream>
#include "symintegrationc++.h"
#include <bits/stdc++.h>
#include <cmath>
#include <chrono>

using namespace std::chrono;

```



```
using namespace std;
using namespace SymbolicConstant;

#define STB_IMAGE_WRITE_IMPLEMENTATION
#include "stb_image_write.h"

int main() {
    int width = 434;
    int height = 679;
    int channels = 4; // RGBA

    // Get starting timepoint
    auto start = high_resolution_clock::now();
    // Allocate memory for the pixel array (width * height * 4 bytes)
    uint8_t* pixels = (uint8_t*)malloc(width * height * channels);
    //int pixels[width * height * channels];
    if (pixels == NULL)
    {
        printf("Memory allocation failed!\n");
        return 1;
    }

    vector<vector<int>> matrixR = loadIntMatrixFromFile("matrixR.txt");
    vector<vector<int>> matrixG = loadIntMatrixFromFile("matrixG.txt");
    vector<vector<int>> matrixB = loadIntMatrixFromFile("matrixB.txt");
    vector<vector<int>> matrixA = loadIntMatrixFromFile("matrixA.txt");

    // Generate the image from RGBA matrices
    for (int y = 0; y < height; y++)
    {
        for (int x = 0; x < width; x++)
        {
            int index = (y * width + x) * channels;
            pixels[index + 0] = (uint8_t)matrixR[y][x]; // Red channel
            pixels[index + 1] = (uint8_t)matrixG[y][x]; // Green channel
            pixels[index + 2] = (uint8_t)matrixB[y][x]; // Blue channel
            pixels[index + 3] = (uint8_t)matrixA[y][x]; // A channel
        }
    }

    // Calculate stride: distance in bytes between the start of consecutive
    rows
    int stride = width * channels;
    stbi_flip_vertically_on_write(true); // tell stb_image.h to flip the saved
    image's on the y-axis.
    // Save the pixel data to a PNG file
    if (stbi_write_png("SineBamBam.png", width, height, channels, pixels,
```

```

        stride))
    {
        printf("Image saved successfully!\n");
    }
    else
    {
        printf("Failed to save image.\n");
    }
    // stbi_write_jpg("output.jpg", width, height, channels, pixels, 100); //
    // 100 is quality (1-100)
    // Get ending timepoint
    auto stop = high_resolution_clock::now();
    auto duration = duration_cast<microseconds>(stop - start);

    cout << "\nTime taken by function: " << duration.count() << " microseconds"
        << endl;
    // Always clean up allocated memory
    free(pixels);
    return 0;
}

```

Code 68: Neural Network/Test Create Image from RGBA Matrices with STB Image/main.cpp

We need to define the correct width and height of the original image at `/sources/Hamzst-lab/LearnOpenGL/textures/6.png`, we are also using `malloc()`, it is inherited from C language and always paired with `free()`, notice that this is the alternative of `new` operator that we have used in the previous project when we perform kernel / convolution matrix on an image, we delete current `activeShader` to create a new `Shader` that will become the current `activeShader` based on keyboard press.

Some things need to remember when using `malloc()`:

- If you use `malloc()` on a C++ class or struct, the constructor will not run. This means internal elements (like `std::string` fields) will remain uninitialized, leading to immediate program crashes.
- Memory allocated via `malloc()` lives forever until the program closes or until you call `free()` this is called memory leaks problem.
- It allocates raw bytes only, while `new` operator calls object constructors.
- It returns `void*` (requires cast), while `new` operator automatically returns correct type.
- It returns `NULL` when it has failure, while `new` operator throws `std::bad_alloc` exception.

The main point of this project example is to extract the RGB matrices, we can leave alone the matrix `A`, since it is sufficient with RGB / 3 channels only for CNN to do image classification.

Suppose we want to classify image based on the images that we have with label on each image as well (so we are not only depend on the MNIST dataset / images), we will need to learn from the two projects (the previous one and this current project), resize all of the images so the input layer will be uniform, then extract the RGB matrices of each image then perform the CNN operation to get the best model weights and biases that has high accuracy to predict the next image.

iii. Convolutional Neural Networks for Handwriting Recognition: Digit Classifier with SymIntegration

[SI*] Handwriting recognition is a classic, nontrivial task in artificial intelligence and machine learning and well-suited for neural networks, yet recognizing hand-written digits is still feasible as a first problem.

Fortunately, a large database of hand-written digits is available as the MNIST (modified National Institute of Standards and Technology) database, which contains 60,000 training images and 10,000 test images. The grayscale images have a size of 28×28 pixels and were digitized from digits written by employees of the United States Census Bureau and American high-school students.

The classification pipeline for the MNIST digit recognition with LeNet-5 architecture:

1. **Data Preparation:** Load the dataset (MNIST) and normalize pixel values from $[0, 255]$ to $[-1, 1]$ so the model converges faster during training.
2. **Compilation:** Select an optimizer between Adam or SGD (Stochastic Gradient Descent), a loss function (e.g. sparse categorical cross-entropy), and metrics.
3. **Input Layer:** Determine the input data, the standard MNIST digit recognition uses 28×28 images, we can choose whether to flatten the image into a vector with 784 elements (using the CSV file format) or to still use the images as 3D matrix (using the idx3-ubyte format), we will break down the differences between these two below.
4. **Convolutional Layers:** Apply filters (kernels) to slide over the image and extract distinct visual features like lines and loops.
5. **Activation function:** Use ReLU / tanh / sigmoid / scaled hyperbolic tangent activation function, it introduces non-linearity so the network can learn complex patterns.
6. **Pooling Layers:** Downsample the data to reduce spatial dimensions and computation while keeping the most critical features.
7. **Flattening:** Converts the multidimensional feature maps into a 1-dimensional array.
8. **Fully Connected Layers (Dense):** Interpret the extracted features and assign probabilities to each digit class (0 through 9).
9. **Output Layer:** Uses a softmax activation function to output the final prediction

$$z_i = w_i \cdot a_i + b_i$$

$$a_2 = \text{Softmax}(z_2)$$

The class with the highest probability is chosen as the predicted label.

Instead of using MNIST dataset, we create our own handwritten digits about 100 images thus in this section we will use our own handwritten digits as the input image, the explanation that is using MNIST dataset is just a model benchmark.

We will feed the training images data into the model in batches across multiple epochs to continuously adjust the weights and improve accuracy, once done we will test the model on unseen data (the test images) to calculate the accuracy and then using SFML (a multimedia C++ library) we can create a GUI to create a canvas where we can write a digit 0 – 9 and then the model will predict our hand-drawn digits.

At the start of making a C++ code for digit recognition with CNN we will have to choose whether to use **idx3-ubyte** file or a **CSV** file. The primary difference is how the data is structured and stored: **idx3-ubyte** is the original compressed binary format, while **CSV** is a plain-text representation created for human readability and easier parsing.

Feature	idx3-ubyte Binary Format	MNIST CSV Format
File Type	Raw Binary	Plain Text (ASCII)
Data Separation	Separated (Images and labels are stored in distinct files)	Unified (The label and its 784 pixels are stored in the same row)
Structure	3D Matrix ($60000 \times 28 \times 28$)	2D Tabular Row (1 label + 784 columns)
File Size	Very Small (± 47 MB unzipped)	Large (± 100 MB unzipped)
Readability	Requires a custom binary parser	Readable with Excel or Text Editors
I/O Speed	Blazing fast to load	Much slower due to text-to-number conversion

Table 10.3: The differences between the MNIST' **idx3-byte** file and an MNIST' **CSV** file.

The explanation of **idx3-ubyte** format:

- **The Name:** **idx** means indexed, **3** denotes it is a 3-dimensional tensor (number of images $\times$ rows $\times$ columns), and **ubyte** means "unsigned byte" (values from 0 to 255).
- **Separation:** To use it, you must download two separate files: **train-images-idx3-ubyte** (the raw pixels) and **train-labels-idx1-ubyte** (the correct digit label).

The explanation of MNIST' **CSV** format:

- **The Structure:** Each row in CSV represents exactly one handwritten digit.
- **Layout:** The very first column / the last column contains the integer **label**. The remaining 784 columns represent the flattened pixel values on the 28×28 image.
- **How to open:** It is highly accessible. It can be opened with a simple text editor like mousepad.

If you want to perform quick data exploration with small datasets you can use **CSV** format, otherwise if you are training heavy Deep Learning networks then **idx3-ubyte** format is the best as it minimizes storage overhead and optimizes GPU data-loading pipelines.

After we are done with the format for the input layer, then we are going to explain about the model architecture for this CNN digit recognition example. A model architecture is the design of the neural network with which we train the parameters in training process. We are using LeNet-5 architecture. It is pretty popular for its minimal structure and easy to train nature.

LeNet-5, a gradient-based learning model, developed in 1998 by Yann LeCun, Léon Bottou, Yoshua Bengio, and Patrick Haffner [17] was developed for the purpose of document recognition. LeNet-5 is a small network and very efficient convolutional neural network for handwritten character recognition and classification of different classes of objects in small resolution images, it contains the basic modules of deep learning: convolutional layer, pooling layer, and full link layer. It is the basis of other deep learning models. LeNet-5 has

a total of seven-layer, that do not comprise an input, each containing trainable parameters; each layer has a plurality of the pap, the feature, a characteristic of each of the input feature map extracted by means of a convolution filter, and then each feature map has multiple neurons.

This arrangement of layers was designed to automatically learn the features from the input images, rather than relying on hand-engineered features, setting a new standard for machine learning applications.




Figure 10.41: The architecture of LeNet-5, a convolutional neural network for digits recognition. Each plane is a feature map, i.e. a set of units whose weights are constrained to be identical.

The model comprises of total 1 input layer + 6 processing layers + 1 output layer.

1. Input Layer

In the data input layer, we make sure that the size of the input image is uniformly normalized to $32 \times 32 \times 1$, the "1" signifies that it is grayscale (single-channel, no color). The original MNIST dataset consists of 28×28 but were padded to 32×32 to ensure that the features at the edges of the image could be captured by the network's filters. Grayscale image still has a pixel value of $[0, 255]$, thus normalization (scales pixel values to a range of $[0, 1]$ or $[-1, 1]$), is also necessary for stable and faster training. This layer does not count as the network structure of LeNet-5. Traditionally, the input layer is not considered as one of the network hierarchy.

2. C1 Convolutional Layer

In LeNet-5, the first layer C1 performs the first convolutional operation on the input image (using 6 convolution kernels of size 5×5) to obtain 6 C1 feature maps. A filter, or kernel, is a small matrix of weights that slides over the image, performing a dot product (hadamard product) at each position, this process is also known as convolution, it allows the network to detect low-level features like horizontal line, edges, corners, and curves.

Input picture: $32 \times 32 \times 1$

it means we analyze one image at a time.

Convolutional kernel size: 5×5

Number of convolutional kernels: 6

Stride: 1

with a stride of 1, the filters/kernels move one pixel at a time.

Padding: 0

Output feature map size: 28×28

the output spatial dimension formula: $O = \frac{I-K+2P}{S} + 1$, with $I = 32, K = 5, P = 0, S = 1$

we will obtain $\frac{32-5+2(0)}{1} + 1 = 28$.

Number of neurons: $28 \times 28 \times 6 = 4,704$

Trainable parameters: $(5 \times 5 + 1) \times 6 = 156$

there are 25 unit parameters (the size of the convolutional kernel is $5 \times 5 = 25$) and one bias parameter per kernel/filter(+1 indicates a kernel has a bias), a total of 6 filters.

Number of connections: $(5 \times 5 + 1) \times 28 \times 28 \times 6 = 122,304$

each pixel in C1 is connected to 5×5 pixels and 1 bias in the input image, so there are 122,304 connections in total, but we only need to learn 156 parameters, mainly through weight sharing.

Output: $28 \times 28 \times 6$ feature maps

The mathematical formula for the C1 layer, for filter / kernel k ($k \in \{1, 2, \dots, 6\}$), the output position (i, j) is defined as:

$$y_{i,j}^k = f \left(\sum_{m=0}^{K-1} \sum_{n=0}^{K-1} w_{m,n}^k \cdot x_{i+m,j+n} + b^k \right)$$

where:

- x is the 2D input matrix or previous layer's feature map.
- w^k is the k -th filter/kernel of size $K \times K$ (in LeNet-5, $K = 5$).
- b^k is the learnable bias term for the k -th filter.
- $f(\cdot)$ is the activation function (LeNet-5 originally uses the scaled hyperbolic tangent function).
- $y_{i,j}^k$ is the pixel value at output coordinate (i, j) for the k -th feature map.

3. S2 Pooling Layer

The next layer is a pooling layer, which is used to reduce the spatial dimensions of the feature maps. Unlike a convolutional layer, a pooling layer does not learn a parameters. It is purely a deterministic mathematical function, we only train the parameters after the pooling operation (e.g., max pooling, average pooling, original LeNet-5 pooling). This has two key benefits: it reduces the number of parameters and computations in the network, and it helps to make the learned features more robust to small translations in the input image.

In LeNet-5, the S2 layer uses average pooling. It performs downsampling by taking the average of 2×2 non-overlapping windows from the previous layer (C1), multiplying by a trainable weight, adding a trainable bias, and applying a sigmoid activation function.

Input: $28 \times 28 \times 6$ feature maps

number of neurons from C1.

Input feature map size: 28×28

Sampling area: 2×2

Stride: 2

Sampling method: 4 inputs are added, multiplied by a trainable parameter, plus a trainable offset.

Number of feature maps / kernels: 6

Output feature map size: 14×14

pooling is performed using a 2×2 kernels, the output feature map size for a pooling (subsampling) layer in the LeNet-5 architecture is $O = \frac{I-K}{S} + 1 = \frac{28-2}{2} + 1 = 14$, K is the kernel size, if it is 2×2 then $K = 2$, S is the stride size there is no padding applied during LeNet-5 pooling stages, thus 6 feature maps of 14×14 are obtained. Each unit

connected to a 2×2 neighborhood in C1.

Number of neurons: $14 \times 14 \times 6 = 1,176$

Trainable parameters: $(1 + 1) \times 6 = 12$ (one weight and one bias per feature map)
(trainable weight / coefficient + bias) $\times$ filters

Number of connection: $(2 \times 2 + 1) \times 14 \times 14 \times 6 = 5,880$

Output: $14 \times 14 \times 6$ feature maps

The mathematical formula for each of the S2 layer's output feature map y_k , where $k \in \{1, 2, \dots, 6\}$, the operation is computed as:

$$y_k = f \left(w_k \cdot \frac{1}{4} \sum_{i=1}^4 x_i + b_k \right)$$

where:

- x_i represents the activations in the 2×2 receptive field of the previous C1 layer map.
- w_k is the trainable weight for that specific feature map.
- b_k is the trainable bias for that feature map.
- $f(\cdot)$ is the activation function (sigmoid or scaled hyperbolic tangent function)

4. C3 Convolutional Layer

The third layer for LeNet-5. The C3 layer is a convolutional layer that performs cross-correlation using 5×5 kernels. Its defining mathematical feature is a custom connectivity map where not every output map connects to every input map, reducing computation and enforcing varied feature extraction. Each feature map in C3 connected to all 6 or several feature maps in S2, indicating that the feature map of this layer is a different combination of the feature maps extracted from the previous layer, these are the combinations :

- **First 6 feature maps:** Connect to any **3 contiguous** subset of S2.
- **Next 6 feature maps:** Connect to any **4 contiguous** subset of S2.
- **Next 3 feature maps:** Connect to any **4 discontinuous** subset of S2.
- **Last 1 feature map:** Connects to all **6** input maps from S2.

for those who wonder why we obtain $10 \times 10 \times 16$ feature maps (16 feature maps of size 10×10) in C3 from $14 \times 14 \times 6$ feature maps in S2. The C3 layer's 16 feature maps are not fully connected to all 6 input feature maps from the previous layer (S2). This was originally done to limit computation and break symmetry in order to learn diverse, higher-level combinations of basic shapes found in C1 layer.

Thus, instead of simply applying every kernel/filter to every previous feature map, this layer combines subsets of the input feature maps from S2 / it uses sparse (partial) connectivity. This means individual kernels in C3 are only connected to specific combinations of feature maps from the previous layer.

In C3, the 16 kernels / filters scan the $14 \times 14 \times 6$ output from the preceding pooling layer (S2), capturing more complex visual representations by combining the basic features.

To find all the contiguous and discontinuous combinations of k elements from n available elements you can check the function `contiguousCombinations(int n, int k, int start_idx)` and `discontiguousCombinations(int n, int k, int start_idx)`, these functions are located in `src/operationsresearch.cpp` and explained in this book at chapter 5 Operations Research Computation with SymIntegration.

The first 12 feature maps are fixed because the contiguous combinations for choosing 3 elements from 6 available elements are 6 in total, the same as the contiguous combinations for choosing 4 elements from 6 available elements are also 6 in total, while the discontinuous combinations for choosing 4 elements from 6 available elements are 9 in total, and we only need 3, thus we will use a roulette wheel method, generating 4 random integers from 0 to 8 that are unique to each other. This roulette wheel method is adapted from genetic algorithms. The last feature map is also fix since it is a combination of all the 6 previous feature maps from S2.

Thus, in order to generate 4 unique integer random numbers that have uniform distribution with range $[0, 8]$ (C++ index starts from 0; 9 elements means the index will ends at 8) we can use this function in SymIntegration: `vrandn_uniqueinteger(0, 8, 4)`

```
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Operations Research/Test SymIntegration Unique Random Number Integer ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -gdb main.o -lstdc++ -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Operations Research/Test SymIntegration Unique Random Number Integer ]# ./main
3
8
0
Time taken by function: 610 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Operations Research/Test SymIntegration Unique Random Number Integer ]# ./main
5
2
7
Time taken by function: 563 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Operations Research/Test SymIntegration Unique Random Number Integer ]# ./main
3
7
0
Time taken by function: 575 microseconds
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Operations Research/Test SymIntegration Unique Random Number Integer ]# []
```

Figure 10.42: The 3 unique integer generated with `vrandn_uniqueinteger(0, 8, 3)` that can be used to determine the next 3 feature maps (after the first 12) in C3 that connect to any 4 discontinuous subset of S2, it is using a modified roulette wheel method. (*SymIntegration/Examples/Operations Research/Test SymIntegration Unique Random Number Integer/main.cpp*).

The roulette wheel method that is able to produce a set of unique integers is using `std::unordered_set<int>`, this is a standard C++ associative container that stores unique elements in no particular order. So if the set already contains the number 1, it cannot be inserted another 1 that is generated by the `std::uniform_int_distribution<>` with Mersenne Twister engine, it will generate again another integer random number until it obtains a number that is unique to all the numbers that are already in the set.

All discontiguous index combinations of 4 out of 6 are

```

0  1  2  4  → 0
0  1  3  4  → 1
0  1  3  5  → 2
0  2  3  4  → 3
0  2  3  5  → 4
0  2  4  5  → 5
1  2  3  5  → 6
1  2  4  5  → 7
1  3  4  5  → 8

```

So if we choose 3,7,0 from the roulette wheel method the index of feature maps in S2 that are chosen for the next 3 feature maps (after the first 12) in C3 are

```

0  1  2  4  → 0
0  2  3  4  → 3
1  2  4  5  → 7

```

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
0	X				X	X	X			X	X	X	X		X	X
1	X	X				X	X	X			X	X	X	X		X
2	X	X	X				X	X	X			X		X	X	X
3		X	X	X			X	X	X	X			X		X	X
4			X	X	X			X	X	X	X		X	X		X
5				X	X	X			X	X	X	X		X	X	X

Figure 10.43: The combinations to create feature maps in C3, each column indicates which feature map in S2 are combined to create a particular feature map in C3.

Output Maps (C3)	Inputs Connected (S2 Maps)	Number of Input Maps
Maps 0 to 5	Indices {0,1,2}, {1,2,3}, {2,3,4}, {3,4,5}, {4,5,0}, {5,0,1}	3
Maps 6 to 11	Indices {0,1,2,3}, {1,2,3,4}, {2,3,4,5}, {3,4,5,0}, {4,5,0,1}, {5,0,1,2}	4
Maps 12 to 14	Indices {0,1,3,4}, {1,2,4,5}, {2,3,5,0}	4
Maps 15	All indices {0,1,2,3,4,5}	6

Table 10.4: The input combinations for creating the feature maps in C3 layer, this is the C3 connectivity map. In the original LeNet-5 paper, the 16 output maps of C3 take input from the 6 maps of S2 using a custom subset table to form varying depths.

Input size: $14 \times 14 \times 6$

Input feature map size: 14×14

Convolutional kernel size: 5×5

Number of convolutional kernels: 16

Stride: 1

with a stride of 1, the filters/kernels move one pixel at a time.

Padding: 0

Output feature map size: 10×10

the output spatial dimension formula: $O = \frac{I-K+2P}{S} + 1$, with $I = 14, K = 5, P = 0, S = 1$ we will obtain $\frac{14-5+2(0)}{1} + 1 = 10$.

Number of neurons: $10 \times 10 \times 16 = 1,600$

Trainable parameters: $(5 \times 5) \times (6 \times 10) + 16 = 1,516$

the number of trainable parameters varies since C3 is not connected to every feature map of the previous layer. Instead, each feature map of C3 is connected to a subset of the previous maps. There are 16 kernels' receptive fields that are divided into 6 of 3 S2' features, 9 of 4 S2' features, and 1 of 6' S2 features (the size of the C3 convolutional kernel is $5 \times 5 = 25$) and one bias parameter per kernel/filter (+16 indicates that one kernel has only one bias).

Number of connections: $(10 \times 10) \times 1,516 = 151,600$

the partial connectivity reduces parameters and encourages diverse feature learning.

Output: $10 \times 10 \times 16$ feature maps

The mathematical formula for each of the C3 layer's output feature map y_k , where $k \in \{1, 2, \dots, 16\}$, the operation is computed as:

$$y_k = f \left(\sum_{m \in M_c} x_m * w_{k,m} + b_k \right)$$

where:

- y_k is the output feature map of the k -th channel in the C3 layer.
- $*$ denotes the 2D spatial cross-correlation operator.
- x_m is the m -th input feature map from the previous layer (S2).
- $w_{k,m}$ is the learnable 5×5 convolution kernel connecting the m -th input map to the k -th output map.
- b_k is the trainable bias for the k -th output map.
- $f(\cdot)$ is the activation function (typically the scaled hyperbolic tangent function).
- M_k is the subset of input maps from the S2 layer that are connected to the k -th output map.

5. S4 Pooling Layer

The next layer is another pooling layer. In LeNet-5, the S4 layer is a subsampling (average pooling) layer. It takes the 10×10 feature maps from the C3 layer and reduces them. For each of the 16 feature maps, S4 computes the average of 2×2 non-overlapping blocks, multiplies it by a trainable weight, adds a bias, and passes it through an activation function.

Input: $10 \times 10 \times 16$

Input feature map size: 10×10

Sampling area: 2×2

Stride: 2

Sampling method: 4 inputs are added, multiplied by a trainable parameter, plus a

trainable offset.

Number of feature maps / kernels: 16

Output feature map size: $\frac{10}{2} \times \frac{10}{2} = 5 \times 5$

pooling is performed using a 2×2 kernels, 16 feature maps of 10×10 are obtained. Each unit connected to a 2×2 neighborhood in C3.

Number of neurons: $5 \times 5 \times 16 = 400$

Trainable parameters: $(1 + 1) \times 16 = 32$ (one weight and one bias per feature map)
(coefficient + bias) $\times$ filters

Number of connection: $(2 \times 2 + 1) \times 5 \times 5 \times 16 = 2,000$

output size = $5 \times 5 \times 16$, the size of each feature map in S4 is $\frac{1}{2}$ of the size of the feature map in C3.

Output: $5 \times 5 \times 16$ feature maps

The mathematical formula for each of the S4 layer's output feature map y_k , where $k \in \{1, 2, \dots, 16\}$, the operation is computed as:

$$y_k = f \left(w_k \cdot \frac{1}{4} \sum_{i=1}^4 x_i + b_k \right)$$

where:

- y_k is the output of the subsampled feature map.
- x_i is the four input values from the 2×2 receptive field in the preceding C3 layer.
- w_k is a trainable weight, specific to the feature map.
- b_k is a trainable bias, specific to the feature map.
- f is the activation function (historically a scaled hyperbolic tangent, or tanh, in the original LeNet-5).

We can also write the forward pass equation for an average pooled unit from C3 to S4 k -th feature map as:

$$x_{i,j,k}^{S4} = f \left(w_k^{C3} \cdot \frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 x_{2i-2+m, 2j-2+n}^{C3} + b_k^{C3} \right)$$

with $i, j = 0, 1, 2, 3, 4$.

6. C5 Convolutional Layer

The next layer is a convolutional layer again. In LeNet-5, the C5 layer is a fully connected convolutional layer. It acts as the bridge between the convolutional/pooling features and the traditional dense neural network. The C5 layer has 120 featured maps, with each feature map measuring 1×1 . It connects to the output of the previous layer, S4, which consists of 16 feature maps, each measuring 5×5 , it then convolves them with 120 filters of size 5×5 . Since the size of the preceding layer's feature map (S4) precisely matches the size of the filter/kernel in C5 (5×5), there is no spatial reduction, then the output will be a feature map of size 1×1 / exactly 1 scalar value is produced per filter.

Input: $5 \times 5 \times 16$

Input feature map size: 5×5

Convolutional kernel size: 5×5

Output feature map size: 120×1

Trainable parameters: $((5 \times 5 \times 16) \times 120) + 120 = 48,120$
(input size $\times$ output feature map size + bias)

Number of connection: Parameters $\times$ output area = $48,120 \times 1 \times 1 = 48,120$ fully connected to all previous feature maps.

Output: $1 \times 1 \times 120$ a vector

The fundamental mathematical formula for the C5 layer's forward pass consists of two operations: **convolution** and **activation**.

(a) **The Convolution Operation**

For each output feature map o_k where $k \in \{1, 2, \dots, 120\}$, the operation is computed as:

$$o_k = \sum_{c=1}^{16} (w_{k,c} * x_c) + b_k$$

where:

- $*$ denotes the 2D discrete convolution operator.
- x_c is the c -th input feature map from the 16 feature maps of layer S4.
- $w_{k,c}$ is the 5×5 convolutional kernel connecting the c -th input map to the k -th output map.
- b_k is the learnable bias term for the k -th filter.

(b) **The Activation Function**

The resulting pre-activation from the convolution step is passed through a non-linear activation function (f). LeNet-5 originally used a scaled hyperbolic tangent ($\tanh$) activation:

$$y_k = f(o_k) = 1.7159 \tanh\left(\frac{2}{3}o_k\right)$$

7. F6 Fully Connected Layer

The next layer is a fully connected layer, which might remind you to feed-forward neural network. The LeNet-5 F6 layer processes the 120-dimensional output vector from the preceding C5 layer and outputs an 84-dimensional vector, this 84 nodes corresponding to a 7×12 ASCII bitmap matrix used to decode and recognize stylistically similar alphanumeric characters, -1 means white, 1 means black, so the black and white of the bitmap of each symbol corresponds to a code.

Input size: 120

Output size: 84

Trainable parameters: $(120 \times 84) + 84 = 10,164$
(input size $\times$ output size + bias)

Number of connection: fully connected to all previous neurons in the previous layer C5.

The mathematical formula for each of the F6 layer's output y_k , where $k \in \{1, 2, \dots, 84\}$,

the operation is computed as:

$$y_k = f \left(\sum_{j=0}^{119} w_{kj} x_j + b_k \right)$$

where:

- w_{kj} is the weight connecting the j -th input from C5 layer to the k -th neuron in the F6 layer.
- x_j is the j -th element of the 120-dimensional input vector from the C5 layer.
- b_k is the trainable bias for the k -th neuron in the F6 layer.
- f is the activation function, in the original LeNet-5 formulation, the F6 layer uses a scaled hyperbolic tangent function to introduce non-linearity and keep the network's values within a specific range:

$$f(x) = A \tanh(Sx)$$

with $A = 1.7159$ and $S = \frac{2}{3}$.

8. Output Layer

The output layer is also a fully connected layer, with a total of 10 nodes, which respectively represent the numbers 0 to 9, and if the value of node i is 0, the result of network recognition is the number i (if we are using RBF / radial basis function) For the case of digit recognition we will have the output layer of size 10.

$$\vec{y} = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \end{bmatrix}$$

Input size: 84

Output size: 10

Trainable parameters: $(84 \times 10) + 10 = 850$

Number of connection: 850 (fully connected to all previous neurons in the previous layer F6)

In the original LeNet-5 architecture, the output layer utilizes Radial Basis Function (RBF) units. Modern implementations, however, usually substitute this with a softmax layer for multi-class classification.

Radial Basis Function

Each class is represented by a Radial Basis Function (RBF) unit, where the output depends on the Euclidean distance between the input and its parameter vector, with larger distances indicating poorer fit. In the 1998 paper, each of the $i = 10$ output nodes (representing digits 0 through 9) evaluates how closely the 84-dimensional feature

vector x from the previous layer matches the predefined, hand-coded target bitmaps vector of that digit class.

The formula to compute the output of each RBF unit y_i is the squared Euclidean distance:

$$y_i = \sum_{j=1}^{84} (x_j - w_{ij})^2 \quad (10.85)$$

where:

- x_j represents the j -th element of the input vector from layer F6.
- w_{ij} is a fixed, non-trainable weight representing the j -th pixel value of a stylized 7×12 binary bitmap for character class i .
- The target class selected by the network is $\min_i(y_i)$, meaning the class with the minimum distance is the predicted digit.

Softmax

Most modern deep learning frameworks adapt LeNet-5 by replacing RBF layer with a standard Dense/Fully connected layer followed by a softmax activation function.

The output vector $\hat{y}$ is defined as

$$\hat{y}_i = \frac{e^{z_i}}{\sum_{k=0}^9 e^{z_k}}$$

where the raw logit z_i for class i is calculated via standard matrix multiplication:

$$z_i = \sum_{j=1}^{84} x_j w_{ij} + b_i$$

where:

- x_j is the j -th element of the input vector from layer F6.
- w_{ij} are the learnable weights of the j -th input vector from layer F6 to the i -th output vector in the output layer.
- b_i is the learnable bias of the output layer.
- z_k represents the logits for all 10 possible classes.

Backpropagation and Weights Update for LeNet-5

Backpropagation is a fundamental training algorithm in CNNs that propagates errors backwards through the network to optimise learning. It enables the model to adjust internal parameters and improve performance over time. In CNN convolutional layer, backpropagation updates the filter weights by calculating the gradient of the loss function with respect to each weight element using the cross-correlation between the input feature map and the upstream gradient map.

To define the formulas clearly, we track a single forward and backward pass with these variables:

- X as the input feature map to the convolutional layer.
- W as the filter (kernel) representing the shared weights.
- Z as the output feature map before activation ($Z = X * W + b$).
- L as the final scalar loss function value.

- $\frac{\partial L}{\partial Z}$ as the upstream gradient matrix received from the subsequent layer.

For a standard 2D convolution (mathematically implemented as valid cross-correlation), the value at position (i, j) of the output map Z is calculated as:

$$z_{i,j} = \left(\sum_m \sum_n x_{i+m,j+n} \cdot w_{m,n} \right) + b$$

During the backpropagation, we apply the calculus chain rule. Because a single weight $w_{m,n}$ is reused across every receptive field during the forward pass, its total impact on the loss is the sum of its impacts across all output positions:

$$\frac{\partial L}{\partial w_{m,n}} = \sum_i \sum_j \frac{\partial L}{\partial z_{i,j}} \cdot \frac{\partial z_{i,j}}{\partial w_{m,n}}$$

Since $\frac{\partial z_{i,j}}{\partial w_{m,n}} = x_{i+m,j+n}$, substituting this yields the final Weight Gradient Formula:

$$\frac{\partial L}{\partial w_{m,n}} = \sum_i \sum_j \frac{\partial L}{\partial z_{i,j}} \cdot x_{i+m,j+n}$$

The explicit sum describes a valid cross-correlation operation where the input feature map X acts as the larger matrix and the upstream gradient $\frac{\partial L}{\partial Z}$ acts as the sliding filter:

$$\frac{\partial L}{\partial W} = X * \frac{\partial L}{\partial Z}$$

To continue passing gradients to earlier layers, you must also compute how the loss changes with respect to the input matrix X . This requires a full convolution (padding the upstream gradient matrix) using a 180-degree rotated version of the weight filter:

$$\frac{\partial W}{\partial X} = \text{pad} \left(\frac{\partial L}{\partial Z} \right) * \text{rot}_{180}(W)$$

Once the weight gradient matrix $\frac{\partial L}{\partial W}$ is isolated, the shared weights are updated using gradient descent:

$$W^{\text{new}} = W^{\text{old}} - \eta \cdot \frac{\partial L}{\partial W}$$

with η as the learning rate and $\frac{\partial L}{\partial W}$ as the gradient with respect to weights.

In a convolutional neural network like LeNet-5, computing the weight gradient does not involve rotating the weight matrix. Instead, we compute the gradient with respect to the weights by performing a valid 2D convolution between the input features and the error gradients from the subsequent layer.

1. Weight Gradient (Filter/Kernel Update)

Operation: Valid 2D convolution (for C1, C3, C5 layer).

Method: To calculate the gradient for a specific filter, you perform a standard convolution (or cross-correlation) using the input patch as the feature map and the upstream error gradient (δ^{L+1}) from the next layer as the kernel.

Rotation Required: No rotation is required. We do not rotate the weight matrix to find the weight gradient. We simply correlate the incoming error with the layer's input.

2. Input Gradient / Error Propagation (δ) to Previous Layers

Operation: Full 2D convolution (for S2 and S4 layer).

Method: When passing error gradients backwards to the previous layers, we need to convolve the upstream error gradient (δ) with the layer's weights.

Rotation Required: Yes. To make the convolution mathematically align during this backward pass, the weight matrix (filter) must be rotated (or flipped) by 180° (flipped vertically and horizontally). We only perform full convolution on S2 and S4 layer because when we perform forward pass from S2 to C3 or S4 to C5 we use weights kernel of size $k \times k$.

Feature	Valid Convolution	Full Convolution
Padding	None	Implicitly pads the input
Output Size	Smaller than the input	Larger than the input
Formula	Output = $N - K + 1$	Output = $N + K - 1$
Pros	Highly accurate because all calculations are performed on real input pixels, not artificial zeros	Every pixel-including the original corner pixels-is visited exactly the same number of times by the filter.
Cons	The feature map shrinks with every layer. If your network is very deep, the data can shrink down too fast to extract meaningful hierarchical features.	The output includes elements that are calculated partially over "fake" zero-padded values, which can introduce artifacts along the edges.
Use Case	Standard CNN layers to extract features without "fake" padded data	Signal processing, polynomial multiplication, or inverting operations.

Table 10.5: Key differences between 2D valid convolution and 2D full convolution.

Now we will start the backpropagation / training the parameters from the last layer / output layer to the C1 layer in LeNet-5 architecture:

1. Output Layer

In LeNet-5, if we are using softmax in the output layer for classification tasks, then we will use cross-entropy loss to calculate the Loss:

$$L = - \sum y \log(\hat{y})$$

with y as the true label and $\hat{y}$ as the predicted label.

We can compute the delta (δ) at this output layer with this formula:

$$\delta_j^{\text{output layer}} = \frac{\partial L}{\partial a_j^{\text{output layer}}} f'(z_j^{\text{output layer}}) = f'(s_j^{\text{output layer}}) (y_j - \hat{y}_j)$$

with $\delta_j^{\text{output layer}}$ as the delta error term for the j -th neuron in the output layer, L is the loss or cost function (MSE or cross-entropy), $a_j^{\text{output layer}}$ is the actual output / activation

of the j -th neuron, $z_j^{output\ layer}$ is the weighted sum of inputs for the j -th neuron before activation ($z = \sum_i w_{ij}x_i + b_j$), $f'(z_j^{output\ layer})$ is the derivative of the activation function for the output layer, $f'(s_j^{output\ layer})$ as the derivative of softmax, which is a Jacobian matrix, $j = 1, 2, \dots, n_{classification}$.

MSE & Linear Activation

For an output node with a linear activation function, the delta combines the prediction error with the derivative of the identity function:

$$\delta_j^{output\ layer} = (\hat{y}_j^{output\ layer} - y_j)1$$

with y_j as the actual target value. This simplification is often used in frameworks because it is computationally cheaper and more numerically stable than calculating the full Jacobian matrix separately.

Cross-Entropy Loss & Softmax Activation

LeNet-5's original design uses Euclidean Radial Basis Functions (RBF) in the output layer, but modern implementations of LeNet-5 frequently swap this for a softmax output layer combined with cross-entropy loss. For the j -th neuron:

$$\delta_j^{output\ layer} = a_j^{output\ layer} - y_j$$

with $y_j = 1$ if the true class is j , and 0 otherwise.

After we have the $\delta_j^{output\ layer}$ then we can compute the weight update for the output layer:

$$w_{ij}^{new} = w_{ij}^{old} - \eta \delta_j^{output\ layer} y_i^{F6}$$

with η as the learning rate, $\delta_j^{output\ layer}$ as the j -th error term in the output layer, y_i^{F6} is the activated output from i -th neuron in F6 layer, $i = 1, 2, \dots, 84$, $j = 1, 2, \dots, 10$.

The bias update formula for the output layer is

$$b_j^{new} = b_j^{old} - \eta \delta_j^{output\ layer}$$

2. F6 (10,164 trainable parameters)

In LeNet-5, F6 layer is a fully connected layer with 84 neurons that uses a scaled hyperbolic tangent activation function. Since F6 layer has no weight sharing, every single parameter acts as a distinct connection between the source nodes (120 outputs of C5) and the destination nodes (84 neurons in F6). The weight update that is connecting the i -th neuron from C5 layer to the j -th neuron in the F6 layer:

$$w_{ij}^{new} = w_{ij}^{old} - \eta \frac{\partial L}{\partial w_{ij}} = w_{ij}^{old} - \eta \delta_j^{F6} a_i^{C5}$$

with η as the learning rate, δ_j^{F6} as the error gradient calculated for the j -th neuron in the F6 layer, a_i^{C5} is the activated output of the i -th neuron from the C5 layer,

$$i = 1, 2, \dots, 120, j = 1, 2, \dots, 84.$$

The bias update formula for the F6 layer is

$$b_j^{new} = b_j^{old} - \eta \delta_j^{F6}$$

Once the output delta is calculated, you can propagate the error backward to the preceding fully connected layer (F6).

The formula to compute the delta error term (δ_j^{F6}) during backpropagation is using the chain rule, which multiplies the downstream gradients by the derivative of the activation function, the formula is

$$\delta_j^{F6} = \left(\sum_k^{10} \delta_k^{output \ layer} w_{jk} \right) f'(z_j^{F6})$$

with δ_j^{F6} as the delta error term for the j -th neuron in F6 layer, $\sum_k^{10} \delta_k^{output \ layer} w_{jk}$ is the backpropagated error from the next layer (the output layer), this is the sum of the delta terms of all subsequent neurons k , weighted by their respective connection weights (w_{jk}) to neuron j in F6 layer, $f'(z_j^{F6})$ is the derivative of the scaled hyperbolic tangent activation function with respect to the pre-activation z_j .

The original LeNet-5 utilizes the following activation function in F6 layer

$$f(z) = 1.7159 \tanh \left(\frac{2}{3} z \right)$$

its derivative with respect to the pre-activation z is

$$f'(z) = \frac{(1.7159)(2)}{3} \text{sech}^2 \left(\frac{2}{3} z \right) \approx 1.1439 \left[1 - \tanh^2 \left(\frac{2}{3} z \right) \right]$$

3. C5 (48,120 trainable parameters)

In LeNet-5, the C5 layer is functionally treated as the first fully connected layer during the weight update process. Weights (W) are updated during the backward pass using standard gradient descent and the chain rule.

To update any weight w_{ij} in the C5 layer, the formula is

$$w_{ij}^{new} = w_{ij}^{old} - \eta \frac{\partial L}{\partial w_{ij}}$$

with η as the learning rate and $\frac{\partial L}{\partial w_{ij}}$ as the gradient of the loss function L (cross-entropy) with respect to that specific weight, $i = 1, 2, \dots, 400, j = 1, 2, \dots, 84$.

The bias update formula for the C5 layer is

$$b_j^{new} = b_j^{old} - \eta \delta_j^{C5}$$

Since C5 is a fully connected layer, a weight's gradient depends on the downstream error (from the F6 layer) and the upstream activation (from the S4 layer). The formula is

$$\frac{\partial L}{\partial w_{ij}} = \frac{\partial L}{\partial z_j} \frac{\partial z_j}{\partial w_{ij}} = \delta_j^{C5} a_i^{S4}$$

with w_{ij} as the weight connecting i -th neuron in S4 layer to the j -th neuron in C5 layer, δ_j^{C5} as the j -th neuron error term in C5 layer, and a_i^{S4} is the i -th neuron activation output in the previous S4 layer, z_j is the forward pass calculation from S4 layer to C5 layer $z_j = f(\sum_i w_{ij}x_i + b_j)$.

Remember that S4 layer has $16 \times 5 \times 5$ feature maps / 400 inputs that is fully connected to the 120 neurons in C5, the number of weights is $400 \times 120 = 48,000$ and the number of biases is 120.

Knowing that C5 layer acting as a convolutional layer with 5×5 kernels mapped over the entire S4 feature map. It contains 120 feature maps that collapse to a 1D vector of length 120.

For backpropagation when we arrive in layer C5, even if it is a convolutional layer, we do not use a full convolution operation between the error gradient δ^{F6} and the weight kernel in C5 to compute the input gradient (delta) for the C5 layer. Instead, we compute a standard matrix multiplication (vector-matrix product). Full convolutions are only used during backpropagation when we are passing gradients backward through a spatial convolutional layer (where the stride is 1) to restore the original dimensions of the input feature maps. Because the spatial dimensions of C5 and F6 are already reduced to 1×1 , there is no spatial structure to convolve over, making full convolution unnecessary.

The delta error term (δ_j^{C5}) for neuron / node j in the C5 layer represents the derivative of the activation function with respect to the layer's inactivated weighted input (z_j^{C5}). It is calculated by taking the sum of the propagated delta terms from the subsequent layer (F6).

Since C5 is fully connected to the subsequent layer F6, the error is propagated backward using matrix multiplication:

$$\delta_j^{C5} = \left(\sum_k \delta_k^{F6} w_{kj}^{F6} \right) f'(z_j^{C5})$$

where:

- δ_j^{C5} is the delta error term for the j -th neuron in layer C5.
- k is the indices of the nodes in the next layer (F6) connected to the j -th neuron in C5 layer.
- δ_k^{F6} is the delta error term of the k -th neuron in layer F6.
- w_{kj}^{F6} is the weight connecting the j -th neuron in C5 to the k -th neuron in F6.
- $f'(z_j^{C5})$ is the derivative of the pre-activated j -th neuron in C5 layer. Depending on the activation function that is used in C5 layer, if the activation function is a sigmoid then $f'(z_j^{C5}) = a_j(1 - a_j)$, with a_j as the activated j -th neuron in C5.

If the activation function is hyperbolic tangent then $f'(z_j^{C5}) = 1 - \tanh^2(z_j^{C5}) = 1 - a_j^2$, with a_j as the activated j -th neuron in C5.

$a_j = f(z_j)$ is the actual output / activation of the j -th neuron in C5 layer.

Once you compute and obtain 120 δ_j^{C5} , it is directly used to calculate how much the weights and biases in layer C5 contributed to the overall network error, allowing gradient descent to adjust them. With valid convolution the 120 δ^{C5} will become 48,000

δ^{C5} that will be used for computing weights gradient.

Gradient for weights

The mathematical formula to calculate the gradient of the loss with respect to a weight kernel tensor element is a convolution between the forward-pass input (from S4 to C5) and the error sensitivity (δ) from the subsequent layer:

$$\frac{\partial L}{\partial w} = 2Dvalidconv(X_k^{S4}, \delta_j^{C5})$$

Where:

- X_k^{S4} is the 5×5 input feature map from layer S4, $k = 1, 2, \dots, 16$.
- δ_j^{C5} is the 1×1 error gradient map / delta at layer C5 with $j = 1, 2, \dots, 120$.
- $2Dvalidconv()$ represents the valid convolution operation, the result will be the required 5×5 weight gradient matrix for C5 ($N - K + 1 = 5 - 1 + 1 = 5$).

Gradient for biases

The bias gradient is very simple, it is the sum of all error gradient in the corresponding feature map, we have 120 feature maps in C5, and the size is 1×1 for each feature map, thus

$$\frac{\partial L}{\partial b_j} = \delta_j^{C5}$$

with $j = 1, 2, \dots, 120$.

The computation for δ and weight gradient until layer C5 are still easy and quite similar with FNN. After this, the CNN technique begins, it gets more difficult to comprehend on how to compute δ and weight gradient for layer S4 to layer C1.

4. S4 (32 trainable parameters)

In LeNet-5 architecture, S4 layer is a trainable average pooling layer. Its weights are updated using the standard gradient descent backpropagation rule, multiplied by a learning rate η .

Because S4 has 16 feature maps, this gives exactly 32 trainable parameters. The 32 trainable parameters consists of 16 weights for each feature map, and 16 bias for each feature map in S4 layer.

There are exactly 400 deltas (or gradients) in the flattened S4 layer of LeNet-5, we will compute this 400 deltas first. Remember that the S4 layer is a subsampling (pooling) layer that consists of 16 feature maps, each with a spatial dimension of 5×5 , after flattened it becomes 400 neurons.

In backpropagation, there is exactly one delta (error gradient) δ computed for every activation or neuron in the layer. Computing delta in CNN is special, especially in the S4 pooling (subsampling) layer that is connected from C3 layer, and will be connected to C5 layer (C3 and C5 are convolutional layer).

To compute the delta (the error gradient, δ) for a layer during backpropagation in a neural network, you backpropagate the error from the layer above it. In the LeNet-5 architecture, S4 is an average subsampling (pooling) layer, the mathematical formulation

for computing its delta relies solely on upsampling the deltas of the following layer (C5) and applying the derivative of S4's activation function.

The exact mathematical formulation relies on whether you are analyzing LeNet-5 based on its original Yann LeCun publication specifications or adapting it for modern deep learning practices.

The δ (delta) error term represents the local gradient, or the derivative of the loss with respect to a neuron's pre-activation. For the subsampling/pooling layer (like S4 or S2), the delta is computed by backpropagating the errors from the downstream convolutional layer (C5) and upscaling them using a kronecker product (or upscale matrix).

Original LeNet-5 Specification

In the original leNet-5 architecture, the S4 average pooling operation is defined as a linear function with trainable coefficients, followed by a scaled hyperbolic tangent activation function.

The delta error term (δ_j^{S4}) for the j -th neuron in the S4 layer that is flattened is computed as

$$\delta_{flattened,j}^{S4} = \left[\sum_k \delta_k^{C5} w_{jk}^{S4 \rightarrow C5} \right] f'(z_j^{S4})$$

where:

- $\delta_{flattened,j}^{S4}$ is the delta error term of the j -th feature map in the S4 layer, $j = 0, 1, \dots, 399$.
- δ_k^{C5} is the delta error term of the k -th neuron in the C5 layer, $k = 0, 1, 2, \dots, 119$.
- w_{jk} is the weights connecting the j -th neuron in the pooling layer S4 to the k -th neuron in the next layer C5 (C5 is fully connected to S4, meaning each S4 output feeds into every C5 neuron).
- z_j^{S4} is the pre-activation input of the S4 layer.
- $f'(z_j^{S4})$ is the derivative of the activation function with respect to the pre-activated j -th neuron in S4 layer.

Computing δ^{S4} with SymIntegration

There are a lot of ways to compute the δ^{S4} , in SymIntegration we have created 2 arrays that represents the trainable weights for S4 layer:

s4_weights_kernel[16] (to represent the trainable weights for each feature map)

s4_bias[16] (to represent the trainable biases for each feature map)

Since S4 is connected to C5 layer, and in order to compute the delta (error gradient, δ) for the S4 layer we will need to propagate the gradients backward from the C5 layer. To connect S4 layer to C5 layer we create this array that represents the deltas of the flattened S4 layer and the C5 deltas:

c5_weights_kernel[400][120]

Flattening the $16 \times 5 \times 5$ S4 layer into a 400×1 vector does not change the 32 trainable parameters, the 32 parameters belong to the S4 downsampling operation, not the flattening process.

Before computing δ^{S4} , we must already have computed the deltas for the C5 layer (120 δ^{C5}). Since C5 is effectively fully connected to S4 (each C5 neuron takes input from all 400 neurons from flattened S4 layer), the error propagates backwards from the 120 C5 neurons.

To compute the 400 δ^{S4} we are using this formula:

$$\delta_{k,x,y}^{S4} = \left[\sum_{j=1}^{120} 2Dfullconv(\delta_j^{C5}, \text{rot180}(w_{jk}^{S4 \rightarrow C5})) \right] \oplus f'(z_{k,x,y}^{S4})$$

Where:

- δ_j^{C5} is the delta error term for the j -th neuron in layer C5.
- $\text{rot180}(w_{jk}^{S4 \rightarrow C5})$ is the rotated 5×5 filter/kernel connecting the k -th map in S4 layer to the j -th map of C5 layer, we have total of 48,000 trainable weights, which means in C5 there are 1,920 distinct weights kernels.
- $f'(z_{k,x,y}^{S4})$ is the derivative of the activation function in S4 layer with respect to the pre-activated neurons in S4 k -th map at spatial position (x, y) , $k = 1, 2, \dots, 16$, $x, y = 1, 2, \dots, 5$.
- $2Dfullconv(\dots)$ is the full convolution operation across all 120 feature maps of C5 layer for the padded δ_j^{C5} with the rotated weight kernel $\text{rot180}(w_{jk}^{S4 \rightarrow C5})$, we didn't write it in the formula explicitly, but this is the equivalent:

$$2Dfullconv(\delta_j^{C5}, \text{rot180}(w_{jk}^{S4 \rightarrow C5})) = 2Dvalidconv(\text{pad}(\delta_j^{C5}, 4), \text{rot180}(w_{jk}^{S4 \rightarrow C5}))$$

pad(...,n) is the notation to pad each 1×1 C5 delta map with 4 rows and columns of zeros on all sides (making it 9×9). This step ensures the backward convolution outputs an array the same size as the S4 map.

- $\oplus$ represents the element-wise multiplication (Hadamard product).

When propagating error, computing δ^{L-1} from δ^L for layer S4 and S2 we do need to rotate the weight matrix, the rotated weight matrix is used to convolve with the error gradients to pass the error signal back to the previous layer. Propagating error requires a mathematical operation that reverses the spatial dimensions to match the error gradient's direction back to the input space (which is why a 180^0 rotation is needed).

About Rotated 180^0 Matrix rot180 is the notation for 180 degree 2D matrix rotation, in backpropagation the weights must be rotated 180^0 for proper convolution gradient alignment. To perform a 180^0 rotation on a 2D matrix you reverse the rows and reverse the columns of the matrix. For a given weight matrix W , the rotated matrix W' satisfies $w'_{m,n} = w_{k-m, k-n}$, $k = 5$, $m, n = 1, 2, 3, 4, 5$.

$$W = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$W' = \text{rot180}(W) = \begin{bmatrix} i & h & g \\ f & e & d \\ c & b & a \end{bmatrix}$$

After we have obtained 400 δ^{S4} from backpropagating 120 δ^{C5} we can then shrink this 400 error gradient into 16 δ^{S4} that represents each feature map in S4 pooling layer.

In the original pooling layer S4 we have 16 features map with size of 5×5 each, thus if it is flattened then the first 25 entries will represents the first feature map, till the last 25 entries that represents the last feature map. In the flattened vector of size 400 we will partition every 25 δ^{S4} to represents each feature map and it will be used to compute the weight updates in S4 layer.

Gradient for weights

$$\frac{\partial L}{\partial w_{ji}} = \sum_{pixels} \frac{\partial L}{\partial z^{S4}} \frac{\partial z^{S4}}{\partial u^{S4}} \frac{\partial u^{S4}}{\partial w_{ji}}$$

with z^{C3} as the pre-activated feature map values from the previous layer (C3) corresponding to the receptive field, δ^{S4} is the error term propagated backward from the C5 layer to the S4 layer, x and y are the spatial coordinates within the feature map. $\frac{\partial L}{\partial z^{S4}}$ is the gradient propagated from the C5 layer, $\frac{\partial z^{S4}}{\partial u^{S4}} = f'(u)$ is the derivative of the activation function with respect to the pre-activated u^{S4} , $\frac{\partial u^{S4}}{\partial w_{ji}} = \frac{1}{4} \sum_{l=1}^4 x_l^{C3}$ is the sum of the 4 inputs in the 2×2 receptive field from C3 layer, multiplied by $\frac{1}{4}$ for average pooling.

Once the 400 delta δ^{S4} are computed for S4 layer, the gradients for the S4 layer's trainable weights are computed by taking the sum over the batch of the element-wise product between the delta and the C3 layer's output feature maps.

Thus the formula to compute the total gradient for the k -th feature map' weight become:

$$\frac{\partial L}{\partial w_k} = \sum_{i=1}^5 \sum_{j=1}^5 \delta_{k,i,j}^{S4} \left(\frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 x_{k,2i-2+m,2j-2+n}^{C3} \right)$$

with $i, j = 1, 2, 3, 4, 5$ as the pixel that represents the spatial position for the 5×5 size of each feature map in S4 layer.

Once the gradients are computed, the weight update in the S4 layer using standard gradient descent formula is

$$\begin{aligned} w_k^{new} &= w_k^{old} - \eta \frac{\partial L}{\partial w_k} \\ &= w_k^{old} - \eta \frac{\partial L}{\partial a_{k,i,j}} \frac{\partial a_{k,i,j}}{\partial w_k} \\ &= w_k^{old} - \eta \sum_{i,j \in \text{map } k} \delta_{k,i,j}^{S4} \left(\frac{1}{4} \sum_{l=1}^4 x_l^{C3} \right) \end{aligned}$$

with η as the learning rate and $\frac{\partial L}{\partial w_j}$ as the gradient of the loss function L (cross-entropy) with respect to that specific weight for the k -th feature map in S4 layer, $i, j = 1, 2, \dots, 5$, $k = 1, 2, \dots, 16$, a_i^{S4} is the output activation for the i -th neuron in the S4

layer $\frac{\partial a_{k,i,j}^{S4}}{\partial w_k} = f'(w_k \cdot \frac{1}{4} \sum_{l=1}^4 x_l^{C3} + b_k) \cdot \frac{1}{4} \sum_{l=1}^4 x_l^{C3}$, and $\frac{\partial a_{k,i,j}^{S4}}{\partial w_k} = \frac{1}{4} \sum_{l=1}^4 x_l^{C3}$, x_l are the four activations in the 2×2 pooling window from the preceding C3 convolutional layer.

Thus

$$\frac{\partial L}{\partial w_k} = \sum_{i=1}^5 \sum_{j=1}^5 \delta_{k,i,j}^{S4} \cdot x_{k,i,j}^{S4} = \sum_{i=1}^5 \sum_{j=1}^5 \delta_{k,i,j}^{S4} \left(\frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 x_{k,2i-2+m,2j-2+n}^{C3} \right)$$

where:

- k is the index of the feature map for S4 layer, $k = 1, 2, \dots, 16$.
- (i, j) is the spatial position for each feature map in S4, $i, j = 1, 2, \dots, 5$.
- $x_{k,i,j}^{S4}$ is the original forward-pass activations of the k -th feature map in S4 at spatial position (i, j) , $a_{k,i,j}^{S4} = f \left(w_k * \left(\sum_{m=1}^2 \sum_{n=1}^2 x_{m,2i-2+m,2j-2+n}^{C3} \right) + b_k \right)$, $x_{k,i,j}^{S4} = \left(\sum_{m=1}^2 \sum_{n=1}^2 x_{m,2i-2+m,2j-2+n}^{C3} \right)$.

For weight gradients we do not need to rotate or flip any matrices, we simply compute the unrotated convolution of the input volume (from the previous layer) and the incoming error gradients δ (from the next layer). Weight updating merely measures how much each specific weight contributed to the loss. This is achieved by element-wise multiplication (dot product) of the incoming gradient matrix and the exact receptive field it interacted with during the forward pass. Because we are doing cross-correlation to compute the weight gradient, the weight kernel stays in the original orientation. Cross-correlation is a standard valid convolution without rotating any matrices / filter matrix.

Gradient for biases

Thus the formula to compute the total gradient for the k -th feature map' bias become:

$$\frac{\partial L}{\partial b_k} = \sum_{i=1}^5 \sum_{j=1}^5 \delta_k^{S4}$$

with $i, j = 1, 2, 3, 4, 5$ as the pixel that represents the spatial position for the 5×5 size of each feature map in S4 layer.

The bias update formula for the S4 layer is

$$b_k^{new} = b_k^{old} - \eta \left(\sum_{i=1}^5 \sum_{j=1}^5 \delta_k^{S4} \right)$$

with η as the learning rate and $k = 1, 2, \dots, 16$. Since the bias is added uniformly as a scalar constant to the pre-activation sum before passing through the activation function, its direct partial derivative is the sum of the gradients flowing back from the downstream activation. Because the bias term b is additive, its rate of change relative to a_j is exactly 1, $\frac{\partial a_j^{S4}}{\partial b_j} = 1$.

Important variables and dimensions when computing delta in S4:

- **S4 Output:** 16 feature maps of size 5×5 , flattened into a 400-element array.

- **C5 Layer:** 120 fully-connected neurons.
- w_{ji} is the weight connecting the j -th neuron in S4 to the i -th neuron in the C5 layer (dimension: 400×120).
- δ_i^{C5} is the error term of the i -th neuron in the C5 layer.
- w_j^{S4}, b_j^{S4} are the trainable weights and bias for the S4 average pooling layer (32 total parameters: 16 weights and 16 biases).
- a_j^{S4} is the activation (output) of the j -th node in layer S4.

5. C3 (1,516 trainable parameters)

In LeNet-5, the C3 layer's weight change formula relies on gradient descent and backpropagation. The weight update formula formula is

$$w^{new} = w^{old} - \eta \frac{\partial L}{\partial w}$$

with η as the learning rate, $\frac{\partial L}{\partial w}$ as the gradient of the error with respect to the specific weight.

Using the chain rule, the gradient for a specific weight in C3 is the sum over all pixels in the C3 feature map of the local error gradient (delta, δ^{C3}) multiplied by the corresponding input pixel from the previous layer (S2).

The C3 layer produces exactly 16 feature maps. During backpropagation, one delta (error gradient) matrix is generated for each feature map. Therefore, the C3 layer produces 16 delta matrices, each with dimensions of 10×10 .

For a C3 feature map k and S2 feature map j :

$$\frac{\partial L}{\partial w_{kj}} = \sum_x \sum_y \delta_k(x, y) \cdot a_j(x + m, y + l)$$

where:

- $\delta_k(x, y)$ is the error gradient at output coordinate (x, y) of C3 feature map k .
- a_j is the activation / output of the S2 feature map j that is connected to C3 feature map k .
- m, l are the spatial offsets of the 5×5 kernel mask.

Weight Sharing in C3

Because C3 is a convolutional layer, it utilizes weight sharing. A single weight value is applied to multiple connections across the feature map. Therefore, the total gradient for a single weight is the sum of all individual gradients calculated for every spatial position where that weight is shared.

Unlike a standard modern convolutional layer, in LeNet-5 C3 is only partially connected. Out of $16 \times 6 = 96$ possible combinations between the 6 S2 feature maps and 16 C3 feature maps, only 60 connections are active (the rest are zeroed out). The gradient is only calculated and updated for the 1,516 active weights and 16 biases in this layer.

Computing the delta (δ) for the C3 layer in LeNet-5 involves calculating the error gradient $\frac{\partial L}{\partial z}$ with respect to the pre-activation value z . This is done via backpropagation, where error gradients are passed backward from the next layer (S4) to C3. Because C3 is a convolutional layer, this requires propagating the deltas backward from the subsequent S4 pooling layer, adjusted for C3's unique partially-connected feature map

scheme.

Passing δ^{S4} to C3 Layer

Because S4 layer performs average pooling, the delta from the m -th feature map of S4 (δ_m^{S4}) gets upsampled to match the C3 dimension before being propagated backwards. If S4 has a 2×2 pooling window, we distribute (broadcast) the error to the four corresponding units in the C3 feature map.

Subsampling for S4 Delta

Since S4 is an average pooling layer with trainable weights and biases. During backpropagation, its delta is mapped as follows:

- **Error Upsampling:** The errors flowing back from S4 (which has a 5×5 spatial dimension) must be mapped back to the 10×10 dimensions of C3. Since pooling condenses the image by taking the average (e.g., a 2×2 pool is unrolled into 4 identical values), the incoming delta is upsampled.

Because S4 uses average pooling with a 2×2 window and stride of 2, forward pooling takes an average of 2×2 block. During backpropagation when we compute delta for C3, to reverse this, we must distribute the error term from a single S4 neuron equally across the 2×2 block (a technique often called Kronecker product or "upsampling" the error gradient).

For a 2D feature map matrix x of shape $H \times W$ and a subsampling (pooling) factor t , the upsampled matrix $\text{upsample}(x)$ is computed via the Kronecker product:

$$\text{upsample}(x) \equiv \frac{1}{t * t} \cdot x \otimes \mathbf{1}_{t \times t}$$

with $\mathbf{1}_{t \times t}$ as a $t \times t$ matrix filled with 1s.

For example if we have

$$\delta_k^{S4} = 2.3$$

then the Kronecker product is

$$2.3 \xrightarrow{\text{upsampling}} \frac{1}{4} \begin{bmatrix} 2.3 & 2.3 \\ 2.3 & 2.3 \end{bmatrix}$$

Applying the Kronecker product upsamples it back by mapping each individual pixel into a 2×2 block of identical values (given that the current pooling layer is an average pooling).

- **Scaling / Apply the Layer's Scale Weight**

After upsampling the delta from S4 layer, multiply them with the corresponding weight from S4 (the 16 trainable weights for each feature map in S4) at that current feature map.

- **Gradient Application:** The upscaled delta will be 1,600, and it will be multiplied (element-wise / Hadamard product) by the derivative of the C3 layer's activation function.

Formula to Compute δ^{C3}

During backpropagation, the delta for a specific neuron or spatial position (i, j) in a

feature map k of the C3 layer is computed using the chain rule. It is the sum of the upsampled upstream deltas from all connected feature maps in the next layer (S4) multiplied by the derivative of the activation function:

$$\delta_{k,i,j}^{C3} = w_k^{S4} \cdot f'(z_{k,i,j}^{C3}) \oplus \text{upsampled}(\delta_{k,i,j}^{S4})$$

where:

- $\delta_{k,i,j}^{C3}$ is the error gradient for the k -th feature map in layer C3 in feature map k with spatial position i, j .
- $f'(z_{k,i,j}^{C3})$ is the derivative of the activation function with respect to the pre-activation z_k^{C3} at layer C3 in feature map k with spatial position i, j .
- $\delta_{k,i,j}^{S4}$ is the upsampled error gradient of the k -th S4 feature map with spatial position i, j .
- w_k^{S4} is the trainable weight from S4 layer, $k = 1, 2, \dots, 16$.
- $\oplus$ is the Hadamard (element-wise) product.

To get the final delta for layer C3 and C1, that has pooling layer afterwards, we only need to multiply the upsampled δ^{S4} (for C3) or δ^{S2} (for C1), then multiply it by the trainable weight w_k^{S4} (for C3) or w_k^{S2} (for C1). Because a single S4 value is derived from four C3 values, distributing the gradient backward requires scaling it by the exact same shared weight used during the forward pass.

We do not add the S4 bias during the delta computation in backpropagation. In neural networks, biases are strictly used during the forward pass to shift activations. During backpropagation, the derivative of a constant bias with respect to the loss is zero, meaning it does not contribute to the error gradient flowing backward to the previous layer.

The Special Case of C3 Sparse Connections

The C3 layer in LeNet-5 is not fully connected to the previous layer, S2. Instead each of the 16 feature maps in C3 selectively receives inputs from a subset of S2's 6 feature maps (sparse connections). So when backpropagating, the error for an S2 feature map is the sum of the gradients flowing back through all filters in C3 that were connected to it (the gradients from upstream layers are strictly routed backward only through these pre-defined mapped connections).

When updating $w_{ij}^{S2 \rightarrow C3}$ (weights for the convolution kernel from S2 to C3), you only calculate and update the weights for the valid, allowed connections as defined by the classic LeNet-5 adjacency table (Table 10.4), setting non-existent connections to zero.

Gradient for weights

Once you have the delta δ_j^{C3} for every node, you can calculate the gradients to update the C3 layer's trainable parameters (the weights):

$$\frac{\partial L}{\partial w_{ij}} = \sum_{x,y} \delta_j^{C3} * x_i^{S2}$$

To calculate the weight gradients for the C3 convolutional layer in LeNet-5, you use the standard backpropagation chain rule. The gradient for any specific weight is the

sum of the partial derivatives of the loss with respect to that weight across all spatial positions where the weight was applied (due to weight sharing).

To perform this calculation, you need the following standard variables from both the forward and backward passes:

- m, n as the spatial dimensions of the C3 feature maps (10×10).
- δ_k^{C3} as the error gradient matrix (delta) for the k -th feature map in layer C3.
- X_j^{S2} as the input feature map from the previous layer (S2) into the C3 convolutional filter.
- W_{kj} as the weight tensor connecting the j -th feature map in S2 to the k -th feature map in C3 (5×5).
- L as the loss or cost function.

Because C3 layer in LeNet-5 is a convolutional layer, the same set of weights is slid across the entire image. Therefore, the total gradient for a specific weight w is the sum of its gradients over all coordinates (x, y) in the feature map:

$$\frac{\partial L}{\partial W_{kj}} = \sum_{x=1}^m \sum_{y=1}^n \frac{\partial L}{\partial z_{k,x,y}^{C3}} \cdot \frac{\partial z_{k,x,y}^{C3}}{\partial W_{kj}}$$

where:

- $\frac{\partial L}{\partial z_{k,x,y}^{C3}}$ is the delta $\delta_{k,x,y}^{C3}$ for the specific neuron at map k and position (x, y) .
- $\frac{\partial z_{k,x,y}^{C3}}{\partial W_{kj}}$ is the local receptive field of the previous layer's activations at that position.

In practice, you do not calculate the gradient for each pixel individually. Instead, you compute it as a full convolution between the input activations and the error deltas:

(a) **Calculate the 1600 deltas for C3**

First, backpropagate the errors from the next layer (S4) into C3. This gives you the error matrix δ^{C3} , there will be $16 \times 10 \times 10$ matrix of deltas.

To backpropagate the error delta from an average pooling layer S4 to a convolutional layer C3 in LeNet-5, you must reverse the forward-pass operations by: **upsample the S4 deltas** and pass them through the derivative of the activation function used in C3.

i. **Upscale the S4 Deltas**

Layer S4 has 16 feature maps of size 5×5 . The pooling operation in S4 is an average pooling, which downsampled the C3 layer (S4 pools a 2×2 neighborhood into a single pixel). During the backward pass, each individual delta in the 5×5 S4 maps is "unpooled" (upsampled) into a 2×2 region. This spreads the error back out/ distribute the error uniformly backward to match the original C3 shape, expanding the resolution to 10×10 . Each delta value in S4 is scaled by $\frac{1}{M+N}$ and broadcast to a 2×2 block in the corresponding C3 feature map.

The formula is:

$$\text{upsample}(\delta_i^{S4}) = \delta_i^{S4} \times \frac{1}{4}$$

Because S4 layer is an average pooling layer, its deltas are scaled by $\frac{1}{4}$ as they are mapped backwards, the scale formula is $\frac{1}{M+N}$ for 2×2 pooling $M = 2, N = 2$.

The result will be $16 \times 10 \times 10$ matrices filled with upsampled δ^{S4} .

ii. **Multiply by S4 Trainable Weight**

Multiply the upsampled error matrix by the S4 layer's specific weight w_k^{S4} .

iii. **Apply the Activation Function Derivative**

In LeNet-5, the activation function for C3 is the hyperbolic tangent (tanh). The gradient (derivative) must be applied to the upscaled deltas for each neuron.

$$\tanh'(x) = 1 - \tanh^2(x)$$

The backpropagation computation is to multiply the upscaled delta maps element-wise by the derivative of the C3 layer output evaluated during the forward pass (pre-activated output).

$$\delta_{k,x,y}^{C3} = w_k^{S4} \cdot f'(z_{k,x,y}^{C3}) \oplus \text{upsampled}(\delta_{k,x,y}^{S4})$$

the 1,600 deltas in C3 can be represented with the above formula, at feature map k , spatial position (x, y) , notice that the **upsampled** is different than **upsample** notation, the **upsampled** notation means that the 400 δ^{S4} is already upsampled from $16 \times 5 \times 5$ into $16 \times 10 \times 10$ size that match C3 layer / the 400 deltas of S4 are map directly onto the 16 corresponding feature maps of C3.

After upsampling δ^{S4} from 16 feature maps of 5×5 dimensions to 10×10 dimensions, scale / multiply with the trainable weight in S4 layer (16 trainable weights) and multiply the upscaled delta maps with derivative of the pre-activated output in C3 layer we will obtain 1,600 backpropagated deltas for C3 (δ^{C3}) / error delta matrix for C3 of size $16 \times 10 \times 10$. Upsample is also called Kronecker product operation.

(b) **2D Convolution**

Perform a valid convolution operation on the input activations X_j^{S2} with the C3 layer's unrotated deltas δ_k^{C3} with $k = 1, 2, \dots, 16$, a matrix delta for each feature map in C3.

The convolution of the unrotated 180 degree delta map with the corresponding 14×14 input feature map from S2 is used to compute the spatial gradient correctly to obtain the weight gradient for the 1,516 trainable parameters. The resulting 5×5 matrix of the convolution operation will represent the gradient $\frac{\partial L}{\partial w_{kj}}$ for that specific kernel. In S4 layer we perform a **rot180** towards the weight kernel $W^{S4 \rightarrow C5}$, thus in this C3 layer when computing δ^{C3} we do not need to rotate the matrix delta, instead we use upsample operation to obtain δ^{C3} .

Alternatively, you can calculate this efficiently as a cross-correlation: Keep the delta map unflipped, but slide the 14×14 S2 input matrix over the 10×10 delta map, computing the sum of element-wise products.

(c) **Compute the Weight Gradient**

Now, we can compute the 1,516 weight gradients by convolution operation formula:

$$\frac{\partial L}{\partial W_k} = 2D\text{validconv}(X_j^{S2}, \delta_k^{C3})$$

$2Dvalidconv$ is a valid convolution operation which don't add artificial zeros for the matrices, no padding. The kernel that will slide over the matrix X_j^{S2} is the δ_k^{C3} matrix. The resulting matrix will be of size $N - k + 1 = 14 - 10 + 1 = 5$ thus we will obtain sixteen 5×5 weight gradients matrix.

The weight update in the C3 layer using standard gradient descent formula is

$$w_{kj}^{new} = w_{kj}^{old} - \eta \left(2Dvalidconv(X_j^{S2}, \delta_k^{C3}) \right)$$

Gradient for biases

The bias gradient for a specific feature map is simply the sum of all deltas in that feature map:

$$\frac{\partial L}{\partial b_k} = \sum_{x=1}^{10} \sum_{y=1}^{10} \delta_{k,x,y}^{C3}$$

x, y is the spatial position in the k -th feature map.

The bias update in the C3 layer using standard gradient descent formula is

$$b_k^{new} = b_k^{old} - \eta \left(\sum_{x=1}^{10} \sum_{y=1}^{10} \delta_{k,x,y}^{C3} \right)$$

6. S2 (12 trainable parameters)

Layer S2 has 6 feature maps, and each feature map has exactly 1 trainable weight (coefficient) for the sum of the 4 inputs (2×2 receptive field) and 1 trainable bias (offset). An average pooling layer does not use complex, multi-dimensional trainable filter weights. Layer S2 and S4 are average pooling layers in the original LeNet-5. Many modern variations replace this with max pooling, which generally yields sharper edge detection features and speeds up training convergence. Average pooling layer on the other side assigns equal weight to all features in the window, avoiding the loss of background or texture information. Average pooling was chosen due to its feature smoothing, it acts as a low-pass filter, creating smoother feature maps that help the network recognize general shapes without over-indexing on localized sharp edges. While average pooling is foundational to the original design, most modern CNNs prefer max pooling.

- Max pooling selects the maximum activation within the window, effectively isolating the strongest features(e.g., the most distinct stroke of a handwritten digit).
- Because max pooling selects the most prominent feature, it produces larger gradients during back-propagation, which can help the network train faster and achieve better classification accuracy.

For the S2 average pooling layer, we compute the delta by propagating the gradients backwards from the upstream layer. The mathematical operations required to compute and propagate the delta depend on the type of pooling (max / average / sum only pooling), the sum only pooling adds the four inputs in a 2×2 window, multiplies them by a trainable weight, adds a trainable bias, and passes the result through tanh activation function, the average pooling will take the average of the four inputs in a 2×2 window, and the max pooling will take the largest value as the output of the four inputs in a 2×2 window.

Delta Calculation

From S2 to C3 layer, the forward pass performs a 2D convolution using a set of weights / kernels. During the backward pass, to calculate the error gradients for the previous layer, you must perform a cross-correlation. The mathematical equivalent of doing this is running a standard convolution using a flipped (180-degree rotated) version of the weight matrix. Rotating the weights allows the network to smoothly propagate the derivative of the error backward through the exact spatial reverse of the forward pass.

S2 layer has 1,176 neurons from its $6 \times 14 \times 14$ feature map, thus we will compute 1,176 delta first. To find the delta for a specific S2 map, sum the full 2D convolution of the C3 deltas with the flipped C3 weights kernel. Do this only for the C3 maps that are connected to that S2 map.

This is just a food for thought: if you take a look at the S2 to C3 connectivity map, excluding the feature map 13-15 in C3 layer that connect to 4 discontinuous feature maps in S2 layer, you will realize then that each feature map in S2 from $k = 1, 2, \dots, 6$ will appear 8 times each in the first 12 feature maps and the last feature map of C3 layer.

The formula for the delta of the j -th feature map in S2 layer is

$$\delta_{k,x,y}^{S2} = f'(z_{k,x,y}^{S2}) \oplus \left(\sum_{k \in \text{Connected}(j)} 2D\text{validconv}(\text{pad}(\delta_j^{C3}, 4), \text{rot180}(W_{jk}^{S2 \rightarrow C3})) \right)$$

where:

- $\delta_{k,x,y}^{C3}$ is the delta map (size 10×10) of the k -th feature map in C3.
- W_{jk} is the 5×5 weight kernel connecting the k -th map in S2 to the j -th map in C3 layer.
- **rot180(...)** is the notation to rotate the 5×5 weight matrix 180 degree to align it for the backward pass.
- **2Dvalidconv** is the notation to perform 2D valid convolution between two matrices, the padded δ^{C3} and the flipped W_{ij} , the flipped / rotated weight will be the kernel matrix that will slide over the delta matrix. If we do not want to pad the delta manually first we can use the **2Dfullconv(...)** it will pad the delta matrix automatically while running the operation.
- **pad(...,n)** is the notation to pad each 10×10 C3 delta map with 4 rows and columns of zeros on all sides (making it 18×18). This step ensures the backward convolution outputs an array the same size as the S2 map.
- $\sum_{k \in \text{Connected}(j)}$ / sum maps means for each S2 map, sum the results of all convolutions coming from its connected C3 maps. This gives you exactly 6 S2 delta feature maps, ready to be passed to the S2 to C1 backward pass.
- $\oplus$ is the element-wise (Hadamard) multiplication.
- $f'(z_{k,x,y}^{S2})$ is the derivative of the activation function with respect to the pre-activated neuron in k -th feature map of S2 layer at spatial position (x, y) , $k = 1, 2, \dots, 6$, $x, y = 1, 2, \dots, 14$.

Gradient for weights

During training, the change in the weight ΔW is proportional to the negative gradient of the loss function (L) with respect to W , scaled by the learning rate (η).

$$\Delta W = -\eta \frac{\partial L}{\partial W}$$

Using the chain rule, $\frac{\partial L}{\partial W}$ expands to:

$$\frac{\partial L}{\partial W} = \sum \delta^{S2} \frac{\partial a^{S2}}{\partial W}$$

where:

- δ^{S2} is the error term propagated from the next layer.
- $\frac{\partial a^{S2}}{\partial W}$ is the derivative of the S2 activation.

$$\frac{\partial a^{S2}}{\partial W} = f'(W \cdot \text{avg}(a^{C1}) + b) \cdot \text{avg}(a^{C1})$$

Because the S2 layer uses average pooling, the gradient of the loss function L with respect to a specific weight w_k (for k -th feature map, $k = 1, 2, \dots, 6$) requires you to reverse the pooling operation

- Sum of the Gradients:** For each of the 6 feature maps, sum all the $14 \times 14 = 196$ corresponding delta values (δ_k^{S2}) in the S2 layer.
- Weight Gradient** $\left(\frac{\partial L}{\partial w_k}\right)$: Multiply this sum by the average of the 2×2 pixel region from the previous layer (C1).

Mathematically, for a single feature map, the weight gradient is:

$$\frac{\partial L}{\partial w_k^{S2}} = \sum_{i=1}^{14} \sum_{j=1}^{14} \delta_{k,i,j}^{S2} \left(\frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 x_{k,2i-2+m,2j-2+n}^{C1} \right)$$

with $k = 1, 2, \dots, 6$.

To apply the weight for the next iteration, we apply standard gradient descent. The weight update in the S2 layer using standard gradient descent formula is:

$$\begin{aligned} W^{new} &= W^{old} - \eta \frac{\partial L}{\partial W} \\ &= W^{old} - \eta \left(\sum_{i=1}^{14} \sum_{j=1}^{14} \delta_{k,i,j}^{S2} \left(\frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 x_{k,2i-2+m,2j-2+n}^{C1} \right) \right) \end{aligned}$$

Gradient for biases

The gradient for the bias is simply the sum of all deltas in that specific feature map:

$$\frac{\partial L}{\partial b_k^{S2}} = \sum_{i=1}^{14} \sum_{j=1}^{14} \delta_{k,i,j}^{S2}$$

The bias update in the S2 layer using standard gradient descent formula is

$$b_k^{new} = b_k^{old} - \eta \left(\sum_{i=1}^{14} \sum_{j=1}^{14} \delta_{k,i,j}^{S2} \right)$$

7. C1 (156 trainable parameters)

To backpropagate the error delta from an average pooling layer (S2) to a convolutional layer (C1) in LeNet-5, you must reverse the forward-pass operations: upsample the S2 deltas and pass them through the derivative of the activation function used in C3.

(a) Upsample the S2 Deltas

Layer S2 has 6 feature maps of size 14×14 . The pooling operation in S2 is an average pooling, which downsampled the C1 layer. During the backward pass, each individual delta in the 14×14 maps is "unpooled" (upsampled) into a 2×2 region. This spreads the error back out, expanding the resolution to 28×28 .

$$\text{upsample}(\delta_i^{S2}) = \frac{1}{4} \delta_i^{S2}$$

the resulting array will be six 28×28 matrices.

(b) Scale / Multiply by S2 Trainable Weight

Multiply the upsampled error matrix by the S2 layer's specific weight w_k^{S2} .

(c) Apply the Activation Function Derivative

In LeNet-5, the activation function for C1 is the hyperbolic tangent (tanh). The gradient (derivative) must be applied to the upscaled deltas for each neuron.

$$\tanh'(x) = 1 - \tanh^2(x)$$

Backpropagation: Multiply the upscaled delta maps element-wise by the derivative of the C1 layer output evaluated during the forward pass.

(d) Result: 4,704 Deltas

This upsampling of 6 feature maps of 14×14 dimensions to 28×28 dimensions perfectly accounts for the 4,704 values across C1 (6 maps $\times$ 28 rows $\times$ 28 columns). Thus the formula to compute the 4,704 δ^{C1} is

$$\delta_{k,i,j}^{C1} = (\text{upsample}(\delta_{k,x,y}^{S2})) \cdot w_k^{S2} \cdot f'(z_{k,i,j}^{C1})$$

Gradient for weights

The weight update in the C1 layer using standard gradient descent formula is

$$W^{new} = W^{old} - \eta \frac{\partial L}{\partial W}$$

Generally, the delta (error term) for any convolutional neuron i is calculated by multiplying the upstream error by the derivative of its activation function:

$$\delta_i = \frac{\partial L}{\partial u_i} = \frac{\partial L}{\partial a_i} \frac{\partial a_i}{\partial u_i}$$

where:

- $\frac{\partial L}{\partial a_i}$ is the summed gradient flowing backwards from all upstreams neurons connected to neuron i .
- $\frac{\partial a_i}{\partial u_i} = f'(u_i)$ is the derivative of the activation function with respect to the pre-activation u_i

The weight gradient for C1 layer is computed by performing a valid 2D convolution between the input image pixels (at C1) and the delta map (δ^{C1}).

$$\frac{\partial L}{\partial w_{ij}^{C1}} = 2D\text{validconv}(X^{input}, \delta_k^{C1})$$

$2Dvalidconv(X^{input}, \delta_k^{C1})$ will produce a 6 matrices with size $N - K + 1 = 32 - 28 + 1 = 5$ / six 5×5 weight gradient matrix.

Thus the weight update formula for C1 layer is

$$w_{k,i,j}^{new} = w_{k,i,j}^{old} - \eta \left(2Dvalidconv(X^{input}, \delta_k^{C1}) \right)$$

Gradient for biases

Once you have the delta (δ^{C1}), you use it to find the gradients for C1's biases, the gradient of the loss with respect to a bias for a specific feature map in C1 is the sum of all deltas in that corresponding feature map.

$$\frac{\partial L}{\partial b_k^{C1}} = \sum_{i=1}^{28} \sum_{j=1}^{28} \delta_{k,i,j}^{C1}$$

with $k = 1, 2, \dots, 6$.

The bias update in the C1 layer using standard gradient descent formula is

$$b_k^{new} = b_k^{old} - \eta \left(\frac{\partial L}{\partial b_k^{C1}} \right) = b_k^{old} - \eta \left(\sum_{i=1}^{28} \sum_{j=1}^{28} \delta_{k,i,j}^{C1} \right)$$

We have summarized forward pass, delta backpropagation computation, the weight gradient, and bias gradient formula in tables (10.7), (10.8), (10.9), and (10.10).

LeNet-5 has seven layer in total, does not comprise an input layer, each containing a trainable parameters; each layer has a plurality of the feature map, a characteristic of each of the input feature map extracted by means of convolution filter, and then in each there are multiple neurons.




Figure 10.44: The LeNet-5 architecture & pipeline for digit recognition. It is a very efficient convolutional neural network for handwritten character recognition.

Layer	Feature Maps	Size	Kernel	Stride	Activation
Input	1	32×32	-	-	-
Conv1 (C1)	6	28×28	5×5	1	Leaky ReLU ($\alpha = 0.8$)
Avg Pool (S2)	6	14×14	2×2	2	Leaky ReLU ($\alpha = 0.8$)
Conv2 (C3)	16	10×10	5×5	1	Leaky ReLU ($\alpha = 0.8$)
Avg Pool (S4)	16	5×5	2×2	2	Leaky ReLU ($\alpha = 0.8$)
Conv3 (C5)	120	1×1	5×5	1	Leaky ReLU ($\alpha = 0.8$)
FC (F6)	84	-	-	-	Leaky ReLU ($\alpha = 0.8$)
Output	10	-	-	-	softmax

Table 10.6: Summary of LeNet-5 architecture.

LeNet-5's significance in deep learning cannot be overstated. It was one of the first demonstrations that convolutional neural networks could be successfully applied to visual pattern recognition. LeNet-5 introduced several key concepts that are now standard in CNN architectures, including the use of multiple convolutional and pooling layers, local receptive fields, shared weights, and the backpropagation algorithm for training the network. The development of LeNet was also driven by the need for automation in industries that dealt with large volumes of handwritten documents, such as banks and postal services, where automatic digit recognition could significantly speed up processing times and reduce errors. These innovations have paved the way for the development of more complex and deeper networks, which are the backbone of modern artificial intelligence systems in various applications ranging from autonomous vehicles to medical diagnosis. This architecture leads to the development of more sophisticated deep learning frameworks that continue to push the boundaries of what machines can learn and achieve.

The concept of using localized receptive fields, shared weights, and spatial hierarchies in neural networks was inspired by studies of the visual cortex in animals, suggesting that these biological processes could be mimicked to improve machine perception.

Layer	δ	Trainable Parameters	Forward Pass Formula
Conv1 (C1)	4,704	156 (150w + 6b)	$a_k^{C1} = f(\text{conv2D}(W_k^{C1}, X^{input}) + b_k^{C1}); k = 1, 2, \dots, 6$
Avg Pool (S2)	1,176	12 (6w + 6b)	$a_{i,j,k}^{S2} = f\left(w_k^{S2} \cdot \frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 a_{2i-2+m, 2j-2+n}^{C1} + b_k^{S2}\right); i, j = 1, 2, \dots, 28; k = 1, 2, \dots, 6$
Conv2 (C3)	1,600	1,516 (1,500w + 16b)	$a_k^{C3} = f\left(\sum_{i \in \text{connection}} \text{conv2D}(X_i^{S2}, W_i^{S2 \rightarrow C3}) + b_k^{C3}\right); k = 1, 2, \dots, 16$
Avg Pool (S4)	400	32 (16w + 16b)	$a_{i,j,k}^{S4} = f\left(w_k^{S4} \cdot \frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 x_{2i-2+m, 2j-2+n}^{C3} + b_k^{S4}\right); i, j = 1, 2, \dots, 10; k = 1, 2, \dots, 16$
Conv3 (C5)	120	48,120 (48,000w + 120b)	$a_k^{C5} = f\left(\sum_{j=1}^{16} \text{conv2D}(W_{j,k}^{C5}, X_j^{S4}) + b_k^{C5}\right); k = 1, 2, \dots, 120, j = 1, 2, \dots, 16$
FC (F6)	84	10,164 (10,080w + 84b)	$a_k^{F6} = f\left(\sum_{j=1}^{120} (W_{j,k}^{F6}, X_j^{C5}) + b_k^{F6}\right); k = 1, 2, \dots, 84, j = 1, 2, \dots, 120$
Output	10	850 (840w + 10b)	$\hat{y}_i = \frac{e^{z_i}}{\sum_{k=0}^9 e^{z_k}}; z_i = \sum_{j=1}^{84} x_j w_{ij} + b_i$

Table 10.7: Summary of LeNet-5 architecture forward pass formula.

Layer	δ	δ Backpropagation Formula
Output	10	$\delta_j^{output} = \hat{y}_j - y_j; j = 1, 2, \dots, 10$
FC (F6)	84	$\delta_j^{F6} = f'(z_j^{F6}) \left(\sum_{k=1}^{10} \delta_k^{output} w_{kj}^{F6 \rightarrow output} \right); j = 1, 2, \dots, 84$
Conv3 (C5)	120	$\delta_j^{C5} = f'(z_j^{C5}) \left(\sum_{k=1}^{84} \delta_k^{F6} w_{kj}^{C5 \rightarrow F6} \right); j = 1, 2, \dots, 120$
Avg Pool (S4)	400	$\delta_{k,x,y}^{S4} = \left[\sum_{j=1}^{120} 2Dfullconv(\delta_j^{C5}, \text{rot180}(w_{jk}^{S4 \rightarrow C5})) \right] \oplus f'(z_{k,x,y}^{S4}); k = 1, 2, \dots, 16, x, y = 1, 2, \dots, 5$
Conv2 (C3)	1,600	$\delta_{k,i,j}^{C3} = w_k^{S4} \cdot f'(z_{k,i,j}^{C3}) \oplus \text{upsample}(\delta_{k,x,y}^{S4}); k = 1, 2, \dots, 16; x, y = 1, 2, \dots, 5; i, j = 1, 2, \dots, 10$
Avg Pool (S2)	1,176	$\delta_{k,x,y}^{S2} = f'(z_{k,x,y}^{S2}) \oplus \left(\sum_{k \in \text{Connected}(j)} 2Dfullconv(\delta_j^{C3}, \text{rot180}(W_{jk}^{S2 \rightarrow C3})) \right); x, y = 1, 2, \dots, 14; k = 1, 2, \dots, 6$
Conv1 (C1)	4,704	$\delta_{k,i,j}^{C1} = w_k^{S2} \cdot f'(z_{k,i,j}^{C1}) \oplus \text{upsample}(\delta_{k,x,y}^{S2}); k = 1, 2, \dots, 6; x, y = 1, 2, \dots, 14; i, j = 1, 2, \dots, 28$

Table 10.8: Summary of LeNet-5 architecture backpropagation formula, z_j^L means pre-activated neuron in layer L.

Layer	Trainable Parameters	Weight Gradient Formula
Output	840w + 10b	$\frac{\partial L}{\partial w_{ij}^{output}} = \delta_j^{output} * f(z_i^{F6}); i = 1, 2, \dots, 84$
FC (F6)	10,080w + 84b	$\frac{\partial L}{\partial w_{ij}^{F6}} = \delta_j^{F6} * f(z_i^{C5}); i = 1, 2, \dots, 120$
Conv3 (C5)	48,000w + 120b	$\frac{\partial L}{\partial w_{ij}^{C5}} = 2Dvalidconv(X_k^{S4}, \delta_j^{C5}); j = 1, 2, \dots, 120 \quad k = 1, 2, \dots, 16$
Avg Pool (S4)	16w + 16b	$\frac{\partial L}{\partial w_{k,i,j}^{S4}} = \sum_{i=1}^5 \sum_{j=1}^5 \delta_{k,i,j}^{S4} \left(\frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 x_{k,2i-2+m,2j-2+n}^{C3} \right); k = 1, 2, \dots, 16$
Conv2 (C3)	1,500w + 16b	$\frac{\partial L}{\partial w_{jk}^{C3}} = 2Dvalidconv(X_j^{S2}, \delta_k^{C3}); j = 1, 2, \dots, 6, k = 1, 2, \dots, 16$
Avg Pool (S2)	6w + 6b	$\frac{\partial L}{\partial w_{k,i,j}^{S2}} = \sum_{i=1}^{14} \sum_{j=1}^{14} \delta_{k,i,j}^{S2} \left(\frac{1}{4} \sum_{m=1}^2 \sum_{n=1}^2 x_{k,2i-2+m,2j-2+n}^{C1} \right); k = 1, 2, \dots, 6$
Conv1 (C1)	150w + 6b	$\frac{\partial L}{\partial w_{ij}^{C1}} = 2Dvalidconv(X^{input}, \delta_k^{C1}); k = 1, 2, \dots, 6, i, j = 1, 2, \dots, 28$

Table 10.9: Summary of LeNet-5 architecture weight update formula.

Layer	Trainable Parameters	Bias Gradient
Output	840w + 10b	$\frac{\partial L}{\partial b^{output}} = \delta_j^{output}; j = 1, 2, \dots, 10$
FC (F6)	10,080w + 84b	$\frac{\partial L}{\partial b^{F6}} = \delta_j^{F6}; j = 1, 2, \dots, 84$
Conv3 (C5)	48,000w + 120b	$\frac{\partial L}{\partial b^{C5}} = \delta_j^{C5}; j = 1, 2, \dots, 120$
Avg Pool (S4)	16w + 16b	$\frac{\partial L}{\partial b_k^{S4}} = \sum_{i=1}^5 \sum_{j=1}^5 \delta_{k,i,j}^{S4}$
Conv2 (C3)	1,500w + 16b	$\frac{\partial L}{\partial b_k^{C3}} = \sum_{x=1}^{10} \sum_{y=1}^{10} \delta_{k,x,y}^{C3}; k = 1, 2, \dots, 16$
Avg Pool (S2)	6w + 6b	$\frac{\partial L}{\partial b_k^{S2}} = \sum_{i=1}^{14} \sum_{j=1}^{14} \delta_{k,i,j}^{S2}; k = 1, 2, \dots, 6$
Conv1 (C1)	150w + 6b	$\frac{\partial L}{\partial b_k^{C1}} = \sum_{i=1}^{28} \sum_{j=1}^{28} \delta_{k,i,j}^{C1}; k = 1, 2, \dots, 6$

Table 10.10: Summary of LeNet-5 architecture bias update formula.

[SI*] We have learned a nice C++ code for digit recognition using SFML from here:

<https://github.com/AdamYuan/SimpleNN>

The input vector is flattened to become a vector of 784 elements.

The good thing is this code has a feature to make user interface handwriting with SFML library, so the neural network that is being used will be trained and create around 300,000 parameters, that is way too many compared to LeNet-5 architecture, afterwards with the trained parameters / weights and biases it will be the model to predict our handwritten digits from 0 to 9.

The downside of that code is the documentation / the book that cover the formula being used is not complete or established nicely, thus instead of spending time to figure out the formula behind that code, we create our own code based on LeNet-5 architecture, that we will explain below.

[SI*] We have created a CNN C++ code for digit recognition, modifying from the one that we have used in FNN, this code can predict the classification with softmax output and train the training images with steepest descent by using **CNN_LeNet5** class.

When we finished with the code, we learned that the loss keeps increasing, this is due to the learning rate that we set at first is too high at $\eta = 0.5$, for CNN with LeNet-5 architecture $\eta = 0.01$ is the highest / the cap for starter, but since we are using our own digit handwriting with 60 training images thus we set $\eta = 0.1$ then the loss is decreasing, if any readers facing the same problem, these are the cause that makes the loss increases when backpropagating / training in CNN:

1. The Learning Rate is Too High

If the learning rate is set too high, the gradients calculated during backpropagation will adjust the network's weights by massive amounts. This causes the model to step completely over the lowest point (optimum) of the loss function, making the loss get progressively worse with each training step. To fix this try lowering the learning rate by an order of magnitude (e.g., reduce from 0.1 to 0.01 or 0.0001) or implement a learning

scheduler to dynamically decrease the rate during training.

2. Exploding Gradients

During the backward pass in deep CNNs, error gradients can multiply exponentially as they travel backward through the layers. This causes weight updates to become overwhelmingly large, rendering the network unstable. To fix this try to implement gradient clipping, a technique that caps the maximum value of gradients during backpropagation.

3. Vanishing Gradients

Conversely, the gradients can become infinitesimally small as they propagate backward if we are using tanh or sigmoid activation function. This halts the network's ability to update its weights in the earlier, feature-extracting convolutional layers, while later layers may overfit. To fix this try to use ReLU (or its variants) in the hidden layers to mitigate this issue. If the network is very deep, consider incorporating residual connections (like those used in ResNet).

4. Incorrect Loss Function Implementation

The loss function quantifies how wrong the CNN predictions are. If the custom loss function or the framework's loss criterion is misaligned with the final activation function, the backpropagated gradients will push the network in the wrong direction. To fix this ensure that the final layer's activation matches the loss function. For example if the output's activation is softmax then the loss function is categorical cross-entropy.

5. Incorrect Data or Label Mismatch

If the gradients and loss calculations are mathematically sound, the issue may be in the data pipeline. If the training labels are noisy, misaligned with the images, or the input data pipeline is unnormalized, the model cannot learn meaningful patterns. To fix this double-check the data augmentation logic, ensure batches are properly shuffled, and verify that the target labels match the expected ground truth.

We have written our own handwriting digits, about 100 handwriting digits from 0 to 9. It is not like MNIST that has 60,000 handwriting digits, the main purpose is for educational only, how CNN LeNet-5 architecture works, will it works with our own handwritten digits even if it is only 100? Based on theory, it should work.




Figure 10.45: The handwritten digits that we have created on a paper.




Figure 10.46: We process this by uploading the photo to computer, crop each digit written, naming each of them nicely so it is well clustered (give unique label to each), then convert it to **idx3-ubyte** file and **idx1-ubyte** file.

1. Preparing the Input Images and Convert Them to IDX3-Ubyte

The reader can use their own handwriting or the one that we have created.

First crop each digit and name the file that match the label for the resizing, grayscaling and converting purpose till the CNN phase.

Then all we have to do is to resize all the images and turn them into grayscale, **STB Image** is very handy for this purpose. You can take a look at **SymIntegration/Exam-**

ples/Neural Networks/Test Resize and Convert to Grayscale Images with STB Image/ put all the images there and compile the **main.cpp**, from the current working directory open xterm / terminal then type:

```
make  
./main
```

All the images will be resized to 32×32 and become grayscale images, the name is also written with **gray_(originalname)**.

Then we move all the resized and grayscaled images to a new directory that will process them and convert the images into **idx3-ubyte** file, it is in **SymIntegration/Examples/Neural Networks/Test Create idx3-ubyte Files with STB Image/**, after you copy the images you can open terminal there and type:

```
bash createimageslist
```

(this will create **images.txt** that will store the images name)

You still need to create manually **labels.txt** to store the label name for each images, but if you can do it automatically it is good then. Afterwards, type again

```
make  
./main
```

The images have been successfully converted to **idx3-ubyte** file along with the corresponding labels to **idx1-ubyte** file.

The function:

```
convert_to_idx(images, labels, "train-images-idx3-ubyte", "train-labels-idx1-ubyte",  
target_width, target_height)
```

will create the **idx3-ubyte** to represents the images and **idx1-ubyte** to represents the corresponding labels for the images.

We have finished preparing for the input, move the **train-images-idx3-ubyte** and **train-labels-idx1-ubyte** to the directory for the CNN training in **SymIntegration/Examples/Neural Networks/Test CNN using Class with Steepest Descent**.

2. Normalize the Input Image to the $[-1, 1]$ Range

After we moved the **train-images-idx3-ubyte** and **train-labels-idx1-ubyte** files to this directory **SymIntegration/Examples/Neural Networks/Test CNN using Class with Steepest Descent** all we have to do is normalize the grayscale input image. Normalizing grayscale images to the $[0, 1]$ range is significantly better for training a LeNet-5 CNN than leaving the raw pixel values at $[0, 255]$. But since we are using the LeNet-5 with tanh activation functions we will proceed to normalizing into $[-1, 1]$ range, it is slightly better than $[0, 1]$ because the inputs are strictly positive with $[0, 1]$ this forces the first layer of tanh activations to operate entirely in its upper quadrant, introducing a massive bias, causing earlier gradient saturation, and slowing down the overall optimization process. Here is why centering the data around zero helps the neural network train faster and more accurately:

- **Faster Convergence:** Neural networks optimize weights by calculating gradients. When input data is centered around zero, the gradients flow more evenly. This helps the model find the optimal weights in fewer training steps. Large input values (like 255) cause large weight updates, making the training process unstable

or causing the model to diverge.

- **Better Gradient Flow:** Prevents numerical instability and ensures that gradients propagate properly to the earlier layers of the network during backpropagation. Large, positive-only inputs like $[0, 255]$ can cause weight updates to push neurons into saturation, particularly with sigmoid and tanh activation functions. Saturated neurons have near-zero gradients, which effectively stops the network from learning. The derivative of $\tanh(x)$ is highest near 0 ($f'(0) = 1$) and saturates (flattens out) as values approach $+1$ or -1 . Inputs scaled to $[-1, 1]$ fall neatly into high-sensitivity zone, preventing the vanishing gradient problem in early training stages.
- **Symmetry:** LeNet-5 traditionally uses tanh activation functions in its hidden layer. Since tanh outputs values between $[-1, 1]$, having inputs on the same scale helps the network map features much more effectively. Leaky ReLU activation function can also work with normalized input that has range $[-1, 1]$.
- **Zero-Centered Input:** The tanh function has inflection point at exactly 0. Normalizing the inputs to $[-1, 1]$ creates a zero-centered dataset. This prevents gradients from systematically pushing in the same direction during backpropagation, resulting in faster and more stable convergence.

The formula to normalize to $[-1, 1]$:

$$\text{Normalized Pixel} = \frac{\text{Pixel Value} - 127.5}{127.5}$$

Under standard training conditions on the 60,000 images MNIST dataset, this configuration with normalized grayscale images to a $[-1, 1]$ typically yields a test accuracy between 98.5% and 99.0%.

shuffledXtrain.txt	input_images.txt
1 = 1	
2 129 129 129 129 130 130 130 129 129 129 130 129 129 130 132 132 132 132 134 134 133 133 133 133 134 134 134 134 135	
3 130 130 130 131 130 130 129 129 130 131 130 130 131 131 131 132 132 134 134 133 132 133 133 133 133 134 134 133 135 134	
4 131 131 131 132 131 130 130 131 130 131 131 130 131 132 132 132 133 133 133 134 133 133 133 133 133 134 134 134 134 134	
5 131 131 132 132 132 131 130 131 131 131 131 132 132 132 132 133 133 133 133 135 134 134 134 134 134 134 134 134 135 133 135	
6 131 130 132 131 131 130 130 131 132 131 131 132 132 131 131 134 134 132 133 134 134 135 135 134 134 134 134 134 134 135 134 135	
7 130 131 131 131 132 132 132 132 132 132 132 133 132 132 132 135 135 135 134 135 135 135 135 135 134 133 134 134 135 135 136	
8 131 132 131 132 132 132 132 131 131 132 132 132 132 133 132 134 134 136 135 135 135 135 135 136 135 135 134 134 136 135 135	
9 132 132 131 132 131 132 132 131 131 132 132 132 131 132 132 133 135 136 136 136 136 136 136 136 136 136 136 135 136 136 136	
10 133 133 132 132 132 131 131 132 132 132 132 132 133 133 133 133 134 135 135 136 136 136 135 135 136 136 136 136 136 136	
11 132 132 131 131 131 131 132 132 131 132 132 132 133 134 133 133 135 135 136 137 137 137 136 136 136 137 137 137 137 137 138	
12 133 132 132 132 131 132 133 133 132 133 133 133 132 133 135 134 134 134 135 136 135 136 135 135 136 136 137 137 137 137 137	
13 133 132 132 132 132 132 133 132 133 133 133 133 132 133 134 134 134 134 135 136 135 136 135 135 136 136 137 136 137 136	
14 133 132 132 132 131 133 132 133 132 133 133 133 133 134 135 135 130 118 110 115 131 136 137 137 137 137 137 137 136 137 137	
15 133 134 133 132 131 132 133 135 134 133 133 132 133 133 133 127 115 102 87 84 84 102 129 136 136 136 137 138 137 136 137 138	
16 134 134 133 133 132 134 133 134 134 133 133 132 132 134 129 97 84 95 113 121 112 86 102 134 136 136 137 138 138 136 137 137	
17 134 134 134 134 134 135 134 134 134 135 133 133 133 134 130 92 101 127 134 136 136 109 84 127 136 137 136 138 138 138 138 138	
18 134 134 134 135 135 134 134 134 135 135 135 135 136 134 123 127 135 135 136 137 123 85 118 136 136 136 138 138 138 138 138 137	
19 135 135 135 135 135 134 135 135 136 135 135 135 135 136 135 136 137 137 138 125 88 117 136 136 137 137 138 138 139 137	
20 136 136 136 136 136 135 136 135 135 135 135 136 135 136 136 137 137 135 138 137 124 90 121 137 138 137 138 138 138 137 137	
21 135 136 136 136 136 136 135 136 135 135 134 135 136 136 136 136 137 138 137 138 138 124 92 126 138 138 137 137 137 138 138	
22 135 136 136 135 135 135 135 135 135 135 136 136 136 137 137 137 137 138 137 139 120 94 130 138 137 137 138 138 138 138 138	
23 136 136 135 135 135 134 135 136 137 135 136 137 137 137 137 137 135 137 138 138 137 112 100 131 136 137 136 137 137 138 138	
24 136 136 135 137 135 135 136 136 136 136 137 136 137 138 136 136 138 138 133 99 114 135 137 137 138 139 139 139 139 138	
25 136 135 135 136 135 136 137 136 135 136 136 137 137 137 137 138 138 137 126 119 109 91 120 137 137 137 138 139 139 140 139	
26 137 136 136 136 137 135 135 136 135 136 134 137 137 136 138 138 131 110 94 89 84 81 91 123 128 133 138 139 139 140 140 140	
27 137 137 137 137 136 135 137 136 136 137 137 137 137 138 131 105 91 107 112 93 105 83 86 91 121 138 139 139 141 140	
28 138 138 137 137 137 136 137 137 136 137 137 138 138 138 126 89 114 126 104 108 133 121 101 111 131 138 138 139 140 141	
29 137 139 138 137 137 137 137 138 137 137 136 137 137 138 133 106 99 102 103 132 137 136 136 136 137 139 139 140 140 141 141	
30 138 138 138 138 136 136 137 137 138 137 136 137 137 135 131 114 116 132 137 138 138 138 138 138 138 138 139 141 141 141	
31 138 138 139 139 138 137 137 136 138 139 138 138 138 138 138 138 139 139 140 139 140 139 140 140 140 140 140 141 141	
32 139 140 139 140 139 138 138 138 139 139 139 139 138 140 141 140 141 142 141 141 141 140 140 140 140 140 140 141 141	
33 138 139 139 139 139 138 139 140 140 140 140 140 140 140 141 141 143 141 140 140 140 140 140 140 141 141 141 141 141	
34	

Figure 10.47: The original input image is in standard grayscale range $[0, 255]$.


```

{
    double epoch_loss = 0.0;
    for (size_t iter = 0; iter < X.size(); ++iter)
    {

        // Normalize the input image
        for(int i = 0; i < n_input; ++i)
        {
            for(int j = 0; j < n_input; ++j)
            {
                X[iter][i][j] = divisiond(X_input[iter][i][j]
                    ]-127.5,127.5);
            }
        }

        ...
    }
}
...
}

```

Code 69: *src/artificialneuralnetworks.cpp*

The convolutional neural networks training will be starting, it takes more time compared to FNN training, because we are having a lot of matrices operation, including convolutional operation, pooling operation, forward pass and backpropagation, all requires more cost compared to plain FNN with iris dataset.

We will explain the code that is located in **src/artificialneuralnetworks.cpp** so the reader can comprehend and able to modify it to create their own CNN network, or replicate other famous network like MobileNet or VGG.

In the **include/symintegral/artificialneuralnetworks.h** we will define the class **CNN_LeNet5**, in **private** we will declare the variables that will be used as weights (array), and we declare a **vector<int> vec_uniqueinteger** to store the integers that will be generated randomly to choose the 3 feature maps in S3 layer that connect to any 4 discontiguous feature maps of S2 layer.

```

...

#ifndef CNN_LENET5_H
#define CNN_LENET5_H

class CNN_LeNet5 {
private:

    double c1_weights_kernel[6][5][5]; // first convolutional layer, 6
        feature maps
    double s2_weights_kernel[6];
    double c3_weights_kernel_first6[450]; // second convolutional
        layer, first 6 feature maps

```

```
double c3_weights_kernel_next6[600]; // second convolutional layer
    , next 6 feature maps
double c3_weights_kernel_next3[300]; // second convolutional layer
    , next 3 feature maps
double c3_weights_kernel_last1[150]; // second convolutional layer
    , last 1 feature map
double s4_weights_kernel[16];
double c5_weights_kernel[400][120]; // third convolutional layer,
    120 feature maps, kernel size of 5x5
double f6_weights_kernel[120][84]; // the sixth layer F6
double output_weights[84][10];

double c1_bias[6];
double s2_bias[6];
double c3_bias[16];
double s4_bias[16];
double c5_bias[120];
double f6_bias[84];
double output_bias[10];

vector<int> vec_uniqueinteger;

double learning_rate;

public:
CNN_LeNet5(); // Constructor

vector<double> predict(const vector<vector<double>> &inputs);

void train(vector<vector<vector<int>>> &X, vector<int> &y, int
    epochs);

void save_model(const string &filename) ; // Void function
    declaration
void load_model(const string& filename);
};

#endif

...
```

Code 70: *include/symintegral/artificialneuralnetworks.h*

In the `src/artificialneuralnetworks.cpp` we define the `vector<double> CNN_LeNet5::predict(const vector<vector<double>> &inputs)` function that is used to produce the output vector (activated by softmax) from the input of `vector<vector<double>>` that is a normalized input image with range of $[-1, 1]$.

```
vector<double> CNN_LeNet5::predict(const vector<vector<double>> &inputs)
    // become vector<vector<double>>
```

```

{
    int n_kernelSize = 5;
    int n_classification = 10;
    int pooling_kernelSize = 2;
    int C1_stride = 1, C3_stride = 1, C5_stride = 1;
    int S2_stride = 2, S4_stride = 2;
    int padding = 0;
    vector<vector<vector<double>>> C1_conv, C1_Matrices, S2_Matrices,
        C3_Matrices, S4_Matrices;

    int n_input = inputs.size();
    int r_C1 = (n_input - n_kernelSize + 2*padding)/(C1_stride) + 1 ;
    int c_C1 = r_C1;
    int r_S2 = (r_C1 - pooling_kernelSize)/(S2_stride) + 1;
    int c_S2 = r_S2;
    double alpha = 0.8;

    // First convolutional layer
    for (int k = 0 ; k < 6; ++k)
    {
        vector<vector<double>> c1_weights(n_kernelSize,vector<
            double>(n_kernelSize,0.0));
        for (int i = 0; i < n_kernelSize; ++i)
        {
            for (int j = 0; j < n_kernelSize; ++j)
            {
                c1_weights[i][j] = c1_weights_kernel[k][i][j]
                ;
            }
        }

        vector<vector<double>> C1_result =
            CNN_2DConvolutionOperation(inputs,c1_weights, C1_stride
            );
        C1_conv.push_back(C1_result);
        //cout <<"\ncl k: "<< k << endl;

        for (int i = 0; i < r_C1; ++i)
        {
            for (int j = 0; j < c_C1; ++j)
            {
                if(C1_result[i][i] != 0)
                {
                    C1_result[i][j] += c1_bias[k] ; // add
                    a bias
                }
                C1_result[i][j] = LeakyReLUActivationfunction
                    (C1_result[i][j], alpha); // apply the

```

```
        activation function
    }
}

// The pooling layer S2
vector<vector<double>> S2_result = CNN_2DAveragePooling(
    C1_result, pooling_kernelSize, S2_stride) ; // average
    pooling

for (int i = 0; i < r_S2; ++i)
{
    for (int j = 0; j < c_S2; ++j)
    {
        S2_result[i][j] *= s2_weights_kernel[k] ; //
            multiply with a weight
        if(S2_result[i][j] != 0)
        {
            S2_result[i][j] += s2_bias[k] ; // add
                a bias
        }
        S2_result[i][j] = LeakyReLUActivationfunction
            (S2_result[i][j], alpha); // apply the
            activation function
    }
}

C1_Matrices.push_back(C1_result);
S2_Matrices.push_back(S2_result);
}

// Continuing on the second convolutional layer C3 then S4
vector<vector<int>> contiguous_combinations35 =
    contiguousCombinations(5,3,0);
vector<vector<int>> contiguous_combinations45 =
    contiguousCombinations(5,4,0);
vector<vector<int>> discontinuous_combinations45 =
    discontinuousCombinations(5,4,0);

...
...

vector<double> f6_output;
for (int i = 0; i < 84; ++i)
{
    double f6_sum = 0;

    for (int j = 0; j < 120; ++j)
    {
```

```

        f6_sum += f6_weights_kernel[j][i]*c5_output[j] ;
    }

    f6_output.push_back(f6_sum);
    if(f6_output[i] != 0)
    {
        f6_output[i] += f6_bias[i]; // add a bias
    }
    //f6_output[i] = TanhActivationfunction(f6_output[i]); // apply
    the activation function
    f6_output[i] = LeakyReLUActivationfunction(f6_output[i], alpha);
    // apply the activation function
}

// The output layer in LeNet-5, we use softmax instead of RBF
vector<double> outputs; // to compute the raw logit z_{i}
for (int i = 0; i < n_classification; ++i)
{
    double output_sum = 0;

    for (int j = 0; j < 84; ++j)
    {
        output_sum += output_weights[j][i]*f6_output[j] ;
    }

    outputs.push_back(output_sum);
    outputs[i] += output_bias[i];
}
// apply the activation function
vector<double>predicted_output_softmax =
    SoftMax_vectorresult_activationfunction(outputs);

return predicted_output_softmax;
}

```

Code 71: *src/artificialneuralnetworks.cpp*

It can be read fully and understood by the reader when reading the source code and matching it with the forward pass formula from Table (10.7).

For the training of the convolutional neural networks, we use a function:

void CNN_LeNet5::train(vector<vector<vector<int>>> &X_input, vector<int> &y, int epochs)

The input here is a 3D integer matrix to represent the input images with order that are shuffled first before training (**vector<vector<vector<int>>> &X_input**). At each epoch, each of the input image will be normalized to have the range of $[-1, 1]$.


```
void CNN_LeNet5::train(vector<vector<vector<int>>> &X_input, vector<int>
    &y, int epochs)
{
    cout << BLUE << BOLD << "\nTraining Progress:\n" << RESET << endl;

    int bar_width = 50;
    int n_classification = 10;
    int n_F6 = 84;
    int n_C5 = 120;
    //int n_C2 = 6;
    int n_kernelSize = 5;
    int pooling_kernelSize = 2;
    int C1_stride = 1, C3_stride = 1, C5_stride = 1;
    int S2_stride = 2, S4_stride = 2;
    int padding = 0;

    ...
    ...

    for (int epoch = 0; epoch < epochs; ++epoch)
    {
        double epoch_loss = 0.0;
        for (size_t iter = 0; iter < X_input.size(); ++iter)
        {
            // Normalize the input image
            for(int i = 0; i < n_input; ++i)
            {
                for(int j = 0; j < n_input; ++j)
                {
                    X[iter][i][j] = divisiond(X_input[iter][i][j]
                        ]-127.5,127.5);
                    if (X[iter][i][j] > 0)
                    {
                        X[iter][i][j] = 0;
                    }
                }
            }

            vector<vector<vector<double>>> C1_Matrices, S2_Matrices,
                S2_Matrices_xi, C3_Matrices, S4_Matrices,
                S4_Matrices_xi;
            vector<vector<vector<double>>> C3_Distinct_Kernels;
            vector<double> delta_outputlayer(n_classification,0.0);
            vector<double> outputs = predict(X[iter]);
            vector<double> target(n_classification, 0.0);
            target[y[iter]] = 1.0;
```

```

std::vector<double> errors(n_classification, 0.0);
for (int k = 0; k < n_classification; ++k)
{
    errors[k] = -target[k] * log(outputs[k]); //
    Categorical Cross-Entropy (CCE) Loss
    epoch_loss += errors[k];
}

...

```

Code 72: *src/artificialneuralnetworks.cpp*

We will do the forward pass once again in the train code in order to compute δ for all layers from the output layer back to the C1 layer.

```

...

// We will do forward pass once again here to help computing delta from
// F6 back to C1
// First convolutional layer
for (int k = 0; k < 6; ++k)
{
    vector<vector<double>> C1_weights(n_kernelSize,vector<double>(
        n_kernelSize,0.0));
    for (int i = 0; i < n_kernelSize; ++i)
    {
        for (int j = 0; j < n_kernelSize; ++j)
        {
            C1_weights[i][j] = c1_weights_kernel[k][i][j] ;
        }
    }

    vector<vector<double>> C1_result = CNN_2DConvolutionOperation(X[
        iter],C1_weights, C1_stride);

    for (int i = 0; i < r_C1; ++i)
    {
        for (int j = 0; j < c_C1; ++j)
        {
            if(C1_result[i][j] != 0)
            {
                C1_result[i][j] += c1_bias[k] ; // add a bias
            }
            C1_result[i][j] = LeakyReLUActivationfunction(
                C1_result[i][j], alpha); // apply the activation
            function
        }
    }
}

```

```
// The pooling layer S2

...
}
// Continuing on the second convolutional layer C3 then S4
vector<vector<int>> contiguous_combinations35 = contiguousCombinations
    (5,3,0);
vector<vector<int>> contiguous_combinations45 = contiguousCombinations
    (5,4,0);
vector<vector<int>> discontinuous_combinations45 =
    discontinuousCombinations(5,4,0);

// The first 6: connect to any 3 contiguous feature maps of S2
for (int k = 0 ; k < 6; ++k)
{
    vector<vector<double>> C3_result(r_C3, vector<double>(c_C3,0.0));

    for (int k3 = 0 ; k3 < 3 ; ++k3)
    {
        vector<vector<double>> c3_weights(n_kernelSize,vector<
            double>(n_kernelSize,0.0));
        for (int i = 0; i < n_kernelSize; ++i)
        {
            for (int j = 0; j < n_kernelSize; ++j)
            {
                c3_weights[i][j] = c3_weights_kernel_first6
                    [75*k + 25*k3 + 5*i + j] ; // The first
                    three contiguous feature maps will cover
                    75 neurons, each feature map from S2 will
                    cover 25 neurons
            }
        }

        int n_comb1 = contiguous_combinations35[k][k3];

        vector<vector<double>> S2(r_S2, vector<double>(c_S2, 0.0));
        for (int i = 0; i < r_S2; ++i)
        {
            for (int j = 0; j < c_S2; ++j)
            {
                S2[i][j] = S2_Matrices[n_comb1][i][j];
            }
        }
        vector<vector<double>> C3_result_temp =
            CNN_2DConvolutionOperation(S2,c3_weights, C3_stride);

        for (int i = 0; i < r_C3; ++i)
        {
```

```

        for (int j = 0; j < c_C3; ++j)
        {
            C3_result[i][j] += C3_result_temp[i][j]; //
            sum of the indices combination
        }
    }

    for (int i = 0; i < r_C3; ++i)
    {
        for (int j = 0; j < c_C3; ++j)
        {
            if(C3_result[i][j] != 0)
            {
                C3_result[i][j] += c3_bias[k] ; // add a bias
            }
            C3_result[i][j] = LeakyReLUActivationfunction(
                C3_result[i][j], alpha); // apply the activation
            function
        }
    }
    C3_Matrices.push_back(C3_result);

    vector<vector<double>> S4_result = CNN_2DAveragePooling(C3_result,
        pooling_kernelSize, S4_stride) ; // average pooling
    vector<vector<double>> S4_xi(r_S4, vector<double>(c_S4,0.0));
    for (int i = 0; i < r_S4; ++i)
    {
        for (int j = 0; j < c_S4; ++j)
        {
            S4_xi[i][j] = S4_result[i][j];
            S4_result[i][j] *= s4_weights_kernel[k] ; //
            multiply with a weight
            if(S4_result[i][j] != 0)
            {
                S4_result[i][j] += s4_bias[k] ; // add a bias
            }
            S4_result[i][j] = LeakyReLUActivationfunction(
                S4_result[i][j], alpha); // apply the activation
            function
        }
    }
    S4_Matrices_xi.push_back(S4_xi);
    S4_Matrices.push_back(S4_result);
}

...

```

Code 73: *src/artificialneuralnetworks.cpp*

Afterwards we will compute the δ starting from the output layer, backpropagating to C1 layer. The formula to compute the δ can be seen in Table (10.8).

```
...

for (int k_row = 0; k_row < n_classification; ++k_row)
{
    delta_outputlayer[k_row] = outputs[k_row] - target[k_row];
}

// Delta for F6 layer, output_weights connects the F6 to output layer
vector<double> delta_F6(n_F6, 0.0);
for (int j = 0; j < n_F6; ++j)
{
    for (int i = 0; i < n_classification; ++i)
    {
        delta_F6[j] += delta_outputlayer[i] * output_weights[j][i];
    }
    delta_F6[j] *= LeakyReLUderivative(f6_output[j], alpha);
}
// Delta for C5 layer, f6_weights_kernel connects the C5 to F6 layer
vector<double> delta_C5(n_C5, 0.0);
for (int j = 0; j < n_C5; ++j)
{
    for (int i = 0; i < n_F6; ++i)
    {
        delta_C5[j] += delta_F6[i] * f6_weights_kernel[j][i];
    }
    delta_C5[j] *= LeakyReLUderivative(c5_output[j], alpha);
}
vector<vector<vector<double>>> S4_deltamap;

// Compute delta for S4, 1920 distinct kernels
for (int k = 0; k < 16; ++k) // 16 x 5 x 5
{
    vector<vector<double>> S4_deltamap_k(5, vector<double>(5, 0.0));
    for (int k5 = 0; k5 < 120; ++k5)
    {
        int iter25 = 0;
        vector<vector<double>> c5_weights(5, vector<double>(5, 0.0));
        for (int i = 0; i < n_kernelSize; ++i)
        {
            for (int j = 0; j < n_kernelSize; ++j)
            {
                c5_weights[i][j] = c5_weights_kernel[25*k +
                    iter25][k5] ;
                iter25 += 1;
            }
        }
    }
}
```

```

        vector<vector<double>>> C5_M(1, vector<double>(1, delta_C5[
            k5]));
        vector<vector<double>>> kernel_C5_rotated = rotate180(
            c5_weights); // rotate 180 = transpose weight matrix
                           from C5 to F6 or F6 to output to compute delta
        vector<vector<double>>> delta_C5_padded = CNN_2DpadBorder(
            C5_M,4);

        vector<vector<double>>> fullconv_S4_result =
            CNN_2DConvolutionOperation(delta_C5_padded,
            kernel_C5_rotated, C5_stride);
        for (int i = 0; i < n_kernelSize; ++i)
        {
            for (int j = 0; j < n_kernelSize; ++j)
            {
                S4_deltamap_k[i][j] += fullconv_S4_result[i][
                    j];
            }
        }
        for (int i = 0; i < n_kernelSize; ++i)
        {
            for (int j = 0; j < n_kernelSize; ++j)
            {
                S4_deltamap_k[i][j] *= LeakyReLUderivative(
                    S4_Matrices[k][i][j], alpha);
            }
        }
        S4_deltamap.push_back(S4_deltamap_k);
    }

    vector<double> delta_S4_final(16, 0.0);
    vector<double> delta_S4_final_bias(16, 0.0);
    for (int k = 0; k < 16; ++k)
    {
        for (int i = 0; i < n_kernelSize; ++i)
        {
            for (int j = 0; j < n_kernelSize; ++j)
            {
                delta_S4_final[k] += S4_deltamap[k][i][j] *
                    S4_Matrices_xi[k][i][j];
                delta_S4_final_bias[j] += S4_deltamap[k][i][j];
            }
        }
    }

    vector<vector<vector<double>>>> upsampled_delta_S4 =
        CNN_2DaverageUpsample3DMatrix(S4_deltamap,2,2);

```

```
vector<vector<vector<double>>> delta_C3(16,vector<vector<double>>(10,
    vector<double>(10,0.0)));
for (int k = 0 ; k < 16; ++k)
{
    for (int i = 0; i < 10 ; ++i)
    {
        for (int j = 0; j < 10; ++j)
        {
            //delta_C3[k][i][j] = upsamped_delta_S4[k][i][j] *
            s4_weights_kernel[k] * TanhDerivative(
            C3_Matrices[k][i][j]);
            delta_C3[k][i][j] = upsamped_delta_S4[k][i][j] *
            s4_weights_kernel[k] * LeakyReLUDerivative(
            C3_Matrices[k][i][j], alpha);
        }
    }
}

for (int k = 0 ; k < 6; ++k)
{
    for (int k3 = 0 ; k3 < 3 ; ++k3)
    {
        int n_comb1 = contiguous_combinations35[k][k3];

        vector<vector<double>> S2(r_S2, vector<double>(c_S2, 0.0));
        for (int i = 0; i < r_S2; ++i)
        {
            for (int j = 0; j < c_S2; ++j)
            {
                S2[i][j] = S2_Matrices[n_comb1][i][j];
            }
        }
        vector<vector<double>> delta_C3_matrix(10, vector<double>
            >(10, 0.0));
        for (int i = 0; i < 10; ++i)
        {
            for (int j = 0; j < 10; ++j)
            {
                delta_C3_matrix[i][j] = delta_C3[k][i][j];
            }
        }
        vector<vector<double>> distinct_kernels =
            CNN_2DConvolutionOperation(S2,delta_C3_matrix,
            C3_stride);

        C3_Distinct_Kernels.push_back(distinct_kernels);
    }
}
```

```

}
k1 = 0;
for (int k = 6 ; k < 12; ++k)// The next 24 distinct kernels that connect
    to any 4 contiguous feature maps of S2
{
    for (int k3 = 0 ; k3 < 4 ; ++k3)
    {
        int n_comb1 = contiguous_combinations45[k1][k3];

        vector<vector<double>> S2(r_S2, vector<double>(c_S2, 0.0));
        for (int i = 0; i < r_S2; ++i)
        {
            for (int j = 0; j < c_S2; ++j)
            {
                S2[i][j] = S2_Matrices[n_comb1][i][j];
            }
        }
        vector<vector<double>> delta_C3_matrix(10, vector<double>
            >(10, 0.0));
        for (int i = 0; i < 10; ++i)
        {
            for (int j = 0; j < 10; ++j)
            {
                delta_C3_matrix[i][j] = delta_C3[k][i][j];
            }
        }
        vector<vector<double>> distinct_kernels =
            CNN_2DConvolutionOperation(S2,delta_C3_matrix,
            C3_stride);

        C3_Distinct_Kernels.push_back(distinct_kernels);
    }
    k1 += 1;
}

k1 = 0;
for (int k = 12 ; k < 15; ++k) // The next 12 distinct kernels that
    connect to any 4 discontinuous feature maps of S2
{
    for (int k3 = 0 ; k3 < 4 ; ++k3)
    {
        int n1 = vec_uniqueinteger[k1];

        int n_comb1 = discontinuous_combinations45[n1][k3];
        vector<vector<double>> S2(r_S2, vector<double>(c_S2, 0.0));
        for (int i = 0; i < r_S2; ++i)
        {

```



```
        for (int j = 0; j < c_S2; ++j)
        {
            S2[i][j] = S2_Matrices[n_comb1][i][j];
        }
    }
    vector<vector<double>> delta_C3_matrix(10, vector<double>
        >(10, 0.0));
    for (int i = 0; i < 10; ++i)
    {
        for (int j = 0; j < 10; ++j)
        {
            delta_C3_matrix[i][j] = delta_C3[k][i][j];
        }
    }
    vector<vector<double>> distinct_kernels =
        CNN_2DConvolutionOperation(S2,delta_C3_matrix,
            C3_stride);

    C3_Distinct_Kernels.push_back(distinct_kernels);
}
k1 +=1;
}

for (int k = 15 ; k < 16; ++k) // The last 6 distinct kernels that
    connect to all 6 feature maps in S2
{
    for (int k3 = 0 ; k3 < 6 ; ++k3)
    {
        vector<vector<double>> S2(r_S2, vector<double>(c_S2, 0.0));
        for (int i = 0; i < r_S2; ++i)
        {
            for (int j = 0; j < c_S2; ++j)
            {
                S2[i][j] = S2_Matrices[k3][i][j];
            }
        }
        vector<vector<double>> delta_C3_matrix(10, vector<double>
            >(10, 0.0));
        for (int i = 0; i < 10; ++i)
        {
            for (int j = 0; j < 10; ++j)
            {
                delta_C3_matrix[i][j] = delta_C3[k][i][j];
            }
        }
        vector<vector<double>> distinct_kernels =
            CNN_2DConvolutionOperation(S2,delta_C3_matrix,
                C3_stride);
```

```

        C3_Distinct_Kernels.push_back(distinct_kernels);
    }
}
...

```

Code 74: *src/artificialneuralnetworks.cpp*

In the C3 layer, we have to compute 1600 δ for the 16 feature maps of size 10×10 then we will use the deltas to compute the weight update for the 1,500 trainable weights and 16 trainable biases in C3 layer. The 1,500 trainable weights can be updated if we know the vector gradients for the corresponding trainable weights, thus we will have to compute 60 distinct kernels that corresponds to the 16 feature maps in C3 layer: the first 18 distinct kernels connect to any 3 contiguous feature maps of S2 layer (the first 6 feature maps in C3), then the next 24 distinct kernels connect to any 4 contiguous feature maps in S2 layer (the next 6 feature maps in C3), then the next 12 distinct kernels connect to any 4 discontiguous feature maps in S2 layer (the next 3 feature maps in C3), finally the last 6 distinct kernels connect to all feature maps in S2 (the last feature map in C3).

We store 60 distinct kernels that will be responsible for the weight update for C3 layer in a 3D matrix `vector<vector<vector<double>>>C3_Distinct_Kernels`.

- `c3_weights_kernel_first6[450]` to represent the weights for the first 6 feature maps will use the first 18 distinct kernels for the weight update.
- `c3_weights_kernel_next6[600]` to represent the weights for the next 6 feature maps will use the next 24 distinct kernels for the weight update.
- `c3_weights_kernel_next3[300]` to represent the weights for the next 3 feature maps will use the next 12 distinct kernels for the weight update.
- `c3_weights_kernel_last1[150]` to represent the weights for the last feature map will use the last 6 distinct kernels for the weight update.

To compute the δ for the S2 layer it is as complex as the C3 layer since it is partially connected thus we need longer code. The formula to compute the δ^{S2} can be seen in table (10.8) in which we have to rotate weight matrix from S2 to C3 180° and perform full convolution with the delta matrix (to perform full convolution: pad the delta matrix of C3 layer of size 10×10 with 4 rows and columns of zeros on all sides making it 18×18 , afterwards do the convolution with the rotated weight matrix).

First we will declare / create 3D matrices to represent the delta map for S2 from 1 to 6, since we are using sparse connectivity then we perform the full convolution between the padded δ^{C3} matrix and the rotated weight matrix from S2 to C3 from the first 6 feature maps in C3, to the next 6 feature maps in C3, to the next 3 feature maps in C3, and the last feature map in C3.

Each time we perform convolution operation we know which feature map in S2 that is used from the functions `contiguous_combinations35` `contiguous_combinations45`, `discontiguous_combinations45`, thus we create `if.. else if..` conditional to store the result of the full convolution operation to the 3D matrix `vector<vector<vector<double>>>`

S2_deltamap_i ($i = 1, 2, \dots, 6$) that represents the feature map in S2 layer that is used for the current full convolution operation.

In the end, we will obtain 60 results from the full convolution operation between the rotated weight matrix from S2 to C3 and the padded δ^{C3} , each of these 60 matrices will represent feature map of S2, each could belong to feature map 1,2,3,4,5, or 6. After we already have them clustered from the operations above, then we will use a class **Matrix3Dto2D** to sum over the depth in 3D matrix at index $[i][j]$ into a 2D matrix. These 2D matrices are the delta map (1,176 δ^{S2} that represents $6 \times 14 \times 14$ feature maps in S2) that will be used to compute the weight gradients for S2 weights update.

```
...

vector<vector<vector<double>>> S2_deltamap_1, S2_deltamap_2,
    S2_deltamap_3, S2_deltamap_4, S2_deltamap_5, S2_deltamap_6; // Store
    delta map for S2 with sparse connectivity
// Computation for delta map for S2 layer:
// The first 6 feature maps in S3 that connect to any 3 contiguous
    feature maps of S2
for (int k = 0 ; k < 6; ++k)
{
    for (int k3 = 0 ; k3 < 3 ; ++k3)
    {
        vector<vector<double>> delta_C3_matrix(10, vector<double>
            >(10, 0.0));
        vector<vector<double>> S2_C3_kernelmatrix(5, vector<double>
            >(5, 0.0));
        for (int i = 0; i < 10; ++i)
        {
            for (int j = 0; j < 10; ++j)
            {
                delta_C3_matrix[i][j] = delta_C3[k][i][j];
            }
        }
        vector<vector<double>> padded_delta = CNN_2DpadBorder(
            delta_C3_matrix,4);
        for (int i = 0; i < 5; ++i)
        {
            for (int j = 0; j < 5; ++j)
            {
                S2_C3_kernelmatrix[i][j] =
                    c3_weights_kernel_first6[75*k + 25*k3 +
                    5*i + j] ;
            }
        }
        vector<vector<double>> rotated_W = rotate180(
            S2_C3_kernelmatrix);

        int n_comb1 = contiguous_combinations35[k][k3];
```

```

        vector<vector<double>> deltamap_kernel =
            CNN_2DConvolutionOperation(padded_delta, rotated_W,
            C3_stride);

        if(n_comb1 == 0)
        {
            S2_deltamap_1.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 1)
        {
            S2_deltamap_2.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 2)
        {
            S2_deltamap_3.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 3)
        {
            S2_deltamap_4.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 4)
        {
            S2_deltamap_5.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 5)
        {
            S2_deltamap_6.push_back(deltamap_kernel);
        }
    }
}

// The next 6 feature maps in S3 that connect to any 4 contiguous feature
// maps of S2
k1 = 0;
for (int k = 6 ; k < 12; ++k)
{
    for (int k3 = 0 ; k3 < 4 ; ++k3)
    {
        vector<vector<double>> delta_C3_matrix(10, vector<double>
            >(10, 0.0));
        vector<vector<double>> S2_C3_kernelmatrix(5, vector<double>
            >(5, 0.0));
        for (int i = 0; i < 10; ++i)
        {
            for (int j = 0; j < 10; ++j)
            {

```

```
        delta_C3_matrix[i][j] = delta_C3[k][i][j];
    }
}
vector<vector<double>> padded_delta = CNN_2DpadBorder(
    delta_C3_matrix,4);
for (int i = 0; i < 5; ++i)
{
    for (int j = 0; j < 5; ++j)
    {
        S2_C3_kernelmatrix[i][j] =
            c3_weights_kernel_next6[100*k1 +25*k3 +
                5*i + j] ;
    }
}
vector<vector<double>> rotated_W = rotate180(
    S2_C3_kernelmatrix);

int n_comb1 = contiguous_combinations45[k1][k3];

vector<vector<double>> deltamap_kernel =
    CNN_2DConvolutionOperation(padded_delta, rotated_W,
        C3_stride);

if(n_comb1 == 0)
{
    S2_deltamap_1.push_back(deltamap_kernel);
}
else if(n_comb1 == 1)
{
    S2_deltamap_2.push_back(deltamap_kernel);
}
else if(n_comb1 == 2)
{
    S2_deltamap_3.push_back(deltamap_kernel);
}
else if(n_comb1 == 3)
{
    S2_deltamap_4.push_back(deltamap_kernel);
}
else if(n_comb1 == 4)
{
    S2_deltamap_5.push_back(deltamap_kernel);
}
else if(n_comb1 == 5)
{
    S2_deltamap_6.push_back(deltamap_kernel);
}
}
```

```

        k1 += 1;
    }

    // The next 3 feature maps in S3 that connect to any 4 discontiguous
    // feature maps of S2
    k1 = 0;
    for (int k = 12 ; k < 15; ++k)
    {
        for (int k3 = 0 ; k3 < 4 ; ++k3)
        {
            vector<vector<double>> delta_C3_matrix(10, vector<double>
                >(10, 0.0));
            vector<vector<double>> S2_C3_kernelmatrix(5, vector<double>
                >(5, 0.0));
            for (int i = 0; i < 10; ++i)
            {
                for (int j = 0; j < 10; ++j)
                {
                    delta_C3_matrix[i][j] = delta_C3[k][i][j];
                }
            }
            vector<vector<double>> padded_delta = CNN_2DpadBorder(
                delta_C3_matrix,4);
            for (int i = 0; i < 5; ++i)
            {
                for (int j = 0; j < 5; ++j)
                {
                    S2_C3_kernelmatrix[i][j] =
                        c3_weights_kernel_next3[100*k1 + 25*k3+
                            5*i + j] ;
                }
            }
            vector<vector<double>> rotated_W = rotate180(
                S2_C3_kernelmatrix); // rotate 180 = transpose weight
            // matrix from C5 to F6 or F6 to output to compute delta
            int n1 = vec_uniqueinteger[k1];

            int n_comb1 = discontiguous_combinations45[n1][k3];

            vector<vector<double>> deltamap_kernel =
                CNN_2DConvolutionOperation(padded_delta, rotated_W,
                    C3_stride);

            if(n_comb1 == 0)
            {
                S2_deltamap_1.push_back(deltamap_kernel);
            }
            else if(n_comb1 == 1)

```

```
        {
            S2_deltamap_2.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 2)
        {
            S2_deltamap_3.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 3)
        {
            S2_deltamap_4.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 4)
        {
            S2_deltamap_5.push_back(deltamap_kernel);
        }
        else if(n_comb1 == 5)
        {
            S2_deltamap_6.push_back(deltamap_kernel);
        }
    }
    k1 += 1;
}

// The last feature map in S3 that connect to all 6 discontiguous feature
maps of S2
for (int k = 15 ; k < 16; ++k)
{
    for (int k3 = 0 ; k3 < 6 ; ++k3)
    {
        vector<vector<double>> delta_C3_matrix(10, vector<double>
            >(10, 0.0));
        vector<vector<double>> S2_C3_kernelmatrix(5, vector<double>
            >(5, 0.0));
        for (int i = 0; i < 10; ++i)
        {
            for (int j = 0; j < 10; ++j)
            {
                delta_C3_matrix[i][j] = delta_C3[k][i][j];
            }
        }
        vector<vector<double>> padded_delta = CNN_2DpadBorder(
            delta_C3_matrix,4);
        for (int i = 0; i < 5; ++i)
        {
            for (int j = 0; j < 5; ++j)
            {
                S2_C3_kernelmatrix[i][j] =
                    c3_weights_kernel_last1[25*k3+ 5*i + j] ;
            }
        }
    }
}
```

```

    }
}
vector<vector<double>> rotated_W = rotate180(
    S2_C3_kernelmatrix); // rotate 180 = transpose weight
                           matrix from C5 to F6 or F6 to output to compute delta

vector<vector<double>> deltamap_kernel =
    CNN_2DConvolutionOperation(padded_delta, rotated_W,
    C3_stride);

if(k3 == 0)
{
    S2_deltamap_1.push_back(deltamap_kernel);
}
else if(k3 == 1)
{
    S2_deltamap_2.push_back(deltamap_kernel);
}
else if(k3 == 2)
{
    S2_deltamap_3.push_back(deltamap_kernel);
}
else if(k3 == 3)
{
    S2_deltamap_4.push_back(deltamap_kernel);
}
else if(k3 == 4)
{
    S2_deltamap_5.push_back(deltamap_kernel);
}
else if(k3 == 5)
{
    S2_deltamap_6.push_back(deltamap_kernel);
}
}
}

int n_S2_map1 = S2_deltamap_1.size();
int n_S2_map2 = S2_deltamap_2.size();
int n_S2_map3 = S2_deltamap_3.size();
int n_S2_map4 = S2_deltamap_4.size();
int n_S2_map5 = S2_deltamap_5.size();
int n_S2_map6 = S2_deltamap_6.size();

// Sum over the depth at [i][j] into a 2D matrix with class Matrix3DTo2D
// 14 rows, 14 columns, n_S2_map_k depth layers
Matrix3DTo2D S2_deltamap_1_final(14, 14, n_S2_map1);
Matrix3DTo2D S2_deltamap_2_final(14, 14, n_S2_map2);

```



```
Matrix3DTo2D S2_deltamap_3_final(14, 14, n_S2_map3);
Matrix3DTo2D S2_deltamap_4_final(14, 14, n_S2_map4);
Matrix3DTo2D S2_deltamap_5_final(14, 14, n_S2_map5);
Matrix3DTo2D S2_deltamap_6_final(14, 14, n_S2_map6);

// To compute the 1,176 deltas for S2 layer (6 x 14 x 14)
for (int k = 0; k < n_S2_map1; ++k)
{
    for (int i = 0; i < 14; ++i)
    {
        for (int j = 0; j < 14; ++j)
        {
            double index = S2_deltamap_1[k][i][j];
            S2_deltamap_1_final.set(i, j, k, index);
        }
    }
}
for (int k = 0; k < n_S2_map2; ++k)
{
    for (int i = 0; i < 14; ++i)
    {
        for (int j = 0; j < 14; ++j)
        {
            double index = S2_deltamap_2[k][i][j];
            S2_deltamap_2_final.set(i, j, k, index);
        }
    }
}
for (int k = 0; k < n_S2_map3; ++k)
{
    for (int i = 0; i < 14; ++i)
    {
        for (int j = 0; j < 14; ++j)
        {
            double index = S2_deltamap_3[k][i][j];
            S2_deltamap_3_final.set(i, j, k, index);
        }
    }
}
for (int k = 0; k < n_S2_map4; ++k)
{
    for (int i = 0; i < 14; ++i)
    {
        for (int j = 0; j < 14; ++j)
        {
            double index = S2_deltamap_4[k][i][j];
            S2_deltamap_4_final.set(i, j, k, index);
        }
    }
}
```

```

    }
}
for (int k = 0; k < n_S2_map5; ++k)
{
    for (int i = 0; i < 14; ++i)
    {
        for (int j = 0; j < 14; ++j)
        {
            double index = S2_deltamap_5[k][i][j];
            S2_deltamap_5_final.set(i, j, k, index);
        }
    }
}
for (int k = 0; k < n_S2_map6; ++k)
{
    for (int i = 0; i < 14; ++i)
    {
        for (int j = 0; j < 14; ++j)
        {
            double index = S2_deltamap_6[k][i][j];
            S2_deltamap_6_final.set(i, j, k, index);
        }
    }
}
vector<double> S2_Delta_Final1 = S2_deltamap_1_final.collapseTo2D();
vector<double> S2_Delta_Final2 = S2_deltamap_2_final.collapseTo2D();
vector<double> S2_Delta_Final3 = S2_deltamap_3_final.collapseTo2D();
vector<double> S2_Delta_Final4 = S2_deltamap_4_final.collapseTo2D();
vector<double> S2_Delta_Final5 = S2_deltamap_5_final.collapseTo2D();
vector<double> S2_Delta_Final6 = S2_deltamap_6_final.collapseTo2D();
vector<vector<vector<double>>> S2_deltamap(6,vector<vector<double>>(14,
    vector<double>(14,0.0)));
for (int i = 0; i < 14; ++i)
{
    for (int j = 0; j < 14; ++j)
    {
        S2_deltamap[0][i][j] = S2_Delta_Final1[i*14 + j];
        S2_deltamap[1][i][j] = S2_Delta_Final2[i*14 + j];
        S2_deltamap[2][i][j] = S2_Delta_Final3[i*14 + j];
        S2_deltamap[3][i][j] = S2_Delta_Final4[i*14 + j];
        S2_deltamap[4][i][j] = S2_Delta_Final5[i*14 + j];
        S2_deltamap[5][i][j] = S2_Delta_Final6[i*14 + j];
    }
}

// to compute the delta for each feature map in S2 layer
vector<double> delta_S2_final(6, 0.0);
vector<double> delta_S2_final_bias(6, 0.0);

```

```
for (int k = 0; k < 6; ++k)
{
    double delta_sum = 0;
    for (int i = 0; i < 14; ++i)
    {
        for (int j = 0; j < 14; ++j)
        {
            delta_S2_final[k] += S2_deltamap[k][i][j] *
                S2_Matrices_xi[k][i][j] * LeakyReLUDerivative(
                    S2_Matrices[k][i][j], alpha);
            delta_sum += S2_deltamap[k][i][j];
        }
    }
    delta_S2_final_bias[k] = delta_sum;
}
```

Code 75: *src/artificialneuralnetworks.cpp*

Finally, we will compute the δ^{C1} , we follow the formula from Table (10.8) and we will obtain 4,704 δ^{C1} that represents $6 \times 28 \times 28$ feature maps in C1. The weight update for C1 is performing valid convolution between the input image (that is normalized) and the δ_k^{C1} with $k = 1, 2, \dots, 6$.

```
...

// To Compute the delta for each feature map in C1 layer
vector<vector<vector<double>>> upsampled_delta_S2 =
    CNN_2DaverageUpsample3DMatrix(S2_deltamap, 2, 2);

vector<vector<vector<double>>> delta_C1(6, vector<vector<double>>(28,
    vector<double>(28, 0.0)));
for (int k = 0; k < 6; ++k)
{
    for (int i = 0; i < 28; ++i)
    {
        for (int j = 0; j < 28; ++j)
        {
            //delta_C1[k][i][j] = upsampled_delta_S2[k][i][j] *
                s2_weights_kernel[k] * TanhDerivative(
                    C1_Matrices[k][i][j]);
            delta_C1[k][i][j] = upsampled_delta_S2[k][i][j] *
                s2_weights_kernel[k] * LeakyReLUDerivative(
                    C1_Matrices[k][i][j], alpha);
        }
    }
}

// Now we have obtained 4704 deltas in each neuron in 6x28x28 feature map
for C1 layer.
vector<vector<vector<double>>> C1_weights_update;
for (int k = 0; k < 6; ++k)
```

```

{
    vector<vector<double>> weights_update = CNN_2DConvolutionOperation
        (X[iter],delta_C1[k], C1_stride);
    C1_weights_update.push_back(weights_update);
}

...

```

Code 76: *src/artificialneuralnetworks.cpp*

Adaptive Moment Estimation (Adam) Optimizer An alternative in neural networks besides using steepest descent / gradient descent method is using Adam optimizer. The Adam optimizer is a highly efficient, first-order gradient-based optimization algorithm used to train deep learning models. It dynamically adjusts the learning rate for each network weight individually by combining the principles of Momentum and RMSprop. Adam is favored as the default optimizer for many machine learning and deep learning tasks because it requires minimal tuning. Default values typically perform well across a wide variety of datasets. To help LeNet-5 converge faster and avoid getting stuck in local minima we will use Adam optimizer. We will apply Adam optimizer to the weights of the feature maps in the convolutional layers (C1, C3, and C5), and the fully connected layers (F6 and output) and their respective biases. We will maintain separate, adaptive learning rates for each layer's parameters.

Adam optimizer is applied after the backpropagation phase calculates the gradients of the loss function with respect to the network's weights and biases. Adam combines Momentum (first moment) and RMSProp (second moment) to dynamically adjust the learning rate for each specific weight and bias in the convolutional and fully connected layers.

During backpropagation at time step t , we first compute the gradient of the loss L with respect to a specific parameter (weight w , or bias b), represented here as $g_t = \nabla_{\theta} L_t$.

(a) **Calculate the First and Second Moments**

Adam calculates the exponential moving averages of the gradient (first moment) and the squared gradient (second moment).

The first moment vector (mean / momentum) tracks the average of past gradients / acts like a moving average of the past gradients to maintain velocity, it provides direction and smooths out the path of descent, providing "momentum" to speed up training:

$$m_t = \beta_1 m_{t-1} + (1 - \beta_1) g_t$$

The second moment vector (uncentered variance) acts like RMSProp, calculating the exponential moving average of the squared gradients to scale the learning rate, increasing it (taking larger steps) for sparse features (noisy gradients) and decreasing it (more cautious steps) for frequent ones (complex areas):

$$v_t = \beta_2 v_{t-1} + (1 - \beta_2) g_t^2$$

Where:

- β_1 is the exponential decay rate for the first moment (usually 0.9).
- β_2 is the exponential decay rate for the second moment (usually 0.999).
- g_t is the gradient at time t .
- m_t is the first moment vector.
- v_t is the second moment vector.

(b) Apply Bias Correction

Because m_t and v_t are initialized to zero, they are biased toward zero early in training. Bias correction is used to adjust for this.

Bias-corrected first moment:

$$\hat{m}_t = \frac{m_t}{1 - \beta_1^t}$$

Bias-corrected second moment:

$$\hat{v}_t = \frac{v_t}{1 - \beta_2^t}$$

(c) Update the Parameters

Finally, the weights and biases of the LeNet-5 network are updated using the bias-corrected moments.

$$\theta_t = \theta_{t-1} - \frac{\eta}{\sqrt{\hat{v}_t} + \epsilon} \hat{m}_t$$

Where:

- θ represents the parameter being updated (a weight w or a bias b).
- η is the learning rate.
- ϵ is a very small number added to prevent division by zero (e.g., $\epsilon 10^{-8}$)

(d) Update Exponential Decay Rates (β_1 and β_2)

In standard Adam, β_1 and β_2 are static hyperparameters. However, for specialized training (like fine-tuning a LeNet-5), you can update or decay them over time using the following techniques:

- **Exponential Decay (for Schedule):** You can decay the decay rate itself over time to narrow down the search space as training converges. For instance, at each epoch t , you can update β_1 as:

$$\beta_1^t = \beta_1^0 \times \lambda^t$$

and update β_2 as:

$$\beta_2^t = \beta_2^0 \times \lambda^t$$

with λ as a decay rate constant (e.g., $\lambda = 0.99$), and t is the epoch number.

- **Decay Rate for the Learning Rate**

If you wish to decay the learning rate instead to prevent oscillation near the minimum, you can dynamically adjust η per epoch:

$$\eta_t = \frac{\eta_0}{\sqrt{t}}$$

Each weight in the network gets its own specific learning rate, which is incredibly useful for processing sparse data. Adam corrects for early-stage biases during the initial

iterations to ensure precise gradient approximations. It requires very little memory and works incredibly well on a wide range of architectures, including convolutional neural networks and recurrent neural networks.

Adam optimizer can also get stuck in local minima primarily because its learning rate diminishes too quickly, its momentum becomes trapped by directional shifts, or the batch size removes the necessary exploration noise. While Adam combines momentum and adaptive learning rates to escape shallow traps, specific mathematical dynamics can lock it into suboptimal basins.

(a) **Bad Weight Initialization**

Adam may converge to the nearest poor local minimum.

The Mechanism: Poor weights initialization will place the starting weights far from optima. Zero-initialization breaks symmetry and stalls learning entirely.

(b) **Vanishing Learning Rate via the Second Moment (v_t)**

Adam tracks the moving average of squared gradients (v_t) to scale updates for each parameter.

The Mechanism: In a deep or wide local minimum, gradients drop significantly.

The Trap: The historical accumulation of v_t forces individual parameter learning rates to shrink exponentially. This leaves the optimizer with too little "energy" to step back up the slope and escape.

(c) **Situational Oscillations and Momentum Cancellation**

Adam utilizes a first-moment vector (m_t) to act like physics-based momentum.

The Mechanism: When entering a local minimum, the optimizer relies on past inertia to carry it through.

The Trap: If the true loss landscape requires a sharp change in direction to find the exit, the calculated step stalls. The opposing forces of historical momentum and the new uphill gradient can cancel each other out, causing Adam to oscillate back and forth at the basin floor until it settles there completely.

(d) **Too Large of a Batch Size (Lack of Stochastic Noise)**

Adam is heavily reliant on stochasticity (randomness) to jump out of local wells.

The Mechanism: Small batch training creates a high-variance, "noisy" loss landscape that naturally bounces the optimizer out of narrow local traps.

The Trap: If your batch size is too large, the loss landscape becomes incredibly smooth and deterministic. Without random perturbations, Adam follows the exact mathematical gradient straight to the bottom of the nearest local minimum and freezes.

(e) **Poor Learning Rate Scheduling**

The Mechanism: Models often start with a high learning rate to explore the landscape and lower it over time to fine-tune.

The Trap: If the initial learning rate (η) is set too low, Adam never develops the speed required to clear initial shallow valleys. Conversely, if the learning rate schedule decays too aggressively, it cuts off Adam's stepping power before it can escape early traps.

(f) **Saturated Activation Functions**

Original LeNet-5 uses sigmoid or tanh activation functions.

The Mechanism: Deep layers can cause vanishing gradients problem.

The Trap: The weights stop updating because gradients approach zero.

(g) **Small Batch Sizes**

Tiny batches introduce extreme gradient noise.

The Mechanism: The Adam optimizer moves erratically across the loss landscape.

The Trap: It can get trapped in sharp, sub-optimal valleys.

(h) **Loss Landscape of LeNet-5**

LeNet-5 has 7 layers in total, it is a shallow, classic architecture.

The Mechanism: It processes an image through 3 convolutional layers and 2 pooling layers then a fully dense connected layer and an output.

The Trap: Shallow networks like LeNet-5 have more high-loss local minima.

How to help Adam escape local minima traps:

(a) **Use Activation Function that Fix Vanishing Gradients Problem:** Replace sigmoid or tanh to ReLU or Leaky ReLU activation functions. The original LeNet-5 uses tanh, which suffer heavily from vanishing gradients. If using standard ReLU, then dead neurons could still stall the updates. Swap out ReLU for LeakyReLU($\alpha = 0.01$) to ensure backpropagation always maintains a small, healthy gradient flow.

(b) **Implement Batch Normalization:** LeNet-5 was originally built without Batch Normalization (BN), but modern implementations benefit greatly from it. Without it, vanishing or exploding internal covariate shifts can stall the gradients. Insert a BN layer on the input image that still have value in the range of $[0, 255]$ to become $[-1, 1]$.

(c) **Increase Batch Size:** Increase the number of input images processed together in a batch during training to 64 or 128.

(d) **Implement a Learning Rate Warmup and Decay Schedule:** Start with a learning rate then scale it down under certain conditions to prevent v_t from locking up prematurely. High learning rate cause Adam to overshoot, thus making the algorithm skips over the global minimum.

In LeNet-5 particularly with MNIST dataset, we can tune the learning rate within these ranges:

- **0.001 (Recommended baseline):** The default learning rate for Adam optimizes in major architecture. It provides the best balance between fast convergence and stable training.

Adam's default learning rate (0.001) can sometimes be too large for a simple architecture like LeNet-5, causing the weights to oscillate back and forth over a narrow minimum valley. To fix this drop the learning rate to 0.0001 or 0.0005.

- **0.0005 to 0.0001 (Conservative / Fine-Tuning):** Use this if your loss curve oscillates wildly or if you notice overfitting happening too early.
- **0.002 to 0.005 (Aggressive):** Use this only if you are using large batch sizes (e.g., 256 or higher) and want to speed up training.

For LeNet-5 a learning rate of 0.01 or higher will cause Adam to overshoot, resulting in unstable training or failing to converge entirely.

(e) **Reduce Batch Size:** Force more stochastic noise into the training routine to shake the optimizer out of shallow basins.

(f) **Switch to Cosine Annealing with Restarts:** Periodically spike the learning rate back up during training to forcefully push the model out of local minima.

(g) **Use He/Xavier Initialization:** To initialize weights that is near the optima. **Use a Modern Architecture:** If accuracy plateaus, try to use modern CNN architecture such as ResNet, GoogleNet, EfficientNet, DenseNet, or ConvNeXt.

When LeNet-5 model that is using Adam optimizer hits a stagnant training loss, it means the parameter updates have dropped too low to escape a local minimum, a saddle point, or a flat plateau. The stagnated loss in LeNet-5 with Adam is primarily caused by uncalibrated learning rates or vanishing gradient trajectories.

5. Results

The hyperparameters for the implementation of LeNet-5 architecture for digit recognition are as follows:

- Data = 100 images
Training data 60%, testing data 100%
- Epochs = 20
- Learning rate

$$\eta = \begin{cases} 0.1, & \text{if } i = 1 \\ 0.06, & \text{if } \text{Loss}[i] > 2 \text{ \& } |\text{Loss}[i] - \text{Loss}[i-1]| < 0.1 \\ 0.025, & \text{if } 1 \leq \text{Loss}[i] < 2 \\ 0.01, & \text{if } \text{Loss}[i] < 1 \end{cases}$$

with $i = 1, 2, \dots, 20$. We start with $\eta = 0.1$ and after the first epoch if the loss difference is less than 0.1 then η will be 0.06 and keep decreasing as the loss decreases at certain point. This scheme of learning rate decreasing as the loss decreases really helps and it still depends on the initial weights that are initialized because it can sometimes overshoot and the loss become very high. If the loss become high, we reset and train again.




Figure 10.49: The training loss of LeNet-5 as plotted over 20 epochs along with the learning rate at each epoch. The training loss is monotonically decreasing indicating the network is effectively learning to differentiate between the ten classes in the dataset (*Hamzstplot/Examples with Makefile/Neural Networks/Plot 2D Line for Loss and Learning rate for CNN/main.cpp*).

Here are two results of training the CNN with LeNet-5 architecture:

```
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Neural Networks/Test CNN using Class with Steepest Descent ]# ./
===== Convolutional Neural Network =====
Loading dataset...
Successfully loaded 100 images!
Dimension of Image : 32x32
DEBUG: Number of Labels: 100
DEBUG: Label names:
DEBUG: - 0
DEBUG: - 1
DEBUG: - 2
DEBUG: - 3
DEBUG: - 4
DEBUG: - 5
DEBUG: - 6
DEBUG: - 7
DEBUG: - 8
DEBUG: - 9
DEBUG: Dataset loaded. Size: 100 images.
DEBUG: Number of labels: 100
DEBUG: Train set size: 60
DEBUG: Test set size: 40

Initializing Convolutional Neural Network with LeNet5 Architecture for Digit Recognition...

Starting Training Process

Training Progress:

[=====] 100% Epoch 20/20 - Loss: 0.6317 - Learning Rate: 0.025000000
```

Figure 10.50: Training the convolutional neural networks with LeNet-5 architecture for handwritten digit recognition, we are using configuration in Table (10.6), all activation functions are Leaky ReLU ($\alpha = 0.8$) and the output layer is using softmax. The whole data is 100 images, we train 60% of the data, we use 20 epochs and it took 1093 seconds till it finishes, we haven't optimize or make the code faster and more efficient by replacing some vectors and matrices operations with either *class* or *struct* (*SymIntegration/Examples/Neural Networks/Test CNN using Class with Steepest Descent/main.cpp*).

```
DEBUG: - 7
DEBUG: - 8
DEBUG: - 9
DEBUG: Dataset loaded. Size: 100 images.
DEBUG: Number of labels: 100
DEBUG: Train set size: 60
DEBUG: Test set size: 40

Initializing Convolutional Neural Network with LeNet5 Architecture for Digit Recognition...

Starting Training Process

Training Progress:

[=====] 100% Epoch 20/20 - Loss: 0.6317 - Learning Rate: 0.025000000

Training completed in 1093.48092652 seconds.

Evaluating the model on the test set:
[=====] 100%

Accuracy: 73.00%

Confusion Matrix:

Predicted >
0      0      1      2      3      4      5      6      7      8      9
0      8      0      0      0      1      0      0      0      1      0
1      0      9      0      0      1      0      0      0      0      0
2      0      0      5      1      2      0      1      1      0      0
3      0      0      0      7      1      0      0      1      1      0
4      0      0      0      0      6      0      1      1      0      2
5      0      0      0      1      0      9      0      0      0      0
6      0      0      0      0      1      2      6      0      1      0
7      0      0      0      0      1      0      0      9      0      0
8      0      0      0      0      0      0      1      0      8      1
9      0      0      0      1      1      0      1      0      1      6

Precision, Recall, and F1-score:
```

Figure 10.51: The accuracy after training the convolutional neural networks with LeNet-5 architecture for handwritten digit recognition is 73%, the confusion matrix is also generated after the training is finished, the testing data for confusion matrix is the whole 100 images of handwritten digit (*SymIntegration/Examples/Neural Networks/Test CNN using Class with Steepest Descent/main.cpp*).

```

Initializing Convolutional Neural Network with LeNet5 Architecture for Digit Recognition...
Starting Training Process
Training Progress:
[=====] 100% Epoch 20/20 - Loss: 0.1597 - Learning Rate: 0.02500000
Training completed in 589.67249413 seconds.
Evaluating the model on the test set:
[=====] 100%
Accuracy: 76.00%
Confusion Matrix:
Predicted >
0          0          1          2          3          4          5          6          7          8          9
0          9          1          0          0          0          0          0          0          0          0
1          0         10          0          0          0          0          0          0          0          0
2          0          1          5          1          2          1          0          0          0          0
3          0          0          0          8          1          0          1          0          0          0
4          0          0          1          0          5          0          0          2          2          0
5          0          0          0          0          0          9          0          0          1          0
6          0          0          0          0          1          0          9          0          0          0
7          0          0          0          0          0          0          0          10         0          0
8          0          0          0          0          0          0          0          0          9          0
9          0          1          0          0          2          3          0          0          2          2

Precision, Recall, and F1-score:
0:
Precision: 1.00
Recall:    0.90
F1-score: 0.95

```

Figure 10.52: We try to perform another training for the convolutional neural networks with LeNet-5 architecture for handwritten digit recognition, we obtain the accuracy of 76 %. The digit 9 is wrongly predicted as 4, some is predicted as 5 or as 8, we can see this happens because the shape is almost similar. Confusion matrix plotting is another way of model evaluation. It is used for graphical representation of performance of the model. It shows the performance of model in predicting every class (digit 0 to 9) (*SymIntegration/Examples/Neural Networks/Test CNN using Class with Steepest Descent/main.cpp*).

In a convolutional neural network, accuracy measures how often the model is correct, while loss measures the penalty for incorrect predictions. We use the optimizer like Adam to minimize the loss during training.

We have obtained accuracy of 73% and 76%, when we searched and read on internet and paper, they claim to have 98% of accuracy or even 99%, it is quite misleading. An accuracy of 70-80% on unseen data is almost always better than hitting almost 100%, because the latter usually indicates overfitting, meaning the model has simply memorized the training data rather than actually learning to recognize patterns.

When a CNN achieves near 100% training accuracy but fails in the real world, it has usually memorized the specific noise, lighting conditions, or artifacts of the training dataset. When presented with new, unseen data, it struggles to adapt. While 70-80% accuracy indicates the model has learned general, underlying patterns and can successfully generalize to new situations. Pushing the accuracy to 99% or 100% requires training the model on edge cases, outliers, and dataset anomalies, which actively hurts its ability to perform well on standard, everyday inputs. Rather than obsessing over a raw accuracy percentage, it is often more important to analyze a confusion matrix to understand where the model is making mistakes.

VIII. Deep Learning: Convolutional Neural Network

[illegible]

Figure 10.53: The $C1$ matrix that represents the first feature map of $C1$ at $i = 1$.

Input_images.txt	S2_matrix.txt	shuffledytrain.txt
1 i = 1		
2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		
3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		
4 0 0 0 0 0 0 -0.640991 -0.67187 -0.859323 -0.973114 -0.856043 -0.689521 0		
5 0 0 0 0 0 -0.656539 -0.790359 -0.950822 -0.815535 -0.977147 -1.0433 -0.728682 0		
6 0 0 0 0 0 -0.789662 -0.976421 -0.721181 -0.551489 -1.07007 -1.10696 -0.50344 0		
7 0 0 0 0 0 -0.606191 -0.882354 -0.849124 -1.05706 -1.04658 -0.749792 -0.487221 0		
8 0.504319 0.504319 0.504319 0.504319 0.504319 0.504319 0.705344 0.571616 0.751127 0.835718 0.428994 0.48441 0.606927 0.504319		
9 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.110857 -0.0490013 -0.0423389 -0.0532883 -0.371561 -0.209952 -0.0780112 -0.116935		
10 0 0 0 0 0 0 0 0 -0.597848 -0.949244 -0.695614 -0.6315 0		
11 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935 -0.116935		
12 0.504319 0.504319 0.504319 0.504319 0.504319 0.504319 0.504319 0.496757 0.335243 0.185846 0.0715519 -0.160985 0.61476 0.504319 0.504319		
13 0.504319 0.504319 0.504319 0.504319 0.504319 0.504319 0.504319 0.508505 0.409631 0.130953 0.367484 0.052985 0.792957 0.504319 0.504319		
14 0 0 0 0 0 -0.624618 -0.508061 -0.455606 -0.399022 -0.477459 -0.597692 0		
15 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		
16		
17		
18 i = 2		
19 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0		
20 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0		
21 -0 -0 -0 -0 -0 -0 -0.305412 -0.240972 -0.118565 -0.0417917 -0.242303 -0.264694 -0		
22 -0 -0 -0 -0 -0 -0.26108 -0.157004 -0.0664739 0.0346655 0.444999 0.0616858 -0.25382 -0		
23 -0 -0 -0 -0 -0 -0.0721619 0.0692259 0.10375 0.255627 1.08347 0.471182 -0.127588 -0		
24 -0 -0 -0 -0 -0 -0.0195733 0.342044 0.270639 0.231972 0.952651 0.507031 -0.1695 -0		
25 -0.52045 -0.52045 -0.52045 -0.52045 -0.52045 -0.52045 -0.310752 0.00801914 0.032264 0.251727 0.752012 0.0688095 -0.437859 -0.52045		
26 -0.414124 -0.414124 -0.414124 -0.414124 -0.414124 -0.414124 -0.340198 -0.20767 -0.123953 0.323786 -0.033496 -0.377352 -0.414124		
27 -0 -0 -0 -0 -0 -0 -0 -0.145425 0.434612 0.038635 -0.304603 -0		
28 -0.414124 -0.414124 -0.414124 -0.414124 -0.414124 -0.414124 -0.193709 0.365106 -0.0349484 -0.414124 -0.414124		
29 -0.52045 -0.52045 -0.52045 -0.52045 -0.52045 -0.52045 -0.518269 -0.354217 -0.343877 -0.0126615 0.233618 -0.191972 -0.52045 -0.52045		
30 -0.52045 -0.52045 -0.52045 -0.52045 -0.52045 -0.52045 -0.489242 -0.294626 -0.132121 0.0694017 0.065242 -0.268918 -0.52045 -0.52045		
31 -0 -0 -0 -0 -0 -0.303198 -0.146304 0.122487 0.151459 -0.0955583 -0.283415 -0 -0		
32 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0 -0		

Figure 10.54: The S2 feature maps after performing average pooling on C1 layer.




Figure 10.55: The C3 feature maps after performing convolution operation with sparse connectivity with S2 layer.





Figure 10.56: The output vector after we use softmax activation function.

All the results from C1 to output layer can be saved or choose not to be saved by commenting them in the `src/artificialneuralnetworks.cpp` or commenting them out. The δ computation while training / backpropagating can also be seen and saved as `.txt` files so we can figure out whether the CNNs are training well or have vanishing gradients problem or others.




```

output_vector.txt
1 4.36339e-05
2 -0.000129459
3 9.54815e-06
4 4.08578e-05
5 9.4857e-05
6 3.87842e-05
7 -2.83785e-05
8 2.22647e-05
9 3.65675e-05
10 8.26934e-05
11 -1.97492e-05
12 -1.78704e-05
13 -9.2854e-06
14 3.92866e-06
15 -7.68817e-05
16 -2.00093e-05
17 3.76887e-05
18 -1.22074e-05
19 2.76977e-05
20 -1.09376e-05
21 -0.00010593
22 2.34821e-05
23 3.73629e-05
24 -8.27391e-05
25 9.95176e-05
26 6.11887e-05
27 -2.67277e-05
28 6.37315e-06
29 -6.71053e-05
30 -9.90484e-05
31 8.02027e-05

```

Figure 10.57: The δ^{F6} .



```

S4_deltamap_final.txt
1 i = 1
2 6.32029e-06 8.95793e-05 -0.00010703 -0.000173552 1.60147e-05
3 0.000182448 5.66686e-05 -4.33069e-07 7.4924e-05 3.225e-05
4 4.20137e-05 -3.48339e-05 7.88585e-05 -4.42406e-05 -0.000181813
5 -9.03876e-05 -3.00092e-05 7.60825e-05 0.000150513 9.20057e-05
6 0.000203196 -0.00022227 -0.000144309 7.47829e-05 -9.90843e-05
7
8
9 i = 2
10 -5.42072e-05 5.24281e-06 3.56729e-05 0.000193734 -3.63205e-05
11 3.73557e-05 -0.000124641 2.59496e-05 -1.64415e-05 -7.37138e-05
12 0.00011741 1.27629e-05 2.36809e-05 4.49779e-05 -0.000113744
13 3.74159e-05 -0.000131021 5.12295e-05 -5.45604e-06 -6.24822e-05
14 -7.59466e-05 4.85688e-05 -6.52174e-05 -1.4011e-05 0.000118302
15
16
17 i = 3
18 2.17248e-05 2.96477e-05 7.5694e-05 3.765e-05 9.6928e-05
19 -5.69775e-05 -5.69638e-05 0.00012873 -1.5534e-05 5.12692e-05
20 -2.56431e-05 1.68554e-05 9.07676e-06 1.55472e-06 3.37963e-05
21 -3.42228e-05 -7.01426e-06 1.21498e-05 8.4814e-05 2.64127e-05
22 -6.47918e-05 -2.5804e-05 -2.54201e-05 5.58348e-05 -5.11937e-06
23
24
25 i = 4
26 -0.000210311 -0.00014297 -0.000246277 -0.000206578 -0.000101722
27 -2.98123e-05 9.21134e-06 4.89934e-05 8.53895e-05 -7.73975e-05
28 -7.64518e-05 0.000132129 -1.19056e-05 -7.53743e-05 4.84581e-05
29 4.51766e-05 0.00023555 7.20006e-05 8.09797e-05 6.49746e-05
30 8.84252e-05 6.09046e-05 -9.68885e-05 3.98228e-05 3.48768e-06
31
32
33 i = 5
34 -0.00012443 -0.000219234 -8.10631e-05 -8.22525e-05 -6.02224e-05
35 -7.71172e-05 3.4899e-06 2.40098e-05 -2.01917e-05 1.46021e-05
36 -4.3721e-05 -5.73209e-05 4.61834e-05 0.00011961 3.25934e-05
37 -8.29863e-05 -4.0672e-05 -9.72248e-06 2.3081e-05 -4.80831e-05
38 -4.34243e-05 -5.76859e-05 -2.32531e-05 9.0909e-06 5.03772e-05
39

```

Figure 10.58: The δ^{S4} .

iv. Functions in SymIntegration Related to Convolutional Neural Network

[SI*] We will list the basic functions that can be used for computing convolutional neural networks:

CNN_2DConvolutionOperation(vector<vector<int>>& input, vector<vector<double>>& kernel)

this void function will perform a convolution operation of the *input* matrix of size $R \times C$ with *kernel* matrix of size $k \times k, k < \min(R, C)$, the stride is set as 1.

CNN_2DConvolutionOperation(vector<vector<int>>& input, vector<vector<double>>& kernel, int stride)

this function will perform a convolution operation of the *input* matrix of size $R \times C$ with *kernel* matrix of size $k \times k, k < \min(R, C)$, the stride number can be set in this function, the domain for stride is $\in [1, M]$ with M as the $\min(R, C)$.

CNN_2DpadBorder(const vector<vector<int>>& matrix, int pad_size)

this function will return a 2D matrix of **integer** data type from the padding operation toward *matrix*, we can set the padding size here at *pad_size*.

CNN_2DmaxPooling(const vector<vector<int>>& input, int kernelSize, int stride)

this function returns a 2D matrix of **integer** data type from performing max pooling operation on *input* matrix, we can define the kernel size for the max pooling here at *kernelSize* and the number of stride at *stride*.

CNN_2DaveragePooling(const vector<vector<int>>& input, int kernelSize, int stride)

this function returns a 2D matrix of **integer** data type from performing average pooling operation on *input* matrix, we can define the kernel size for the max pooling here at *kernelSize* and the number of stride at *stride*.

```
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Neural Networks/Test
st CNN Max Pooling, Average Pooling, and Padding Operation ]# make
g++ -c -o main.o main.cpp
g++ -o main -g -std=c++11 -lsymintegration
root [ ~/SourceCodes/CPP/C++ Create Library/SI Test Clustered/Neural Networks/Test CNN Max Pooling, Average Pooling, and Padding Operation
]# ./main

Matrix A :
      10      15      20      25
      30      35      40      45
       5      10      15      20
      25      30      35      40

Max Pooling :
      35      45
      30      40

Average Pooling :
      22.500000      32.500000
      17.500000      27.500000

Padding Matrix A :
      0      0      0      0      0      0
      0      10     15     20     25     0
      0      30     35     40     45     0
      0       5     10     15     20     0
      0      25     30     35     40     0
      0      0      0      0      0      0

Time taken by function: 1841 microseconds
```

Figure 10.59: Performing max pooling, average pooling, and padding operation on input matrix (*SymIntegration/Examples/Neural Networks/Test CNN Max Pooling, Average Pooling, and Padding Operation/main.cpp*).

CNN_2DUpsample3DMatrix(const vector<vector<vector<double>>>& input, int M, int N)

this function returns a 3D upsampled matrix of **double** data type from *input* matrix.

CNN_2DAverageUpsample3DMatrix(const vector<vector<vector<double>>>& input, int M, int N)

this function returns a 3D average upsampled matrix of **double** data type from *input* matrix and then multiply each entry by $\frac{1}{M*N}$.

```
Matrix A :
--- Slice (Depth) 0 ---
1   2   3
4   5   6
7   8   9

--- Slice (Depth) 1 ---
10  11  12
13  14  15
16  17  18

Upsample Matrix A :
--- Slice (Depth) 0 ---
1   1   2   2   3   3
1   1   2   2   3   3
4   4   5   5   6   6
4   4   5   5   6   6
7   7   8   8   9   9
7   7   8   8   9   9

--- Slice (Depth) 1 ---
10  10  11  11  12  12
10  10  11  11  12  12
13  13  14  14  15  15
13  13  14  14  15  15
16  16  17  17  18  18
16  16  17  17  18  18

Average Upsample Matrix A :
--- Slice (Depth) 0 ---
0.25 0.25 0.5 0.5 0.75 0.75
0.25 0.25 0.5 0.5 0.75 0.75
1   1   1.25 1.25 1.5 1.5
1   1   1.25 1.25 1.5 1.5
1.75 1.75 2   2   2.25 2.25
1.75 1.75 2   2   2.25 2.25

--- Slice (Depth) 1 ---
2.5 2.5 2.75 2.75 3   3
2.5 2.5 2.75 2.75 3   3
3.25 3.25 3.5 3.5 3.75 3.75
3.25 3.25 3.5 3.5 3.75 3.75
4   4   4.25 4.25 4.5 4.5
4   4   4.25 4.25 4.5 4.5

Time taken by function: 2034 microseconds
```

Figure 10.60: Performing upsampling and average upsampling on input matrix (*SymIntegration/Examples/Neural Networks/Test Upsample 3D Matrix/main.cpp*).

v. Functions in SymIntegration Related to Image Manipulation with STB Image

The source code in SymIntegration that handles these functions is in **src/stbimage.cpp** with its corresponding header **include/symintegral/stbimage.h**.

[SI*] We will list the basic functions that can be used for image manipulation that will be very useful in convolutional neural network (CNN), the engine behind this is the STB image library (it is a header only library so it is quite light) :

ResizeAndGrayscale(const std::string& inputPath, const std::string& outputPath, int newWidth, int newHeight)

this **void** function resize the images that we have listed as **string** vector then turn them into grayscale. This is the basic to create MNIST dataset for digit recognition, if we would like to recreate 60,000 train images of digits from 0 to 9 with our own handwriting or our peers handwriting this function will be very handy.




Figure 10.61: Converting several images with different sizes to uniform size of 28×28 then convert them to grayscale (SymIntegration/Examples/Neural Networks/Test Resize and Convert to Grayscale Images with STB Image/main.cpp).

swap_endian(uint32_t val)

this function convert little-endian to big-endian, it is required to create **idx3-ubyte** or **idx1-ubyte** format.

To create an **idx3-ubyte** file (the standard format for MNIST-style image datasets) in C++, you must write a specific binary header followed by the raw pixel data. The format uses big-endian byte ordering, which requires reversing the byte order of integers on most modern (little-endian) systems.

convert_to_idx(const vector<string>& image_paths, const vector<uint8_t>& labels, const string& images_out_path, const string& labels_out_path, int width, int height)

this function will create the **idx3-ubyte** format for the training images, and **idx1-ubyte** for the labels of the images, it corresponds one-to-one, order matters. The input is a **string** vector to represents the path to the images, one by one, we can call them from a **.txt** file or manually write them in the **main.cpp**, then the labels is a **uint8_t** vector to represents the label, for digit recognition we can use the label directly as 0 to 0, 1 to 1, but if it is an image classification that we are creating, then we can represents 0 as "cat", 1 as "dog", 2 as "horse", 3 as "monkey", 4 as "dragon", 5 as "snake", 6 as "rooster", etc.




Figure 10.62: An example to create **idx3-ubyte** and **idx1-ubyte** files from two grayscale images of size 28×28 (SymIntegration/Examples/Neural Networks/Test Create idx3-ubyte Files with STB Image/main.cpp).

An **idx3-ubyte** file consists of:

- **Magic Number (4 bytes):** Always **0x00000803** for 3D unsigned byte tensors.
- **Number of Items (4 bytes):** The total count of images.
- **Number of Rows (4 bytes):** Height of each image.
- **Number of Columns (4 bytes):** Width of each image.
- **Data:** Raw pixel bytes (0-255) for all images, ordered sequentially.

* Pixel values should be unsigned 8-bit integers (**uint8_t**) ranging from 0 to 255.

the **idx** file format is a method for storing numerical data in the form of vectors and multi-dimensional matrices.

loadIDX3(const std::string& filename)

this function will load idx3-ubyte format and convert it to become 3D matrix of **integer** data type (**vector<vector<vector<int>>>**).

loadIDX1(const std::string& filename)

this function will load idx1-ubyte format and convert it to become a vector of **integer** data type (**vector<int>**).



```

1 | = 1
2 229 235 232 233 234 232 231 231 233 234 234 232 231 234 230 233 231 231 234 232 230 235 240 242 239 236 237
3 154 157 155 159 177 181 184 185 185 184 179 177 170 171 167 168 176 183 187 189 170 167 161 161 159 157 174
4 50 50 50 50 55 101 112 115 113 110 115 114 107 97 103 80 84 98 114 128 141 98 60 50 50 50 48 85
5 50 50 50 51 56 96 106 108 110 107 109 105 97 104 98 98 101 104 115 117 126 102 68 50 50 50 48 87
6 50 50 50 50 55 85 92 93 89 90 95 95 99 111 95 106 113 111 118 117 104 99 55 49 50 50 48 86
7 50 50 50 50 53 71 74 73 72 76 79 80 86 91 81 88 94 93 92 89 78 66 62 52 50 50 48 87
8 50 50 50 50 51 57 58 58 58 61 64 66 79 66 54 67 76 82 77 61 70 60 79 55 50 50 48 87
9 50 50 50 50 51 52 50 51 62 72 73 73 87 84 84 83 87 81 90 118 105 87 93 54 50 50 48 87
10 50 49 50 50 54 75 86 88 84 79 81 84 92 102 100 94 75 69 97 101 88 89 78 53 50 50 48 87
11 50 50 50 50 58 118 133 139 120 78 73 73 83 133 92 86 76 68 92 73 75 76 65 52 50 50 48 87
12 50 50 50 50 64 156 182 182 157 83 70 75 79 143 160 142 116 95 124 84 84 94 88 55 50 50 48 87
13 50 50 50 50 64 156 181 184 169 141 107 86 77 127 179 176 176 173 171 145 132 112 94 54 50 50 48 87
14 50 50 50 50 65 163 189 179 100 117 163 151 112 80 131 171 174 159 162 167 141 102 81 53 50 50 48 87
15 50 50 50 50 68 186 209 196 102 71 110 109 103 66 80 140 148 127 118 153 138 106 91 53 50 50 48 87
16 50 50 50 50 70 196 215 211 164 80 88 89 93 72 77 99 128 133 140 142 132 102 90 53 50 50 48 87
17 50 50 50 50 70 198 217 216 216 147 81 114 86 103 111 109 119 103 102 117 113 92 85 53 50 50 48 87
18 49 50 50 50 70 198 223 226 224 199 138 142 119 111 98 98 91 85 70 109 133 103 83 53 50 50 48 87
19 50 50 50 50 70 199 226 226 224 220 152 95 111 80 87 89 80 87 100 173 200 174 125 55 50 50 48 87
20 50 49 50 72 213 235 220 226 222 163 114 113 84 119 124 98 99 148 204 208 198 161 58 50 50 48 87
21 50 50 50 72 213 234 224 225 223 141 84 75 101 192 155 140 177 207 210 205 217 176 60 50 50 48 87
22 50 50 51 50 71 207 233 231 230 215 135 108 112 143 179 180 201 207 199 164 184 214 178 61 50 50 48 87
23 50 49 50 70 200 227 227 222 199 129 118 127 153 144 132 143 144 138 131 165 217 179 61 50 50 48 87
24 50 50 50 68 186 210 215 215 179 129 124 126 128 116 120 130 133 135 132 147 211 193 62 50 50 48 87
25 50 50 50 67 190 218 208 204 164 106 116 123 120 116 121 132 132 134 129 137 193 178 62 50 50 48 87
26 50 50 50 69 199 221 212 199 180 116 83 115 122 120 121 131 128 113 87 132 198 180 61 50 50 48 87
27 51 50 50 70 204 223 218 202 190 170 115 97 117 119 123 125 106 88 109 166 209 182 61 50 50 48 87
28 48 48 47 47 67 199 222 220 217 203 188 171 131 119 113 117 117 115 143 170 193 205 183 58 48 47 85
29 93 93 93 92 104 205 233 230 231 227 215 208 199 185 170 164 169 181 199 202 209 196 101 92 93 91 121
30
31
32 | = 2
33 227 235 231 232 234 232 231 231 232 234 234 233 230 231 231 231 231 232 234 231 232 229 239 241 243 237 238
34 123 127 125 125 129 154 162 166 166 165 165 163 160 149 149 145 143 147 162 165 177 165 141 135 128 128 127
35 48 48 48 48 51 94 110 112 112 107 111 113 106 96 102 88 80 86 106 115 135 132 76 56 48 47 49 48
36 50 50 50 50 53 89 105 106 109 105 106 106 96 104 103 93 107 100 115 112 122 116 94 57 50 50 50 50
37 50 50 50 51 53 80 89 90 86 85 91 93 94 106 107 91 119 104 116 115 111 98 80 46 50 50 50 50
38 51 50 50 50 52 67 71 71 70 73 77 77 81 87 81 73 91 86 94 79 82 68 63 63 50 50 50 50
39 50 50 50 50 51 55 55 56 56 59 61 63 75 76 56 64 75 79 83 70 74 67 71 78 50 50 50 50
40 50 50 50 50 53 51 51 61 74 76 78 83 91 86 85 87 82 82 107 122 95 94 79 49 50 50 50
41 50 50 50 50 52 80 93 97 91 80 78 81 84 104 101 99 84 67 81 102 91 84 88 69 50 50 50 50

```

Figure 10.63: An example to create a 3D integer matrix from **idx3-ubyte** format and an integer vector from **idx1-ubyte** format then save them as separated text files (*SymIntegration/Examples/Neural Networks/Test Create 3D Matrix from idx3-ubyte File and Vector from idx1-ubyte/main.cpp*).

Chapter 11

Computational Fluid Dynamics

"Om Mani Padme Hum." - Tibetan Buddhist Chanting

Mankind has been seeking to harness the power of fluids since the early ages in its attempts to utilize it for various utilities such as irrigation in the form of the aqueducts built by the Romans, windmills for milling, ballistics in artillery, and so on. However, engineering these systems relied on rudimentary techniques, which were often trial-and-error experiments based on highly simplified models and empirical observations. Advancing to the 18th century the mathematical model for fluid flow was conceived by David Bernoulli, laying the foundation for studying fluid flows in a systematic and quantitative manner.

This led to various innovations in engineering, helping build hydraulic systems such as pumps and turbines, designing ventilation systems, early aircraft systems, etc. Although designing such systems was now feasible, they relied heavily on manual calculations, experimental data, and simplified assumptions. This required lots of tedious work and manpower, leading to inefficiencies and multiple points of vulnerability. The Wright brothers required wind tunnel testing due to the idealized noninviscid assumption of Bernoulli's equations, whereas flow analysis for pipe networks in the real world required empirical formulas such as the Darcy-Weisback equations, since Bernoulli's equations neglected losses due to friction. Until the mid-20th century, many such drawbacks pushed back on innovation, which was, until the advent of computer technology. The rise of computing technology combined with mathematical models describing the underlying physical phenomena of fluid flows, kick-started the field of computational fluid dynamics (CFD) [1].

I. INTRODUCTION

[SI*] CFD is an amalgamation of various disciplines such as numerical methods, computational algorithms, and principles of physics, for the study and simulation of fluid flows. Over the past few decades, complex systems such as jet turbines and internal combustion engines greatly benefited from the enhancements in this multidisciplinary field, leading to increased efficiencies in performance and energy utilization. Early models, such as the finite difference method (FDM), were used to discretize the governing equations into solvable numerical grids in various applications. The finite element method (FEM) was later developed for solving fluid problems and structural mechanics in complex geometries, where the structure under study was subdivided into simpler elements like triangular or quadrilateral sections. In the 1960s the finite volume method (FVM) transformed the field of CFD by subdividing

the problem into small cells and applying the laws of conservation across these volumes. With complex models such as Reynolds-averaged Navier-Stokes equations, chaotic turbulent models could be comprehensively studied.

Leaps in this domain helped humanity in achieving significant milestones: in 1960s NASA uses CFD in modeling parts of the Apollo spacecraft; in 1970s Boeing adopted CFD for aerodynamic analysis and design of aircraft, decreasing their reliance on wind tunnel systems; and supersonic and hypersonic flows were modeled by CFD to aid in the development of Lockheed Martin's SR-71 aircraft, etc. Evidently, CFD has been crucial in scientific breakthroughs, and these systems are still extensively used in the analysis and design of fluid systems. With advances in parallel computing in the 1990s, large-scale simulations could be tackled with methods such as large eddy simulation (LES) and direct numerical simulation (DNS). This led to applications not only in aerospace but in the automotive industry for combustion simulation and engine cooling, medicine for simulation of blood flow in arteries and design of microfluid devices, environmental science for weather pattern prediction and climate change modeling, etc.

Employing CFD, industries have been able to optimize the design workflows, enhance performance, and streamline costs across automotive, aerospace, energy, and environmental sectors. However, as the simulation models increase in complexity and the demand for higher computational power grows, traditional methods become inefficient. Therefore data-driven techniques, or so-called artificial intelligence (AI) methods, have entered into play by providing faster and often accurate solutions.

Bibliography

- [1] Ahmad, Kamarul Arifin, Jawaid, Mohammad, Singh, Balbir (2026) Artificial Intelligence for Computational Fluid Dynamics, Academic Press, London, England.
- [2] Alpaydin, Ethem (2014) Introduction to Machine Learning 3rd Edition, MIT Press, London, England.
- [3] Arbenz, Peter (2018) Lecture Notes on Solving Large Scale Eigenvalue Problems, <https://people.inf.ethz.ch/arbenz/ewp/Lnotes/lsevp.pdf>, ETH Zürich, Switzerland.
- [4] Bazaraa, Mokhtar S., Jarvis, John J., Sherali, Hanif D. (1990) Linear Programming and Network Flows, John Wiley & Sons, New Jersey, United States.
- [5] Bishop, Christopher M. (2006) Pattern Recognition and Machine Learning, Springer, New York, United States.
- [6] Boyce, William E., DiPrima, Richard C. (2010) Elementary Differential Equations and Boundary Value Problems 9th Edition, John Wiley & Sons, Hoboken, New Jersey, United States.
- [7] Burden, Richard L., Faires, J. Douglas (2001) Numerical Analysis 7th Edition, Brooks/Cole, Pacific Grove, California, United States.
- [8] Colley, Susan Jane (2012) Vector Calculus 4th Edition, Pearson, Boston, MA, United States.
- [9] Dorf, Richard C., Bishop, Robert H. (2010) Modern Control Systems 12th Edition, Pearson, New Jersey, United States.
- [10] Freya, Glanzsche, DS (2025) DianFreya Math Physics Simulator, Berlin-Sentinel Academy of Science, AliceGard.
- [11] Gen, Mitsuo, Cheng, Runwei (1997) Genetic Algorithms and Engineering Design, John Wiley & Sons, New York, United States.
- [12] Goldberg, David E. (2025) Genetic Algorithms in Search, Optimization and Machine Learning, Addison-Wesley, Canada.
- [13] Grother, Patrick J. Blue, James L. (1992) Training Feed Forward Neural Networks Using Conjugate Gradients, Gaithersburg, United States.
- [14] Halak, Basel (2026) Convolutional Neural Network Accelerators, Springer, Switzerland.
- [15] Heitzinger, Clemens (2022) Algorithms with JULIA, Springer, Switzerland.
- [16] Lasthrim, Glanzsche, DS, Freya (2025) Lasthrim Projection 1st Edition, Berlin-Sentinel Academy of Science, AliceGard.

- [17] LeCun, Y., Bottou, L., Haffner, P., Bengio, Y. (1998). Gradient-based learning applied to document recognition. *Proceedings of the IEEE*, 86(11), 2278–2324.
- [18] Miettinen, K., Neittaanmäki, P., Mäkelä, M.M. Periaux, J., Evolutionary Algorithms in Engineering and Computer Science, John Wiley & Sons, West Sussex, England, 1999.
- [19] Nazarathy, Yoni, Klok, Hayden, Statistics with Julia, Springer, Cham, Switzerland, 2021.
- [20] Nelder J.A, Mead R. (1965) A simplex method for function minimization *The computer journal*, 7(4), 308-313.
- [21] Nocedal, Jorge, Wright, Stephen J. (1999) Numerical Optimization, Springer, New York, United States.
- [22] Panju, Maysum, Iterative Methods for Computing Eigenvalues and Eigenvectors, The Waterloo Mathematics Review, University of Waterloo, Canada.
- [23] Purcell, Rigdon, Varberg (2007) Calculus 9th Edition, Pearson, Upper Saddle River, New Jersey, United States.
- [24] Rao, SS (2006) Optimization Theory and Applications 2nd Edition, Wiley Eastern Limited, New Delhi, India.
- [25] Kenneth H. Rosen (2006) Discrete Mathematics and its Applications 6th Edition, McGraw-Hill, Boston, United States.
- [26] Anton, Rorres (2006) Elementary Linear Algebra with Supplemental Applications 10th Edition, Wiley, Boston, United States.
- [27] Anton, Howard, Rorres, Chris, Elementary Linear Algebra with Supplemental Applications 12th edition, Wiley, Hoboken, New Jersey, United States.
- [28] Trefethen, Lloyd N., III, David Bau (1997) Numerical Linear Algebra, SIAM, 3600 University City Science Center, Philadelphia, United States.
- [29] Vries, Joey de (2020) Learn OpenGL - Graphics Programming, Kendall & Welling, United States.
- [30] Walpole, Ronald E., Myers, Raymond H., Myers, Sharon L., Ye, Keying (2012) Probability & Statistics 9th Edition, Pearson, Boston, United States.
- [31] Yousif, O. O. O. , Ziadi, R., Almaymuni, A. Z., Saleh, M.A. (2025) Decent Directions Generator Conjugate Gradient Method With Its Application To Train A Two-Layer Neural Network Model, *AIMS Mathematics* 2025, Volume 10, Issue 11: 26844-26866. doi: 10.3934/math.20251180 (<https://www.aimspress.com/article/doi/10.3934/math.20251180>)
- [32] Zaki, Mohammed J., Meira Jr., Wagner (2006) Data Mining and Analysis: Fundamental Concepts and Algorithms, Wiley Eastern Limited, New Delhi, India.