

**ELEC 221 Lecture 24**  
**The Laplace transform and feedback systems; introducing the  $z$ -transform**

Thursday 1 December 2022

# Announcements

- Midterms available for pickup
- Assignment 6 (computational) due tonight at 23:59 - submit via e-mail, but still fill out contributions on PL
- Assignment 7 available, due Tuesday at 23:59

# Last time


We explored various properties of the Laplace transform.

**TABLE 9.1** PROPERTIES OF THE LAPLACE TRANSFORM

Section	Property	Signal	Laplace Transform	ROC
		$x(t)$ $x_1(t)$ $x_2(t)$	$X(s)$ $X_1(s)$ $X_2(s)$	$R$ $R_1$ $R_2$
9.5.1	Linearity	$ax_1(t) + bx_2(t)$	$aX_1(s) + bX_2(s)$	At least $R_1 \cap R_2$
9.5.2	Time shifting	$x(t - t_0)$	$e^{-st_0}X(s)$	$R$
9.5.3	Shifting in the $s$ -Domain	$e^{s_0 t}x(t)$	$X(s - s_0)$	Shifted version of $R$ (i.e., $s$ is in the ROC if $s - s_0$ is in $R$ )
9.5.4	Time scaling	$x(at)$	$\frac{1}{ a }X\left(\frac{s}{a}\right)$	Scaled ROC (i.e., $s$ is in the ROC if $s/a$ is in $R$ )
9.5.5	Conjugation	$x^*(t)$	$X^*(s^*)$	$R$
9.5.6	Convolution	$x_1(t) * x_2(t)$	$X_1(s)X_2(s)$	At least $R_1 \cap R_2$
9.5.7	Differentiation in the Time Domain	$\frac{d}{dt}x(t)$	$sX(s)$	At least $R$
9.5.8	Differentiation in the $s$ -Domain	$-tx(t)$	$\frac{d}{ds}X(s)$	$R$

## Last time

We used the ROC to reason about the stability and causality of systems with rational Laplace transforms.



## Last time


We saw how to compute  $H(s)$  for systems described by linear constant-coefficient ODEs.

Learning outcomes:

- compute system functions for feedback systems
- use the Laplace transform to design an inverse system
- define the z-transform and compute it and its ROC for basic signals

# Feedback systems

An important application of Laplace transforms is the analysis of **feedback systems**.




Some important applications of feedback include:

- reducing sensitivity to disturbance and errors
- stabilizing unstable systems
- designing tracking systems

# Feedback systems

An important application of Laplace transforms is the analysis of **feedback systems**.




- $H(s)$  is the system function of the forward path
- $G(s)$  is the system function of the feedback path
- the combined function  $Q(s)$  is the closed-loop system function

Try it: compute  $Q(s)$  in terms of  $H(s)$  and  $G(s)$ .



# Feedback systems




Solution: from the convolution property, know that

From the diagram, find that


Thus:

## Application of feedback: constructing inverse systems

Suppose we have some LTI system



Let's use it as part of a larger system:




where the transfer function  $K$  is simply gain of strength  $K$ .

Exercise: What is  $Q(s)$ , and under what conditions can it act as the inverse of  $P(s)$ ?

## Application of feedback: constructing inverse systems

Solution: we can directly apply the expression for the closed-loop system function here




## Application of feedback: stabilizing an unstable system

Consider a system described by the first order DE


$$\frac{dy(t)}{dt} - ay(t) = bx(t)$$

Exercise: compute the system function and draw the ROC. Under what conditions is it stable?



## Application of feedback: stabilizing an unstable system


Suppose we have this setup ( $a > 0$ ):



How can we make it stable?

## Application of feedback: stabilizing an unstable system

Show that the following system will move the pole (under certain conditions on  $K$ ):



Called a *proportional feedback system* since feeding back in a rescaled version of the output.

# The z-transform

CT

Fourier series  
coefficients

$$C_k = \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt$$

Fourier transform  
(spectrum)

$$X(j\omega) = \int_{-\infty}^{\infty} x(t) e^{-j\omega t} dt$$

Laplace transform

$$X(s) = \int_{-\infty}^{\infty} x(t) e^{-st} dt$$

DT

Fourier series  
coefficients

$$C_k = \frac{1}{N} \sum_{n=\langle N \rangle} x[n] e^{-jk\frac{2\pi n}{N}}$$

Fourier transform

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

Z-transform

$$X(z) = \sum_{n=-\infty}^{\infty} x[n] z^{-n}$$

# The z-transform

Consider a DT complex exponential signal

If we put this in a system with impulse response  $h[n]$ , obtain

where

- $z = e^{j\omega}$ : discrete-time Fourier transform
- $z = re^{j\omega}$ : z-transform



## The z-transform

For a general signal  $x[n]$ ,

Just like in CT, this can be expressed with a DTFT involving  $x[n]$ :

## The z-transform

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

Exercise: compute the z-transform of

$$x[n] = a^n u[n]$$

For what values of  $z$  does it converge?

## The z-transform

Exercise: compute the z-transform of

$$x[n] = a^n u[n]$$

For what values of  $z$  does it converge?

Must be the case that  $|z| > |a|$ , or  $|z| < |a|$ .

## The z-transform

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

Exercise: compute the z-transform of

$$x[n] = -a^n u[-n - 1]$$

For what values of  $z$  does it converge?

## The z-transform

Exercise: compute the z-transform of



$$x[n] = -a^n u[-n - 1]$$

For what values of  $z$  does it converge?

Must have  $|z| > |a|$ , or  $|z| < |a|$ . Then can write

## Regions of convergence

To represent the ROC of the z-transform, we will use the z-plane and pole-zero plots:



Unit circle  $z = e^{j\omega}$  ( $|z| = 1$ ) corresponds to the DTFT case (like the vertical axis  $s = j\omega$  for CT).

Exercise: compute the z-transform for

$$x[n] = 7 \left(\frac{1}{3}\right)^n u[n] - 6 \left(\frac{1}{2}\right)^n u[n]$$


and sketch the pole-zero plot of its ROC.

## Regions of convergence

Exercise: compute the z-transform for

$$x[n] = 7 \left(\frac{1}{3}\right)^n u[n] - 6 \left(\frac{1}{2}\right)^n u[n]$$

and sketch the pole-zero plot of its ROC.







# Regions of convergence

ROC of the z-transform has many properties:

- if ROC doesn't contain unit circle, DTFT doesn't converge
- it is a ring in the z-plane centred around origin (for  $z = re^{j\omega}$ , does not depend on  $\omega$ , only  $r$ )
- it does not contain any poles



If a signal  $x[n]$  is of finite duration, its ROC is the entire  $z$ -plane *except possibly*  $z = 0$  and/or  $z = \infty$ .

Exercise: compute the  $z$ -transform and ROC of

1.  $z[n] = \delta[n]$
2.  $z[n] = \delta[n - 1]$
3.  $z[n] = \delta[n + 1]$

## Regions of convergence

If a signal  $x[n]$  is of finite duration, its ROC is the entire  $z$ -plane *except possibly*  $z = 0$  and/or  $z = \infty$ .

Exercise: compute the  $z$ -transform and ROC of

1.  $x[n] = \delta[n]$
2.  $x[n] = \delta[n - 1]$
3.  $x[n] = \delta[n + 1]$

Solution:

Right-sided signal:  $X(z)$  has the form


$$X(z) = \sum_{n=N_1}^{\infty} x[n]z^{-n}$$

This may or may not include  $\infty$  depending on the structure of the signal (in particular, if  $N_1 < 0$ , terms will become unbounded).


If  $|z| = r_0$  is in the ROC for right-sided signal, then so are all *finite*  $z$  where  $|z| > r_0$ .

Similar argument for left-sided signals and the zero point.


# Regions of convergence



Right-sided



Left-sided



Two-sided


## Regions of convergence

Exercise: how many signals could have produced the z-transform

$$X(z) = \frac{1}{(1 - \frac{1}{3}z^{-1})(1 - 2z^{-1})}$$

Draw the pole-zero plot and determine the possible ROCs.

*Hint: this function has 2 zeros; express it in a different way to find them.*




## Regions of convergence

Exercise: how many signals could have produced the z-transform

$$X(z) = \frac{1}{(1 - \frac{1}{3}z^{-1})(1 - 2z^{-1})}$$

Solution:



## Inverse z-transforms

When the z-transform can be expressed as a rational function, we can compute the inverse using partial fractions. We still need the ROC to help us.

Exercise: compute the inverse z-transform of

$$X(z) = \frac{1}{(1 - \frac{1}{3}z^{-1})(1 - 2z^{-1})}$$

if ROC is specified to be  $|z| > 2$ .




## Inverse z-transforms

Exercise: compute the inverse z-transform of

$$X(z) = \frac{1}{(1 - \frac{1}{3}z^{-1})(1 - 2z^{-1})}$$

if ROC is specified to be  $|z| > 2$ .



Use partial fractions:

From ROC, signal is right-sided:

## Inverse z-transforms

Take a closer look at the structure of  $X(z)$ :

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

This is a *power series in z*. If we can do the expansion, we can recover  $x[n]$  from the coefficients.

Exercise 1: what is the inverse z-transform of

$$X(z) = 3z^2 - 1 + 2z^{-3}, \quad 0 < |z| < \infty$$

Solution:

## Inverse z-transforms

Particularly helpful for non-linear cases.

Exercise 2 (Oppenheim 10.63a): what is the inverse z-transform of

$$X(z) = \log(1 - 2z), \quad |z| < \frac{1}{2}$$

Hint:

$$\log(1 - w) = - \sum_{i=1}^{\infty} \frac{w^i}{i}, \quad |w| < 1$$

Solution:

Learning outcomes:

- compute system functions for feedback systems
- use the Laplace transform to design an inverse system
- define the  $z$ -transform and compute it and its ROC for basic signals

Oppenheim practice problems: 9.48, 11.1-11.4, 10.1-10.8, 10.21-10.23, 10.26

## For next time

### Content:

- more properties of z-transforms
- systems described by difference equations
- z-transforms and feedback system analysis

### Action items:

1. Assignment 6 due tonight at 23:59
2. Assignment 7 due Tuesday at 23:59

### Recommended reading:

- From this class: Oppenheim 9.7, 11.0-11.2, 10.1-10.3
- For next class: 10.5-10.7, 11.2