

Circular sets and powers of two

by [G. Lathoud](#), May 2017

Contents

- [Disclaimer](#)
- [Motivation](#)
- [Definition: circular set](#)
- [Filling the set: an example \(\$N = 8\$ \)](#)
- [Success and failure](#)
- [Main result](#)
- [\(O1\): Filling, seen as a permutation](#)
- [\(O2\): About other filling methods](#)
- [\(A1\) Show that \$N\$ power of 2 \$\Rightarrow P_N\$ true](#)
- [\(A2\) Show that \$N\$ not a power of 2 \$\Rightarrow P_N\$ false](#)

Disclaimer


This paper only presents a mathematical result and its demonstration — nothing more. I'd be happy to receive information about related work, and include it as good as I can.

Motivation

Investigate a particular way to fill a circular set without repetition.


Definition: circular set

Take N holes arranged in a circle:




Example circular set with $N = 8$

...pick one as the first hole #0, chose a direction (e.g. counterclockwise), and name the following holes accordingly #1, #2, ..., #($N - 1$):




Circular set ($N = 8$) with direction and numbers


Filling the set: an example ($N = 8$)




Fill the first hole #0




Move $i = 1$ hole forward, and fill the destination hole **#1**






Move $i = 2$ holes forward, and fill the destination hole **#3**



Move $i = 3$ holes forward, and fill the destination hole **#6**



Move $i = 4$ holes forward, and fill the destination hole **#2**




Last steps: We repeat the process, moving forward:

- $i = 5$ holes (to **#7**),
- then $i = 6$ holes (to **#5**),
- and finally $i = 7$ holes (to **#4**).

At this point all holes have been filled, so we stop. The order in which we filled the holes: $[\#0, \#1, \#3, \#6, \#2, \#7, \#5, \#4]$ can be seen as a permutation of the first $N = 8$ non-negative integers.

Success and failure

Repetitions are not accepted, i.e., whenever we land on a hole that has already been filled, we call that a failure. Example: $N = 3$:



Looking at the first few values of N :

$N=2$	success
$N=3$	failure
$N=4$	success
$N=5$	failure
$N=6$	failure
$N=7$	failure
$N=8$	success
$N=9$	failure
\dots	failure
$N=15$	failure
$N=16$	success
$N=17$	failure

Successes seem to correspond to powers of two: $N = 2^q$ where $q \in \mathbb{N}_+^*$

Main result

Formally: we define the property:

$P_N \triangleq$ "for the circular set of N holes, for each step $i = 1 \dots N$, we land on an empty hole (and thus after the N steps all holes are filled)".

The main result of the present paper is:

$$N \text{ is a power of two} \Leftrightarrow P_N \text{ true}$$

where " N is a power of two" means $\log_2(N) \in \mathbb{N}_+$ or equivalently:
 $\exists q \in \mathbb{N}_+ \text{ s.t. } N = 2^q$.

Appendix [\(A1\)](#) demonstrates that N power of 2 $\Rightarrow P_N$ true.

Appendix [\(A2\)](#) demonstrates that N not a power of 2 $\Rightarrow P_N$ false.

Openings

(O1): Filling, seen as a permutation

The order in which we fill the holes, e.g. for $N = 8$: `#0, #1, #3, #6, #2, #7, #5, #4` can be seen as a permutation of the first $N = 8$ non-negative integers.

What happens if we repeat this permutation?

$N=8$ (2^3)

step	0	current	0,1,2,3,4,5,6,7
step	1	current	0,1,3,6,2,7,5,4
step	2	current	0,1,6,5,3,4,7,2
step	3	current	0,1,5,7,6,2,4,3
step	4	current	0,1,7,4,5,3,2,6
step	5	current	0,1,4,2,7,6,3,5
step	6	current	0,1,2,3,4,5,6,7

=> period: 6

So we can observe a periodicity. What about other values of $N = 2^q$?

$N=$	4 (2^2)	period:	2
$N=$	8 (2^3)	period:	6
$N=$	16 (2^4)	period:	14
$N=$	32 (2^5)	period:	30
$N=$	64 (2^6)	period:	2280
$N=$	128 (2^7)	period:	18480
$N=$	256 (2^8)	period:	2964
$N=$	512 (2^9)	period:	10248

N=1024 (2^{10}) period: 6036022

These results were obtained using this JavaScript [code](#), running it in a browser console.

Open questions: Do all power of two $N = 2^q$ have a periodicity? If yes, can someone derive a formula giving the period as a function of N , i.e. the series (2,6,14,30,2280,...)?

(O2): About other filling methods

This paper investigated the particular filling method, where at each step i we move $j = i$ holes forward, thus defining the series:

$$(j)_i = (1, 2, 3, \dots, N - 1)$$

Besides the obvious "uniform" filling method, where we move 1 hole forward each time:

$$(j)_i = (1, 1, 1, \dots, 1)$$

...are there other "non-uniform" filling methods without repetition, at least for N being a power of two?

For example, for $N = 2^2 = 4$ the answer is yes. Besides the filling method investigated so far:

$$(j)_i = (1, 2, 3)$$

there is also:

$$(j)_i = (3, 2, 1)$$

which is equivalent to invert the direction. If we additionally restrict $(j)_i$ being itself a permutation of $(1, 2, 3, \dots, N - 1)$, these are the only two possibilities for $(j)_i$ for $N = 4$.

Open question: Are there other methods $(j)_i$ to fill without repetition, which work for all N powers of two? Especially when we restrict the series $(j)_i$ being itself a permutation of $(i)_i = (1, 2, 3, \dots, N - 1)$?

Appendices

(A1) Show that N power of 2 $\Rightarrow P_N$ true

Let us assume H_1 and H_2 :

H_1

N is a power of two: $\exists q \in \mathbb{N}_+^* \text{ s.t. } N = 2^q$

H_2

P_N false, i.e. in at least one of the N steps $i = 1 \dots N$, we land on a hole that has already been filled:

$$\exists (a, b) \in \mathbb{N}^2 \text{ s.t. } 0 \leq a < b < N \text{ and } V_a \equiv V_b [N]$$

where:

- V_i is, at step i , the total number of holes we've been moving since the beginning:

$$V_i \triangleq \sum_{j=1}^i j = \frac{i(i+1)}{2}$$

- $V_a \equiv V_b [N]$ means congruence modulo N :

$$\exists k \in \mathbb{N} \text{ s.t. } V_b - V_a = k \cdot N$$

H_2 implies:

$$\exists k \in \mathbb{N} \text{ s.t. } \frac{b(b+1)}{2} - \frac{a(a+1)}{2} = k \cdot N$$

which we can rewrite:

$$b^2 - a^2 + b - a = 2 \cdot k \cdot N$$

$$(b-a) \cdot (b+a+1) = 2 \cdot k \cdot N \quad (\alpha)$$

Observations: $b-a$ and $b+a$ have same parities, hence $b-a$ and $b+a+1$ have opposite parities, i.e. one is odd and the other one is even.

On the right hand side, $2 \cdot N = 2^{q+1}$ is a power of two, thus k must be odd. Moreover, since $(b-a)$, $(b+a+1)$ and $2 \cdot N$ are all non-negative, we have $k > 0$.

Summarized: $k \geq 1$, and k is the only odd term on the right hand side.

Let us assume $k = 1$. Because the two terms on the left hand side have opposite parities, then either $a+b+1 = 1$ (impossible), or $b-a = 1$ i.e. $b = a+1$ so (α) can be written: $1 \cdot 2 \cdot b = 1 \cdot 2 \cdot N$, and thus $b = N$, which is impossible as well.

Therefore: k is odd and $k \geq 3$.

(A1.a) Let us assume $b - a = k$

(α) can be rewritten:

$$k \cdot (1 + 2a + k) = 2 \cdot k \cdot N$$

$$1 + 2a + k = 2 \cdot N$$

$$a = N - \frac{k + 1}{2}$$

$$b = a + k = N + \frac{2k - k - 1}{2}$$

$$b = N + \frac{k - 1}{2}$$

Since $k \geq 3$, we have $b > N$, which is impossible.

(A1.b) Let us assume $a + b + 1 = k$

i.e.

$$b - a = b - (k - (b + 1)) = 2b - k + 1$$

(α) can be rewritten:

$$(2b - k + 1) \cdot k = 2 \cdot k \cdot N$$

$$2b - k + 1 = 2N$$

$$b = N + \frac{k - 1}{2}$$

Since $k \geq 3$, we have $b > N$, which is impossible.

Conclusion of (A1)

For $q \in \mathbb{N}_+^*$ and $N = 2^q$, assuming P_N false leads to a contradiction, therefore P_N is true. $P_{2^0} = P_1$ is obvious, therefore:

$$\forall q \in \mathbb{N}_+ \ P_{2^q} \text{ true}$$

(A2) Show that N not a power of 2 $\Rightarrow P_N$ false

Formally: we want to show that:

$$\forall N \in \mathbb{N}_+^* \text{ s.t. } \log_2 N \notin \mathbb{N}$$

$$\exists (a, b) \in \mathbb{N}^2 \quad 0 \leq a < b < N \quad \text{s.t.} \quad V_a \equiv V_b [N]$$

where $V_i \triangleq \frac{i(i+1)}{2}$ is the number of holes we've moved since the beginning.

For $(a, b) \in \mathbb{N}^2$, we define the property:

$$T_{a,b} \triangleq 0 \leq a < b < N \text{ and } V_a \equiv V_b [N]$$

To prove the result, we need to find at least one value of (a, b) that verifies $T_{a,b}$

(A2.1) Case: $N = 2p + 1$ where $p \in \mathbb{N}_+^*$

Since $V_i \triangleq \frac{i(i+1)}{2}$ we can write:

$$V_{p+1} - V_{p-1} = p + 1 + p = N$$

Thus, $a = p - 1$ and $b = p + 1$ verify $T_{a,b}$.

Transition

It remains to find (a, b) verifying $T_{a,b}$ when $N = 2p$ is not a power of two.

(A2.2) Case: N even but not a power of two

i.e.

$$\exists (p, q) \in (\mathbb{N}_+^*)^2 \quad N = 2^q \cdot (2p + 1)$$

e.g.

$N = 6 = 2 \cdot 3$	$q:1$	$p:1$
$N = 10 = 2 \cdot 5$	$q:1$	$p:2$
$N = 12 = 4 \cdot 3$	$q:2$	$p:1$
$N = 14 = 2 \cdot 7$	$q:1$	$p:3$

(A2.2.1) When $p \geq 2^q$

Let $a = p - 2^q$ and $b = p + 2^q$.

We have $0 \leq a$ and $a < b$ and $N - b = 2^q \cdot 2p - p = p(2^{q+1} - 1) > 0$

therefore we have $0 \leq a < b < N$.

Besides,

$$\begin{aligned} V_b - V_a &= \sum_{j=1}^{p+2^q} j - \sum_{j=1}^{p-2^q} j \\ &= \sum_{j=p-2^q}^{p+2^q} j - (p - 2^q) \\ &= p \cdot (2 \cdot 2^q + 1) - (p - 2^q) \\ &= 2^q \cdot (2p + 1) \\ &= N \end{aligned}$$

(a, b) verify $T_{a,b}$.

(A2.2.2) When $p < 2^q$

Let $a = 2^q - p - 1$ and $b = 2^q + p$.

We can show, as in (A2.2.1), that $0 \leq a < b < N$.

Besides,

$$\begin{aligned} V_b - V_a &= \sum_{j=1}^{2^q+p} j - \sum_{j=1}^{2^q-p-1} j \\ &= \sum_{j=2^q-p}^{2^q+p} j = 2^q \cdot (2p + 1) \\ &= N \end{aligned}$$

Thus (a, b) verify $T_{a,b}$.