

# Circular sets and powers of two

by [G. Lathoud](#), May 2017

## Contents

- [Disclaimer](#)
- [Motivation](#)
- [Definition: circular set](#)
- [Filling the set: an example \( \$N = 8\$ \)](#)
- [Success and failure](#)
- [Main result](#)
- [\(O1\): Filling, seen as a permutation](#)
- [\(O2\): About other filling methods](#)
- [\(A1\) Show that  \$N\$  power of 2  \$\Rightarrow P\_N\$  true](#)
- [\(A2\) Show that  \$N\$  not a power of 2  \$\Rightarrow P\_N\$  false](#)

## Disclaimer


This paper only presents a mathematical result and its demonstration — nothing more.  
I'd be happy to receive information about related work, and include it as good as I can.

## Motivation

Investigate a particular way to fill a circular set without repetition.


## Definition: circular set

Take  $N$  holes arranged in a circle:




Example circular set with  $N = 8$

...pick one as the first hole #0, chose a direction (e.g. counterclockwise), and name the following holes accordingly #1, #2, ..., #(N - 1):




Circular set ( $N = 8$ ) with direction and numbers


## Filling the set: an example ( $N = 8$ )




Fill the first hole #0




Move  $i = 1$  hole forward, and fill the destination hole **#1**




Move  $i = 2$  holes forward, and fill the destination hole **#3**



Move  $i = 3$  holes forward, and fill the destination hole **#6**



Move  $i = 4$  holes forward, and fill the destination hole **#2**




Last steps: We repeat the process, moving forward:

- $i = 5$  holes (to **#7**),
- then  $i = 6$  holes (to **#5**),
- and finally  $i = 7$  holes (to **#4**).

At this point all holes have been filled, so we stop. The order in which we filled the holes:  $[ \#0, \#1, \#3, \#6, \#2, \#7, \#5, \#4 ]$  can be seen as a permutation of the first  $N = 8$  non-negative integers.

## Success and failure

Repetitions are not accepted, i.e., whenever we land on a hole that has already been filled, we call that a failure. Example:  $N = 3$ :



Failure for  $N = 3$

Looking at the first few values of  $N$ :

$N=2$	success
$N=3$	failure
$N=4$	success
$N=5$	failure
$N=6$	failure
$N=7$	failure
$N=8$	success
$N=9$	failure
...	failure
$N=15$	failure
$N=16$	success
$N=17$	failure

Successes seem to correspond to powers of two:  $N = 2^q$  where  $q \in \mathbb{N}_+^*$

## Main result

Formally: we define the property:

$P_N \triangleq$  "for the circular set of  $N$  holes, for each step  $i = 1 \dots N$ , we land on an empty hole (and thus after the  $N$  steps all holes are filled)".

The main result of the present paper is:

$N$  is a power of two  $\Leftrightarrow P_N$  true

where " $N$  is a power of two" means  $\log_2(N) \in \mathbb{N}_+$  or equivalently:  $\exists q \in \mathbb{N}_+ \text{ s.t. } N = 2^q$ .

Appendix [\(A1\)](#) demonstrates that  $N$  power of 2  $\Rightarrow P_N$  true.

Appendix [\(A2\)](#) demonstrates that  $N$  not a power of 2  $\Rightarrow P_N$  false.

N=1024 (2^10)      period: 6036022

These results were obtained using this JavaScript [code](#), running it in a browser console.

*Open questions:* Do all power of two  $N = 2^q$  have a periodicity? If yes, can someone derive a formula giving the period as a function of  $N$ , i.e. the series (2,6,14,30,2280,...)?

## Openings

### (O1): Filling, seen as a permutation

The order in which we fill the holes, e.g. for  $N = 8$ : [ #0, #1, #3, #6, #2, #7, #5, #4 ] can be seen as a permutation of the first  $N = 8$  non-negative integers.

What happens if we repeat this permutation?

N=8 (2^3)

```
step 0  current  0,1,2,3,4,5,6,7
step 1  current  0,1,3,6,2,7,5,4
step 2  current  0,1,6,5,3,4,7,2
step 3  current  0,1,5,7,6,2,4,3
step 4  current  0,1,7,4,5,3,2,6
step 5  current  0,1,4,2,7,6,3,5
step 6  current  0,1,2,3,4,5,6,7
```

=> period: 6

So we can observe a periodicity. What about other values of  $N = 2^q$ ?

N= 4 (2^2)	period: 2
N= 8 (2^3)	period: 6
N= 16 (2^4)	period: 14
N= 32 (2^5)	period: 30
N= 64 (2^6)	period: 2280
N= 128 (2^7)	period: 18480
N= 256 (2^8)	period: 2964
N= 512 (2^9)	period: 10248

### (O2): About other filling methods

This paper investigated the particular filling method, where at each step  $i$  we move  $j = i$  holes forward, thus defining the series:

$$(j)_i = (1, 2, 3, \dots, N-1)$$

Besides the obvious "uniform" filling method, where we move 1 hole forward each time:

$$(j)_i = (1, 1, 1, \dots, 1)$$

...are there other "non-uniform" filling methods without repetition, at least for  $N$  being a power of two?

For example, for  $N = 2^2 = 4$  the answer is yes. Besides the filling method investigated so far:

$$(j)_i = (1, 2, 3)$$

there is also:

$$(j)_i = (3, 2, 1)$$

which is equivalent to invert the direction. If we additionally restrict  $(j)_i$  being itself a permutation of  $(1, 2, 3, \dots, N-1)$ , these are the only two possibilities for  $(j)_i$  for  $N = 4$ .

*Open question:* Are there other methods  $(j)_i$  to fill without repetition, which work for all  $N$  powers of two? Especially when we restrict the series  $(j)_i$  being itself a permutation of  $(i)_i = (1, 2, 3, \dots, N-1)$ ?

## Appendices

## (A1) Show that $N$ power of 2 $\Rightarrow P_N$ true

Let us assume  $H_1$  and  $H_2$ :

$H_1$

$N$  is a power of two:  $\exists q \in \mathbb{N}_+^* \text{ s.t. } N = 2^q$

$H_2$

$P_N$  false, i.e. in at least one of the  $N$  steps  $i = 1 \dots N$ , we land on a hole that has already been filled:

$$\exists (a, b) \in \mathbb{N}^2 \text{ s.t. } 0 \leq a < b < N \text{ and } V_a \equiv V_b [N]$$

where:

- $V_i$  is, at step  $i$ , the total number of holes we've been moving since the beginning:

$$V_i \triangleq \sum_{j=1}^i j = \frac{i(i+1)}{2}$$

- $V_a \equiv V_b [N]$  means congruence modulo  $N$ :

$$\exists k \in \mathbb{N} \text{ s.t. } V_b - V_a = k \cdot N$$

$H_2$  implies:

$$\exists k \in \mathbb{N} \text{ s.t. } \frac{b(b+1)}{2} - \frac{a(a+1)}{2} = k \cdot N$$

which we can rewrite:

$$\begin{aligned} b^2 - a^2 + b - a &= 2 \cdot k \cdot N \\ (b-a) \cdot (b+a+1) &= 2 \cdot k \cdot N \quad (\alpha) \end{aligned}$$

Observations:  $b-a$  and  $b+a$  have same parities, hence  $b-a$  and  $b+a+1$  have opposite parities, i.e. one is odd and the other one is even.

On the right hand side,  $2 \cdot N = 2^{q+1}$  is a power of two, thus  $k$  must be odd. Moreover, since  $(b-a)$ ,  $(b+a+1)$  and  $2 \cdot N$  are all non-negative, we have  $k > 0$ .

Summarized:  $k \geq 1$ , and  $k$  is the only odd term on the right hand side.

Let us assume  $k = 1$ . Because the two terms on the left hand side have opposite parities, then either  $a+b+1 = 1$  (impossible), or  $b-a = 1$  i.e.  $b = a+1$  so  $(\alpha)$  can be written:  $1 \cdot 2 \cdot b = 1 \cdot 2 \cdot N$ , and thus  $b = N$ , which is impossible as well.

Therefore:  $k$  is odd and  $k \geq 3$ .

## (A1.a) Let us assume $b-a = k$

$(\alpha)$  can be rewritten:

$$k \cdot (1 + 2a + k) = 2 \cdot k \cdot N$$

$$1 + 2a + k = 2 \cdot N$$

$$a = N - \frac{k+1}{2}$$

$$b = a + k = N + \frac{2k - k - 1}{2}$$

$$b = N + \frac{k-1}{2}$$

Since  $k \geq 3$ , we have  $b > N$ , which is impossible.

## (A1.b) Let us assume $a+b+1 = k$

i.e.

$$b-a = b - (k - (b+1)) = 2b - k + 1$$

$(\alpha)$  can be rewritten:

$$(2b - k + 1) \cdot k = 2 \cdot k \cdot N$$

$$2b - k + 1 = 2N$$

$$b = N + \frac{k-1}{2}$$

Since  $k \geq 3$ , we have  $b > N$ , which is impossible.

## Conclusion of (A1)

For  $q \in \mathbb{N}_+^*$  and  $N = 2^q$ , assuming  $P_N$  false leads to a contradiction, therefore  $P_N$  is true.  $P_{2^0} = P_1$  is obvious, therefore:

$$\forall q \in \mathbb{N}_+ \ P_{2^q} \text{ true}$$

## (A2) Show that $N$ not a power of 2 $\Rightarrow P_N$ false

Formally: we want to show that:

$$\boxed{\begin{array}{l} \forall N \in \mathbb{N}_+^* \text{ s.t. } \log_2 N \notin \mathbb{N} \\ \exists (a, b) \in \mathbb{N}^2 \quad 0 \leq a < b < N \quad \text{s.t.} \quad V_a \equiv V_b [N] \end{array}}$$

where  $V_i \triangleq \frac{i(i+1)}{2}$  is the number of holes we've moved since the beginning.

For  $(a, b) \in \mathbb{N}^2$ , we define the property:

$$T_{a,b} \triangleq 0 \leq a < b < N \quad \text{and} \quad V_a \equiv V_b [N]$$

To prove the result, we need to find at least one value of  $(a, b)$  that verifies  $T_{a,b}$

**(A2.1) Case:  $N = 2p + 1$  where  $p \in \mathbb{N}_+^*$**

Since  $V_i \triangleq \frac{i(i+1)}{2}$  we can write:

$$V_{p+1} - V_{p-1} = p + 1 + p = N$$

Thus,  $a = p - 1$  and  $b = p + 1$  verify  $T_{a,b}$ .

### Transition

It remains to find  $(a, b)$  verifying  $T_{a,b}$  when  $N = 2p$  is not a power of two.

**(A2.2) Case:  $N$  even but not a power of two**

i.e.

$$\exists (p, q) \in (\mathbb{N}_+^*)^2 \quad N = 2^q \cdot (2p + 1)$$

e.g.

N = 6	= 2*3	q:1	p:1
N = 10	= 2*5	q:1	p:2
N = 12	= 4*3	q:2	p:1
N = 14	= 2*7	q:1	p:3

**(A2.2.1) When  $p \geq 2^q$**

Let  $a = p - 2^q$  and  $b = p + 2^q$ .

We have  $0 \leq a$  and  $a < b$  and  $N - b = 2^q \cdot 2p - p = p(2^{q+1} - 1) > 0$

therefore we have  $0 \leq a < b < N$ .

Besides,

$$\begin{aligned} V_b - V_a &= \sum_{j=1}^{p+2^q} j - \sum_{j=1}^{p-2^q} j \\ &= \sum_{j=p-2^q}^{p+2^q} j - (p - 2^q) \\ &= p \cdot (2 \cdot 2^q + 1) - (p - 2^q) \\ &= 2^q \cdot (2p + 1) \\ &= N \end{aligned}$$

$(a, b)$  verify  $T_{a,b}$ .

**(A2.2.2) When  $p < 2^q$**

Let  $a = 2^q - p - 1$  and  $b = 2^q + p$ .

We can show, as in (A2.2.1), that  $0 \leq a < b < N$ .

Besides,

$$\begin{aligned} V_b - V_a &= \sum_{j=1}^{2^q+p} j - \sum_{j=1}^{2^q-p-1} j \\ &= \sum_{j=2^q-p}^{2^q+p} j = 2^q \cdot (2p + 1) \\ &= N \end{aligned}$$

Thus  $(a, b)$  verify  $T_{a,b}$ .