

- Alpaga: A Python package for automated analysis of Second Harmonic Generation polarization experiments
- **3** Fabien Rondepierre ¹*, Maxime Fery¹, Oriane Bonhomme ¹*, and
- 4 Guillaume Le Breton [□] ^{1*¶}
- ⁵ 1 Institut Lumière Matière, UMR5306, Villeurbanne, France ¶ Corresponding author * These authors
- 6 contributed equally.

DOI: 10.xxxxx/draft

Software

- Review 🗗
- Repository 2
- Archive ♂

Editor: Open Journals ♂ Reviewers:

@openjournals

Submitted: 01 January 1970 Published: unpublished

License

Authors of papers retain copyright and release the work under a 19 Creative Commons Attribution 4.0 International License (CC BY 4.0).

Summary

Alpaga (AnaLyse en PolArisation de la Génération de second hArmonique) is a Python package designed for the automated analysis of Second Harmonic Generation (SHG) experimental acquisitions. The software provides a comprehensive workflow for processing spectroscopic measurements from Surface Second Harmonic Generation (SSHG) Tran et al. (2017) and Second Harmonic Scattering (SHS) experiments, which are crucial techniques in surface science and nonlinear optics research.

The package implements a robust automated procedure that extracts Gaussian peak intensities from spectral measurements through three main steps: automatic file detection and organization, spectral cleaning and averaging with removing non-physical artifacts, and Gaussian fitting for intensity extraction. This automated approach significantly reduces the time and potential human error associated with manual data processing while providing consistent and reproducible analysis results.

Statement of need

SHG experiments involve long acquisition times, which generate large volumes of spectroscopic data often affected by significant noise, requiring careful processing to extract meaningful physical parameters. Researchers typically face several challenges when analyzing SHG data:
(1) handling numerous acquisition files with varying experimental parameters, (2) removing non-physical artifacts, (3) averaging multiple acquisitions to improve signal-to-noise ratios, and (4) consistently fitting Gaussian profiles to extract peak intensities. These tasks are often performed manually or with custom scripts, leading to inconsistencies between research groups and potential analysis errors.

Alpaga addresses these challenges by providing a standardized, automated workflow specifically designed for SHG polarization analysis. The software is particularly valuable for research groups working with SSHG and SHS experiments, where systematic analysis of polarization-dependent measurements is essential for understanding surface properties and molecular orientation. Python enables Alpaga to provide a user-friendly interface while leveraging efficient numerical libraries for computationally intensive operations.

Alpaga was designed to be used by both experienced researchers in nonlinear optics and students learning SHG analysis techniques. The automated nature, supported by comprehensive documentation (wiki) and tutorials, make the workflow accessible to newcomers while providing the reliability and consistency required for research applications.



Key Features and Implementation

- Alpaga is built on established Python scientific libraries including NumPy, SciPy, and Matplotlib,
- 41 providing reliable numerical operations and visualization capabilities. The software architecture
- follows a modular design with several key components:
- 43 Automated File Management: The software automatically identifies and organizes spectroscopic
- data files based on experimental parameters, streamlining the analysis workflow for large datasets
- with consistent naming conventions.
- 46 Spectral Cleaning and Averaging: Detection and removal of electronic noise and other non-
- 47 physical spikes from spectra. The cleaning procedure includes configurable parameters for
- spike detection sensitivity and handles the averaging of multiple acquisitions with identical
- experimental parameters to improve signal-to-noise ratios.
- 50 Gaussian Fitting with Multiple Options: The package offers various fitting approaches for
- extracting peak intensities from Gaussian profiles. Users can select from different fitting
- ₅₂ algorithms and configure parameters such as baseline handling (fluorescence backround) to
- optimize results based on their specific experimental conditions.
- Domain-Specific Analysis Tools: Dedicated modules for SSHG and SHS analysis provide spe-
- 55 cialized functionality for extracting physical parameters relevant to surface science applications,
- including polarization-dependent analysis and orientation parameter extraction.








Figure 1: Caption for example figure.

Related Work

- While several general-purpose spectroscopic analysis packages exist, such as scikit-spectra
- and rampy (?), Alpaga fills a specific niche in the SHG community by providing specialized
- tools tailored to the unique requirements of polarization-dependent SHG measurements. Unlike
- general spectroscopy packages, Alpaga incorporates domain knowledge about SHG experiments,
- 62 including understanding of typical artifact patterns, polarization conventions, and the specific
- mathematical frameworks used in SSHG and SHS analysis .
- 64 The automated nature of Alpaga's workflow distinguishes it from manual analysis approaches
- commonly used in the field, providing both consistency and efficiency improvements for SHG
- 66 research groups.



Ju Usage and Impact

- The Alpaga project started in 2022 at the Institut Lumière Matière (France) laboratory to merge all the different analysis tools already established in our experimental group. The four authors of this publication participated on the code development and usage. Up-to-date, about a dozen scientists have used Alpaga to treat experimental data. The code have been successfully used in multiple scientific communications and publications already Rondepierre (2025). The software has already enabled more efficient and consistent data analysis workflows for research groups working with SSHG and SHS experiments (at ILM or maybe in other labs thanks to the automated file management procedure), and contributes to improved reproducibility in SHG research.
- The package includes comprehensive documentation with detailed examples and parameter explanations, making it accessible to both experienced researchers and newcomers to SHG analysis. Installation is straightforward through standard Python package management tools, and the software is distributed under the LGPL v2.1 license to ensure broad accessibility.

Acknowledgements

We acknowledge the Institut Lumière Matière (ILM) for supporting the development of this software and the broader SHG research community for valuable feedback and testing. We also thank the contributors who have helped improve the software through bug reports and feature suggestions.

References

- Fery, M. (2025). Sonder la structure de liquides par des méthodes optiques non linéaires: Des électrolytes concentrés en volume aux liquides confinés [PhD thesis]. Université Claude Bernard-Lyon I.
- Le Breton, G. (2022). Second harmonic generation of non-resonant liquid: From the molecular response to the experimental measurement [PhD thesis]. Université Claude Bernard-Lyon I.
- Le Breton, G., Bonhomme, O., Benichou, E., & Loison, C. (2023). Liquid water: When
 hyperpolarizability fluctuations boost and reshape the second harmonic scattering intensities.
 The Journal of Physical Chemistry Letters, 14(18), 4158–4163.
- Le Breton, G., Loison, C., Vynck, K., Benichou, E., & Bonhomme, O. (2024). Microscopic view on the polarization-resolved s-SHG intensity of the vapor/liquid interface of pure water. *The Journal of Chemical Physics*, 161(15).
- Rondepierre, F. (2025). Corrélations orientationnelles des liquides explorées par diffusion de second harmonique [PhD thesis]. Université Claude Bernard Lyon I.
- Shen, Y. (1989). Surface properties probed by second-harmonic and sum-frequency generation.

 Nature, 337(6207), 519–525.
- Tran, R. J., Sly, K. L., & Conboy, J. C. (2017). Applications of surface second harmonic generation in biological sensing. *Annual Review of Analytical Chemistry*, 10(1), 387–414.