

The Impact of Vaccine Resistance on Long-Term Epidemic Outcomes

Glenn Ledder

University of Nebraska-Lincoln

Joint work with Yi (Jacky) Jiang, Kristin Kurianski,
Jane HyoJin Lee, Daniel Cicala, and Yanping Ma


(REUF group at AIM, Summer 2022, 2023)

UNL MathBio Seminar, Oct 3, 2024.

Overview

- ▶ Uniform vaccination can reduce the basic reproduction number of a disease from $\mathcal{R}_0 > 1$ to \mathcal{R}_v .
 - If $\mathcal{R}_v < 1$, the epidemic ends.
- ▶ A significant fraction of the population might be unwilling or unable to be vaccinated. This can help the disease become endemic.
- ▶ The unwilling fraction can change over time.
 - It should generally increase as the disease prevalence decreases.
- ▶ How does this influence epidemic outcomes?

Standard SIR with Vaccination and Loss of Immunity



- ▶ The rate constant ϕ can be reduced to reflect vaccine hesitancy.
 - But even if ϕ is small, everyone gets vaccinated eventually.
- ▶ If everyone gets boosters, the rate constant θ is reduced.
- ▶ Our experience with COVID-19 shows:
 - Many people never get vaccinated.
 - Most people stop getting boosters.

Outline

Model Formulation

General Results

When Vaccination Acceptance is Monotone Increasing

Vaccine Acceptance Peaks at $Y = Y_m$

Conclusions

Model Formulation


General Results

When Vaccination Acceptance is Monotone Increasing

Vaccine Acceptance Peaks at $Y = Y_m$


Conclusions

SIR-WU Model Assumptions




- ▶ Newborns are initially unable to be vaccinated.
- ▶ Willing Susceptibles are vaccinated at rate ϕS_W .
- ▶ Unwilling Recovereds lose immunity at rate θR_U .
- ▶ Unwilling become Willing with rate coefficient $\Omega(I)$.
- ▶ Willing become Unwilling with rate coefficient $\Psi(I)$.

Simplifying Assumptions




1. No disease-induced mortality.
 2. Infection confers total immunity for 100% of patients.
 3. Willing Recovereds remain recovered through regular boosters.
- These assumptions make disease eradication easier.
- A more realistic model would add unnecessary complications.

Initial Formulation



- Population total is constant ($N = 1$); we need 5 differential equations for the 6 variables.
- $S + I + R = 1$, so we can use S and I as dependent variables.
- $W + U = 1$, so we can use W as a dependent variable.
- We use S_W and I_W as the remaining dependent variables.

Initial Formulation



$$\frac{dI}{dt} = -\mu I + \beta S I - \gamma I$$


$$\frac{dI_W}{dt} = -\mu I_W + \beta S_W I - \gamma I_W + \Omega(I)(I - I_W) - \Psi(I)I_W$$

$$\frac{dS}{dt} = \mu - \mu S - \beta S I - \phi S_W + \theta ([1 - S - I] - R_W)$$

$$\frac{dS_W}{dt} = -\mu S_W - \beta S_W I - \phi S_W + \Omega(I)(S - S_W) - \Psi(I)S_W$$

$$\frac{dW}{dt} = -\mu W + \Omega(I)(1 - W) - \Psi(I)W$$

Two Time Scales in Disease Models



- ▶ **The fast time scale (days) shows the epidemic phase.**
 - Infectious population fractions are significant.
 - Plots on the fast time scale show no clue to endemic behavior.
 - Demographic changes (birth, natural death, etc) are negligible.
- ▶ **The slow time scale (years) shows the long-term behavior.**
 - Infectious population fractions are very small.
 - On the slow scale, the epidemic behavior appears at $t = 0$.
 - Both demographic and disease processes are important.

Scaling for the Endemic Phase

- ▶ Time is referred to the demographic time scale $1/\mu$.
- ▶ Infectious populations are rescaled via $I = \epsilon Y$, $I_W = \epsilon Z$.
 - *This is necessary so that all variables are $O(1)$ as $t \rightarrow \infty$.*
- ▶ Slow process parameters are referred to a slow time scale.
 - Loss of immunity and behavior changes are slow.
 - $1/\mu$ is the principal slow time scale (≈ 80 years).

$$h = \frac{\theta}{\mu}, \quad \omega = \frac{\Omega}{\mu}, \quad \Sigma = \frac{\Omega + \Psi}{\mu}.$$

- ▶ Fast process parameters are referred to a fast time scale.
 - Infection and vaccination are fast.
 - $1/(\gamma + \mu)$ is the principal fast time scale (≈ 2 weeks).

$$\mathcal{R}_0 = \frac{\beta}{\gamma + \mu}, \quad v = \frac{\phi}{\gamma + \mu}, \quad \epsilon = \frac{\mu}{\gamma + \mu} \ll 1.$$

Final Formulation

$$\epsilon Y' = (\mathcal{R}_0 S - 1) Y,$$

$$S' = \bar{h}(1 - S) - P - hW - \mathcal{R}_0 SY + O(\epsilon),$$

$$\epsilon v^{-1} P' = \omega(Y)S - P + O(\epsilon),$$

$$W' = \omega(Y) - \bar{\Sigma}(Y)W,$$

$$P = vS_W, \quad \bar{h} = h + 1, \quad \bar{\Sigma} = \Sigma + 1.$$

Base Parameters and Values:

- ▶ $\mathcal{R}_0 = 4$: infectiousness is between influenza and Covid
- ▶ $v = 0.5$: vaccination takes a mean of one month
- ▶ $h = 10$: loss of immunity has a mean of 8 years
- ▶ $\Sigma = 8$: opinion change has a mean of 10 years
(can be taken as constant)

Model Formulation

General Results

When Vaccination Acceptance is Monotone Increasing

Vaccine Acceptance Peaks at $Y = Y_m$

Conclusions

Disease-Free Equilibrium (DFE)


1. There is a unique disease-free equilibrium having susceptible fraction $S^0 < 1$.
2. There is a critical value $\omega_{cr} > 0$ such that the DFE is stable if and only if $\omega(0) > \omega_{cr}$.

Interpretation

- ▶ If the infectious population drops to 0, very few Unwilling will become Willing.
 - Then $\omega(0) < \omega_{cr}$.
- ▶ **Optional vaccination means the disease will persist.**

Endemic Disease Equilibrium (EDE) ($\mathcal{R}_0 > 1$)

1. EDEs are solutions of $\omega(Y) = \omega_{cr} \left(1 - \frac{\mathcal{R}_0 Y}{(\mathcal{R}_0 - 1)h}\right)$.
 - $\omega(0) < \omega_{cr}$, $\omega' > 0$ ($\forall Y$): guarantees a unique EDE.
2. There exist functions $\Upsilon_1(Y) > 0$ and $\Upsilon_2(Y) > 0$ such that an EDE is stable if and only if $-\Upsilon_2(Y) < \omega' < \Upsilon_1(Y)$.



- **Extreme sensitivity of attitude to prevalence (large $|\omega'|$) is destabilizing.**

Model Formulation

General Results

When Vaccination Acceptance is Monotone Increasing

Vaccine Acceptance Peaks at $Y = Y_m$


Conclusions

EDE for monotone ω (more disease \rightarrow more vaccination)

► If $\omega(0) < \omega_{cr}$, then \exists a unique EDE:


$$0 < Y^* < (\mathcal{R}_0 - 1)(h + 1).$$

Example: $\omega = \Sigma (1 - e^{-aY})$



EDE for monotone ω (more disease \rightarrow more vaccination)


- ▶ The EDE is unstable if $\omega'(Y^*) > \Upsilon_1(Y^*)$.
 - $\omega = \Sigma(1 - e^{-aY})$: Stable for smaller a .



EDE Stability for monotone ω (more disease \rightarrow more vaccination)


Example: $\omega = \Sigma (1 - e^{-aY})$

- ▶ Σ is the mean number of opinion changes per lifetime.
- ▶ a is the sensitivity of the switch to vaccination.




Numerical Solutions: Example 1, Stable Solution

$$R_0 = 4, \quad h = 10, \quad v = 0.5, \quad \Sigma = 4, \quad a = 1$$



Numerical Solutions: Example 2, Limit Cycle

$$R_0 = 4, \quad h = 10, \quad v = 0.5, \quad \Sigma = 8, \quad a = 2$$



Model Formulation

General Results


When Vaccination Acceptance is Monotone Increasing

Vaccine Acceptance Peaks at $Y = Y_m$

Conclusions

Non-monotone ω (high disease \rightarrow waning confidence)



Example: $\omega = \Sigma d a Y e^{-aY}$



- ▶ Small d : large stable solution
- ▶ Moderate d : small maybe, middle unstable, large stable
- ▶ Large d : small unstable solution

Non-monotone ω (high disease \rightarrow waning confidence)

Example: $\omega = \Sigma daY e^{-aY}$



- ▶ Above/right of black curve: three solutions.
 - Middle unstable, largest stable.
- ▶ Above blue curve: smallest Y is unstable.
 - There is a stable limit cycle.

Model Formulation

General Results

When Vaccination Acceptance is Monotone Increasing


Vaccine Acceptance Peaks at $Y = Y_m$

Conclusions

Conclusions

- ▶ We proposed a disease spread model that includes flexible ideology on vaccination, where the transition of the opinion is a function of the disease spread.
- ▶ There is a unique endemic disease equilibrium for monotone increasing willingness, with a result that may be stable or unstable.
- ▶ By tuning the parameters carefully, we were able to observe limit cycles in the phase plane, that is, oscillatory endemic states.
- ▶ Multiple equilibria are possible with a willingness function that increases to a peak and then decreases.
 - There is a large Y stable equilibrium and either a small Y stable equilibrium or a small Y limit cycle.

Acknowledgement: We would like to thank



REUF is supported by the NSF grants to AIM and ICERM.

