

A cross-platform-replicated activated-fibroblast signature in ligamentum flavum hypertrophy nominates locally-deliverable anti-fibrotic repurposing candidates

Glen Charles Ritschel (Ritschel Research) · Claude (Anthropic)

Preprint · discovery-stage computational analysis · released openly (CC-BY)

2026

Abstract

Ligamentum flavum (LF) hypertrophy is a fibrotic driver of degenerative lumbar spinal stenosis for which there is no approved pharmacotherapy. Using public human single-cell RNA-seq of LF (GSE294458), we derived a within-tissue *activated-fibroblast* transcriptional program — a matrix-depositing state contrast — and tested its reproducibility in an independent bulk LF cohort profiled on a different platform (GSE113212, Agilent microarray). The activated-fibroblast UP program replicated strongly across platforms (18 of 20 genes concordant; rank-enrichment $p = 1.5 \times 10^{-7}$), led by fibrillar collagens, small leucine-rich proteoglycans, HTRA1, MFGE8, POSTN and FN1. Connectivity-based signature reversal (LINCS L1000) was uninformative, returning cytotoxic and anti-proliferative perturbagens rather than anti-fibrotic candidates — an expected limitation for signatures dominated by structural matrix genes. We therefore nominate candidates mechanistically: the approved anti-fibrotics **nintedanib** (PDGFR/FGFR/VEGFR inhibitor) and **pirfenidone** (TGF- β 1 modulator) act on the pathways upstream of the replicated program and are amenable to local intra-ligamentous delivery. We position these against existing prior art, distinguishing degenerative LF hypertrophy from post-surgical epidural fibrosis (for which both drugs have precedent), and release the analysis openly. This is a discovery-stage, in-silico hypothesis intended to motivate wet-lab evaluation; it is not preclinical validation.

1. Introduction

Lumbar spinal stenosis, a leading cause of disability and spine surgery in older adults, is driven in part by hypertrophy of the ligamentum flavum (LF): a chronic, degenerative thickening of the ligament that narrows the spinal canal. LF hypertrophy is fundamentally a fibrotic process — loss of elastic fibres, accumulation of type I/III collagen, fibroblast-to-myofibroblast conversion, and TGF- β 1/Smad-driven matrix deposition. Despite a well-mapped mechanism, there is no approved drug that slows or reverses LF hypertrophy; management remains surgical decompression once stenosis is established.

Two obstacles have limited computational drug-repurposing approaches here. First, human single-cell data for LF is scarce. Second, disease-versus-control transcriptional contrasts derived from one small cohort frequently fail to reproduce in another. We address both by (i) building the disease signature as a within-tissue cell-state contrast rather than a case-versus-control contrast, and (ii) requiring the signature to reproduce in an independent cohort on a different platform before it is used. We then identify repurposing candidates by mechanism rather than by connectivity mapping, which we show is ill-suited to this signature, and we assess the resulting hypothesis against the existing patent and literature landscape.

2. Methods

2.1 Data

Single-cell discovery cohort: GSE294458 (human LF, 10x Genomics; one hypertrophic and one non-hypertrophic donor). Independent replication cohort: GSE113212 (human LF, Agilent SurePrint G3 microarray; four elderly/hypertrophic versus four young/non-hypertrophic samples). Both are public; no unpublished or author-provided data were used.

2.2 Single-cell processing and signature definition

Standard quality control was applied, including an explicit hemoglobin-fraction filter to remove erythrocyte contamination, and hemoglobin and mitochondrial genes were excluded from feature selection. Cells were clustered and annotated to compartments (fibroblast, macrophage, endothelial, and others) by canonical markers, with an erythrocyte label to exclude residual red-blood-cell clusters. Within the fibroblast compartment, cells were sub-clustered and scored for a myofibroblast/activation program; the disease signature was defined as the differential expression between the highest-scoring (activated) and lowest-scoring (resting) fibroblast states, pooling both donors so the contrast reflects fibroblast state rather than the single-donor difference. The ranked signature was scrubbed of ribosomal, translation-factor, mitochondrial, hemoglobin and immunoglobulin genes, which are technical noise. Because only the activated (UP) pole reproduced (see Results), the signature was locked as UP-anchored.

2.3 Cross-platform replication

In GSE113212, per-gene differential expression (hypertrophic versus young) was computed by Welch t-test with log2 fold-change. Replication of the single-cell signature was assessed by (i) the fraction of UP genes with positive bulk fold-change and (ii) a rank-enrichment test of the UP set against the full bulk ranking (one-sided Mann-Whitney).

2.4 Candidate identification and prior-art assessment

Connectivity-based reversal was attempted with LINCS L1000 (L1000CDS2), including an approved-drug filter via the Broad Drug Repurposing Hub. As this proved uninformative (Results 3.3), candidates were instead nominated by matching approved anti-fibrotic mechanisms to the pathways upstream of the replicated program. The resulting hypothesis was assessed against issued and pending patents (Google Patents, USPTO, WIPO) and the peer-reviewed literature.

3. Results

3.1 An activated-fibroblast program in hypertrophic LF

The within-fibroblast activated-versus-resting contrast yielded a coherent matrix-depositing program: fibrillar collagens (COL1A1, COL1A2, COL3A1, COL5A2), small leucine-rich proteoglycans (LUM, ASPN, OGN, DCN), the matrix-remodeling genes HTRA1 and MFGE8, and the fibroblast-activation marker S100A4/FSP1. This is the effector program of fibroblast-to-myofibroblast activation and is consistent with the established histopathology of LF hypertrophy.

3.2 The signature replicates across platforms

Tested in the independent bulk microarray cohort, the activated-fibroblast UP program reproduced strongly despite bulk-tissue dilution and an age-based contrast: 18 of 20 genes showed positive fold-

change in hypertrophic LF, with a rank-enrichment $p = 1.5 \times 10^{-7}$ (Figure 1). The canonical fibrosis core was elevated and mostly significant — FN1 (+3.1), POSTN (+2.5), HTRA1 (+1.76), LUM (+1.89), and every fibrillar collagen (+1.3 to +1.75). This is cross-platform (single-cell to microarray) confirmation that the signature reflects LF disease biology rather than a single-cohort artifact. The resting (DOWN) pole did not reproduce, as expected — its inflammatory/secretory genes rise with age and inflammation and are not lower in aged bulk tissue — and it was therefore dropped.




Figure 1. The single-cell-discovered activated-fibroblast signature, tested in an independent bulk LF cohort (GSE113212). Bars show bulk log2 fold-change (hypertrophic vs young); teal = $p < 0.05$. 18/20 genes are up; rank-enrichment $p = 1.5 \times 10^{-7}$.

3.3 Connectivity mapping was uninformative

Signature-reversal querying of LINCS L1000 returned perturbation signatures dominated by cytotoxic and anti-proliferative agents (tubulin, HDAC, topoisomerase, Chk1, and PI3K/mTOR inhibitors) and chemical-probe tool compounds; after filtering to approved, non-oncology drugs, no credible anti-fibrotic candidate remained. This is expected: a signature dominated by structural matrix genes is nominally “reversed” by anything that halts secretory or proliferating cells, and the gentle, pleiotropic anti-fibrotics of therapeutic interest produce weak L1000 signatures that connectivity ranking disfavors. Connectivity mapping is therefore the wrong instrument for this problem, which motivates mechanism-based nomination.

3.4 Mechanism-matched candidates

Two approved anti-fibrotics map directly onto the replicated program. **Nintedanib** (approved for idiopathic pulmonary fibrosis, systemic-sclerosis-associated ILD, and progressive fibrosing ILD) inhibits PDGFR, FGFR and VEGFR — receptors driving fibroblast activation, and, via VEGFR, the angiogenic coupling implicated in LF hypertrophy — all upstream of the collagen/matrix effector program observed here.

Pirfenidone (approved for IPF) suppresses TGF- β 1 and collagen synthesis, directly upstream of the replicated collagen program. Both are amenable to local intra-ligamentous or peri-ligamentous depot delivery, consistent with the localization principle for tissue-restricted anti-fibrotic action. We nominate nintedanib as the lead on mechanistic breadth, with pirfenidone as an in-class companion.

3.5 Prior-art positioning

No published study has tested nintedanib or pirfenidone in LF or LF hypertrophy, and no patent claims either drug for any spinal indication. The nearest art is *post-surgical epidural fibrosis*, where both drugs have precedent (a locally-delivered pirfenidone spinal hydrogel; rat laminectomy studies of both drugs). Epidural fibrosis is a distinct pathology — acute, iatrogenic scar in the epidural space — whereas LF hypertrophy is chronic degenerative thickening of the ligament body; the delivery target (intra-ligamentous) and disease context also differ. A pending application (US 2024/0026359, University of Pittsburgh) claims the LF-hypertrophy indication with local delivery but only for a microRNA (miR-29a) agent, not small molecules. The mechanism these drugs share with LF fibrosis is what makes them plausible, and also what limits the novelty of the concept.

4. Discussion

Two results are worth emphasizing. First, a disease-relevant fibroblast-activation signature can be recovered from a single, minimal single-cell LF dataset and shown to reproduce across platforms, provided it is built as a within-tissue cell-state contrast and validated by rank-based replication rather than trusted from one cohort. Second, this replicated signature points, by mechanism, to two approved, locally-deliverable anti-fibrotics as repurposing hypotheses for a disease with no approved pharmacotherapy.

The limitations are substantial and stated plainly. The single-cell evidence rests on one cohort of one donor per group; the replication cohort is bulk tissue with an age-based rather than pathology-graded contrast; and the entire analysis is computational, with illustrative rather than measured pharmacology. Connectivity mapping failed to nominate candidates, so the drug hypotheses are mechanistic inferences, not data-driven hits. Nintedanib and pirfenidone are the archetypal anti-fibrotics acting on the very pathway that defines the disease, so applying them to another fibrotic tissue is a rational but modest inductive step; the shared mechanism that makes them plausible also makes the concept unsurprising.

We release this openly rather than seeking patent protection. The therapeutic concept is mechanistically obvious and adjacent prior art exists in epidural fibrosis; a defensible claim would require specific formulation and efficacy data that a computational program cannot generate. Open publication establishes the finding, invites the wet-lab groups who can test it, and serves the translational goal better than a narrow filing. We hope the replicated signature and the candidate rationale are useful to laboratories positioned to evaluate local anti-fibrotic therapy for ligamentum flavum hypertrophy.

Data and code availability

Analyses use public data (GSE294458, GSE113212). Processing notebooks (single-cell signature, cross-platform replication, and reversal analyses) are available from the authors and will be deposited in a public repository alongside this preprint.

Competing interests

None declared. No patent has been filed on this work; it is released openly.

Acknowledgements

Prepared by Ritschel Research in collaboration with Claude (Anthropic). This is discovery-stage computational research and is not medical or legal advice.

References

1. Sun C, et al. Ligamentum flavum fibrosis and hypertrophy: molecular pathways, cellular mechanisms, and future directions. FASEB J, 2020. pubmed.ncbi.nlm.nih.gov/32608536/
2. Fibroblast heterogeneity and myofibroblast conversion in ligamentum flavum hypertrophy (single-cell). The Spine Journal, 2025. pubmed.ncbi.nlm.nih.gov/39653186/
3. Lineage-specific alterations of fibroblasts and endothelial cells in LF hypertrophy (GSE294458). Front Immunol, 2025. ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE294458
4. Gene expression of ligamentum flavum, young vs elderly (GSE113212). Agilent microarray. ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE113212
5. Decorin antagonizes TGF- β 1 in ligamentum flavum hypertrophy. Exp Mol Med, 2023. nature.com/articles/s12276-023-01023-y
6. Angiogenesis couples mechanical stress and ligamentum flavum hypertrophy. 2015. pubmed.ncbi.nlm.nih.gov/25850600/
7. Pirfenidone-loaded HAMA hydrogel prevents epidural adhesions after laminectomy. Drug Deliv Transl Res, 2023. link.springer.com/article/10.1007/s13346-022-01236-0
8. Nintedanib reduces epidural fibrosis in a rat laminectomy model. NASSJ, 2026. sciencedirect.com/science/article/pii/S2590139726000347
9. US 2024/0026359 A1, Methods of treating spinal stenosis and ligamentum flavum hypertrophy (miR-29a), University of Pittsburgh. patents.google.com/patent/US20240026359A1/en
10. Subramanian A, et al. A next generation connectivity map: L1000 platform. Cell, 2017; Broad Drug Repurposing Hub. clue.io