

FINAL REPORT

2017 HIV/AIDS KNOWLEDGE ATTITUDES AND BEHAVIOR SURVEY, JAMAICA

*Report of Findings from HIV/AIDS Knowledge Attitudes and Behavior Survey,
Jamaica 2017*



THE MINISTRY OF HEALTH





ACKNOWLEDGEMENTS

Funding for the project was provided by the Global Fund.

The investigators wish to thank the respondents who participated in the survey and the Training and Field Staff who worked hard to preserve the rigour of the methodology and the integrity of the data during data collection.

We also wish to acknowledge the assistance provided Dr. Nicola Skyers, Senior Medical Officer and Dr. Kelly-Ann Gordon-Johnson, Research Officer from the Ministry of Health who provided direct supervision of the project throughout.

Special thanks to Dr. Sharlene Jarrett, Senior Director, Monitoring and Evaluation, National HIV/STI Control Programme, who provided direct supervision of the project throughout.



Contents

A. INTRODUCTION.....	8
B. EXECUTIVE SUMMARY	10
C. METHODOLOGY	16
CHAPTER 1: REVIEW OF SPECIFIC RISK BEHAVIOUR	22
CHAPTER 2: BEHAVIOUR BY MARITAL STATUS.....	42
CHAPTER 3: ACCESS & ATTITUDES TO CONDOMS.....	49
CHAPTER 4: KNOWLEDGE	54
CHAPTER 5: STIGMA & DISCRIMINATION	57
CHAPTER 6: ADVERTISING MESSAGE RECALL AND IMPACT	61
CHAPTER 7: CHILDBEARING AND ANTENATAL CARE	66
CHAPTER 8: WORKPLACE INTERVENTION.....	67
CHAPTER 9: INTIMATE PARTNER VIOLENCE	68
CHAPTER 10: GENERAL ATTITUDES	74
CHAPTER 11: PRE-EXPOSURE PROPHYLAXIS	76
CHAPTER 12: HIV SELF TESTING	78
CHAPTER 13: RESULTS OF PROGRAMME INDICATORS	79
Bibliography	86

INDEX OF TABLES

Table 1: Age of Sexual Initiation 2017	22
Table 2: Sexual Activity 2017 vs 2012.....	23
Table 3: Married/Cohabiting Status by Age & Gender YR 2017 vs 2012	24
Table 5: Condom use Last Time by Relationship Status / Gender & Age.....	25
Table 6: Multiple Partnerships by Relationship Status YR 2017 vs YR 2012.....	26
Table 7: Multiple Partner Relationships: Condom Used at Last Sex by Length of Relationship, Age & Gender - 2017	27
Table 8: Mean Number of Multiple Partners & Frequency of Condom Use among Persons with Multiple Partnerships	28
Table 9: Perception of Persons with Multiple Partners Analyzed by Condom Non-user YR 2017 vs 2012	29
Table 10: Risk Perception of Persons with Multiple Partners Analyzed by Sometimes and Most Times Condom User YR 2017 vs 2012	30
Table 11: Transactional Sex by Socio-Demographic Variables YR 2012 vs YR 2017	31
Table 12: Condom Use by Persons Reporting Transactional Sex YR 2017 vs YR 2012.....	32
Table 13: Likelihood of using a condom next time sex had by Persons Reporting Transactional Sex in Last 12 months	32
Table 14: Coercive Intercourse (past 12 months) by Socio-Demographic Variables YR 2017 vs YR 2012	33
Table 15: Casual Partnerships by Socio-Demographic Variables YR 2017 vs YR 2012.....	34
Table 16: Same sex Intercourse by Socio-Demographic Variables YR 2017vs YR 2012	35
Table 17: Males STI Incidence YR 2017 vs YR 2012 VS YR 2008.....	36
Table 18: Females STI Incidence YR 2017 vs YR 2012 vs YR 2008.....	36
Table 19: Risky Partnerships Reported by Persons Reporting an STI; YR 2017 vs YR 2012.....	37
Table 20: Had an HIV Test in Past Year.....	38
Table 21: HIV testing by Condom usage of those engaging in Multiple Partnerships in last year YR 2017 vs YR 2012	38
Table 22: Reason for doing HIV Test by Age, Sex and level of Sexual Activity.....	39
Table 23: Know Result of HIV Test done in Past Year	40
Table 24: Reasons for NOT doing an HIV Test	40
Table 25: Participated in Workshop on HIV/ AIDS in Past Year	41
Table 26: Where participated in workshop on HIV/ AIDS in Past Year	41
Table 27: 2017 Length of Time Living with Married/Cohabiting Partner by Age & Gender.....	42
Table 28: 2017 Frequency of Condom Use with Married/Cohabiting by Age & Gender	43
Table 29: Condom Used at Last Sex with Married/Cohabiting by Age & Gender YR 2017 VS YR 2012	43
Table 30: Multiple Partnerships within Married/Cohabiting Unions by Age & Gender	44
Table 31: Condom Use Every Time & Last time with Main Partner Where Multiple Partners Exist YR 2017 VS 2012	44
Table 32: Condom Use with Multiple Partners other than Married/Cohabiting by Age & Gender	44
Table 33: Sexually active Visiting/Casual/Non-Cohabiting by Age & Gender YR 2017 VS YR 2012	45
Table 34: Sexually active Having a Main Partner by Age & Gender YR 2017 VS YR 2012	45
Table 35: Condom Use at Last Sex (With Main Partner) by Age & Gender 2017 VS 2012	46



Table 36: 2017 Frequency of Condom Use with Main Partner by Age & Gender.....	46
Table 37: 2017 Incidence of Multiple Partnerships (with or without main partner)	47
Table 38: 2017 Incidence of Condom Use at Last Sex in Multiple Partnerships	47
Table 39: Mean number of partners in Visiting/Casual/ Relationships in the past 12 months by Age & Gender 2012 vs. 2017	48
Table 40a: Where condoms usually bought by Age YR 2017	49
Table 40b: Where condoms usually bought by Gender YR 2017.....	50
Table 41a: Extent to Which Persons Generally Have a Condom by Gender YR 2017.....	50
Table 41b: Extent to Which Persons Usually Have a Condom by Age YR 2017.....	51
Table 42: Reasons for not using condoms all of the last 10 times sex had YR 2017.....	52
Table 43: General Attitudes to Condoms and Contraceptives	53
Table 44: Details Correct Knowledge of Protective Factors & Myths.....	55
Table 45: Where Obtained Information on HIV.....	56
Table 46: Spoken about AIDS and Safe Sex.....	56
Table 47: Attitudes to People Living with HIV/AIDS by Gender YR 2017.....	57
Table 48: Attitudes to People Living with HIV/AIDS by Age YR 2017.....	58
Table 49: Knowledge of a Person living/dying from HIV/ AIDS.....	59
Table 50: Attitude to Persons with HIV.....	59
Table 51: Attitude to Persons with HIV by Age & Gender.....	60
Table 52: Advertising Message Recall (unprompted) YR 2017.....	61
Table 53: Specific Message recalled (prompted) YR 2017.....	62
Table 54: Behaviour Change as a result of Mass Media Campaign YR 2017.....	63
Table 55: Profiling Persons impacted by message YR 2017.....	64
Table 56: Percentage who have discussed message of campaign with sex partner and friends YR 2017	65
Table 57: Percentage of Females Being Pregnant YR 2017.....	66
Table 58: Percentage of Pregnant Females Making Antenatal Care Visits YR 2017.....	66
Table 59: The Workplace as a point for disseminating information on HIV/AIDS YR 2017.....	67
Table 60: Types of Violence Experienced by Ever Married or Partnered Females in Past 12 Months: YR 2017.....	68
Table 61: Sex of Aggressor by Age Group & Gender.....	69
Table 62: Person who committed violent acts YR 2017.....	69
Table 63: Consider Themselves a Victim of Physical Violence.....	70
Table 64: Do you feel you deserve it.....	70
Table 65: Respondent committing violent acts YR 2017.....	70
Table 66: Statements Indicative of Emotional Health.....	71
Table 67: General Attitudes.....	74
Table 68: Knowledge of and Willingness to take PrEP.....	76
Table 69: Reasons for Unwillingness to take PrEP.....	77
Table 70: Reasons for Willingness to take PrEP.....	77
Table 71: Knowledge of HIV Self Tests.....	79



INDEX OF CHARTS

Chart 1: Percentage able to Obtain Condoms Almost Immediately YR 2017	49
Chart 2: Correct knowledge of all 5 (protective factors and myths)	54
Chart 3: Perception of Persons with HIV.....	60



Table of Abbreviations

AIDS	Acquired Immune Deficiency Syndrome
FSW	Female sex workers
HIV	Human Immunodeficiency Virus
KMR	Kingston Metropolitan Region
MSM	Men who have sex with men
NGO	Non-governmental Organization
NISP	National Integrated Strategic Plan for Sexual and Reproductive Health and HIV
NSUM	Network Scale-Up Method
PrEP	Pre- exposure Prophylaxis
PPS	Probability Proportional to Size
PSU	Primary Sampling Units
RHS	Reproductive Health Surveys
SA	Sexually Active
STIs	Sexually Transmitted Infections
TGP	Transgender



A. INTRODUCTION

The Caribbean accounts for a small percentage of the global epidemic, nevertheless, its HIV prevalence of 1.6% is the second highest in the world after Africa. In 2009, AIDS was said to be the leading cause of death among 25-44 year-old's in the Caribbean. Further to this, women accounted for half of all infections, with younger women having significantly higher prevalence than their male counterparts.

Jamaica, the third largest island in the Caribbean with an estimated 32,000 persons living with HIV, had an adult prevalence of 1.7% in 2011 (UNAIDS 2018). In 2016, Jamaica had 1700 new HIV infections and 1300 AIDS-related deaths. There were also 30,000 people living with HIV in 2016, among whom 35% were accessing antiretroviral therapy.

Through cross sectional studies the key populations identified in Jamaica as being most affected by HIV are:

- Men who have sex with men (MSM), with an HIV prevalence of 32.8%
- Female sex workers (FSW), with an HIV prevalence of 2.9%

However, notable changes have been made in the fight against HIV/AIDS, as since 2010 new HIV infections have decreased by 9%. Additionally, >95% pregnant women living with HIV have accessed treatment or prophylaxis to prevent transmission of HIV to their children. Jamaica also has the highest proportion of people living who know their status in the English-speaking Caribbean—an achievement attributed to community-led prevention and testing programmes and wide spread of provider-initiated testing (UNAIDS 2018).

The response to HIV in Jamaica is guided by a National Integrated Strategic Plan for Sexual and Reproductive Health and HIV (NISP) 2014–2019. NISP was developed to direct the implementation of an integrated approach to Sexual and Reproductive Health (SRH) and HIV planning and programming in Jamaica.

NISP 2014–2019 states that, approximately 74% of all AIDS cases occur in the age group 20-49. Additionally, the 2008 Jamaica's Reproductive Health Surveys (RHS) has flagged the vulnerability of young girls to forced sex and exploitation by older males. Child sexual exploitation, transactional and age disparate sex are also common during this period and both have been linked to poverty, income inequality and inadequate enforcement of protection of children from sexual exploitation.

Key to the integrated HIV prevention and SRH outreach component is to provide a comprehensive package of services for those most vulnerable and at elevated risk for HIV/STI transmission, unplanned pregnancies and unmet family planning needs. Specifically, the component also develops, promotes



and supports health promotion and behaviour change strategies that impact sexual risk behaviours and cultural norms such as multiple partnerships in order to reduce transmission of HIV and other STIs and prevent unplanned pregnancies.

The Ministry of Health has over the years used the National Knowledge, Attitude, Behavior and Practices (KABP) survey to track the attitude and behavior of the general public to the epidemic and monitor the impact of its HIV/AIDS interventions. Continuous monitoring of the knowledge, attitudes, behaviors and practices is pivotal to understanding and identifying the barriers to change and developing the appropriate strategies to manage this epidemic.

B. EXECUTIVE SUMMARY

i. Conclusions

Sexual initiation: Fewer persons initiate early

- **Median age at which persons reported first having sex is maintained at 2012 levels**

Median age for first sex is maintained at age 15 years for males and age 17 years for females as obtained in the 2012 survey.

- **Sex before age 15: Notably fewer males initiate sex before age 15**

In respect of the percentage of young women and men, aged 15-24, who have had sexual intercourse before the age of 15, a marginal improvement is recorded in total. When reviewed by gender however it is seen that there is notable improvement among the males (from 49% in 2012 to 41.6% in 2017). No change is recorded among the females.

2017 Total: 15-24:	Male 41.6%	Female 12.4%	Total 30.7%
2012 Total: 15-24:	Male 49%	Female 12.5%	Total 31.6%

- **Percentage of young adults, 15 to 19 years old, who have never had sex increases**

This improvement in early sexual debut is also reflected in more persons 15-19 who report having never had sex.

2017	15-19	Male 29.7%	Female 54.1%	Total 38.1%
2012	15-19	Male 23.7%	Female 41.8%	Total 31.7%

Correct knowledge of HIV transmission declines versus 2012

The knowledge indicator reviews the percentage of young women and men aged 15–24 who correctly identify ways of preventing the sexual transmission of HIV (100% condom use and one faithful partner) and who reject major misconceptions about HIV transmission (sharing food, healthy looking person and mosquito bites). Among the cohort of persons 15-24 years, it was 32.9% who were able to correctly identify both ways of preventing transmission while rejecting major misconceptions. This represents a decline in correct knowledge over 2012 among both males and females but more so among females. While females were more likely to have debunked the myths, they were less likely than their male counterparts to endorse the correct preventive practices. It was 74.2% of the female cohort, 15-19 years who endorsed 100% condom use, compared to 82.7% of their male peers. (*Table 44*).

2017	Total: 15-24:	Male 31.1%	Female 36%	Total 32.9%
2012	Total: 15-24:	Male 35.6%	Female 42.7%	Total 38.5%

Risk Behaviour

- **Multiple partners in past 12 months increases marginally**

This indicator reviews the percentage of adults aged 15–49 who have had sexual intercourse with more than one partner in the past 12 months. Multiple partner relationships increased marginally in 2017 versus 2012. A deep dive into the data further reveals that multiple partnerships remained fairly stable versus 2012 among females, but increased, although not significantly, among males in the 2017 survey. Most notable within the age subgroups was an increase among the 15-24 years male cohort while the most notable decrease was among the 25-49 years cohort. (*Table 37*)

- **Mean number of partners increases**

While fewer females had multiple partners, they along with all subgroups showed an increase in the mean number of partners. (*Table 39*).

2017	Total: 15-49:	Male 48.9%	Female 13.8%	Total 31.5%
2012	Total: 15-49:	Male 47.2%	Female 13.6%	Total 30.4%

- **Condom Use at last sex among persons reporting multiple partners in the past 12 months decreases versus 2012**

This indicator reviews the percentage of adults aged 15–49 who had more than one sexual partner in the past 12 months and who report the use of a condom during their last intercourse. The results show a decline in the percentage of high risk males and females that protected themselves at their last sex act.

2017 Total: 15-49:	Male 63.7%	Female 40.9%	Total 58.7%
2012 Total: 15-49:	Male 66.2%	Female 43.4%	Total 61.1%

Of note is the fact that among those who protected themselves there may also be elements of inconsistent protection. This is so as when asked about the last 10 times sex was had, significantly fewer persons in this high-risk cohort of multiple partners protected themselves (17.1% in 2017 never used a condom last 10 times sex was had, compared to 9.7% in 2012). (*Table 8*). This reduced condom use is evident irrespective of the type of relationship, whether married/cohabiting or non-cohabiting (*Table 8*).

This decline in condom use is also evident in casual partnerships. Just over a fifth (21%) reported having never used a condom in the last 10 times sex was had and over a half (56.1%) of that cohort indicated that they are not likely to use a condom to protect themselves next time either. (*Table 13*). Fortunately casual partnerships showed significant decline in 2017 versus 2012. (*Table 15*).

The risk of inconsistent condom use in risky sexual encounters is underscored where persons reporting being diagnosed with an STI were significantly more likely to also have reported multiple partners in the last 12 months, as also transactional sex. Both these risk factors showed an increase over 2012, although not significantly so. (*Table 19*).

- **Last time condom use with a non-regular partner increases**

Percentage of young women and men aged 15-24 reporting the use of a condom the last time they had sex with a non-regular partner increases versus 2012 and is highest among persons 15-19 years.

2017 Total: 15-24:	Male 80.1%	Female 65.2%	Total 78.8%
2012 Total: 15-24:	Male 79.2%	Female 57.1%	Total 75.3%

2017: 15-19:	Male 80.2%	Female 83.3%	80.3%
20-24:	Male 80.0%	Female 58.8%	77.5%

- **Percentage of adults aged 15–49 reporting transactional sex declines significantly**

The proportion reporting transactional sex also declined significantly in 2017 versus 2012 (31% in 2017 versus 39% in 2012). This decline was evidenced in both gender, age and relationship status. Where we see an uptick (but not a significant one) in transactional sex is among those whose main partner was still virtually “new” (<1 year) (*Table 11*).

The cohort of persons admitting to transactional sex were more likely to use a condom more times on average within the last 10 times sex had (6.41 cohort reporting transactional sex versus 5.68 for those not reporting transactional sex). *(Table 12)*

Just over a fifth (21%) reported having never used a condom in the last 10 times sex had. Over a half (56.1%) of that cohort indicated that they are not likely to use a condom to protect themselves next time either *(Table 13)*.

2017 Total: 15-49:	Male 45.4%	Female 16.5%	Total 31.1%
2012 Total: 15-49:	Male 53.8%	Female 23.6%	Total 39.0%

- **The Heart rules the Head in opting against condom use**

Against the general decline in condom use we note that the fact of loving or trusting their partner (51%) was the main reason for not using condoms during all of the last 10 times sex was had. These numbers show an increase compared to the 2012 results (46.9%). *(Table 42)*.

The proportion indicating they do not like using a condom also significantly increased in 2017 to 20% from 14.7% in 2012. This was primarily so among the 15-19 years cohort (34.1% compared to 19.4% and 18.7% for the 20-24 and 25-49 years groups respectively). *(Table 42)*

- **Majority know the results of HIV tests done**

This indicator reviews the percentage of women and men aged 15-49 who did an HIV test in the past 12 months and know their results. Among the total sample it is 55% who had done an HIV test in the past year *(Table 20)* and the vast majority (96.8%) report knowing the results of the test.

Among the high risk cohort of inconsistent condom users with multiple partners (condom non-users and sometimes users), it is 46.7% and 28.6% respectively who both did an HIV test and returned for the results *(Table 21)*.

- **Voluntary HIV testing in Antenatal Care accepted by all pregnant women in previous 2 years**

This indicator reviews the percentage of women who were counselled and offered voluntary HIV testing during antenatal care for their most recent pregnancy. They would have accepted an offer of testing and received their results.

Majority of women who had given birth in the past two years or were at least 6 months pregnant had attended an antenatal clinic (90.4%) and private doctor (17.2%). In fact 7.6% had attended both. All had done HIV tests during their visits and the vast majority had been counselled (84.4%) and received their results (96.5%). (*Table 58*).

- **Proportion of persons, 15-49 years, expressing accepting attitudes towards people with HIV/AIDS declines**

This indicator is evaluated based on responses to three statements as indicated:

- If I knew that a shopkeeper or food seller has HIV, I would definitely still buy food or fresh vegetables from them (Strongly Agree and Agree)
- If a teacher has HIV but is not sick, he/she should be allowed to continue to teaching in school (Strongly Agree and Agree)
- If a member of family got infected with HIV I would want them to keep it a secret (Strongly Disagree and Agree)

Accepting attitudes to PLWAs not only remained low but declined significantly in 2017 (11.6%) versus 2012 (14.3%). The main area of concern remains a willingness to purchase fresh vegetables from a vendor known to be HIV+. (*Table 49*).

Just under a half of the sample (48%) knew of someone living with or who has died from AIDS. For about a half of them, this person was a close relative or friend. This further supports the need to foster more accepting attitudes to PLWA to support them and enable them to cope (*Table 52*).

- **Intimate partner violence among women, 15-49 years, in partnered relationships increases significantly**

This indicator reviews ever married or partnered women 15-49 years who experienced physical or sexual violence from a male intimate partner in the previous 12 month period.

Intimate partner violence against married or partnered women, 15-49 years, became considerably more prevalent in 2017 with 14.8% of women reporting same versus 9.3% in 2012. This was notably so for women younger than 25 years. Most prevalent violent acts experienced were 'Pushed or Shoved you' (17.7%), 'Slapped you/ Threw something that could hurt you' (16.8%) and 'Hit you with a fist/ Something that could hurt you' (15.6%). For the most part, these violent acts were experienced to the same extent despite age differences. (*Table 60*)



ii. Recommendations

- Sexual and Reproductive Health programs should seek to build on the gains made in the reduction of early sexual activity in an attempt to make this normative and even lower.
- Efforts must be made to address the decline in knowledge of correct preventive practices among persons 15-24 years and in particular among females. The decline in their endorsement of 100% condom use needs to be reversed as this, coupled with their self-report of intimate partner violence, and could make them less likely to negotiate condom use with their partner.
- The growing tendency of not using condoms because of love or trust needs to once again be aggressively addressed to promote greater protection in relationships.
- An attempt needs to be made to make condom use appealing and attractive as this is also a main reason for non-use.
- Attempts need also to be aggressively continued to reduce stigma and discrimination to someone living with HIV or AIDS among the population. This is particularly so as just under a half of the sample knew someone who has or who has died from AIDS and in a half of these cases it was a close friend or relative.

C. METHODOLOGY

Data was collected from a cross-sectional, household-based, survey among a randomly selected sample of 2000 Persons Island wide. Respondents represented persons aged 15-49yrs.

Sample Design and Selection

The sample design reflected the following multi-staged approach:

1. The island was stratified into 14 parishes with Kingston and St. Andrew treated as two parishes to ensure that the inner-city areas of Kingston in particular, were fully represented in the sample.
2. Each parish was further stratified into constituencies.
3. Each constituency was stratified into two areas, namely:
 - a. Parish capitals and main towns
 - b. Rural areas
4. Each of the two areas comprising the constituencies was then divided into primary sampling units (PSU's) or Enumeration Districts.
5. A random sample of PSUs was then selected with probability proportional to size (PPS). This statistical technique was designed to ensure that the larger PSUs were selected with a greater probability while at the same time; each household was selected with equal probability irrespective of the PSU from which it came. Kingston Metropolitan Region (KMR) and St. James were purposively selected.
6. 80 EDs were selected as follows:
 - 25 EDs in KMR and Montego Bay
 - 28 EDs in other urban areas
 - 27 EDs in rural areas
7. Twenty-five households were then systematically selected from each ED and one person identified within each household as the person to be interviewed. Interviewers identified the households to be included using a map of the area, a random starting point and a predetermined sampling interval. Within the household one (1) respondent was then randomly selected to participate using the Kish card method.
8. The sample was quota controlled for age and gender.

The sample size was estimated to enable results projectable +/- 5% at a 90% confidence level.

- Data Collection:

Data was collected in face-to-face confidential interviews by trained interviewers. Answers to sensitive questions on sexual behavior were however not told to the interviewer but answered by the respondent himself on separate cards provided for the purpose. These cards were then dropped into a large sealed reinforced envelope by the respondent.

The instrument was first pretested for flow, comprehension and to identify areas that would need specific attention in administering. Following refinement of the instrument, trained interviewers



from Hope Caribbean traveled to each enumeration district, selected the household and the person to be interviewed then administered the questionnaire to available and consenting persons.

Selected persons who were at home at the time of the survey and consented to being interviewed, after being read the informed consent form, were interviewed. In urban areas where the selected respondent was not at home at the time of the survey, interviewers requested an appointment on a convenient day and time when they could be interviewed. Based on suggested return days and times, a second visit to the urban ED was made.

If respondent was still unavailable at the time of the second visit they were removed from the sample and a new respondent selected from the specific household. This substitute respondent was the person who answered the door when the interviewer approached the house, provided their demographic quota had not yet been filled. If this quota had been filled then the household was removed from the sample and no interview conducted.

The interviewer then substituted the entire household with the household to the right of the “non-response” household (provided this household had not already been included in the survey)¹ and began selection of a new respondent. Respondent was selected using the Kish card method. If the selected respondent was not at home then the interviewer substituted with the person who first answered the door, provided their demographic quota had not been filled. Only one call/attempt was made to any substitute household. Interviewer continued substitution until a qualifying interview was gained.

Interviewers were rigorously trained over a five-day period with two days devoted to field practice. Interviewers travelled to each ED with a supervisor (for the purpose of on-site validation).

Informed consent was obtained from each respondent before proceeding with the interview. Interviewers assured participants of their anonymity and the confidentiality of the information. No identifiers (name, address, etc.) were included on the questionnaires.

The data collection instrument utilized indicator measures and definitions consistent with UNGASS, where appropriate existing indicators (similar to those used in YR 2012) were used to ensure comparability with previous surveys. Fieldwork was conducted between August and October, 2017.

¹ If household has been included then substitution was made with the nearest household that has not been included

Results of Fieldwork

- A total of 2818 households were called and 68% of initially selected respondents were completed.
- 32% of respondents were substituted.
- Refusals accounted for 4% of initial calls.

Absence from home at time of first visit was highest among older females of employment age and younger males as shown in table below.

Table i: Profile of respondents absent on first visit to household

Profile of selected respondent absent from household at time of first visit (n=140)	
	%
Males 15-24yrs	31.4
Females 15-24yrs	23.5
Males 25-49yrs	9.2
Females 25-49yrs	35.7

Questionnaire Design

The questionnaire was designed to accommodate both face-to-face interviewing and respondent completed sections. This was handled as follows:

Questionnaires were each given a unique identifier number which was also recorded on the self-administered cards for each questionnaire. For the self-administered portion respondents were given Card A if they were married/cohabiting and Card B if they were sexually active in the last year but neither married nor cohabiting. Card C on STI and Health Issues and Card D on Emotional Health were also self-administered but given to all respondents. The questions were read by the interviewer and the respondent asked to fill in the appropriate response on the card without the interviewer seeing. On completion, the respondent was asked to fold the card and drop it into a large sealed and reinforced envelope carried by the interviewer. This served as the receptacle carried by each interviewer into which all cards collected for the day were dropped. Envelopes had a slot cut into them through which the completed cards were dropped. Envelopes were then brought back to the office where they were opened and the cards attached to the correct questionnaire based on the assigned identification number.

The following areas were self-administered:

- Sexual behaviour with main partner based on being in a cohabiting union and not being in a Cohabiting union
- Multiple partnerships
- High risk sexual activities including transactional, coercion, casual partners and commercial sex
- Condom use based on partner type
- Last time condom use
- Substance use
- Sexually transmitted infections
- Emotional Health

The following areas were covered in the face to face portion of the interview:

- Demographics
- Union status
- HIV/AIDS knowledge and protective practices
- Stigma and discrimination
- Condom attitudes and access
- Voluntary counselling and testing
- Risk perception
- Exposure to intervention
- NHP Campaign recall
- Pre-exposure Prophylaxis
- Self- testing
- Parenting
- Knowledge of individuals, MSMs, TGP, FSW and clients of FSW

NOTE: Wherever the self-reported response varied from the interviewed response, the self-reported was taken as the truth and used in analyzing the data and arriving at measurements for indicators. For those questions dealing with sexual behavior which were self-administered, a few persons did not always respond resulting in a slight change in the denominator in some tables.

The 2017 questionnaire included questions on respondent's knowledge of individuals whether by sight, name, phone or internet. Further to this, respondents were asked of their knowledge of Men Sex Men (MSMs), Transgender (TGP), Female Sex Workers (FSW) and clients of FSW and their ages. This was done to facilitate the network scale-up method (NSUM), to estimate the population sizes of MSMs, TGP, FSW and clients of FSW. The NSUM was done based on the assumption that an individual's social network, or the people that they know, is an average representation of the general population that they belong to.



The benefit of the NSUM is that the data should be less prone to social desirability bias as persons will be able to maintain distance and anonymity. This is so as persons are more likely to speak about persons that they know without fear of stigma, than their own behavior.

Sample

Demographics:

The sample comprised 50/50 split of persons aged 15 – 24 years and 25 – 49 years. 50.4% males and 49.6% females were interviewed (*Table i*).

The lower (D) income group accounted for the largest proportion of the sample (58.8%), while the working class (C2) and upper/middle (ABC1) income groups accounted for another 30.1% and 9.8% of the sample respectively. Less than a half of the sample (47.6%) were employed that is 33.7% employed on a full-time basis and while 13.9% reported being employed part time. Unemployed persons and students accounted for 52.4% of the sample (*Table ii*).

Table ii: Socio-Demographic Characteristics of Sample by Age and Gender

Socio-Demographic Characteristics	15-19yrs (%)			20-24yrs (%)			25-49yrs (%)			Total (%)		
Base	M (370)	F (194)	Total (564)	M (258)	F (178)	Total (436)	M (380)	F (620)	Total (1000)	M (1008)	F (992)	Total (2000)
Location												
Urban	78.9	68.6	75.4	73.6	62.4	69.0	60.3	59.4	59.7	70.5	61.7	66.1
Rural	21.1	31.4	24.6	26.4	37.6	31.0	39.7	40.5	40.2	29.5	38.2	33.8
SES Group												
Upper/middle income (ABC1)	13.2	9.8	12.1	10.1	7.9	9.2	7.4	9.5	8.7	10.2	9.3	9.8
Working class (C2)	24.1	28.9	25.7	31.4	37.6	33.9	31.6	30.5	30.9	28.8	31.5	30.1
Lower income (DE)	60.5	59.8	60.3	56.6	53.9	55.5	60.0	59.2	59.5	59.3	58.4	58.8
Refused	2.2	1.5	2.0	1.9	0.6	1.4	1.1	0.8	0.9	1.7	0.9	1.3
Education												
Primary/All-Age	23.2	24.7	23.8	17.8	21.3	19.3	26.3	26.1	26.2	23.0	25.0	24.0
Secondary Level/ Vocational	60.3	58.2	59.6	67.1	66.3	66.7	61.6	59.8	60.5	62.5	60.7	61.6
College/ Associate Degree	5.9	8.8	6.9	7.8	7.3	7.6	7.1	7.3	7.2	6.8	7.6	7.2
University Undergraduate	5.4	2.1	4.3	4.3	3.9	4.1	3.4	5.0	4.4	4.4	4.2	4.3
University Graduate	2.7	2.1	2.5	1.2	0.6	0.9	0.8	0.6	0.7	1.6	0.9	1.2
Not Sure/ No Answer	2.5	4.1	3.0	2.0	0.6	1.4	0.8	1.1	1.0	1.7	1.6	1.7
Employment Status												
Employed full time	6.8	3.6	5.7	41.5	20.8	33.0	64.7	40.6	49.8	37.5	29.8	33.7
Employed part time	6.8	2.1	5.1	19.0	8.4	14.7	23.2	15.6	18.5	16.1	11.7	13.9
Unemployed	30.0	35.1	31.7	29.5	62.9	43.1	11.6	42.4	30.7	22.9	44.7	33.7
Student	56.5	59.3	57.4	10.1	7.9	9.2	0.5	1.3	1.0	23.5	13.8	18.7
Marital Status												
Married/Cohabiting	2.4	15.9	5.8	14.6	37.9	23.8	43.6	59.2	52.9	24.0	50.6	36.8
Visiting /Casual	97.6	84.1	94.2	85.4	62.2	76.2	56.4	40.6	47.0	76.0	49.2	63.1
Sexual Activity												
Not S.A. in last 12 months	20.4	22.5	20.9	10.8	10.8	10.8	9.4	18.2	14.9	13.0	17.3	15.1
"Who raised you"												
Biological Mother	88.9	85.6	87.7	83.3	84.8	83.9	77.1	79.5	78.6	83.0	81.7	82.3
Biological Father	48.4	43.3	46.6	48.1	38.2	44.0	42.9	34.7	37.8	46.2	37.0	41.6
Adopted Parent(s)	0.3	-	-	1.2	1.1	1.1	0.5	0.8	0.7	0.6	0.7	0.6
Grandparents or relatives	24.6	25.8	25.0	21.3	26.4	23.3	30.3	28.7	29.3	25.9	27.7	26.8
Family friends/ Non Relatives	2.2	0.5	1.6	0.8	4.5	2.2	-	2.6	1.6	1.0	2.5	1.75
The State/	-	0.5	-	-	0.6	-	0.5	-	0.2	0.2	0.2	0.2
Siblings	2.2	1.5	1.9	1.6	3.9	2.5	2.9	1.9	2.3	2.3	2.2	2.2
Other	3.0	1.0	2.3	2.7	2.8	2.7	2.6	2.7	2.7	2.8	2.4	2.6

CHAPTER 1: REVIEW OF SPECIFIC RISK BEHAVIOUR

By age 12, 22.3% of boys compared to 2.2% of girls reported having engaged in their first act of sexual intercourse. The median age for first sex among boys is 15years versus 17years for girls. (*Table 1*).

Table 1: Age of Sexual Initiation 2017

	Male	Female
	2017	2017
<u>Median Age of First Sex</u>	15yrs	17yrs
<u>Age: 15-19</u>	(N=260)	(N=89)
≤12yrs.	22.3	2.2
13-19 yrs.	70.0	92.1
<u>Age: 20-24</u>	(N=250)	(N=166)
≤12 yrs.	23.6	5.4
13-19 yrs.	62.0	83.1
20-24 yrs.	2.0	4.2
<u>Age 25-49</u>	(N=371)	(N=614)
≤12 yrs.	17.0	3.1
13-19 yrs.	60.1	75.1
20-24 yrs.	3.5	8.8
25-33 yrs.	0.8	1.6
<u>Total</u>	(N=881)	(N=869)
≤12 yrs.	20.4	3.5
13-19 yrs.	63.6	78.4
20-24 yrs.	2.2	7.0
25-33 yrs.	0.3	1.2

Among males there is a significant reduction in the proportion who had ever had sex (87.5% vs 91% previously) or had sex in the previous 12 months (76.1% in 2017 vs 80.3% in 2012). Among women, sexual activity, in terms of ever had sex or sexually active in the past 12 months, declined marginally. (Table 2).

Table 2: Sexual Activity 2017 vs 2012

	Male		% Points change	Female		% Points change
	2017	2012		2017	2012	
Age: 15-19	(N=370)	(N= 270)		(N= 194)	(N= 215)	
Ever sexually active	70.3	76.3	-6.0	45.9	58.1	-12.2
Sexually active in last 12 months	55.9	60.0	-4.1	35.5	40.0	-4.5
Age: 20-24	(N=258)	(N=185)		(N=178)	(N=201)	
Ever sexually active	97.3	93.0	+4.3	93.3	95.5	-2.2
Sexually active in last 12 months	86.8	80.0	+6.8	83.1	80.0	+3.1
Age: 25-49	(N=380)	(N=446)		(N=620)	(N=482)	
Ever sexually active	97.6	99.6	-2.0	99.0	99.2	-0.2
Sexually active in last 12 months	88.4	90.0	-1.6	80.9	80.0	+0.9
Total Sample	(N=1008)	(N=902)		(N=992)	(N=898)	
Ever sexually active	87.5*	91.0	-3.5	87.6	89.0	-1.4
Sexually active in last 12 months	76.1*	80.3	-4.2	72.5	73.5	-1.0

*p<.05

Additionally, there were significantly fewer committed relationships (married/living together) reported among men while the opposite was true among the females. It was among the older cohort of males (25-49 years) that we see a noticeable, although not significant, decline in the more committed relationships (Table 3). The younger female cohort also displayed higher levels of abstinence in 2017 with more reporting no sexual activity in the past 12 months. (Table 3)

Table 3: Married/Cohabiting Status by Age & Gender YR 2017 vs 2012

	Married/cohabiting		% Points change	Not sexually active in last 12mths		% Points change
	2017	2012		2017	2012	
Male (n=1008)	17.6*	23.2	-5.6	23.6	19.7	3.9
Female (n=992)	36.3*	31.0	5.3	27.5	26.5	1.0
15-24yrs (n=1000)	10.1	9.5	0.6	34.8	32.7	2.1
25-49yrs (n=1000)	43.6	43.5	0.1	16.3	14.1	2.2
Male (15-24yrs); (n=629)	5.4	4.8	0.6	30.9	31.2	-0.3
Female (15-24y); (n=372)	18.0	14.7	3.3	41.4	34.4	7.0
Male (25-49yrs); (n=379)	37.7	41.8	-4.1	11.6	8.1	3.5
Female (25-49); (n=620)	47.3	45.0	2.3	19.2	19.7	-0.05

*p<.05

While the proportion of men involved in visiting or casual relationships remained similar, a statistically significant decline in these relationships was reported by females 15-24yrs. (Table 4).

Table 4: Visiting/Casual Relationship Status by Age & Gender YR 2017 vs 2012

	Visiting or casual relationships		% points change
	YR2017	YR2012	
Male (n=1008)	58.8	57.1	1.7
Female (n=992)	36.2*	42.5	-6.3
15-24yrs (n=1000)	55.1	57.7	-2.6
25-49yrs (n=1000)	40.0	42.4	-2.4
Male (15-24yrs); (n=629)	63.8	64.0	-0.2
Female (15-24y); (n=372)	40.6*	51.0	-10.4
Male (25-49yrs); (n=379)	50.7	50.1	0.6
Female (25-49); (n=620)	33.5	35.3	-1.8

*p<.05

CONDOM USE LAST TIME

Among younger males, last time condom use decreased significantly in relationships of 1 year old or less and increased in relationship > 1 year.

Specifically, among males 15-19yrs, last time condom use in relationships a year old or less, declined between YR2017 and YR 2012 (45.9% 2017 vs 72.3% in 2012). In contrast, there was an almost doubling of the portion of males, 15-19 years, who reported last time condom use in relationships more than a year old (54.1% 2017 vs 27.7% in 2012).

Generally, condoms continue to be used more in Casual/ Visiting Relationships than in Married or Cohabiting relationships for both male and female respondents. Among females, reported last time condom use increased in married/cohabiting relationships (37.7% 2017 vs 5.7% 2012) but declined in visiting/ casual relationships (62.3% 2017 vs 94.3% 2012). (Table 5)

Table 4: Condom use Last Time by Relationship Status / Gender & Age

	Male		% Points change	Female		% Points change
	2017	2012		2017	2012	
Age: 15-19	(N= 148)	(N= 94)		(N=32)	(N=52)	
Condom used last time in relationship ≤1yr	45.9	72.3	-26.4	68.8	71.2	-2.4
Condom used last time in relationship >1yr	54.1	27.7	+26.4	31.2	28.8	+2.4
Age: 20-24	(N=134)	(N= 86)		(N=52)	(N=20)	
Condom Use last time in Married/cohabiting relationship	6.0	4.7	+1.3	28.8	0.0	+28.8
Condom Use last time in Visiting/Casual relationship	94.0	95.3	-1.3	71.2	100.0	-28.8
Age: 25-49	(N=159)	(N=122)		(N=131)	(N=14)	
Condom Use last time in Married/cohabiting relationship	20.1	23.0	-2.9	48.1	21.4	+26.7
Condom Use last time in Visiting/Casual relationship	79.9	77.0	+2.9	51.9	78.6	-26.7
Total Sample	(N=441)	(N=292)		(N=215)	(N=53)	
Condom Use last time in Married/cohabiting relationship	9.1	12.0	-2.9	37.7	5.7	+32.0
Condom Use last time in Visiting/Casual relationship	90.9	88.0	+2.9	62.3	94.3	-32.0

MULTIPLE PARTNERSHIPS

Overall males and females in multiple partnerships were more likely to be single and sexually active than married/cohabiting. Despite this when compared to YR 2012, YR 2017 showed an increase in females in multiple partnerships also reporting being married or cohabiting. (Table 6)

Among persons with multiple partners, the length of the main relationship increased with respondent age. Thus the 15-19yr cohort with multiple partners were most likely to be in a main relationship which was less than 12 months old. In contrast, the 25-49yr old cohort with multiple partners were most likely to be in a main relationship which was more than 12 months old. (Table 6)

Table 5: Multiple Partnerships by Relationship Status YR 2017 vs YR 2012

	Male		% Points change	Female		% Points change
	2017	2012		2017	2012	
Age: 15-19	(n=163)	(n=105)		(n=24)	(n=34)	
Married/cohabiting	1.8	3.5	-1.7	16.7	2.9	+13.8
Single & Sexually active in last 12 months	98.2	96.2	+2.0	83.3	97.1	-13.8
Have multiple partners	(n=103)	(n=77)		(n=19)	(n=28)	
Main relationship >12 mths	79.6	67.5	+12.1	94.7	75.0	+19.7
Main relationship <12 mths	20.4	32.5	-12.1	5.3	25.0	-19.7
Age: 20-24	(n=155)	(n= 111)		(n=43)	(n= 45)	
Married/cohabiting	8.4	5.4	+3.0	14.0	6.7	+7.3
Single & Sexually active in last 12 months	91.6	94.6	-3.0	86.0	93.3	-7.3
Have multiple partners	(n=105)	(n=81)		(n=38)	(n=39)	
Main relationship >12 mths	59.0	55.6	+3.4	52.6	64.1	-11.5
Main relationship <12 mths	41.0	44.4	-3.0	47.4	35.9	+11.5
Age: 25-49	(n=173)	(n=210)		(n=70)	(n=43)	
Married/cohabiting	32.4	31.0	+1.4	44.3	34.9	+9.4
Single & Sexually active in last 12 months	67.6	69.0	-1.4	55.7	65.1	-9.4
Have multiple partners	(n=124)	(n=153)		(n=63)	(n=36)	
Main relationship >12 mths	25.8	36.6	-10.8	27.0	50.0	-23.0
Main relationship <12 mths	74.2	63.4	+10.8	73.0	50.0	+23.0

Table 6 (cont'd): Multiple Partnerships by Relationship Status YR 2017 vs YR 2012

	Male		% Points change	Female		% Points change
	2017	2012		2017	2012	
Total sample	(n=491)	(n=426)		(n=137)	(n=122)	
Married/cohabiting	14.7	17.6	-3.0	29.9	15.6	+14.3
Single & Sexually active in last 12 months	85.3	82.4	+2.9	70.1	84.4	-14.3
Have multiple partners	(n=332)	(n=311)		(n=120)	(n=103)	
Main relationship >12 mths	53.0	49.2	+3.8	45.8	62.1	-16.3
Main relationship <12 mths	47.0	50.8	-3.8	54.2	37.9	+16.3

Table 6: Multiple Partner Relationships: Condom Used at Last Sex by Length of Relationship, Age & Gender - 2017

Condom used at last sex			
	M (n=114)	F (n=11)	Total (N=125)
Age: 15-19			
Relationship >1yr			
Relationship <1yr	50.9	90.9	54.4
Relationship >1yr	49.1	9.1	45.6
Age: 20-24	(n=101)	(n=17)	(N=118)
Relationship <1yr	39.6	58.8	42.4
Relationship >1yr	60.4	41.2	57.6
Age: 25-49	(n=98)	(n=28)	(N=126)
Relationship <1yr	17.3	32.1	20.6
Relationship >1yr	82.7	67.9	79.4
Total	(n=313)	(n=56)	(N=144)
Relationship <1yr	36.7	51.8	39.0
Relationship >1yr	63.3	48.2	61.0

Among persons with multiple partners, in YR 2017 when compared to YR 2012, males reported fewer mean number of partners over the previous 12 months while females recorded a marginal increase. (Table 8).

Interestingly however, significantly less persons in this high-risk cohort of multiple partners protected themselves the last ten times sex was had (17.1% in 2017 never used a condom last 10 times sex was had, compared to 9.7% in 2012). (Table 8)

Reduced condom use is evidenced irrespective of the type of relationship, whether married/cohabiting or non-cohabiting (Table 8).

**Table 7: Mean Number of Multiple Partners & Frequency of Condom Use among Persons with Multiple Partnerships
YR 2017 vs YR 2012**

RESPONDENTS WITH MULTIPLE PARTNERS			
	%		
	YR 2017	YR 2012	
Mean number of partners had in last 12 months			
• Males with multiple partners	5.52 (SD6.49)	6.19 (SD7.17)	
• Females with multiple partners	2.31 (SD1.45)	2.24(SD1.4)	
• Total persons in multiple partnerships	4.82 (SD5.93)	5.31(SD5.57)	
Frequency of condom use last 10 times sex had:			
	(n=627)	(n=545)	
Never (0 times)	17.1*	9.7	
Sometimes (1-7 times)	32.4	32.3	
Most times (8 times)	11.6	15.4	
Every time (10 times)	38.9	42.6	
	50.5%	58.0%	
Condom used last time sex had:			
	56.2 *(n=658)	63.0 (n=548)	
Condom used last time sex had by relationship status			
• Married and used condom	31.4 (n=121)	40.4 (n=94)	
• Visiting/casual and used a condom	61.8 (n=537)	67.6 (n=454)	

RISK PERCEPTION

Risk perception has remained virtually unchanged over the period with just under three-quarters (73%) of those with multiple partners perceiving they have little or no risk of HIV infection.

In total, it is approximately 4 in 10 for both periods (39.4% in 2017 and 40.8% in 2012) who are correctly assessing their risk (*use a condom all the time*).

Approximately, 31.5% do seem to be cognizant of the need to use a condom but do this selectively (sometimes). A few (7.4%) are also opening themselves to risk by claiming that their partners “are clean” so they trust them.

This is corroborated in Table 10 where 62% of persons with multiple partners, who use a condom sometimes, perceive they have little or no chance of infection because of their occasional condom use.

Table 8: Perception of Persons with Multiple Partners Analyzed by Condom Non-user YR 2017 vs 2012

	YR 2017 Total (N=630) %	YR 2012 Total (N=545) %	YR 2017 Non- user (n=107) %	YR 2012 Non- user (n=53) %
No chance	42.0	32.8	33.6	24.5
Little chance	31.0	41.0	31.8	37.7
Moderate chance	10.5	9.2	15.9	13.2
Good chance	6.0	6.4	11.2	13.2
Unsure	10.5	9.9	7.5	9.4
Reasons for <u>little or no</u> chance:	(n=459)	(n=404)	(n=70)	(n=33)
- Use a condom all the time	39.4	40.8	2.9	3.0
- Use a condom sometimes	31.5	35.0	25.7	33.4
- Have sex with spouse only	9.8	8.6	21.4	18.2
- Trust partner/partners are clean	7.4	2.2	18.6	6.1
- Partner may be unfaithful	4.6	4.7	10.0	9.1
- Get check-ups regularly	3.3	0.5	2.9	-
- Condom can burst, not 100% safe	2.2	6.5	4.3	6.1
- No blood transfusions	0.7	0.5	1.4	0.0



Table 9: Risk Perception of Persons with Multiple Partners Analyzed by Sometimes and Most Times Condom User YR 2017 vs 2012

	YR 2017 Sometimes User (n=203) %	YR 2017 Most Times User (n=73) %	YR 2017 Total (N=630) %	YR 2012 Sometimes User (n=176) %	YR 2012 Most Times User (n=316) %	YR 2012 Total (N=545) %
No chance	28.6	38.4	41.9	22.2	40.2	32.8
Little chance	33.5	31.5	31.0	42.0	41.1	41.1
Moderate chance	12.8	12.3	10.5	10.8	7.6	9.2
Good chance	7.9	9.6	6.0	8.5	4.1	6.4
Unsure	17.3	8.2	10.5	16.4	7.0	9.9
Reasons for little or no chance:	(n=126)	(n=51)	(n=459)	(n=113)	(n=257)	(n=370)
- Use a condom all the time	7.9	21.6	39.4	7.1	55.6	40.8
- Use a condom sometimes	55.6	49.0	29.8	63.7	19.5	33.0
- Have sex with spouse only	13.5	11.8	9.8	11.5	7.4	8.6
- Get check-ups regularly	5.6	3.9	3.3	-	-	0.5
- Condom can burst, not 100% safe	0.8	0.0	2.2	1.8	8.6	6.5

TRANSACTIONAL SEX

The proportion reporting transactional sex also declined significantly in 2017 versus 2012 (31% in 2017 versus 39% in 2012). This decline was evidenced in both gender, age and relationship status. Where we see an uptick (but not a significant one) in transactional sex is among those whose main partner was still virtually “new” (<1 year) (*Table 11*).

Table 10: Transactional Sex by Socio-Demographic Variables YR 2012 vs YR 2017

SAMPLE SIZE BY DEMOGRAPHICS FOR TRANSACTIONAL SEX		Total Transactional sex %	
	YR 2017	YR 2017 %	YR 2012 %
Total	(n=2000)	31.0* ↓	39.0
Male	(n=1008)	45.4 ↓	53.8
Female	(n=992)	16.5 ↓	23.6
15-24yrs	(n=1000)	33.6 ↓	42.6
25-49yrs	(n=1000)	28.6 ↓	36.3
Married/cohabiting;	(n=537)	26.4 ↓	29.6
Visiting or casual in last 12 months but not married;	(n=952)	50.4 ↓	54.3
Length of primary relationship (whether married or not)			
Less than a year	(n=244)	57.4 ↑	53.1
More than a year	(n=1756)	27.4 ↓	38.1

*p<0.05;

The cohort of persons admitting to transactional sex were more likely to use a condom more times on average within the last 10 times sex had (6.41 cohort reporting transactional sex versus 5.68 for those not reporting transactional sex). (*Table 12*)

When disaggregated just over a fifth (21%) reported having never used a condom in the last 10 times sex had. Over a half (56.1%) of that cohort indicated that they are not likely to use a condom to protect themselves next time either (*Table 13*).

Table 11: Condom Use by Persons Reporting Transactional Sex YR 2017 vs YR 2012

	YR 2017 Total Transactional sex %	YR 2012 Total Transactional sex %
Average number of times condom used, of last 10 times sex had		
-Had transactional sex	6.41 (SD8.3)	6.24 (SD3.9)
-Had no transactional sex	5.68 (SD12.5)	4.77 (SD4.3)
Frequency of condom use last 10 times sex had:	(n=618)	(n=628)
Never (0 times)	21.0	17.2
Sometimes (1-7 times)	31.9	31.7
Most times (8 times)	12.0	12.6
Every time (10 times)	35.1	38.5
Condom used last time sex had	55.8 (n=622)	58.8 (n=612)

***p< 0.001

Table 12: Likelihood of using a condom next time sex had by Persons Reporting Transactional Sex in Last 12 months

	YR 2017 Non-user (n=130) %	YR 2017 Sometimes User (n=197) %	YR 2017 Most times user (n=74) %	YR 2017 Total (N=618) %
Very likely	15.4	43.1	66.2	57.8
Somewhat Likely	13.8	27.4	18.9	14.9
Neither likely nor unlikely	4.6	4.1	1.4	2.4
Unlikely	16.9	12.2	2.7	8.1
Not at all likely	39.2	6.1	4.1	11.0

COERCIVE SEX

Incidence of coercive sex, whether pressured or forced to have sex, also significantly declined in total and in all sub groups (*Table 14*). However, the percentage of respondents who reported being the aggressor/ forcing their partner into having sex have increased slightly, especially amongst the female cohort. (*Table 14*)

Table 13: Coercive Intercourse (past 12 months) by Socio-Demographic Variables YR 2017 vs YR 2012

	BEEN FORCED %		FORCED/AGGRESSOR %	
	YR 2017	YR 2012	YR 2017	YR 2012
Total	6.7 * (n=1486) ↓	10.3 (n=1384)	10.6 (n=1486)	8.1 (n=1384)
Male	7.7 * (n=767) ↓	13.7 (n=724)	13.4 (n=767)	12.2 (n=724)
Female	5.6 (n=719) ↓	6.7 (n=660)	7.5 ↑ (n=719)	3.6 (n=660)
15-19yrs	6.9 (n=276)	12.8 (n=266)	4.0 (n=276)	8.6 (n=266)
20-24yrs	5.4 (n=372)	11.9 (n=320)	7.0 (n=372)	10.3 (n=320)
25-49yrs	6.7 (n=838)	8.9 (n=798)	5.3 (n=838)	7.0 (n=798)

*p<0.05;

CASUAL PARTNERSHIPS

Casual partnerships are those in which there is no main partner but there is someone they see occasionally and have sex with. These partnerships declined significantly in total and in all sub groups but once again we observe a decline in condom use in last 10 sexual encounters with these partners (*never use 15.4% in YR 2017 versus 10.6% in YR 2012*). (Table 15).

Table 14: Casual Partnerships by Socio-Demographic Variables YR 2017 vs YR 2012

SAMPLE SIZE BY DEMOGRAPHICS FOR HIGH RISK PARTNERS				
	YR 2017	YR 2017	YR 2012	YR 2012
Total	(n=1486)	↓ 29.0 *	(n=1384)	39.2
Male	(n=767)	↓ 46.2 *	(n=724)	57.5
Female	(n=719)	↓ 10.7 *	(n=660)	19.1
15-24yrs	(n=648)	↓ 43.1 *	(n=586)	52.6
25-49yrs	(n=838)	↓ 18.1 *	(n=798)	29.3
Married/cohabiting:	(n=537)	↓ 12.5	(n=487)	19.7
Visiting/casual: Sexually active in last 12 months	(n=948)	↓ 38.4	(n=897)	49.7
Frequency of condom use last 10 times sex had:	(n=429)		(n=537)	
Never (0 times)		↑ 15.4		10.6
Sometimes (1-7 times)		31.0		33.5
Most times (8 times)		11.4		13.8
Every time (10 times)		42.2		42.1
Condom used last time sex had: (n=523)	(n=431)	62.2	(n=523)	65.2

***p<0.001

SAME SEX INTERCOURSE

Persons reporting same sex intercourse in the previous 12 months was 3% of those sexually active during the period. This reflected a doubling of self-reported same sex intercourse for men (1% in 2012 to 2% in 2017) while women remained at 4% self-reporting.

This is so far the only type of relationship in which condom use at last sex increased. Despite this *Never use* and *sometimes use* (of the last 10 times sex had) increased from 44% in 2012 to 67% in 2017.

Table 15: Same sex Intercourse by Socio-Demographic Variables YR 2017vs YR 2012

	YR 2017 Had Same Sex Intercourse (n=1489) %	YR 2012 Had Same Sex Intercourse (n=1333) %
Total sample	3.0	2.4
Male	2.0	1.0
Female	4.0	4.0
Profiling Same Sex Relationships		
	(n=45)	(n=32)
Male	35.6	25.0
Female	64.4	75.0
15-24yrs	51.1	46.9
25-49yrs	48.9	53.1
Married/cohabiting	44.4	31.2
Sexually active in last 12 months but not married	55.6	68.8
Frequency of condom use last 10 times sex had:		
Never (0 times)	37.8	28.1
Sometimes (1-7 times)	28.9	15.6
Most times (8 times)	11.1	9.4
Every time (10 times)	22.2	40.6
Condom used last time sex had	40.0 ↑	37.5

*p < 0.05

STI INCIDENCE

Based on self-reports, males reported a significantly increased incidence of STIs in their lifetime (*Ever had an STI increased from 6.6% in 2012 to 10.5% in 2017 for males 15-24; Table 17*) while females reported a slightly decreased incidence (*Table 18*).

Self-reported genital discharge in both males and females in the past 12 months also significantly reduced over 2012.

Table 16: Males STI Incidence YR 2017 vs YR 2012 VS YR 2008

	MALES 15-24 YRS			MALES 25-49 YRS		
	YR 2017 (n=512) %	YR 2012 (n=377) %	YR 2008 (n=274) %	YR 2017 (n=370) %	YR 2012 (n=442) %	YR 2008 (n=398) %
Ever had an STI	10.5* ↑	6.6	8.0	17.3	16.5	34.9
Had genital discharge in past 3 months	3.1 ↓	4.8	-	1.4	4.1	-
Had genital discharge in last 12 months	5.3* ↓	9.8	6.6	4.1	9.1	4.3
Had genital ulcer in last 12 months	1.8 ↓	2.1	2.9	2.2	3.2	1.5

***p < 0.001

Table 17: Females STI Incidence YR 2017 vs YR 2012 vs YR 2008

	FEMALES 15-24 YRS			FEMALES 25-49 YRS		
	YR 2017 (n=255) %	YR 2012 (n=315) %	YR 2008 (n=299) %	YR 2017 (n=614) %	YR 2012 (n=477) %	YR 2008 (n=367) %
Ever had an STI	8.2 ↓	10.5	14.4**	10.9	12.2	17.2
Had genital discharge in past 3 months	8.6* ↓	12.7	-	7.2	7.8	-
Had genital discharge in last 12 months	12.2* ↓	25.7***	18.1***	7.0	17.7	16.1
Had genital ulcer in last 12 months	2.7 ↑	2.5	2.0	1.5	1.5	1.4

** p<0.005;*** p< 0.001

Persons reporting being diagnosed with an STI were significantly more likely to also have reported multiple partners in the last 12 months, as also transactional sex. Both these risk factors showed an increase over 2012, although not significantly so. (*Table 19*).

Table 18: Risky Partnerships Reported by Persons Reporting an STI; YR 2017 vs YR 2012

	2017	2012
Had multiple partners in last 12 months; (n=630)	17.3***	14.8
No multiple partners; (n=859)	9.7	11.0
Had transactional sex in last 12 months; (n=622)	18.8***	16.0
No transactional sex; (n=1129)	7.9	9.1
Had casual partners in last 12 months; (n=432)	15.7	14.0
No casual partners; (n=1319)	10.5	11.1

***p<0.001

KNOWING ONE'S HIV STATUS

Among the total sample it is 55% who have done an HIV test in the past year. Among sexually active, persons from the youngest cohort, 15-24 years were the most likely to have done the test (63.5%, *Table 20*). The youngest age cohort was the most likely to have done an HIV test in the past year (*Table 20*).

Among the higher risk cohort of inconsistent condoms users with multiple partners (condom non-users and sometimes users), it is 46.7% and 28.6% respectively who both did an HIV test and returned for the results (*Table 21*).

Table 19: Had an HIV Test in Past Year

	Total (n=1201) %	15-24 yrs. (n=386) %	25-49 yrs. (n=815) %	Male (n=451) %	Female (n=750) %	Sexually Active (n=1179) %	Not Sexually Active (n=22) %
Yes	54.8	63.5	50.7	53.2	55.7	54.5	68.2
No	44.4	35.5	48.6	45.9	43.5	44.7	27.3
No Answer	0.8	1.0	0.7	0.9	0.8	0.8	4.5

*Base= Persons who have ever done an HIV test

Table 20: HIV testing by Condom usage of those engaging in Multiple Partnerships in last year YR 2017 vs YR 2012

	YR 2017 Non-user (n=107) %	YR 2012 Non-user (n=53) %	YR 2017 Sometimes User (n=203) %	YR 2012 Sometimes User (n=176) %	YR 2017 Most times user (n=73) %	YR 2012 Most times user (n=316) %
Ever had an HIV test done	69.2 ↓	71.7	57.1	60.2	52.1	50.6
HIV test done in last year and know the results	46.7	50.9	28.6	35.2	30.1	32.3

Reasons for having done an HIV test were largely personal (55%) or medical / pregnancy related (37.4%). Males (70.7%) and persons 15-24 years (60.4%) were more likely to have done the test for personal reasons.

Table 21: Reason for doing HIV Test by Age, Sex and level of Sexual Activity

Demographics							
	Total (n=1201) %	15-24 yrs. (n=386) %	25-49 yrs. (n=815) %	Male (n=451) %	Female (n=750) %	Sexually Active (n=1179) %	Not Sexually Active (n=22) %
Personal Decision	55.0	60.4	52.4	70.7	45.5	55.0	54.5
Had Baby	22.2	19.4	23.5	0.7	35.2	22.6	-
Medical Reasons	15.2	12.4	16.4	14.9	8.2	15.0	27..2
Employment	2.4	1.8	2.7	4.2	1.3	2.5	-
To donate blood	1.8	2.1	1.7	4.4	0.3	1.8	4.5
Immigration/ Farm Work	1.4	0.8	1.7	2.4	0.8	1.4	-
For Insurance	0.7	0.3	1	0.4	0.9	0.8	-
Other Reason	0.6	0.5	0.6	1.3	0.1	0.6	-
Part of regular check up	0.3	0.3	0.4	0.2	0.4	0.3	4.5
For School	0.6	1.6	0.1	0.7	0.5	0.4	9.1
Wanted to know Status	0.2	0.3	0.2	0.2	0.3	0.3	-
Partner/ Family suggested it	0.8	1.6	0.5	1.8	0.3	0.8	-
No Answer	0.7	0.8	0.7	0.2	1.1	0.8	-

*Base= Persons who have ever done an HIV test

Table 22: Know Result of HIV Test done in Past Year

	Total (n=1201) %	15-24 yrs. (n=386) %	25-49 yrs. (n=815) %	Male (n=451) %	Female (n=750) %	S/Active (n=1179) %	Not S/Active (n=22) %
Yes	96.8	96.6	96.8	96.2	97.1	96.9	86.4
No	2.5	2.6	2.6	2.7	2.4	2.3	13.6
D/K	0.7	0.8	0.6	1.1	0.5	0.8	-

**Base=Persons who have ever done an HIV test*

Table 23: Reasons for NOT doing an HIV Test

	Demographics						
	Total (n=799) %	15-24 yrs. (n=614) %	25-49 yrs. (n=185) %	Male (n=557) %	Female (n=242) %	S A (n=572) %	Not SA (n=227) %
Not at Risk	24.5	24.1	25.9	23.3	27.3	23.6	26.9
Sure of HIV- status (I know I don't have it)	20.8	19.4	25.4	23.2	15.3	25.2	9.7
Never had sex/not SA	15.9	19.5	3.8	10.1	29.3		54.6
No specific reason/ Never thought to do one	12.0	11.4	14.1	12.9	9.9	14.9	4.8
Don't want to know status	9.3	9	10.3	10.8	5.8	11.7	3.1
Fear of needles	5.9	5.5	7	5.2	7.4	7.3	2.2
Not sure where to do one	3.9	4.1	3.2	3.8	4.1	4.4	2.6
Have not gotten a chance to do it/ No time	2.8	2.4	3.8	3.2	1.7	3.8	-
No sign of infection or sickness	1.5	1.5	1.6	2.2	-	1.9	0.4
Partner always does the test/ Do test when pregnant	1.1	0.5	3.2	1.6	-	1.6	-
Always uses protection	0.8	1.0	-	0.9	0.4	1.0	-
Can't afford it	0.3	0.2	0.5	0.2	0.4	0.3	-
Don't know how to go about getting one done	0.6	0.7	0.5	0.8	0.4	0.9	-
Always do other blood tests	0.1	-	0.5	-	0.4	0.2	-
Have only one partner/ Trust Partner	0.6	0.5	1.1	0.5	0.8	0.9	-
Not necessary	1.4	1.3	1.6	1.8	0.4	1.7	0.4
Don't know	3.0	3.2	2.2	3.4	2.0	3.8	1.3

**Base= Persons who have never done an HIV test*

***percentages exceed 100% due to multiple response*

WORKSHOPS ON HIV /AIDS IN PREVIOUS 12 MONTHS

It was a quarter of the population (25.8%) that had attended workshops on HIV/AIDS in the previous 12 months. These were mainly among persons 15-24 years with workshops being held mainly at schools. (Tables 25 & 26).

Table 24: Participated in Workshop on HIV/ AIDS in Past Year

	Demographics						
	Total (n=2000) %	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %	S/A (n=1179) %	Not S/A (n=22) %
Yes	25.8	31.4	20.2	23.5	28.1	23.7	40.2
No	74.1	68.5	79.7	76.3	71.9	76.2	59.4
Don't Know/ No Answer	0.1	0.1	0.1	0.2	-	0.1	0.4

Table 25: Where participated in workshop on HIV/ AIDS in Past Year

	Demographics						
	Total (n=516) %	15-24 yrs. (n=315) %	25-49 yrs. (n=201) %	Male (n=238) %	Female (n=278) %	S/A (n=417) %	Not S/A (n=99) %
School	44.2	64.1	12.9	50.8	38.5	35.0	82.8
Clinic	26	15.2	42.8	8.8	40.6	31.2	4
Church	8.3	7	10.4	8	8.6	8.6	7.1
In their community	12.6	8.6	18.9	18.5	7.6	13.4	9.1
Youth Club	3.9	4.4	3	3.8	4	3.1	7.1
Sports Club	1.2	10.0	1.5	2.5	-	1.4	-
On the Job	8.3	2.9	16.9	10.9	6.1	10.3	-
Hotel	1	0.6	1.5	0.4	1.4	1	1
Other	4.1	4.1	4	5.5	2.9	4.6	2.0
In another community	0.4	0.6	-	0.4	0.4	0.5	-
Hospital	0.8	0.6	1	0.8	0.7	1	-
No Answer	1.2	1	1.5	0.8	1.4	1.4	-

***percentages exceed 100% due to multiple response*

CHAPTER 2: BEHAVIOUR BY MARITAL STATUS

Fewer women reported living with their partner for <1 year compared to 2012 (*Table 27*).

Table 26: 2017 Length of Time Living with Married/Cohabiting Partner by Age & Gender

	2012		2017			
	<1yr	< 1yr	One yr.	2 to 4 yrs.	5 to 9 yrs.	9+yrs
Male (n=177)	7.8	15.3 ↑	7.9	31.1	25.4	19.8
Female (n=360)	8.9	7.5 ↓	8.9	21.9	25.8	35.3
15-24yrs (n=101)	19.8	21.8 ↑	18.8	38.6	19.8	1.0
25-49yrs (n=436)	6.1	7.3 ↑	6.2	21.8	27.1	36.9
Male (15-24yrs); (n=34)	22.7	35.3 ↑	17.6	35.3	11.8	0.0
Female (15-24yrs); (n=67)	18.6	14.9 ↓	19.4	40.3***	23.9	1.5
Male (25-49yrs); (n=143)	6.0	10.5 ↑	5.6	30.1***	28.7	24.5
Female (25-49yrs); (n=293)	6.1	5.8 ↓	6.5	17.7	26.3	43.0***
TOTAL	16.7	22.8	16.8	53.0	51.2	55.1

***p<0.001

Although the length of the sexual relationships among those who are now married or living together is less risky, in that more have been in a sexual relationship for more than 1 year, we note that the incidence of never using a condom in these relationships is increasing (*Table 28*).

Supporting this is the finding that the incidence of last time condom use declined among every group except females, 15-24 years (*Table 29*). In fact, the decline among men and persons 25-49 years is statistically significant.

Table 27: 2017 Frequency of Condom Use with Married/Cohabiting by Age & Gender

	2017				2012
	Every time	Most times	Occasionally	Never use	Never
Male (n=177)	4.0	18.1	33.3	42.4 ↑	39.0
Female (n=360)	8.9	17.5	35.0	38.3 ↑	35.2
15-24yrs (n=101)	7.9	23.8	32.7	35.6 ↑	29.3
25-49yrs (n=436)	7.1	16.3	34.9	40.6 ↑	38.4
Male (15-24yrs); (n=34)	5.9	26.5	23.5	44.1 ↑	22.7
Female (15-24yrs); (n=67)	9.0	22.4	37.3	31.3	31.7
Male (25-49yrs); (n=143)	3.5	16.1	35.7	42.0 ↑	41.0
Female (25-49yrs); (n=293)	8.9	16.4	34.5	39.9 ↑	36.2

Table 28: Condom Used at Last Sex with Married/Cohabiting by Age & Gender YR 2017 VS YR 2012

Condom use at last sex with married/live in partner				
	YEAR 2017		YEAR 2012	
Male	↓ 15.3*	(n=177)	24.5	(n=204)
Female	↓ 22.5	(n=360)	26.5	(n=272)
15-24yrs	↓ 22.8	(n=101)	25.6	(n=82)
25-49yrs	↓ 19.5*	(n=436)	25.6	(n=394)
Male (15-24yrs)	↓ 20.6	(n=34)	40.9	(n=22)
Female (15-24yrs)	23.9	(n=67)	20.2	(n=60)
Male (25-49yrs)	↓ 14.0	(n=143)	22.5	(n=182)
Female (25-49yrs)	↓ 22.2	(n=293)	28.3	(n=212)

***p<0.001

Another type of sexual practice that contributes to risk in a married/cohabiting relationship is the practice of having other partners simultaneously. Unfortunately, this incidence increased among men and women overall in the 2017 survey (*Table 30*).

**Table 29: Multiple Partnerships within Married/Cohabiting Unions by Age & Gender
YR 2017 VS YR 2012**

Multiple Partnerships				
	YEAR 2017		YEAR 2012	
Male	↑ 40.7***	(n=177)	35.5	(n=203)
Female	↑ 11.4	(n=359)	5.5	(n=274)
Male (15-24yrs)	↑ 47.1	(n=34)	31.8	(n=22)
Female (15-24yrs)	↑ 14.9	(n=67)	5.0	(n=60)
Male (25-49yrs)	↑ 39.2	(n=143)	35.9	(n=181)
Female (25-49yrs)	↑ 10.6	(n=293)	5.6	(n=214)

** p< 0.005; ***p<0.001

**Table 30: Condom Use Every Time & Last time with Main Partner Where Multiple Partners Exist YR
2017 VS 2012**

SAMPLE SIZE BY DEMOGRAPHICS FOR CONDOM USE		Incidence of Condom Use Every time		Incidence of condom Use Last Time	
	YR 2017	YR 2017	YR 2012	YR 2017	YR 2012
Male	(n=72)	5.6	9.7	18.1	22.2
Female	(n=41)	9.8	20.0	31.7	33.3
15-24yrs	(n=26)	3.8	10.0	23.1	20.0
25-49yrs	(n=87)	8.0	11.7	23.0	25.0

**Table 31: Condom Use with Multiple Partners other than Married/Cohabiting by Age & Gender
YR 2017 VS YR 2012**

Incidence of Condom Use in Multiple Partnerships				
	YR 2017		YR 2012	
Male	65.3	(n=72)	79.2***	(n=72)
Female	41.5	(n=41)	42.9	(n=14)
15-24yrs	61.5	(n=26)	77.8***	(n=9)
25-49 yrs.	55.2	(n=87)	72.7	(n=77)
Male (15-24yrs)	81.2	(n=16)	85.7***	(n=7)
Female (15-24yrs)	30.0	(n=10)	50.0	(n=2)
Male (25-49yrs)	60.7	(n=56)	78.5*	(n=65)
Female (25-49yrs)	45.2	(n=31)	41.7	(n=12)

*p< 0.05; ***p<0.001

Sexual activity in the previous 12 months among those who were unmarried/non-cohabiting, declined or remained fairly stable. Declines versus 2012 were significant among females and in particular, those 15-24 years (Table 33).

Table 32: Sexually active Visiting/Casual/Non-Cohabiting by Age & Gender YR 2017 VS YR 2012

<i>Sexually active and unmarried/cohabiting</i>		
	YR 2017	YR 2012
Male	58.8 (n=1008)	57.1** (n=902)
Female	36.2 * (n=992)	42.5 (n=898)
15-24yrs	55.1(n=1000)	57.7 (n=871)
25-49 yrs.	40.0 (n=1000)	42.4 (n=929)
Male (15-24yrs)	63.8*** (n=629)	64.0 (n=455)
Female (15-24yrs)	40.6* (n=372)	51.0 (n=416)
Male (25-49yrs)	50.7*** (n=379)	50.1* (n=447)
Female (25-49yrs)	33.5 (n=620)	35.3 (n=482)

* p< 0.05; **p< 0.005; ***p<0.001

The incidence of persons who were not married or co-habiting but who had a main partner, declined among males and persons 15-24 years but increased among females in particular those 25-49 years.

Table 33: Sexually active Having a Main Partner by Age & Gender YR 2017 VS YR 2012

<i>Proportion with Main Partner</i>		
	YR 2017 %	YR 2012 %
Male	↓ 62.7*** (n=593)	66.5 (n=507)
Female	↑ 85.2 (n=359)	82.4 (n=374)
15-24yrs	↓ 69.0 (n=552)	74.8 (n=497)
25-49 yrs.	↑ 74.2 (n=400)	71.1 (n=384)
Male (15-24yrs)	↓ 63.8 (n=401)	68.4 (n=288)
Female (15-24yrs)	↑ 82.8 (n=151)	83.7 (n=209)
Male (25-49yrs)	↓ 60.4 (n=192)	63.9 (n=219)
Female (25-49yrs)	↑ 87.0 (n=208)	80.6 (n=165)

***p<0.001

Among those of that cohort who had a main partner, condom use at last sex declined (*Table 35*). The majority of respondents reported to using condoms 'Most times' when engaging in sex (*Table 36*).

Table 34: Condom Use at Last Sex (With Main Partner) by Age & Gender 2017 VS 2012

<i>Condom use at last sex with Main Partner</i>		
	<i>%</i>	
	YR 2017	YR 2012
Male	↓ 59.5*** (n=395)	64.0*** (n=333)
Female	↓ 37.5 (n=315)	47.1 (n=306)
15-24yrs	↓ 57.4*** (n=399)	62.9 (n=369)
25-49yrs	↓ 39.9 (n=311)	46.3 (n=270)
Male (15-24yrs)	↓ 63.2 (n=269)	72.8 (n=195)
Female (15-24yrs)	↓ 45.4 (n=130)	51.7 (n=174)
Male (25-49yrs)	51.6 (n=126)	51.4 (n=138)
Female (25-49yrs)	↓ 31.9 (n=185)	40.9 (n=132)

***p<0.001

Table 35: 2017 Frequency of Condom Use with Main Partner by Age & Gender

	Every time	Most times	Occasionally	Never
Male (n=389)	0.3	45.5	21.9	32.4
Female (n=309)	0.0	50.8	29.4	19.7
15-24yrs (n=389)	0.3	47.6	22.9	29.3
25-49yrs (n=309)	0.0	48.2	28.2	23.6
Male (15-24yrs); (n=264)	0.4	47.0	19.7	33.0
Female (15-24yrs); (n=125)	0.0	48.8	29.6	21.6
Male (25-49yrs); (n=125)	0.0	42.4	26.4	31.2
Female (25-49yrs); (n=184)	0.0	52.2	29.3	18.5

In respect of sexual partner behavior, multiple partnerships declined marginally among females but basically held firm among males in the 2017 survey. Most notable within the subgroups was an increase among the 15-24 years male cohort while the most notable decrease was among the 25-49 years cohort. (Table 37).

Table 36: 2017 Incidence of Multiple Partnerships (with or without main partner)

	2012 Had multiple partners %	2017 Had multiple partners %	2017 No main partner %
Male	70.1	71.0 (n=593)	74.1 (n=220)
Female	28.5	↓ 26.7 (n=359)	37.7 (n=53)
15-24yrs	58.1	↑ 65.3 (n=551)	71.6 (n=169)
25-49yrs	45.6	↓ 39.2 (n=401)	59.6 (n=104)
Male (15-24yrs)	73.0	↑ 75.8 (n=400)	76.2 (n=143)
Female (15-24yrs)	37.1	37.3 (n=151)	46.2 (n=26)
Male (25-49yrs)	66.2	↓ 61.1 (n=193)	70.1 (n=77)
Female (25-49yrs)	17.5	18.8 (n=208)	29.6 (n=27)

Table 37: 2017 Incidence of Condom Use at Last Sex in Multiple Partnerships

	2012 Condom used last time (Have main partner) %	2017 Condom use last time (Have main partner) %	2012 Condom used last time (No main partner) %	2017 Condom Use last time (No main partner) %
Male	72.1 (n=215)	↓ 63.3 (n=258)	78.3 (n=129)	↓ 77.9 (n=163)
Female	45.6 (n=79)	43.4 (n=76)	63.6 (n=22)	↓ 40.0 (n=20)
15-24yrs	70.1 (n=194)	↓ 62.8 (n=239)	79.5 (n=83)	↓ 71.1 (n=121)
25-49yrs	55.0 (n=100)	↓ 49.5 (n=95)	72.1 (n=68)	↑ 79.0 (n=62)
Male (15-24yrs)	78.7 (n=136)	↓ 66.5 (n=194)	82.4 (n=68)	↓ 75.2 (n=109)
Female (15-24yrs)	50.0 (n=58)	↓ 46.7 (n=45)	66.7 (n=15)	↓ 33.3 (n=12)
Male (25-49yrs)	60.8 (n=79)	↓ 54.7 (n=64)	73.8 (n=61)	↑ 83.3 (n=54)
Female (25-49yrs)	33.3 (n=21)	↑ 38.7 (n=31)	57.1 (n=7)	↓ 50.0 (n=8)

While fewer females had multiple partners, the mean number of partners increased in every group (Table 39).


Table 38: Mean number of partners in Visiting/Casual/ Relationships in the past 12 months by Age & Gender 2012 vs. 2017

	2017			2012	
Male	5.50	↑	(n=592)	4.82	(n=495)
Female	3.37	↑	(n=359)	1.36	(n=363)
15-24yrs	5.87	↑	(n=551)	3.86	(n=481)
25-49yrs	3.07	↑	(n=400)	2.72	(n=377)
Male (15-24yrs)	6.46	↑	(n=400)	5.53	(n=279)
Female (15-24yrs)	4.32	↑	(n=151)	1.55	(n=202)
Male (25-49yrs)	3.49		(n=192)	3.91	(n=216)
Female (25-49yrs)	2.67	↑	(n=208)	1.12	(n=161)

CHAPTER 3: ACCESS & ATTITUDES TO CONDOMS

Most persons had no problems accessing condoms readily. Access to condoms was higher among the male population. (Chart 1).

Chart 1: Percentage able to Obtain Condoms Almost Immediately YR 2017



The shop and clinic significantly increased as access points for condoms (33.3% in 2017 versus 28.7% in 2012). The shop was specifically favoured for those aged 15-24 years while the Pharmacy was the access point for the 25-49 year old cohort (Table 40a).

Table 39a: Where condoms usually bought by Age YR 2017

Source	2012	2017	15-24yrs (n=746) %	25-49yrs (n=967) %
	Total (n=1575) %	Total (n=1713) %		
Shop	28.7	33.3*	41.2	27.3
Pharmacy	24.3	17.9	12.7	21.9
Clinic/don't buy	9.2	14.0*	15	13.2
Partner provides it	13.9	9.2	9.5	9
Supermarket	12.0	13.7	9.1	17.2
Gas Station	2.2	4.6	4.4	4.8
Wholesale	2.0	3.3	3.5	3.1
Bar	.9	1.6	1.9	1.3
Anywhere	0.3	0.1	0.1	0.1
Vendor	0.1	0.5	0.5	0.5

*P <.05

The shop was the place of choice not only for the younger, 15-24 years cohort, but specifically for males while the pharmacy or clinic were the important access points for the women (*Table 40b*).

Table 40b: Where condoms usually bought by Gender YR 2017

Source	2012	2017		
	Total (n=1575) %	Total (n=1713) %	Males (n=867) %	Females (n=846) %
Shop	28.0	↑ 33.3*	48.4	17.9
Pharmacy	24.3	17.9	14.8	21.2
Clinic/don't buy	9.2	↑ 14.0*	6.8	21.4
Partner provides it	13.9	9.2	1.3	17.4
Supermarket	12.	13.	12.9	14.4
Gas Station	2.2	4.	6.7	2.5
Wholesale	2.0	3.	3.9	2.6
Bar	0.9	1.	2.5	0.6
Anywhere	0.3	0.	0.2	0
Vendor	0.1	0.	0.7	0.4

The proportion of persons who never have a condom readily available on them, significantly increased in 2017 to 38.6% from 31.2% in 2012. (*Table 41a*). It was the older age group, 25-49 years, and women, that were most likely to never have a condom on them (*Tables 41a & 41b*).

The extent to which persons were likely to have a condom available in the house however showed negligible change from 2012. (*Table 41a & 41b*).

Table 41a: Extent to Which Persons Generally Have a Condom by Gender YR 2017

Source	2012	2017		
	Total (n=1575) %	Total (n=1714) %	Males (n=868) %	Females (n=846) %
Extent to which usually have a condom on you				
- Every time	24.7	22.9	30.6	14.9
- Most times	17.5	13.7	18.3	8.9
- Sometimes	13.6	14.6	19.6	9.5
- Rarely	10.5	10.3	10.6	10.0
- Never	31.2	↑ 38.6*	20.9	56.7
Extent to which usually have a condom in the house:				
- Every time	55.2	55.4	65.0	45.5
- Most times	12.4	12.1	12.4	11.7
- Sometimes	12.3	10.3	8.8	11.8
- Rarely	5.0	5.3	4.1	6.4
- Never	14.7	16.9	9.6	24.5

Table 41b: Extent to Which Persons Usually Have a Condom by Age YR 2017

Source	Total (n=1714) %	15-24yrs (n=747) %	25- 49yrs (n=967)
Extent to which usually have a condom on self:			
- Every time	22.9	24.4	21.7
- Most times	13.7	17.9	10.3
- Sometimes	14.6	19.5	10.8
- Rarely	10.3	9.1	11.3
- Never	38.6	29.0	45.9
Extent to which usually have a condom in the house:			
- Every time	55.4	57.4	53.8
- Most times	12.1	12.4	11.8
- Sometimes	10.3	10.3	10.2
- Rarely	5.3	4.4	5.9
- Never	16.9	15.1	18.3

The fact of loving or trusting their partner (51%) was the main reason for not using condoms during all of the last 10 times sex was had. These numbers show an increase compared to the 2012 results (46.9%). (Table 42).

The proportion indicating they do not like using a condom also significantly increased in 2017 to 20% from 14.7% in 2012. This was primarily so among the 15-19 years cohort (34.1% compared to 19.4% and 18.7% for the 20-24 and 25-49 years groups respectively). (Table 42)

Table 42: Reasons for not using condoms all of the last 10 times sex had YR 2017

	% of Respondents 2017 (n=601)	% of Respondents 2012 (n=1024)	15-19 yr. old (n=44) %	20-24 yr. old (n=124) %	25-49 yr. old (n=433) %
Love/trust partner	↑ 51.0	46.9	31.8	42.7	54.5
Don't like using condom	20.0*	14.7	34.1	19.4	18.7
Partner does not like	11.8	8.0	2.3	9.7	13.4
Allergic to	10.3	3.9	-	10.5	11.3
Only one	5.0	0.9	2.3	3.2	5.5
No particular reasons/No	4.7	.5	6.8	6.5	3.0
Trying to have a child	3.7	3.8	2.3	4.0	3.7
Other	3.0	1.1	6.8	4.0	2.3
No answer	2.5	3	6.8	2.4	2.1
Get regular check-ups; no	1.3	-	0.0	1.6	1.4
Did not have one on me	↓ 1.5	15.1	2.3	4.0	0.2
Did HIV test/Get check-	0.8	1.3	4.5	0.8	0.5
Partner gets regular	0.7	-	-	0.8	0.7
Now pregnant/Didn't	0.7	0.6	-	0.8	0.7
Pressured/Coerced by	0.5	0.5	-	-	0.7
Partner/self on other	0.3	-	-	0.8	0.2
Don't know how to use a	0.2	0.5	-	0.8	-
Can't get pregnant again	0.2	-	-	-	0.2
Partner will think I am	0.2	-	-	0.8	-
No money to buy	0.2	-	2.3	-	-

Overall attitudes to condom use was mixed. While most (69.9%) agreed that sex could be enjoyed even when using a condom, at least a third thought that condoms negatively impacted the experience by reducing the quality and spontaneity. A third also felt condoms were not necessary where the partner was trusted. (Table 43).

Males were more likely to agree that: condoms “reduce the quality of sex so it is better to not use at all” (43.7%), condoms reduced the spontaneity and heat (38%) and were unnecessary when the partner was trusted (38.8%).(Table 43).

Irrespective of how they felt about using condoms, the majority reported having ready access to condoms. (Table 43)

Table 40: General Attitudes to Condoms and Contraceptives

Strongly Agree/ Agree					
	Total (n=1751*) %	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %
Attitude to condom use					
A condom reduces the quality of sex so much that it is better not to use it at all	36.2	40.3	32.9	43.7	28.6
It is possible to enjoy sex when you use a condom	69.9	71.6	68.7	69.5	70.4
Contraceptives ruin the spontaneity and heat of sexual intercourse	32.7	32.3	32.9	38.0	27.3
It is not necessary to use condoms if you trust the person	35.9	33.9	37.5	38.8	33
Access to condoms					
I can get a condom around here easily during the night	87.2	72.2	72	79.7	64.4
I can get a condom around here easily during the day	87.2	87.4	87.2	93.3	81.1
Free condoms distributed by the Ministry of Health are easy to get when I need them	57	52.5	60.4	52.1	61.8
Shops nearby here always have condoms for sale	79.7	80.0	79.4	85.8	73.4
Condom brands and quality					
Condoms you get for free are just as good as the ones available in the shops	50.8	42.6	57.1	46.1	55.6
The brand of condom really does not matter to me	43.6	42.1	44.8	43.1	44.1
Some condoms are better than others	77.7	77.7	77.7	85.1	70.2

*BASE; Persons who are sexually active & who responded

CHAPTER 4: KNOWLEDGE

Overall 40.2% of respondents, 15-49 years, were able to provide correct answers to five statements of preventive practices and myths relating to HIV transmission. These were 100% condom use and one faithful partner as preventative methods and the rejection of myths about transmission (sharing food, healthy looking person and mosquito bites). Correct knowledge was generally higher among females and in particular among females 20yrs and older. (*Chart 2*)

Among persons 15-24 years it was 32.9% who were able to correctly identify both ways of preventing transmission while rejecting major misconceptions.

Chart 2: Correct knowledge of all 5 (*protective factors and myths*)




Table 44. Details Correct Knowledge of Protective Factors & Myths

	Correctly Endorsing preventive measures		Rejecting misconceptions about transmission			
	100% condom use %	One faithful partner %	A healthy looking person %	Mosquito Bites %	Sharing food %	Correctly answering all 5 %
Males 15-19 (N=370)	82.7	74.1	94.1	57.0	66.5	28.1
Females 15-19 (N=194)	74.2	73.2	95.9	63.4	76.3	29.4
Males 20-24 (N=258)	88.0	83.7	98.8	58.9	70.5	35.3
Females 20-24 (N=178)	82.0	82.6	96.6	68.5	79.8	43.3
Males 25-49 (N=380)	88.9	85.8	98.4	60.5	75.3	42.9
Females 25-49 (N=620)	88.2	84.8	98.1	70.3	84.5	50.3
Males 15-49 (N=1008)	86.4	81.0	96.9	58.8	70.8	35.5
Females 15-49 (N=992)	84.4	82.2	97.4	68.6	82.1	45.0
TOTAL Sample	85.4	81.6	97.2	63.7	76.4	40.2

Traditional media and the internet were the top two sources for information in the previous 12 months. Traditional media comprising television, radio and print was used by approximately two-thirds of the sample (65.1%) followed by the internet (45.6%). (Table 45).

Table 45: Where Obtained Information on HIV/ AIDS

<i>In the past 12 months, where have you gotten information about AIDS?...</i>	Total (n=2000) %	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %
Media	65.1	61.8	68.4	61.7	68.5
Internet	45.6	48.4	42.9	43.2	48.2
Doctor/Clinic	34.2	28.3	40.2	23.1	45.6
Friends	26.7	28.1	25.3	26.2	27.2
Home/Family	24.4	28.5	20.4	22.5	26.4
School	22.6	38.5	6.6	23.4	21.7
Community	21.8	22.5	21	25	18.4
Church	9.4	8.7	10	7.3	11.4
Workplace	8.7	5.5	11.9	10.4	7.0
Other	5.8	4.2	7.1	0	13.6

Table 46: Spoken about AIDS and Safe Sex

<i>Spoken about AIDS?...</i>	Total (n=2000) %	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %
Most Recent Partner	56.6	42.7	70.6	49.9	63.4
The Previous Partner	46.2	36.8	55.6	45.1	47.2
Family Members	51.2	45.3	57.2	44.5	58.0
Friends	61.6	56.6	66.6	60.8	62.4
Someone at church	18.0	12.5	23.6	14.4	21.8

CHAPTER 5: STIGMA & DISCRIMINATION

Note: The 2012 percentages for accepting attitudes to PLWAs have been recalculated to facilitate comparison with 2017. In 2012, 4 statements were identified as comprising this indicator but was reduced in 2017 to 3 statements. The statement which has now been removed is “Do you think children living with HIV should be able to attend school...”

Accepting attitudes to PLWA not only remained low but declined significantly in 2017 versus 2012. The main area of concern remains a willingness to purchase fresh vegetables from a vendor known to be HIV+. (Table 47).

Table 47: Attitudes to People Living with HIV/AIDS by Gender YR 2017

<i>Top 2 Box: Strongly Agree & Agree</i>				
	YR 2017 Male (N=1000) %	YR 2017 Female (N=1000) %	Total 2017 (N=2000) %	Total 2012 (N=1800) %
Accepting attitudes to PLWA (3 components)	10.3	12.9	↓ 11.6 *	14.3
- I would definitely purchase food/veg. from a shopkeeper who have HIV				
- Should a teacher who have HIV but not sick be allowed to continue teaching in school?				
- Would want family member to keep being infected with HIV a secret				
Willing to care for a family member who becomes sick with the AIDS virus	79.2	78.5	78.8	81.8
Willing to buy fresh vegetables from a vendor whom they knew was HIV+	21.3	26.2	23.7	28.9
Agree that a teacher who is HIV+ but not sick should be allowed to continue teaching in school	68	78	72.8	76.3
Agree that they would not want to keep the HIV status of a family member a secret	29.2	33	31.1	31.3
Persons who get AIDS have gotten what they deserve	9.9	4.1	7.1	6.3
When a person contracts AIDS they let their family down	24.3	16.5	20.5	23.4
Attitudes to disclosure of HIV status:				
- Status should be disclosed to best friend	47.8	35.6	41.8	42.7
- Status should be disclosed to partner	78.5	84.2	81.3	81.7
- Status should be disclosed to parent	83.5	85.6	84.6	82.9
- Status should be disclosed to church member	41.6	33.6	37.6	-
- Status should be disclosed to co-workers	36.3	27.8	32.1	33.7
- Status should be disclosed the employer/boss	46.3	40	43.2	39.6

Table 48: Attitudes to People Living with HIV/AIDS by Age YR 2017

	Age of Respondent		Total Sample	
	15-24 yrs. (N=1000) %	25-49 y. (N=1000) %	YR 2017 (N=2000) %	YR 2012 (N=1800) %
<i>Accepting attitudes to PLWA (3 components)</i>	8.7	14.5	11.6	14.3
- I would definitely purchase food/veg. from a shopkeeper who have HIV				
- Should a teacher who have HIV but not sick be allowed to continue teaching in school?				
- Would want family member to keep being infected with HIV a secret				
<u>Top 2 Box: Strongly Agree & Agree</u>				
Willing to care for a family member who becomes sick with the AIDS virus	78.2	79.5	78.8	81.8
Willing to buy fresh vegetables from a vendor whom they knew was HIV+	18.9	28.4	23.7	28.9
Agree that a teacher who is HIV+ but not sick should be allowed to continue teaching in school	65.2	80.6	72.8	76.3
Agree that they would not want to keep the HIV status of a family member a secret	32.9	29.3	31.1	31.3
Persons who get AIDS have gotten what they deserve	9.8	4.3	7.1	6.3
When a person contracts AIDS they let their family down	26.2	14.7	20.5	23.4
<u>Attitudes to disclosure of HIV status:</u>				
– Status should be disclosed to best friend	40.9	42.6	41.8	42.7
– Status should be disclosed to partner	78.6	84.0	81.3	81.7
– Status should be disclosed to parent	84.8	84.3	84.6	82.9
– Status should be disclosed to church member	34.1	41.1	37.6	-
– Status should be disclosed to co-workers	29.1	35.1	32.1	33.7
– Status should be disclosed the employer/boss	39.3	47.1	43.2	39.6

Just under a half of the sample (48%) knew of someone living with or who has died from AIDS. For about a half of them, this person was a close relative or friend. This further supports the need to foster more accepting attitudes to PLWA to support them and enable them to cope (*Table 49*).


Table 49: Knowledge of a Person living/dying from HIV/ AIDS

	Total (n=2000) %	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %
Yes, a close relative or friend	23.1	15.6	30.6	19.8	26.4
Yes, but not a close relative or friend	24.8	20.5	29.2	26.2	23.5
No, I don't know anyone	48.9	60.6	37.1	50.7	47.1
Not sure/can't recall	3.0	3.0	2.9	3.0	2.9
No Answer	0.2	0.3	0.1	0.3	0.1

Interestingly, when given specific statements to respond to, the responses would suggest that the majority do not ostracize a PLWA. It is the minority (5-12%) who thinks that PLWAs are dirty, cursed, and should be ashamed. It is males and persons of the younger cohort who are most likely to have these opinions (*Table 50*).

Table 50: Attitude to Persons with HIV

Strongly Agree/ Agree					
	Total (n=2000) %	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %
People who have AIDS are dirty	12.0	15.3	8.7	15.7	8.3
People who have AIDS are cursed	5.3	6.9	3.6	7.6	2.9
People who have AIDS should be ashamed	11.8	16.6	7.1	16	7.7
It is not safe for People who have AIDS to work with children	33.5	30.9	36.1	33.1	34
People who have AIDS must expect some restriction on freedom	45.5	49.6	41.4	49.5	41.4
People who have AIDS have done something wrong and deserve to be punished	8.3	10.9	5.6	11.4	5

Chart 3: Perception of Persons with HIV

Table 51: Attitude to Persons with HIV by Age & Gender

YR 2017
% Reporting Strongly Agree/ Agree

	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %
A man who has HIV should not seek to have children	61.2	66.5	61.8	66.0
A woman who has HIV should not seek to have children	60.5	66.1	61.9	64.8
It is a NOT waste of money to train or give a promotion to someone with HIV/AIDS	83.8	86.5	83.0	87.3
A person would be foolish to marry a person who is living with HIV/AIDS	43.4	39.1	46.5	36.0

CHAPTER 6: ADVERTISING MESSAGE RECALL AND IMPACT

Majority of respondents (83.0%) indicated that they had heard or seen advertisements on condom use or multiple partnerships. Similarly, 79.6% reported that they had heard or seen advertisements on human rights or HIV. The main medium through which respondents were exposed to these kinds of ads were televisions (92.1%; *Table 52*).

However, message recall did not necessarily translate into change in behavior, with a third (36.2%) reporting that what they *'saw or heard made them think or behave differently'* (*Table 54*). Main behavior change reported in regards to ads recalled were 'Use protection when having sex' (34.9%) and 'Use a condom every time' (27.1%). Interestingly, men were more likely than their female counterparts to report these behavior changes 'Use protection when having sex' (36.5% male vs. 33.3% females) and 'Use a condom every time' (30.2% males vs. 24.0% females; *Table 54*).

Table 52: Advertising Message Recall (unprompted) YR 2017

	(N=2000) %
Q. Recall seeing of hearing messages or advertising on condom use or multiple partners	83.0
Q. Recall seeing of hearing messages or advertising on human rights or HIV	79.6
Q. Where did you hear the advertisement?	(n=1743)
- Television	92.1
- Radio	29.6
- Internet/ social Media	21.5
- Poster	14.8
- Back of Bus	3.3
- Super board	2.9
- Cinema	0.2

Main messages recalled were ‘Use a condom every time’ (94.7%) and ‘Pinch leave an inch and roll’ (93.7%). Messages with the least recall were ‘Get it got’ recalled by 39.2% and “Choose two: Use dual protection’ (33.6%; *Table 53*).

Table 53: Specific Message recalled (prompted) YR 2017
(Top 12 responses)

	(N=1746) %
Q. What do you recall seeing or hearing?	
Use a condom every time	94.7
Pinch, leave an inch and roll	93.7
Real men nuh ride without condom	79.4
Protect yourself when having sex	78.3
Live up, love, respect	76.2
Know your partner status	74.7
More partners more risk	65.2
Smart women always buy, carry and use condoms	63.1
We are Jamaicans	46.0
Yes, I can support someone living with HIV	42.5
Got it? Get it	39.2
Choose two: Use dual protection	33.6

Just over a third of persons reported having changed their behavior because of the messaging and for 62% of these persons, the change was to practice safe sex and use condoms (*Table 54*).

Table 54: Behaviour Change as a result of Mass Media Campaign YR 2017

	Total (n=1744) %	15-24 yr. old (n=857) %	25-49 yr. old (n=887) %	Male (n=863) %	Female (n=881) %
Q. Did any of the things you saw or heard make you think or behave differently					
Yes	36.2	33.0	39.2	37.7	34.7
Q. In what way did you think or behave differently?					
	Total (n=627) %	15-24 yr. old (n=281) %	25-49 yr. old (n=346) %	Male (n=315) %	Female (n=312) %
Use protection when having sex	34.9	38.8	31.8	36.5	33.3
Use a condom every time	27.1	33.1	22.3	30.2	24.0
Get an HIV test	11.6	8.5	14.2	9.2	14.1
Have only one partner/Reduce number of partners	10.8	5.7	15.0	10.5	11.2
Makes you think about life and steps to take/Be careful	8.3	7.8	8.7	8.9	7.7
Abstain from having sex	7.3	11.4	4.0	4.1	10.6
Ask partner to get an HIV test	6.1	5.0	6.9	4.4	7.7

Persons who reported being impacted by the messages were more likely to be male (37.7% vs 34.7%, females) and older (39.2% of persons 25-49 years vs 33% among those 15-24 years). Reported impact was highest among persons practicing risk behavior. These were persons with multiple partners in last 12 months (41.7%) and persons with high risk partnerships (43%). (*Table 55*).

Table 55: Profiling Persons impacted by message YR 2017

Has impacted behaviour	%
Total (n=1744)	36.2
15-24 yrs. (n=857)	33.0
25-49 yrs. (n=887)	39.2
Male (n=863)	37.7
Female (n=881)	34.7
Ever Had Sex (n=1537)	37.4
Never Had Sex (n=207)	25.1
Married/ Cohabiting (n=480)	36.0
Not married but sexually active (n=835)	39.0
Multiple partners in last 12 months (n=540)	41.7
High Risk Partners (new partner, One night stand, someone met at bar or club) (n=370)	43.0

The Mass Media campaign seems to have resonated most with the older cohorts as well as females. This was deemed to be so as the older age group as well as females were more likely to discuss the message of the campaigns with their friends and partner. (*Table 56*).

Those in high risk and multiple partnerships were more likely to discuss the campaign message with friends than with any of their partners (*Table 56*).

Table 416: Percentage who have discussed message of campaign with sex partner and friends YR 2017

	Discuss message with partner %	Discuss message with friends %
Total (n=1744)	41.6	50.3
15-24 yrs. (n=857)	30.8	48.2
25-49 yrs. (n=887)	52.1	52.4
Male (n=863)	38.4	51.9
Female (n=881)	44.8	48.8
Ever Had Sex (n=1537)	47.1	51.4
Never Had Sex (n=207)	-	42.5
Married/ Cohabiting (n=480)	53.5	51.0
Not married but sexually active (n=835)	47.3	53.9
Multiple partners in last 12 months (n=540)	43.1	50.7
High Risk Partners (new partner, One night stand, someone met at bar or club) (n=370)	45.7	51.9

CHAPTER 7: CHILDBEARING AND ANTENATAL CARE

Approximately a quarter of the females in the sample (24.9%) had either given birth in the last two years or were at least 6 months pregnant. (*Table 57*).

Table 57: Percentage of Females Being Pregnant YR 2017

	REONDENTS (n=872) %
Given Birth in last 2 yrs.	21.8
At least 6 months pregnant	3.1
Neither	75.1

Majority of women who had given birth in the past two years or were at least 6 months pregnant had attended an antenatal clinic (90.4%) and private doctor (17.2%). In fact 7.6% had attended both. All had done HIV tests during their visits and the vast majority had been counselled and received their results (*Table 58*).

Table 58: Percentage of Pregnant Females Making Antenatal Care Visits YR 2017

	REONDENTS (n=217) %
Attended Antenatal Clinic	90.4
Attended Private Doctor	17.2
Counseled about HIV	84.4
Counseled about STI	87.3
HIV Test offered during visits	93.4
HIV Test done during visits	100.0
HIV Test results received	96.5

CHAPTER 8: WORKPLACE INTERVENTION

It was 18.1% of those employed full-time and 6.8% of those employed part-time who recalled receiving information on HIV/AIDS at their workplace during 2017. Among those who had received information, it was just over a quarter (26.4%) of those employed fulltime and 7.5% of those employed part-time who had participated in a workshop accordingly (*Table 59*).

Table 59: The Workplace as a point for disseminating information on HIV/AIDS YR 2017

	Employed Full Time (n=674) %	Employed Part Time (n=278) %
Gotten information on HIV at Workplace	18.1	6.8
	Employed Full Time (n=140) %	Employed Part Time (n=53) %
Participated in workshop on HIV/AIDS/STD on the Job	26.4	7.5

CHAPTER 9: INTIMATE PARTNER VIOLENCE

This indicator looks at the proportion of ever-married or partnered women aged 15-49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months. Results indicate that 14.8% of this cohort of females had experienced physical or sexual violence from a male partner in the previous 12 months.

Most prevalent violent acts experienced were 'Pushed or Shoved you' (17.7%), 'Slapped you/ Threw something that could hurt you' (16.8%) and 'Hit you with a fist/ Something that could hurt you' (15.6%). For the most part, these violent acts were experienced to the same extent despite age differences. (Table 60)

**Table 60: Types of Violence Experienced by Ever Married or Partnered Females in Past 12 Months:
YR 2017**

	Ever Married or Partnered Female in Past 12 Months		
	Total (n=719) %	15-24 yrs. (n=218) %	25-49 yrs. (n=501) %
Pushed or Shoved you	17.7	18.8	17.2
Slapped You/ Threw something that could hurt you	16.8	16.5	17.0
Hit you with a fist/ Something that could hurt you	15.6	15.6	15.6
Kick, drag or beat you up	11.7	10.5	12.2
Choke or burn you	5.8	6.0	5.8
Threaten or use a Knife, Gun or weapon	6.8	7.3	6.6
Forced you to do something degrading/humiliating	3.4	3.7	3.2
Made you afraid of what they would do if you did not have sex with them	3.7	5.1	3.2

In 75% of cases, the aggressor was a male. This was similar for both age cohorts. (*Table 61*).

Table 61: Sex of Aggressor by Age Group & Gender

Gender of Aggressor	Ever Married or partnered Female in Past 12 Months		
	Total	15-24 yrs.	25-49 yrs.
	(n=174) %	(n=56) %	(n=118) %
Male	75.3	75.0	75.4
Female	0.6	0.0	0.8

Majority of violent acts were committed by a partner (63.6%) of the victim (*Table 62*)

Table 62: Person who committed violent acts YR 2017

	Ever Married or partnered Female in Past 12 Months		
	Total	15-24 yrs.	25-49 yrs.
	(n=162) %	(n=54) %	(n=108) %
Boyfriend/Girlfriend	63.6	66.7	62.0
Social acquaintance	5.6	5.6	5.6
Do not know the person	3.1	2.8	3.1
Family/Relative	1.2	1.9	0.9
Other	1.2	0.0	1.9
Not applicable	23.5	20.4	25.0
Declined to answer	1.9	1.9	1.9

Approximately 3 in 10 (28.9%) of females who were either married or been in a partnership and experienced intimate partner violence considered themselves as victims of physical violence. (Table 63).

Majority of these respondents (72.4%) noted that they themselves had not committed any violent acts. The 15-24 year olds were more likely to admit to committing violent acts themselves (28.1% vs 15.8% for the 25-49 year olds). (Table 65).

Table 423: Consider Themselves a Victim of Physical Violence

Ever Married or partnered Female in Past 12 Months			
Response	Total (n=173) %	15-24 yrs. (n=56) %	25-49 yrs. (n=117) %
Yes	28.9	30.4	28.2
No	61.3	58.9	62.4
Not Applicable	8.1	8.9	7.7
Declined to answer	1.7	1.8	1.7

Table 64: Do you feel you deserve it?

Response	Ever Married or partnered Female in Past 12 Months		
	Total (n=50) %	15-24 yrs. (n=17) %	25-49 yrs. (n=33) %
Yes	6.0	5.9	6.1
No	94.0	94.1	93.9

Table 65: Respondent committing violent acts YR 2017

Ever Married or partnered Female in Past 12 Months			
	Total (n=177) %	15-24 yrs. (n=57) %	25-49 yrs. (n=120) %
Yes	26.6	28.1	15.8
No	72.4	70.2	83.3
Declined to answer	1.0	1.8	0.8

EMOTIONAL HEALTH

Females were much more likely than their male counterparts, to report having emotional problems as follows:

- 47% of females compared to 38% of males reported having had little or no interest or pleasure in doing things at some time in the previous two weeks;
- 54% of females compared to 39% of males, felt down, depressed or hopeless at some time in the previous two weeks;
- 35% of females compared to 30.1% of males reported feeling nervous, anxious or on edge, at some time over the previous 2 weeks.
- 44% of females compared to 28% of males indicated they had not been able to stop or control worrying in the previous 2 weeks.
- And it was 13% of females and 7.5% of males who had experienced thoughts of being better off dead or hurting themselves in some way over the previous 12 months.

Table 66: Statements Indicative of Emotional Health

	Not at all	Several days	More than half the days	Nearly every day
<i>Over the past 2 weeks how often have you: Had Little Interest Or Pleasure In Doing Things</i>				
Males 15-19 (N=370)	59.5	24.1	8.4	8.1
Females 15-19 (N=194)	56.7	20.6	7.2	15.5
Males 20-24 (N=258)	58.5	20.9	10.9	8.9
Females 20-24 (N=178)	50.6	27.0	13.5	9.0
Males 25-49 (N=380)	66.1	23.2	5.3	5.5
Females 25-49 (N=620)	53.2	28.3	10.2	8.2
TOTAL MALES (N=1008)	61.7	22.9	7.8	7.3
TOTAL FEMALES (N=992)	53.3	26.6	10.2	9.8

Table 66 (cont'd): Statements Indicative of Emotional Health

<i>Over the past 2 weeks how often have you Felt Down, Depressed Or Hopeless</i>	Not at all	Several days	More than half the days	Nearly every day
Males 15-19 (N=370)	66.5	17.6	7.8	8.1
Females 15-19 (N=194)	45.9	28.4	11.9	13.9
Males 20-24 (N=258)	58.9	28.7	8.9	3.1
Females 20-24 (N=178)	43.8	33.7	11.2	11.2
Males 25-49 (N=380)	56.6	29.7	9.5	3.9
Females 25-49 (N=620)	46.0	32.8	11.6	9.4
TOTAL MALES (N=1008)	60.8	25.0	8.7	5.3
TOTAL FEMALES (N=992)	45.7	32.1	11.6	10.6
<i>Over the past 2 weeks how often have you <u>Felt Nervous, Anxious Or On Edge</u></i>	Not at all	Several days	More than half the days	Nearly every day
Males 15-19 (N=370)	69.5	18.1	8.6	3.8
Females 15-19 (N=194)	59.3	20.1	12.9	7.2
Males 20-24 (N=258)	69.8	19.8	6.2	3.9
Females 20-24 (N=178)	59.6	24.2	9.6	6.7
Males 25-49 (N=380)	70.5	20.3	6.6	2.4
Females 25-49 (N=620)	67.9	8.1	4.2	.3
TOTAL MALES (N=1008)	69.9	19.3	7.2	3.3
TOTAL FEMALES (N=992)	64.7	20.5	9.3	5.2

Table 66 (cont'd): Statements Indicative of Emotional Health

Over the past 2 weeks how often have you <u>Not Been Able To Stop Or Control Worrying</u>	Not at all	Several days	More than half the days	Nearly every day
Males 15-19 (N=370)	77.0	14.3	5.1	3.5
Females 15-19 (N=194)	54.1	25.8	6.2	13.4
Males 20-24 (N=258)	70.2	18.2	7.0	4.3
Females 20-24 (N=178)	53.4	23.6	8.4	14.6
Males 25-49 (N=380)	67.9	22.4	6.3	3.2
Females 25-49 (N=620)	57.2	23.7	9.7	9.0
TOTAL MALES (N=1008)	71.8	18.4	6.1	3.6
TOTAL FEMALES (N=992)	55.8	24.2	8.8	10.9
Over the past 12 months, how often did you have thoughts of <u>being better off dead or hurting self in some way</u>	Not at all %	Several days %	More than half the days %	Nearly every day %
Males 15-19 (N=370)	92.4	5.4	2.2	0
Females 15-19 (N=194)	72.2	22.2	5.7	0
Males 20-24 (N=258)	89.5	8.9	1.2	0
Females 20-24 (N=178)	83.1	12.4	4.5	0
Males 25-49 (N=380)	94.5	3.4	1.8	0
Females 25-49 (N=620)	92.2	5.3	2.4	0
TOTAL MALES (N=1008)	92.5	5.6	1.8	0
TOTAL FEMALES (N=992)	86.7	9.9	3.4	0

CHAPTER 10: GENERAL ATTITUDES

The sample was asked to respond to suggested policies regarding HIV, access to condoms and anal sex. The most endorsed of these policies was to “Develop a policy to guarantee access to HIV testing for sexually active adolescents between ages 10-16’ (91.4%). On the other hand ‘Decriminalizing Anal Sex was the least endorsed (11.6%), this was especially true among males (9.5%). Further to this, the majority were of the view that ‘men who have sex with men are dirty’ (86.6%), ‘men who have sex with men should be ashamed’ (88.8%) and ‘it is not safe for men who have sex with men to work with children’ (83.2%). In all cases men were more likely to strongly agree/agree. (Table 67).

Interesting to note is that respondents appear to be more accepting/tolerant of the transgender individual than MSMs as fewer i.e. 6%-13% strongly agree/agree with transgender statements (Table 67).

Table 67: General Attitudes

	Strongly Agree/ Agree				
	Total (n=2000) %	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %
Attitudes to Policies regarding HIV, Access to Condom and Anal Sex					
Develop a policy to guarantee access to HIV testing for sexually active adolescents between ages 10-16	91.4	90.5	92.3	90.2	92.6
Develop a policy to guarantee access to condoms to sexually active adolescents aged 10-16	77.3	77.7	76.8	77.4	77.2
Decriminalize Anal Sex	11.6	11.2	11.8	9.5	13.6
Attitudes towards Homosexuality					
Men who have sex with men are dirty	86.6	87.5	85.7	91.9	81.2
Men who have sex with men are cursed	66.5	66.2	66.7	76.2	56.6
Men who have sex with men should be ashamed	88.8	88.6	89	92.7	84.7
It is not safe for Men who have sex with men to work with children	83.2	80.9	85.5	85	81.4
Men who have sex with men are doing something wrong and deserve to be punished	69.4	73.1	65.7	78.4	60.3

Table 67 (cont'd): General Attitudes

Attitudes towards Transgender Identity					
	Strongly Agree/ Agree				
	Total (n=2000) %	15-24 yrs. (n=1000) %	25-49 yrs. (n=1000) %	Male (n=1008) %	Female (n=992) %
Transgender individuals are dirty	73.6	74.6	72.4	82.5	64.3
Transgender individuals are cursed	61.6	60.1	62.9	70.7	52.1
Transgender individuals should be ashamed	78.5	78.1	79	85	72.1
It is not safe for Transgender individuals to work with children	76.8	73.9	79.8	85.6	74.2
Transgender individuals are doing something wrong and deserve to be punished	59.5	62.8	56.1	69.2	49.5

CHAPTER 11: PRE-EXPOSURE PROPHYLAXIS

The PrEP concept was read to all to ascertain level of awareness of the medication and their response to it. Very few persons (10.4%) were aware of it but many (50.4%) indicated a willingness to take one pill daily to reduce their chances of getting HIV. The Health Centre was the preferred place from which to source this new medication (64%). (*Table 68*)

Table 68: Knowledge of and Willingness to take PrEP

	Total (n=2000) %
Knowledge of PrEP	10.4
Willingness to take PrEP (<i>Very or moderately willing</i>)	50.4
<i>Q. How would you like to get the medication? Top 6 responses</i>	Total (n=1009) %
Health Centre	64.0
NGO or Health Department Outreach Team	14.0
Pharmacy	12.6
From a Peer in their area	5.3
Private Doctor	3.4
Hospital	1.0
Doesn't matter	0.2

Reasons for lack of interest in the medication were equally divided between persons concerned about side effects or not having enough information about it and those who perceive they would not need it since they are not practicing risk behavior (47% each; Table 69).

Table 69: Reasons for Unwillingness to take PrEP

Q. Why would you not be willing to take PrEP?

<i>Top 6 responses</i>	Total (n=826) %
Concerned about side effects	41.0
Don't trust/ not sure that it is effective/Need more information	6.0
Not at risk (faithful to one partner)	24.5
Not at risk (abstinent)	15.7
Not at risk (Consistent condom use)	7.5
Don't take pills/medication	10.4

47.7%

Responses to why persons were willing to take PrEP, all indicate what seems to be a rational calculation of their risk and a willingness to protect themselves (Table 70).

Table 70: Reasons for Willingness to take PrEP

Q: Why would you be willing to take PrEP?

<i>Top 6 responses</i>	Total (n=1009) %
To reduce chances of getting HIV/Precautionary measures	61.8
I know/suspect my partner has other partners and does not consistently use condoms	10.4
I have multiple sex partners and don't consistently use condoms	10.1
I do not know my partner's HIV status	9.8
No answer/ Not Sure	6.6
I have a partner who migrates for work	2.1

CHAPTER 12: HIV SELF TESTING

Very few persons (12.8%) currently have knowledge of HIV Self-Tests (*Table 79*) and 2% have administered same. The majority (69.1%) of the sample however indicated a likelihood that they could consider using this method of testing in the future (*Table 71*).

Table 71: Knowledge of HIV Self Tests

Q. Have you heard of home HIV self-testing?	%	
Know about HIV Self Tests (N=2000)	12.8	
Have administered an HIV test to his/herself (n=256 <i>persons aware</i>)	2.0	
<i>Where Self-test kit obtained:</i>		
Pharmacy	2 persons	
Clinic	1 person	
Doctor’s office	1 person	
Declined to answer	1 person	
Q. How likely are you to use an HIV self-test in the future?	Total (n=2000) %	
Very Likely	49.3	69.1%
Somewhat likely	19.8	
Neutral	4.4	
Somewhat unlikely	3.8	
Very Unlikely	18.0	
Don’t Know	4.2	
Declined to answer	0.5	



CHAPTER 13: RESULTS OF PROGRAMME INDICATORS

List of key GARPR Indicators included in KABP survey.

1. Percentage of young women and men aged 15–24 who correctly identify ways of preventing the sexual transmission of HIV and who reject major misconceptions about HIV transmission
2. Percentage of young women and men aged 15–24 who have had sexual intercourse before the age of 15
3. Percentage of adults aged 15–49 who have had sexual intercourse with more than one partner in the past 12 months
4. Percentage of adults aged 15–49 who had more than one sexual partner in the past 12 months and who report the use of a condom during their last intercourse*
5. Percentage of women and men aged 15–49 who received an HIV test in the past 12 months and know their results
6. Proportion of ever-married or partnered women aged 15–49 who experienced physical or sexual violence from a male intimate partner in the past 12 months
7. Percentage of young adults, 15 to 19 years old, who have never had sex
8. Percentage of young women and men aged 15–24 reporting the use of a condom the last time they has sex with a non-regular partner
9. Median age at which persons reported first having sex
10. Percentage of women who were counselled and offered voluntary HIV test during antenatal care for their most recent pregnancy, accepted an offer of testing and received their results, of all women who were pregnant in the 2 years preceding the survey
11. Percentage of adults aged 15–49 reporting transactional sex
12. Percentage of people 15–49 years expressing accepting attitudes towards people with HIV/AIDS



Indicator –NHP Calculation	Male %	Female %	Total %
1. Percentage of young women and men aged 15–24 who correctly identify both ways of preventing the sexual transmission of HIV and who reject major misconceptions about HIV transmission	31.1	36.0	32.9
Numerator: # of respondents aged 15–24 who gave the correct answer to all five knowledge questions - one partner, 100% condom use, healthy looking person, mosquito bites, sharing food.	195	134	329
Denominator: #Number of all respondents age 15-24	628	372	1000
15-19 <u>Indicator:</u>	28.1	29.4	28.5
<u>Numerator:</u>	104	57	161
<u>Denominator:</u>	370	194	564
20-24 <u>Indicator:</u>	35.3	43.3	38.5
<u>Numerator:</u>	91	77	168
<u>Denominator:</u>	258	178	436
2. Sex before the age of 15	41.6	12.4	30.7
Numerator: # of respondents age 15 -24 report the age at which they first had sexual intercourse as under 15	261	46	307
Denominator: Number of all respondents aged 15-24	628	372	1000
15-19 <u>Indicator:</u>	39.5	11.9	30.0
<u>Numerator:</u>	146	23	169
<u>Denominator:</u>	370	194	564
20-24 <u>Indicator:</u>	44.6	12.9	31.7
<u>Numerator:</u>	115	23	138
<u>Denominator:</u>	258	178	436



3. Multiple Sex Partners in past 12 months	48.9	13.8	31.5
Numerator: # of Respondents aged 15 – 49 who have had sexual intercourse with more than one partner in the last 12 months	493	137	630
Denominator: # of all respondents aged 15 - 49	1008	992	2000
15-19 <u>Indicator:</u>	44.3	12.3	33.3
<u>Numerator:</u>	164	24	188
<u>Denominator:</u>	370	194	564
20-24 <u>Indicator:</u>	60.0	24.1	45.4
<u>Numerator:</u>	155	43	198
<u>Denominator:</u>	258	178	436
25-49 <u>Indicator:</u>	45.7	11.2	24.4
<u>Numerator:</u>	174	70	244
<u>Denominator:</u>	380	620	1000
4. Multiple Sex Partners in past 12 months + condom use at last sex	63.7	40.9	58.7
Numerator: # of respondents (aged 15-49) who reported having had more than one sexual partner in the last 12 months who also reported that a condom was used the last time they had sex	314	56	370
Denominator: Number of respondents (15-49) who reported having had more than one sexual partner in the last 12 months	493	137	630
15-19 <u>Indicator:</u>	70.1	45.8	67.0
<u>Numerator:</u>	115	11	126
<u>Denominator:</u>	164	24	188
20-24 <u>Indicator:</u>	65.2	39.5	59.6
<u>Numerator:</u>	101	17	118
<u>Denominator:</u>	155	43	198
25-49 <u>Indicator:</u>	56.3	40.0	51.6
<u>Numerator:</u>	98	28	126
<u>Denominator:</u>	174	70	244



5. Percentage of women and men aged 15-49 who received an HIV test in the past 12 months and know their results.	23.0	40.7	31.8
Numerator: # of respondents age 15-49 who have been tested for HIV during the last 12 months and who know their results	232	404	636
Denominator: Number of all respondents age 15-49	1008	992	2000
15-19 <u>Indicator:</u>	9.7	18.6	12.8
<u>Numerator:</u>	36	36	72
<u>Denominator:</u>	370	194	564
20-24 <u>Indicator:</u>	29.1	52.8	38.8
<u>Numerator:</u>	75	94	169
<u>Denominator:</u>	258	178	436
25-49 <u>Indicator:</u>	31.8	44.2	39.5
<u>Numerator:</u>	121	274	395
<u>Denominator:</u>	380	620	1000
6. Proportion of ever-married or partnered woman aged 15 – 49 who experienced physical or sexual violence from a male intimate partner in the past 12 months		14.8	14.8
Numerator: Ever married or partnered women aged 15-49 include women who have ever been married or had an intimate partner. An intimate partner is defined as a cohabiting partner, whether or not they had been married at the time. These women are asked if they experienced physical or sexual violence from a male intimate partner in the past 12 months. Those reporting at least one incident corresponding to any one of these items the last 12 months are included in the numerator		106	106
Denominator: Total women surveyed aged 15-49 who currently have or had an intimate partner.		719	719
15-19 <u>Indicator:</u>		20.2	20.2
<u>Numerator:</u>		14	14
<u>Denominator:</u>		69	69
20-24 <u>Indicator:</u>		20.8	20.8
<u>Numerator:</u>		31	31
<u>Denominator:</u>		149	149
25-49 <u>Indicator:</u>		12.1	12.1
<u>Numerator:</u>		61	61
<u>Denominator:</u>		501	501
7. Percentage of young adults, 15-19 years old, who have never had sex	29.7	54.1	38.1



<u>Numerator:</u> # of persons 15-19 yrs who have never had sex	110	105	215
<u>Denominator:</u> # of all respondents 15-19 yrs	370	194	564
8. Percentage of young women and men aged 15-24 reporting the use of a condom the last time they has sex with a non-regular partner	80.1	65.2	78.8
<u>Numerator:</u> # of persons 15-24yrs who used a condom the last time had sex with a non-regular partner	201	15	216
<u>Denominator:</u> # of persons 15-24yrs who had sex with a non-regular partner	251	23	274
15-19 <u>Indicator:</u>	80.2	83.3	80.3
<u>Numerator:</u>	101	5	106
<u>Denominator:</u>	126	6	132
20-24 <u>Indicator:</u>	80.0	58.8	77.5
<u>Numerator:</u>	100	10	110
<u>Denominator:</u>	125	17	142
9. Median age at which persons reported first having sex	14yrs	17yrs	16yrs
Base= Sexually active population 15-49yrs who could recall age of first sex			
15-19	14yrs	16yrs	15yrs
20-24	15yrs	16yrs	16yrs
25-49	15yrs	17yrs	17yrs



10. Percentage of women who were counselled and offered voluntary HIV test during antenatal care for their most recent pregnancy, accepted an offer of testing and received their results, of all women who were pregnant in the 2 years preceding the survey		76.0	76.0
<u>Numerator:</u> # of women 15-49 who were counselled and offered voluntary HIV testing during antenatal care for their most recent pregnancy, accepted an offer of testing and received their test results		165	165
<u>Denominator:</u> # of all woman aged 15-49yrs who were pregnant at any time in the 2 years preceding the survey		217	217
15-19 <u>Indicator:</u>		60.0	60.0
<u>Numerator:</u>		12	12
<u>Denominator:</u>		20	20
20-24 <u>Indicator:</u>		82.1	82.1
<u>Numerator:</u>		55	55
<u>Denominator:</u>		67	67
25-49 <u>Indicator:</u>		75.4	75.4
<u>Numerator:</u>		98	98
<u>Denominator:</u>		130	130
11. Percentage of adults aged 15–49 reporting transactional sex	45.4	16.5	31.1
<u>Numerator:</u> # of persons 15-49yrs reporting transactional sex in last 12 months	458	164	622
<u>Denominator:</u> # of all respondents aged 15-49yrs	1008	992	2000
15-19 <u>Indicator:</u>	31.6	10.8	24.5
<u>Numerator:</u>	117	21	138
<u>Denominator:</u>	370	194	564
20-24 <u>Indicator:</u>	57.4	28.1	45.4
<u>Numerator:</u>	148	50	198
<u>Denominator:</u>	258	178	436
25-49 <u>Indicator:</u>	50.8	15.0	28.6
<u>Numerator:</u>	193	93	286
<u>Denominator:</u>	380	620	1000
12. Percentage of women and men aged 15–49 who report accepting attitudes towards people living with HIV (3 components as per 2012 KABP study)	10.4	12.9	11.6



Numerator: Number of respondents (aged 15–49) who respond positively to either of the three questions - If I knew that a shopkeeper or food seller has HIV, I would definitely still buy food or fresh vegetables from them (Strongly Agree and Agree) If a teacher has HIV but is not sick, he/she should be allowed to continue to teaching in school (Strongly Agree and Agree) and If a member of family got infected with HIV I would want them to keep it a secret (Strongly Disagree and Agree)		104	127	231
Denominator: Number of all respondents (aged 15–49) who have heard of HIV.		1004	988	1992
15-19	<u>Indicator:</u>	6.5	7.7	6.9
	<u>Numerator:</u>	24	15	39
	<u>Denominator:</u>	368	194	562
20-24	<u>Indicator:</u>	8.5	14.6	11.0
	<u>Numerator:</u>	22	26	48
	<u>Denominator:</u>	258	178	436
25-49	<u>Indicator:</u>	15.3	14.0	14.5
	<u>Numerator:</u>	58	86	144
	<u>Denominator:</u>	378	616	994

* “Do you think children living with HIV should be able to attend school...” has been eliminated from 2017 indicators as it was not a part of the 2017 KABP questionnaire. Hence this portion of the indicators has been revised to reflect previous 2012 KABP indicators.



Bibliography

NISP. (n.d.). *National Integrated Strategic Plan for Sexual and Reproductive Health and HIV 2014-2019*.

Retrieved from Ministry of Health: <http://moh.gov.jm/wp-content/uploads/2016/05/Final-National-Integrated-Strategic-Plan-2014-2019-with-SRH-Validation.pdf>

UNAIDS. (2018). *UNAIDS Fact Sheet: Caribbean* . Retrieved from UNAIDS:

http://www.who.int/hiv/mediacentre/200605-FS_Caribbean_en.pdf

USAID. (n.d.). *HIV/AIDS Health Profile*. Retrieved from USAID:

http://pdf.usaid.gov/pdf_docs/pdacu642.pdf