

## 1. VOEVODSKY'S CONNECTIVITY THEOREM FOR $\mathbb{P}^1$ -SPECTRA

The following is from an email from Aravind to me concerning the proof of the connectivity theorem for  $\mathbb{P}^1$  spectra.

Most of the proof appears in Morel's Trieste notes (in the section on the homotopy t-structure for  $P^1$ -spectra). First you prove a connectivity result for  $P^1$ -stable homotopy sheaves by using Morel's  $S^1$ -stable connectivity theorem and studying what happens under  $G_m$  loops and  $G_m$ -suspension: suspension preserves connectivity, and Morel shows that taking  $G_m$ -loops has the effect of making a "contraction". Then, you globalize this.

Our goal is to prove theorem 4.14 of [Voev98], which should be restated in terms of  $\mathbb{P}^1$ -spectra.

**Theorem 1.1.** Let  $X$  be a pointed smooth scheme over  $\mathrm{Spec}(k)$  where  $k$  is an infinite field. Let  $\mathcal{Y}$  be a pointed space. Then for any  $n > \dim(X)$ , and any integer  $m$

$$\mathcal{SH}(k)(\Sigma^\infty X, S^n \wedge \mathbb{G}_m^m \wedge \Sigma^\infty \mathcal{Y}) = 0.$$

*Remark 1.* We can formulate the theorem by using homotopy sheaves. Indeed, if we can show that for any pointed space  $\mathcal{Y}$  the homotopy sheaves  $\pi_{n+\alpha m}^{\mathbb{A}^1}(\mathcal{Y})$  are  $-1$  connected, the result will follow, as

$$\begin{aligned} \mathcal{SH}(k)(\Sigma^\infty X, S^n \wedge \mathbb{G}_m^m \wedge \Sigma^\infty \mathcal{Y}) &= \mathcal{SH}(k)(S^{-n} \wedge \mathbb{G}_m^{-m} \wedge \Sigma^\infty X, \Sigma^\infty \mathcal{Y}) \\ &= \pi_{-n-m\alpha}^{\mathbb{A}^1}(X). \end{aligned}$$

and by the connectivity theorem, this will vanish whenever  $-n < 0$ , i.e., whenever  $n > 0$ . This is a stronger statement than [?, theorem 4.14].

The line of attack is then to show that for a pointed space  $\mathcal{Y}$ , the  $S^1$  suspensions spectrum  $\Sigma_s^\infty \mathcal{Y}$  in the stable model category is  $-1$  connected. Then we establish the first connectivity theorem which ensures the  $\mathbb{A}^1$  localization of  $\Sigma_s^\infty \mathcal{Y}$  is  $-1$  connected. Finally, we show that inverting  $\mathbb{G}_m$  does not affect the connectivity, i.e.,  $\Sigma_m^\infty \Sigma_s^\infty \mathcal{Y}$  is again  $-1$  connected.

## 2. ASSUMPTIONS FROM PREVIOUS LECTURES

**2.1. Facts about Nisnevich topology.** Points and neighborhoods. Distinguished squares determine the topology. Relation to étale, zariski, and fpqc topology.

**Proposition 2.1.** [Mor04, 2.4.1] Let  $M$  be a sheaf of abelian groups on  $\mathrm{Sm}/k$ , and let  $X \in \mathrm{Sm}/k$  with Krull dimension  $d$ . Then whenever  $n > d$ ,  $H_{Nis}^n(X; M) = 0$ .

**Proposition 2.2.** [Mor04, 2.4.1] For any  $X \in \mathrm{Sm}/k$ , and for any  $x \in X(k)$ , there is an isomorphism of pointed sheaves of sets in the Nisnevich topology

$$X/(X - \{x\}) \cong \mathbb{A}^n/(\mathbb{A}^n - \{0\}).$$

**2.2. Unstable model category  $\Delta^{op}\mathrm{Shv}(\mathrm{Sm}/k, Nis)$ .**

- (1) Don't forget the adjunction for sheafification  $a_{Nis} \dashv U$ . i.e., to give a map  $a_{Nis}\mathcal{X} \rightarrow \mathcal{Y}$ , of sheaves, it is equivalent to just give a map  $\mathcal{X} \rightarrow U\mathcal{Y}$ .
- (2) Give the unstable model category the injective model structure to start. (This is Morel's choice, so we stick with it)
- (3) Weak equivalences:  $\mathcal{X} \rightarrow \mathcal{Y}$  iff for all  $U$  the map  $\mathcal{X}(U) \rightarrow \mathcal{Y}(U)$  a simplicial w.e.

- (4) Cofibs:  $\mathcal{X} \rightarrow \mathcal{Y}$  is a cofib. iff for any  $U \in \text{Sm}/k$  the map  $\mathcal{X}(U) \rightarrow \mathcal{Y}(U)$  is a monomorphism.
- (5) Fibrations: what they need to be.
- (6) The ass. htpy. cat. to the injective model category structure is denoted  $\mathcal{H}_s(k)$ . The pointed htpy. cat. is denoted  $\mathcal{H}_{s,\bullet}(k)$ .
- (7) Important properties/constructions in these categories: Cartesian closed, i.e., have smash/tensor  $\times$  and  $\wedge$ , with right adjoints  $\underline{\text{Hom}}$  and  $\underline{\text{Hom}}_\bullet$ . Representable sheaf functor and constant simplicial set functor.

**Definition 2.3.** Internal hom. (Ref: P. Pelaez)

Let  $U \in \text{Sm}/k$ . Let  $\Delta_U^n$  denote the simplicial sheaf given by

$$(V, m) \in \text{Sm}/k \times \Delta^{op} \rightarrow \text{Sm}/k(V, U) \times \Delta_m^n.$$

In other words,  $\Delta_U^n = (rU) \times c\Delta^n$ .

Let  $\mathcal{X}$  and  $\mathcal{Y}$  be spaces. The internal hom in the unpointed category is given by the formula

$$(U, m) \in \text{Sm}/k \times \Delta \rightarrow \text{Hom}_{\Delta^{op}\text{Shv}}(X \times \Delta_U^m, Y).$$

How to describe the adjunction?

$$\text{Hom}(\mathcal{X} \times \mathcal{Y}, \mathcal{Z}) \cong \text{Hom}(\mathcal{Y}, \underline{\text{Hom}}(\mathcal{X}, \mathcal{Z}))$$

Certainly we may send a map  $g \in \text{Hom}(\mathcal{Y}, \underline{\text{Hom}}(\mathcal{X}, \mathcal{Z}))$  to the map  $\eta(g)$  given by

$$\begin{aligned} \eta(g)(U, n) : \mathcal{X}_n(U) \times \mathcal{Y}_n(U) &\rightarrow \mathcal{Z}_n(U) \\ (a, b) &\rightarrow g(U, n)(U, n)(a, \text{id}, \text{id}). \end{aligned}$$

Why is this a bijection?

**Definition 2.4.** Internal hom in the pointed category. Consider  $\mathcal{X}$  and  $\mathcal{Y}$  pointed spaces. For a point  $x \in \mathcal{X}$ , there is an evaluation map  $ev_x : \underline{\text{Hom}}(\mathcal{X}, \mathcal{Y}) \rightarrow \mathcal{Y}$ , where at  $(U, n) \in (\text{Sm}/k \times \Delta)^{op}$  we send  $g : \mathcal{X} \times \Delta_U^n \rightarrow \mathcal{Y}$  to  $g(U, n)(x, \text{id}, \text{id}) \in \mathcal{Y}_n(U)$ . This makes sense as we have the map

$$g(U, n) : \mathcal{X}_n(U) \times \text{Sm}/k(U, U) \times \Delta_n^n \rightarrow \mathcal{Y}_n(U).$$

The pointed internal hom  $\underline{\text{Hom}}_\bullet(\mathcal{X}, \mathcal{Y})$  is the fiber  $ev_x^{-1}(y)$ .

### 2.3. $\mathbb{A}^1$ localization.

- (1) See [?, Prop 2.3.3.] for details on the various properties of fibrant objects in the unstable motivic category.
- (2)

**Definition 2.5.** A space  $\mathcal{X}$  is called  $\mathbb{A}^1$  local if for any smooth scheme  $U$ , the canonical map

$$\text{Hom}(rU, \mathcal{X}) \rightarrow \text{Hom}(rU \times \mathbb{A}^1, \mathcal{X})$$

is a bijection.

- (3)

**Definition 2.6.** A map  $f : \mathcal{X} \rightarrow \mathcal{Y}$  is an  $\mathbb{A}^1$  weak equivalence if

$$\text{Hom}(\mathcal{Y}, \mathcal{Z}) \rightarrow \text{Hom}(\mathcal{X}, \mathcal{Z})$$

is a bijection for every  $\mathbb{A}^1$  local space  $\mathcal{Z}$ .

- (4) The unstable motivic homotopy category is obtained by left Bousfield localization of the injective model category structure on spaces with respect to the class of maps  $W = W_{\mathbb{A}^1} = \{U \times \mathbb{A}^1 \rightarrow U \mid U \in \mathbf{Sm}/k\}$ .
- (5) The general theory of Bousfield localization then gives a localization functor  $L_{\mathbb{A}^1} : \mathcal{H}_s(k) \rightarrow L_W \mathcal{H}_s(k)$ ; denote the category  $L_W \mathcal{H}_s(k)$  by  $\mathcal{H}(k)$ . The localization functor sends  $\mathbb{A}^1$  weak equivalences to isomorphisms.

See [?, Definition 3.3.1]. Keep the same underlying category, the weak equivalences are the  $\mathbb{A}^1$ -local weak equivalences, do not change the cofibrations from the ones in the injective model structure on  $\Delta^{op}\mathbf{Shv}(\mathbf{Sm}/k, \mathbf{Nis})$ , and let the fibrations be what they need to be.

- (6) Morel writes  $L_{\mathbb{A}^1}$  for the (derived) functor which sends a space (without base point)  $\mathcal{X}$  to an  $\mathbb{A}^1$ -localization.
- (7) Morel gives an explicit construction which takes a pointed space  $\mathcal{Y}$  and produces a space  $L^\infty \mathcal{Y}$  which is  $\mathbb{A}^1$  local and a map  $\mathcal{Y} \rightarrow L^\infty \mathcal{Y}$  which is an  $\mathbb{A}^1$  weak equivalence.

Let  $\mathcal{Y}_f$  be the functorial fibrant replacement of  $\mathcal{Y}$  with respect to the injective model structure. Consider  $\mathbb{A}^1$  to be pointed at 0. Morel defines  $L^{(1)}(\mathcal{Y})$  to be the cone of the map  $ev_1 : \underline{Hom}_\bullet(\mathbb{A}^1, \mathcal{Y}_f) \rightarrow \mathcal{Y}_f$ . So there is a map  $\mathcal{Y} \rightarrow L^{(1)}(\mathcal{Y})$  obtained from the trivial cofibration  $\mathcal{Y} \rightarrow \mathcal{Y}_f$  and the defining map  $\mathcal{Y}_f \rightarrow L^{(1)}(\mathcal{Y})$ .

So  $L^{(1)}(-)$  is a functor with a natural transformation  $\eta : \text{id} \rightarrow L^{(1)}(-)$ . Define by induction  $L^{(n)}(\mathcal{Y}) = L^{(1)}(L^{(n-1)}(\mathcal{Y}))$ . There is thus a directed system  $L^{(n-1)}(\mathcal{Y}) \rightarrow L^{(n)}(\mathcal{Y})$ . Denote the direct limit/hocolim of this directed system by  $L^\infty(\mathcal{Y})$ .

**Proposition 2.7.** The natural morphism  $\mathcal{Y} \rightarrow L^\infty(\mathcal{Y})$  is an  $\mathbb{A}^1$  weak equivalence, and  $L^\infty(\mathcal{Y})$  is  $\mathbb{A}^1$  local.

(8)

**Definition 2.8.** Let  $\mathcal{X}$  be a space. Define  $\pi_0(\mathcal{X})$  to be the sheaf on  $\mathbf{Sm}/k$  associated to  $U \rightarrow \pi_0(\mathcal{X}(U))$ . A space  $\mathcal{X}$  is called 0-connected if and only if  $\pi_0(\mathcal{X})$  is the trivial sheaf.

Let  $(\mathcal{X}, x)$  be a pointed space. Define  $\pi_n(\mathcal{X})$  to be the sheafification of the presheaf on  $\mathbf{Sm}/k$  given by

$$U \rightarrow \pi_n(\mathcal{X}(U)).$$

A pointed space  $\mathcal{X}$  is called  $n$ -connected if it is 0-connected and for all  $i \leq n$ , the sheaves  $\pi_i(\mathcal{X})$  are trivial.

(9)

**Proposition 2.9.** Let  $\mathcal{X}$  be a 0-connected simplicial sheaf. Then  $L^\infty \mathcal{X}$  is also 0-connected.

*Proof.* M.V. IHES paper? Sketch of argument? It shouldn't be too hard by chasing components around.  $\square$

- (10) For a sheaf of abelian groups  $M$  on  $\mathbf{Sm}/k$  and a natural number  $n$ , a Dold-Kan construction gives a simplicial presheaf  $K(M, n)$ . It is called the Eilenberg-MacLane spectrum of type  $(M, n)$  and has homotopy sheaves as expected:

$$\pi_m(K(M, n)) = \begin{cases} 0 & \text{if } m \neq n \\ M & \text{if } m = n \end{cases}.$$

Important equation for this construction. For  $X \in \mathbf{Sm}/k$ ,  $M$  a sheaf of Abelian groups,

$$\mathcal{H}_s(k)(rX, K(M, n)) \cong H_{Nis}^n(X; M).$$

- (11) Use square brackets to denote the maps in the unstable pointed (motivic) homotopy category, i.e.,  $[\mathcal{X}, \mathcal{Y}] = \mathcal{H}_\bullet(k)(\mathcal{X}, \mathcal{Y})$ .

Use  $\pi_n^{\mathbb{A}^1}(\mathcal{X})$  for the sheaf of homotopy groups in the motivic category, i.e.,  $\pi_n^{\mathbb{A}^1}(\mathcal{X}) = \pi_n(L^\infty \mathcal{X})$ . This is also obtained by sheafifying the presheaf given by

$$U \in \mathbf{Sm}/k \mapsto [S^n \wedge U_+, \mathcal{X}].$$

## 2.4. Homotopy purity, connectedness calculations.

**Proposition 2.10.** Weak  $n$ -connectedness is equivalent to  $n$ -connectedness

*Proof.* Details? □

**Proposition 2.11.** If  $V$  is an irreducible, smooth  $k$ -scheme, and  $U \subseteq V$  is a dense open subset, the space  $L_{\mathbb{A}^1}(V/U)$  is 0-connected.

*Proof.* Seems to use that local structure of spaces is given by  $\mathbb{A}^n/(\mathbb{A}^n - \{0\}) \wedge L_+$ . How do all the reductions work? Also uses some results specific to working over perfect fields. □

## 2.5. $S^1$ -spectra.

**Definition 2.12.** Let  $\mathbf{Spt}_s(k)$  denote the category of  $S^1$ -spectra of spaces  $\Delta^{op}\mathbf{Shv}(\mathbf{Sm}/k)$ . We first endow this category with the projective model structure, i.e., a map  $f : E \rightarrow F$  is a weak equivalence iff for any  $n$  the map  $f_n : E_n \rightarrow F_n$  is a w.e.; a map  $f : E \rightarrow F$  is a fibration iff for all  $n$  the map  $f_n : E_n \rightarrow F_n$  is a fibration. The cofibrations are characterized by the property that  $f : E \rightarrow F$  is a cofib iff  $f_0 : E_0 \rightarrow F_0$  is a cofib and for any  $n \geq 1$

$$\begin{array}{ccc} S^1 \wedge E_{n-1} & \xrightarrow{\sigma_{n-1}} & E_n \\ S^1 \wedge f_{n-1} \downarrow & & \downarrow \\ S^1 \wedge F_{n-1} & \longrightarrow & P.O. \\ & \searrow \sigma_{n-1} & \downarrow f_n \\ & & F_n \end{array}$$

This model structure does not actually invert  $S^1 \wedge -$ . To accomplish this, we must localize with respect to the stable equivalences.

**Definition 2.13.** A map  $f : E \rightarrow F$  of  $S^1$ -spectra is a stable equivalence iff for any  $n \in \mathbb{Z}$  the induced map of homotopy sheaves  $\pi_n(f) : \pi_n(E) \rightarrow \pi_n(F)$  is an isomorphism.

The stable model category structure on  $\mathbf{Spt}_s(k)$  is given by declaring the weak equivalences to be the stable weak equivalences, and the cofibrations to be the same as those for the projective model structure. This is indeed a left Bousfield localization, but we will not describe it further as such.

**Definition 2.14.** Let  $\mathbf{Spt}_s^{\mathbb{A}^1}(k)$  denote the category of  $S^1$  spectra endowed with the stable model category structure localized at the collection of maps  $\{\Sigma^\infty U_+ \wedge \mathbb{A}^1 \rightarrow \Sigma^\infty U_+ \mid U \in \mathbf{Sm}/k\}$ .

Let  $\mathcal{SH}^{S^1}(k)$  denote the homotopy category associated to  $\mathrm{Spt}_s^{\mathbb{A}^1}(k)$ . We will use  $\mathcal{H}_s^{S^1}(k)$  to denote the homotopy category of  $\mathrm{Spt}_s(k)$ .

*Remark 2.* There is a functor  $L^\infty$  on the category of  $S^1$  spectra which is similar to the unstable construction.

So we can use the functor  $L^\infty$  as an  $\mathbb{A}^1$  localization functor. To be precise, we let  $\mathrm{Spt}_s^{\mathbb{A}^1 \mathrm{loc}}(k)$  be the subcategory of  $\mathrm{Spt}_s(k)$  consisting of the  $\mathbb{A}^1$  local spectra. We may equip  $\mathrm{Spt}_s^{\mathbb{A}^1 \mathrm{loc}}(k)$  with a model structure with weak equivalences the  $\mathbb{A}^1$  weak equivalences, the cofibrations are the stable cofibrations, and the fibrations are what they have to be.

The functor  $L^\infty : \mathrm{Spt}_s(k) \rightarrow \mathrm{Spt}_s^{\mathbb{A}^1 \mathrm{loc}}(k)$  is left adjoint to the inclusion functor, and is a left Quillen functor. The homotopy category of  $\mathrm{Spt}_s^{\mathbb{A}^1 \mathrm{loc}}(k)$  is categorically equivalent to  $\mathcal{SH}^{S^1}(k)$ . [Mor05, Corollary 4.2.3].

For spectra  $E$  and  $F$ , we may compute  $[E, F]^{\mathbb{A}^1} := \mathcal{SH}^{S^1}(k)(E, F)$  by calculating  $[L^\infty E, L^\infty F]^{\mathbb{A}^1}$ . Note that this is  $\mathcal{H}_s^{S^1}(k)(E, L^\infty F)$  by using the adjunction. If we assume  $E$  is cofibrant and  $L^\infty F$  is fibrant, we get the formula

$$[E, F]^{\mathbb{A}^1} = \mathrm{Spt}_s(k)(E, L^\infty F).$$

**Definition 2.15.** Let  $E$  be an  $S^1$  spectrum of spaces. Let  $\pi_n$  denote the sheaf obtained by taking the colimit of the directed system  $\pi_{n+r}(E_r)$  in  $\underline{\mathrm{Ab}}(\mathrm{Sm}/k, \mathrm{Nis})$ . That is,

$$\pi_n(E) = \mathrm{colim}_r \pi_{n+r}(E_r).$$

In particular, for a  $U \in \mathrm{Sm}/k$ , we have

$$\pi_n(E)(U) = \mathrm{colim}_r \pi_{n+r}(E_r)(U).$$

**Definition 2.16.** An  $S^1$ -spectrum  $E$  is said to be  $n$ -connected if for any  $m \leq n$ , the homotopy sheaves  $\pi_m(E)$  are trivial.

**Definition 2.17.** There is a left Quillen functor  $\Sigma_s^\infty : \mathrm{Spc}_\bullet \rightarrow \mathrm{Spt}_s(k)$  given by  $(\Sigma^\infty \mathcal{Y})_n = (S^1)^{\wedge n} \wedge \mathcal{Y}$  with the evident bonding maps. The right adjoint to this functor is given by “evaluation at 0”, i.e.,  $\Omega^\infty(E) = E_0$ .

*Remark 3.* The right derived functor  $R\Omega^\infty : \mathcal{SH}_s^{S^1}(k) \rightarrow \mathcal{H}_\bullet(k)$  is given by the formula

$$R\Omega^\infty(E) = \mathrm{colim}_i \Omega_s^i E_i.$$

This comes from the fact that fibrant  $S^1$  spectra are exactly the  $\Omega$  spectra, and the description of the fibrant replacement functor.

*Remark 4.* We also get a left Quillen functor  $\Sigma_s^\infty : \mathrm{Spc}_\bullet^{\mathbb{A}^1}(k) \rightarrow \mathrm{Spt}_s^{\mathbb{A}^1}(k)$  given by the same formula as above.

What is the fibrant replacement functor in  $\mathrm{Spt}_s^{\mathbb{A}^1}(k)$ ?

*Remark 5.* The stable homotopy category is symmetric monoidal, with smash product  $\wedge$  and internal hom  $\underline{\mathrm{Hom}}$ . Using symmetric spectra, one can give these constructions on the category of spectra.

The stable homotopy category is a triangulated category.

## 2.6. $t$ structure.

**Definition 2.18.** Let  $\mathfrak{C}$  be a triangulated category. A  $t$ -structure on  $\mathfrak{C}$  is a pair of full subcategories  $(\mathfrak{C}_{\geq 0}, \mathfrak{C}_{\leq 0})$  which satisfies

- (1) For any  $X \in \mathfrak{C}_{\geq 0}$  and any  $Y \in \mathfrak{C}_{\leq 0}$ ,  $\mathrm{Hom}_{\mathfrak{C}}(X, Y[-1]) = 0$ .
- (2)  $\mathfrak{C}_{\geq 0}[1] \subseteq \mathfrak{C}_{\geq 0}$  and  $\mathfrak{C}_{\leq 0}[-1] \subseteq \mathfrak{C}_{\leq 0}$
- (3) for any  $X \in \mathfrak{C}$  there exists a distinguished triangle

$$Y \rightarrow X \rightarrow Z \rightarrow Y[1]$$

for which  $Y \in \mathfrak{C}_{\geq 0}$ ,  $Z \in \mathfrak{C}_{\leq 0}[-1]$ .

The heart of a  $t$ -structure is the full subcategory given by  $\mathfrak{C}_{\geq 0} \cap \mathfrak{C}_{\leq 0}$ .

**Definition 2.19** ( $t$ -structure on  $\mathcal{SH}_s^{S^1}(k)$ ). Define  $\mathcal{SH}_s^{S^1}(k)_{\geq 0}$  to be the full subcategory of  $\mathcal{SH}_s^{S^1}(k)$  consisting of objects  $E$  such that  $\pi_n(E) = 0$  whenever  $n < 0$ .

Define  $\mathcal{SH}_s^{S^1}(k)_{\leq 0}$  to be the full subcategory of  $\mathcal{SH}_s^{S^1}(k)$  consisting of objects  $E$  such that  $\pi_n(E) = 0$  whenever  $n > 0$ .

**Theorem 2.20.** The triple  $(\mathcal{SH}_s^{S^1}(k), \mathcal{SH}_s^{S^1}(k)_{\geq 0}, \mathcal{SH}_s^{S^1}(k)_{\leq 0})$  is a  $t$ -structure on  $\mathcal{SH}_s^{S^1}(k)$ .

**Proposition 2.21.** [Mor03, Lemma4.2.4] The functor  $L^\infty : \mathrm{Spt}_s^{S^1}(k) \rightarrow \mathrm{Spt}_{s, \mathbb{A}^1}^{S^1}(k)$  identifies the  $\mathbb{A}^1$ -localized  $S^1$  stable homotopy category with the homotopy category of  $\mathbb{A}^1$ -local  $S^1$  spectra.

**Theorem 2.22** ( $S^1$  stable connectivity theorem). Let  $E \in \mathcal{SH}_s^{S^1}(k)$ , and suppose that whenever  $n < 0$  the sheaf  $\pi_n E = 0$ . Then for all  $n < 0$ ,  $\pi_n L_{\mathbb{A}^1} E = 0$ .

**Theorem 2.23.** The pair  $(\mathcal{SH}_{\geq 0}^{S^1}(k), \mathcal{SH}_{\leq 0}^{S^1}(k))$  is a  $t$ -structure on the category  $\mathcal{SH}^{S^1}(k)$ .

**Definition 2.24.** Strictly  $\mathbb{A}^1$  invariant sheaf of Abelian groups.

If  $M$  is strictly  $\mathbb{A}^1$  invariant sheaf of groups, define the Eilenberg-MacLane spectrum  $HM$  associated to it.

**Proposition 2.25.**  $HM$  is  $\mathbb{A}^1$  local iff  $M$  is strictly  $\mathbb{A}^1$  invariant.

**Proposition 2.26.** The heart of the homotopy  $t$  structure is equivalent to the category of strictly  $\mathbb{A}^1$  invariant sheaves.

## 3. INVERTING $\mathbb{G}_m$ ; $\mathbb{P}^1$ SPECTRA

**3.1.  $\mathbb{G}_m$  suspension and loops.** We always consider  $\mathbb{G}_m$  to be pointed at 1 unless otherwise specified.

**Definition 3.1.** On the category  $\mathrm{Spt}_s(k)$  equipped with the motivic stable model category structure, there is a functor  $\Sigma_m(-) = \mathbb{G}_m \wedge -$  given by  $\Sigma_m(E)_n = \mathbb{G}_m \wedge E_n$  with the evident structure maps.

Smashing with  $\mathbb{G}_m$  is also a functor on the unstable category of pointed spaces, and we give it the same name  $\Sigma_m$ .

**Definition 3.2.** The functor  $\Sigma_m$  on  $\mathrm{Spc}_\bullet(k)$  has a right adjoint denoted  $\Omega_m$ . It is given by the formula  $\Omega_m \mathcal{X} = \underline{\mathrm{Hom}}_\bullet(\mathbb{G}_m, \mathcal{X})$ .

The functor  $\Sigma_m$  on  $\mathrm{Spt}_s^{S^1}(k)$  also has a right adjoint  $\Omega_m$  given by the internal hom functor, i.e.,  $\Omega_m E = \underline{\mathrm{Hom}}(\Sigma^\infty \mathbb{G}_m, E)$ .

**Proposition 3.3.** The functor  $\Sigma_m$  is a left Quillen functor on  $\text{Spt}_s(k)$  and on  $\text{Spt}_s^{\mathbb{A}^1}(k)$ . Furthermore,  $\Sigma_m$  is an exact functor on  $\mathcal{SH}^{S^1}(k)$ .

### 3.2. Contraction in $\underline{\text{Ab}}(\text{Sm}/k, \text{Nis})$ .

**Definition 3.4.** Let  $G$  be sheaf of pointed sets on  $\text{Sm}/k$ . The contraction of  $G$  is the sheaf  $G_{-1} = G_{\text{con}}$  given by the formula

$$U \in \text{Sm}/k \mapsto \ker(G(X \times \mathbb{G}_m) \xrightarrow{ev_1} G(X))$$

Where the map  $ev_1$  is the map induced by  $ev_1 : \text{Spec}(k) \rightarrow \mathbb{G}_m$ , i.e.,  $k[x, x^{-1}] \rightarrow k$  given by  $x \mapsto 1$ .

Note that indeed  $G_{-1}$  is a sheaf since it is the kernel of the morphism of sheaves  $G(-) \rightarrow G(- \times \mathbb{G}_m)$ .

**Proposition 3.5.** If  $G$  is the trivial sheaf of abelian groups, then so is its contraction  $G_{-1}$ .

**Proposition 3.6.** Contraction is an exact functor on the category  $\underline{\text{Ab}}(\text{Sm}/k, \text{Nis})$ .

*Proof.* Functoriality can be established by using the universal property of kernels.

To show exactness, let  $F \rightarrow G \rightarrow H$  be a short exact sequence of sheaves. We must show  $F_{-1} \rightarrow G_{-1} \rightarrow H_{-1}$  is still exact. It suffices to check exactness at the level of stalks.

Since  $\text{id} = ev_1 \circ \pi : G(X) \rightarrow G(X \times \mathbb{G}_m) \rightarrow G(X)$ , we have  $ev_1 : G(-) \rightarrow G(- \times \mathbb{G}_m)$  is a surjection on all smooth schemes. Hence  $ev_1$  is an epimorphism. For a Nisnevich point  $x$  of  $X$ , The following commutative diagram then establishes the exactness of the contraction by the 3x3 lemma.

$$\begin{array}{ccccc} F_{-1,x} & \longrightarrow & G_{-1,x} & \longrightarrow & H_{-1,x} \\ \downarrow & & \downarrow & & \downarrow \\ F_{(x,1)} & \twoheadrightarrow & G_{(x,1)} & \twoheadrightarrow & H_{(x,1)} \\ \downarrow & & \downarrow & & \downarrow \\ F_x & \twoheadrightarrow & G_x & \twoheadrightarrow & H_x \end{array}$$

□

### 3.3. Homotopy sheaves of $\underline{\text{Hom}}(\mathbb{G}_m, E)$ .

**Proposition 3.7.** If  $G$  is a sheaf of Abelian groups, then  $G_{-1} \cong \underline{\text{Hom}}_{\bullet}(\mathbb{G}_m, G)$ . Hence contraction is right adjoint to  $- \wedge \mathbb{G}_m$ . The claim is also true for pointed sheaves of sets.

*Proof.* For this category,  $\underline{\text{Hom}}_{\bullet}(\mathbb{G}_m, G)$  and  $G_{-1}$  both have sections at  $X$  given by  $\ker(ev_1 : G(X \times \mathbb{G}_m) \rightarrow G(X))$ . See description of pointed internal hom for this. □

*Remark 6.* Construction of canonical map  $\pi_n(\underline{\text{Hom}}(\mathbb{G}_m, E)) \rightarrow \pi_n(E)_{-1}$  for an  $S^1$  spectrum  $E$ .

**Lemma 3.8.** For  $E \in \pi(k)$  in the heart of the homotopy  $t$ -structure, the canonical morphism  $\pi_n(\underline{\text{Hom}}(\mathbb{G}_m, E)) \rightarrow \pi_n(E)_{-1}$  is an isomorphism for all  $n \in \mathbb{Z}$ .

*Proof.* As  $E \in \pi(k)$ , there is some strictly  $\mathbb{A}^1$  invariant sheaf  $M$  for which  $E = HM$ . We show that  $\underline{\text{Hom}}(\mathbb{G}_m, HM) \rightarrow H(M_{-1})$  is a weak equivalence. To do this, we show that the

map induces an isomorphism on the sections of any field extension  $K$  of finite type over  $k$ .  
[Mor03, Cor 4.2.8]

Details to come...

□

**Lemma 3.9.** If  $M$  is a strictly  $\mathbb{A}^1$  invariant sheaf of abelian groups, then

$$\underline{Hom}(\mathbb{G}_m, HM) \cong H(M_{-1})$$

**Lemma 3.10.** When  $n \neq 0$ ,

$$(1) \quad [\Sigma^\infty \mathbb{G}_m, HM[n]]_s^{S^1} = 0.$$

The following map is an iso.

$$(2) \quad [\Sigma^\infty \mathbb{G}_m, HM]_s^{S^1} \rightarrow [S^0, H(M_{-1})]_s^{S^1}$$

**Proposition 3.11.** If  $E \in \text{Spt}_s(k)$  is an  $\mathbb{A}^1$ -local spectrum, then for any  $n \in \mathbb{Z}$  the canonical map

$$\pi_n(\underline{Hom}(\mathbb{G}_m, E)) \rightarrow \pi_n(E)_{-1}$$

is an isomorphism.

*Proof.*

□

**3.4.  $\mathbb{P}^1$  spectra,  $(S^1, \mathbb{G}_m)$  bi-spectra.** Goal is to invert  $-\wedge \mathbb{G}_m$  on  $\text{Spt}_s(k)$ . Several different approaches to doing this.

Bigraded homotopy sheaves, bigraded spheres

Show connectivity of  $S^1$  spectra is preserved by taking  $\Sigma_m^\infty$ .

Prove that [Mor03, Definition 5.2.1] gives a  $t$ -structure on  $\mathcal{SH}(k)$ .

Now show/realize that Voevodsky's connectivity theorem holds.

## REFERENCES

- [A-1974] Adams, J.F., *Stable Homotopy and Generalized Homology*. Chicago Lectures in Mathematics, (1974).
- [B] Blander, Benjamin, *Local Projective Model Structures on Simplicial Presheaves*. K-Theory, **24** (2001) 283–301.
- [DHI] Dugger, Dan; Hollander, Sharon; Isaksen, Dan, *Hypercovers and simplicial presheaves*.
- [DLØRV] Dundas, B.; Levine, M.; Østvær, P.; Röndigs, O.; Voevodsky, V., *Motivic Homotopy Theory*. Springer (2000).
- [Hir] Phillip, Hirschhorn, *Model Categories and Their Localization*. AMS (2003).
- [H-Mod] Hovey, Mark, *Model Categories*. online preprint (1991).
- [H-Spt] Hovey, Mark, *Spectra and symmetric spectra in general model categories*. journal? (2001).
- [J] Jardine, J.F., *Simplicial presheaves*. Journal of Pure and Applied Algebra, **47** (1987) 35–87.
- [Mor03] Morel, Fabien, *An introduction to  $\mathbb{A}^1$  homotopy theory*.
- [Mor04] Morel, Fabien, *On the motivic  $\pi_0$  of the sphere spectrum*. NATO science series.
- [Mor05] Morel, Fabien, *The stable  $\mathbb{A}^1$  connectivity theorems*. preprint (2004).
- [Voev98] Voevodsky, Vladimir.  $\mathbb{A}^1$ -Homotopy Theory. Doc. Math. J., (1998) pp. 579–604.