

Numerical Integration

Basics and Newton-Cotes

Sachin Shanbhag

Department of Scientific Computing
Florida State University,
Tallahassee, FL 32306.



References

- ▶ S. Chapra and R. Canale, Numerical Methods for Engineers
- ▶ Carnahan and Wilkes, Applied Numerical Methods, University of Michigan, Class Notes, 1996.
- ▶ Pal, Numerical Analysis for Scientists and Engineers, 2007.
- ▶ Holistic Numerical Methods Institute webpage
- ▶ Samir Al-Amer, Class Notes
- ▶ wikipedia.org

Motivation: Why numerical integration?

- Numerical evaluation of an analytical solution may be problematic

$$\int_0^{\infty} \frac{dx}{\sqrt{x} \cosh(x)} = 2\sqrt{\pi} \sum_{k=0}^{\infty} \frac{(-1)^k}{\sqrt{2k+1}}$$

- Many integrals cannot be evaluated analytically


$$I = \int_a^b e^{-x^2} dx$$

This is related to the so-called “error function”

Motivation: Why numerical integration?

- Integral may be multidimensional/have complicated boundaries

$$I = \iint_{x^2 < \sin(y), y^2 < \cos(x)} e^{\sqrt{x^2+y^2}} dx dy$$




Riemann Sum

In the differential limit, an integral is equivalent to a summation

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=0}^n f(x_i)\Delta x \approx \sum_{i=0}^{N-1} f(x_i)\Delta x$$

where N is a large, but finite, number



Orientation

- ▶ Newton-Cotes
 - ▶ interpolating polynomials
 - ▶ trapezoidal, Simpson's rules, Bode's rule
 - ▶ Multistep rules
- ▶ Romberg Integration
 - ▶ extrapolation technique
 - ▶ successive refinement
- ▶ Gaussian Quadrature
 - ▶ optimize grid spacing to obtain higher order accuracy
 - ▶ unevenly spaced points
- ▶ Singularities
- ▶ Monte Carlo Integration

Newton-Cotes

- The function to be integrated is approximated by a polynomial of order n , in the interval $[a, b]$

$$f(x) \approx p_n(x) = \sum_{i=0}^n a_i x^i$$

This can be done by using our knowledge of interpolating polynomials

- Getting the integral of the polynomial is easy

$$\int_a^b f(x) dx \approx \sum_{i=0}^n a_i \int_a^b x^i dx = \sum_{i=0}^n a_i \frac{b^{i+1} - a^{i+1}}{i+1}$$


Newton-Cotes: Trapezoid Rule

First ordered polynomial

$$f(x) \approx p_1(x) = a_0 + a_1x = f(a) + \frac{f(b) - f(a)}{b - a}(x - a)$$

This implies that the area of the gray shaded region:

$$\int_a^b f(x)dx \approx (b-a) \frac{f(b) + f(a)}{2}$$



Trapezoid Rule: Error

From Taylor series expansion:

$$\begin{aligned}\epsilon &= \int_a^b f(x)dx - \int_a^b p_n(x)dx \\ &= \frac{1}{n+1!} \int_a^b f^{(n+1)}(\xi) \left(\prod_{l=0}^n (x - x_l) \right) dx.\end{aligned}$$

For trapezoidal rule $n = 1$,


$$\begin{aligned}\epsilon &= \frac{1}{2} \int_a^b f''(\xi)(x-a)(x-b)dx \\ &= \frac{1}{2} f''(\eta) \int_a^b (x-a)(x-b)dx \\ &= -\frac{(b-a)^3}{12} f''(\eta)\end{aligned}$$

where $\eta \in [a, b]$, and the mean-value theorem is applied.

Multistep Trapezoidal

Divide interval $[a, b]$ into n segments, which may not necessarily be equal:

$$a = x_0 < x_1 < x_2 < \dots < x_n = b$$



Composite formula

$$\int_a^b f(x)dx \approx \sum_{i=0}^n (x_{i+1} - x_i) \frac{f(x_{i+1}) + f(x_i)}{2}$$

Multistep Trapezoidal: Error

Equally spaced points: $h = x_{i+1} - x_i$

$$\int_a^b f(x)dx \approx h \left[\frac{f(x_0) + f(x_n)}{2} + \sum_{i=1}^{n-1} f(x_i) \right]$$

From Taylor series expansion, for equally spaced case, the error can be shown to be:


$$\epsilon = \frac{(b-a)^3}{12n^2} \max_{\xi \in [a,b]} f''(\xi)$$

Trapezoidal Rule: Error

Consider the example:


$$I = \int_0^{\pi/2} \cos(x) dx = 1$$

with $n = 2, 4, 8, 16, 32, 64, 128$.



Simpson's 1/3 Rule

In the single step version, we use a second order polynomial (quadratic) to pass through three points in the domain.




If points are equispaced, then

$$h = \frac{(b - a)}{2}$$

Simpson's 3/8 Rule

In the single step version, we use a third order polynomial (cubic) to pass through 4 points in the domain.



If points are equispaced, then

$$h = \frac{(b - a)}{3}$$

Error: Single Step

Assuming equispaced points $h = x_{i+1} - x_i$ with $a = x_0$ and $b = x_n$ the endpoints, and $f_i = f(x_i)$, we can summarize the first few Newton-Cotes formulas:

Degree	Common name	Formula	Error
1	Trapezoid rule	$\frac{b-a}{2}(f_0 + f_1)$	$-\frac{(b-a)^3}{12}f^{(2)}(\xi)$
2	Simpson's 1/3 rule	$\frac{b-a}{6}(f_0 + 4f_1 + f_2)$	$-\frac{(b-a)^5}{2880}f^{(4)}(\xi)$
3	Simpson's 3/8 rule	$\frac{b-a}{8}(f_0 + 3f_1 + 3f_2 + f_3)$	$-\frac{(b-a)^5}{6480}f^{(4)}(\xi)$
4	Boole's rule	$\frac{b-a}{90}(7f_0 + 32f_1 + 12f_2 + 32f_3 + 7f_4)$	$-\frac{(b-a)^7}{1935360}f^{(6)}(\xi)$

Note that there is something special about Simpson's 1/3 rule.

Simpson's 1/3 Multistep

Assuming equally spaced points divide interval $[a, b]$ into **even** number of strips ($n = 2, 4, 6$, etc.)

$$\int_a^b f(x) dx \approx \frac{b-a}{6} \left[f(x_0) + 2 \sum_{j=1}^{n/2-1} f(x_{2j}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n) \right]$$

With $n = 6$ for example

$$\int_a^b f(x) dx \approx \frac{b-a}{6} \left[f_0 + 2f_1 + 4f_2 + 2f_3 + 4f_4 + 2f_5 + f_6 \right]$$

Multistep Error Bounds

- ▶ Trapezoidal

$$e = \frac{(b-a)^3}{12n^2} \max_{x \in [a,b]} f''(x)$$

- ▶ Simpson's 1/3

$$e = \frac{(b-a)^5}{2880n^4} \max_{x \in [a,b]} f^{(4)}(x)$$

- ▶ Simpson's 3/8

$$e = \frac{(b-a)^5}{6480n^4} \max_{x \in [a,b]} f^{(4)}(x)$$

Newton Cotes

- ▶ Prefer odd-point (= even degree polynomial) formulas over even point
Note: Simpson's 1/3 rule popularity
- ▶ Very high order schemes are undesirable large weights with alternating signs
- ▶ Essentially the same reason that interpolation fails with high order schemes
- ▶ In practice n is almost never greater than 8.
- ▶ Summary: In Newton-Cotes you have two choices: the order of the polynomial (first = trapezoidal, second = Simpson's 1/3, etc.), and the number of strips you want to discretize the domain into (n).

Romberg Integration

- ▶ Trapezoid formula with n subintervals gives error $\epsilon \sim 1/n^2$
- ▶ Say, I repeat the calculation with $2n$ subintervals
- ▶ Can I combine to get a better estimate?

Error in trapezoidal:

$$E = \frac{(b-a)^3}{12n^2} f''(\xi), \text{ for some } \xi \in [a, b]$$

If we assume $f''(\xi)$ is independent of n

$$E = \frac{C}{n^2}$$

Romberg Integration

- ▶ If we assume that C is roughly constant $I_{true} = \text{true value of the integral}$ $I_n = \text{numerical approximation using } n \text{ steps}$,
- ▶ We have

$$I_{true} = I_n + E(n)$$

- ▶ Therefore,

$$\begin{aligned} E(n) &\approx \frac{C}{n^2} = I_{true} - I_n \\ E(2n) &\approx \frac{C}{4n^2} = I_{true} - I_{2n} \end{aligned}$$


Eliminate C/n^2 ,

$$I_{true,est} = I_{2n} + \frac{I_{2n} - I_n}{3}$$

Romberg Integration: Example

- Consider the integral

$$I = \int_0^1 \frac{4}{1+x^2} dx = \pi.$$



- Let us numerically compute the solution using trapezoidal rule with different number of steps, n

Example

Using multistep trapezoidal rule:

n	I_n	ϵ
1	3.0000	1.4159e-01
2	3.1000	4.1593e-02
4	3.1312	1.0416e-02
8	3.1390	2.6042e-03
16	3.1409	6.5104e-04
32	3.1414	1.6276e-04
64	3.1416	4.0690e-05
128	3.1416	1.0173e-05
256	3.1416	2.5431e-06
512	3.1416	6.3578e-07

You can see that the error goes down roughly by a factor of 4 with every doubling of n

Example

- ▶ We can refine an estimate by using say, $I_2 = 3.1000$ and $I_4 = 3.1312$ as

$$I_{true,est} = I_4 + \frac{I_4 - I_2}{3} = 3.1313$$

which pulls the estimate in the direction of the true solution for essentially no extra work.

- ▶ Note: I_2 uses a subset of the same function evaluations as I_4 . That is, the grid points x_i that are used in the calculation of I_4 include all of the grid points that are used in the evaluation of I_2 .
- ▶ This is where a majority of the cost-savings comes from.

Romberg

Trapezoidal			Romberg	
n	$I_n^{(1)}$	ϵ	$I_n^{(2)}$	ϵ
1	3.0000	1.4159e-01		
2	3.1000	4.1593e-02	3.1004	4.1200e-02
4	3.1312	1.0416e-02	3.1313	1.0294e-02
8	3.1390	2.6042e-03	3.1390	2.5735e-03
16	3.1409	6.5104e-04	3.1409	6.4338e-04

Typically, this itself leads to better estimates. But that is not all. We can squeeze out some more performance relatively cheaply.

Error Estimate

- For trapezoidal rule, it can be shown that

$$E = \frac{C_1}{n^2} + \frac{C_2}{n^4} + \frac{C_3}{n^6} + \dots$$

- For the scheme

$$I_{true,est} = I_{2n} + \frac{I_{2n} - I_n}{3}$$

the leading error is $\mathcal{O}(1/n^4)$ (like Simpson's rule).

- In fact, in some places, this is how Simpson's is implemented (Numerical Recipes, Press et al.)

Successive Refinement

- ▶ General Expression for Romberg Integration

$$I_{2n}^{(k)} = \frac{4^{k-1} I_{2n}^{(k-1)} - I_n^{(k-1)}}{4^{k-1} - 1}$$

- ▶ index k represents order of extrapolation
- ▶ $I_n^{(1)}$ represents regular trapezoidal with n segments
- ▶ $k = 2$ represents values reported above using one application of Romberg integration
- ▶ $I_n^{(k)}$ has an error of order $1/n^{2k}$

Example: Successive Refinement

n	$I_n^{(1)}$	$I_n^{(2)}$	$I_n^{(3)}$	$I_n^{(4)}$	$I_n^{(5)}$
1	3.0000				
2	3.1000	3.1333			
4	3.1312	3.1416	3.1421		
8	3.1390	3.1416	3.1416	3.1416	
16	3.1409	3.1416	3.1416	3.1416	3.1416