
Nuclear Fuel Cycle

NUGN506 - LME1

By

GUILLAUME L'HER



Department of Nuclear Engineering
COLORADO SCHOOL OF MINES

LME submitted for the Nuclear Fuel Cycle class at the Colorado School of Mines.

FALL 2017

TABLE OF CONTENTS

	Page
1 Learning Measurement 1	1
1.1 Extra Credit	1
1.1.1 Problem	1
1.1.2 Solution	1
Bibliography	3

LEARNING MEASUREMENT 1

The extra-credit problems for the first learning measurement exercise is tackled in this document.

1.1 Extra Credit

1.1.1 Problem

Thermal neutron fission of U-235 produces nuclides of the $A = 132$ isobar in 4.31% of all fissions. This isobar includes the radioactive doubly-magic nuclide Sn-132 with $Z = 50$, $N = 82$ and a half-life of 39.7 seconds, but Te-132 has the highest independent fission yield for $A = 132$. Why aren't significant amounts of Cs-132 and Ba-132 (or Cs-131 or Ba-133 for the $A = 131$ or $A = 133$ isobars, respectively) observed in the cumulative fission yield when neighboring nuclides like Cs-133 and Ba-134 are? [Hint: It has nothing to do with magic numbers as Te-132 is the most probable fission product.]

1.1.2 Solution

The cumulative fission yield is different from the independent fission yield. The cumulative fission yield also consider the decay of the precursors to form a nuclide of interest. In our case, we know that Te-132 has the highest independent fission yield for $A = 132$. According the the chart of the nuclides, it beta decay to I-132, I-132 then can beta decay to a stable Xe-132, or lose its excited state. Consequently, this decay chain does not produce any element with more protons than Xenon, and thus no Cs-133 or Ba-133. Similarly for $A = 131$ or $A = 133$, the decay chains stop at the elements Xe-131 and Cs-133 respectively.

Consequently, the cumulative fission yield for A132-isobars Cs-132 and Ba-132 are "only" their independent fission yield. In the meantime, Cs-133 and Ba-134 can also be created by the β - decay chains:



BIBLIOGRAPHY