

Numerical simulations of LZ78 for Markovian sources

I Conditions of the experiment

1 Simulation details

Simulations in this report follow this experimental process :


- Generating a random Markov chain of size 2 of matrix

$$\begin{pmatrix} p_{00} & p_{01} \\ p_{10} & p_{11} \end{pmatrix}$$


- Generating $n_{\text{exp}} \sim 10^3$ words of length n (or n_{word}), with $n \sim 10^6$ or 10^7
- Applying LZ78 on each of these words to estimate, for each n , the number of phrases M_n . A simple histogram of these values can be seen in figure 1.
- From this sampling of the random variable M_n and other parameters such as the entropy of the Markov chain, computing
 - the empirical mean (μ) and the empirical variance (σ^2)
 - different theoretical expressions for the variance
- Using these expressions to standardize M_n in different ways, plotting
 - the probability distribution of M_n (standardized)
 - the cumulative distribution function of M_n (standardized)
- Finally, comparing the different theoretical expressions for the variance by plotting their differences for a large range of values of n , and a constant number of experiments n_{exp} .

2 Empirical normalization

Using the empirical mean (μ) and variance (σ^2) of the dataset to normalize M_n , this is a plot of the normalized distribution, compared to the normal distribution in red :



and its cumulative distribution function in green, compared to the normal one in red :



These simulations and figures strongly indicate that the general distribution of M_n respects the central limit theorem. We now experiment with candidates for the variance of M_n : V_n

II Validating variance candidates

1 A first expression

As it will be used in the next section, this is the detail of the expression from [2] :

$$V_n = \frac{H^3 \sigma^2 n}{\log_2^2(n)}$$

$$\sigma^2 = \sigma_0^2 + \sigma_1^2$$

where

$$\sigma_i^2 = \frac{\pi_i p_{i0} p_{i1}}{H^3} \left(\log_2 \left(\frac{p_{i0}}{p_{i1}} \right) + \frac{H_1 - H_0}{p_{01} + p_{10}} \right)^2$$

with

$$\pi_0 = \frac{p_{10}}{p_{10} + p_{01}} \quad \pi_1 = \frac{p_{01}}{p_{10} + p_{01}}$$

and

$$H_i = -p_{i0} \log_2(p_{i0}) - p_{i1} \log_2(p_{i1}) \quad H = \pi_0 H_0 + \pi_1 H_1$$

2 Using the Frobenius eigenvalue of $P(s)$

An expression which seems to be succesful for the variance is :

$$V_n = \left(\ddot{\lambda}(-1) - \dot{\lambda}(-1)^2 \right) \frac{n}{\ln^2 n}$$

Let's compute $\ddot{\lambda}(-1)$ with a Markov chain of order 1.

In the paper, $\ddot{\lambda}(-1) = \pi \ddot{P}(-1) \psi + 2 \dot{\pi}(-1) \dot{P}(-1) \psi - 2 \dot{\lambda}(-1) \dot{\pi}(-1) \psi$

However, the relations defining $\pi(s)$:

$$\begin{cases} \pi(s) P(s) &= \lambda(s) \pi(s) \\ P(s) \psi(s) &= \lambda(s) \psi(s) \\ \pi(s) \psi(s) &= \lambda(s) \end{cases}$$

did not seem to allow me to directly compute $\dot{\pi}(s)$ (it seemed like I need one more). Therefore, I computed $\lambda(s)$ as the greatest eigenvalue of $P(s)$. Let χ the characteristic polynomial of $P(s)$, and Δ its discriminant

$$\chi = (X - p_{00}^{-s})(X - p_{11}^{-s}) - (p_{01} p_{10})^{-s}$$

and

$$\begin{aligned}\Delta &= (p_{00}^{-s} + p_{11}^{-s})^2 - 4[(p_{00} p_{11})^{-s} - (p_{01} p_{10})^{-s}] \\ &= p_{00}^{-2s} + p_{11}^{-s} - 2(p_{00} p_{11})^{-s} + 4(p_{01} p_{10})^{-s}\end{aligned}$$

Informally, we have this expression for $\lambda(s)$ where we need to decide which sign is the correct one :

$$\lambda(s) = \frac{p_{00}^{-s} + p_{11}^{-s} \pm \sqrt{\Delta(s)}}{2}$$

Since

$$\Delta(-1) = (p_{00} + p_{11})^2 - 2p_{00}p_{11} + 4p_{01}p_{10} = (p_{00} + p_{11} - 2)^2$$

then $\sqrt{\Delta(-1)} = 2 - p_{00} - p_{11} = p_{01} + p_{10}$. Thus, picking the $+$ sign in the former expression, we verify that

$$\lambda(-1) = \frac{p_{00} + p_{11} + \sqrt{\Delta(-1)}}{2} = 1$$

Derivating

$$\dot{\lambda}(s) = \frac{1}{2} \left(-\ln p_{00} p_{00}^{-s} - \ln p_{11} p_{11}^{-s} + \frac{\Delta'(s)}{2\sqrt{\Delta(s)}} \right)$$

and the expression for $\Delta'(s)$

$$\Delta'(s) = -2 \ln p_{00} p_{00}^{-2s} - 2 \ln p_{11} p_{11}^{-2s} + 2 \ln(p_{00} p_{11}) (p_{00} p_{11})^{-s} - 4 \ln(p_{01} p_{10}) (p_{01} p_{10})^{-s}$$

gives

$$\Delta'(-1) = -2 \ln p_{00} p_{00}^2 - 2 \ln p_{11} p_{11}^2 + 2 \ln(p_{00} p_{11}) (p_{00} p_{11}) - 4 \ln(p_{01} p_{10}) (p_{01} p_{10})$$



allowing to compute $\dot{\lambda}(-1)$. Numerically, we verified that $\dot{\lambda}(-1) = h$. Derivating again yields

$$\ddot{\lambda}(s) = \frac{1}{2} \left(\ln^2 p_{00} p_{00}^{-s} + \ln^2 p_{11} p_{11}^{-s} + \frac{\Delta''(s)\sqrt{\Delta(s)} - \Delta'(s) \cdot \Delta'(s)/2\sqrt{\Delta(s)}}{2\Delta(s)} \right)$$


with $\Delta''(s) = 4 \ln^2 p_{00} p_{00}^{-2s} + 4 \ln^2 p_{11} p_{11}^{-2s} - 2 \ln^2(p_{00} p_{11}) (p_{00} p_{11})^{-s} + 4 \ln^2(p_{01} p_{10}) (p_{01} p_{10})^{-s}$

$$\text{Finally } \ddot{\lambda}(-1) = \frac{1}{2} \left(\ln^2 p_{00} p_{00} + \ln^2 p_{11} p_{11} + \frac{\Delta''(-1)\sqrt{\Delta(-1)} - \Delta'(-1)^2/2\sqrt{\Delta(-1)}}{2\Delta(-1)} \right)$$

The simulations using this coefficient for the variance are quite good. It also seems that this formula for the variance is equivalent to the one used in the unpublished paper *Probabilistic Analysis of Lempel-Ziv Parsing for Markov Sources* by Leckey, Wormald and Neininger, but our two ways of deriving it differs. Numerical instability might account for the tiny differences found for high n values (10^7), although this hasn't been verified.

Empirical standard deviation (σ) and theoretical ones ($\sigma_{\text{Neininger}}$, σ_S), $n_{\text{exp}} = 400$ Difference between standard deviations, $n_{\text{exp}} = 400$ 

Now, for some distributions of very long words that were normalized using our theoretical standard deviations, and empirical means. The blue plot is a gaussian fit for the simulation results, which also appear as a blue histogram. The two sets of figures are identical but obtained using different expressions.



III Conclusion

Similar results were obtained for a variety of randomly generated Markov sources, which seem to indicate that this formula for the variance could be proven theoretically correct.

Limits of this work

The figures suffer from imprecision over the computation of the empirical variance : this is due to the difficulties encountered in computing large amounts of long words (of size over 10^6). The figure in appendix A is an example of this limitation : with only 100 experiments, the empirical variance varies a lot. Possible ideas of improvement might come from parallelization, rewriting functions in a computational language such as Julia, or using/devising a datastructure specific to the task of building very long words.

Another problem is that the space of random Markov chains (here : stochastic matrices of size 2) is not sampled thoroughly. Sampling a small number of Markov chains uniformly according to their entropy might be interesting as a representation of the space, because otherwise it would

be hard to sample a large number of stochastic matrices due to the necessity of computing large words for each of them.

Finally, the difference between empirical V_n and our expression seems to be growing very slowly with n . This might be a term that is negligible (*i.e.* of order lower than $n/\log_2^2(n)$), or a small detail in the formula of V_n . For example, the natural base logarithm (in $n/\ln^2(n)$) in the eigenvalue expression works slightly better than the one in base 2 (in $n/\log_2^2(n)$), with the inverse situation happening for the first expression ($\sigma_{\text{Neininger}}$). Anyway, the figures obtained seem to indicate that we are close to the exact solution.


The code for these experiments, with detailed procedures for reproducibility, is hosted on GitHub in one of my private repositories (which I plan to make public eventually). Another (probably bad) way of computing $\lambda(s)$ is in appendix B.

Références

- [1] JACQUET, SZPANKOWSKI, TANG, *Average profile of the Lempel-Ziv parsing scheme for a Markovian source*
- [2] LECKET, WORMALD, NEININGER, *Probabilistic Analysis of Lempel-Ziv Parsing for Markov Sources*

Appendices

A Much longer words



B Another (more complicated) computation of $\ddot{\lambda}(-1)$

This expression gives the same numerical results as the first one, but is more complex to compute for no apparent gain other than having yet another similar way of computing $\ddot{\lambda}(-1)$. Computing $\delta(s)$, a complex root of $\Delta(s)$, writing Δ as :

$$\begin{aligned} \Delta = & \underbrace{p_{00}^{-2\operatorname{Re}(s)} \cos(2 \ln(p_{00}) \operatorname{Im}(s))}_{a_0(s)} \\ & + \underbrace{p_{11}^{-2\operatorname{Re}(s)} \cos(2 \ln(p_{11}) \operatorname{Im}(s))}_{a_1(s)} \\ & - \underbrace{2(p_{00} p_{11})^{-\operatorname{Re}(s)} \cos(\ln(p_{00} p_{11}) \operatorname{Im}(s))}_{a_2(s)} \\ & + \underbrace{4(p_{01} p_{10})^{-\operatorname{Re}(s)} \cos(\ln(p_{01} p_{10}) \operatorname{Im}(s))}_{a_3(s)} \\ & + i \operatorname{Im}(\Delta) \end{aligned}$$

where $\operatorname{Im}(\Delta) = b_0(s) + b_1(s) + b_2(s) + b_3(s)$, with each $b_i(s)$ being the same term as $a_i(s)$ with \cos replaced by \sin . Writing

$$\Delta = \alpha(s) + i\beta(s)$$

and searching for $\delta = x(s) + iy(s)$, meaning that

$$\begin{cases} x^2 - y^2 &= \alpha \\ 2xy &= \beta \\ x^2 + y^2 &= \sqrt{\alpha^2 + \beta^2} \end{cases}$$

This yields

$$\begin{cases} x &= \pm \sqrt{\frac{1}{2}(\sqrt{\alpha^2 + \beta^2} + \alpha)} \\ y &= \pm \sqrt{\frac{1}{2}(\sqrt{\alpha^2 + \beta^2} - \alpha)} \end{cases}$$

and since $2xy = \beta$, there is $\varepsilon \in \{-1, 1\}$ such that

$$\delta = \pm(x + i\varepsilon y)$$

so

$$\lambda(s) = \frac{p_{00}^{-s} + p_{11}^{-s} \pm (x + i\varepsilon y)}{2}$$

i.e.

$$\ddot{\lambda}(-1) = \frac{p_{00} \ln^2(p_{00}) + p_{11} \ln^2(p_{11}) \pm (\ddot{x}(-1) + i\varepsilon \ddot{y}(-1))}{2}$$

where we'll have to find what is ε and which sign to pick.

But first, computing the derivatives of $x(s) = \sqrt{f(s)}$:

$$\dot{x}(s) = \frac{f'(s)}{2x(s)}$$

and

$$\ddot{x}(s) = \frac{f''(s)x(s) - f'(s) \cdot \frac{f'(s)}{2x(s)}}{2x^2(s)}$$

and then computing $f(s)$:

$$f(s) = \frac{1}{2}(\sqrt{\alpha^2 + \beta^2} + \alpha)$$

$$f'(s) = \frac{1}{2} \left[\frac{\overbrace{\dot{\alpha}\alpha + \dot{\beta}\beta}^{\gamma(s)}}{\underbrace{\sqrt{\alpha^2 + \beta^2}}_{\kappa(s)}} + \dot{\alpha} \right]$$

with

$$\dot{\alpha} = \dot{a}_0 + \dot{a}_1 + \dot{a}_2 + \dot{a}_3$$

As for $f''(s)$, it is

$$f''(s) = \frac{1}{2} \left[\frac{\dot{\gamma}(s)\kappa(s) - \gamma(s)\dot{\kappa}(s)}{\kappa^2(s)} + \ddot{\alpha}(s) \right]$$

with

$$\dot{\gamma}(s) = \ddot{\alpha}\alpha + \dot{\alpha}^2 + \ddot{\beta}\beta + \dot{\beta}^2$$

$$\dot{\kappa}(s) = \frac{2\alpha\dot{\alpha} + 2\beta\dot{\beta}}{2\sqrt{\alpha^2 + \beta^2}}$$

Derivating according to s amounts to derivating according to $\text{Re}(s)$, so in $s = -1$:

$$\dot{\alpha}(-1) = -2 \ln p_{00} a_0(-1) - 2 \ln p_{11} a_1(-1) - \ln q_0 a_2(-1) - \ln q_1 a_3(-1)$$

and

$$\ddot{\alpha}(-1) = 4 \ln^2 p_{00} a_0(-1) + 4 \ln^2 p_{11} a_1(-1) + \ln^2 q_0 a_2(-1) + \ln^2 q_1 a_3(-1)$$

At this point we have fully determined $\ddot{x}(s)$, and we realize two things:

1. In $s = -1$, since $\text{Im}(-1) = 0$ and because of the sinus function, all the β terms, including derivatives, are equal to 0. This will simplify the expression for $\ddot{x}(-1)$.
2. Furthermore, it also means that $\ddot{y}(-1) = 0$, so

$$\ddot{\lambda}(-1) = \frac{p_{00} \ln^2(p_{00}) + p_{11} \ln^2(p_{11}) + \ddot{x}(-1)}{2}$$

where the $+$ comes from the fact that $\lambda(s)$ is the highest eigenvalue (and $\ddot{x}(-1) > 0$, so by continuity the expression around $s = -1$ retained the same sign)

The final expression of $\ddot{\lambda}$ (as well as $\dot{\lambda}(-1)$) can be fully expressed with $\alpha(-1)$, $\dot{\alpha}(-1)$ and $\ddot{\alpha}(-1)$. I empirically verified that $\dot{\lambda}(-1) = h$, and the final result is the same as with the first method of computation.