

Menstruation as a Vital Sign in Mumbai

Lauren C. Houghton and MYNA

Statement of Purpose

Menstruation in India is highly stigmatized and is often perceived as a disease.¹ However, there is growing interest in the de-stigmatization of menstruation in India. The framing of menstruation as a vital sign²—that is a clinical marker to be considered when assessing health— may be an additional lever to aid in the de-stigmatization process. The purpose of this proposal is to do engage with Mumbai-based stakeholders to understand if “menstruation as a vital sign” should be a public health agenda to pursue across multiple levels.

Background and Rationale

In collaboration with the Columbia Global Centers | Mumbai, I have identified gaps in menstrual health knowledge in the populational health, educational, and medical sectors in India. At the populational level, the established demographic averages for adolescent menstrual cycle characteristics are outdated and based on data from North American cohorts born in the 1960s;³ even less has been established within the Indian context. Only a handful of cross-sectional studies have assessed average age at menarche (the first menstrual period) in Indian girls and they suggest menarche is occurring earlier in India following secular trends.⁴⁻⁶

Within the educational sector, 71% of Indian girls have no knowledge of menstruation before their first period.⁷⁻⁹ School health programs and community-based initiatives primarily focus on the practical aspects of menstrual hygiene, with limited attention to the broader understanding of menstruation as an integral aspect of health across the lifecourse.^{10,11} The lifecourse approach to menstrual health is important because the hormones that drive the cyclic nature of menstruation are shaped by factors that occur early in life, particularly during childhood and adolescents. There is international variation in the age at menarche and the hormones underlying healthy menstrual cycles.¹² This variation should be considered normal rather than pathological in terms of fertility.¹³ However, high levels of menstrual cycle hormones are also associated with increased risk of chronic disease including breast cancer,^{14,15} cardiovascular disease¹⁶ and polycystic ovary syndrome.¹⁷ Despite this robust evidence, within the medical sector, biomedical guidelines have yet to incorporate menstruation as a vital sign into pediatric and gynecological practice as means to track women’s health across the lifecourse.

Health seeking behavior is also poor during adolescence due to a lack of scientific knowledge about menstruation. 75% of girls experience menstrual problems during adolescence that may affect their health during adulthood.² Yet, only one-third of adolescents seek treatment for their menstrual problems.^{18,19} Providing education to adolescents is not enough. It is also important to sensitize clinicians, particularly pediatricians, on addressing adolescents’ menstrual health needs in a safe, positive and friendly environment to improve health outcomes.

Aim 1: Characterize menstrual cycle characteristics (cycle length, period length, flow, symptoms) in mothers and daughters (n=20 dyads) and compare between two settings, informal settlements and private schools, through menstrual diaries and apps

Aim 2: Characterize salivary estrogen and progesterone over the menstrual cycle in mothers and daughters (n=20 dyads) and compare between two settings, informal settlements and private schools

Aim 3: Discuss menstrual health knowledge and needs through focus groups and key informant interviews with children, educators, and pediatricians.

Methods

Recruitment: We will recruit 20 mother/daughter dyads through MYNA an NGO and through X school

We will engage and pay citizen scientists to help conduct the work.

Participants will be compensated for each part of the study: x for questionnaire, x for menstrual diary, x for app use, x for a month of saliva samples

Questionnaire: At the first study visit, study staff will administer a questionnaire including pubertal development scale, menstrual cycle characteristics, diet and age.

Anthropometrics: Study staff will measure height and weight.

Menstrual Diary: Participants will track their menstrual cycles for 1 month using a paper calendar, tracking days of bleeding and other symptoms such as cramping, cravings, mood.

Menstrual Apps: Participants will also track their cycles using the MYNA app.

Saliva collection (freezers?): At the initial study visit, we will collect 2ml of saliva using the passive drool method. We will provide a month's worth of collection supplies for the participants to take home. We will ask the participants to provide a daily saliva sample (2ml) and either keep the samples in their freezer or use a preservative. At the second study visit, study staff will collect the samples and corresponding menstrual tracking data. We will also collect a final 2ml sample at the second study visit. Saliva will be measured in a lab in Delhi using Salimetrics kits

Focus Groups: Two focus groups with 6-8 children each one in informal settlements and one in the school

Key informant interviews: 2 with pediatricians and 4 educators in schools and NGOs

Budget

Research Assistants: INR?

Participant Incentives: 40 x INR?

Collection Supplies: \$3820

Hormone Analysis: TBD

1. FSG and BMGF. *Menstrual Health in India | Country Landscape Analysis.*; 2016. www.fsg.org. Accessed August 4, 2022.
2. Menstruation in Girls and Adolescents: Using the Menstrual Cycle as a Vital Sign. In: *Pediatric Clinical Practice Guidelines & Policies*. American Academy of Pediatrics; 2021:1517-1517. doi:10.1542/9781610020862-part06-menstruation
3. APTER D, VIHKO R. Early Menarche, a Risk Factor for Breast Cancer, Indicates Early Onset of Ovulatory Cycles*. *J Clin Endocrinol Metab*. 1983;57(1):82-86. doi:10.1210/jcem-57-1-82
4. Dambhare DG, Wagh S V., Dudhe JY. Age at menarche and menstrual cycle pattern among school adolescent girls in Central India. *Glob J Health Sci*. 2012;4(1):105-111. doi:10.5539/GJHS.V4N1P105
5. Gupta N, Singh MP, Dhillon BS, Saxena NC. Preparing for adulthood--patterns of physical growth, sexual maturity and menarche of adolescent girls in selected urban slums and rural areas. *J Indian Med Assoc*. 2007;105(3):119-122,126.
6. Agarwal DK, Agarwal KN, Upadhyay SK, Mittal R, Prakash R, Rai S. Physical and sexual growth pattern of affluent Indian children from 5 to 18 years of age. *Indian Pediatr*. 1992;29(10):1203-1282.
7. Omidvar S, Amiri F, Bakhtiari A, Begum K. A study on menstruation of Indian adolescent girls in an urban area of South India. *J Fam Med Prim Care*. 2018;7(4):698. doi:10.4103/JFMPC.JFMPC_258_17
8. Chandra-Mouli V, Patel SV. Mapping the knowledge and understanding of menarche, menstrual hygiene and menstrual health among adolescent girls in low- and middle-income countries. *Reprod Health*. 2017;14(1):1-16. doi:10.1186/S12978-017-0293-6/TABLES/7
9. Trust K. On! Improving Menstrual Health and Hygiene in India USAID FROM THE AMERICAN PEOPLE. www.dasra.org. Accessed August 4, 2022.
10. Ministry of Health and Family Welfare Government of India Operational Guidelines on School Health Programme under Ayushman Bharat Health and Wellness Ambassadors partnering to build a stronger future A Joint initiative of Ministry of Health & Family Welfare and Ministry of Human Resource & Development, Government of India Ministry of Human Resource Development Government of India School. 2018.
11. Menstrual Hygiene Scheme(MHS) :: National Health Mission. <https://nhm.gov.in/index1.php?lang=1&level=3&sublinkid=1021&lid=391>. Accessed August 4, 2022.
12. Ellison PT. Salivary steroids and natural variation in human ovarian function. *Ann N Y Acad Sci*. 1994;709:287-298. <http://www.ncbi.nlm.nih.gov/pubmed/8154721>. Accessed March 19, 2019.
13. Ellison PT, Panter-Brick C, Lipson SF, O'Rourke MT. The ecological context of human ovarian function. *Hum Reprod*. 1993;8(12):2248-2258. <http://www.ncbi.nlm.nih.gov/pubmed/8150934>. Accessed February 1, 2019.
14. Wallace RB, Sherman BM, Bean JA, Leeper JP, Treloar AE. Menstrual cycle patterns and breast cancer risk factors. *Cancer Res*. 1978;38(11 Pt 2):4021-4024. <http://www.ncbi.nlm.nih.gov/pubmed/698949>. Accessed November 7, 2019.
15. Butler LM, Potischman NA, Newman B, et al. Menstrual risk factors and early-onset breast cancer. *Cancer Causes Control*. 2000;11(5):451-458. <http://www.ncbi.nlm.nih.gov/pubmed/10877338>. Accessed September 2, 2016.
16. Hoyt LT, Falconi AM. Puberty and perimenopause: reproductive transitions and their implications for women's health. *Soc Sci Med*. 2015;132:103-112. doi:10.1016/j.socscimed.2015.03.031
17. Rosenfield RL. Clinical review: Adolescent anovulation: maturational mechanisms and implications. *J*

Clin Endocrinol Metab. 2013;98(9):3572-3583. doi:10.1210/jc.2013-1770

18. S. MH, L. H, S. RT, N. V. Assessment of knowledge and health care seeking behaviour for menstrual health among adolescent school girls in urban slums of Bengaluru: a cross sectional study. *Int J Community Med Public Heal.* 2019;6(11):4881-4886. doi:10.18203/2394-6040.IJCMPH20195073
19. Chauhan S, Kumar P, Patel R, Srivastava S, Jean Simon D, Muhammad T. Association of lifestyle factors with menstrual problems and its treatment-seeking behavior among adolescent girls. *Clin Epidemiol Glob Heal.* 2021;12:2213-3984. doi:10.1016/J.CEGH.2021.100905