

D-SCRAMBLED CANTOR SETS

B. MILLER

ABSTRACT. For all $d \geq 3$, we provide an example of a homeomorphism of Cantor space that has an uncountable d -scrambled set but no d -scrambled Cantor set.

INTRODUCTION

Suppose that X is a metric space and $f: X \rightarrow X$. We say that $x \in X^d$ is *proximal* if $\liminf_{n \rightarrow \infty} \max_{i < j < d} d_X(f^n(x(i)), f^n(x(j))) = 0$, *asymptotic* if $\limsup_{n \rightarrow \infty} \min_{i < j < d} d_X(f^n(x(i)), f^n(x(j))) = 0$, and *Li–Yorke* if it is proximal but not asymptotic. A set $Y \subseteq X$ is *d-scrambled* if every injective sequence in Y^d is Li–Yorke, and f is *Li–Yorke d-chaotic* if there is an uncountable d -scrambled set $Y \subseteq X$.

In [GGM21], it was shown that every Li–Yorke 2-chaotic continuous function on a Polish space has a 2-scrambled Cantor set. Here we show:

Theorem 1 (ZFC). *Suppose that $d \geq 3$. Then there is a Li–Yorke d-chaotic homeomorphism of $2^{\mathbb{N}}$ with no d-scrambled Cantor set.*

A *graph* on X is an irreflexive symmetric set $G \subseteq X \times X$. Let $\text{Hyp}^d(G)$ be the d -ary relation on X of all injective sequences $x \in X^d$ such that $G \cap (x[d] \times x[d]) \neq \emptyset$. An *embedding* of a d -ary relation R on X into a d -ary relation S on Y is an injection $\pi: X \rightarrow Y$ such that $x \in R \iff \pi \circ x \in S$ for all $x \in X^d$. The *saturation* of a set $Y \subseteq X$ under a bijection $T: X \rightarrow X$ is given by $[Y]_T = \bigcup_{n \in \mathbb{Z}} T^n[Y]$. A subset of a topological space is G_δ if it is a countable intersection of open sets and $G_{\delta\sigma}$ if it is a countable union of G_δ sets. The primary new technical result underlying our proof of Theorem 1 is the following:

Theorem 2 (ZF + DC). *Suppose that $d \geq 3$ and G is a $G_{\delta\sigma}$ graph on $2^{\mathbb{N}}$. Then there is a continuous embedding $\pi: 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ of $\text{Hyp}^d(G)$ into the d -ary Li–Yorkeness relation of a homeomorphism $T: 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ for which $2^{\mathbb{N}}$ is the one-point compactification of the T -saturation of $\pi[2^{\mathbb{N}}]$.*

In §1, we establish Theorem 2. In §2, we combine this with known facts to establish Theorem 1 and a related independence phenomenon.

2010 *Mathematics Subject Classification.* Primary 54H20; Secondary 03E15.

Key words and phrases. Li–Yorke chaos, scrambled set, topological dynamics.

1. EMBEDDING HYPERGRAPHS

Define $S: 2^{\mathbb{Z}} \rightarrow 2^{\mathbb{Z}}$ by $S(c)(i) = c(i+1)$ for all $i \in \mathbb{Z}$ and $c \in 2^{\mathbb{Z}}$. As every non-empty compact perfect zero-dimensional metric space is homeomorphic to $2^{\mathbb{N}}$ (see, for example, [Kec95, Proposition 7.4]), Theorem 2 is a consequence of the following fact:

Theorem 1.1 (ZF + DC). *Suppose that $d \geq 3$ and G is a $G_{\delta\sigma}$ graph on $2^{\mathbb{N}}$. Then there exist an S -invariant perfect set $F \subseteq 2^{\mathbb{Z}}$ whose finite sequences are proximal and a continuous embedding $\pi: 2^{\mathbb{N}} \rightarrow F$ of $\text{Hyp}^d(G)$ into the complement of the d -ary asymptoticity relation for which $0^{\mathbb{Z}}$ is the unique point of F not in the S -saturation of $\pi[2^{\mathbb{N}}]$.*

Proof. For all $s \in 2^{<\mathbb{N}}$, define $\mathcal{N}_s = \{c \in 2^{\mathbb{N}} \mid \forall i < |s| \ s(i) = c(i)\}$. For all $S \subseteq 2^{<\mathbb{N}}$, define $\mathcal{N}_S = \bigcup_{s \in S} \mathcal{N}_s$. For all $\ell \in \mathbb{N}$ and $p: 2^\ell \rightarrow d$, define $\ell_p = \ell$ and $\mathcal{N}_p = \prod_{i < d} \mathcal{N}_{p^{-1}(\{i\})}$. We say that p is a *near bijection* if it is surjective and there is a unique $k_p < d$ for which $|p^{-1}(\{k_p\})| > 1$. For $k \neq k_p$, let $p^{-1}(k)$ denote the unique element of $p^{-1}(\{k\})$.

Fix G_δ graphs G_n on $2^{\mathbb{N}}$ for which $G = \bigcup_{n \in \mathbb{N}} G_n$. For all $n \in \mathbb{N}$, fix a decreasing sequence $(G_{m,n})_{m \in \mathbb{N}}$ of open graphs whose intersection is G_n . Put $H_{m,n} = \text{Hyp}^d(G_{m,n})$ and $H_n = \bigcap_{m \in \mathbb{N}} H_{m,n}$, and observe that $\text{Hyp}^d(G) = \bigcup_{n \in \mathbb{N}} H_n$. For all $m, n \in \mathbb{N}$, let $P_{m,n}$ denote the set of all near bijections p for which $\ell_p \geq m$, $p(0^{\ell_p}) = 0$, and $\mathcal{N}_p \subseteq H_{m,n}$. Let $Q_{m,n}$ denote the set of all $q \in P_{m,n}$ for which there is no $p \in P_{m,n}$ such that $k_p = k_q$, $\ell_p < \ell_q$, and $p^{-1}(k) = q^{-1}(k) \upharpoonright \ell_p$ for all $k \in d \setminus \{k_p\}$.

Fix an infinite set $I \subseteq \mathbb{N} \setminus \{0\}$ such that $|I \cap (I - n)| < \aleph_0$ for all $n > 0$, as well as a bijection $\phi: I \rightarrow \mathbb{N} \cup \bigcup_{m,n \in \mathbb{N}} \{m\} \times \{n\} \times Q_{m,n}$ such that $I \cap [i, 2i + (n+1)(d-2)] = \{i\}$ whenever $\phi(i) = (m, n, q)$. Define $\pi: 2^{\mathbb{N}} \rightarrow 2^{\mathbb{Z}}$ by setting $\pi(c)(i) = 1$ if and only if

- (1) $i = 0$;
- (2) $i \in I$, $\phi(i) \in \mathbb{N}$, and $c(\phi(i)) = 1$; or
- (3) $\exists 0 < k < d \exists m, n \in \mathbb{N} \exists q \in Q_{m,n}$
 $((m, n, q) = \phi(i - (k-1)(n+1)) \text{ and } q(c \upharpoonright \ell_q) = k)$.

Clearly π is continuous.

Lemma 1.2. *The function π is injective.*

Proof. Suppose that $c, d \in 2^{\mathbb{N}}$ are distinct. Then there exists $n \in \mathbb{N}$ with $c(n) \neq d(n)$. Fix $i \in I$ for which $\phi(i) = n$. Then $\pi(b)(i) = b(n)$ for all $b \in \{c, d\}$, so $\pi(c)(i) \neq \pi(d)(i)$, thus $\pi(c) \neq \pi(d)$. $\square$

The *product metric* on $2^{\mathbb{Z}}$ is given by $\rho(c, d) = \sum_{i \in \mathbb{Z}} |c(i) - d(i)|/2^{|i|}$ for all $c, d \in 2^{\mathbb{Z}}$. A *homomorphism* from a d -ary relation R on a set Y to a d -ary relation S on a set Z is a function $\pi: Y \rightarrow Z$ such that

$\pi \circ y \in S$ for all $y \in R$. For all $i \in I$, define

$$N_i = \begin{cases} \{i\} & \text{if } \phi(i) \in \mathbb{N} \text{ and} \\ [i, i + (d-2)(n+1)] & \text{if } \phi(i) = (m, n, q). \end{cases}$$

Lemma 1.3. *The function π is a homomorphism from $\text{Hyp}^d(G)$ to the complement of the d -ary asymptoticity relation.*

Proof. Suppose that $c \in \text{Hyp}^d(G)$, and fix $n \in \mathbb{N}$ and $j, k < d$ such that $c(j) G_n c(k)$. By permuting the coordinates of c , we can assume that either $c(0) = 0^\mathbb{N}$ or both $0^\mathbb{N} \notin c[d]$ and $j, k > 0$. Fix $k' \in d \setminus \{j, k\}$ if $c(0) = 0^\mathbb{N}$ and set $k' = 0$ otherwise. If $m \in \mathbb{N}$ is sufficiently large that $\text{proj}_{2^m} \circ c$ is injective and $0^m \in (\text{proj}_{2^m} \circ c)[d] \iff 0^\mathbb{N} \in c[d]$, then there is a least $\ell \geq m$ for which there is a nearly bijective extension $q \in P_{m,n}$ of $(\text{proj}_{2^\ell} \circ c)^{-1}$ such that $k_q = k'$ and $\ell_q = \ell$. Then $q \in Q_{m,n}$. Fix $i \in I$ with $\phi(i) = (m, n, q)$, and observe that $\text{proj}_{2^{N_i}} \circ \pi \circ c$ is injective, so $\min_{j < k < d} \rho(S^i(\pi(c(j))), S^i(\pi(c(k)))) \geq 1/2^{(d-2)(n+1)}$. $\square$

Lemma 1.4. *The function π is a homomorphism from $(2^\mathbb{N})^d \setminus \text{Hyp}^d(G)$ to the d -ary asymptoticity relation.*

Proof. Suppose that $c \in (2^\mathbb{N})^d$ and $\pi \circ c$ is not asymptotic. Fix $\epsilon > 0$ such that $I' = \{i' \in \mathbb{N} \mid \min_{j < k < d} \rho(S^{i'}(\pi(c(j))), S^{i'}(\pi(c(k)))) \geq \epsilon\}$ is infinite, as well as $r \in \mathbb{N}$ such that $\sum_{|i| > r} 1/2^{|i|} < \epsilon$. By throwing out finitely many elements of I' , we can assume that $[i' - r, i' + r]$ intersects at most one set of the form N_i for all $i' \in I'$. Our choice of r then yields a unique such $i \in I$, in which case $\text{proj}_{2^{N_i}} \circ \pi \circ c$ is injective. As $d \geq 3$, there exist $m, n \in \mathbb{N}$, $q \in Q_{m,n}$, and $\sigma \in \text{Sym}(d)$ for which $(d-2)(n+1) \leq 2r$, $\phi(i) = (m, n, q)$, and $c \circ \sigma \in \mathcal{N}_q$. By passing to an infinite subset of I' , we can assume that the values of k_q , n , and σ are independent of $i' \in I'$. As the definition of $Q_{m,n}$ then ensures that at most $4r + 1$ distinct elements of I' correspond to the same value of m , it follows that $c \in \bigcap_{m \in \mathbb{N}} H_{m,n} = H_n \subseteq \text{Hyp}^d(G)$. $\square$

Let F be the closure of the S -saturation of $\pi[2^\mathbb{N}]$. Clearly F is perfect.

Lemma 1.5. *The sequence $0^\mathbb{Z}$ is the unique element of $F \setminus [\pi[2^\mathbb{N}]]_S$.*

Proof. Suppose that $x \in F \setminus [\pi[2^\mathbb{N}]]_S$. Then there exist a sequence $(c_n)_{n \in \mathbb{N}}$ of elements of $2^\mathbb{N}$, $\epsilon \in \{\pm 1\}$, and a strictly increasing sequence $(i_n)_{n \in \mathbb{N}}$ of natural numbers such that $S^{\epsilon i_n}(\pi(c_n)) \rightarrow x$. If $\epsilon = -1$, then clearly $x = 0^\mathbb{Z}$. If $\epsilon = 1$ and $x \neq 0^\mathbb{Z}$, then our choice of I ensures that x is in the S -orbit of $\pi(0^\mathbb{N})$, a contradiction. $\square$

Lemma 1.5 implies that $\bigcup \{\mathbb{Z} \setminus x^{-1}(\{0\}) \mid x \in F'\}$ is contained in a finite union of translates of $\{0\} \cup \bigcup_{i \in I} N_i$ whenever $F' \subseteq F$ is finite. As

the defining property of I ensures that such a set has arbitrarily long gaps, every finite sequence of elements of F is proximal. $\square$

A set $Y \subseteq X$ is a G -clique if all distinct elements of Y are G -related, and G -independent if no elements of Y are G -related.

Proposition 1.6 (ZF + DC). *Suppose that $d \geq 2$, X and Y are Polish spaces, G is a Borel graph on X , $T: Y \rightarrow Y$ is a Borel automorphism, $B \subseteq Y$ is an uncountable Borel d -scrambled set, $\pi: X \rightarrow Y$ is an injective Borel homomorphism from $X^d \setminus \text{Hyp}^d(G)$ to the complement of the d -ary Li–Yorkeness relation, and the saturation of $\pi[X]$ is co-countable. Then G has a perfect clique.*

Proof. Fix $i \in \mathbb{Z}$ for which $B \cap (T^i \circ \pi)[X]$ is uncountable. The perfect set theorem and Galvin’s Theorem (see, for example, [Kec95, Theorems 13.6 and 19.7]) then yield a perfect set $P \subseteq (T^i \circ \pi)^{-1}(B)$ that is either a G -clique or G -independent. But G -independence would contradict the fact that B is d -scrambled. $\square$

2. D -SCRAMBLED SETS

We begin this section with the proof of our main result:

Proof (of Theorem 1). By [KV12, Theorem 2.1] and a simple Baire category argument, there is an F_σ graph G on $2^\mathbb{N}$ with an uncountable clique $Y \subseteq 2^\mathbb{N}$ but no perfect clique. By Theorem 2, there is a continuous embedding $\pi: 2^\mathbb{N} \rightarrow 2^\mathbb{N}$ of $\text{Hyp}^d(G)$ into the d -ary Li–Yorkeness relation of a homeomorphism $T: 2^\mathbb{N} \rightarrow 2^\mathbb{N}$ for which the saturation of $\pi[2^\mathbb{N}]$ is co-countable. Then $\pi[Y]$ is an uncountable d -scrambled set and there is no perfect d -scrambled set by Proposition 1.6. $\square$

Let $\mathfrak{c}$ denote the cardinality of the continuum.

Theorem 2.1 (ZFC + $\neg\text{CH}$). *Suppose that $d \geq 3$. Then it is consistent that there is a homeomorphism $T: 2^\mathbb{N} \rightarrow 2^\mathbb{N}$ with a d -scrambled set of cardinality $\mathfrak{c}$ but no perfect d -scrambled set.*

Proof. By [She99, Theorem 1.13], we can assume that there is an F_σ graph G on $2^\mathbb{N}$ with a clique of cardinality $\mathfrak{c}$ but no perfect clique. The proof of Theorem 1 therefore yields the desired result. $\square$

A *hypergraph* is a permutation-invariant set of injective sequences.

Theorem 2.2 (ZFC + $\neg\text{CH}$). *Suppose that $d \geq 3$. Then it is consistent that every continuous function on a Polish space with a d -scrambled set of cardinality $\aleph_2$ has a d -scrambled perfect set.*

Proof. By [KS03, Proposition 3.4], we can assume that every Borel d -ary hypergraph on a Polish space with a clique of cardinality $\aleph_2$ has a perfect clique. But the d -ary Li–Yorke relation is Borel. $\square$

Remark. Given any set $D \subseteq \mathbb{N} \setminus 3$, the underlying ideas can be interleaved to obtain generalizations of our results to individual sets that are simultaneously d -scrambled for all $d \in D$.

Acknowledgements. I would like to thank Alexander Kechris for encouraging me to utilize ChatGPT-5.6 Sol, which provided substantial assistance in the discovery and development of the results.

REFERENCES

- [GGM21] Stefan Geschke, Jan Grebík, and Benjamin D. Miller, *Scrambled Cantor sets*, Proceedings of the American Mathematical Society **149** (2021), no. 10, 4461–4468.
- [Kec95] Alexander S. Kechris, *Classical Descriptive Set Theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995.
- [KS03] Wiesław Kubiś and Saharon Shelah, *Analytic colorings*, Annals of Pure and Applied Logic **121** (2003), no. 2–3, 145–161.
- [KV12] Wiesław Kubiś and Benjamin Vejnar, *Covering an uncountable square by countably many continuous functions*, Proceedings of the American Mathematical Society **140** (2012), no. 12, 4359–4368.
- [She99] Saharon Shelah, *Borel sets with large squares*, Fundamenta Mathematicae **159** (1999), no. 1, 1–50. MR 1669643

B. MILLER, 1008 BALSABWOOD DRIVE, DURHAM, NC 27705

Email address: glimmeffros@gmail.com

URL: <https://glimmeffros.github.io>