Distance-regular graphs obtained from the Mathieu groups

Dean Crnković (deanc@math.uniri.hr)

Nina Mostarac (nmavrovic@math.uniri.hr)

and

Andrea Švob (asvob@math.uniri.hr)

Department of Mathematics
University of Rijeka

Radmile Matejčić 2, 51000 Rijeka, Croatia

Abstract

In this paper we construct distance-regular graphs admitting a transitive action of the five sporadic simple groups discovered by E. Mathieu, the Mathieu groups M_{11} , M_{12} , M_{22} , M_{23} and M_{24} . From the code spanned by the adjacency matrix of the strongly regular graph with parameters (176,70,18,34) we obtain block designs having the full automorphism groups isomorphic to the Higman-Sims finite simple group. Further, we discuss a possibility of permutation decoding of the codes spanned by the adjacency matrices of the graphs constructed and find small PD-sets for some of the codes.

AMS classification numbers: 05E18, 05E30, 94B05, 05B05.

Keywords: Mathieu group, distance-regular graph, strongly regular graph, block design, permutation decoding, PD-set.

1 Introduction

The main motivation for this paper is to give further contribution to the classification of transitive distance regular graphs (DRGs), especially those admitting a transitive action of a simple group. The research presented in the paper can be seen as a continuation of the work given in [10], in which transitive structures constructed from the Mathieu group M_{11} were described. On the other hand, this is also a continuation of the work by Praeger and Soicher [31], where they study the graphs admitting a sporadic simple group or its automorphism group as a vertex-transitive group of automorphisms of rank at most 5. However, in this paper we are also interested in graphs admitting a vertex-transitive automorphism groups of ranks greater than 5. In the literature, there are also other examples of the research in design and graph theory in which finite groups had important contribution (see [5, 7, 12, 13, 28]). In this paper we are focused on DRGs admitting a transitive action of one of the five Mathieu sporadic simple groups.

For relevant background reading in the group theory we refer the reader to [9, 32], to [5, 34] for the theory of strongly regular graphs, to [7, 14] for the theory of distance-regular graphs, and to [17, 23] for coding theory.

In this paper we study the Mathieu groups M_{11} , M_{12} , M_{22} , M_{23} and M_{24} , which are the sporadic simple groups of orders 7920, 95040, 443520, 10200960 and 244823040, respectively. The Mathieu groups M_{11} , M_{12} , M_{22} , M_{23} and M_{24} were found by Emile Mathieu in [24, 25, 26]. They are multiply transitive permutation groups on 11, 12, 22, 23 or 24 objects, and they were the first sporadic groups discovered (see [35, 36]). The Mathieu groups have been studied so far in various literature and in various settings. For example, t-designs arising from Mathieu groups M_{22} , M_{23} and M_{24} have been studied in a work by Kramer, Magliveras and Mesner (see [20]) and those arising from M_{11} in [10]. Codes connected with Mathieu group M_{11} have been studied in [27] and those connected with M_{12} in [2]. More about small representations (up to rank 5) of these five sporadic simple groups can be found in [31]. We refer the reader to [9, 36] for more details about these groups. Using the method outlined in Section 3, in this paper we constructed and partially classified DRGs (SRGs and DRGs of diameter $d \geq 3$)

from above mentioned simple groups.

We also study codes spanned by the adjacency matrices of the constructed DRGs. Codes with large automorphism groups are suitable for permutation decoding (see [18, 19, 22]), the decoding method developed by Jessie MacWilliams in the early 60s that can be used when a linear code has a sufficiently large automorphism group to ensure the existence of a set of automorphisms, called a PD-set, that has some specific properties. Therefore, the codes constructed in this paper are suitable for permutation decoding, and we found small PD-sets for some of the constructed codes. From the adjacency matrix of the SRG with parameters (176,70,18,34), beside the well-known symmetric block design with parameters (176,50,14) having the full automorphism group isomorphic to the Higman-Sims group, we obtain block designs with parameters 2-(176,56,110), 2-(176,64,540), 2-(176,66,780), 2-(176,70,2760), 2-(176,72,2556), 2-(176,78,37752), 2-(176,80,124030), 2-(176,82,99630), 2-(176,86,87720) and 2-(176,88,210540) as support designs. The full automorphism groups of the new block designs are also isomorphic to the Higman-Sims group HS. Further, one of the constructed SRGs with parameters (144,55,22,20) is new, up to our best knowledge.

To find the graphs and compute their full automorphism groups, and to obtain PD-sets and the corresponding information sets of the codes, we used programmes written for Magma [4] and GAP [15]. The constructed DRGs, including the SRGs, and the obtained PD-sets and the corresponding information sets of the codes can be found at the link:

http://www.math.uniri.hr/~asvob/DRGs_Mathieu_PD.7z.

2 Preliminaries

In this section we begin with the required definitions and notation.

Definition 1. A coherent configuration on a finite non-empty set Ω is an ordered pair (Ω, \mathcal{R}) with $\mathcal{R} = \{R_0, R_1, \dots, R_d\}$ a set of non-empty relations on Ω , such that the following axioms hold.

- (i) $\sum_{i=0}^{t} R_i$ is the identity relation, where $\{R_0, R_1, \dots, R_t\} \subseteq \{R_0, R_1, \dots, R_d\}$.
- (ii) \mathcal{R} is a partition of Ω^2 .
- (iii) For every relation $R_i \in \mathcal{R}$, its converse $R_i^T = \{(y, x) : (x, y) \in R_i\}$ is in \mathcal{R} .
- (iv) There are constants p_{ij}^k known as the intersection numbers of the coherent configuration \mathcal{R} , such that for $(x,y) \in R_k$, the number of elements z in Ω for which $(x,z) \in R_i$ and $(z,y) \in R_j$ equals p_{ij}^k .

We say that a coherent configuration is homogeneous if it contains the identity relation, i.e., if $R_0 = I$. If \mathcal{R} is a set of symmetric relations on Ω , then a coherent configuration is called symmetric. A symmetric coherent configuration is homogeneous (see [8]). Symmetric coherent configurations are introduced by Bose and Shimamoto in [3] and called association schemes. An association scheme with relations $\{R_0, R_1, \ldots, R_d\}$ is called a d-class association scheme.

Let Γ be a graph with diameter d, and let $\delta(u,v)$ denote the distance between vertices u and v of Γ . The ith-neighborhood of a vertex v is the set $\Gamma_i(v) = \{w : \delta(v,w) = i\}$. Similarly, we define Γ_i to be the ith-distance graph of Γ , that is, the vertex set of Γ_i is the same as for Γ , with adjacency in Γ_i defined by the ith distance relation in Γ . We say that Γ is distance-regular if the distance relations of Γ give the relations of a d-class association scheme, that is, for every choice of $0 \le i, j, k \le d$, all vertices v and w with $\delta(v, w) = k$ satisfy $|\Gamma_i(v) \cap \Gamma_j(w)| = p_{ij}^k$ for some constant p_{ij}^k . In a distance-regular graph, we have that $p_{ij}^k = 0$ whenever i + j < k or k < |i - j|. A distance-regular graph Γ is necessarily regular with degree p_{11}^0 ; more generally, each distance graph Γ_i is regular with degree $k_i = p_{ii}^0$.

An equivalent definition of distance-regular graphs is the existence of the constants $b_i = p_{i+1,1}^i$ and $c_i = p_{i-1,1}^i$ for $0 \le i \le d$ (notice that $b_d = c_0 = 0$). The sequence $\{b_0, b_1, \ldots, b_{d-1}; c_1, c_2, \ldots, c_d\}$, where d is the diameter of Γ is called the intersection array of Γ . Clearly, $b_0 = k$, $b_d = c_0 = 0$, $c_1 = 0$.

A regular graph is strongly regular with parameters (v, k, λ, μ) if it has v vertices, degree k, and if any two adjacent vertices are together adjacent to λ vertices, while any two non-adjacent vertices are together adjacent to μ vertices. A strongly regular graph with parameters (v, k, λ, μ) is usually denoted by $SRG(v, k, \lambda, \mu)$. A strongly regular graph is a distance-regular graph with diameter 2 whenever $\mu \neq 0$. The intersection array of an SRG is given by $\{k, k-1-\lambda; 1, \mu\}$.

3 DRGs constructed from the Mathieu groups

Let G be a finite permutation group acting on the finite set Ω . This action induce the action of the group G on the set $\Omega \times \Omega$. For more information see [35]. The orbits of this action are the sets of the form $\{(\alpha g, \beta g) : g \in G\}$. If G is transitive, then $\{(\alpha, \alpha) : \alpha \in \Omega\}$ is one such orbit. If the rank of G is r, then it has r orbits on $\Omega \times \Omega$. Let $|\Omega| = n$ and Δ_i is one of these orbits. We say that the $n \times n$ matrix A_i , with rows and columns indexed by Ω and entries

$$A_i(\alpha, \beta) = \begin{cases} 1, & \text{if } (\alpha, \beta) \in \Delta_i \\ 0, & \text{otherwise.} \end{cases}$$

is called the adjacency matrix for the orbit Δ_i .

Let A_0, \ldots, A_{r-1} be the adjacency matrices for the orbits of G on $\Omega \times \Omega$. These satisfy the following conditions.

- (i) $A_0 = I$, if G is transitive on Ω . If G has s orbits on Ω , then I is a sum of s adjacency matrices.
- (ii) $\sum_{i} A_{i} = J$, where J is the all-one matrix.
- (iii) If A_i is an adjacency matrix, then so is its transpose A_i^T .
- (iv) If A_i and A_j are adjacency matrices, then their product is an integer-linear-combination of adjacency matrices.

If A_i is symmetric, then the corresponding orbit is called self-paired. Further, if $A_i = A_j^T$, then the corresponding orbits are called mutually paired.

The graphs obtained in this paper are constructed using the method described in [11] which can be rewritten in terms of coherent configurations in the following way.

Theorem 1. Let G be a finite permutation group acting transitively on the set Ω and A_0, \ldots, A_d be the adjacency matrices for orbits of G on $\Omega \times \Omega$. Let $\{B_1, \ldots, B_t\} \subseteq \{A_1, \ldots, A_d\}$ be a set of adjacency matrices for the self-paired or mutually paired orbits. Then $M = \sum_{i=1}^t B_i$ is the adjacency matrix of a regular graph Γ . The group G acts transitively on the set of vertices of the graph Γ .

Using this method one can construct all regular graphs admitting a transitive action of the group G. We will be interested only in those regular graphs that are distance-regular, and especially strongly regular.

Remark 1. Because of the large number of possibilities for building the first row of the adjacency matrix of a DRG, the only way to obtain the classification of DRGs given in this paper was with the use of computers. The running time complexity of the algorithm used for the construction of graphs depends on a number of parameters, such as the size of the used subgroup, the number of orbits of a vertex stabilizer, the number of vertices of the graphs and the number of self-paired and mutually paired orbits in a particular case.

3.1 DRGs from the group M_{11}

The Mathieu group M_{11} has the order 7920 and up to conjugation has 39 subgroups. In Table 1 we give the list of all the subgroups $H_i^1 \leq M_{11}$ which lead to the construction of SRGs or DRGs of diameter $d \geq 3$.

Subgroup	Structure	Order	Index	Rank	Primitive
H_1^1	S_5	144	55	3	yes
H_2^1	$Z_9:QD16$	120	66	4	yes
H_3^1	$Z_{11}: Z_5$	55	144	6	no
H_4^1	GL(2,3)	48	165	8	yes
H_5^1	$S_3 \times S_3$	36	220	16	no
H_6^1	S_4	24	330	23	no

Table 1: Subgroups of the group M_{11}

Using the method described in Theorem 1 we obtained all DRGs with at most 2000 vertices and for which the rank of the permutation representation of the group is at most 25, i.e. we gave the classification of such DRGs.

Theorem 2. Up to isomorphism there are exactly five strongly regular graphs and exactly three distance-regular graphs of diameter $d \geq 3$ with at most 2000 vertices and for which the rank of the permutation representation of the group is at most 25, admitting a transitive action of the group M_{11} . The SRGs have parameters (55, 18, 9, 4), (66, 20, 10, 4), (144, 55, 22, 20), (144, 66, 30, 30) and (330, 63, 24, 9), and the DRGs have 165, 220 and 330 vertices, respectively. Details about the obtained strongly regular graphs are given in Table 2 and details about the obtained DRGs with $d \geq 3$ are given in Table 3.

Graph Γ	Parameters	$Aut(\Gamma)$
$\Gamma_1^1 = \Gamma(M_{11}, H_1^1)$	(55,18,9,4)	S_{11}
$\Gamma_2^1 = \Gamma(M_{11}, H_2^1)$	(66,20,10,4)	S_{12}
$\Gamma_3^1 = \Gamma(M_{11}, H_3^1)$	(144,55,22,20)	M_{11}
$\Gamma_4^1 = \Gamma(M_{11}, H_3^1)$	(144,66,30,30)	$M_{12}: Z_2$
$\Gamma_5^1 = \Gamma(M_{11}, H_6^1)$	(330,63,24,9)	S_{11}

Table 2: SRGs constructed from the group M_{11}

Graph Γ	Number of vertices	Diameter	Intersection array	$Aut(\Gamma)$
$\Gamma_6^1 = \Gamma(M_{11}, H_4^1)$	165	3	{24, 14, 6; 1, 4, 9}	S_{11}
$\Gamma_7^1 = \Gamma(M_{11}, H_5^1)$	220	3	$\{27, 16, 7; 1, 4, 9\}$	S_{12}
$\Gamma_8^1 = \Gamma(M_{11}, H_6^1)$	330	4	$\{28, 18, 10, 4; 1, 4, 9, 16\}$	S_{11}

Table 3: DRGs constructed from the group M_{11} , $d \ge 3$

Proof. There are 39 conjugacy classes of subgroups of M_{11} , but only 19 of them lead to a permutation representation of rank at most 25 and of index at most 2000. Applying the method described in Theorem 1 to the permutation representations on cosets of these 19 subgroups we obtain the results. \Box

Remark 2. All SRGs given in Table 2 are isomorphic to the ones constructed in [10]. Since the SRG Γ_3^1 does not yield a partial geometry, it cannot be obtained from an orthogonal array.

Remark 3. The graphs Γ_6^1 , Γ_7^1 and Γ_8^1 are unique graphs with the given intersection arrays, known as Johnson graphs, J(11,3), J(12,3) and J(11,4), respectively (see [7]).

3.2 DRGs from the group M_{12}

The Mathieu group M_{12} has the order 95040 and up to conjugation has 147 subgroups. In Table 4 we give the list of all the subgroups $H_i^2 \leq M_{12}$ which lead to the construction of SRGs or DRGs of diameter $d \geq 3$.

Subgroup	Structure	Order	Index	Rank	Primitive
H_1^2	$(A_6.Z_2):Z_2$	1440	66	3	yes
H_2^2	L(2, 11)	660	144	5	no
H_3^2	$((E_9:Q_8):Z_3):Z_2$	432	220	5	yes
H_4^2	$((E_8:E_4):Z_3):Z_2$	192	495	11	yes
H_5^2	S_5	120	792	15	no

Table 4: Subgroups of the group M_{12}

Using the method described in Theorem 1 we obtained all DRGs with at most 2000 vertices and for which the rank of the permutation representation of the group is at most 20, i.e. we gave the classification of such DRGs.

Theorem 3. Up to isomorphism there are exactly seven strongly regular graphs and exactly three distance-regular graphs of diameter $d \geq 3$ with at most 2000 vertices and for which the rank of the permutation representation of the group is at most 20, admitting a transitive action of the group M_{12} . The SRGs have parameters (66, 20, 10, 4), (144, 66, 30, 30), (144, 55, 22, 20), (144, 22, 10, 2) and (495, 238, 109, 119), and the DRGs have 220, 495 and 792 vertices, respectively. Details about the obtained strongly regular graphs are given in Table 5 and details about the obtained DRGs with $d \geq 3$ are given in Table 6.

Graph Γ	Parameters	$Aut(\Gamma)$
$\Gamma_1^2 = \Gamma(M_{12}, H_1^2)$	(66,20,10,4)	S_{12}
$\Gamma_2^2 = \Gamma(M_{12}, H_2^2)$	(144,66,30,30)	$M_{12}:Z_2$
$\Gamma_3^2 = \Gamma(M_{12}, H_2^2)$	(144,66,30,30)	$M_{12}:Z_2$
$\Gamma_4^2 = \Gamma(M_{12}, H_2^2)$	(144,66,30,30)	M_{12}
$\Gamma_5^2 = \Gamma(M_{12}, H_2^2)$	(144,55,22,20)	$M_{12}:Z_2$
$\Gamma_6^2 = \Gamma(M_{12}, H_2^2)$	(144,22,10,2)	$S_{12} \wr S_2$
$\Gamma_7^2 = \Gamma(M_{12}, H_4^2)$	(495, 238, 109, 119)	$O^-(10,2): Z_2$

Table 5: SRGs constructed from the group M_{12}

Graph Γ	Number of vertices	Diameter	Intersection array	$Aut(\Gamma)$
$\Gamma_8^2 = \Gamma(M_{12}, H_3^2)$	220	3	{27, 16, 7; 1, 4, 9}	S_{12}
$\Gamma_9^2 = \Gamma(M_{12}, H_4^2)$	495	4	$\{32,21,12,5;1,4,9,16\}$	S_{12}
$\Gamma_{10}^2 = \Gamma(M_{12}, H_5^2)$	792	5	$\{35, 24, 15, 8, 3; 1, 4, 9, 16, 25\}$	S_{12}

Table 6: DRGs constructed from the group M_{12} , $d \geq 3$

Proof. There are 147 conjugacy classes of subgroups of M_{12} , but only 31 of them lead to a permutation representation of rank at most 20 and of index at most 2000. Applying the method described in Theorem 1 to the permutation representations on cosets of these 31 subgroups we obtain the results. \Box

Remark 4. The strongly regular graph Γ_1^2 is isomorphic to the triangular graph T(12). The adjacency matrices of non-isomorphic SRGs Γ_2^2 , Γ_3^2 and Γ_4^2 are the incidence matrices of symmetric designs with parameters (144, 66, 30), designs with Menon parameters (related to

a regular Hadamard matrix of order 144). These symmetric designs have been described in [21, 37]. According to Brouwer's table (see [6]), known graphs with the parameters equal to the parameters of the graph Γ_5^2 (not isomorphic to Γ_3^1) are obtainable from orthogonal arrays OA(12,5). Since the SRG Γ_5^2 cannot be obtained from orthogonal array, our graph is new. So far, according to [6], the graphs Γ_3^1 and Γ_5^2 are the only known graphs with these parameters not arising from orthogonal array. The graph Γ_6^2 is unique graph with the given parameters and the graph Γ_7^2 is isomorphic to the $O^-(10,2)$ polar graph. Strongly regular graphs with parameters (144,66,30,30) have been known before (see [5,6]).

Remark 5. The graphs Γ_8^2 , Γ_9^2 and Γ_{10}^2 are unique graphs with the given intersection arrays, known as Johnson graphs, J(12,3), J(12,4) and J(12,5), respectively (see [7]).

3.3 DRGs from the group M_{22}

The Mathieu group M_{22} has the order 443520 and up to conjugation 156 subgroups. In Table 7 we give the list of all the subgroups $H_i^3 \leq M_{22}$ which lead to the construction of SRGs or DRGs of diameter $d \geq 3$.

Subgroup	Structure	Order	Index	Rank	Primitive
H_1^3	$E_{16}: A_6$	5760	77	3	yes
H_2^3	A_7	2520	176	3	yes
H_3^3	$E_{16}: S_5$	1920	231	4	yes
H_4^3	$E_8: L(3,2)$	1344	330	5	yes
H_5^3	L(2, 11)	660	672	6	yes
H_{6}^{3}	$(A_4 \times A_4) : Z_2$	288	1540	22	no

Table 7: Subgroups of the group M_{22}

Using the method described in Theorem 1 we obtained all DRGs with at most 2000 vertices and for which the rank of the permutation representation of the group is at most 30, i.e. we gave the classification of such DRGs.

Theorem 4. Up to isomorphism there are exactly five strongly regular graphs and exactly three distance-regular graphs of diameter $d \geq 3$ with at most 2000 vertices and for which

the rank of the permutation representation of the group is at most 30, admitting a transitive action of the group M_{22} . The SRGs have parameters (77, 16, 0, 4), (176, 70, 18, 34), (231, 30, 9, 3), (231, 40, 20, 4) and (672, 176, 40, 48), and the DRGs have 330, 672 and 1540 vertices, respectively. Details about the obtained strongly regular graphs are given in Table 8 and details about the obtained DRGs with $d \geq 3$ are given in Table 9.

Graph Γ	Parameters	$Aut(\Gamma)$
$\Gamma_1^3 = \Gamma(M_{22}, H_1^3)$	(77, 16, 0, 4)	$M_{22}: Z_2$
$\Gamma_2^3 = \Gamma(M_{22}, H_2^3)$	(176, 70, 18, 34)	M_{22}
$\Gamma_3^3 = \Gamma(M_{22}, H_3^3)$	(231, 30, 9, 3)	$M_{22}: Z_2$
$\Gamma_4^3 = \Gamma(M_{22}, H_3^3)$	(231, 40, 20, 4)	S_{22}
$\Gamma_5^3 = \Gamma(M_{22}, H_5^3)$	(672, 176, 40, 48)	$(U(6,2):Z_2):Z_2$

Table 8: SRGs constructed from the group M_{22}

Graph Γ	Number of vertices	Diameter	Intersection array	$Aut(\Gamma)$
$\Gamma_6^3 = \Gamma(M_{22}, H_4^3)$	330	4	{7, 6, 4, 4; 1, 1, 1, 6}	$M_{22}: Z_2$
$\Gamma_7^3 = \Gamma(M_{22}, H_5^3)$	672	3	{110, 81, 12; 1, 18, 90}	$M_{22}: Z_2$
$\Gamma_8^3 = \Gamma(M_{22}, H_6^3)$	1540	3	$\{57, 36, 17; 1, 4, 9\}$	S_{22}

Table 9: DRGs constructed from the group M_{22} , $d \geq 3$

Proof. There are 156 conjugacy classes of subgroups of M_{22} , but only 21 of them lead to a permutation representation of rank at most 30 and of index at most 2000. Applying the method described in Theorem 1 to the permutation representations on cosets of these 21 subgroups we obtain the results. \Box

Remark 6. The strongly regular graphs Γ_1^3 and Γ_2^3 are unique graphs with these parameters. The graph Γ_3^3 is isomorphic to the SRG known as the Cameron graph. The SRG Γ_4^3 is isomorphic to the triangular graph T(22) and Γ_5^3 is isomorphic to the graph known as U(6,2)-graph. For more information we refer the reader to [5, 6].

Remark 7. The graph Γ_6^3 is isomorphic to the graph known as M_{22} -graph or doubly truncated Witt graph. The graph Γ_7^3 is isomorphic to the one constructed by Soicher in [33]. So far, it

is the only known example of DRG with this intersection array. The graph Γ_8^3 is known as Johnson graph J(22,3). (see [7])

3.4 DRGs from the group M_{23}

The Mathieu group M_{23} has order 10200960 and up to conjugation 204 subgroups. In Table 10 we give the list of all the subgroups $H_i^4 \leq M_{23}$ which lead to the construction of SRGs or DRGs of diameter $d \geq 3$.

Subgroup	Structure	Order	Index	Rank	Primitive
H_1^4	$L(3,4):Z_2$	40320	253	3	yes
H_2^4	$E_{16}: A_7$	40320	253	3	yes
H_3^4	A_8	20160	506	4	yes
H_4^4	M_{11}	7920	1288	4	yes
H_5^4	$E_{16}:(A_5:S_3)$	5760	1771	8	yes

Table 10: Subgroups of the group M_{23}

Using the method described in Theorem 1 we obtained all DRGs with at most 10000 vertices and for which the rank of the permutation representation of the group is at most 20, i.e. we gave the classification of such DRGs.

Theorem 5. Up to isomorphism there are exactly three strongly regular graphs and exactly two distance-regular graphs of diameter $d \geq 3$ with at most 10000 vertices and for which the rank of the permutation representation of the group is at most 20, admitting a transitive action of the group M_{23} . The SRGs have parameters (253, 42, 21, 4), (253, 112, 36, 60) and (1288, 495, 206, 180), and the DRGs have 506 and 1771 vertices, respectively. Details about the obtained strongly regular graphs are given in Table 11 and details about the obtained DRGs with $d \geq 3$ are given in Table 12.

Graph Γ	Parameters	$Aut(\Gamma)$
$\Gamma_1^4 = \Gamma(M_{23}, H_1^4)$	(253,42,21,4)	S_{23}
$\Gamma_2^4 = \Gamma(M_{23}, H_2^4)$	(253,112,36,60)	M_{23}
$\Gamma_3^4 = \Gamma(M_{23}, H_4^4)$	(1288, 495, 206, 180)	M_{24}

Table 11: SRGs constructed from the group M_{23}

Graph Γ	Number of vertices	Diameter	Intersection array	$Aut(\Gamma)$
$\Gamma_4^4 = \Gamma(M_{23}, H_3^4)$	506	3	{15, 14, 12; 1, 1, 9}	M_{23}
$\Gamma_5^4 = \Gamma(M_{23}, H_5^4)$	1771	3	$\{60, 38, 18; 1, 4, 9\}$	S_{23}

Table 12: DRG constructed from the group $M_{23}, d \geq 3$

Proof. There are 204 conjugacy classes of subgroups of M_{23} , but only 14 of them lead to a permutation representation of rank at most 20 and of index at most 10000. Applying the method described in Theorem 1 to the permutation representations on cosets of these 14 subgroups we obtain the results. \square

Remark 8. The graph Γ_1^4 is isomorphic to the triangular graph T(23). The graph Γ_2^4 can be constructed from the group M_{23} as a rank 3 graph, and Γ_3^4 (isomorphic to the graph Γ_2^5) can be constructed from the group M_{24} as a rank 3 graph.

Remark 9. The graph Γ_4^4 is isomorphic to the distance-regular graph that can be obtained from residual design of Steiner system S(5,8,24). The graph Γ_5^4 is known as Johnson graph J(23,3). For more information we refer the reader to [7].

3.5 DRGs from the Mathieu group M_{24}

The Mathieu group M_{24} has order 244823040 and up to conjugation 1529 subgroups. In Table 13 we give the list of all the subgroups $H_i^5 \leq M_{24}$ which lead to the construction of SRGs or DRGs of diameter $d \geq 3$.

Subgroup	Structure	Order	Index	Rank	Primitive
H_1^5	$M_{22}: Z_2$	887040	276	3	yes
H_2^5	$E_{16}: A_8$	322560	759	4	yes
H_3^5	$M_{12}:Z_2$	190080	1288	3	yes
H_4^5	$(L(3,4):Z_3):Z_2$	120960	2024	5	yes

Table 13: Subgroups of the group M_{24}

Using the method described in Theorem 1 we obtained all DRGs with at most 10000 vertices and for which the rank of the permutation representation of the group is at most 20, i.e. we gave the classification of such DRGs.

Theorem 6. Up to isomorphism there are exactly two strongly regular graphs and exactly two distance-regular graphs of diameter $d \geq 3$ with at most 10000 vertices and for which the rank of the permutation representation of the group is at most 20, admitting a transitive action of the group M_{24} . The SRGs have parameters (276, 44, 22, 4) and (1288, 495, 206, 180), and the DRGs have 759 and 2024 vertices, respectively. Details about the obtained strongly regular graphs are given in Table 14 and details about the obtained DRGs with $d \geq 3$ are given in Table 15.

Graph Γ	Parameters	$Aut(\Gamma)$
$\Gamma_1^5 = \Gamma(M_{24}, H_1^5)$	(276, 44, 22, 4)	S_{24}
$\Gamma_2^5 = \Gamma(M_{24}, H_3^5)$	(1288, 495, 206, 180)	M_{24}

Table 14: SRGs constructed from the group M_{24}

Graph Γ	Number of vertices	Diameter	Intersection array	$Aut(\Gamma)$
$\Gamma_3^5 = \Gamma(M_{24}, H_2^5)$	759	3	{30, 28, 24; 1, 3, 15}	M_{24}
$\Gamma_4^5 = \Gamma(M_{24}, H_4^5)$	2024	3	{63, 40, 19; 1, 4, 9}	S_{24}

Table 15: DRGs constructed from the group M_{24} , $d \geq 3$

Proof. There are 1529 conjugacy classes of subgroups of M_{24} , but only 15 of them lead to a permutation representation of rank at most 20 and of index at most 10000. Applying

the method described in Theorem 1 to the permutation representations on cosets of these 15 subgroups we obtain the results. \Box

Remark 10. The graph Γ_1^5 is isomorphic to the triangular graph T(24). The graph Γ_2^5 (isomorphic to the graph Γ_3^4) can be constructed from the group M_{24} as a rank 3 graph.

Remark 11. The graph Γ_3^5 is unique distance-regular graph known as near hexagon which can be obtained from Steiner system S(5,8,24). The graph Γ_4^5 is known as Johnson graph J(24,3). For more information we refer the reader to [7].

4 Codes

A code C of length n over the alphabet Q is a subset $C \subseteq Q^n$. Elements of the code are called codewords. A code C is called a p-ary linear code of dimension m if $Q = \mathbb{F}_p$, for a prime power p, and C is an m-dimensional subspace of the vector space $(\mathbb{F}_p)^n$. For $Q = \mathbb{F}_2$ a code is called binary.

Let $x = (x_1, ..., x_n)$ and $y = (y_1, ..., y_n) \in \mathbb{F}_p^n$. The Hamming distance between the words x and y is the number $d(x, y) = |\{i : x_i \neq y_i\}|$. The minimum distance of the code C is defined by $d = \min\{d(x, y) : x, y \in C, x \neq y\}$. The weight of a codeword x is $w(x) = d(x, 0) = |\{i : x_i \neq 0\}|$. For a linear code the minimum distance equals the minimum weight $d = \min\{w(x) : x \in C, x \neq 0\}$.

A p-ary linear code of length n, dimension k, and minimum distance d is called an $[n, k, d]_p$ code or [n, k, d] code when the size p of the field is not mentioned. A linear [n, k, d] code can detect at most d-1 errors in one codeword and correct at most $t=\left\lfloor\frac{d-1}{2}\right\rfloor$ errors. Two binary linear codes are isomorphic if one can be obtained from the other by permuting the coordinate positions. An automorphism of the code C is an isomorphism from C to C. The generator matrix of a code is a $k \times n$ matrix whose rows are the vectors of a base of the code. Every code is isomorphic to a code with the generator matrix in the standard form, i.e. in the form $[I_k, A]$, where I_k is the identity matrix of order k and A some $k \times (n-k)$ matrix.

4.1 Permutation decoding

Let $C \subseteq \mathbb{F}_p^n$ be a linear [n, k, d] code. For $I \subseteq \{1, ..., n\}$, let $p_I : \mathbb{F}_p^n \to \mathbb{F}_p^{|I|}$, $x \mapsto x|_I$, be the I-projection of \mathbb{F}_p^n . Then I is called an information set for C if |I| = k and $p_I(C) = \mathbb{F}_p^{|I|}$. The set of the first k coordinates for a code with a generator matrix in the standard form is an information set. The first k coordinates are then called information symbols and the last n - k coordinates are the check symbols and they form the corresponding check set.

Let $C \subseteq \mathbb{F}_p^n$ be a linear [n, k, d] code that can correct at most t errors (i.e. t-error-correcting code) and let I be an information set for C. A subset $S \subseteq \operatorname{Aut} C$ is a PD-set for C if every t-set of coordinate positions can be moved by at least one element of S out of the information set I. The property of having a PD-set for a code is not invariant under isomorphism of codes, it depends on the choice of the information set.

The algorithm of permutation decoding (see [23]) uses PD-sets and it is more efficient the smaller the size of a PD-set is. A lower bound on the size of a PD-set is given in the following theorem and it is due to Gordon [16].

Theorem 7. If S is a PD-set for an [n, k, d] code C that can correct t errors, r = n - k, then

$$|S| \ge \left\lceil \frac{n}{r} \left\lceil \frac{n-1}{r-1} \left\lceil \cdots \left\lceil \frac{n-t+1}{r-t+1} \right\rceil \cdots \right\rceil \right\rceil \right\rceil.$$

PD-sets for codes do not always exist. Even if they exist, PD-sets are not easy to find, since they depend on the chosen information set of the code.

Let A be the adjacency matrix of a graph Γ . Then the full automorphism group of Γ is a subgroup of the full automorphism group of the linear code spanned by A over \mathbb{F}_p . Codes with large automorphism groups are likely to have PD-sets, therefore, we were looking for PD-sets for the codes spanned by adjacency matrices of the DRGs constructed in this paper.

For any of the constructed DRG Γ_j^i from the previous section, let C_j^i denote the linear code spanned by the adjacency matrix of the graph Γ_j^i . Sizes of the obtained PD-sets (for specific information sets) for some of these codes are given in Table 16. For the other codes computation of PD-sets was not feasible. We denote by t the error correcting capacity of the

code, and by g the Gordon bound for the size of the PD-set of a code, from Theorem 7. The code C_2^1 is equivalent to the code C_1^2 .

$\mathrm{Code}\; C$	Parameters $[n, k, d]$	Aut(C)	t	g	Size of PD-set
C_1^1	[55,10,10]	S_{11}	4	5	5
C_2^1, C_1^2	[66,10,20]	S_{12}	9	15	55
C_{5}^{1}	[330,286,6]	S_{11}	2	60	420
C_6^1	[165, 120, 4]	S_{11}	1	4	5
C_8^1	[330,120,8]	S_{11}	3	7	22
C_{1}^{3}	[77,20,16]	$M_{22}: Z_2$	7	19	110
C_{5}^{4}	[1771,1540,4]	S_{23}	1	8	23

Table 16: PD-sets for codes from constructed DRGs from Mathieu groups

4.2 Block designs obtained from a code

Let w_i denote the number of codewords of weight i in a code C of length n. The weight distribution of C is the list $[\langle i, w_i : 0 \le i \le n \rangle]$. The support of a nonzero vector $x = (x_1, ..., x_n) \in F_q^n$ is the set of indices of its nonzero coordinates, i.e. $\sup(x) = \{i | x_i \ne 0\}$. The support design of a code of length n for a given nonzero weight w is the design with points the n coordinate indices and blocks the supports of all codewords of weight w.

Here we describe block designs obtained from the code $[176, 22, 50]_2$ spanned by the adjacency matrix of the graph Γ_2^3 .

Some remarkable block designs have been constructed from suport designs of codes, for example the 5-designs constructed by Assmus and Mattson in [1], and 5-designs constructed by V. Pless in [29, 30]. In this paper, from the supports of all codewords of the weights of the code [176, 22, 50]₂ we obtain block designs on 176 points on which the finite simple group Higman-Sims acts as the automorphism group. The support design for the minimum weight is the very well known Higman-Sims design, denoted by D_1 design in Table 17.

The weight distribution of the code $[176, 22, 50]_2$ is:

 $\left[\left< 0,1 \right>, \left< 50,176 \right>, \left< 56,1100 \right>, \left< 64,4125 \right>, \left< 66,5600 \right>, \left< 70,17600 \right>, \left< 72,15400 \right>, \left< 78,193600 \right>, \\ \left< 80,604450 \right>, \left< 82,462000 \right>, \left< 86,369600 \right>, \left< 88,847000 \right>, \left< 90,369600 \right>, \left< 94,462000 \right>, \left< 96,604450 \right>, \\ \left< 98,193600 \right>, \left< 104,15400 \right>, \left< 106,17600 \right>, \left< 110,5600 \right>, \left< 112,4125 \right>, \left< 120,1100 \right>, \left< 126,176 \right>, \left< 176,1 \right> \right]$

In Table 17 we describe the results.

Block design D	Parameters (v, k, λ)	Aut(D)	Block design D	Parameters (v, k, λ)	Aut(D)
D_1	(176,50,14), b=176	HS	D_7	(176,78,37752), b=193600	HS
D_2	(176,56,110), b=1100	HS	D_8	(176,80,124030), b=604450	HS
D_3	(176,64,540), b=4125	HS	D_9	(176,82,99630), b=462000	HS
D_4	(176,66,780), b=5600	HS	D_{10}	(176,86,87720), b=369600	HS
D_5	(176,70,2760), b=17600	HS	D_{11}	(176,88,210540), b=847000	HS
D_6	(176,72,2556), b=15400	HS			

Table 17: Block designs from the code $[176, 22, 50]_2$

Remark 12. The action of the group Higman-Sims is transitive on the points of the constructed designs, but in some cases it is not transitive on the blocks of the designs. The action that is not transitive on blocks appears in the cases of the designs D_7 , D_8 and D_{11} .

Remark 13. In Table 17 we did not include the support designs obtained from the weights 90-176 since they give rise to the complements of the block designs given in Table 17.

Acknowledgement

This work has been fully supported by Croatian Science Foundation under the project 6732. The authors would like to thank Sven Reichard for pointing out that the graphs Γ_3^1 and Γ_5^2 constructed in this paper are not arising from orthogonal arrays.

References

[1] E. F. Assmus jr, H. F. Mattson jr, New 5-designs, J. Combinatorial Theory A 6 (1969), 122–151.

- [2] R. F. Bailey, J. N. Bray, Decoding the Mathieu group M_{12} , Adv. Math. Commun. 1 (2007), 477–487.
- [3] R. C. Bose, T. Shimamoto, Classification and analysis of partially balanced incomplete block designs with two associate classes, J. Amer. Statist. Assoc. 47 (1952), 151–184.
- [4] W. Bosma, J. Cannon, Handbook of Magma Functions, Department of Mathematics, University of Sydney, 1994. http://magma.maths.usyd.edu.au/magma.
- [5] A. E. Brouwer, Strongly Regular Graphs, in: C. J. Colbourn, J. H. Dinitz (Eds.), Handbook of Combinatorial Designs, 2nd ed., Chapman & Hall/CRC, Boca Raton, 2007, pp. 852–868.
- [6] A. E. Brouwer, Parameters of Strongly Regular Graphs, Available at http://www.win.tue.nl/~aeb/graphs/srg/srgtab.html, Accessed on 24/02/2021.
- [7] A. E. Brouwer, A. M. Cohen, A. Neumaier, Distance-Regular Graphs, Springer-Verlag, Berlin, 1989.
- [8] P. J. Cameron, Coherent configurations, association schemes and permutation groups, Groups, combinatorics & geometry (Durham, 2001), 55–71, World Sci. Publ., River Edge, NJ, 2003.
- [9] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, Atlas of Finite Groups, Oxford University Press, Eynsham, 1985.
- [10] D. Crnković, V. Mikulić Crnković, A. Švob, Block designs and strongly regular graphs admitting a transitive action of the Mathieu group M_{11} , Australas. J. Combin. 73 (2019), 149–161.
- [11] D. Crnković, V. Mikulić Crnković, A. Švob, On some transitive combinatorial structures constructed from the unitary group U(3,3), J. Statist. Plann. Inference 144 (2014), 19–40.

- [12] D. Crnković, S. Rukavina, A. Švob, New strongly regular graphs from orthogonal groups $O^+(6,2)$ and $O^-(6,2)$, Discrete Math. 341 (2018), 2723–2728.
- [13] D. Crnković, S. Rukavina, A. Švob, On some distance-regular graphs with many vertices, J. Algebraic Combin. 51 (2020), 641–652.
- [14] E. R. van Dam, J. H. Koolen, H. Tanaka, Distance-Regular Graphs, Electron. J. Combin. (2016), DS22, Dynamic Survey, 156 pp.
- [15] The GAP Group, GAP Groups, Algorithms, and Programming, Version 4.11.0; 2020. (http://www.gap-system.org)
- [16] D. M. Gordon, Minimal permutation sets for decoding the binary Golay codes, IEEE Trans. Inform. Theory 28 (1982), 541–543.
- [17] W. C. Huffman, V. Pless, Fundamentals of Error-Correcting Codes, Cambridge University Press, Cambridge, 2003.
- [18] J. D. Key, T. P. McDonough, V. C. Mavron, Improved partial permutation decoding for Reed-Muller codes, Discrete Math. 340 (2017), 722–728.
- [19] J. D. Key, B. G. Rodrigues, Minimal PD-sets for codes associated with the graphs Q_2^m , m even, Appl. Algebra Engrg. Comm. Comput., to appear.
- [20] E. S. Kramer, S. S. Magliveras, D. M. Kramer, t-designs from large Mathieu groups, Discrete Math. 36 (1981), 171–189.
- [21] W. Lempken, Two new symmetric 2-(144,66,30) designs, preprint, 1999.
- [22] F. J. MacWilliams, Permutation decoding of systematic codes, Bell System Tech. J., 43 (1964), 485–505.
- [23] F. J. MacWilliams, N. J. A. Sloane, The Theory of Error-Correcting Codes. II, North-Holland Mathematical Library, Vol. 16. North-Holland Publishing Co., Amsterdam-New York-Oxford, 1977.

- [24] E. Mathieu, Memoire sur le nombre de valeurs que peut acqu erir une function quand on y permut ses variables de toutes le manière possibles, J. de Math. Pure et App. 5 (1860), 9–42.
- [25] E. Mathieu, Memoire sur l'étude des functions de plusieures quantites, sur la manière des formes et sur les substitutions qui laissent invariables, J. de Math. Pure et App. 6 (1861), 241–323.
- [26] E. Mathieu, Sur la function cinq fois transitive des 24 quantites, J. de Math. Pure et App. 18 (1873), 25–46.
- [27] V. Mikulić Crnković, I. Traunkar, Self-orthogonal codes constructed from weakly self-orthogonal designs invariant under an action of M_{11} , Appl. Algebra Engrg. Comm. Comput., to appear.
- [28] J. Moori, B. G. Rodrigues, A. Saeidi, S. Zandi, Some symmetric designs invariant under the small Ree groups, Comm. Algebra 47 (2019), 2131–2148.
- [29] V. Pless, On a new family of symmetry codes and related new five-designs, Bull. Am. Math. Soc. 75 (1969), 1339–1342.
- [30] V. Pless, Symmetry codes over GF(3) and new five-designs, J. Comb. Theory, Ser. A 12 (1972), 119–142.
- [31] C. E. Praeger, L. H. Soicher, Low rank representations and graphs for sporadic groups, Australian Mathematical Society Lecture Series 8, Cambridge University Press, Cambridge, 1997.
- [32] D. Robinson, A Course in the Theory of groups, Springer-Verlag, New York, Berlin, Heidelberg, 1996.
- [33] L. H Soicher, Yet another distance-regular graph related to Golay code, Electronic J. Combin. 2 (1995), N1, 4pp.

- [34] V. D. Tonchev, Combinatorial Configurations: Designs, Codes, Graphs, John Willey & Sons, New York, 1988.
- [35] H. Wielandt, Finite permutation groups, Academic Press, New York, 1964.
- [36] R. A. Wilson, The finite simple groups, Springer-Verlag, London, 2009.
- [37] W. Wirth, Konstruktion symmetrischer Designs, PhD thesis, Johannes Gutenberg-Universität Mainz, 2000.