TEXAS A&M

BMEN 615: Bioengineered Cellular Systems

Full Backup Reference Document — every piece of content from the interactive web app (Introduction, all four papers, the Evaluation & Frontier comparison, the New Questions lens, the full 48-item Q&A bank, and the glossary) compiled into one standalone, printable document. Use this if the live app is unreachable, the presence/presenter service is down, or a projector/network fails mid-class — everything here is self-contained and needs no server.

Presented by Justin Houck, PhD Student, Department of Animal Science
Generated from the live app's own source data

1. Introduction: The Lymphatic System

A Textbook of Histology, Maximow & Bloom, 1957

⚡ What the Lymphatic System Actually Does

Strip the anatomy away for a second and ask what this system is actually *for*. Blood capillaries are leaky by design — pressure pushes roughly 20 liters of plasma a day out into the tissues, and only about 17 of those liters get pulled straight back in by the venous side. The lymphatic system exists to handle the other ~3 liters, plus everything too large to slip back into a vein unassisted. Four jobs fall out of that one basic leak:

- **Fluid balance (drainage).** Blind-ended lymphatic capillaries pick up the interstitial fluid blood vessels leave behind. When this fails, tissue swells — that failure mode has a name: lymphedema.
- **Protein & macromolecule return.** Plasma proteins that leak out of blood capillaries are too large to re-enter a venule on their own; lymph is their only way back into circulation.
- **Fat absorption.** In the gut specifically, lacteals absorb dietary fat as chylomicrons and carry it to the thoracic duct, bypassing the liver on the first pass.
- **Immune surveillance.** Lymph carries antigen, pathogens, and antigen-presenting cells from tissue into lymph nodes, where patrolling lymphocytes can encounter them and activate — a node is a checkpoint, not just a filter.

None of this moves the way blood does — there is no central pump. Lymph is pushed along passively (by skeletal muscle contracting, and by breathing) and actively by the **lymphangion**: the vessel segment between two one-way valves, lined on the inside with a thin layer of cells and wrapped on the outside in smooth muscle that contracts rhythmically, segment by segment, like a chain of hands passing a bucket forward.

🫓 What Papers 1, 2 & 3 Are Trying to Mimic

None of the three lab-on-a-chip papers try to rebuild the whole circuit above. Each one isolates a single piece of it — the leak, the pump, or the growth of the network itself — and asks whether that piece can be reproduced, measured, and manipulated outside the body.

2022 mimics: the pump

The lymphangion itself — the vessel wall that squeezes to push lymph forward between valves.

Selahi et al. build a single lymphangion: a collagen tube lined with lymphatic-vessel lining cells and wrapped in lymphatic muscle cells. Growing the two cell types together, under fluid shear, shifts the muscle cells to wrap more around the vessel's circumference (mean ~61°, vs. ~31° when the muscle cells are grown alone).

2024 mimics: how the network gets built

Not the adult vessel, but a new, dead-ended capillary sprouting from an existing one.

Tronolone et al. watch lymphatic-vessel lining cells sprout from a parent vessel into a gel over four days, driven by chemical gradients and fluid flow acting together (new lymphatic-vessel growth, or lymphangiogenesis).

2026 mimics: the leak & pickup

The first moment in the live model — fluid and immune cells crossing from the blood side into the lymphatic side.

Gray-Scherr et al. build two separate, never-connecting vessel networks (blood and lymphatic) sharing one continuous gel. The gel filters molecules by size (10 and 70 kDa dextran cross through; 2000 kDa is sharply reduced), and TNF-alpha increases how many T-cells cross from the blood network into the lymphatic network.

🔬 The 1957 Blind Spot: What They Could See But Not Yet Explain

The late 1950s and early 1960s were the final years of "pre-modern" immunology. Four specific claims in Chapter 11 read very differently today:

⚠️ What the 1957 Text Said

- ✕ A lymphocyte is just "a lymphocyte" — no visible way to tell one apart from another under the microscope.
- ✕ The function of the thymus is "largely unknown"; it shrinks with age, so some assumed it was a vestigial "graveyard" for dying cells.
- ✕ The small lymphocyte is described as an undifferentiated stem cell circulating in the blood — a direct extension of Maximow's own unitarian stem-cell theory.
- ✕ The *reticuloendothelial system* (RES): reticular cells and sinus-lining endothelial cells were believed able to transform directly into phagocytic macrophages.

✨ What We Know Now

- ✓ Lymphocytes split into distinct T-cell and B-cell lineages (plus NK cells), established in the mid-1960s — identical under a light microscope, completely different jobs.
- ✓ Jacques Miller's thymectomy experiments (1961) showed the thymus is where T-cells mature — a training school, not a graveyard.
- ✓ A small lymphocyte is a specialized immune cell on patrol, not a generic stem cell.
- ✓ Macrophages are a separate, monocyte-derived lineage from the structural cells of the node — the "mononuclear phagocyte system" replaced the RES concept.

2. 2026: Blood-Lymph

Lab Chip 2026 — 3D Microfluidic Model of Blood & Lymphatic Exchange

Authors: Delaney Gray-Scherr, Terry Ching, Christopher S. Chen, et al.

Why This Paper Matters

First Dual-Network Exchange Chip

The Discovery:

This is the first chip to grow two separate, self-assembled human vessel networks — one blood, one lymphatic — side by side in one shared gel, each fed independently, that stay distinct (never fusing into one network) while still exchanging molecules and immune cells across the gel between them. Using that setup, the paper watched, in real time, how an inflammatory signal (TNF-alpha) increases how many T-cells cross from the blood network into the lymphatic network (3.31-fold for resting T-cells, 5.89-fold for already-activated ones).

Why It Matters:

Before this, studying how immune cells move from blood into lymphatic vessels meant using whole animals or much simpler single-vessel setups. Having both vessel types side by side, built from human cells, with live imaging, gives researchers a way to directly test drugs that affect immune-cell trafficking (e.g., anti-inflammatories) or study how tumors exploit lymphatic drainage to spread — without leaving the dish.

How Long It Works

Day 5 Standard Timepoint

HOW LONG IT WAS TESTED

Day 5 (established timepoint; not reported beyond this)

LENGTH OF EACH MEASUREMENT

3 hrs (drainage test) / 4 hrs (T-cell movement test)

WHY IT STOPPED

Not reported in the paper

Question: How long would this keep working?

The paper settles on Day 5 as its standard testing point, once both vessel networks have finished forming on their own over 5 days. From there, individual tests run a few hours each (drainage: 3 hours; T-cell movement: 4 hours). The paper does not report what happens to the networks after Day 5, and gives no reason they would eventually stop working or how long they could keep going.

⚠️ What Could Cause It to Fail (Biology)

This cannot be grounded in fact in the paper. It does not report a biological reason the system would eventually fail (for example, the gel breaking down, or effects from the missing support cells called pericytes).

⚠️ Physical/Resource Limits

This cannot be grounded in fact in the paper beyond the stated test durations. The small pressure difference (9.6 Pa, created by 40 microliters on the blood side vs. 10 on the lymph side) drives the reported 3-4 hour tests; the paper doesn't say how long it would take for the reservoirs to run out.

🔧 Stated Next Steps

This cannot be grounded in fact as a stated fix in the paper. The paper's own discussion names different future directions: adding a recirculating pump or a more controlled pressure source, together with computer flow simulations, to better define and control the flow.

Figures



Fig 1: Design & Assembly

Separate blood and lymphatic sections, each fed with fluid on its own, with a gel between them.



Fig 2: Vessel Morphology

Blood vs. lymphatic vessel cell-junction protein (VE-cadherin) and channel widths.



Fig 3: Morphometric Analysis

Charts of vessel width and how densely packed the cell junctions are.



Fig 4: Independent Perfusion

Fluid staying within the blood vs. lymph channels under normal flow.



Fig 5: Size Solute Transport

Small tracer molecules (10 and 70 kDa) cross the gel barrier; a much larger one (2000 kDa) is sharply reduced.



Fig 6: T-Cell Trafficking

TNF-alpha increases T-cell crossing into the lymphatic network: 3.31-fold for resting T-cells, 5.89-fold for already-activated ones.

3. 2022: Lymphangion

Lab Chip 2022 — Lymphangion-Chip: Co-Culture & Bidirectional Signaling

Authors: Amirali Selahi, Teshan Fernando, Abhishek Jain, et al.

Why This Paper Matters

First LEC-LMC Crosstalk Model

The Discovery:

No prior lab model had ever grown lymphatic lining cells (LECs) and lymphatic muscle cells (LMCs) together around one vessel. This paper did, and found that doing so changes the biology: the two cell types align differently under flow when grown together vs. alone, and — in the paper's own words — when the muscle cells were co-cultured with the lining cells, "the lymphatic endothelial barrier function was relatively conserved" after exposure to the inflammatory signal TNF-alpha (Fig. 7D), where the lining cells leaked much more when grown without the muscle layer.

Why It Matters:

This is the first direct evidence that lymphatic muscle cells actively protect the vessel lining during inflammation, not just squeeze lymph along mechanically. That points to LMC health, not just LEC health, as a possible target for treating lymphedema and inflammatory vessel leakiness. The fabrication method itself (gravitational lumen patterning) is also a reusable toolbox for building other tunable, round, multi-layered vessel models.

How Long It Works

Up to 96 hr (5 Days) Tracked

HOW LONG IT WAS TESTED

Up to 96 hr / 5 Days (tracked range)

LENGTH OF EACH MEASUREMENT

Continuous fluid flow from a pump; no stated time limit

WHY IT STOPPED

Not reported in the paper

Question: How long would this keep working?

The paper's own experiments track the system for up to 5 days: the vessel lining becomes a complete layer within 60 hours, and the gap between the two cell layers is followed through 96 hours. The paper does not report what happens to the device after Day 5, and does not name a specific reason it would eventually fail.

⚠️ What Could Cause It to Fail (Biology)

This cannot be grounded in fact in the paper. It does not report a specific biological reason for failure, such as the gel shrinking or the cell layers coming apart.

⚠️ Physical/Resource Limits

This cannot be grounded in fact in the paper. It makes no claim about the vessel wearing out from fluid friction, or cell-junction proteins declining, beyond Day 5.

🔧 Stated Next Steps

This cannot be grounded in fact as a lifespan fix in the paper. The paper's stated next steps are about adding valves inside the vessel, a pulsing expansion-and-contraction motion, and linking several vessel segments together — not about extending how long the device lasts.

Figures



Fig 1: Lymphangion Model

The lymphangion (vessel segment between two valves) recreated in a lab channel.



Fig 2: GLP Technique

Turning the channel vertical during fabrication prevents it from sagging unevenly.



Fig 3: Tuning Diameters

Controlling channel width using fluid pressure and how concentrated the collagen gel is.



Fig 4: Dual-Layer Reconstitution

3D microscope image showing a complete layer of lymphatic-vessel lining cells surrounded by muscle



Fig 5: Cellular Cross-Section

How the gap between the vessel lining and the muscle layer changes over 96 hours.

cells.



Fig 6: Shear & Alignment

The muscle cells shift from mostly lengthwise (~31°) toward wrapping more around the vessel (~61°), but only when grown together with lining cells under fluid shear — never reaching a literal 90°.



Fig 7: Barrier Function

A test measuring how much a labeled tracer molecule leaks through the vessel wall.

4. 2024: L-Chip Sprouting

Adv Biol 2024 — Engineering Lymphangiogenesis-on-Chip (L-Chip)

Authors: James J. Tronolone, Nadin Mohamed, Abhishek Jain

Why This Paper Matters

ESM-1 Identified as a Lymphatic-Specific Driver

The Discovery:

The paper found that endocan (ESM-1) alone — without the rest of the growth-factor cocktail normally used — is enough to strongly drive new lymphatic sprouting, and that this effect is specific to lymphatic vessels: the same factor did not drive sprouting the same way in a matched blood-vessel (angiogenesis) chip.

Why It Matters:

ESM-1 was already known as a marker linked to blood-vessel growth and cancer, but it was under-explored as a driver specifically of lymphatic-vessel growth. Pinning down one specific, lymphatic-selective molecular trigger gives researchers a concrete target for drugs meant to encourage new lymphatic growth (e.g., after lymph node removal or in lymphedema) or block it (e.g., to slow tumor-associated lymphatic spread) — without also disturbing blood vessels.

How Long It Works

Fixed 96 hr (4-Day) Endpoint

HOW LONG IT WAS TESTED

96 hr / 4 Days (fixed experimental endpoint)

LENGTH OF EACH MEASUREMENT

96-hour fixed growth period before the tissue is preserved and imaged

WHY IT STOPPED

Not reported in the paper

Question: How long would this keep working?

Every experiment in this paper uses a fixed 96-hour (4-day) growth and measurement period, by which sprouts reach up to ~1 mm. The paper does not report what happens to sprouts after Day 4 and makes no claim about them breaking down, shrinking back, or any limit past that point.

⚠️ What Could Cause It to Fail (Biology)

This cannot be grounded in fact in the paper. It does not report sprouts shrinking back, internal structures collapsing, or cell death beyond Day 4.

⚠️ Physical/Resource Limits

This cannot be grounded in fact in the paper. It does not report how long the chemical gradient would keep working beyond the fixed 96-hour endpoint.

🔧 Stated Next Steps

This cannot be grounded in fact as a lifespan fix in the paper. The paper's own stated next steps are studying exactly how ESM-1 works, and adding supporting cell types currently missing from the model (muscle cells, connective-tissue cells, pericytes, macrophages).

Figures



Fig 1: Sprout Dynamics (Day 1-4)

New capillaries sprout past the tiny retaining pillars, reaching about 1 mm by Day 4.



Fig 2: Conditioning & Gradients

How chemical gradients and fluid flow through the gel work together.



Fig 3: Interstitial Flow Interactions



Fig 4: Conditioning Duration

How pre-treating cells with growth-signaling proteins affects how far sprouts grow.

Sprouts grow fastest when growing against the direction of fluid flow.



Fig 5: Endocan (ESM-1) Role

ESM-1 alone is enough to strongly drive new lymphatic-vessel growth.

5. 1937: Whole-Organ

J Exp Med 1937 — The Culture of Whole Organs (Lindbergh Apparatus)

Authors: Alexis Carrel (perfused with the Lindbergh pump)

Why This Paper Matters

First Whole-Organ Culture Outside the Body

The Discovery:

This paper demonstrated that a whole mammalian (or bird) organ — not just isolated cells, but an intact gland with its own blood vessels — could be kept alive, metabolically active, and even growing new cells for days to weeks entirely outside the body, using nothing but pulsatile fluid pressure (the Lindbergh pump) and a nutrient fluid with no actual blood cells in it. The paper also pinned down a specific cause-and-effect: adding Witte's peptone to that fluid, not perfusion itself, is what drove the glands to grow new cells (hyperplasia).

Why It Matters:

This was 90 years before organ-on-a-chip devices, and it first proved that an organ's physiology — not just its cells in a dish — could be sustained and studied outside the body at all. It is a direct conceptual ancestor of the organ-preservation and ex vivo perfusion techniques used today to keep donor organs viable for transplant.

How Long It Works

3–21 Days (pH-Limited)

HOW LONG IT WAS TESTED

3–21 Days reported; 6 days used for most main figures

LENGTH OF EACH MEASUREMENT

Continuous pulsing, gas-driven perfusion

WHY IT STOPPED

Fluid pH decline (stopping rule: pH below 7)

Question: How long would this keep working?

Carrel reports perfusion runs lasting from 3 to 21 days, though 6 days is the duration used for most of the paper's main histology comparisons (Plates 17-18). The paper's own explicit stopping rule is pH-based: the fluid's pH starts around 7.5 and falls progressively over about a week toward roughly 7.0-7.1, and "when the pH goes below 7, the experiment is interrupted."

⚠️ What Could Cause It to Fail (Biology)

A gradual drop in the fluid's pH (becoming more acidic) is the paper's own stated limiting factor — pH falls from about 7.5 toward 7.0-7.1 over about a week. The paper does not attribute this to a named modern diagnosis (like "metabolic acidosis") — that label is a present-day gloss, not the paper's own language.

⚠️ Physical/Resource Limits

This cannot be grounded in fact as stated. The paper notes small side-vessels are deliberately left open rather than tied off, bleeding into the surrounding tissue and causing swelling — but it states directly that this swelling "has no ill effect on the gland."

🔧 Stated Next Steps

This cannot be grounded in fact in the paper itself. Modern organ-preservation technology (continuous oxygen supply, fluid filtering, and automatic pH control) is a reasonable present-day point of

comparison, but it is not something the 1937 paper discusses.

Figures

Plate 17: Control & 6-Day Perfusion

Normal follicle structure maintained after 6 days outside the body.



Plate 18: Hyperplasia under Peptone

Extra cell growth lining the follicles, with visible signs of the stored colloid substance being reabsorbed.



Plate 19: Explant Viability Outgrowth

Clusters of cells still actively growing after 8–18 days of whole-organ culture outside the body.

6. Evaluation & Frontier

These four studies span 90 years, from keeping a whole organ alive outside the body (1937) to a lab-on-a-chip device with two separate, communicating blood and lymphatic vessel networks (2026).

How the Studies Compare

These four studies span 90 years and show a clear shift in approach. In 1937, Alexis Carrel used the pulsatile glass pump built by Charles Lindbergh to keep whole organs (cat and bird thyroids) alive outside the body for days to weeks, but had no way to see or control what was happening at the level of individual cells. Over the following decades, research moved to smaller scales: first reconstructing a single lymphatic vessel segment under controlled fluid shear (Selahi et al., 2022), then studying how new lymphatic capillaries grow in response to chemical and flow cues (Tronolone et al., 2024), and most recently building two separate, independently-fed blood and lymphatic vessel networks that exchange fluid and immune cells with each other (Gray-Scherr et al., 2026). Modern microfluidic devices give researchers far more control and much better imaging than a whole organ ever could, and use human cells instead of animal tissue — but none of them has been shown to run for more than about a week, and none includes the full mix of supporting cell types and organ-to-organ interactions a real body has. Closing that gap is the open problem this research still needs to address.

Scorecard

What's Compared	Paper 1: Blood-Lymph (2026)	Paper 2: Lymphangion (2022)	Paper 3: L-Chip (2024)	Paper 4: Whole Organ (1937)
Scale & Structure	Two vessel networks side by side (~3 mm chamber, capillaries 20–40 µm wide)	One tube-shaped vessel (750 µm-wide channel, 900 µm across including the wall)	New capillaries growing into a gel (1.5 mm chamber, sprouts up to ~1 mm long)	A whole organ (intact thyroid gland, ~5–10 cm)
Cell Types & Source	Human donor cells: blood-vessel lining cells (glow-tagged for tracking), lymphatic-vessel lining cells, connective-tissue cells, and T-cells from blood	Human lymphatic-vessel lining cells for the inner tube, but rat muscle cells for the outer sheath — a cross-species mix the authors note as a limitation, since human lymphatic muscle cells aren't commercially available	Human lymphatic-vessel lining cells, pre-treated with a 5-factor growth-signal cocktail, compared against ordinary human blood-vessel lining cells	Whole thyroid glands from cats and birds, still attached to their feeding blood vessel
What Drives the Fluid Flow	A small pressure difference from gravity alone (9.6 Pa) pushes fluid through the gel	A pump sets steady fluid friction (shear) on the vessel wall at three levels — none, low, and roughly what's measured in real lymph vessels	Slow fluid seepage through the gel (1–5 µm/s) combined with a chemical gradient	Gas pressure pulses push fluid through the artery in waves, visibly expanding and relaxing the vessel wall with each pulse

What's Compared	Paper 1: Blood-Lymph (2026)	Paper 2: Lymphangion (2022)	Paper 3: L-Chip (2024)	Paper 4: Whole Organ (1937)
How the Results Were Measured	Live microscope imaging while the experiment ran, a dye-drainage test, plus fixed tissue stained afterward	A live dye-leakage test, plus fixed tissue stained for specific muscle and lymphatic marker proteins	Fixed tissue imaged in 3D under a microscope, with software automatically tracing and measuring the new vessel sprouts	Tissue examined only after the animal was euthanized — thin stained slices under a microscope — plus a test of whether small tissue pieces could still grow in a flask afterward
How Long It Ran	Day 5 is the paper's standard timepoint; assays run 3–4 hrs. Nothing beyond Day 5 is reported.	Tracked up to 96 hrs (5 days); no failure mechanism reported beyond that	Fixed 96 hr (4-day) endpoint in every experiment; no outcome beyond Day 4 reported	3–21 days reported (6 days used for most main figures); the experiment was stopped once the fluid's pH dropped below 7
Main Trade-off / Limitation	Built by hand with needles and sealed with grease; no pericytes (vessel-stabilizing cells) included	Stiff plastic walls; just one straight tube; no valves inside the vessel	The new sprouts are dead-ended — no fluid actually flows through the vessels that form	Sealed glass chamber — no way to see inside while running, only after the fact; some minor bleeding caused swelling, but the authors state it didn't harm the gland

What's Compared	Paper 1: Blood-Lymph (2026)	Paper 2: Lymphangion (2022)	Paper 3: L-Chip (2024)	Paper 4: Whole Organ (1937)
What It Could Be Used For	Studying how immune cells move in cancer, testing anti-inflammatory drugs, and modeling lymph drainage	Developing treatments for lymphedema (fluid buildup from poor drainage), restoring normal muscle tone in vessels, and testing how well drugs pass through vessel walls	Finding drugs that encourage or block new lymphatic-vessel growth, understanding how tumors spread, and studying wound healing	A lab method for studying normal and diseased thyroid function (the paper's own stated goal); historically, an early ancestor of today's organ-preservation technology

What Changed Over 90 Years

1937 — Keeping a Whole Organ Alive Outside the Body

Carrel (1937) · J Exp Med · Whole organ (~10 cm)

Milestone: Proved that a whole mammalian organ could be kept alive and responsive outside the body using pulsing fluid pressure alone.
Limitation: Sealed glass apparatus with no way to see inside while it ran; animal tissue only; no way to observe individual cells.

2022 — Growing a Lymph Vessel Segment With Two Cell Types

Selahi et al. · Lab Chip · Microscale (900 µm)

Milestone: Grew human lymphatic-vessel lining cells and muscle cells together around one channel; found that fluid shear stress shifts the muscle cells' orientation further around the vessel's circumference (not a literal 90°, but a measured shift from ~31° to ~53–61°).
Limitation: Just one straight vessel; stiff plastic channel; no valves inside it.

2024 — Guiding New Lymphatic Vessel Growth (L-Chip)

Tronolone et al. · Adv Biol · Mesoscale (1.5 mm chamber)

Milestone: Tracked new capillary growth day-by-day over a 4-day period; found that fluid flow through the gel works together with a growth-factor gradient, and identified ESM-1 as a protein that drives this growth on its own.
Limitation: The new sprouts are dead-ended; no fluid actually flows through the newly grown branches.

2026 — A Chip With Separate Blood and Lymph Networks

Gray-Scherr et al. · Lab Chip · Two connected networks (~3 mm)

Milestone: Built two separately-fed human vessel networks (blood and lymphatic) sharing one gel, with filtering between them and live tracking of immune T-cells moving from one network to the other.
Limitation: Built by hand with needles; fluid drains one-way under gravity rather than recirculating; detailed imaging requires ending the experiment.

✨ What Lab-on-a-Chip Devices Gained

- ✓ Human cells instead of animal tissue: the move from whole cat and chicken organs to cells taken directly from human donors (vessel-lining, connective-tissue, and immune cells).
- ✓ The ability to watch it happen: modern chips can be imaged live under a microscope at very high resolution, tracking individual cells moving and dye molecules flowing in real time — something a sealed glass apparatus never allowed.
- ✓ Testing one variable at a time: fluid shear, pressure, gel stiffness, and chemical signals can each be changed independently, without the rest of a whole animal's physiology changing along with it and confusing the results.
- ✓ Lower material cost per experiment: these chips need only microliters of reagents, compared to the hundreds of milliliters of donor animal serum the 1937 apparatus required.

⚠️ What Got Left Out

- ✗ Long testing windows: the 1937 whole-organ setup was reported alive for up to 21 days. Each of the three modern chip papers stops at a fixed point (4 to 5 days) and doesn't report what happens after — not necessarily because something breaks down, simply because none of them tested further.
- ✗ Supporting vessel-wall cells: none of the three modern chip papers include pericytes — cells that normally wrap around small vessels and help stabilize them — in their cell lists.
- ✗ Connections to the rest of the body: modern chips study one or two tissue types in isolation, leaving out the hormonal, immune, and filtering interactions that normally happen between organs in a whole body.
- ✗ Soft, flexible surroundings: chips are bounded by rigid glass or stiff plastic (PDMS), instead of the soft, pulsing living tissue that surrounds real vessels in the body.

Where This Research Could Go Next

1. Measuring Chips Without Ending the Experiment

Instead of ending the experiment at Day 4 or 5 to take microscope images, future chips could have small sensors built directly into the channel walls — electrical sensors that measure how tightly sealed the vessel lining is, and optical or chemical sensors that track waste buildup and signaling molecules — all while the cells are still alive and the experiment is still running.

2. Building Chips From a Specific Patient's Own Cells

For patients with vessel malformations, lymphedema caused by another condition, or a spreading tumor, a chip built from their own cells could let doctors test which chemotherapy or immune-based treatment is likely to work best for them before actually giving it.

3. Connecting Multiple Chips Into One System

By connecting a blood-lymph exchange chip to a living tumor or liver chip upstream, and a lymph-node chip downstream, researchers could follow the whole path a cancer cell or immune signal takes through the body: entering a vessel, draining through tissue, arriving at a lymph node, and triggering an immune response there.

4. Building Chips by Machine Instead of by Hand

Building each chip by hand is a major reason these devices haven't been adopted more widely in drug development, since no two hand-built chips are exactly identical. A robotic printer could build hundreds of blood-lymph devices at once, with far more consistent, precise dimensions than a person can achieve by hand.

5. Making These Chips Last Longer Than a Few Days

Each modern chip paper stops at a fixed point — 4 to 5 days — not because researchers measured a breakdown and hit a wall, but simply because that's where each study's experiments ended. Whether these devices could keep working longer, and what would actually cause them to fail if pushed further, is still an open question. Possible culprits — the gel breaking down, chemical gradients running out, or fluid never actually flowing through the new dead-ended sprouts — are reasonable guesses, but none of these papers tested or confirmed them. Getting past this limit would likely need a gel that can repair itself under stress, supporting cells wrapped around the

vessels to stabilize them (the way pericytes do in the body), and a way to continuously clear waste from a closed loop — aiming at weeks-to-months of continuous operation instead of days.

7. New Questions — A Reproductive & Stress-Physiology Lens

Old Experiments, New Instruments: Leaky Gut in Stressed Animals

Viewed through stress and reproductive physiology — e.g., Thomas H. Welsh, Jr., Regents Professor, Texas A&M AgriLife Research Faculty Fellow, Physiology of Reproduction — these four systems map onto a well-established pathway in livestock stress research: **stress → a leaky gut barrier → bacterial endotoxin reaches blood and lymph → systemic inflammation → reduced reproductive performance.** That is the same barrier-permeability problem these chips were built to study, just in a different epithelium.

🔗 Why These Four Papers Are Relevant to Leaky Gut

During heat or psychological stress, reduced gut blood flow and elevated cortisol break down the gut epithelial barrier. Endotoxin (LPS) and bacterial products cross into circulation largely via the mesenteric lymphatics, bypassing the liver's first-pass filtering (the "gut-lymph pathway" of stress-induced systemic inflammation) — mechanistically the same event paper 1 was built to watch directly.

🧪 What New Experiments Does This Open Up?

From Paper 1 (2026): watch the leak happen live

Rebuild the dual-network chip with gut epithelium in place of blood-endothelial cells.

Re-run the size-exclusion tracer panel (10 / 70 / 2000 kDa), but challenge the barrier with cortisol or LPS instead of TNF-α — visualizing live what leaky-gut research has only inferred from blood endotoxin assays.

From Paper 2 (2022): test if the protective crosstalk survives chronic stress

Re-run the muscle-cell barrier-rescue assay with a stress hormone.

Does LMC protection against TNF-α also hold against cortisol or LPS? Does it erode under repeated, chronic stress exposure rather than one pulse?

From Paper 3 (2024): ask whether stress rewires drainage capacity itself

Does chronic stress remodel gut lymphatic vessels, not just leak through them?

Does a stress-altered interstitial flow pattern in the gut wall drive abnormal lymphatic growth, permanently changing drainage capacity?

From Paper 4 (1937): isolate one stressor, get a clean causal answer

Carrel's one-organ, one-factor logic, updated for a chip.

Perfuse an isolated gut-epithelium-on-chip with just cortisol, or just LPS, in isolation — separating one variable a live stressed animal never cleanly isolates.

📊 Old Data, New Ruler

The standardized 10 / 70 / 2000 kDa tracer panel and fold-change immune-trafficking analysis from paper 1 could be retrofitted as a quantitative barrier-leak severity scale against archived samples from past heat-stress trials — a sharper readout than cruder in vivo gut-permeability markers typically used today.

8. Full Q&A Reference Bank

All 48 question-and-answer entries, organized by paper — the same bank the live app's "Interrogate the Paper" panel draws from.

2026: Blood-Lymph

BIOLOGICAL SYSTEM & CELL ORIGINS

Where did the cells come from (human vs. animal, cell lines vs. primary)?

All human cells, taken directly from donors (not lab-grown cell lines). The cells lining the blood-vessel network are human skin blood-vessel lining cells, genetically tagged so they glow green under a microscope. The cells lining the lymphatic-vessel network are human skin lymphatic-vessel lining cells (confirmed 99% lymphatic by testing for a marker protein, podoplanin). Supporting connective-tissue cells come from human lung donors. The immune cells are T-cells (a type of white blood cell) isolated from healthy donors' blood, used either "resting" (straight from frozen storage) or after a week of lab treatment that activates them.

Source: Materials & Methods, "Cell culture"

BIOLOGICAL SYSTEM & CELL ORIGINS

What functional tissue systems are analyzed and reconstructed?

Two small vessel-like networks side by side: a continuous blood-vessel network that can be fed fluid on its own, and a separate, dead-ended lymphatic-vessel network next to it. Both are grown inside one shared gel chamber, separated by a thin, cell-free layer of the same gel — thin enough that molecules and cells can cross between the two networks, but thick enough that the vessels themselves never physically connect.

Source: Fig. 1C-D; Results, "The resulting device architecture..."

BIOLOGICAL SYSTEM & CELL ORIGINS

What is the physiological relevance of this model?

It recreates the space between blood and lymphatic vessels where the two work together to move fluid, molecules, and immune cells through tissue — the same process that breaks down in cancer spread, chronic inflammation, and lymphedema (fluid buildup from poor drainage). It specifically lets researchers watch T-cells flow through the blood vessels, squeeze out into the surrounding tissue, and enter the lymphatic vessels — mirroring how immune cells patrol and exit tissue in the body.

Source: Introduction; Results, "Reconstitution of T cell trafficking...", Fig. 6

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

What is the physical process and operation of the device?

The device is built by layering glass, laser-cut plastic, and tape. Thin needles (250 micrometers wide — a few human hairs' width) are threaded through guide channels to mold the shape of the vessel channels. A gel mixed with blood-vessel cells is cast around part of the needles, then a cell-free gel layer, then a gel mixed with lymphatic-vessel cells around the rest. Once the gel sets, the needles are pulled out, leaving open channels behind, and the entry points are sealed with grease. The whole device is then cultured for 5 days on a slowly rocking platform with fresh growth medium added daily, and each experiment begins by adding a set amount of fluid to the blood and lymph reservoirs (40 microliters on the blood side, 10 on the lymph side).

Source: Materials & Methods, "Microfluidic device design and fabrication," "Cell seeding and gel casting"; Fig. 1A-C

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

What hemodynamic forces and shear gradients play a role?

A small pressure difference, created simply by putting more fluid on the blood side than the lymph side (40 vs. 10 microliters), drives flow through the gel — chosen to roughly match the real pressure difference between tissue and lymphatic vessels in the body. By tracking small particles in the fluid, the researchers measured the friction (shear stress) the flow exerts on the vessel walls: roughly 0.1–3 dyne/cm^2 in the blood network (across vessels 25–200 micrometers wide), and much lower in the lymphatic network — about 0.016 dyne/cm^2 on average, dropping even lower in the smallest branches.

Source: Results, "Quantitative drainage assay..." and "Flow characterization by particle tracking velocimetry"; Fig. S8

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

Do fluid acceleration, deceleration, angle, or orientation play a role?

Yes, during the 5-day growth period: the devices sit on a platform that rocks up to a 30° tilt, which creates a gentle back-and-forth flow through the blood-vessel channels (reaching a peak pressure of about 27 Pa, but averaging to zero net push over each rocking cycle, since the dead-ended lymphatic side has no matching reservoir to drive flow against). During the actual measurement experiments (the drainage and T-cell tests), the rocking is turned off, so the pressure difference instead pushes steadily in one direction rather than back and forth.

Source: Materials & Methods, "Cell seeding and gel casting"; Results, Flow characterization section

ENGINEERING CONTROL & SACRIFICES

What was sacrificed to engineer this, and how does that impact relevance?

The paper's own discussion admits its control over fluid flow is not fully worked out: flow comes "primarily through rocker-driven movement and static hydrostatic gradients," and because the channel shapes aren't simple standard geometry, the exact flow pattern and the friction it creates on vessel walls "are not fully characterized." The authors say that adding computer flow simulations and a mechanical pump would be needed to map this out properly against real physiological values. Separately — something the paper doesn't itself call out as a trade-off — the list of cell types used does not include pericytes, the cells that normally wrap around small vessels and help stabilize them.

Source: Discussion, paragraph beginning "The tape-based fabrication workflow..."

ENGINEERING CONTROL & SACRIFICES

How much independent control does the researcher have over variables?

Strong control over the parameters the researchers set directly: how much fluid goes in each reservoir, how many cells are seeded, and the exact dose of chemical signals added, such as the inflammatory protein TNF-alpha. The shape of the needle channels is fixed once the device is built, not something that varies run to run. The actual branching pattern of the vessels is not directly controlled — it emerges on its own as the cells grow together over 5 days, landing on consistent but not researcher-chosen sizes (blood vessels averaging 25.80 ± 0.52 micrometers wide, lymphatic vessels 35.17 ± 1.15 micrometers).

Source: Results, Fig. 3C; Materials & Methods

ENGINEERING CONTROL & SACRIFICES

How long will this work for continuously, and when does the system degrade beyond designed function?

The paper settles on day 5 as its standard testing point, once both vessel networks have finished forming on their own. From there, individual experiments run just a few hours: 3 hours for the drainage test, 4 hours for the T-cell test. This cannot be grounded in fact in the paper: it does not say what happens to the networks after day 5, and gives no explanation for why they would eventually stop working or how long they could keep going.

Source: Results, "...established day 5 as the standard timepoint"; Materials & Methods, drainage and trafficking assay sections

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

In what context was the data collected and interpreted (immunostaining and imaging)?

Two complementary approaches. First, at a fixed end point, the tissue was stained for several marker proteins (markers for lymphatic cells, nuclei, blood-vessel cells, and cell junctions, plus the built-in green glow tag) and imaged in 3D layers under a microscope; custom software then measured vessel width, length, surface area, and how densely packed the network was. Second, live imaging tracked the experiment as it happened: a dye-drainage test measured how much of a labeled molecule moved into the lymphatic region every 30 minutes, and time-lapse video followed individually-dyed T-cells as they moved from the blood vessels into the lymphatic ones — though the paper describes this by how the cells' shape changed (from round to elongated) rather than giving a measured speed.

Source: Materials & Methods, "Immunostaining and imaging," "Vessel network architecture quantification," "T cell trafficking assay and quantification"

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

How can you predict what will happen as a disease progresses over time?

By treating the blood-vessel network with an inflammatory signaling protein (TNF-alpha) for 4 hours beforehand, to model sudden vessel inflammation. The paper's discussion attributes the resulting increase in T-cell movement at least partly to TNF-alpha's well-known effect of making blood- and lymphatic-vessel cells more "sticky" to immune cells, by increasing a docking protein (ICAM-1) on their surface. The measured effect is large: resting T-cells crossed into the lymphatic network 3.31 times more than in the untreated control, and already-activated T-cells crossed 5.89 times more than resting T-cells did.

Source: Results, Fig. 6H-I; Discussion, ICAM-1 paragraph

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

How can it be used in practice, and what are the probable next steps?

The paper's introduction points to diseases where blood- and lymphatic-vessel communication breaks down: cancer spread, chronic inflammation, and lymphedema. For next steps, the paper's own discussion names: adding a recirculating pump or more controlled pressure source (paired with computer flow simulations) to better define and control fluid flow; testing additional growth signals, blood-vessel-blocking drugs, or other supporting cell types; and extending the T-cell experiments to more types and states of immune cells, and to chronic (not just short-term) inflammation.

Source: Introduction; Discussion, final two paragraphs

2022: Lymphangion

BIOLOGICAL SYSTEM & CELL ORIGINS

Where did the cells come from (human vs. animal, cell lines vs. primary)?

Cells from two different species. The cells lining the inner channel are human lymphatic-vessel lining cells, taken from donor skin. The muscle cells wrapped around the outside are from rat intestinal lymphatic vessels — not human. The authors flag this themselves as a limitation, since human lymphatic muscle cells aren't commercially available, which makes the cell-to-cell signaling studied here a mix of two species, not two human cell types.

Source: Discussion, p.8: "LMCs and LECs in this project are derived from rat mesenteric vessels and human dermal tissue, respectively. This is a limitation because human LMCs are not commercially available."

BIOLOGICAL SYSTEM & CELL ORIGINS

What functional tissue systems are analyzed and reconstructed?

A single tube-shaped lymph vessel segment — specifically the portion between two one-way valves. It has an inner channel (averaging about 750 micrometers across in the dual-cell-type devices) lined by a complete layer of lymphatic-vessel cells, surrounded by a collagen gel sheath containing muscle cells (averaging about 150 micrometers thick).

Source: Results §2.2: "the average lumen inner diameter and muscle layer thickness are measured to be nearly 750 μm and 150 μm, respectively"; Fig. 1B, Fig. 4A-B

BIOLOGICAL SYSTEM & CELL ORIGINS

What is the physiological relevance of this model?

Recreates the lymphangion, the pumping unit that helps push lymph forward. The paper shows that growing the muscle cells together with the lining cells, combined with fluid shear, shifts the muscle cells to wrap more around the vessel's circumference (compared to mostly running lengthwise when grown alone), while the lining cells shift to align more lengthwise with the direction of flow — matching how these cells are arranged in the body. The paper shows this shift happens when both cell types are grown together under flow; it does not identify what specific chemical signal between the cells causes it, so claims naming a specific messenger molecule (like nitric oxide) cannot be grounded in fact in this paper.

Source: Results §2.3, Fig. 6A-D

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

What is the physical process and operation of the device?

A mold is used to form a square plastic channel (900 micrometers wide and tall, 5 millimeters long). A mixture of muscle cells and collagen gel is loaded into the channel; the whole device is then turned 90° so the channel runs straight up and down, and a syringe pulls a gentle vacuum at the bottom outlet. This technique (called gravitational lumen patterning, or GLP) uses gravity — now aligned with the direction the channel runs — to keep the hollowed-out center evenly positioned as it forms, giving a round channel with an evenly thick wall all the way around. This avoids the lopsided, sagging channel that forms if the same hollowing-out process is done with the channel lying flat. Lymphatic-vessel lining cells are then added to the hollow channel and grown under flow.

Source: Results §2.1, Fig. 2A-D; Methods §5.1

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

What hemodynamic forces and shear gradients play a role?

Three levels of fluid friction (shear) were tested: none (static fluid), a low level (about 0.1 dyne/cm^2, which the authors describe as representing a lymphedema-like state with poor drainage), and a level matching real rat lymphatic vessels (about 1 dyne/cm^2). When both cell types were grown together, both the lining cells' lengthwise alignment and the muscle cells' circumferential alignment became more pronounced as the shear increased from none to normal. A continuous range from 0.1 to 10 dyne/cm^2 was not tested and cannot be grounded in fact in this paper.

Source: Results §2.3, Fig. 6B-D

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

Do fluid acceleration, deceleration, angle, or orientation play a role?

The angle of the device during building, not flow speeding up or slowing down, is what matters here. Turning the channel 90° to vertical during the GLP process lines gravity up with the direction the channel runs. This prevents the lopsided, sagging channel produced when the same process is done with the channel flat, and instead gives a channel with nearly even wall thickness all the way around.

Source: Results §2.1, Fig. 2B-D

ENGINEERING CONTROL & SACRIFICES

What was sacrificed to engineer this, and how does that impact relevance?

The device models a single straight vessel segment — the part between two valves — without the valves themselves, and without pulsing flow or the muscle layer actively squeezing on its own (the paper notes that future versions may need to add this kind of flow pattern, and that an outside stimulus might be needed to trigger the muscle to contract on its own in the lab). Another limitation the paper states directly: the muscle cells come from rats, not humans, because human lymphatic muscle cells aren't commercially available — a cross-species gap in how directly these co-culture signaling results apply to human biology. Specific claims about sacrificing "bicuspid valves" or "afterload" capacity use terms not found in this paper and cannot be grounded in fact.

Source: Discussion, p.7-8

ENGINEERING CONTROL & SACRIFICES

How much independent control does the researcher have over variables?

Strong, independent control over the vessel's shape: the inner channel's width can be tuned from about 400 to 800 micrometers by changing the pressure applied during the GLP process; increasing how concentrated the collagen gel is reduces the channel width by about 30%, independent of pressure; and the outer size of the whole structure can be set by using different molds. Fluid shear is set directly by a pump's flow rate (none, ~0.1, or ~1 dyne/cm^2 were tested).

Source: Results §2.1, Fig. 3A-E

ENGINEERING CONTROL & SACRIFICES

How long will this work for continuously, and when does the system degrade beyond designed function?

The paper's own experiments run for up to 5 days: the lymphatic-vessel lining becomes a complete layer within 60 hours, and the gap between the two cell layers and how densely the cells pack is tracked through 96 hours (day 5). The paper does not report what happens to the device after day 5, and does not name a specific reason it would eventually fail (like the gel shrinking or the plastic channel coming apart) — a stated "60-120 hour degradation threshold" or named failure mode cannot be grounded in fact in this paper.

Source: Results §2.2, Fig. 5 (confluency and gap-distance time course, up to 96 hr / 5 days)

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

In what context was the data collected and interpreted (immunostaining and imaging)?

Fixed-endpoint microscope imaging of stained tissue, combined with live functional tests. Cell orientation was measured from images of the tissue stained for muscle- and lining-cell marker proteins, using free image-analysis software to measure each cell's angle relative to the vessel — 0° meaning lengthwise along the vessel, 90° meaning fully wrapped around it. The actual measured average angles land between these two extremes (for example, the muscle cells in co-culture averaged roughly 53-61° depending on the flow condition, not a literal 90°). How well the vessel wall blocked leakage was measured live by flowing a glowing dye of several different sizes through it and calculating how much leaked across the wall, before and after treatment.

Source: Methods §5 (Orientation)); Fig. 6B-D; Methods §5.7 (permeability)

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

How can you predict what will happen as a disease progresses over time?

Models two disease-relevant situations. First, reduced fluid shear (about 0.1 dyne/cm^2, which the authors describe as representing a lymphedema-like state with poor drainage) results in weaker cell alignment compared to normal shear. Second, the inflammatory signal TNF-alpha makes the vessel lining leakier whether or not muscle cells are present, but having the muscle cells present leads to a partial recovery of the barrier's leak-resistance after TNF-alpha exposure, compared to vessels with lining cells alone.

Source: Results §2.3 (lymphedema-like shear); Results §2.4, Fig. 7D (TNF-alpha permeability)

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

How can it be used in practice, and what are the probable next steps?

The authors describe it as an adjustable platform for early-stage lymphatic (and blood) vessel research and drug testing, usable alongside molecular and genetic tools to study how lymphatic cells signal each other; they state directly that it "can also be applied in blood vascular models." Specific future additions — like valves inside the vessel, a membrane that expands and contracts to mimic pumping, or linking several of these vessel segments in a row — are not discussed in the paper and cannot be grounded in fact.

Source: Conclusion §4

2024: L-Chip Sprouting

BIOLOGICAL SYSTEM & CELL ORIGINS

Where did the cells come from (human vs. animal, cell lines vs. primary)?

Human cells only, no animal tissue: lymphatic-vessel lining cells from donor skin, grown in the L-Chip device, compared directly against ordinary blood-vessel lining cells (from umbilical cord donors) grown in a matching "A-Chip" used to study blood-vessel growth instead. Before being added to the chip, cells were optionally pre-treated with a cocktail of five growth-signaling proteins known to encourage lymphatic-vessel growth, including the protein ESM-1.

Source: Methods, "Cells and reagents" and "Tissue culture" (HDLEC/HUVEC sourcing and pLF cocktail composition)

BIOLOGICAL SYSTEM & CELL ORIGINS

What functional tissue systems are analyzed and reconstructed?

A starting lymphatic vessel (modeling a 200-micrometer-wide lymphatic capillary) that sends out new sprouts into an adjacent 1.5-millimeter gel chamber, growing toward a source of growth-signaling chemicals, with connecting cells linking each new sprout back to the original vessel.

Source: Fig. 1A (device schematic, 1.5 mm chamber / 200 µm vessel dimensions); Fig. 1D-H (sprout and tube morphology)

BIOLOGICAL SYSTEM & CELL ORIGINS

What is the physiological relevance of this model?

Models lymphangiogenesis — the process by which new lymphatic vessels form. This matters for restoring normal fluid drainage, for how immune cells travel during inflammation and injury, and for how cancer cells spread into lymph nodes. The paper's core finding is that fluid flow through the gel and chemical growth signals work together to guide this process, rather than either one alone.

Source: Introduction (roles of lymphangiogenesis in homeostasis, inflammation, metastasis); Results, "L-Chips reveal that lymphangiogenesis is regulated by biochemical cues and interstitial fluid flow"

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

What is the physical process and operation of the device?

A mold forms a 3-channel plastic device, with the channels separated by rows of tiny pillars close enough together to hold a gel in place by surface tension (the paper doesn't state the exact pillar spacing). A gel is injected into the middle channel and left to set; vessel-lining cells are then added to the channel next to it. A chemical gradient (built from the growth-factor cocktail, or ESM-1 alone in later experiments) and/or a pressure difference driving fluid through the gel are applied, and the amount of new vessel growth is measured after a fixed 96-hour (4-day) period.

Source: Methods, "Microfluidic Device Fabrication" and "A- and L-Chip seeding, culturing, and stimulation"

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

What hemodynamic forces and shear gradients play a role?

Fluid seeping slowly through the gel, driven by a pressure difference, tested at a speed matching the body (1 micrometer/second) and a faster, higher-than-physiological speed (5 micrometers/second). This flow moves from the far channel toward the starting vessel, while the new sprouts grow in the opposite direction — against the flow — and faster flow increased how much the vessels sprouted and branched, compared to no flow at all. The paper does not explain what's happening at the cell level to cause this.

Source: Results, "L-Chips reveal that lymphangiogenesis is regulated by biochemical cues and interstitial fluid flow"; Fig. 2C, Fig. 3

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

Do fluid acceleration, deceleration, angle, or orientation play a role?

This cannot be grounded in fact in the paper. It does not discuss fluid speeding up, slowing down, or channel angle; the only relevant detail is that rows of tiny pillars physically separate the three channels, with no stated spacing or shape given for them.

Source: Not addressed in the paper

ENGINEERING CONTROL & SACRIFICES

What was sacrificed to engineer this, and how does that impact relevance?

The device leaves out other supporting cell types present in real tissue — muscle cells, connective-tissue cells, pericytes, and immune clean-up cells (macrophages) — which the authors themselves note could affect how new vessels grow. The paper doesn't discuss whether or how the new sprouts might shrink back after Day 4, since no experiment runs past that point.

Source: Conclusion ("Some limitations exist, including the lack of other supporting cells... that are present in a tissue microenvironment")

ENGINEERING CONTROL & SACRIFICES

How much independent control does the researcher have over variables?

Strong, independent control: the growth-factor pre-treatment (on or off, and at what dose and duration), the strength of the growth-factor gradient, and the strength of fluid flow through the gel can each be set separately from the others, letting the researchers test every combination.

Source: Fig. 2A-C (independent pLF/GFG/IF comparisons); Fig. 3A-G (combined GFG x IF matrix); Fig. 4D-E (pLF dose and duration sweeps)

ENGINEERING CONTROL & SACRIFICES

How long will this work for continuously, and when does the system degrade beyond designed function?

Every experiment in this paper uses a fixed 96-hour (4-day) culture and measurement endpoint; sprouts reach up to ~1 mm in that time. This cannot be grounded in fact in the paper: the study does not report what happens to sprouts beyond Day 4, and makes no claim about degradation, regression, or an operational limit past that point.

Source: Methods, "A- and L-Chip seeding, culturing, and stimulation" ("Chips were cultured for 96 hr before fixing")

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

In what context was the data collected and interpreted (immunostaining and imaging)?

At the end of the experiment, devices were stained for a protein found specifically on vessel-lining cells (PECAM-1, also called CD31) plus a general stain that lights up cell nuclei, then imaged in 3D layers under a microscope. The images were then run through image-analysis software that automatically measured sprout length, the number of branch junctions, and the number of branches, for statistical comparison between conditions.

Source: Methods, "Immunofluorescence" and "Image processing and data analysis"; Fig. 2 caption (quantified metrics)

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

How can you predict what will happen as a disease progresses over time?

The paper doesn't model how a disease changes over time — every data point is from the same fixed 4-day endpoint. What it does show is that the protein ESM-1 is a specific driver of new lymphatic-vessel growth: leaving ESM-1 out of the growth-factor cocktail reduced sprouting, while ESM-1 by itself was almost as effective as the full five-protein cocktail. This effect was specific to lymphatic vessels — ESM-1 alone did not drive extra growth in the ordinary blood-vessel (A-Chip) system.

Source: Fig. 5A-D; Conclusion ("identified ESM-1 as a prospective specific regulator of lymphangiogenesis, which was not the case... on our A-Chips")

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

How can it be used in practice, and what are the probable next steps?

The authors describe the L-Chip as a standardized platform for lymphatic-vessel-growth research, and a tool for finding drug targets related to either disease-driven or therapeutic vessel growth, with ESM-1 as the current focus. Stated next steps are a deeper look at exactly how ESM-1 works, and gradually adding the supporting cell types currently missing from the model (muscle cells, connective-tissue cells, pericytes, and macrophages).

Source: Conclusion

1937: Whole-Organ

BIOLOGICAL SYSTEM & CELL ORIGINS

Where did the cells come from (human vs. animal, cell lines vs. primary)?

Carrel used intact, non-human organs from etherized cats and rabbits — the primary animals used — and less frequently fowls (chickens). For cats and rabbits, the gland was removed together with a segment of the carotid artery and surrounding connective tissue; for fowls, it was removed with a segment of the aorta instead, since the thyroid in birds lies directly on the great thoracic vessels.

Source: p. 517-518: 'Cats and rabbits are generally used, less frequently, fowls... the gland is removed, together with a segment of carotid artery in cats and rabbits, and of aorta in fowls.'

BIOLOGICAL SYSTEM & CELL ORIGINS

What functional tissue systems are analyzed and reconstructed?

The whole intact thyroid gland — including its hormone-producing follicle cells, the jelly-like substance (colloid) those follicles store, and the small blood vessels feeding it — along with its surrounding connective-tissue capsule. The nerve running alongside the main artery was deliberately cut and removed during preparation, so this preparation explicitly does not preserve the gland's nerve connections.

Source: p. 517-518: vagus nerve 'severed in the lower part of the neck and separated from the artery'; pp. 515, 523-524: colloid and small-vessel/capillary descriptions.

BIOLOGICAL SYSTEM & CELL ORIGINS

What is the physiological relevance of this model?

This was a published demonstration that a whole mammalian (or bird) organ could be kept alive and fed with fluid, looking close to normal under the microscope, entirely outside the body — using only pulsing mechanical pressure and a nutrient fluid that contained no actual blood cells.

Source: Conclusions, p. 524-525: 'The gland remains alive during the period of perfusion, which lasted from 3 to 21 days.'

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

What is the physical process and operation of the device?

The Lindbergh apparatus is a sealed, sterilized, all-glass device. A glass tube is tied into the artery left attached when the gland is removed. The gland and its vessel are stretched flat on a small rectangular holder — a plate with slits to anchor the threads tying the artery in place, and teeth to hang the surrounding tissue on — which then slides into the glass chamber. An outside gas-pressure system, cycling between a set high and low pressure, pushes the fluid (diluted blood serum, or an artificial nutrient mix) through the vessels in pulses; there is no mechanical pump involved. The whole setup sits inside an incubator.

Source: pp. 517-521 ('Preparation of the Thyroid', 'Setting the Gland'); Text-fig. 1 (mica holder diagram, not a perforated glass disc as an earlier version of this answer claimed).

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

What hemodynamic forces and shear gradients play a role?

Pulsing pressure, at a specific setting the paper reports rather than a range: "the pulse rate has generally been 80," cycling between pressures of 160 and 110 (in the paper's own units — it doesn't convert these to a more familiar pressure unit like mmHg). This visibly expands and relaxes the main artery with each pulse, and the paper notes the pulsing "can also be detected in the thyroid artery" itself, pushing fluid into the gland's smallest vessels.

Source: p. 521: 'The pulse rate has generally been 80, the maximum pressure 160, and the minimum pressure 110.' p. 523: 'the carotid artery begins to pulsate. Pulsation can also be detected in the thyroid artery.'

BIOPHYSICAL DYNAMICS & FLUID MECHANICS

Do fluid acceleration, deceleration, angle, or orientation play a role?

Yes, to speeding up and slowing down: the gas-pressure system cycles between a set high and low pressure, which is what makes the artery "begin to pulsate" rather than receiving constant flow. This cannot be grounded in fact as a claim about a specific tilt angle, a gravity-driven vertical design, or avoiding mechanical blades that would damage blood cells — none of that is in the paper, and the fluid described here contains no actual blood cells to damage in the first place.

Source: p. 521, 523 (pulse rate/pressure cycling, artery pulsation); p. 524 ('the circulating fluid contains neither red blood corpuscles nor hemoglobin').

ENGINEERING CONTROL & SACRIFICES

What was sacrificed to engineer this, and how does that impact relevance?

There was no way to see the gland or its cells while the experiment ran — the chamber is sealed glass, so its condition could only be judged indirectly (by the fluid's clarity and color) during the run, and examined directly only afterward, once the gland was removed and prepared for viewing under a microscope. Small side-branches off the main artery were deliberately left open rather than tied off, and allowed to bleed into the surrounding tissue; the authors state directly that this resulting swelling "has no ill effect on the gland," so it was treated as an accepted part of the technique, not a problem that affected the results.

Source: p. 522: fluid examined for 'clearness or opacity'; p. 522: 'These collaterals bleed into the connective tissue, which becomes more or less edematous. Such edema has no ill effect on the gland.'

ENGINEERING CONTROL & SACRIFICES

How much independent control does the researcher have over variables?

Fine control over the gas mixture (generally 80% oxygen, 3-4% carbon dioxide, 16-17% nitrogen), the pulse rate, and the high/low pulsing pressures, all adjustable "according to the nature of the experiment" — plus what went into the perfusing fluid (how diluted the serum was, or whether Witte's peptone, a nutrient supplement, was added). Essentially no control at the level of individual small vessels or cells, which behaved however the gland's own existing anatomy dictated.

Source: p. 516 (gas mixture); p. 521 ('The pulse rate is adjusted, and also the maximum and minimum pressures, according to the nature of the experiment').

ENGINEERING CONTROL & SACRIFICES

How long will this work for continuously, and when does the system degrade beyond designed function?

Across the reported experiments, perfusion "lasted from 3 to 21 days," with 6 days being the duration used for most of the main histology comparison figures. The explicit stopping rule given is pH-based: "When the pH goes below 7, the experiment is interrupted." Fluid pH starts around 7.5 at the beginning of perfusion and falls progressively over about a week to roughly 7.0-7.1 or lower. The paper does not state a single "typical" duration — claiming "6 to 8 days typical" is not something it says; 6 days is simply how long most of the comparison figures happen to run.

Source: Conclusions, p. 524-525: 'lasted from 3 to 21 days.' p. 521 (pH stopping rule). p. 523-524 (pH decline from ~7.5 toward 7.0-7.1).

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

In what context was the data collected and interpreted (immunostaining and imaging)?

Purely after-the-fact examination, not live imaging: glands were chemically fixed, sliced thin, stained, and photographed at a fixed zoom level once removed from the pump. Two of the figure plates compare an untreated control thyroid from one side of an animal against the pump-treated thyroid from the other side of the same animal. Separately, small pieces of already-pumped glands were placed in a lab flask afterward, to see if their cells could still actively grow — shown for both a cat gland and a chicken gland. One inconsistency worth flagging: the paper's own body text and one of its figure captions disagree on how long the chicken sample was perfused — the text says 3 days, the caption says 8 days, with growth developing after 3 days in the flask.

Source: Explanation of Plates, p. 525-526 (Figs. 1-12); body text p. 524-525 vs. Plate 19 caption for Fig. 12 (discrepancy as noted).

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

How can you predict what will happen as a disease progresses over time?

Glands perfused with serum alone stayed close to their untreated control, though the paper notes even serum alone caused "a tendency to hyperplasia and to vacuolization of the colloid" (some extra cell growth and small gaps forming in the stored hormone substance). Adding Witte's peptone (a nutrient supplement) to the fluid produced a clearly stronger effect: noticeably larger cells and pronounced extra growth lining the follicles. The paper never uses the word "goiter" — that is a modern clinical label for this kind of enlargement, not a term the paper itself uses.

Source: p. 524: 'cultivated glands show a tendency to hyperplasia and to vacuolization of the colloid. When the fluid contained some Witte's peptone, considerable increase in the size of the cells was observed... Marked proliferation of the epithelial cells.'

ANALYTICAL CONTEXT, DISEASE & TRANSLATION

How can it be used in practice, and what are the probable next steps?

The paper's own stated conclusion is more modest than a claim about founding modern organ-preservation machines — that connection is a reasonable observation to make today about where this technique sits historically, but it is not something the 1937 paper claims about itself. What the paper actually concludes is that "the technique is simple enough to be used as a routine procedure for physiological or pathological studies of the thyroid gland" — the authors saw it as a practical lab tool for studying how the thyroid works and what goes wrong with it, such as the growth response to peptone shown here.

Source: Conclusions, p. 525, item 3.

9. Key Terms Primer

Lymphangion *lim-FAN-jee-on*

The pumping segment of a lymph vessel, located between two one-way valves.

A lymphangion squeezes rhythmically to push lymph forward against pressure — the way a line of hands passing a bucket keeps it moving in one direction. It is lined on the inside by a single layer of endothelial cells and wrapped on the outside by a layer of smooth muscle running around its circumference.

Paper 2 (Selahi et al.) · Physiology

Interstitial Fluid Flow (IF) *in-ter-STISH-ul flow*

The slow movement of fluid through the gel-like space between cells and vessels (the extracellular matrix).

This flow is driven by pressure differences — for example, fluid leaking out of blood capillaries gets pushed toward the lymphatic vessels that drain it. In Paper 3, this flow works together with chemical signals to encourage new lymphatic vessels to grow.

Paper 1 & Paper 3 · Fluid Mechanics

Gravitational Lumen Patterning (GLP) *GLP*

A fabrication technique where a microfluidic chip is turned vertical (90°) while a gel sets, so gravity pulls straight down along the flow direction and produces a round, evenly-shaped channel.

When a gel is left to set horizontally, gravity makes it sag unevenly, leaving an oval-shaped channel instead of a round one. Turning the chip vertical during setting avoids this sagging and produces a channel shaped more like a real blood or lymph vessel.

Paper 2 (Selahi et al.) · Bioengineering

Anastomosis *uh-nas-tuh-MOH-sis*

The joining of two separate vessel branches into one continuous, connected loop.

This term doesn't actually appear in Paper 3 for what it shows — the paper's own figure captions never say "anastomosis" or describe sprouts fusing tip-to-tip. What the paper does show is that chemical gradients and interstitial flow together cause new sprouts to grow toward the existing vessel by Day 4 — not that they are shown physically fusing into loops at that point.

Paper 1 & Paper 3 · Vascular Biology

Endocan (ESM-1) *EN-doh-kan*

A protein (endothelial cell-specific molecule-1) that Paper 3 identifies as a strong driver of new lymphatic-vessel growth.

Paper 3 found that ESM-1 alone was enough to trigger substantial new-vessel growth in human lymphatic endothelial cells, even without the broader mix of growth factors normally used.

Paper 3 (Tronolone et al.) · Molecular Biology

Extravasation / Transmigration *ex-tra-vuh-SAY-shun*

The process by which immune cells, like T-cells, squeeze through the junctions between blood-vessel wall cells to leave the bloodstream and enter the surrounding tissue.

In Paper 1, treating the blood vessel network with the inflammatory signal TNF-alpha made T-cells stick to the vessel wall more and increased how many crossed into the tissue. The paper reports two separate numbers: resting (naive) T-cells crossed over 3.31 times more than in the untreated control, and already-activated T-cells crossed over 5.89 times more than resting T-cells did.

Paper 1 (Gray-Scherr et al.) · Immunology

Hyperplasia *hy-per-PLAY-zhuh*

An increase in the number of cells in a tissue, which makes the tissue or organ grow larger.

Carrel found that thyroid glands kept alive outside the body in the Lindbergh pump, when fed a nutrient fluid containing Witte's peptone, grew new cells vigorously — evidence the gland was still fully alive and functioning, not just surviving.

Paper 4 (Carrel) · Pathology

Hydrostatic Pressure Head *hy-droh-STAT-ik*

The pressure created by a standing column of fluid due to gravity — the same reason pressure is higher at the bottom of a pool than at the top.

In Paper 1, filling one reservoir with 40 microliters of fluid and a connected one with only 10 microliters creates a small pressure difference (9.6 Pa) between them. That difference alone pushes fluid from the blood side to the lymph side, with no pump or motor needed.

Paper 1 & Paper 3 · Physics

Matrix Remodeling & Fibrinolysis *fy-brin-AHL-ih-sis*

The breakdown of the gel scaffolding (fibrin or collagen) that cells grow in, carried out by enzymes the cells themselves release.

This specific claim — that the gel breaks down on roughly a 5–7 day timescale — is not something either Paper 1 or Paper 3 actually reports. Both papers stop their experiments at a fixed point (Day 5 for Paper 1, Day 4 for Paper 3) and don't say why they stopped there or what would happen if they kept going. Enzymatic gel breakdown is a real process in biology generally, but neither paper measured it or used it to explain their results.

Paper 1 & Paper 3 · Biochemistry

Normothermic Machine Perfusion (NMP) *nor-moh-THUR-mik*

A modern technique for keeping an organ alive outside the body by pumping oxygenated, body-temperature fluid through its blood vessels.

Carrel's 1937 apparatus, built around the pump Charles Lindbergh engineered, is a reasonable historical ancestor of this modern technique, but the paper itself does not claim that connection — its own stated goal is more modest: a method simple enough for routine use in studying the thyroid gland. Modern versions of this technique add filters and artificial lungs (oxygenators) to prevent tissue damage and keep organs usable for longer.

Paper 4 (Carrel) & Frontier · Bioengineering