Frequently Asked Questions – Erectile Dysfunction Treatment

1. What is Erectile Dysfunction?

Erectile Dysfunction (ED) is the persistent or recurring difficulty in achieving or maintaining an erection sufficient for satisfactory sexual activity.

2. What causes Erectile Dysfunction?

ED may be associated with diabetes, high blood pressure, cardiovascular conditions, hormonal changes, stress, anxiety, smoking, or certain medications.

3. Can Erectile Dysfunction be treated?

Yes. Treatment depends on the underlying cause and may include lifestyle changes, counselling, management of medical conditions, and prescribed medication.

4. Can stress and anxiety cause Erectile Dysfunction?

Yes. Stress, performance anxiety, depression, and relationship difficulties can contribute to ED. Physical and psychological factors may also occur together.

5. Is Erectile Dysfunction common in young men?

ED can affect adults of different ages. Persistent erection difficulties in younger men should also be assessed rather than automatically attributed to psychological causes.

6. Are medicines for Erectile Dysfunction safe?

Some prescribed medicines can help, but they are not suitable for everyone. Medicines such as sildenafil and tadalafil can interact dangerously with nitrate medications. Consult a qualified doctor before using them.

7. When should I consult a doctor for Erectile Dysfunction?

Seek medical advice if erection difficulties occur repeatedly, cause distress, or affect your relationships. A consultation can help identify possible causes and appropriate treatment options.

8. How can I book an Erectile Dysfunction consultation at Kelkar Manas Health Care Clinic?

Contact Kelkar Manas Health Care Clinic using the details listed on our website to enquire about sexual health consultations and available treatment options.

Frequently Asked Questions – Premature Ejaculation Treatment

1. What is Premature Ejaculation?

Premature Ejaculation (PE) is a common sexual concern in which ejaculation occurs sooner than desired, often causing distress or difficulty with sexual satisfaction.

2. What causes Premature Ejaculation?

Possible contributing factors include performance anxiety, stress, relationship concerns, heightened sensitivity, and certain medical or psychological conditions.

3. Can Premature Ejaculation be treated?

Yes. Treatment may include behavioural techniques, counselling, prescribed medication, or a combination of approaches depending on the individual's needs.

4. How can I improve ejaculation control?

Behavioural approaches such as the start-stop technique may help some individuals. A healthcare professional can also assess contributing factors and recommend suitable treatment.

5. Is Premature Ejaculation a sign of sexual weakness?

No. PE is a recognised sexual health concern and does not determine a person's masculinity or overall health.

6. Can anxiety make Premature Ejaculation worse?

Yes. Worry about sexual performance can contribute to a cycle of anxiety and early ejaculation. Psychological support may be helpful when anxiety plays a role.

7. Do medicines help with Premature Ejaculation?

Certain prescription medicines may help selected individuals. Their suitability depends on medical history, other medications, side effects, and clinical assessment.

8. When should I seek treatment for Premature Ejaculation?

Consider a consultation if the problem happens repeatedly, causes distress, or affects your sexual relationship.

9. How can I book a consultation at Kelkar Manas Health Care Clinic?

Contact Kelkar Manas Health Care Clinic to enquire about sexual health consultations and appropriate treatment options for premature ejaculation.

Frequently Asked Questions – Sexual Performance Anxiety

1. What is Sexual Performance Anxiety?

Sexual performance anxiety is fear or worry about sexual performance, satisfying a partner, maintaining an erection, or reaching orgasm. It can interfere with sexual enjoyment and confidence.

2. What are the symptoms of Performance Anxiety?

Symptoms may include nervousness before intimacy, excessive worry, difficulty maintaining an erection, reduced sexual desire, rapid ejaculation, or difficulty reaching orgasm.

3. What causes Sexual Performance Anxiety?

It may be associated with fear of failure, previous negative experiences, relationship concerns, stress, low confidence, or unrealistic expectations about sexual performance.

4. Can Performance Anxiety cause Erectile Dysfunction?

Yes. Anxiety can interfere with the physical and emotional processes involved in sexual arousal and contribute to erection difficulties.

5. How can Sexual Performance Anxiety be treated?

Treatment may include psychological counselling, cognitive behavioural therapy (CBT), relaxation techniques, communication strategies, and treatment of any underlying sexual health condition.

6. Can Performance Anxiety be overcome without medication?

Yes. Some individuals benefit from counselling, behavioural strategies, and reducing performance pressure without medication. Treatment depends on the underlying cause.

7. Can relationship problems contribute to Performance Anxiety?

Yes. Communication difficulties, unresolved conflict, and concerns about intimacy can contribute to anxiety. Couple counselling may help when appropriate.

8. When should I seek professional help?

Consider professional support when anxiety repeatedly interferes with intimacy, sexual satisfaction, confidence, or relationships.

9. How can I book a consultation at Kelkar Manas Health Care Clinic?

Contact Kelkar Manas Health Care Clinic through our website to enquire about sexual health counselling and consultation options.

Frequently Asked Questions – Sexually Transmitted Infections (STIs)

1. What are Sexually Transmitted Infections (STIs)?

STIs are infections that can be transmitted through sexual contact. Examples include chlamydia, gonorrhoea, syphilis, genital herpes, human papillomavirus (HPV), and HIV.

2. What are the common symptoms of STIs?

Symptoms may include unusual genital discharge, painful urination, sores, blisters, itching, rashes, or pelvic discomfort. However, many STIs cause no noticeable symptoms.

3. Can STIs occur without symptoms?

Yes. A person can have and transmit certain STIs without experiencing symptoms. Testing may be recommended depending on sexual exposure and individual risk.

4. How are STIs diagnosed?

Diagnosis may involve urine tests, blood tests, swabs, or physical examination, depending on the suspected infection and clinical circumstances.

5. Can Sexually Transmitted Infections be treated?

Many bacterial and parasitic STIs can be cured with appropriate treatment. Viral infections may be managed with medication, although some cannot currently be completely cured.

6. How can I reduce my risk of STIs?

Using condoms correctly, getting appropriate vaccinations, undergoing recommended testing, and discussing sexual health with partners can reduce risk. Condoms do not eliminate every STI risk.

7. Should my partner also be tested?

Depending on the infection and exposure, partner testing and treatment may be important to prevent reinfection and further transmission. Follow medical advice about sexual activity during treatment.

8. When should I seek medical advice for a possible STI?

Seek assessment if you have symptoms, a partner has been diagnosed with an STI, or you believe you may have been exposed. Do not rely on symptoms alone to determine infection status.

9. Is STI consultation confidential?

Sexual health consultations should be conducted respectfully and with appropriate protection of patient privacy, subject to applicable legal and professional requirements.

10. How can I enquire about STI testing or treatment at Kelkar Manas Health Care Clinic?

Contact Kelkar Manas Health Care Clinic to ask about available sexual health assessments, testing arrangements, and appropriate referral options.

Frequently Asked Questions – Precum Concerns

1. What is Precum?

Precum, medically known as pre-ejaculate, is a clear fluid that may be released from the penis during sexual arousal before ejaculation.

2. Is Precum normal?

Yes. Pre-ejaculate is a normal bodily fluid. The amount varies between individuals and may vary from one occasion to another.

3. Can Precum cause pregnancy?

Yes. Pregnancy is possible if pre-ejaculate contains sperm or carries sperm remaining in the urethra into the vagina. The withdrawal method is not completely reliable for preventing pregnancy.

4. Can STIs spread through Precum?

Certain sexually transmitted infections can spread through sexual fluids or direct genital contact. Using condoms correctly can reduce the risk of many STIs.

5. Is excessive Precum a sign of a medical problem?

Not necessarily. The amount of pre-ejaculate varies naturally. However, persistent unusual discharge, pain, burning during urination, or blood should be medically evaluated.

6. Can I reduce the amount of Precum?

Pre-ejaculate is a normal physiological response, and there is generally no need to reduce it. Avoid taking medicines or supplements to change fluid production without medical advice.

7. Is Precum the same as semen?

No. Pre-ejaculate is released before ejaculation and differs from semen, which is released during ejaculation. They have different functions.

8. When should I consult a doctor about Precum?

Seek medical advice if the fluid has an unusual colour or smell, occurs without arousal and persists, or is accompanied by pain, itching, sores, or urinary symptoms.

9. How can I enquire about sexual health concerns at Kelkar Manas Health Care Clinic?

Contact Kelkar Manas Health Care Clinic to enquire about sexual health consultations and assessment of any symptoms that concern you.

Frequently Asked Questions – Nocturnal Emissions

1. What are Nocturnal Emissions?

Nocturnal emissions, commonly called wet dreams, are involuntary ejaculations that occur during sleep. They are a normal physiological experience for many people.

2. Are Nocturnal Emissions normal?

Yes. Wet dreams are common, particularly during adolescence and young adulthood, although they can occur at other ages as well.

3. What causes Nocturnal Emissions?

They may occur naturally during sleep and are associated with normal reproductive physiology. They do not necessarily indicate a sexual or medical problem.

4. Do Nocturnal Emissions cause weakness?

No. Normal nocturnal emissions do not cause lasting physical weakness, infertility, or a loss of masculinity.

5. How often are Nocturnal Emissions normal?

There is no single normal frequency. Some people experience them frequently, others occasionally, and some never experience them.

6. Can Nocturnal Emissions be prevented?

They are a normal bodily process and generally do not require prevention or treatment. Be cautious of products claiming to cure wet dreams without reliable medical evidence.

7. When should I consult a doctor about Nocturnal Emissions?

Seek medical advice if episodes are accompanied by pain, blood in the semen, urinary symptoms, or significant distress, or if you have other concerns about your sexual health.

8. How can I book a consultation at Kelkar Manas Health Care Clinic?

Contact Kelkar Manas Health Care Clinic to enquire about sexual health consultations and professional guidance.

Frequently Asked Questions – Masturbation Habits

1. Is Masturbation normal?

Yes. Masturbation is a common sexual behaviour and is generally considered a normal part of human sexuality.

2. Does Masturbation cause physical weakness?

There is no good evidence that masturbation by itself causes lasting physical weakness, infertility, or a permanent reduction in testosterone.

3. How often is Masturbation considered normal?

There is no fixed normal frequency. It becomes a concern mainly when it causes physical injury, significant distress, or interference with work, relationships, sleep, or other daily activities.

4. Can Masturbation become difficult to control?

Some people may feel unable to control repetitive sexual behaviour, particularly when it interferes with daily responsibilities or continues despite negative consequences. Professional support may help.

5. Can Masturbation affect sexual performance?

Masturbation does not generally cause sexual dysfunction. However, individual habits, anxiety, unrealistic expectations, or specific patterns of stimulation may contribute to concerns for some people.

6. Can I reduce or manage a Masturbation Habit?

If the behaviour is interfering with your life, identifying triggers, establishing healthy routines, reducing exposure to triggering situations, and seeking counselling may help.

7. Does Masturbation cause Erectile Dysfunction or infertility?

Masturbation itself is not generally considered a cause of erectile dysfunction or infertility. Persistent erection or fertility concerns should be assessed by a qualified healthcare professional.

8. When should I seek professional help for Masturbation concerns?

Consider counselling if you experience persistent guilt or distress, repeated unsuccessful attempts to change the behaviour, physical injury, or significant interference with daily life.

9. How can I enquire about counselling at Kelkar Manas Health Care Clinic?

Contact Kelkar Manas Health Care Clinic to enquire about confidential sexual health consultations and counselling options.

Frequently Asked Questions – Sexual Orientation and Counselling

1. What is Homosexuality?

Homosexuality refers to romantic, emotional, or sexual attraction to people of the same sex. Sexual orientation is a natural aspect of human diversity.

2. Is Homosexuality a mental illness?

No. Homosexuality is not a mental disorder and does not require treatment or a cure.

3. What is Sexual Orientation Counselling?

Sexual orientation counselling provides a supportive, confidential space to explore feelings, identity, relationships, family concerns, or social pressures without judgement.

4. Can counselling change a person's sexual orientation?

Sexual orientation is not a disorder that needs to be changed. Attempts to force or pressure someone to change their orientation can be harmful. Ethical counselling focuses on wellbeing, self-understanding, and personal choice.

5. Can I seek counselling if I am confused about my sexual orientation?

Yes. You can seek support to explore your feelings and questions at your own pace. You do not need to adopt a particular label or make an immediate decision about your identity.

6. Can counselling help with family or relationship concerns?

Yes. Supportive counselling may help individuals navigate family reactions, relationship challenges, social stigma, stress, or concerns about disclosure.

7. Is sexual orientation counselling confidential?

Counselling should respect privacy and confidentiality, subject to applicable legal and professional requirements. You can ask the counsellor about confidentiality before beginning sessions.

8. Do I need counselling because I am homosexual?

No. Being homosexual does not mean you need mental health treatment. Counselling is an option if you want support with stress, relationships, identity exploration, or other personal concerns.

9. How can I enquire about supportive counselling at Kelkar Manas Health Care Clinic?

Contact Kelkar Manas Health Care Clinic to ask about available counselling services and whether a suitable, non-judgmental service is available for your concerns.